# On the Fredholm determinant of the confluent hypergeometric kernel with discontinuities

Shuai-Xia Xu<sup>1</sup>, Shu-Quan Zhao<sup>2,\*</sup>, and Yu-Qiu Zhao<sup>2</sup>

<sup>1</sup>Institut Franco-Chinois de l'Energie Nucléaire, Sun Yat-sen University, Guangzhou 510275, China <sup>2</sup>Department of Mathematics, Sun Yat-sen University, Guangzhou 510275, China \*E-mail address: zhaoshq25@mail2.sysu.edu.cn

**Abstract** We consider the determinantal point process with the confluent hypergeometric kernel. This process is a universal point process in random matrix theory and describes the distribution of eigenvalues of large random Hermitian matrices near the Fisher-Hartwig singularity. Applying the Riemann-Hilbert method, we study the generating function of this process on any given number of intervals. It can be expressed as the Fredholm determinant of the confluent hypergeometric kernel with n discontinuities. In this paper, we derive an integral representation for the determinant by using the Hamiltonian of the coupled Painlevé V system. By evaluating the total integral of the Hamiltonian, we obtain the asymptotics of the determinant as the n discontinuities tend to infinity up to and including the constant term. Here the constant term is expressed in terms of the Barnes G-function.

**Keywords** Confluent hypergeometric kernel; Coupled Painlevé V system; Asymptotic expansion; Fredholm determinant; Riemann-Hilbert approach.

# Contents

1	Introduction	<b>2</b>
	1.1 Main results	5
	1.2 Applications	7
<b>2</b>	Model RH problem and the coupled Painlevé V system	9
	2.1 RH problem for the Fredholm determinant	9
	2.2 Model RH problem	10
	2.3 Lax pair and the coupled Painlevé V system	12
	2.4 Differential identity	14
3	Asymptotic Analysis as $t \to 0^+$	15
	3.1 Local parametrix	15
	3.2 Ratio RH problem	16
	3.3 Asymptotic behavior as $t \to 0^+$	17

<b>4</b>	$\textbf{Asymptotic Analysis as } t \to +\infty$	19
	4.1 Normalization	19
	4.2 Opening lenses and deformation	20
	4.3 Global parametrix	21
	4.4 Local parametrices	22
	4.5 Final transformation	23
	4.6 Asymptotic behaviors as $t \to +\infty$	25
<b>5</b>	Proof of main theorems	26
	5.1 Proof of Theorem 1	26
	5.2 Proof of Theorem 2	27
$\mathbf{A}$	The confluent hypergeometric parametrix	29

### 1 Introduction

In random matrix theory, one of the most important problems is to study the local statistics of eigenvalues for large random matrices. Compared with the global perspective in which the limiting average densities of eigenvalues are diverse, the local statistics of eigenvalues for large random matrices display remarkable universality. For instance, for general large unitary invariant random matrices, the local eigenvalue distributions near a regular point in the bulk, near the soft edge and the hard edge can be described by the determinantal point processes associated with the sine kernel, Airy kernel and Bessel kernel, respectively; see [1, 34, 47]. The gap probabilities in these processes can then be expressed in terms of the Fredholm determinants with the kernels associated. It is remarkable that the Fredholm determinants admit integral representations in terms of the Painlevé transcendents; [1, 34, 35, 42, 49, 50].

A different type of local statistics will arise if we consider the unitary ensemble with Fisher-Hartwig singularity as follows [28, 37, 43]:

$$p_n(x_1, \dots, x_n) = \frac{1}{Z_n} \prod_{i=1}^n e^{-V(x_i)} |x_i|^{2\alpha} \chi_\beta(x_i) \prod_{i < j} (x_i - x_j)^2, \qquad (1.1)$$

where  $Z_n$  is the normalization constant, the potential V is a general real valued function with enough increase at infinity. The parameters satisfy  $\alpha > -\frac{1}{2}$ ,  $\beta \in i\mathbb{R}$ , and

$$\chi_{\beta}(x) = \begin{cases} e^{\beta \pi i}, & x < 0, \\ e^{-\beta \pi i}, & x > 0. \end{cases}$$

The root-type singularity of the form  $|x - a|^{2\alpha}$  and the jump-type singularity such as  $\chi_{\beta}(x - a)$  are known as Fisher-Hartwig (FH) singularities in the literature. The asymptotics of the large Toeplitz determinants with these types of singularities were first investigated and some formulas on the asymptotics were conjectured by Fisher and Hartwig in [33] and subsequently in [45] by Lenard, motivated by their applications in Ising models, random walks and one-dimensional Bose gases.

When the limiting average density function of the ensemble (1.1) is positive at the origin, the scaling limit of the correlation function near the origin is now described by the following confluent hypergeometric kernel

$$K^{(\alpha,\beta)}(x,y) = \frac{1}{2\pi i} \frac{\Gamma(1+\alpha+\beta)\Gamma(1+\alpha-\beta)}{\Gamma(1+2\alpha)^2} \frac{\mathbb{A}(x)\mathbb{B}(y) - \mathbb{A}(y)\mathbb{B}(x)}{x-y},$$
(1.2)

where

$$\mathbb{A}(x) = \chi_{\beta}(x)^{1/2} |2x|^{\alpha} e^{-ix} \phi(1 + \alpha + \beta, 1 + 2\alpha, 2ix), \tag{1.3}$$

and  $\mathbb{B}(x)$  is the complex conjugation of  $\mathbb{A}(x)$ . Here  $\phi(a, b, z)$  denotes the confluent hypergeometric function

$$\phi(a,b,z) = 1 + \sum_{k=1}^{\infty} \frac{a(a+1)\dots(a+k-1)z^k}{b(b+1)\dots(b+k-1)k!}.$$
(1.4)

For special parameters, the confluent hypergeometric kernel reduces to the sine kernel and Bessel kernel:

$$K^{(0,0)}(x,y) = K_{\sin}(x,y) = \frac{\sin(x-y)}{\pi(x-y)},$$
(1.5)

$$K^{(\alpha,0)}(x,y) = K_{\text{Bess},1}(x,y) = \frac{|x|^{\alpha}|y|^{\alpha}}{x^{\alpha}y^{\alpha}} \frac{\sqrt{xy}}{2} \frac{J_{\alpha+\frac{1}{2}}(x)J_{\alpha-\frac{1}{2}}(y) - J_{\alpha-\frac{1}{2}}(x)J_{\alpha+\frac{1}{2}}(y)}{x-y}.$$
 (1.6)

Analogously to the sine, Airy and Bessel point processes, the determinantal point process defined by the confluent hypergeometric kernel represents another universality class in random matrix theory. It describes the local statistics of the eigenvalues of general random matrices near the Fisher-Hartwig singularity as shown in (1.1). This process arises also in the studies of the representation theory of the infinite-dimensional unitary group [4, 5].

In the present work, we consider the Fredholm determinant of the confluent hypergeometric kernel (1.2) with discontinuities

$$F(t\vec{r},\vec{\gamma};\alpha,\beta) = \det\left(I - K_{\sigma}^{(\alpha,\beta)}\right),\tag{1.7}$$

where t > 0, and  $K_{\sigma}^{(\alpha,\beta)}$  denotes the integral operator acting on  $L^2(tr_0, tr_n)$  with the confluent hypergeometric kernel with discontinuities  $\sigma(y)K^{(\alpha,\beta)}(x,y)$ . Here the function  $\sigma(x)$  is a step function defined by

$$\sigma(x) = \sum_{k=0}^{n-1} \gamma_k \chi_{(r_k t, r_{k+1} t)}(x).$$
(1.8)

The parameters  $\vec{\gamma} = (\gamma_0, \dots, \gamma_{n-1}) \in [0, 1]^n$ ,  $\vec{r} = (r_0, r_1, \dots, r_n) \in \mathbb{R}^{n+1}$  satisfy  $r_0 < r_1 < \dots < r_n$  and  $r_m = 0$  for some  $0 \le m \le n$ . The determinant represents the moment generating function of the confluent hypergeometric process. To see this, we denote by N(s) the random variable counting the number of particles in this process falling into the interval [0, s] for s > 0 and [s, 0] for s < 0. The determinant (1.7) can be written as the moment generating function

$$F(t\vec{r},\vec{\gamma};\alpha,\beta) = \mathbb{E}\left(\prod_{j=0}^{n} e^{2\pi i h_j N(tr_j)}\right),\tag{1.9}$$

where  $h_j$  is related to the parameters  $\vec{\gamma}$  by  $h_j = \frac{1}{2\pi i} \ln \frac{1-\gamma_{j-1}}{1-\gamma_j}$  for  $m < j \leq n$  and  $h_j = \frac{1}{2\pi i} \ln \frac{1-\gamma_j}{1-\gamma_{j-1}}$  for  $0 \leq j \leq m$  with  $\gamma_{-1} = \gamma_n = 0$ . The Fredholm determinant (1.7), or equivalently the moment generating function (1.9),

The Fredholm determinant (1.7), or equivalently the moment generating function (1.9), contains information about the statistics of the confluent hypergeometric process. For example, by taking n = 2 in (1.9) we have

$$F((r_0t, r_1t, r_2t), \gamma(1, 1); \alpha, \beta) = \det\left(1 - \gamma \left. K^{(\alpha, \beta)} \right|_{(r_0t, r_2t)} \right), \tag{1.10}$$

where  $K^{(\alpha,\beta)}|_{(r_0t,r_2t)}$  denotes the integral operator acting on  $L^2(r_0t, r_2t)$  with the confluent hypergeometric kernel  $K^{(\alpha,\beta)}(x,y)$ . Therefore, the quantity  $F((r_0t, r_1t, r_2t), (1, 1); \alpha, \beta)$  gives the gap probability of finding no particles of the confluent hypergeometric process on the interval  $(r_0t, r_2t)$ . While  $F((r_0t, r_1t, r_2t), \gamma(1, 1); \alpha, \beta)$  is the gap probability of having no particles falling in the interval  $(r_0t, r_2t)$  in the thinned confluent hypergeometric process, where each of the particles in the process is removed independently with the probability  $1 - \gamma \in [0, 1)$  [3]. Generally,  $F((r_0t, 0, r_2t), (\gamma_1, \gamma_2); \alpha, \beta)$  with  $r_0 < 0$  and  $r_2 > 0$ , describes the probability to find no particles in the interval  $(r_0t, r_2t)$  in the thinned confluent hypergeometric process, where the positive particles in the process are removed independently with the probability  $1 - \gamma_2 \in [0, 1)$  and the negative particles in the process are removed independently with the probability  $1 - \gamma_2 \in [0, 1)$  and the

In [30], the large gap asymptotics of the confluent hypergeometric process, which is given by  $F((-t, 0, t), (1, 1); \alpha, \beta)$ , was derived by Deift, Krasovsky, and Vasilevska by considering the asymptotics of the Toeplitz determinant generated by certain symbols with Fisher-Hartwig singularity on the unit circle. Later in [55], the first and third authors of the present paper established an integral representation of the Fredholm determinant  $F((-t, 0, t), (1, 1); \alpha, \beta)$  in terms of the Hamiltonian of a coupled Painlevé V system by considering the double scaling limit of the Toeplitz determinant with merging singularities. By evaluating the total integral of the Hamiltonian, the large gap asymptotics was then derived therein. Recently, the large gap probability asymptotics of the deformed Fredholm determinant  $F((-t, 0, t), (\gamma, \gamma); \alpha, \beta)$  was derived by Dai and Zhai in [25] by evaluating again the total integral of the Hamiltonian of the coupled Painlevé V system. It is also worth mentioning that the integral representations of the gap probabilities and their asymptotics of some other determinantal processes arising in random matrix theory have been considered, such as the Pearcey process and the hard edge Pearcey process in [17, 22, 23, 24], the higher order Airy processes [8, 46, 53] and the process defined by the Painlevé II kernel [7, 54]. The integral representations and the asymptotics of the moment generating functions for the sine, Airy and Bessel point processes are considered in [10, 11, 12, 14, 18].

The studies of the asymptotics of the Fredholm determinants with jump-type FH singularities similar to (1.7) are closely related to the problem of deriving the asymptotics of the Toeplitz and Hankel determinants with FH singularities. The problem has a long history tracing back to the pioneering work of Fisher and Hartwig [33], and has been widely explored due partially to their various important applications of which we have mentioned a few in the previous context. For example, the conjectures proposed in [33] was proved later by Widom in [51]. The asymptotics of the Toeplitz determinants with several separate FH singularities have been obtained by Deift, Its and Krasovsky in [28, 29]. Recently, the asymptotics of the Toeplitz determinants with merging singularities have been derived in [20, 21, 32]. There are also important progresses in the studies of the asymptotics of the Hankel determinants with FH singularities [44, 39, 2, 19, 13, 52, 9, 16, 15].

In the present work, we formulate a Riemann-Hilbert (RH, for short) representation for the determinant (1.7) of the confluent hypergeometric kernel (1.2) with n discontinuities by using the general theory of integrable operator developed in [38]. By exploring the RH problem, we obtain an integral representation for the determinant (1.7) in terms of the Hamiltonian of the coupled Painlevé V system of 2n dimensions. The large gap asymptotics of the determinant up to and including the constant term are then derived by evaluating the integral of the Hamiltonian.

#### 1.1 Main results

To state our main results, it is convenient to introduce the couple Painlevé V system defined by the following Hamiltonian dynamic system

$$\frac{dv_k}{dt} = \frac{\partial H}{\partial u_k}, \quad \frac{du_k}{dt} = -\frac{\partial H}{\partial v_k}, \quad k = 0, 1, \dots, n, \quad k \neq m$$
(1.11)

for any specific m and  $0 \le m \le n$ , where the Hamiltonian H is defined by

$$tH(t) = \sum_{k \neq m} -2itr_k H_V(u_k, v_k, -2itr_k; \alpha, \beta) + \frac{1}{2} \sum_{j \neq k; \ j, k \neq m} u_j u_k(v_j + v_k)(v_j - 1)(v_k - 1), \ (1.12)$$

with  $r_k \neq 0$  for  $k \neq m$ . Here  $H_V$  denotes the Hamiltonian for the Painlevé V equation defined in [36, 41]

$$sH_V(u, v, s; \alpha, \beta) = -suv - \alpha u \left(v^2 - 1\right) - \beta u (v - 1)^2 + u^2 v (v - 1)^2.$$
(1.13)

From (1.11)-(1.13), the coupled Painlevé V system can be expressed explicitly as follows

$$\begin{cases} t \frac{du_k}{dt} = -2itu_k r_k - u_k \sum_{j \neq m} u_j (v_j - 1)^2 - 2u_k v_k \left(\sum_{j \neq m} u_j (v_j - 1)\right) \\ + 2(\alpha + \beta)u_k v_k - 2\beta u_k, \\ t \frac{dv_k}{dt} = 2itv_k r_k + v_k \left(\sum_{j \neq m} u_j (v_j - 1)^2\right) + v_k^2 \left(\sum_{j \neq m} u_j (v_j - 1)\right) \\ - \sum_{j \neq m} u_j v_j (v_j - 1) - \alpha \left(v_k^2 - 1\right) - \beta \left(v_k - 1\right)^2. \end{cases}$$
(1.14)

When n = 2, the above system of equations reduce to the one used in [25, 55].

As the first result, we obtain an integral representation of the determinant (1.7) by using the Hamiltonian associated with a family of solutions of the coupled Painlevé V system (1.14).

**THEOREM 1.** Let  $\alpha > -\frac{1}{2}$ ,  $\beta \in i\mathbb{R}$ ,  $n \ge 1$ ,  $\vec{\gamma} = (\gamma_0, \gamma_1, \dots, \gamma_{n-1}) \in [0, 1)^n$  and  $\vec{r} = (r_0, r_1, \dots, r_n) \in \mathbb{R}^{n+1}$  be such that  $r_0 < \dots < r_m = 0 < \dots < r_n$ , we have the following integral expression of the Fredholm determinant defined in (1.7)

$$\ln F(t\vec{r},\vec{\gamma};\alpha,\beta) = \int_0^t H(s)ds, \qquad (1.15)$$

where the Hamiltonian H is defined by (1.11) and (1.12) subject to the following asymptotics: as  $t \to 0^+$ 

$$H(t) = i \sum_{k=0}^{n} \frac{c_k \cdot \Gamma(1+\alpha-\beta)\Gamma(1+\alpha+\beta) (2|r_j|)^{2\alpha+1}}{(2\alpha+1)\Gamma^2(1+2\alpha)} t^{2\alpha} + \mathcal{O}\left(t^{2\alpha+1}\right), \qquad (1.16)$$

with

$$c_{k} = \begin{cases} \frac{\gamma_{k-1} - \gamma_{k}}{2\pi i} e^{\beta \pi i}, & 0 \le k < m, \\ \frac{\gamma_{k} - \gamma_{k-1}}{2\pi i} e^{-\beta \pi i}, & m < k \le n, \end{cases}$$
(1.17)

and as  $t \to +\infty$ 

$$H(t) = \sum_{k=0}^{n} 2ib_k r_k - \left(\sum_{k=0}^{n} b_k^2 + 2\beta b_m\right) t^{-1} + \mathcal{O}\left(t^{-2}\right), \qquad (1.18)$$

with

$$b_k = \frac{1}{2\pi i} \ln \frac{1 - \gamma_{k-1}}{1 - \gamma_k},\tag{1.19}$$

where  $\gamma_{-1} = \gamma_n = 0$ .

**REMARK 1.** For n = 2, with the special parameter  $r_0 = -1, r_1 = 0, r_2 = 1$  and  $\gamma_0 = \gamma_1 = \gamma$ , (1.8) is reduced into  $\sigma(x) = \gamma_0 \chi_{(-t,0)}(x) + \gamma_1 \chi_{(0,t)}(x) = \gamma \chi_{(-t,t)}(x)$ . The equations (1.15) and (1.18) recover the integral representation of the determinant  $F((-t, 0, t), (\gamma, \gamma); \alpha, \beta)$ , which was derived by Dai and Zhai in [25]. While the determinant  $F((-t, 0, t), (1, 1); \alpha, \beta)$  was expressed in terms of another solution of the coupled Pailevé V system characterized by different asymptotic behavior as  $t \to +\infty$ , as shown by the first and third authors of the paper in [55]. For general n, the integral representation (1.15) is also true when some of the parameters  $\gamma_j = 1$ , but the asymptotic behavior would be quite different from (1.18). We will consider this problem in a separate paper.

**THEOREM 2.** Under the same assumptions as in Theorem 1, we have the asymptotic approximation as  $t \to +\infty$ 

$$\ln F(t\vec{r},\vec{\gamma};\alpha,\beta) = \sum_{k\neq m} 2ib_k r_k t + \sum_{k\neq m} \left(2\beta b_k - 2b_k^2\right) \ln |2r_k t| - \sum_{0\leq j
(1.20)$$

where  $b_k$  is defined by (1.19) and G is the Barnes G-function. The asymptotics (1.20) are uniform for  $\gamma_k$  in compact subsets of [0,1) and uniform for  $r_k$  in compact subsets of  $\mathbb{R}$ , as long as there exists  $\epsilon > 0$  such that  $\min_{0 \le j < k \le n} (r_k - r_j) > \epsilon$ .

Furthermore, (1.20) can be differentiated any k times with respect to  $\gamma_j$  in the form  $\prod_{j=0}^{n-1} \partial_{\gamma_j}^{k_j}$ , where  $k_0, \ldots, k_{n-1} \in \mathbb{N} \cup \{0\}$  and  $k = \sum_{j=0}^{n-1} k_j$ , with the error term replaced by  $\mathcal{O}\left(\frac{(\ln t)^k}{t}\right)$ .

For n = 2 with the special parameter  $r_0 = -1, r_1 = 0, r_2 = 1$  and  $\gamma_0 = \gamma_1 = \gamma \in [0, 1)$ , we recover the asymptotics of the determinant  $F((-t, 0, t), (\gamma, \gamma); \alpha, \beta)$  in [25].

**COROLLARY 1.** (Dai and Zhai [25]) For  $\alpha > -\frac{1}{2}$ ,  $\beta \in i\mathbb{R}$  and  $\gamma \in [0,1)$ , let  $K^{(\alpha,\beta)}|_{(-t,t)}$  be the integral operator acting on  $L^2(-t,t)$  with the confluent hypergeometric kernel defined in (1.2). Then, as  $t \to +\infty$ , we have

$$\ln \det \left( 1 - \gamma \left. K^{(\alpha,\beta)} \right|_{(-t,t)} \right) = -4ct + 2c^2 \ln(4t) + 2\alpha\pi c + 2\ln \left[ G(1+ic)G(1-ic) \right] + \mathcal{O}\left(\frac{1}{t}\right),$$
(1.21)

where  $c = -\frac{1}{2\pi} \ln(1 - \gamma)$ .

While for the special parameter  $\alpha = \beta = 0$ , we recover the asymptotics of the determinant of the sine kernel with several discontinuities in [11].

**COROLLARY 2.** (Charlier [11]) Let  $K_{\sigma}^{(0,0)}$  denotes the integral operator acting on  $L^2(tr_0, tr_n)$  with the sine kernel multiplied by the function  $\sigma$  given in (1.8). Then, as  $t \to +\infty$ , we have

$$\ln \det(I - K_{\sigma}^{(0,0)}) = \sum_{k \neq m} 2ib_k r_k t - \sum_{k \neq m} 2b_k^2 \ln |2r_k t| - \sum_{0 \le j < k \le n; \ j,k \neq m} 2b_j b_k \ln \left| \frac{2r_j r_k t}{r_k - r_j} \right| + \sum_{k=0}^n \ln \left[ G\left(1 + b_k\right) G\left(1 - b_k\right) \right] + \mathcal{O}\left(\frac{1}{t}\right).$$
(1.22)

#### 1.2 Applications

Consider the counting function N(t) of the number of particles in the confluent hypergeometric process falling into the interval [0, t] for t > 0 and [t, 0] for t < 0. As shown in (1.9), the moment generating function for the counting functions can be expressed in terms of the confluent hypergeometric-kernel determinant. By using Theorem 2, we derive the asymptotics for the mean, variance and covariance of the counting functions.

**COROLLARY 3.** For  $r_2 > r_1 > 0$  fixed, as  $t \to +\infty$ , we have

$$\mathbb{E}\left(N(tr_1)\right) = \mu_{\alpha}(tr_1) + \frac{\beta}{i\pi}\delta(tr_1) + \theta_{\alpha,\beta,1} + \mathcal{O}\left(\frac{\ln t}{t}\right), \qquad (1.23)$$

$$\mathbb{E}\left(N(-tr_1)\right) = \mu_{\alpha}(tr_1) - \frac{\beta}{i\pi}\delta(tr_1) - \theta_{\alpha,\beta,1} + \mathcal{O}\left(\frac{\ln t}{t}\right), \qquad (1.24)$$

$$\operatorname{Var}(N(\pm tr_1)) = \frac{\delta(tr_1) - (\ln G)''(1)}{\pi^2} + \theta_{\alpha,\beta,2} + \mathcal{O}\left(\frac{(\ln t)^2}{t}\right), \qquad (1.25)$$

$$\operatorname{Cov}\left(N(tr_1), N(tr_2)\right) = \Sigma(tr_1, tr_2) + \theta_{\alpha, \beta, 2} + \mathcal{O}\left(\frac{(\ln t)^2}{t}\right), \qquad (1.26)$$

$$\operatorname{Cov}\left(N(tr_1), N(-tr_2)\right) = -\Sigma(tr_1, tr_2) - \theta_{\alpha, \beta, 2} + \mathcal{O}\left(\frac{(\ln t)^2}{t}\right), \qquad (1.27)$$

where

$$\mu_{\alpha}(x) = \frac{x}{\pi} - \frac{\alpha}{2}, \quad \delta(x) = \ln 2x, \quad \Sigma(x, y) = \frac{1}{2\pi^2} \ln \frac{2xy}{|x - y|}, \quad \text{for } x, y > 0.$$
(1.28)

Here the parameters

$$\theta_{\alpha,\beta,1} = \frac{1}{2\pi i} \left( (\ln G)' (1 + \alpha - \beta) - (\ln G)' (1 + \alpha + \beta) \right), \tag{1.29}$$

$$\theta_{\alpha,\beta,2} = -\frac{1}{4\pi^2} \left( (\ln G)^{''} (1 + \alpha + \beta) + (\ln G)^{''} (1 + \alpha - \beta) \right), \tag{1.30}$$

and  $-(\ln G)^{''}(1) - 1 = \gamma_E \approx 0.5772$  is Euler's constant.

*Proof.* From (1.9) with n = 1 and the parameters  $0 = r_0 < r_1$ ,  $b_m = b_0 = -b$ , we obtain the following equation

$$F(t(0,r_1), 1 - e^{2\pi i b}; \alpha, \beta) = \mathbb{E}\left(e^{2\pi i b N(tr_1)}\right).$$
(1.31)

Thus, we derive that

$$\mathbb{E}(N(tr_1)) = \frac{1}{2\pi i} \partial_b \ln F(t(0, r_1), 1 - e^{2\pi i b}; \alpha, \beta) \Big|_{b=0}, \qquad (1.32)$$

and

$$\operatorname{Var}(N(tr_1)) = -\frac{1}{4\pi^2} \left. \partial_b^2 \ln F(t(0, r_1), 1 - e^{2\pi i b}; \alpha, \beta) \right|_{b=0}.$$
 (1.33)

Similarly, we compute the covariance between  $N(tr_1)$  and  $N(tr_2)$  as follows

$$\operatorname{Cov}\left(N(tr_{1}), N(tr_{2})\right) = -\frac{1}{8\pi^{2}} \left.\partial_{b}^{2} \ln\left(\frac{F\left(t(0, r_{1}, r_{2}), (1 - e^{4\pi i b}, 1 - e^{2\pi i b}); \alpha, \beta\right)}{F\left(t(0, r_{1}), 1 - e^{2\pi i b}; \alpha, \beta\right)F\left(t(0, r_{2}), 1 - e^{2\pi i b}; \alpha, \beta\right)}\right)\right|_{b=0} (1.34)$$

By using the definition of the Barnes G-function ([48, (5.17.3)]), we have

$$G(1+z) = (2\pi)^{z/2} \exp\left(-\frac{1}{2}z(z+1) - \frac{1}{2}\gamma_E z^2\right) \prod_{k=1}^{\infty} \left(\left(1 + \frac{z}{k}\right)^k \exp\left(-z + \frac{z^2}{2k}\right)\right).$$
(1.35)

From Theorem 2 and (1.32)-(1.35), we obtain after some direct calculations the mean, variance of  $N(r_1t)$  and the covariance between  $N(tr_1)$  and  $N(tr_2)$  as given in (1.23), (1.25) and (1.26).

The mean and variance of  $N(-r_1t)$  in (1.24) and (1.25) can be derived in the same manner by considering (1.9) with n = 1, specifying the parameters  $t(-r_1, 0)$  and  $b_m = b_1 = -b$ . The formula (1.27) can be derived similarly by considering  $F((-tr_2, 0, tr_1), (1 - e^{2\pi i b}, 1 - e^{2\pi i b}); \alpha, \beta)$ like (1.34). We complete the proof of the corollary.

From the definition of the counting function N(t), the number of particles in the confluent hypergeometric process falling into the interval [-t, t] is given by  $N_0(t) := N(t) + N(-t)$ . Combining (1.23), (1.25), (1.24) and (1.27), we immediately derive the asymptotics of the mean and variance of the counting function  $N_0(t)$ , which recovers the one established by Dai and Zhai in [25].

**COROLLARY 4.** (Dai and Zhai [25]) We have as  $t \to +\infty$ ,

$$\mathbb{E}(N_0(t)) = \frac{2t}{\pi} - \alpha + \mathcal{O}\left(\frac{\ln t}{t}\right), \quad \operatorname{Var}(N_0(t)) = \frac{\ln 4t + 1 + \gamma_E}{\pi^2} + \mathcal{O}\left(\frac{(\ln t)^2}{t}\right), \quad (1.36)$$

where  $\gamma_E \approx 0.5772$  is Euler's constant.

The rest of the paper is organized as follows. In Section 2, we formulate an RH problem for the Fredholm determinant defined in (1.7) by using the general theory of integrable operator developed in [38]. Making use of the confluent hypergeometric functions, we transform the RH problem to a model RH problem  $Y(\eta, t)$  with constant jumps. We then derive a Lax pair for  $Y(\eta, t)$  and the compatibility conditions of the Lax pair lead us to the coupled Painlevé V system. The Hamiltonian formulation of the system and several useful differential identities are also established in this section. In Sections 3 and 4, we derive the asymptotic approximations of a family of solutions to the coupled Painlevé V system and the associated Hamiltonian as  $t \to 0^+$  and  $t \to +\infty$  by performing Deift-Zhou nonlinear steepest descent analysis [26, 27] of the RH problem for  $Y(\eta, t)$ . The results are stated in Proposition 4 and 5 at the end of Sections 3 and 4, respectively. The last section, Section 5, is devoted to the proofs of Theorems 1 and 2. For the convenience of the reader, a brief description of the confluent hypergeometric parametrix is put in the Appendix.

# 2 Model RH problem and the coupled Painlevé V system

In this section, we first relate the determinant (1.7) to the solution of an RH problem by using the general theory of integrable operator developed in [38]. Then, we derive from the RH problem the Lax pair and the Hamiltonian for the coupled Painlevé V system (1.14). These results in the present section will be used in Section 5 to prove Theorem 1.

#### 2.1 RH problem for the Fredholm determinant

The kernel of the operator  $K_{t\vec{r},\vec{\gamma},\alpha,\beta}$  can be expressed in the following from

$$\sigma(y)K^{(\alpha,\beta)}(x,y) = \frac{\vec{f}(x)^T \vec{g}(y)}{x-y},$$
(2.1)

where

$$\vec{f}(z) = \frac{1}{\sqrt{2\pi i}} \begin{bmatrix} \frac{\Gamma(1+\alpha-\beta)}{\Gamma(1+2\alpha)} \mathbb{B}(z) \\ \frac{\Gamma(1+\alpha+\beta)}{\Gamma(1+2\alpha)} \mathbb{A}(z) \end{bmatrix}, \quad \vec{g}(z) = \frac{\sigma(z)}{\sqrt{2\pi i}} \begin{bmatrix} -\frac{\Gamma(1+\alpha+\beta)}{\Gamma(1+2\alpha)} \mathbb{A}(z) \\ \frac{\Gamma(1+\alpha-\beta)}{\Gamma(1+2\alpha)} \mathbb{B}(z) \end{bmatrix}, \quad (2.2)$$

with  $\mathbb{A}(z)$  and  $\mathbb{B}(z)$  defined in (1.3) and (1.4). Therefore, the kernel is integrable in the sense of [38]. By standard properties of trace class operators, we have

$$\frac{d}{dt}\ln F = \frac{d}{dt}\operatorname{tr}\left(\ln\left(1 - K_{\sigma}^{(\alpha,\beta)}\right)\right) 
= -\operatorname{tr}\left(\left(1 - K_{\sigma}^{(\alpha,\beta)}\right)^{-1}\frac{d}{dt}K_{\sigma}^{(\alpha,\beta)}\right) 
= \sum_{k=0}^{n} r_{k}\left(\lim_{z \to (r_{k}t)^{+}} R(z,z) - \lim_{z \to (r_{k}t)^{-}} R(z,z)\right),$$
(2.3)

where R(x, y) denotes the kernel of the resolvent operator  $(1 - K_{\sigma}^{(\alpha,\beta)})^{-1}K_{\sigma}^{(\alpha,\beta)}$ . According to [38] and [31], the kernel is also of the integrable form

$$R(x,y) = \frac{\left(\vec{F}(x), \vec{G}(y)\right)}{x - y},\tag{2.4}$$

with

$$\vec{F}(z) = \left(1 - K_{\sigma}^{(\alpha,\beta)}\right)^{-1} \vec{f}(z), \quad \vec{G}(z) = \left(1 - K_{\sigma}^{(\alpha,\beta)}\right)^{-1} \vec{g}(z).$$
(2.5)

Furthermore,  $\vec{F}$  and  $\vec{G}$  can be expressed as

$$\vec{F}(z) = m_{+}(z)\vec{f}(z), \quad \vec{G}(z) = m_{+}(z)^{-T}\vec{g}(z).$$
 (2.6)

Here m(z) solves the following RH problem.

#### **RH problem for** m(z)

- (1) m(z) analytic in  $\mathbb{C} \setminus [r_0 t, r_n t],$ (2)  $m_+(z) = m_-(z) \left( I - 2\pi i \vec{f}(z) \vec{g}(z)^T \right), \text{ on } \cup_{k=0}^{n-1} (r_k t, r_{k+1} t),$
- (3)  $m(z) \to I$ , as  $z \to \infty$ .

Conversely, we have [38]

$$m(z) = I - \int_{r_0 t}^{r_n t} \frac{\vec{F}(s)\vec{g}(s)^T}{s - z} ds,$$
(2.7)

with  $\vec{F}(s)$  defined by using the inverse of the operator  $I - K_{\sigma}^{(\alpha,\beta)}$  as given in (2.5). The determinant det $(1 - K_{\sigma}^{(\alpha,\beta)})$  is strictly positive since it represents the gap probability of the thinned confluent hypergeometric process. Therefore, the operator  $I - K_{\sigma}^{(\alpha,\beta)}$  is invertible and hence the existence of solution to the RH for Y(z) is justified.

#### 2.2 Model RH problem

Next, we transform the RH problem for m(z) to a new RH problem with constant jumps. To begin with, we note that the function  $\vec{f}(z)$  and  $\vec{g}(z)$  defined in (2.2) can be expressed as

$$\vec{f}(z) = e^{\frac{\beta}{2}\pi i\sigma_3} \Phi(2z) e^{\pm \frac{\alpha}{2}\pi i\sigma_3} e^{-\frac{\beta}{2}\pi i\sigma_3} \left(\frac{1}{\sqrt{2\pi i}}\right)^{\sigma_3} \begin{bmatrix} 1\\0 \end{bmatrix},$$
(2.8)

and

$$\vec{g}(z) = \sigma(z)e^{-\frac{\beta}{2}\pi i\sigma_3} \left(\Phi(2z)\right)^{-T} e^{\mp \frac{\alpha}{2}\pi i\sigma_3} e^{\frac{\beta}{2}\pi i\sigma_3} \left(\frac{1}{\sqrt{2\pi i}}\right)^{-\sigma_3} \begin{bmatrix} 0\\1 \end{bmatrix},$$
(2.9)

for  $z \in \Omega_{\Phi,1}$  and  $\pm \operatorname{Re} z > 0$ . Here  $\Phi(z)$  is the solution of the RH problem for the confluent hypergeometric parametrix given in the Appendix.

We introduce the transformation

$$Y(z,t) = (2t)^{\beta\sigma_3} e^{-\frac{\beta}{2}\pi i\sigma_3} m(tz) e^{\frac{\beta}{2}\pi i\sigma_3} \begin{cases} \Phi(2tz), & z \in \Omega_{Y,j}, \ j = 2, 5, \\ \widehat{\Phi}_j(2tz), & z \in \Omega_{Y,j}, \ j = 1, 3, 4, \end{cases}$$
(2.10)

where  $\Phi(z)$  is the solution of the RH problem for the confluent hypergeometric parametrix given in the Appendix. The functions  $\widehat{\Phi}_j(z)$  denote the analytic extension of  $\Phi(z)$  from the regions  $\Omega_{\Phi,j}$  to  $\Omega_{Y,j}$ , for j = 1, 3, 4, respectively; see Figures 1 and 5 for the regions.

Then Y(z) = Y(z, t) satisfies the following RH problem.

#### The model RH problem for Y(z)

(1) Y(z) is analytic for  $z \in \mathbb{C} \setminus \bigcup_{j=1}^{7} \Gamma_j$ , where the oriented contour are given by

$$\Gamma_1 = r_n + e^{\frac{\pi i}{4}} \mathbb{R}^+, \quad \Gamma_2 = r_0 + e^{\frac{3\pi i}{4}} \mathbb{R}^+, \quad \Gamma_3 = r_0 + e^{-\frac{3\pi i}{4}} \mathbb{R}^+, \quad \Gamma_4 = e^{-\frac{\pi i}{2}} \mathbb{R}^+,$$
  
$$\Gamma_5 = r_n + e^{-\frac{\pi i}{4}} \mathbb{R}^+, \quad \Gamma_6 = \bigcup_{k=0}^{m-1} (r_k, r_{k+1}), \quad \Gamma_7 = \bigcup_{k=m}^{n-1} (r_k, r_{k+1}),$$

as shown in Figure 1.



Figure 1: The jump contours and regions of the model RH problem for Y

(2)  $Y_+(z) = Y_-(z)J_Y(z)$  on  $\cup_{j=1}^7 \Gamma_j$ , where

$$J_{Y}(z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ e^{-(\alpha-\beta)\pi i} & 1 \end{bmatrix}, & z \in \Gamma_{1}, \\ \begin{bmatrix} 1 & 0 \\ e^{(\alpha-\beta)\pi i} & 1 \end{bmatrix}, & z \in \Gamma_{2}, \\ \begin{bmatrix} 1 & -e^{-(\alpha-\beta)\pi i} \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{3}, \\ e^{2\beta\pi i\sigma_{3}}, & z \in \Gamma_{4}, \\ \begin{bmatrix} 1 & -e^{(\alpha-\beta)\pi i} \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{5}, \\ \begin{bmatrix} 0 & -e^{-(\alpha-\beta)\pi i} \\ e^{(\alpha-\beta)\pi i} & 1 - \sigma(zt) \\ e^{-(\alpha-\beta)\pi i} & 1 - \sigma(zt) \end{bmatrix}, & z \in \Gamma_{6}, \\ \begin{bmatrix} 0 & -e^{(\alpha-\beta)\pi i} \\ e^{-(\alpha-\beta)\pi i} & 1 - \sigma(zt) \end{bmatrix}, & z \in \Gamma_{7}. \end{cases}$$

(3) As  $z \to \infty$ , we have for  $\arg z \in \left(-\frac{\pi}{2}, \frac{3\pi}{2}\right)$ 

$$Y(z) = \left(I + \frac{Y_1(t)}{z} + \frac{Y_2(t)}{z^2} + \mathcal{O}\left(z^{-3}\right)\right) z^{-\beta\sigma_3} e^{-izt\sigma_3}.$$
 (2.12)

(4) As  $z \to r_k$ ,  $k \neq m$ , we have

$$Y(z) = Y_0^{(k)}(t) \left( I + Y_1^{(k)}(t) \left( z - r_k \right) + Y_2^{(k)}(t) \left( z - r_k \right)^2 + \dots \right) \begin{bmatrix} 1 & \tilde{c}_k \ln \left( z - r_k \right) \\ 0 & 1 \end{bmatrix},$$
(2.13)

where  $\tilde{c}_k = \operatorname{sgn}(r_k)c_k e^{\operatorname{sgn}(r_k)\alpha\pi i}$  and  $c_k$  is defined in (1.17) and  $z \in \Omega_{Y,1}$ . The branch for  $\ln (z - r_k)$  is taken to be  $\operatorname{arg}(z - r_k) \in (0, 2\pi)$  if  $r_k < 0$ , or  $\operatorname{arg}(z - r_k) \in (-\pi, \pi)$  if  $r_k > 0$ . The behavior of Y(z) as  $z \to r_k$  for z in other regions is determined by using the jump condition (2.11).

(5) As  $z \to 0$  , we have

$$Y(z) = Y_0^{(m)}(t) \left( I + Y_1^{(m)}(t)z + Y_2^{(m)}(t)z^2 + \dots \right) z^{\alpha\sigma_3} \begin{cases} C_{m,1}, 2\alpha \notin \mathbb{Z}^+ \cup \{0\}, \\ C_{m,2}, 2\alpha \in \mathbb{Z}^+ \cup \{0\}, \end{cases}$$
(2.14)

where  $z \in \Omega_{Y,1}$ ,

$$C_{m,1} = \begin{bmatrix} 1 & \frac{c_m}{2i\sin 2\alpha\pi} \\ 0 & 1 \end{bmatrix}, \quad C_{m,2} = \begin{bmatrix} 1 & \frac{e^{-2\alpha\pi i}}{2\pi i}c_m\ln z \\ 0 & 1 \end{bmatrix},$$

and

$$c_m = (1 - \gamma_{m-1}) e^{(\alpha + \beta)\pi i} - (1 - \gamma_m) e^{-(\alpha + \beta)\pi i}.$$

Both  $z^{\alpha}$  and  $\ln z$  take the principal branches. The behavior of Y(z) as  $z \to 0$  for z in other regions is determined by using the jump condition (2.11).

#### 2.3 Lax pair and the coupled Painlevé V system

In this section, we derive a Lax pair from the above model RH problem, of which the compatibility condition gives us the coupled Painlevé V system. The Hamiltonian for the coupled Painlevé V system is also derived.

#### **PROPOSITION 1.** We have the Lax pair

$$\frac{\partial}{\partial z}Y(z,t) = \mathcal{A}(z,t)Y(z,t), \quad \frac{\partial}{\partial t}Y(z,t) = \mathcal{B}(z,t)Y(z,t), \quad (2.15)$$

where

$$\mathcal{A}(z,t) = \sum_{k=0}^{n} \frac{A_k(t)}{z - r_k} - it\sigma_3,$$
(2.16)

$$\mathcal{B}(z,t) = -iz\sigma_3 + B(t), \qquad (2.17)$$

with the coefficients being given as

$$A_m(t) = \begin{bmatrix} -\beta - \sum_{k \neq m} u_k(t) v_k(t) & \left(\alpha + \beta + \sum_{k \neq m} u_k(t) v_k(t)\right) y(t) \\ \frac{\alpha - \beta - \sum_{k \neq m} u_k(t) v_k(t)}{y(t)} & \beta + \sum_{k \neq m} u_k(t) v_k(t) \end{bmatrix},$$
(2.18)

$$A_{k}(t) = \begin{bmatrix} -u_{k}(t)v_{k}(t) & u_{k}(t)y(t) \\ -\frac{u_{k}(t)v_{k}^{2}(t)}{y(t)} & u_{k}(t)v_{k}(t) \end{bmatrix}, \quad k \neq m,$$
(2.19)

and

$$B(t) = \frac{1}{t} \begin{bmatrix} 0 & d_1(t)y(t) \\ \frac{d_2(t)}{y(t)} & 0 \end{bmatrix},$$
 (2.20)

where

$$d_{1}(t) = \alpha + \beta - \sum_{k \neq m} u_{k}(t) (v_{k}(t) - 1),$$
  

$$d_{2}(t) = \alpha - \beta - \sum_{k \neq m} u_{k}(t) v_{k}(t) (v_{k}(t) - 1).$$
(2.21)

The compatibility condition of the Lax pair is expressed as the coupled Painlevé V system (1.14)and the following differential equation for y(t)

$$t\frac{d}{dt}y = y(d_1 - d_2).$$
 (2.22)

 $Defining \ the \ Hamiltonian \ by$ 

$$tH = 2it(Y_1)_{11} - (\alpha^2 - \beta^2), \qquad (2.23)$$

where  $(Y_1)_{11}$  is the (1,1)-entry of the coefficient matrix  $Y_1$  in the asymptotic approximation (2.12) as  $z \to \infty$ , then we have

$$tH = \sum_{k \neq m} -2itr_k H_V(u_k, v_k, -2itr_k; \alpha, \beta) + \frac{1}{2} \sum_{j \neq k; \ j,k \neq m} u_j u_k(v_j + v_k)(v_j - 1)(v_k - 1), \ (2.24)$$

where  $H_V$  is the classical Hamiltonian for the Painlevé V equation defined in (1.13). The Hamiltonian system (1.11) is equivalent to the coupled Painlevé V system (1.14).

*Proof.* Observing that the jumping matrices in (2.11) are independent of both the variables z and t, Y(z,t),  $\frac{d}{dz}Y(z,t)$  and  $\frac{d}{dt}Y(z,t)$  have the same jump conditions. Thus,  $\mathcal{A}$  and  $\mathcal{B}$  are meromorphic functions in the z-plane with the only possible isolated singularities at  $z = r_k$ ,  $k = 0, \ldots, n$ . Taking (2.12),(2.13) and (2.14) into account, we then obtain that  $\mathcal{A}$  and  $\mathcal{B}$  are of the form given in (2.16) and (2.17).

Next, we derive the coefficients in the  $\mathcal{A}$  and  $\mathcal{B}$ . It is seen from (2.12), (2.13)) and (2.14) that

$$\det A_m = -\alpha^2, \qquad \det A_k = 0, \ k \neq m.$$
(2.25)

Moreover, in view of (2.12), we obtain by comparing the coefficients of 1/z on both sides of the first equation in (2.15)

$$\sum_{k=0}^{n} A_k = \begin{bmatrix} -\beta & 2it(Y_1)_{12} \\ -2it(Y_1)_{21} & \beta \end{bmatrix}.$$
 (2.26)

Therefore,  $A_k$ , k = 0, ..., n can be parametrized as (2.18) and (2.19). Similarly, we obtain (2.20) by substituting (2.12) into the second equation of (2.15) and using (2.26).

The compatibility condition of the Lax pair gives

$$\frac{d}{dt}A_k(t) = \left[-ir_k\sigma_3 + B(t), A_k(t)\right], \quad k \neq m,$$
(2.27)

and

$$\frac{d}{dt}A_m(t) = [B(t), A_m(t)].$$
(2.28)

We then have from (2.27)

$$\begin{cases} t \frac{d}{dt} (u_k v_k) = d_1 u_k v_k^2 + d_2 u_k, & k \neq m, \\ t \frac{d}{dt} (u_k y) = -2itr_k u_k y + 2d_1 u_k v_k y, & k \neq m. \end{cases}$$
(2.29)

While the equation (2.28) gives us

$$\begin{cases} t \frac{d}{dt} (A_m)_{11} = (A_m)_{11} (d_1 + d_2) + \alpha (d_1 - d_2), \\ t \frac{d}{dt} y = y (d_1 - d_2), \end{cases}$$
(2.30)

with  $d_1$  and  $d_2$  given in (2.21). Combining (2.29) and (2.30), we arrive at the coupled Painlevé V system (1.14) and (2.22).

To derive the expression of the Hamiltonian, we compare the coefficient of  $\frac{1}{z^2}$  in the large z expansion of the first equation in (2.14) and obtain

$$[-it\sigma_3 Y_1, Y_1] + [Y_2, -it\sigma_3] - \beta [Y_1, \sigma_3] - Y_1 = \sum_{k=0}^n A_k r_k.$$
(2.31)

A combination of (2.23) and (2.31) then gives (2.24). From (2.24), it is direct to see that the Hamiltonian system (1.11) is equivalent to the coupled Painlevé V system (1.14). This completes the proof of the proposition.

For later use, we also collect several relations as what follows.

**PROPOSITION 2.** The function y(t) appears in the coefficient (2.18) is related to the solution to the RH problem for Y(z) via

$$y(t) = \frac{\left(Y_0^{(m)}\right)_{11}}{\left(Y_0^{(m)}\right)_{21}},\tag{2.32}$$

where  $Y_0^{(m)}$  is given in the asymptotic expansion of Y(z) in (2.14). Moreover, let

$$d(t) = 2\alpha \left(Y_0^{(m)}\right)_{11} \left(Y_0^{(m)}\right)_{21},$$
(2.33)

then d(t) satisfies the differential equation

$$t\frac{d}{dt}d = d(d_1 + d_2), \qquad (2.34)$$

with  $d_1$  and  $d_2$  given in (2.21).

*Proof.* From (2.18), we have

$$y = \frac{(A_m)_{11} + \alpha}{(A_m)}.$$
 (2.35)

Then, by (2.14), (2.25) and (2.35), we obtain (2.32).

To derive (2.34), we obtain from (2.14) and the second equation in the Lax pair (2.15) that

$$\frac{d}{dt}Y_0^{(m)}(t) = B(t)Y_0^{(m)}(t).$$
(2.36)

This, together with (2.33) and (2.22), implies (2.34).

#### 2.4 Differential identity

Recently, the authors in [6, 40] have developed a new method to evaluate the total integral of the Hamiltonian. In this method, the differential identities for Hamiltonian play an important role. In the present subsection, we derive the differential identities for the Hamiltonian which will be used in Section 5, there we applied the method developed in [6, 40] to calculate the constant term in the large gap asymptotics.

**PROPOSITION 3.** We have the relation

$$H = \left(\sum_{k \neq m} u_k \frac{dv_k}{dt} - H\right) + \frac{d}{dt} \left(tH + \alpha \ln d - \beta \ln y - 2\left(\alpha^2 - \beta^2\right) \ln t\right), \qquad (2.37)$$

and the following differential identities

$$\frac{d}{d\gamma_j} \left( \sum_{k \neq m} u_k \frac{dv_k}{dt} - H \right) = \frac{d}{dt} \left( \sum_{k \neq m} u_k \frac{dv_k}{d\gamma_j} \right), \qquad (2.38)$$

with j = 0, ..., n - 1.

*Proof.* By using the Hamiltonian dynamic system (1.11), (2.22) and (2.34), we have

$$\sum_{k \neq m} u_k \frac{dv_k}{dt} - H = \sum_{k \neq m} u_k \frac{\partial H}{\partial u_k} - H$$
  
=  $\frac{1}{t} \sum_{k \neq m} u_k^2 v_k (v_k - 1)^2 + \frac{1}{t} \sum_{j < k} u_j u_k (v_j + v_k) (v_j - 1) (v_k - 1),$  (2.39)

and

$$\frac{d}{dt} (tH + \alpha \ln d(t) - \beta \ln y(t) - 2 (\alpha^2 - \beta^2) \ln t) = 2i \sum_{k \neq m} u_k v_k r_k - \frac{\alpha}{t} \sum_{k \neq m} u_k (v_k^2 - 1) - \frac{\beta}{t} \sum_{k \neq m} u_k (v_k - 1)^2.$$
(2.40)

Combining (2.24) with (2.39) and (2.40), we obtain (2.37).

Applying again the Hamiltonian dynamic system (1.11), we derive the differential identity with respect to the parameters

$$\frac{d}{d\gamma_j} \left( \sum_{k \neq m} u_k \frac{dv_k}{dt} - H \right) = \sum_{k \neq m} \left( \frac{du_k}{d\gamma_j} \frac{dv_k}{dt} + u_k \frac{d^2 v_k}{d\gamma_j dt} - \frac{\partial H}{\partial u_k} \frac{du_k}{d\gamma_j} - \frac{\partial H}{\partial v_k} \frac{dv_k}{d\gamma_j} \right) 
= \frac{d}{dt} \left( \sum_{k \neq m} u_k \frac{dv_k}{d\gamma_j} \right).$$
(2.41)

This completes the proof of the proposition.

# 3 Asymptotic Analysis as $t \to 0^+$

In this section, we derive the asymptotic approximations of a family of solutions to the coupled Painlevé V system (1.14) and the associated Hamiltonian as  $t \to 0^+$  by performing the Deift-Zhou nonlinear steepest descent analysis [26] of the RH problem for Y(z).

For the purpose, we consider

$$X(z) = (2t)^{-\beta\sigma_3} Y\left(\frac{z}{2t}, t\right).$$
(3.1)

It is readily seen from the RH problem for Y(z) that X(z) shares the same jump matrices with  $\Phi(z)$  given in the Appendix, for  $|z| > \max \{2 |r_0t|, 2 |r_nt|\}$ . Therefore, the solution to the RH problem for X(z) can be approximated by  $\Phi(z)$  for  $|z| > \max \{2 |r_0t|, 2 |r_nt|\}$  bounded away from the origin as  $t \to 0$ .

In the next subsection, we construct a local parametrix L(z) which has the same jump condition as X(z) in a neighborhood of the origin containing the interval  $[2r_0t, 2r_nt]$ .

#### 3.1 Local parametrix

#### **RH** problem for L(z)

(1) L(z) is analytic for  $z \in U(0, \varepsilon) \setminus \Sigma_X$  with some constant  $\varepsilon > 0$ .

(2) We have the jump relation

$$L_{+}(z) = L_{-}(z)J_{Y}(z/2t)$$
(3.2)

for  $z \in \Sigma_X \cap U(0, \varepsilon)$ , where  $J_Y(z)$  is given in (2.11).

(2) As  $t \to 0^+$ , we have

$$L(z)\Phi(z)^{-1} = I + O(t^{2\alpha+1}), \qquad (3.3)$$

where the error term is uniform for  $z \in \partial U(0, \varepsilon)$ .

Inspired by the asymptotic behavior of  $\Phi(z)$  near the origin given in (A.6) and (A.7), we construct L(z) of the following form

$$L(z) = \Phi^{(0)}(z) \begin{bmatrix} 1 & t^{2\alpha}k\left(\frac{z}{t}\right) \\ 0 & 1 \end{bmatrix} z^{\alpha\sigma_3} \begin{cases} \begin{bmatrix} 1 & \frac{\sin(\alpha+\beta)\pi}{\sin 2\alpha\pi} \\ 0 & 1 \end{bmatrix} C_j, & 2\alpha \notin \mathbb{N}, \\ \begin{bmatrix} 1 & (-1)^{2\alpha}\frac{\sin(\alpha+\beta)\pi}{\pi}\ln z \\ 0 & 1 \end{bmatrix} C_j, & 2\alpha \in \mathbb{N}, \end{cases}$$
(3.4)

where  $\alpha > -\frac{1}{2}$  and the branches for  $z^{\alpha}$  and  $\ln z$  are chosen such that  $\arg z \in (-\frac{\pi}{2}, \frac{3}{2}\pi)$ . Here the function k(z) is analytic for  $z \in \mathbb{C} \setminus [2r_0, 2r_n]$  with the following jumps on  $(2r_0, 2r_n)$ 

$$k_{+}(x) - k_{-}(x) = -|x|^{2\alpha} \sigma\left(\frac{xt}{2}\right) e^{(\alpha+\beta)\pi i}, \quad 2r_{0} < x < 0,$$
(3.5)

and

$$k_{+}(x) - k_{-}(x) = -x^{2\alpha}\sigma\left(\frac{xt}{2}\right)e^{(\alpha-\beta)\pi i}, \quad 0 < x < 2r_n,$$
(3.6)

in which  $\sigma(x)$  is the step function defined in (1.8). Applying the Plemelj formula, we have

$$k(z) = -\frac{e^{(\alpha+\beta)\pi i}}{2\pi i} \int_{2r_0}^0 \frac{|x|^{2\alpha}\sigma\left(\frac{xt}{2}\right)}{x-z} dx - \frac{e^{(\alpha-\beta)\pi i}}{2\pi i} \int_0^{2r_n} \frac{x^{2\alpha}\sigma\left(\frac{xt}{2}\right)}{x-z} dx, \quad z \in \mathbb{C} \setminus [2r_0, 2r_n].$$
(3.7)

It is seen from the construction (3.4) and the jump relations (3.5) and (3.6) that the jump condition (3.2) is fulfilled.

To show (3.3), we expand k(z) as  $z \to \infty$ :

$$k(z) = -e^{\alpha \pi i} \sum_{k=0}^{n} \frac{c_k}{2\alpha + 1} \left(2|r_k|\right)^{2\alpha + 1} \frac{1}{z} + \mathcal{O}\left(\frac{1}{z^2}\right).$$
(3.8)

Substituting (3.8) into (3.4), we have as  $t \to 0$ 

$$L(z)\Phi(z)^{-1} = I - e^{\alpha\pi i} \sum_{k=0}^{n} \frac{c_k}{2\alpha + 1} \left(2|r_k|\right)^{2\alpha + 1} \frac{t^{2\alpha + 1}}{z} \Phi^{(0)}(z) \begin{bmatrix} 0 & 1\\ 0 & 0 \end{bmatrix} \Phi^{(0)}(z)^{-1} + \mathcal{O}\left(t^{2\alpha + 2}\right),$$
(3.9)

where the error term is uniform for  $z \in \partial U(0, \varepsilon)$ . Therefore, we have the matching condition (3.3).

#### 3.2 Ratio RH problem

We define the following ratio RH problem

$$E(z) = \begin{cases} X(z)L(z)^{-1}, & |z| < \varepsilon, \\ X(z)\Phi(z)^{-1}, & |z| > \varepsilon. \end{cases}$$
(3.10)

Then, E(z) satisfies the following RH problem.

#### **RH problem for** E(z)

- (1) E(z) is analytic for  $z \in \mathbb{C} \setminus U(0, \varepsilon)$  with some constant  $\varepsilon > 0$ .
- (2) E(z) satisfies the jump condition

$$E_{+}(z) = E_{-}(z)J_{E}(z), \quad J_{E}(z) = \Phi(z)L(z)^{-1},$$
(3.11)

for  $z \in \partial U(0, \varepsilon)$  with the orientation clockwise.

(3)  $E(z) = I + \mathcal{O}(1/z)$  as  $z \to \infty$ .

Using (3.3) and a standard argument for the small norm RH problem as given in [26] , we have the expansion

$$E(z) = I - e^{\alpha \pi i} \sum_{k=0}^{n} \frac{c_k}{2\alpha + 1} \frac{(2|r_k|t)^{2\alpha + 1}}{z} \Phi^{(0)}(0) \begin{bmatrix} 0 & 1\\ 0 & 0 \end{bmatrix} \Phi^{(0)}(0)^{-1} + \mathcal{O}\left(t^{2\alpha + 2}\right), \quad |z| > \varepsilon, \quad (3.12)$$

and

$$E(z) = I + \mathcal{O}\left(t^{2\alpha+1}\right) \tag{3.13}$$

for  $|z| < \varepsilon$ .

### **3.3** Asymptotic behavior as $t \to 0^+$

As a consequence of nonlinear steepest descent analysis of the RH problem for Y(z), we obtain the asymptotics of a family of special solutions to the coupled Painlevé V system and the Hamiltonian associated.

**PROPOSITION 4.** For  $\alpha > -\frac{1}{2}$ ,  $\beta \in i\mathbb{R}$  and  $\gamma_k \in [0,1)$ , there exist solutions to the coupled Painlevé V system (1.14) with the following asymptotic behaviors as  $t \to 0^+$ 

$$\begin{cases} u_k(t) = \frac{\operatorname{sgn}(r_k)c_k\Gamma(1+\alpha-\beta)\Gamma(1+\alpha+\beta)(2|r_k|t)^{2\alpha}}{\Gamma^2(1+2\alpha)} \left(1 + \mathcal{O}\left(t^{2\alpha+1}\right)\right), \\ v_k(t) = 1 + \mathcal{O}\left(t^{2\alpha+1}\right), \end{cases}$$
(3.14)

with  $c_k$  given in (1.17) for  $k \neq m$ . Furthermore, the Hamiltonian H(t) associated with the solutions has the asymptotic behavior (1.16) as  $t \to 0^+$ .

*Proof.* Tracing back the invertible transformations  $Y \to X \to E$  given in (3.1) and (3.10), we obtain

$$Y(z/2t) = \frac{1}{2t} (2t)^{\beta \sigma_3} E(z) \begin{cases} L(z), & |z| < \varepsilon, \\ \Phi(z), & |z| > \varepsilon, \end{cases}$$
(3.15)

where L(z),  $\Phi(z)$  and E(z) are given in (3.4), (3.12), (3.13) and (A.8).

Comparing the large-z expansion on both sides of the second equation in (3.15), we have

$$Y_1 = \frac{1}{2t} (2t)^{\beta \sigma_3} (E_1 + \Phi_1) (2t)^{-\beta \sigma_3}, \qquad (3.16)$$

where  $Y_1$ ,  $E_1$  and  $\Phi_1$  are the coefficients of 1/z in the large-z expansion of Y(z), E(z) and  $\Phi(z)$ , respectively. Substituting (3.12) and (A.5) into (3.16) yields

$$(Y_1)_{11} = \sum_{k=0}^{n} \frac{c_k \Gamma(1+\alpha-\beta) \Gamma(1+\alpha+\beta) \left(2|r_k|\right)^{2\alpha+1}}{2(2\alpha+1)\Gamma^2(1+2\alpha)} t^{2\alpha} + \frac{\alpha^2-\beta^2}{2it} + \mathcal{O}\left(t^{2\alpha+1}\right), \quad (3.17)$$

as  $t \to 0^+$ . This, together with (2.23), implies the asymptotics of the Hamiltonian as given in (1.16).

To derive the asymptotics of y(t) and d(t), we consider the asymptotic expansion of Y(z) as  $t \to 0^+$  for z near the origin. Plugging (2.14), (3.4) and (3.13) into the first equation in (3.15), we have

$$Y_0^{(m)} \begin{bmatrix} 1\\0 \end{bmatrix} = (2t)^{\alpha} (2t)^{\beta \sigma_3} \left( I + \mathcal{O} \left( t^{2\alpha+1} \right) \right) \Phi^{(0)}(0) \begin{bmatrix} 1\\0 \end{bmatrix}, \qquad (3.18)$$

where  $\Phi^{(0)}(0)$  is given in (A.6). Using (A.8), we have from (3.18) that

$$Y_0^{(m)} = \begin{bmatrix} (2t)^{\alpha+\beta} e^{-\frac{\alpha+\beta}{2}\pi i} \frac{\Gamma(1+\alpha-\beta)}{\Gamma(1+2\alpha)} & * \\ (2t)^{\alpha-\beta} e^{-\frac{\alpha-\beta}{2}\pi i} \frac{\Gamma(1+\alpha+\beta)}{\Gamma(1+2\alpha)} & * \end{bmatrix} \left(I + \mathcal{O}(t^{2\alpha+1})\right),$$
(3.19)

for  $\alpha > -1/2$ ,  $2\alpha \notin \mathbb{N}$  and  $\beta \in i\mathbb{R}$ . The case  $2\alpha \in \mathbb{N}$  can be derived in a similar way and we skip the details. Recalling (2.32) and (2.33), we derive the asymptotics of y(t) and d(t) as  $t \to 0^+$ :

$$y(t) = \frac{\Gamma(1+\alpha-\beta)}{\Gamma(1+\alpha+\beta)} e^{-\beta\pi i} (2t)^{2\beta} \left(1 + \mathcal{O}\left(t^{2\alpha+1}\right)\right), \qquad (3.20)$$

$$d(t) = 2\alpha \frac{\Gamma(1+\alpha-\beta)\Gamma(1+\alpha+\beta)}{\Gamma^2(1+2\alpha)} e^{-\alpha\pi i} (2t)^{2\alpha} \left(1+\mathcal{O}\left(t^{2\alpha+1}\right)\right).$$
(3.21)

Finally, we derive the asymptotics of  $u_k$  and  $v_k$  for  $k \neq m$  by using (3.20) and the following relations

$$v_k(t) = \frac{\left(Y_0^{(k)}(t)\right)_{21}}{\left(Y_0^{(k)}(t)\right)_{11}} y(t), \qquad (3.22)$$

and

$$u_k(t) = \frac{\tilde{c}_k \left(Y_0^{(k)}\right)_{11} \left(Y_0^{(k)}\right)_{21}}{v(t)},\tag{3.23}$$

which can be seen from Proposition 1 and (2.13). Similarly, we obtain after substituting (3.4) and (3.13) into the first equation in (3.15) that

$$Y(z) = (2t)^{\beta\sigma_3} \left( I + \mathcal{O}\left(t^{2\alpha+1}\right) \right) \Phi^{(0)}(z) \begin{bmatrix} 1 & t^{2\alpha}k\left(\frac{z}{t}\right) \\ 0 & 1 \end{bmatrix} z^{\alpha\sigma_3} \begin{bmatrix} 1 & \frac{\sin(\alpha+\beta)\pi}{\sin 2\alpha\pi} \\ 0 & 1 \end{bmatrix}.$$
(3.24)

From the integral expression of k(z) in (3.7), we have

$$k(z) \sim \operatorname{sgn}(r_k) c_k e^{\alpha \pi i} (2|r_k|)^{2\alpha} \ln (z - 2r_k),$$
 (3.25)

as  $z \to 2r_k$  for  $k \neq m$ . Comparing (2.13) with (3.24) and (3.25), we obtain

$$Y_0^{(k)} = \begin{bmatrix} (2t)^{\alpha+\beta} e^{-\frac{\alpha+\beta}{2}\pi i} \frac{\Gamma(1+\alpha-\beta)}{\Gamma(1+2\alpha)} r_k^{\alpha} & * \\ (2t)^{\alpha-\beta} e^{-\frac{\alpha-\beta}{2}\pi i} \frac{\Gamma(1+\alpha+\beta)}{\Gamma(1+2\alpha)} r_k^{\alpha} & * \end{bmatrix} (I + \mathcal{O}(t^{2\alpha+1}))$$
(3.26)

for  $r_k > 0$ , and

$$Y_0^{(k)} = \begin{bmatrix} (2t)^{\alpha+\beta} e^{-\frac{\alpha+\beta}{2}\pi i} \frac{\Gamma(1+\alpha-\beta)}{\Gamma(1+2\alpha)} \left( |r_k| e^{\pi i} \right)^{\alpha} & * \\ (2t)^{\alpha-\beta} e^{-\frac{\alpha-\beta}{2}\pi i} \frac{\Gamma(1+\alpha+\beta)}{\Gamma(1+2\alpha)} \left( |r_k| e^{\pi i} \right)^{\alpha} & * \end{bmatrix} (I + \mathcal{O}(t^{2\alpha+1}))$$
(3.27)

for  $r_k < 0$ . Using (3.20), (3.22), (3.23), (3.26) and (3.27), we have (3.14). This completes the proof of the proposition.

# 4 Asymptotic Analysis as $t \to +\infty$

In this section, we carry out the Deift-Zhou nonlinear steepest descent analysis of the RH problem for Y(z) as  $t \to +\infty$ . Based on this, we obtain the asymptotic behavior of a family of solutions to the coupled Painlevé V system and the associated Hamiltonian as  $t \to +\infty$ .

#### 4.1 Normalization

In the first transformation, we normalize the behavior of Y(z) at infinity.

$$T(z) = Y(z,t)e^{izt\sigma_3},\tag{4.1}$$

Then, T(z) satisfies the following RH problem.

### **RH** problem for T(z)

- (1) T(z) is analytic in  $\mathbb{C} \setminus \bigcup_{j=1}^{7} \Gamma_j$ . The jump contours are shown in Figure 1.
- (2) T(z) satisfies the jump condition

$$T_{+}(z) = T_{-}(z)J_{T}(z), \qquad (4.2)$$

where

$$J_{T}(z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ e^{-(\alpha-\beta)\pi i}e^{2izt} & 1 \end{bmatrix}, & z \in \Gamma_{1}, \\ \begin{bmatrix} 1 & 0 \\ e^{(\alpha-\beta)\pi i}e^{2izt} & 1 \end{bmatrix}, & z \in \Gamma_{2}, \\ \begin{bmatrix} 1 & -e^{-(\alpha-\beta)\pi i}e^{-2izt} \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{3}, \\ e^{2\beta\pi i\sigma_{3}}, & z \in \Gamma_{4}, \\ \begin{bmatrix} 1 & -e^{(\alpha-\beta)\pi i}e^{-2izt} \\ 0 & 1 \end{bmatrix}, & z \in \Gamma_{5}, \\ \begin{bmatrix} 0 & -e^{-(\alpha-\beta)\pi i}e^{-2izt} \\ e^{(\alpha-\beta)\pi i}e^{2izt} & 1 - \sigma(zt) \\ 0 & -e^{(\alpha-\beta)\pi i}e^{-2izt} \\ e^{-(\alpha-\beta)\pi i}e^{2izt} & 1 - \sigma(zt) \end{bmatrix}, & z \in \Gamma_{6}, \\ \begin{bmatrix} 0 & -e^{(\alpha-\beta)\pi i}e^{-2izt} \\ e^{-(\alpha-\beta)\pi i}e^{2izt} & 1 - \sigma(zt) \\ e^{-(\alpha-\beta)\pi i}e^{2izt} & 1 - \sigma(zt) \end{bmatrix}, & z \in \Gamma_{7}. \end{cases}$$

(3) As  $z \to \infty$ ,

$$T(z) = \left(I + \frac{Y_1(t)}{z} + \mathcal{O}\left(\frac{1}{z^2}\right)\right) z^{-\beta\sigma_3}.$$
(4.4)

(4) The behavior of T(z) near  $r_k$ , k = 0, ..., n, are the same as that of Y(z), as given in (2.13) and (2.14).

#### 4.2 Opening lenses and deformation

It is readily observed that the off-diagonal entries in the jump matrices on  $\Gamma_6$  and  $\Gamma_7$  are highly oscillatory as  $t \to +\infty$ . To eliminate the oscillation, we factorize the jump matrices on  $\Gamma_6$  and  $\Gamma_7$  as follow:

$$\begin{bmatrix} 0 & -e^{-(\alpha-\beta)\pi i}e^{-2izt} \\ e^{(\alpha-\beta)\pi i}e^{2izt} & 1-\sigma \end{bmatrix} = \\ 1 & -\frac{1}{1-\sigma}e^{-(\alpha-\beta)\pi i}e^{-2izt} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{1-\sigma} & 0 \\ 0 & 1-\sigma \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \frac{1}{1-\sigma}e^{(\alpha-\beta)\pi i}e^{2izt} & 1 \end{bmatrix}, \quad z \in \Gamma_6,$$

$$(4.5)$$

$$\begin{bmatrix} 0 & -e^{(\alpha-\beta)\pi i}e^{-2izt} \\ e^{-(\alpha-\beta)\pi i}e^{2izt} & 1-\sigma \end{bmatrix} =$$

$$\begin{bmatrix} 1 & -\frac{1}{1-\sigma}e^{(\alpha-\beta)\pi i}e^{-2izt} \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{1-\sigma} & 0 \\ 0 & 1-\sigma \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \frac{1}{1-\sigma}e^{-(\alpha-\beta)\pi i}e^{2izt} & 1 \end{bmatrix}, \quad z \in \Gamma_7.$$

$$(4.6)$$

We also deform the contour  $\Gamma_6$  and  $\Gamma_7$  into lens-shape regions as shown in Figure 2.



Figure 2: Jump contours and regions for S

Based on the factorization of jumps and deformation of contours, we introduce the second transformation

$$S(z) = T(z) \begin{cases} \begin{bmatrix} 1 & 0 \\ -\frac{1}{1-\sigma}e^{(\alpha-\beta)\pi i}e^{2izt} & 1 \end{bmatrix}, & z \in H_k, 0 \le k < m, \\ \begin{bmatrix} 1 & 0 \\ -\frac{1}{1-\sigma}e^{-(\alpha-\beta)\pi i}e^{2izt} & 1 \end{bmatrix}, & z \in H_k, m \le k < n, \\ \begin{bmatrix} 1 & -\frac{1}{1-\sigma}e^{-(\alpha-\beta)\pi i}e^{-2izt} \\ 0 & 1 \end{bmatrix}, & z \in L_k, 0 \le k < m, \\ \begin{bmatrix} 1 & -\frac{1}{1-\sigma}e^{(\alpha-\beta)\pi i}e^{-2izt} \\ 0 & 1 \end{bmatrix}, & z \in L_k, m \le k < n, \\ I & \text{elsewhere.} \end{cases}$$
(4.7)

Then, S(z) satisfies the following RH problem.

### **RH** problem for S(z)

(1) S(z) is analytic in  $\mathbb{C}\backslash\Gamma_S$  as shown in Figure 2.

(2) S(z) satisfies the jump condition

$$S_{+}(z) = S_{-}(z)J_{S}(z), \qquad (4.8)$$

where

$$J_{S}(z) = \begin{cases} J_{T}(z), & z \in \bigcup_{j=1}^{5} \Gamma_{S,j}, \\ \begin{bmatrix} 1 & 0 \\ \frac{1}{1-\gamma_{k}} e^{(\alpha-\beta)\pi i} e^{2izt} & 1 \end{bmatrix}, & \text{on } \Gamma_{S,8,k}, k < m, \\ \begin{bmatrix} 1 & 0 \\ \frac{1}{1-\gamma_{k}} e^{-(\alpha-\beta)\pi i} e^{2izt} & 1 \end{bmatrix}, & \text{on } \Gamma_{S,8,k}, k \ge m, \\ \begin{bmatrix} 1 & -\frac{1}{1-\gamma_{k}} e^{-(\alpha-\beta)\pi i} e^{-2izt} \\ 0 & 1 \end{bmatrix}, & \text{on } \Gamma_{S,9,k}, k < m, \\ \begin{bmatrix} 1 & -\frac{1}{1-\gamma_{k}} e^{(\alpha-\beta)\pi i} e^{-2izt} \\ 0 & 1 \end{bmatrix}, & \text{on } \Gamma_{S,9,k}, k \ge m, \\ \begin{bmatrix} 1 & -\frac{1}{1-\gamma_{k}} e^{(\alpha-\beta)\pi i} e^{-2izt} \\ 0 & 1 \end{bmatrix}, & \text{on } \Gamma_{S,9,k}, k \ge m, \\ \begin{bmatrix} 1 & -\frac{1}{1-\gamma_{k}} e^{(\alpha-\beta)\pi i} e^{-2izt} \\ 0 & 1 \end{bmatrix}, & \text{on } \Gamma_{S,9,k}, k \ge m, \\ \begin{bmatrix} 1 & -\frac{1}{1-\gamma_{k}} e^{(\alpha-\beta)\pi i} e^{-2izt} \\ 0 & 1 \end{bmatrix}, & \text{on } \Gamma_{S,9,k}, k \ge m, \end{cases}$$

(3) As  $z \to \infty$ , we have

$$S(z) = \left(I + \frac{Y_1(t)}{z} + \mathcal{O}\left(\frac{1}{z^2}\right)\right) z^{-\beta\sigma_3}.$$
(4.10)

(4) The asymptotics of S(z) are the same as T(z) near  $r_k$ , k = 0, ..., n.

#### 4.3 Global parametrix

The jumps for S(z) tend to the identity matrix exponentially fast except the ones on the real axis as  $t \to \infty$ . Therefore, we arrive at the following approximate RH problem as  $t \to +\infty$ .

# **RH** problem for $P^{(\infty)}(z)$

- (1)  $P^{(\infty)}(z)$  is analytic for  $z \in \mathbb{C} \setminus \{ [r_0, r_n] \cup \Gamma_{S,4} \}.$
- (2)  $P^{(\infty)}(z)$  satisfies the following jump condition

$$P_{+}^{(\infty)}(z) = P_{-}^{(\infty)}(z) \begin{cases} \left[ \begin{array}{cc} \frac{1}{1-\sigma} & 0\\ 0 & 1-\sigma \end{array} \right], & z \in (r_0, r_n), \\ e^{2\beta\pi i\sigma_3}, & z \in \Gamma_{S,4}, \end{cases}$$
(4.11)

(3) As  $z \to \infty$ ,

$$P^{(\infty)}(z) = \left(I + \mathcal{O}\left(\frac{1}{z}\right)\right) z^{-\beta\sigma_3}.$$
(4.12)

The solution  $P^{(\infty)}(z)$  can be constructed explicitly as below

$$P^{(\infty)}(z) = \prod_{k=0}^{n-1} \left(\frac{z - r_{k+1}}{z - r_k}\right)^{-\frac{\ln(1 - \gamma_k)}{2\pi i}\sigma_3} z^{-\beta\sigma_3},$$
(4.13)

where  $\left(\frac{z-r_{k+1}}{z-r_k}\right)^{-\frac{\ln(1-\gamma_k)}{2\pi i}}$  takes the principal branch for  $0 \le k \le n-1$  and the branch of  $z^{-\beta}$  is taken such that  $\arg z \in \left(-\frac{\pi}{2}, \frac{3\pi}{2}\right)$ .

#### 4.4 Local parametrices

Near the points  $r_k$ , k = 0, ..., n, the solution of the RH problem S(z) can not be approximated by the global parametrix  $P^{(\infty)}(z)$ . In this section, we construct the local parametrices in the neighborhoods of  $r_k$ , k = 0, ..., n.

# **RH** problem for $P^{(r_k)}(z)$

- (1)  $P^{(r_k)}(z)$  is analytic for  $z \in U(r_k, \varepsilon) \setminus \Gamma_S$  with some constant  $\varepsilon > 0$ .
- (2)  $P^{(r_k)}(z)$  has the same jumps as S(z) for  $z \in \Gamma_S \cap U(r_k, \varepsilon)$ .
- (3) As  $t \to +\infty$ , we have

$$S(z) = \left(I + O\left(\frac{1}{t}\right)\right) P^{(r_k)}(z), \qquad (4.14)$$

where the error term is uniform for  $z \in \partial U(r_k, \varepsilon)$ .

For  $k \neq m,$  the solution can be constructed by using the confluent hypergeometric parametrix  $\Phi^{\rm (CHF)}$ 

$$P^{(r_{k})}(z) = E^{(r_{k})}(z)\Phi^{(CHF)}(2it(z-r_{k});0,b_{k})\left[(1-\gamma_{k})(1-\gamma_{k-1})\right]^{-\frac{\sigma_{3}}{4}} \\ \times \begin{cases} e^{-\operatorname{sgn}(r_{k})\frac{\alpha-\beta}{2}\pi i\sigma_{3}}e^{izt\sigma_{3}}, & \operatorname{Im} z > 0, \\ 0 & 1 \\ -1 & 0 \end{cases} e^{-\operatorname{sgn}(r_{k})\frac{\alpha-\beta}{2}\pi i\sigma_{3}}e^{izt\sigma_{3}}, & \operatorname{Im} z < 0, \end{cases}$$
(4.15)

where

$$E^{(r_k)}(z) = \prod_{j \neq k} (z - r_j)^{-b_j \sigma_3} z^{-\beta \sigma_3} (2it)^{b_k \sigma_3} e^{-ir_k t \sigma_3} e^{\operatorname{sgn}(r_k) \frac{\alpha - \beta}{2} \pi i \sigma_3} \\ \times \begin{cases} (1 - \gamma_{k-1})^{-\frac{\sigma_3}{4}} (1 - \gamma_k)^{\frac{3}{4} \sigma_3}, & \operatorname{Im} z > 0, \\ [(1 - \gamma_k) (1 - \gamma_{k-1})]^{-\frac{\sigma_3}{4}}, & \operatorname{Im} z < 0, \end{cases}$$
(4.16)

with  $\gamma_{-1} = \gamma_n = 0$ . While for  $r_m = 0$ , we construct  $P^{(r_m)}$  as follows

$$P^{(r_m)}(z) = E^{(r_m)}(z)\Phi^{(\text{CHF})}(2itz;\alpha,\beta+b_m) \left[ (1-\gamma_{m-1})(1-\gamma_m) \right]^{-\frac{\sigma_3}{4}} \\ \times \begin{cases} e^{\frac{\alpha}{2}\pi i\sigma_3} e^{izt\sigma_3}, & \arg z \in (0,\frac{\pi}{2}), \\ e^{-\frac{\alpha}{2}\pi i\sigma_3} e^{izt\sigma_3}, & \arg z \in (\frac{\pi}{2},\pi), \\ e^{\frac{\alpha+2\beta}{2}\pi i\sigma_3} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} e^{izt\sigma_3}, & \arg z \in (-\pi,\frac{\pi}{2}), \\ e^{-\frac{\alpha+2\beta}{2}\pi i\sigma_3} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} e^{izt\sigma_3}, & \arg z \in (-\frac{\pi}{2},0), \end{cases}$$
(4.17)

where

$$E^{(r_m)}(z) = \prod_{j \neq m} (z - r_j)^{-b_j \sigma_3} (2it)^{(\beta + b_m)\sigma_3} \\ \times \begin{cases} e^{-\beta \pi i \sigma_3} (1 - \gamma_{m-1})^{-\frac{\sigma_3}{4}} (1 - \gamma_m)^{\frac{3}{4}\sigma_3}, & \arg z \in (0, \pi), \\ e^{\beta \pi i \sigma_3} [(1 - \gamma_{m-1}) (1 - \gamma_m)]^{-\frac{\sigma_3}{4}}, & \arg z \in (-\pi, -\frac{\pi}{2}), \\ e^{-\beta \pi i \sigma_3} [(1 - \gamma_{m-1}) (1 - \gamma_m)]^{-\frac{\sigma_3}{4}}, & \arg z \in (-\frac{\pi}{2}, 0). \end{cases}$$
(4.18)

The branch for  $z^{\beta}$  is taken such that  $\arg z \in (-\frac{\pi}{2}, \frac{3\pi}{2})$ . While the branches of the other power functions in (4.15), (4.16), (4.17) and (4.18) take the principal branches.

#### 4.5 Final transformation

In the final transformation, we define

$$R(z) = \begin{cases} S(z)P^{(r_k)}(z)^{-1}, & z \in U(r_k,\varepsilon), \quad 0 \le k \le n, \\ S(z)P^{(\infty)}(z)^{-1} & \text{elsewhere.} \end{cases}$$
(4.19)

#### **RH** problem for R(z)

(1) The jump contours for R are shown in Figure 3



Figure 3: Jump contours  $\Gamma_R$  and regions for R

(2) R(z) satisfies the jump condition

$$R_{+}(z) = R_{-}(z)J_{R}(z), \qquad (4.20)$$

where

$$J_R(z) = \begin{cases} P^{(r_k)}(z)P^{(\infty)}(z)^{-1}, & z \in \partial U(r_k,\varepsilon), \quad 0 \le k \le n, \\ P^{(\infty)}(z)J_S(z)P^{(\infty)}(z)^{-1} & \text{elsewhere.} \end{cases}$$
(4.21)

(3) As  $z \to \infty$ ,

$$R(z) = I + \mathcal{O}(\frac{1}{z}). \tag{4.22}$$

From (4.9) and (4.14), we have

$$J_R(z) = I + \mathcal{O}\left(\frac{1}{t}\right),\tag{4.23}$$

where the error term is uniform for z on the jump contours for R(z),  $\gamma_j$  in any compact subset of [0, 1), and  $r_0 < \cdots < r_m = 0 < \cdots < r_n$  in any compact subset of  $\mathbb{R}$  and bounded away from each other. Therefore, we have as  $t \to +\infty$ 

$$R(z) = I + \mathcal{O}\left(\frac{1}{t}\right),\tag{4.24}$$

where the error term is uniform for z in the complex plane. For later use, we derive from (4.13), (4.15), (4.16) and (A.5) the following expansion

$$J_R(z) = I + \frac{J_{R,1}(z)}{t} + \mathcal{O}\left(\frac{1}{t^2}\right), \qquad (4.25)$$

where

$$J_{R,1}(z) = -\frac{b_k^2}{2i(z-r_k)} E^{(r_k)}(z) \begin{bmatrix} 1 & \frac{\Gamma(-b_k)}{\Gamma(1+b_k)} \\ \frac{-\Gamma(b_k)}{\Gamma(1-b_k)} & -1 \end{bmatrix} E^{(r_k)}(z)^{-1},$$
(4.26)

for  $z \in \partial U(r_k, \varepsilon), k \neq m$  and

$$J_{R,1}(z) = \frac{\alpha^2 - (\beta + b_m)^2}{2iz} E^{(r_m)}(z) \begin{bmatrix} 1 & \frac{\Gamma(\alpha - \beta - b_m)}{\Gamma(1 + \alpha + \beta + b_m)} \\ \frac{-\Gamma(\alpha + \beta + b_m)}{\Gamma(1 + \alpha - \beta - b_m)} & -1 \end{bmatrix} E^{(r_m)}(z)^{-1}, \quad (4.27)$$

for  $z \in \partial U(0, \varepsilon)$ . Therefore, we have as  $t \to +\infty$ 

$$R(z) = I + \frac{R_1(z)}{t} + \mathcal{O}\left(\frac{1}{t^2}\right), \qquad (4.28)$$

where

$$R_1(z) = \frac{1}{2\pi i} \sum_{k=0}^n \int_{\partial U(r_k,\xi)} \frac{J_{R,1}(x)}{x-z} dx.$$
(4.29)

After some residue calculation, we have

$$(R_{1}(z))_{11} = \begin{cases} -\sum_{j=0}^{n} \frac{b_{j}^{2}}{2i(z-r_{j})} + \frac{\alpha^{2}-\beta^{2}-2\beta b_{m}}{2iz}, & z \in \mathbb{C} \setminus \bigcup_{j=0}^{n} U(r_{j},\varepsilon), \\ -\sum_{j \neq k} \frac{b_{j}^{2}}{2i(z-r_{j})} + \frac{\alpha^{2}-\beta^{2}-2\beta b_{m}}{2iz}, & z \in U(r_{k},\varepsilon), \ k \neq m, \\ -\sum_{j \neq m} \frac{b_{j}^{2}}{2i(z-r_{j})}, & z \in U(r_{m},\varepsilon). \end{cases}$$
(4.30)

For later use, we derive the following estimate of derivatives of  $J_R(z)$  with respect to  $\gamma_j$  as  $t \to +\infty$ ,

$$\left\| \left( \prod_{j=0}^{n-1} \partial_{\gamma_j}^{k_j} \right) (J_R(z) - I) \right\|_{L^2(\Gamma_R) \cap L^\infty(\Gamma_R)} \le c \frac{(\ln t)^k}{t}, \tag{4.31}$$

where  $k = \sum_{j=0}^{n-1} k_j$  and the constant c > 0. The factor  $(\ln t)^k$  is due to the factors  $t^{\pm b_j}$  in the entries of  $J_R(z)$ , cf. (4.15)-(4.18) and (4.21). It is worth pointing out that the estimate holds uniformly for  $\gamma_j$  in any compact subset of [0, 1), and uniformly for  $r_0 < \cdots < r_m = 0 < \cdots < r_n$  in any compact subset of  $\mathbb{R}$  and bounded away from each other. Then, by (4.29), we obtain

$$\left(\prod_{j=0}^{n-1} \partial_{\gamma_j}^{k_j}\right) R_1(z) = \mathcal{O}((\ln t)^k), \quad \text{as } t \to +\infty.$$
(4.32)

#### 4.6 Asymptotic behaviors as $t \to +\infty$

Using the results of the nonlinear steepest descent analysis of the RH problem for Y(z), we obtain the asymptotics of a family of special solutions to the coupled Painlevé V system and the Hamiltonian associated as  $t \to +\infty$ .

**PROPOSITION 5.** Under the same conditions as in Proposition 4, the family of solutions of the coupled Painlevé V system (1.14) with the asymptotic behaviors as  $t \to 0^+$  given in Proposition 4 have the asymptotics behaviors as  $t \to +\infty$ 

$$\begin{cases} u_{k}(t) = \operatorname{sgn}(r_{k}) \ c_{k} \cdot \frac{\Gamma^{2} \left(1 - b_{k}\right) \Gamma(1 + \alpha + \beta + b_{m})}{\Gamma(1 + \alpha - \beta - b_{m})} \prod_{j \neq m, k} \left(\frac{r_{k} - r_{j}}{r_{m} - r_{j}}\right)_{+}^{-2b_{j}} |r_{k}|^{2(b_{k} - b_{m} - \beta)} \\ [(1 - \gamma_{m-1}) \left(1 - \gamma_{m}\right)]^{-\frac{1}{2}} e^{\operatorname{sgn}(r_{k})\pi i(b_{k} + b_{m} + \alpha + \beta)} (2t)^{2(b_{k} - b_{m} - \beta)} e^{-2itr_{k}} (1 + \mathcal{O}(t^{-1})), \\ v_{k}(t) = \operatorname{sgn}(r_{k}) \ \frac{(\gamma_{k} - \gamma_{k-1})}{2\pi i c_{k}} \frac{\Gamma(1 + \alpha - \beta - b_{m}) \Gamma(1 + b_{k})}{\Gamma(1 + \alpha + \beta + b_{m}) \Gamma(1 - b_{k})} \prod_{j \neq m, k} \left(\frac{r_{k} - r_{j}}{r_{m} - r_{j}}\right)_{+}^{2b_{j}} |r_{k}|^{2(\beta + b_{m} - b_{k})} \\ \left[\frac{(1 - \gamma_{m-1}) \left(1 - \gamma_{m}\right)}{(1 - \gamma_{k-1}) \left(1 - \gamma_{k}\right)}\right]^{\frac{1}{2}} e^{-\operatorname{sgn}(r_{k})\pi i(b_{k} + b_{m} + \alpha + \beta)} (2t)^{2(\beta + b_{m} - b_{k})} e^{2itr_{k}} \left(1 + \mathcal{O}\left(t^{-1}\right)\right), \\ (4.33)$$

for  $k \neq m$  and  $c_k$  is defined in (1.17). Here the power functions take the principal branches. Moreover, the Hamiltonian associated with this family of solutions has the asymptotics behavior as  $t \to +\infty$ 

$$H(t) = \sum_{k=0}^{n} 2ib_k r_k - \frac{\sum_{k=0}^{n} b_k^2 + 2\beta b_m}{t} + \mathcal{O}\left(t^{-2}\right).$$
(4.34)

*Proof.* Tracing back the transformations  $Y \mapsto N \mapsto S \mapsto R$ , we can derive the asymptotics of Y(z) uniformly for z in several different regions covering the whole complex plane as  $t \to +\infty$ . As  $t \to +\infty$ , we obtain from (4.1), (4.7), (4.13), (4.19) and (4.28) that

$$Y(z) = R(z)P^{(\infty)}(z)e^{-izt\sigma_{3}}$$
  
=  $\left(I + \frac{R_{1}(z)}{t} + \mathcal{O}\left(\frac{1}{t^{2}}\right)\right)\prod_{k=0}^{n-1}\left(\frac{z - r_{k+1}}{z - r_{k}}\right)^{-\frac{\ln(1-\gamma_{k})}{2\pi i}\sigma_{3}}z^{-\beta\sigma_{3}}e^{-izt\sigma_{3}},$  (4.35)

where the error term is uniform for  $|z - r_k| > \varepsilon$ , k = 1, ..., n. Here, the branch for  $z^{\beta}$  is taken such that  $\arg z \in (-\frac{\pi}{2}, \frac{3\pi}{2})$  and the other power functions take the principal branches. Using(2.12), (4.30) and (4.35), we have

$$(Y_1)_{11} = -\frac{1}{2\pi i} \sum_{k=0}^{n-1} (r_k - r_{k+1}) \ln(1 - \gamma_k) + \frac{\alpha^2 - \beta^2 - 2\beta b_m - \sum_{k=0}^n b_k^2}{2it} + \mathcal{O}\left(\frac{1}{t^2}\right).$$
(4.36)

Substitute (4.36) into (2.23), we obtain (4.34).

From (4.1), (4.7), (4.17), (4.19) and (4.28), we derive as  $t \to +\infty$ 

$$Y(z) = R(z)P^{(r_m)}(z)e^{-izt\sigma_3} = \left(I + \frac{R_1(z)}{t} + \mathcal{O}\left(\frac{1}{t^2}\right)\right) E^{(r_m)}(z)\Phi\left(2itz;\alpha,\beta+b_m\right)$$
(4.37)  
$$[(1 - \gamma_{m-1})(1 - \gamma_m)]^{-\frac{\sigma_3}{4}}e^{\frac{\alpha}{2}\pi i\sigma_3},$$

where  $\arg z \in (0, \frac{\pi}{2})$  and the error term is uniform for  $|z| < \varepsilon$ . Using (2.14), (4.18) and (4.30), we obtain from (4.37) that

$$\begin{pmatrix} Y_0^{(m)} \end{pmatrix}_{11} = \frac{\Gamma\left(1 + \alpha - \beta - b_m\right)}{\Gamma(1 + 2\alpha)} \prod_{j \neq m} (-r_j)_+^{-b_j} e^{-i\frac{\pi}{2}(\alpha + \beta + b_m)} (2t)^{\alpha + \beta + b_m} \left(1 + \mathcal{O}\left(\frac{1}{t}\right)\right),$$

$$\begin{pmatrix} Y_0^{(m)} \end{pmatrix}_{21} = \frac{\Gamma\left(1 + \alpha + \beta + b_m\right)}{\Gamma(1 + 2\alpha)} \prod_{j \neq m} (-r_j)_+^{b_j} e^{-i\frac{\pi}{2}(\alpha - \beta - b_m)} (2t)^{\alpha - \beta - b_m}$$

$$[(1 - \gamma_{m-1})(1 - \gamma_m)]^{-\frac{1}{2}} \left(1 + \mathcal{O}\left(\frac{1}{t}\right)\right).$$

$$(4.38)$$

Therefore, we have from (2.32), (2.33) and (4.38) that as  $t \to +\infty$ 

$$y(t) = \frac{\Gamma(1+\alpha-\beta-b_m)}{\Gamma(1+\alpha+\beta+b_m)} \prod_{j\neq m} (-r_j)_+^{-2b_j} e^{-(\beta+b_m)\pi i} (2t)^{2(\beta+b_m)} [(1-\gamma_{m-1})(1-\gamma_m)]^{\frac{1}{2}} \left(1+\mathcal{O}\left(\frac{1}{t}\right)\right)$$
(4.39)

and

$$d(t) = 2\alpha \frac{\Gamma\left(1 + \alpha - \beta - b_m\right)\Gamma\left(1 + \alpha + \beta + b_m\right)}{\Gamma^2(1 + 2\alpha)} e^{-\alpha\pi i} (2t)^{2\alpha} \left[\left(1 - \gamma_m\right)\left(1 - \gamma_{m-1}\right)\right]^{-\frac{1}{2}} \left(1 + \mathcal{O}\left(\frac{1}{t}\right)\right)$$

$$(4.40)$$

Similarly, we have from (4.1), (4.7) and (4.19) that as  $t \to +\infty$ 

$$Y(z) = R(z)P^{(r_k)}(z)e^{-izt\sigma_3},$$
(4.41)

where the error term is uniform for  $|z - r_k| < \varepsilon$ . From (4.1), (4.7), (4.17), (4.18) and (4.28)), we obtain

$$\begin{pmatrix} Y_0^{(k)}(t) \end{pmatrix}_{11} = \Gamma \left(1 - b_k\right) \prod_{j \neq k} \left(r_k - r_j\right)_+^{-b_j} r_k^{-\beta} e^{-\frac{b_k}{2}\pi i} (2t)^{b_k} e^{-itr_k} (1 + \mathcal{O}(\frac{1}{t})),$$

$$\begin{pmatrix} Y_0^{(k)}(t) \end{pmatrix}_{21} = \Gamma \left(1 + b_k\right) \prod_{j \neq k} \left(r_k - r_j\right)_+^{b_j} r_k^{\beta} e^{\frac{b_k}{2}\pi i} (2t)^{-b_k} e^{-\operatorname{sgn}(r_k)(\alpha - \beta)\pi i}$$

$$[(1 - \gamma_{k-1}) (1 - \gamma_k)]^{-\frac{1}{2}} (1 + \mathcal{O}(\frac{1}{t})).$$

$$(4.42)$$

From (3.22), (3.23) and (4.42), we have (4.33). This completes the proof of the proposition.  $\Box$ 

# 5 Proof of main theorems

#### 5.1 Proof of Theorem 1

From (2.3), (2.4), (2.6), (2.10) and (2.13), we have

$$\frac{d}{dt}\ln F = \frac{1}{t} \sum_{k=0}^{n} \tilde{c}_k r_k \left(Y_1^{(k)}\right)_{21},\tag{5.1}$$

where  $r_m = 0$ ,  $Y_1^{(k)}$  is the coefficient appears in the expansion (2.13) for k = 0, ..., n. Here the coefficients  $\tilde{c}_k = c_k e^{\operatorname{sgn}(r_k)(\alpha+1)\pi i}$  and  $c_k$  is defined in (1.17). By (2.13) and (2.15), we have

$$\left(\frac{d}{dt}Y_1^{(k)}\right)_{21} = 2i\left(Y_0^{(k)}\right)_{11}\left(Y_0^{(k)}\right)_{21} = \frac{2iu_k v_k}{\tilde{c}_k}.$$
(5.2)

Therefore, we have

$$\frac{d}{dt}\left(t\frac{d}{dt}\ln F\right) = 2i\sum_{k=1}^{n}r_{k}u_{k}v_{k}.$$
(5.3)

On the other side, we get from (1.13), (1.12) (1.11) that

$$\frac{d}{dt}(tH(t)) = 2i\sum_{k=1}^{n} r_k u_k v_k,$$
(5.4)

with  $r_m = 0$ . From (1.16), (5.3) and (5.4), we have

$$\frac{d}{dt}\ln F = H(t). \tag{5.5}$$

By (1.16), the Hamiltonian function H is integrable near zero. By using the fact that F(0) = 1, we obtain the integral expression (1.15) by taking integration on both sides of the above equation. This completes the proof of the theorem.

#### 5.2 Proof of Theorem 2

From the asymptotic behavior of the Hamiltonian function (1.18), we obtain the asymptotics of the integral except the constant term as  $t \to +\infty$ 

$$\int_{0}^{t} H(s)ds = \left(\sum_{k=0}^{n} 2ib_{k}r_{k}\right)t - \left(\sum_{k=0}^{n} b_{k}^{2} + 2\beta b_{m}\right)\ln t + C_{1} + \mathcal{O}\left(\frac{1}{t}\right),$$
(5.6)

where  $C_1 = C_1(\alpha, \beta, \vec{\gamma})$  is some constant depending on the parameters.

To determine the constant  $C_1$ , we integrate on both sides of (2.37)

$$\int_{0}^{t} H(s)ds = \int_{0}^{t} \sum_{k \neq m} u_{k} \frac{dv_{k}}{ds} - H(s)ds + (sH(s) + \alpha \ln d(s) - \beta \ln y(s) - 2(\alpha^{2} - \beta^{2}) \ln s) \Big|_{0}^{t}.$$
(5.7)

Substituting the asymptotics (1.16), (3.20), (3.21), (4.34), (4.39) and (4.40) into the second term on the right-hand side of the above equation, we have

$$\left(sH + \alpha \ln d - \beta \ln y - 2\left(\alpha^2 - \beta^2\right) \ln s\right)\Big|_{0}^{t}$$

$$= \sum_{k=0}^{n} 2ib_k r_k t - \sum_{k=0}^{n} b_k^2 - 2\beta b_m + \alpha \ln \frac{\Gamma\left(1 + \alpha - \beta - b_m\right)\Gamma\left(1 + \alpha + \beta + b_m\right)}{\Gamma\left(1 + \alpha - \beta\right)\Gamma\left(1 + \alpha + \beta\right)}$$

$$- \beta \ln \frac{\Gamma\left(1 + \alpha - \beta - b_m\right)\Gamma\left(1 + \alpha + \beta\right)}{\Gamma\left(1 + \alpha + \beta + b_m\right)\Gamma\left(1 + \alpha + \beta\right)} + \beta \sum_{j \neq m} 2b_j \ln |r_j| - 2\beta b_m \ln(2t)$$

$$- \frac{\alpha}{2} \ln \left[\left(1 - \gamma_{m-1}\right)\left(1 - \gamma_m\right)\right] + \mathcal{O}\left(\frac{1}{t}\right).$$

$$(5.8)$$

Comparing (5.6) with (5.7) and (5.8), we have

$$\int_{0}^{t} \sum_{k \neq m} u_{k} \frac{dv_{k}}{ds} - H ds = -\sum_{k=0}^{n} b_{k}^{2} \ln t + C_{2} + \mathcal{O}\left(\frac{1}{t}\right),$$
(5.9)

with

$$C_{1} = C_{2} + \alpha \ln \frac{\Gamma \left(1 + \alpha - \beta - b_{m}\right) \Gamma \left(1 + \alpha + \beta + b_{m}\right)}{\Gamma (1 + \alpha - \beta) \Gamma (1 + \alpha + \beta)} - \beta \ln \frac{\Gamma \left(1 + \alpha - \beta - b_{m}\right) \Gamma (1 + \alpha + \beta)}{\Gamma \left(1 + \alpha + \beta + b_{m}\right) \Gamma (1 + \alpha - \beta)} + \beta \sum_{k \neq m} 2b_{k} \ln |r_{k}| - 2\beta b_{m} \ln(2) - \frac{\alpha}{2} \ln \left[\left(1 - \gamma_{m-1}\right) \left(1 - \gamma_{m}\right)\right] - \Sigma b_{k}^{2} - 2\beta b_{m}.$$

$$(5.10)$$

Therefore, the remaining task is to derive the constant  $C_2$ . It is more convenient to evaluate the integral constant  $C_2$  since the derivatives of the integrand in (5.9) with respect to the parameters  $\vec{\gamma}$  is totally differential in t as shown in (2.38).

Actually, we obtain from (2.38) that

$$\frac{d}{db_j}C_2 = \lim_{t \to +\infty} \left( \sum_{k \neq m} u_k(t) \frac{dv_k(t)}{db_j} + 2b_j \ln t - 2b_m \ln t \right) - \lim_{t \to 0^+} \left( \sum_{k \neq m} u_k(t) \frac{dv_k(t)}{db_j} + 2b_j \ln t - 2b_m \ln t \right)$$
(5.11)

Substituting into the above formula the asymptotics (3.14), (3.21), (4.33) and (4.40), we have

$$\frac{d}{db_j}C_2 = b_j \left(\partial_{b_j} \ln \frac{\Gamma\left(1+\alpha-\beta-b_m\right)}{\Gamma\left(1+\alpha+\beta+b_m\right)} + \partial_{b_j} \ln \frac{\Gamma\left(1+b_j\right)}{\Gamma\left(1-b_j\right)} - 4\ln|2r_j|\right) \\
+ \sum_{k \neq m,j} b_k \left(\partial_{b_j} \ln \frac{\Gamma\left(1+\alpha-\beta-b_m\right)}{\Gamma\left(1+\alpha+\beta+b_m\right)} - 2\ln\left|\frac{2r_jr_k}{r_j-r_k}\right|\right),$$
(5.12)

with  $j \neq m$  and  $b_m = -\sum_{j\neq m} b_j$ .

When the parameters  $\vec{b} = \vec{0}$ , we have  $\ln F(s) = 0$  and the constant  $C_1(0, \ldots, 0; \alpha, \beta) = C_2(0, \ldots, 0; \alpha, \beta) = 0$ . Using this initial value and the differential identity (5.12), we evaluate the constant  $C_2$  by taking integration with respect to the variables  $b_j$ ,  $j \neq m$  successively. For the purpose, we recall the integral representation for the Barnes G-function (see [48])

$$\ln G(z+1) = \frac{1}{2}z\ln(2\pi) - \frac{1}{2}z(z+1) + z\ln\Gamma(z+1) - \int_0^z\ln\Gamma(t+1)dt.$$
 (5.13)

Using (5.13), we obtain by integrating by parts that

$$\int_{0}^{b_{j}} x \partial_{x} \ln \frac{\Gamma(1+x)}{\Gamma(1-x)} dx = b_{j}^{2} + \ln \left[ G \left( 1 + b_{j} \right) G \left( 1 - b_{j} \right) \right].$$
(5.14)

In the *j*-th step, taking  $b_{j+1} = \cdots = b_n = 0$ , we have  $b_m = -s_j$  with  $s_j = \sum_{l=0, l \neq m}^j b_l$ . Then, we have from integration by parts and (5.13) that

$$\int_{0}^{b_{j}} (s_{j}+x)\partial_{x} \ln \frac{\Gamma\left(1+\alpha-\beta+s_{j-1}+x\right)}{\Gamma\left(1+\alpha+\beta-s_{j-1}-x\right)} dx$$

$$= \left[x \ln \frac{\Gamma(1+\alpha-\beta+x)}{\Gamma(1+\alpha+\beta-x)}\right]_{s_{j-1}}^{s_{j}} - \int_{s_{j-1}}^{s_{j}} \partial_{x} \ln \frac{\Gamma(1+\alpha-\beta+x)}{\Gamma(1+\alpha+\beta-x)} dx$$

$$= b_{j}^{2} + 2b_{j}s_{j-1} - (\alpha+\beta) \ln \frac{\Gamma(1+\alpha+\beta-s_{j})}{\Gamma\left(1+\alpha+\beta-s_{j-1}\right)} - (\alpha-\beta) \ln \frac{\Gamma(1+\alpha-\beta+s_{j})}{\Gamma\left(1+\alpha-\beta+s_{j-1}\right)}$$

$$+ \ln \frac{G\left(1+\alpha-\beta+s_{j}\right)G\left(1+\alpha+\beta-s_{j-1}\right)}{G(1+\alpha-\beta+s_{j-1})G(1+\alpha+\beta-s_{j-1})}.$$
(5.15)

By (5.12), (5.15), (5.14) and the initial value  $C_2(0, \ldots, 0; \alpha, \beta) = 0$ , we derive

$$C_{2}(\vec{b};\alpha,\beta) = \sum_{k \neq m} b_{k}^{2} + 2\beta b_{m} + \alpha \ln \frac{\Gamma(1+\alpha+\beta)\Gamma(1+\alpha-\beta)}{\Gamma(1+\alpha+\beta+b_{m})\Gamma(1+\alpha-\beta-b_{m})}$$

$$+ \beta \ln \frac{\Gamma(1+\alpha+\beta)\Gamma(1+\alpha-\beta-b_{m})}{\Gamma(1+\alpha-\beta)\Gamma(1+\alpha+\beta+b_{m})} + \sum_{j \neq m} \ln \left[G\left(1+b_{j}\right)G\left(1-b_{j}\right)\right]$$

$$+ \ln \frac{G\left(1+\alpha+\beta+b_{m}\right)G\left(1+\alpha-\beta-b_{m}\right)}{G(1+\alpha+\beta)G(1+\alpha-\beta)} - \sum_{\substack{0 \leq j \neq k \leq n \\ j,k \neq m}} 2b_{j}b_{k} \ln \left|\frac{2r_{j}r_{k}}{r_{j}-r_{k}}\right|$$

$$(5.16)$$

$$- \sum_{j \neq m} 2b_{j}^{2} \ln |2r_{j}|.$$

This, together with (5.10), implies

$$C_{1} = -\sum_{k \neq m} 2b_{k}^{2} \ln|2r_{k}| + \sum_{k \neq m} 2b_{k}\beta \ln|2r_{k}| - \sum_{\substack{0 \leq j < k \leq n \\ j,k \neq m}} 2b_{j}b_{k}\ln\left|\frac{2r_{j}r_{k}}{r_{j} - r_{k}}\right| + \sum_{k \neq m} \ln\left[G\left(1 + b_{j}\right)G\left(1 - b_{j}\right)\right] + \ln\frac{G\left(1 + \alpha + \beta + b_{m}\right)G\left(1 + \alpha - \beta - b_{m}\right)}{G(1 + \alpha + \beta)G(1 + \alpha - \beta)}$$
(5.17)  
$$- \frac{\alpha}{2}\ln\left[\left(1 - \gamma_{m-1}\right)\left(1 - \gamma_{m}\right)\right].$$

Substitute (5.17) into (5.6), we obtain (1.20).

Finally, from (1.15), (2.23), (2.38), (4.32) and (4.35), we see that (1.20) can be be differentiated with respect to  $\gamma_0, \ldots, \gamma_{n-1}$  at the cost of increasing the error term by a factor  $\ln t$  for each differentiation. This completes the proof of the theorem.

# A The confluent hypergeometric parametrix

# RH problem for $\Phi^{\rm (CHF)}$

(1)  $\Phi^{(CHF)}(z; \alpha, \beta)$  ( $\Phi^{(CHF)}(z)$ , for short) is analytic in  $\mathbb{C} \setminus \{\bigcup_{j=1}^{8} \Sigma_j\}$ , where the contours  $\Sigma_j$ ,  $j = 1, \ldots, 8$ , are shown in Fig. 4.

(2)  $\Phi^{(CHF)}(z)$  satisfies the jump condition

$$\left(\Phi^{(\mathrm{CHF})}\right)_{+}(z) = \left(\Phi^{(\mathrm{CHF})}\right)_{-}(z)J_{j}(z), \quad z \in \Sigma_{i}, \quad j = 1, \dots, 8,$$
(A.1)

where

$$J_{1}(z) = \begin{bmatrix} 0 & e^{-\pi i\beta} \\ -e^{\pi i\beta} & 0 \end{bmatrix}, \quad J_{2}(z) = \begin{bmatrix} 1 & 0 \\ e^{\pi i(\beta - 2\alpha)} & 1 \end{bmatrix}, \quad J_{3}(z) = J_{7}(z) = e^{\pi i\alpha\sigma_{3}},$$
$$J_{4}(z) = \begin{bmatrix} 1 & 0 \\ e^{-\pi i(\beta - 2\alpha)} & 1 \end{bmatrix}, \quad J_{5}(z) = \begin{bmatrix} 0 & e^{\pi i\beta} \\ -e^{-\pi i\beta} & 0 \end{bmatrix}, \quad J_{6}(z) = \begin{bmatrix} 1 & 0 \\ e^{-\pi i(\beta + 2\alpha)} & 1 \end{bmatrix},$$

and

$$J_8(z) = \begin{bmatrix} 1 & 0 \\ e^{\pi i(\beta + 2\alpha)} & 1 \end{bmatrix}.$$



Figure 4: The jump contours and regions of the RH problem for  $\Phi^{(CHF)}(z)$ .

(3) As  $z \to \infty$ ,  $\Phi^{(CHF)}(z)$  satisfies the asymptotic behavior

$$\Phi^{(\text{CHF})}(z) = \left(I + \frac{\alpha^2 - \beta^2}{z} \begin{bmatrix} 1 & \frac{\Gamma(\alpha - \beta)}{\Gamma(1 + \alpha + \beta)} \\ -\frac{\Gamma(\alpha + \beta)}{\Gamma(1 + \alpha - \beta)} & -1 \end{bmatrix} + \dots \right) z^{-\beta\sigma_3} e^{-\frac{1}{2}z\sigma_3}$$

$$\begin{cases} e^{-\frac{1}{2}\alpha\pi i\sigma_3} e^{\beta\pi i\sigma_3}, & \frac{\pi}{2} < \arg z < \pi, \\ e^{\frac{1}{2}\alpha\pi i\sigma_3} e^{\beta\pi i\sigma_3}, & \pi < \arg z < \frac{3\pi}{2}, \\ \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} e^{-\frac{1}{2}\alpha\pi i\sigma_3}, & -\frac{\pi}{2} < \arg z < 0, \\ \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} e^{\frac{1}{2}\alpha\pi i\sigma_3}, & 0 < \arg z < \frac{\pi}{2}. \end{cases}$$
(A.2)

(4) As  $z \to 0$ ,  $\Phi^{(CHF)}(z)$  has the following asymptotic behaviors

$$\Phi^{(\text{CHF})}(z) = \begin{bmatrix} O(|z|^{\alpha}) & O(|z|^{-|\alpha|}) \\ O(|z|^{\alpha}) & O(|z|^{-|\alpha|}) \end{bmatrix}, \quad \alpha > 0,$$
$$\Phi^{(\text{CHF})}(z) = \begin{bmatrix} O(1) & O(\ln|z|) \\ O(1) & O(\ln|z|) \end{bmatrix}, \quad \alpha = 0,$$

and

$$\Phi^{(\mathrm{CHF})}(z) = \begin{bmatrix} O(|z|^{\alpha}) & O(|z|^{|\alpha|}) \\ O(|z|^{\alpha}) & O(|z|^{|\alpha|}) \end{bmatrix}, \quad \alpha < 0.$$

For the application in the present paper, we introduce a matrix-valued function  $\Phi(z)$  defined

via  $\Phi^{(CHF)}(z)$  by the transformation:

$$\Phi(z) = e^{-\frac{1}{2}\beta\pi i\sigma_3} \Phi^{(\text{CHF})}(e^{\frac{1}{2}\pi i}z;\alpha,\beta) \begin{cases} e^{\frac{\alpha}{2}\pi i\sigma_3}, & \arg z \in \left(0,\frac{\pi}{2}\right), \\ e^{-\frac{\alpha}{2}\pi i\sigma_3}, & \arg z \in \left(\frac{\pi}{2},\pi\right), \\ e^{\frac{\alpha+2\beta}{2}\pi i\sigma_3} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, & \arg z \in \left(-\pi, -\frac{\pi}{2}\right), \\ e^{-\frac{\alpha+2\beta}{2}\pi i\sigma_3} \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, & \arg z \in \left(-\frac{\pi}{2}, 0\right). \end{cases}$$
(A.3)

Then  $\Phi(z)$  satisfies the following RH problem.

(1)  $\Phi(z)$  is analytic in  $\mathbb{C} \setminus \bigcup_{j=1}^{5} \Sigma_{\Phi,j}$ , where the oriented contours

$$\Sigma_{\Phi,1} = e^{\frac{\pi i}{4}} \mathbb{R}^+, \quad \Sigma_{\Phi,2} = e^{\frac{3\pi i}{4}} \mathbb{R}^+, \quad \Sigma_{\Phi,3} = e^{-\frac{3\pi i}{4}} \mathbb{R}^+,$$
$$\Sigma_{\Phi,4} = e^{-\frac{\pi i}{2}} \mathbb{R}^+, \quad \Sigma_{\Phi,5} = e^{-\frac{\pi i}{4}} \mathbb{R}^+,$$

are shown in Figure 5.



Figure 5: The jump contours and regions of the RH problem for  $\Phi(z)$ .

(2)  $\Phi(z)$  satisfies the jump condition  $\Phi_+(z) = \Phi_-(z)J_{\Phi}(z)$  on  $\cup_{j=1}^5 \Sigma_{\Phi,j}$ , where

$$J_{\Phi}(z) = \begin{cases} \begin{bmatrix} 1 & 0 \\ e^{-\pi i (\alpha - \beta)} & 1 \end{bmatrix}, & z \in \Sigma_{\Phi, 1}, \\ \begin{bmatrix} 1 & 0 \\ e^{\pi i (\alpha - \beta)} & 1 \end{bmatrix}, & z \in \Sigma_{\Phi, 2}, \\ \begin{bmatrix} 1 & -e^{-\pi i (\alpha - \beta)} \\ 0 & 1 \end{bmatrix}, & z \in \Sigma_{\Phi, 3}, \\ e^{2\pi i \beta \sigma_3}, & z \in \Sigma_{\Phi, 4}, \\ \begin{bmatrix} 1 & -e^{\pi i (\alpha - \beta)} \\ 0 & 1 \end{bmatrix}, & z \in \Sigma_{\Phi, 5}. \end{cases}$$
(A.4)

(3) As  $z \to \infty$ , we have

$$\Phi(z) = \left(I + \frac{\Phi_1}{z} + \mathcal{O}\left(\frac{1}{z^2}\right)\right) z^{-\beta\sigma_3} e^{-\frac{i}{2}z\sigma_3}, \tag{A.5}$$
$$- \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ - \left(\alpha^2 - \beta^2\right) i - e^{-\pi i\beta} \frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)} \\ -$$

where 
$$\Phi_1 = \begin{bmatrix} -(\alpha^2 - \beta^2)i & -e^{-\pi i\beta}\frac{\Gamma(1+\alpha-\beta)}{i\Gamma(\alpha+\beta)}\\ e^{\pi i\beta}\frac{\Gamma(1+\alpha+\beta)}{i\Gamma(\alpha-\beta)} & (\alpha^2 - \beta^2)i \end{bmatrix}$$
, and arg  $z \in (-\frac{\pi}{2}, \frac{3\pi}{2})$ 

(4) If  $\alpha > -\frac{1}{2}$  and  $2\alpha \notin \mathbb{N}$ , we have as  $z \to 0$ 

$$\Phi(z) = \Phi^{(0)}(z) z^{\alpha \sigma_3} C_j, \quad z \to 0, \ z \in \Omega_{\Phi,j}, \ j = 1, 2, 3, 4, 5,$$
(A.6)

where  $\Phi^{(0)}(z)$  is analytic near the origin, the regions  $\Omega_{\Phi,j}$  are shown in Figure 5, and the branch for  $z^{\alpha\sigma_3}$  is chosen such that  $\arg z \in (-\pi/2, 3\pi/2)$ . The constant matrix

$$C_1 = \begin{bmatrix} 1 & \frac{\sin(\pi(\alpha + \beta))}{\sin(2\pi\alpha)} \\ 0 & 1 \end{bmatrix}$$

and the constant matrices  $C_j$ , j = 2, 3, 4, 5 are determined by  $C_1$  and the jump condition. If  $2\alpha \in \mathbb{N}$ , we have as  $z \to 0$ 

$$\Phi(z) = \widehat{\Phi}^{(0)}(z) z^{\alpha \sigma_3} \begin{bmatrix} 1 & (-1)^{2\alpha} \frac{\sin(\pi(\alpha+\beta))}{\pi} \ln z \\ 0 & 1 \end{bmatrix} \widehat{C}_j, \ z \in \Omega_{\Phi,j}, \ j = 1, 2, 3, 4, 5,$$
(A.7)

where  $\widehat{\Phi}^{(0)}(z)$  is analytic near the origin, the regions  $\Omega_{\Phi,j}$  are shown in Figure 5, and the branches for  $z^{\alpha\sigma_3}$  and  $\ln z$  are chosen such that  $\arg z \in (-\pi/2, 3\pi/2)$ , the constant matrix  $\widehat{C}_1$  is the identity matrix and the other constant matrices are determined by  $\widehat{C}_1$  and the jump condition.

The solution to the RH problem can be constructed explicitly by using the confluent hypergeometric function

$$\Phi(z) = e^{-\frac{iz}{2}} \begin{bmatrix} e^{-\frac{\pi i (\alpha + \beta)}{2}} \frac{\Gamma(1 + \alpha - \beta)}{\Gamma(1 + 2\alpha)} \phi(\alpha + \beta, 1 + 2\alpha, iz) & -e^{\frac{\pi i (\alpha - \beta)}{2}} \frac{\Gamma(2\alpha)}{\Gamma(\alpha + \beta)} \phi(-\alpha + \beta, 1 - 2\alpha, iz) \\ e^{-\frac{\pi i (\alpha - \beta)}{2}} \frac{\Gamma(1 + \alpha + \beta)}{\Gamma(1 + 2\alpha)} \phi(1 + \alpha + \beta, 1 + 2\alpha, iz) & e^{\frac{\pi i (\alpha + \beta)}{2}} \frac{\Gamma(2\alpha)}{\Gamma(\alpha - \beta)} \phi(1 - \alpha + \beta, 1 - 2\alpha, iz) \\ z^{\alpha \sigma_3} \begin{bmatrix} 1 & \frac{\sin(\pi(\alpha + \beta))}{\sin(2\pi\alpha)} \\ 0 & 1 \end{bmatrix} \end{bmatrix}$$
(A.8)

for  $z \in \Omega_{\Phi,1}$ ,  $\alpha > -1/2$  and  $2\alpha \notin \mathbb{N}$ ; see [20, 39]. Here,  $\phi(a, b, z)$  is the confluent hypergeometric function defined in (1.4). The expression of  $\Phi(z)$  in the other regions is then determined by using (A.8) and the jump condition (A.4). The case  $2\alpha \in \mathbb{N}$  can be constructed in a similar way and particularly the first column of  $\Phi(z)$  is given by the same formula in (A.8).

### Acknowledgements

The work of Shuai-Xia Xu was supported in part by the National Natural Science Foundation of China under grant numbers 11971492 and 12371257, and by Guangdong Basic and Applied Basic Research Foundation (Grant No. 2022B1515020063). Yu-Qiu Zhao was supported in part by the National Natural Science Foundation of China under grant numbers 11971489 and 12371077.

### References

- G. Akemann, J. Baik and P. Di Francesco, *The Oxford Handbook of Random Matrix Theory*, Oxford University Press, Oxford, 2011.
- [2] A. Bogatskiy, T. Claeys and A. Its, Hankel determinant and orthogonal polynomials for a Gaussian weight with a discontinuity at the edge, *Comm. Math. Phys.*, 347 (2016), 127–162.
- [3] O. Bohigas and M.P. Pato, Missing levels in correlated spectra, *Phys. Lett. B*, 595 (2004), 171–176.
- [4] A. Borodin and P. Deift, Fredholm determinants, Jimbo-Miwa-Ueno  $\tau$ -functions, and representation theory, *Comm. Pure Appl. Math.*, **223** (2002), 1160–1230.
- [5] A. Borodin and G. Olshanski, Infinite random matrices and ergodic measures, Comm. Pure Appl. Math., 223 (2001), 87–123.
- [6] T. Bothner, A. Its, A. Prokhorov, The analysis of incomplete spectra in random matrix theory through an extension of the Jimbo-Miwa-Ueno differential, Adv. Math., 345 (2019), 483–551.
- [7] T. Bothner and A. Its, Asymptotics of a Fredholm determinant corresponding to the first bulk critical universality class in random matrix models. *Comm. Math. Phys.*, **328** (2014), 155–202.
- [8] M. Cafasso, T. Claeys and M. Girotti, Fredholm determinant solutions of the Painlevé II hierarchy and gap probabilities of determinantal point processes, *Int. Math. Res. Not.*, 2021 (2021), 2437–2478.
- [9] C. Charlier, Asymptotics of Hankel determinants with a one-cut regular potential and Fisher-Hartwig singularities, *Int. Math. Res. Not.*, **2019**, 7515–7576.
- [10] C. Charlier, Exponential moments and piecewise thinning for the Bessel point process. Int. Math. Res. Not., 2021 (2021), 16009–16073.
- [11] C. Charlier, Large gap asymptotics for the generating function of the sine point process, P. Lond. Math. Soc., 123 (2021), 103–152.

- [12] C. Charlier and T. Claeys, Large gap asymptotics for Airy kernel determinants with discontinuities, Comm. Math. Phys., 375 (2020), 1299–1339.
- [13] C. Charlier and A. Deaño, Asymptotics for Hankel determinants associated to a Hermite weight with a varying discontinuity, SIGMA Symmetry Integrability Geom. Methods Appl., 14 (2018), 018.
- [14] C. Charlier and A. Doeraene, The generating function for the Bessel point process and a system of coupled Painlevé V equations, *Random Matrices Theory Appl.*, 8 (2019), 1950008.
- [15] C. Charlier, B. Fahs, C. Webb and M.-D. Wong, Asymptotics of Hankel determinants with a multi-cut regular potential and Fisher-Hartwig singularities. arXiv:2111.08395.
- [16] C. Charlier and R. Gharakhloo, Asymptotics of Hankel determinants with a Laguerre-type or Jacobi-type potential and Fisher-Hartwig singularities, Adv. Math., 383 (2021), 107672.
- [17] C. Charlier and P. Moreillon, On the generating function of the Pearcey process, Ann. Appl. Probab., 33 (2023), 3240–3277.
- [18] T. Claeys and A. Doeraene, The generating function for the Airy point process and a system of coupled Painlevé II equations, Stud. Appl. Math., 140 (2018), 403–437.
- [19] T. Claeys and B. Fahs, Random matrices with merging singularities and the Painlevé V equation, SIGMA Symmetry Integrability Geom. Methods Appl., 12 (2016), 031.
- [20] T. Claeys, A. Its and I. Krasovsky, Emergence of a singularity for Toeplitz determinants and Painlevé V, Duke Math. J., 160 (2011), 207–262.
- [21] T. Claeys and I. Krasovsky, Toeplitz determinants with merging singularities, Duke Math. J., 164 (2015), 2897–2987.
- [22] D. Dai, S.-X. Xu and L. Zhang, Asymptotics of Fredholm determinant associated with the Pearcey kernel, *Comm. Math. Phys.*, **382** (2021), 1769–1809.
- [23] D. Dai, S.-X. Xu and L. Zhang, On the deformed Pearcey determinant, Adv. Math., 400 (2022), 108291.
- [24] D. Dai, S.-X. Xu and L. Zhang, Gap probability for the hard edge Pearcey process, Ann. Henri Poincaré, 24 (2023), 2067–2136.
- [25] D. Dai and Y. Zhai, Asymptotics of the deformed Fredholm determinant of the confluent hypergeometric kernel, Stud. Appl. Math., 49 (2022), 1032–1085.
- [26] P. Deift and X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation, Ann. Math., 137 (1993), 295–368.
- [27] P. Deift, Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach, Courant Lecture Notes 3, New York University, New York, 1999.
- [28] P. Deift, A. Its and I. Krasovsky, Asymptotics of Toeplitz, Hankel, and Toeplitz+Hankel determinants with Fisher-Hartwig singularities, Ann. Math., 174 (2011), 1243–1299.

- [29] P. Deift, A. Its and I. Krasovsky, On the asymptotics of a Toeplitz determinant with singularities, *Random Matrix Theory, Interacting Particle Systems, and Integrable Systems*, 93–146, Math. Sci. Res. Inst. Publ., **65**, Cambridge University Press, New York, 2014.
- [30] P. Deift, I. Krasovsky and J. Vasilevska, Asymptotics for a determinant with a confluent hypergeometric kernel, Int. Math. Res. Not., 2011 (2011), 2117–2160.
- [31] P. Deift, A. Its and X. Zhou, A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics, Ann. Math., 146 (1997), 149–235.
- [32] B. Fahs, Uniform asymptotics of Toeplitz determinants with Fisher-Hartwig singularities, Comm. Math. Phys., 383 (2021), 685–730.
- [33] M.E. Fisher and R.E. Hartwig, Toeplitz determinants: Some applications, theorems, and conjectures, Advan. Chem. Phys., 15 (1968), 333–353.
- [34] P. J. Forrester, *Log-gases and Random Matrices (LMS-34)*, Princeton University Press, Princeton, 2010.
- [35] P.J. Forrester and N.S. Witte, Application of the τ-function theory of Painlevé equations to random matrices: PIV, PII and the GUE, Comm. Math. Phys., 219 (2001), 357–398.
- [36] P.J. Forrester and N.S. Witte, Application of the  $\tau$ -function theory of Painlevé equations to random matrices: PV, PIII, the LUE, JUE and CUE, *Comm. Pure Appl. Math.*, **55** (2002), 679–727.
- [37] A. Foulquié Moreno, A. Martínez-Finkelshtein and V.L. Sousa, Asymptotics of orthogonal polynomials for a weight with a jump on [-1, 1], *Constr. Approx.*, **33** (2011), 219–263.
- [38] A. Its, A. Izergin, V. Korepin, and N. Slavnov, Differential equations for quantum correlation functions, Int. J. Modern Phys. B, 4 (1990), 1003–1037.
- [39] A. Its, I. Krasovsky, Hankel determinant and orthogonal polynomials for the Gaussian weight with a jump, *Contemp. Math.*, 458 (2008), 215–248.
- [40] A. Its, O. Lisovyy and A.Prokhorov, Monodromy dependence and connection formulae for isomonodromic tau functions, *Duke Math. J.*, 167 (2018), 1347–1432.
- [41] M. Jimbo and T. Miwa, Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. II. Phys. D, 2 (1981), 407–448.
- [42] M. Jimbo, T. Miwa, Y. Môri and M. Sato, Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent, *Phys. D*, 1 (1980), 80–158.
- [43] A.B.J. Kuijlaars and M. Vanlessen, Universality for eigenvalue correlations at the origin of the spectrum, *Comm. Math. Phys.*, 243 (2003), 163–191.
- [44] I. Krasovsky, Correlations of the characteristic polynomials in the Gaussian unitary ensemble or a singular Hankel determinant, Duke Math. J., 139 (2007), 581–619.
- [45] A. Lenard, Some remarks on large Toeplitz determinants, Pacific J. Math., 42 (1972), 137–145.

- [46] P. Le Doussal, S.N. Majumdar and G. Schehr, Multicritical edge statistics for the momenta of fermions in nonharmonic traps, *Phys. Rev. Lett.*, **121** (2018), 030603.
- [47] M.L. Mehta, Random Matrices, 3rd ed., Elsevier/Academic Press, Amsterdam, 2004.
- [48] F.W.J. Olver, A.B. Olde Daalhuis, D.W. Lozier, B.I. Schneider, R.F. Boisvert, C.W. Clark, B.R. Miller, B.V. Saunders, H.S. Cohl, and M.A. McClain, eds., *NIST digital library of mathematical functions*, https://dlmf.nist.gov/, Release 1.1.10 of 2023-06-15.
- [49] C. Tracy and H. Widom, Level spacing distributions and the Airy kernel, Comm. Math. Phys., 159 (1994), 151–174.
- [50] C. Tracy and H. Widom, Level spacing distributions and the Bessel kernel, Comm. Math. Phys., 161 (1994), 289–309.
- [51] H. Widom. Toeplitz determinants with singular generating functions. Amer. J. Math., 95 (1973), 333–383.
- [52] X.-B. Wu, S.-X. Xu and Y.-Q. Zhao, Gaussian unitary ensemble with boundary spectrum singularity and  $\sigma$ -form of the Painlevé II equation, *Stud. Appl. Math.*, **140** (2018), 221–251.
- [53] J. Xia, Y.-F. Hao, S.-X. Xu, L. Zhang and Y.-Q. Zhao, Asymptotics of the deformed higher order Airy-kernel determinant and applications, *Nonlinearity*, 36 (2023), 4384–4424.
- [54] S.-X. Xu and D. Dai, Tracy-Widom distributions in critical unitary random matrix ensembles and the coupled Painlevé II system, Comm. Math. Phys., 365 (2019), 515–567.
- [55] S.-X. Xu and Y.-Q. Zhao, Gap probability of the circular unitary ensemble with a Fisher-Hartwig singularity and the coupled Painlevé V system, *Comm. Math. Phys.*, **337** (2020), 1545–1596.