

# Two Algorithms for Computing Rational Univariate Representations of Zero-Dimensional Ideals with Parameters

Dingkang Wang<sup>a,b</sup>, Jingjing Wei<sup>c</sup>, Fanghui Xiao<sup>d</sup>, Xiaopeng Zheng<sup>a,b</sup>

<sup>a</sup>KLMM, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China

<sup>b</sup>School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>c</sup>Beijing ETown Academy, Beijing 100176, China

<sup>d</sup>MOE-LCSM, School of Mathematics and Statistics, Hunan Normal University, Hunan 410081, China

---

## Abstract

Based on the partition of parameter space, two algorithms for computing the rational univariate representation of zero-dimensional ideals with parameters are presented in the paper. Unlike the rational univariate representation of zero-dimensional ideals without parameters, the number of zeros of zero-dimensional ideals with parameters under various specializations is different, which leads to choosing and checking the separating element, the key to computing the rational univariate representation, is difficult. In order to pick out the separating element, we first ensure that under each branch the ideal has the same number of zeros by partitioning the parameter space. Subsequently two ideas are given to choose and check the separating element. One idea is that by extending the subresultant theorem to parametric cases, we utilize the extended subresultant theorem to choose the separating element with the further partition of parameter space and then with the help of parametric greatest common divisor theory compute rational univariate representations. Another one is that we go straight to choose and check the separating element by the computation of parametric greatest common divisors, then immediately get the rational univariate representations. Based on these, we design two different algorithms for computing rational univariate representations of zero-dimensional ideals with parameters. Furthermore, the algorithms have been implemented on *Singular* and the performance comparison are presented.

*Keywords:* Rational univariate representation, Parametric zero-dimensional ideal, Comprehensive Gröbner system

---

## 1. Introduction

Solving multivariate polynomial equations has always been a classical algebraic problem, which plays an important role in many fields. The rapid development of computer technology makes the function of computer algebra system perfect. Thus, it provides a new opportunity for the solution of polynomial equations. Many new methods for solving polynomial equations have emerged in computer algebra and algebraic geometry, such as Wu's method (Wu, 1984), Gröbner basis method (Becker and Weispfenning, 1993), resultant-based method (Cox et al., 2005) and

---

*Email addresses:* dwang@mmrc.iss.ac.cn (Dingkang Wang), weijingjing18@mailsucas.ac.cn (Jingjing Wei), xiaofanghui@hnnu.edu.cn (Fanghui Xiao), zhengxiaopeng@amss.ac.cn (Xiaopeng Zheng)

*Preprint submitted to ...*

July 25, 2024

eigenvalue methods (Auzinger and Stetter, 1988; Stetter, 1996). These new theories and methods inject new vitality into the solution of polynomial equations, making them more and more widely used in computer-aided design, computer vision, cybernetics, robot trajectory design, curve and surface design and modeling.

Rouillier (1999) proposed the rational univariate representation (RUR) to solve zero-dimensional polynomial systems and presented an efficient algorithm for computing rational univariate representations. The rational univariate representation consists in expressing all the coordinates of the roots for zero-dimensional ideal  $I \subset k[x_1, \dots, x_n]$  (where  $k$  is a field of characteristic 0) as rational functions of the roots of an univariate polynomial. That is, the roots of  $I$  can be represented in the following way:

$$\mathbb{V}_L(I) = \left\{ \left( \frac{g_1(\beta)}{g(\beta)}, \dots, \frac{g_n(\beta)}{g(\beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}(T)) \right\},$$

where  $\mathcal{X}, g, g_1, \dots, g_n \in k[T]$  are univariate polynomials,  $L$  is an algebraic closure of  $k$ ,  $\mathbb{V}_L(I)$  is the variety of  $I$  in  $L^n$ . Since then, it has been extensively studied. Noro and Yokoyama (1999) used modular method to compute the rational univariate representation of zero-dimensional ideal. Ouchi and Keyser (2008) based on toric resultants gave the computation of the rational univariate representation. Tan and Zhang (2009) presented an improved algorithm for finding separating elements of zero-dimensional ideals. Zeng and Xiao (2010) proposed a method for computing the rational univariate representation by Wu's method. Ma et al. (2012) presented an approach to compute the rational univariate representation via properties of Gröbner basis.

Tan (2009); Tan and Zhang (2010) generalized the rational univariate representation theory to high-dimensional polynomial systems and proposed the rational representation theory. Along this, Shang et al. (2017) proposed a simplified rational representation and Xiao et al. (2021) presented an improvement of the rational representation by introducing minimal Dickson basis proposed by Kapur et al. (2013). Similar to rational representation, Schost (2003) proposed parametric geometric resolution and Safey El Din et al. (2018) used rational parametrizations to represent all irreducible components of real algebraic sets.

Although the rational representation of polynomial systems has formed a relatively perfect theory, the research on the rational representation of polynomial systems with parameters as the method of solving parametric multivariate polynomial equations is still blank. In this paper, we extend the rational univariate representation of zero-dimensional ideals to the parametric case. Given an ideal  $I = \langle f_1(U, X), \dots, f_s(U, X) \rangle \subset k[U, X]$  with variables  $X$  and parameters  $U$ . The considered problem is to find finite sets  $E_i, N_i \subset k[U]$ ,  $i \in \{1, \dots, s\}$  such that for all  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $I(\bar{u}) = \langle f_1(\bar{u}, X), \dots, f_s(\bar{u}, X) \rangle \subset L[X]$  is a zero-dimensional ideal and give an RUR of  $I(\bar{u})$  to express the variety:

$$\mathbb{V}_L(I(\bar{u})) = \left\{ \left( \frac{g_{i1}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)}, \dots, \frac{g_{in}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}_i(\bar{u}, T)) \right\},$$

where  $g_i, g_{i1}, \dots, g_{in}, \mathcal{X}_i \in k(U)[T]$ . That is to compute a finite set:

$$\{(E_1, N_1, \mathcal{X}_1, g_1, g_{11}, \dots, g_{1n}), \dots, (E_s, N_s, \mathcal{X}_s, g_s, g_{s1}, \dots, g_{sn})\}.$$

Two ideas based on the partition of parameter space are given. First, by means of comprehensive Gröbner systems (see Weispfenning (1992); Kalkbrener (1997); Montes (2002); Suzuki and Sato (2006); Nabeshima (2007); Kapur et al. (2013)) and Finiteness theorem, we pick out the zero-dimensional branches which satisfy the ideal is zero-dimensional under parametric specializations. Different from the rational univariate representation of zero-dimensional ideals without

parameters, the number of zeros for zero-dimensional ideals with parameters under various specializations is different, which leads to choosing and checking the separating element, the premise and the key to computing the rational univariate representation, is difficult. Thus, the second step is to determine the number of zeros by partitioning of the parameter space such that each branch of parameter space has the same number of zeros. Subsequently, one idea is that we directly use extended subresultant theorem (Mishra, 1993) for parametric cases to choose the separating element corresponding to each branch with the further partition of parameter space and then with the help of parametric greatest common divisor theory (Nagasaka, 2017; Kapur et al., 2018) compute a finite set of which each branch shares the same expression of rational univariate representation. Another one is that we go straight to check the separating elements and get the rational univariate representations by the computation of parametric greatest common divisors, which actually uses the extended subresultant theorem potentially. As a result, we present two algorithms for computing rational univariate representations of zero-dimensional ideals with parameters.

This paper is an extension of Wang et al. (2022b), and new contributions are as follows. 1) Unlike previous study which divides the parameter space by directly utilizing the extended subresultant theorem for parametric cases for choosing the separating element, we give a new idea to choose and check the separating element based on parametric greatest common divisors, then present the different algorithm for computing rational univariate representations of zero-dimensional ideals with parameters. 2) The two algorithms (the algorithm (Wang et al., 2022b) and new algorithm) have been implemented on the computer algebra system *Singular*. Experimental data and their performance comparison are reported in the paper. Moreover, the codes and examples are available on the web: <http://www.mmrc.iss.ac.cn/~dwang/software.html>.

The rest of the paper is organized as follows. In Section 2, we introduce some notations and definitions, and extend the subresultant theorem to parametric cases. The main content is presented in Section 3. We use comprehensive Gröbner systems, the extended subresultant theorem and parametric greatest common divisors to give the rational univariate representation of zero-dimensional ideals with parameters, and two algorithm are presented. In Section 4, we give an example to illustrate the algorithms. The implementation of the algorithms and the performance comparison are presented in Section 5. We end with some concluding remarks in Section 6.

## 2. Preliminaries

In this section we will introduce some notations and definitions to prepare for the discussion of this article.

Let  $k$  be a field of characteristic 0, and  $L$  be its algebraic closure.  $k[X]$  is the polynomial ring over  $k$  in the variables  $X = \{x_1, \dots, x_n\}$  and  $k[U, X]$  is the parametric polynomial ring with the parameters  $U = \{u_1, \dots, u_m\}$  and variables  $X$ .  $I$  is an ideal of  $k[X]$  and  $\mathbb{V}_L(I)$  is the variety of  $I$  in  $L^n$ . For a set  $N \subset k[X]$ ,  $p \in k[X]$ , we have the notation  $N \times p = \{f \cdot p \mid f \in N\}$ .

### 2.1. Rational univariate representation

Let  $I \subset k[X]$  be a zero-dimensional ideal. Since  $I$  is zero-dimensional,  $k[X]/I$  is a linear space over  $k$  by Finiteness theorem. For all  $t \in k[X]$ , we denote by  $m_t$  the  $k$ -linear map:

$$m_t : k[X]/I \rightarrow k[X]/I$$

$$\bar{f} \mapsto \overline{tf},$$

where  $\bar{f}$  denotes the residue class in  $k[X]/I$  of any polynomial  $f \in k[X]$ .

We denote by  $M_t$  the matrix representation of  $m_t$  w.r.t. a basis in quotient ring  $k[X]/I$ . And we call  $m_t$  the multiplication map and  $M_t$  the multiplication matrix.

**Theorem 1 (Rouillier (1999)).** *Let  $I \subset k[X]$  be a zero-dimensional ideal,  $t \in k[X]$  and  $m_t$  be the multiplication map. Then the eigenvalues of  $M_t$  are  $\{t(p) : p \in \mathbb{V}_L(I)\}$ . More specifically, the characteristic polynomial of  $m_t$  is*

$$\mathcal{X}_t(T) = \prod_{p \in \mathbb{V}_L(I)} (T - t(p))^{\mu(p)},$$

where  $\mu(p)$  is the multiplicity of  $p$  in  $\mathbb{V}_L(I)$ .

Similarly, given the polynomial  $h \in k[X]$ , we can construct a bilinear form

$$S_h(f, g) = \text{Tr}(m_{hf} \cdot m_g) = \text{Tr}(m_{hfg}).$$

Suppose that  $B = \{X^{\alpha_1}, \dots, X^{\alpha_s}\}$  is a basis of  $k[X]/I$ . Let  $Q_h$  be the matrix of  $S_h$  w.r.t.  $B$ , i.e.,  $Q_h = (S_h(X^{\alpha_i}, X^{\alpha_j}))_{1 \leq i, j \leq s} \in k^{s \times s}$ .  $Q_h$  is called the Hermite's quadratic form associated  $h$  w.r.t.  $B$  in quotient ring  $k[X]/I$ .

**Theorem 2 (Cox et al. (2006), Page 71, Theorem 5.2).** *Let  $I \subset k[X]$  be a zero-dimensional ideal. Then the number of points in  $\mathbb{V}_L(I)$  is equal to the rank of  $Q_1$ .*

Next, we review the definition of separating elements which plays an important role in rational univariate representation theory.

**Definition 3.** *Let  $I \subset k[X]$  be a zero-dimensional ideal. A polynomial  $t \in k[X]$  separates  $\mathbb{V}_L(I)$ , if*

$$p_1, p_2 \in \mathbb{V}_L(I), p_1 \neq p_2 \Rightarrow t(p_1) \neq t(p_2).$$

We also call  $t$  a separating element of  $I$ .

**Definition 4.** *Let  $I \subset k[X]$  be a zero-dimensional ideal, and  $t \in k[X]$ .  $\mathcal{X}_t$  is the characteristic polynomial of multiplication map  $m_t$ . For any  $v \in k[X]$ , we define:*

$$g(v, T) = \sum_{\alpha \in \mathbb{V}_L(I)} \mu(\alpha)v(\alpha) \prod_{y \neq t(\alpha), y \in \mathbb{V}_L(\mathcal{X}_t)} (T - y).$$

For any  $t \in k[X]$ , the  $t$ -representation of  $I$  is the  $(n + 2)$ -tuple:

$$\{\mathcal{X}_t(T), g(1, T), g(x_1, T), \dots, g(x_n, T)\}.$$

If  $t$  separates  $\mathbb{V}_L(I)$ , the  $t$ -representation of  $I$  is called the **Rational Univariate Representation (RUR)** of  $I$  associated to  $t$ .

In Rouillier (1999), Rouillier proposed an algorithm to compute the RUR of a zero-dimensional ideal.

### Algorithm for Computing the RUR

**Input:**  $I = \langle f_1, \dots, f_l \rangle \subset k[X]$  is a zero-dimensional ideal.

1. Compute the Gröbner basis of  $I$  with a monomial order  $<$  and a basis  $B$  of  $k[X]/I$  over  $k$ .
2. Compute the rank of  $Q_1$  with respect to  $B$ .
3. Set  $d = \text{rank}(Q_1)$  and choose  $t \in S = \{x_1 + ix_2 + \dots + i^{n-1}x_n, i = 1, \dots, nd(d-1)/2\}$  randomly, then compute the characteristic polynomial  $\mathcal{X}_t$  of  $m_t$ .
4. Let  $\overline{\mathcal{X}}_t = \mathcal{X}_t / \gcd(\mathcal{X}'_t, \mathcal{X}_t)$ . If  $\deg(\overline{\mathcal{X}}_t) \neq \text{rank}(Q_1)$ , then goto (3).
5. Suppose that  $\overline{\mathcal{X}}_t = \sum_{j=0}^d a_j T^{d-j}$ . Compute

$$g(T) = \sum_{i=0}^{d-1} \sum_{j=0}^{d-i-1} \text{Tr}(m_{t^i}) a_j T^{d-i-j-1}, \quad (1)$$

$$g_k(T) = \sum_{i=0}^{d-1} \sum_{j=0}^{d-i-1} \text{Tr}(m_{x_k t^i}) a_j T^{d-i-j-1}, k = 1, \dots, n. \quad (2)$$

**Output:**  $\{\mathcal{X}_t(T), g(T), g_1(T), \dots, g_n(T)\} \subset k[T]$ .

By the above algorithm, the variety of  $I$  can be represented as

$$\mathbb{V}_L(I) = \left\{ \left( \frac{g_1(\beta)}{g(\beta)}, \dots, \frac{g_n(\beta)}{g(\beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}_t(T)) \right\}.$$

There are some remarks:

- The Step (4) is aimed to check whether  $t$  is a separating element of  $I$ . According to Definition 3, Theorem 1 and Theorem 2,  $t$  is a separating element of  $I$  if and only if  $\deg(\overline{\mathcal{X}}_t) = \text{rank}(Q_1)$ . In Rouillier (1999), it has been proved that  $S$  contains at least one element that separates  $\mathbb{V}_L(I)$ .
- We can use Formula (1) and (2) to compute  $g(T)$  and  $g_i(T)$  only if  $t$  is a separating element.
- The RUR of  $I$  computed by this algorithm is not unique. It depends on the choice of the separating element.

## 2.2. Comprehensive Gröbner System and Parametric Greatest Common Divisor

In order to deal with the polynomial system with parameters, we will introduce the comprehensive Gröbner system. Fix a monomial order  $<_X$  in  $k[X]$  and monomial order  $<_U$  in  $k[U]$ . Let  $<_{X,U}$  be a block order,  $U \ll X$ , within  $<_U$  and  $<_X$ . For convenience, we often omit writing the order if there is no confusion.

**Definition 5 (Weispfenning (1992)).** Let  $I = \langle f_1, \dots, f_l \rangle \subset k[U, X]$ ,  $S$  be a subset of  $L^m$ ,  $G_i = \{g_{i1}, \dots, g_{ii}\} \subset k[U, X]$  and  $E_i, N_i \subset k[U]$  for  $i = 1, \dots, s$  such that  $S = \bigcup_{i=1}^l \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ . A finite set

$$\mathcal{G} = \{(E_1, N_1, G_1), \dots, (E_s, N_s, G_s)\}$$

is called a comprehensive Gröbner system (CGS) on  $S$  for  $I$  if for every  $i \in \{1, \dots, s\}$ ,  $G_i(\bar{u}) = \{g_{i1}(\bar{u}, X), \dots, g_{ii}(\bar{u}, X)\}$  is a Gröbner basis of  $I(\bar{u}) = \langle f_1(\bar{u}, X), \dots, f_l(\bar{u}, X) \rangle \subset L[X]$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ . Each  $(E_i, N_i, G_i)$  is called a branch of  $\mathcal{G}$ . In particular, if  $S = L^m$ , then  $\mathcal{G}$  is called a comprehensive Gröbner system for  $I$ .

**Definition 6.** A comprehensive Gröbner system  $\mathcal{G} = \{(E_1, N_1, G_1), \dots, (E_s, N_s, G_s)\}$  on  $S$  for  $I$  is said to be minimal, if for each  $i = 1, \dots, s$ ,

1.  $\mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i) \neq \emptyset$ , and furthermore, whenever  $i \neq j$ ,

$$\mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i) \cap \mathbb{V}_L(E_j) \setminus \mathbb{V}_L(N_j) = \emptyset;$$

2.  $G_i(\bar{u})$  is a minimal Gröbner basis for  $I(\bar{u}) \subset L[X]$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ;  
3. for each  $g \in G_i$ , assume that  $c_1 = \text{LC}_X(g)$  is the leading coefficient of  $g$  w.r.t.  $X$ , then  $c_1(\bar{u}) \neq 0$  for any  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

For computing the minimal comprehensive Gröbner system, we can refer to the algorithm proposed by [Kapur et al. \(2013\)](#).

Based on the comprehensive Gröbner system, [Kapur et al. \(2018\)](#) proposed an efficient algorithm for computing the greatest common divisor of polynomials with parameters.

**Definition 7.** For  $F = \{f_1, \dots, f_l\} \subset k[U][X]$  and  $S \subset L^m$ , we call  $\{(E_1, N_1, g_1), \dots, (E_r, N_r, g_r)\}$  a parametric GCD of  $F$  on  $S$ , if for each  $i = 1, \dots, r$ ,  $g_i(\bar{u}, X) \in L[X]$  is a greatest common divisor of  $\{f_1(\bar{u}, X), \dots, f_l(\bar{u}, X)\} \subset L[X]$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ , where  $E_i, N_i \subset k[U]$ ,  $g_i \in k[U][X]$  for  $i = 1, \dots, r$  and  $S = \bigcup_{i=1}^r \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ . If  $S = k^m$ , we simply call it a parametric GCD of  $F$ .

### 2.3. Subresultant and Common Divisor

In the process of computing the RUR, we can see that it is necessary to compute  $\text{gcd}(\mathcal{X}'_t, \mathcal{X}_t)$ . Therefore, if the polynomial system is with parameters, we need to discuss the relationship between  $\text{gcd}(\mathcal{X}'_t, \mathcal{X}_t)$  and the value of parameters. In the following, we will build the connection between the degree of  $\text{gcd}(\mathcal{X}'_t, \mathcal{X}_t)$  and the parameters  $U$  by using the subresultant theory.

First, we review the theorem of the subresultant.

**Theorem 8 (Mishra (1993)).** Let  $S$  be a unique factorization domain with identity, and  $A(x), B(x) \in S[x]$  be univariate polynomials of positive degrees  $m$  and  $n$ , respectively. Then for all  $0 \leq i < \min(m, n)$ , the following three statements are equivalent:

1.  $A(x)$  and  $B(x)$  have a common divisor with degree  $> i$ ;
2.  $(\forall j \leq i) [\text{SubRes}_j(A, B) = 0]$ ;
3.  $(\forall j \leq i) [\text{PSC}_j(A, B) = 0]$ .

where  $\text{SubRes}_j(A, B)$  denotes the  $i$ th subresultant of  $A$  and  $B$ ,  $\text{PSC}_j(A, B)$  denotes the leading coefficient of  $\text{SubRes}_j(A, B)$ .

Now we extend this result to the case for parameters and obtain the relationship between the degree of  $\text{gcd}(\mathcal{X}'_t, \mathcal{X}_t)$  and the values of parameters.

**Lemma 9.** Let  $\mathcal{X}(U, T) = \sum_{i=0}^l c_i(U)T^i \in k[U, T]$  and  $E, N \subset k[U]$ . Suppose that for  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,  $c_l(\bar{u}) \neq 0$ , then there exist  $E_0, \dots, E_{l-1}, N_0, \dots, N_{l-1} \subset k[U]$  such that for each  $0 \leq i \leq l-1$  and for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,

$$\deg(\text{gcd}(\mathcal{X}(\bar{u}, T), \mathcal{X}'(\bar{u}, T))) = i \iff \bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i).$$

where  $\mathcal{X}' \in k[U, T]$  is the derivation of  $\mathcal{X}$  w.r.t.  $T$ .

*Proof.* We regard  $\mathcal{X}$  and  $\mathcal{X}'$  as polynomials in  $S[T]$ , where  $S = k[U]$ . Then  $\text{PSC}_j(\mathcal{X}, \mathcal{X}') \in k[U]$ . For  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,  $c_l(\bar{u}) \neq 0$ , so

$$\deg_T(\mathcal{X}(U, T)) = \deg(\mathcal{X}(\bar{u}, T)) \text{ and } \deg_T(\mathcal{X}'(U, T)) = \deg(\mathcal{X}'(\bar{u}, T)).$$

It implies that the degree of  $\mathcal{X}(U, T)$  and  $\mathcal{X}'(U, T)$  w.r.t.  $T$  does not change under  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . By the definition, for  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,

$$\text{PSC}_j(\mathcal{X}, \mathcal{X}')(\bar{u}) = \text{PSC}_j(\mathcal{X}(\bar{u}, T), \mathcal{X}'(\bar{u}, T)), j = 0, 1, \dots, l-1.$$

Let  $\lambda = l-1$  and

$$E_i = E \cup \{\text{PSC}_j(\mathcal{X}, \mathcal{X}') \mid j < i\}, 0 \leq i \leq \lambda,$$

$$N_i = N \times \{\text{PSC}_i(\mathcal{X}, \mathcal{X}')\}, 0 \leq i < \lambda, \text{ and } N_\lambda = N,$$

where  $N \times \{f\} = \{gf \mid g \in N\}$ . By Theorem 8, for each  $0 \leq i < \lambda$  and  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ , the following statements are equivalent:

1.  $\deg(\gcd(\mathcal{X}(\bar{u}, T), \mathcal{X}'(\bar{u}, T))) = i$ ;
2.  $(\forall j < i) [\text{PSC}_j(\mathcal{X}(\bar{u}, T), \mathcal{X}'(\bar{u}, T)) = 0]$  and  $\text{PSC}_i(\mathcal{X}(\bar{u}, T), \mathcal{X}'(\bar{u}, T)) \neq 0$ ;
3.  $(\forall j < i) [\text{PSC}_j(\mathcal{X}, \mathcal{X}')(\bar{u}) = 0]$  and  $\text{PSC}_i(\mathcal{X}, \mathcal{X}')(\bar{u}) \neq 0$ ;
4.  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

Moreover, it is easy to verify that the conclusion is also true when  $i = \lambda$ . □

### 3. RUR of zero-dimensional ideal with parameters

In the section, we will consider the RUR of zero-dimensional ideal with parameters  $U$ . Let  $I = \langle f_1(U, X), \dots, f_l(U, X) \rangle \in k[U, X]$  and  $E, N \subset k[U]$ . For  $f \in k[U]$ ,  $N \times \{f\} = \{gf \mid g \in N\}$ . For every  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ , the ideal generated by  $f_1(\bar{u}, X), \dots, f_l(\bar{u}, X)$  is denoted by  $I(\bar{u})$ . We want to find finite sets  $E_i, N_i \subset k[U]$ ,  $i \in \{1, \dots, s\}$  such that for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $I(\bar{u})$  is a zero-dimensional ideal and give an RUR of  $I(\bar{u})$  to express

$$\mathbb{V}_L(I(\bar{u})) = \left\{ \left( \frac{g_{i1}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)}, \dots, \frac{g_{im}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}_i(\bar{u}, T)) \right\},$$

where  $g_i, g_{i1}, \dots, g_{im}, \mathcal{X}_i \in k(U)[T]$ .

The ideas are based on the partition of parameter space. The first idea contains four steps and another contains three steps, among which the first two steps are same.

The first idea is as follows.

**Step 1.** Pick zero-dimensional branches. That is to compute  $E_i, N_i \subset k[U]$ ,  $i = 1, \dots, s$ , such that  $I(\bar{u})$  is a zero-dimensional ideal if and only if  $\bar{u}$  belongs to some  $\mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

**Step 2.** Determine the number of zeros. For each  $E_i, N_i$  in Step 1, compute  $E_{ij}, N_{ij} \subset k[U]$  such that  $\mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i) = \bigcup_{j=1}^{s_1} \mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij})$  and for each  $j \in \{1, \dots, s_1\}$ ,  $I(\bar{u})$  has the same number of zeros for all  $\bar{u} \in \mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij})$ .

**Step 3.** Choose and check the separating element. For each  $E_{ij}, N_{ij}$  in Step 2, compute  $E_{ijk}, N_{ijk} \subset k[U]$  such that  $\mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij}) = \bigcup_{k=1}^{s_2} \mathbb{V}_L(E_{ijk}) \setminus \mathbb{V}_L(N_{ijk})$  and for each  $k \in \{1, \dots, s_2\}$ ,  $I(\bar{u})$  shares the same separating element for all  $\bar{u} \in \mathbb{V}_L(E_{ijk}) \setminus \mathbb{V}_L(N_{ijk})$ .

**Step 4.** Give the RUR for each branch. For each  $E_{ijk}, N_{ijk}$  in Step 3, compute  $E_{ijkl}, N_{ijkl} \subset k[U]$  such that  $\mathbb{V}_L(E_{ijk}) \setminus \mathbb{V}_L(N_{ijk}) = \bigcup_{l=1}^{s_3} \mathbb{V}_L(E_{ijkl}) \setminus \mathbb{V}_L(N_{ijkl})$  and for each  $l \in \{1, \dots, s_3\}$ , for all  $\bar{u} \in \mathbb{V}_L(E_{ijkl}) \setminus \mathbb{V}_L(N_{ijkl})$ ,  $I(\bar{u})$  shares the same expression of rational univariate representation.

The second idea is to skip the third step and go straight to check the separating elements and get the RURs by computing GCDs of  $\mathcal{X}_i$  and  $\mathcal{X}'_i$ . That is to say,

**Step 3<sup>\*</sup>.** Choose the separating element and compute GCDs of  $\mathcal{X}_r$  and  $\mathcal{X}'_r$ . For each  $E_{ij}, N_{ij}$  in Step 2, compute  $E_{ijk}, N_{ijk} \subset k[U]$  such that  $\mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij}) = \bigcup_{k=1}^{s_2} \mathbb{V}_L(E_{ijk}) \setminus \mathbb{V}_L(N_{ijk})$ , and for each  $k \in \{1, \dots, s_2\}$ , the GCD of  $\mathcal{X}_r$  and  $\mathcal{X}'_r$  has the same expression, and  $I(\bar{u})$  shares the same separating element as well as the same expression of rational univariate representation for all  $\bar{u} \in \mathbb{V}_L(E_{ijk}) \setminus \mathbb{V}_L(N_{ijk})$ .

### 3.1. Algorithm 1

Based on four steps described above, we can design the first algorithm to compute RURs of zero-dimensional ideal with parameters. Next, we will explain these steps in detail, and then present the algorithm.

#### 3.1.1. Partition according to the Gröbner basis

In fact, Step 1 is easy to be achieved by comprehensive Gröbner system and Finiteness Theorem.

**Theorem 10** (Finiteness Theorem [Cox et al. \(2006\)](#)). *Let  $I \subset k[X]$  be an ideal. Then the following conditions are equivalent:*

1. *The algebra  $A = k[x_1, \dots, x_n]/I$  is finite-dimensional over  $k$ .*
2. *The variety  $\mathbb{V}_L(I) \subset L^n$  is a finite set.*
3. *If  $G$  is a Gröbner basis for  $I$ , then for each  $1 \leq i \leq n$ , there is an  $m_i \geq 0$  such that  $x_i^{m_i} = \text{LM}(g)$  for some  $g \in G$ , where  $\text{LM}(g)$  is the leading monomial of  $g$ .*

An ideal satisfying any of the above conditions is said to be zero-dimensional.

**Theorem 11.** *Let  $I = \langle f_1(U, X), \dots, f_l(U, X) \rangle \subset k[U, X]$ . There exists a finite set  $\{(E_1, N_1, G_1), \dots, (E_s, N_s, G_s)\}$ , such that:*

1.  *$I(\bar{u})$  is a zero-dimensional ideal if and only if*

$$\bar{u} \in \bigcup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i).$$

2. *For  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $G_i(\bar{u}) = \{g_{i1}(\bar{u}, X), \dots, g_{ie}(\bar{u}, X)\}$  is a Gröbner basis of  $I(\bar{u})$ .*

*Proof.* We only need to compute the minimal comprehensive Gröbner system  $\mathcal{G} = \{(E_1, N_1, G_1), \dots, (E_{s_0}, N_{s_0}, G_{s_0})\}$  for  $I$ . Choose the branches  $\{(E_{i_1}, N_{i_1}, G_{i_1}), \dots, (E_{i_l}, N_{i_l}, G_{i_l})\}$  which satisfy for each  $1 \leq j \leq n$ ,  $k[x_j] \cap \text{LM}_X(G_{i_v}) \neq \emptyset$  ( $v = 1, \dots, l$ ). By Finiteness Theorem and the definition of minimal comprehensive Gröbner system,  $\{(E_{i_1}, N_{i_1}, G_{i_1}), \dots, (E_{i_l}, N_{i_l}, G_{i_l})\}$  satisfies the conclusion.  $\square$

**Remark 12.** *For each branch  $(E_i, N_i, G_i)$  in Theorem 11, and for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,*

$$\text{LM}(G_i(\bar{u})) = \{\text{LM}(g_{i1}(\bar{u}, X)), \dots, \text{LM}(g_{ie}(\bar{u}, X))\}.$$

*Therefore, for all  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $L[X]/I(\bar{u})$  shares the same basis:*

$$B_i = \{X^{\alpha_1}, \dots, X^{\alpha_r} \mid X^{\alpha_j} \notin \langle \text{LM}_X(G_i) \rangle, j = 1, \dots, r\},$$

*where  $\text{LM}_X(G)$  is a set of the leading monomials of all elements in  $G$  w.r.t.  $X$ .*



### 3.1.2. Partition according to the number of zeros

Actually, Step 2 is a preparation for Step 3. For non-parametric cases,  $t \in k[X]$  is a separating element of  $I$  if and only if  $\deg(\overline{X}_t) = \text{rank}(Q_1)$ , where  $\overline{X}_t = X_t / \gcd(X'_t, X_t)$ . As for parametric cases,  $Q_1$  and  $X_t$  are both with parameters. Therefore, choosing a  $t \in k[X]$ , it should discuss when  $\deg(\overline{X}_t) = \text{rank}(Q_1)$  hold with the change of the parameters. First, we need to determine  $\text{rank}(Q_1)$ , equally the number of zeros for  $I$ , under parametric specializations.

Let  $(E, N, G)$  be one of the branches obtained in Theorem 11 and  $G = \{g_1, \dots, g_e\} \subset k[U, X]$ . Then  $(E, N, G)$  satisfies that for any  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,

1.  $I(\bar{u})$  is zero-dimensional and  $G(\bar{u})$  is a Gröbner basis of  $I(\bar{u})$ ;
2. for each  $g \in G$ ,  $c_1(\bar{u}) \neq 0$ , where  $c_1$  is the leading coefficient of  $g$  w.r.t.  $X$ .

Assume that  $B = \{X^{\alpha_1}, \dots, X^{\alpha_r}\}$  be a basis of  $L[X]/I(\bar{u})$ . For  $t \in k[X]$ , let  $R_i$  be a remainder of  $t \cdot X^{\alpha_i}$  on division by  $G$  in  $k(U)[X]$ , i.e.,

$$t \cdot X^{\alpha_i} = \sum_{j=1}^e q_{ij} g_j + R_i, \quad i = 1, \dots, r.$$

where  $q_{ij}, R_i \in k(U)[X]$  and their denominators are the products of some factors of the leading coefficient of  $g_j \in G$  w.r.t.  $X$ . So they are not equal to zero under  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . Therefore, we have

$$t \cdot X^{\alpha_i} = \sum_{j=1}^e q_{ij}(\bar{u}, X) g_j(\bar{u}, X) + R_i(\bar{u}, X), \quad i = 1, \dots, r.$$

The remainder of  $t \cdot X^{\alpha_i}$  on division by  $G(\bar{u})$  in  $L[X]$  is equal to  $R_i(\bar{u}, X)$ . Let  $M_t^I \in k(U)^{r \times r}$  be the matrix satisfying

$$(R_1, \dots, R_r) = B \cdot M_t^I,$$

For any  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,

$$(R_1(\bar{u}, X), \dots, R_r(\bar{u}, X)) = B \cdot M_t^I(\bar{u}).$$

Assume that  $M_t^{I(\bar{u})}$  is the multiplication matrix of  $t$  w.r.t.  $B$  in quotient ring  $L[X]/I(\bar{u})$ . Then  $M_t^I(\bar{u}) = M_t^{I(\bar{u})}$ . Therefore, we obtain the following lemma.

**Lemma 13.** *Let  $(E, N, G)$ ,  $B$  and  $M_t^I \in k(U)^{r \times r}$  be as defined above. For any  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,*

1. *The multiplication matrix  $M_t^{I(\bar{u})}$  of  $t$  w.r.t.  $B$  in quotient ring  $L[X]/I(\bar{u})$  is equal to  $M_t^I(\bar{u})$ .*
2. *Assume that the characteristic polynomial of  $M_t^I$  over  $k(U)$  is  $X_t(U, T)$ , then the characteristic polynomial of  $M_t^{I(\bar{u})}$  over  $L$  is  $X_t(\bar{u}, T)$ .*
3. *Let  $h \in k[X]$ ,  $Q_h^I \triangleq (\text{Tr}(M_{hX^{\alpha_i}X^{\alpha_j}}^I))_{1 \leq i, j \leq r}$  be a  $r \times r$  matrix over  $k(U)$ . Then the Hermite's quadratic form associated  $h$  w.r.t.  $B$  in quotient ring  $L[X]/I(\bar{u})$  is equal to  $Q_h^I(\bar{u})$ .*

By the above analysis, first we compute  $Q_1^I \in k(U)^{r \times r}$  defined above. For all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,  $Q_1^{I(\bar{u})} = Q_1^I(\bar{u})$  by Lemma 13 and the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is equal to  $\text{rank}(Q_1^{I(\bar{u})})$  by Theorem 2.

**Theorem 14.** *Let  $(E, N, G)$  be one of the branches in Theorem 11. Then there exists a finite set*

$$\{(E_1, N_1, k_1), \dots, (E_s, N_s, k_s)\}, \quad E_i, N_i \subset k[U], \quad k_i \in \mathbb{Z}_{>0},$$

*such that  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \cup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$  and for each  $i \in \{1, \dots, s\}$  and all  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ , the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is equal to  $k_i$ .*

*Proof.* We first compute  $Q_1^I \in k(U)^{r \times r}$  as defined above. It suffices to discuss  $\text{rank}(Q_1^I(\bar{u}))$  for  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . Let  $D$  is the least common multiple of all denominators of entries in  $Q_1^I$ . Then  $D \in k[U]$  and  $D(\bar{u}) \neq 0$  for any  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . Let  $Q = D \cdot Q_1^I \in k[U]^{r \times r}$ . It suffices to discuss  $\text{rank}(Q(\bar{u}))$ .

Assume that  $Q = (f_{ij})_{1 \leq i, j \leq r}$ . Using the method in [Ma et al. \(2017\)](#), we construct polynomial  $f_i = \sum_{j=1}^r f_{ij} z_j \in k[U][Z]$ ,  $Z = \{z_1, \dots, z_r\}$ ,  $i = 1, \dots, r$ . Compute the minimal comprehensive Gröbner system  $\mathcal{G} = \{(E_1, N_1, G_1), \dots, (E_s, N_s, G_s)\}$  of  $F = \{f_1, \dots, f_r\}$  where  $U$  is the parameters. By Theorem 1 and Theorem 2 in [Ma et al. \(2017\)](#), for any  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ , the rank of  $Q(\bar{u})$  is equal to the number of elements in  $G_i$ . Let  $k_i$  denote the number of elements in  $G_i$ ,  $\{(E_1, N_1, k_1), \dots, (E_s, N_s, k_s)\}$  satisfies the conditions we need.  $\square$

### 3.1.3. Partition according to separating elements

Choosing the separating element is the key to obtain the rational univariate representation. In the following, we will analyze each branch in Theorem 14. Assume that for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ , the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is the same, denoted by  $k_0$ .

If  $k_0 = 1$ , we can choose  $t = 1$  which is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . Otherwise, choose randomly a  $t \in \{x_1 + ix_2 + \dots + i^{n-1}x_n, i = 1 \dots nk_0(k_0 - 1)/2\}$ .  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$  can be divided into two parts,  $\mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $i = 1, 2$ , where

1.  $t$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_1) \setminus \mathbb{V}_L(N_1)$ .
2.  $t$  is not a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_2) \setminus \mathbb{V}_L(N_2)$ .

We will prove in fact the second part is "small enough" in numbers of points. Thus we first find a separating element for the most part of  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . For the small part, we will choose another  $t \in k[X]$ . Sequentially, it does a recursive procedure.

**Theorem 15.** *Let  $I = \langle f_1(U, X), \dots, f_l(U, X) \rangle \subset k[U, X]$ ,  $E, N \subset k[U]$ ,  $G \subset k[U, X]$ . Assume that for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,*

1.  $I(\bar{u})$  is zero-dimensional and  $G(\bar{u})$  is a Gröbner basis of  $I(\bar{u})$ ;
2. for each  $g \in G$ ,  $c_1(\bar{u}) \neq 0$ , where  $c_1$  is the leading coefficient of  $g$  w.r.t.  $X$ ;
3. the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is equal to  $k_0 \neq 1$ .

Suppose that  $t \in k[X]$ ,  $\mathcal{X}_t(U, T) \in k(U)[T]$  is the characteristic polynomial as defined in Lemma 13. Let  $\mathcal{X}(U, T) \in k[U][T]$  (briefly,  $\mathcal{X}$ ) be the product of  $\mathcal{X}_t(U, T)$  and the least common multiple of all denominators of  $\mathcal{X}_t(U, T)$ . Then  $t$  is separating element of  $I(\bar{u})$  if and only if  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N \times \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}) \neq \emptyset$ .

*Proof.* By Definition 3 and Theorem 1, for  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,  $t$  is a separating element of  $I(\bar{u})$  if and only

$$\deg\left(\frac{\mathcal{X}(\bar{u}, T)}{\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))}\right) = k_0,$$

which is equivalent to

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) = \deg(\mathcal{X}(\bar{u}, T)) - k_0.$$

Since  $\deg_T(\mathcal{X}) = \deg(\mathcal{X}(\bar{u}, T))$ , the above equation becomes

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) = \deg_T(\mathcal{X}) - k_0.$$

Let  $d = \deg_T(\mathcal{X}) - k_0$ . By Theorem 1, for any  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,  $\mathcal{X}(\bar{u}, T)$  has at most  $k_0$  distinct roots. It implies that

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) \geq d.$$

Since  $k_0 \neq 1$ ,  $\deg_T(\mathcal{X}') = \deg_T(\mathcal{X}) - 1 > \deg_T(\mathcal{X}) - k_0 = d$ . By Theorem 8 and the proof of Lemma 9,

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) \leq d \iff \text{PSC}_d(\mathcal{X}, \mathcal{X}')(\bar{u}) \neq 0.$$

Let  $N_1 = N \times \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}$ . Then,

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) = d \iff \bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N_1) \neq \emptyset.$$

Thus,  $t$  is separating element of  $I(\bar{u})$  if and only if  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N \times \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}) \neq \emptyset$ .  $\square$

**Remark 16.** 1. For non-parametric cases, choosing  $t$  from  $S$  randomly,  $t$  is a separating element of some zero-dimensional ideal with probability 1. Therefore, it is easy to choose a  $t$  satisfying Theorem 15.

2. For  $E, N \in k[U]$ , we can use the method in [Kapur et al. \(2013\)](#) to check whether  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \emptyset$ .

**Theorem 17.** Let  $I = \langle f_1(U, X), \dots, f_l(U, X) \rangle \subset k[U, X]$ ,  $E, N \subset k[U]$ ,  $G \subset k[U, X]$ . Assume that for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,

1.  $I(\bar{u})$  is zero-dimensional and  $G(\bar{u})$  is a Gröbner basis of  $I(\bar{u})$ ;
2. for each  $g \in G$ ,  $c_1(\bar{u}) \neq 0$ , where  $c_1$  is the leading coefficient of  $g$  w.r.t.  $X$ ;
3. the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is equal to  $k_0$ .

Then there is a finite set

$$\{(E_1, N_1, t_1), \dots, (E_s, N_s, t_s)\}, \quad E_i, N_i \subset k[U], \quad t_i \in k[X],$$

such that  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \cup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$  and for each  $i \in \{1, \dots, s\}$ ,  $t_i$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

*Proof.* If  $k_0 = 1$ , then  $t_1 = 1$ ,  $E_1 = E$ ,  $N_1 = N$  and it is done. When  $k_0 \neq 1$ , we choose  $t \in S = \{x_1 + ix_2 + \dots + i^{n-1}x_n, i = 1, \dots, nk_0(k_0 - 1)/2\}$ . By Theorem 15, if  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N \times \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}) = \emptyset$ , pick another  $t$  from  $S$ ; otherwise, let  $t_1 = t$ ,  $E_1 = E$ ,  $N_1 = N \times \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}$ ,  $E_{10} = E \cup \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}$ ,  $N_{10} = N$ . Then

$$\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \mathbb{V}_L(E_1) \setminus \mathbb{V}_L(N_1) \cup \mathbb{V}_L(E_{10}) \setminus \mathbb{V}_L(N_{10})$$

and  $t_1$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_1) \setminus \mathbb{V}_L(N_1)$ . If  $\mathbb{V}_L(E_{10}) \setminus \mathbb{V}_L(N_{10}) = \emptyset$ , it is done. Otherwise, it implies from  $\mathbb{V}_L(E_1) \setminus \mathbb{V}_L(N_1) \neq \emptyset$  that  $\text{PSC}_d(\mathcal{X}, \mathcal{X}') \notin \langle E \rangle$ . Thus  $\langle E \rangle \subsetneq \langle E_{10} \rangle$ . Continually, using Theorem 15 to  $(E_{10}, N_{10})$  and choosing  $t_2 \in S$ , we have

$$\mathbb{V}_L(E_{10}) \setminus \mathbb{V}_L(N_{10}) = \mathbb{V}_L(E_2) \setminus \mathbb{V}_L(N_2) \cup \mathbb{V}_L(E_{20}) \setminus \mathbb{V}_L(N_{20})$$

and  $t_2$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_2) \setminus \mathbb{V}_L(N_2)$ . If  $\mathbb{V}_L(E_{20}) \setminus \mathbb{V}_L(N_{20}) = \emptyset$ , it is done. Otherwise,  $\langle E_{10} \rangle \subsetneq \langle E_{20} \rangle$ . Go on like this, we get strictly ascending ideal chains:

$$E \subsetneq E_{10} \subsetneq E_{20} \subsetneq E_{30} \subsetneq \dots$$

According to the ascending chains condition, it must stop in finite steps. Thus, we obtain a finite set

$$\{(E_1, N_1, t_1), \dots, (E_s, N_s, t_s)\}, \quad E_i, N_i \subset k[U], \quad t_i \in k[X],$$

such that  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \cup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$  and for each  $i \in \{1, \dots, s\}$ ,  $t_i$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .  $\square$

### 3.1.4. Partitions according to GCDs of $\mathcal{X}_t$ and $\mathcal{X}'_t$

Suppose that  $(E, N, t)$  is one of branches in Theorem 17. That is,  $t$  is a separating element of  $I(\bar{u})$  for any  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ . We know if one obtain the RUR then you have to compute the GCD of  $\mathcal{X}_t$  and  $\mathcal{X}'_t$ . However, for different  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ , the GCD of  $\mathcal{X}_t(\bar{u}, T)$  and  $\mathcal{X}'_t(\bar{u}, T)$  may be different. In order to obtain the RUR with parameters, one needs to use parametric GCD algorithm (Kapur et al., 2018; Wang et al., 2020, 2022a).  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$  will be divided into finite branches. In each branch the GCD of  $\mathcal{X}_t$  and  $\mathcal{X}'_t$  has the same expression, then by adjusting Formula (1), (2) in Section 2.1 to  $k(U)[T]$ , we can compute  $g(U, T), g_1(U, T), \dots, g_n(U, T)$  and obtain an RUR for each branch.

**Theorem 18.** Let  $I = \langle f_1(U, X), \dots, f_l(U, X) \rangle \subset k[U, X]$ ,  $E, N \subset k[U]$ ,  $G \subset k[U, X]$ ,  $t \in k[X]$ . Assume that for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,

1.  $I(\bar{u})$  is zero-dimensional and  $G(\bar{u})$  is a Gröbner basis of  $I(\bar{u})$ ;
2. for each  $g \in G$ ,  $c_1(\bar{u}) \neq 0$ , where  $c_1$  is the leading coefficient of  $g$  w.r.t.  $X$ ;
3.  $t$  is a separating element of  $I(\bar{u})$ .

Let  $\mathcal{X}_t(U, T) \in k(U)[T]$  be the characteristic polynomial as defined in Lemma 13. Then there is a finite set

$$\{(E_1, N_1, \mathcal{X}_t, g_1, g_{11}, \dots, g_{1n}), \dots, (E_s, N_s, \mathcal{X}_t, g_s, g_{s1}, \dots, g_{sn})\},$$

such that  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \bigcup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$  and for each  $i \in \{1, \dots, s\}$  and for each  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,

$$\mathbb{V}_L(I(\bar{u})) = \left\{ \left( \frac{g_{i1}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)}, \dots, \frac{g_{in}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}_t(\bar{u}, T)) \right\},$$

where  $E_i, N_i \in k[U]$ ,  $\mathcal{X}_t, g_i, g_{i1}, \dots, g_{in} \in k(U)[T]$ .

*Proof.* Let  $\mathcal{X}(U, T) \in k[U][T]$  be the product of  $\mathcal{X}_t(U, T)$  and the least common multiple of all denominators of  $\mathcal{X}_t(U, T)$ . By the parametric GCD algorithm, we can obtain the parametric GCDs of  $\mathcal{X}(U, T)$  and  $\mathcal{X}'(U, T)$  on  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$  and assume that it is

$$\{(E_1, N_1, d_1), \dots, (E_s, N_s, d_s)\},$$

where  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \bigcup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $d_1, \dots, d_s \in k[U][T]$  and for every  $i = 1, \dots, s$ ,  $d_i(\bar{u}, T)$  is a greatest common divisor of  $\mathcal{X}(\bar{u}, T)$  and  $\mathcal{X}'(\bar{u}, T)$  for any  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ . Moreover, the leading coefficient of  $d_i$  w.r.t.  $T$  is not zero under  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ . Therefore, we can divide  $d_i$  by the leading coefficient of  $d_i$  and make  $d_i \in k(U)[T]$  monic. For  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ , we have

$$\gcd(\mathcal{X}_t(\bar{u}, T), \mathcal{X}'_t(\bar{u}, T)) = \gcd(\mathcal{X}(\bar{u}, T), \mathcal{X}'(\bar{u}, T)) = d_i(\bar{u}, T).$$

For each  $i = 1, \dots, s$ , using divide algorithm in  $k(U)[T]$ , we get

$$\mathcal{X}_t(U, T) = q_i(U, T)d_i(U, T) + r_i(U, T), \quad \deg_T(r_i) < \deg_T(d_i),$$

Since  $d_i(\bar{u}, T)$  divides  $\mathcal{X}_t(\bar{u}, T)$ , for all  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $r_i(U, T) = 0$ . Therefore,

$$\frac{\mathcal{X}_t(\bar{u}, T)}{\gcd(\mathcal{X}'_t(\bar{u}, T), \mathcal{X}_t(\bar{u}, T))} = q_i(\bar{u}, T).$$

Let  $\overline{\mathcal{X}}_i(U, T) = q_i(U, T) = \sum_{j=0}^d a_j T^{d-j} \in k(U)[T]$ . Compute

$$g_i(U, T) = \sum_{i=0}^{d-1} \sum_{j=0}^{d-i-1} \text{Tr}(M_{\mu_i}^T) a_j T^{d-i-j-1}, \quad (3)$$

$$g_{ij}(U, T) = \sum_{i=0}^{d-1} \sum_{j=0}^{d-i-1} \text{Tr}(M_{x_k^i}^T) a_j T^{d-i-j-1}, j = 1, \dots, n. \quad (4)$$

Then

$$\{\mathcal{X}_i(\bar{u}, T), g_i(\bar{u}, T), g_{i1}(\bar{u}, T), \dots, g_{in}(\bar{u}, T)\}$$

is an RUR of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .  $\square$

We are now ready to give the first algorithm for computing the rational univariate representations with parameters of an ideal in  $k[U, X]$ , where we deliberately avoid tricks and optimizations, such as the selection of separating elements, reducing  $\mathcal{X}$  or  $g_i$  by  $E_i$ .

### Algorithm 1 for computing RURs with parameters

**Input:**  $I = \langle f_1, \dots, f_r \rangle \subset k[U, X]$ .

1. Compute a comprehensive Gröbner system of  $I$  with  $\prec_{X,U}$ . Choose the zero-dimensional branches by Theorem 11 and denote them by  $\mathcal{G} = \{(E_1, N_1, G_1), \dots, (E_s, N_s, G_s)\}$ .
2.  $B_2 := \emptyset$ . For each branch  $(E_i, N_i, G_i) \in \mathcal{G}$  do:
  - (a) Compute  $Q_1^i = (p_{ij}/q_{ij}) \in k(U)^{r \times r}$  defined in Lemma 13. Let  $Q = D \cdot Q_1^i$ , where  $D = \text{lcm}\{q_{ij} : i, j = 1, \dots, r\}$ .
  - (b) Assume that  $Q = (f_{ij})_{1 \leq i, j \leq r}$ . Construct polynomial  $f_i = \sum_{j=1}^r f_{ij} z_j$ . Compute the minimal Comprehensive Gröbner system  $\mathcal{G} = \{(E_{i1}, N_{i1}, G_{i1}), \dots, (E_{is_i}, N_{is_i}, G_{is_i})\}$  of  $\{f_1, \dots, f_r\}$ . Set  $A_i := \{(E_{i1}, N_{i1}, k_{i1}), \dots, (E_{is_i}, N_{is_i}, k_{is_i})\}$ , where  $k_{ij}$  is equal to number of element in  $G_{ij}$ ,  $j = 1, \dots, s_i$ .
  - (c)  $B_2 := B_2 \cup A_i$ .
3.  $B_3 := \emptyset$ . For each branch  $(E_{ij}, N_{ij}, k_{ij}) \in B_2$  do:
  - (a) If  $k_{ij} = 1$  then  $A_{ij} := \{(E_{ij}, N_{ij}, 1)\}$  and goto (3.d); else  $A_{ij} := \emptyset$ ,  $S := \{x_1 + ix_2 + \dots + i^{n-1} x_n, i = 1, \dots, nk_{ij}(k_{ij} - 1)/2\}$ .
  - (b) Randomly choose  $t \in S$ . Compute  $M_t^i \in k(U)^{r \times r}$ ,  $\mathcal{X}_t \in k(U)[T]$ . Let  $\mathcal{X} = D' \cdot \mathcal{X}_t$ . Compute  $\text{PSC}_d(\mathcal{X}, \mathcal{X}')$ , where  $D'$  is the least common multiple of all denominators of  $\mathcal{X}_t$  and  $d = \deg_T(\mathcal{X}) - k_{ij}$ .
  - (c) If  $\mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij} \times \text{PSC}_d(\mathcal{X}, \mathcal{X}')) = \emptyset$ , then  $S := S \setminus \{t\}$  and goto (3.b); else,  $A_{ij} := A_{ij} \cup \{(E_{ij}, N_{ij} \times \text{PSC}_d(\mathcal{X}, \mathcal{X}'), t)\}$ . Update  $E_{ij} := E_{ij} \cup \{\text{PSC}_d(\mathcal{X}, \mathcal{X}')\}$  and  $S := S \setminus \{t\}$ . If  $\mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij}) \neq \emptyset$ , then goto (3.b).
  - (d)  $B_3 := B_3 \cup A_{ij}$ .
4.  $B_4 := \emptyset$ . For each branch  $(E_{ijk}, N_{ijk}, t_{ijk}) \in B_3$  do:
  - (a) Compute parametric GCDs of  $\mathcal{X}$  and  $\mathcal{X}'$ , denoted by

$$\{(E_{ijkl}, N_{ijkl}, d_{ijkl}) : l = 1, \dots, v\},$$

where  $\mathcal{X}_t \in k(U)[T]$  and  $\mathcal{X} \in k[U][T]$  are the polynomials computed in (3.b) corresponding to  $t_{ijk}$ .

- (b) For each  $(E_{ijkl}, N_{ijkl}, d_{ijkl})$  do:  
 $A_{ijkl} := \emptyset$ . Compute  $\overline{\mathcal{X}}_t$ . By Formula (3), (4) compute  $g_{ijkl}(U, T)$ ,  $g_{ijkl1}(U, T)$ ,  $\dots$ ,  $g_{ijkln}(U, T)$ .  $A_{ijkl} := A_{ijkl} \cup \{(E_{ijkl}, N_{ijkl}, \mathcal{X}_t, g_{ijkl}, g_{ijkl1}, \dots, g_{ijkln})\}$   
(c)  $B_4 := B_4 \cup A_{ijkl}$ .

**Output:** a finite set  $B_4$  of the rational univariate representations with parameters, renumbering it by

$$\{(E_1, N_1, \mathcal{X}_1, g_1, g_{11}, \dots, g_{1n}), \dots, (E_s, N_s, \mathcal{X}_s, g_s, g_{s1}, \dots, g_{sn})\}.$$

For each  $i \in \{1, \dots, s\}$ , it satisfies that : for all  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,

$$\mathbb{V}_L(I(\bar{u})) = \left\{ \left( \frac{g_{i1}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)}, \dots, \frac{g_{in}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}_i(\bar{u}, T)) \right\},$$

where  $I(\bar{u}) = \langle f_1(\bar{u}, X), \dots, f_l(\bar{u}, X) \rangle \subset k[X]$ ,  $\mathcal{X}_i(\bar{u}, T) \in k[T]$ .

**Theorem 19.** *The above algorithm works correctly and terminates in a finite number of steps.*

*Proof.* The correctness and the termination of the algorithm directly follows from Theorems 11, 14, 17 and 18.  $\square$

### 3.2. Algorithm 2

Based on three steps of the second idea, we can design another algorithm different from Algorithm 1. Since the first two steps are same, we will only elaborate on Step 3 $^\clubsuit$ .

Assume that for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ , the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is the same, denoted by  $k_0$ . Choose randomly a  $t \in \{x_1 + ix_2 + \dots + i^{n-1}x_n, i = 1, \dots, nk_0(k_0 - 1)/2\}$  and compute the characteristic polynomial  $\mathcal{X}_t(U, T) \in k(U)[T]$  as defined in Lemma 13. Let  $\mathcal{X}(U, T) \in k[U][T]$  be the product of  $\mathcal{X}_t(U, T)$  and the least common multiple of all denominators of  $\mathcal{X}_t(U, T)$ . By the parametric GCD algorithm, we can obtain the parametric GCDs of  $\mathcal{X}(U, T)$  and  $\mathcal{X}'(U, T)$  on  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$  and assume that it is

$$\{(E_1, N_1, d_1), \dots, (E_r, N_r, d_r)\},$$

where  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \cup_{i=1}^r \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,  $d_1, \dots, d_r \in k[U][T]$  and for every  $i = 1, \dots, r$ ,  $d_i(\bar{u}, T)$  is a greatest common divisor of  $\mathcal{X}(\bar{u}, T)$  and  $\mathcal{X}'(\bar{u}, T)$  for any  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ . Moreover, the leading coefficient of  $d_i$  w.r.t.  $T$  is not zero under  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

As stated in the proof of Theorem 15, for  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,  $t$  is a separating element of  $I(\bar{u})$  if and only

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) = \deg_T(\mathcal{X}(U, T)) - k_0.$$

Since

$$\deg(\gcd(\mathcal{X}'(\bar{u}, T), \mathcal{X}(\bar{u}, T))) \geq \deg_T(\mathcal{X}(U, T)) - k_0,$$

combining with Remark 16, it implies that there exists a unique  $i \in \{1, \dots, r\}$  such that  $t$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

Then we obtain the following theorem.

**Theorem 20.** *Let  $I = \langle f_1(U, X), \dots, f_l(U, X) \rangle \subset k[U, X]$ ,  $E, N \subset k[U]$ ,  $G \subset k[U, X]$ . Assume that for all  $\bar{u} \in \mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$ ,*

1.  $I(\bar{u})$  is zero-dimensional and  $G(\bar{u})$  is a Gröbner basis of  $I(\bar{u})$ ;

2. for each  $g \in G$ ,  $c_1(\bar{u}) \neq 0$ , where  $c_1$  is the leading coefficient of  $g$  w.r.t.  $X$ ;
3. the number of points in  $\mathbb{V}_L(I(\bar{u}))$  is equal to  $k_0$ .

Then there is a finite set

$$\{(E_1, N_1, t_1, d_1), \dots, (E_s, N_s, t_s, d_s)\}, E_i, N_i \subset k[U], t_i \in k[X], d_i \in k[X][T],$$

such that  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \cup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$  and for each  $i \in \{1, \dots, s\}$ ,  $t_i$  is a separating element of  $I(\bar{u})$  and  $d_i(\bar{u}, T)$  is a greatest common divisor of  $\mathcal{X}(\bar{u}, T)$  and  $\mathcal{X}'(\bar{u}, T)$  for any  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .

*Proof.* We choose  $t \in S = \{x_1 + ix_2 + \dots + i^{n-1}x_n, i = 1, \dots, nk_0(k_0 - 1)/2\}$ . As mentioned above, we can compute the parametric GCDs of  $\mathcal{X}(U, T)$  and  $\mathcal{X}'(U, T)$  on  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N)$  and assume that it is

$$\{(\tilde{E}_1, \tilde{N}_1, \tilde{d}_1), \dots, (\tilde{E}_r, \tilde{N}_r, \tilde{d}_r)\}.$$

If  $\deg(\tilde{d}_i) > \deg_T(\mathcal{X}(U, T)) - k_0$  for every  $i \in \{1, \dots, r\}$ , pick another  $t$  from  $S$ ; otherwise, there exists a unique  $i \in \{1, \dots, r\}$  such that  $\deg(\tilde{d}_i) = \deg_T(\mathcal{X}(U, T)) - k_0$ , then  $t$  is a separating element of  $I(\bar{u})$  for  $\bar{u} \in \mathbb{V}_L(\tilde{E}_i) \setminus \mathbb{V}_L(\tilde{N}_i)$ . Without loss of generality, we assume  $i = 1$ , let  $E_1 = \tilde{E}_1, N_1 = \tilde{N}_1, t_1 = t, d_1 = \tilde{d}_1$ . If  $r = 1$ , it is done. Otherwise, it implies from Theorem 15 that  $\text{PSC}_{\deg_T(\mathcal{X}(U, T)) - k_0}(\mathcal{X}, \mathcal{X}') \in \sqrt{\langle \tilde{E}_i \rangle}$  for  $i = 2, \dots, r$ . Since  $\mathbb{V}_L(E_1) \setminus \mathbb{V}_L(N_1) \neq \emptyset$ ,  $\text{PSC}_{\deg_T(\mathcal{X}(U, T)) - k_0}(\mathcal{X}, \mathcal{X}') \notin \sqrt{\langle E \rangle}$ . Thus  $\sqrt{\langle E \rangle} \subsetneq \sqrt{\langle \tilde{E}_i \rangle}$  for  $i = 2, \dots, r$ . Now, for the remaining  $r - 1$  branches, choosing  $t_2 \in S$  and computing the parametric GCDs of  $\mathcal{X}(U, T)$  and  $\mathcal{X}'(U, T)$  on these branches, we can obtain  $(E_2, N_2, t_2, d_2)$ . Go on like this, we get strictly ascending ideal chains:

$$E \subsetneq \tilde{E}_i \subsetneq \tilde{E}_{ij} \subsetneq \tilde{E}_{ijk} \subsetneq \dots,$$

where  $i, j, k \neq 1$ . According to the ascending chains condition, it must stop in finite steps. Thus, we obtain a finite set

$$\{(E_1, N_1, t_1, d_1), \dots, (E_s, N_s, t_s, d_s)\}, E_i, N_i \subset k[U], t_i \in k[X], d_i \in k[X][T],$$

such that  $\mathbb{V}_L(E) \setminus \mathbb{V}_L(N) = \cup_{i=1}^s \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$  and for each  $i \in \{1, \dots, s\}$ ,  $t_i$  is a separating element of  $I(\bar{u})$  and  $d_i(\bar{u}, T)$  is a greatest common divisor of  $\mathcal{X}(\bar{u}, T)$  and  $\mathcal{X}'(\bar{u}, T)$  for any  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ .  $\square$

According to Theorem 18, we further compute  $g(U, T), g_1(U, T), \dots, g_n(U, T)$  and obtain an RUR for each branch.

Now we design another algorithm for computing the rational univariate representations with parameters of an ideal in  $k[U, X]$ .

### Algorithm 2 for computing RURs with parameters

**Input:**  $I = \langle f_1, \dots, f_l \rangle \subset k[U, X]$ .

1. Compute a comprehensive Gröbner system of  $I$  with  $\prec_{X,U}$ . Choose the zero-dimensional branches by Theorem 11 and denote them by  $\mathcal{G} = \{(E_1, N_1, G_1), \dots, (E_s, N_s, G_s)\}$ .
2.  $B_2 := \emptyset$ . For each branch  $(E_i, N_i, G_i) \in \mathcal{G}$  do:
  - (a) Compute  $Q_1^i = (p_{ij}/q_{ij}) \in k(U)^{r \times r}$  defined in Lemma 13. Let  $Q = D \cdot Q_1^i$ , where  $D = \text{lcm}(\{q_{ij} : i, j = 1, \dots, r\})$ .

- (b) Assume that  $Q = (f_{ij})_{1 \leq i, j \leq r}$ . Construct polynomial  $f_i = \sum_{j=1}^r f_{ij} z_j$ . Compute the minimal Comprehensive Gröbner system  $\mathcal{G}' = \{(E_{i1}, N_{i1}, G_{i1}), \dots, (E_{is_i}, N_{is_i}, G_{is_i})\}$  of  $\{f_1, \dots, f_r\}$ . Set  $A_i := \{(E_{i1}, N_{i1}, k_{i1}), \dots, (E_{is_i}, N_{is_i}, k_{is_i})\}$ , where  $k_{ij}$  is equal to number of element in  $G_{ij}$ ,  $j = 1, \dots, s_i$ .
- (c)  $B_2 := B_2 \cup A_i$ .
3.  $B_3 := \emptyset$ . For each branch  $(E_{ij}, N_{ij}, k_{ij}) \in B_2$  do:
- (a) Randomly choose  $t \in S$ . Compute  $M_t^i \in k(U)^{r \times r}$ ,  $\mathcal{X}_t \in k(U)[T]$ . Let  $\mathcal{X} = D' \cdot \mathcal{X}_t$ . Compute parametric GCDs of  $\mathcal{X}$  and  $\mathcal{X}'$ , denoted by

$$\{(E_{ijk}, N_{ijk}, d_{ijk}) : k = 1, \dots, v\},$$

where  $D'$  is the least common multiple of all denominators of  $\mathcal{X}_t$ .

- (b) For each  $(E_{ijk}, N_{ijk}, d_{ijk})$  do:  
 $A_{ijk} := \emptyset$ . If  $\deg(d_{ijk}) > \deg_T(\mathcal{X}(U, T)) - k_{ij}$ , then  $S := S \setminus \{t\}$  and goto (3.a); else, compute  $\overline{\mathcal{X}}_t$ . By Formula (3), (4) compute  $g_{ijk}(U, T)$ ,  $g_{ijk1}(U, T)$ ,  $\dots$ ,  $g_{ijkn}(U, T)$ .  
 $A_{ijk} := A_{ijk} \cup \{(E_{ijk}, N_{ijk}, \mathcal{X}_t, g_{ijk}, g_{ijk1}, \dots, g_{ijkn})\}$
- (c)  $B_3 := B_3 \cup A_{ijk}$ .

**Output:** a finite set  $B_3$  of the rational univariate representations with parameters, renumbering it by

$$\{(E_1, N_1, \mathcal{X}_1, g_1, g_{11}, \dots, g_{1n}), \dots, (E_s, N_s, \mathcal{X}_s, g_s, g_{s1}, \dots, g_{sn})\}.$$

For each  $i \in \{1, \dots, s\}$ , it satisfies that : for all  $\bar{u} \in \mathbb{V}_L(E_i) \setminus \mathbb{V}_L(N_i)$ ,

$$\mathbb{V}_L(I(\bar{u})) = \left\{ \left( \frac{g_{i1}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)}, \dots, \frac{g_{in}(\bar{u}, \beta)}{g_i(\bar{u}, \beta)} \right) \mid \beta \in \mathbb{V}_L(\mathcal{X}_i(\bar{u}, T)) \right\},$$

where  $I(\bar{u}) = \langle f_1(\bar{u}, X), \dots, f_i(\bar{u}, X) \rangle \subset k[X]$ ,  $\mathcal{X}_i(\bar{u}, T) \in k[T]$ .

**Theorem 21.** *The above algorithm works correctly and terminates in a finite number of steps.*

*Proof.* The correctness and the termination of the algorithm directly follows from Theorems 11, 14, 20.  $\square$

#### 4. Example

We use the following simple example to illustrate the steps in the above proposed algorithms.

**Example 22.** *Let  $I = \langle u_1 x_1^2 + u_2 x_2 + u_2, u_2 x_2^2 + u_1 x_2 + u_1 \rangle \subset \mathbb{C}[u_1, u_2, x_1, x_2]$  with  $<_U$  and  $<_X$  being graded lexicographic order. We see the process of Algorithm 1:*

(1) *Compute the minimal comprehensive Gröbner system for  $I$ :*

$$\begin{aligned} \mathcal{G} = & \{(\{0\}, \{u_1 u_2\}, \{u_2 x_2^2 + u_1 x_2 + u_1, u_1 x_1^2 + u_2 x_2 + u_2\}), \\ & (\{u_1\}, \{u_2, u_2 u_1\}, \{1\}), (\{u_2\}, \{u_1\}, \{u_1 x_1^2, u_1 x_2 + u_1\}), \\ & (\{u_1, u_2\}, \{1\}, \{0\})\}. \end{aligned}$$

*By Finiteness Theorem,  $(\{0\}, \{u_1 u_2\}, \{u_2 x_2^2 + u_1 x_2 + u_1, u_1 x_1^2 + u_2 x_2 + u_2\})$  and  $(\{u_2\}, \{u_1\}, \{u_1 x_1^2, u_1 x_2 + u_1\})$  are two zero-dimensional branches. For brevity, we only consider the first branch.*



(2) For  $(\{0\}, \{u_1u_2\}, \{u_2x_2^2 + u_1x_2 + u_1, u_1x_1^2 + u_2x_2 + u_2\})$ , we compute  $Q_1^I(U)$  as defined in Theorem 14:

$$Q_1^I(U) = \begin{bmatrix} 4 & -\frac{2u_1}{u_2} & 0 & 0 \\ \frac{2u_1}{u_2} & \frac{2u_1(u_1-2u_2)}{u_2^2} & 0 & 0 \\ 0 & 0 & \frac{2(u_1-2u_2)}{u_1} & -\frac{2(-3u_2+u_1)}{u_2} \\ 0 & 0 & \frac{2(3u_2-u_1)}{u_2} & \frac{2(u_1^2-4u_1u_2+2u_2^2)}{u_2^2} \end{bmatrix}.$$

Then  $D = u_1u_2^2$  be the least common multiple of all denominators of  $Q_1^I$ . Let  $Q = D \cdot Q_1^I$ . Denote the entries of  $Q$  by  $f_{ij}$  and set  $f_i = \sum_{j=1}^4 f_{ij}z_j$ ,  $i = 1, \dots, 4$ . We compute the minimal comprehensive Gröbner system of  $\{f_1, f_2, f_3, f_4\}$  on  $\mathbb{C}^2 \setminus \mathbb{V}_{\mathbb{C}}(u_1u_2)$ :  $(\{0\}, \{(u_1 - 4u_2)u_2u_1\}, \{(u_1^3u_2 - 4u_1^2u_2^2)z_2, (u_1^2u_2^3 - 4u_1u_2^4)z_4, 2u_1u_2^2z_1 - u_1^2u_2z_2, 2u_2^4z_3 - u_1u_2^3z_4\}), (\{u_1 - 4u_2\}, \{u_1u_2\}, \{u_2^3z_1 - 2u_2^3z_2, u_2^3z_3 - 2u_2^3z_4\})$ . Therefore,

$$\begin{aligned} B_2 &= \{(E_{11}, N_{11}, k_{11}), (E_{12}, N_{12}, k_{12})\} \\ &= (\{0\}, \{u_1u_2(u_1 - 4u_2)\}, 4), (\{u_1 - 4u_2\}, \{u_1u_2\}, 2). \end{aligned}$$

(3) Let  $B_3 = \emptyset$ . For the branch  $(E_{11}, N_{11}, k_{11}) = (\{0\}, \{u_1u_2(u_1 - 4u_2)\}, 4)$ , let  $A_{11} = \emptyset$ . Randomly choose  $t = x_1 \in S$ . By computation,

$$M_t^I = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -\frac{u_2}{u_1} & -\frac{u_2}{u_1} & 0 & 0 \\ 1 & \frac{u_1-u_2}{u_1} & 0 & 0 \end{bmatrix}, \quad X_t = T^4 - \frac{(u_1 - 2u_2)T^2}{u_1} + \frac{u_2^2}{u_1^2}.$$

Then  $\mathcal{X} = u_1^2T^4 - u_1(u_1 - 2u_2)T^2 + u_2^2$ ,  $\mathcal{X}' = 4u_1^2T^3 - 2u_1(u_1 - 2u_2)T$ ,  $PSC_0(\mathcal{X}, \mathcal{X}') = 16u_1^0u_2^2(u_1 - 4u_2)^2$ . By check,  $\mathbb{C}^2 \setminus \mathbb{V}_{\mathbb{C}}(u_1u_2(u_1 - 4u_2) \cdot PSC_0(\mathcal{X}, \mathcal{X}')) \neq \emptyset$ ,  $x_1$  is a separating element of  $I(\bar{u})$  under  $\bar{u} \in \mathbb{C}^2 \setminus \mathbb{V}_{\mathbb{C}}(16u_1^1u_2^3(u_1 - 4u_2)^3)$ . Now

$$A_{11} := A_{11} \cup \{(\{0\}, \{16u_1^1u_2^3(u_1 - 4u_2)^3\}, x_1)\}.$$

Update  $E_{11} = E_{11} \cup \{PSC_0(\mathcal{X}, \mathcal{X}')\}$ . We can check  $\mathbb{V}_{\mathbb{C}}(E_{11}) \setminus \mathbb{V}_{\mathbb{C}}(N_{11}) = \emptyset$ . Then this branch has been done and

$$B_3 := B_3 \cup A_{11} = \{(\{0\}, \{16u_1^1u_2^3(u_1 - 4u_2)^3\}, x_1)\}.$$

For the branch  $(E_{12}, N_{12}, k_{12}) = (\{u_1 - 4u_2\}, \{u_1u_2\}, 2)$ , let  $A_{12} = \emptyset$ . We also choose  $t = x_1$ . Then  $M_t^I, \mathcal{X}_t, \mathcal{X}, \mathcal{X}'$  are the same as above. By computation,

$$PSC_2(\mathcal{X}, \mathcal{X}') = -8u_1^5(u_1 - 2u_2).$$

We can check that  $\mathbb{V}_{\mathbb{C}}(u_1 - 4u_2) \setminus \mathbb{V}_{\mathbb{C}}(u_1u_2 \cdot PSC_2(\mathcal{X}, \mathcal{X}')) \neq \emptyset$ , then

$$A_{12} := A_{12} \cup \{(\{u_1 - 4u_2\}, \{-8u_1^5u_2(u_1 - 2u_2)\}, x_1)\}.$$

Update  $E_{12} = E_{12} \cup \{PSC_2(\mathcal{X}, \mathcal{X}')\} = \{u_1 - 4u_2, -8u_1^5(u_1 - 2u_2)\}$ . By check,  $\mathbb{V}_{\mathbb{C}}(E_{12}) \setminus \mathbb{V}_{\mathbb{C}}(N_{12}) = \emptyset$ . This branch has been done and

$$\begin{aligned} B_3 &:= B_3 \cup A_{12} \\ &= (\{0\}, \{u_1^1u_2^3(u_1 - 4u_2)^3\}, x_1), (\{u_1 - 4u_2\}, \{-8u_1^5u_2(u_1 - 2u_2)\}, x_1). \end{aligned}$$

(4) For the branch  $(\{0\}, \{u_1^1u_2^3(u_1 - 4u_2)^3\}, x_1)$ , we compute the parametric GCDs of  $\mathcal{X}$  and  $\mathcal{X}'$  corresponding to  $t = x_1$ . That is  $\{(\{0\}, \{u_2u_1(u_1 - 4u_2)\}, 1)\}$ . Then  $\bar{\mathcal{X}}_t = \mathcal{X}_t$ . By Formula (3), (4),

$g_1 = 4T^3 - \frac{2(u_1-2u_2)T}{u_1}$ ,  $g_{11} = \frac{2(u_1-2u_2)}{u_1}T^2 - \frac{4u_2^2}{u_1^2}$ ,  $g_{12} = -\frac{2u_1}{u_2}T^3 + 2T$ . Let  $\mathcal{X}_1 = \mathcal{X}_t = T^4 - \frac{u_1-2u_2}{u_1}T^2 + \frac{u_2^2}{u_1^2}$ , we obtain the RUR with parameters of this branch is  $(\{0\}, \{u_1u_2(u_1-4u_2)\}, \mathcal{X}_1, g_1, g_{11}, g_{12})$ .

For the branch  $(\{u_1-4u_2\}, \{-8u_1^6u_2(u_1-2u_2)\}, x_1)$ , we compute the parametric GCDs of  $\mathcal{X}, \mathcal{X}'$  corresponding to  $t = x_1$ . That is

$$(\{u_1-4u_2\}, \{u_1u_2\}, 4T^2u_2^2 - u_2^2)$$

By computation,  $\overline{\mathcal{X}}_t = T^2 - \frac{3u_1-8u_2}{4u_1}$ ,  $g_2 = 4T$ ,  $g_{21} = \frac{2(u_1-2u_2)}{u_1}$ ,  $g_{22} = -\frac{2u_1T}{u_2}$ . Let  $\mathcal{X}_2 = \mathcal{X}_t$ . In this case, we can reduce  $\mathcal{X}_2, g_2, g_{21}, g_{22}$  by the relation  $u_1-4u_2=0$ . Then we have  $\mathcal{X}_2 = T^4 - \frac{1}{2}T^2 + \frac{1}{16}$ ,  $g_2 = 4T$ ,  $g_{21} = 1$ ,  $g_{22} = -8T$ . Thus, the RUR with parameters of this branch is  $(\{u_1-4u_2\}, \{(u_1-2u_2)u_1u_2\}, \mathcal{X}_2, g_2, g_{21}, g_{22})$ .

In conclusion, we obtain two RURs of  $I$  under the parameter branch  $(\{0\}, \{u_1u_2\})$ . That is  $(\{0\}, \{u_1u_2(u_1-4u_2)\}, \mathcal{X}_1, g_1, g_{11}, g_{12})$  and  $(\{u_1-4u_2\}, \{(u_1-2u_2)u_1u_2\}, \mathcal{X}_2, g_2, g_{21}, g_{22})$ . Therefore,

(i) For all  $\bar{u} = (\bar{u}_1, \bar{u}_2) \in \mathbb{C}^2 \setminus \mathbb{V}_{\mathbb{C}}(u_1u_2(u_1-4u_2))$ ,

$$\begin{aligned} \mathbb{V}_{\mathbb{C}}(I(\bar{u})) &= \left\{ \left( \frac{g_{11}(\bar{u}, \beta)}{g_1(\bar{u}, \beta)}, \frac{g_{12}(\bar{u}, \beta)}{g_1(\bar{u}, \beta)} \right) \mid \beta \in \mathbb{V}_{\mathbb{C}} \left( T^4 - \frac{\bar{u}_1-2\bar{u}_2}{\bar{u}_1}T^2 + \frac{\bar{u}_2^2}{\bar{u}_1^2} \right) \right\} \\ &= \left\{ \left( \frac{(\bar{u}_1^2-2\bar{u}_1\bar{u}_2)\beta^2-2\bar{u}_2^2}{2\beta^3\bar{u}_1^2-(\bar{u}_1-2\bar{u}_2)\bar{u}_1\beta}, \frac{-\bar{u}_1^2\beta^2+\bar{u}_1\bar{u}_2}{2\beta^2\bar{u}_1\bar{u}_2-\bar{u}_2(\bar{u}_1-2\bar{u}_2)} \right) \mid \right. \\ &\quad \left. \beta \in \mathbb{V}_{\mathbb{C}} \left( T^4 - \frac{\bar{u}_1-2\bar{u}_2}{\bar{u}_1}T^2 + \frac{\bar{u}_2^2}{\bar{u}_1^2} \right) \right\}. \end{aligned}$$

(ii) For all  $\bar{u} = (\bar{u}_1, \bar{u}_2) \in \mathbb{V}_{\mathbb{C}}(u_1-4u_2) \setminus \mathbb{V}_{\mathbb{C}}(u_2)$ ,

$$\mathbb{V}_{\mathbb{C}}(I(\bar{u})) = \left\{ \left( \frac{1}{4\beta}, -2 \right) \mid \beta \in \mathbb{V}_{\mathbb{C}} \left( T^4 - \frac{1}{2}T^2 + \frac{1}{16} \right) \right\},$$

**Example 23.** Let  $I$  be the same as above. We see the process of Algorithm 2. For brevity, we also consider the first branch:

$$(\{0\}, \{u_1u_2\}, \{u_2x_2^2 + u_1x_2 + u_1, u_1x_1^2 + u_2x_2 + u_2\}).$$

Step 2 is the same as Algorithm 1, then we get

$$\begin{aligned} B_2 &= \{(E_{11}, N_{11}, k_{11}), (E_{12}, N_{12}, k_{12})\} \\ &= (\{0\}, \{u_1u_2(u_1-4u_2)\}, 4), (\{u_1-4u_2\}, \{u_1u_2\}, 2). \end{aligned}$$

For branch  $(\{0\}, \{u_1u_2(u_1-4u_2)\}, 4)$ . We choose  $t = x_1$ . Compute  $\mathcal{X} = u_1^2T^4 - u_1(u_1-2u_2)T^2 + u_2^2$  and  $\mathcal{X}' = 4u_1^2T^3 - 2u_1(u_1-2u_2)T$ . Then we compute the parametric GCDs of  $\mathcal{X}$  and  $\mathcal{X}'$  over  $\mathbb{C}^2 \setminus \mathbb{V}_L(u_1u_2(u_1-4u_2))$  immediately. That is

$$(\{0\}, \{u_1u_2(u_1-4u_2), 1\}).$$

By the computation of parametric GCDs, we know that, in fact,  $t = x_1$  is separate element for all  $I(\bar{u})$  when  $\bar{u} \in \mathbb{C}^2 \setminus \mathbb{V}_L(u_1u_2(u_1-4u_2))$ . Then the computations of  $g_1, g_{11}$  and  $g_{12}$  are the same as Example 22. We can get  $g_1 = 4T^3 - \frac{2(u_1-2u_2)T}{u_1}$ ,  $g_{11} = \frac{2(u_1-2u_2)}{u_1}T^2 - \frac{4u_2^2}{u_1^2}$ ,  $g_{12} = -\frac{2u_1}{u_2}T^3 + 2T$ .

For branch  $(\{u_1 - 4u_2\}, \{u_1u_2\}, 4)$ . We choose  $t = x_1$ . Then  $\mathcal{X}$  and  $\mathcal{X}'$  are the same as above. We compute the parametric GCDs of  $\mathcal{X}$  and  $\mathcal{X}'$  over  $\mathbb{V}_L(u_1 - 4u_2) \setminus \mathbb{V}_L(u_1u_2)$  immediately. That is

$$\{(\{u_1 - 4u_2\}, \{u_1u_2\}, 4T^2u_2^2 - u_2^2)\}.$$

Then the computations of  $g_2$ ,  $g_{21}$  and  $g_{22}$  are the same as Example 22. We can get  $g_2 = 4T$ ,  $g_{21} = 1$ ,  $g_{22} = -8T$ . Finally, the result is the same as Example 22.

**Remark 24.** In Algorithm 2, we skip the computation of  $\text{PSC}_d(\mathcal{X}, \mathcal{X}')$ , so it seems to be more effective than Algorithm 1. But in fact, by the computation of  $\text{PSC}_d(\mathcal{X}, \mathcal{X}')$ , we can divide the parameter space into smaller one, and this may accelerate the computation of parametric GCD. Therefore, it is hard to tell which algorithm is better.

## 5. Implementation and Comparative Performance

The algorithms for computing RURs of zero-dimensional ideals with parameters have been implemented on the computer algebra system *Singular*. The codes and examples are available on the web: <http://www.mmrc.iss.ac.cn/~dwang/software.html>.

We try all examples in Manubens and Montes (2006) and Nabeshima (2007), addition with some other examples from Lebrun (2011), Robbiano and Torrente (2013) and Ayad et al. (2012). Specifically, all examples are listed in appendix.

For all these examples, we compute the Gröbner basis of ideals under the block order  $\prec_{U,X}$  with  $U \ll X$ ; within each block,  $\prec_U$  and  $\prec_X$  are graded reverse lexicographic orders. The following tables show the comparison of our two algorithms in the number of branches and running time in seconds. Timings were obtained on Intel(R) Core(TM) i5-8250U CPU @ 1.60GHz 1.80 GHz with 8GB Memory running Windows 10.

Some remarks about the implementation issues are listed in the following:

- To speed up, in Algorithm 1, Step 3.(c), we first factorize  $\text{PSC}_d(\mathcal{X}, \mathcal{X}')$ . Suppose that  $\text{PSC}_d(\mathcal{X}, \mathcal{X}') = p_1^{r_1} \cdots p_s^{r_s}$ . Then  $\mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij})$  can be divided into

$$\mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij}) = \mathbb{V}_L(E_{ij}) \setminus \mathbb{V}_L(N_{ij} \times (p_1 \cdots p_s)) \cup \bigcup_{j=1}^s \mathbb{V}_L(E_{ij}, p_j) \setminus \mathbb{V}_L(N_{ij} \times (p_1 \cdots p_{j-1})).$$

Let  $A_{ij} := A_{ij} \cup \{(E_{ij}, N_{ij} \times (p_1 \cdots p_s), t)\}$  and  $S := S \setminus \{t\}$ . For each  $\mathbb{V}_L(E_{ij}, p_j) \setminus \mathbb{V}_L(N_{ij} \times (p_1 \cdots p_{j-1}))$ , if  $\mathbb{V}_L(E_{ij}, p_j) \setminus \mathbb{V}_L(N_{ij} \times (p_1 \cdots p_{j-1})) \neq \emptyset$ , do the step (3.b). This process may increase the number of branches, but for each branch the polynomials are simpler.

- The choice of separating element may affect the performance of the algorithm significantly. How to choose a proper separating element is a complicated issue, so we do not discuss here. In our implementation, we set a variate called **Sep\_ele\_set**. The user can choose the separating elements manually or automatically. If the user set **Sep\_ele\_set** be an empty set, then the separating elements will be chosen orderly in  $\{x_1, x_1 + x_2 + \cdots + x_n, x_1 + 2x_2 + \cdots + 2^{n-1}x_n, x_1 + 3x_2 + \cdots + 3^{n-1}x_n, \dots\}$ . If **Sep\_ele\_set** is not empty, we will test the element in **Sep\_ele\_set** preferentially. In the examples, we set  $\text{Sep\_ele\_set} = \{x_1, x_3, x_2 + x_3, x_1 - x_3, x_1 + x_3, x_2 + x_3, x_2 - x_3\}$  in  $F_7$ ,  $\text{Sep\_ele\_set} = \{x_4, x_1 + x_4, x_3 + x_4, x_1 - x_4, x_3 - x_4\}$  in  $F_8$ ,  $\text{Sep\_ele\_set} = \{x_1, x_1 + 2x_2, x_2 + 2x_3, x_1 + 2x_3\}$  and  $\text{Sep\_ele\_set} = \{x_2\}$  in  $S_{15}$ . In other examples, the separating elements are chosen automatically.

Examples	Number of 0-dim branches in Alg.1	Number of 0-dim branches in Alg.2	Time of Alg. 1	Time of Alg. 2
$F_1$	0	0	0.048	0.040
$F_2$	4	4	0.181	0.158
$F_3$	0	0	0.090	0.089
$F_4$	14	14	0.718	0.680
$F_5$	4	4	0.255	0.196
$F_6$	9	9	0.442	0.433
$F_7$	10	-	36.693	>1h
$F_8$	31	32	56.298	55.797
$S_1$	3	3	0.105	0.096
$S_2$	1	1	0.053	0.048
$S_3$	0	0	0.019	0.019
$S_4$	1	1	0.045	0.040
$S_5$	43	39	2.768	2.392
$S_6$	4	4	0.259	0.215
$S_7$	8	8	2.064	0.994
$S_8$	1	1	0.068	0.061
$S_9$	3	3	0.223	0.243
$S_{10}$	13	17	0.625	0.779
$S_{12}$	13	14	0.798	0.909
$S_{13}$	4	4	0.205	0.204
$S_{14}$	0	0	0.076	0.075
$S_{15}$	15	15	1.044	1.105
$S_{16}$	4	4	0.433	0.426
$C_1$	30	30	3.382	2.646
$R_1$	7	7	0.233	0.248
$E_1$	2	2	0.075	0.062
$E_2$	2	2	0.277	0.291
$E_3$	4	4	0.171	0.155

- In implementation, we also output the conditions when the system has no solution and the conditions when the system is positive dimensional. In this paper, the zero-dimensional branch does not include the branch which corresponds to a system with no solutions. If the parametric polynomial system has no zero-dimensional branches, we will output: There are no zero-dimensional branches. (For example,  $F_1, F_3, S_3$  and  $S_{14}$ ).
- In formula 3 and 4, we store the values of  $\text{Tr}(M_{i^l}^l), \text{Tr}(M_{x_{k^l}^l}^l)$  after computation since they may be used in other branches.
- In practice, in Step 2.(b), we compute the minimal comprehensive Gröbner system of the module generated by the row of  $Q$  instead of introducing new variables  $z_i$ . It is more efficient to compute over module than introducing new variables.

## 6. Concluding remarks

The rational univariate representation of zero-dimensional ideals with parameters has been considered in the paper. Because the number of zeros for zero-dimensional ideals with parameters under parametric specializations is different, the choosing of separating elements which is the premise and the key to computing the rational univariate representation is quite difficult. By means of comprehensive Gröbner systems to divide the parameter space, we make the ideal under each branch have the same number of zeros. Moreover, we extended the subresultant theorem to parametric cases, and based on it we proposed two ideas to choose and check the separating element corresponding to each branch, which further divides the parameter space. As a result, we obtained a finite set of which each branch shares the same expression of rational univariate representation and proposed two algorithms for computing rational univariate representations of parametric zero-dimensional ideals.

## Acknowledgments

This research was supported by the National Natural Science Foundation of China under Grant Nos. 12171469 and 12201210, and the National Key Research and Development Project 2020YFA0712300.

## References

- Auzinger, W., Stetter, H., 1988. An elimination algorithm for the computation of all zeros of a system of multivariate polynomial equations. *International Series of Numerical Mathematics* 86, 11–30.
- Ayad, A., Fares, A., Ayyad, Y., 2012. An algorithm for solving zero-dimensional parametric systems of polynomial homogeneous equations. *Journal of Nonlinear Sciences and Applications* 5 (6, Special issue), 426–438.
- Becker, T., Weispfenning, V., 1993. *Gröbner Bases*. Springer-Verlag.
- Cox, D., Little, J., O’Shea, D., 2005. *Using Algebraic Geometry*. Springer.
- Cox, D. A., Little, J., O’Shea, D., 2006. *Using algebraic geometry*. Vol. 185. Springer Science & Business Media.
- Kalkbrenner, M., 1997. On the stability of gröbner bases under specializations. *Journal of Symbolic Computation* 24 (1), 51–58.
- Kapur, D., Lu, D., Monagan, M., Sun, Y., Wang, D., 2018. An efficient algorithm for computing parametric multivariate polynomial gcd. In: *Proceedings of the 2018 International Symposium on Symbolic and Algebraic Computation*. pp. 239–246.
- Kapur, D., Sun, Y., Wang, D., 2013. An efficient algorithm for computing a comprehensive gröbner system of a parametric polynomial systems. *Journal of Symbolic Computation* 49, 27–44.
- Lebrun, J., 2011. *Normal forms in statistical signal processing*. De Gruyter, Berlin, Boston, pp. 107–126.
- Ma, X., Sun, Y., Wang, D., 2012. Computing polynomial univariate representations of zero-dimensional ideals by gröbner basis. *Science China Mathematics* 55 (6), 1293–1302.
- Ma, X., Sun, Y., Wang, D., Xue, Y., 2017. On checking linear dependence of parametric vectors. In: *Intelligent Computing Theories and Application*. pp. 188–196.
- Manubens, M., Montes, A., 2006. Improving the DISPGB algorithm using the discriminant ideal. *Journal of Symbolic Computation* 41 (11), 1245–1263.
- Mishra, B., 1993. *Algorithmic algebra*. Springer-Verlag, New York.
- Montes, A., 2002. A new algorithm for discussing gröbner basis with parameters. *Journal of Symbolic Computation* 33, 183–208.
- Nabeshima, K., 2007. A speed-up of the algorithm for computing comprehensive gröbner systems. In: *Proceedings of the 2007 International Symposium on Symbolic and Algebraic Computation*. pp. 299–306.
- Nagasaka, K., 2017. Parametric greatest common divisors using comprehensive gröbner systems. In: *Proceedings of the 2017 International Symposium on Symbolic and Algebraic Computation*. pp. 341–348.
- Noro, M., Yokoyama, K., 1999. A modular method to compute the rational univariate representation of zero-dimensional ideals. *Journal of Symbolic Computation* 28 (1-2), 243–263.

- Ouchi, K., Keyser, J., 2008. Rational univariate reduction via toric resultants. *Journal of Symbolic Computation* 43 (11), 811–844.
- Robbiano, L., Torrente, M., 2013. Zero-dimensional families of polynomial systems. *Matematiche (Catania)* 68 (1), 137–164.
- Rouillier, F., 1999. Solving zero-dimensional systems through the rational univariate representation. *Applicable Algebra in Engineering Communication and Computing* 9 (5), 433–461.
- Safey El Din, M., Yang, Z., Zhi, L., 2018. On the complexity of computing real radicals of polynomial systems. In: *Proceedings of the 2018 International Symposium on Symbolic and Algebraic Computation*. pp. 351–358.
- Schost, E., 2003. Computing parametric geometric resolutions. *Applicable Algebra in Engineering Communication and Computing* 13 (5), 349–393.
- Shang, B., Zhang, S., Tan, C., Xia, P., 2017. A simplified rational representation for positive-dimensional polynomial systems and shepwm equation solving. *Journal of Systems Science and Complexity* 30, 1470–1482.
- Stetter, H., 1996. Matrix eigenproblem are at the heart of polynomial system solving. *ACM SIGSAM Bull.* 30, 27–36.
- Suzuki, A., Sato, Y., 2006. A simple algorithm to compute comprehensive gröbner bases using gröbner bases. In: *Proceedings of the 2006 International Symposium on Symbolic and Algebraic Computation*. pp. 326–331.
- Tan, C., 2009. The rational representation for solving polynomial systems (in chinese). Ph.D. thesis.
- Tan, C., Zhang, S., 2009. Separating element computation for the rational univariate representation with short coefficients in zero-dimensional algebraic varieties. *Journal of Jilin University (Science Edition)* 47, 174–178.
- Tan, C., Zhang, S., 2010. Computation of the rational representation for solutions of high-dimensional systems. *Communications in Mathematical Research* 26 (2), 119–130.
- Wang, D., Wang, H., Wei, J., Xiao, F., 2022a. An extended gcd algorithm for parametric univariate polynomial matrices and application to parametric smith form. *Journal of Symbolic Computation*.
- Wang, D., Wang, H., Xiao, F., 2020. An extended gcd algorithm for parametric univariate polynomials and application to parametric smith normal form. In: *Proceedings of the 45th International Symposium on Symbolic and Algebraic Computation*. pp. 442–449.
- Wang, D., Wei, J., Xiao, F., Zheng, X., 2022b. Rational univariate representation of zero-dimensional ideals with parameters. In: *Proceedings of the 2022 International Symposium on Symbolic and Algebraic Computation*. pp. 217–224.
- Weispfenning, V., 1992. Comprehensive gröbner bases. *Journal of Symbolic Computation* 14 (1), 1–29.
- Wu, W., 1984. *Basic Principles of Mechanical Theorem Proving in Geometries (in Chinese)*, Vol. I: Part of Elementary Geometries. Science Press.
- Xiao, F., Lu, D., Ma, X., Wang, D., 2021. An improvement of the rational representation for high-dimensional systems. *Journal of Systems Science and Complexity* 34 (6), 2410–2427.
- Zeng, G., Xiao, S., 2010. Computing the rational univariate representations for zero-dimensional systems by wu’s method. *Science China Mathematics(Chinese)* 40 (10), 999–1016.

## Appendix: Examples.

### 1. Examples in Nabeshima (2007):

$$\begin{aligned}
 F_1 &= \{u_1x_1^4x_2 + x_1x_2^2 + u_2x_1, x_1^3 + 2x_1x_2, u_2x_1^2 + x_1^2x_2\}. \\
 F_2 &= \{u_1x_1^2x_2^3 + u_2x_2 + x_2, x_1^2x_2^2 + x_1x_2 + 2, u_1x_1^2 + u_2x_2 + 2\}. \\
 F_3 &= \{u_1x_1^4 + u_3x_1^2 + u_2, u_2x_1^3 + x_1^2 + 2, u_3x_1^2 + u_4x_1\}. \\
 F_4 &= \{u_1x_1^3x_2 + u_3x_1x_2^2, x_1^4x_2 + 3u_4x_2, u_3x_1^2 + u_2x_1x_2, x_1^2x_2^2 + u_1x_1^2, x_1^5 + x_2^5\} \\
 F_5 &= \{u_1x_1^2x_2 + u_2x_1 + x_2^3, u_1x_1^2x_2 + u_2x_1x_2, x_2^2 + u_2x_1^2x_2 + u_3x_1x_2\}. \\
 F_6 &= \{x_1^4 + u_1x_1^3 + u_2x_1^2 + u_3x_1 + u_4, 4x_1^3 + 3u_1x_1^2 + 2u_2x_1 + u_3\}. \\
 F_7 &= \{x_1^3 - u_1, x_2^4 - u_2, x_1 + x_2 - u_1x_3\}. \\
 F_8 &= \{u_1x_1^2 + u_2x_3, u_3x_4^2 + x_2, (x_1 - x_2)^2 + (x_3 - x_4)^2, 2u_4x_1x_4 - 2u_2x_3\}.
 \end{aligned}$$

2. Examples in [Manubens and Montes \(2006\)](#):

$$S_1 = \{u_1(x_1 + x_2), u_2(x_1 + x_2), x_1^2 + u_1x_1\}.$$

$$S_2 = \{x_1^2, x_1x_2, x_1x_3^2, u_1x_1 + x_2, x_2x_3 - x_3^2, u_1x_2, x_3^3, u_1x_3^2, u_1^2\}.$$

$$S_3 = \{x_1^3 - u_1x_1x_2, x_1^2x_2 - 2x_2^2 + x_1\}.$$

$$S_4 = \{u_1x_1 + x_2 - 1, u_2x_1 + x_2 - 2, 2x_1 + u_1x_2, u_2x_1 + u_1x_2 + 1\}.$$

$$S_5 = \{x_4 - (u_4 - u_2), x_1 + x_2 + x_3 + x_4 - (u_1 + u_3 + u_4), x_1x_3x_4 - u_1u_3u_4, \\ x_1x_3 + x_1x_4 + x_2x_3 + x_3x_4 - (u_1u_4 + u_1u_3 + u_3u_4)\}.$$

$$S_6 = \{u_2x_1x_2 + u_1x_1^2 + x_1, u_1x_2^2 + x_1^2\}.$$

$$S_7 = \{x_2^2 - x_1x_2x_3 + x_1^2 + x_3 - 1, x_1x_2 + x_3^2 - 1, x_1^2 + x_2^2 + x_3^2 - u_1^2\}.$$

$$S_8 = \{u_1 - u_2 + (u_1x_1x_2 - u_2x_1^2x_2 - 3u_1)^3 + (x_1x_2u_2 - 3x_1u_2 - 5u_2)^4, \\ u_1x_1x_2 - u_2x_1^2x_2 - 3u_1, u_2x_1x_2 - 3u_2x_1 - 5u_2\}.$$

$$S_9 = \{x_1 + u_3x_2 + u_2x_3 + u_1, u_3x_1 + x_2 + u_1x_3 + u_2, u_2x_1 + u_1x_2 + x_3 + u_3\}.$$

$$S_{10} = \{x_1^2 + x_3^2 - 1, x_2^2 + x_4^2 - 1, u_3(x_1x_2 - x_3x_4) - x_3 + u_1, u_3(x_1x_4 + x_3x_2) + x_1 - u_2\}.$$

$$S_{12} = \{u_1 - u_3x_2 - u_4x_1, u_2 - u_3x_4 - u_4x_3, x_1^2 + x_3^2 - 1, x_2^2 + x_4^2 - 1\}.$$

$$S_{13} = \{u_1x_1^2x_2 + u_13u_2^2, u_1(u_2 - u_3)x_1x_2 + u_1u_2x_1 + 5u_3\}.$$

$$S_{14} = \{x_4^3 - u_1x_1x_4^2 - x_1x_2^2 - x_1x_3^2, x_4^3 - u_1x_2x_4^2 - x_2x_1^2 - x_2x_4^2, x_4^3 - u_1x_3x_4^2 - x_3x_1^2 - x_3x_2^2\}.$$

$$S_{15} = \{u_1 + u_4x_1, u_2 - u_4x_4, u_5x_5 + u_6x_6 - u_4, u_5x_2 + u_6x_3 - u_3, x_1^2 + x_4^2 - 1, x_2^2 + x_5^2 - 1, x_3^2 + x_6^2 - 1\}.$$

$$S_{16} = \text{See Section 5.2 in [Manubens and Montes \(2006\)](#)}.$$

3. Other examples:  $C_1$  is from [Lebrun \(2011\)](#),  $R_1$  is from [Ayad et al. \(2012\)](#),  $E_1, E_2, E_3$  are from [Robbiano and Torrente \(2013\)](#).

$$C_1 = \{u_1 - 3x_1^2 - 4x_1x_2 - 2x_2^2 - 2x_1x_3 - 2x_2x_3 - x_3^2, u_2 - 2x_1^2 - 3x_1x_2 - x_1x_3 - x_2^2 - x_2x_3, u_3 - x_1^2 - x_1x_2 - x_1x_3\}.$$

$$R_1 = \{x_1 + x_2 - u_1, x_3 + x_4 - u_2, x_1^2 + x_3^2 - 1, x_2^2 + x_4^2 - 1\}.$$

$$E_1 = \{x_1^2 + u_1x_2^2 - 1, x_2^2 + u_2x_2, -2u_1u_2x_2 + 4x_1x_2\}.$$

$$E_2 = \{u_1x_1x_2 + u_2, u_3x_1^2 + u_4x_2^2 + u_5, -2u_1u_3x_1^2 + 2u_1u_4x_2^2\}.$$

$$E_3 = \{x_1x_2 + u_1x_1 + 1, x_1^2 + x_2^2 + u_2, -2x_1^2 + 2x_2^2 + 2u_1x_2\}.$$