## Cluster state as a non-invertible symmetry protected topological phase

Sahand Seifnashri<sup>1</sup> and Shu-Heng Shao<sup>2</sup>

<sup>1</sup>School of Natural Sciences, Institute for Advanced Study, Princeton, NJ <sup>2</sup>C. N. Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook, NY

We show that the standard 1+1d  $\mathbb{Z}_2 \times \mathbb{Z}_2$  cluster model has a non-invertible global symmetry, described by the fusion category  $\operatorname{Rep}(D_8)$ . Therefore, the cluster state is not only a  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry protected topological (SPT) phase, but also a non-invertible SPT phase. We further find two new commuting Pauli Hamiltonians for the other two  $\operatorname{Rep}(D_8)$  SPT phases on a tensor product Hilbert space of qubits, matching the classification in field theory and mathematics. We identify the edge modes and the local projective algebras at the interfaces between these non-invertible SPT phases. Finally, we show that there does not exist a symmetric entangler that maps between these distinct SPT states.

17

#### CONTENTS

I.	Introduction	1
II.	Non-invertible symmetry of the cluster model	2
III.	Non-invertible SPT phases A. Gaugings and the KT transformation B. Three $\operatorname{Rep}(D_8)$ SPT phases	2 3 3
IV.	Edge modes	5
V.	No symmetric entangler for non-invertible SPT phases	6
VI.	Conclusions and outlook	7
	Acknowledgments	8
Α.	Lattice gauging 1. KW <sup>⊗2</sup> : untwisted lattice gauging ( <b>S</b> ) 2. KT: twisted lattice gauging ( <b>TST</b> )	8 8 9
В.	Non-invertible symmetry under gauging 1. $\operatorname{Rep}(D_8) \to \hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$ 2. Type III anomaly and the Levin-Gu model 3. Gauging $\operatorname{Rep}(D_8)$ in steps	9 9 10 11
С.	SPT phases as invertible field theories 1. <b>S</b> gauging in the continuum 2. <b>TST</b> gauging in the continuum	12 12 12
D.	Interface between the cluster and product states	13
Ε.	Action of D on the SPT states  1. $D   \text{cluster} \rangle = 2   \text{cluster} \rangle$ 2. $D   \text{odd} \rangle = 2(-1)^{L/4}   \text{odd} \rangle$ 3. Action on the interface states	13 13 13 14
F.	More on the absence of a $\operatorname{Rep}(D_8)$ -symmetric entangler	15
G.	Boundary conditions for the cluster model	16

References

#### I. INTRODUCTION

Symmetry protected topological (SPT) phases [1-6] are some of the most fundamental quantum phases of matter. Without imposing any global symmetry, these phases are gapped with a unique, non-degenerate ground state, and are completely featureless. However, they become distinct topological states when we impose a symmetry G, in the sense there cannot be a continuous G-symmetric deformation connecting these states without a phase transition. See [7-9] for reviews.

The simplest example of an SPT phase is the 1+1d cluster Hamiltonian [10, 11]:

$$H_{\text{cluster}} = -\sum_{j=1}^{L} Z_{j-1} X_j Z_{j+1} \,.$$
 (1)

We assume the space is a closed periodic chain of L qubits with even L. It has a  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry generated by:

$$\eta^{\mathrm{e}} = \prod_{j: \text{even}} X_j, \quad \eta^{\mathrm{o}} = \prod_{j: \text{odd}} X_j.$$
(2)

The Hamiltonian is gapped with a unique ground state, known as the cluster state, satisfying  $Z_{j-1}X_jZ_{j+1}$  |cluster $\rangle = |\text{cluster}\rangle$ . Indeed, it is a commuting projector of Pauli operators with no relation, and therefore the ground state degeneracy is one.

The cluster state is in a distinct  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  SPT phase compared to the product state  $|++\cdots+\rangle$  (which is the ground state of the trivial Hamiltonian  $H_{\text{trivial}} = -\sum_{j=1}^L X_j$ ). Explicitly, it is given by

$$|\text{cluster}\rangle = V|++\cdots+\rangle$$
, where  $V = \prod_{j=1}^{L} CZ_{j,j+1}$ . (3)

Here  $CZ_{j,j+1} = \frac{1+Z_j+Z_{j+1}-Z_jZ_{j+1}}{2}$ . Since V, known as the cluster entangler, is a finite-depth circuit, the cluster

<sup>&</sup>lt;sup>1</sup> Our convention is that  $X \mid \pm \rangle = \pm \mid \pm \rangle$ ,  $Z \mid 0 \rangle = \mid 0 \rangle$ ,  $Z \mid 1 \rangle = - \mid 1 \rangle$ .

state is in the same phase as the product state if we do not impose any global symmetry. However, if we impose the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry, the cluster state is a distinct SPT state compared to the (trivial) product state [9]. Indeed, while V globally is  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric, the individual gates  $CZ_{j,j+1}$  are not.

In recent years, the notion of global symmetry has been generalized in many different directions [12], leading to many new topological phases. In particular, there has been a lot of progress on a novel kind of symmetry, known as non-invertible symmetries, in quantum field theory and condensed matter theory. Non-invertible symmetries are implemented by conserved operators without inverses, and therefore are not described by group theory. Nonetheless, they lead to new selection rules, new constraints on the phase diagram, and more. See [13–17] for recent reviews.

The critical Ising lattice model serves as the prototypical example of a gapless system with a non-invertible symmetry, associated with the Kramers-Wannier duality [18–24]. However, this non-invertible symmetry has a Lieb-Schultz-Mattis-type constraint, implying that it is incompatible with a unique gapped ground state [24].<sup>2</sup> It is then natural to ask what is the simplest non-degenerate gapped phase protected by a non-invertible symmetry.

In this paper we show that the standard cluster Hamiltonian (1) has a non-invertible symmetry. Therefore, the cluster state is not only a  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  SPT phase, but also a non-invertible SPT phase, i.e., a non-degenerate gapped phase invariant under a non-invertible symmetry. We furthermore find two other distinct SPT states protected by the same non-invertible symmetry, matching the expectation from category theory [33] and field theory [26].

# II. NON-INVERTIBLE SYMMETRY OF THE CLUSTER MODEL

The key observation is that the cluster Hamiltonian is invariant under the transformation  $^3$ 

$$X_j \rightsquigarrow Z_{j-1}Z_{j+1}, \qquad Z_{j-1}Z_{j+1} \rightsquigarrow X_j, \qquad (4)$$

which defines an automorphism of the algebra of  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  invariant operators. However, suppose this transformation were implemented by an invertible operator U such that  $UX_jU^{-1}=Z_{j-1}Z_{j+1}$ , then  $U\eta^{\rm e}U^{-1}=\prod_{j:{\rm even}}Z_{j-1}Z_{j+1}=1$ , which is a contradiction. Instead,

this transformation is implemented by the following conserved operator

$$D = TD^{e}D^{o}, (5)$$

where

$$D^{e} = e^{\frac{2\pi i L}{16}} \frac{1 + \eta^{e}}{\sqrt{2}} \frac{1 - iX_{L}}{\sqrt{2}} \cdots \frac{1 - iZ_{4}Z_{2}}{\sqrt{2}} \frac{1 - iX_{2}}{\sqrt{2}},$$

$$D^{o} = e^{\frac{2\pi i L}{16}} \frac{1 + \eta^{o}}{\sqrt{2}} \frac{1 - iX_{L-1}}{\sqrt{2}} \cdots \frac{1 - iZ_{3}Z_{1}}{\sqrt{2}} \frac{1 - iX_{1}}{\sqrt{2}},$$
(6)

are the Kramers-Wannier operators on the even and odd sites in [23, 24]. Here T is the lattice translation by one site which acts on local operators as  $TX_jT^{-1}=X_{j+1},\,TZ_jT^{-1}=Z_{j+1}$  and satisfies  $T^L=1$ . We have  $T\mathsf{D}^e=\mathsf{D}^oT$  and  $T\mathsf{D}=\mathsf{D}T$ . However,  $\mathsf{D}$  is not a conventional symmetry operator because it has a kernel—it annihilates every state that is not  $\mathbb{Z}_2^e\times\mathbb{Z}_2^o$  symmetric. It implements (4) in the sense that

$$DX_j = Z_{j-1}Z_{j+1}D$$
,  $DZ_{j-1}Z_{j+1} = X_jD$ . (7)

The operators  $\eta^{e}$ ,  $\eta^{o}$ , and D commutes with the Hamiltonian and satisfy the following algebra

$$D^{2} = 1 + \eta^{e} + \eta^{o} + \eta^{e} \eta^{o},$$
  

$$\eta^{e} D = D \eta^{e} = \eta^{o} D = D \eta^{o} = D.$$
(8)

We have an (internal) non-invertible symmetry in the cluster model that does not mix with the lattice translation. Therefore, the cluster state is a topological phase protected by a non-invertible symmetry.

In Appendix B 3, we show that this non-invertible symmetry is described by the fusion category  $Rep(D_8)$ , whose fusion algebra is given by the tensor product of the irreducible representations of the group  $D_8$ . (See [34] for a review of fusion category, which is built on [35, 36].) We prove this by gauging the non-invertible symmetry and find a dual  $D_8$  symmetry.

Finally, we note that the cluster Hamiltonian is also invariant under another non-invertible symmetry  $\tilde{\mathsf{D}} = \mathsf{D}^{\mathsf{e}}\mathsf{D}^{\mathsf{o}}$ , which obeys the algebra  $\tilde{\mathsf{D}}^2 = (1 + \eta^{\mathsf{e}} + \eta^{\mathsf{o}} + \eta^{\mathsf{e}} \eta^{\mathsf{o}})T^{-2}$ . Similar to the Kramers-Wannier symmetry of the critical Ising model [23],  $\tilde{\mathsf{D}}$  mixes with the lattice translation, therefore it does not form a fusion category on the lattice. In the continuum, it can flow to a Rep(H<sub>8</sub>) fusion category symmetry. Note that breaking the translation symmetry T in (1) also breaks  $\tilde{\mathsf{D}}$ , but preserves  $\mathsf{D}$ .

### III. NON-INVERTIBLE SPT PHASES

We have shown that the cluster state is a non-invertible SPT phase. Are there other  $\operatorname{Rep}(D_8)$  SPT phases? As we discuss in Section V, there is no notion of a trivial SPT phase for non-invertible symmetries. Also note that

<sup>&</sup>lt;sup>2</sup> For non-invertible symmetries in continuum field theory, the incompatibility with a non-degenerate gapped phase is often defined as a generalized 't Hooft anomaly [25–29]. See [30–32] for the relation between this definition and the obstruction to gauging the non-invertible global symmetry.

<sup>&</sup>lt;sup>3</sup> Throughout the paper, we use  $\leadsto$  to denote a transformation implemented by a non-invertible operator, and  $\mapsto$  for a conventional symmetry transformation implemented by a unitary operator.

D does not act on-site and the product state is not invariant under it.<sup>4</sup> Mathematically, a non-invertible SPT phase corresponds to a fiber functor of the fusion category. It is known [33] that there are 3 distinct fiber functors for Rep(D<sub>8</sub>), labeled by the 3 non-trivial elements  $\eta^e$ ,  $\eta^o$ ,  $\eta^d = \eta^e \eta^o$  of  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ . See [26, 28, 38–40] for a field theory exposition of these SPT phases.

What are the lattice models for the other two  $\operatorname{Rep}(D_8)$  SPT phases? To find them, we partially gauge the non-invertible symmetry and look for different spontaneous symmetry breaking patterns in the gauged theory. We first review this method in the case of  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ .

### A. Gaugings and the KT transformation

For  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$ , there are only two distinct SPT phases: the product state  $|++\cdots+\rangle$  and the cluster state. One way to distinguish these two phases is to gauge the  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  symmetry. Gauging the symmetry via minimal coupling amounts to doing Kramers-Wannier transformations on even and odd sites separately. See Appendix A 1 for a review. Up to a lattice translation, this is the same transformation as in (4). The Hamiltonians of these two phases after gauging are

$$H'_{\text{trivial}} = -\sum_{j=1}^{L} Z'_{j-1} Z'_{j+1},$$

$$H'_{\text{cluster}} = -\sum_{j=1}^{L} Z'_{j-1} X'_{j} Z'_{j+1}.$$
(9)

The two SPT phases are distinct in that they are mapped to a symmetry breaking phase and a symmetry preserving phase under gauging. In particular, the cluster Hamiltonian is invariant under gauging.<sup>5</sup>

For later convenience, we will choose to distinguish these two SPT phases by a twisted gauging, reviewed in Appendix A 2. In the condensed matter literature, it is known as the Kennedy-Tasaki (KT) transformation [44, 45] (see also [22, 46]):

$$X_j \rightsquigarrow \hat{X}_j$$
,  $Z_{j-1}Z_{j+1} \rightsquigarrow \hat{Z}_{j-1}\hat{X}_j\hat{Z}_{j+1}$ . (10)

(In the field theory context, the untwisted and twisted gaugings are respectively referred to as the  $\bf S$  and  $\bf TST$ 

TABLE I. The KT transformation, corresponding to a twisted gauging labeled by **TST**, maps the product state and the cluster state to a symmetry preserving phase and a symmetry breaking phase, respectively.

gaugings, representing different elements of the modular group.) If we identify the hatted operators with the unhatted ones, then the KT transformation is implemented by the non-invertible operator VDV [46]:

$$\begin{aligned} & (\mathsf{VDV}) X_j = X_j (\mathsf{VDV}) \,, \\ & (\mathsf{VDV}) Z_{j-1} Z_{j+1} = Z_{j-1} X_j Z_{j+1} (\mathsf{VDV}) \,. \end{aligned} \tag{11}$$

The two SPT phases after twisted gauging become

$$\hat{H}_{\text{trivial}} = -\sum_{j} \hat{X}_{j},$$

$$\hat{H}_{\text{cluster}} = -\sum_{j} \hat{Z}_{j-1} \hat{Z}_{j+1}.$$
(12)

The trivial phase is invariant under twisted gauging and the cluster state goes to the symmetry breaking phase of the dual  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o$  global symmetry, generated by  $\hat{\eta}^e = \prod_{j:\text{even}} \hat{X}_j$ ,  $\hat{\eta}^o = \prod_{j:\text{odd}} \hat{X}_j$ . See Table I for a summary and Appendix C for the (twisted) gauging the two SPT phases in continuum field theory.

# B. Three $Rep(D_8)$ SPT phases

Now we are ready to study the SPT phases of the non-invertible symmetry. As we show in Appendix B, after the twisted gauging of  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ , the non-invertible symmetry (8) becomes an anomalous  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$  symmetry generated by

$$\hat{\eta}^{\mathrm{e}} = \prod_{j:\mathrm{even}} \hat{X}_j, \quad \hat{\eta}^{\mathrm{o}} = \prod_{j:\mathrm{odd}} \hat{X}_j, \quad \hat{\mathsf{V}} = \prod_j \hat{\mathrm{CZ}}_{j,j+1}. \quad (13)$$

In other words, applying the transformation (10) to any Hamiltonian invariant under (8) results in a Hamiltonian with the  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$  symmetry. The 't Hooft anomaly of this symmetry is described by the 2+1d invertible field theory  $(-1)^{\int \hat{A}^e \cup \hat{A}^o \cup \hat{A}^V}$ , known as the type III anomaly [47, 48]. (See Appendix B 2. Here  $\hat{A}^e$ ,  $\hat{A}^o$ , and  $\hat{A}^V$  are discrete background gauge fields.) We summarize this relation as

$$\operatorname{Rep}(D_8) \xrightarrow{\mathbf{TST}} \underbrace{\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^{\mathsf{V}}}_{\text{type III anomaly}} \tag{14}$$

The three non-invertible SPT phases can be distinguished by the symmetry breaking patterns of the this dual symmetry after the KT transformation. Equivalently, these

<sup>&</sup>lt;sup>4</sup> Even though non-invertible symmetries do not act on-site, they can sometimes leave the product state invariant. For instance, by performing the similarity transformation V we find the non-invertible symmetry operator VDV that does leave the product state invariant. See also [37], where the authors considered SPT phases protected by the non-invertible symmetry  $G \times \text{Rep}(G)$ . In their lattice model, the product state is invariant under their non-invertible symmetry.

<sup>&</sup>lt;sup>5</sup> Following the half-gauging argument in [24, 25, 41, 42], the invariance under gauging a global symmetry implies the existence of the non-invertible global symmetry, which is precisely the operator D. This will be explored in [43].

phases are distinguished by certain non-local, string order parameters (see, e.g., [39, 49]).

Given any of the three  $\operatorname{Rep}(D_8)$  SPT phases, the fact that it has the non-invertible symmetry D implies the invariance under the untwisted gauging of  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  [24, 43]. As discussed above, the cluster state is invariant under the untwisted gauging but the product state is not. Therefore, any  $\operatorname{Rep}(D_8)$  SPT must be in the same phase as the cluster state as far as the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry is concerned.

This implies that after the twisted gauging, the dual  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o$  symmetry must be completely broken, with a single  $\mathbb{Z}_2$  subgroup of  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$  preserved.<sup>6</sup> A priori, there are four options for this unbroken  $\mathbb{Z}_2$  subgroup, generated by  $\hat{V}$ ,  $\hat{V}\hat{\eta}^e$ ,  $\hat{V}\hat{\eta}^o$ , or  $\hat{V}\hat{\eta}^e\hat{\eta}^o$ . However, the diagonal subgroup generated by  $\hat{V}\hat{\eta}^e\hat{\eta}^o$  cannot be preserved since it is anomalous, which is obtained from  $(-1)^{\int \hat{A}^e \cup \hat{A}^o \cup \hat{A}^V}$  by setting  $\hat{A}^e = \hat{A}^o = \hat{A}^V$ . This leaves us with three possible symmetry breaking patterns in Table II.

a.  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$  preserving phase: The original cluster Hamiltonian after gauging is given by (12). The order parameters of the gauged theory are given by  $\hat{Z}_0$  and  $\hat{Z}_1$  which are both invariant under  $\hat{\mathsf{V}}$ . Thus the cluster state preserves the  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$  subgroup.

b.  $diag(\hat{\mathbb{Z}}_2^{\mathsf{V}} \times \hat{\mathbb{Z}}_2^{\mathsf{e}})$  preserving phase: For the second option, we propose the following Hamiltonian for the gauged system<sup>7</sup>

$$\hat{H}_{\text{odd}} = \sum_{n=1}^{L/2} \hat{Z}_{2n-1} \hat{Z}_{2n+1}$$

$$- \sum_{n=1}^{L/2} \hat{Y}_{2n} \hat{Y}_{2n+2} (1 + \hat{Z}_{2n-1} \hat{Z}_{2n+3}),$$
(15)

and take the number of sites L to be a multiple of 4. (The subscript "odd" will become clear later.) This is a commuting Pauli Hamiltonian and thus is exactly solvable.<sup>8</sup> Note that the projection factor in the second term does not affect the ground states. This is because for the ground states  $\hat{Z}_{2n-1}\hat{Z}_{2n+1}=-1$  and thus  $\hat{Z}_{2n-1}\hat{Z}_{2n+3}=1$ . The ground space is determined by L-2 independent constraints  $\hat{Z}_{2n-1}\hat{Z}_{2n+1}=-1$  and  $\hat{Y}_{2n}\hat{Y}_{2n+2}=1$ . This leads to a four-fold degeneracy with ground states given by  $\hat{Z}_1=-\hat{Z}_3=\cdots=-\hat{Z}_{L-1}=\pm 1$  and  $\hat{Y}_0=\hat{Y}_2=\cdots=\hat{Y}_{L-2}=\pm 1$ . The two order parameters are subscripted as  $\hat{Y}_0=\hat{Y}_0=\hat{Y}_0=0$ .

$Rep(D_8)$ SPT states	Symmetry breaking pattern of $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$		
$ cluster\rangle$	$\hat{\mathbb{Z}}_2^{V}$ unbroken		
$ { m odd} angle$	$\underline{\text{TST}} \rightarrow \operatorname{diag}(\hat{\mathbb{Z}}_2^{V} \times \hat{\mathbb{Z}}_2^{e}) \text{ unbroken}$		
$ { m even}\rangle$	$\operatorname{diag}(\hat{\mathbb{Z}}_2^{V} \times \hat{\mathbb{Z}}_2^{o})$ unbroken		

TABLE II. The three Rep(D<sub>8</sub>) SPT phases are distinguished by the symmetry breaking patterns of the dual  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^v \times \hat{\mathbb{Z}}$ 

rameters can be taken to be  $\hat{Z}_1$  and  $\hat{Y}_2(1-\hat{Z}_1\hat{Z}_3)$ , which are both invariant under  $\hat{V}\hat{\eta}^e$ .

Next, we undo the twisted gauging of  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  to find the Hamiltonian for this Rep(D<sub>8</sub>) SPT phase:

$$H_{\text{odd}} = \sum_{n=1}^{L/2} Z_{2n-1} X_{2n} Z_{2n+1} - \sum_{n=1}^{L/2} Y_{2n} X_{2n+1} Y_{2n+2} + \sum_{n=1}^{L/2} Z_{2n-1} Z_{2n} X_{2n+1} Z_{2n+2} Z_{2n+3}.$$
(16)

This Hamiltonian has a unique ground state, denoted by  $|\text{odd}\rangle$ , stabilized by the following L generators:

$$-Z_{2n-1}X_{2n}Z_{2n+1} = 1$$
,  $Y_{2n}X_{2n+1}Y_{2n+2} = 1$ . (17)

Note that while  $-Z_{2n-1}X_{2n}Z_{2n+1}$  is invariant under D,  $Y_{2n}X_{2n+1}Y_{2n+2}$  is mapped to a product of terms in (17). Similarly, the individual terms in the first sum of  $H_{\text{odd}}$  are invariant under D, while the terms in the second and the third sums are mapped to each other.

The |odd\rangle state is related to the product state as

$$|\text{odd}\rangle = \prod_{n=1}^{L/2} \text{CZ}_{2n-1,2n+1} \prod_{j=1}^{L} \text{CZ}_{j,j+1} |--\cdots-\rangle$$
. (18)

In Appendix E, we will show  $D | \text{odd} \rangle = 2(-1)^{L/4} | \text{odd} \rangle$ , while  $D | \text{cluster} \rangle = 2 | \text{cluster} \rangle$ .

c.  $diag(\hat{\mathbb{Z}}_2^{\mathsf{V}} \times \hat{\mathbb{Z}}_2^{\mathsf{o}})$  preserving phase: The third option is related to the previous one by exchanging  $e \leftrightarrow o$ . More precisely, the Hamiltonian for this state, denoted as  $|\text{even}\rangle$ , is obtained by conjugating  $H_{\text{odd}}$  with the lattice translation, i.e.,  $H_{\text{even}} = TH_{\text{odd}}T^{-1}$ .

We have thus identified three  $Rep(D_8)$  SPT phases,  $|cluster\rangle$ ,  $|odd\rangle$ , and  $|even\rangle$ . The SPT states  $|odd\rangle$  and  $|even\rangle$  are related to the cluster state by the finite-depth circuits

$$\prod_{j=1}^{L} Z_j \prod_{n=1}^{L/2} \operatorname{CZ}_{2n-1,2n+1} \quad \text{and} \quad \prod_{j=1}^{L} Z_j \prod_{n=1}^{L/2} \operatorname{CZ}_{2n,2n+2},$$
(19)

<sup>&</sup>lt;sup>6</sup> The entire  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$  cannot be completely broken. This would have led to 8-fold degenerate ground states which cannot arise from gauging  $\mathbb{Z}_2 \times \mathbb{Z}_2$  of a model with a single gapped ground state.

<sup>&</sup>lt;sup>7</sup> In our conventions for indices  $j=1,\cdots,L$  and  $n=1,\cdots,L/2$ . <sup>8</sup> Crucially, the Hamiltonian is frustration-free in the sense that even though different terms in the Hamiltonian are not linearly independent, it is consistent to minimize all of them simultaneously

<sup>&</sup>lt;sup>9</sup> We have used the fact that L is a multiple of 4.

respectively. The gates of the circuits can be made  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ -symmetric:

$$|\text{odd}\rangle = \prod_{n=1}^{L/2} Z_{2n} Z_{2n+2} e^{\frac{\pi i}{4}(Z_{2n-1} Z_{2n+1} - 1)} |\text{cluster}\rangle$$
. (20)

This shows that the three SPT states are in the same  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  SPT phase. However, they are different SPT phases for Rep(D<sub>8</sub>). Indeed the circuits in (19) do not commute with the non-invertible symmetry operator D (see Section V for more discussions).

#### IV. EDGE MODES

In the previous section, we distinguished the three SPT phases by their different symmetry breaking patterns after gauging a part of the non-invertible symmetry. Here, we discuss the edge modes on the boundary of these phases. For ordinary on-site symmetries, non-trivial edge modes appear on the boundary between a (nontrivial) SPT phase and the product state. As emphasized before, there is no canonical notion of a trivial SPT phase for non-invertible symmetries, and we are instructed to identify the edge modes at the interface between two distinct SPT phases.

Here we focus on the interface between the SPT states  $|\text{cluster}\rangle$  and  $|\text{odd}\rangle$ . Consider a closed chain of L sites, where the cluster state is on half of the chain between sites 1 and  $\ell$  with  $1 < \ell < L$ , and the  $|\text{odd}\rangle$  state is in the region between sites  $\ell$  and  $\ell$ . We assume  $\ell = \ell$  to be a multiple of 4 and  $\ell$  to be even. Consider the Rep(D<sub>8</sub>)-symmetric interface Hamiltonian:

$$H_{\text{cluster}|\text{odd}} = -\sum_{j=1}^{\ell} Z_{j-1} X_j Z_{j+1} + \sum_{n=\frac{\ell}{2}+1}^{\frac{L}{2}} Z_{2n-1} X_{2n} Z_{2n+1}$$

$$-\sum_{n=\frac{\ell}{2}+1}^{\frac{L}{2}-2} Y_{2n} X_{2n+1} Y_{2n+2} \left(1 + Z_{2n-1} X_{2n} X_{2n+2} Z_{2n+3}\right).$$
(21)

The ground space  $\mathcal{H}_{\text{cluster}|\text{odd}}$  is stabilized by the following 2L-2 generators:

$$Z_{j-1}X_{j}Z_{j+1} = 1 \quad \text{for } j = 1, \dots, \ell,$$

$$-Z_{2n-1}X_{2n}Z_{2n+1} = 1 \quad \text{for } n = \ell/2 + 1, \dots, L/2,$$

$$Y_{2n}X_{2n+1}Y_{2n+2} = 1 \quad \text{for } n = \ell/2 + 1, \dots, L/2 - 2.$$
(22)

Hence the ground space is four-fold degenerate, signaling edge modes at the interfaces.

Before identifying the edge modes, we first discuss the projective algebra of the non-invertible symmetry at each of the interfaces, which protects the edge modes. As shown in Appendix E 3, the action of symmetry operators on any state  $|\psi\rangle \in \mathcal{H}_{\text{cluster}|\text{odd}}$  factorizes into local factors

at the interfaces:

$$D |\psi\rangle = (-1)^{\frac{L-\ell}{4}} \left( D_{L}^{(1)} D_{R}^{(1)} + D_{L}^{(2)} D_{R}^{(2)} \right) |\psi\rangle ,$$
  

$$\eta^{\circ} |\psi\rangle = \eta_{L}^{\circ} \eta_{R}^{\circ} |\psi\rangle ,$$
  

$$\eta^{e} |\psi\rangle = |\psi\rangle ,$$
(23)

where

$$\begin{aligned}
\mathsf{D}_{\mathrm{L}}^{(1)} &= Y_{L-2} Y_{L-1} Z_{L} \,, & \mathsf{D}_{\mathrm{R}}^{(1)} &= Z_{\ell+1} \,, \\
\mathsf{D}_{\mathrm{L}}^{(2)} &= Z_{L-1} \,, & \mathsf{D}_{\mathrm{R}}^{(2)} &= Z_{\ell} Y_{\ell+1} Y_{\ell+2} \,, \\
\eta_{\mathrm{L}}^{\circ} &= Y_{L-2} X_{L-1} Z_{L} \,, & \eta_{\mathrm{R}}^{\circ} &= Z_{\ell} X_{\ell+1} Y_{\ell+2} \,.
\end{aligned} \tag{24}$$

Here L and R stand for the left and right interfaces around sites j=L and  $j=\ell$ , respectively; see Figure 1. We find that the *local* factors of D are charged under  $\eta^{\circ}$  (but not under  $\eta^{\circ}$ ):

$$\eta^{\rm o}\,\mathsf{D}_{\rm L}^{(I)} = -\mathsf{D}_{\rm L}^{(I)}\,\eta^{\rm o}\,, \quad \ \eta^{\rm o}\,\mathsf{D}_{\rm R}^{(I)} = -\mathsf{D}_{\rm R}^{(I)}\,\eta^{\rm o}\,, \quad I = 1,2\,. \eqno(25)$$

Therefore, there is a projective algebra at each interface between the local factors:

$$\eta_{\rm L}^{\rm o} {\sf D}_{\rm L}^{(I)} = - {\sf D}_{\rm L}^{(I)} \eta_{\rm L}^{\rm o} \,, \quad \, \eta_{\rm R}^{\rm o} {\sf D}_{\rm R}^{(I)} = - {\sf D}_{\rm R}^{(I)} \eta_{\rm R}^{\rm o} \,, \quad I = 1, 2 \,, \eqno(26)$$

whereas  $\eta_{\rm L}^{\rm e} \mathsf{D}_{\rm L}^{(I)} = \mathsf{D}_{\rm L}^{(I)} \eta_{\rm L}^{\rm e}$  and  $\eta_{\rm R}^{\rm e} \mathsf{D}_{\rm R}^{(I)} = \mathsf{D}_{\rm R}^{(I)} \eta_{\rm R}^{\rm e}$ . This projective algebra matches with the continuum discussion in [26, (3.62)]. Importantly, the *global*  $\mathsf{D}, \eta^{\rm e}, \eta^{\rm o}$  operators realize the algebra (8) linearly.

On the left interface, the operators  $\eta_{\rm L}^{\rm o}$  and  ${\rm D}_{\rm L}^{(2)}$  form a Pauli algebra acting on a qubit localized around site L, and similarly for the operators  $\eta_{\rm R}^{\rm o}$  and  ${\rm D}_{\rm R}^{(1)}$  at the right interface. These operators commute with the interface Hamiltonian and form a complete basis of operators acting on its ground space. These edge modes are protected by the projective algebra (25) and cannot be lifted by symmetric deformations of the Hamiltonian near the interfaces.

In general, the three SPT phases are distinguished by the projective algebra involving the local factors of D and  $\eta^{\rm e}, \eta^{\rm o}$ . The projective algebra implies that the *defect* of the non-invertible symmetry D carries a non-trivial charge under  $\eta^{\rm e}, \eta^{\rm o}$ , or  $\eta^{\rm d}$ , which will be explored in [43]. In the continuum, these charges are denoted by  $\nu({\rm e}), \nu({\rm o}), \nu({\rm d})$  in [26, 33, 38]. Their values for the three SPT phases are<sup>10</sup>

$$\begin{split} \nu_{\rm cluster}(e) &= +1, & \nu_{\rm cluster}(o) = +1, & \nu_{\rm cluster}(d) = -1\,, \\ \nu_{\rm odd}(e) &= +1, & \nu_{\rm odd}(o) = -1, & \nu_{\rm odd}(d) = +1\,, \\ \nu_{\rm even}(e) &= -1, & \nu_{\rm even}(o) = +1, & \nu_{\rm even}(d) = +1\,. \end{split}$$

<sup>&</sup>lt;sup>10</sup> These invariants explain our terminology of 'odd' and 'even' SPT phases and suggest that the cluster state can be called the 'diagonal' SPT.




FIG. 1. Localization of the non-invertible operator D at the interfaces between SPT phases. The interface system (21) is locally in the cluster state in the region  $j=1,2,\cdots,\ell$ , and locally in the  $|{\rm odd}\rangle$  state in the complement region. The operator D factorizes into local factors  ${\rm D}_{\rm L}^{(1)}{\rm D}_{\rm R}^{(1)}+{\rm D}_{\rm L}^{(2)}{\rm D}_{\rm R}^{(2)}$  (multiplied by  $(-1)^{(L-\ell)/4}$ ) on the ground space .

The projective signs (25) capture the relative charges between |cluster $\rangle$  and |odd $\rangle$ , corresponding to  $\nu_{\rm cluster}(o)\nu_{\rm odd}(o)=-1$ .

Similarly, the interface between |cluster $\rangle$  and |even $\rangle$  is obtained by conjugating the previous interface system by the lattice translation T. For this interface, we find that the local factors of D are charged under  $\eta^{\rm e}$ , leading to a projective algebra involving their local factors. This projective algebra is captured by  $\nu_{\rm cluster}({\rm e})\nu_{\rm even}({\rm e})=-1$ . We leave the interface between |even $\rangle$  and |odd $\rangle$  for the future.

### V. NO SYMMETRIC ENTANGLER FOR NON-INVERTIBLE SPT PHASES

Relative classification of ordinary SPT phases: For ordinary invertible global symmetries, without knowing the microscopic details, it is only meaningful to discuss the relative SPT phase between two systems. In continuum field theory, this corresponds to the ambiguity of adding a counterterm at short distances.

In lattice systems with on-site symmetries, one often declares that the product state is the trivial SPT state. This is because, in the microscopic spin variables, the product state has trivial entanglement and a non-trivial SPT state has a short-range entanglement that cannot be destroyed without breaking the symmetry. However, from the macroscopic point of view, the notion of trivial SPT is only a choice and can be changed by doing a change of basis in the microscopic variables.

Indeed, there is usually a (locality-preserving) unitary operator U that commutes with the symmetry operators and maps the product state to any other SPT state. This operator U is called the *entangler*, since it maps the product state to a short-range entangled state. <sup>11</sup> A non-trivial

entangler U must have a mixed 't Hooft anomaly with the original symmetry. The simplest example is the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric cluster entangler V in (3), which has a type III anomaly with  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ . <sup>12</sup>

Stacking of ordinary SPT phases: Given two G-SPT states for an invertible, finite symmetry G, their tensor product state is another SPT state with respect to the diagonal subgroup of  $G \times G$ . Hence, ordinary G-SPT phases can be multiplied via stacking. For instance, the 1+1d bosonic SPT phases form a torsor over the abelian group  $H^2(G, U(1))$ , which means that there is no canonical trivial element (identity). Microscopically, the stacking is implemented by the symmetric entangler.

No stacking of non-invertible SPT phases: However, for non-invertible symmetries, there is no notion of stacking operation for the following reason [26]. Starting with two systems with the same non-invertible symmetry, the tensor product system does not have a "diagonal" non-invertible symmetry of the same type [51]. For instance, taking two systems with symmetries generated by  $\eta^e, \eta^o, D$ , the "diagonal" symmetry in the tensor product theory contains  $D \otimes D$ , which obeys a different algebra since  $(D \otimes D)^2 \neq (1 + \eta^e \otimes \eta^e)(1 + \eta^o \otimes \eta^o)$ . In fact, the "diagonal" symmetry is generally not closed under fusion, i.e., it is not a subcategory. Thus, there is no notion of invertibility for SPT phases protected by non-invertible symmetries.<sup>14</sup>

In summary, stacking exists for invertible symmetries, so the notion of a trivial (invertible) SPT phase exists, but not in a canonical way. Indeed, the classification is relative, and declaring the product state as trivial is a *choice*. Stacking does not exist for non-invertible symmetries, so the notion of a trivial non-invertible SPT phase does not exist, even if we specify a microscopic description.

No symmetric entangler between non-invertible SPT phases: This suggests that there is no Rep(D<sub>8</sub>)-symmetric entangler between the three different SPT states  $|cluster\rangle$ ,  $|odd\rangle$ ,  $|even\rangle$ .

We first prove a weaker statement that there cannot be a  $\operatorname{Rep}(D_8)$ -symmetric *Clifford* operator U such that, say,  $U | \operatorname{cluster} \rangle = | \operatorname{odd} \rangle$ . By definition, a Clifford unitary operator maps the Pauli group to itself. Therefore, if

<sup>&</sup>lt;sup>11</sup> A symmetric entangler might not always be realizable on a tensor-factorized Hilbert space. Its existence is related to the bulk-boundary correspondence between SPT states and anomalous symmetries on the boundary [50, Appendix A].

<sup>&</sup>lt;sup>12</sup> While V is  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric, in the sense that it commutes with  $\eta^e$  and  $\eta^o$ , the individual gates,  $CZ_{j,j+1}$ , are not.

This is analogous to the space of spin structures of an orientable Riemannian manifold M, which is a torsor over  $H^1(M, \mathbb{Z}_2)$ . This means that while there is no canonical trivial spin structure, the difference between any two spin structures is an element of  $H^1(M, \mathbb{Z}_2)$ , i.e., a  $\mathbb{Z}_2$  connection. Similarly, while SPT phases have a relative classification, the differences between the SPT phases, i.e., the stacking action implemented by the entanglers, have an absolute classification.

Therefore, the adjective "non-invertible" in "non-invertible symmetry protected topological phase" is attached to the word "symmetry", rather than to the word "phase." (This is similar to the use of the term "subsystem symmetry protected topological phase.")

there were such a Clifford operator, it must map between the two stabilizer groups:

$$\begin{aligned}
& \mathsf{U}\langle Z_{j-1}X_{j}Z_{j+1}\rangle\mathsf{U}^{-1} \\
&= \langle -Z_{2n-1}X_{2n}Z_{2n+1}, Y_{2n}X_{2n+1}Y_{2n+2}\rangle.
\end{aligned} (28)$$

This in particular implies that  $Y_{2n}X_{2n+1}Y_{2n+2}$  can be written as a product of  $UZ_{j-1}X_jZ_{j+1}U^{-1}$ . However, this is a contradiction since the former does not commute with D (as discussed around (17)), while the latter does.

We now argue for the more general statement without assuming U is Clifford. The idea it to show that if such a (unitary) symmetric entangler U exists, then there must exist a unitary operator  $\hat{U}$  in the **TST** gauged/KT transformed system such that it is invariant under the dual symmetry  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^v \times \hat{\mathbb{Z}}_2^v$  and maps the two symmetry breaking phases to each other. However, this is impossible since the two SPT phases correspond to two different patterns of symmetry breaking after gauging and hence cannot be related by a symmetric unitary operator.

In other words, we claim that the existence of an entangler  $\mathsf{U}$  satisfying

$$\begin{split} \mathsf{UD} &= \mathsf{DU} \,, & \mathsf{U} \eta^\mathrm{e} &= \eta^\mathrm{e} \mathsf{U} \,, \\ \mathsf{U} \, |\mathrm{cluster} \rangle &= |\mathrm{odd} \rangle \,\,, & \mathsf{U} \eta^\mathrm{o} &= \eta^\mathrm{o} \mathsf{U} \,, \end{split} \tag{29}$$

implies the existence of  $\hat{U}$  satisfying

$$\hat{\mathbf{U}}\hat{\mathbf{V}} = \hat{\mathbf{V}}\hat{\mathbf{U}}, \qquad \hat{\mathbf{U}}\hat{\eta}^{e} = \hat{\eta}^{e}\hat{\mathbf{U}}, 
\hat{\mathbf{U}} |\widehat{\text{cluster}}\rangle = |\widehat{\text{odd}}\rangle, \qquad \hat{\mathbf{U}}\hat{\eta}^{o} = \hat{\eta}^{o}\hat{\mathbf{U}}.$$
(30)

Here,  $|\widehat{\mathrm{cluster}}\rangle$  and  $|\widehat{\mathrm{odd}}\rangle$  are the symmetric ground states of  $\hat{H}_{\mathrm{cluster}}$  and  $\hat{H}_{\mathrm{odd}}$  in finite volume. In infinite volume, each of these two states splits into 4 superselection sectors, which break the  $\hat{\mathbb{Z}}_2^{\mathrm{e}} \times \hat{\mathbb{Z}}_2^{\mathrm{o}}$  symmetry spontaneously. We prove this claim in Appendix F by showing that (30) is the image of (29) under the KT transformation, which maps U, D,  $|\operatorname{cluster}\rangle$ ,  $|\operatorname{odd}\rangle$  to  $(\hat{\mathsf{U}},\hat{\mathsf{V}},|\operatorname{cluster}\rangle$ ,  $|\operatorname{odd}\rangle$ , respectively. However, (30) cannot be true since  $|\operatorname{cluster}\rangle$  represents a  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$ -preserving phase, while  $|\operatorname{odd}\rangle$  represents a spontaneous  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$ -breaking phase (see Section III B).

#### VI. CONCLUSIONS AND OUTLOOK

In the recent development of generalized symmetries, sometimes new symmetries are found in known quantum systems, and some other times they point to new systems.

In this paper we see an interesting interplay of these two scenarios. A new non-invertible symmetry D is discovered in the standard cluster model  $H_{\rm cluster}$ , which,

together with the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry, forms a Rep(D<sub>8</sub>) fusion category. This symmetry leads to two new commuting Pauli Hamiltonians  $H_{\text{odd}}$  and  $H_{\text{even}}$ , whose ground states  $|\text{odd}\rangle$  and  $|\text{even}\rangle$  are SPT states protected by Rep(D<sub>8</sub>). They are distinguished from the cluster Hamiltonian by their symmetry breaking patterns after the KT transformation/**TST** gauging.

While  $|\text{cluster}\rangle$ ,  $|\text{odd}\rangle$ ,  $|\text{even}\rangle$  are in the same SPT phase as far as the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry is concerned, they are distinguished by the non-invertible symmetry. This is reflected in the edge modes from the local projective algebra of the non-invertible symmetry. In contrast, the product state is not invariant under D, and there is no canonical notion of a trivial non-invertible SPT phase.

Finally, we argue that there is no  $\operatorname{Rep}(D_8)$  symmetric entangler relating these three SPT states. This observation is related to the fact that there is no stacking operation for systems with non-invertible symmetries. Moreover, it suggests that the classification of non-invertible SPT phases is not relative. This means that two distinct non-invertible SPT phases correspond to two different continuum field theories at long distances that do not differ just by a local counterterm. This is unlike the case of ordinary invertible symmetries, where the individual SPT phases have the same behavior at long distances, and their distinction is only visible from the edge modes on the interface between two such phases.

Our lattice construction provides some of the simplest examples of non-invertible SPT phases. First, our Hilbert space is a tensor product of qubits, and the Hamiltonians are written in terms of the Pauli operators. Second, the  $Rep(D_8)$  fusion category is one of the simplest (non-anomalous) non-invertible symmetries that admit SPT phases, and we constructed the lattice models for all of its possible SPT phases.

Our models are complementary to the non-invertible SPT models of [37, 52], which employ more general local Hilbert spaces than qubits. In particular, our  $\operatorname{Rep}(D_8)$  symmetry of the standard cluster model is different from the  $G \times \operatorname{Rep}(G)$  symmetry of the generalized cluster model discussed in [37] . When  $G = \mathbb{Z}_2$ , their model reduces to the standard cluster model (1), but their category symmetry reduces to the ordinary  $\mathbb{Z}_2 \times \operatorname{Rep}(\mathbb{Z}_2) \simeq \mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry, which is an invertible subgroup of our  $\operatorname{Rep}(D_8)$  fusion category.

There are several future directions:

- The key to our construction is that many of the non-invertible symmetries become (anomalous) invertible symmetries after gauging. Thus, the different non-invertible SPT phases are distinguished by the conventional spontaneous symmetry breaking patterns in the gauged systems. This construction can be readily generalized to many other non-invertible symmetries.
- It would be interesting to explore the anomaly inflow for non-invertible symmetry by the explicit non-invertible SPT lattice models.

<sup>&</sup>lt;sup>15</sup> Recall that KT transformation is implemented by a non-invertible operator that acts invertibly on the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  invariant operators/states.

- The Rep(D<sub>8</sub>) fusion category, along with its two other cousins Rep(Q<sub>8</sub>) and Rep(H<sub>8</sub>), are some of the simplest non-invertible symmetries that can be gauged. In Appendix B 3 we gauge Rep(D<sub>8</sub>) in steps, but it would be desirable to directly gauge the entire category on the lattice, mirroring the results of [32, 53, 54].
- Generalization to higher dimensions.

#### ACKNOWLEDGMENTS

We thank Nati Seiberg for discussions which initiated this project. We are grateful to Yichul Choi, Abhinav Prem, Wilbur Shirley, Nikita Sopenko, Nathanan Tantivasadakarn, Ruben Verresen, Tzu-Chieh Wei, Carolyn Zhang, and Yunqin Zheng for interesting discussions. We thank Da-Chuan Lu, Nikita Sopenko, and Yifan Wang for comments on the draft. SS gratefully acknowledges support from the U.S. Department of Energy grant DE-SC0009988, the Sivian Fund, and the Paul Dirac Fund at the Institute for Advanced Study. SHS is supported by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651444, SHS). The authors of this paper were ordered alphabetically.

#### Appendix A: Lattice gauging

In this appendix we review some standard facts about the untwisted (S) and twisted gauging (TST) of a  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry (2) on the lattice. These two gaugings are also known as (the square of) the Kramers-Wannier transformation and the KT transformation, respectively. Our discussion follows Appendix A of [24] closely.

We work with a closed chain of L sites, with the Pauli operators  $X_j, Z_j$  acting on each site. We assume a general finite-range Hamiltonian H that is  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric (such as the trivial Hamiltonian  $H_{\text{trivial}}$  or the cluster Hamiltonian  $H_{\text{cluster}}$ ).

## 1. KW<sup>⊗2</sup>: untwisted lattice gauging (S)

To gauge the  $\mathbb{Z}_2^{\rm e}$  symmetry, we introduce a gauge field qubit on every odd site. We denote the corresponding Pauli operators by  $\sigma_{2n-1}^x$ ,  $\sigma_{2n-1}^z$ . Similarly, to gauge the  $\mathbb{Z}_2^{\rm e}$  symmetry, we introduce a gauge field qubit on every even site, and denote the corresponding Pauli operators by  $\sigma_{2n}^x$ ,  $\sigma_{2n}^z$ . We impose the  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm e}$  Gauss law at every site:

$$G_i = \sigma_{i-1}^z X_i \sigma_{i+1}^z = 1.$$
 (A1)

We then couple the gauge fields to the original Hamiltonian H minimally by replacing every  $Z_{j-1}Z_{j+1}$  by

 $Z_{j-1}\sigma_j^x Z_{j+1}$ . This leads to a gauged Hamiltonian H', written in terms of  $X_j, Z_j, \sigma_j^x, \sigma_j^z$ .

Not every Pauli operator is gauge invariant and commutes with the Gauss law constraint. To proceed, we define a new set of gauge-invariant variables  $X'_j, Z'_j$  that commute with the Gauss law:

$$X'_{i} = Z_{j-1}\sigma_{i}^{x}Z_{j+1}, \quad Z'_{i} = \sigma_{i}^{z}.$$
 (A2)

Clearly,  $X'_{i}, Z'_{i}$  obey the standard Pauli algebra.

Now we can write the gauged Hamiltonian in terms of new, gauge-invariant variables. The minimal coupling term becomes  $Z_{j-1}\sigma_j^x Z_{j+1} = X_j'$ , and the operator  $X_j$  is replaced by  $Z_{j-1}' Z_{j+1}'$  using the Gauss law. All in all, the untwisted gauging maps the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  local operators as

$$\mathbf{S}: \quad X_j \leadsto Z'_{j-1} Z'_{j+1} \,, \quad Z_{j-1} Z_{j+1} \leadsto X'_j \,. \tag{A3}$$

We use  $\rightsquigarrow$  to emphasize that this transformation is not implemented by a unitary operator. Indeed, if we identify the new variables with the original ones  $X'_j = X_j, Z'_j = Z_j$ , then (A3) is implemented by the non-invertible operator D as in (7):

$$DX_j = Z_{j-1}Z_{j+1}D$$
,  $DZ_{j-1}Z_{j+1} = X_jD$ . (A4)

In the gauged Hamiltonian H', the original  $\mathbb{Z}_2^{\mathrm{e}} \times \mathbb{Z}_2^{\mathrm{o}}$  global symmetry  $\eta^{\mathrm{e}}, \eta^{\mathrm{o}}$  in (2) is gauged and is no longer a global symmetry of H'. Rather, the gauged Hamiltonian is invariant under a dual global  $\mathbb{Z}_2^{\mathrm{e}'} \times \mathbb{Z}_2^{\mathrm{o}'}$  symmetry generated by

$$\eta^{e'} = \prod_{j:\text{even}} X'_j, \quad \eta^{o'} = \prod_{j:\text{odd}} X'_j.$$
(A5)

This is the lattice version of the dual symmetry (also known as the quantum symmetry or the orbifold symmetry) in [55].

Applying the untwisted gauging on the trivial and the cluster Hamiltonians

$$H_{\text{trivial}} = -\sum_{j=1}^{L} X_j,$$

$$H_{\text{cluster}} = -\sum_{j=1}^{L} Z_{j-1} X_j Z_{j+1},$$
(A6)

we find the two gauged Hamiltonians in (9)

$$H'_{\text{trivial}} = -\sum_{j=1}^{L} Z'_{j-1} Z'_{j+1},$$

$$H'_{\text{cluster}} = -\sum_{j=1}^{L} Z'_{j-1} X'_{j} Z'_{j+1}.$$
(A7)

We see that the trivial phase becomes a  $\mathbb{Z}_2^{e'} \times \mathbb{Z}_2^{o'}$  spontaneously symmetry broken phase, while the cluster phase remains invariant and preserves the global symmetry.

Via gauging, one can distinguish the two SPT phases by the string order parameters. In the two gauged models, the local order parameters are  $\langle Z'_{2n-1} \rangle$  and  $\langle Z'_{2n} \rangle$ , which distinguish the spontaneous symmetry breaking phase from the unbroken phase. For large l,  $\langle Z'_{0}Z'_{2l} \rangle \sim \langle Z'_{0} \rangle \langle Z'_{2l} \rangle$  is equal to 0 and 1 in the symmetry preserving and breaking phases, respectively, up to exponentially small corrections. Before gauging, these local, order parameters come from certain non-local, string-like order parameters. More specifically, doing an inverse of (4), we find the string order parameter  $\langle X_{j}X_{j+2}\cdots X_{j+2l} \rangle$  that distinguishes the trivial state from the cluster state. Their correlation functions are

$$\langle X_j X_{j+2} \cdots X_{j+2l} \rangle \sim \begin{cases} 0 & \text{product state} \\ \text{constant} & \text{cluster state} \end{cases}$$
 (A8)

up to corrections exponentially small in l.

### 2. KT: twisted lattice gauging (TST)

The twisted gauging **TST** is related to the untwisted one **S** by conjugating with the cluster entangler  $V = \prod_{j} CZ_{j,j+1}$ , which implements the **T** transformation:

$$\mathbf{T}: \quad \begin{array}{ll} X_j \mapsto \mathsf{V} X_j \mathsf{V}^{-1} = Z_{j-1} X_j Z_{j+1} \,, \\ Z_j \mapsto \mathsf{V} Z_j \mathsf{V}^{-1} = Z_j \,. \end{array} \tag{A9}$$

Let us break down the **TST** gauging in steps. First, we conjugate the Hamiltonian by V. Second, we perform the untwisted gauging to map it to a gauged Hamiltonian as in Appendix A 1. We denote the gauge-invariant variables in the gauged Hamiltonian as  $\hat{X}_j$ ,  $\hat{Z}_j$ , to distinguish from the primed variables for the untwisted gauging. Third, we conjugate the gauged Hamiltonian by

$$\hat{V} = \prod_{j=1}^{L} \hat{CZ}_{j,j+1} = \prod_{j=1}^{L} \frac{1 + \hat{Z}_j + \hat{Z}_{j+1} - \hat{Z}_j \hat{Z}_{j+1}}{2}$$
(A10)

This sequence of operation results in the following gauging map, commonly denoted as **TST** in the field theory literature, on the  $\mathbb{Z}_2^{\text{e}} \times \mathbb{Z}_2^{\text{o}}$  symmetric local operators:

$$\mathbf{TST}: \quad X_j \leadsto \hat{X}_j \,, \quad Z_{j-1} Z_{j+1} \leadsto \hat{Z}_{j-1} \hat{X}_j \hat{Z}_{j+1} \,. \tag{A11}$$

Again, we use  $\rightsquigarrow$  to emphasize that this transformation is not implemented by a unitary operator. Indeed, if we identify the new variables with the original ones  $\hat{X}_j = X_j, \hat{Z}_j = Z_j$ , then (A3) is implemented by the non-invertible operator VDV [46]:

$$\begin{aligned} & (\mathsf{VDV}) X_j = X_j (\mathsf{VDV}) \,, \\ & (\mathsf{VDV}) Z_{j-1} Z_{j+1} = Z_{j-1} X_j Z_{j+1} (\mathsf{VDV}) \,. \end{aligned} \tag{A12}$$

This is the KT transformation.

The twisted gauging maps the original Hamiltonian H of  $X_j, Z_j$  to a gauged Hamiltonian  $\hat{H}$  of  $\hat{X}_j, \hat{Z}_j$ . Again,

operation	transformation	operator
S gauging	$X_j \leadsto Z'_{j-1} Z'_{j+1}$	D
$(Kramers-Wannier^{\otimes 2})$	$Z_{j-1}Z_{j+1} \leadsto X'_j$	
TST gauging	$X_j \leadsto \hat{X}_j$	VDV
(Kennedy-Tasaki)	$Z_{j-1}Z_{j+1} \leadsto \hat{Z}_{j-1}\hat{X}_j\hat{Z}_{j+1}$	
<b>T</b> : stacking with SPT	$X_j \mapsto Z_{j-1} X_j Z_{j+1}$	V
(Cluster entangler)	$Z_j \mapsto Z_j$	

TABLE III. The table discusses two different gaugings of a  $\mathbb{Z}_2 \times \mathbb{Z}_2$  global symmetry, which in the continuum field theory literature are often denoted as **S** and **TST**, as well as the stacking by the cluster SPT phase (**T**). The middle column shows the action of these transformations on the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetric local operators. The rightmost column gives the lattice operators implementing these transformations if we identify the gauged and ungauged variables by dropping the primes/hats.

the gauged Hamiltonian is invariant under a dual global  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o$  symmetry generated by

$$\hat{\eta}^{\mathrm{e}} = \prod_{j:\mathrm{even}} \hat{X}_j, \quad \hat{\eta}^{\mathrm{o}} = \prod_{j:\mathrm{odd}} \hat{X}_j.$$
 (A13)

As examples, applying the twisted gauging on  $H_{\text{trivial}}$  and  $H_{\text{cluster}}$  gives (12).

We summarize these operations in Table III.

# Appendix B: Non-invertible symmetry under gauging

So far we have only assumed the original Hamiltonian H is invariant under an invertible  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  global symmetry. The twisted gauged Hamiltonian  $\hat{H}$  is then invariant under a dual  $\hat{\mathbb{Z}}_2^{\rm e} \times \hat{\mathbb{Z}}_2^{\rm o}$  symmetry. Now we further assume H commutes with the non-invertible global symmetry D. What happens to D under the twisted gauging **TST**?

1. Rep(D<sub>8</sub>) 
$$\rightarrow \hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$$

Below we argue that the non-invertible symmetry D of H becomes an invertible  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$  symmetry, generated by  $\hat{\mathsf{V}}$  in (A10) after the twisted gauging. More precisely, we will show that any Hamiltonian that is invariant under  $\eta^e, \eta^o, \mathsf{D}$  (which forms a  $\mathrm{Rep}(\mathsf{D}_8)$  fusion category) is mapped to a gauged Hamiltonian  $\hat{H}$  that is invariant under a dual  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^\mathsf{V}$  symmetry generated by  $\hat{\eta}^e, \hat{\eta}^o, \hat{\mathsf{V}}$  under the twisted gauging.

For instance, the double-Ising Hamiltonian

$$H_{\text{Ising}^2} = -\sum_{j=1}^{L} (Z_{j-1}Z_{j+1} + X_j)$$
 (B1)

is invariant under  $D, \eta^e, \eta^o$ . Applying the **TST** gauging turns it into

$$\hat{H}_{LG} = -\sum_{j=1}^{L} \left( \hat{Z}_{j-1} \hat{X}_{j} \hat{Z}_{j+1} + \hat{X}_{j} \right),$$
 (B2)

which indeed is invariant under  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$ . This Hamiltonian (up to a unitary transformation) is the boundary theory for the 2+1d Levin-Gu SPT phase labeled by  $H^3(\mathbb{Z}_2, U(1)) = \mathbb{Z}_2$  [56].

To prove this generally, it suffices to focus on the action of D on the local terms in the Hamiltonian H, which are all  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric. On the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric sector, the KT transformation (A11) is invertible, and we can conjugate the action of D in (A4) by the KT transformation. This results in a transformation in the gauged model that acts in the same way as  $\hat{V}$ :

$$\hat{\mathsf{V}}: \quad \begin{array}{l} \hat{X}_{j} \mapsto \hat{Z}_{j-1} \hat{X}_{j} \hat{Z}_{j+1} \,, \\ \hat{Z}_{j-1} \hat{Z}_{j+1} \mapsto \hat{Z}_{j-1} \hat{Z}_{j+1} \,. \end{array} \tag{B3}$$

This shows at the level of the Hamiltonians that D becomes  $\hat{V}$  after the twisted gauging.

The fact that D becomes  $\hat{V}$  can be understood more intuitively if we identify the hatted variables in the gauged Hamiltonian with the original one as  $\hat{X}_j = X_j$ ,  $\hat{Z}_j = Z_j$  and  $\hat{V} = V$ . Under this identification, the twisted gauging/KT transformation is implemented by VDV in the sense of (A12). More generally, given any  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric operator  $O = O(X_j, Z_j)$ , its image under the KT transformation is given by  $\hat{O} = \hat{O}(X_j, Z_j)$  satisfying

$$(VDV) O = \hat{O}(VDV). \tag{B4}$$

Now the fact that  $\mathsf{D}$  becomes  $\mathsf{V}$  under the KT transformation follows from the following identity

$$(VDV)D = 2V(VDV). (B5)$$

This relation is reminiscent of the relation TSTS = ST in the modular group.

To show this relation, we first note that both the left and right hand sides of the equation above have the same action, in the sense of (B4), on  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric operators. This implies that the restriction of these two operators to the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric sector is the same up to an overall constant, i.e.,

$$\Pi (VDVD) \Pi = c \Pi (2DV) \Pi,$$
 (B6)

where  $\Pi = \frac{1+\eta^e}{2} \frac{1+\eta^o}{2}$  and  $c \in \mathbb{C}$ . Since  $\Pi$  commutes with V and  $\Pi D = D\Pi = D$ , we find that  $VDVD = c \, 2DV$ .

To fix the overall constant we compare the action of these operators on the cluster state. We begin by

$$DV | cluster \rangle = D | + + \cdots + \rangle$$

$$= D^{e}D^{o}T | + + \cdots + \rangle$$

$$= |00\rangle + |01\rangle + |10\rangle + |11\rangle ,$$
(B7)

where  $|z_{\rm e}z_{\rm o}\rangle = |z_{\rm e}z_{\rm e}\cdots z_{\rm e}\rangle_{\rm even}\otimes |z_{\rm o}z_{\rm o}\cdots z_{\rm o}\rangle_{\rm odd}$  for  $z_{\rm e},z_{\rm o}=0,1.$  This follows from [24]

$$\begin{array}{l} D^{\rm e} \left| + + \cdots + \right\rangle_{\rm even} = \left| 00 \cdots 0 \right\rangle_{\rm even} + \left| 11 \cdots 1 \right\rangle_{\rm even} \; , \\ D^{\rm o} \left| + + \cdots + \right\rangle_{\rm odd} = \left| 00 \cdots 0 \right\rangle_{\rm odd} + \left| 11 \cdots 1 \right\rangle_{\rm odd} \; . \end{array} \tag{B8}$$

Next, we consider the action of VDVD:

$$\begin{aligned} \mathsf{VDVD} \left| \mathsf{cluster} \right\rangle &= 2\mathsf{VD} \left| + + \dots + \right\rangle \\ &= 2\mathsf{V} \left( \left| 00 \right\rangle + \left| 01 \right\rangle + \left| 10 \right\rangle + \left| 11 \right\rangle \right) \\ &= 2 \left( \left| 00 \right\rangle + \left| 01 \right\rangle + \left| 10 \right\rangle + \left( -1 \right)^L \left| 11 \right\rangle \right). \end{aligned} \tag{B9}$$

Here we have used the relation  $D | \text{cluster} \rangle = 2 | \text{cluster} \rangle$ , proven in Appendix E, and the fact that

$$CZ_{i,j+1} | z_e z_o \rangle = (-1)^{z_e z_o} | z_e z_o \rangle$$
. (B10)

Since L is taken to be even throughout, comparing (B7) with (B9) fixes c = 1. We have thus proved (B5).

### 2. Type III anomaly and the Levin-Gu model

We now show that the dual  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$  symmetry generated by (A10) and (A13) has an 't Hooft anomaly of type III.

While the anomaly of the symmetry operators  $\hat{\eta}^{e}$ ,  $\hat{\eta}^{o}$ ,  $\hat{V}$  does not depend on the choice of the symmetric Hamiltonian, for concreteness, we consider the Levin-Gu Hamiltonian in (B2). (We assume L to be a multiple of 4 for simplicity.)

It is known that the diagonal  $\mathbb{Z}_2$  symmetry generated by

$$\hat{\eta}^{e} \hat{\eta}^{o} \hat{\mathsf{V}} = \prod_{j=1}^{L} \hat{X}_{j} \,\hat{\mathsf{CZ}}_{j,j+1} \,. \tag{B11}$$

has an 't Hooft anomaly described by  $e^{i\pi\int\hat{A}\cup\hat{A}\cup\hat{A}}$  [57] (see also [58]). Below we show that the full  $\hat{\mathbb{Z}}_2^{\rm e}\times\hat{\mathbb{Z}}_2^{\rm o}\times\hat{\mathbb{Z}}_2^{\rm v}$  symmetry has an 't Hooft anomaly described by the 2+1d invertible field theory

$$e^{i\pi\int \hat{A}^{e}\cup\hat{A}^{o}\cup\hat{A}^{V}},$$
 (B12)

where  $\hat{A}^{\rm e}$ ,  $\hat{A}^{\rm o}$ ,  $\hat{A}^{\rm V}$  are background  $\mathbb{Z}_2$  gauge fields. This is referred to as a type III anomaly [47, 48] because it involves all three generators of  $\hat{\mathbb{Z}}_2^{\rm e} \times \hat{\mathbb{Z}}_2^{\rm v} \times \hat{\mathbb{Z}}_2^{\rm v}$ . As a consistency check, this reduces to the Levin-Gu anomaly for the diagonal symmetry by setting  $\hat{A}^{\rm e} = \hat{A}^{\rm o} = \hat{A}^{\rm v}$ . One feature of this type III anomaly is that out of the 7  $\mathbb{Z}_2$  subgroup of  $\hat{\mathbb{Z}}_2^{\rm e} \times \hat{\mathbb{Z}}_2^{\rm o} \times \hat{\mathbb{Z}}_2^{\rm v}$ , only the one generated by  $\hat{\eta}^{\rm e}\hat{\eta}^{\rm o}\hat{\mathsf{V}}$  has an anomaly.

One diagnostic of the type III anomaly is the projective algebra of two of the  $\mathbb{Z}_2$  operators in the Hamiltonian twisted by the third  $\mathbb{Z}_2$ . See [59, 60] (and also [61]) for general discussions of anomalies on the lattice. Consider

the Levin-Gu Hamiltonian twisted by the  $\hat{\eta}^{e}$  symmetry at site 1:

$$\hat{H}_{\eta^{e}} = \dots - \left( \hat{X}_{L} + \hat{Z}_{L-1} \hat{X}_{L} \hat{Z}_{1} \right) - \left( \hat{X}_{1} - \hat{Z}_{L} \hat{X}_{1} \hat{Z}_{2} \right) - \left( \hat{X}_{2} + \hat{Z}_{1} \hat{X}_{2} \hat{Z}_{3} \right) - \dots,$$
(B13)

where  $\cdots$  represents terms of the form  $\hat{Z}_{j-1}\hat{X}_j\hat{Z}_{j+1}+\hat{X}_j$ . The symmetry operators in the  $\hat{\eta}^e$ -twisted problem are

$$\hat{\eta}_{\mathrm{e}}^{\mathrm{e}} = \prod_{j:\mathrm{even}} \hat{X}_{j} \,, \quad \hat{\eta}_{\mathrm{e}}^{\mathrm{o}} = \prod_{j:\mathrm{odd}} \hat{X}_{j} \,, \quad \hat{\mathsf{V}}_{\mathrm{e}} = \hat{Z}_{1} \hat{\mathsf{V}} \,, \quad \text{(B14)}$$

where the subscript e means that these are symmetry operators that commute with the  $\hat{\eta}^{\text{e}}$ -twisted Hamiltonian  $\hat{H}_{\eta^{\text{e}}}$ . We find a projective algebra in this twisted problem:

$$\hat{\eta}_{e}^{o} \hat{V}_{e} = -\hat{V}_{e} \hat{\eta}_{e}^{o}, \qquad (B15)$$

which is the hallmark of the type III anomaly. Related projective algebras have also been discussed in [48].

To summarize, the twisted gauging maps a non-invertible symmetry  $\operatorname{Rep}(D_8)$  to an anomalous invertible symmetry  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V$ :

$$\operatorname{Rep}(D_8) \xrightarrow{\mathbf{TST}} \underbrace{\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^V}_{\text{type III anomaly}}$$
(B16)

This fact was known in the continuum [62, 63], and here we give a lattice derivation. It is also consistent with the fact that the two fusion categories  $\operatorname{Rep}(D_8)$  and  $\operatorname{Vec}_{\mathbb{Z}_2^3}^\omega$  (with  $\omega \in H^3(\mathbb{Z}_2^3, U(1))$  chosen to be the type III cocycle [47]) have the same Drinfeld center, i.e., the same 2+1d topological order (see, e.g., [40, 64]). More generally, it is common that an anomalous invertible symmetry is mapped to a non-invertible symmetry under gauging, and vice versa [65]. For instance, the non-invertible Kramers-Wannier symmetry of the critical Ising model arises from the chiral fermion parity [23, 66, 67], and the non-invertible chiral symmetry of the 3+1d quantum electrodynamics arises from the Adler-Bell-Jackiw anomaly [68, 69].

## 3. Gauging Rep( $D_8$ ) in steps

The algebra (8) implies that  $D, \eta^e, \eta^o$  form one of the four  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  Tambara-Yamagami (TY) fusion categories [70] (see also [26, 51, 54]). Among these four TY categories, three of them admit fiber functors (i.e, the mathematical description of the non-invertible SPT phases), which are  $\operatorname{Rep}(D_8)$ ,  $\operatorname{Rep}(Q_8)$ , and  $\operatorname{Rep}(H_8)$ . Here  $Q_8$  is the quaternion group of order 8 and  $H_8$  is a non-grouplike Hopf algebra of dimension 8 known as the Kac-Paljutkin algebra. These three fusion categories are free of generalized 't Hooft anomaly both in the sense that they can be gauged [32, 53, 54], and they are compatible with a

trivially gapped phase, i.e., a non-invertible SPT phase. More specifically,  $Rep(D_8)$  admits three fiber functors, while  $Rep(Q_8)$ , and  $Rep(H_8)$  each has a unique fiber functor [26, 38, 70].

Here we show that our lattice symmetry  $D, \eta^e, \eta^o$  form the Rep(D<sub>8</sub>). To prove this, we use the following fact. Gauging a finite G global symmetry in 1+1d gives rise to a dual global symmetry described by Rep(G) [51] (see also [71]). Below we gauge the non-invertible symmetry generated by  $D, \eta^e, \eta^o$  and show that the gauged theory has a dual  $D_8$  symmetry. This determines the fusion category of the ungauged system to be Rep(D<sub>8</sub>).

Instead of gauging the non-invertible symmetry  $\operatorname{Rep}(D_8)$  in one shot, we do it in two steps, with each step only consisting of gauging ordinary invertible symmetries. In the first step, we twisted gauge the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry, i.e., the **TST** gauging, i.e., the KT transformation. This was done in Appendix B, which turns the non-invertible symmetry (8) into an anomalous  $\hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^v$  symmetry generated by  $\hat{\eta}^e$ ,  $\hat{\eta}^o$ , and  $\hat{V}$ .

In the second step, we gauge the  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$  symmetry generated by  $\hat{\mathsf{V}}$ . To do this gauging, we introduce gauge fields  $\tilde{X}_i$  on sites and impose the Gauss law constraints

$$G_{j,j+1} = \tilde{X}_j \hat{C} Z_{j,j+1} \tilde{X}_{j+1}$$
 (B17)

to obtain the Hilbert space of the (second) gauged model. In this model, we find a dual  $D_8$  symmetry generated by

$$\tilde{\eta}^{e} = \prod_{j:\text{even}} \hat{X}_{j} \prod_{j:\text{odd}} \frac{1 + \hat{Z}_{j} + \tilde{Z}_{j} - \hat{Z}_{j} \tilde{Z}_{j}}{2},$$

$$\tilde{\eta}^{o} = \prod_{j:\text{odd}} \hat{X}_{j} \prod_{j:\text{even}} \frac{1 + \hat{Z}_{j} + \tilde{Z}_{j} - \hat{Z}_{j} \tilde{Z}_{j}}{2},$$

$$\tilde{V} = \prod_{j} \tilde{Z}_{j}.$$
(B18)

One can check the operators above commute with the Gauss's law constraints and generate a  $D_8$  symmetry since  $(\tilde{\eta}^e)^2 = (\tilde{\eta}^o)^2 = \tilde{V}^2 = 1$  and  $\tilde{\eta}^e \tilde{\eta}^o = \tilde{V} \hat{\eta}^o \hat{\eta}^e$ .

As a concrete example, we again start with the double-Ising Hamiltonian in (B1) which has a Rep(D<sub>8</sub>) symmetry. The first step of **TST** gauging turns it into the Levin-Gu model  $\hat{H}_{LG}$  in (B2). In the second step, gauging  $\hat{\mathbb{Z}}_2^{\mathsf{V}}$  turns  $\hat{H}_{LG}$  into

$$\tilde{H} =$$

$$-\sum_{j=1}^{L} X_{j} \left( \frac{1+\tilde{Z}_{j}}{2} (1+\hat{Z}_{j-1}\hat{Z}_{j+1}) + \frac{1-\tilde{Z}_{j}}{2} (\hat{Z}_{j-1}+\hat{Z}_{j+1}) \right)$$
(B19)

which indeed is invariant under the  $D_8$  symmetry generated by (B18).

To conclude, we gauge the symmetry generated by  $\eta^{e}$ ,  $\eta^{o}$ , D of a general Hamiltonian in two steps:

$$\operatorname{Rep}(D_8) \xrightarrow{\operatorname{gauge} \mathbb{Z}_2^e \times \mathbb{Z}_2^o} \hat{\mathbb{Z}}_2^e \times \hat{\mathbb{Z}}_2^o \times \hat{\mathbb{Z}}_2^{\mathsf{V}} \xrightarrow{\operatorname{gauge} \hat{\mathbb{Z}}_2^{\mathsf{V}}} D_8$$
(B20)

In the end, we find an ordinary global symmetry  $D_8$  in the gauged Hamiltonian. This implies that the original non-invertible symmetry is  $Rep(D_8)$ .

### Appendix C: SPT phases as invertible field theories

In this appendix we review the continuum description of the  $\mathbb{Z}_2 \times \mathbb{Z}_2$  SPT phases as invertible field theories (see, e.g., [72]). We then gauge these SPT phases in two different ways. Our discussion follows [22, 26] closely.

We first review the gauging of a non-anomalous  $\mathbb{Z}_2$  global symmetry in 1+1d quantum field theory (QFT). The generalization to gauging a  $\mathbb{Z}_2 \times \mathbb{Z}_2$  symmetry will be straightforward. Consider a general 1+1d QFT with a  $\mathbb{Z}_2$  global symmetry. Let the partition function on a closed spacetime 2-manifold  $\Sigma$  be Z[A], where  $A \in H^1(\mathbb{Z}_2, U(1))$  is the background  $\mathbb{Z}_2$  gauge field. The gauged QFT has a dual symmetry  $\mathbb{Z}'_2$  [55]. Let A' be the background gauge field for  $\mathbb{Z}'_2$ , then the partition function of the gauged theory is

$$Z'[A'] = \frac{1}{\sqrt{|H^1(\Sigma, \mathbb{Z}_2)|}} \sum_{a \in H^1(\Sigma, \mathbb{Z}_2)} Z[a] \exp\left[i\pi \oint_{\Sigma} a \cup A'\right].$$
(C1)

The normalization is up to the ambiguity of an Euler counterterm  $\lambda^{\chi(\Sigma)}$  with  $\lambda > 0$ . From (C1), it is clear that if we further gauge the dual  $\mathbb{Z}'_2$  global symmetry of the gauged QFT, we retrieve the original QFT.

The continuum description of a product state  $|++...+\rangle$  can be chosen to be the trivial QFT, whose partition function is 1 on every closed spacetime 2-manifold  $\Sigma$ :

$$Z_{\text{trivial}}[A^{\text{e}}, A^{\text{o}}] = 1.$$
 (C2)

Once this choice is made, the continuum description for the cluster state is given by a nontrivial invertible field theory of the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^e$  background gauge fields  $A^e$ ,  $A^o$ :

$$Z_{\text{cluster}}[A^{\text{e}}, A^{\text{o}}] = \exp\left[i\pi \oint_{M} A^{\text{e}} \cup A^{\text{o}}\right]$$
 (C3)

Note that  $A^{e}$ ,  $A^{o}$  are classical, background  $\mathbb{Z}_{2}$  gauge fields, and there is no path integral over them. Therefore, the partition function of this invertible field theory is always a phase on every closed 2-manifold  $\Sigma$  [73].

### 1. S gauging in the continuum

We now gauge the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  global symmetry of these two invertible field theories. This (untwisted) gauging is referred to as the **S** gauging in the literature (e.g., [12, 22, 74, 75]), representing the corresponding element in the modular group.

We promote the background gauge fields to be dynamical and denote them by  $a^{\rm e}, a^{\rm o}$ . We also introduce the

background gauge fields  $A^{e'}, A^{o'}$  for the dual  $\mathbb{Z}_2^{e'} \times \mathbb{Z}_2^{o'}$  global symmetry. Gauging the trivial theory gives:

$$Z'_{\text{trivial}}[A^{e'}, A^{o'}] = \frac{1}{\sqrt{|H^1(\Sigma, \mathbb{Z}_2 \times \mathbb{Z}_2)|}} \times \sum_{(a^e, a^o) \in H^1(\Sigma, \mathbb{Z}_2 \times \mathbb{Z}_2)} \exp \left[ i\pi \oint_{\Sigma} (a^e \cup A^{e'} + a^o \cup A^{o'}) \right]$$

$$= \sqrt{|H^1(\Sigma, \mathbb{Z}_2 \times \mathbb{Z}_2)|} \ \delta(A^{e'}) \delta(A^{o'}) ,$$
(C4)

where  $\delta(A)=1$  if A=0, and  $\delta(A)=0$  otherwise. The gauged theory is a 1+1d topological quantum field theory (TQFT) describing the spontaneous symmetry breaking (SSB) of  $\mathbb{Z}_2^{e'} \times \mathbb{Z}_2^{o'}$  to nothing. Indeed, when  $\Sigma=T^2$  and  $A^{e'}=A^{o'}=0$ ,  $Z'_{\text{trivial}}=4$ , which is the number of superselection sectors in a  $\mathbb{Z}_2^{e'} \times \mathbb{Z}_2^{o'}$  broken phase.

On the other hand, the **S** gauging of  $Z_{\text{cluster}}$  gives back the original partition function:

$$Z'_{\text{cluster}}[A^{e'}, A^{o'}] = \frac{1}{\sqrt{|H^1(\Sigma, \mathbb{Z}_2 \times \mathbb{Z}_2)|}} \sum_{(a^e, a^o) \in H^1(\Sigma, \mathbb{Z}_2 \times \mathbb{Z}_2)}$$

$$\exp\left[i\pi \oint_{\Sigma} (a^e \cup a^o + a^e \cup A^{e'} + a^o \cup A^{o'})\right]$$

$$= \exp\left[i\pi \oint_{\Sigma} A^{e'} \cup A^{o'}\right],$$
(C5)

where the sum over  $a^{e}$  sets  $a^{o} = A^{e'}$ . We conclude that

$$\mathbf{S}: \quad \begin{array}{c} \text{trivial} \rightarrow \text{SSB} \\ \text{cluster} \rightarrow \text{cluster} \end{array}$$
 (C6)

This reproduces the lattice gauging in (A7). Since  $Z_{\rm cluster}$  is invariant under the **S** gauging, the half-gauging argument in [41, 42] leads to a non-invertible global symmetry in the cluster SPT phase. (See [17] for a review of half-gauging.) In [43], we will gauge in half of the lattice to obtain the duality defect in the cluster model, extending the discussion in [24, 76] to the cluster model.

#### 2. TST gauging in the continuum

Next, we perform the twisted gauging labeled by **TST**. Here **T** corresponds to stacking the partition function by the invertible field theory  $Z_{\text{cluster}}$  in (C3):

$$\mathbf{T}: Z[A^{e}, A^{o}] \mapsto \exp\left[i\pi \oint_{\Sigma} A^{e} \cup A^{o}\right] Z[A^{e}, A^{o}].$$
 (C7)

The  $\mathbf{TST}$  gauging maps the trivial and the cluster theories to

$$\mathbf{TST}: \quad \begin{array}{c} \text{trivial} & \rightarrow \text{trivial} \\ \text{cluster} & \rightarrow \text{SSB} \end{array}$$
 (C8)

matching the lattice results in (12).

# Appendix D: Interface between the cluster and product states

We review the edge modes for the cluster model. For convenience of generalization to the case of non-invertible symmetries, we view the boundary condition for the cluster model as an interface between the cluster model and a trivial paramagnet. More specifically, instead of working with an open chain, our space is a closed chain of L sites. The interface Hamiltonian is

$$H_{\text{interface}} = -Z_1 X_2 Z_3 - Z_2 X_3 Z_4 - \dots - Z_{\ell-2} X_{\ell-1} Z_{\ell} - X_{\ell+1} - \dots - X_L.$$

Here  $1 < \ell < L$  and we assume  $\ell$  and L to be both even for simplicity. In other words, we have the cluster Hamiltonian in the region  $j=1,\cdots,\ell$  and the trivial paramagnet Hamiltonian in the region  $j=\ell+1,\cdots,L$ . The interface Hamiltonian enjoys the  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  symmetry generated by  $\eta^{\rm e} = \prod_{j=1,j:{\rm even}}^L X_j, \eta^{\rm o} = \prod_{j=1,j:{\rm odd}}^L X_j$  on a closed chain. The advantage of this approach is that we never need to cut open the symmetry operators.

A state  $|\psi\rangle$  in the ground space of  $H_{\text{interface}}$  satisfies

$$Z_1X_2Z_3 = Z_2X_3Z_4 = \dots = Z_{\ell-2}X_{\ell-1}Z_{\ell} = 1$$
  
 $X_{\ell+1} = \dots = X_L = 1$ . (D2)

Since there are only L-2 commuting Pauli constraints for L qubits, the ground space is 4-dimensional, signaling the edge modes at the interfaces.

For any state  $|\psi\rangle$  in the ground space, we can simplify the action of the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry using (D2). The global  $\eta^e, \eta^o$  operators factorize into products of local operators on the two interfaces:

$$\begin{split} &\eta^{\mathrm{e}} \left| \psi \right\rangle = \eta_{\mathrm{L}}^{\mathrm{e}} \eta_{\mathrm{R}}^{\mathrm{e}} \left| \psi \right\rangle \;, \quad \eta_{\mathrm{L}}^{\mathrm{e}} = Z_{1} \;, \quad \eta_{\mathrm{R}}^{\mathrm{e}} = Z_{\ell-1} X_{\ell} \;, \\ &\eta^{\mathrm{o}} \left| \psi \right\rangle = \eta_{\mathrm{L}}^{\mathrm{o}} \eta_{\mathrm{R}}^{\mathrm{o}} \left| \psi \right\rangle \;, \quad \eta_{\mathrm{L}}^{\mathrm{o}} = X_{1} Z_{2} \;, \quad \eta_{\mathrm{R}}^{\mathrm{o}} = Z_{\ell} \;. \end{split} \tag{D3}$$

Here L and R stand for the left and right interfaces around j=1 and  $j=\ell$ , respectively. While  $\eta^{\rm e}$  and  $\eta^{\rm o}$  act linearly on the ground space (i.e.,  $\eta^{\rm e}\eta^{\rm o}=\eta^{\rm o}\eta^{\rm e}$ ), their local factors obey a projective algebra:

$$\eta_{\mathrm{L}}^{\mathrm{o}} \eta_{\mathrm{L}}^{\mathrm{e}} = -\eta_{\mathrm{L}}^{\mathrm{e}} \eta_{\mathrm{L}}^{\mathrm{o}}, 
\eta_{\mathrm{R}}^{\mathrm{o}} \eta_{\mathrm{R}}^{\mathrm{e}} = -\eta_{\mathrm{R}}^{\mathrm{e}} \eta_{\mathrm{R}}^{\mathrm{o}}.$$
(D4)

Equivalently, the left and right local factors  $\eta_{L,R}^{o}$  are charged under  $\eta^{e}$ , and vice versa. This local projective algebra signals the relative difference in their SPT phases between the cluster state and the product state.

#### Appendix E: Action of D on the SPT states

In this appendix, we compute the action of the non-invertible symmetry D on various SPT and interface states. We begin with the three SPT states.

1. 
$$D | \mathbf{cluster} \rangle = 2 | \mathbf{cluster} \rangle$$

To find the action of D on the cluster state, we use the expression  $D = TD^{e}D^{o}$ , where

$$D^{e} = e^{\frac{2\pi i L}{16}} \frac{1 + \eta^{e}}{\sqrt{2}} \frac{1 - iX_{L}}{\sqrt{2}} \cdots \frac{1 - iZ_{4}Z_{2}}{\sqrt{2}} \frac{1 - iX_{2}}{\sqrt{2}},$$

$$D^{o} = e^{\frac{2\pi i L}{16}} \frac{1 + \eta^{o}}{\sqrt{2}} \frac{1 - iZ_{1}Z_{L-1}}{\sqrt{2}} \cdots \frac{1 - iX_{3}}{\sqrt{2}} \frac{1 - iZ_{3}Z_{1}}{\sqrt{2}}.$$
(E1)

Here we have used a different presentation of D<sup>o</sup> than the one in (6).

We first find the action of  $D^o$  on the cluster states as  $D^o | \text{cluster} \rangle =$ 

$$\sqrt{2}e^{\frac{2\pi iL}{16}}\frac{1-iX_2}{\sqrt{2}}\frac{1-iZ_2Z_4}{\sqrt{2}}\cdots\frac{1-iX_L}{\sqrt{2}}|\text{cluster}\rangle$$
, (E2)

where we have repeatedly used  $Z_{j-1}X_jZ_{j+1}$  |cluster $\rangle$  = |cluster $\rangle$  to move the local factors from the right to the left. We also used  $\eta^{\circ}$  |cluster $\rangle$  = |cluster $\rangle$ , which follows writing  $\eta^{\circ} = \prod_{j:\text{odd}} Z_{j-1}X_jZ_{j+1}$ . Using (E2), we find

$$D^{e}D^{o} | \text{cluster} \rangle =$$

$$= 2e^{\frac{2\pi iL}{8}} (-iX_{2})(-iX_{4}) \cdots (-iX_{L}) | \text{cluster} \rangle$$

$$= 2\eta^{e} | \text{cluster} \rangle = 2 | \text{cluster} \rangle .$$
(E3)

Putting everything together, we find

$$D | \text{cluster} \rangle = 2T | \text{cluster} \rangle = 2 | \text{cluster} \rangle$$
, (E4)

where the eigenvalue 2 is the quantum dimension of D. The equality  $T | \text{cluster} \rangle = | \text{cluster} \rangle$  follows from (3) and the facts that  $T \mathsf{V} = \mathsf{V} T$  and  $T | + + \dots + \rangle = | + + \dots + \rangle$ .

2. 
$$\mathsf{D} |\mathbf{odd}\rangle = 2(-1)^{L/4} |\mathbf{odd}\rangle$$

We begin with the relation (20), which we repeat here for convenience. Namely,

$$|\text{odd}\rangle = U_{\text{odd}} |\text{cluster}\rangle$$
, (E5)

where

$$U_{\text{odd}} = \prod_{n=1}^{L/2} Z_{2n} Z_{2n+2} e^{\frac{\pi i}{4} (Z_{2n-1} Z_{2n+1} - 1)}.$$
 (E6)

We assume L to be a multiple of 4 when discussing  $|\text{odd}\rangle$ , while other values of L correspond to inserting defects.

To find the action of D on the SPT state, we use the following commutation relation between D and  $U_{\text{odd}}$ :

$$D U_{\text{odd}} = \left( \prod_{n=1}^{L/2} e^{\frac{\pi i}{4} (X_{2n} - 1)} \prod_{k=1}^{L/4} X_{4k-1} \right) D,$$

$$= U_{\text{odd}} \left( \prod_{n=1}^{\frac{L}{2}} e^{-\frac{\pi i}{4} (Z_{2n-1} Z_{2n+1} + X_{2n})} \prod_{k=1}^{\frac{L}{4}} h_{4k-1} \right) D,$$
(E7)

where

$$h_j = Z_{j-1} X_j Z_{j+1}.$$
 (E8)

Using this relation and the facts that  $D | \text{cluster} \rangle = 2 | \text{cluster} \rangle$  and  $h_j | \text{cluster} \rangle = | \text{cluster} \rangle$ , we find

$$D |odd\rangle = U_{odd} \left( \prod_{n=1}^{\frac{L}{2}} e^{-\frac{\pi i}{2} Z_{2n-1} Z_{2n+1}} \right) 2 |cluster\rangle ,$$

$$= 2(-i)^{\frac{L}{2}} U_{odd} |cluster\rangle = 2(-1)^{\frac{L}{4}} |odd\rangle .$$
(E9)

Even though the model is defined for all  $L=0 \mod 4$ , here we see an interesting dependence on  $L \mod 8$ . This is reminiscent of the mod 4 dependence in the SO(3) Heisenberg chain discussed in [59]. We leave the study of this interesting sign for future investigations.

Finally, we have  $D | \text{even} \rangle = 2(-1)^{L/4} | \text{even} \rangle$ , which follows from TD = DT and  $T | \text{odd} \rangle = | \text{even} \rangle$ .

#### 3. Action on the interface states

Here, we find the action of the non-invertible symmetry on the edge modes localized on the interfaces between different SPT phases. We focus on the interface Hamiltonian  $H_{\rm cluster|odd}$  (21) for the cluster and 'odd' SPT states.

Define

$$X^{(\mathrm{L})} = Y_{L-2} X_{L-1} Z_L , \qquad X^{(\mathrm{R})} = Z_{\ell} X_{\ell+1} Y_{\ell+2} ,$$
  
 $Z^{(\mathrm{L})} = Z_{L-1} , \qquad Z^{(\mathrm{R})} = Z_{\ell+1} ,$  (E10)

which are localized on the left and right interfaces and form a complete basis of operators acting on the ground space. We define the 4 interface ground states by  $|\varepsilon_{\rm L}, \varepsilon_{\rm R}\rangle$ , with  $\varepsilon_{\rm L}, \varepsilon_{\rm R} = \pm$ , where they satisfy

$$\begin{split} X^{(\mathrm{L})} \left| \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle &= \varepsilon_{\mathrm{L}} \left| \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle, \quad X^{(\mathrm{R})} \left| \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle = \varepsilon_{\mathrm{R}} \left| \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle, \\ Z^{(\mathrm{L})} \left| \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle &= \left| -\varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle, \quad Z^{(\mathrm{R})} \left| \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \right\rangle = \left| \varepsilon_{\mathrm{L}}, -\varepsilon_{\mathrm{R}} \right\rangle, \end{split} \tag{E11}$$

and are stabilized by:

$$h_{j} = 1 \quad \text{for } j = 1, \dots, \ell,$$

$$-h_{2n} = 1 \quad \text{for } n = \ell/2 + 1, \dots, L/2,$$

$$Y_{2n}X_{2n+1}Y_{2n+2} = 1 \quad \text{for } n = \ell/2 + 1, \dots, L/2 - 2.$$
(E12)

The goal of this appendix is to find the action of the non-invertible symmetry on the ground space. We start with the  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  symmetry.

We first rewrite the  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  symmetry operators as products of the generators of the stabilizer (E12):

$$\eta^{e} = \prod_{n=1}^{\ell/2} h_{2n} \prod_{n=\ell/2+1}^{L/2} (-h_{2n}),$$

$$\eta^{o} = X^{(L)} X^{(R)} \prod_{n=1}^{\frac{\ell}{2}} h_{2n-1} \prod_{n=\frac{\ell}{2}+2}^{\frac{L}{2}-1} (Y_{2n-2} X_{2n-1} Y_{2n}).$$
(E13)

(Recall  $L - \ell = 0 \mod 4$  and L is even.) We have

$$\eta^{e} | \varepsilon_{L}, \varepsilon_{R} \rangle = | \varepsilon_{L}, \varepsilon_{R} \rangle, 
\eta^{o} | \varepsilon_{L}, \varepsilon_{R} \rangle = X^{(L)} X^{(R)} | \varepsilon_{L}, \varepsilon_{R} \rangle.$$
(E14)

Now we study the action of D. Since  $X^{(L)}D = -DX^{(L)}(-h_{L-2})$  and  $X^{(R)}D = -DX^{(R)}(-h_{\ell+2})$ , we find

$$\begin{split} X^{(\mathrm{L})} \mathsf{D} & | \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \rangle = - \mathsf{D} X^{(\mathrm{L})} & | \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \rangle \;, \\ X^{(\mathrm{R})} \mathsf{D} & | \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \rangle = - \mathsf{D} X^{(\mathrm{R})} & | \varepsilon_{\mathrm{L}}, \varepsilon_{\mathrm{R}} \rangle \;. \end{split} \tag{E15}$$

Moreover,  $Z^{(L)}Z^{(R)} D = -DZ^{(L)}Z^{(R)} \prod_{n=\ell/2+1}^{L/2-1} (-h_{2n})$ , which leads to

$$Z^{(L)}Z^{(R)}D|\varepsilon_{L}, \varepsilon_{R}\rangle = -DZ^{(L)}Z^{(R)}|\varepsilon_{L}, \varepsilon_{R}\rangle$$
. (E16)

Finally, since  $D^2 = (1 + \eta^e)(1 + \eta^o)$  we find

$$\mathsf{D}^{2}\left|\varepsilon_{\mathsf{L}},\varepsilon_{\mathsf{R}}\right\rangle = 2\left(1 + X^{(\mathsf{L})}X^{(\mathsf{R})}\right)\left|\varepsilon_{\mathsf{L}},\varepsilon_{\mathsf{R}}\right\rangle. \tag{E17}$$

Putting equations (E15), (E16), and (E17) together, we find the action of D on the ground space  $\mathcal{H}_{\text{cluster}|\text{odd}}$ , up to a sign  $\epsilon = \pm$ , to be:

$$\mathsf{D}\left|\varepsilon_{\mathrm{L}},\varepsilon_{\mathrm{R}}\right\rangle = i\epsilon Z^{(\mathrm{L})}Z^{(\mathrm{R})}\left(X^{(\mathrm{L})} + X^{(\mathrm{R})}\right)\left|\varepsilon_{\mathrm{L}},\varepsilon_{\mathrm{R}}\right\rangle \ . \ \ (\text{E}18)$$

To see this, we first note that the most general operator constructed from (E10) that anti-commutes with  $X^{(L)}$ ,  $X^{(R)}$ , and  $Z^{(L)}Z^{(R)}$ , have the form  $Z^{(L)}Z^{(R)}(\alpha X^{(L)}+\beta X^{(R)})$  for some  $\alpha,\beta\in\mathbb{C}$ . Further requiring that this operator squares to  $2(1+X^{(L)}X^{(R)})$ , leads to  $\alpha=\beta=\pm i$ . Thus we have found the action of D up to a sign  $\epsilon$ . To fix  $\epsilon$ , below we compute the action of D on the interface state  $|+,+\rangle$ . The method is similar to the one we used in Appendix E 2.

We note that the interface state  $|+,+\rangle$  can be prepared by acting with a finite-depth circuit on |cluster $\rangle$  on a closed chain. Namely, we have

$$|+,+\rangle = U^{++} |\text{cluster}\rangle,$$
 (E19)

where

$$U^{++} = \prod_{n=\ell/2+1}^{L/2} Z_{2n} \prod_{n=\ell/2+1}^{L/2-1} e^{-\frac{\pi i}{4} Z_{2n-1} Z_{2n+1}}.$$
 (E20)

The circuit  $U^{++}$  is a truncation of the circuit in (E6) to the region between sites  $\ell$  and L. One can verify this by studying the action of the circuit above on the operators  $h_j = Z_{j-1}X_jZ_{j+1}$ , which stabilize the cluster state. Indeed, the circuit  $U^{++}$  acts as

$$h_{2n} \mapsto \begin{cases} h_{2n} & \text{for } 1 \le n \le \frac{\ell}{2} \\ -h_{2n} & \text{for } \frac{\ell}{2} + 1 \le n \le \frac{L}{2} \end{cases}$$

$$h_{2n-1} \mapsto \begin{cases} h_{2n-1} & 1 \le n \le \frac{\ell}{2} \\ (-h_{\ell+2})X^{(R)} & n = \frac{\ell}{2} + 1 \\ h_{2n-2}(Y_{2n-2}X_{2n-1}Y_{2n})h_{2n} & \frac{\ell+4}{2} \le n \le \frac{L-2}{2} \\ (-h_{L-2})X^{(L)} & n = \frac{L}{2} \end{cases}$$
(E21)

To study the action of D on the interface, we first find the action of D on  $U^{++}$  as:

$$D U^{++} = \left( \prod_{n=\ell/2+1}^{L/2-1} e^{-\frac{\pi i}{4} X_{2n}} \prod_{k=1}^{\frac{L-\ell}{4}} X_{\ell+4k-1} \right) D,$$

$$= U^{++} \left( \prod_{n=\frac{\ell}{2}+1}^{\frac{L}{2}-1} e^{\frac{\pi i}{4} (Z_{2n-1} Z_{2n+1} + X_{2n})} \prod_{k=1}^{\frac{L-\ell}{4}} h_{\ell+4k-1} \right) D.$$
(E22)

Using this relation and the facts that  $D | \text{cluster} \rangle = 2 | \text{cluster} \rangle$  and  $h_j | \text{cluster} \rangle = | \text{cluster} \rangle$ , we find

$$D |+,+\rangle = U^{++} \left( \prod_{n=\frac{\ell}{2}+1}^{\frac{L}{2}-1} e^{\frac{\pi i}{2} Z_{2n-1} Z_{2n+1}} \right) 2 |\text{cluster}\rangle ,$$

$$= 2 \left( i^{\frac{L-\ell}{2}-1} Z_{\ell+1} Z_{L-1} \right) U^{++} |\text{cluster}\rangle .$$

$$= -2i(-1)^{\frac{L-\ell}{4}} Z_{\ell+1} Z_{L-1} |+,+\rangle .$$
(E23)

Hence,  $\epsilon = -(-1)^{\frac{L-\ell}{4}}$ .

Similarly, for the  $|-,-\rangle$  state we have  $|-,-\rangle = U^{--} |\text{cluster}\rangle$ , where

$$U^{--} = \prod_{n=\ell/2+1}^{L/2} Z_{2n} \prod_{n=\ell/2+1}^{L/2-1} e^{+\frac{\pi i}{4} Z_{2n-1} Z_{2n+1}} . \quad (E24)$$

This relation leads to

$$D |-,-\rangle = +2i(-1)^{\frac{L-\ell}{4}} Z_{\ell+1} Z_{L-1} |-,-\rangle$$
 (E25)

Equations (E23) and (E25) are compatible with (E18) and fix the action of D on the whole ground space. In summary, we find

$$\begin{split} \mathsf{D} \left| \varepsilon_{\mathsf{L}}, \varepsilon_{\mathsf{R}} \right\rangle &= \left( -1 \right)^{\frac{L-\ell}{4}} \left( Y^{(\mathsf{L})} Z^{(\mathsf{R})} + Z^{(\mathsf{L})} Y^{(\mathsf{R})} \right) \left| \varepsilon_{\mathsf{L}}, \varepsilon_{\mathsf{R}} \right\rangle, \\ \eta^{\mathsf{o}} \left| \varepsilon_{\mathsf{L}}, \varepsilon_{\mathsf{R}} \right\rangle &= X^{(\mathsf{L})} X^{(\mathsf{R})} \left| \varepsilon_{\mathsf{L}}, \varepsilon_{\mathsf{R}} \right\rangle, \\ \eta^{\mathsf{e}} \left| \varepsilon_{\mathsf{L}}, \varepsilon_{\mathsf{R}} \right\rangle &= \left| \varepsilon_{\mathsf{L}}, \varepsilon_{\mathsf{R}} \right\rangle. \end{split} \tag{E26}$$

# Appendix F: More on the absence of a $Rep(D_8)$ -symmetric entangler

Here, we fill in the steps in the argument of the non-existence of a  $Rep(D_8)$  symmetric entangler in Section V. There we claimed that the existence of an entangler U satisfying (29) implies the existence of the unitary operator  $\hat{U}$  satisfying (30). Below we provide an explicit argument for it.

The idea is to show that (30) is the image of (29) under the KT transformation. To simplify the discussion in this appendix, we identify the hatted operators with the unhatted operators. That is,  $\hat{X}_j = X_j$  and  $\hat{Z}_j = Z_j$ , which implies  $\hat{V} = V$ ,  $\hat{\eta}^e = \eta^e$ , and  $\hat{\eta}^o = \eta^o$ . Under this identification, the KT transformation is implemented by the non-invertible operator VDV as in (A12).

In this simplified notation, we will show that

$$\begin{split} \mathsf{UD} &= \mathsf{DU} \,, & \mathsf{U} \eta^\mathrm{e} &= \eta^\mathrm{e} \mathsf{U} \,, \\ \mathsf{U} \, |\mathrm{cluster} \rangle &= |\mathrm{odd} \rangle \,\,, & \mathsf{U} \eta^\mathrm{o} &= \eta^\mathrm{o} \mathsf{U} \,, \end{split} \tag{F1}$$

implies the existence of the unitary operator  $\hat{\mathsf{U}} = \hat{\mathsf{U}}(X_j, Z_j)$  satisfying

$$\hat{\mathbf{U}}\mathbf{V} = \mathbf{V}\hat{\mathbf{U}}, \qquad \hat{\mathbf{U}}\eta^{e} = \eta^{e}\hat{\mathbf{U}}, 
\hat{\mathbf{U}}|\widehat{\text{cluster}}\rangle = |\widehat{\text{odd}}\rangle, \qquad \hat{\mathbf{U}}\eta^{o} = \eta^{o}\hat{\mathbf{U}}.$$
(F2)

Here,  $|\text{cluster}\rangle$  and  $|\text{odd}\rangle$  are, respectively, the *unique* ground states of  $\hat{H}_{\text{cluster}}$  and  $\hat{H}_{\text{odd}}$  with eigenvalue +1 under  $\eta^{\text{e}}, \eta^{\text{o}}, \mathsf{V}$ . For instance, using the convention used around (B7), we have

$$|\widehat{\text{cluster}}\rangle = \frac{1}{2}(|00\rangle + |01\rangle + |10\rangle + |11\rangle).$$
 (F3)

In infinite volume, each of these four short-range entangled states  $|z_e z_o\rangle$  becomes a superselection sector that preserves  $\mathbb{Z}_2^{\mathsf{v}}$  and spontaneously breaks  $\mathbb{Z}_2^{\mathsf{e}} \times \mathbb{Z}_2^{\mathsf{e}}$ . Similarly,  $|\widehat{\mathrm{odd}}\rangle$  is the symmetric superposition of the 4 short-range entangled ground states of  $\hat{H}_{\mathrm{odd}}$ . In infinite volume, each of these short-range entangled states becomes a superselection sector that only preserves  $\mathrm{diag}(\mathbb{Z}_2^{\mathsf{e}} \times \mathbb{Z}_2^{\mathsf{v}})$ . These two spontaneous symmetry breaking patterns are distinguished by the following order parameters:

$$\langle \widehat{\text{cluster}} | Z_j Z_{j+2n} | \widehat{\text{cluster}} \rangle = 1, \text{ for } j = 0, 1,$$
 (F4)

and

$$\begin{split} &\langle \widehat{\mathrm{odd}} | \left( Y_0 \frac{1 - Z_{-1} Z_1}{2} \right) \left( Y_{4k} \frac{1 - Z_{4k-1} Z_{4k+1}}{2} \right) | \widehat{\mathrm{odd}} \rangle = 1 \,, \\ &\langle \widehat{\mathrm{odd}} | \, Z_1 Z_{4k+1} \, | \widehat{\mathrm{odd}} \rangle = 1 \,. \end{split}$$

Moreover, we will show that the action of KT transformation on  $U, D, |\text{cluster}\rangle, |\text{odd}\rangle$  is given by  $\hat{U}, V, |\widehat{\text{cluster}}\rangle, |\widehat{\text{odd}}\rangle$ , in the sense that

$$\begin{split} & (\mathsf{VDV})\,\mathsf{U} = \hat{\mathsf{U}}\,(\mathsf{VDV})\,, \quad \mathsf{VDV}\,|\mathrm{cluster}\rangle = 2\,|\widehat{\mathrm{cluster}}\rangle\,, \\ & (\mathsf{VDV})\,\mathsf{D} = 2\mathsf{V}\,(\mathsf{VDV})\,, \quad \mathsf{VDV}\,|\mathrm{odd}\rangle = 2\,|\widehat{\mathrm{odd}}\rangle\,. \end{split}$$

The relation (VDV)D = 2V(VDV) is derived in Appendix B 1, and VDV |cluster $\rangle = 2$  |cluster $\rangle$  follows from equation (B9) of Appendix B 1.

Existence of  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric  $\hat{\mathsf{U}}$ : To argue for the existence of a  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric  $\hat{\mathsf{U}}$ , we note that the invariance of  $\mathsf{U}$  under  $\mathsf{D}$  means that the operator  $\mathsf{U}$  remains invariant under the untwisted gauging  $\mathsf{S}$ . This implies that there is no mixed anomaly involving  $\mathsf{U}$  and  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ . Therefore, the image of  $\mathsf{U}$  under  $\mathsf{TST} / \mathsf{KT}$ 

transformation is an order 2, invertible,  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric operator  $\hat{U}$ , which satisfies

$$(VDV) U = \hat{U} (VDV).$$
 (F7)

We note that (F7) fixes the operator  $\hat{U}$  up to a factor of  $\eta^{\rm e}$  and  $\eta^{\rm o}$ . This is because multiplying (F7) from the right by VDV implies (VDV)U(VDV) =  $\hat{U}(1+\eta^{\rm e})(1+\eta^{\rm o})$ . In other words, if  $\hat{U}$  satisfies (F7) (and (F2)) then  $\hat{U}\eta^{\rm e}$ ,  $\hat{U}\eta^{\rm o}$ , and  $\hat{U}\eta^{\rm e}\eta^{\rm o}$  also satisfy those relations.

**Action of KT on |odd**>: The relation

$$(VDV) H_{odd} = \hat{H}_{odd} (VDV)$$
 (F8)

implies that the  $\eta^{\rm e}=\eta^{\rm o}=1$  spectrum of  $H_{\rm odd}$  and  $\hat{H}_{\rm odd}$  are related by the operator VDV. In particular,  $|{\rm odd}\rangle$  and  $|{\rm odd}\rangle$  are the unique ground states of these Hamiltonians with eigenvalues  $\eta^{\rm e}=\eta^{\rm o}=1$ . Hence, they are related by the operator VDV:

$$VDV |odd\rangle = 2 |\widehat{odd}\rangle. \tag{F9}$$

The overall normalization is fixed by requiring the states to be normalized.

**Deriving** (F2): Now that we have verified (F6), we are ready to derive (F2). Applying VDV to  $U | \text{cluster} \rangle = | \text{odd} \rangle$ , and using (F6) we find

$$\hat{U} (VDV) | \text{cluster} \rangle = (VDV) | \text{odd} \rangle , 
\Rightarrow 2 \, \hat{U} | \widehat{\text{cluster}} \rangle = 2 | \widehat{\text{odd}} \rangle .$$
(F10)

Thus, we have shown that  $\hat{U}$  maps the ground states of  $\hat{H}_{\text{cluster}}$  to those of  $\hat{H}_{\text{odd}}$ .

It remains to show that  $\hat{U}$  commutes with V. Intuitively, this is follows from the fact that U and D commute, and their images under the KT transformation are  $\hat{U}$  and V, respectively. In the following, we show this algebraically.

From UD = DU and (F6) we find

$$\hat{\mathsf{U}}\mathsf{V}(\mathsf{VDV}) = \mathsf{V}\hat{\mathsf{U}}(\mathsf{VDV}). \tag{F11}$$

By multiplying this equation on the right-hand-side by  $\mathsf{VDV}$  we find

$$\hat{\mathsf{U}}\mathsf{V}(1+\eta^{\rm e})(1+\eta^{\rm o}) = \mathsf{V}\hat{\mathsf{U}}(1+\eta^{\rm e})(1+\eta^{\rm o}).$$
 (F12)

A priori, as we show below, this is only possible if

$$\hat{\mathsf{U}}\mathsf{V}\hat{\mathsf{U}}^{-1} = \mathsf{V}, \, \mathsf{V}\eta^{\mathrm{e}}, \, \mathsf{V}\eta^{\mathrm{o}}, \, \mathsf{V}\eta^{\mathrm{e}}\eta^{\mathrm{o}} \,. \tag{F13}$$

**Proof of** (F13): Define

$$W = \hat{U}V\hat{U}^{-1}V^{-1}.$$
 (F14)

Since we assume the entangler U to be locality-preserving, so is its image  $\hat{U}$ , and therefore also W. Next, since W commutes with  $\eta^{\rm e}, \eta^{\rm o}$ , it can be diagonalized to the block-diagonal form  ${\rm diag}(W_{++}, W_{+-}, W_{-+}, W_{--})$ ,

where  $W_{\epsilon^e\epsilon^o}$  is the restriction of W to the eigenspace  $\mathcal{H}_{\epsilon^e\epsilon^o}$  with  $\eta^e=\epsilon^e$  and  $\eta^o=\epsilon^o$ . Then (F12) implies  $W_{++}=1$ . Below we show that this implies that W acts trivially on all  $\mathbb{Z}_2^e\times\mathbb{Z}_2^o$  symmetric local *operators*.

Since W acts trivially on the states in  $\mathcal{H}_{++}$ , given any local operator  $O_j$  that commutes with  $\eta^e$ ,  $\eta^o$ , we have

$$O_j |\psi\rangle = \mathsf{W} O_j \mathsf{W}^{-1} |\psi\rangle$$
, for all  $|\psi\rangle \in \mathcal{H}_{++}$ . (F15)

Given that  $O_j$  and  $WO_jW^{-1}$  are localized around site j, we can insert  $Z_{j'}$ 's far away from site j and conclude that the relation above holds not only in  $\mathcal{H}_{++}$ , but on the entire Hilbert space. Hence,  $O_j = WO_jW^{-1}$  for any  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetric local operator  $O_j$ .

Next, consider the local operators  $O_j$  that satisfy  $\eta^{\rm e}O_j\eta^{\rm e}=\epsilon^{\rm e}O_j$  and  $\eta^{\rm e}O_j\eta^{\rm e}=\epsilon^{\rm e}O_j$  for some fixed  $\mathbb{Z}_2^{\rm e}\times\mathbb{Z}_2^{\rm e}$  charges  $\epsilon^{\rm e},\epsilon^{\rm o}=\pm$ . Multiplying two such operators,  $O_j$  and  $O'_{j'}$  that are well-separated, we find  $O_jO'_{j'}$  that is  $\mathbb{Z}_2^{\rm e}\times\mathbb{Z}_2^{\rm e}$  symmetric. Hence, it must commute with W:

$$(WO_jW^{-1})(WO'_{j'}W^{-1}) = O_jO'_{j'}.$$
 (F16)

Since  $\mathsf{W}O_j\mathsf{W}^{-1}$  and  $\mathsf{W}O'_{j'}\mathsf{W}^{-1}$  are well-separated, the equality above is only possible if  $\mathsf{W}O_j\mathsf{W}^{-1} = \alpha(\epsilon^\mathrm{e},\epsilon^\mathrm{o})O_j$  and  $\mathsf{W}O'_{j'}\mathsf{W}^{-1} = \alpha(\epsilon^\mathrm{e},\epsilon^\mathrm{o})O'_{j'}$  for  $\alpha(\epsilon^\mathrm{e},\epsilon^\mathrm{o}) = \pm$  that only depends on  $\epsilon^\mathrm{e}$  and  $\epsilon^\mathrm{o}$ . Furthermore,  $\alpha(\epsilon^\mathrm{e},\epsilon^\mathrm{o})$  is a bicharacter, i.e.,  $\alpha(+,+) = +$  and  $\alpha(-,-) = \alpha(+,-)\alpha(-,+)$ . This implies that the action of

W and 
$$(\eta^{e})^{\frac{1-\alpha(-,+)}{2}} (\eta^{o})^{\frac{1-\alpha(+,-)}{2}}$$
 (F17)

on local operators is the same. We have thus proved (F13).

Finally, we show that only the first option in (F13) is compatible with anomalies. The last option is not possible since V is anomaly free but  $\nabla \eta^e \eta^o$  is anomalous. To rule out the second possibility, we note that  $\hat{U}\hat{V}\hat{U}^{-1} = \nabla \eta^e$  implies  $\hat{U}\nabla \eta^o \hat{U}^{-1} = \nabla \eta^e \eta^o$ , which is not possible since  $\nabla \eta^o$  is anomaly free but  $\nabla \eta^e \eta^o$  is anomalous. Similarly, we can also rule out the third possibility. We conclude that  $\hat{U}$  must commute with V. This completes the argument for (F2) if there were a symmetric entangler, which subsequently led to the contradiction in Section V.

# Appendix G: Boundary conditions for the cluster model

In this appendix we discuss symmetric boundary conditions for the cluster model.

The standard boundary condition for the cluster Hamiltonian on an open chain with  $\ell$  sites (with even  $\ell$ ) is:

$$H_{\text{open}} = -\sum_{j=1}^{\ell-2} Z_j X_{j+1} Z_{j+2},$$
 (G1)

(This can be obtained from (D1) by fixing the qubits on  $j = \ell + 1, \dots, L$  to be all  $|+\rangle$ .) This Hamiltonian is invariant under the symmetry operators

$$\eta_{\text{open}}^{\text{e}} = \prod_{j=2}^{\ell} X_j, \quad \eta_{\text{open}}^{\text{o}} = \prod_{j=1}^{\ell-1} X_j.$$
(G2)

The ground space is 4-dimensional, with each boundary being 2-fold degenerate. The Hamiltonian  $H_{\text{open}}$  represents a  $\mathbb{Z}_2^{\text{e}} \times \mathbb{Z}_2^{\text{o}}$  symmetric boundary condition with edge modes.

However, in a general quantum system (such as a continuum field theory or a lattice model without a tensor product Hilbert space), there may not be a canonical way to cut open the symmetry operator U(g) of a global symmetry G at the boundary. The notion of a "symmetric boundary" therefore requires clarifications. One definition is the following  $^{16}$ 

- A pair of boundary conditions are characterized by a Hamiltonian  $H_{\rm open}$  on an interval. The local terms of this Hamiltonian away from the boundaries are identical to the local terms in the Hamiltonian on a closed chain.
- There exist conserved operators  $U_{\text{open}}(g)$  with  $g \in G$  on an interval that commute with  $H_{\text{open}}$  and realize the symmetry G linearly, i.e.,  $U_{\text{open}}(g_1)U_{\text{open}}(g_2) = U_{\text{open}}(g_1g_2)$ . The local factors of  $U_{\text{open}}(g)$  away from the boundaries are identical to the local factors of the conserved operator U(g) on a closed chain.

Note that in this definition, the local factors of  $U_{\text{open}}(g)$  near the boundaries might be different from the local factors in the interior.

It is generally expected [31, 77–79] that given a quantum system with an ordinary global symmetry G free of any 't Hooft anomaly (i.e., G can be gauged and is compatible with a trivially gapped phase), there should exist a such symmetric boundary condition with no edge mode.

Since the  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry of the cluster model is free of 't Hooft anomalies, we expect there to be such a  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$ -symmetric boundary condition in the above sense. This is achieved by considering a different Hamiltonian on the open chain:

$$ar{H}_{ ext{open}} = -X_1 Z_2 - \left( \sum_{j=1}^{\ell-2} Z_j X_{j+1} Z_{j+2} \right) - Z_{\ell-1} X_{\ell} \,.$$

(G3)

Since the Hamiltonian consists of L commuting Pauli operators, the ground space is non-degenerate. In particular, there is no local degeneracy at the boundary. This Hamiltonian on an open chain is also invariant under a  $\mathbb{Z}_2^e \times \mathbb{Z}_2^o$  symmetry, generated by

$$\bar{\eta}_{\text{open}}^{\text{e}} = Z_1 \prod_{j=2}^{\ell} X_j, \quad \bar{\eta}_{\text{open}}^{\text{o}} = \left(\prod_{j=1}^{\ell-1} X_j\right) Z_\ell, \quad (G4)$$

which obey  $[\bar{\eta}_{\text{open}}^{\text{e}}, \bar{H}_{\text{open}}] = 0$ ,  $[\bar{\eta}_{\text{open}}^{\text{o}}, \bar{H}_{\text{open}}] = 0$ . Let us summarize some features of this Hamiltonian  $\bar{H}_{\text{open}}$ :

- The local terms (i.e.,  $Z_j X_{j+1} Z_{j+2}$ ) in  $\bar{H}_{\text{open}}$  in the interior of the chain are identical to the cluster Hamiltonian on a closed chain.
- The local terms (i.e.,  $X_j$ ) of the  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  symmetry operators  $\bar{\eta}_{\rm open}^{\rm e}$ ,  $\bar{\eta}_{\rm open}^{\rm o}$  in the interior of the chain are identical to  $\eta^{\rm e}$ ,  $\eta^{\rm o}$  on a closed chain.

In this sense,  $\bar{H}_{\rm open}$  represents a  $\mathbb{Z}_2^{\rm e} \times \mathbb{Z}_2^{\rm o}$  symmetric boundary condition for the cluster model with no edge mode.

This is not in contradiction with the usual statement in the literature that any symmetric boundary condition of the cluster model must have edge modes. This statement assumes the symmetry to take a strict on-site form in (G2). However, this requirement does not have a generalization to non-onsite symmetries (such as our non-invertible symmetry D), or to continuum field theories. Here we relax this requirement by allowing a local modification of the symmetry operators near the boundaries.

<sup>[1]</sup> F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Symmetry protection of topological phases in one-dimensional quantum spin systems, Phys. Rev. B 85, 075125 (2012), arXiv:0909.4059 [cond-mat.str-el].

<sup>[2]</sup> X. Chen, Z. C. Gu, and X. G. Wen, Local unitary transformation, long-range quantum entanglement, wave func-

<sup>&</sup>lt;sup>16</sup> The different notions of symmetric boundary conditions for non-invertible symmetries are discussed in [31], where such a boundary condition is referred to as a weakly symmetric boundary condition.

tion renormalization, and topological order, Phys. Rev. B 82, 155138 (2010), arXiv:1004.3835 [cond-mat.str-el].

<sup>[3]</sup> X. Chen, Z.-C. Gu, and X.-G. Wen, Classification of gapped symmetric phases in one-dimensional spin systems, Phys. Rev. B 83, 035107 (2011), arXiv:1008.3745 [cond-mat.str-el].

<sup>[4]</sup> L. Fidkowski and A. Kitaev, Topological phases of fermions in one dimension, Phys. Rev. B 83, 075103 (2011), arXiv:1008.4138 [cond-mat.str-el].

<sup>[5]</sup> N. Schuch, D. Pérez-García, and I. Cirac, Classifying quantum phases using matrix product states and projected entangled pair states, Phys. Rev. B 84, 165139

- (2011).
- [6] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Symmetry protected topological orders and the group cohomology of their symmetry group, Phys. Rev. B 87, 155114 (2013), arXiv:1106.4772 [cond-mat.str-el].
- [7] T. Senthil, Symmetry Protected Topological phases of Quantum Matter, Ann. Rev. Condensed Matter Phys. 6, 299 (2015), arXiv:1405.4015 [cond-mat.str-el].
- [8] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, Rev. Mod. Phys. 88, 035005 (2016), arXiv:1505.03535 [cond-mat.mes-hall].
- [9] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, Quantum Information Meets Quantum Matter: From Quantum Entanglement to Topological Phases of Many-Body Systems, Quantum Science and Technology (Springer, 2019) arXiv:1508.02595 [cond-mat.str-el].
- [10] W. Son, L. Amico, and V. Vedral, Topological order in 1d cluster state protected by symmetry, Quantum Information Processing 11, 1961 (2012).
- [11] M. A. Nielsen, Cluster-state quantum computation, Reports on Mathematical Physics 57, 147 (2006), arXiv:quant-ph/0504097 [quant-ph].
- [12] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, Generalized Global Symmetries, JHEP 02, 172, arXiv:1412.5148 [hep-th].
- [13] J. McGreevy, Generalized Symmetries in Condensed Matter 10.1146/annurev-conmatphys-040721-021029 (2022), arXiv:2204.03045 [cond-mat.str-el].
- [14] C. Cordova, T. T. Dumitrescu, K. Intriligator, and S.-H. Shao, Snowmass White Paper: Generalized Symmetries in Quantum Field Theory and Beyond, in *Snowmass 2021* (2022) arXiv:2205.09545 [hep-th].
- [15] T. D. Brennan and S. Hong, Introduction to Generalized Global Symmetries in QFT and Particle Physics, (2023), arXiv:2306.00912 [hep-ph].
- [16] S. Schafer-Nameki, ICTP Lectures on (Non-)Invertible Generalized Symmetries, (2023), arXiv:2305.18296 [hep-th].
- [17] S.-H. Shao, What's Done Cannot Be Undone: TASI Lectures on Non-Invertible Symmetry, (2023), arXiv:2308.00747 [hep-th].
- [18] U. Grimm and G. M. Schutz, The Spin 1/2 XXZ Heisenberg chain, the quantum algebra U(q)[sl(2)], and duality transformations for minimal models, J. Statist. Phys. 71, 921 (1993), arXiv:hep-th/0111083.
- [19] M. Oshikawa and I. Affleck, Boundary conformal field theory approach to the critical two-dimensional Ising model with a defect line, Nucl. Phys. B 495, 533 (1997), arXiv:cond-mat/9612187.
- [20] D. Aasen, R. S. K. Mong, and P. Fendley, Topological Defects on the Lattice I: The Ising model, J. Phys. A 49, 354001 (2016), arXiv:1601.07185 [cond-mat.stat-mech].
- [21] D. Aasen, P. Fendley, and R. S. K. Mong, Topological Defects on the Lattice: Dualities and Degeneracies, (2020), arXiv:2008.08598 [cond-mat.stat-mech].
- [22] L. Li, M. Oshikawa, and Y. Zheng, Noninvertible duality transformation between symmetry-protected topological and spontaneous symmetry breaking phases, Phys. Rev. B 108, 214429 (2023), arXiv:2301.07899 [cond-mat.strel].
- [23] N. Seiberg and S.-H. Shao, Majorana chain and Ising model – (non-invertible) translations, anomalies, and emanant symmetries, SciPost Phys. 16, 064 (2024),

- arXiv:2307.02534 [cond-mat.str-el].
- [24] N. Seiberg, S. Seifnashri, and S.-H. Shao, Non-invertible symmetries and LSM-type constraints on a tensor product Hilbert space, (2024), arXiv:2401.12281 [condmat.str-el].
- [25] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, Topological Defect Lines and Renormalization Group Flows in Two Dimensions, JHEP 01, 026, arXiv:1802.04445 [hep-th].
- [26] R. Thorngren and Y. Wang, Fusion Category Symmetry I: Anomaly In-Flow and Gapped Phases, (2019), arXiv:1912.02817 [hep-th].
- [27] C. Zhang and C. Córdova, Anomalies of (1+1)D categorical symmetries, (2023), arXiv:2304.01262 [cond-mat.strel].
- [28] C. Cordova, P.-S. Hsin, and C. Zhang, Anomalies of Non-Invertible Symmetries in (3+1)d, (2023), arXiv:2308.11706 [hep-th].
- [29] A. Antinucci, F. Benini, C. Copetti, G. Galati, and G. Rizi, Anomalies of non-invertible self-duality symmetries: fractionalization and gauging, (2023), arXiv:2308.11707 [hep-th].
- [30] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, Symmetries and strings of adjoint QCD<sub>2</sub>, JHEP 03, 103, arXiv:2008.07567 [hep-th].
- [31] Y. Choi, B. C. Rayhaun, Y. Sanghavi, and S.-H. Shao, Remarks on boundaries, anomalies, and noninvertible symmetries, Phys. Rev. D 108, 125005 (2023), arXiv:2305.09713 [hep-th].
- [32] Y. Choi, D.-C. Lu, and Z. Sun, Self-duality under gauging a non-invertible symmetry, (2023), arXiv:2310.19867 [hep-th].
- [33] D. Tambara, Representations of tensor categories with fusion rules of self-duality for abelian groups, Israel Journal of Mathematics 118, 29 (2000).
- [34] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor Categories*, Mathematical Surveys and Monographs (American Mathematical Society, 2016).
- [35] G. W. Moore and N. Seiberg, Classical and Quantum Conformal Field Theory, Commun. Math. Phys. 123, 177 (1989).
- [36] G. W. Moore and N. Seiberg, Naturality in Conformal Field Theory, Nucl. Phys. B 313, 16 (1989).
- [37] C. Fechisin, N. Tantivasadakarn, and V. V. Albert, Non-invertible symmetry-protected topological order in a group-based cluster state, (2023), arXiv:2312.09272 [cond-mat.str-el].
- [38] K. Inamura, Topological field theories and symmetry protected topological phases with fusion category symmetries, JHEP 05, 204, arXiv:2103.15588 [cond-mat.str-el].
- [39] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schafer-Nameki, Gapped Phases with Non-Invertible Symmetries: (1+1)d, (2023), arXiv:2310.03784 [hep-th].
- [40] L. Bhardwaj, D. Pajer, S. Schafer-Nameki, and A. Warman, Hasse Diagrams for Gapless SPT and SSB Phases with Non-Invertible Symmetries, (2024), arXiv:2403.00905 [cond-mat.str-el].
- [41] Y. Choi, C. Cordova, P.-S. Hsin, H. T. Lam, and S.-H. Shao, Noninvertible duality defects in 3+1 dimensions, Phys. Rev. D 105, 125016 (2022), arXiv:2111.01139 [hep-th].
- [42] Y. Choi, C. Cordova, P.-S. Hsin, H. T. Lam, and S.-H. Shao, Non-invertible Condensation, Duality, and Triality Defects in 3+1 Dimensions, Commun. Math. Phys. 402,

- 489 (2023), arXiv:2204.09025 [hep-th].
- [43] S. Seifnashri and S.-H. Shao, to appear.
- [44] T. Kennedy and H. Tasaki, Hidden symmetry breaking and the haldane phase in s= 1 quantum spin chains, Communications in mathematical physics 147, 431 (1992).
- [45] T. Kennedy and H. Tasaki, Hidden z<sub>2</sub>×z<sub>2</sub> symmetry breaking in haldane-gap antiferromagnets, Phys. Rev. B 45, 304 (1992).
- [46] A. Parayil Mana, Y. Li, H. Sukeno, and T.-C. Wei, Kennedy-Tasaki transformation and non-invertible symmetry in lattice models beyond one dimension, (2024), arXiv:2402.09520 [cond-mat.str-el].
- [47] M. D. F. de Wild Propitius, Topological interactions in broken gauge theories, Ph.D. thesis, Amsterdam U. (1995), arXiv:hep-th/9511195.
- [48] J. Wang, L. H. Santos, and X.-G. Wen, Bosonic Anomalies, Induced Fractional Quantum Numbers and Degenerate Zero Modes: the anomalous edge physics of Symmetry-Protected Topological States, Phys. Rev. B 91, 195134 (2015), arXiv:1403.5256 [cond-mat.str-el].
- [49] L. Bhardwaj, L. E. Bottini, D. Pajer, and S. Schafer-Nameki, Categorical Landau Paradigm for Gapped Phases, (2023), arXiv:2310.03786 [cond-mat.str-el].
- [50] D. V. Else and C. Nayak, Classifying symmetry-protected topological phases through the anomalous action of the symmetry on the edge, Phys. Rev. B 90, 235137 (2014), arXiv:1409.5436 [cond-mat.str-el].
- [51] L. Bhardwaj and Y. Tachikawa, On finite symmetries and their gauging in two dimensions, JHEP 03, 189, arXiv:1704.02330 [hep-th].
- [52] K. Inamura, On lattice models of gapped phases with fusion category symmetries, JHEP 03, 036, arXiv:2110.12882 [cond-mat.str-el].
- [53] O. Diatlyk, C. Luo, Y. Wang, and Q. Weller, Gauging Non-Invertible Symmetries: Topological Interfaces and Generalized Orbifold Groupoid in 2d QFT, (2023), arXiv:2311.17044 [hep-th].
- [54] A. Perez-Lona, D. Robbins, E. Sharpe, T. Vandermeulen, and X. Yu, Notes on gauging noninvertible symmetries, part 1: Multiplicity-free cases, (2023), arXiv:2311.16230 [hep-th].
- [55] C. Vafa, Quantum Symmetries of String Vacua, Mod. Phys. Lett. A 4, 1615 (1989).
- [56] M. Levin and Z.-C. Gu, Braiding statistics approach to symmetry-protected topological phases, Phys. Rev. B 86, 115109 (2012), arXiv:1202.3120 [cond-mat.str-el].
- [57] X. Chen, Z.-X. Liu, and X.-G. Wen, Two-dimensional symmetry-protected topological orders and their protected gapless edge excitations, Phys. Rev. B 84, 235141 (2011), arXiv:1106.4752 [cond-mat.str-el].
- [58] J. Wang, X.-G. Wen, and E. Witten, Symmetric Gapped Interfaces of SPT and SET States: Systematic Constructions, Phys. Rev. X 8, 031048 (2018), arXiv:1705.06728 [cond-mat.str-el].
- [59] M. Cheng and N. Seiberg, Lieb-Schultz-Mattis, Luttinger, and 't Hooft anomaly matching in lattice systems, SciPost Phys. 15, 051 (2023), arXiv:2211.12543 [cond-mat.str-el].
- [60] S. Seifnashri, Lieb-Schultz-Mattis anomalies as obstructions to gauging (non-on-site) symmetries, (2023), arXiv:2308.05151 [cond-mat.str-el].
- [61] Y. Yao and M. Oshikawa, Twisted boundary con-

- dition and Lieb-Schultz-Mattis ingappability for discrete symmetries, Phys. Rev. Lett. **126**, 217201 (2021), arXiv:2010.09244 [cond-mat.str-el].
- [62] Y. Tachikawa, On gauging finite subgroups, SciPost Phys. 8, 015 (2020), arXiv:1712.09542 [hep-th].
- [63] J. Kaidi, E. Nardoni, G. Zafrir, and Y. Zheng, Symmetry TFTs and anomalies of non-invertible symmetries, JHEP 10, 053, arXiv:2301.07112 [hep-th].
- [64] M. Iqbal et al., Non-Abelian topological order and anyons on a trapped-ion processor, Nature 626, 505 (2024), arXiv:2305.03766 [quant-ph].
- [65] J. Kaidi, K. Ohmori, and Y. Zheng, Kramers-Wannier-like Duality Defects in (3+1)D Gauge Theories, Phys. Rev. Lett. 128, 111601 (2022), arXiv:2111.01141 [hep-th].
- [66] W. Ji, S.-H. Shao, and X.-G. Wen, Topological Transition on the Conformal Manifold, Phys. Rev. Res. 2, 033317 (2020), arXiv:1909.01425 [cond-mat.str-el].
- [67] Y.-H. Lin and S.-H. Shao, Duality Defect of the Monster CFT, J. Phys. A 54, 065201 (2021), arXiv:1911.00042 [hep-th].
- [68] Y. Choi, H. T. Lam, and S.-H. Shao, Noninvertible Global Symmetries in the Standard Model, Phys. Rev. Lett. 129, 161601 (2022), arXiv:2205.05086 [hep-th].
- [69] C. Cordova and K. Ohmori, Noninvertible Chiral Symmetry and Exponential Hierarchies, Phys. Rev. X 13, 011034 (2023), arXiv:2205.06243 [hep-th].
- [70] D. Tambara and S. Yamagami, Tensor categories with fusion rules of self-duality for finite abelian groups, Journal of Algebra 209, 692 (1998).
- [71] W. Ji and X.-G. Wen, Categorical symmetry and non-invertible anomaly in symmetry-breaking and topological phase transitions, Phys. Rev. Res. 2, 033417 (2020), arXiv:1912.13492 [cond-mat.str-el].
- [72] D. S. Freed, Short-range entanglement and invertible field theories, (2014), arXiv:1406.7278 [cond-mat.str-el].
- [73] D. S. Freed and G. W. Moore, Setting the quantum integrand of M-theory, Commun. Math. Phys. 263, 89 (2006), arXiv:hep-th/0409135.
- [74] E. Witten, SL(2,Z) action on three-dimensional conformal field theories with Abelian symmetry, in From Fields to Strings: Circumnavigating Theoretical Physics: A Conference in Tribute to Ian Kogan (2003) pp. 1173–1200, arXiv:hep-th/0307041.
- [75] L. Bhardwaj, Y. Lee, and Y. Tachikawa, SL(2, Z) action on QFTs with Z<sub>2</sub> symmetry and the Brown-Kervaire invariants, JHEP 11, 141, arXiv:2009.10099 [hep-th].
- [76] M. Sinha, F. Yan, L. Grans-Samuelsson, A. Roy, and H. Saleur, Lattice Realizations of Topological Defects in the critical (1+1)-d Three-State Potts Model, (2023), arXiv:2310.19703 [hep-th].
- [77] K. Jensen, E. Shaverin, and A. Yarom, 't Hooft anomalies and boundaries, JHEP 01, 085, arXiv:1710.07299 [hepth].
- [78] T. Numasawa and S. Yamaguchi, Mixed global anomalies and boundary conformal field theories, JHEP 11, 202, arXiv:1712.09361 [hep-th].
- [79] R. Thorngren and Y. Wang, Anomalous symmetries end at the boundary, JHEP 09, 017, arXiv:2012.15861 [hepth].