A nonperturbative test of nucleation calculations for strong phase transitions

Oliver Gould*

School of Physics and Astronomy, University of Nottingham, Nottingham NG7 2RD, U.K.

Anna Kormu^{\dagger} and David J. Weir^{\ddagger}

Department of Physics and Helsinki Institute of Physics, PL 64, FI-00014 University of Helsinki, Finland

(Dated: April 2, 2024)

Nucleation rate computations are of broad importance in particle physics and cosmology. Perturbative calculations are often used to compute the nucleation rate Γ , but these are incomplete. We perform nonperturbative lattice simulations of nucleation in a scalar field theory with a treelevel barrier, computing a final result extrapolated to the thermodynamic and continuum limits. Although the system in question should be well-described by a complete one-loop perturbative calculation, we find only qualitative agreement with the full perturbative result, with a 20% discrepancy in $|\log \Gamma|$. Our result motivates further testing of the current nucleation paradigm.

I. INTRODUCTION

First order phase transitions in the early universe have been of sustained interest. A primordial phase transition could help to explain the matter-antimatter asymmetry of the universe [1–6]. The phase transition itself, as well as any resulting nonequilibrium physics of the primordial plasma, would produce a stochastic gravitational wave background that could potentially be detected [7].

Hints of a gravitational wave background have been seen by the various Pulsar Timing Array (PTA) collaborations [8–11]. Such a gravitational wave background is expected to be due to supermassive black holes [12], but it is not possible to rule out other new physics such as a phase transition [13]. Future missions such as LISA will be well placed to look for gravitational waves from electroweak-scale phase transitions [14–16], but the theoretical uncertainties in the predicted gravitational wave power spectrum will need to be constrained.

Thanks in part to large-scale simulations [17, 18], considerable progress has been made in modelling the resulting gravitational wave power spectrum [19, 20]. At the same time, the accuracy of the quantities which parametrise the power spectrum – including the nucleation rate – have faced growing scrutiny [21–26].

Experimental tests of nucleation theory in condensed matter systems show a mixed picture. For the AB transition in superfluid ³He, there is a longstanding and puzzling discrepancy [27–29], whereas good agreement was found for nucleation in a ferromagnetic superfluid [30]. Recently, there have been proposals to test bubble nucleation in ultracold atomic gases [31–37].

There is a need for reliable predictions with controlled systematic uncertainties so that we can test particle physics models against gravitational wave observations, as well as for comparison with analogue experiments. In this paper we focus on calibrating the accuracy of nucleation rate calculations in relativistic finite-temperature field theory, comparing perturbative semiclassical methods against direct nonperturbative numerical simulation.

At some critical temperature T_c , the effective potential energy develops two or more degenerate minima as a function of some order parameter, but the system remains in the higher energy metastable phase even after it is energetically less favourable, as the temperature continues to fall. Eventually this will lead into bubble nucleation, where bubbles of the stable phase form.

In the field theoretical context – at zero temperature – a metastable vacuum state can decay via quantum tunnelling [38]. A complete one-loop calculation of the rate of this process was first carried out by Coleman and Callan [39, 40], using a saddlepoint approximation of the path integral. This result has since been rederived from a number of perspectives and formalisms [41–43].

At high temperature, bubbles may also nucleate classically. Within classical field theory, a complete oneloop calculation for this process was first carried out by Langer [44–46], extending the tree-level theory of Cahn and Hilliard [47]. However, the analogous calculation within quantum field theory is much less clear. Early papers by Linde [48] and Affleck [49] gave slightly different expressions in the high temperature regime. Neither agrees with the classical result of Langer.

The above methods all depend on a semiclassical picture of the bubble and its fluctuations. A fully numerical, non-perturbative lattice calculation of the bubble nucleation rate was introduced in Ref. [50]. At the time, the focus was on the physics of the minimal Standard Model, where a first-order phase transition would have arisen from radiative corrections to the quartic Higgs potential. In Ref. [51] a toy model with similar features was studied. More recently, in Ref. [52] a more complete study of nucleation in the minimal Standard Model was carried out, motivated by the idea that any sufficiently heavy new particles could be integrated out [22].

Another approach has been to study thermal nucleation directly as a real time process [53–60]. This involves evolving the lattice in time, waiting until a growing bub-

^{*} oliver.gould@nottingham.ac.uk

[†] anna.kormu@helsinki.fi

[‡] david.weir@helsinki.fi

ble appears and recording the time taken for the bubble to appear. Since the nucleation rate can vary over many orders of magnitude, this approach is only viable over a narrow temperature range.

We consider what we call the real scalar theory [61], with Lagrangian

$$\mathscr{L} = -\frac{1}{2}\partial_{\mu}\varphi\partial^{\mu}\varphi - V(\varphi) - J_{1}\varphi - J_{2}\varphi^{2}, \quad (1)$$

$$V(\varphi) = \sigma\varphi + \frac{1}{2}m^2\varphi^2 + \frac{1}{3!}g\varphi^3 + \frac{1}{4!}\lambda\varphi^4, \qquad (2)$$

where σ , m^2 , g and λ are the model parameters, J_1 and J_2 represent operators of other fields, and we have used mostly plus signature. Such a scalar field may couple to the Standard Model, through Higgs portal interactions (with $J_1 = \kappa_1 H^{\dagger} H$ and $J_2 = \kappa_2 H^{\dagger} H/2$) or serve as a toy model in itself to test key ideas.

Compared with radiatively-induced transitions, the presence of the tree-level barrier in equation (2) is expected to allow for stronger transitions [62]. The perturbative expansion is also simpler [63, 64], allowing us to focus on bubble nucleation without extraneous details.

We will focus on the dynamics of a single bubble and its real time evolution from the metastable phase to the stable phase. The bubbles arise as long wavelength thermal fluctuations of the scalar field. These fluctuations are highly occupied in a thermal bath, and so their effective evolution is classical [65–68]. The effects of short wavelength fluctuations appear both through screening of the effective parameters, and as noise and damping in the evolution equations [68–70].

Thermal fluctuations significantly modify the scalar effective potential for temperatures $T^2 \sim m^2/\lambda$ and above. At such temperatures, modes with energies of order $m \sim \sqrt{\lambda}T$ satisfy the following Langevin equation,

$$\partial_t \phi(t, \mathbf{x}) = \pi(t, \mathbf{x}),\tag{3}$$

$$\partial_t \pi(t, \mathbf{x}) = -\frac{\delta H_{\text{eff}}}{\delta \phi} - \gamma \pi(t, \mathbf{x}) + \xi(t, \mathbf{x}), \qquad (4)$$

where (after absorbing factors of T) ϕ is the effective field, π is its canonical momentum, γ is a damping parameter and ξ is a local Gaussian noise term, satisfying

$$\langle \xi(t, \mathbf{x})\xi(t', \mathbf{x}') \rangle = 2\gamma \delta(t - t')\delta^{(3)}(\mathbf{x} - \mathbf{x}').$$
 (5)

The effective Hamiltonian, $H_{\rm eff}$, is constructed to match the long-wavelength equal-time correlation functions of the full quantum theory [71, 72]. This matching reduces the long-wavelength equilibrium thermodynamics of a four-dimensional (4d) quantum field theory to that of a 3d classical statistical field theory, and hence is known as high-temperature dimensional reduction (see Section I in the Supplemental Material for details). Importantly, for a scalar field theory, an appropriate choice of γ also leads to the matching of the unequal time correlation functions [65–68].

II. SADDLEPOINT APPROXIMATION

Perturbative calculations of the bubble nucleation rate are based on saddlepoint approximations to the probability current between phases. To arrive at this from Eqs. (3) and (4), one introduces the probability density $P(\phi, \pi)$, which satisfies a Fokker-Planck equation [45, 73]. If the metastable state is sufficiently long-lived, the calculation can be set up as a static problem: one imposes boundary conditions that the metastable phase is populated thermally $- P(\phi, \pi) \propto e^{-H_{\text{eff}}[\phi, \pi]}$ – and the stable phase is unpopulated. There is then a constant, small flux of probability over the barrier between phases, which does not appreciably deplete the metastable state nor appreciably populate the stable state [74].

For field theories, the computation of this probability current was described by Langer [44–46]. The relationship of Langer's formalism to high-temperature quantum field theories has been studied in Refs. [75, 76], and generalised to higher orders in Ref. [77]. A crucial result is that the nucleation rate factorises

$$\Gamma = A_{\rm dyn} \times A_{\rm stat},\tag{6}$$

where A_{dyn} is the *dynamical* factor and A_{stat} is the *sta*tistical factor. The latter is a purely time-independent quantity, equal to the vacuum nucleation rate in 2+1 dimensions. At leading (one-loop) order, $2\pi A_{dyn}$ is equal to the initial growth rate of the critical bubble. Note that while this split into dynamical and statistical parts mirrors the split in our lattice approach, there are nevertheless differences.

The statistical part takes the form of a semiclassical path integral. Its computation requires first solving for a radially symmetric saddlepoint of the 3d tree-level action defined in Section I of the Supplemental Material,

$$\frac{\mathrm{d}^2\phi}{\mathrm{d}r^2} + \frac{2}{r}\frac{\mathrm{d}\phi}{\mathrm{d}r} = \frac{\mathrm{d}V_3(\phi)}{\mathrm{d}\phi}.$$
(7)

The relevant solution, the critical bubble, has spherical symmetry. This reduces the problem to solving a onedimensional boundary value problem, which be solved using the shooting method [39].

With the critical bubble $\phi_{\rm b}$ in hand, as well as the position of the metastable phase ϕ_0 , the statistical part of the nucleation rate is proportional to the relative Boltzmann-weighted phase-space volume of these two stationary points,

$$A_{\rm stat} = \sqrt{\left|\frac{\det(S''[\phi_0]/2\pi)}{\det'(S''[\phi_{\rm b}]/2\pi)}\right|} \left(\frac{\Delta S[\phi_{\rm b}]}{2\pi}\right)^{3/2} e^{-\Delta S[\phi_{\rm b}]},$$
(8)

where S'' denotes the second functional derivative of the action, the prime on det' denotes that zero modes are not to be included, and $\Delta S[\phi_{\rm b}] \equiv S[\phi_{\rm b}] - S[\phi_0]$. The computation of the functional determinants can be carried out using the Gelfand-Yaglom theorem [78, 79].

For the dynamical factor, one finds that

$$A_{\rm dyn} = \frac{1}{2\pi} \left(\sqrt{|\lambda_-| + \frac{\gamma^2}{4} - \frac{\gamma}{2}} \right), \tag{9}$$

where λ_{-} is the single negative eigenvalue of $S''[\phi_b]$ and γ is the damping rate of equation (4).

For the present theory, the computation of both A_{stat} and A_{dyn} can be carried out using the numerical package BubbleDet [80], or the numerical fits from Ref. [81]. This yields our full one-loop rate, $\Gamma_{\text{one-loop}}$.

However, for applications of nucleation theory, especially for more complicated models, neither the functional determinant nor the dynamical prefactor are typically computed. In Section II in the Supplemental Material we show two common lower-order approximations: in the first, $\Gamma_{\text{tree-level}}$, the functional determinant and dynamical prefactor are estimated as T^4 , and in the second, Γ_{LPA} , a local potential approximation is made for the functional determinant, together with a prescription for how to drop unwanted imaginary parts.

III. LATTICE SIMULATIONS

Given the limitations of directly simulating the Hamiltonian system of Eqs. (3) and (4), our nonperturbative calculation of the nucleation rate closely follows the approach introduced in Refs. [50, 51].

The lattice discretisation of the action takes the form

$$S_{\text{lat}} = \sum_{x} a^{3} \left[-\frac{1}{2} Z_{\phi} \phi_{x} (\nabla_{\text{lat}}^{2} \phi)_{x} + \sigma_{\text{lat}} \phi_{x} + \frac{1}{2} Z_{\phi} Z_{m} m_{\text{lat}}^{2} \phi_{x}^{2} + \frac{1}{4!} Z_{\phi}^{2} \lambda_{\text{lat}} \phi_{x}^{4} \right], \quad (10)$$

where the sum over x extends over the 3d lattice sites. We have removed the cubic term by a constant shift in the field. One can derive exact lattice-continuum relations for this model within lattice perturbation theory [82, 83], and improvements up to $O(a^2)$ [51, 84, 85]. See Section III in the Supplemental Material for more information about our lattice action.

There are two stages: lattice Monte Carlo simulations which generate configurations, some of which lie close to the separatrix (sometimes known as the 'transition surface' [45, 75]) of field configurations between the two phases. Selected near-separatrix configurations are then evolved numerically in a thermal bath to determine whether they tunnel or not.

A. Critical bubble probability

The separatrix configurations are suppressed by $\sim e^{-65}$ at the temperature we simulate. Our lattice Monte Carlo simulations therefore use the multicanonical method [86, 87] to overcome this suppression and



FIG. 1. Probability distribution $P(\theta_{\rm op})$ of the order parameter at some temperature below $T_{\rm c}$. The metastable and stable phases are separated by an exponentially suppressed area, the mixed critical bubble. The separatrix configurations are drawn from the narrow range ϵ around the critical bubble.

generate a sufficient number of near-critical bubble configurations (see Ref. [61] for its application to the current model). The order parameter θ_{op} is measured, yielding a histogram that (below the critical temperature) will resemble Fig. 1. If the choice of order parameter, system volume and geometry admit, then there will be a local minimum between the metastable and stable phase peaks that we identify with the separatrix. From this local minimum, an estimate of θ_{op} corresponding to critical bubble configurations, $\theta_{\rm c}$, can be inferred. Configurations sufficiently near the separatrix $\theta_{\rm op} \in [\theta_{\rm c} - \frac{\epsilon}{2}, \theta_{\rm c} + \frac{\epsilon}{2}]$ are used as the initial conditions for evolution of the the lattice versions of the stochastic evolution equations (3) and (4), along with momenta drawn from a Gaussian distribution. The final result is independent of the exact choice of ϵ , but it should be sufficiently small that the sampled configurations correspond to near-critical bubbles.

From the selected near-separatrix field configurations we can determine the probabilities $P(\theta_{\rm op})$ for the critical bubble and metastable phases and we normalise the probability of being in the critical configuration to that of the metastable phase,

$$P_c^{\text{normalised}} = \frac{P(|\theta_{\text{op}} - \theta_{\text{c}}| < \epsilon/2)}{\epsilon P(\theta_{\text{op}} < \theta_{\text{c}})}.$$
 (11)

B. Effective tunnelling fraction

Following Ref. [51], the dynamical information can be separated into two parts, the flux – the rate of change of the order parameter as it crosses the separatrix – and \mathbf{d} – the ratio of tunnelled configurations determined by real time trajectories. To determine whether a configuration tunnels or not, we directly evolve the stochastic Hamiltonian equations, (3) and (4), both forwards and backwards in time, with the initial momenta reversed for the backwards evolution (see Fig. 2). We use a timestep $\Delta t\lambda_3 = 0.01$ much smaller than the lattice spacing.

If a given trajectory begins and ends in different phases (either metastable \rightarrow stable or vice versa), we set $\delta_{\text{tunnel}} = 1$, otherwise $\delta_{\text{tunnel}} = 0$. The ratio of tunnelled



FIG. 2. Snapshots of a nucleating bubble with lattice size $L\lambda_3 = 72$ and spacing $a\lambda_3 = 1.5$. We remove ultraviolet fluctuations by performing eight steps of nearest neighbour averaging. Here, $t = 0\Delta t$ corresponds to the initial separatrix configuration and the negative time direction describes the backward time evolution. Note that the snapshots are not evenly spaced in time.

trajectories is an observable defined for each trajectory as

$$\mathbf{d} = \frac{\delta_{\text{tunnel}}}{N_{\text{crossings}}},\tag{12}$$

where $N_{\text{crossings}}$ is the number of times the given trajectory crosses θ_c . Lastly, we must add a factor of $\frac{1}{2}$ to account for metastable \rightarrow stable trajectories only, to avoid double counting the tunnelled trajectories.

The full nucleation rate on the lattice is a combination of the above three elements,

$$\Gamma \mathcal{V} = P_c^{\text{normalised}} \left\langle \frac{1}{2} \text{flux} \times \mathbf{d} \right\rangle, \qquad (13)$$

where \mathcal{V} is the lattice volume.

For field-theory systems with many degrees of freedom, the flux is dominated by uncorrelated short-time fluctuations, whereas the global behaviour of the trajectories depends on longer-time correlations. These are therefore assumed to be uncorrelated and we approximate the rate by,

$$\Gamma \mathcal{V} \approx P_c^{\text{normalised}} \frac{1}{2} \langle \text{flux} \rangle \langle \mathbf{d} \rangle.$$
 (14)

These formulae are order parameter independent, but the choice of order parameter affects the viability of this method. At larger volumes, the contribution of bulk fluctuations about the metastable minimum will in general grow relative to the critical bubble. The physical volume occupied by the critical bubble is fixed, so that eventually bulk fluctuations dominate.

The flux through the critical separatrix surface is order parameter dependent. It can be solved analytically due to the Gaussianity of the momentum field [51]

$$\langle \text{flux} \rangle = \left\langle \left| \frac{\Delta \theta_{\text{op}}}{\Delta t} \right|_{\theta_{\text{c}}} \right\rangle = \sqrt{\frac{8}{\pi \mathcal{V}} (\theta_{\text{c}} + A^2)}, \quad (15)$$

where our order parameter $\theta_{\rm op} = \bar{\phi}^2 - 2A\bar{\phi}$ and A is a constant which we choose to be the peak of the histogram of ϕ in metastable phase. We have used overbar to mean volume averaging and angle brackets for statistical averaging. We also considered $\theta'_{\rm op} = \bar{\phi}$ (with corresponding flux $\sqrt{2/(\pi \mathcal{V})}$) but found that bulk fluctuations affected our ability to go to large volumes with this choice.

C. Results

In Figure 3, we show how the obtained nucleation rate varies with lattice spacing and volume. The dependence on both is rather mild, allowing us to take controlled continuum extrapolations.

Given our use of an $O(a^2)$ improved lattice discretisation, the leading dependence on lattice spacing arises at $O(a^3)$. For a fixed volume of $L\lambda_3 = 42$ we vary the lattice spacing and perform a least-squares fit $f(a) = b + ca^3$ to the logarithm of the rate. The best fit line, with $\chi^2/dof = 5.53$, is plotted in the figure. Jackknife resampling this fit yields $\log(\Gamma/\lambda_3^4) = -73.26(10)$, where the error is dominated by our largest lattice spacing $a\lambda_3 = 3.5$. This is approximately the inverse screening mass of the system, at which point the cubic form is expected to break down. A one-loop estimate of the screening mass yields $m_s^{\rm PT}/\lambda_3 = 0.294(1)$ [61]. From the figure we deduce that lattice artefacts are comparable to statistical uncertainties already with $\lambda_3 a = 1.5$.

For reference we also show the nucleation rate computed using the linear order parameter $\theta_{\rm op} = \phi_{\rm lin}$ for one lattice spacing. We find good agreement (within errors) with our final results, however our final choice of order parameter shows substantially reduced errors, which we attribute to the suppression of the metastable phase bulk fluctuations by our new choice of order parameter.

We then show the dependence on volume at constant lattice spacing $\lambda_3 a = 1.5$. Away from the second-order phase transition, this 3d Euclidean model has no massless modes, so long-range correlations die off exponentially with distance. Motivated by this, we fit the logarithm of the rate to an exponential $f(L) = b + c \exp(-m_s L)$, finding $m_s/\lambda_3 = 0.287(12)$, consistent with the perturbative screening mass. We show the best fit line in the figure, with $\chi^2/\text{dof} = 1.54$. Jackknife resampling this fit yields $\log(\Gamma/\lambda_3^4) = -74.08(5)$ for the infinite-volume extrapolation. Note that fitting $f(L) = b + c/L^n$ with $n \in \{1, 2, 3\}$ to our finite-volume data yields much poorer fits with $\chi^2/\text{do.f.} \sim 10^2$.

Even with our improved choice of order parameter, we are limited in the largest volume we can reach for a number of reasons. For sufficiently large volumes, critical



FIG. 3. The zero lattice spacing (top) and infinite volume (bottom) extrapolations, together with cubic and exponential fits respectively. The vertical line in the top plot marks a perturbative estimate of the correlation length.

bubble configurations are again buried under the bulk fluctuations of the metastable phase. Nevertheless, we see that the bubble is well resolved for the volumes we simulate in two ways: both geometrically determining which interface geometry (bubble, cylinder or slab) is favoured in the thin-wall approximation [50], and also visual inspection of configurations show a well-resolved spherical bubble configuration, see Fig. 2.

IV. DISCUSSION

The main result of this work is the logarithm of the nucleation rate, reliably computed to high precision for a real scalar quantum field theory at high temperature. Statistical uncertainties are much smaller than discrepancies with other methods, as seen in Fig. 4. Systematic uncertainties related to the continuum extrapolation are also well under control, as evidenced by Fig. 3. This has been made possible by use of efficient multicanonical algorithms [50, 51, 86, 87], and exact and $O(a^2)$ improved lattice-continuum relations [51, 82–85].

The benchmark parameter point that we simulated was chosen with the hope of testing perturbation theory in a regime where it is expected to work well. At the critical temperature, the tree-level approximation to the dis-



FIG. 4. The nucleation rate as a function of temperature. Uncertainty bands for the tree-level and one-loop perturbative results are based on varying the renormalisation scale over $\mu_3/\lambda_3 \in \{0.5, 1, 2\}$. The uncertainty estimate on the LPA relflects different choices for removing imaginary parts of the potential. The lattice points are continuum extrapolated; the red triangle highlights the temperature actually simulated, while the orange circles have utilised reweighting. The orange continuous line is the reweighted result for $a\lambda_3 = 1.5$, $L\lambda_3 =$ 60. The results in this figure are tabulated at https://doi. org/10.5281/zenodo.10891523.

continuity in the background field value agrees with the lattice to 10% accuracy, and the one-loop approximation agrees to better than 1% [61].

Yet, for $|\log \Gamma|$ at the temperature simulated, we have found that the lattice result is 100% larger than the treelevel approximation and almost 20% larger than the oneloop approximation. It may be that this is explained by the growth of the loop expansion parameter as the system supercools, which becomes almost twice as large at the temperature we have simulated than at the critical temperature. Note that the discrepancy in $|\log \Gamma|$ cannot be explained by a shift of the critical temperature between orders, as in this model the critical temperature is fixed by the Z_2 symmetry between phases, and hence is exact already at tree-level within the 3d EFT [61].

However, what if the discrepancy between the one-loop and lattice results cannot be explained simply by the next loop order? While the critical bubble is a saddlepoint of the path integral, there may be other relevant saddlepoints [88, 89]. Further, for such weakly-damped systems, it has been argued that the saddlepoint approximation itself breaks down for the dynamical part of the nucleation rate [77, 90]. Alternatively, a number of suggestions have been put forward to resolve the longstanding discrepancy between theory and experiment for the nucleation rate between the A and B phases of superfluid ³He, including resonant tunnelling and tunnelling via intermediate solitonic configurations [91, 92].

As far as we are aware, there has never been a calculation of the thermal nucleation rate of any quantum field theory beyond one-loop order. To really test whether perturbation theory provides a reliable guide to bubble nucleation, a complete two-loop calculation at our benchmark point provides a clear target for future work. Further, nonperturbative results over a range of couplings could test convergence.

For phenomenological studies of cosmological phase transitions, and their gravitational wave signals, our results demonstrate that the widely-used tree-level and local potential approximations give relatively poor accuracy for the logarithm of the bubble nucleation rate. The full one-loop approximation fares significantly better. This calculation can be carried out in simple models using existing numerical tools [80], but is still out of reach for e.g. scalar extensions of the Standard Model.

While our lattice simulations give reliable and precise predictions for the nucleation rate, their calculation is slow. To use such lattice simulations for phenomenological studies of cosmological phase transitions, faster sampling of the relevant bubble configurations is needed. We have identified one direction of progress, through the development of order parameters optimised to enhance the sampling of tunnelling configurations. Our quadratic order parameter $\theta_{\rm op} = \bar{\phi}^2 - 2A\bar{\phi}$ was crucial for efficiently simulating the larger volumes. In a subsequent paper we will explore this further and also set out how our computations can be applied to predictions of the gravitational wave spectrum.

ACKNOWLEDGMENTS

The authors would like to thank Kari Rummukainen for his insights and advice at a number of points throughout this study. O.G. (ORCID ID 0000-0002-7815-3379) was supported by the Research Funds of the University of Helsinki, U.K. Science and Technology Facilities Council (STFC) Consolidated Grant ST/T000732/1, a Research Leadership Award from the Leverhulme Trust, and a Royal Society Dorothy Hodgkin Fellowship. A.K. (OR-CID ID 0000-0002-0309-3471) was supported by Research Council of Finland grant no. 328958 and a travel grant from the Jenny and Antti Wihuri Foundation. D.J.W. (ORCID ID 0000-0001-6986-0517) was supported by Research Council of Finland grant nos. 324882, 328958, 349865 and 353131. The authors also wish to acknowledge CSC - IT Center for Science, Finland, for computational resources.

DATA ACCESS STATEMENT

Data supporting this paper is available from Zenodo at https://doi.org/10.5281/zenodo.10891523.

- V. A. Kuzmin, V. A. Rubakov, and M. E. Shaposhnikov, On the Anomalous Electroweak Baryon Number Nonconservation in the Early Universe, Phys. Lett. B 155, 36 (1985).
- [2] J. M. Cline, A. Friedlander, D.-M. He, K. Kainulainen, B. Laurent, and D. Tucker-Smith, Baryogenesis and gravity waves from a UV-completed electroweak phase transition, Phys. Rev. D 103, 123529 (2021), arXiv:2102.12490 [hep-ph].
- [3] J. Ellis, M. Lewicki, M. Merchand, J. M. No, and M. Zych, The scalar singlet extension of the Standard Model: gravitational waves versus baryogenesis, JHEP 01, 093, arXiv:2210.16305 [hep-ph].
- [4] S. Bruggisser, B. von Harling, O. Matsedonskyi, and G. Servant, Status of electroweak baryogenesis in minimal composite Higgs, JHEP 08, 012, arXiv:2212.11953 [hep-ph].
- [5] S. Kanemura and Y. Mura, Electroweak baryogenesis via top-charm mixing, JHEP 09, 153, arXiv:2303.11252 [hep-ph].
- [6] C. Idegawa and E. Senaha, Electron electric dipole moment and electroweak baryogenesis in a complex singlet extension of the Standard Model with degenerate scalars, Phys. Lett. B 848, 138332 (2024), arXiv:2309.09430 [hep-ph].
- [7] M. B. Hindmarsh, M. Lüben, J. Lumma, and M. Pauly, Phase transitions in the early universe, SciPost Phys. Lect. Notes 24, 1 (2021), arXiv:2008.09136 [astroph.CO].
- [8] G. Agazie et al. (NANOGrav), The NANOGrav 15 yr Data Set: Evidence for a Gravitational-wave

Background, Astrophys. J. Lett. **951**, L8 (2023), arXiv:2306.16213 [astro-ph.HE].

- [9] J. Antoniadis *et al.* (EPTA, InPTA:), The second data release from the European Pulsar Timing Array - III. Search for gravitational wave signals, Astron. Astrophys. **678**, A50 (2023), arXiv:2306.16214 [astro-ph.HE].
- [10] H. Xu *et al.*, Searching for the Nano-Hertz Stochastic Gravitational Wave Background with the Chinese Pulsar Timing Array Data Release I, Res. Astron. Astrophys. **23**, 075024 (2023), arXiv:2306.16216 [astroph.HE].
- [11] D. J. Reardon *et al.*, Search for an Isotropic Gravitational-wave Background with the Parkes Pulsar Timing Array, Astrophys. J. Lett. **951**, L6 (2023), arXiv:2306.16215 [astro-ph.HE].
- [12] G. Agazie *et al.* (NANOGrav), The NANOGrav 15 yr Data Set: Constraints on Supermassive Black Hole Binaries from the Gravitational-wave Background, Astrophys. J. Lett. **952**, L37 (2023), arXiv:2306.16220 [astroph.HE].
- [13] A. Afzal *et al.* (NANOGrav), The NANOGrav 15 yr Data Set: Search for Signals from New Physics, Astrophys. J. Lett. **951**, L11 (2023), arXiv:2306.16219 [astroph.HE].
- [14] C. Caprini *et al.*, Science with the space-based interferometer eLISA. II: Gravitational waves from cosmological phase transitions, JCAP **04**, 001, arXiv:1512.06239 [astro-ph.CO].
- [15] C. Caprini *et al.*, Detecting gravitational waves from cosmological phase transitions with LISA: an update, JCAP **03**, 024, arXiv:1910.13125 [astro-ph.CO].

- [16] P. Auclair *et al.* (LISA Cosmology Working Group), Cosmology with the Laser Interferometer Space Antenna, Living Rev. Rel. **26**, 5 (2023), arXiv:2204.05434 [astro-ph.CO].
- [17] M. Hindmarsh, S. J. Huber, K. Rummukainen, and D. J. Weir, Shape of the acoustic gravitational wave power spectrum from a first order phase transition, Phys. Rev. D 96, 103520 (2017), [Erratum: Phys.Rev.D 101, 089902 (2020)], arXiv:1704.05871 [astro-ph.CO].
- [18] R. Jinno, T. Konstandin, H. Rubira, and I. Stomberg, Higgsless simulations of cosmological phase transitions and gravitational waves, JCAP 02, 011, arXiv:2209.04369 [astro-ph.CO].
- [19] T. Konstandin, Gravitational radiation from a bulk flow model, JCAP 03, 047, arXiv:1712.06869 [astro-ph.CO].
- [20] M. Hindmarsh and M. Hijazi, Gravitational waves from first order cosmological phase transitions in the Sound Shell Model, JCAP **12**, 062, arXiv:1909.10040 [astroph.CO].
- [21] K. Kainulainen, V. Keus, L. Niemi, K. Rummukainen, T. V. I. Tenkanen, and V. Vaskonen, On the validity of perturbative studies of the electroweak phase transition in the Two Higgs Doublet model, JHEP 06, 075, arXiv:1904.01329 [hep-ph].
- [22] O. Gould, J. Kozaczuk, L. Niemi, M. J. Ramsey-Musolf, T. V. I. Tenkanen, and D. J. Weir, Nonperturbative analysis of the gravitational waves from a first-order electroweak phase transition, Phys. Rev. D 100, 115024 (2019), arXiv:1903.11604 [hep-ph].
- [23] A. Ekstedt and J. Löfgren, A Critical Look at the Electroweak Phase Transition, JHEP 12, 136, arXiv:2006.12614 [hep-ph].
- [24] D. Croon, O. Gould, P. Schicho, T. V. I. Tenkanen, and G. White, Theoretical uncertainties for cosmological first-order phase transitions, JHEP 04, 055, arXiv:2009.10080 [hep-ph].
- [25] O. Gould and T. V. I. Tenkanen, On the perturbative expansion at high temperature and implications for cosmological phase transitions, JHEP 06, 069, arXiv:2104.04399 [hep-ph].
- [26] A. Ekstedt, Convergence of the nucleation rate for firstorder phase transitions, Phys. Rev. D 106, 095026 (2022), arXiv:2205.05145 [hep-ph].
- [27] P. Schiffer, D. Osheroff, and A. Leggett, Nucleation of the AB transition in superfluid ³He: experimental and theoretical considerations, in *Progress in Low Temperature Physics*, Vol. 14 (Elsevier, 1995) pp. 159–211.
- [28] Y. Tian *et al.*, Supercooling of the A phase of ³He, Nature Commun. **14**, 148 (2023), arXiv:2208.14981 [condmat.other].
- [29] M. Hindmarsh *et al.*, A-B transition in superfluid ³He and cosmological phase transitions (2024) arXiv:2401.07878 [cond-mat.supr-con].
- [30] A. Zenesini, A. Berti, R. Cominotti, C. Rogora, I. G. Moss, T. P. Billam, I. Carusotto, G. Lamporesi, A. Recati, and G. Ferrari, Observation of false vacuum decay via bubble formation in ferromagnetic superfluids, Nature Physics, 1 (2023), arXiv:2305.05225 [hep-ph].
- [31] O. Fialko, B. Opanchuk, A. I. Sidorov, P. D. Drummond, and J. Brand, Fate of the false vacuum: towards realization with ultra-cold atoms, EPL 110, 56001 (2015), arXiv:1408.1163 [cond-mat.quant-gas].
- [32] O. Fialko, B. Opanchuk, A. I. Sidorov, P. D. Drummond, and J. Brand, The universe on a table top: engi-

neering quantum decay of a relativistic scalar field from a metastable vacuum, J. Phys. B **50**, 024003 (2017), arXiv:1607.01460 [cond-mat.quant-gas].

- [33] T. P. Billam, R. Gregory, F. Michel, and I. G. Moss, Simulating seeded vacuum decay in a cold atom system, Phys. Rev. D 100, 065016 (2019), arXiv:1811.09169 [hep-th].
- [34] T. P. Billam, K. Brown, and I. G. Moss, Bubble nucleation in a cold spin 1 gas, New J. Phys. 25, 043028 (2023), arXiv:2212.03621 [cond-mat.quant-gas].
- [35] T. P. Billam, K. Brown, and I. G. Moss, Falsevacuum decay in an ultracold spin-1 Bose gas, Phys. Rev. A 105, L041301 (2022), arXiv:2108.05740 [condmat.quant-gas].
- [36] A. C. Jenkins, J. Braden, H. V. Peiris, A. Pontzen, M. C. Johnson, and S. Weinfurtner, Analog vacuum decay from vacuum initial conditions, Phys. Rev. D 109, 023506 (2024), arXiv:2307.02549 [cond-mat.quant-gas].
- [37] A. C. Jenkins, I. G. Moss, T. P. Billam, Z. Hadzibabic, H. V. Peiris, and A. Pontzen, Generalized cold-atom simulators for vacuum decay, (2023), arXiv:2311.02156 [cond-mat.quant-gas].
- [38] I. Y. Kobzarev, L. B. Okun, and M. B. Voloshin, Bubbles in Metastable Vacuum, Yad. Fiz. 20, 1229 (1974).
- [39] S. R. Coleman, The Fate of the False Vacuum. 1. Semiclassical Theory, Phys. Rev. D 15, 2929 (1977), [Erratum: Phys.Rev.D 16, 1248 (1977)].
- [40] C. G. Callan, Jr. and S. R. Coleman, The Fate of the False Vacuum. 2. First Quantum Corrections, Phys. Rev. D 16, 1762 (1977).
- [41] A. D. Plascencia and C. Tamarit, Convexity, gaugedependence and tunneling rates, JHEP 10, 099, arXiv:1510.07613 [hep-ph].
- [42] A. Andreassen, D. Farhi, W. Frost, and M. D. Schwartz, Direct Approach to Quantum Tunneling, Phys. Rev. Lett. 117, 231601 (2016), arXiv:1602.01102 [hep-th].
- [43] W.-Y. Ai, B. Garbrecht, and C. Tamarit, Functional methods for false vacuum decay in real time, JHEP 12, 095, arXiv:1905.04236 [hep-th].
- [44] J. S. Langer, Theory of the condensation point, Annals Phys. 41, 108 (1967).
- [45] J. S. Langer, Statistical theory of the decay of metastable states, Annals Phys. 54, 258 (1969).
- [46] J. Langer, Metastable states, Physica **73**, 61 (1974).
- [47] J. W. Cahn and J. E. Hilliard, Free energy of a nonuniform system. III. Nucleation in a two-component incompressible fluid, The Journal of Chemical Physics **31**, 688 (1959).
- [48] A. D. Linde, Fate of the False Vacuum at Finite Temperature: Theory and Applications, Phys. Lett. B 100, 37 (1981).
- [49] I. Affleck, Quantum Statistical Metastability, Phys. Rev. Lett. 46, 388 (1981).
- [50] G. D. Moore and K. Rummukainen, Electroweak bubble nucleation, nonperturbatively, Phys. Rev. D 63, 045002 (2001), arXiv:hep-ph/0009132.
- [51] G. D. Moore, K. Rummukainen, and A. Tranberg, Nonperturbative computation of the bubble nucleation rate in the cubic anisotropy model, JHEP 04, 017, arXiv:hep-lat/0103036.
- [52] O. Gould, S. Güyer, and K. Rummukainen, Firstorder electroweak phase transitions: A nonperturbative update, Phys. Rev. D 106, 114507 (2022), arXiv:2205.07238 [hep-lat].

- [53] D. Y. Grigoriev and V. A. Rubakov, Soliton Pair Creation at Finite Temperatures. Numerical Study in (1+1)-dimensions, Nucl. Phys. B 299, 67 (1988).
- [54] O. T. Valls and G. F. Mazenko, Nucleation in a timedependent Ginzburg-Landau model: A numerical study, Physical Review B 42, 6614 (1990).
- [55] M. G. Alford, H. Feldman, and M. Gleiser, Thermal nucleation of kink-antikink pairs, Phys. Rev. Lett. 68, 1645 (1992).
- [56] M. G. Alford, H. Feldman, and M. Gleiser, Thermal activation of metastable decay: Testing nucleation theory, Phys. Rev. D 47, R2168 (1993).
- [57] M. G. Alford and M. Gleiser, Metastability in twodimensions and the effective potential, Phys. Rev. D 48, 2838 (1993), arXiv:hep-ph/9304245.
- [58] S. Borsanyi, A. Patkos, J. Polonyi, and Z. Szep, Fate of the classical false vacuum, Phys. Rev. D 62, 085013 (2000), arXiv:hep-th/0004059.
- [59] L. Batini, A. Chatrchyan, and J. Berges, Real-time dynamics of false vacuum decay, Phys. Rev. D 109, 023502 (2024), arXiv:2310.04206 [hep-th].
- [60] D. Pîrvu, M. C. Johnson, and S. Sibiryakov, Bubble velocities and oscillon precursors in first order phase transitions, (2023), arXiv:2312.13364 [hep-th].
- [61] O. Gould, Real scalar phase transitions: a nonperturbative analysis, JHEP 04, 057, arXiv:2101.05528 [hep-ph].
- [62] D. J. H. Chung, A. J. Long, and L.-T. Wang, 125 GeV Higgs boson and electroweak phase transition model classes, Phys. Rev. D 87, 023509 (2013), arXiv:1209.1819 [hep-ph].
- [63] A. Ekstedt, O. Gould, and J. Löfgren, Radiative firstorder phase transitions to next-to-next-to-leading order, Phys. Rev. D 106, 036012 (2022), arXiv:2205.07241 [hep-ph].
- [64] O. Gould and T. V. I. Tenkanen, Perturbative effective field theory expansions for cosmological phase transitions, JHEP 01, 048, arXiv:2309.01672 [hep-ph].
- [65] G. Aarts and J. Smit, Finiteness of hot classical scalar field theory and the plasmon damping rate, Phys. Lett. B 393, 395 (1997), arXiv:hep-ph/9610415.
- [66] G. Aarts and J. Smit, Classical approximation for time dependent quantum field theory: Diagrammatic analysis for hot scalar fields, Nucl. Phys. B 511, 451 (1998), arXiv:hep-ph/9707342.
- [67] D. Bodeker, Classical real time correlation functions and quantum corrections at finite temperature, Nucl. Phys. B 486, 500 (1997), arXiv:hep-th/9609170.
- [68] C. Greiner and B. Muller, Classical fields near thermal equilibrium, Phys. Rev. D 55, 1026 (1997), arXiv:hepth/9605048.
- [69] M. Morikawa, Classical Fluctuations in Dissipative Quantum Systems, Phys. Rev. D 33, 3607 (1986).
- [70] F. Gautier and J. Serreau, On the Langevin description of nonequilibrium quantum fields, Phys. Rev. D 86, 125002 (2012), arXiv:1209.1827 [hep-th].
- [71] K. Kajantie, M. Laine, K. Rummukainen, and M. E. Shaposhnikov, Generic rules for high temperature dimensional reduction and their application to the standard model, Nucl. Phys. B 458, 90 (1996), arXiv:hepph/9508379.
- [72] E. Braaten and A. Nieto, Effective field theory approach to high temperature thermodynamics, Phys. Rev. D 51, 6990 (1995), arXiv:hep-ph/9501375.

- [73] A. Berera, J. Mabillard, B. W. Mintz, and R. O. Ramos, Formulating the Kramers problem in field theory, Phys. Rev. D 100, 076005 (2019), arXiv:1906.08684 [hep-ph].
- [74] H. A. Kramers, Brownian motion in a field of force and the diffusion model of chemical reactions, Physica 7, 284 (1940).
- [75] O. Gould and J. Hirvonen, Effective field theory approach to thermal bubble nucleation, Phys. Rev. D 104, 096015 (2021), arXiv:2108.04377 [hep-ph].
- [76] J. Hirvonen, Nucleation Rate in a High-Temperature Quantum Field Theory with Hard Particles, (2024), arXiv:2403.07987 [hep-ph].
- [77] A. Ekstedt, Bubble nucleation to all orders, JHEP 08, 115, arXiv:2201.07331 [hep-ph].
- [78] I. M. Gelfand and A. M. Yaglom, Integration in functional spaces and it applications in quantum physics, J. Math. Phys. 1, 48 (1960).
- [79] J. Baacke and V. G. Kiselev, One loop corrections to the bubble nucleation rate at finite temperature, Phys. Rev. D 48, 5648 (1993), arXiv:hep-ph/9308273.
- [80] A. Ekstedt, O. Gould, and J. Hirvonen, BubbleDet: a Python package to compute functional determinants for bubble nucleation, JHEP 12, 056, arXiv:2308.15652 [hep-ph].
- [81] A. Ekstedt, Higher-order corrections to the bubblenucleation rate at finite temperature, Eur. Phys. J. C 82, 173 (2022), arXiv:2104.11804 [hep-ph].
- [82] M. Laine, Exact relation of lattice and continuum parameters in three-dimensional SU(2) + Higgs theories, Nucl. Phys. B 451, 484 (1995), arXiv:hep-lat/9504001.
- [83] M. Laine and A. Rajantie, Lattice continuum relations for 3-D SU(N) + Higgs theories, Nucl. Phys. B 513, 471 (1998), arXiv:hep-lat/9705003.
- [84] P. B. Arnold and G. D. Moore, Monte Carlo simulation of O(2) φ^4 field theory in three-dimensions, Phys. Rev. E **64**, 066113 (2001), [Erratum: Phys.Rev.E 68, 049902 (2003)], arXiv:cond-mat/0103227.
- [85] X.-p. Sun, Monte Carlo studies of three-dimensional O(1) and O(4) φ^4 theory related to BEC phase transition temperatures, Phys. Rev. E **67**, 066702 (2003), arXiv:hep-lat/0209144.
- [86] B. A. Berg and T. Neuhaus, Multicanonical algorithms for first order phase transitions, Phys. Lett. B 267, 249 (1991).
- [87] B. A. Berg and T. Neuhaus, Multicanonical ensemble: A New approach to simulate first order phase transitions, Phys. Rev. Lett. 68, 9 (1992), arXiv:hep-lat/9202004.
- [88] A. Andreassen, D. Farhi, W. Frost, and M. D. Schwartz, Precision decay rate calculations in quantum field theory, Phys. Rev. D 95, 085011 (2017), arXiv:1604.06090 [hep-th].
- [89] J. Braden, M. C. Johnson, H. V. Peiris, A. Pontzen, and S. Weinfurtner, New Semiclassical Picture of Vacuum Decay, Phys. Rev. Lett. **123**, 031601 (2019), [Erratum: Phys.Rev.Lett. **129**, 059901 (2022)], arXiv:1806.06069 [hep-th].
- [90] P. Hanggi, P. Talkner, and M. Borkovec, Reaction-Rate Theory: Fifty Years After Kramers, Rev. Mod. Phys. 62, 251 (1990).
- [91] D. K. Hong, Q Balls in Superfluid ³He, J. Low Temp. Phys. **71**, 483 (1988).
- [92] S. H. H. Tye and D. Wohns, Resonant Tunneling in Superfluid Helium-3, Phys. Rev. B 84, 184518 (2011), arXiv:1106.3075 [cond-mat.other].

- [93] E. J. Weinberg and A.-q. Wu, Understanding complex perturbative effective potentials, Phys. Rev. D 36, 2474 (1987).
- [94] C. L. Wainwright, CosmoTransitions: Computing Cosmological Phase Transition Temperatures and Bubble Profiles with Multiple Fields, Comput. Phys. Commun. 183, 2006 (2012), arXiv:1109.4189 [hep-ph].
- [95] C. Delaunay, C. Grojean, and J. D. Wells, Dynamics of Non-renormalizable Electroweak Symmetry Breaking, JHEP 04, 029, arXiv:0711.2511 [hep-ph].
- [96] P. Athron, C. Balazs, A. Fowlie, L. Morris, G. White, and Y. Zhang, How arbitrary are perturbative calculations of the electroweak phase transition?, JHEP 01, 050, arXiv:2208.01319 [hep-ph].
- [97] E. Forest and R. D. Ruth, Fourth Order Symplectic Integration, Physica D 43, 105 (1990).
- [98] A. M. Horowitz, A Generalized guided Monte Carlo algorithm, Phys. Lett. B 268, 247 (1991).
- [99] R. M. Neal, Handbook of Markov Chain Monte Carlo (2011) arXiv:1206.1901 [stat.CO].
- [100] H. Yoshida, Construction of higher order symplectic integrators, Phys. Lett. A 150, 262 (1990).

I. HIGH-TEMPERATURE EFFECTIVE FIELD THEORY

At high temperatures, the equilibrium thermodynamics of a weakly coupled 3+1 dimensional quantum field theory is captured by a 3 dimensional effective field theory containing only the light bosonic fields [71, 72]. The parameters of this effective theory depend on those of the full theory and on the temperature. The precise matching relations can be computed order-by-order in powers of couplings.

In the vicinity of a phase transition, the effective mass of the field undergoing the transition becomes small, and hence the longest wavelength modes may often be described by an effective theory for this field alone, with all other fields integrated out. We will assume that this is the case for the real scalar field, in which case the effective 3d Lagrangian is

$$\mathscr{L}_{\text{eff}} = \frac{1}{2} \partial_i \phi \partial_i \phi + V_3(\phi), \qquad (16)$$

where
$$V_3(\phi) = \sigma_3 \phi + \frac{m_3^2}{2} \phi^2 + \frac{g_3}{3!} \phi^3 + \frac{\lambda_3}{4!} \phi^4$$
, (17)

where i = 1, 2, 3 runs over the spatial dimensions. Note that the field has mass dimension 1/2, being canonically normalised in 3 dimensions. To leading order, it corresponds to the zero Matsubara mode of the 4d field φ divided by \sqrt{T} .

For the real-scalar extended Standard Model, the parameters of the effective action are, at leading order,

$$\sigma_3(T) = \frac{\sigma}{\sqrt{T}} + \frac{1}{24}(g + 4\kappa_1)T^{3/2},$$
 (18)

$$m_3^2(T) = m^2 + \frac{1}{24}(\lambda + 4\kappa_2)T^2,$$
(19)

$$g_3(T) = \sqrt{Tg},\tag{20}$$

$$\lambda_3(T) = T\lambda. \tag{21}$$

At this leading order, the dimensional reduction relations take the same form for a wide range of interactions to the real scalar. For example, coupling to a Dirac fermion through a Yukawa interaction, $J_1 = y\bar{\psi}\psi$, instead of to the Standard Model Higgs, would simply replace $\kappa_1 \rightarrow$ $ym_{\psi}, \kappa_2 \rightarrow y^2$ in Eqs. (18) to (21), where m_{ψ} is the tree-level mass of the fermion [75].

High-temperature dimensional reduction describes only the equilibrium thermodynamics of our model. The real-time dynamics of ϕ are described by the Langevin equations given in equation (3) and (4) [65–68]. The relevant effective Hamiltonian is

$$H_{\text{eff}} = \int \mathrm{d}^3 x \left[\frac{1}{2} \pi^2 + \frac{1}{2} \partial_i \phi \partial_i \phi + V_3(\phi) \right].$$
(22)

The parameter γ in equation (4) describes the damping and fluctuations that ϕ experiences due to hard thermal fluctuations. This parameter can be found by ensuring that the Langevin equation reproduces the longwavelength real-time correlation functions of the underlying quantum field theory. The result is that $\gamma = O(\lambda^{3/2}T/\pi)$ [68], so that the damping and noise terms are subdominant in equation (4). At leading order γ may therefore be taken to be zero.

However, a nonzero positive value for γ can help reduce finite size effects due to the heating of the lattice upon bubble nucleation. Thus, in our simulations we follow Ref. [51] and take $\gamma = 1/L$, thereby reaching $\gamma \to 0_+$ in the infinite volume limit.

For our benchmark point of the xSM, we have chosen

$$M_{\phi} = 240 \text{ GeV}, \qquad \sin \theta = 0.1, \qquad \kappa_2 = 1.5,$$

 $q = -223.75 \text{ GeV}, \qquad \lambda = 1.5489,$

where M_{ϕ} is the physical scalar mass, and θ the Higgs mixing angle, both evaluated at tree-level; for details, see Appendix A of Ref. [22]. The corresponding Lagrangian parameters are $\sigma = -5.1340 \times 10^5 \text{ GeV}^3$, $\kappa_1 = 16.937 \text{ GeV}$ and m = 108.23 GeV.

At the critical temperature (in this case $T_c = 98.513 \text{ GeV}$), this goes through the second benchmark point studied in Ref. [61]. The critical temperature is exact within the 3d EFT, as it is protected by a Z_2 symmetry. The equilibrium thermodynamics at the critical temperature was found to be under good perturbative control at one-loop order, at which point the dimensionless loop-expansion parameter within the EFT is $\alpha_3 \approx 0.13$ [61]. The temperature of spinodal decomposition, where the mass in the metastable phase goes through zero, is $T_s = 89.920$ GeV at tree-level, though this is corrected by loops within the EFT.

Our lattice simulations are carried out within the 3d effective theory. The majority of our simulations were carried out at the parameter point corresponding to T = 93.121 GeV, where we have used Eqs. (18) to (21) to fix the corresponding 3d effective parameters. The dimensionless loop-expansion parameter at this temperature is $\alpha_3 \approx 0.23$ [61]. Note that, while we have used only leading order dimensional reduction matching relations, our calculations within the 3d effective theory are valid nonperturbatively at the values of the effective parameters studied.

$T \; (\text{GeV})$	μ_3/λ_3	$\sigma_3/\lambda_3^{5/2}$	m_3^2/λ_3^2	$g_3/\lambda_3^{3/2}$	$\lambda_3 ~({\rm GeV})$
93.121	1	-0.016687	-0.082770	0	144.23

TABLE I. Parameters of our lattice simulations. Note that neither T nor λ_3 explicitly enter the simulations: the temperature is used to fix the 3d parameters through dimensional reduction, and λ_3 can be scaled out by dimensional analysis.

II. LOWER ORDER SEMICLASSICAL APPROXIMATIONS

A common approximation to the nucleation rate is to take

$$\Gamma_{\text{tree-level}} = T^4 \left(\frac{\Delta S[\phi_{\text{b}}]}{2\pi}\right)^{3/2} e^{-\Delta S[\phi_{\text{b}}]}, \qquad (23)$$

where T^4 replaces the functional determinant and dynamical prefactor.

The local potential approximation (LPA) is an alternative and widely-used approximation, utilising the oneloop effective potential when solving for the critical bubble,

$$V_{\text{one-loop}}(\phi) = -\frac{1}{12\pi} V_3''(\phi)^{3/2}.$$
 (24)

This is imaginary where $V_3''(\phi) < 0$, such as on the treelevel potential barrier between phases. Such field values are absolutely unstable (as opposed to metastable), and the imaginary part of the potential gives the corresponding decay rate, which, unlike bubble nucleation, is not exponentially suppressed [93]. The imaginary part arises because the computation of the effective potential assumes the background field is constant, yet for the critical bubble this assumption fails [24, 75]. To bypass this complication, one throws away the imaginary part in some ad-hoc way, for example one of

$$V_{\rm LPA}(\phi) = V_3(\phi) + \begin{cases} \frac{-1}{12\pi} |V_3''(\phi)|^{3/2}, \\ \frac{-1}{12\pi} \operatorname{Re}\left(V_3''(\phi)^{3/2}\right). \end{cases}$$
(25)

Note that these different choices will yield different physical results. The former choice [94] is negative for $V_3''(\phi) < 0$, while the latter choice [95, 96] is zero. One then solves equation (7) using $V_{\text{LPA}}(\phi)$ as the potential, and inserts the resulting action into equation (23) to yield the LPA rate, Γ_{LPA} .

III. LATTICE DETAILS

Here we give further details of our lattice discretisation. We outline the purely spatial action used in the Monte-Carlo simulations, as well as details of our realtime update algorithm. Our simulation code is available at https://bitbucket.org/og113/scalnuc/. For the discretised action we use, Eq (10), possible discretisations of the lattice Laplacian include

$$(\nabla_{\text{lat}}^2 \phi)_x^{(1)} = \sum_i \frac{1}{a^2} \left(\phi_{x+i} + \phi_{x-i} - 2\phi_x \right), \qquad (26)$$

$$(\nabla_{\text{lat}}^2 \phi)_x^{(2)} = \sum_i \frac{1}{a^2} \left(-\frac{1}{12} \phi_{x+2i} + \frac{4}{3} \phi_{x+i} - \frac{5}{2} \phi_x + \frac{4}{3} \phi_{x-i} - \frac{1}{12} \phi_{x-2i} \right), \quad (27)$$

where *i* runs over the three directions of the lattice. These expressions have $O(a^2)$ and $O(a^4)$ errors accurate respectively. For smooth field configurations, such as one encounters in classical field theory in the absence of fluctuations, one can simply set the lattice parameters equal to their renormalised continuum counterparts. However, in the presence of thermal fluctuations the lattice action requires renormalisation, and the relationships between lattice and renormalised continuum coefficients are modified as $\kappa_{\text{lat}} = \kappa_{\overline{\text{MS}}} + \delta \kappa$.

The interaction terms of this 3d real scalar theory all have coefficients with positive mass dimension. As a consequence the way that physical quantities can depend on the lattice spacing a is tied to the way that they depend on the couplings. In the approach to the continuum limit, the leading a dependence is determined by dimensional analysis up to a small number of constants, which can be computed in lattice perturbation theory. Further, the lattice regularisation scheme can be related to other regularisation schemes, such as $\overline{\text{MS}}$, by matching perturbative computations of the vacuum energy [82, 83].

For the real scalar theory, shifting the field by a constant so that $g_{\text{lat}} = 0$ ensures that the coefficients of odd powers of the field are not renormalised, i.e. $\delta \sigma_3 = 0$ and $\delta g_3 = 0$. One can then extract the lattice-continuum relations from Refs. [51, 84, 85],

$$\delta m_3^2 = -\frac{\Sigma\lambda_3}{2(4\pi)a} + \frac{\lambda_3^2}{(4\pi)^2} \left[\log\frac{6}{a\mu_3} + C_3 - \Sigma\xi \right], \quad (28)$$

$$\delta\lambda_3 = \frac{3\xi\lambda_3^2a}{2(4\pi)} + \frac{\lambda_3^3a^2}{(4\pi)^3} \left(\frac{3}{4}\xi^2 - 3C_1 - \frac{C_2}{3}\right),\tag{29}$$

$$Z_{\phi} = 1 + \frac{C_2 \lambda_3^2 a^2}{6(4\pi)^2},\tag{30}$$

$$Z_m = 1 + \frac{\xi \lambda_3 a}{2(4\pi)} + \frac{\lambda_3^2 a^2}{(4\pi)^2} \left(\frac{\xi^2}{4} - \frac{C_1}{2} - \frac{C_2}{6}\right), \quad (31)$$

which all receive corrections at $O(a^3)$, except the squared mass parameter, which receives corrections already at O(a). In these equations Σ , ξ , C_1 , C_2 and C_3 are numerical constants which were computed in Ref. [51]. Their values depend on the scalar propagator, and hence on the choice of lattice Laplacian. For the $O(a^4)$ accurate Laplacian, which we have used for our final results, their values are

$$\Sigma = 2.75238391130752, \tag{32}$$

$$\xi = -0.083647053040968, \tag{33}$$

$$C_1 = 0.0550612, (34)$$

$$C_2 = 0.0334416, \tag{35}$$

$$C_3 = -0.86147916. \tag{36}$$

Note that if the $O(a^2)$ terms in Eqs. (28)-(31) are retained, one must use the higher order accurate Laplacian, Eq. (27) for the kinetic term to have the same accuracy.

Eqs. (28)-(31) give the relationship between bare and $\overline{\text{MS}}$ parameters up to cubic $O(a^3)$ corrections, except for the mass, which still has linear O(a) corrections. However, the residual lattice-spacing dependence of the mass squared is of the form $c_3\lambda_3^3a + c_4\lambda_3^4a^2$ with c_3 and c_4 pure dimensionless numbers. These corrections are independent of the mass. As a consequence, they cancel when considering differences such as $m_3^2 - m_{3,c}^2$, a trick which can be used to ensure faster convergence to the continuum limit for physical quantities.

The real time evolution is a mix of the Forest-Ruth algorithm [97] and momentum refresh (sometimes referred as partial momentum refreshment or partial momentum Monte Carlo) [51, 98, 99]. Forest-Ruth is a symplectic and fourth order accurate algorithm, which can be constructed starting from the second order accurate leapfrog aglorithm [100]. In our case, in order to make measurements at every timestep, we utilise the kick-drift-kick form of the leapfrog algorithm. For the field $\phi_{t,x}$ and momentum $\pi_{t,x}$ this latter update algorithm is

$$\pi_{t+\frac{1}{2},x} = \pi_{t,x} - \frac{\partial H_{\text{eff}}}{\partial \phi_{t,x}} \frac{\Delta t}{2}, \qquad (37)$$

$$\phi_{t+1,x} = \phi_{t,x} + a^3 \pi_{t+\frac{1}{2},x} \Delta t, \qquad (38)$$

$$\pi_{t+1,x} = \pi_{t+\frac{1}{2},x} - \frac{\partial H_{\text{eff}}}{\partial \phi_{t+1,x}} \frac{\Delta t}{2}.$$
(39)

Forest-Ruth then consists of three successive kick-driftkick updates, with timesteps, Δt_1 , Δt_2 and Δt_3 , given by

$$\Delta t_1 = (2 - 2^{1/3})^{-1} \Delta t, \qquad (40)$$

$$\Delta t_2 = -2^{1/3} (2 - 2^{1/3})^{-1} \Delta t, \qquad (41)$$

$$\Delta t_3 = \Delta t_1. \tag{42}$$

Being symplectic, for $\gamma = 0$ this algorithm conserves energy for long times, with errors of order $O(\Delta t/a)^5$. For our choice of timestep $\Delta t \lambda_3 = 0.01$, we have demonstrated energy convergence at the level of $O(10^{-6})$. We have also demonstrated that for $\gamma = 0$ the algorithm is reversible to numerical precision.

In order to reproduce the damping γ and Gaussian random noise ξ of the Langevin Eqs. (3) and (4), we use the momentum refresh. In our case the update reads

$$\pi_{t+0,x} = \sqrt{1 - \vartheta^2} \pi_{t-0,x} + \vartheta \xi_{t,x},$$
 (43)

$$\vartheta = 1 - \exp(-2a^3\gamma\Delta t). \tag{44}$$

The momentum refresh is applied after each Forest-Ruth evolution step, together completing one full time evolution iteration.

This complete algorithm conserves the thermal distribution of the momenta with errors of order $O(\Delta t/a)^4$ [51]. For our parameter choices, we found experimentally that this level of accuracy was crucial for the real-time and Monte-Carlo parts of the code to agree sufficiently precisely on the position of the separatrix.