# Cool matter distribution in inner solar corona from 2023 total solar eclipse observation

Z.Q. Qu<sup>1,2</sup>, H. Su<sup>3</sup>, Y. Liang<sup>4</sup>, Z.Xu<sup>1</sup>, R.Y. Zhou<sup>5</sup>

1. Yunnan Astronomical Observatory, CAS, Tianwentai road, Guandu district, Kunming, Yunnan 650216, China

2. School of Astronomy and Space Sciences, University of Chinese Academy of Sciences, Yanxihu, Huairou District, Beijing, China

3. Yunnan Amateur Astronomers Association, No.2506 Jida Square, Panlong district, Kunming, Yunnan, China

4. Shanghai Astronomical Observatory, CAS, Xuhui District, Shanghai, China

5. Department of Physics, Chinese University of Hong Kong, Shatin, New Territories, Hong Kong SAR, China

### ABSTRACT

Solar corona has been judged to consist of free electrons and highly ionized ions with extremely high temperature as a widely accepted knowledge. This view is changed by our eclipse observations. Distributions of cool matter represented by neutral iron atoms in hot inner solar corona are presented via derived global maps of solar Fraunhofer(F-) and Emission(E-) coronae, compared with those of continuum(Kontinuierlich, K-) corona formed by free electrons. The maps are obtained from simultaneous observations of dual filtering bands centered respectively at 659.4nm and 660.1nm, performed from twin telescopes during the total solar eclipse on April 20, 2023 at Com town of East Timor, assisted for judgement via spectral images obtained by a portable spectrograph. They show respectively presences of these neutral iron atoms yielding 659.3nm and 659.4nm lines in both the quiet sun and active regions. The distribution of the cool matter in form of line depression forms an inner F-corona, different from that of the cool matter in form of line enhancement. Both the distributions show a crucial difference from that of the free electrons represented by the Kcorona map. It is also found that intensities of the F-corona and the E-corona induced by these neutral atoms are only small fractions of the K-corona, and the diffusion can be seen clearly in all these maps. They uncover also that the coronal heating resources do not distribute pervasively but likely form a thermodynamic griddle where minor photospheric neutral atoms can escape from the heating into the corona globally.

Key Words Sun:corona; Sun: Fraunhofer corona

#### 1. Introduction

Solar corona has been being observed during total solar eclipses since ancient times with human naked eves. Giovanni Cassini coined the name after his observation of 1706 solar eclipse. However, it had been naturally regarded to be cooler than its underneath atmospheric layers such as solar photosphere with an average temperature of several thousand degrees since it is farther away from the solar core, the only heat resource recognized then. Temperature of the corona was first reliably diagnosed and found to be hotter than one million Kelvin since Hardness and Young recorded the green coronal line in the corona during 1869 solar eclipse, and Grotrian (Grotrian, 1939) in 1939 and later Edlen(Edlen, 1943) in 1943 realized that this spectral line is caused by transition from upper energy levels in thirteen times ionized iron ions, whose metastable presence necessary for yielding observable radiation needs an environment of more than one million degrees in the atmosphere. Since then, the opposite view has been accepted that the quiet corona unconfined by the magnetic fields or in open field line region, was regarded to be consisted of completely highly ionized ions and free electrons. This forms the basis of the solar coronal physics(Aschwanden,2015). On the other hand, beyond a height of three solar radii above solar limb, dust was found scattering the Fraunhofer lines originally yielded in the photosphere(Morgan et al., 2007; Stenborg et al., 2021). It is called F-corona. And according to space observations, the dust is distributed in an elliptic space around the sun (Burtovoi et al., 2022; Lamy et al., 2022). However, conscious detection for presence of a cool component has not yet been explored in the inner solar corona, where the dust should be sublimated (Anderson, 1924; Russell, 1929), probably due to its potential confliction with the aforementioned view.

In astrophysics, temperature of a mass of electromagnetically structured particles can be diagnosed from the specific spectral lines that are produced with a favorite temperature, causing the maximum occupation number of the involved energy levels of the transition, and thus giving the birth to the most strong line intensity. In detail, this kind of 'formation' temperature can be roughly estimated from the excitation potentials of these levels excited from the ground ones. For instance, for our used neutral iron line at 659.3nm and 659.4nm, the excitation energies of their lower levels are respectively 2.73eV and 2.43eV, and those of their upper levels are respectively 4.61eV and 4.31eV. As a contrast, the excitation and ionization energy needed for producing the green coronal line FeXIV530.3nm is greater than 394.6eV(from NIST, https : //physics.nist.gov /PhysRefData/ASD/lines \_form.html). Therefore, the environment in which the neutral atoms yielding these two lines can survive is two order colder than that producing the green coronal line, provided that the excitation and ionization energy *E* is supplied by the collisions of thermal random motions. That is, E = 3/2kT may be approximately applied, where *T* means the temperature and *k* the Boltzmann constant. Accordingly, the specific spectral lines can be temperature proxies of mass of these atoms or ions.

It was noteworthy that our spectropolarimetry of upper solar atmosphere during 2013 Gabon eclipse led to the finding of the neutral atoms in the inner solar corona(Qu et al., 2024). As pointed out previously, one way to identify the existence of the cool matter represented by the neutral atoms is to detect the presence of the specific spectral lines yielded by these atoms and then to exclude the possibility that the presence is caused by the dust scattering due to projective effect along the lineof-sight. The first step can be taken by performance of simultaneous observations of two bands with the same-width containing respectively the suitable lines and their adjacent continua, using two same telescopes and detectors. It is basically understood that if absolute value of difference between the intensities of the two bands is well above the detection noise, a definite evidence for the presence of the relevant atoms producing the specific spectral lines is implied. Furthermore, if the intensity difference is definitely positively greater than the noise level, it indicates that there are spectral lines detected above the continuum by the neutral atoms producing the emission lines via transitions from the upper energy levels to the lower ones and/or scattering these emission lines. And if the difference is negative and its absolute value well above the noise level, it can be judged that there are the specific lines whose intensities are depressed due to the neutral atoms yielding the absorption via transitions from the lower energy levels to the upper ones, and/or scattering the Fraunhofer lines predominantly formed in the photosphere. The aforementioned criterion is the reasoning base in the following.

## 2. Observations and Results

It becomes well known that total solar eclipses provide us uniquely precious chances to get the accurate measurement of the quiet solar corona on the ground with minimum light pollution from the telluric atmosphere and instrument during the totality(Pasachoff, 2017). We performed the observation during a total solar eclipse occurred on April 20, 2023, at Com town, East Timor. The geographical coordinates of the site read as E127°3'42" S8°21'38". The totality last about 76 seconds from 04:21:34UT to 04:22:50UT. Our observational goal is to survey the simultaneously available the F-corona described by the selected Fraunhofer line intensity depression and the E-corona formed by the same specific line enhancement in the inner corona, compared with the K-corona formed by the continuum intensity source distribution. The Fraunhofer lines are selected, as stated above, to be two neutral iron lines, FeI 659.29nm  $(3d^7(^2G)4s \rightarrow 3d^7(^4F)4p)$  and 659.38nm  $(3d^64s^2 \rightarrow 3d^7({}^4F)4p)$ . They are contained in a 659.4nm-centered filtering band with 0.3nm bandpass. The images are obtained from the twin apochromatic refractive telescopes named APO80. Both the telescopes have the same primary optical parameters (Aperture 80mm and focal ratio f/6) except two different filters mounted respectively in front of two same electronic detectors Sony IMX174. Thus, one telescope provides via a filter(type: Alluxa 7337,659.4-0.3nm) images of the band 659.25-659.55nm containing the two lines mixed with continuum, and the other genders

images of the continuum via a filter(type: Aulluxa 7340, 660.1-0.3nm) with the same bandpass width 0.3nm but centered at 660.1nm. These twin telescopes are installed symmetrically on one dovetail plate attached to an equatorial mount. Finally, in order to check whether additional lines are present within the two band thus possibly distort the intensity difference between the two bands, a portable spectrograph named Sol'Ex with a guiding optics of 8cm aperture refractive telescope was put into actual observation, whose observational band covers the two filtering bands as well as the famous H $\alpha$  line.

Before we carried out the observation, an integrating sphere with a stable light in our laboratory had been used for intensity calibration between the two telescopes and test for readout linearity of the detector. Further intensity calibration between the two channels was performed in a way that their intensity difference should be minimized in the dark disk formed by the lunar occultation during the totality after the data reduction of dark current and flat-fielding. And since the two kinds of images were recorded at the different parts of the detector chips and influenced by seeing-induced image movement, the location correlation registration is carried out before the further data reduction operation. Finally, the intensity differences at those points are set to be zero whose intensities obtained from both the two bands are saturated.

The top two panels in Fig.1 show the sample images of the two bands obtained respectively from the twin telescopes with the same cadence and exposure times of 0.1 second for each frame during the total solar eclipse. Both of these images are respectively superpositions of thirteen raw frames after the data reduction. The left one is the filtering image within the band centered at 659.4nm containing the two lines, with intensity indicated by  $I_{c,l}$ . And the K-corona(continuum intensity) image is plotted in the right panel acquired within the band centered at 660.1nm, with its intensity indicated with  $I_c$ . It is intuitively very hard to distinguish one image from the other. However, quantitative comparison will reveal their delicate difference and this is just our observational goal. It is notable that the regions with strong radiation are regarded to be active ones.

The middle panel below the raw images is the map of the intensity difference  $I_{c,l} - I_c$  between the intensity of 659.4nm-centered band  $I_{c,l}$  and that of the 660.1nm-centered band  $I_c$ . All the intensities are in unit of the detector readout counts(cts). According to a simple calculation, the absolute value of the normalized difference  $(I_{c,l} - I_c)/I_c$  is found to be less than 0.3% in all these spatial points with intensities well above the noise level. This means that the neutral iron atoms cannot be easily observed at these two lines. As a contrast, we have found that more than 20% line depression in many Fraunhofer lines including those of the neutral iron, magnesium and chromium atoms in the band 516.3nm-532.6nm was distorted in a very small coronal region(Qu et al., 2024). And more popular is the several thousandth line depression. Therefore, even the same kind of neutral atoms but yielding different spectral lines can be distributed differently in solar corona. This reflects different temperature distributions. The weak correlation between the derived map and the



Fig. 1.— Eclipse images. Top panels from left to right: superposed maps obtained respectively from 659.4nmcentered filtering observation and from 660.1nm-centered observation, performed via the twin telescopes during the total solar eclipse on 20 Apr 2023. The 660.1nm-centered filtering image(on the top right) represents the K-corona at this band. It is very hard to distinguish these two images from each other on the top panels, since their difference is very small. Middle panel: the map of intensity difference between the two bands. Bottom panels: the map of the line depression(left) and the map of the line enhancement(right). They are respectively the maps of the inner F-corona and E-corona.

K-corona one on the top right means that the distribution of the neutral iron atoms is very different from that of the free electrons. On the whole, the line depression seems overwhelming, while the line enhancement is more narrowly distributed. This reflects again the different temperature distributions.

It is noteworthy that there was no detectable exotic lines found in the two bands, as witnessed in Fig.2, where the spectra obtained from the Sol'Ex portable spectrograph are presented. The spectrum acquired before the second contact shows that the H $\alpha$  absorption line is the strongest and broadest in the band, while the pair of neutral iron lines FeI659.4nm and FeI659.3nm are relatively strong Fraunhofer lines compared to the other lines, and no spectral lines can be detected within the band centered at 660.1nm with width 0.3nm. In the bottom panel is shown the spectrum acquired during totality. Now, the H $\alpha$  line intensity becomes prominent and even saturate in those points close to the limb(the dark part boarder), but the pair of the neutral iron lines now become so weak that the 659.3nm line can be seen and the 659.4nm line can be distinguished only by very carefully view. It is turned to be even much weaker than other medium strong lines. This specifies that the lines are not produced by the dust scattering, or the relative intensities among these Fraunhofer lines should be kept. On the other hand, there are no lines appeared within 0.3nm around 660.1nm. All of these promise that the difference between intensities of these two bands respectively centered at 659.4nm and 660.1nm is not caused by other lines emerged during the totality.

The middle panel map is composed of the line depression(dark) and the line enhancement(bright) simultaneously. It contains also those noisy points with their intensity differences not well above the noise level. In order to separate the line depression and the line enhancement thus to get the F-corona and E-corona maps, we execute an operation to divide the difference map into two. The F-corona map depicted on the bottom left panel is obtained to collect those points with their  $(I_c - I_{c,l}) > 3\sigma$ , and  $\sigma$  indicates the standard deviation evaluated at the solar disk occulted by the moon during the totality, while the E-corona map shown in the bottom right panel is attained if requirement  $(I_{c,l} - I_c) > 3\sigma$  is met.

It is impressive that the line depression can be strong in both the quiet sun and active regions. Further, in the quiet sun region around the south pole, the line depression is spread much more widely than that in the domain around the prominence close to the north pole, though the line depression can be found in the central thin axis vertical to the local limb. In fact, the line depression is much more concentrated in the active regions of the west hemisphere, while the line enhancement locates more densely in the hemisphere divided by a direction from the northwest to the southeast. The diffusion motions of these neutral atoms can be seen above both the active and quiet sun regions, indicated by the rapid decrease of the line depression and enhancement with height above the limb. Meanwhile, in the active region, the magnetic constraint can indirectly influence the diffusion via collision of the neutral atoms with these confined free electrons and ions. However, the line depression can neither be viewed in the main body around the line depression axis of the prominence, nor in the jet occurred close to the east limb, and hardly observed in other domains dominated by the line enhancement. Therefore, both of them are not homogeneously spread around the solar disk occulted by the moon. Finally, it seems that the line enhancement can detectable in higher regions, as witnessed from the space above the prominence.

The non-similarity between the line depression and enhancement maps means the different formation temperature distribution of the neutral iron atoms due to different excitation potentials of the lower and upper energy levels involved in the transition. Furthermore, it is easy to distinguish these two kinds of maps from that of the K-corona depicted in the top right panel of Fig.1. It is witnessed that though the K-coronal radiative intensity is much stronger, it experiences much steeper decrease with the heliocentric height than the E-corona or the F-corona.

It becomes clear that both the two kinds of distributions are neither homogeneous nor symmetric. The F-corona map reflects the distribution of the Fraunhofer line intensity depression, proportional to net amount of photons taken away from the continuum by particles staying at the lower levels of the involved transitions, via scattering along the line-of-sight (Zanstra, 1941) and/or absorption overwhelming emission. The E-corona map, on the other hand, presents the line enhancement, proportional to net amount of photons created by the atoms stayed at the upper energy levels of the transitions above the continuum, via scattering and/or emission overwhelming absorption. The K-corona map depicted on the top right panel of Fig.1 provides quantitatively the distribution of the free electrons predominantly responsible for the continuum radiation. Therefore, till now, we can judge the presence of atoms or ions responsible for the production of the specific spectral lines within the inner corona concerned here. It is noteworthy that the F-corona map induced by the neutral iron atoms presented here looks very different from those caused from scattering by the dust which exists above 3.0 solar radius height, distributed in an elliptic space around the sun (Lamy et al., 2022; Burtovoi, 2022). Furthermore, the line depression in these upper solar atmospheric layers is evidently connected to those in the photosphere below, because no line depression faultage is found in the chromosphere or the transition zone.

## 3. Conclusion

After we obtain the F-corona and E-corona maps based on the line depression and enhancement of the FeI659.3nm and 659.4nm lines, it is judged that the neutral iron atoms are widely but sparsely distributed in the inner solar corona as a cool component, with help of the spectral images obtained by the portable Sol'Ex spectrograph. The F-corona induced from the neutral iron atoms are very different from that F-corona caused by dust scattering the photospheric Fraunhofer lines, in aspects of components, distribution patterns and heliocentric height ranges. Therefore,





Fig. 2.— Spectrum samples acquired from the portable Sol'Ex spectrograph. Top panel: the spectrum acquired before the second contact. The H $\alpha$  absorption line is the strongest and broadest line in the band, while the pair of neutral iron lines FeI659.4nm and FeI659.3nm are strong Fraunhofer lines among the other lines, while no spectral lines can be seen within the band centered at 660.1nm with width 0.3nm. Bottom panel: the spectrum acquired during totality. The H $\alpha$  line becomes dominant emission one and saturated in spatial points close to the limb, but the pair of the neutral iron lines become so weak that the 659.3nm line can be hardly seen and the 659.4nm line can be visible by careful checking in the lower part and turned to be much weaker than other medium strong lines. This specifies the lines are not produced by the dust scattering. On the other hand, there are no lines appeared around 660.1nm within 0.3nm. This promises that the difference between intensities of these two subbands respectively centered at 659.4nm and 660.1nm is not caused by other lines emerged during the totality.

the new F-corona forms the inner F-corona. The F-corona and E-corona maps differ greatly from each other, and further both of them are distinguished from K-corona critically. This reflects the different distributions of particles with different temperatures.

The finding of the neutral atoms additional to the highly ionized ions in the inner corona reveals a new nature of the solar corona. The more micro atomic energy states of iron found leads to a conclusion that the entropy estimated in the inner corona would be much greater than that in the photosphere. Therefore, the coronal heating can be understood according to the second law of the thermodynamics. On the other hand, due to the presence of the neutral atoms and thus ion-atom collision, the Cowling dissipation will play a significant role in coronal heating(Cowling, 1957), since the dissipation rate will become larger by many orders than the rate of dissipation caused by electron-ion collisions. It also provides a constraint on the coronal heating mechanism. That is, the heating is not pervasive since a small fraction of neutral iron atoms can escape from the heating into the corona, companying upflows of other heated particles(Tian et al., 2021), through the chromosphere and transition zone from the photosphere, where the Fraunhofer lines are predominantly formed. Finally, the cool matter injected into magnetic reconnection region can evidently promote the energy release efficiencies(Ni et al., 2015; Liu et al., 2023;). Therefore, the simulations of multi-fluid formed by the free electrons, ions and neutral atoms sound more reasonable(Leake et al., 2012) than the two-fluid model.

This work is sponsored by National Science Foundation of China (NSFC) under the grant numbers 11078005, U1931206 and 12003066.

### REFERENCES

- Aschwanden, M.J. 2015, Physics of the solar corona. Springer, Published in association with Praxis Publishing Chichester, UK, ISBN: 3-540-22321-5
- Anderson, W., 1924, Z Phys. 28,299
- Burtovoi, A. Naletto, G., Dolei, S., Spadaro, D., Romoli, M., F. Landini, F., and De Leo, Y. ,2022, *Astron. & Astrophys.* **659**, A50
- Cowling, T.G., Magnetohydrodynamics, Interscience Publishers, New York, 1957
- Edlen, Bengt, 1943, Zeitschrift fur Astrophysik, 22, 30
- Grotrian, Walter, 1939, Naturwissenschaften, 27, 214
- Lamy, P.L., Gilardy, H., and LIebaria, A., 2022, Space Sci. Rev., 218, 53
- Leake, J.E., Lukin, V.S., Linton, M.G., and Meier, E.T., 2012, ApJ, 760:109
- Liu Mingyu, Ni Lei, Cheng Guan-chong, et al., 2023, Research in Astronomy and Astrophysics, 23: 035006
- Morgan, H., and Habbal, S.R., 2007, A&A, 471:L47
- Ni Lei, Kliem, B., Lin Jun and Wu Ning, 2015, ApJ, 799:79
- Pasachoff, J.M., 2017, Nature Astronomy, 1, 0190
- Qu, Z.Q., Chang, L., Dun, G.T., et al., 2024, Spectropolarimetry of Fraunhofer lines in local upper solar atmosphere, arXiv:2403.16570vl, also submitted to *Astrophys.J*
- Russell, H.N., 1929, Astrophys.J., 69,49
- Stenborg, G., Howard, R.A., Hess, P., Gallagher, B., 2021, 650:28
- Tian, Hui, and Harra, Louise, 2021, Solar Phys. 296, 47
- Zanstra, H., 1941, MNRAS, 101, 250

This preprint was prepared with the AAS LATEX macros v5.2.