

Intersection Theorem for DG-modules

Xiaoyan Yang

School of Science, Zhejiang University of Science and Technology, Hangzhou 310023, China

E-mail: yangxy@zust.edu.cn

Abstract

Let A be a commutative noetherian local DG-ring with bounded cohomology. The Intersection Theorem for DG-modules is examined and some of its applications are provided. The first is to prove the DG-setting of the amplitude inequality, New Intersection Theorem and Krull's principle ideal theorem. The second is to solve completely the Minamoto's conjecture in [Israel J. Math. 242 (2021) 1–36]. The third is to show the DG-version of the Bass conjecture about Cohen-Macaulay rings and the Vasconcelos conjecture about Gorenstein rings.

Keywords: Intersection Theorem; cohomological dimension; depth

2020 Mathematics Subject Classification: 13D22, 13C15, 16E45

1. Introduction

The New Intersection Theorem (NIT) of Peskine and Szpiro [14], Hochster [8] and Roberts [16] is a central result in the homological theory of commutative noetherian rings, which has a variety of applications (see [6, 17]), and was extended and polished by commutative algebra practitioners (see [1, 5, 7, 9, 10, 15]).

Let A be a commutative noetherian local ring and M, N two finitely generated A -modules with $\text{projdim}_A M < \infty$. The NIT yields an inequality, i.e., Intersection Theorem (IT)

$$(\dagger) \quad \dim_A N \leq \dim_A(M \otimes_A N) + \text{projdim}_A M$$

By applying the inequality (\dagger) with $N = A$, one has $\dim A \leq \dim_A M + \text{projdim}_A M$. By the Auslander-Buchsbaum Formula, the inequality is equivalent to $\text{cmd} A \leq \text{cmd}_A M$, where $\text{cmd}_A M = \dim_A M - \text{depth}_A M$ is the Cohen-Macaulay defect of M which determines the failure of M to be Cohen-Macaulay. Further, Yassemi [27] showed the inequalities

$$(\ddagger) \quad \dim_A N \leq \dim_A \text{RHom}_A(M, N) \leq \dim_A(M \otimes_A N) + \text{projdim}_A M$$

By [4, Proposition 5.2.6], $\text{depth}_A \text{RHom}_A(M, A) = \text{depth} A$, it follows by (\ddagger) that $\text{cmd} A \leq \text{cmd}_A \text{RHom}_A(M, A) \leq \text{cmd}_A M$. Also $\text{grade}_A M + \dim_A M \leq \dim A$, so if M is perfect then $\text{cmd} A = \text{cmd}_A \text{RHom}_A(M, A) = \text{cmd}_A M$.

On the other hand, Foxby [6] generalized the NIT for complexes, for bounded complexes X, Y with finite cohomologies, showed the IT of complexes

$$\dim_A Y \leq \dim_A(X \otimes_A^L Y) + \text{projdim}_A X$$

whenever $\text{projdim}_A X < \infty$. In 2005, Dibaei and Yassemi [5] continued to study the IT of complexes and proved the following inequalities

$$\dim_A Y - \sup X \leq \dim_A \text{RHom}_A(X, Y) \leq \dim_A(X \otimes_A^L Y) + \text{projdim}_A X - \inf X,$$

which generalizes the above inequalities (‡).

An *associative DG-ring* A is a \mathbb{Z} -graded ring $A = \bigoplus_{i \in \mathbb{Z}} A^i$ with a differential of degree 1, that satisfies the graded Leibniz rule. A DG-ring A is called *commutative* if $b \cdot a = (-1)^{i \cdot j} a \cdot b$ for all $a \in A^i$ and $b \in A^j$, and $a^2 = 0$ if i is odd. A DG-ring A is called *non-positive* if $A^i = 0$ for all $i > 0$. A non-positive DG-ring A is called *noetherian* if the ring $H^0(A)$ is noetherian and the $H^0(A)$ -module $H^i(A)$ is finitely generated for all $i < 0$. If A is a noetherian DG-ring and $(H^0(A), \bar{\mathfrak{m}}, \bar{\kappa})$ is a local ring, then we say that $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ is *local noetherian*. **Unless stated to the contrary we assume throughout this paper that A is a commutative and noetherian DG-ring.**

The derived category of DG-modules over A will be denoted by $D(A)$, and write $- \otimes_A^L -$ for its monoidal product, $\text{RHom}_A(-, -)$ for the internal hom. The full subcategory which consists of DG-modules whose cohomology is above (resp. below, bounded) is denoted by $D^-(A)$ (resp. $D^+(A)$, $D^b(A)$), denote by $D_f(A)$ the full subcategory of $D(A)$ consisting of DG-modules with finite cohomology, and set $D_f^-(A) = D^-(A) \cap D_f(A)$, $D_f^+(A) = D^+(A) \cap D_f(A)$, $D_f^b(A) = D^b(A) \cap D_f(A)$. For a DG-module X , we set $\inf X := \inf\{n \in \mathbb{Z} | H^n(X) \neq 0\}$, $\sup X := \sup\{n \in \mathbb{Z} | H^n(X) \neq 0\}$ and $\text{amp} X := \sup X - \inf X$.

DG-rings allow us to use techniques of homological algebra of ordinary rings in a much wider context. Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring. Shaul [21] extended the definitions of Krull dimensions and depth of complexes to DG-setting. The *local cohomology Krull dimension* of a DG-module $X \in D^-(A)$ is the number

$$\text{lc.dim}_A X := \sup\{\dim_{H^0(A)} H^\ell(X) + \ell | \ell \in \mathbb{Z}\},$$

where $\dim_{H^0(A)} H^\ell(X)$ is the usual Krull dimension of the $H^0(A)$ -module $H^\ell(X)$. The *depth* of a DG-module $Y \in D^+(A)$ is the number

$$\text{depth}_A Y := \inf \text{RHom}_A(\bar{\kappa}, Y).$$

The theory of classical Cohen-Macaulay rings, Cohen-Macaulay modules and Gorenstein rings has also been extended in [21, 22]. This paper focuses on local cohomological Krull dimension of the DG-modules $\text{RHom}_A(X, Y)$ and $X \otimes_A^L Y$ for appropriate X and Y in $D(A)$. We wish to extend the above inequalities to local DG-rings and obtains some characterizations about local Cohen-Macaulay and local Gorenstein DG-rings.

The paper is organized as follows.

In Section 2, we gather some preliminaries and results about local cohomological Krull dimension of DG-modules, and generalize the DG-version of Grothendieck's local duality theorem in [19, Theorem 7.26] to a more general case. In Section 3, the DG-setting of IT is proved when A is a local DG-ring with bounded cohomology (see Theorem 3.1), a bit more

generally, for $0 \not\cong Y \in D_f^-(A)$ and $X \in D^-(A)$ with $\text{flatdim}_A X < \infty$ and $\text{depth}_A X < \infty$, we show the following inequality:

$$(\dagger') \quad \text{lc.dim}_A Y \leq \text{lc.dim}_A(X \otimes_A^L Y) + \text{flatdim}_A X$$

Moreover, some upper and lower bounds of $\text{lc.dim}_A(X \otimes_A^L Y)$ and $\text{lc.dim}_A \text{RHom}_A(X, Y)$ for some DG-modules X and Y are provided (see Theorem 3.5). When the DG-ring A is sequence-regular and $X, Y \in D_f^b(A)$, we improve the inequality (\dagger') in Proposition 3.7, prove the following stronger inequalities:

$$\text{lc.dim}_A X + \text{lc.dim}_A Y \leq \text{lc.dim}_A A + \text{lc.dim}_A(X \otimes_A^L Y).$$

In view of the inequalities (\dagger') , we extend in Section 4 some classical results about amplitude inequality, Cohen-Macaulay rings and Gorenstein rings to local DG-rings with bounded cohomology. For $0 \not\cong X, Y \in D_f^b(A)$ with $\text{projdim}_A X < \infty$, Jørgensen [11, Theorem 3.1] showed the amplitude inequality $\text{amp}(X \otimes_A^L Y) \geq \text{amp} X$ when A is a sufficiently nice local DG-ring with $\text{amp} A < \infty$. The first application of our results is to eliminate the hypothesis on A . We improve the Jørgensen's result (see Proposition 4.1), and prove the DG-version of NIT and Krull's principle ideal theorem (see Proposition 4.4). Bass [2] raised the question whether existence of a finitely generated module of finite injective dimension would imply Cohen-Macaulayness of the ring, and later known as the Bass conjecture. For $0 \not\cong X \in D_f^b(A)$ with $\text{injdim}_A X < \infty$, Minamoto [12, Conjecture 2.36] conjectured that $\text{amp} X \geq \text{amp} A$. Shaul [21, Theorem 5.22(2)] showed the DG-version of Bass Conjecture and solved this conjecture under the noetherian model assumption, and conjectured in [21, Remark 5.25] that this assumption is redundant. As the second application, Minamoto's Conjecture or Shaul's Conjecture is solved (see Proposition 4.6). Vasconcelos [24] conjectured that rings of type one are Gorenstein. The last application is to prove the DG-version of the Vasconcelos conjecture (see Proposition 4.8).

2. Preliminaries and some facts

This section is devoted to recalling some notions and basic facts.

Projective, injective and flat dimensions. For a DG-module $X \in D(A)$, we recall the definition of the projective, injective and flat dimensions of X introduced by Bird, Shaul, Sridhar and Williamson in [3], which are different from the definition, $\text{pd} X$, $\text{id} X$ and $\text{fd} X$, introduced by Yekutieli [28].

The *projective dimension* of X is defined by

$$\text{projdim}_A X = \inf\{n \in \mathbb{Z} \mid H^i(\text{RHom}_A(X, Y)) = 0 \text{ for any } Y \in D^b(A) \text{ and } i > n + \sup Y\}.$$

The *injective dimension* of X is defined by

$$\text{injdim}_A X = \inf\{n \in \mathbb{Z} \mid H^i(\text{RHom}_A(Y, X)) = 0 \text{ for any } Y \in D^b(A) \text{ and } i > n - \inf Y\}.$$

The *flat dimension* of X is defined by

$$\text{flatdim}_A X = \inf\{n \in \mathbb{Z} \mid H^{-i}(Y \otimes_A^L X) = 0 \text{ for any } Y \in D^b(A) \text{ and } i > n - \inf Y\}.$$

For $X \in D^-(A)$ and $Y \in D^+(A)$, it follows by [13, Theorems 2.22, 3.21, 4.13] that

$$\begin{aligned} \text{projdim}_A X &= \text{pd}X - \sup X, \\ \text{injdim}_A Y &= \text{id}Y + \inf Y, \\ \text{flatdim}_A X &= \text{fd}X - \sup X. \end{aligned}$$

Lemma 2.1. *Let A be a DG-ring with $\text{amp}A < \infty$, $X \in D^-(A)$ and $Y \in D^+(A)$.*

- (1) *If $\text{projdim}_A X < \infty$, then $\inf X \geq \inf A - \text{projdim}_A X$, i.e. $X \in D^b(A)$.*
- (2) *If $\text{injdim}_A Y < \infty$, then $\sup Y \leq -\inf A + \text{injdim}_A Y$, i.e. $Y \in D^b(A)$.*
- (3) *If $\text{flatdim}_A X < \infty$, then $\inf X \geq \inf A - \text{flatdim}_A X$, i.e. $X \in D^b(A)$.*

Proof. We just prove (1) since (2) follows by duality.

(1) We use induction on $n = \text{projdim}_A X + \sup X$. If $n = 0$, then $X \in \mathcal{P}[-\sup X]$ by [13, Lemma 2.14] and so $\inf X \geq \inf A - \text{projdim}_A X$. Assume that $n \geq 1$. By [13, Lemma 2.13], there is an exact triangle $Z \rightarrow P \rightarrow X \rightsquigarrow$ so that $P \in \mathcal{P}[-\sup X]$ and $\text{projdim}_A Z + \sup Z \leq \text{projdim}_A Z + \sup X = n - 1$, it follows by induction that $\inf X \geq \inf A - \text{projdim}_A X$.

(3) For $H^0(A)$ -module $\text{Hom}_{\mathbb{Z}}(H^0(A), \mathbb{Q}/\mathbb{Z})$, it follows by [19, Theorem 5.7] that there is a DG-module E so that $H^0(E) = \text{Hom}_{\mathbb{Z}}(H^0(A), \mathbb{Q}/\mathbb{Z})$. Hence [19, Theorem 4.10] and (2) imply that $\inf X = -\sup \text{RHom}_A(X, E) \geq \inf A - \text{injdim}_A \text{RHom}_A(X, E) = \inf A - \text{flatdim}_A X$. \square

Lemma 2.2. ([3]). *Let A be a DG-ring with $\text{amp}A < \infty$ and $X, Y, Z \in D(A)$.*

(1) *If either $X \in D_f^-(A)$, $Y \in D^+(A)$ and $\text{injdim}_A Z < \infty$, or $X \in D_f^b(A)$ and $\text{projdim}_A X < \infty$, then the natural map*

$$X \otimes_A^L \text{RHom}_A(Y, Z) \rightarrow \text{RHom}_A(\text{RHom}_A(X, Y), Z)$$

is an isomorphism in $D(A)$.

(2) *If either $X \in D_f^-(A)$, $Y \in D^+(A)$ and $\text{flatdim}_A Z < \infty$, or $X \in D_f^b(A)$ and $\text{projdim}_A X < \infty$, then the natural map*

$$\text{RHom}_A(X, Y) \otimes_A^L Z \rightarrow \text{RHom}_A(X, Y \otimes_A^L Z)$$

is an isomorphism in $D(A)$.

Derived torsion and completion. Let $\bar{\mathfrak{a}} \subseteq H^0(A)$ be an ideal. Recall that an $H^0(A)$ -module \bar{X} is called $\bar{\mathfrak{a}}$ -torsion if for any $\bar{x} \in \bar{X}$ there exists $n \in \mathbb{N}$ such that $\bar{\mathfrak{a}}^n \bar{x} = 0$. The category $D_{\bar{\mathfrak{a}}\text{-tor}}(A)$ consisting of DG-modules X such that the $H^0(A)$ -module $H^n(X)$ is $\bar{\mathfrak{a}}$ -torsion for every n , is a triangulated subcategory of $D(A)$. Following [20], the inclusion functor $D_{\bar{\mathfrak{a}}\text{-tor}}(A) \hookrightarrow D(A)$ has a right adjoint $D(A) \rightarrow D_{\bar{\mathfrak{a}}\text{-tor}}(A)$, and composing this right adjoint with the inclusion, one obtains a triangulated functor

$$\text{R}\Gamma_{\bar{\mathfrak{a}}} : D(A) \rightarrow D(A),$$

which is called the *derived $\bar{\mathfrak{a}}$ -torsion functor*. The *derived $\bar{\mathfrak{a}}$ -completion functor* $\mathrm{L}\Lambda_{\bar{\mathfrak{a}}} : \mathrm{D}(A) \rightarrow \mathrm{D}(A)$ is defined by $\mathrm{L}\Lambda_{\bar{\mathfrak{a}}}(X) := \mathrm{RHom}_A(\mathrm{R}\Gamma_{\bar{\mathfrak{a}}}(A), X)$, which is left adjoint to $\mathrm{R}\Gamma_{\bar{\mathfrak{a}}}$ as endofunctors on $\mathrm{D}(A)$. The derived $\bar{\mathfrak{a}}$ -adic completion $\mathrm{L}\Lambda_{\bar{\mathfrak{a}}}(A)$, denoted by $\mathrm{L}\Lambda(A, \bar{\mathfrak{a}})$, is itself a commutative noetherian DG-ring.

Given a commutative ring A and $a \in A$, the telescope complex associated to A and a is

$$0 \rightarrow \bigoplus_{n=0}^{\infty} A \rightarrow \bigoplus_{n=0}^{\infty} A \rightarrow 0$$

in degrees 0, 1, with the differential being defined by

$$d(e_i) = \begin{cases} e_0 & i = 0, \\ e_{i-1} - ae_i & i \geq 1, \end{cases}$$

where e_0, e_1, \dots denote the standard basis of the countably generated free A -module $\bigoplus_{n=0}^{\infty} A$. We denote this complex by $\mathrm{Tel}(A; a)$. Given a sequence $\mathbf{a} = a_1, \dots, a_n \in A$, we set

$$\mathrm{Tel}(A; \mathbf{a}) := \mathrm{Tel}(A; a_1) \otimes_A \cdots \otimes_A \mathrm{Tel}(A; a_n),$$

which is a bounded complex of free A -modules, called the telescope complex associated to A and \mathbf{a} .

For an ideal $\bar{\mathfrak{a}} \subseteq \mathrm{H}^0(A)$, if $\mathbf{a} = (a_1, \dots, a_n)$ is a sequence in A^0 , whose image in $\mathrm{H}^0(A)$ generates $\bar{\mathfrak{a}}$, then for $X \in \mathrm{D}(A)$, there are natural isomorphisms

$$\mathrm{R}\Gamma_{\bar{\mathfrak{a}}}(X) \cong \mathrm{Tel}(A^0; \mathbf{a}) \otimes_{A^0} A \otimes_A X, \quad \mathrm{L}\Lambda_{\bar{\mathfrak{a}}}(X) \cong \mathrm{Hom}_A(\mathrm{Tel}(A^0; \mathbf{a}) \otimes_{A^0} A, X).$$

Let $(A, \bar{\mathfrak{m}})$ be a local DG-ring. For $X \in \mathrm{D}_{\bar{\mathfrak{f}}}^-(A)$ and $Y \in \mathrm{D}^+(A)$, it follows by [21, Theorem 2.15] and [21, Proposition 3.3] that $\mathrm{lc.dim}_A X = \sup \mathrm{R}\Gamma_{\bar{\mathfrak{m}}}(X)$ and $\mathrm{depth}_A Y = \inf \mathrm{R}\Gamma_{\bar{\mathfrak{m}}}(Y)$.

Localization. Let $\pi_A : A \rightarrow \mathrm{H}^0(A)$ be the canonical surjection and $\pi_A^0 : A^0 \rightarrow \mathrm{H}^0(A)$ be its degree 0 component. Given a prime ideal $\bar{\mathfrak{p}} \in \mathrm{Spec} \mathrm{H}^0(A)$, let $\mathfrak{p} = (\pi_A^0)^{-1}(\bar{\mathfrak{p}}) \in \mathrm{Spec} A^0$, and define $A_{\bar{\mathfrak{p}}} := A \otimes_{A^0} A_{\mathfrak{p}}^0$. More generally, given $X \in \mathrm{D}(A)$, we define

$$X_{\bar{\mathfrak{p}}} := X \otimes_A A_{\bar{\mathfrak{p}}} = X \otimes_{A^0} A_{\mathfrak{p}}^0 \in \mathrm{D}(A_{\bar{\mathfrak{p}}}).$$

A prime ideal $\bar{\mathfrak{p}} \in \mathrm{Spec} \mathrm{H}^0(A)$ is called an *associated prime* of X if $\mathrm{depth}_{A_{\bar{\mathfrak{p}}}} X_{\bar{\mathfrak{p}}} = \inf X_{\bar{\mathfrak{p}}}$. The set of associated primes of X is denoted by $\mathrm{Ass}_A X$. The *support* of X is the set

$$\mathrm{Supp}_A X := \{\bar{\mathfrak{p}} \in \mathrm{Spec} \mathrm{H}^0(A) \mid X_{\bar{\mathfrak{p}}} \neq 0\}.$$

For $\bar{\mathfrak{p}} \in \mathrm{Spec} \mathrm{H}^0(A)$, denote by $E(A, \bar{\mathfrak{p}})$ the DG-module corresponding to the injective hull $E(\mathrm{H}^0(A), \bar{\mathfrak{p}})$ of the residue field $\mathrm{H}^0(A)_{\bar{\mathfrak{p}}}/\bar{\mathfrak{p}}\mathrm{H}^0(A)_{\bar{\mathfrak{p}}}$.

Next we bring several necessary lemmas and their proofs.

Lemma 2.3. (1) For $X \in \mathrm{D}_{\bar{\mathfrak{f}}}^-(A)$ and $Y \in \mathrm{D}^+(A)$, one has

$$\inf \mathrm{RHom}_A(X, Y) = \inf \{\inf \mathrm{RHom}_A(\mathrm{H}^\ell(X), Y) - \ell \mid \ell \in \mathbb{Z}\}.$$

(2) For $X \in \mathrm{D}_{\bar{\mathfrak{f}}}^-(A)$ and $Z \in \mathrm{D}^-(A)$, one has

$$\sup(X \otimes_A^L Z) = \sup \{\sup(\mathrm{H}^\ell(X) \otimes_A^L Z) + \ell \mid \ell \in \mathbb{Z}\}.$$

(3) Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring. For $X, Y \in \mathrm{D}_{\bar{\mathfrak{f}}}^-(A)$, one has

$$\sup(X \otimes_A^L Y) = \sup X + \sup Y.$$

Proof. (1) We first show that $\text{infRHom}_A(X, Y) = \inf\{\text{depth}_{A_{\bar{\mathfrak{p}}}}Y_{\bar{\mathfrak{p}}} - \text{sup}X_{\bar{\mathfrak{p}}}\mid \bar{\mathfrak{p}} \in \text{Spec}H^0(A)\}$. For each $\bar{\mathfrak{p}} \in \text{Spec}H^0(A)$ one has the next (in)equalities

$$\begin{aligned} r := \text{infRHom}_A(X, Y) &\leq \text{infRHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, Y_{\bar{\mathfrak{p}}}) \\ &\leq \text{depth}_{A_{\bar{\mathfrak{p}}}}\text{RHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, Y_{\bar{\mathfrak{p}}}) \\ &= \text{depth}_{A_{\bar{\mathfrak{p}}}}Y_{\bar{\mathfrak{p}}} - \text{sup}X_{\bar{\mathfrak{p}}}, \end{aligned}$$

where the first inequality is by [12, Proposition 2.4], the second inequality is by [21, Proposition 3.3] and the equality is by [26, Proposition 4.9] and the Nakayama's lemma. If $\bar{\mathfrak{p}} \in \text{Ass}_{H^0(A)}H^r(\text{RHom}_A(X, Y))$, then the claimed equality holds by [21, Proposition 3.3].

Set $i = \text{infRHom}_A(X, Y)$ and $j = \inf\{\text{infRHom}_A(H^\ell(X), Y) - \ell \mid \ell \in \mathbb{Z}\}$. For $\bar{\mathfrak{p}} \in \text{Spec}H^0(A)$, set $s = \text{sup}X_{\bar{\mathfrak{p}}}$. By the preceding proof, we have

$$\text{depth}_{A_{\bar{\mathfrak{p}}}}Y_{\bar{\mathfrak{p}}} - \text{sup}X_{\bar{\mathfrak{p}}} = \text{depth}_{A_{\bar{\mathfrak{p}}}}Y_{\bar{\mathfrak{p}}} - s \geq \text{infRHom}_A(H^s(X), Y) - s \geq j,$$

and hence $i \geq j$. On the other hand, let $\ell \in \mathbb{Z}$ and $H(\text{RHom}_A(H^\ell(X), Y)) \neq 0$. Choose $\bar{\mathfrak{p}} \in \text{Spec}H^0(A)$ such that $\text{infRHom}_A(H^\ell(X), Y) = \text{depth}_{A_{\bar{\mathfrak{p}}}}Y_{\bar{\mathfrak{p}}}$. Then

$$\text{RHom}_A(H^\ell(X), Y) - \ell \geq \text{depth}_{A_{\bar{\mathfrak{p}}}}Y_{\bar{\mathfrak{p}}} - \text{sup}X_{\bar{\mathfrak{p}}} \geq i.$$

Therefore, the equality holds.

(2) Set $E = \bigoplus_{\bar{\mathfrak{m}} \in \text{Max}H^0(A)} E(A, \bar{\mathfrak{m}})$. We have the following equalities

$$\begin{aligned} \text{sup}(X \otimes_A^L Z) &= -\text{infRHom}_A(X, \text{RHom}_A(Z, E)) \\ &= -\inf\{\text{infRHom}_A(H^\ell(X), \text{RHom}_A(Z, E)) - \ell \mid \ell \in \mathbb{Z}\} \\ &= \text{sup}\{\ell - \text{infRHom}_A(H^\ell(X) \otimes_A^L Z, E) \mid \ell \in \mathbb{Z}\} \\ &= \text{sup}\{\ell + \text{sup}(H^\ell(X) \otimes_A^L Z) \mid \ell \in \mathbb{Z}\}, \end{aligned}$$

where the first and the last ones are by [19, Theorem 4.10], the second one is by (1).

(3) As $H^{\text{sup}X}(X)$ and $H^{\text{sup}Y}(Y)$ are finitely generated $H^0(A)$ -modules,

$$H^{\text{sup}X + \text{sup}X}(\bar{\kappa} \otimes_A^L X \otimes_A^L Y) \cong \bar{\kappa} \otimes_{H^0(A)} H^{\text{sup}X}(X) \otimes_{\bar{\kappa}} \bar{\kappa} \otimes_{H^0(A)} H^{\text{sup}Y}(Y) \neq 0,$$

which implies the desired equality. □

By the equality (2.14) of [21, Proposition 2.13], for $X \in D^-(A)$, one has

$$\text{lc.dim}_A X = \text{sup}\{\text{dim}H^0(A)/\bar{\mathfrak{p}} + \text{sup}X_{\bar{\mathfrak{p}}}\mid \bar{\mathfrak{p}} \in \text{Spec}H^0(A)\}.$$

Lemma 2.4. *For $X \in D_f^-(A)$ and $Y \in D^-(A)$, there exists an equality*

$$\text{lc.dim}_A(X \otimes_A^L Y) = \text{sup}\{\text{lc.dim}_A(H^\ell(X) \otimes_A^L Y) + \ell \mid \ell \in \mathbb{Z}\}.$$

Proof. We have the following computations

$$\begin{aligned}
\text{lc.dim}(X \otimes_A^L Y) &= \sup\{\dim H^0(A)/\bar{\mathfrak{p}} + \sup(X \otimes_A^L Y)_{\bar{\mathfrak{p}}} \mid \bar{\mathfrak{p}} \in \text{Spec} H^0(A)\} \\
&= \sup\{\dim H^0(A)/\bar{\mathfrak{p}} + \sup\{\sup(H^\ell(X_{\bar{\mathfrak{p}}}) \otimes_{A_{\bar{\mathfrak{p}}}^L} Y_{\bar{\mathfrak{p}}}) + \ell \mid \ell \in \mathbb{Z}\} \mid \bar{\mathfrak{p}} \in \text{Spec} H^0(A)\} \\
&= \sup\{\dim H^0(A)/\bar{\mathfrak{p}} + \sup(H^\ell(X) \otimes_A^L Y)_{\bar{\mathfrak{p}}} + \ell \mid \bar{\mathfrak{p}} \in \text{Spec} H^0(A), \ell \in \mathbb{Z}\} \\
&= \sup\{\text{lc.dim}_A(H^\ell(X) \otimes_A^L Y) + \ell \mid \ell \in \mathbb{Z}\},
\end{aligned}$$

where the second equality follows by Lemma 2.3(2), as desired. \square

Dualizing DG-module. Following [19], a dualizing DG-module R over a DG-ring A is a DG-module $R \in D_f^+(A)$ such that $\text{injdim}_A R < \infty$ and the natural map $A \rightarrow \text{RHom}_A(R, R)$ is an isomorphism in $D(A)$. If A is local then dualizing DG-module over A is unique up to shift. We say that R is normalized if $\text{inf} R = -\dim H^0(A)$. The next result is a DG-version of Grothendieck's local duality, which is a nice generalization of [19, Theorem 7.26].

Theorem 2.5. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp} A < \infty$ and R a normalized dualizing DG-module over A .*

(1) *For every $X \in D_f(A)$, one has an isomorphism in $D(A)$*

$$\text{R}\Gamma_{\bar{\mathfrak{m}}}(X) \simeq \text{RHom}_A(\text{RHom}_A(X, R), E(A, \bar{\mathfrak{m}})).$$

(2) *For every $Y \in D_f^-(A)$, one has an isomorphism in $D(A)$*

$$\text{L}\Lambda^{\bar{\mathfrak{m}}}(Y) \simeq \text{RHom}_A(\text{RHom}_A(Y, E(A, \bar{\mathfrak{m}})), R).$$

Proof. (1) Set $E = E(A, \bar{\mathfrak{m}})$ and $\text{Tel}(A; \bar{\mathfrak{m}}) = \text{Tel}(A^0; \bar{\mathfrak{m}}) \otimes_{A^0} A$. First assume that $X \in D_f^-(A)$. It follows by Lemma 2.2 and [19, Proposition 7.25] that

$$X \otimes_A^L \text{Tel}(A, \bar{\mathfrak{m}}) \simeq X \otimes_A^L \text{RHom}_A(R, R \otimes_A^L \text{Tel}(A, \bar{\mathfrak{m}})) \simeq \text{RHom}_A(\text{RHom}_A(X, R), E).$$

Now assume that $X \in D_f(A)$, set $X' = \cdots \rightarrow X^{n-2} \xrightarrow{d_{n-2}} X^{n-1} \xrightarrow{d_{n-1}} \text{im} d^{n-1} \rightarrow 0$ and $X'' = 0 \rightarrow X^n / \text{im} d^{n-1} \xrightarrow{\bar{d}_n} X^{n+1} \xrightarrow{\bar{d}_{n+1}} X^{n+2} \rightarrow \cdots$. The exact triangle $X' \rightarrow X \rightarrow X'' \rightsquigarrow$ induces a commutative diagram of exact triangles in $D(A)$:

$$\begin{array}{ccccc}
\text{R}\Gamma_{\bar{\mathfrak{m}}}(X') & \longrightarrow & \text{R}\Gamma_{\bar{\mathfrak{m}}}(X) & \longrightarrow & \text{R}\Gamma_{\bar{\mathfrak{m}}}(X'') \\
\downarrow & & \downarrow & & \downarrow \\
\text{RHom}_A(\text{RHom}_A(X', R), E) & \longrightarrow & \text{RHom}_A(\text{RHom}_A(X, R), E) & \longrightarrow & \text{RHom}_A(\text{RHom}_A(X'', R), E)
\end{array}$$

As $X' \in D_f^-(A)$ and $X'' \in D_f^+(A)$, it follows by [19, Theorem 7.26] and the preceding proof that the first and the third vertical morphisms are isomorphisms in $D(A)$. Consequently, the middle vertical morphism is an isomorphism in $D(A)$, as desired.

(2) Since $Y \simeq \text{RHom}_A(\text{RHom}_A(Y, R), R)$, $\text{L}\Lambda^{\bar{\mathfrak{m}}}(Y) \simeq \text{RHom}_A(\text{R}\Gamma_{\bar{\mathfrak{m}}}(\text{RHom}_A(Y, R)), R) \simeq \text{RHom}_A(\text{RHom}_A(Y, \text{R}\Gamma_{\bar{\mathfrak{m}}}(R)), R) \simeq \text{RHom}_A(\text{RHom}_A(Y, E(A, \bar{\mathfrak{m}})), R)$ by Lemma 2.2(2) and [19, Proposition 7.25]. \square

Corollary 2.6. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$ and R a dualizing DG-module for A . For $X \in D_{\bar{\mathfrak{f}}}^-(A)$ and $Y \in D_{\bar{\mathfrak{f}}}^+(A)$, one has*

$$\begin{aligned} \inf \text{RHom}_A(X, R) &= \text{depth}_A R - \text{lc.dim}_A X, & \text{depth}_A \text{RHom}_A(X, R) &= \text{depth}_A R - \sup X, \\ \sup \text{RHom}_A(Y, R) &= \text{depth}_A R - \text{depth}_A Y, & \text{lc.dim}_A \text{RHom}_A(Y, R) &= \text{depth}_A R - \inf Y, \\ \text{injdim}_A \text{RHom}_A(X, R) &= \text{projdim}_A X + \text{depth}_A R, \\ \text{projdim}_A \text{RHom}_A(Y, R) &= \text{injdim}_A Y - \text{depth}_A R. \end{aligned}$$

In particular, if R is normalized then $\text{depth}A = -\sup R$ and $\text{injdim}_A R = 0 = \text{depth}_A R$.

Proof. By Lemma 2.2(1), we have

$$\bar{\kappa} \otimes_A^L \text{RHom}_A(R, R) \simeq \text{RHom}_A(\text{RHom}_A(\bar{\kappa}, R), R) \simeq \text{RHom}_{\bar{\kappa}}(\text{RHom}_A(\bar{\kappa}, R), \text{RHom}_A(\bar{\kappa}, R)),$$

it follows by [12, Corollary 2.3] that $0 = -\inf(\bar{\kappa} \otimes_A^L \text{RHom}_A(R, R)) = -\inf \text{RHom}_A(\bar{\kappa}, R) + \sup \text{RHom}_A(\bar{\kappa}, R) = -\text{depth}_A R + \text{injdim}_A R$. Set $R' = R[-\inf R - \text{lc.dim}A]$. Then R' is a normalized dualizing DG-module over A . Hence we have the following equalities

$$\begin{aligned} \inf \text{RHom}_A(X, R) &= \inf \text{RHom}_A(X, R') + \inf R + \text{lc.dim}A \\ &= -\sup \text{RHom}_A(\text{RHom}_A(X, R'), E(A, \bar{\mathfrak{m}})) + \inf R + \text{lc.dim}A \\ &= -\sup \text{R}\Gamma_{\bar{\mathfrak{m}}}(X) + \inf R + \text{lc.dim}A \\ &= -\text{lc.dim}_A X + \text{injdim}_A R \\ &= \text{depth}_A R - \text{lc.dim}_A X, \end{aligned}$$

where the second one is by [19, Theorem 4.10], the third one is by Theorem 2.5, the fourth one is by [3, Lemma 2.7]. We also have the following equalities

$$\begin{aligned} \sup \text{RHom}_A(Y, R) &= \sup \text{RHom}_A(Y, R') + \inf R + \text{lc.dim}A \\ &= -\inf \text{RHom}_A(\text{RHom}_A(Y, R'), E(A, \bar{\mathfrak{m}})) + \inf R + \text{lc.dim}A \\ &= -\inf \text{R}\Gamma_{\bar{\mathfrak{m}}}(Y) + \text{injdim}_A R \\ &= \text{depth}_A R - \text{depth}_A Y, \end{aligned}$$

where the second one is by [19, Theorem 4.10], the third one is by Theorem 2.5. Therefore, $\text{lc.dim}_A \text{RHom}_A(Y, R) = \text{depth}_A R - \inf Y$ and $\text{depth}_A \text{RHom}_A(X, R) = \text{depth}_A R - \sup X$.

By [12, Corollary 2.31], one has the following equalities:

$$\begin{aligned} \text{injdim}_A \text{RHom}_A(X, R) &= \sup \text{RHom}_A(\bar{\kappa}, \text{RHom}_A(X, R)) \\ &= \sup \text{RHom}_{\bar{\kappa}}(\bar{\kappa} \otimes_A^L X, \text{RHom}_A(\bar{\kappa}, R)) \\ &= \sup \text{RHom}_A(\bar{\kappa}, R) - \inf(\bar{\kappa} \otimes_A^L X) \\ &= \text{projdim}_A X + \text{depth}_A R. \end{aligned}$$

By [12, Corollary 2.3] and Lemma 2.2(1), one has the following equalities:

$$\begin{aligned}
\text{projdim}_A \text{RHom}_A(Y, R) &= -\inf(\bar{\kappa} \otimes_A^L \text{RHom}_A(Y, R)) \\
&= -\inf \text{RHom}_{\bar{\kappa}}(\text{RHom}_A(\bar{\kappa}, Y), \text{RHom}_A(\bar{\kappa}, R)) \\
&= -\inf \text{RHom}_A(\bar{\kappa}, R) + \sup \text{RHom}_A(\bar{\kappa}, Y) \\
&= \text{injdim}_A Y - \text{depth}_A R.
\end{aligned}$$

We obtain the equalities we seek. \square

3. Intersection Theorem

In this section, we examine the Intersection Theorem (IT) for DG-modules and study the local cohomology Krull dimension of the DG-modules $\text{RHom}_A(X, Y)$ and $X \otimes_A^L Y$, which extends some classical results proved by Foxby and Christensen in [6, 4].

We begin with the following main theorem of this section, which is an extension of IT in [6, 18.7] and [5, Theorem 2.6] in DG-setting.

Theorem 3.1. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp} A < \infty$.*

(1) *Let $X \in D^-(A)$ with $\text{flatdim}_A X < \infty$ and $0 \not\cong Y \in D_f^-(A)$. If $\text{depth}_A X < \infty$, then*

$$\text{lc.dim}_A Y \leq \text{lc.dim}_A(X \otimes_A^L Y) + \text{flatdim}_A X.$$

(2) *For $0 \not\cong X, Y \in D_f^-(A)$ with $\text{projdim}_A X < \infty$, one has an inequality*

$$\text{lc.dim}_A Y \leq \text{lc.dim}_A(X \otimes_A^L Y) + \text{projdim}_A X.$$

Proof. (1) A straightforward computation based on Lemma 2.4:

$$\begin{aligned}
\text{lc.dim}_A(X \otimes_A^L Y) &= \sup\{\text{lc.dim}_A(X \otimes_A^L H^\ell(Y)) + \ell \mid \ell \in \mathbb{Z}\} \\
&= \sup\{\text{lc.dim}_{H^0(A)}(X \otimes_A^L H^0(A) \otimes_{H^0(A)}^L H^\ell(Y)) + \ell \mid \ell \in \mathbb{Z}\} \\
&\geq \sup\{\dim_{H^0(A)} H^\ell(Y) + \ell \mid \ell \in \mathbb{Z}\} - \text{flatdim}_{H^0(A)}(X \otimes_A^L H^0(A)) \\
&= \text{lc.dim}_A Y - \text{flatdim}_A X,
\end{aligned}$$

where the inequality follows by [10, Theorem 2.3] and [6, 19.7], the last equality follows from [3, Corollary 1.5].

(2) As $\text{projdim}_A X < \infty$, it follows by [26, Theorem 4.5] that $\text{depth}_A X < \infty$. Hence (1) yields the desired inequality. \square

Corollary 3.2. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp} A < \infty$ and $Y \in D^-(A)$ with $\text{depth}_A Y < \infty$. If either $0 \not\cong X \in D_f^-(A)$ or $\text{depth}_A X < \infty$, then there is an inequality,*

$$\text{flatdim}_A Y \geq \text{depth}_A X - \text{lc.dim}_A(Y \otimes_A^L X).$$

Proof. One can assume that $\text{flatdim}_A Y < \infty$.

If $0 \not\cong X \in D_f^-(A)$, then we have the next inequalities

$$\begin{aligned}
\text{flatdim}_A Y &\geq \text{lc.dim}_A X - \text{lc.dim}_A(Y \otimes_A^L X) \\
&\geq \sup \text{R}\Gamma_{\bar{\mathfrak{m}}}(X) - \text{lc.dim}_A(Y \otimes_A^L X) \\
&\geq \inf \text{R}\Gamma_{\bar{\mathfrak{m}}}(X) - \text{lc.dim}_A(Y \otimes_A^L X) \\
&= \text{depth}_A X - \text{lc.dim}_A(Y \otimes_A^L X),
\end{aligned}$$

where the first one is by Theorem 3.1(1), the second one is by [21, Theorem 2.15], the third one is by $\text{R}\Gamma_{\bar{\mathfrak{m}}}(X) \neq 0$ and the equality is by [21, Proposition 3.3].

If $\text{depth}_A X < \infty$, then $\text{depth}_A(Y \otimes_A^L X) < \infty$ and so $\text{depth}_A(Y \otimes_A^L X) \leq \text{lc.dim}_A(Y \otimes_A^L X)$. By [26, Theorem 4.8], one has $\text{flatdim}_A Y = \text{depth}_A X - \text{depth}_A(Y \otimes_A^L X)$, which implies the desired inequality. \square

Corollary 3.3. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$, and let $X \in \text{D}^-(A)$ with $\text{flatdim}_A X < \infty$ and $0 \neq Y \in \text{D}_f^-(A)$. If either $\text{depth}_A X < \infty$ or $0 \neq X \in \text{D}_f(A)$, then there exists an inequality,*

$$\text{lc.dim}_A(X \otimes_A^L Y) - \text{depth}_A(X \otimes_A^L Y) \geq \text{lc.dim}_A Y - \text{depth}_A Y.$$

In particular, $\text{lc.dim}_A X - \text{depth}_A X \geq \text{lc.dim}A - \text{depth}A$.

Proof. By [12, Corollary 2.31] and [3, Proposition 6.14], one has $\text{flatdim}_A X = -\inf(\bar{\kappa} \otimes_A^L X)$. If $X \in \text{D}_f^-(A)$, then $\text{projdim}_A X = \text{flatdim}_A X$ by [12, Proposition 2.2]. As $\text{R}\Gamma_{\bar{\mathfrak{m}}}(\bar{\kappa}) \simeq \bar{\kappa}$, it follows by [26, Theorem 4.8] that $\inf(\bar{\kappa} \otimes_A^L X) = \inf \text{R}\Gamma_{\bar{\mathfrak{m}}}(\bar{\kappa} \otimes_A^L X) = \text{depth}_A(\bar{\kappa} \otimes_A^L X) = \text{depth}_A X - \text{depth}A$. By Theorem 3.1 and [26, Theorem 4.8], one has

$$\text{lc.dim}_A(X \otimes_A^L Y) - \text{depth}_A(X \otimes_A^L Y) \geq \text{lc.dim}_A Y - \text{flatdim}_A X + \text{depth}A - \text{depth}_A X - \text{depth}_A Y.$$

Thus we obtain the desired inequality. \square

Following [21], a local DG-ring $(A, \bar{\mathfrak{m}})$ with $\text{amp}A < \infty$ is called *local Cohen-Macaulay* if $\text{amp} \text{R}\Gamma_{\bar{\mathfrak{m}}}(A) = \text{amp}A$. A DG-module $X \in \text{D}_f^b(A)$ is called *local Cohen-Macaulay* if

$$\text{amp}X = \text{amp}A = \text{amp} \text{R}\Gamma_{\bar{\mathfrak{m}}}(X).$$

Remark 3.4. Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$.

(1) The next example shows that the condition $\text{depth}_A X < \infty$ or $X \in \text{D}_f^-(A)$ can not be dropped in Corollaries 3.2 and 3.3.

Set $(A, \bar{\mathfrak{m}}) = (R, \mathfrak{m})$ be a local ring, $\mathbf{x} = x_1, \dots, x_n$ an R -regular sequence in \mathfrak{m} and \mathfrak{p} a prime ideal such that $\mathbf{x} \notin \mathfrak{p}$. Set $X = E(R/\mathfrak{p})$ and $Y = R/\mathbf{x}$. Then $X \notin \text{D}_f^-(R)$ and $\text{depth}_R X = \infty$. Thus $\text{flatdim}_R Y = n < \text{depth}_R X - \text{lc.dim}_R(Y \otimes_R^L X)$ and $\text{lc.dim}_R X - \text{depth}_R X = -\infty < \text{lc.dim}R - \text{depth}R$.

(2) Let $X \in \text{D}^-(A)$ with $\text{depth}_A X < \infty$. One can assume that $\text{flatdim}_A X < \infty$. Then $\text{flatdim}_A X = -\text{depth}_A(\bar{\kappa} \otimes_A^L X) = \text{depth}A - \text{depth}_A X$, it follows by Corollary 3.3 that $\text{flatdim}_A X \geq \text{lc.dim}A - \text{lc.dim}_A X$. If $X \in \text{D}_f^-(A)$, then $\text{depth}_A X < \infty$ by [21, Proposition 3.5], and so $\text{projdim}_A X \geq \text{lc.dim}A - \text{lc.dim}_A X$.

(3) Let $0 \neq X \in D^-(A)$ with $\text{flatdim}_A X < \infty$. If $X \simeq R\Gamma_{\bar{\mathfrak{m}}}(X)$, then $\text{lc.dim}_A X = \sup X$ and $\text{depth}_A X = \inf X$, and hence $\text{amp} X \geq \text{lc.dim} A - \text{depth} A$.

(4) If there is a local Cohen-Macaulay DG-module $X \in D_f^b(A)$ with $\text{projdim}_A X < \infty$, then $\text{amp} R\Gamma_{\bar{\mathfrak{m}}}(X) = \text{lc.dim}_A X - \text{depth}_A X \geq \text{lc.dim} A - \text{depth} A = \text{amp} R\Gamma_{\bar{\mathfrak{m}}}(A)$ by Corollary 3.3. As $\text{amp} R\Gamma_{\bar{\mathfrak{m}}}(A) \geq \text{amp} A$ by [21, Theorem 4.1(1)], it implies that $\text{amp} R\Gamma_{\bar{\mathfrak{m}}}(A) = \text{amp} A$ and A is local Cohen-Macaulay.

(5) For $0 \neq X, Y \in D_f^b(A)$ with $\text{projdim}_A X < \infty$, if $\text{Supp}_A X \cap \text{Supp}_A Y = \{\bar{\mathfrak{m}}\}$, then $\text{lc.dim}_A(X \otimes_A^L Y) = \sup\{\dim H^0(A)/\bar{\mathfrak{p}} + \sup(X \otimes_A^L Y)_{\bar{\mathfrak{p}}} \mid \bar{\mathfrak{p}} \in \text{Supp}_A X \cap \text{Supp}_A Y\} = \sup X + \sup Y$. Thus $\text{lc.dim}_A Y - \sup Y \leq \text{projdim}_A X + \sup X$ by Theorem 3.1(2).

The next result is a generalization of [27, Theorem 3.7] and [5, Theorems 2.1 and 2.6].

Theorem 3.5. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp} A < \infty$, and let $X, Y \in D(A)$ such that $\text{RHom}_A(X, Y) \neq 0$.*

(i) *If $X \in D_f^-(A)$ with $\text{projdim}_A X < \infty$ and $Y \in D^b(A)$, then there is an inequality*

$$\text{lc.dim}_A \text{RHom}_A(X, Y) \leq \text{lc.dim}_A Y + \text{projdim}_A X.$$

(ii) *If $X, Y \in D_f^b(A)$ with $\text{projdim}_A X < \infty$, then there are inequalities*

$$\text{lc.dim}_A Y - \sup X \leq \text{lc.dim}_A \text{RHom}_A(X, Y) \leq \text{lc.dim}_A(X \otimes_A^L Y) + \text{projdim}_A X - \inf X.$$

(iii) *If $X, Y \in D_f^b(A)$ with either $\text{projdim}_A X < \infty$ or $\text{injdim}_A Y < \infty$, then*

$$\text{lc.dim}_A \text{RHom}_A(X, Y) \leq \text{lc.dim} A - \inf X + \sup Y.$$

Proof. (i) By [21, Remark 2.3], $\text{lc.dim}_A \text{RHom}_A(X, Y) \leq \dim H^0(A) + \sup \text{RHom}_A(X, Y) < \infty$. One can choose $\bar{\mathfrak{p}} \in \text{Supp}_A X \cap \text{Supp}_A Y$ such that

$$\text{lc.dim}_A \text{RHom}_A(X, Y) = \dim H^0(A)/\bar{\mathfrak{p}} + \sup \text{RHom}_A(X, Y)_{\bar{\mathfrak{p}}}.$$

By Lemmas 2.1(1) and 2.2(2), we have $\text{RHom}_A(X, Y)_{\bar{\mathfrak{p}}} \simeq \text{RHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, A_{\bar{\mathfrak{p}}}) \otimes_{A_{\bar{\mathfrak{p}}}}^L Y_{\bar{\mathfrak{p}}}$ and $\sup \text{RHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, A_{\bar{\mathfrak{p}}}) \leq \sup \text{RHom}_A(X, A) = \text{projdim}_A X$. Hence

$$\begin{aligned} \dim H^0(A)/\bar{\mathfrak{p}} + \sup \text{RHom}_A(X, Y)_{\bar{\mathfrak{p}}} &\leq \dim H^0(A)/\bar{\mathfrak{p}} + \sup Y_{\bar{\mathfrak{p}}} + \sup \text{RHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, A_{\bar{\mathfrak{p}}}) \\ &\leq \text{lc.dim}_A Y + \text{projdim}_A X. \end{aligned}$$

(ii) By the proof of (i), one has $\text{lc.dim}_A \text{RHom}_A(X, Y) \leq \dim H^0(A)/\bar{\mathfrak{p}} + \sup Y_{\bar{\mathfrak{p}}} + \text{projdim}_A X$ for some $\bar{\mathfrak{p}} \in \text{Supp}_A X \cap \text{Supp}_A Y$. By Lemma 2.3(3), we have

$$\begin{aligned} \text{lc.dim}_A \text{RHom}_A(X, Y) &\leq \dim H^0(A)/\bar{\mathfrak{p}} + \sup(X \otimes_A^L Y)_{\bar{\mathfrak{p}}} - \sup X_{\bar{\mathfrak{p}}} + \text{projdim}_A X \\ &\leq \text{lc.dim}_A(X \otimes_A^L Y) + \text{projdim}_A X - \inf X. \end{aligned}$$

For the left-hand side of (ii), as $\text{RHom}_A(X, A) \in D_f^b(A)$ and $\text{projdim}_A \text{RHom}_A(X, A) < \infty$, it follows by Theorem 3.1(2) that

$$\begin{aligned}
\text{lc.dim}_A Y &\leq \text{lc.dim}_A(\text{RHom}_A(X, A) \otimes_A^L Y) + \text{projdim}_A \text{RHom}_A(X, A) \\
&= \text{lc.dim}_A \text{RHom}_A(X, Y) + \text{supRHom}_A(\text{RHom}_A(X, A), A) \\
&= \text{lc.dim}_A \text{RHom}_A(X, Y) + \text{sup}X,
\end{aligned}$$

where the first equality is by [26, Proposition 4.4(3)] and the second one follows from the isomorphism $X \simeq \text{RHom}_A(\text{RHom}_A(X, A), A)$, as claimed.

(iii) Fix a point $\bar{\mathfrak{p}} \in \text{Spec}H^0(A)$. Assume that $\text{projdim}_A X < \infty$. Then the following computations hold:

$$\begin{aligned}
\dim H^0(A)/\bar{\mathfrak{p}} + \text{supRHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, Y_{\bar{\mathfrak{p}}}) &\leq \dim H^0(A)/\bar{\mathfrak{p}} + \text{projdim}_{A_{\bar{\mathfrak{p}}}} X_{\bar{\mathfrak{p}}} + \text{sup}Y_{\bar{\mathfrak{p}}} \\
&= \dim H^0(A)/\bar{\mathfrak{p}} + \text{depth}_{A_{\bar{\mathfrak{p}}}} - \text{depth}_{A_{\bar{\mathfrak{p}}}} X_{\bar{\mathfrak{p}}} + \text{sup}Y_{\bar{\mathfrak{p}}} \\
&\leq \dim H^0(A)/\bar{\mathfrak{p}} + \dim H^0(A)_{\bar{\mathfrak{p}}} + \text{inf}A_{\bar{\mathfrak{p}}} - \text{depth}_{A_{\bar{\mathfrak{p}}}} X_{\bar{\mathfrak{p}}} + \text{sup}Y_{\bar{\mathfrak{p}}} \\
&\leq \dim H^0(A) - \text{inf}X_{\bar{\mathfrak{p}}} + \text{sup}Y_{\bar{\mathfrak{p}}} \\
&\leq \text{lc.dim}A - \text{inf}X + \text{sup}Y,
\end{aligned}$$

where the equality is by [26, Theorem 4.5], the second inequality is by [21, Corollary 3.6] and the third one is by [21, Proposition 3.3].

Assume that $\text{injdim}_A Y < \infty$. Then the next computations hold:

$$\begin{aligned}
\dim H^0(A)/\bar{\mathfrak{p}} + \text{supRHom}_{A_{\bar{\mathfrak{p}}}}(X_{\bar{\mathfrak{p}}}, Y_{\bar{\mathfrak{p}}}) &\leq \dim H^0(A)/\bar{\mathfrak{p}} + \text{injdim}_{A_{\bar{\mathfrak{p}}}} Y_{\bar{\mathfrak{p}}} - \text{inf}X_{\bar{\mathfrak{p}}} \\
&= \dim H^0(A)/\bar{\mathfrak{p}} + \text{depth}_{A_{\bar{\mathfrak{p}}}} + \text{sup}Y_{\bar{\mathfrak{p}}} - \text{inf}X_{\bar{\mathfrak{p}}} \\
&\leq \dim H^0(A)/\bar{\mathfrak{p}} + \dim H^0(A)_{\bar{\mathfrak{p}}} + \text{inf}A_{\bar{\mathfrak{p}}} + \text{sup}Y_{\bar{\mathfrak{p}}} - \text{inf}X_{\bar{\mathfrak{p}}} \\
&\leq \dim H^0(A) + \text{sup}Y_{\bar{\mathfrak{p}}} - \text{inf}X_{\bar{\mathfrak{p}}} \\
&\leq \text{lc.dim}A + \text{sup}Y - \text{inf}X,
\end{aligned}$$

where the equality is by [26, Theorem 4.7].

Therefore, we obtain the desired inequality. \square

Following [21], a local DG-ring $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ with $\text{amp}A < \infty$ is called *Gorenstein* if $\text{injdim}_A A < \infty$. The following characterization of Gorenstein DG-rings is a DG-version of Auslander-Bridger characterization of classical Gorenstein rings.

Corollary 3.6. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$. Then the following are equivalent:*

- (1) A is local Gorenstein;
- (2) $\text{lc.dim}_A \text{RHom}_A(X, A) < \infty$ for all $X \in D_{\bar{\mathfrak{f}}}^b(A)$;
- (3) $\text{lc.dim}_A \text{RHom}_A(T, A) < \infty$ for all finitely generated $H^0(A)$ -modules T ;
- (4) $\text{lc.dim}_A \text{RHom}_A(\bar{\kappa}, A) < \infty$.

Proof. (1) \Rightarrow (2) follows from Theorem 3.5(iii) and (2) \Rightarrow (3) \Rightarrow (4) are trivial.

(4) \Rightarrow (1) One has the following equalities

$$\sup \mathrm{RHom}_A(\bar{\kappa}, A) = \sup \mathrm{RHom}_A(\bar{\kappa}, A)_{\bar{\mathfrak{m}}} = \mathrm{lc. dim}_A \mathrm{RHom}_A(\bar{\kappa}, A) < \infty.$$

Hence [12, Corollary 2.31] implies that $\mathrm{injdim}_A A < \infty$ and A is local Gorenstein. \square

Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring and $X \in \mathrm{D}^+(A)$. An element $\bar{x} \in \bar{\mathfrak{m}}$ is called *X-regular* if it is $\mathrm{H}^{\mathrm{inf}X}(X)$ -regular, that is, if the multiplication map

$$\bar{x} : \mathrm{H}^{\mathrm{inf}X}(X) \rightarrow \mathrm{H}^{\mathrm{inf}X}(X)$$

is injective. Inductively, we say that a sequence $\bar{x}_1, \dots, \bar{x}_n \in \bar{\mathfrak{m}}$ is *X-regular* if \bar{x}_1 is *X-regular* and the sequence $\bar{x}_2, \dots, \bar{x}_n$ is $X//\bar{x}_1 X$ -regular. Here, $X//\bar{x}_1 X$ is the cone of the map $\bar{x}_1 : X \rightarrow X$ in $\mathrm{D}(A)$. Following [23], a local DG-ring $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ with $\mathrm{amp}A < \infty$ is called *sequence-regular* if $\bar{\mathfrak{m}}$ is generated by an *A-regular* sequence. If A is a regular local ring and M, N two finitely generated *A*-modules, then there is an inequality

$$\dim_A M + \dim_A N \leq \dim A + \dim_A(M \otimes_A N),$$

which is due to Serre and is called strong Intersection Theorem (see [6, 20.4]). We next provide the DG-version of this theorem.

Proposition 3.7. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a sequence-regular DG-ring and $X, Y \in \mathrm{D}_f^b(A)$. Then there exists an inequality*

$$\mathrm{lc. dim}_A X + \mathrm{lc. dim}_A Y \leq \mathrm{lc. dim} A + \mathrm{lc. dim}_A(X \otimes_A^L Y).$$

Proof. One has the following (in)equalities

$$\begin{aligned} \mathrm{lc. dim}_A(X \otimes_A^L Y) &= \sup_{s,t \in \mathbb{Z}} \{ \mathrm{lc. dim}_A(\mathrm{H}^s(X) \otimes_A^L \mathrm{H}^t(Y)) + s + t \} \\ &= \sup_{s,t \in \mathbb{Z}} \{ \dim_{\mathrm{H}^0(A)}(\mathrm{H}^s(X) \otimes_{\mathrm{H}^0(A)} \mathrm{H}^t(Y)) + s + t \} \\ &\geq \sup_{s,t \in \mathbb{Z}} \{ \dim_{\mathrm{H}^0(A)} \mathrm{H}^s(X) + s + \dim_{\mathrm{H}^0(A)} \mathrm{H}^t(Y) + t \} - \dim \mathrm{H}^0(A) \\ &= \mathrm{lc. dim}_A X + \mathrm{lc. dim}_A Y - \mathrm{lc. dim} A, \end{aligned}$$

where the first equality is by Lemma 2.4, the second one is by the equality of the above of Lemma 2.4, the inequality is by [6, 20.4] as $\mathrm{H}^0(A)$ is regular by [23, Theorem 3.4]. \square

4. Applications

In this section, we explore some implications of our previous results. More precisely, the amplitude inequality of DG-modules is improved; the DG-version of NIT, Krull's principle ideal theorem, Bass Conjecture about Cohen-Macaulay rings and Vasconcelos Conjecture about Gorenstein rings are proved; the Minamoto's conjecture is solved completely.

The next proposition improves the Jørgensen's results, that is to say, eliminate the hypothesis on local DG-rings in [11, Theorem 3.1].

Proposition 4.1. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$ and $0 \neq X, Y \in D_f^b(A)$ with $\text{projdim}_A X < \infty$. One has an inequality*

$$\text{amp}(X \otimes_A^L Y) \geq \text{amp}Y.$$

In particular, we have $\text{amp}X \geq \text{amp}A$.

Proof. Let $B = \text{L}\Lambda(A, \bar{\mathfrak{m}})$. Then $X \otimes_A^L B, Y \otimes_A^L B \in D_f^b(B)$ with $\text{projdim}_B(X \otimes_A^L B) < \infty$ by [3, Proposition 3.7] and $\text{amp}(X \otimes_A^L Y) = \text{amp}((X \otimes_A^L B) \otimes_B^L (Y \otimes_A^L B))$, $\text{amp}Y = \text{amp}(Y \otimes_A^L B)$. Thus we may assume that A is derived $\bar{\mathfrak{m}}$ -adic complete, and A has a normalized dualizing DG-module R by [19, Proposition 7.21]. We have the following isomorphisms

$$\begin{aligned} \text{RHom}_A(X \otimes_A^L Y, R) &\simeq \text{RHom}_A(X, \text{RHom}_A(Y, R)) \\ &\simeq \text{RHom}_A(\text{RHom}_A(\text{RHom}_A(X, A), A), \text{RHom}_A(Y, R)) \\ &\simeq \text{RHom}_A(X, A) \otimes_A^L \text{RHom}_A(Y, R). \end{aligned}$$

where the second one is by $X \simeq \text{RHom}_A(\text{RHom}_A(X, A), A)$ and the third one is by Lemma 2.2(1) as $\text{projdim}_A \text{RHom}_A(X, A) < \infty$. So we have the next (in)equalities

$$\begin{aligned} \text{amp}(X \otimes_A^L Y) &= \text{lc.dim}_A \text{RHom}_A(X \otimes_A^L Y, R) - \text{depth}_A \text{RHom}_A(X \otimes_A^L Y, R) \\ &\geq \text{lc.dim}_A \text{RHom}_A(Y, R) - \text{depth}_A \text{RHom}_A(Y, R) \\ &= \text{amp}Y, \end{aligned}$$

where the equalities are by Corollary 2.6 and the inequality is by the above isomorphism and Corollary 3.3, as desired. \square

Remark 4.2. Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$.

(1) If $\text{amp}A > 0$, then $\bar{\kappa}$ never has finite projective dimension. Thus there are DG-modules in $D_f(A)$ which are not compact.

(2) Let $\bar{\mathfrak{x}}$ be a sequence in $\bar{\mathfrak{m}}$ and $X \in D_f^b(A)$ with $\text{amp}X = \text{amp}A$. If $\text{projdim}_A X < \infty$, then $\text{amp}X // \bar{\mathfrak{x}} = \text{amp}(A // \bar{\mathfrak{x}} \otimes_A^L X) \geq \text{amp}A // \bar{\mathfrak{x}} \geq \text{amp}A$ by Proposition 4.1, it follows by [12, Lemma 2.13] that $\text{amp}A // \bar{\mathfrak{x}} = \text{amp}A$ and $\bar{\mathfrak{x}}$ is A -regular. This is the DG-version of Auslander's zerodivisor conjecture.

(3) The example after [11, Theorem 0.2] shows that the condition $\text{amp}A < \infty$ is crucial.

The next result is an extend of the strong amplitude inequality (see [6, 20.8]).

Proposition 4.3. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a sequence-regular DG-ring and $X, Y \in D_f^b(A)$ with $\text{projdim}_A X < \infty$. One has an inequality*

$$\text{amp}(X \otimes_A^L Y) \geq \text{amp}X + \text{amp}Y + \text{inf}A.$$

Proof. Let $B = \text{L}\Lambda(A, \bar{\mathfrak{m}})$. It follows by [23, Corollary 4.5] that A is sequence-regular if and only if B is sequence-regular. Also $\text{projdim}_B(X \otimes_A^L B) < \infty$, we may assume that A is derived $\bar{\mathfrak{m}}$ -adic complete and has a dualizing DG-module R . Then $\text{RHom}_A(X \otimes_A^L Y, R) \simeq$

$\mathrm{RHom}_A(\mathrm{RHom}_A(\mathrm{RHom}_A(X, A), A), \mathrm{RHom}_A(Y, R)) \simeq \mathrm{RHom}_A(X, A) \otimes_A^L \mathrm{RHom}_A(Y, R)$ by Lemma 2.2(1) as $\mathrm{projdim}_A \mathrm{RHom}_A(X, A) < \infty$. Note that $X \simeq \mathrm{RHom}_A(R, R \otimes_A^L X)$, one has $\inf(R \otimes_A^L X) \leq \inf X + \sup R$ and $\sup(R \otimes_A^L X) = \sup X + \sup R$ by Lemma 2.3(3). Also $\mathrm{RHom}_A(X, A) \simeq \mathrm{RHom}_A(X \otimes_A^L R, R)$, we have the next (in)equalities

$$\begin{aligned} \mathrm{amp}(X \otimes_A^L Y) &= \mathrm{lc.dim}_A \mathrm{RHom}_A(X \otimes_A^L Y, R) - \mathrm{depth}_A \mathrm{RHom}_A(X \otimes_A^L Y, R) \\ &= \mathrm{lc.dim}_A \mathrm{RHom}_A(X, A) \otimes_A^L \mathrm{RHom}_A(Y, R) - \mathrm{depth}_A \mathrm{RHom}_A(X, A) \otimes_A^L \mathrm{RHom}_A(Y, R) \\ &\geq \mathrm{amp}(X \otimes_A^L R) + \mathrm{amp} Y + \inf A \\ &\geq \mathrm{amp} X + \mathrm{amp} Y + \inf A, \end{aligned}$$

where the first inequality is by Proposition 3.7, [26, Theorem 4.8] and Corollary 2.6, the second one is by the preceding proof. \square

We denote by A^\natural the graded ring obtained by forgetting the differential of the DG-ring A , and X^\natural denotes the graded A^\natural -module obtained by forgetting the differential of X in $\mathrm{D}(A)$. We now prove the DG-version of NIT and Krull's principle ideal theorem.

Proposition 4.4. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\mathrm{amp} A < \infty$.*

(1) *Let F be a semi-free DG-module with $F^\natural = \coprod_{-n \leq j \leq 0} A^\natural[-j]^{(\beta_j)}$, where each β_j is finite. If $F \not\cong 0$ and $\dim_{\mathrm{H}^0(A)} \mathrm{H}^i(F) \leq -i$ holds for all i , then $\mathrm{lc.dim} A \leq n$.*

(2) *Let $\bar{x}_1, \dots, \bar{x}_n$ be a sequence in $\mathrm{H}^0(A)$ and $\bar{\mathfrak{p}}$ be minimal among the prime ideals containing $(\bar{x}_1, \dots, \bar{x}_n)$. Then $\mathrm{lc.dim} A_{\bar{\mathfrak{p}}} \leq n$.*

Proof. (1) By induction on n , one can obtain that $\mathrm{projdim}_A F \leq n$. As $\dim_{\mathrm{H}^0(A)} \mathrm{H}^i(F) \leq -i$ for all i , one has $\mathrm{lc.dim}_A F \leq 0$. Hence Theorem 3.1(2) yields that $n \geq \mathrm{projdim}_A F \geq \mathrm{lc.dim} A - \mathrm{lc.dim}_A F \geq \mathrm{lc.dim} A$.

(2) Let $X = A_{\bar{\mathfrak{p}}} // (\bar{x}_1/1, \dots, \bar{x}_n/1)$. Then $\mathrm{projdim}_{A_{\bar{\mathfrak{p}}}} X \leq n$ by [13, Theorem 2.19]. Also $\mathrm{lc.dim}_{A_{\bar{\mathfrak{p}}}} X = 0$, so the asserted inequality follows by Theorem 3.1(2). \square

Let A be a local Cohen-Macaulay DG-ring and $\bar{\mathfrak{x}}$ a maximal A -regular sequence. Then the DG-module $X = A // \bar{\mathfrak{x}}$ has finite projective dimension, $\mathrm{H}(X)$ has finite length and $\mathrm{amp} A = \mathrm{amp} X$. The next corollary proves that its opposite is also true.

Corollary 4.5. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\mathrm{amp} A < \infty$ and $0 \not\cong X \in \mathrm{D}^b(A)$. If $\mathrm{projdim}_A X < \infty$ and $\mathrm{H}(X)$ is finite length, then $\mathrm{lc.dim} A \leq \mathrm{projdim}_A X + \sup X$. In addition, if $\mathrm{amp} X = \mathrm{amp} A$ then A is local Cohen-Macaulay.*

Proof. Note that $\mathrm{projdim}_A X[\sup X] = \mathrm{projdim}_A X + \sup X$ and $\sup X[\sup X] = 0$, it follows that $X[\sup X]$ has a minimal semi-free resolution F with $F^\natural = \coprod_{-n \leq j \leq 0} A^\natural[-j]^{(\beta_j)}$, where $n = \mathrm{projdim}_A X[\sup X]$ and each β_j is finite. As $\mathrm{H}(F) \cong \mathrm{H}(X)$ is finite length, it follows by Proposition 4.4(1) that $\mathrm{lc.dim} A \leq n = \mathrm{projdim}_A X + \sup X$.

If $\text{amp}X = \text{amp}A$, then $\text{lc.dim}A \leq \text{projd}_A X + \text{sup}X = \text{depth}A - \text{depth}_A X - \text{inf}A + \text{inf}X = \text{depth}A - \text{inf}A$, and so A is local Cohen-Macaulay. \square

The affirmative answer to the DG-version of Bass conjecture is the special case of the following proposition, which solves affirmatively conjecture of Minamoto (see [12, Conjecture 2.36]) or Shaul (see [21, Remark 5.25]).

Proposition 4.6. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$, and let $0 \neq X \in D_f^+(A)$ with $\text{injdim}_A X < \infty$ and $0 \neq Y \in D_f^b(A)$. One has an inequality,*

$$\text{ampRHom}_A(Y, X) \geq \text{lc.dim}_A Y - \text{depth}_A Y.$$

In particular, A is local Cohen-Macaulay if and only if there is such X with $\text{amp}X = \text{amp}A$.

Proof. Let $B = \text{L}\Lambda(A, \bar{\mathfrak{m}})$. As $H^0(B) = H^0(A) \otimes_A^L B$ by [18, Lemma 6.3], we have

$$\text{RHom}_B(H^0(B)/\bar{\mathfrak{m}}H^0(B), X \otimes_A^L B) \simeq \text{RHom}_A(H^0(A)/\bar{\mathfrak{m}}, X) \otimes_A^L B,$$

it follows by [12, Corollary 2.31] that $\text{injdim}_B(X \otimes_A^L B) < \infty$. By [21, Propositions 2.10 and 3.3 and Theorem 2.15], one has $\text{depth}_A Y = \text{depth}_B(Y \otimes_A^L B)$ and $\text{lc.dim}_A Y = \text{lc.dim}_B(Y \otimes_A^L B)$. Also $\text{ampRHom}_A(Y, X) = \text{amp}(\text{RHom}_A(Y, X) \otimes_A^L B)$ by [3, Proposition 1.14]. Thus we may assume that A is derived $\bar{\mathfrak{m}}$ -adic complete, and A has a normalized dualizing DG-module R . Then $\text{RHom}_A(X, R) \in D_f^b(A)$ by [28, Proposition 7.2] and Lemma 2.1(2) and $\text{projd}_A \text{RHom}_A(X, R) < \infty$ by Corollary 2.6. Also $\text{RHom}_A(Y, X) \in D_f^b(A)$ and $\text{RHom}_A(X, R) \otimes_A^L Y \simeq \text{RHom}_A(\text{RHom}_A(Y, X), R)$, it follows that

$$\begin{aligned} \text{lc.dim}_A Y - \text{depth}_A Y &\leq \text{lc.dim}_A(\text{RHom}_A(X, R) \otimes_A^L Y) - \text{depth}_A(\text{RHom}_A(X, R) \otimes_A^L Y) \\ &= \text{ampRHom}_A(Y, X). \end{aligned}$$

where the inequality is by Corollary 3.3 and the equality is by Corollary 2.6.

Let A be local Cohen-Macaulay and $\bar{\mathfrak{x}}$ a maximal A -regular sequence. Then the DG-module $X = \text{Hom}_A(A//\bar{\mathfrak{x}}, E(A, \bar{\mathfrak{m}}))$ has finite injective dimension and $\text{amp}A = \text{amp}X$. Conversely, one has $\text{amp}A \leq \text{lc.dim}A - \text{depth}A \leq \text{amp}X = \text{amp}A$ by [21, Theorem 4.1(1)]. Thus A is local Cohen-Macaulay. \square

The next corollary is a DG-setting of Bass characterization of Cohen-Macaulay rings.

Corollary 4.7. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$. Then the following are equivalent:*

- (1) *A is local Cohen-Macaulay;*
- (2) *There exists $0 \neq Y \in D_f^b(A)$ with $\text{amp}Y = \text{amp}A$ so that $\text{lc.dim}_A \text{RHom}_A(X, Y) < \infty$ for all $X \in D_f^b(A)$;*
- (3) *There is $0 \neq Y \in D_f^b(A)$ with $\text{amp}Y = \text{amp}A$ so that $\text{lc.dim}_A \text{RHom}_A(\bar{\kappa}, Y) < \infty$.*

Proof. (1) \Rightarrow (2) By [21, Theorem 5.22(1)], there exists $0 \neq Y \in D_{\mathfrak{f}}^b(A)$ such that $\text{amp}Y = \text{amp}A$ and $\text{injd}im_A Y < \infty$. Hence Theorem 3.5(iii) yields the statement (2).

(2) \Rightarrow (3) is trivial.

(3) \Rightarrow (1) Since $\text{supRH}om_A(\bar{\kappa}, Y) = \text{lc.d}im_A \text{RH}om_A(\bar{\kappa}, Y) < \infty$, $\text{injd}im_A Y < \infty$ by [12, Corollary 2.31]. Hence Proposition 4.6 implies that A is local Cohen-Macaulay. \square

Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring. For $X \in D(A)$, set

$$\mu_A^n(X) = \text{rank}_{\bar{\kappa}} H^n(\text{RH}om_A(\bar{\kappa}, X)).$$

Vasconcelos [24] conjectured the following result for a local ring.

Proposition 4.8. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$ and set $d = \text{dim}H^0(A)$. If $\mu_A^{d+\text{inf}A}(A) = 1$, then A is local Gorenstein.*

Proof. We use induction on d . If $d = 0$, then $\text{inf}A = \text{depth}A$ by assumption and [21, Propositions 3.3] and so A is local Cohen-Macaulay. Consequently, $\text{injd}im_A A = \text{supRH}om_A(\bar{\kappa}, A) = \text{inf}A$ by [25, Propositions 4.7] and A is local Gorenstein. Now assume that $d > 0$. By [21, Theorem 5.18], there is an A -regular element $\bar{x} \in \bar{\mathfrak{m}}$ such that $\text{lc.d}im A//\bar{x} = \text{dim}H^0(A)/(\bar{x}) = d - 1$. Also by [12, Lemma 2.9], one has an isomorphism

$$\text{RH}om_{A//\bar{x}}(\bar{\kappa}, A//\bar{x})[-1] \simeq \text{RH}om_A(\bar{\kappa}, A)$$

which implies that $\mu_{A//\bar{x}}^{d-1+\text{inf}A//\bar{x}}(A//\bar{x}) = 1$ as $\text{inf}A//\bar{x} = \text{inf}A$. By induction, $A//\bar{x}$ is local Gorenstein, and so $\text{lc.d}im_A \text{RH}om_{A//\bar{x}}(\bar{\kappa}, A//\bar{x}) = \text{lc.d}im_{A//\bar{x}} \text{RH}om_{A//\bar{x}}(\bar{\kappa}, A//\bar{x}) < \infty$ by Corollary 3.6. Thus $\text{lc.d}im_A \text{RH}om_A(\bar{\kappa}, A) < \infty$ and A is local Gorenstein. \square

Corollary 4.9. *Let $(A, \bar{\mathfrak{m}}, \bar{\kappa})$ be a local DG-ring with $\text{amp}A < \infty$ and set $d = \text{dim}H^0(A)$ and $E = E(A, \bar{\mathfrak{m}})$. Then the following are equivalent:*

- (1) A is local Gorenstein;
- (2) $\mu_A^{d+\text{inf}A}(A) = 1$;
- (3) $\text{RH}om_A(E, \bar{\kappa}) \simeq \bar{\kappa}[-d - \text{inf}A]$;
- (4) $\bar{\kappa} \otimes_A^L E \simeq \bar{\kappa}[d + \text{inf}A]$.

Proof. (1) \Leftrightarrow (2) follows from Proposition 4.8 and [12, Theorem 3.9].

(1) \Leftrightarrow (4) By [12, Theorem 3.9] and the isomorphism $\bar{\kappa} \otimes_A^L E \simeq \text{RH}om_A(\text{RH}om_A(\bar{\kappa}, A), E)$.

(3) \Leftrightarrow (4) follows from the isomorphism $\text{RH}om_A(E, \bar{\kappa}) \simeq \text{RH}om_A(\bar{\kappa} \otimes_A^L E, E)$. \square

References

- [1] L.L. Avramov, S.B. Iyengar, A. Neeman, Big Cohen-Macaulay modules, morphisms of perfect complexes, and intersection theorems in local algebra, *Doc. Math.* **23** (2018) 1601–1619.
- [2] H. Bass, On the ubiquity of Gorenstein rings, *Math. Z.* **82** (1963) 8–28.
- [3] I. Bird, L. Shaul, P. Sridhar and J. Williamson, Finitistic dimension over commutative DG-rings, arXiv:2204.06865v2.

- [4] L.W. Christensen, H.-B. Foxby, *Hyperhomological Algebra with Applications to Commutative Rings*, 2006.12.
- [5] M.T. Dibaei and S. Yassemi, Generalized local cohomology and the intersection theorem, *Comm. Algebra* **33** (2005) 899–908.
- [6] H.-B. Foxby, *Hyperhomological algebra and commutative algebra*, Notes in preparation.
- [7] H.-B. Foxby, S.B. Iyengar, Depth and amplitude for unbounded complexes. *Contemp. Math.* 331 (2003) 119–137.
- [8] M. Hochster, *Topics in the homological theory of modules over commutative rings*, CBMS Regional Conf. Ser. Math., vol. 24, Amer. Math. Soc. 1975.
- [9] S.B. Iyengar, Depth for complexes, and intersection theorems, *Math. Z.* **230** (1999) 545–567.
- [10] S.B. Iyengar, L.Q. Ma, K. Schwede, M.E. Walker, Maximal Cohen-Macaulay complexes and their uses: a partial survey. In: *Commutative algebra*, pp. 475–500. Springer, 2021.
- [11] P. Jørgensen, Amplitude inequalities for differential graded modules, *Forum Math.* **22** (2010) 941–948.
- [12] H. Minamoto, Homological identities and dualizing complexes of commutative differential graded algebras, *Israel J. Math.* **242** (2021) 1–36.
- [13] H. Minamoto, Resolutions and homological dimensions of DG-modules commutative DG-algebras, *Israel J. Math.* **245** (2021) 409–454.
- [14] C. Peskine, L. Szpiro, Dimension projective finie et cohomologie locale, *Inst. Hautes Études Sci. Publ. Math.* **42** (1973) 47–119.
- [15] G. Piepmeyer, M.E. Walker, A new proof of the New Intersection Theorem, *J. Algebra* **322** (2009) 3366–3372.
- [16] P. Roberts, Two applications of dualizing complexes over local rings, *Ann. Sci. École Norm. Sup.* **9** (1976) 103–106.
- [17] T. Sharif, S. Yassemi, Special homological dimensions and Intersection Theorem, *Math. Scand.* **96** (2005) 161–168.
- [18] L. Shaul, The twisted inverse image pseudofunctor over commutative DG rings and perfect base change, *Adv. Math.* **320** (2017) 279–328.
- [19] L. Shaul, Injective DG-modules over non-positive DG-rings, *J. Algebra* **515** (2018) 102–156.
- [20] L. Shaul, Completion and torsion over commutative DG rings, *Israel J. Math.* **232** (2019) 531–588.
- [21] L. Shaul, The Cohen-Macaulay property in derived commutative algebra, *Trans. Amer. Math. Soc.* **373** (2020) 6095–6138.
- [22] L. Shaul, Koszul complexes over Cohen-Macaulay rings, *Adv. Math.* **386**, 107806 (2021).
- [23] L. Shaul, Sequence-regular commutative DG-rings. *J. Algebra* **647** (1994) 400–435.
- [24] W.V. Vasconcelos, *Divisor theory in module categories*, North-Holl and Publishing Co., Amsterdam, 1974.
- [25] X.Y. Yang, Y.J. Li, Local Cohen-Macaulay DG-Modules *Appl. Categor. Struct.* 31 (8) (2023) 1–13.
- [26] X.Y. Yang, L. Wang, Homological invariants over non-positive DG-rings, *J. Algebra Appl.* 2050153 (2020) (18 pages).
- [27] S. Yassemi, Generalized section functors, *J. Pure Appl. Algebra* **95** (1994) 103–119.
- [28] A. Yekutieli, Duality and tilting for commutative DG rings, arXiv:1312.6411v4 (2016).