

Asymptotics of Saran's hypergeometric function F_K

Peng-Cheng Hang¹ and Min-Jie Luo^{2*}

¹*Department of Mathematics, School of Mathematics and Statistics,
Donghua University, Shanghai 201620,
People's Republic of China.
E-mail: mathroc618@outlook.com*

²*Department of Mathematics, School of Mathematics and Statistics,
Donghua University, Shanghai 201620,
People's Republic of China.
E-mail: mathwinnie@live.com, mathwinnie@dhu.edu.cn*

Abstract

In this paper, we first establish asymptotic expansions of the Humbert function Ψ_1 for one large variable. The resulting expansions are then used to derive an asymptotic expansion of Saran's hypergeometric function F_K when two of its variables become simultaneously large.

Keywords: Asymptotic expansion, Humbert function, Saran's function.

Mathematics Subject Classification (2010): Primary 33C65, 33C70, Secondary 33C05,

1 Introduction

Saran's hypergeometric function F_K is defined by [22, p. 294, Eq. (2.4)]

$$F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = \sum_{m, n, p=0}^{\infty} \frac{(\alpha_1)_m (\alpha_2)_{n+p} (\beta_1)_{m+p} (\beta_2)_n}{(\gamma_1)_m (\gamma_2)_n (\gamma_3)_p} \frac{x^m y^n z^p}{m! n! p!}, \quad (1.1)$$

where $(x, y, z) \in \mathbb{D}_K := \{(x, y, z) \in \mathbb{C}^3 : |x| < 1, |y| < 1, |z| < (1 - |x|)(1 - |y|)\}$. It has been shown in [16, p. 1] that \mathbb{D}_K is a complete Reinhardt domain. Using the series manipulation technique, we can obtain the following equivalent forms of (1.1):

$$\begin{aligned} & F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] \\ &= \sum_{p=0}^{\infty} \frac{(\alpha_2)_p (\beta_1)_p}{p! (\gamma_3)_p} {}_2F_1 \left[\begin{matrix} \beta_1 + p, \alpha_1 \\ \gamma_1 \end{matrix}; x \right] {}_2F_1 \left[\begin{matrix} \alpha_2 + p, \beta_2 \\ \gamma_2 \end{matrix}; y \right] z^p \\ &= \sum_{m=0}^{\infty} \frac{(\alpha_1)_m (\beta_1)_m}{m! (\gamma_1)_m} F_2[\alpha_2, \beta_2, \beta_1 + m; \gamma_2, \gamma_3; y, z] x^m, \end{aligned} \quad (1.2)$$

*Corresponding author

where ${}_2F_1$ denotes the familiar Gauss hypergeometric function and F_2 denotes the second Appell hypergeometric function (see [24, p. 23, Eq. (3)]). Over the past few years, more and more attention has been given to various properties and applications of this function (see [7], [8], [15] and [16] and the references therein).

As a special function in several complex variables, the asymptotic behaviour of F_K is rather complicated. Recently, Luo and Raina [15] obtain the complete asymptotic formulas for F_K when one of its variables is large. So it is natural to study the asymptotic behaviour of F_K when two variables are large. In fact, as a special case, Garcia and López [10] have deduced the behaviour of F_2 when $x, y \rightarrow \infty$ with $|x|/|y|$ bounded. Their approach is based on the asymptotic method introduced in [14] and a suitable Laplace integral representation for F_2 . In view of the expression (1.2), where y and z appear together as variables of F_2 , we shall study the asymptotic behaviour of F_K when $y, z \rightarrow \infty$ with $|y|/|z|$ bounded. Note also that since

$$F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = F_K[\beta_2, \beta_1, \beta_1, \alpha_2, \alpha_1, \alpha_2; \gamma_2, \gamma_1, \gamma_3; y, x, z],$$

the behaviour for large x, z is equivalent to that for large y, z .

To accomplish this, we need to first study the asymptotics of the Humbert function Ψ_1 defined by [24, p. 26, Eq. (21)]

$$\Psi_1[a, b; c, c'; x, y] := \sum_{m, n=0}^{\infty} \frac{(a)_{m+n} (b)_m}{(c)_m (c')_n} \frac{x^m y^n}{m! n!} \quad (|x| < 1, |y| < \infty). \quad (1.3)$$

The motivation of considering Ψ_1 is that F_K possesses the following Laplace integral representation [23, p. 134, Eq. (3)]:

$$\begin{aligned} & F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] \\ &= \frac{1}{\Gamma(\alpha_2)} \int_0^{\infty} e^{-s} s^{\alpha_2-1} {}_1F_1 \left[\begin{matrix} \beta_2 \\ \gamma_2 \end{matrix}; ys \right] \Psi_1[\beta_1, \alpha_1; \gamma_1, \gamma_3; x, zs] ds, \end{aligned} \quad (1.4)$$

where Ψ_1 is given by (1.3) and ${}_1F_1$ is Kummer's confluent hypergeometric function. The convergence condition of (1.4)

$$\Re(\alpha_2) > 0, \Re((1-x)(1-y)) > \Re(z),$$

given in Saran's paper, is incorrect for it is derived from an incomplete estimate about Ψ_1 (see [23, p. 143, Eq. (3)]). To clarify the convergence condition, we must derive the correct estimate of Ψ_1 .

The paper is organized as follows:

In Section 2, we provide some necessary information about the Gauss hypergeometric function ${}_2F_1$, the Humbert function Ψ_1 and the Kampé de Fériet function $F_{0:1;2}^{1:1;2}$.

Section 3 devotes to the asymptotics of Ψ_1 . We first establish a contour integral representation for Ψ_1 , from which an asymptotic expansion of Ψ_1 for large y is obtained in the left half-plane. Then, by making use of a result of Olver [19], we obtain under certain restrictions the complete asymptotic expansions of Ψ_1 for large y in the right half-plane. Finally, we obtain the asymptotic expansion of Ψ_1 for large x .

Section 4 offers a comprehensive proof of the main result. First we convert the integral representation (1.4) into the form of the Mellin convolution. The desired approximation of F_K is then obtained by applying López's theorem [14] to that Mellin convolution integral.

In Appendix A, we compute an important Mellin transform of a product of ${}_1F_1$ and Ψ_1 . Finding different proofs of it may have independent interest.

2 Preliminaries

2.1 Asymptotics of the Gauss hypergeometric function ${}_2F_1$

Wagner [25, p. 2, Satz 1] proved that

$$\begin{aligned} \frac{1}{\Gamma(c)} {}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix}; z \right] &= \frac{(-az)^{-b}}{\Gamma(c-b)} [1 + o(1)] \\ &+ \frac{e^{\pm \pi i(b-c)}}{\Gamma(b)} (1-z)^{c-a-b} (-az)^{b-c} [1 + o(1)], \text{ as } |a| \rightarrow \infty \end{aligned} \quad (2.1)$$

with b and c ($c \notin \mathbb{Z}_{\leq 0}$) fixed complex numbers, $z \neq 0$ and $|\arg(1-z)| < \pi$. The choice of the sign \pm in the exponent and the determination of the range of $\arg(-az)$ depend on the conditions satisfied by the parameters. For convenience, we present some corollaries of Wagner's formula (2.1) which we shall use in the sequel. Similar analysis of [15, Eq. (12)] gives that for $n \in \mathbb{Z}_{>0}$,

$$\left| {}_2F_1 \left[\begin{matrix} a+n, b \\ c \end{matrix}; z \right] \right| = \mathcal{O}(n^{-\omega} \rho_z^n), \text{ as } n \rightarrow \infty, \quad (2.2)$$

where

$$\omega := \min\{\Re(b), \Re(c-b)\} \text{ and } \rho_z := \max\{1, |1-z|^{-1}\}. \quad (2.3)$$

We use Landau's notation $f = \mathcal{O}(g)$ to mean that $|f| \leq C|g|$ for a suitable positive number C , which may be absolute or depend on various parameters, in which case the dependence may be indicated in a subscript.

We need the following estimate [15, p. 4]:

$$\left| {}_2F_1 \left[\begin{matrix} a+it, b \\ c \end{matrix}; z \right] \right| = \mathcal{O}\left(|t|^{-\omega} e^{|t| \max\{0, \mp \arg(1-z)\}}\right), \quad t \rightarrow \pm\infty, \quad (2.4)$$

where ω is given in (2.3). We also need the behaviour of

$$\left| {}_2F_1 \left[\begin{matrix} a+s, b \\ c \end{matrix}; z \right] \right|, \quad s \in \mathfrak{C},$$

where \mathfrak{C} is the semi-circle $s = (N+1/2)e^{i\theta}$ ($|\theta| \leq \pi/2$) and $N \rightarrow \infty$ is taken through integral values.

Note that for $s \in \mathfrak{C}$,

$$|(1-z)^{-s}| = e^{(N+1/2)(\sin\theta \arg(1-z) - \cos\theta \log|1-z|)}.$$

This together with (2.1) implies that for $s \in \mathfrak{C}$,

$$\left| {}_2F_1 \left[\begin{matrix} a+s, b \\ c \end{matrix}; z \right] \right| = \mathcal{O}\left(N^{-\omega} e^{(N+1/2)(|\sin\theta| \arg(1-z) - \cos\theta \log(1-|z|))}\right), \quad N \rightarrow \infty, \quad (2.5)$$

where $|z| < 1$ and ω is given in (2.3).

2.2 Properties of the Humbert function Ψ_1

We have (see [12, p. 75] and [4, p. 118, Eq. (83)])

$$\Psi_1[a, b; c, c'; x, y] = \sum_{n=0}^{\infty} \frac{(a)_n}{(c')_n} {}_2F_1 \left[\begin{matrix} a+n, b \\ c \end{matrix}; x \right] \frac{y^n}{n!}, \quad |x| < 1, |y| < \infty. \quad (2.6)$$

Taking (2.2) into account, the summand in (2.6) has the order of magnitude

$$\mathcal{O} \left(n^{\Re(a-c')-\omega} \frac{(\rho_x |y|)^n}{n!} \right), \quad n \rightarrow \infty,$$

which implies that the series converges absolutely in the region

$$\mathbb{D}_{\Psi_1} := \{(x, y) \in \mathbb{C}^2 : x \neq 1, |\arg(1-x)| < \pi, |y| < \infty\}.$$

So the series in (2.6) provides an analytic continuation of Ψ_1 to \mathbb{D}_{Ψ_1} .

We also have [4, p. 115, Eq. (38)]

$$\Psi_1[a, b; c, c'; x, y] = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 u^{b-1} (1-u)^{c-b-1} (1-ux)^{-a} {}_1F_1 \left[\begin{matrix} a \\ c' \end{matrix}; \frac{y}{1-ux} \right] du, \quad (2.7)$$

which is valid for $\Re(c) > \Re(b) > 0$ and $(x, y) \in \mathbb{D}_{\Psi_1}$. The integral representation (2.7) follows from the series representation (2.6) and Euler integral representation for ${}_2F_1$ (see [20, p. 388, Eq. (15.6.1)]).

We shall also use the Kummer transformation of Ψ_1 [6, Eq. (2.54)]

$$\Psi_1[a, b; c, c'; x, y] = (1-x)^{-a} \Psi_1 \left[a, c-b; c, c'; \frac{x}{x-1}, \frac{y}{1-x} \right]. \quad (2.8)$$

There are some useful identities about Ψ_1 in the literature (see [4], [6], [12] and [17]). But we still know very little about its asymptotics. Joshi and Arya [13] established a set of bilateral inequalities for Ψ_1 under certain rather restrictive conditions. By using a Taubarian theorem for Laplace transform, Wald and Henkel [26] found the leading asymptotic behaviour of the Humbert functions Φ_2 , Φ_3 and Ξ_2 when the absolute values of the two independent variables become simultaneously large. They also considered Ψ_1 and pointed out that their technique can not handle Ψ_1 (see [26, p. 99]). Therefore, our work in Section 3 may help advance the research in this direction.

2.3 Kampé de Fériet function $F_{0:1;2}^{1:1;2}$

We need the following special case of the Kampé de Fériet function [24, p. 27]:

$$F_{0:1;2}^{1:1;2} \left[\begin{matrix} a_1 : b_1; d_1, d_2 \\ - : c_1; e_1, e_2 \end{matrix}; x, y \right] = \sum_{m,n=0}^{\infty} \frac{(a_1)_{m+n} (b_1)_m (d_1)_n (d_2)_n}{(c_1)_m (e_1)_n (e_2)_n} \frac{x^m}{m!} \frac{y^n}{n!}, \quad |x| + |y| < 1. \quad (2.9)$$

In view of $(a_1)_{n+m} = (a_1)_n (a_1+n)_m$, we have from (2.9) that

$$F_{0:1;2}^{1:1;2} \left[\begin{matrix} a_1 : b_1; d_1, d_2 \\ - : c_1; e_1, e_2 \end{matrix}; x, y \right] = \sum_{n=0}^{\infty} \frac{(a_1)_n (d_1)_n (d_2)_n}{(e_1)_n (e_2)_n} {}_2F_1 \left[\begin{matrix} a_1+n, b_1 \\ c_1 \end{matrix}; x \right] \frac{y^n}{n!}. \quad (2.10)$$

Denote by F_n the summand of the series (2.10). Using the estimate (2.2) gives

$$|F_n| = \mathcal{O} \left(n^{\Re(a_1+d_1+d_2-e_1-e_2)-\omega'-1} (\rho_x |y|)^n \right),$$

where $\omega' := \min\{\Re(b_1), \Re(c_1 - b_1)\}$. Then (2.10) converges absolutely in the region

$$\mathbb{D}_F := \{(x, y) \in \mathbb{C}^2 : x \neq 1, |\arg(1-x)| < \pi, |y| < \min\{1, |1-x|\}\},$$

and thus gives an analytic continuation of $F_{0:1;2}^{1:1;2}$ to \mathbb{D}_F .

Replacing ${}_2F_1$ with its Euler integral representation in (2.10) and then interchanging the order of summation and integration we obtain (initially with $\Re(c_1) > \Re(b_1) > 0$ and $|x| + |y| < 1$)

$$F_{0:1;2}^{1:1;2} \left[\begin{matrix} a_1 : b_1, d_1, d_2 \\ - : c_1, e_1, e_2 \end{matrix}; x, y \right] = \frac{\Gamma(c_1)}{\Gamma(b_1)\Gamma(c_1 - b_1)} \int_0^1 u^{b_1-1} (1-u)^{c_1-b_1-1} \\ \times (1-ux)^{-a_1} {}_3F_2 \left[\begin{matrix} d_1, d_2, a_1 \\ e_1, e_2 \end{matrix}; \frac{y}{1-ux} \right] du, \quad (2.11)$$

where

$$\Re(c_1) > \Re(b_1) > 0, |\arg(1-x)| < \pi \text{ and } \left| \arg\left(1 - \frac{y}{1-ux}\right) \right| < \pi \ (\forall u \in [0, 1]).$$

A simpler condition under which (2.11) holds is given by

$$\Re(c_1) > \Re(b_1) > 0, x < 1, |\arg(1-y)| < \pi \text{ and } \left| \arg\left(1 - \frac{y}{1-x}\right) \right| < \pi. \quad (2.12)$$

The integral (2.11) provides another analytic continuation of $F_{0:1;2}^{1:1;2}$.

2.4 Other functions

The asymptotic expansions of ${}_1F_1$ are given by ([2, p. 193])

$${}_1F_1 \left[\begin{matrix} a \\ c \end{matrix}; z \right] \sim \frac{\Gamma(c)}{\Gamma(c-a)} \sum_{n=0}^{\infty} \frac{(a)_n (1+a-c)_n}{n!} (-z)^{-a-n}, \text{ as } z \rightarrow \infty \text{ with } \Re(z) < 0; \quad (2.13)$$

$${}_1F_1 \left[\begin{matrix} a \\ c \end{matrix}; z \right] \sim \frac{\Gamma(c)}{\Gamma(a)} e^z \cdot \sum_{n=0}^{\infty} \frac{(1-a)_n (c-a)_n}{n!} z^{a-c-n}, \text{ as } z \rightarrow \infty \text{ with } \Re(z) > 0. \quad (2.14)$$

As $z \rightarrow \infty$ in the sector $|\arg(z)| \leq \pi - \delta (< \pi)$,

$$\Gamma(az + b) \sim \sqrt{2\pi} e^{-az} (az)^{az+b-(1/2)}, \quad (2.15)$$

where $a (> 0)$ and $b (\in \mathbb{C})$ are both fixed (see [20, p. 141, Eq. (5.11.7)]). When $y \rightarrow \pm\infty$,

$$|\Gamma(x + iy)| \sim \sqrt{2\pi} |y|^{x-(1/2)} e^{-\pi|y|/2}, \quad (2.16)$$

uniformly for bounded real values of x (see [20, p. 141, Eq. (5.11.9)]).

The Laurent expansion of $\Gamma(z)$ at its pole $z = -n$ ($n \in \mathbb{Z}_{\geq 0}$) is given by ([9, p. 29])

$$\Gamma(z - n) = \frac{(-1)^n}{n!} [z^{-1} + \psi(n+1)] + \mathcal{O}(z), \text{ as } z \rightarrow 0,$$

where $\psi(x)$ is the digamma function. Hence for $m, n \in \mathbb{Z}_{\geq 0}$, we have

$$\lim_{z \rightarrow 0} [a\Gamma(z - m) + b\Gamma(-z - n)] = \frac{(-1)^m}{m!} \psi(m+1) a + \frac{(-1)^n}{n!} \psi(n+1) b, \quad (2.17)$$

with a and b chosen such that this limit exists.

Luo and Raina [15] extended Saran's function F_K to the region

$$\mathbb{V}_K = \left\{ (x, y, z) \in \mathbb{C}^3 : \begin{array}{l} x \neq 1, y \neq 1, |\arg(1-x)| < \pi, |\arg(1-y)| < \pi, \\ z \neq 0, |\arg(-z)| < \pi, \\ \max\{0, \mp \arg(1-x)\} + \max\{0, \mp \arg(1-y)\} < \pi \pm \arg(-z) \end{array} \right\}.$$

3 Asymptotics of Ψ_1

3.1 Asymptotics for large y

Following [15], we shall use the Mellin-Barnes contour integral representation of Ψ_1 to obtain the asymptotic expansion of Ψ_1 for large y . The starting point is (2.6). Define

$$\lambda(n) := \frac{\Gamma(a+n)\Gamma(c')}{\Gamma(a)\Gamma(c'+n)} {}_2F_1 \left[\begin{matrix} a+n, b \\ c \end{matrix}; x \right]$$

so that

$$\Psi_1[a, b; c, c'; x, -y] = \sum_{n=0}^{\infty} \frac{\lambda(n)}{n!} (-y)^n.$$

Using Ramanujan's master theorem [1, p. 107], we formally have the following Mellin transform of the function Ψ_1 with respect to y :

$$\int_0^{\infty} y^{s-1} \Psi_1[a, b; c, c'; x, -y] dy = \Gamma(s) \lambda(-s). \quad (3.1)$$

Applying the inverse Mellin transform to (3.1), we arrive at

$$\Psi_1[a, b; c, c'; x, -y] = \frac{1}{2\pi i} \frac{\Gamma(c')}{\Gamma(a)} \int_{\sigma-i\infty}^{\sigma+i\infty} {}_2F_1 \left[\begin{matrix} a-s, b \\ c \end{matrix}; x \right] \frac{\Gamma(a-s)}{\Gamma(c'-s)} \Gamma(s) y^{-s} ds,$$

where $\sigma := \Re(s)$ satisfies: (i) if $\Re(a) > 0$ then $0 < \sigma < \Re(a)$; (ii) if $\Re(a) \leq 0$ then $\sigma = 0$.

Theorem 3.1. *Let*

$$\mathbb{V}_{\Psi_1} = \left\{ (x, y) \in \mathbb{C}^2 : \begin{array}{l} x \neq 1, y \neq 0, |\arg(1-x)| < \pi, |\arg(-y)| < \pi, \\ \max\{0, \mp \arg(1-x)\} < \frac{\pi}{2} \pm \arg(-y) \end{array} \right\}.$$

Then

$$\Psi_1[a, b; c, c'; x, y] = \frac{1}{2\pi i} \frac{\Gamma(c')}{\Gamma(a)} \int_{\mathcal{L}_{i\sigma\infty}} {}_2F_1 \left[\begin{matrix} a+s, b \\ c \end{matrix}; x \right] \frac{\Gamma(a+s)}{\Gamma(c'+s)} \Gamma(-s) (-y)^s ds, \quad (3.2)$$

where the path of integration $\mathcal{L}_{i\sigma\infty}$, starting at $\sigma - i\infty$ and ending at $\sigma + i\infty$, is a straight line parallel to the imaginary axis intended if necessary to separate the poles of $\Gamma(a+s)$ from the poles of $\Gamma(-s)$.

Proof. The proof will be divided into two parts. Firstly, we prove the integral exists for $(x, y) \in \mathbb{V}_{\Psi_1}$. Secondly, we show by using Cauchy's residue theorem that the integral is equal to Ψ_1 when x and y are appropriately restricted.

Denote by $\Psi(s)$ the integrand in (3.2), namely,

$$\Psi(s) := {}_2F_1 \left[\begin{matrix} a+s, b \\ c \end{matrix}; x \right] \frac{\Gamma(a+s)}{\Gamma(c'+s)} \Gamma(-s) (-y)^s. \quad (3.3)$$

It follows from (2.4) and (2.16) that as $t \rightarrow \pm\infty$,

$$\begin{aligned} \left| \frac{\Gamma(a+\sigma+it)}{\Gamma(c'+\sigma+it)} \Gamma(-\sigma-it) (-y)^{\sigma+it} \right| &= \mathcal{O} \left(|t|^{\Re(a-c')-\sigma-\frac{1}{2}} e^{-\left(\frac{\pi}{2} \pm \arg(-y)\right)|t|} \right), \\ \left| {}_2F_1 \left[\begin{matrix} a+\sigma+it, b \\ c \end{matrix}; x \right] \right| &= \mathcal{O} \left(|t|^{-\omega} e^{|t| \max\{0, \mp \arg(1-x)\}} \right), \end{aligned} \quad (3.4)$$

where $\omega = \min\{\Re(b), \Re(c-b)\}$. Now we can conclude that when

$$\max\{0, \mp \arg(1-x)\} < \frac{\pi}{2} \pm \arg(-y), \quad (3.5)$$

the integrand $|\Psi(s)|$ ($s = \sigma + it$) decays exponentially as $|s| \rightarrow \infty$ along the path $\mathcal{L}_{i\sigma\infty}$.

The inequality (3.5) implies that $|\arg(1-x) + \arg(-y)| < \pi/2$. Thus, if we define

$$\tilde{\mathbb{V}}_{\Psi_1} := \left\{ (x, y) \in \mathbb{C}^2 : |x| < 1, y \neq 0, |\arg(1-x)| + |\arg(-y)| < \frac{\pi}{2}, |y| < \frac{1}{e}(1-|x|) \right\},$$

then we have $\tilde{\mathbb{V}}_{\Psi_1} \subset \mathbb{V}_{\Psi_1}$.

Next we show that when $(x, y) \in \tilde{\mathbb{V}}_{\Psi_1}$, the integral in (3.2) yields the series representation (2.6) of Ψ_1 . We close the contour by the semi-circle \mathcal{C} in the right-hand side of the complex plane. The semi-circle \mathcal{C} is parametrized by $s = (N+1/2)e^{i\theta}$ ($|\theta| \leq \pi/2$) and $N \rightarrow \infty$ through integral values.

From (2.15), we have immediately

$$\left| \frac{\Gamma(a+s)}{\Gamma(c'+s)} \right| = \mathcal{O}\left(N^{\Re(a-c')}\right),$$

as $N \rightarrow \infty$. Since $|\sin \pi s| \geq |\sinh \Im(\pi s)| = |\sinh(\pi(N+1/2)\sin\theta)|$, we have

$$|\Gamma(-s)| = \left| \frac{\pi}{\sin \pi s} \frac{1}{\Gamma(1+s)} \right| = \mathcal{O}\left(N^{-\frac{1}{2}-(N+\frac{1}{2})\cos\theta} e^{(N+\frac{1}{2})(\cos\theta+\theta\sin\theta-\pi|\sin\theta|)}\right),$$

as $N \rightarrow \infty$. Thus

$$\left| \frac{\Gamma(a+s)}{\Gamma(c'+s)} \Gamma(-s) (-y)^s \right| = \mathcal{O}\left(N^{\Re(a-c')-\frac{1}{2}-(N+\frac{1}{2})\cos\theta} e^{(N+\frac{1}{2})(\cos\theta+\theta\sin\theta+\cos\theta\ln|y|-\sin\theta\arg(-y)-\pi|\sin\theta|)}\right),$$

as $N \rightarrow \infty$. Combining this with (2.5), we obtain

$$\begin{aligned} \Psi(s) &= \mathcal{O}\left(N^{\Re(a-c')-\frac{1}{2}-(N+\frac{1}{2})\cos\theta} e^{(N+\frac{1}{2})(\cos\theta+|\theta||\sin\theta|+\cos\theta\ln|y|+|\sin\theta||\arg(-y)|-\pi|\sin\theta|)}\right) \\ &\quad \cdot \mathcal{O}\left(N^{-\omega} e^{(N+1/2)(|\sin\theta||\arg(1-x)|-\cos\theta\ln(1-|x|))}\right) \\ &= \mathcal{O}\left(N^{\Re(a-c')-\omega-\frac{1}{2}-(N+\frac{1}{2})\cos\theta} e^{(N+\frac{1}{2})(\delta_1|\sin\theta|+\delta_2\cos\theta)}\right), \end{aligned}$$

where $\delta_1 := |\theta| + |\arg(-y)| + |\arg(1-x)| - \pi < 0$ and $\delta_2 := 1 + \ln|y| - \ln(1-|x|) < 0$. It follows that the integral on the semi-circle \mathcal{C} tends to zero as $N \rightarrow \infty$.

Finally, by Cauchy's residue theorem and the assertion that ([2, p. 7])

$$\operatorname{Res}_{s=-n} \Gamma(-s) = \frac{(-1)^n}{n!},$$

we get the representation (2.6). This completes the proof. \square

Corollary 3.2. *When $(x, y) \in \mathbb{V}_{\Psi_1}$ and $|y| \rightarrow \infty$, then*

$$\Psi_1[a, b; c, c'; x, y] \sim \frac{\Gamma(c')}{\Gamma(c'-a)} \sum_{n=0}^{\infty} {}_2F_1 \left[\begin{matrix} -n, b \\ c \end{matrix}; x \right] \frac{(a)_n (1+a-c')_n}{n!} (-y)^{-n-a}. \quad (3.6)$$

Proof. Let I denote the right-hand side of (3.2), and let $\Psi(s)$ denote the integrand as defined in (3.3). Choose the positive integer $M \geq \max\{1, \Re(-a)\}$ and shift the integration contour to the left (which is permissible on account of the exponential decay of the integrand).

Note that $s = -a - n$ ($n \in \mathbb{Z}_{\geq 0}$) are the only poles of $\Psi(s)$ because the principal branch of ${}_2F_1[a, b; c; z]$ is an entire function in a (see [20, p. 384]). Therefore,

$$\operatorname{Res}_{s=-a-n} \Psi(s) = {}_2F_1 \left[\begin{matrix} -n, b \\ c \end{matrix}; x \right] \frac{\Gamma(a+n)}{\Gamma(c'-a-n)} \frac{(-1)^n}{n!} (-y)^{-a-n}.$$

By Cauchy's residue theorem, we obtain

$$\frac{\Gamma(a)}{\Gamma(c')} I = \sum_{n=0}^M {}_2F_1 \left[\begin{matrix} -n, b \\ c \end{matrix}; x \right] \frac{\Gamma(a+n)}{\Gamma(c'-a-n)} \frac{(-1)^n}{n!} (-y)^{-n-a} + R_M(y), \quad (3.7)$$

where

$$R_M(y) := \frac{1}{2\pi i} \int_{C_M} \Psi(s) ds \quad (3.8)$$

and C_M denotes the vertical line $\Re(s) = \Re(-a) - M - 1/2$. Using (3.4) can get

$$|\Psi(s)| = \mathcal{O} \left(|y|^{\Re(-a) - M - \frac{1}{2}} |t|^{\Re(2a - c') + M - \omega} e^{-\chi|t|} \right), \quad t = \Im(s),$$

where ω is given in (2.3) and

$$\chi := \frac{\pi}{2} \pm \arg(-y) - \max\{0, \mp \arg(1-x)\} > 0.$$

Thus the integral (3.8) converges when $(x, y) \in \mathbb{V}_{\Psi_1}$. It follows that

$$R_M(y) = \mathcal{O}_M \left(|y|^{\Re(-a) - M - \frac{1}{2}} \right),$$

which shows that (3.7) is an asymptotic expansion of Poincaré type. Simplifying (3.7) with the help of the identities $\Gamma(a+n) = \Gamma(a) (a)_n$ and

$$\Gamma(c' - a - n) = (-1)^n \frac{\Gamma(c' - a)}{(1 + a - c')_n}$$

gives the desired expansion (3.6). □

Remark 3.3. *The region \mathbb{V}_{Ψ_1} has an equivalent form:*

$$\mathbb{V}_{\Psi_1} = \left\{ (x, y) \in \mathbb{C}^2 : x \neq 1, y \neq 0, |\arg(-y)| < \frac{\pi}{2}, |\arg(1-x) + \arg(-y)| < \frac{\pi}{2} \right\},$$

which suggests that (3.6) is an asymptotic expansion for y on the left half-plane. If we further let $x = 0$ in (3.6), we get the expansion of ${}_1F_1[a; c'; y]$ for $\Re(y) < 0$, viz. the formula (2.13).

In the rest of this subsection, we establish the asymptotics of Ψ_1 for y on the right half-plane by applying Olver's result [19, Theorem 1], which adapts Laplace's method to the complex parameter.

We restrict ourselves to the following three cases:

- (i) $\Re(c) > \Re(b) > 0, x = 0, y \in \mathbb{S}$;
- (ii) $\Re(c) > \Re(b) > 0, x < 0, y \in \mathbb{S}$;
- (iii) $\Re(c) > \Re(b) > 0, 0 < x < 1, y \in \mathbb{S}$,

where $\mathbb{S} := \{y \in \mathbb{C} : |\arg y| \leq \frac{\pi}{2} - \delta\}$ with $\delta \in (0, \frac{\pi}{2})$ fixed.

3.1.1 Case (i)

In this case, the asymptotics follows from

$$\Psi_1[a, b; c, c'; 0, y] = {}_1F_1\left[\begin{matrix} a \\ c' \end{matrix}; y\right]$$

and (2.14).

3.1.2 Case (ii)

In this case, if $u \in (0, 1)$ then $1 - ux \in (1, 1 - x)$. Denote

$$\Psi_1 \equiv \Psi_1[a, b; c, c'; x, y], \quad g(u) := u^{b-1} (1-u)^{c-b-1} (1-ux)^{-a}.$$

The expansion (2.14) implies that there exists a constant $C > 0$ such that

$$\left| {}_1F_1\left[\begin{matrix} a \\ c' \end{matrix}; \frac{y}{1-ux}\right] - h(u) \right| \leq CR(u), \quad y \in \mathbb{S}, \quad (3.9)$$

where

$$h(u) = \frac{\Gamma(c')}{\Gamma(a)} e^{\frac{y}{1-ux}} \left(\frac{y}{1-ux}\right)^{a-c'}, \quad R(u) = e^{\frac{\Re(y)}{1-ux}} \left(\frac{|y|}{1-ux}\right)^{\Re(a-c')-1}.$$

For $\Re(\alpha) > 0$ and $\Re(\beta) > 0$, we define

$$I_y(\alpha, \beta, \gamma) := \int_0^1 q(u) e^{-yp(u)} du,$$

where $q(u) := u^{\alpha-1} (1-u)^{\beta-1} (1-ux)^\gamma$ and $p(u) := 1/(ux-1)$. Obviously, $p(u)$ and $q(u)$ satisfy the conditions [19, p. 229]. Then applying Olver's result to $I_y(\alpha, \beta, \gamma)$, we obtain

$$I_y(\alpha, \beta, \gamma) \sim \Gamma(\alpha) y^{-\alpha} e^y \cdot \sum_{n=0}^{\infty} \frac{(\alpha)_n \lambda_n}{y^n}, \quad y \rightarrow \infty, \quad y \in \mathbb{S}, \quad (3.10)$$

where $\lambda_0 = (-x)^{-\alpha}$ and

$$\lambda_n := \operatorname{Res}_{u=0} \frac{q(u)}{(p(u)+1)^{\alpha+n}}, \quad n = 0, 1, 2, \dots. \quad (3.11)$$

Take the leading term of the expansion (3.10) and then

$$I_y(\alpha, \beta, \gamma) = \Gamma(\alpha) (-x)^{-\alpha} y^{-\alpha} e^y [1 + \mathcal{O}(y^{-1})], \quad y \rightarrow \infty, \quad y \in \mathbb{S}.$$

Using (2.7) and (3.9), we get that as $y \rightarrow \infty$ in \mathbb{S} ,

$$\begin{aligned} \Psi_1 &= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 g(u) h(u) du + \mathcal{O}\left(\int_0^1 |g(u)| R(u) du\right) \\ &= \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(b)\Gamma(c-b)} y^{a-c'} I_y(b, c-b, c'-2a) + \mathcal{O}\left(|y|^{\Re(a-c')-1} I_{\Re(y)}(\Re(b), \Re(c-b), \Re(c'-2a)+1)\right) \\ &= \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} (-x)^{-b} y^{a-b-c'} e^y + \mathcal{O}_x\left(|y|^{\Re(a-b-c')-1} e^{\Re(y)}\right). \end{aligned}$$

Hence, we have the following theorem.

Theorem 3.4. *When $x < 0$ and $y \rightarrow \infty$ with $y \in \mathbb{S}$, then*

$$\Psi_1[a, b; c, c'; x, y] \sim \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} (-x)^{-b} y^{a-b-c'} e^y.$$

3.1.3 Case (iii)

In this case, the pair

$$(x', y') = \left(\frac{x}{x-1}, \frac{y}{1-x} \right)$$

satisfies **Case (ii)**. We can apply the Kummer transformation (2.8) to the pair (x', y') . Hence we get the following corollary.

Corollary 3.5. *When $0 < x < 1$ and $y \rightarrow \infty$ with $y \in \mathbb{S}$, then*

$$\Psi_1[a, b; c, c'; x, y] \sim \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(b)} x^{b-c} (1-x)^{c'+2(c-a-b)} y^{a+b-c-c'} e^{\frac{y}{1-x}}. \quad (3.12)$$

We can obtain the complete asymptotic expansion of Ψ_1 in **Case (ii)**. Indeed, in (3.10), take $h(u)$ as the first $N+1$ terms of (2.14) and take $R(u)$ as the corresponding reminder term. Using (2.7) and (3.10), we get that as $y \rightarrow \infty$ in \mathbb{S} ,

$$\begin{aligned} \Psi_1 &= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 g(u)h(u)du + \mathcal{O}\left(\int_0^1 |g(u)|R(u)du\right) \\ &= \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(b)\Gamma(c-b)} y^{a-c'} \sum_{n=0}^N \frac{(1-a)_n(c'-a)_n}{n!y^n} I_y(b, c-b, n+c'-2a) + \mathcal{O}_x\left(|y|^{\Re(a-b-c')-N-1} e^{\Re(y)}\right) \\ &= \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} y^{a-b-c'} e^y \sum_{n=0}^N \frac{(1-a)_n(c'-a)_n}{n!y^n} \sum_{n=0}^N \frac{(b)_n \lambda_n}{y^n} + \mathcal{O}_x\left(|y|^{\Re(a-b-c')-N-1} e^{\Re(y)}\right) \\ &= \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} y^{a-b-c'} e^y \left\{ \sum_{n=0}^N \psi_n y^{-n} + \mathcal{O}_x(|y|^{-N-1}) \right\}, \end{aligned}$$

where

$$\psi_n := \sum_{k=0}^n \frac{(1-a)_k(c'-a)_k}{k!} (b)_{n-k} \lambda_{n-k}, \quad n = 0, 1, 2, \dots, \quad (3.13)$$

and $\lambda_n = \lambda_n(x)$ is given by (3.11). In particular, $\psi_0 := (-x)^{-b}$. Hence, we get the following theorem.

Theorem 3.6. *When $x < 0$ and $y \rightarrow \infty$ with $y \in \mathbb{S}$, the complete expansion of Ψ_1 is given by*

$$\Psi_1[a, b; c, c'; x, y] \sim \frac{\Gamma(c)\Gamma(c')}{\Gamma(a)\Gamma(c-b)} y^{a-b-c'} e^y \sum_{n=0}^{\infty} \psi_n y^{-n}, \quad (3.14)$$

where $\psi_n = \psi_n(x)$ is given by (3.13). In particular, $\psi_0 = (-x)^{-b}$.

Remark 3.7. *The coefficients in Laplace's method, such as λ_n in (3.11), were explicitly determined by G. Nemes; see [18, p. 477, Eq. (2.6)]. To get the complete asymptotic expansion of Ψ_1 in **Case (iii)**, we may insert (3.14) into (2.8), but the resulting expansion is terribly complicated.*

3.2 Asymptotics for large x

In this subsection, we consider the behaviour of Ψ_1 for large x and derive the complete asymptotic expansion without using the contour integrals and the residue theorem.

The connection formula for the Gauss hypergeometric function is given by ([20, p. 390, Eq. (15.8.2)])

$$\begin{aligned} {}_2F_1 \left[\begin{matrix} a, b \\ c \end{matrix}; z \right] &= f_c(b, a) (-z)^{-a} {}_2F_1 \left[\begin{matrix} 1-c+a, a \\ 1-b+a \end{matrix}; \frac{1}{z} \right] \\ &\quad + f_c(a, b) (-z)^{-b} {}_2F_1 \left[\begin{matrix} 1-c+b, b \\ 1-a+b \end{matrix}; \frac{1}{z} \right], \quad |\arg(-z)| < \pi. \end{aligned} \quad (3.15)$$

where

$$f_\gamma(a, b) := \frac{\Gamma(\gamma)\Gamma(a-b)}{\Gamma(a)\Gamma(\gamma-b)}. \quad (3.16)$$

Applying (3.15) to (2.6), we get

$$\Psi_1[a, b; c, c'; x, y] = f_c(b, a) (-x)^{-a} \mathbf{U}_1(x, y) + f_c(a, b) (-x)^{-b} \mathbf{U}_2(x, y), \quad (3.17)$$

where

$$\begin{aligned} \mathbf{U}_1(x, y) &:= \sum_{n=0}^{\infty} \frac{(a)_n}{n!(c')_n} \frac{(1-c+a)_n}{(1-b+a)_n} {}_2F_1 \left[\begin{matrix} 1-c+a+n, a+n \\ 1-b+a+n \end{matrix}; \frac{1}{x} \right] \left(-\frac{y}{x}\right)^n, \\ \mathbf{U}_2(x, y) &:= \sum_{n=0}^{\infty} \frac{(a-b)_n}{n!(c')_n} {}_2F_1 \left[\begin{matrix} 1-c+b, b \\ 1-a-n+b \end{matrix}; \frac{1}{x} \right] y^n \end{aligned}$$

and $f_\gamma(a, b)$ is given by (3.16).

The series \mathbf{U}_1 is a special case of the Kampé de Fériet function. In fact, we have

$$\begin{aligned} \mathbf{U}_1(x, y) &= \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(a)_{n+m} (1-c+a)_{n+m}}{(1-b+a)_{n+m} (c')_n m! n!} \frac{(-y)^n}{x^{n+m}} \\ &= F_{1:0;1}^{2:0;0} \left[\begin{matrix} a, 1-c+1: -; - \\ 1-b+a: -; c' \end{matrix}; \frac{1}{x}, -\frac{y}{x} \right], \end{aligned}$$

which suggests that the double series converges absolutely for $|x| > 1$ and $|y| < \infty$. Further calculation gives

$$\begin{aligned} \mathbf{U}_1(x, y) &= \sum_{k=0}^{\infty} \frac{(a)_k (1-c+a)_k}{(1-b+a)_k k!} \frac{1}{x^k} \sum_{n=0}^k \frac{(-k)_n}{(c')_n} \frac{y^n}{n!} \\ &= \sum_{k=0}^{\infty} {}_1F_1 \left[\begin{matrix} -k \\ c' \end{matrix}; y \right] \frac{(a)_k (1-c+a)_k}{(1-b+a)_k k!} \frac{1}{x^k}, \quad |x| > 1, |y| < \infty. \end{aligned} \quad (3.18)$$

For the series \mathbf{U}_2 , we first note that

$$\begin{aligned} \mathbf{U}_2(x, y) &= \sum_{\ell, n=0}^{\infty} \frac{(1-c+b)_\ell (b)_\ell}{(1-a+b-n)_\ell \ell!} \frac{(a-b)_n}{(c')_n} \frac{1}{x^\ell} \frac{y^n}{n!} \\ &= \sum_{n, \ell=0}^{\infty} (a-b)_{n-\ell} \frac{(1-c+b)_\ell (b)_\ell}{(c')_n} \frac{(-1/x)^\ell}{\ell!} \frac{y^n}{n!} \\ &= H_{11} \left[\begin{matrix} a-b, b, 1-c+b, c' \\ a-b, b, 1-c+b, c' \end{matrix}; y, -\frac{1}{x} \right], \quad |x| > 1, |y| < \infty, \end{aligned}$$

where H_{11} defined by

$$H_{11}[a, b, c, d; x, y] := \sum_{m, n=0}^{\infty} \frac{(a)_{m-n} (b)_n (c)_n}{(d)_m} \frac{x^m}{m!} \frac{y^n}{n!}, \quad |x| < \infty, |y| < 1$$

is one member of *Horn's list* (see [9, Eq. (39), p. 227] and [5]). In addition, we have

$$\begin{aligned} \mathbf{U}_2(x, y) &= \sum_{\ell, n=0}^{\infty} \frac{(1-c+b)_\ell (b)_\ell}{(1-a+b)_\ell \ell!} \frac{1}{x^\ell} \cdot \frac{(a-b-\ell)_n}{(c')_n} \frac{y^n}{n!} \\ &= \sum_{\ell=0}^{\infty} {}_1F_1 \left[\begin{matrix} a-b-\ell \\ c' \end{matrix}; y \right] \frac{(1-c+b)_\ell (b)_\ell}{(1-a+b)_\ell \ell!} \frac{1}{x^\ell}, \quad |x| > 1, |y| < \infty. \end{aligned} \quad (3.19)$$

The previous derivations are summarized in the following theorem.

Theorem 3.8. *When $|\arg(-x)| < \pi$ and $|x| \rightarrow \infty$, we have*

$$\Psi_1[a, b; c, c'; x, y] \sim \mathfrak{f}_c(b, a) (-x)^{-a} + \mathfrak{f}_c(a, b) (-x)^{-b} {}_1F_1 \left[\begin{matrix} a-b \\ c' \end{matrix}; y \right],$$

where \mathfrak{f}_γ is given by (3.16). The complete expansion is obtained by combining (3.17), (3.18) and (3.19).

4 Asymptotics of F_K

Using (2.13) and (3.6), we get a sufficient condition of convergence for the integral (1.4):

$$\Re(\alpha_2) > 0, \quad |\arg(-y)| < \frac{\pi}{2} \quad \text{and} \quad (x, z) \in \mathbb{V}_{\Psi_1}.$$

Under this condition, we also have $(x, y, z) \in \mathbb{V}_K$. So the integral (1.4) makes sense.

After the change of the variables $s = \tilde{y}t$, with $\tilde{y} = 1/|y|$, and defining $\gamma := \tilde{z}/\tilde{y} := |y|/|z|$, the integral (1.4) reads

$$\begin{aligned} F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] \\ = \frac{\tilde{y}}{\Gamma(\alpha_2)} \int_0^\infty e^{-\tilde{y}t} (\tilde{y}t)^{\alpha_2-1} {}_1F_1 \left[\begin{matrix} \beta_2 \\ \gamma_2 \end{matrix}; t e^{i\alpha} \right] \Psi_1[\beta_1, \alpha_1; \gamma_1, \gamma_3; x, t e^{i\beta}/\gamma] dt, \end{aligned} \quad (4.1)$$

where α and β are the principal arguments of the respective variables y and z : $y = |y|e^{i\alpha}$ and $z = |z|e^{i\beta}$.

We are interested in the approximation of F_K for large y and z uniformly in γ with

$$0 < \varrho_1 < \gamma < \varrho_2 < \infty.$$

This means that \tilde{y} is small, γ is bounded, and the remaining parameters $\alpha_1, \alpha_2, \beta_1, \beta_2, \gamma_1, \gamma_2, \gamma_3, x$ are fixed. In order to apply López's theorem [14], we also require that

$$\alpha_2 > 0 \quad \text{and} \quad \beta_1 + \beta_2 > 0.$$

Throughout this section, we define

$$f(t) := {}_1F_1 \left[\begin{matrix} \beta_2 \\ \gamma_2 \end{matrix}; t e^{i\alpha} \right] \Psi_1[\beta_1, \alpha_1; \gamma_1, \gamma_3; x, t e^{i\beta}/\gamma], \quad h(t) := e^{-t} t^{\alpha_2-1}. \quad (4.2)$$

Then the integral (4.1) is of the form of the Mellin integrals considered in [14]:

$$F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = \frac{\tilde{y}}{\Gamma(\alpha_2)} \int_0^\infty h(\tilde{y}t) f(t) dt. \quad (4.3)$$

The functions $f(t)$ and $h(t)$ are locally integrable on $[0, \infty)$, a property required by the technique introduced in [14] to obtain an asymptotic expansion of the integral (4.3) for small \tilde{y} . That technique also requires the asymptotic behaviours of these functions given below and is valid for $\Re(y) < 0$ and $\Re(z) < 0$ ($\pi/2 < |\alpha|, |\beta| < \pi$):

(i) A power asymptotic expansion of $h(t)$ at $t = 0$:

$$h(t) = \sum_{k=0}^{m-1} A_k t^{k+\alpha_2-1} + h_m(t), \quad A_k := \frac{(-1)^k}{k!},$$

with $h_m(t) = \mathcal{O}(t^{m+\alpha_2-1})$ when $t \rightarrow 0^+$;

(ii) An inverse power asymptotic expansion of $f(t)$ at $t = \infty$:

$$f(t) = \sum_{k=0}^{n-1} B_k t^{-k-\beta_1-\beta_2} + f_n(t),$$

with $B_k = (-y\tilde{y})^{-k-\beta_1-\beta_2} \tilde{B}_k$,

$$\begin{aligned} \tilde{B}_k := & \frac{\Gamma(\gamma_2)\Gamma(\gamma_3)}{\Gamma(\gamma_2-\beta_2)\Gamma(\gamma_3-\beta_1)} \left(\frac{y}{z}\right)^{\beta_1} \\ & \times \sum_{j=0}^k \frac{(\beta_2)_j (1+\beta_2-\gamma_2)_j (\beta_1)_{k-j} (1+\beta_1-\gamma_3)_{k-j}}{j!(k-j)!} {}_2F_1 \left[\begin{matrix} j-k, \alpha_1 \\ \gamma_1 \end{matrix}; x \right] \left(\frac{y}{z}\right)^{k-j}, \end{aligned}$$

and $f_n(t) = \mathcal{O}(t^{-\beta_1-\beta_2-n})$ when $t \rightarrow \infty$. The above expansion follows from (2.13) and (3.6);

(iii) $h(t) = \mathcal{O}(t^{-N})$ when $t \rightarrow \infty$ for any positive integer N and $f(t) = 1 + \mathcal{O}(t)$ when $t \rightarrow 0^+$.

Following [14], we denote by $\mathfrak{M}[g; s]$ the Mellin transform of g . Clearly, $\mathfrak{M}[h; s] = \Gamma(s + \alpha_2 - 1)$. Using (A.3), we can readily get

$$\begin{aligned} \mathfrak{M}[f; s] = & (-y\tilde{y})^{-s} \left\{ C_1^*(s) \left(\frac{y}{z}\right)^{s-\beta_2} F_{0:1;2}^{1:1;2} \left[\begin{matrix} \beta_1 + \beta_2 - s \\ \beta_2 + 1 - s, \beta_2 + \gamma_3 - s \end{matrix}; \begin{matrix} \alpha_1 \\ \gamma_1 \end{matrix}; x, -\frac{z}{y} \right] \right. \\ & \left. + C_2^*(s) F_{0:1;2}^{1:1;2} \left[\begin{matrix} \beta_1 \\ - \end{matrix}; \begin{matrix} \alpha_1 \\ \gamma_1 \end{matrix}; s, s - \gamma_2 + 1 \\ \gamma_3, s - \beta_2 + 1 \end{matrix}; x, -\frac{z}{y} \right] \right\}, \end{aligned}$$

where

$$C_1^*(s) := \frac{\Gamma(\gamma_2)\Gamma(\gamma_3)\Gamma(s-\beta_2)\Gamma(\beta_1+\beta_2-s)}{\Gamma(\beta_1)\Gamma(\gamma_2-\beta_2)\Gamma(\beta_2+\gamma_3-s)}, \quad C_2^*(s) := \frac{\Gamma(\gamma_2)\Gamma(s)\Gamma(\beta_2-s)}{\Gamma(\beta_2)\Gamma(\gamma_2-s)}, \quad (4.4)$$

$\Re(y) < 0$, $(x, z) \in \mathbb{V}_{\Psi_1}$ and $\rho_x < |y|/|z|$.

Let us summarize the conditions needed in our further derivation as follows

$$\alpha_2 > 0, \beta_1 + \beta_2 > 0, |\arg(-y)| < \frac{\pi}{2}, (x, z) \in \mathbb{V}_{\Psi_1} \text{ and } \rho_x < \frac{|y|}{|z|} < \rho_2 < \infty. \quad (4.5)$$

4.1 The nonlogarithmic case

When $\beta_1 + \beta_2 - \alpha_2 \notin \mathbb{Z}$, from [14, Theorem 1] we have the following asymptotic expansion of F_K for small \tilde{y} . For any $n \in \mathbb{Z}_{>0}$,

$$\begin{aligned} & \Gamma(\alpha_2) F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] \\ &= \sum_{k=0}^{n-1} B_k \mathfrak{M}[h; 1 - k - \beta_1 - \beta_2] \tilde{y}^{k+\beta_1+\beta_2} + \sum_{k=0}^{m-1} A_k \mathfrak{M}[f; k + \alpha_2] \tilde{y}^{k+\alpha_2} + R_n(\tilde{y}), \end{aligned} \quad (4.6)$$

where $m = n + \lfloor 1 - \alpha_2 + \beta_1 + \beta_2 \rfloor$ and the reminder $R_n(\tilde{y}) = \mathcal{O}(\tilde{y}^{n+\beta_1+\beta_2})$ when $\tilde{y} \rightarrow 0$.

Computing the Mellin transforms involved in (4.6), we obtain that, under the condition (4.5),

$$F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = \sum_{k=0}^{n-1} \frac{\widehat{B}_k(y/z)}{y^{k+\beta_1+\beta_2}} + \sum_{k=0}^{m-1} \frac{\widehat{A}_k(y/z)}{y^{k+\alpha_2}} + R_n(y), \quad (4.7)$$

with

$$\begin{aligned} \widehat{A}_k(y/z) &:= \frac{(-1)^{-k-\alpha_2}}{\Gamma(\alpha_2)} \Gamma(\gamma_2) A_k \left\{ \frac{\Gamma(\gamma_3) \Gamma(k + \alpha_2 - \beta_2) \Gamma(\beta_1 + \beta_2 - \alpha_2 - k)}{\Gamma(\beta_1) \Gamma(\gamma_2 - \beta_2) \Gamma(\beta_2 + \gamma_3 - \alpha_2 - k)} \left(\frac{y}{z} \right)^{k+\alpha_2-\beta_2} \right. \\ &\quad \times F_{0:1;2}^{1:1;2} \left[\begin{matrix} \beta_1 + \beta_2 - \alpha_2 - k : \alpha_1; & \beta_2, \beta_2 - \gamma_2 + 1 \\ \beta_2 + 1 - \alpha_2 - k, \beta_2 + \gamma_3 - \alpha_2 - k; & x, -\frac{z}{y} \end{matrix} \right] \\ &\quad \left. + \frac{\Gamma(k + \alpha_2) \Gamma(\beta_2 - \alpha_2 - k)}{\Gamma(\beta_2) \Gamma(\gamma_2 - \alpha_2 - k)} F_{0:1;2}^{1:1;2} \left[\begin{matrix} \beta_1 : \alpha_1; & k + \alpha_2, k + \alpha_2 - \gamma_2 + 1 \\ \text{---} : \gamma_1; & \gamma_3, k + \alpha_2 - \beta_2 + 1 \end{matrix} ; x, -\frac{z}{y} \right] \right\} \end{aligned} \quad (4.8)$$

and

$$\widehat{B}_k(y/z) := \frac{\Gamma(\alpha_2 - \beta_1 - \beta_2 - k)}{\Gamma(\alpha_2)} (-1)^{-k-\beta_1-\beta_2} \widetilde{B}_k. \quad (4.9)$$

The reminder $R_n(y)$ verifies $R_n(y) = \mathcal{O}(y^{-n-\beta_1-\beta_2})$ when $|y| \rightarrow \infty$ uniformly in $|z|$ with y/z bounded.

4.2 The logarithmic case I

When $1 + \beta_1 + \beta_2 - \alpha_2 \in \mathbb{Z}_{>0}$, from [14, Theorem 1] we have the following asymptotic expansion for small \tilde{y} . For any $n \in \mathbb{Z}_{>0}$,

$$\begin{aligned} & \Gamma(\alpha_2) F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = \sum_{k=0}^{\beta_1+\beta_2-\alpha_2-1} A_k \mathfrak{M}[f; k + \alpha_2] \tilde{y}^{k+\alpha_2} \\ & \quad + \sum_{k=0}^{n-1} \tilde{y}^{k+\beta_1+\beta_2} \left\{ -B_k A_{k-\alpha_2+\beta_1+\beta_2} \log \tilde{y} + \lim_{s \rightarrow 0} \left(B_k \mathfrak{M}[h; s + 1 - k - \beta_1 - \beta_2] \right. \right. \\ & \quad \left. \left. + A_{k-\alpha_2+\beta_1+\beta_2} \mathfrak{M}[f; s + k + \beta_1 + \beta_2] \right) \right\} + R_n(\tilde{y}), \end{aligned} \quad (4.10)$$

where $m = n - \alpha_2 + \beta_1 + \beta_2$ and the reminder $R_n(\tilde{y}) = \mathcal{O}(\tilde{y}^{n+\beta_1+\beta_2} \log \tilde{y})$ when $\tilde{y} \rightarrow 0$.

To compute the limit involved in (4.10), we follow [14, Eq. (9) and (10)]. Write

$$\mathfrak{M}[f; s + k + \beta_1 + \beta_2] = C_f - \frac{B_k}{s}, \quad \mathfrak{M}[h; s + 1 - k - \beta_1 - \beta_2] = C_h + \frac{A_{k-\alpha_2+\beta_1+\beta_2}}{s}, \quad \text{as } s \rightarrow 0.$$

Thus, that limit exists. Further, observing that only the factor $\Gamma(\beta_1 + \beta_2 - s)$ of $C_1^*(s)$ (see (4.4)) and $\mathfrak{M}[h; s]$ have poles at $1 - k - \beta_1 - \beta_2$, that limit can be reduced to the form as described in (2.17). Therefore, using (2.17) and after some calculations, we obtain that, under the condition (4.5),

$$\begin{aligned} F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] \\ = \sum_{k=0}^{\beta_1 + \beta_2 - \alpha_2 - 1} \frac{\widehat{A}_k(y/z)}{y^{k+\alpha_2}} + \sum_{k=0}^{n-1} \frac{\widehat{C}_k(y/z) + B_k^\times(y/z) \log|y|}{y^{k+\beta_1+\beta_2}} + R_n(y), \end{aligned} \quad (4.11)$$

with $\widehat{A}_k(y/z)$ given in (4.8),

$$B_k^\times(y/z) := \frac{(-1)^{-k-\beta_1-\beta_2}}{\Gamma(\alpha_2)} \widetilde{B}_k A_{k-\alpha_2+\beta_1+\beta_2} \quad (4.12)$$

and

$$\begin{aligned} \widehat{C}_k(y/z) := & \frac{(-1)^{-k-\beta_1-\beta_2}}{\Gamma(\alpha_2)} \left\{ \frac{\widetilde{B}_k (-1)^{k-\alpha_2+\beta_1+\beta_2} \psi(k - \alpha_2 + \beta_1 + \beta_2 + 1)}{(k - \alpha_2 + \beta_1 + \beta_2)!} \right. \\ & + A_{k-\alpha_2+\beta_1+\beta_2} \Gamma(\gamma_2) \left[\frac{(y/z)^{k+\beta_1} (-1)^k \psi(k+1) \Gamma(\gamma_3) \Gamma(k+\beta_1)}{\Gamma(\beta_1) \Gamma(\gamma_2 - \beta_2) \Gamma(\gamma_3 - \beta_1 - k) k!} \right. \\ & \times F_{0:1;2}^{1:1;2} \left[\begin{matrix} -k: & \alpha_1; & \beta_2, \beta_2 - \gamma_2 + 1 \\ \text{---} & \gamma_1; & 1 - \beta_1 - k, \gamma_3 - \beta_1 - k; \end{matrix} x, -\frac{z}{y} \right] \\ & \left. \left. + \frac{\Gamma(-\beta_1 - k) \Gamma(k + \beta_1 + \beta_2)}{\Gamma(\beta_2) \Gamma(\gamma_2 - \beta_1 - \beta_2 - k)} F_{0:1;2}^{1:1;2} \left[\begin{matrix} \beta_1: & \alpha_1; & k + \beta_1 + \beta_2, k + \beta_1 + \beta_2 - \gamma_2 + 1; \\ \text{---} & \gamma_1; & \gamma_3, k + \beta_1 + 1 \end{matrix} x, -\frac{z}{y} \right] \right] \right\}, \end{aligned} \quad (4.13)$$

where $\psi(s)$ is the digamma function.

4.3 The logarithmic case II

When $\alpha_2 - \beta_1 - \beta_2 \in \mathbb{Z}_{>0}$, from [14, Theorem 1] we have the following asymptotic expansion for small \tilde{y} . For any $n \in \mathbb{Z}_{>0}$,

$$\begin{aligned} \Gamma(\alpha_2) F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = & \sum_{k=0}^{\alpha_2 - \beta_1 - \beta_2 - 1} B_k \mathfrak{M}[h; 1 - k - \beta_1 - \beta_2] \tilde{y}^{k+\beta_1+\beta_2} \\ & + \sum_{k=0}^{m-1} \tilde{y}^{k+\alpha_2} \left\{ -A_k B_{k+\alpha_2-\beta_1-\beta_2} \log \tilde{y} + \lim_{s \rightarrow 0} \left(A_k \mathfrak{M}[f; s + k + \alpha_2] \right. \right. \\ & \left. \left. + B_{k+\alpha_2-\beta_1-\beta_2} \mathfrak{M}[h; s + 1 - \alpha_2 - k] \right) \right\} + R_n(\tilde{y}), \end{aligned} \quad (4.14)$$

where $n = m + \alpha_2 - \beta_1 - \beta_2$ and the remainder $R_n(\tilde{y}) = \mathcal{O}(\tilde{y}^{n+\beta_1+\beta_2} \log \tilde{y})$ when $\tilde{y} \rightarrow 0$. Computing the limit involved in (4.14) and after some calculations, we obtain that, under the condition (4.5),

$$\begin{aligned} F_K[\alpha_1, \alpha_2, \alpha_2, \beta_1, \beta_2, \beta_1; \gamma_1, \gamma_2, \gamma_3; x, y, z] = & \sum_{k=0}^{\alpha_2 - \beta_1 - \beta_2 - 1} \frac{\widehat{B}_k(y/z)}{y^{k+\beta_1+\beta_2}} \\ & + \sum_{k=0}^{m-1} \frac{\widehat{C}_{k+\alpha_2-\beta_1-\beta_2}(y/z) + B_{k+\alpha_2-\beta_1-\beta_2}^\times(y/z) \log|y|}{y^{k+\alpha_2}} + R_n(y), \end{aligned} \quad (4.15)$$

with $\widehat{B}_k(y/z)$, $B_k^\times(y/z)$ and $\widehat{C}_k(y/z)$ given in (4.9), (4.12) and (4.13), respectively. In (4.11) and (4.15), the reminder term $R_n(y) = \mathcal{O}(y^{-m-\alpha_2} \log y)$ when $|y| \rightarrow \infty$ uniformly in $|z|$ with y/z bounded.

Remark 4.1. When $x < 1$, in view of (2.12), the condition (4.5) attached to the functions f and h defined in (4.2) can be replaced by

$$\begin{aligned} \alpha_2 > 0, \beta_1 + \beta_2 > 0, |\arg(-y)| < \frac{\pi}{2}, |\arg(-z)| < \frac{\pi}{2}, \\ \Re(\gamma_1) > \Re(\alpha_1) > 0, x < 1, \left| \arg\left(1 - \frac{y}{1-x}\right) \right| < \pi, 0 < \varrho_1 < \frac{|y|}{|z|} < \varrho_2 < \infty. \end{aligned}$$

Under this condition, the asymptotic expansions (4.7), (4.11) and (4.15) and the corresponding error estimates remain true. In addition, by letting $x = 0$ in (4.3), we obtain

$$F_2[\alpha_2, \beta_2, \beta_1; \gamma_2, \gamma_3; y, z] = \frac{\tilde{y}}{\Gamma(\alpha_2)} \int_0^\infty h(\tilde{y}t) f(t) dt,$$

where

$$f(t) = {}_1F_1 \left[\begin{matrix} \beta_2 \\ \gamma_2 \end{matrix}; t e^{i\alpha} \right] {}_1F_1 \left[\begin{matrix} \beta_1 \\ \gamma_3 \end{matrix}; t e^{i\beta} / \gamma \right], \quad h(t) = e^{-t} t^{\alpha_2 - 1}$$

and F_2 is the second Appell hypergeometric function. The integral shares the same form with the Mellin integral considered by Garcia and López in [10, Eq. (1.4)]. Therefore, our results contain Garcia and López's results as special cases.

A Calculation of a Mellin transform

Here we calculate a Mellin transform needed in Section 4. As in Section 4, the symbol $\mathfrak{M}[g; s]$ denotes the Mellin transform of g when it exists, or its analytic continuation as a function of s .

When $\Re(\lambda) < 0$ and $(x, \mu) \in \mathbb{V}_{\Psi_1}$, we define

$$\Psi_F(t) := {}_1F_1 \left[\begin{matrix} \xi \\ \eta \end{matrix}; \lambda t \right] \Psi_1[a, b; c, c'; x, \mu t].$$

In view of (2.13) and (3.6), $\Psi_F(t) = 1 + o(1)$ as $t \rightarrow 0^+$ and Ψ_F has an inverse power asymptotic expansion as $t \rightarrow +\infty$. Moreover, $\Psi_F \in L^1_{\text{loc}}(0, \infty)$. Thanks to [14, p. 278, Remark 1], the Mellin transform $\mathfrak{M}[\Psi_F; s]$ exists and defines a meromorphic function of s in the half plane $\Re(s) > 0$.

Using (2.7) and swapping the order of integrations, we obtain

$$\begin{aligned} \mathfrak{M}[\Psi_F; s] &= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 u^{b-1} (1-u)^{c-b-1} (1-ux)^{-a} \\ &\quad \times \mathfrak{M} \left[{}_1F_1 \left[\begin{matrix} \xi \\ \eta \end{matrix}; \lambda t \right] {}_1F_1 \left[\begin{matrix} a \\ c' \end{matrix}; \frac{\mu t}{1-ux} \right]; s \right] du. \end{aligned} \quad (\text{A.1})$$

The Mellin transform in right-hand side of (A.1) is given by [3, p. 436, Eq. (1)]

$$\begin{aligned} \mathfrak{M} \left[{}_1F_1 \left[\begin{matrix} a \\ b \end{matrix}; -\omega t \right] {}_1F_1 \left[\begin{matrix} c \\ d \end{matrix}; -\sigma t \right]; s \right] &= \omega^{-s} \frac{\Gamma(b)\Gamma(s)\Gamma(a-s)}{\Gamma(a)\Gamma(b-s)} {}_3F_2 \left[\begin{matrix} c, s, s-b+1 \\ d, s-a+1 \end{matrix}; -\frac{\sigma}{\omega} \right] \\ &\quad + \sigma^{a-s} \omega^{-a} \frac{\Gamma(b)\Gamma(d)\Gamma(s-a)\Gamma(a+c-s)}{\Gamma(d)\Gamma(b-a)\Gamma(a+d-s)} {}_3F_2 \left[\begin{matrix} a, a-b+1, a+c-s \\ a-s+1, a+d-s \end{matrix}; -\frac{\sigma}{\omega} \right], \end{aligned} \quad (\text{A.2})$$

where $\Re(\sigma) > 0$, $\Re(\omega) > 0$ and $0 < \Re(s) < \Re(a + c)$. In order to apply (A.2) to (A.1), we need the restriction $\Re(\mu/(1 - ux)) < 0$. Here for convenience, we require that $x < 1$ so that

$$\Re\left(\frac{\mu}{1 - ux}\right) = \frac{1}{1 - ux}\Re(\mu) = -\frac{1}{1 - ux}|\mu|\cos(\arg(-\mu)) < 0.$$

Taking $\omega = -\lambda$ and $\sigma = \mu/(ux - 1)$ in (A.2) and inserting (A.2) into (A.1), we obtain

$$\mathfrak{M}[\Psi_F; s] = C_1(s)(-\mu)^{-s}\left(\frac{\mu}{\lambda}\right)^\xi I_1(s) + C_2(s)(-\lambda)^{-s}I_2(s),$$

where

$$\begin{aligned} C_1(s) &:= \frac{\Gamma(\eta)\Gamma(c')\Gamma(s - \xi)\Gamma(a + \xi - s)}{\Gamma(a)\Gamma(\eta - \xi)\Gamma(\xi + c' - s)}, & C_2(s) &:= \frac{\Gamma(\eta)\Gamma(s)\Gamma(\xi - s)}{\Gamma(\xi)\Gamma(\eta - s)}, \\ I_1(s) &:= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c - b)} \int_0^1 u^{b-1} (1 - u)^{c-b-1} (1 - ux)^{s-a-\xi} \\ &\quad \times {}_3F_2\left[\begin{matrix} \xi, \xi - \eta + 1, a + \xi - s \\ \xi + 1 - s, \xi + c' - s \end{matrix}; \frac{\mu}{\lambda} \frac{1}{ux - 1}\right] du \end{aligned}$$

and

$$\begin{aligned} I_2(s) &:= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c - b)} \int_0^1 u^{b-1} (1 - u)^{c-b-1} (1 - ux)^{-a} \\ &\quad \times {}_3F_2\left[\begin{matrix} a, s, s - \eta + 1 \\ c', s - \xi + 1 \end{matrix}; \frac{\mu}{\lambda} \frac{1}{ux - 1}\right] du. \end{aligned}$$

With the help of (2.11), $I_1(s)$ and $I_2(s)$ can be evaluated in terms of the Kampé de Fériet function $F_{0:1;2}^{1:1;2}$.

Theorem A.1.

$$\begin{aligned} \mathfrak{M}[\Psi_F; s] &= C_1(s)(-\mu)^{-s}\left(\frac{\mu}{\lambda}\right)^\xi F_{0:1;2}^{1:1;2}\left[\begin{matrix} a + \xi - s : b; & \xi, \xi - \eta + 1 \\ - : c; & \xi + 1 - s, \xi + c' - s \end{matrix}; x, -\frac{\mu}{\lambda}\right] \\ &\quad + C_2(s)(-\lambda)^{-s} F_{0:1;2}^{1:1;2}\left[\begin{matrix} a : b; & s, s - \eta + 1 \\ - : c; & c', s - \xi + 1 \end{matrix}; x, -\frac{\mu}{\lambda}\right]. \end{aligned} \quad (\text{A.3})$$

By continuation of $F_{0:1;2}^{1:1;2}$, (A.3) is valid for

$$\Re(\lambda) < 0, (x, \mu) \in \mathbb{V}_{\Psi_1} \text{ and } \left(x, -\frac{\mu}{\lambda}\right) \in \mathbb{D}_F. \quad (\text{A.4})$$

Remark A.2. *There are several different proofs of (A.2), which makes it of particular interest. A classical approach is to replace the second ${}_1F_1$ function with its Mellin-Barnes integral representation and then interchange the order of integration to obtain a single contour integral. The use of Cauchy's residue theorem will do the rest. A more interesting proof requires the method of brackets (see, for example, [1] and [11]).*

Acknowledgement

The research of the second author is supported by National Natural Science Foundation of China (Grant No. 12001095).

References

- [1] T. Amdeberhan, O. Espinosa, I. Gonzalez, M. Harrison, V.H. Moll, A. Straub, Ramanujan's master theorem, *Ramanujan J.* 29 (2012) 103–120.
- [2] G.E. Andrews, R. Askey, R. Roy, *Special Functions*, Cambridge Univ Press, 1999.
- [3] Y.A. Brychkov, O.I. Marichev, N.V. Svischenko, *Handbook of Mellin transforms*. CRC Press, 2018.
- [4] Y.A. Brychkov, N. Saad, On some formulas for the Appell function $F_2(a, b, b'; c, c'; w, z)$, *Integral Transforms Spec. Funct.* 25(2) (2014) 111–123.
- [5] Y.A. Brychkov, N.V. Svischenko, On some formulas for the confluent Horn functions $H_{10}^{(c)}(a; c; w, z)$ and $H_{11}^{(c)}(a, c, c'; d; w, z)$ *Integral Transforms Spec. Funct.* 34(12) (2023) 915–930.
- [6] J. Choi, A. Hasanov, Applications of the operator $H(\alpha, \beta)$ to the Humbert double hypergeometric functions, *Comput. Math. Appl.* 61 (3) (2011) 663–671.
- [7] T. Antonova, R. Dmytryshyn, V. Goran, On the analytic continuation of Lauricella-Saran hypergeometric function $F_K(a_1, a_2, b_1, b_2; a_1, b_2, c_3; \mathbf{z})$, *Mathematics* 2023, 11, 4487.
- [8] R. Dmytryshyn, V. Goran, On the analytic extension of Lauricella-Saran's hypergeometric function F_K to symmetric domains, *Symmetry* 2024, 16, 220.
- [9] A. Erdélyi, W. Magnus, F. Oberhettinger, F. Tricomi, *Higher Transcendental Functions*, Vol. I. McGraw-Hill, New York, 1981.
- [10] E. Garcia, J. L. López, The Appell's function F_2 for large values of its variables, *Q. Appl. Math.* 68 (4) (2010) 701–712.
- [11] I. Gonzalez, I. Kondrashuk, V.H. Moll, L.M. Recabarren, Mellin-Barnes integrals and the method of brackets, *Eur. Phys. J. C* 82, 28 (2022).
- [12] P. Humbert, The confluent hypergeometric functions of two variables, *Edinb. R. S. Proc.* 41 (1919), 73–96.
- [13] C.M. Joshi, J.P. Arya, Inequalities for certain confluent hypergeometric functions of two variables, *Indian J. Pure Appl. Math.* 13 (1982), 491–500.
- [14] J.L. López, Asymptotic expansions of Mellin convolution integrals, *SIAM Rev.* 50 (2) (2008) 275–293.
- [15] M.-J. Luo, R.K. Raina, On certain results related to the hypergeometric function F_K , *J. Math. Anal. Appl.* 504 (2) (2021), 125439.
- [16] M.-J. Luo, M.-H. Xu, R.K. Raina, On certain integrals related to Saran's hypergeometric function F_K , *Fractal Frac.* 6 (3) (2022).
- [17] T.H. Nguyen, S. Yakubovich, *The Double Mellin-Barnes Type Integrals and Their Applications to Convolution Theory*, World Scientific, Singapore, 1992.
- [18] G. Nemes, An explicit formula for the coefficients in Laplace's method. *Constr. Approx.* 38 (3) (2013) 471–487 .
- [19] F.W.J. Olver, Why steepest descents?. *SIAM Rev.* 12 (2) (1970) 228–247.
- [20] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clark (Eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010.
- [21] S. Saran, Hypergeometric functions of three variables, *Ganita* 5 (1954) 77–91.
- [22] S. Saran, Transformations of certain hypergeometric functions of three variables, *Acta Math.* 93 (1955) 293–312.
- [23] S. Saran, Integral representations of Laplace type for certain hypergeometric functions of three variables, *Riv., Mat. Univ. Parma* 8 (1957) 133–143.
- [24] H.M. Srivastava, P.W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Ellis Horwood Ltd., Chichester, 1985.
- [25] E. Wagner, Asymptotische Darstellungen der hypergeometrischen Funktionen für große Werte eines Parameters, *Z. Anal. Anwend.* 1(3) (1982) 1–11.
- [26] S. Wald, M. Henkel, On integral representations and asymptotics of some hypergeometric functions in two variables, *Integral Transforms Spec. Funct.* 29(2) (2018) 95–112.