

COMPETING BOOTSTRAP PROCESSES ON THE RANDOM GRAPH $G(n, p)$

BY MICHELE GARETTO^{1,a}, EMILIO LEONARDI^{2,b} AND GIOVANNI LUCA TORRISI^{3,c}

¹*Università di Torino, Torino, Italy.*, ^amichele.garetto@unito.it

²*Politecnico di Torino, Torino, Italy.*, ^bemilio.leonardi@polito.it

³*CNR-IAC, Roma, Italy.*, ^cgiovanniluca.torrise@cnr.it

We consider a generalization of classic bootstrap percolation in which two competing processes concurrently evolve on the same graph $G(n, p)$. Nodes can be in one of three states, conveniently represented by different colors: red, black and white. Initially, a given number a_R of active red nodes (red seeds) are selected uniformly at random among the n nodes. Similarly, a given number a_B of active black nodes (black seeds) are selected uniformly at random among the other $n - a_R$ nodes. All remaining nodes are initially white (inactive). White nodes wake up at times dictated by independent Poisson clocks of rate 1. When a white node wakes up, it checks the state of its neighbors: if the number of red (black) neighbors exceeds the number of black (red) neighbors by a fixed amount $r \geq 2$, the node becomes an active red (black) node, and remains so forever. The parameters of the model are, besides r (fixed) and n (tending to ∞), the numbers a_R (a_B) of initial red (black) seeds, and the edge existence probability $p = p(n)$. We study the size A_R^* (A_B^*) of the final set of active red (black) nodes, identifying different regimes which are analyzed under suitable time-scales, allowing us to obtain detailed (asymptotic) temporal dynamics of the two concurrent activation processes.

1. Introduction. Bootstrap percolation, in its classical form, is a simple activation process on a graph that starts with a given number of initially active nodes (called seeds) and evolves as follows. Every inactive node that has at least $r \geq 2$ active neighbors is activated, and remains so forever (an irreversible activation process). The process stops when no more nodes can be activated, and unfolds over discrete rounds (or generations): in each round, all susceptible vertices (i.e., vertices that can be activated) become active together (i.e., they are synchronously activated).

As many percolation processes, bootstrap percolation exhibits an “all-or-nothing” behavior: either the process percolates to (nearly) all vertices in the graph, or it stops very soon with a final number of active vertices that is not much larger than the starting set. The process is said to almost percolate if the final number of active nodes is $n - o(n)$.

Historically, bootstrap percolation was first introduced in [1] on a Bethe lattice, and successively investigated on regular grids and trees [2, 4]. More recently, bootstrap percolation has been studied on random graphs and random trees, motivated by the increasing interest in large-scale complex systems such as technological, biological and social networks.

A milestone in this direction is the paper by Janson et al. [8], where the authors have provided a detailed analysis of the bootstrap percolation process on the Erdős–Rényi random graph $G(n, p)$. Specifically, in [8] authors study the critical size a_0 of the starting set and show that, for $1/n \ll p \ll n^{-1/r}$, there exists a threshold $g(n, p, r)$ such that, for every $\epsilon > 0$,

MSC2020 subject classifications: Primary 60K35; secondary 05C80.

Keywords and phrases: Bootstrap Percolation, Random Graphs.

a.s. the process almost percolates for $a_0 \geq (1 + \epsilon)g$, whereas the final number of active nodes is $O(g)$ when $a_0 \leq (1 - \epsilon)g$.

We emphasize that in [8] seeds are chosen uniformly at random. Later, it has been shown that the critical number of seeds triggering percolation can be significantly reduced if the selection of seeds is optimized (in the form of so-called contagious sets) [24, 25].

Somehow related to ours is the work in [23], where authors study a variant of the classical bootstrap percolation process on the $G(n, p)$ graph with two types of vertices: excitatory and inhibitory. The activation spreads to vertices for which the number of excitatory active neighbors exceeds the number of inhibitory active neighbors by a certain amount. When more than half of the vertices are inhibitory, they discover, in the traditional (round-based) model, curious non-monotonous effects on the final size, which disappear in a continuous-time setting in which some exponential (i.i.d.) transmission delays are added on the edges. Note that we also consider a continuous-time setting, but we put exponential delays on nodes, rather than on edges. Moreover, our process is very different from the one in [23], since we study the competition between two opposite activation processes. Another variant of classic bootstrap percolation, somehow related to our work, is majority bootstrap percolation [17], in which a node becomes active if at least half of its neighbors are active.

Large deviations of classic bootstrap percolation in $G(n, p)$ have also been studied: in [26] authors calculate the rate function for the event that a small (subcritical) set of initially active nodes eventually infects an unexpected number of vertices, and identify the least-cost trajectory realizing such a large deviation. Large deviations in the super-critical regime have instead been fully characterized in our previous work [21].

Bootstrap percolation has also been analyzed on random regular graphs [5], on random graphs with given vertex degrees [9], on Galton–Watson random trees [6], on random geometric graphs [15], on Chung–Lu random graphs [10, 11] (which notably permit considering the case of power-law node degree distribution), on small-world random graphs [12, 13] and on Barabasi–Albert random graphs [14]. In [22] we have analyzed the bootstrap percolation process on the stochastic block model (SBM), a natural extension of the Erdős–Rényi random graph that incorporates the community structure observed in many real systems.

Instead of considering yet another underlying graph, in this paper we open a new (to the best of our knowledge) direction in the theory of bootstrap percolation, where nodes can be in three states, and two competing bootstrap-like processes evolve in parallel over continuous time. We analyze this process on the simple $G(n, p)$ graph, leaving to future work the extension of the analysis to different graphs.

2. Model description and main results.

2.1. Model description. In this paper we consider a generalization of the bootstrap percolation process on the Erdős–Rényi random graph $G(n, p_n) = (\mathcal{V}^{(n)}, \mathcal{E}^{(n)})$, $n \in \mathbb{N} := \{1, 2, \dots\}$, introduced in [8]. Here $\mathcal{V}^{(n)} := \{1, \dots, n\}$ is the set of nodes and $\mathcal{E}^{(n)}$ is the set of edges, which are independently added with probability $p_n \in (0, 1)$. Our model is defined as follows:

- Nodes can be red (R), black (B) or white (W). In the following we will refer to either R nodes or B nodes as active, and to W nodes as inactive.
- At time 0, an arbitrary number $a_R^{(n)}$ of nodes (selected uniformly at random among the n nodes) are set R , an arbitrary number $a_B^{(n)}$ of nodes (selected uniformly at random among the remaining $n - a_R^{(n)}$ nodes) are set B , and all the other nodes are set W . Nodes already active at time 0 are called “seeds”.¹

¹As seeds are selected uniformly at random in a $G(n, p)$ graph, the order in which the two seed sets are created is not relevant, i.e., it has no impact on the process evolution.

- Each node W periodically wakes up according to a Poissonian clock, i.e., the sequence of times at which a node W wakes up forms a Poisson process with intensity 1. The Poissonian clocks attached to different white nodes are stochastically independent. Upon wake up, a node W activates by taking a color $S \in \{R, B\}$ if the difference between the number of its neighbors of color S and the number of its neighbors of color $S^c \in \{R, B\} \setminus S$ is bigger than or equal to a given threshold $r \in \mathbb{N} \setminus \{1\}$, otherwise the node remains W . Throughout this paper we will refer such condition as “threshold condition with respect to S ”.
- Active nodes never deactivate (i.e. change color), hence the number of nodes R (or B) is non-decreasing over time.
- The process stops when no more nodes can be activated, i.e., no white node satisfies the “threshold condition with respect to either R or B ”.

The main objective of this paper is to study the asymptotic behavior of the final number $A_R^{*(n)}$ ($A_B^{*(n)}$) of nodes R (B), as n grows large. As a common practice in the theory of large random graphs, in general we will omit the dependence on n of quantities, writing e.g. G in place of $G(n, p_n)$, p in place of p_n , a_S in place of $a_S^{(n)}$, A_S^* in place of $A_S^{*(n)}$, $S \in \{R, B\}$, and so on. We will explicit such a dependence only when needed.

Moreover, in the following, we always assume that:

$$(2.1) \quad \frac{1}{n} \ll p \ll \frac{1}{n^{1/r} \log n}.$$

(2.1) is slightly tighter than the corresponding condition in [8], (i.e., $\frac{1}{n} \ll p \ll \frac{1}{n^{1/r}}$). This is justified by the fact that our results are tighter (i.e., we prove almost sure convergence) than those in [8] (where convergence in probability is shown). Furthermore, we assume:

$$(2.2) \quad a_R/q \rightarrow \alpha_R, \quad a_B/q \rightarrow \alpha_B, \quad \text{for some positive constants } \alpha_R, \alpha_B > 0,$$

where the sequence $\{q_n\}$ is chosen in such a way that or

$$(2.3) \quad \begin{aligned} & i) \quad q = g := \left(1 - \frac{1}{r}\right) \left(\frac{(r-1)!}{np^r}\right)^{\frac{1}{r-1}} \text{ with } pg \rightarrow 0 \quad \text{or} \\ & ii) \quad g \ll q \ll p^{-1} \quad \text{or} \quad iii) \quad q = p^{-1} \quad \text{or} \quad iv) \quad p^{-1} \ll q \ll n. \end{aligned}$$

REMARK 2.1. In contrast to the bootstrap percolation process considered in [8], where the order in which nodes activate has no impact on the final size of active nodes (see Proposition 4.1 in [22]), in our case the order in which nodes activate is important, as one can check on toy examples. Poissonian clocks have been introduced as a naturally way to solve this problem: by so doing, essentially we consider a system in which, at any given time, the next node to activate is chosen uniformly at random among the nodes that satisfy the threshold condition with respect to either R or B .

REMARK 2.2. When $a_{S^c} = 0$, our process reduces to an asynchronous version of the classic bootstrap percolation process, in which not yet active nodes, i.e. W nodes, become S -active at the times of a suitably thinned unit-rate Poisson process. Therefore A_S^* equals the final number of active nodes of the standard bootstrap percolation process on the Erdős-Renyi random graph G with threshold $r \geq 2$ and number of seeds a_S , see [8].

2.2. *Main results.* To state our results we need to introduce the following function $\beta_S : [0, \infty)^2 \rightarrow \mathbb{R}$, $S \in \{R, B\}$:

$$(2.4) \quad \beta_S(x_R, x_B) := \begin{cases} r^{-1}(1-r^{-1})^{r-1}(x_S + \alpha_S)^r - x_S & \text{if } q = g \\ \frac{1}{r!}(x_S + \alpha_S)^r & \text{if } g \ll q \ll p^{-1} \\ \sum_{r'=r}^{\infty} \sum_{r''=0}^{r'-r} \frac{(x_S + \alpha_S)^{r'}}{r'!} \frac{(x_{Sc} + \alpha_{Sc})^{r''}}{r''!} e^{-(x_R + x_B + \alpha_R + \alpha_B)} & \text{if } q = p^{-1} \\ \mathbf{1}_{[0, \infty)} \left(\frac{x_S + \alpha_S}{x_R + \alpha_R + x_B + \alpha_B} - \frac{1}{2} \right) & \text{if } p^{-1} \ll q \ll n. \end{cases}$$

Roughly speaking, $\beta_S(x_R, x_B)$ represents a suitably scaled asymptotic estimate of the average number of nodes satisfying the threshold condition with respect to S , when $x_R q$ nodes have become R -active and $x_B q$ nodes have become B -active (see Lemma E.4 in Appendix). As it will become clear in the following, the asymptotic behavior of the R and B activation processes on time-scale q (i.e., as long as the number of active nodes is $\Theta(q)$) is tightly related to the properties of function β_S .

REMARK 2.3. Suppose that $q = g$. Function $\beta_S(x_S)$ is strictly positive for $\alpha_S > 1$, whereas it has two strictly positive zeros for $\alpha_S < 1$ (we denote by z_S the smallest zero in this case). When either $g \ll q \ll p^{-1}$ or $q = p^{-1}$, β_S is strictly positive, while β_S is non-negative for $p^{-1} \ll q \ll n$.

Assume (2.1), $a_R/g \rightarrow \alpha_R$ and $a_B = 0$. It follows from the main results in [8] that $A^*/g \rightarrow z_R + \alpha_R$ almost surely, provided that $\alpha_R < 1$; instead, $A^*/n \rightarrow 1$ almost surely when $\alpha_R > 1$. This means that there exists a critical number of seeds under which the bootstrap percolation process basically does not evolve, and above which it percolates the entire graph almost completely. This well known behavior of the classical bootstrap percolation process suggests us to adopt the following terminology. Restricting ourselves, without lack of generality, to the case $\alpha_R > \alpha_B$, we say that the system is in the sub-critical regime whenever $q = g$ and $\alpha_R < 1$; we say that the system is in the super-critical regime whenever either $g \ll q$, or $q = g$ and $\alpha_R > 1$.

Consider the system evolution in the sub-critical regime. One would expect that, for the effect of competition, the asymptotic final sizes of S -active nodes ($S \in \{R, B\}$) might be smaller than those achieved in the absence of competition (e.g., when $a_{Sc} = 0$). However this is not the case, as stated by the following theorem.

THEOREM 2.4. Assume $q = g$ with $\alpha_B < \alpha_R < 1$. Then

$$\frac{A_R^*}{q} \rightarrow z_R + \alpha_R, \quad \frac{A_B^*}{q} \rightarrow z_B + \alpha_B \quad \mathbb{P}\text{-a.s.}$$

where z_S is the smallest zero of β_S (see Remark 2.3).

Theorem 2.4 states that, in the sub-critical regime, the two competing processes essentially do not interact with each other. Indeed, A_S^*/q converges exactly to the same value it would converge to, when $a_{Sc} = 0$.

Consider now the more interesting super-critical regime.

THEOREM 2.5. (i) when $q = g$ and $\alpha_R > 1$, then

$$(2.5) \quad \frac{A_R^*}{n} \rightarrow 1 \quad \text{and} \quad \frac{A_B^*}{q} \rightarrow g_B(\kappa_g) + \alpha_B, \quad \mathbb{P}\text{-a.s.}$$

(ii) when $g \ll q \ll n$, then

$$\frac{A_R^*}{n} \rightarrow 1 \quad \text{and} \quad \frac{A_B^*}{n} \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

where $\kappa_{\mathbf{g}}$ and $g_B(\kappa_{\mathbf{g}}) := \lim_{x \uparrow \kappa_{\mathbf{g}}} g_B(x)$ are defined as follows:

DEFINITION 2.6. (Underlying Cauchy problem). Let $\mathbf{g}(x) = (g_R(x), g_B(x))$ denote the maximal solution of the Cauchy problem:

$$(2.6) \quad \mathbf{g}'(x) = \beta(\mathbf{g}(x)), \quad x \in [0, \kappa_{\mathbf{g}}), \quad \mathbf{g}(0) = (0, 0)$$

where $\beta := (\beta_R, \beta_B)$.

In words, Theorem 2.5 states that, in the super-critical regime, the R -activation process percolates the entire graph almost completely, causing an “early stop” of the competing activation process, even when $\alpha_B > 1$, i.e., when the B -activation process would potentially percolate in the absence of competition. Observe that, while in the sub-critical regime the process stops when $O(q)$ nodes are active, in the super-critical regime almost all nodes become active (i.e., the final size of active nodes is $n - o(n)$). A numerical illustration of our results is provided in Appendix B.

3. Notation and Preliminary Analysis.

3.1. *Main variables and their relations.* In this subsection we introduce the random quantities in terms of which we will describe the dynamics over G of our competing bootstrap percolation processes and quantify their final sizes. All the random variables considered hereafter are defined on an underlying probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Let $\mathcal{V}_W \subset \mathcal{V}$ be the set of non-seed nodes, and set $n_W := |\mathcal{V}_W| = n - (a_R + a_B)$. Here, given a finite set A , we denote by $|A|$ its cardinality. We attach, independently to every node $v \in \mathcal{V}_W$, a unit rate Poissonian clock, whose ordered points represent the successive wake-up times of node v . More formally, we define a sequence $\{N'_v\}_{v \in \mathcal{V}_W}$ of independent Poisson processes on $[0, \infty) \times \mathcal{V}_W$ with N'_v having mean measure $dt\delta_v(d\ell)$, where $\delta_v(\cdot)$ is the Dirac measure on \mathcal{V}_W concentrated at $v \in \mathcal{V}_W$. As it is well-known, the point process

$$(3.1) \quad N' := \sum_{v \in \mathcal{V}_W} N'_v$$

is still a Poisson process on $[0, \infty) \times \mathcal{V}_W$ with intensity measure $n_W dt \mathbb{U}(dv)$, where \mathbb{U} is the uniform law on \mathcal{V}_W . We denote by $\{(T'_k, V'_k)\}_{k \in \mathbb{N}}$ the points of N' (with ordered first coordinates): here T'_k is the time at which the k -th wake-up event occurs and V'_k is the corresponding node. We denote by N_S , $S \in \{R, B\}$, the S -activation point process on $[0, \infty) \times \mathcal{V}_W$, i.e., for any $t > 0$ and any $L \subseteq \mathcal{V}_W$, $N_S([0, t] \times L)$ is the number of S -active nodes in $L \subseteq \mathcal{V}_W$ at time t . Let (T_k^S, V_k^S) denote the k -th point of N_S . By construction T_k^S is the “activation time” of node V_k^S , i.e., the time at which node V_k^S becomes S -active (by taking color S). Note that white node V' , which wakes up at time T' , can become S -active if and only if it satisfies the “threshold condition with respect to S ”. Therefore, the point process N_S can be constructed by thinning $\{(T'_k, V'_k)\}_{k \in \mathbb{N}}$ as follows: we retain only those couples (T'_k, V'_k) , $k \in \mathbb{N}$, for which, at time $(T'_k)^-$, the white node V'_k satisfies the “threshold condition with respect to S ”.

We set $N := N_R + N_B$ and denote by (T_k, V_k) , $k \in \mathbb{N}$, the points of N . Throughout this paper we refer to N as the (global) activation process. In the following we will use $N_S(t)$ and $N(t)$ as a shorthand notation for $N_S([0, t] \times \mathcal{V}_W)$ and $N([0, t] \times \mathcal{V}_W)$, respectively. Hereafter, we denote by $\mathcal{V}_S(t) \subset \mathcal{V}_W$, $t \geq 0$, the set of non-seed nodes which are S -active at time t , i.e.,

$$\mathcal{V}_S(t) = \{V_k^S\}_{k: T_k^S \in [0, t]} \quad \text{with} \quad \mathcal{V}_S(0) = \emptyset$$

and with $\mathcal{V}_W(t) \subset \mathcal{V}_W$, $t \geq 0$, the set of non-seed nodes which are still W at time t , i.e.,

$$\mathcal{V}_W(t) := \mathcal{V}_W \setminus (\mathcal{V}_R(t) \cup \mathcal{V}_B(t)) \quad \text{with} \quad \mathcal{V}_W(0) := \mathcal{V}_W.$$

Let $\{E_i^{R,(v)}\}_{i \in \mathbb{N}}$, $\{E_i^{B,(v)}\}_{i \in \mathbb{N}}$, $v \in \mathcal{V}_W$, be two independent sequences of independent and identically distributed random variables with Bernoulli's law and mean p , independent of $\{(T'_k, V'_k)\}_{k \in \mathbb{N}}$. The random variable $E^{S,(v)}$ indicates the presence (or not) of an edge between node $v \in \mathcal{V}_W$ and an S -active node. We often refer to the random variables $E^{S,(v)}$ as S -marks and define the quantities

$$(3.2) \quad D_R^{(v)}(t) := \sum_{i=1}^{N_R(t)+a_R} E_i^{R,(v)} \quad \text{and} \quad D_B^{(v)}(t) := \sum_{i=1}^{N_B(t)+a_B} E_i^{B,(v)}, \quad v \in \mathcal{V}_W,$$

which represent the number of neighbors of node v whose color is, respectively, R or B at time t . We also define the random variables:

$$\mathcal{V}_S(t^-) = \{V_k^S\}_{k: T_k^S \in [0, t)}, \quad \mathcal{V}_W(t^-) := \mathcal{V}_W \setminus (\mathcal{V}_R(t^-) \cup \mathcal{V}_B(t^-)),$$

$$D_R^{(v)}(t^-) := \sum_{i=1}^{N_R([0, t) \times \mathcal{V}_W) + a_R} E_i^{R,(v)} \quad \text{and} \quad D_B^{(v)}(t^-) := \sum_{i=1}^{N_B([0, t) \times \mathcal{V}_W) + a_B} E_i^{B,(v)}, \quad v \in \mathcal{V}_W.$$

Moreover, the set of S -susceptible nodes at t is defined as

$$(3.3) \quad \mathcal{S}_S(t) = \{v \in \mathcal{V}_W : D_S^{(v)}(t) - D_{S^c}^{(v)}(t) \geq r\},$$

and similarly the set of S -susceptible nodes at time t^- is defined as $\mathcal{S}_S(t^-) := \{v \in \mathcal{V}_W : D_S^{(v)}(t^-) - D_{S^c}^{(v)}(t^-) \geq r\}$.

Lastly, we denote by

$$\mathcal{S}(t) := \mathcal{S}_R(t) \cup \mathcal{S}_B(t) \quad \text{and} \quad \mathcal{S}(t^-) := \mathcal{S}_R(t^-) \cup \mathcal{S}_B(t^-)$$

the set of susceptible nodes at time t and t^- , respectively.

$$\mathcal{S}(t) := \mathcal{S}_R(t) \cup \mathcal{S}_B(t)$$

the set of susceptible nodes at time t . The final number of active nodes is clearly given by

$$A^* := A_R^* + A_B^*, \quad \text{where} \quad A_S^* := N_S([0, \infty)) + a_S.$$

Furthermore, defined

$$K^* := \min\{k \in \mathbb{N} : \mathcal{S}(T_{k-1}) \cap \mathcal{V}_W(T_{k-1}) = \emptyset\}, \quad (\text{we conventionally set } T_0 := 0)$$

we have

$$A^* = K^* + a_R + a_B - 1.$$

Note that the overall activation process N naturally stops at time T_{K^*-1} , since no node becomes active after T_{K^*-1} . For the moment, we conventionally define $T_k := \infty$ on the event $\{K^* \leq k\}$, and note that $T_{K^*} = \infty$. We mention that, for technical reasons, in Section 3.4 we will ‘‘artificially’’ extend, in a suitable way, the activation process beyond T_{K^*-1} , redefining T_k on the event $\{K^* \leq k\}$. Of course, such extension will not have any impact on the dynamics of the activation process until T_{K^*-1} . We remark that random graphs $G(n, p_n)$, as well as the dynamical processes evolving on top of them, are independent for different values of n .

REMARK 3.1. During the evolution of the activation process, every edge $(v, w) \in \mathcal{E}$ is unveiled potentially twice, i.e., when both v and w get active; consistency between the two choices is not guaranteed, nevertheless this has no effect on the dynamical process, similarly to what happens in the bootstrap percolation process studied in [8]. Indeed, assuming that v gets active before w , the mark potentially added to v (already active), when w gets active, has no impact on the activation process.

3.2. *Further notation.* Throughout this paper, all the unspecified limits are taken as $n \rightarrow \infty$. Given two numerical sequences $\{f(n)\}_{n \in \mathbb{N}}$ and $\{g(n)\}_{n \in \mathbb{N}}$, we write: $f(n) \ll g(n)$ if $f(n) = o(g(n))$, i.e., $f(n)/g(n) \rightarrow 0$; $f(n) = O(g(n))$ if $\limsup_{n \rightarrow \infty} \left| \frac{f(n)}{g(n)} \right| < \infty$; $f(n) = \Theta(g(n))$ if both $f(n) = O(g(n))$ and $g(n) = O(f(n))$, $f(n) \sim g(n)$ if $f(n)/g(n) \rightarrow 1$. We denote with $\|\cdot\|$ the euclidean norm, and with $\lfloor \cdot \rfloor$ and $\lceil \cdot \rceil$ the floor and the ceiling functions, respectively. Given set \mathcal{A} , we denote with \mathcal{A}^c the complementary set. Moreover, given a sequence of real-valued random variables $\{X_n\}_{n \in \mathbb{N}}$, we write $X_n = o_{\text{a.s.}}(f(n))$ if $\mathbb{P}\left(\lim \left| \frac{X_n}{f(n)} \right| = 0\right) = 1$. Given two real-valued random variables X and Y , we denote by $X \leq_{st} Y$ the usual stochastic order, i.e., we write $X \leq_{st} Y$ if $\mathbb{P}(X > z) \leq \mathbb{P}(Y > z)$, $z \in \mathbb{R}$. $\text{Bin}(m, \theta)$, $\text{Po}(\lambda)$ and $\text{EXP}(\lambda)$ respectively denote random variables distributed according to a binomial law with parameters (m, θ) , a Poisson law and an exponential law (both with parameter $\lambda > 0$). Symbol $\stackrel{\text{L}}{=}$ denotes the identity in law. At last, throughout this paper we will use several times the function

$$(3.4) \quad H(x) := 1 - x + x \log x, \quad x > 0, \quad H(0) := 1.$$

3.3. *Markovianity of the process.* As consequence of our assumptions, the considered system can be described by a Markovian process² specified by the following proposition

PROPOSITION 3.2. The stochastic process

$$\mathbf{X} = \{\mathbf{X}(t)\}_{t \geq 0} = \{(\mathbf{1}_{\{v \in \mathcal{V}_R(t)\}}, \mathbf{1}_{\{v \in \mathcal{V}_B(t)\}}, D_R^{(v)}(t), D_B^{(v)}(t))_{v \in \mathcal{V}_W}\}_{t \geq 0}$$

is a regular-jump continuous time homogeneous Markov chain, i.e., a continuous time homogeneous Markov chain such that, for almost all ω , the cardinality of $\text{Disc}(\omega) \cap [0, c]$ is finite for any $c \geq 0$. Here $\text{Disc}(\omega)$ denotes the set of discontinuity points of the mapping $t \mapsto \mathbf{X}(t, \omega)$. The state space of \mathbf{X} , say \mathcal{X} , is contained in $(\{0, 1\} \times \{0, 1\} \times \{0, \dots, n\} \times \{0, \dots, n\})^{|\mathcal{V}_W|}$, with diagonal elements of the transition-rate matrix:

$$q_{\mathbf{x}} := \lim_{h \rightarrow 0} \frac{1 - \mathbb{P}(\mathbf{X}(h) = \mathbf{x} \mid \mathbf{X}(0) = \mathbf{x})}{h} = |\mathcal{V}_W(\mathbf{x}) \cap \mathcal{S}(\mathbf{x})| \geq 0, \quad \mathbf{x} \in \mathcal{X}.$$

Regarding notation in the r.h.s., observe that, for any $t \geq 0$, both $\mathcal{V}_W(t)$ and $\mathcal{S}(t)$ are, by construction, $\sigma\{\mathbf{X}(t)\}$ -measurable. Therefore, we will conveniently denote $\mathcal{V}_W(t)$ and $\mathcal{S}(t)$ also with $\mathcal{V}_W(\mathbf{X}(t))$ and $\mathcal{S}(\mathbf{X}(t))$, respectively.

Observe that the sequence of transition times of \mathbf{X} coincides, by construction, with the sequence of activation times $\{T_k\}_{k \geq 0}$ of nodes.³ Let $\mathcal{F}_t^{\mathbf{X}} := \sigma\{\mathbf{X}(s) : s \leq t\}$ be the natural filtration of the Markov chain \mathbf{X} and $\{\mathbf{X}_k\}_{k \in \mathbb{N} \cup \{0\}}$ the embedded chain defined by $\mathbf{X}_k = \mathbf{X}(T_k)$. On the event $\{K^* - 1 \leq k\}$ we have $\mathbf{X}_k = \mathbf{X}(\infty) \in \Delta := \{\mathbf{x} \in \mathcal{X} : q_{\mathbf{x}} = 0\}$, while $\{\mathbf{X}_k\}_{0 \leq k < K^* - 1} \in \mathcal{X} \setminus \Delta = \{\mathbf{x} \in \mathcal{X} : q_{\mathbf{x}} > 0\}$. Moreover, given K^* and

²Our suggested reference for Markov Chains is [3].

³We recall that conventionally we have $T_0 = 0$.

$\{\mathbf{X}_k\}_{0 \leq k < K^* - 1}$, the random variables $\{W_k\}_{0 \leq k < K^* - 1}$, $W_k := T_{k+1} - T_k$ are independent and W_k is exponentially distributed with mean $1/q_{\mathbf{x}_k}$. In particular, for an arbitrary finite sequence of states $\{\mathbf{x}_h\}_{0 \leq h \leq k} \subset \mathcal{X} \setminus \Delta$ and an arbitrary finite sequence of positive numbers $\{a_h\}_{0 \leq h \leq k} \subset (0, \infty)$, we have

$$(3.5) \quad \mathbb{P} \left(\bigcap_{0 \leq h < k} \{\mathbf{X}_h = \mathbf{x}_h\} \cap \{W_h > a_h\} \right) = \mathbb{P}(\mathbf{X}_0 = \mathbf{x}_0) \prod_{0 \leq h < k} p_{\mathbf{x}_h \mathbf{x}_{h+1}} e^{-q_{\mathbf{x}_h} a_h},$$

where $(p_{\mathbf{x}\mathbf{y}})$ denotes the transition matrix of $\{\mathbf{X}_k\}_k$.

3.3.1. Discrete time notation. To study the evolution of the system at the points $\{T_k\}_{k \in \mathbb{N}}$, it is convenient to introduce some discrete time notation. For $k \in \mathbb{N} \cup \{0\}$, we set ⁴

$$N_S[k] := N_S(T_k), \quad \mathcal{S}_S[k] := \mathcal{S}_S(T_k), \quad \mathcal{V}_W[k] := \mathcal{V}_W(T_k), \quad D_S^{(v)}[k] := D_S^{(v)}(T_k).$$

Since all these random variables are $\sigma\{\mathbf{X}_k\}$ -measurable, therefore, when convenient to highlight the dependence on the state, we write

$$N_S[k] = N_S(\mathbf{X}_k), \quad \mathcal{S}_S[k] = \mathcal{S}_S(\mathbf{X}_k), \quad \mathcal{V}_W[k] = \mathcal{V}_W(\mathbf{X}_k), \quad D_S^{(v)}[k] = D_S^{(v)}(\mathbf{X}_k).$$

Moreover, we define

$$U_{k+1}^R = U^R(\mathbf{X}_k) := \frac{|\mathcal{V}_W(\mathbf{X}_k) \cap \mathcal{S}_R(\mathbf{X}_k)|}{|\mathcal{V}_W(\mathbf{X}_k) \cap \mathcal{S}(\mathbf{X}_k)|}, \quad U_{k+1}^B = 1 - U_{k+1}^R, \quad \forall k \in \mathbb{N} \cup \{0\},$$

where conventionally we put $0/0 := 1/2$. Finally, we note that

$$(3.6) \quad N_S[k+1] = N_S[k] + M_{k+1}^S, \quad S \in \{R, B\}$$

with

$$M_{k+1}^S := \mathbf{1}_{\{V_{k+1} \in \mathcal{V}_W[k] \cap \mathcal{S}_S[k]\}} \mathbf{1}_{\{K^* - 1 > k\}} = \mathbf{1}_{\{V_{k+1} \in \mathcal{V}_W[k] \cap \mathcal{S}_S[k]\}} \mathbf{1}_{\{\mathbf{x}_k \in \mathcal{X} \setminus \Delta\}}$$

where we used that $\mathbf{1}_{\{K^* - 1 > k\}} = \mathbf{1}_{\{\mathbf{x}_k \in \mathcal{X} \setminus \Delta\}}$ by construction. Note that $M_{k+1}^S \in \sigma\{\mathbf{X}_k, \mathbf{X}_{k+1}\}$. Hereafter, we set $\mathcal{H}_k := \sigma\{\mathbf{X}_h : 0 \leq h \leq k\}$. The following proposition holds.

PROPOSITION 3.3. For $S \in \{R, B\}$ and $k \in \mathbb{N} \cup \{0\}$, we have

$$\begin{aligned} \mathbb{P}(M_{k+1}^S = 1 \mid \mathcal{H}_k) &= \mathbb{P}(M_{k+1}^S = 1 \mid \mathbf{X}_k) = \mathbb{P}(V_{k+1} \in \mathcal{V}_W(\mathbf{X}_k) \cap \mathcal{S}_S(\mathbf{X}_k), \mathbf{X}_k \in \mathcal{X} \setminus \Delta \mid \mathbf{X}_k) \\ &= U_{k+1}^S \mathbf{1}_{\{\mathbf{x}_k \in \mathcal{S} \setminus \Delta\}} = U_{k+1}^S \mathbf{1}_{\{K^* - 1 > k\}}. \end{aligned}$$

Here, the first equality is a consequence of the Markovianity of $\{\mathbf{X}_k\}$, while the last one follows from elementary properties of Poisson processes: indeed, given \mathbf{X}_k , V_{k+1} is uniformly distributed over the set $\mathcal{V}_W(\mathbf{X}_k) \cap \mathcal{S}(\mathbf{X}_k)$ whenever $\mathcal{V}_W(\mathbf{X}_k) \cap \mathcal{S}(\mathbf{X}_k) \neq \emptyset$. \square

For $u := m_1/m_2$, where $m_1 \in \{0, 1, \dots, m_2\}$ and $m_2 \in \{1, \dots, n_W\}$, define \mathcal{X}_u as the set of states \mathbf{x} such that $U^R(\mathbf{x}) = u$, i.e.

$$\mathcal{X}_u := \{\mathbf{x} \in \mathcal{X} : U^R(\mathbf{x}) = u\}.$$

Note that by construction: $\{U_{k+1}^R = u\} = \{U^R(\mathbf{X}_k) = u\} = \{\mathbf{X}_k \in \mathcal{X}_u\}$. Moreover, we define $\widehat{\mathcal{X}}_m := \{\mathbf{x} \in \mathcal{X} : q_{\mathbf{x}} = m\}$ for any $m \in \{0, 1, \dots, n_W\}$.

As direct consequence of Markovianity, the following two propositions, whose proofs is reported in Appendix O.

⁴By construction, on the event $\{K^* \leq k\}$, we have $\mathcal{S}_S[k] = \mathcal{S}_S[K^* - 1]$, $\mathcal{V}_W[k] = \mathcal{V}_W[K^* - 1]$, etc.

PROPOSITION 3.4. For any $S \in \{R, B\}$ and $k \in \mathbb{N} \cup \{0\}$, given $\{\mathbf{X}_k \in \mathcal{X}_u\}$, the random variable M_{k+1}^S is independent of the sequence $\{M_h^S\}_{1 \leq h \leq k}$.

PROPOSITION 3.5. For $k \in \mathbb{N} \cup \{0\}$, let $\{m_h\}_{0 \leq h \leq k} \subseteq \{1, \dots, n_W\}$. Given the event $\bigcap_{0 \leq h \leq k} \{\mathbf{X}_h \in \widehat{\mathcal{X}}_{m_h}\}$, the sojourn-time random variables $\{W_h\}_{0 \leq h \leq k}$ are independent. Moreover, for every $0 \leq h \leq k$, W_h is exponentially distributed with parameter m_h .

3.4. *Prolonging the process N beyond T_{K^*-1} .* Since $\mathcal{V}_W[K^* - 1] \cap \mathcal{S}[K^* - 1] = \emptyset$, we have $N((T_{K^*-1}, \infty) \times \mathcal{V}_W) = 0$. To simplify the analysis, it is convenient to extend N beyond T_{K^*-1} by activating nodes that are not susceptible. Hereon, we still denote by N_S and N the activation processes extended beyond T_{K^*-1} and by $\{(T_k^S, V_k^S)\}_{k \geq 1}$ and $\{(T_k, V_k)\}_{k \geq 1}$ their points respectively. Points (T_{K^*+j}, V_{K^*+j}) , for $j \geq 0$, are defined by thinning the point process $\{(T'_{k'}, V'_{k'})\}_{k': T'_{k'} > T_{K^*-1}}$ and retaining only those couples $(T'_{k'}, V'_{k'})$ such that $V'_{k'}$ is still W . More precisely, given points $\{(T_h, V_h)\}_{1 \leq h \leq k}$, on $\{K^* - 1 \leq k\}$, we set $(T_{k+1}, V_{k+1}) := (T'_{\ell_{k+1}}, V'_{\ell_{k+1}})$ with

$$(3.7) \quad \ell_k := \{k' : T'_{k'} = T_k\}, \quad \ell_{k+1} := \min\{k' > \ell_k : V'_{k'} \in \mathcal{V}_W(T'_{\ell_k})\}.$$

Then we assign to V_{k+1} either color R or color B (regardless of the fact that V_{k+1} is R -susceptible or B -susceptible) by flipping a biased coin. I.e., we define the processes N_S for $S \in \{R, B\}$ on the event $\{K^* \leq k\}$ as:

$$(3.8) \quad (T_{N_S[k]+1}^S, V_{N_S[k]+1}^S) := (T_{k+1}, V_{k+1}) \quad \text{with probability } U_{k+1}^S$$

where

$$U_{K^*} := \frac{1}{2}, \quad U_{k+1}^S := \frac{|Q_{k+1}^S|}{|Q_{k+1}^R| + |Q_{k+1}^B|}, \quad \text{on } \{K^* \leq k\}$$

$$(3.9) \quad \begin{aligned} Q_{k+1}^S &:= |\mathcal{S}_S[k]| - N_S[k] + |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| \\ &\quad - |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| \end{aligned}$$

and $Q_1^S := |\mathcal{S}_S(0)|$. In the expression of the random variable Q_{k+1}^S , the first addend is the number of S -susceptible nodes, at time T_k ; the second addend is the number of S -active nodes (excluding seeds), at time T_k ; the third addend is the number of S -active nodes (excluding seeds), at time T_k , which have at least r S -active neighbors, but are not S -susceptible; the fourth addend is the number of S^c -active nodes (excluding the seeds), at time T_k , which have at least r S^c -active neighbors and are S -susceptible. Note that also the extended process stops when all the nodes have got a color, i.e., when k equals the number of nodes that were originally white, i.e., $k = n_W := |\mathcal{V}_W| = n - a_R - a_B$. Finally, note that, (3.2) holds for any $0 \leq k \leq n_W$, since also on the event $\{K^* - 1 \leq k\}$ nodes that become S -active distribute S -marks to all of their neighbors. As before, we conventionally set $T_k = \infty$ for every $k > n_W$. We emphasize that the definition of random variable Q_{k+1}^S is purely instrumental to guarantee that on the event $\{k \leq K^* - 1\}$ we have $Q_{k+1}^S = |\mathcal{V}_W[k] \cap \mathcal{S}_S[k]|$, as stated by the following lemma, which is proved in Appendix C.

LEMMA 3.6. We have:

$$(3.10) \quad Q_{k+1}^S \mathbf{1}_{\{K^* > k\}} = |\mathcal{V}_W[k] \cap \mathcal{S}_S[k]| \mathbf{1}_{\{K^* > k\}},$$

$$(3.11) \quad |\mathcal{S}_S[k]| - k \leq Q_{k+1}^S \leq |\mathcal{S}_S[k]|, \quad k \in \mathbb{N} \cup \{0\}$$

PROPOSITION 3.7. Note that, after its extension, the stochastic process

$$\mathbf{X} = \mathbf{X}(t) = \{(\mathbf{1}_{\{v \in \mathcal{V}_R(t)\}}, \mathbf{1}_{\{v \in \mathcal{V}_B(t)\}}, D_R^{(v)}(t), D_B^{(v)}(t))_{v \in \mathcal{V}_W}, (\mathbf{1}_{T_{K^*} \leq t})\}_{t \geq 0}$$

is still a regular-jump continuous time homogeneous Markov chain.

Similarly to before, we denote by \mathcal{X} the space state of \mathbf{X} and with Δ the set of absorbing states, which correspond to the states in which all nodes are either R or B . Recall that $\{\mathbf{X}_k\}_{k \in \mathbb{N} \cup \{0\}}$ is the embedded chain defined by $\mathbf{X}_k = \mathbf{X}(T_k)$. Note that, conditionally on $\{\mathbf{X}_k = \mathbf{x} \in \mathcal{X} \setminus \Delta\}$, the sojourn time $W_k = T_{k+1} - T_k$ in state \mathbf{X}_k is exponentially distributed with parameter

$$(3.12) \quad q_{\mathbf{x}} := R(\mathbf{x}) = R^R(\mathbf{x}) + R^B(\mathbf{x})$$

where

$$(3.13) \quad R^S(\mathbf{x}) := Q^S(\mathbf{x}) \mathbf{1}_{\{K^*-1 > k\}}(\mathbf{x}) + (n_W - N(\mathbf{x})) U^S(\mathbf{x}) \mathbf{1}_{\{K^*-1 \leq k\}}(\mathbf{x}).$$

$R^S(\mathbf{x})$ is the aggregate rate of all transitions from state \mathbf{x} induced by the S -activation of a new node. Propositions 3.3, 3.4 and 3.5 naturally extend also to the prolonged process, by redefining for $k \in \mathbb{N} \cup \{0\}$:

$$M_{k+1}^S := \mathbf{1}_{\{V_{k+1} \in \mathcal{V}_W[k] \cap \mathcal{S}_S[k]\}} \mathbf{1}_{\{K^*-1 > k\}} + \mathbf{1}_{\{K^*-1 \leq k\}} \mathbf{1}_{\{L_{k+1} < U_{k+1}^S\}} \mathbf{1}_{\{\mathbf{x}_k \in \mathcal{X} \setminus \Delta\}}$$

where L_{k+1} is a random variable uniformly distributed on $[0, 1]$, independent of \mathcal{H}_k . In particular, since, as already noticed, $\mathbf{1}_{\{\mathbf{x}_k \in \mathcal{X} \setminus \Delta\}} = \mathbf{1}_{\{k < n_W\}}$, we have for $S \in \{R, B\}$:

$$\text{PROPOSITION 3.8. } \mathbb{P}(M_{k+1}^S = 1 \mid \mathcal{H}_k) = U_{k+1}^S \mathbf{1}_{\{\mathbf{x}_k \in \mathcal{X} \setminus \Delta\}} = U_{k+1}^S \mathbf{1}_{\{k < n_W\}}.$$

3.4.1. *Properties of the extended process.* By definition (see (3.3)), we have

$$|\mathcal{S}_S[k]| = \sum_{v \in \mathcal{V}_W} \mathbf{1}_{\{D_S^{(v)}[k] - D_{S^c}^{(v)}[k] \geq r\}}, \quad k \in \mathbb{N} \cup \{0\},$$

and recalling (3.2) it follows

$$(3.14) \quad |\mathcal{S}_S[k]| \mid \{\mathbf{N}[k] = \mathbf{k}\} \stackrel{L}{=} \text{Bin}(n_W, \pi_S(\mathbf{k})),$$

where $\mathbf{k} := (k_R, k_B) \in (\mathbb{N} \cup \{0\})^2$, $k := k_R + k_B \leq n_W$, $\mathbf{N}[k] := (N_R[k], N_B[k])$, and

$$(3.15) \quad \pi_S(\mathbf{k}) := \mathbb{P}(\text{Bin}(k_S + a_S, p) - \text{Bin}(k_{S^c} + a_{S^c}, p) \geq r).$$

Moreover defined for $k \leq n_W$ and $h \leq k$:

$$\mathcal{N}_{h,k} := \{N_R[k] \geq k - h, N_B[k] \leq h\} = \{N_R[k] \geq k - h\} = \{N_B[k] \leq h\},$$

LEMMA 3.9. It holds:

$$|\mathcal{S}_R[k]| \mid \mathcal{N}_{h,k} \geq_{st} \text{Bin}(n_W, \pi_R(k - h, h)); \quad |\mathcal{S}_B[k]| \mid \mathcal{N}_{h,k} \leq_{st} \text{Bin}(n_W, \pi_B(k - h, h)).$$

The proof of Lemma 3.9 is elementary (it is reported in Appendix D for completeness).

REMARK 3.10. Note that, for $q \ll p^{-1}$, $\beta_S(x_R, x_B) = \beta_S(x_S)$, i.e. β_S does not depend on x_{S^c} ; while, for $q = p^{-1}$ and $p^{-1} \ll q$, the function β_S depends on both x_R and x_B . The fact that, for $q \ll p^{-1}$, β_S does not depend on x_{S^c} expresses formally that the two activation processes N_R and N_B evolve essentially independently on time-scales q' which are asymptotically less than p^{-1} . On the other hand, the fact that, for $q = p^{-1}$ and $p^{-1} \ll q$, β_S

depends on both x_R and x_B expresses formally that the two activation processes N_R and N_B interact on time-scales which are comparable with p^{-1} or are asymptotically bigger than p^{-1} . Indeed, roughly speaking, given that x_{Sq} nodes have been S -active, by (3.2) we have that the number of S -marks collected by a node $v \in \mathcal{V}_W$, $D_S^{(v)}(x_{Sq})$, is binomially distributed and the average $x_{Sq}p$ tends to 0, whenever $q \ll p^{-1}$. Therefore, only a negligible fraction of the S -susceptible nodes (i.e. nodes $v \in \mathcal{V}_W$, for which $D_S^{(v)}(x_{Sq}) - D_{S^c}^{(v)}(x_{Sq}) \geq r$) got more than the minimum number of marks, (i.e., r marks of color S and 0 of color S^c , as shown in the proof of Lemma E.4). In conclusion, the number of S -susceptible nodes is not significantly impacted by the presence of S^c -marks. In other words, the two activation processes evolve without significantly interfering, because they insist on different sets of nodes. Instead, when q gets comparable with p^{-1} , both the number of S -susceptible nodes, as well as, the number of S -marks that have been distributed turns out to be of order n . This implies that the fraction of S -susceptible nodes that received marks of color S^c is not anymore negligible, and the activation processes N_R and N_B start interacting.

3.5. Brief overview of main proofs. As a guide to the reader, we briefly describe, at high level, the strategy of the proofs. First, we analyze the activation process on time-scale q , i.e., we analyze the asymptotic behavior of $\mathbf{N}_S[\lfloor xq \rfloor]/q$ for bounded values of x .

The main result on time-scale q is provided by Theorem 4.2, which shows that a suitable regularized version of the trajectories $\mathbf{N}_S[\lfloor xq \rfloor]/q$ converges \mathbb{P} -a.s to the (deterministic) solution of the Cauchy Problem (CP) stated in Definition 4.1. To prove convergence of above trajectories, we exploit their structural properties, resorting to Ascoli-Arzelà' theorem to claim their \mathbb{P} -a.s. pointwise convergence to a weak limit (i.e., we show the convergence of some sub-sequence). Then, we provide sufficiently tight upper and lower bounds to the incremental ratio of trajectories within a neighborhood of a fixed point. By so doing we show that limit trajectories are derivable (with assigned derivative), and therefore solution of the CP formulated in Definition 4.1. As side effect, given the uniqueness of CP solutions, we are able to strengthen previous convergence result showing a \mathbb{P} -a.s. pointwise convergence for the whole sequence. Theorem 4.7 complements previous results showing that suitably normalized versions of both $T_{\lfloor xq \rfloor}$ and $T_{\lfloor xq \rfloor}^S$ converge almost surely to deterministic quantities.

When the activation processes of nodes do not stop at time-scale q , (i.e., in the supercritical regime) we complement previous study considering time-scales larger than q . In this case, analyzing the solutions of the Cauchy problem (defined in 4.2) we show that the ratio $N_B[\lfloor xq \rfloor]/N[\lfloor xq \rfloor]$ becomes arbitrarily small as x grows large.

The analysis at time-scales $q' \gg q$ is based on the observation that the number of S -susceptible nodes, $|S_S(t)|$, is sufficiently concentrated around its average, which in turn depends super-linearly on the number of active nodes $N_S(t)$. Therefore, as shown in Theorems 5.1, 5.2 and 5.3, the ratio between the rates at which the two competing activation processes evolve tends quickly to diverge, letting the advantaged R -process to percolate before the competing B -process has managed to activate a non negligible fraction of nodes. In particular, for the case $q = g$ we can show that $A_B^* = O(g)$. This is done: (i) by first analyzing the dynamics of an auxiliary process, the stopped process, where the R -activation process is stopped at a given point and only the B -activation process is allowed to go on; (ii) by then inferring properties on the original process, exploiting a simple coupling argument (8.1).

4. Analysis at time-scale q : main results. In this section we report the main findings of our analysis about the activation process N_S , $S \in \{R, B\}$, when $N = \Theta(q)$, i.e., it is of the same order of the number of seeds. We remark that in the following we will always assume, without lack of generality, $\alpha_R > \alpha_B$. Proofs of results stated in this section are reported in Sec. 7.

Hereon, we start introducing the linear interpolation of N_S defined by:

$$(4.1) \quad \tilde{N}_S(xq) := N_S[\lfloor xq \rfloor] + (xq - \lfloor xq \rfloor) (N_S[\lceil xq \rceil] - N_S[\lfloor xq \rfloor]), \quad x \geq 0$$

with $\tilde{\mathbf{N}}(xq) = (\tilde{N}_R(xq), \tilde{N}_B(xq))$, and the sequence $\{\mathbf{F}_n(x)\}_{n \in \mathbb{N}}$ given by:

$$\mathbf{F}_n(x) := (F_{R,n}(x), F_{B,n}(x)) \quad \text{with} \quad F_{S,n}(x) := \frac{\tilde{N}_S(xq_n)}{q_n}.$$

As usual, when no confusion arises, we omit the dependence on n of \mathbf{F}_n and $F_{S,n}$. It turns out (see Theorem 4.2) that, under suitable assumptions, \mathbf{F} converges to a vectorial function \mathbf{f} , which is the solution of the following Cauchy problem:

DEFINITION 4.1. (Cauchy problem). Let $\mathbf{f}(x) = (f_R(x), f_B(x))$ denote the maximal solution of the Cauchy problem:

$$(4.2) \quad \mathbf{f}'(x) = \frac{\boldsymbol{\beta}(\mathbf{f}(x))}{\beta_R(\mathbf{f}(x)) + \beta_B(\mathbf{f}(x))}, \quad x \in (0, \kappa_{\mathbf{f}}), \quad \mathbf{f}(0) = (0, 0),$$

with $\boldsymbol{\beta}(\mathbf{x}) := \boldsymbol{\beta}(x_R, x_B) := (\beta_R(x_R, x_B), \beta_B(x_R, x_B))$.

THEOREM 4.2. Assume (2.1) and (2.2) with $\alpha_R > \alpha_B$ and let \mathbf{f} be the solution of the Cauchy problem (4.2). Then:

(i) If $q = g$ with $\alpha_R < 1$, then

$$(4.3) \quad \text{For any compact set } \mathbb{K} \subset [0, z_R + z_B], \sup_{x \in \mathbb{K}} \|\mathbf{F}(x) - \mathbf{f}(x)\| \rightarrow 0, \quad \mathbb{P}\text{-a.s.},$$

where z_S is the first zero of $\beta_S(x_S)$.

(ii) If $\alpha_R > 1$ and either $q = g$ or $q \gg g$, then

$$(4.4) \quad \text{For any compact set } \mathbb{K} \subset [0, \infty), \sup_{x \in \mathbb{K}} \|\mathbf{F}(x) - \mathbf{f}(x)\| \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

As immediate consequence of previous theorem we have:

COROLLARY 4.3. For every $\kappa < \kappa_{\mathbf{f}}$ and $S \in \{R, B\}$:

$$(4.5) \quad \lim_{q \rightarrow \infty} \frac{\tilde{N}_S(\kappa q)}{q} = f_S(\kappa), \quad \mathbb{P}\text{-a.s.}$$

4.1. *On the solution of the Cauchy problem (4.2).* Recalling that \mathbf{g} is the maximal solution of Cauchy problem (2.6), the following proposition, whose proof is reported in Appendix A, holds:

PROPOSITION 4.4. (i) For $q = g$ and $\alpha_R < 1$, \mathbf{f} is defined on $(0, z_R + z_B)$ (i.e., $\kappa_{\mathbf{f}} = z_R + z_B$) and

$$f_R(x) \uparrow z_R, \quad f_B(x) \uparrow z_B, \quad \text{as } x \uparrow z_R + z_B.$$

(ii) For $q = g$ and $\alpha_R > 1$, or $g \ll q \ll p^{-1}$, then \mathbf{f} is defined on $(0, \infty)$ (i.e. $\kappa_{\mathbf{f}} = \infty$) and

$$f_R(x) \uparrow +\infty, \quad f_B(x) \uparrow g_B(\kappa_{\mathbf{g}}) < \infty \quad \text{as } x \uparrow +\infty,$$

with $\kappa_{\mathbf{g}} := \int_0^\infty \frac{dx}{\beta_R(x)} \in (0, \infty)$, and $g_B(\kappa_{\mathbf{g}}) := \lim_{x \uparrow \kappa_{\mathbf{g}}} g_B(x)$.

(iii) For $q = p^{-1}$, then \mathbf{f} is defined on $(0, \infty)$ and

$$f_R(x) \uparrow +\infty, \quad f_B(x) \uparrow \bar{f}_B, \quad \text{as } x \uparrow +\infty,$$

for some constant $\bar{f}_B \in (0, \infty)$.

(iv) For $p^{-1} \ll q \ll n$ and $\alpha_R > \alpha_B$, then \mathbf{f} is defined on $(0, \infty)$ and is given by

$$f_R(x) := x, \quad f_B(x) := 0.$$

Moreover if $q = g$ and $\alpha_B \leq 1 < \alpha_R$ then $g_B(\kappa_{\mathbf{g}}) < z_B$.

4.2. *Analysis of K^* and A_B^* .* The following two theorems complement previous results.

THEOREM 4.5. (i) For every $\kappa < \kappa_{\mathbf{f}}$:

$$(4.6) \quad \liminf \frac{K^*}{q} > \kappa, \quad \mathbb{P}\text{-a.s.}$$

Moreover (ii) if either $q = g$ and $\alpha_R > 1$ or $g \ll q \ll p^{-1}$, then

$$(4.7) \quad \liminf \frac{A_B^*}{q} \geq g_B(\kappa_{\mathbf{g}}) + \alpha_B, \quad \mathbb{P}\text{-a.s.}$$

where $g_B(\kappa_{\mathbf{g}})$ and $\kappa_{\mathbf{g}}$ are given in Proposition 4.4 (ii).

THEOREM 4.6. Assume $q = g$ and $\alpha_S < 1$. Then, for any $S \in \{R, B\}$,

$$\limsup \frac{A_S^*}{q} \leq z_S + \alpha_S, \quad \mathbb{P}\text{-a.s.}$$

4.3. *Analysis of the sequences $\{T_k\}_{k \in \mathbb{N}}$ and $\{T_k^S\}_{k \in \mathbb{N}}$ at time-scale q .* The next result provides the asymptotic behavior of $T_{\lfloor \kappa q \rfloor}$ and $T_{\lfloor \kappa_S q \rfloor}^S$, for suitable $\kappa > 0$ and κ_S . Let

$$(4.8) \quad \eta := \begin{cases} 1 & \text{for } q = g \\ \frac{n(qp)^r}{q} & \text{for } g \ll q \ll p^{-1} \\ \frac{n}{q} & \text{for either } q = p^{-1} \text{ or } q \gg p^{-1}. \end{cases}$$

THEOREM 4.7. (i) Let $\kappa < \kappa_{\mathbf{f}}$ (with $\kappa_{\mathbf{f}}$ defined in Proposition 4.4). Then:

$$(4.9) \quad \eta T_{\lfloor \kappa q \rfloor} \rightarrow \int_0^\kappa \frac{1}{\beta_R(\mathbf{f}(y)) + \beta_B(\mathbf{f}(y))} dy, \quad \mathbb{P}\text{-a.s.}$$

(ii) Let $\kappa_S \in (0, \lim_{x \rightarrow \kappa_{\mathbf{f}}} f_S(x))$ for $S \in \{R, B\}$. Then:

$$(4.10) \quad \eta T_{\lfloor \kappa_S q \rfloor}^S \rightarrow \int_0^{f_S^{-1}(\kappa_S)} \frac{1}{\beta_R(\mathbf{f}(y)) + \beta_B(\mathbf{f}(y))} dy.$$

When $q \ll p^{-1}$, from (4.2) it follows that: $\int_0^{f_S^{-1}(\kappa_S)} \frac{1}{\beta_R(\mathbf{f}(y)) + \beta_B(\mathbf{f}(y))} dy = \int_0^{\kappa_S} \frac{1}{\beta_S(y)} dy$.

5. Analysis at time-scales greater than q : main results. In this section we study the joint evolution of $\mathbf{N}[\cdot]$ and $(|\mathcal{S}_R[\cdot]|, |\mathcal{S}_B[\cdot]|)$, at time-scales $q' \gg q$, i.e., for arguments asymptotically greater than the number of seeds. Recalling that function g_B and constant $\kappa_{\mathbf{g}}$ are given in Proposition 4.4(ii), the following theorems, whose proofs are given in Sec. 8, hold.

THEOREM 5.1. If either $q = g$ and $\alpha_R > 1$, or $g \ll q \ll p^{-1}$, then $\forall \varepsilon > 0$, we have:

$$(5.1) \quad \mathbb{P}(\liminf \{N_B[\lfloor f(n)p^{-1} \rfloor]\} \leq \lfloor (g_B(\kappa_{\mathbf{g}}) + \varepsilon)q \rfloor \cap \{K^* - 1 \geq \lfloor f(n)p^{-1} \rfloor\}) = 1,$$

where f is a generic function such that $f(n) \rightarrow \infty$ and $f(n)p^{-1} = o(n)$ in case (i); $f(n) := c_0/(qp)^{r-1} \rightarrow \infty$, for a sufficiently small positive constant c_0 in case (ii).

THEOREM 5.2. Assume $q = g$ and $\alpha_R > 1$. Then, $\forall \varepsilon > 0$ and $c \in (0, 1)$ we have:

$$(5.2) \quad \mathbb{P}(\liminf \{N_B[K^* - 1] \leq \lfloor (g_B(\kappa_{\mathbf{g}}) + \varepsilon)g \rfloor \cap \{K^* - 1 \geq \lfloor cn \rfloor\}) = 1.$$

THEOREM 5.3. Assume $g \ll q \ll n$. Then, $\forall c \in (0, 1)$, we have

$$(5.3) \quad \mathbb{P} \left(\liminf \{K^* - 1 \geq \lfloor cn \rfloor\} \cap \left\{ \lim \frac{N_B[\lfloor cn \rfloor]}{cn} = 0 \right\} \right) = 1.$$

6. Proofs of Theorems 2.4 and 2.5. The proofs of Theorems 2.4 and 2.5 are immediate consequences of previously mentioned results.

6.1. *Proof of Theorem 2.4.* By Theorems 4.6 and 4.5(i) (with $\kappa_f = z_R + z_B$), we have

$$\begin{aligned} z_S + \alpha_S &\geq \limsup \frac{A_S^*}{q} \geq \liminf \frac{A_S^*}{q} \geq \liminf \left(\frac{A^*}{q} - \frac{A_{S^c}^*}{q} \right) \geq \liminf \frac{A^*}{q} + \liminf \left(-\frac{A_{S^c}^*}{q} \right) \\ &\geq z_R + z_B + \alpha_R + \alpha_B - \limsup \frac{A_{S^c}^*}{q} \geq z_S + \alpha_S. \quad \square \end{aligned}$$

6.2. *Proof of Theorem 2.5.* We start focusing on the case (i). By Theorem 5.2, for any $\varepsilon > 0$ \mathbb{P} -a.s. there exists a $n'(\omega)$ such that $N_B[K^* - 1] \leq (g_B(\kappa_g) + \varepsilon)q$, $\forall n > n'(\omega)$. Therefore

$$\limsup \frac{A_B^*}{q} = \limsup \frac{N_B[K^* - 1]}{q} + \alpha_B \leq g_B(\kappa_g) + \alpha_B, \quad \mathbb{P}\text{-a.s.}$$

The second relation in (2.5) descends immediately from this upper bound and the matching lower bound (4.7). As far as the first relation in (2.5) is concerned, we note that (5.2) implies that, for any $c \in (0, 1)$ we have \mathbb{P} -a.s.,

$$\liminf \frac{A_R^*}{n} = \liminf \frac{N_R[K^* - 1]}{n} = \liminf \frac{K^* - N_B[K^* - 1]}{n} \geq c,$$

here, the final inequality follows from: (5.2), indeed, \mathbb{P} -a.s., a $n'(\omega)$ can be found such that jointly $K^* - 1 \geq \lfloor cn \rfloor$ and $N_B[K^* - 1] \leq (g_B(\kappa_g) + \varepsilon)g$ for all $n > n'(\omega)$. The claim immediately follows by the arbitrariness of $c \in (0, 1)$.

Now turning our attention to case (ii), second inequality; we recall that, by construction, $N_B[K^* - 1] - N_B[\lfloor cn \rfloor] \leq \max(0, K^* - 1 - \lfloor cn \rfloor) \leq n - \lfloor cn \rfloor$, therefore \mathbb{P} -a.s. we have

$$\limsup \frac{A_B^*}{n} = \limsup \frac{N_B[K^* - 1]}{n} \leq \limsup \frac{(1-c)n}{n} + \frac{N_B[\lfloor cn \rfloor]}{n} = 1 - c,$$

given that, by (5.3), $\frac{N_B[\lfloor cn \rfloor]}{n} \rightarrow 0$. The result follows from the arbitrariness of c . The first inequality can be proved exactly as in previous case (i). \square

7. Proofs of Theorems 4.2, 4.5, 4.6 and 4.7. The proofs of Theorems 4.2- 4.7 are based on some ancillary preliminary concentration results. In this section we limit ourselves to state these results and postpone their (rather standard) proofs to Appendices C, E, F, G.

7.1. *Preliminaries.* Let $\mathbf{k} := (k_R, k_B) \in (\mathbb{N} \cup \{0\})^2$ and $k = k_R + k_B$, we define the following sets:

$$(7.1) \quad \mathbb{I}_k := \{\mathbf{k} : k_R + k_B = k\}, \quad k \in \mathbb{N} \cup \{0\}$$

Recalling the definition of κ_f in Proposition 4.4, for any $\kappa < \kappa_f$, we define:

$$\mathbb{T}(\kappa) := \left\{ \begin{array}{l} \{\mathbf{k} : k_R + k_B \leq \kappa q\} = \bigcup_{0 \leq k < \kappa q} \mathbb{I}_k \quad \text{for } q \ll p^{-1} \text{ or } q = p^{-1} \\ \{\mathbf{k} : k_R + k_B \leq \kappa q, \frac{k_R + \alpha_R q}{k_B + \alpha_B q} \leq \frac{1}{2} + \frac{\alpha_R}{2\alpha_B}\} \quad \text{for } q \gg p^{-1}. \end{array} \right.$$

Hereafter, we consider the following random variables

$$(7.2) \quad \Upsilon_S(\kappa) := \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} Y_S(\mathbf{k}), \quad \bar{\Upsilon}_S(\kappa) := \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \bar{Y}_S(\mathbf{k}), \quad \mathfrak{N}_S(\kappa) := \frac{\sup_{j \leq \kappa q} |\widehat{N}_S[j]|}{q}$$

where

$$Y_S(\mathbf{k}) := \mathbf{1}_{\{\mathbf{N}(k)=\mathbf{k}\}} \left| U_{k+1}^S - \frac{|\beta_S(\mathbf{k}/q)|}{|\beta_R(\mathbf{k}/q)| + |\beta_B(\mathbf{k}/q)|} \right|,$$

$$\bar{Y}_S(\mathbf{k}) := \mathbf{1}_{\{\mathbf{N}(k)=\mathbf{k}\}} |Q_{k+1}^S - \eta \beta_S(\mathbf{k})q|$$

with η defined as in (4.8).

$$(7.3) \quad \widehat{N}_S[j] := N_S[j] - J_S[j], \quad J_S[j] := \sum_{h=1}^{\min(j, n_w)} U_h^S \quad \forall j \in \mathbb{N}, \quad \widehat{N}(0) := 0.$$

Furthermore, letting $\mathbf{x} = (x_R, x_B) \in [0, \infty)^2$, we introduce the compact sets:

$$(7.4) \quad \mathbb{T}'(\kappa) := \begin{cases} \{\mathbf{x} : x_R + x_B \leq \kappa\} & \text{for } q \ll p^{-1} \text{ or } q = p^{-1} \\ \left\{ \mathbf{x} : x_R + x_B \leq \kappa \text{ and } \frac{x_R + \alpha_R}{x_B + \alpha_B} \geq \frac{1}{2} + \frac{\alpha_R}{2\alpha_B} \right\} & \text{for } q \gg p^{-1}. \end{cases}$$

Moreover, given κ as before, let $z > 0$ be a constant such that $2z < \kappa$ and $\ell \in \mathbb{T}(\kappa - 2z)$, and

$$(7.5) \quad \mathbb{L}_\ell(\kappa, z) := \{\mathbf{x} : x_R \geq \ell_R - z/2, x_B \geq \ell_B - z/2, x_R + x_B \leq \ell_R + \ell_B + 2z\}.$$

At last, if either $q \ll p^{-1}$ or $q = p^{-1}$, we set

$$(7.6) \quad \bar{\beta}_{S, \mathbb{L}_\ell(\kappa, z)} := \max_{\mathbf{x} \in \mathbb{L}_\ell(\kappa, z)} \frac{|\beta_S(\mathbf{x})|}{|\beta_R(\mathbf{x})| + |\beta_B(\mathbf{x})|}, \quad \underline{\beta}_{S, \mathbb{L}_\ell(\kappa, z)} := \min_{\mathbf{x} \in \mathbb{L}_\ell(\kappa, z)} \frac{|\beta_S(\mathbf{x})|}{|\beta_R(\mathbf{x})| + |\beta_B(\mathbf{x})|}$$

whereas for $q \gg p^{-1}$, we set

$$(7.7) \quad \bar{\beta}_{S, \mathbb{L}_\ell(\kappa, z)} := \max_{\mathbf{x} \in \mathbb{L}_\ell(\kappa, z)} \frac{|\beta_S(\mathbf{x})|}{|\beta_R(\mathbf{x})| + |\beta_B(\mathbf{x})|} \mathbf{1}_{\{\mathbb{L}_\ell(\kappa, z) \subseteq \mathbb{T}'(\kappa)\}} + \mathbf{1}_{\{\mathbb{L}_\ell(\kappa, z) \not\subseteq \mathbb{T}'(\kappa)\}}$$

and

$$(7.8) \quad \underline{\beta}_{S, \mathbb{L}_\ell(\kappa, z)} := \min_{\mathbf{x} \in \mathbb{L}_\ell(\kappa, z)} \frac{|\beta_S(\mathbf{x})|}{|\beta_R(\mathbf{x})| + |\beta_B(\mathbf{x})|} \mathbf{1}_{\{\mathbb{L}_\ell(\kappa, z) \subseteq \mathbb{T}'(\kappa)\}}.$$

The proof of Theorem 4.2 makes use of the following Propositions 7.1 and 7.2, whose proofs are rather standard and reported in the Appendices E and F, respectively.

PROPOSITION 7.1. Given η in (4.8) we have:

$$(7.9) \quad \max\{\Upsilon_S(\kappa), (\eta q)^{-1} \bar{\Upsilon}_S(\kappa), \mathfrak{N}_S(\kappa)\} \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

Hereafter, for $\kappa > 0$, we set

$$(7.10) \quad \Omega_\kappa := \{\omega \in \Omega : \max\{\Upsilon_S(\kappa)(\omega), (\eta q)^{-1} \bar{\Upsilon}_S(\kappa)(\omega), \mathfrak{N}_S(\kappa)(\omega)\} \rightarrow 0\}.$$

PROPOSITION 7.2. For any $y, z > 0$ such that $y + 2z \leq \kappa < \kappa_f$ (with κ_f given in Proposition 4.4), we have:

$$(7.11) \quad z \liminf \sum_{\mathbf{k} \in \mathbb{I}_{[yq]}} \underline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} \mathbf{1}_{\{\mathbf{N}[\lfloor yq \rfloor] = \mathbf{k}\}} \leq \liminf \frac{\tilde{N}_S(yq + zq) - \tilde{N}_S(yq)}{q}$$

$$\leq \limsup \frac{\tilde{N}_S(yq + zq) - \tilde{N}_S(yq)}{q} \leq z \limsup \sum_{\mathbf{k} \in \mathbb{I}_{[yq]}} \bar{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} \mathbf{1}_{\{\mathbf{N}[\lfloor yq \rfloor] = \mathbf{k}\}}$$

7.2. *Proof of Theorem 4.2.* We divide the proof in two steps. In the first step we assume either $q \ll p^{-1}$ or $q = p^{-1}$. In the second step we extend the proof to the case $p^{-1} \ll q \ll n$. *Step 1.* Since by Proposition 7.1 we have $\mathbb{P}(\Omega_\kappa) = 1$, it suffices to prove that limit (4.3) holds for all $\omega \in \Omega_\kappa$. For $x_1, x_2 \in [0, \kappa]$ such that $x_1 > x_2$ and $\omega \in \Omega_\kappa$, we have

$$\begin{aligned} F_S(x_1, \omega) - F_S(x_2, \omega) &= q^{-1} \left(\tilde{N}_S(x_1 q)(\omega) - \tilde{N}_S(x_2 q)(\omega) \right) \\ &\leq q^{-1} \left(x_1 q - \lfloor x_1 q \rfloor + N_S[\lfloor x_1 q \rfloor](\omega) - N_S[\lfloor x_2 q \rfloor](\omega) + \lfloor x_2 q \rfloor - x_2 q \right) \\ &\leq q^{-1} (x_1 q - \lfloor x_1 q \rfloor + \lfloor x_1 q \rfloor - \lfloor x_2 q \rfloor + \lfloor x_2 q \rfloor - x_2 q) \\ &= x_1 - x_2, \end{aligned}$$

where we have used the inequality $N_S[j_1] - N_S[j_2] \leq j_1 - j_2$, for any $j_1 \geq j_2$, $j_1, j_2 \in \mathbb{N} \cup \{0\}$. So, for $x_1, x_2 \in [0, \kappa]$ and $\omega \in \Omega_\kappa$,

$$|F_S(x_1, \omega) - F_S(x_2, \omega)| \leq |x_1 - x_2|.$$

Moreover, for any $x \in [0, \kappa]$,

$$(7.12) \quad F_S(x, \omega) = \frac{\tilde{N}_S(xq)(\omega)}{q} \leq q^{-1}(xq) = x \leq \kappa.$$

Thus, for any $\omega \in \Omega_\kappa$, the functions $F_S(\cdot, \omega)$ are Lipschitz continuous with Lipschitz constant less than or equal to 1 and uniformly bounded. Consequently, by the Ascoli-Arzelá theorem there exists a subsequence $\{F_{S, n'}(\cdot, \omega)\}_{n'}$ converging uniformly on $[0, \kappa]$ to some function $f_S(\cdot, \omega)$ (which, clearly, is also Lipschitz continuous with Lipschitz constant less than or equal to 1 and bounded by κ). From now on, to avoid confusion, we explicit the dependence on n . We now prove that $f_S(\cdot, \omega)$ is differentiable on $(0, \kappa)$ and compute its derivative. Note that, for an arbitrarily fixed $x \in (0, \kappa)$ and $z \in (x, \frac{\kappa+x}{2})$, we have

$$\begin{aligned} f_S(z, \omega) - f_S(x, \omega) &= \lim_{n' \rightarrow \infty} [F_{S, n'}(z, \omega) - F_{S, n'}(x, \omega)] \\ &= \limsup_{n' \rightarrow \infty} q_{n'}^{-1} [\tilde{N}_S(xq_{n'} + (z-x)q_{n'})(\omega) - \tilde{N}_S(xq_{n'})(\omega)] \\ (7.13) \quad &\leq (z-x) \lim_{n' \rightarrow \infty} \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor xq_{n'} \rfloor}} \bar{\beta}_{S, \mathbb{I}_{\mathbf{k}/q_{n'}}}(\kappa, z-x) \mathbf{1}_{\{\mathbf{N}[\lfloor xq_{n'} \rfloor](\omega) = \mathbf{k}\}} \end{aligned}$$

where the last inequality descends from Proposition 7.2 (we refer the reader to (7.5) for the definition of the set $\mathbb{I}_{\cdot}(\cdot, \cdot)$). Now, defined $x_{n'} := \frac{\lfloor xq_{n'} \rfloor}{q_{n'}}$, we have

$$N[\lfloor xq_{n'} \rfloor] = \tilde{N}_S(\lfloor xq_{n'} \rfloor)(\omega) = \tilde{N}_S(x_{n'} q_{n'})(\omega) = F_{S, n'}(x_{n'}, \omega) q_{n'}.$$

and

$$F_{S, n'}(x, \omega) - \frac{1}{q_{n'}} \leq F_{S, n'}(x, \omega) - (x - x_{n'}) \leq F_{S, n'}(x_{n'}, \omega) \leq F_{S, n'}(x, \omega),$$

from which it follows:

$$(7.14) \quad \lim_{n' \rightarrow \infty} \mathbf{F}_{n'}(x_{n'}, \omega) = \lim_{n' \rightarrow \infty} \mathbf{F}_{n'}(x, \omega) = \mathbf{f}(x, \omega).$$

Therefore, for any $\omega \in \Omega_\kappa$, we have

$$\begin{aligned} f_S(z, \omega) - f_S(x, \omega) &\leq (z-x) \limsup_{n' \rightarrow \infty} \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor xq_{n'} \rfloor}} \bar{\beta}_{S, \mathbb{I}_{\mathbf{k}/q_{n'}}}(\kappa, z-x) \mathbf{1}_{\{\mathbf{N}[x_{n'} q_{n'}](\omega) = \mathbf{k}\}} \\ (7.15) \quad &= (z-x) \limsup_{n' \rightarrow \infty} \bar{\beta}_{S, \mathbb{I}_{\mathbf{F}_{n'}(x_{n'}, \omega)}}(\kappa, z-x) = (z-x) \bar{\beta}_{S, \mathbb{I}_{\mathbf{f}(x, \omega)}}(\kappa, z-x), \end{aligned}$$

where the second equality follows by (7.14) and the continuity of the function $u \mapsto \bar{\beta}_{S, \mathbb{L}_u(\kappa, z-x)}$. Similarly, exploiting the relation (7.11), for any $\omega \in \Omega_\kappa$, we have

$$f_S(z, \omega) - f_S(x, \omega) \geq (z-x) \underline{\beta}_{S, \mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)}, \quad \forall z \in \left(x, \frac{\kappa+x}{2}\right).$$

Thus, for any $\omega \in \Omega_\kappa$, any $x \in (0, \kappa)$ and any $z \in \left(\frac{\kappa+x}{2}, \kappa\right)$, we have

$$(7.16) \quad \frac{f_S(z, \omega) - f_S(x, \omega)}{z-x} \leq \bar{\beta}_{S, \mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)}, \quad \frac{f_S(z, \omega) - f_S(x, \omega)}{z-x} \geq \underline{\beta}_{S, \mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)}.$$

Since the set $\mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)$ is compact, it holds

$$\bar{\beta}_{S, \mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)} = \frac{|\beta_S(\mathbf{v})|}{|\beta_R(\mathbf{v})| + |\beta_B(\mathbf{v})|} \quad \text{and} \quad \underline{\beta}_{S, \mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)} = \frac{|\beta_S(\mathbf{w})|}{|\beta_R(\mathbf{w})| + |\beta_B(\mathbf{w})|},$$

for some $\mathbf{v} = (v_R, v_B), \mathbf{w} = (w_R, w_B) \in \mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)$. By the definition of the set $\mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x)$ it follows that

$$(7.17) \quad v_R, w_R \rightarrow f_R(x, \omega) \quad \text{and} \quad v_B, w_B \rightarrow f_B(x, \omega), \quad \text{as } z \downarrow x.$$

Therefore, taking the limsup as $z \downarrow x$ in (7.16) and the liminf as $z \downarrow x$ in (7.16), by (7.17) and the continuity of β_S , we have that the right-hand derivative of $f_S(\cdot, \omega)$ at $x \in (0, \kappa)$ is:

$$(7.18) \quad f_S^+(x, \omega) = \varphi_S(x, \omega) := \frac{\beta_S(f_R(x, \omega), f_B(x, \omega))}{\beta_R(f_R(x, \omega), f_B(x, \omega)) + \beta_B(f_R(x, \omega), f_B(x, \omega))}.$$

Since, for fixed $\omega \in \Omega_\kappa$, the functions $f_S(\cdot, \omega)$ and $\varphi_S(\cdot, \omega)$ are continuous on $[0, \kappa]$, and $\varphi_S(\cdot, \omega)$ by (7.18) is the right-hand derivative of $f_S(\cdot, \omega)$ on $(0, \kappa)$, we have that $f_S^+(0, \omega) = \varphi_S(0, \omega)$. Moreover $\varphi_S(\cdot, \omega)$ is the derivative of $f_S(\cdot, \omega)$ on $(0, \kappa)$ (see e.g. Theorem A22 p. 541 of [7]). At last, given that by construction $\mathbf{f}(0, \omega) = (0, 0)$, we conclude that $\mathbf{f}(\cdot, \omega) = \mathbf{f}(\cdot)$ is the unique solution of the Cauchy problem (4.2).

Due to the uniqueness of the solution of the Cauchy problem (4.2), for any $\omega \in \Omega_\kappa$, the whole sequence $\{\mathbf{F}_n(\cdot, \omega)\}_n$ converges to $\mathbf{f}(\cdot)$. Indeed by repeating previous arguments, it can be immediately shown that any converging sub-sequence $\{\mathbf{F}_{n''}(\cdot, \omega)\}_{n''}$ of $\{\mathbf{F}_n(\cdot, \omega)\}_n$ (among which the sub-sequences achieving the limsup and liminf) must converge to the unique solution of the Cauchy problem (4.2), $\mathbf{f}(\cdot)$. Finally, since $\mathbf{F}(\cdot, \omega)$ and $\mathbf{f}(\cdot)$ are both Lipschitz continuous on $[0, \kappa]$, \mathbb{P} -a.s., the convergence $\mathbf{F}(\cdot, \omega) \rightarrow \mathbf{f}(\cdot)$ is uniform on $[0, \kappa]$.

Step 2. Since the function $\beta_S(\mathbf{x})$ is discontinuous at the points (x_R, x_B) such that $\frac{x_R + \alpha_R}{x_B + \alpha_B} = 1$, we have that the mapping $u \mapsto \bar{\beta}_{S, \mathbb{L}_u(\kappa, z-x)}$ is not always continuous. However, the function $u \mapsto \bar{\beta}_{S, \mathbb{L}_u(\kappa, z-x)}$ is continuous on $\mathbb{T}'(\kappa)$ (as defined in (7.4)) if $z-x$ is such that $\mathbb{L}_u(\kappa, z-x) \subseteq \mathbb{T}'(\kappa)$. By Proposition 4.4(iv) we have $\mathbf{f}(x) \in \mathbb{T}'(\kappa)$ if $x < \kappa$. Moreover, as long as $\mathbf{f}(x) \in \mathbb{T}'(\kappa)$ we can always make $z-x$ so small that $\mathbb{L}_{\mathbf{f}(x, \omega)}(\kappa, z-x) \subseteq \mathbb{T}'(\kappa)$. Hence we can again obtain (7.15) and then proceed as before. \square

7.3. *Proof of Theorem 4.5.* We start proving (4.6). Let \mathbf{f} as in (4.2) and define

$$b(\kappa) := \min_{x \in [0, \kappa]} \max\{\beta_R(\mathbf{f}(x)), \beta_B(\mathbf{f}(x))\} > 0$$

where the strict positivity of $b(\kappa)$ stems immediately from Remark 2.3 and Proposition 4.4. For $\delta > 0$ arbitrarily fixed, define:

$$(7.19) \quad \mathbb{B}'_{\mathbf{f}}(\kappa, \delta) := \{\mathbf{x} = (x_R, x_B) : \mathbf{x} \in [0, \kappa]^2 \quad \text{and} \quad \|\mathbf{x} - \mathbf{f}(x_R + x_B)\| \leq \delta\},$$

Denoted with $\overset{\circ}{\mathbb{T}}'(\kappa)$ the interior of $\mathbb{T}'(\kappa)$ as defined in (7.4), by construction $\mathbf{f}(x_R + x_B) \in \overset{\circ}{\mathbb{T}}'(\kappa)$ for every $\mathbf{x} = (x_R, x_B) \in [0, \kappa]^2$. Therefore, since β is uniformly continuous on $\mathbb{T}'(\kappa)$ we can always choose δ_0 sufficiently small so that $\mathbb{B}'_{\mathbf{f}}(\kappa, \delta_0) \subset \overset{\circ}{\mathbb{T}}'(\kappa)$ and

$$\max_{\mathbf{x} \in \mathbb{B}'_{\mathbf{f}}(\kappa, \delta_0)} \|\beta(\mathbf{x}) - \beta(\mathbf{f}(x_R + x_B))\| < b(\kappa)/4$$

from which we immediately get:

$$(7.20) \quad \min_{\mathbf{x} \in \mathbb{B}'_{\mathbf{f}}(\kappa, \delta_0)} \max\{\beta_R(\mathbf{x}), \beta_B(\mathbf{x})\} \geq 3b(\kappa)/4.$$

Now, by Proposition 7.1 and Theorem 4.2, setting $\bar{\Upsilon}(\kappa) := \max\{\bar{\Upsilon}_R(\kappa), \bar{\Upsilon}_B(\kappa)\}$, we have

$$\bar{\Upsilon}(\kappa)/(\eta q) \rightarrow 0, \quad \text{and} \quad \sup_{x \in [0, \kappa]} \|\mathbf{F}(x) - \mathbf{f}(x)\| \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

Therefore, for \mathbb{P} -a.e. $\omega \in \Omega$, there exists $n_0(\omega)$ such that for all $n > n_0(\omega)$:

$$(7.21) \quad \mathbf{F}(x, \omega) \in \mathbb{B}'_{\mathbf{f}}(\kappa, \delta_0) \quad \forall x \in [0, \kappa] \quad \text{and} \quad \bar{\Upsilon}(\kappa)/(\eta q) < b(\kappa)/4.$$

Combining (7.21) with (7.20) it follows that for \mathbb{P} -a.e. $\omega \in \Omega$, there exists $n_0(\omega)$ such that for all $n > n_0(\omega)$:

$$(7.22) \quad \min_{x \in [0, \kappa]} \max \left\{ \beta_R \left(\frac{\mathbf{N}[\lfloor xq \rfloor](\omega)}{q} \right), \beta_B \left(\frac{\mathbf{N}[\lfloor xq \rfloor](\omega)}{q} \right) \right\} \\ \geq \min_{x \in [0, \kappa]} \max \left\{ \beta_R \left(\frac{\tilde{\mathbf{N}}(xq)(\omega)}{q} \right), \beta_B \left(\frac{\tilde{\mathbf{N}}(xq)(\omega)}{q} \right) \right\} \geq 3b(\kappa)/4.$$

From the second relation in (7.21), the uniform continuity of $\beta_S(\cdot)$ on $\mathbb{T}'(\kappa)$, for an arbitrarily fixed $x \in [0, \kappa]$ we have that for \mathbb{P} -a.e. $\omega \in \Omega$, there exists $n_1(\omega)$ such that for all $n > n_1(\omega)$

$$(\eta q)^{-1} \left| Q_{\lfloor xq \rfloor + 1}^S(\omega) - \eta q \beta_S(\tilde{\mathbf{N}}(xq)(\omega)/q) \right| < b(\kappa)/4 \quad S \in \{R, B\}$$

and therefore by (7.22) we have that for \mathbb{P} -a.e. $\omega \in \Omega$, and $n > \max\{n_0(\omega), n_1(\omega)\}$ it holds:

$$(\eta q)^{-1} Q_{\lfloor xq \rfloor + 1}^R(\omega) > b(\kappa)/2 \quad \text{or} \quad (\eta q)^{-1} Q_{\lfloor xq \rfloor + 1}^B(\omega) > b(\kappa)/2,$$

which implies:

$$(7.23) \quad \min_{k \in [0, \kappa q]} \frac{\max\{Q_{k+1}^R(\omega), Q_{k+1}^B(\omega)\}}{(\eta q)} > b(\kappa)/2.$$

By the definition of K^* and Lemma 3.6 we have $Q_{K^*}^R = Q_{K^*}^B = 0$. Then (7.23) implies that $K^*(\omega) > \kappa q$, for \mathbb{P} -a.s. $\omega \in \Omega$ and for $n > \max\{n_0(\omega), n_1(\omega)\}$. Then (4.6) follows.

7.3.1. Proof of (4.7).

$$\liminf \frac{A_B^*}{q} = \liminf \frac{N_B[K^* - 1]}{q} + \alpha_B \geq \liminf \frac{\tilde{N}_B(\kappa q)}{q} + \alpha_B \quad \forall \kappa > 0 \quad \mathbb{P}\text{-a.s.},$$

where the final inequality descends from (4.6) and the monotonicity of $N_B(\cdot)$. Therefore, recalling Corollary 4.3 and Proposition 4.4, we have

$$\liminf \frac{N_B[K^* - 1]}{q} \geq \lim_{\kappa \rightarrow \infty} \liminf \frac{\tilde{N}_B(\kappa q)}{q} = \lim_{\kappa \rightarrow \infty} f_B(\kappa) = g_B(\kappa_{\mathbf{g}}). \quad \square$$

7.4. *Proof of Theorem 4.6*. The proof of Theorem 4.6 exploits Lemma 7.3, which is introduced in the next subsection.

7.4.1. *Preliminaries: comparing activation processes*. The next lemma compares the final number of S -active nodes for two activation processes (denoted by label 1 and 2 respectively), which start from different initial conditions (i.e. a different number of R and B seeds). For instance, given $h \in \{1, 2\}$, we denote by $a_{S,h}$ the number of S -seeds of process h , and by $A_{S,h}^*$ its corresponding final size of S -active nodes. Both processes are defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

LEMMA 7.3. If $a_{R,1} \leq a_{R,2}$ and $a_{B,1} \geq a_{B,2}$, then

$$A_{R,1}^* \leq_{st} A_{R,2}^* \quad \text{and} \quad A_{B,2}^* \leq_{st} A_{B,1}^*.$$

The proof exploits a standard coupling argument. We report it in Appendix G.

7.4.2. *Proof of Theorem 4.6*. The proof follows from a comparison between the dynamics of the original system (say system 1) and a companion system (say system 2) in which we set $a_{S,2} = 0$, while we keep $a_{S,2} = a_{S,1}$. As already noticed in Remark 2.2 the final size of S -active nodes in the companion system, say $A_{S,2}^*$, equals the final size of active nodes of the bootstrap percolation process studied in [8], see also [21, 22]. By Lemma 7.3 and Theorem 3.2 in [22], we have that for any $\delta > 0$ there exist $c(\delta) > 0$ and n_δ such that, for any $n \geq n_\delta$,

$$\mathbb{P}\left(\frac{A_{S,1}^*}{q} > z_S + \alpha_S + \delta\right) \leq \mathbb{P}\left(\frac{A_{S,2}^*}{q} > z_S + \alpha_S + \delta\right) = O(\exp(-c(\delta)q)).$$

The claim then follows by a standard application of the Borel-Cantelli lemma. \square

7.5. *Proof of Theorem 4.7*. Part (i). Denote with $\mathbf{k} = (k_R, k_B) \in (\mathbb{N} \cup \{0\})^2$ and $\mathbf{x} = (x_R, x_B) \in [0, \infty)^2$; and set:

$$\mathbb{C}_{\mathbf{f}}(k, \varepsilon) := \{\mathbf{k} : k_R + k_B = k, \|\mathbf{k}/q - \mathbf{f}(k/q)\| \leq \varepsilon\}, \quad \mathbb{C}'_{\mathbf{f}}(k, \varepsilon) := \{\mathbf{x} : \|\mathbf{x} - \mathbf{f}(k/q)\| \leq \varepsilon\}.$$

By Theorem 4.5, Proposition 7.1 and Theorem 4.2 we have that, for any $\omega \in \Omega_\kappa$ and $\varepsilon \in (0, 1)$ there exists $n_0(\omega, \varepsilon)$ such that for any $n > n_0(\omega, \varepsilon)$ we jointly have:

$$(7.24) \quad K^*(\omega) - 1 > \lfloor \kappa q \rfloor, \quad \sup_{0 \leq k \leq \lfloor \kappa q \rfloor} \|\mathbf{N}[k](\omega)/q - \mathbf{f}(k/q)\| < \varepsilon$$

and

$$(7.25) \quad \mathbf{1}_{\{\mathbf{N}[k](\omega) = \mathbf{k}\}} \eta q \beta_S(\mathbf{k}/q) (1 - \varepsilon) < \mathbf{1}_{\{\mathbf{N}[k](\omega) = \mathbf{k}\}} Q_{k+1}^S(\omega) < \mathbf{1}_{\{\mathbf{N}[k](\omega) = \mathbf{k}\}} \eta q \beta_S(\mathbf{k}/q) (1 + \varepsilon) \quad \forall \mathbf{k} : k_R + k_B < \lfloor \kappa q \rfloor.$$

As long as $k < \lfloor \kappa q \rfloor$, by choosing ε sufficiently small, we can always guarantee that $\mathbb{C}'_{\mathbf{f}}(k, \varepsilon) \subset \mathbb{T}'(\kappa)$, with $\mathbb{T}'(\kappa)$ in (7.4). Therefore by (7.25) and the uniform continuity of $\beta_S(\cdot)$ over $\mathbb{C}'_{\mathbf{f}}(k, \varepsilon)$, we have

$$\begin{aligned} & (1 - \varepsilon) \sum_{\mathbf{k} \in \mathbb{C}_{\mathbf{f}}(k, \varepsilon)} \mathbf{1}_{\{\mathbf{N}[k](\omega) = \mathbf{k}\}} \min_{\mathbf{x} \in \mathbb{C}'_{\mathbf{f}}(k, \varepsilon)} \eta q \beta_S(\mathbf{x}) \\ & < \sum_{\mathbf{k} \in \mathbb{C}_{\mathbf{f}}(k, \varepsilon)} \mathbf{1}_{\{\mathbf{N}[k](\omega) = \mathbf{k}\}} Q_{k+1}^S < (1 + \varepsilon) \sum_{\mathbf{k} \in \mathbb{C}_{\mathbf{f}}(k, \varepsilon)} \mathbf{1}_{\{\mathbf{N}[k](\omega) = \mathbf{k}\}} \max_{\mathbf{x} \in \mathbb{C}'_{\mathbf{f}}(k, \varepsilon)} \eta q \beta_S(\mathbf{x}). \end{aligned}$$

Note that since $\|\mathbf{N}[k](\omega)/q - \mathbf{f}(k/q)\| < \varepsilon$ implies $\mathbf{N}[k](\omega) \in \mathbb{C}_f(k, \varepsilon)$, then from the inequality on the right of (7.24), we have:

$$\sum_{\mathbf{k} \in \mathbb{C}_f(k, \varepsilon)} \mathbf{1}_{\{\mathbf{N}[k](\omega)=\mathbf{k}\}} = \mathbf{1}_{\{\mathbf{N}[k](\omega) \in \mathbb{C}_f(k, \varepsilon)\}} = 1 \quad \text{for } \omega \in \Omega_\kappa \text{ and } n > n_0(\omega, \varepsilon).$$

Moreover, recalling (3.13) we have

$$\{K^*(\omega) - 1 > \lfloor \kappa q \rfloor\} \subseteq \{R_{k+1}^S = Q_{k+1}^S, \forall k < \lfloor \kappa q \rfloor, \forall S \in \{R, B\}\}.$$

Summarizing, we proved that, for any $\omega \in \Omega_\kappa$ and $\varepsilon \in (0, 1)$, there exists $n_0(\omega, \varepsilon)$ such that for any $n > n_0(\omega, \varepsilon)$, it holds

$$0 < (1 - \varepsilon) \min_{\mathbf{x} \in \mathbb{C}'_f(k, \varepsilon)} \eta q \beta_S(\mathbf{x}) < R_{k+1}^S < (1 + \varepsilon) \max_{\mathbf{x} \in \mathbb{C}'_f(k, \varepsilon)} \eta q \beta_S(\mathbf{x}) < \infty,$$

for any $k < \lfloor \kappa q \rfloor$. By the regularity of functions $\beta_S(\cdot)$ and $f_S(\cdot)$, it follows that there exists a $c' \in (0, \infty)$ such that, for any $k < \lfloor \kappa q \rfloor$,

$$(7.26) \quad \beta_S(\mathbf{f}(k/q)) - c'\varepsilon < \min_{\mathbf{x} \in \mathbb{C}'_f(k, \varepsilon)} \beta_S(\mathbf{x}) \leq \max_{\mathbf{x} \in \mathbb{C}'_f(k, \varepsilon)} \beta_S(\mathbf{x}) \leq \beta_S(\mathbf{f}(k/q)) + c'\varepsilon.$$

So, for any $\omega \in \Omega_\kappa$ and $\varepsilon \in (0, 1)$, there exists $n_0(\omega, \varepsilon)$ such that for any $n > n_0(\omega, \varepsilon)$,

$$\underline{R}_{k+1}^S(\varepsilon) := (1 - \varepsilon) \eta q (\beta_S(\mathbf{f}(k/q)) - c'\varepsilon) \leq R_{k+1}^S \leq \overline{R}_{k+1}^S(\varepsilon) := (1 + \varepsilon) \eta q (\beta_S(\mathbf{f}(k/q)) + c'\varepsilon),$$

for any $k < \lfloor \kappa q \rfloor$. Note that the upper and lower bounds on R_{k+1}^S are deterministic.

By Proposition 3.5, and its extension mentioned in Section 3.4, we have that the sojourn times $\{W_k\}_{1 \leq k \leq \lfloor \kappa q \rfloor}$ are conditionally independent given $\{(R_k^R, R_k^B) = (q_k^R, q_k^B)\}_{1 \leq k \leq \lfloor \kappa q \rfloor}$ and W_k is distributed according to the exponential law with mean $(q_k^R + q_k^B)^{-1}$. On Ω_κ , for $1 \leq k \leq \lfloor \kappa q \rfloor$, we define the random variables

$$\underline{W}_k^{(\varepsilon)} := \eta \frac{R_k^R + R_k^B}{\underline{R}_k^R(\varepsilon) + \underline{R}_k^B(\varepsilon)} W_k \quad \text{and} \quad \overline{W}_k^{(\varepsilon)} := \eta \frac{R_k^R + R_k^B}{\overline{R}_k^R(\varepsilon) + \overline{R}_k^B(\varepsilon)} W_k.$$

It is immediate to verify that

$$(7.27) \quad \overline{W}_k^{(\varepsilon)} \mid \{(R_k^R, R_k^B) = (q_k^R, q_k^B)\} \stackrel{\text{L}}{=} \text{EXP} \left(\frac{\overline{R}_k^R(\varepsilon) + \overline{R}_k^B(\varepsilon)}{\eta} \right),$$

$$(7.28) \quad \underline{W}_k^{(\varepsilon)} \mid \{(R_k^R, R_k^B) = (q_k^R, q_k^B)\} \stackrel{\text{L}}{=} \text{EXP} \left(\frac{\underline{R}_k^R(\varepsilon) + \underline{R}_k^B(\varepsilon)}{\eta} \right).$$

By (7.24) and (7.25), for every $\varepsilon > 0$ and $\omega \in \Omega_\kappa$, there exists $n_0(\omega, \varepsilon)$ such that for any $n > n_0(\omega, \varepsilon)$ we have

$$(7.29) \quad \underline{W}_k^{(\varepsilon)} < \eta W_k < \overline{W}_k^{(\varepsilon)} \quad 1 \leq k \leq \lfloor \kappa q \rfloor.$$

Since random variables $\{W_k\}_{1 \leq k \leq \lfloor xq \rfloor}$ are conditionally independent given $\{(R_k^R, R_k^B) = (q_k^R, q_k^B)\}_{1 \leq k \leq \lfloor xq \rfloor}$ and $W_k \mid \{(R_k^R, R_k^B) = (q_k^R, q_k^B)\}$ follows the exponential law with mean $(q_k^R + q_k^B)^{-1}$, a standard computation shows that sequences $\{\underline{W}_k^{(\varepsilon)}\}_{1 \leq k \leq \lfloor xq \rfloor}$ and $\{\overline{W}_k^{(\varepsilon)}\}_{1 \leq k \leq \lfloor xq \rfloor}$ are independent. The proof is reported in Appendix K. Furthermore, relations (7.27) and (7.28) imply that

$$\underline{W}_k^{(\varepsilon)} \stackrel{\text{L}}{=} \text{EXP} \left(\frac{\underline{R}_k^R(\varepsilon) + \underline{R}_k^B(\varepsilon)}{\eta} \right) \quad \text{and} \quad \overline{W}_k^{(\varepsilon)} \stackrel{\text{L}}{=} \text{EXP} \left(\frac{\overline{R}_k^R(\varepsilon) + \overline{R}_k^B(\varepsilon)}{\eta} \right),$$

as can be checked by un-conditioning with respect to the random variables (R_k^R, R_k^B) . By (7.29), recalling that $W_k := T_{k+1} - T_k$, for every $\varepsilon > 0$ and $\omega \in \Omega_\kappa$, there exists $n_0(\omega, \varepsilon)$ such that for any $n > n_0(\omega, \varepsilon)$ we have

$$\sum_{k=0}^{\lfloor \kappa q \rfloor - 1} W_k^{(\varepsilon)} < \eta T_{\lfloor \kappa q \rfloor} < \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \bar{W}_k^{(\varepsilon)}.$$

Therefore, for every $\varepsilon > 0$ and $\omega \in \Omega_\kappa$,

$$\liminf \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} W_k^{(\varepsilon)} \leq \liminf \eta T_{\lfloor \kappa q \rfloor} \leq \limsup \eta T_{\lfloor \kappa q \rfloor} \leq \limsup \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \bar{W}_k^{(\varepsilon)}.$$

Denoted with $\mu_*(\kappa) := \int_0^\kappa \frac{1}{\sum_S \beta_S(\mathbf{f}(y))} dy$, due to the arbitrariness of $\varepsilon > 0$, the claim immediately follows if we prove that

$$\liminf \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} W_k^{(\varepsilon)} \geq \mu_*(\kappa) - \gamma(\varepsilon), \quad \limsup \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \bar{W}_k^{(\varepsilon)} \leq \mu_*(\kappa) + \gamma(\varepsilon), \quad \mathbb{P}\text{-a.s.}$$

with $\gamma(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. To prove these relations, since $\sum_{k=1}^{\lfloor \kappa q \rfloor} \bar{W}_k^{(\varepsilon)}$ and $\sum_{k=1}^{\lfloor \kappa q \rfloor} W_k^{(\varepsilon)}$ are sum of independent and exponentially distributed random variables, we apply the exponential tail bounds provided in [16] and reported in Appendix K and the Borel-Cantelli lemma to infer that

$$\frac{\sum_{k=0}^{\lfloor \kappa q \rfloor - 1} W_k^{(\varepsilon)} - \underline{\mu}^{(\varepsilon)}(\kappa)}{\underline{\mu}^{(\varepsilon)}(\kappa)} \rightarrow 0 \quad \text{and} \quad \frac{\sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \bar{W}_k^{(\varepsilon)} - \bar{\mu}^{(\varepsilon)}(\kappa)}{\bar{\mu}^{(\varepsilon)}(\kappa)} \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

where

$$(7.30) \quad \underline{\mu}^{(\varepsilon)}(\kappa) := \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \frac{\eta}{\bar{R}_k^R(\varepsilon) + \underline{R}_k^R(\varepsilon)} \quad \text{and} \quad \bar{\mu}^{(\varepsilon)}(\kappa) := \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \frac{\eta}{\underline{R}_k^R(\varepsilon) + \bar{R}_k^R(\varepsilon)}.$$

Now, define

$$\underline{\beta}_S(\mathbf{x}, \varepsilon) := (\beta_S(\mathbf{x}) - c'\varepsilon)(1 - \varepsilon), \quad \bar{\beta}_S(\mathbf{x}, \varepsilon) := (\beta_S(\mathbf{x}) + c'\varepsilon)(1 + \varepsilon) \quad \text{and} \quad \Delta := 1/q,$$

with c' defined just before (7.26) and $\varepsilon > 0$ chosen sufficiently small that $\underline{\beta}(x, \varepsilon)$ is strictly positive. By the definition of Riemann's integral we have

$$\underline{\mu}^{(\varepsilon)}(\kappa) = \sum_{k=0}^{\lfloor \kappa q \rfloor - 1} \frac{\eta}{\bar{R}_k^R(\varepsilon) + \underline{R}_k^R(\varepsilon)} = \sum_{\substack{k \in \mathbb{N} \cup \{0\} \\ 0 \leq k < \kappa/\Delta}} \frac{\Delta}{\sum_S \bar{\beta}_S(\mathbf{f}(k\Delta), \varepsilon)} \xrightarrow{n \rightarrow \infty} \int_0^\kappa \frac{1}{\sum_S \bar{\beta}_S(\mathbf{f}(x), \varepsilon)} dx$$

and $\bar{\mu}^{(\varepsilon)}(\kappa) \xrightarrow{n \rightarrow \infty} \int_0^\kappa \frac{1}{\sum_S \underline{\beta}_S(\mathbf{f}(x), \varepsilon)} dx$. To conclude the proof of (4.9), we note that $\underline{\beta}(y, \varepsilon)$ and $\bar{\beta}(y, \varepsilon)$ tend to $\beta(y)$, as $\varepsilon \rightarrow 0$, uniformly in $x \in [0, \kappa]$, and so

$$\int_0^\kappa \frac{1}{\sum_S \underline{\beta}_S(\mathbf{f}(x), \varepsilon)} dx \downarrow \mu_*(\kappa) \quad \text{and} \quad \int_0^\kappa \frac{1}{\sum_S \bar{\beta}_S(\mathbf{f}(x), \varepsilon)} dx \uparrow \mu_*(\kappa), \quad \text{as } \varepsilon \downarrow 0. \quad \square$$

The proof of Part (ii) follows similar lines.

8. Proof of theorems 5.1, 5.2 and 5.3. We start introducing a new auxiliary dynamical process.

8.1. *Preliminaries: the R -stopped activation process.* Let $\mathbf{N}(t)$ be an activation process and Z_{stop} be either a fixed time or a point of N , i.e., $Z_{\text{stop}} = T_k$ for some fixed k . We define the R -stopped activation process N_R^{stop} as follows: up to time Z_{stop} the R -stopped process coincides with the original process. On the event $\{T_{K^*-1} > Z_{\text{stop}}\}$, at Z_{stop} the R -activation process stops (no further nodes becomes R -active), while the B -activation process goes on according to the usual rules: upon wake-up every B -susceptible nodes which is still white becomes B -active. The process terminates when no jointly white and B -susceptible nodes are found. More formally, exactly as in the original process, for $t \leq Z_{\text{stop}}$ points of N_S^{stop} are obtained by thinning $\{(T'_k, V'_k)\}_{k \in \mathbb{N}}$, retaining only those couples (T'_k, V'_k) , $k \in \mathbb{N}$, for which, at time $(T'_k)^-$, the white node V'_k satisfies the ‘‘threshold condition with respect to S ’’. For $t > Z_{\text{stop}}$ we retain in N_B^{stop} only those couples (T'_k, V'_k) , $k \in \mathbb{N}$, for which, at time $(T'_k)^-$, the white node V'_k satisfies the ‘‘threshold condition with respect to B ’’. No points are added to N_R^{stop} for $t > Z_{\text{stop}}$, i.e., $N_R^{\text{stop}}(t) = N_R(\min(t, Z_{\text{stop}}))$.

The R -stopped activation process can be prolonged beyond its natural termination along similar lines as for the original process N . From now on we shall refer to the prolonged process. To distinguish variables associated to the stopped and the original processes, we add a superscripts/subscript ‘‘stop’’ to the former. Through a standard coupling argument we have

$$(8.1) \quad A_B^{*,\text{stop}} \geq A_B^* \quad \text{and} \quad T_k^{B,\text{stop}} \leq T_k^B, \quad \forall k \in \mathbb{N} \cup \{0\} \quad \mathbb{P}\text{-a.s.}$$

The proof of (8.1) is reported in Appendix H. More details on the R -stopped process are in Appendix M. Lastly we state a lemma, whose proof follows the same lines of Theorem 4.7.

LEMMA 8.1. Assume $q \ll p^{-1}$ and $Z_{\text{stop}} \leq T_{\lfloor \kappa q \rfloor}^R$ \mathbb{P} -a.s., for some $\kappa \geq 0$, then it holds:

$$(8.2) \quad \eta T_{\lfloor \kappa_B q \rfloor}^{B,\text{stop}} \rightarrow \int_0^{\kappa_B} \frac{1}{\beta_B(y)} dy, \quad \mathbb{P}\text{-a.s.}$$

for any $\kappa_B \in (0, z_B)$, when $q = g$ and $\alpha_B \leq 1$, and any $\kappa_B \in (0, \infty)$ otherwise.

8.2. *Proof of Theorem 5.1.* By Theorem 4.7, for any $\kappa \in (0, \infty)$, we have

$$(8.3) \quad \eta T_{\lfloor \kappa q \rfloor} \rightarrow \tau := \int_0^\kappa \frac{1}{\sum_{S \in \{R, B\}} \beta_S(\mathbf{f}(y))} dy < \infty \quad \mathbb{P}\text{-a.s.}$$

Moreover, by (8.1) and Lemma 8.1, given an arbitrary $\varepsilon > 0$, we have \mathbb{P} -a.s.

$$(8.4) \quad T_{\lfloor (g(\kappa_g) + \varepsilon)q \rfloor}^{B,\text{stop}} \leq T_{\lfloor (g(\kappa_g) + \varepsilon)q \rfloor}^B \quad \text{and} \quad \eta T_{\lfloor g_B(\kappa_g) + \varepsilon \rfloor}^{B,\text{stop}} \rightarrow \psi := \int_0^{(g_B(\kappa_g) + \varepsilon)} \frac{1}{\beta_B(y)} dy$$

We shall show in Appendix N.2 that $\psi > \tau$ either when $q = g$ or $g \ll q \ll p^{-1}$. Denoted with $m_1 := \frac{\psi + 2\tau}{3\tau}$, $m_2 := \frac{2\psi + \tau}{3\tau}$ and $\kappa_B := \lfloor (g_B(\kappa_g) + \varepsilon)q \rfloor$, we introduce the event:

$$(8.5) \quad \mathcal{A}_0 := \{T_{\kappa_B}^B > m_2\tau, T_{\lfloor \kappa q \rfloor} \leq m_1\tau, K^* - 1 > \lfloor \kappa q \rfloor\}, \quad \text{with} \quad \mathbb{P}(\limsup(\mathcal{A}_0^c) = 0),$$

as immediate consequences of (8.3), (8.4) and Theorem 4.5 (i). Now let $[Z_i, Z_{i+1})$, for $0 \leq i < i_1 := \lceil \log_2 \frac{\lfloor f(n)p^{-1} \rfloor}{\lfloor \kappa q \rfloor} \rceil$ a set of intervals, defined by:

$$(8.6) \quad Z_0 := T_{\lfloor \kappa q \rfloor}, \quad Z_{i+1} := \min(T_{2N(Z_i)}, Z_i + \Delta_i) \quad \text{with} \quad \Delta_i = \frac{2^i \kappa q}{\lambda_i} \quad \text{and}$$

$$\lambda_i := \begin{cases} 2^{ri} \kappa^r g \left[\frac{e^{-1}}{2^{r(1-1/r)r^{r-1}} - 2\kappa^{1-r}} \right], & q = g \quad \text{and} \quad 0 \leq i < i_0 := \lfloor \log_2 \frac{p^{-1}}{\lfloor 2\kappa q \rfloor} \rfloor \\ 2^{ir} n(\kappa q p)^r \frac{e^{-1}}{6r!}, & g \ll q \ll p^{-1} \quad \text{and} \quad 0 \leq i < i_0 \\ c_1 n/3, & i_0 \leq i < i_1 \end{cases}$$

where c_1 is a suitable strictly positive constant. From now on, for the case $q = g$ we assume that κ is a chosen sufficiently large to guarantee $\frac{e^{-1}}{2^{r(1-1/r)^{r-1}}} > 2\kappa^{1-r}$. Define $K_i := N(Z_i)$ for $0 \leq i < i_1$, and introduce the events:

$$\mathcal{D}_i := \{K_{i+1} = 2^{i+1} \lfloor \kappa q \rfloor\}, \quad \text{and} \quad \mathcal{E}_i := \{N_B[h] \leq \kappa_B \ \forall h \in [K_i, K_{i+1}]\}.$$

Note that, as immediate consequence of the definition of K_i , we have $T_{K_i} = Z_i$. Moreover $K_{i+1} \leq 2K_i$, from which we obtain $K_i \leq 2^i \lfloor \kappa q \rfloor$ and $\mathcal{D}_{i+1} \subseteq \mathcal{D}_i$. Moreover by (3.11)

$$(8.7) \quad Q_{k+1}^R \geq |S_R[k]| - k.$$

In addition, note that

$$(8.8) \quad \mathcal{E}_i \cap \{k \in [K_i, K_{i+1}]\} \subseteq \mathcal{G}^{(k)} = \{N_B[k] \leq \kappa_B\}.$$

Moreover by Lemma 3.9, for any $k \in [2^i \lfloor \kappa q \rfloor, 2^{i+1} \lfloor \kappa q \rfloor)$, we have:

$$(8.9) \quad |S_R[k]| | \mathcal{G}^{(k)} \geq_{st} \text{Bin}(n_W, \pi_S(k - \kappa_B, \kappa_B)) \geq_{st} \text{Bin}(n_W, \pi_S(2^i \lfloor \kappa q \rfloor - \kappa_B, \kappa_B))$$

At last, note that for any i such that $2^i \lfloor \kappa q \rfloor < p^{-1}$ we have:

$$(8.10) \quad \begin{aligned} \pi_S(2^i \lfloor \kappa q \rfloor - \kappa_B, \kappa_B) &\geq \mathbb{P}(\text{Bin}(2^i \lfloor \kappa q \rfloor - \kappa_B + a_R, p) = r) \mathbb{P}(\text{Bin}(\kappa_B + a_B) = 0) \\ &= \binom{2^i \lfloor \kappa q \rfloor - \kappa_B + a_R}{r} p^r (1-p)^{2^i \lfloor \kappa q \rfloor - \kappa_B + a_R - r} (1-p)^{\kappa_B + a_B} \\ &= \frac{[(2^i \lfloor \kappa q \rfloor - \kappa_B + a_R)p]^r}{r!} e^{-2^i \lfloor \kappa q \rfloor p} (1 + o(1)) \\ &> \frac{[(2^i \lfloor \kappa q \rfloor - \kappa_B + a_R)p]^r}{r!} e^{-1} (1 + o(1)), \end{aligned}$$

$$(8.11) \quad \begin{aligned} \pi_S(2^i \lfloor \kappa q \rfloor - \kappa_B, -\kappa_B) &\geq \mathbb{P}(\text{Bin}(2^i \lfloor \kappa q \rfloor - \kappa_B + a_R, p) \geq r) \mathbb{P}(\text{Bin}(\kappa_B + a_B, p) = 0) \\ &= [\mathbb{P}(\text{Po}((2^i \lfloor \kappa q \rfloor - \kappa_B + a_R)p) \geq r) + O(p)] \mathbb{P}(\text{Bin}(\kappa_B + a_B, p) = 0) \\ &= [\mathbb{P}(\text{Po}((2^i \lfloor \kappa q \rfloor - \kappa_B + a_R)p) \geq r) + O(p)] (1 + o(1)) > c_1, \end{aligned}$$

for a sufficiently small constant c_1 and any i such that $2^i \lfloor \kappa q \rfloor \geq p^{-1}/2$. Then, defining

$$(8.12) \quad \gamma_i := \begin{cases} \frac{e^{-1}}{2} n \frac{(2^i \lfloor \kappa q \rfloor - \kappa_B + a_R)^r}{r!}, & 0 \leq i < i_0 \\ c_1 n / 2, & i_0 \leq i < i_1 \end{cases}$$

and $\mathcal{X}_i := \{|S_R[h]| > \gamma_i \ \forall h \in [K_i, K_{i+1}]\}$, for any $0 \leq i < i_1$, we have

$$\mathcal{X}_i \cap \{k \in [K_i, K_{i+1}]\} \subseteq \mathcal{X}_i^{(k)} := \{|S_R[k]| > \gamma_i\} \quad \text{and}$$

$$(8.13) \quad \mathcal{X}_i^c = \{\exists k \in [K_i, K_{i+1}] : |S_R[k]| \leq \gamma_i\} = \bigcup_k [(\mathcal{X}_i^{(k)})^c \cap \{k \in [K_i, K_{i+1}]\}]$$

By (8.8) and (8.9), setting conventionally $\mathcal{D}_{-1} := \Omega$, it follows

$$(8.14) \quad \begin{aligned} \mathbb{P}\left((\mathcal{X}_i^{(k)})^c \cap \mathcal{E}_i \cap \mathcal{D}_{i-1} \cap \{k \in [K_i, K_{i+1}]\}\right) &\leq \mathbb{P}\left((\mathcal{X}_i^{(k)})^c \cap \mathcal{G}^{(k)}\right) \\ &\leq \mathbb{P}\left((\mathcal{X}_i^{(k)})^c \mid \mathcal{G}^{(k)}\right) = \mathbb{P}\left(|S_R[k]| \leq \gamma_i \mid \mathcal{G}^{(k)}\right) \leq \mathbb{P}\left(\text{Bin}(n_W, \pi_S(2^i \lfloor \kappa q \rfloor - \kappa_B, \kappa_B)) \leq \gamma_i\right). \end{aligned}$$

$$(8.14) \quad \leq h_i := \begin{cases} e^{-ne^{-1} \frac{(2^i \kappa q p)^r}{3r!}} H\left(\frac{1}{2}\right) & 0 \leq i \leq i_0 \\ e^{-\frac{c_1 n}{2}} H\left(\frac{1}{2}\right) & i_0 < i \leq i_1 \end{cases}$$

where the last inequality descends by (8.10) and (8.11), and the application of concentration inequality for binomials (I.2). In addition, by (8.13), (8.14), the definition of \mathcal{D}_{i-1} and the fact that $K_{i+1} \leq 2K_i$, we have

$$\begin{aligned}
\mathbb{P}(\mathcal{K}_i^c \cap \mathcal{E}_i \cap \mathcal{D}_{i-1}) &= \mathbb{P}(\cup_k [(\mathcal{K}_i^{(k)})^c \cap \{k \in [K_i, K_{i+1}]\}] \cap \mathcal{E}_i \cap \mathcal{D}_{i-1}) \\
&= \mathbb{P}(\cup_k [(\mathcal{K}_i^{(k)})^c \cap \{k \in [K_i, 2K_i]\}] \cap \{k \in [K_i, K_{i+1}]\} \cap \mathcal{E}_i \cap \mathcal{D}_{i-1}) \\
&= \mathbb{P}(\cup_k [(\mathcal{K}_i^{(k)})^c \cap \{k \in [2^i \lfloor \kappa q \rfloor, 2^{i+1} \lfloor \kappa q \rfloor]\}] \cap \{k \in [K_i, K_{i+1}]\} \cap \mathcal{E}_i \cap \mathcal{D}_{i-1}) \\
&= \mathbb{P}(\cup_{k=2^i \lfloor \kappa q \rfloor}^{2^{i+1} \lfloor \kappa q \rfloor - 1} [(\mathcal{K}_i^{(k)})^c \cap \{k \in [K_i, K_{i+1}]\}] \cap \mathcal{E}_i \cap \mathcal{D}_{i-1}) \\
&\leq \sum_{k=2^i \lfloor \kappa q \rfloor}^{2^{i+1} \lfloor \kappa q \rfloor - 1} \mathbb{P}((\mathcal{K}_i^{(k)})^c \cap \mathcal{G}^{(k)}) \leq (2^i \kappa q) h_i,
\end{aligned}$$

where in the final inequality we have applied sub-additivity of probability along with (8.14). Now, for any $0 \leq i < i_0$, from (3.11), by assuming κ sufficiently large, we have

$$\begin{aligned}
(8.15) \quad Q_{k+1}^R \mid \mathcal{K}_i \cap \mathcal{D}_{i-1} \cap \{k \in [K_i, K_{i+1}]\} &\geq \frac{e^{-1}}{2} n \frac{[(2^i \kappa q - \kappa_B + a_R)p]^r}{r!} - 2^{i+1} \kappa q \\
&\geq \frac{e^{-1}}{2} n \frac{[(2^i - 1/2)(\kappa q p)]^r}{r!} - 2^{i+1} \kappa q, \\
\frac{e^{-1}}{2} n \frac{[(2^i - 1/2)(\kappa q p)]^r}{r!} - 2^{i+1} \kappa q &\geq (2^i - 1/2)^r \kappa^r g \frac{e^{-1}}{2^r (1 - 1/r)^{r-1}} - 2^{i+1} \kappa q \geq \lambda_i.
\end{aligned}$$

Now in the case $q = g$,

$$\begin{aligned}
(8.16) \quad \frac{e^{-1}}{2} n \frac{[(2^i - 1/2)(\kappa g p)]^r}{r!} - 2^{i+1} \kappa g &\geq (2^i - 1/2)^r \kappa^r g \frac{e^{-1}}{2^r (1 - 1/r)^{r-1}} - 2^{i+1} \kappa g \\
&\geq \lambda_i := 2^{ri} \kappa^r g \left[\frac{e^{-1}}{2^r r (1 - 1/r)^{r-1}} - 2\kappa^{-(r-1)} \right],
\end{aligned}$$

where we have used the identity $r(1 - \frac{1}{r})^{r-1} \frac{n(gp)^r}{r!} = g$. For the case

$$\begin{aligned}
(8.17) \quad \frac{e^{-1}}{2} n \frac{[(2^i \kappa q - \kappa_B + a_R)p]^r}{r!} - 2^{i+1} \kappa q &\geq \frac{e^{-1}}{3} n \frac{[(2^i - 1/2)(\kappa q p)]^r}{r!} \\
&\geq \lambda_i := 2^{ir} n (\kappa q p)^r \frac{e^{-1}}{6r!}
\end{aligned}$$

where we have exploited that $q \ll n(gp)^r$. Note also that

$$(8.18) \quad Q_{k+1}^R \mid \mathcal{K}_i \cap \mathcal{D}_{i-1} \cap \{k \in [K_i, K_{i+1}]\} \geq c_1 n / 2 - (2^{i+1} \kappa + z_B) g \geq \lambda_i := c_1 n / 3$$

for all n sufficiently large, $i_0 < i \leq i_1$. Define $\mathcal{Z}_i := \{T_{K_{i+1}} - T_{K_i} < \Delta_i\}$ and note that $\mathcal{Z}_i \subseteq \{K_{i+1} = 2K_i\}$, from which we immediately get that:

$$(8.19) \quad \cap_{j < i} \mathcal{Z}_j \subseteq \{K_i = 2^i \lfloor \kappa q \rfloor\} = \mathcal{D}_{i-1} \text{ and } \cap_{i=0}^{i_1-1} \mathcal{Z}_i \subseteq \{K_{i_1} = 2^{i_1} \lfloor \kappa q \rfloor \geq \lfloor f(n)p^{-1} \rfloor\}$$

In addition since $K_i \mid \mathcal{D}_{i-1} = 2^i \lfloor \kappa q \rfloor$ and $K_{i+1} \leq 2K_i$:

$$\begin{aligned}
\mathbb{P}(\mathcal{Z}_i^c \mid \mathcal{K}_i \cap \mathcal{D}_{i-1}) &\leq \mathbb{P}(T_{2^{i+1} \lfloor \kappa q \rfloor} - T_{2^i \lfloor \kappa q \rfloor} > \Delta_i \mid \mathcal{K}_i \cap \mathcal{D}_{i-1}), \quad \text{with} \\
T_{2^{i+1} \lfloor \kappa q \rfloor} - T_{2^i \lfloor \kappa q \rfloor} &= \sum_{h=0}^{2^i \lfloor \kappa q \rfloor - 1} [T_{2^i \lfloor \kappa q \rfloor + h + 1} - T_{2^i \lfloor \kappa q \rfloor + h}] = \sum_{h=0}^{2^i \lfloor \kappa q \rfloor - 1} W_{2^i \lfloor \kappa q \rfloor + h + 1}
\end{aligned}$$

where $\{W_{K_i+h+1}\}_h$ are conditionally independent given $\{R_{2^i \lfloor \kappa q \rfloor + h+1}\}_h$ with $W_{2^i \lfloor \kappa q \rfloor + h+1} | \{R_{2^i \lfloor \kappa q \rfloor + h+1} = m\} \stackrel{L}{=} \text{EXP}(m)$. Then, proceeding similarly as in the proof of Theorem 4.7, for any $0 \leq i < i_1$ we can define a sequence of r.v.'s $\widehat{W}_h^{(i)} := \frac{R_{2^i \lfloor \kappa q \rfloor + h+1}}{\lambda_i} W_{2^i \lfloor \kappa q \rfloor + h+1}$, which turn out to be conditionally independent, with law $\exp(\lambda_i)$, given the sequence of visited states $\{\mathbf{X}_{2^i \lfloor \kappa q \rfloor + h+1}\}_h$, as well as independent of $\mathcal{H}_{2^i \lfloor \kappa q \rfloor}$. Moreover $\widehat{W}_h^{(i)} > W_{2^i \lfloor \kappa q \rfloor + h+1}^{(i)}$ on $\{R_{2^i \lfloor \kappa q \rfloor + h+1} > \lambda_i\}$. Since for an arbitrary $k \leq 2^{i_1} \kappa q$ we have $n - N[k] \geq n - 2^{i_1} \kappa q > \lambda_i$, it follows from (3.12), (3.13) and (3.10) that $\{R_{k+1} \leq \lambda_i\} \subseteq \{R_{k+1} = Q_{k+1}^R + Q_{k+1}^B, Q_{k+1}^R \geq 0, Q_{k+1}^B \geq 0\}$, hence $\{R_{k+1} \leq \lambda_i\} \subseteq \{Q_{k+1}^R \leq \lambda_i\}$. Now, recalling (8.15), we have $\mathcal{K}_i \cap \mathcal{D}_{i-1} \subseteq \{Q_{k+1}^R > \lambda_i \forall k \in [K_i, K_{i+1}), K_i = 2^i \lfloor \kappa q \rfloor\} \subseteq \{R_{k+1} > \lambda_i \forall k \in [K_i, K_{i+1}), K_i = 2^i \lfloor \kappa q \rfloor\} \subseteq \{W_{2^i \lfloor \kappa q \rfloor + h+1}^{(i)} \leq \widehat{W}_h^{(i)}, \forall h \in [0, K_{i+1} - 2^i \lfloor \kappa q \rfloor]\}$, and:

$$\begin{aligned}
(8.20) \quad \mathbb{P}(T_{K_{i+1}} - T_{K_i} > \Delta_i | \mathcal{K}_i \cap \mathcal{D}_{i-1}) &= \mathbb{P}(T_{K_{i+1}} - T_{2^i \lfloor \kappa q \rfloor} > \Delta_i | \mathcal{K}_i \cap \mathcal{D}_{i-1}) \\
&\leq \mathbb{P}\left(\sum_{h=0}^{K_{i+1} - 2^i \lfloor \kappa q \rfloor - 1} \widehat{W}_h^{(i)} > \Delta_i | \mathcal{K}_i \cap \mathcal{D}_{i-1}\right) \\
&\leq \mathbb{P}\left(\sum_{h=0}^{2^i \lfloor \kappa q \rfloor - 1} \widehat{W}_h^{(i)} > \Delta_i | \mathcal{K}_i \cap \mathcal{D}_{i-1}\right) = \mathbb{P}\left(\sum_{h=0}^{2^i \lfloor \kappa q \rfloor - 1} \widehat{W}_h^{(i)} > \Delta_i\right) \\
&= \mathbb{P}(\text{Po}(\lambda_i \Delta_i) < 2^i \lfloor \kappa q \rfloor) < \exp(-\lambda_i \Delta_i H(1/2)),
\end{aligned}$$

where the latter inequality follows from (I.3). Now, since

$$\mathcal{A}_0 \cap (\cap_{i=0}^{i_1-1} (\mathcal{K}_i \cap \mathcal{Z}_i)) \subseteq \mathcal{A}_0 \cap (\cap_{i=0}^{i_1-1} \mathcal{Z}_i) \subseteq \left\{T_{\lfloor f(n)p^{-1} \rfloor}^R \leq m_1 \tau + \sum_{i=0}^{i_1-1} \Delta_i\right\}$$

recalling that $T_{K_i} \leq T_{K_0} + \sum_{j=0}^{i-1} \Delta_j$, and observing that, for n and κ sufficiently large, we can always assume $\sum_i \Delta_i < (m_2 - m_1)\tau$, it results that

$$(8.21) \quad \mathcal{A}_0 \subseteq \cap_{i=0}^{i_1-1} \mathcal{E}_i = \{T_{\kappa_B}^B > T_{K_{i_1}}\},$$

by (8.19) we have ⁵ $\mathcal{A}_0 \cap (\cap_{j < i} (\mathcal{K}_j \cap \mathcal{Z}_j)) \subseteq (\cap_{i=0}^{i_1-1} \mathcal{E}_i) \cap \mathcal{D}_{i-1} \subseteq \mathcal{E}_i \cap \mathcal{D}_{i-1}$, and:

$$\begin{aligned}
(8.22) \quad \mathbb{P}(\mathcal{A}_0^c \cup (\cup_{i=0}^{i_1-1} (\mathcal{K}_i^c \cup \mathcal{Z}_i^c))) &= \mathbb{P}(\mathcal{A}_0^c) + \mathbb{P}(\mathcal{A}_0 \cap [\cup_{i=0}^{i_1-1} (\mathcal{K}_i^c \cap [\cap_{j < i} (\mathcal{K}_j \cap \mathcal{Z}_j)])]) \\
&\quad \mathbb{P}(\mathcal{A}_0 \cap [\cup_{i=0}^{i_1-1} ((\mathcal{Z}_i^c \cap \mathcal{K}_i) \cap [\cap_{j < i} (\mathcal{K}_j \cap \mathcal{Z}_j)])]) \\
&\leq \mathbb{P}(\mathcal{A}_0^c) + \mathbb{P}(\cup_i (\mathcal{K}_i^c \cap \mathcal{D}_{i-1} \cap \mathcal{E}_i)) + \mathbb{P}(\cup_i (\mathcal{Z}_i^c \cap \mathcal{K}_i \cap \mathcal{D}_{i-1})) \\
&\leq \mathbb{P}(\mathcal{A}_0^c) + \sum_i \mathbb{P}(\mathcal{K}_i^c \cap \mathcal{D}_i \cap \mathcal{E}_i) + \sum_i \mathbb{P}(\mathcal{Z}_i^c | \mathcal{K}_i \cap \mathcal{D}_{i-1}).
\end{aligned}$$

At last observe that by (8.19) right inclusion, (8.21) and the inclusion:

$$\mathcal{A}_0 \cap (\cap_{i=0}^{i_1-1} \mathcal{K}_i) \subseteq \{K^* > K_0, Q_{k+1}^R > 0, \forall k \in [K_0, K_{i_1})\} \subseteq \{K^* > K_{i_1}\}.$$

given that by construction $Q_{K^*}^R = 0$, therefore

$$\mathcal{A}_0 \cap [\cap_{i=0}^{i_1-1} (\mathcal{K}_i \cap \mathcal{Z}_i)] \subseteq \mathcal{B} := \{N_B(\lfloor f(n)p^{-1} \rfloor) < \kappa_B\} \cap \{K^* - 1 \geq \lfloor f(n)p^{-1} \rfloor\}$$

⁵We conventionally set $\cap_{j < 0} \mathcal{Z}_j = \cap_{j < 0} \mathcal{K}_j = \cap_{j < 0} (\mathcal{K}_j \cap \mathcal{Z}_j) = \Omega$.

Note that, by the application of the Borel-Cantelli lemma, we can claim that $\limsup \mathbb{P}(\mathcal{B}^c) = 0$ provided that we show $\sum_n \mathbb{P}(\mathcal{B}^c) \leq \sum_n \mathbb{P}(\mathcal{A}_0^c \cup (\cup_{i=0}^{i_1-1} (\mathcal{K}_i^c \cup \mathcal{Z}_i^c))) < \infty$. Now, we can first apply (8.22) to the r.h.s and then (??) and (8.20) to show that the resulting latter two infinite sums are converging. Lastly, $\sum_n \mathbb{P}(\mathcal{A}_0^c)$ converges due to the (second) Borel-Cantelli lemma and (8.5) (we recall, indeed, that $\mathcal{A}_0^{(n)}$ are independent). \square

8.3. *Proof of Theorem 5.2*. The proof of Theorem 5.2 exploits the following lemma, whose proof is given in Appendix N.1.

LEMMA 8.2. Let $\{X_n\}_{n \geq 1}$ and $\{Y_n\}_{n \geq 1}$ be two sequences of non-negative random variables and $\{f_n\}_{n \geq 1}$ a sequence of non-negative numbers. If $X_n \leq_{st} Y_n$ for every n , $Y_n/f_n \rightarrow 0$ a.s., as $n \rightarrow \infty$, and $\{Y_n\}_{n \geq 1}$ are independent, then $X_n/f_n \rightarrow 0$ a.s., as $n \rightarrow \infty$.

Part I: proof of $\mathbb{P}(\liminf \{N_B[K^* - 1] \leq \lfloor (g_B(\kappa_g) + \varepsilon)q \rfloor\}) = 1$. Let $f(n)$ be as in Theorem 5.1(i). and consider the R -stopped process with $Z_{\text{stop}} = T_{\lfloor f(n)p^{-1} \rfloor}$. Set $\kappa_0 := \lfloor f(n)p^{-1} \rfloor$, $\kappa_1 := \kappa_0 + \lfloor \varepsilon g \rfloor$, with $\varepsilon > 0$ arbitrarily fixed, $\kappa_B^{(0)} := \lfloor (g_B(\kappa_g) + \varepsilon)g \rfloor$ and $\kappa_B^{(1)} := \lfloor (g_B(\kappa_g) + 2\varepsilon)g \rfloor$, and define the events:

$$\mathcal{B}_0 := \{T_{\kappa_0} \leq T_{\kappa_B^{(0)}}^B\} = \{N_B[\kappa_0] \leq \kappa_B^{(0)}\} \quad \text{and} \quad \mathcal{C}_0 := \{K^* - 1 \geq \kappa_0\}.$$

By (3.11), we have $Q_{\kappa_0+1}^{B,\text{stop}} = Q_{\kappa_0+1}^B \leq |\mathcal{S}_B[\kappa_0]|$. Now invoking Lemma 3.9, we have

$$\begin{aligned} |\mathcal{S}_B^{\text{stop}}[\kappa_0]| | \mathcal{B}_0 &= |\mathcal{S}_B[\kappa_0]| | \mathcal{B}_0 \leq_{st} \text{Bin}(n_W, \pi_B(\kappa_0 - \kappa_B^{(0)}, \kappa_B^{(0)})) \quad \text{with} \\ (8.23) \quad \text{Bin}(n_W, \pi_B(\kappa_0 - \kappa_B^{(0)}, \kappa_B^{(0)}))/g &\rightarrow 0, \quad \mathbb{P}\text{-a.s.;} \end{aligned}$$

indeed it is rather immediate to check that $n_W \pi_B(\kappa_0 - \kappa_B^{(0)}, \kappa_B^{(0)})/g \rightarrow 0$. Then applying concentration inequality (I.1), we have $\mathbb{P}(\text{Bin}(n_W, \pi_B(\kappa_0 - \kappa_B^{(0)}, \kappa_B^{(0)})) > \varepsilon g) < \exp(-\frac{\varepsilon g}{2})$, for n sufficiently large and $\varepsilon > 0$. The claim (8.23) follows by a standard application of Borel-Cantelli lemma. Similarly, one has $|\mathcal{S}_B^{\text{stop}}[\kappa_1]| | \mathcal{B}_0 \leq_{st} \text{Bin}(n_W, \pi_B(\kappa_1 - \kappa_B^{(1)}, \kappa_B^{(1)}))$ with $\text{Bin}(n_W, \pi_B(\kappa_1 - \kappa_B^{(1)}, \kappa_B^{(1)}))/g \rightarrow 0$ \mathbb{P} -a.s.. Therefore, by Lemma 8.2 we have (recalling that the above r.v.s for different n are independent):

$$(8.24) \quad |\mathcal{S}_B^{\text{stop}}[\kappa_0]| | \mathcal{B}_0 = o_{a.s.}(g) \quad \text{and} \quad |\mathcal{S}_B^{\text{stop}}[\kappa_1]| | \mathcal{B}_0 = o_{a.s.}(g).$$

Now, consider the quantity $Q_{\kappa_1+1}^{B,\text{stop}}$. First note that

$$(8.25) \quad \mathcal{S}_B[\kappa_0] = \mathcal{S}_B^{\text{stop}}[\kappa_0] \subseteq \mathcal{S}_B^{\text{stop}}[\kappa_0 + k] \subseteq \mathcal{S}_B^{\text{stop}}[\kappa_1], \quad \forall k \leq \lfloor \varepsilon g \rfloor$$

since, focusing on the R -stopped process, no node becomes R -active after $T_{\kappa_0}^{\text{stop}}$, and therefore the number of B -susceptible nodes after $T_{\kappa_0}^{\text{stop}}$ is monotonically increasing. Of course it holds:

$$(8.26) \quad \mathcal{V}_B^{\text{stop}}[\kappa_0] = \mathcal{V}_B[\kappa_0] \quad \text{and} \quad \mathcal{V}_R^{\text{stop}}[\kappa_0 + k] = \mathcal{V}_R^{\text{stop}}[\kappa_0] = \mathcal{V}_R[\kappa_0], \quad \forall k \leq \lfloor \varepsilon g \rfloor.$$

Recalling that *i*) up to time T_{K^*-1} only S -susceptible nodes becomes S -active; *ii*) a node can be S -susceptible only if it has collected at least r S -marks, i.e., $\{v \in \mathcal{S}_S(t)\} \subseteq \{D_S^v(t) \geq r\}$; *iii*) since $D_S^v[k]$, i.e., the number of S -marks collected by a node v , is non-decreasing in k , we have

$$\begin{aligned} & \mathbf{1}_{\mathcal{C}_0} |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_1]) \cap \mathcal{V}_B[\kappa_0] \cap \{v : D_B^{v,\text{stop}}[\kappa_1] \geq r\}| \\ (8.27) \quad & \stackrel{(a)}{=} \mathbf{1}_{\mathcal{C}_0} |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_1]) \cap \mathcal{V}_B[\kappa_0]| \leq \stackrel{(b)}{=} \mathbf{1}_{\mathcal{C}_0} |(\mathcal{V}_W \setminus \mathcal{S}_B[\kappa_0]) \cap \mathcal{V}_B[\kappa_0]|. \end{aligned}$$

Here equation (a) follows from the fact that, conditionally on \mathcal{C}_0 , due to the previously mentioned properties *i*), *ii*) and *iii*) we have: $\mathcal{V}_B[\kappa_0] \subseteq \{v : D_B^{v,\text{stop}}[\kappa_0] \geq r\} \subseteq \{v : D_B^{v,\text{stop}}[\kappa_1] \geq r\}$ and inequality (b) follows from (8.25). Then, noticing that: *i*) $N_B^{\text{stop}}[\kappa_1] = N_B[\kappa_0] + \lfloor \varepsilon g \rfloor$; *ii*) conditionally on \mathcal{C}_0 we have $\mathcal{V}_R^{\text{stop}}[\kappa_0] \subseteq \{v : D_R^{v,\text{stop}}[\kappa_0] \geq r\} = \{v : D_R^{v,\text{stop}}[\kappa_1] \geq r\}$; and recalling (3.9), (8.25), (8.26) and (8.27) we have

$$\begin{aligned}
Q_{\kappa_1+1}^{B,\text{stop}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} &= \left[|\mathcal{S}_B^{\text{stop}}[\kappa_1]| - N_B^{\text{stop}}[\kappa_1] - |\mathcal{S}_B^{\text{stop}}[\kappa_1] \cap \mathcal{V}_R^{\text{stop}}[\kappa_1] \cap \{v : D_R^{v,\text{stop}}[\kappa_1] \geq r\}| \right. \\
&\quad \left. + |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_1] \cap \mathcal{V}_B^{\text{stop}}[\kappa_1] \cap \{v : D_B^{v,\text{stop}}[\kappa_1] \geq r\})| \right] \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \\
&= \left[|\mathcal{S}_B^{\text{stop}}[\kappa_1]| - N_B^{\text{stop}}[\kappa_1] + |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_0]) \cap \mathcal{V}_B[\kappa_0]| \right. \\
&\quad \left. + |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_1]) \cap (\mathcal{V}_B^{\text{stop}}[\kappa_1] \setminus \mathcal{V}_B[\kappa_0]) \cap \{v : D_B^{v,\text{stop}}[\kappa_1] > r\}| \right. \\
&\quad \left. - |\mathcal{S}_B^{\text{stop}}[\kappa_1] \cap \mathcal{V}_R[\kappa_0]| \right] \\
&\leq \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \left[|\mathcal{S}_B^{\text{stop}}[\kappa_1]| - |\mathcal{S}_B[\kappa_0]| + |\mathcal{S}_B[\kappa_0]| - N_B[\kappa_0] - \lfloor \varepsilon g \rfloor \right. \\
&\quad \left. - |\mathcal{S}_B[\kappa_0] \cap \mathcal{V}_R[\kappa_0]| + |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_0]) \cap \mathcal{V}_B[\kappa_0]| \right. \\
&\quad \left. + |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_1]) \cap (\mathcal{V}_B^{\text{stop}}[\kappa_1] \setminus \mathcal{V}_B[\kappa_0]) \cap \{v : D_B^{v,\text{stop}}[\kappa_1] > r\}| \right] \\
&\leq \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \left[Q_{\kappa_0+1}^{B,\text{stop}} + |\mathcal{S}_B^{\text{stop}}[\kappa_1]| - \lfloor \varepsilon g \rfloor \right. \\
(8.28) \quad &\quad \left. + |(\mathcal{V}_W \setminus \mathcal{S}_B^{\text{stop}}[\kappa_1]) \cap (\mathcal{V}_B^{\text{stop}}[\kappa_1] \setminus \mathcal{V}_B[\kappa_0]) \cap \{v : D_B^{v,\text{stop}}[\kappa_1] > r\}| \right],
\end{aligned}$$

where the last addend in (8.28) is not larger than $\lfloor \varepsilon g \rfloor$, since

$$|(\mathcal{V}_B^{\text{stop}}[\kappa_1] \setminus \mathcal{V}_B[\kappa_0])| = N_B^{\text{stop}}[\kappa_1] - N_B[\kappa_0] = \lfloor \varepsilon g \rfloor.$$

Moreover, the last addend in (8.28) is different from 0 only on the event $\{K^{*,\text{stop}} \leq \kappa_1\}$. Indeed, for any k be such $\kappa_0 < k \leq \kappa_1$, upon $\{K^{*,\text{stop}} > \kappa_1\}$ we have $V_k^{\text{stop}} \in \mathcal{S}_B^{\text{stop}}(k)$ with $\mathcal{S}_B^{\text{stop}}(k) \subseteq \mathcal{S}_B^{\text{stop}}[\kappa_1]$. In other words, $\{K^{*,\text{stop}} > \kappa_1\} \subseteq \{(\mathcal{V}_B^{\text{stop}}[\kappa_1] \setminus \mathcal{V}_B^{\text{stop}}[\kappa_0]) \in \mathcal{S}_B^{\text{stop}}[\kappa_1]\}$. Consequently, we have

$$Q_{\kappa_1+1}^{B,\text{stop}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \leq \left[Q_{\kappa_0+1}^{B,\text{stop}} + |\mathcal{S}_B^{\text{stop}}[\kappa_1]| - \lfloor \varepsilon g \rfloor \mathbf{1}_{\{K^{*,\text{stop}} > \kappa_1\}} \right] \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0}, \quad \mathbb{P}\text{-a.s.}$$

Combining inequality (3.11) and (8.24), we have

$$(8.29) \quad \mathbf{1}_{\{K^{*,\text{stop}} > \kappa_1\}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \leq \frac{-Q_{\kappa_1+1}^{B,\text{stop}} + o_{a.s.}(g)}{\lfloor \varepsilon g \rfloor} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0}.$$

By construction $\{Q_{\kappa_1+1}^{B,\text{stop}} < 0\} \subseteq \{K^{*,\text{stop}} \leq \kappa_1\}$, and so

$$(8.30) \quad \mathbf{1}_{\{K^{*,\text{stop}} > \kappa_1\}} \leq \mathbf{1}_{\{Q_{\kappa_1+1}^{B,\text{stop}} \geq 0\}}.$$

By (8.29) and (8.30), after multiplying both sides by $\mathbf{1}_{\{Q_{\kappa_1+1}^{B,\text{stop}} \geq 0\}}$ we immediately have

$$\mathbf{1}_{\{K^{*,\text{stop}} > \kappa_1\}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \leq \frac{-Q_{\kappa_1+1}^{B,\text{stop}} + o_{a.s.}(g)}{\lfloor \varepsilon g \rfloor} \mathbf{1}_{\{Q_{\kappa_1+1}^{B,\text{stop}} \geq 0\}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0}.$$

Now, noticing that

$$\limsup \frac{-Q_{\kappa_1+1}^{B,\text{stop}} + o_{a.s.}(g)}{\lfloor \varepsilon g \rfloor} \mathbf{1}_{\{Q_{\kappa_1+1}^{B,\text{stop}} \geq 0\}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \leq \limsup \frac{o_{a.s.}(g)}{g} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} = 0 \quad \mathbb{P}\text{-a.s.},$$

we deduce $\mathbf{1}_{\{K^{*,\text{stop}} > \kappa_1\}} \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} \rightarrow 0$ \mathbb{P} -a.s., which implies $\mathbf{1}_{\{K^{*,\text{stop}} > \kappa_1\}} \rightarrow 0$ \mathbb{P} -a.s., since $\lim \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} = \liminf \mathbf{1}_{\mathcal{B}_0 \cap \mathcal{C}_0} = 1$ \mathbb{P} -a.s., by Theorem 5.1. Now $\mathcal{B}_0 \cap \{K^{*,\text{stop}} \leq \kappa_1\} \subseteq \{A_B^{*\text{stop}} \leq \kappa_B^{(0)} + \lfloor \varepsilon g \rfloor + a_B\}$ and the claim follows by the arbitrariness of ε and (8.1).

Part II: proof of $\mathbb{P}(\liminf \{K^ - 1 \geq \lfloor cn \rfloor\}) = 1$.* Set $k_i = \min\{2^i \kappa_0, \lfloor cn \rfloor\}$ for $0 \leq i \leq i_1 := \lceil \log_2 \frac{\lfloor cn \rfloor}{\lfloor f(n)p^{-1} \rfloor} \rceil$, and define $\mathcal{A}_0 := \{K^* - 1 \geq \kappa_0\} \cap \{N_B[K^* - 1] \leq \kappa_B^{(0)}\}$, with $\mathbb{P}(\limsup(\mathcal{A}_0^c) = 0)$ by Theorem 5.1 and Part I. Then for any $0 \leq i < i_1$, we have:

$$\begin{aligned} \{K^* - 1 \in [k_i, k_{i+1}]\} \cap \mathcal{A}_0 &\subseteq \{\exists k \in [k_i, k_{i+1}] \text{ s.t. } Q_{k+1}^R = 0, N_B[k] \leq \kappa_B^{(0)}\} \quad \text{with} \\ \mathbb{P}(\exists k \in [k_i, k_{i+1}] \text{ s.t. } Q_{k+1}^R = 0, N_B[k] \leq \kappa_B^{(0)}) &\leq \sum_{k=k_i}^{k_{i+1}-1} \mathbb{P}(Q_{k+1}^R = 0, N_B[k] \leq \kappa_B^{(0)}) \end{aligned}$$

and so by (3.11), Lemma 3.9 and concentration inequality (I.2), it follows

$$\begin{aligned} \mathbb{P}(K^* - 1 \in [k_i, k_{i+1}], \mathcal{A}_0) &\leq \sum_{k=k_i}^{k_{i+1}-1} \mathbb{P}\left(S_R[k] \leq k, N_R[k] \geq k - \kappa_B^{(0)}, N_B[k] \leq \kappa_B^{(0)}\right) \\ &\leq \sum_{k=k_i}^{k_{i+1}-1} \mathbb{P}\left(\text{Bin}(n_w, \pi_R(k - \kappa_B^{(0)}, \kappa_B^{(0)}) < k)\right) \\ &\leq 2^i \kappa_0 \mathbb{P}\left(\text{Bin}(n_w, \pi_R(k_i - \kappa_B^{(0)}, \kappa_B^{(0)})) < k_{i+1}\right) < \exp\left(-cnH\left(\frac{c}{\frac{1}{2} + \frac{c}{2}}\right)\right), \end{aligned}$$

for any $0 \leq i < i_1$ and n large enough. The claim, as in Theorem 5.1, follows by a joint application of the two Borel-Cantelli lemmas ($\mathcal{A}_0^{(n)}$ are independent), by observing that:

$$\mathbb{P}(K^* - 1 \geq \lfloor cn \rfloor) \leq \mathbb{P}(\mathcal{A}_0^c) + \sum_{i=0}^{i_1-1} \mathbb{P}(K^* - 1 \in [k_i, k_{i+1}], \mathcal{A}_0). \quad \square$$

8.4. *Proof of Theorem 5.3*. Let $f(n)$ be the function considered in the statement of Theorem 5.1(ii), for the case $g \ll q \ll p^{-1}$. Define

$$(8.31) \quad \kappa_0 := \begin{cases} \lfloor f(n)p^{-1} \rfloor & \text{if } g \ll q \ll p^{-1} \\ \lfloor \kappa p^{-1} \rfloor & \text{if } q = p^{-1} \\ \lfloor \kappa q \rfloor & \text{if } p^{-1} \ll q \ll n \end{cases} \quad \kappa_B^{(0)} := \begin{cases} \lfloor p^{-1} \rfloor & \text{if } g \ll q \ll p^{-1} \\ \lfloor \bar{f}_B p^{-1} \rfloor & \text{if } q = p^{-1} \\ \lfloor q \rfloor & \text{if } p^{-1} \ll q \ll n \end{cases}$$

where κ is arbitrary and \bar{f}_B is defined in Proposition 4.4(iii). Due to the arbitrariness of κ note that ratio $\kappa_0/\kappa_B^{(0)}$ can be assumed arbitrarily large for values of n large enough. Define:

$$(8.32) \quad \mathcal{C}_0 := \{T_{\kappa_0} < T_{\kappa_B^{(0)}}^B\} = \{N_B(\kappa_0) < \kappa_B^{(0)}\}, \quad \mathcal{D}_0 := \{T_{K^*-1} \geq T_{\kappa_0}\} = \{K^* - 1 \geq \kappa_0\},$$

$$(8.33) \quad Z_0 := \min(T_{\kappa_0}, T_{\kappa_B^{(0)}}^B), \quad Z_{i+1} := \min\{T_{4^{i+1}\kappa_0}, T_{2^{i+1}\kappa_B^{(0)}}^B, T_{\lfloor cn \rfloor}\}, \quad i \geq 0.$$

$$(8.34) \quad I := \min\{i : N(Z_{i+1}) = \lfloor cn \rfloor\} = \min\{i : Z_{i+1} = T_{\lfloor cn \rfloor}\}$$

We can bound I as follows:

$$(8.35) \quad \{\underline{i} \leq I \leq \bar{i}\} = \Omega, \quad \text{with } \underline{i} := \left\lfloor \log_4 \frac{\lfloor cn \rfloor}{8\kappa_0} \right\rfloor, \quad \bar{i} := \left\{ \left\lceil \log_4 \frac{\lfloor cn \rfloor}{\kappa_0} \right\rceil + \left\lceil \log_2 \frac{\lfloor cn \rfloor}{\kappa_B^{(0)}} \right\rceil \right\}$$

for all n sufficiently large to guarantee that all previous expressions are meaningful. Letting $K_i = N(Z_i)$, note that by construction:

$$(8.36) \quad K_i := N(Z_i) \leq \min(4^i \kappa_0, \lfloor cn \rfloor) \quad \text{and} \quad N_B(Z_i) \leq 2^i \kappa_B^{(0)}.$$

Finally, consider the events

$$(8.37) \quad \mathcal{A}_i := \{Z_{i+1} = \min\{T_{4^{i+1}\kappa_0}, T_{\lfloor cn \rfloor}\}\} = \{K_{i+1} = \min\{4^{i+1}\kappa_0, \lfloor cn \rfloor\}\}, \quad i \geq 0,$$

$$(8.38) \quad \mathcal{B}_i := \{Q_{h+1}^R > \lambda_i \text{ and } Q_{h+1}^B \leq \phi_i \forall h \in [K_i, K_{i+1}), i \leq I\} \cup \{i > I\}, \quad i \geq 0,$$

Here

$$\lambda_i := n(1 - \delta) - \min\{4^{i+1}\kappa_0 + 2^{i+1}\kappa_B^{(0)}, cn\}, \quad \text{with } \delta \in (0, 1 - c) \text{ arbitrarily fixed,}$$

$$\phi_i := \max\left\{18ne^{-4^i \kappa_0 p \min\{(1-\varepsilon)H(1/8), \frac{1}{18} \log(\frac{1}{18\varepsilon})\}}, g\right\}$$

where $\varepsilon > 0$ is arbitrarily small. Now, we can check that for n large enough:

$$(8.39) \quad \mathcal{A}_i \cap \mathcal{C}_0 = \mathcal{B}_i \cap \mathcal{C}_0 = \mathcal{C}_0, \quad \text{for } i > \bar{i}.$$

Indeed choose $i > \bar{i}$, then by (8.35), $\mathcal{C}_0 \subseteq \{i > I\} = \Omega = \mathcal{B}_i$. This proves $\mathcal{B}_i \cap \mathcal{C}_0 = \mathcal{C}_0$, for $i > \bar{i}$. Now, note that by construction $Z_{I(\omega)+j}(\omega) = T_{\lfloor cn \rfloor}(\omega)$, for any $j \geq 1$. Therefore, for any $\omega \in \mathcal{C}_0$ and $I(\omega) \leq \bar{i} < i$, we have $Z_{i+1}(\omega) = Z_{I(\omega)+j}(\omega) = T_{\lfloor cn \rfloor}(\omega)$ (with $j = i + 1 - I(\omega) > 1$), then

$$Z_{i+1}(\omega) = T_{\lfloor cn \rfloor}(\omega) = \min\{T_{4^{i+1}\kappa_0}(\omega), T_{2^{i+1}\kappa_B^{(0)}}^B(\omega), T_{\lfloor cn \rfloor}(\omega)\} = \min\{T_{4^{i+1}\kappa_0}(\omega), T_{\lfloor cn \rfloor}(\omega)\}$$

which finally yields $\omega \in \mathcal{A}_i$, for all n large enough and $i > \bar{i}$, and the proof of (8.39) is completed. We shall show that there exists \bar{n} such that for any $n \geq \bar{n}$:

$$(8.40) \quad \sup_{0 \leq i \leq \bar{i}} \mathbb{P}(\mathcal{B}_i^c \cap (\cap_{0 \leq j \leq i-1} \mathcal{A}_j) \cap \mathcal{C}_0) \leq n^3 \left(e^{-n(1-\frac{\delta}{2})H(\frac{1-\delta}{1-\delta/2})} + e^{-\frac{\phi_0}{2} \log 8} \right),$$

as well as,

$$(8.41) \quad \sup_{0 \leq i \leq \bar{i}} \mathbb{P}(\mathcal{A}_i^c \cap \mathcal{B}_i \cap (\cap_{0 \leq j \leq i-1} \mathcal{A}_j \cap \mathcal{B}_j) \cap \mathcal{C}_0 \cap \mathcal{D}_0) \leq e^{-\frac{\kappa_B^{(0)}}{2} \log(10)};$$

note that for the case $i = 0$ we conventionally set $(\cap_{0 \leq j \leq -1} \mathcal{A}_j) := \Omega$. Now we prove the relation

$$(8.42) \quad \begin{aligned} & (\cap_{i \geq 0} \mathcal{B}_i) \cap \mathcal{D}_0 \cap \mathcal{C}_0 \subseteq \{T_{K^*-1} \geq Z_0, Q_{k+1}^R > 0 \forall k \in [K_0, K_{I+1}]\} \\ & \subseteq \{T_{K^*} > Z_{I+1} = T_{\lfloor cn \rfloor}\}; \end{aligned}$$

observe, indeed, that we have

$$(8.43) \quad \begin{aligned} \cap_{i \geq 0} \mathcal{B}_i &= \cup_j \left((\cap_{i \geq 0} \mathcal{B}_i) \cap \{I = j\} \right) = \cup_j \left(((\cap_{i \leq j} \mathcal{B}_i) \cap (\cap_{i > j} \mathcal{B}_i)) \cap \{I = j\} \right) \\ &\supseteq \cup_j \left((\cap_{i \leq j} \mathcal{B}_i) \cap (\cap_{i > j} \{i > I\}) \cap \{I = j\} \right) = \cup_j \left((\cap_{i \leq j} \mathcal{B}_i) \cap \{I = j\} \right), \end{aligned}$$

where the set inclusion is a consequence of the relation $\mathcal{B}_i \supseteq \{i > I\}$. Comparing the second and last terms in (8.43), we immediately have

$$(8.44) \quad \cap_{i \geq 0} \mathcal{B}_i = \cup_j \left((\cap_{i \geq 0} \mathcal{B}_i) \cap \{I = j\} \right) = \cup_j \left((\cap_{i \leq j} \mathcal{B}_i) \cap \{I = j\} \right).$$

By the definition of \mathcal{B}_i , we have $\mathcal{B}_i \subseteq \{Q_{k+1}^R > 0, \forall k : k \in [K_i, K_{i+1}), i \leq I\} \cup \{i > I\}$ and so $(\cap_{i \leq j} \mathcal{B}_i) \cap \{I = j\} \subseteq \{Q_{k+1}^R > 0, \forall k : k \in [K_0, K_{I+1}), I = j\}$, which, combined with (8.44), yields $\cap_{i \geq 0} \mathcal{B}_i \subseteq \{Q_{k+1}^R > 0, \forall k : k \in [K_0, K_{I+1})\}$. Similarly we have

$$(8.45) \quad \cap_{j \leq i} \mathcal{B}_j \subseteq \{Q_{k+1}^R > 0 \forall k \in [K_0, \min(K_{i+1}, K_{I+1})\}.$$

Considering the intersection with the set $\mathcal{D}_0 \cap \mathcal{C}_0$, we finally have (8.42), where the last inclusion therein follows by noticing that $Q_{K^*}^R = 0$. Now we prove the relation:

$$(8.46) \quad (\cap_{i \geq 0} \mathcal{A}_i) \cap \mathcal{C}_0 \subseteq \{N(Z_{I+1}) \geq N(Z_I) \geq 4^I \kappa_0, N_B(Z_{I+1}) \leq 2^{I+1} \kappa_B^{(0)}\}.$$

By (8.33), the definition of \mathcal{C}_0 and \mathcal{A}_i , and (8.34) for any $\omega \in \mathcal{A}_i \cap \mathcal{C}_0 \cap \{I(\omega) = j\}$, with $i < j$, we have $Z_{i+1}(\omega) = T_{4^{i+1} \kappa_0}(\omega)$, $T_{4^{i+1} \kappa_0}(\omega) \leq T_{2^{i+1} \kappa_B^{(0)}}^B(\omega)$ and $T_{4^{i+1} \kappa_0}(\omega) < T_{\lfloor cn \rfloor}(\omega)$; while, for any $\omega \in \mathcal{A}_j \cap \mathcal{C}_0 \cap \{I(\omega) = j\}$ we have $T_{\lfloor cn \rfloor}(\omega) \leq T_{4^{j+1} \kappa_0}(\omega)$ and $T_{\lfloor cn \rfloor}(\omega) \leq T_{2^{j+1} \kappa_B^{(0)}}^B(\omega)$. In particular, for $\omega \in \mathcal{A}_{j-1} \cap \mathcal{A}_j \cap \mathcal{C}_0 \cap \{I(\omega) = j\}$, we have $Z_j(\omega) = Z_I(\omega) = T_{4^I \kappa_0}(\omega) \leq Z_{j+1}(\omega) = Z_{I+1}(\omega) = T_{\lfloor cn \rfloor}(\omega) \leq T_{2^{I+1} h_0}^B(\omega)$. Claim (8.46) easily follows taking the union over all values j that I assumes. Combining (8.42) with (8.46), we have:

$$(8.47) \quad \{(\cap_{i \geq 0} (\mathcal{A}_i \cap \mathcal{B}_i)) \cap \mathcal{C}_0 \cap \mathcal{D}_0\} \subseteq \mathcal{J} := \left\{ T_{K^*-1} \geq T_{\lfloor cn \rfloor}, \frac{N_B[\lfloor cn \rfloor]}{\lfloor cn \rfloor} \leq 2^{-\underline{i}+1} \right\},$$

where \underline{i} is defined by (8.35). We shall show later on that

$$(8.48) \quad \sum_{n \geq 1} \mathbb{P}([\cap_{i \geq 0} (\mathcal{A}_i \cap \mathcal{B}_i)] \cap \mathcal{C}_0 \cap \mathcal{D}_0)^c < \infty,$$

therefore by Borel-Cantelli lemma combined with (8.47), we obtain $\mathbb{P}(\limsup \mathcal{J}^c) = 0$, which immediately implies (5.3). To prove (8.48), we observe that by (8.39), we have $(\cap_{i \geq 0} (\mathcal{A}_i \cap \mathcal{B}_i)) \cap \mathcal{C}_0 \cap \mathcal{D}_0 = (\cap_{i=0}^{\underline{i}} (\mathcal{A}_i \cap \mathcal{B}_i)) \cap \mathcal{C}_0 \cap \mathcal{D}_0$, and so

$$(8.49) \quad \begin{aligned} \mathbb{P}([\cap_{i \geq 0} (\mathcal{A}_i \cap \mathcal{B}_i)] \cap \mathcal{C}_0 \cap \mathcal{D}_0)^c &= \mathbb{P}(\cup_{0 \leq i \leq \underline{i}} (\mathcal{A}_i^c \cup \mathcal{B}_i^c) \cup \mathcal{C}_0^c \cup \mathcal{D}_0^c) \\ &= \mathbb{P}(\cup_{0 \leq i \leq \underline{i}} (\mathcal{A}_i^c \cup \mathcal{B}_i^c) \cap \mathcal{C}_0 \cap \mathcal{D}_0) + \mathbb{P}(\mathcal{C}_0^c \cup \mathcal{D}_0^c). \end{aligned}$$

Now we have $\mathbb{P}(\liminf \mathcal{C}_0 \cap \mathcal{D}_0) = 1$, as immediate consequence of Theorem 5.1(i) for the case $g \ll q \ll p^{-1}$, (recalling that for n sufficiently large $h_0 := \lfloor p^{-1} \rfloor > \lfloor (g_B(\kappa_g) + \varepsilon)q \rfloor$) and of Theorem 4.5 (i) and Corollary 4.3 in the remaining cases. Therefore, by the second Borel-Cantelli lemma it follows $\sum_n \mathbb{P}(\mathcal{C}_0^c \cup \mathcal{D}_0^c) < \infty$, (events $\mathcal{C}_0^c \cup \mathcal{D}_0^c$ are independent for different values of n), and so, thanks to (8.49), to get (8.48) we need to show that

$$(8.50) \quad \sum_{n \geq 1} \mathbb{P}(\cup_{0 \leq i \leq \underline{i}} (\mathcal{A}_i^c \cup \mathcal{B}_i^c) \cap \mathcal{C}_0 \cap \mathcal{D}_0) < \infty.$$

To this aim, we note that proceeding similarly to (8.22), we have

$$\begin{aligned} \mathbb{P}\left(\cup_{0 \leq i \leq \bar{i}} (\mathcal{A}_i^c \cup \mathcal{B}_i^c) \cap \mathcal{C}_0 \cap \mathcal{D}_0\right) &\leq \sum_{i=0}^{\bar{i}} \mathbb{P}\left(\left(\mathcal{B}_i^c \cap \left(\cap_{0 \leq j < i} \mathcal{A}_j\right)\right) \cap \mathcal{C}_0\right) \\ &\quad + \sum_{i=0}^{\bar{i}} \mathbb{P}\left(\left(\mathcal{A}_i^c \cap \left(\mathcal{B}_i \cap \left(\cap_{0 \leq j < i} (\mathcal{A}_j \cap \mathcal{B}_j)\right)\right)\right) \cap \mathcal{C}_0 \cap \mathcal{D}_0\right). \end{aligned}$$

Now given that $\bar{i} = O(\log_2(np))$, (8.50) follows by (8.40), (8.41) and (2.1). Now to conclude the proof of the Theorem we verify (8.40) and (8.41). \square

8.4.1. *Proof of (8.40).* By (8.37) and (8.38), it follows

$$(8.51) \quad \mathcal{B}_i^c \cap \{i > I\} = \emptyset \quad \text{and} \quad \mathcal{A}_i^c \cap \{i \geq I\} = \emptyset, \quad \text{for all } n \text{ large enough.}$$

The first relation is an immediate consequence of (8.38). As far as the second relation is concerned, by (8.34) we have $Z_{I+1} = T_{[cn]}$ that implies $T_{[cn]} \leq T_{4^{I+1}\kappa_0} \leq T_{4^{i+1}\kappa_0}$ and $T_{[cn]} \leq T_{2^{I+1}\kappa_B^{(0)}} \leq T_{2^{i+1}\kappa_B^{(0)}}$ on $\{i > I\}$ by (8.33). Hence again by (8.33) we have $Z_{i+1} = T_{[cn]} = \min\{T_{4^{i+1}\kappa_0}, T_{[cn]}\}$ on $\{i > I\}$. Then, the second relation in (8.51) immediately follows from (8.37). Now, set

$$(8.52) \quad \mathcal{E}_{i-1} := \left(\cap_{1 \leq j \leq i-1} \mathcal{A}_j\right) \cap \mathcal{C}_0 \cap \{i \leq I\}.$$

Then, recalling the definition of κ_0 and $\kappa_B^{(0)}$ in (8.31) we have

$$(8.53) \quad \mathcal{E}_{i-1} \subseteq \mathcal{N}_i := \{N(h) \geq 4^i \kappa_0 \text{ and } N_B(h) \leq 2^{i+1} \kappa_B^{(0)}, \forall h \in [K_i, K_{i+1}]\}.$$

Indeed, if $\omega \in \mathcal{E}_{i-1}$, then

$$(8.54) \quad \omega \in \mathcal{A}_{i-1} \cap \mathcal{C}_0 \cap \{i \leq I\} \subseteq \{Z_i = T_{4^i \kappa_0}\} = \{K_i = 4^i \kappa_0\},$$

which implies $N[k](\omega) \geq 4^i \kappa_0$, for any k such that $T_k(\omega) \in [Z_i(\omega), Z_{i+1}(\omega)]$. Furthermore, (8.33) implies $Z_{i+1}(\omega) \leq T_{2^{i+1}\kappa_B^{(0)}}(\omega)$, which yields $N_B[k](\omega) \leq 2^{i+1} \kappa_B^{(0)}$, for any $k \in [K_i(\omega), K_{i+1}(\omega)]$. Moreover,

$$\begin{aligned} \mathcal{N}_i \cap \{k \in [K_i, K_{i+1}]\} &\subseteq \{N[k] \geq 4^i \kappa_0, N_B[k] \leq 2^{i+1} \kappa_B^{(0)}\} \\ &\subseteq \mathcal{M}_i^{(k)} := \{N_R[k] \geq 4^i \kappa_0 - 2^{i+1} \kappa_B^{(0)}, N_B[k] \leq 2^{i+1} \kappa_B^{(0)}\}, \end{aligned}$$

then recalling (8.53) it follows

$$(8.55) \quad \mathcal{E}_{i-1}^{(k)} := \mathcal{E}_{i-1} \cap \{k \in [K_i, K_{i+1}]\} \subseteq \mathcal{M}_i^{(k)}.$$

Then, setting $\underline{k}_R^{(i)} := 4^i \kappa_0 - 2^{i+1} \kappa_B^{(0)}$ and $\bar{k}_B^{(i)} = 2^{i+1} \kappa_B^{(0)}$, by Lemma 3.9 (with $k = \underline{k}_R^{(i)} + \bar{k}_B^{(i)} = 4^i \kappa_0$ and $h = \bar{k}_B^{(i)}$), we have

$$(8.56) \quad |\mathcal{S}_R[k]| \mid \mathcal{M}_i^{(k)} \geq_{st} \text{Bin}(n_W, \pi_R(\underline{k}_R^{(i)}, \bar{k}_B^{(i)})), \quad |\mathcal{S}_B[k]| \mid \mathcal{M}_i^{(k)} \leq_{st} \text{Bin}(n_W, \pi_B(\underline{k}_R^{(i)}, \bar{k}_B^{(i)})).$$

Now, note that for any $z \geq r$ and any $S \in \{R, B\}$, it holds

$$\begin{aligned} \pi_S(k_S, k_{S^c}) &= \mathbb{P}(\text{Bin}(k_S + a_S, p) - \text{Bin}(k_{S^c} + a_{S^c}, p) \geq r) \\ &\geq \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq z, \text{Bin}(k_{S^c} + a_{S^c}, p) \leq z - r) \\ (8.57) \quad &\geq 1 - \mathbb{P}(\text{Bin}(k_S + a_S, p) < z) - \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) > z - r). \end{aligned}$$

Note also that for n sufficiently large, by assuming $\kappa_B^{(0)}/\kappa_0 < (\kappa_B^{(0)} + a_B)/\kappa_0 < \varepsilon/2$, we have

$$\mathbb{E}[\text{Bin}(\underline{k}_R^{(i)} + a_R, p)] \geq \mathbb{E}[\text{Bin}(\underline{k}_R^{(i)}, p)] \geq 4^i \kappa_0 p \left(1 - \frac{2\kappa_B^{(0)}}{2^i \kappa_0}\right) \geq 4^i \kappa_0 p (1 - \varepsilon)$$

$$\mathbb{E}[\text{Bin}(\bar{k}_B^{(i)} + a_B, p)] \leq 2^{i+1} (\kappa_B^{(0)} + a_B) p.$$

Therefore, taking $z = 4^i \kappa_0 p / 9$ by (8.57) and the concentration inequalities in Appendix I, for any i and sufficiently large n , it holds

$$(8.58) \quad \begin{aligned} \pi_R(\underline{k}_R^{(i)}, \bar{k}_B^{(i)}) &\geq 1 - e^{-4^i \kappa_0 p (1-\varepsilon) H(1/8)} - e^{-\frac{4^i}{18} \kappa_0 p \log(2^{i-1} \cdot \frac{1}{9\varepsilon})} \\ \pi_B(\underline{k}_R^{(i)}, \bar{k}_B^{(i)}) &\leq e^{-4^i \kappa_0 p (1-\varepsilon) H(1/8)} + e^{-\frac{4^i}{18} \kappa_0 p \log(2^{i-1} \cdot \frac{1}{9\varepsilon})}. \end{aligned}$$

Combining (8.58) with (8.56), we have

$$\mathbb{E}[|\mathcal{S}_R[k]| \mid \mathcal{M}_i^{(k)}] \geq n_W \left(1 - e^{-\kappa_0 p (1-\varepsilon) H(1/8)} - e^{-\frac{1}{18} \kappa_0 p \log(\frac{1}{18\varepsilon})}\right),$$

where we have exploited the monotonicity with respect to i of the r.h.s. of (8.58). Note that for n large enough we can always assume $\kappa_0 p$ to be sufficiently large that $\mathbb{E}[|\mathcal{S}_R[k]| \mid \mathcal{M}_i^{(k)}] \geq n(1 - \frac{\delta}{2})$ for an arbitrary $\delta > 0$. Now, applying again the concentration inequality reported in Appendix I, for any i and all n large enough, we have

$$(8.59) \quad \mathbb{P}(|\mathcal{S}_R[k]| \leq (1 - \delta)n \mid \mathcal{M}_i^{(k)}) < e^{-n(1 - \frac{\delta}{2})H(\frac{1-\delta}{1-\frac{\delta}{2}})}.$$

Similarly, given (8.56) and (8.58), for any i and all n sufficiently large, we have

$$\mathbb{E}[|\mathcal{S}_B[k]| \mid \mathcal{M}_i^{(k)}] \leq 2n e^{-4^i \kappa_0 p \min\{(1-\varepsilon)H(1/8), \frac{1}{18} \log(\frac{1}{18\varepsilon})\}} := \bar{\mu}_i^B.$$

Setting $\phi_i := \max(9\bar{\mu}_i^B, g)$, for all i and all n large enough, we have

$$(8.60) \quad \mathbb{P}\left(|\mathcal{S}_B[k]| \geq \phi_i \mid \mathcal{M}_i^{(k)}\right) \leq e^{-\frac{\phi_i}{2} \log 8}.$$

By (8.38), for any i and all n large enough, we have

$$\begin{aligned} \mathbb{P}(\mathcal{B}_i^c \cap \mathcal{E}_{i-1}) &= \mathbb{P}\left(\bigcup_k \left(\{k \in [K_i, K_{i+1}), Q_{k+1}^R \leq \lambda_i \text{ or } Q_{k+1}^B > \phi_i\} \cap \mathcal{E}_{i-1}\right)\right) \\ &= \mathbb{P}\left(\bigcup_k \left(\{Q_{k+1}^R \leq \lambda_i \text{ or } Q_{k+1}^B > \phi_i\} \cap \mathcal{E}_{i-1}^{(k)}\right)\right) \\ &= \sum_{k_i, k_{i+1}} \mathbb{P}\left(\{K_i = k_i, K_{i+1} = k_{i+1}\} \cap \left(\bigcup_{k=k_i}^{k_{i+1}-1} [\{Q_{k+1}^R \leq \lambda_i\} \cup \{Q_{k+1}^B > \phi_i\}] \cap \mathcal{E}_{i-1}^{(k)}\right)\right) \\ &\stackrel{(a)}{\leq} \sum_{k_i, k_{i+1}} \sum_{k=k_i}^{k_{i+1}-1} \mathbb{P}\left(\{K_i = k_i, K_{i+1} = k_{i+1}\} \cap \left(\{Q_{k+1}^R \leq \lambda_i\} \cup \{Q_{k+1}^B > \phi_i\}\right) \cap \mathcal{M}_i^{(k)}\right) \\ &\leq \sum_{k_i, k_{i+1}} \sum_{k=k_i}^{k_{i+1}-1} \mathbb{P}(\{Q_{k+1}^R \leq \lambda_i\} \cup \{Q_{k+1}^B > \phi_i\}) \cap \mathcal{M}_i^{(k)} \end{aligned}$$

$$\begin{aligned}
&\stackrel{(b)}{\leq} \sum_{k_i, k_{i+1}} \sum_{k=k_i}^{k_{i+1}-1} \mathbb{P}(\{|\mathcal{S}_R[k]| \leq \lambda_i + k\} \cup \{|\mathcal{S}_B[k]| > \phi_i\} \mid \mathcal{M}_i^{(k)}) \\
(8.61) \quad &\stackrel{(c)}{\leq} n^3 \left(e^{-n(1-\frac{\delta}{2})H\left(\frac{1-\delta}{1-\frac{\delta}{2}}\right)} + e^{-\frac{\phi_0}{2} \log 8} \right),
\end{aligned}$$

where indices k_i and k_{i+1} in the sums span over the support of K_i and K_{i+1} , respectively. Here inequality (a) follows from (8.55), inequality (b) follows from the relations (3.11), inequality (c) follows from the relations (8.59) and (8.60) (note that $\lambda_i + k \leq (1 - \delta)n$, the union bound, the fact that K_i , for every i , takes values in $\{0, \dots, n_W\}$ and the monotonicity of ϕ_i w.r.t. i . Relation (8.40) follows from (8.61), noticing that, due to the first equality in (8.51), we have

$$\mathbb{P}(\mathcal{B}_i^c \cap (\cap_{1 \leq j \leq i-1} \mathcal{A}_j) \cap \mathcal{C}_0) = \mathbb{P}(\mathcal{B}_i^c \cap (\cap_{1 \leq j \leq i-1} \mathcal{A}_j) \cap \mathcal{C}_0 \cap \{i \leq I\}) = \mathbb{P}(\mathcal{B}_i^c \cap \mathcal{E}_{i-1}). \quad \square$$

8.4.2. *Proof of (8.41)*. Since $Q_{K^*}^R = 0$, by (8.32), (8.45) and (8.52), we have that

$$\mathcal{D}_0 \cap (\cap_{j \leq i} \mathcal{B}_j) \cap \mathcal{E}_{i-1} \subseteq \mathcal{C}_0 \cap \mathcal{D}_0 \cap (\cap_{j \leq i} \mathcal{B}_j) \subseteq \{T_{K^*} \geq Z_{i+1}\}.$$

Therefore, by Lemma 3.6, on $\mathcal{D}_0 \cap (\cap_{j \leq i} \mathcal{B}_j) \cap \mathcal{E}_{i-1}$, for $S \in \{R, B\}$, we have $Q_{k+1}^S = |\mathcal{V}_W[k] \cap \mathcal{S}_S[k]|, \forall k \in [K_0, K_{i+1})$. In addition, by (3.6) we have $N_S(K_{i+1}) = N_S(K_i) + \sum_{k=1}^{K_{i+1}-K_i} M_{K_i+k}^S$. Recalling that variables $\{U_k^S\}_{k \in \mathbb{N}}$ have a finite support, and letting u denote an arbitrary element in it, by proposition 3.3 we have that $M_k^S \mid \{U_k^S = u\}$ is Bernoulli distributed with mean u . Again by Proposition 3.4 random variable $M_k^S \mid \{U_k^S = u\}$ is independent of \mathcal{H}_{k-1} . Consequently we can define the following sequence of random variables:

$$\overline{M}_k^B := \begin{cases} 1 & \text{on } \{M_k^B = 1\} \cup \{U_k^B > \frac{\phi_i}{\lambda_i + \phi_i}\} \\ \text{Be} \left(\frac{\frac{\phi_i}{\lambda_i + \phi_i} - u}{1-u} \right) & \text{on } \{M_k^B = 0\} \cap \{U_k^B = u \leq \frac{\phi_i}{\lambda_i + \phi_i}\} \end{cases} \quad \underline{M}_k^R := 1 - \overline{M}_k^B$$

Clearly $\overline{M}_k^B \geq M_k^B$ and $\underline{M}_k^R \leq M_k^R$ \mathbb{P} -a.s.. Moreover it is of immediate verification that:

$$(8.62) \quad \overline{M}_k^B := \text{Be} \left(\frac{\phi_i}{\lambda_i + \phi_i} \right), \quad \text{on the event } \left\{ U_k^B = u \leq \frac{\phi_i}{\lambda_i + \phi_i} \right\},$$

Note that random variables $\overline{M}_k^B \mid \{U_k^B = u\}$ and $\underline{M}_k^R \mid \{U_k^B = u\}$ are independent of \mathcal{H}_{k-1} . By (8.52) and (8.54), we have $\mathcal{E}_{i-1} \subseteq \mathcal{A}_{i-1} \cap \mathcal{C}_0 \cap \{i \leq I\} \subseteq \{K_i = 4^i \kappa_0\}$, therefore,

$$(8.63) \quad \mathcal{G}_i := \mathcal{D}_0 \cap (\cap_{j < i} \mathcal{B}_j) \cap \mathcal{E}_{i-1} \subseteq \{K_i = 4^i \kappa_0\}.$$

In addition, recalling (8.38) we have

$$\mathcal{B}_i \cap \{i \leq I\} \subseteq \tilde{\mathcal{B}}_i := \left\{ U_{k+1}^B < \frac{\phi_i}{\lambda_i + \phi_i} \forall k : k \in [K_i, K_{i+1}) \right\}.$$

Setting $\mathbf{u}_h := (u_1, \dots, u_h)$, where u_m , for $1 \leq m \leq h$, denotes an arbitrary element in the support of $\{U_k^S\}_{k \in \mathbb{N} \cup \{0\}}$ and $\mathcal{U}_B(\mathbf{u}_h) := \cap_{m=1}^h \{U_{4^i \kappa_0 + m}^B = u_m\}$, we have

$$\tilde{\mathcal{B}}_i \cap \{K_{i+1} - K_i = h\} \cap \{K_i = 4^i \kappa_0\} = \bigcup_{\mathbf{u}_h < \frac{\phi_i}{\phi_i + \lambda_i} \mathbf{1}} \mathcal{U}_B(\mathbf{u}_h) \cap \{K_{i+1} - K_i = h\} \cap \{K_i = 4^i \kappa_0\}.$$

with $\mathbf{1} := (1, \dots, 1) \in \mathbb{R}^h$ and $|\{\mathbf{u}_h : \mathbf{u}_h < \frac{\phi_i}{\phi_i + \lambda_i} \mathbf{1}\}| < \infty$, as oblivious consequence of the fact that the support of $\{U_k^S\}_{k \in \mathbb{N} \cup \{0\}}$ is finite. Setting $\Delta_{\max} := \min\{4^{i+1}\kappa_0, \lfloor cn \rfloor\} - 4^i\kappa_0$, by (8.63), and the fact that $\mathcal{E}_{i-1} \subseteq \mathcal{C}_0 \cap \{i \leq I\}$ (see (8.52)), we have:

$$\begin{aligned}
\{K_{i+1} < \Delta_{\max} + K_i\} \cap \mathcal{B}_i \cap \mathcal{G}_i &\subseteq \bigcup_{h=0}^{\Delta_{\max}-1} \{K_{i+1} = h + 4^i\kappa_0\} \cap \tilde{\mathcal{B}}_i \cap \mathcal{G}_i \\
&= \bigcup_{h=0}^{\Delta_{\max}-1} \bigcup_{\mathbf{u}_h < \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \{K_{i+1} = h + 4^i\kappa_0\} \cap \mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i \\
(8.64) \quad &\subseteq \bigcup_{h=0}^{\Delta_{\max}-1} \bigcup_{\mathbf{u}_h < \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \left\{ \sum_{m=1}^h M_{4^i\kappa_0+m}^B \geq 2^i\kappa_B^{(0)} \right\} \cap \mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i
\end{aligned}$$

$$(8.65) \quad \subseteq \bigcup_{h=0}^{\Delta_{\max}-1} \bigcup_{\mathbf{u}_h < \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \left\{ \sum_{m=1}^h \overline{M}_{4^i\kappa_0+m}^B \geq 2^i\kappa_B^{(0)} \right\} \cap \mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i.$$

Here, recalling (8.36), the inclusion (8.64) stems from the fact that, for any $0 \leq h < \Delta_{\max}$,

$$\begin{aligned}
\{K_{i+1} = h + 4^i\kappa_0\} \cap \mathcal{G}_i &= \{K_{i+1} - K_i = h\} \cap \mathcal{G}_i \\
(8.66) \quad &\subseteq \{N_B[K_{i+1}] = N_B[K_i + h] = 2^{i+1}\kappa_B^{(0)}\} \cap \mathcal{G}_i
\end{aligned}$$

$$\begin{aligned}
(8.67) \quad &= \{N_B[K_i + h] - N_B[K_i] = 2^{i+1}\kappa_B^{(0)} - N_B[K_i], N_B[K_i] \leq 2^i\kappa_B^{(0)}\} \cap \mathcal{G}_i \\
&\subseteq \{N_B[K_i + h] - N_B[K_i] \geq 2^i\kappa_B^{(0)}\} \cap \mathcal{G}_i = \left\{ \sum_{m=1}^h M_{4^i\kappa_0+m}^B \geq 2^i\kappa_B^{(0)} \right\} \cap \mathcal{G}_i,
\end{aligned}$$

where relation (8.67) is a consequence of (8.36) and the inclusion (8.66) follows from the definition of Z_{i+1} in (8.33). Indeed, since $h < \Delta_{\max}$ we have that $K_{i+1} < \min\{4^{i+1}\kappa_0, \lfloor cn \rfloor\}$, i.e., $Z_{i+1} < \min\{T_{4^{i+1}\kappa_0}, T_{\lfloor cn \rfloor}\}$, and so $Z_{i+1} = T_{2^{i+1}\kappa_B^{(0)}}^B$, i.e. $N_B(K_{i+1}) = 2^{i+1}\kappa_B^{(0)}$. By (8.33) and the definition of \mathcal{A}_i in (8.37), we have

$$\mathcal{A}_i^c = \{K_{i+1} < \min\{4^{i+1}\kappa_0, \lfloor cn \rfloor\}\},$$

and so by (8.65) we have

$$\begin{aligned}
\mathbb{P}(\mathcal{A}_i^c \cap \mathcal{B}_i \cap \mathcal{G}_i) &\leq \mathbb{P}(\mathcal{A}_i^c \cap \tilde{\mathcal{B}}_i \cap \mathcal{G}_i) = \mathbb{P}(\{K_{i+1} < \Delta_{\max} + 4^i\kappa_0\} \cap \tilde{\mathcal{B}}_i \cap \mathcal{G}_i) \\
&\leq \sum_{h=1}^{\Delta_{\max}-1} \sum_{\mathbf{u}_h < \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \mathbb{P}\left(\left\{ \sum_{m=1}^h \overline{M}_{4^i\kappa_0+m}^B \geq 2^i\kappa_B^{(0)} \right\} \cap \mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i\right) \\
&= \sum_{h=1}^{\Delta_{\max}-1} \sum_{\mathbf{u}_h < \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \mathbb{P}\left(\sum_{m=1}^h \overline{M}_{4^i\kappa_0+m}^B \geq 2^i\kappa_B^{(0)} \mid \mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i\right) \mathbb{P}(\mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i) \\
&\stackrel{(a)}{=} \sum_{h=1}^{\Delta_{\max}-1} \sum_{\mathbf{u}_h < \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \mathbb{P}\left(\sum_{m=1}^h \overline{M}_{4^i\kappa_0+m}^B \geq 2^i\kappa_B^{(0)} \mid \mathcal{U}_B(\mathbf{u}_h)\right) \mathbb{P}(\mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i)
\end{aligned}$$

$$\begin{aligned}
&\stackrel{(b)}{=} \sum_{h=1}^{\Delta_{\max}-1} \mathbb{P} \left(\text{Bin} \left(h, \frac{\phi_i}{\lambda_i + \phi_i} \right) \geq 2^i \kappa_B^{(0)} \right) \sum_{\mathbf{u}_h \leq \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}} \mathbb{P} (\mathcal{U}_B(\mathbf{u}_h) \cap \mathcal{G}_i) \\
&\leq \sum_{h=1}^{\Delta_{\max}-1} \mathbb{P} \left(\text{Bin} \left(h, \frac{\phi_i}{\lambda_i + \phi_i} \right) \geq 2^i \kappa_B^{(0)} \right) \mathbb{P} (\mathcal{G}_i) \\
&\leq (\Delta_{\max} - 1) \mathbb{P} \left(\text{Bin} \left(\Delta_{\max} - 1, \frac{\lambda_i}{\lambda_i + \phi_i} \right) \geq 2^i \kappa_B^{(0)} \right) \mathbb{P} (\mathcal{G}_i) \\
&\leq (3 \cdot 4^i \kappa_0) \mathbb{P} \left(\text{Bin} \left(3 \cdot 4^i \kappa_0, \frac{\phi_i}{\lambda_i + \phi_i} \right) \geq 2^i \kappa_B^{(0)} \right) \mathbb{P} (\mathcal{G}_i).
\end{aligned}$$

Here equality (a) follows from recalling that, given $\mathcal{U}_B(\mathbf{u}_h)$, $\mathbf{u}_h \leq \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}$, random variables $\{\overline{M}_{4^i \kappa_0 + m}^B\}_{1 \leq m \leq h}$ are independent of $\mathcal{H}_{4^i \kappa_0}$, and therefore of \mathcal{G}_i , given that $\mathcal{G}_i = \mathcal{D}_0 \cap (\cap_{j < i} \mathcal{B}_j) \cap \mathcal{E}_{i-1} \in \mathcal{H}_{4^i \kappa_0}$. Equality (b) is a consequence of the fact that, given $\mathcal{U}_B(\mathbf{u}_h)$, with $\mathbf{u}_h \leq \frac{\phi_i}{\lambda_i + \phi_i} \mathbf{1}$, random variables $\{\overline{M}_{4^i \kappa_0 + m}^B\}_{1 \leq m \leq h}$ follow the Bernoulli law with mean $\frac{\phi_i}{\lambda_i + \phi_i}$ by (8.62). Finally, recalling that $\Delta_{\max} \leq 3 \cdot 4^i \kappa_0$ and applying the usual concentration inequality for the binomial law (see Appendix I), for all n large enough, we have

$$\mathbb{P} \left(\text{Bin} \left(3 \cdot 4^i \kappa_0, \frac{\phi_i}{\lambda_i + \phi_i} \right) \geq 2^i \kappa_B^{(0)} \right) \leq e^{-2^{i-1} \kappa_B^{(0)} \log(10)}.$$

Indeed, from the definition of ϕ_i and λ_i we get:

$$\begin{aligned}
&\frac{\phi_i}{\lambda_i + \phi_i} 3 \cdot 4^i \kappa_0 \leq 3 \cdot 4^i \kappa_0 \frac{\phi_i}{\lambda_i} \\
(8.68) \quad &\leq \frac{3 \cdot 4^i \kappa_0}{1 - \delta - c} \max \left\{ 18e^{-4^i \kappa_0 p \min\{(1-\varepsilon)H(1/8), \frac{1}{18} \log(\frac{1}{18\varepsilon})\}}, \frac{g}{n} \right\} < \frac{\kappa_B^{(0)}}{20}
\end{aligned}$$

where we recall that we can assume $\kappa_0 p$ to be sufficiently large. \square

REFERENCES

- [1] CHALUPA J., LEATH, P. L. and REICH G. R. (1979). Bootstrap percolation on a Bethe lattice. *Journal of Physics C*, **12** 31–35.
- [2] BALOGH J. and BOLLOBÁS, B. (2006). Bootstrap percolation on the hypercube. *Probability Theory and Related Fields*, **134** 624–648.
- [3] BRÉMAUD P. (2020). *Markov Chains. Gibbs Fields, Monte Carlo Simulation and Queues*. Springer, New York.
- [4] BALOGH J., PERES Y. and PETE G. (2006). Bootstrap percolation on infinite trees and non-amenable groups. *Combinatorics, Probability and Computing*, **15** 715–730.
- [5] BALOGH J. and PITTEL B. G. (2007). Bootstrap percolation on the random regular graph. *Random Structures and Algorithms*, **30** 257–286.
- [6] BOLLOBÁS B., GUNDERSON K., HOLMGREN C., JANSON S. and PRZYKUCKI M. (2014). Bootstrap percolation on Galton-Watson trees. *Electronic Journal of Probability*, **19** 1–27.
- [7] BILLINGSLEY, P. (1995). *Probability and Measure*, 2nd ed. Wiley, New York.
- [8] JANSON S., ŁUCZAK T., TUROVA T. S. and VALLIER T. (2012). Bootstrap percolation on the random graph $G_{n,p}$. *Annals of Applied Probability*, **22**(5) 1989–2047.
- [9] AMINI H. (2010). Bootstrap percolation and diffusion in random graphs with given vertex degrees. *Electronic Journal of Combinatorics*, **17** 1–20.
- [10] AMINI H. and FOUNTOLAKIS N. (2014). Bootstrap percolation in power-law random graphs. *Journal of Statistical Physics*, **155** 72–92.
- [11] FOUNTOLAKIS N., KANG M., KOCH C. and MAKAI T. (2018). A phase transition regarding the evolution of bootstrap processes in inhomogeneous random graphs. *Annals of Applied Probability*, **28**(2) 990–1051.

- [12] JANSON S., KOZMA R., RUSZINKÓ M. and SOKOLOV Y. (2019). A modified bootstrap percolation on a random graph coupled with a lattice. *Discrete Applied Mathematics*, **258** 152–165.
- [13] TUROVA T. S. and VALLIER T. (2015). Bootstrap percolation on a graph with random and local connections. *Journal of Statistical Physics*, **160** 1249–1276.
- [14] FOUNTOULAKIS N. and ABDULLAH A. (2018). A phase transition in the evolution of bootstrap percolation processes on preferential attachment graphs. *Random Structures and Algorithms*, **52(3)** 379–418.
- [15] BRADONJIĆ M. and SANIEE I. (2014). Bootstrap percolation on random geometric graphs. *Probability in the Engineering and Informational Sciences*, **28** 169–181.
- [16] JANSON S. (2018). Tail Bounds for sums of geometric and exponential random variables. *Statistics & Probability Letters*, **135** 1–6.
- [17] HOLMGREN C., JUŠKEVIČIUS Z. and Kettle N. (2017). Majority bootstrap percolation on $G_{n,p}$. *Electronic Journal of Combinatorics*, **24(1)**.
- [18] LAST G. and PENROSE M. (2017). *Lectures on the Poisson Point Process*. Cambridge University Press, Cambridge, UK.
- [19] PENROSE M. (2004). *Random Geometric Graphs*. Oxford University Press, Oxford, UK.
- [20] SCALIA-TOMBA G. P. (1985). Asymptotic final-size distribution for some chain-binomial processes. *Advances in Applied Probability*, **17** 477–495.
- [21] TORRISI G. L., GARETTO M. and LEONARDI E. (2019). A large deviation approach to super-critical bootstrap percolation on the random graph $G_{n,p}$. *Stochastic Processes and their Applications*, **129(6)** 1873–1902.
- [22] TORRISI G. L., GARETTO M. and LEONARDI E. (2023). Bootstrap percolation on the stochastic block model. *Bernoulli* **29(1)** 696–724.
- [23] EINARSSON H., LENGLER J., MOUSSET F., PANAGIOTOU K. and STEGER A. (2019). Bootstrap percolation with inhibition. *Random Structures and Algorithms*, **55(4)** 881–925.
- [24] FEIGE U., KRIVELEVICH M. and REICHMAN D. (2016). Contagious sets in random graphs. *Annals of Applied Probability*, **27(5)** 2675–2697.
- [25] GARBE F., MYCROFT R. and MCDOWELL A. (2018). Contagious sets in a degree-proportional bootstrap percolation process. *Random Structures and Algorithms*, **53(4)** 638–651.
- [26] ANGEL O. and Kolesnik B. (2021) Large Deviations for Subcritical Bootstrap Percolation on the Erdős–Rényi Graph. *Journal of Statistical Physics*, **185(8)**.

APPENDIX A: PROOF OF PROPOSITION 4.4

To prove Propositions 4.4(i)–(ii) we start establishing a relation between the solutions of Cauchy Problems (4.2) and (2.6).

LEMMA A.1. Assume $\beta_S(x_R, x_B) = \beta_S(x_S)$, $S \in \{R, S\}$, and that the Cauchy problem (2.6) has a unique maximal solution \mathbf{g} on $(0, \kappa_{\mathbf{g}})$ with g_R and g_B strictly increasing. Then the Cauchy problem (4.2) has a unique maximal solution on $(0, \kappa_{\mathbf{f}})$, with $\kappa_{\mathbf{f}} := z(\kappa_{\mathbf{g}})$, where $z := g_R + g_B$, given by

$$\mathbf{f}(x) = \mathbf{g}(z^{-1}(x)).$$

For Cauchy’s problems of the form (2.6), recalling that we always assume $\alpha_R > \alpha_B$, the following propositions hold:

PROPOSITION A.2. Under the assumption that $q = g$:

(i) If $\alpha_R < 1$, then Cauchy problem (2.6) has a unique solution \mathbf{g} defined on $(0, \infty)$ and

$$g_R(x) \uparrow z_R, \quad g_B(x) \uparrow z_B, \quad \text{as } x \uparrow +\infty.$$

(ii) If $\alpha_R > 1$, then Cauchy problem (2.6) has a unique solution \mathbf{g} defined on $(0, \kappa_{\mathbf{g}})$, with

$$g_R(x) \uparrow +\infty, \quad g_B(x) \uparrow g_B(\kappa_{\mathbf{g}}) \in (0, \infty), \quad \text{as } x \uparrow \kappa_{\mathbf{g}}. \quad \text{and}$$

$$\kappa_{\mathbf{g}} := \int_0^{\infty} \frac{dx}{-x + r^{-1}(1 - r^{-1})^{r-1}(\alpha_R + x)^r} < \infty,$$

If $\alpha_B < 1$ then $g_B(\kappa_{\mathbf{g}}) < z_B$.

PROPOSITION A.3. Under the assumption that $g \ll q \ll p^{-1}$, Cauchy problem (2.6) admits the following unique solution \mathbf{g} on $(0, \kappa_{\mathbf{g}})$:

$$g_S(x) = \frac{1}{(\alpha_S^{1-r} - \frac{r-1}{r!}x)^{1/(r-1)}} - \alpha_S \quad S \in \{R, B\}, \quad \kappa_{\mathbf{g}} := \frac{r!}{(r-1)\alpha_R^{r-1}},$$

Propositions 4.4(i)-(ii) immediately follow from Propositions A.2 and A.3 and Lemma A.1. Proposition 4.4(iii) is a simple consequence of the next Lemma A.4 and the relation

$$\frac{\beta_B(x_R, x_B)}{\beta_R(x_R, x_B)} \leq \left(\frac{x_B + \alpha_B}{x_R + \alpha_R} \right)^r, \quad \text{for } x_B + \alpha_B < x_R + \alpha_R.$$

LEMMA A.4. Let $\mathbf{f}(x)$ be the solution of Cauchy problem 4.2 for $q = p^{-1}$ and let $\tilde{\mathbf{f}}(x)$ be the solution of the Cauchy problem:

$$(A.1) \quad \tilde{\mathbf{f}}'(x) = \frac{\tilde{\beta}(\tilde{\mathbf{f}}(x))}{\tilde{\beta}_R(\tilde{f}_R(x)) + \tilde{\beta}_B(\tilde{f}_B(x))}, \quad x \in (0, \kappa_{\mathbf{f}}), \quad \mathbf{f}(0) = (0, 0)$$

where $\tilde{\beta}_S(\mathbf{x}) = (x_S + \alpha_S)^r$, $S \in \{R, B\}$. Then $f_R(x) > \tilde{f}_R(x)$ and $f_B(x) < \tilde{f}_B(x)$, for every $x \in (0, \kappa_{\mathbf{f}})$.

The proofs of the above Lemmas and Propositions are elementary. They are reported for completeness in Appendix J.

APPENDIX B: NUMERICAL ILLUSTRATION OF THE RESULTS FOR $q = g$

For the purpose of numerical illustration of our results, we consider the case $r = 2$, which allows closed-form solutions of many quantities of interest. We focus on the super-critical case $\alpha_R > 1$ (super-critical R -activation process) and consider either the case $\alpha_B < 1$ or $\alpha_B > 1$. Whenever $\alpha_R > 1$, with $r = 2$ we get from Proposition 4.4

$$(B.1) \quad \kappa_{\mathbf{g}} := \int_0^\infty \frac{dx}{\frac{(x+\alpha_R)^2}{4} - x} = \frac{2}{\sqrt{\alpha_R - 1}} \left(\frac{\pi}{2} - \arctan \left(\frac{\frac{\alpha_R}{2} - 1}{\sqrt{\alpha_R - 1}} \right) \right)$$

Note that $\kappa_{\mathbf{g}}$ can be interpreted as the physical time at which the red activation process produces a number of susceptible red nodes of the order of n . As expected, with $\alpha_R \downarrow 1$ we have that $\kappa_{\mathbf{g}}$ tends to infinite. This can be understood by the fact that the activation process becomes increasingly slow while approaching the percolation transition.

In the case $\alpha_B < 1$, recall that $\beta_B(x) = \frac{(x+\alpha_B)^2}{4} - x$ has two zeros at $x_a = 2 - \alpha_B - 2\sqrt{1 - \alpha_B}$ and $x_b = 2 - \alpha_B + 2\sqrt{1 - \alpha_B}$.

In this case, we also get a closed-form expression for the generic integral:

$$(B.2) \quad \int_0^x \frac{du}{\beta_B(u)} = \int_0^x \frac{du}{\frac{(u+\alpha_B)^2}{4} - u} = \frac{1}{4\sqrt{1 - \alpha_B}} \log \left(\frac{x_a(x_b - x)}{x_b(x_a - x)} \right)$$

(note that, as expected, the above integral diverges as $x \uparrow x_a$).

Let

$$\psi(\alpha_R, \alpha_B) := \exp \left(2\sqrt{\frac{1 - \alpha_B}{\alpha_R - 1}} \left(\frac{\pi}{2} - \arctan \left(\frac{\frac{\alpha_R}{2} - 1}{\sqrt{\alpha_R - 1}} \right) \right) \right).$$

Exploiting the results in Theorem 4.7, after some simple algebra, we can compute the asymptotic behavior of the (normalized) final number of active black nodes as:

$$(B.3) \quad \frac{A_B^*}{q} \sim \alpha_B + \frac{\alpha_B^2(\psi - 1)}{(2 - \alpha_B)(\psi - 1) + 2\sqrt{1 - \alpha_B}(\psi + 1)}.$$

When $\alpha_R \downarrow 1$ we have that ψ diverges to ∞ , and we recover the well known result of the classical subcritical bootstrap percolation process with $r = 2$, for which the final (normalized) number of active nodes is $2 - 2\sqrt{1 - \alpha_B}$.

In the case $\alpha_B > 1$, we define instead the constant:

$$\psi'(\alpha_R, \alpha_B) := \arctan\left(\frac{\frac{\alpha_B - 1}{2}}{\sqrt{\alpha_B - 1}}\right) + \sqrt{\frac{\alpha_B - 1}{\alpha_R - 1}} \left(\frac{\pi}{2} - \arctan\left(\frac{\frac{\alpha_B - 1}{2}}{\sqrt{\alpha_B - 1}}\right)\right).$$

Then the (normalized) final number of active black nodes:

$$(B.4) \quad \frac{A_B^*}{q} \sim 2 + 2\sqrt{\alpha_B - 1} \tan(\psi').$$

As expected, with $\alpha_B \downarrow 1$ this number tends to 2, matching the same figure obtained (in the case $\alpha_B < 1$) when $\alpha_B \uparrow 1$. One can also check that, for increasing values of α_R , A_B^*/q approaches α_B (meaning that the infection of black nodes essentially does not evolve, being immediately stopped by the red infection). Instead, as $\alpha_R \downarrow \alpha_B$, A_B^*/q diverges (note indeed that in this case $\psi' \uparrow \frac{\pi}{2}$).




FIG 1. $\alpha_B < 1$: A_B^*/q as function of α_R , for different values of α_B .




FIG 2. $\alpha_B > 1$: A_B^*/q as function of $\alpha_R > \alpha_B$, for different values of α_B .

APPENDIX C: PROOF OF LEMMA 3.6

For every k , we have

$$\begin{aligned} |\mathcal{V}_W[k] \cap \mathcal{S}_S[k]| &= |(\mathcal{V}_W \setminus (\mathcal{V}_R[k] \cup \mathcal{V}_B[k]) \cap \mathcal{S}_S[k])| \\ &= |\mathcal{S}_S[k] \setminus ((\mathcal{V}_R[k] \cup \mathcal{V}_B[k]) \cap \mathcal{S}_S[k])| = |\mathcal{S}_S[k]| - |\mathcal{S}_S[k] \cap \mathcal{V}_R[k]| - |\mathcal{S}_S[k] \cap \mathcal{V}_B[k]| \end{aligned}$$

and

$$|\mathcal{S}_S[k] \cap \mathcal{V}_S[k]| = |\mathcal{V}_S[k]| - |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k]| = N_S[k] - |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k]|.$$

Therefore

$$(C.1) \quad |\mathcal{V}_W[k] \cap \mathcal{S}_S[k]| = |\mathcal{S}_S[k]| - N_S[k] + |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k]| - |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k]|.$$

On the event $\{k < K^*\}$, we have $\mathcal{V}_S[k] \equiv \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}$, due to the monotonicity of paths of $D_S^v(\cdot)$ and the fact that a node becomes S -active only if it is S -susceptible

$$(C.2) \quad |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k]| = |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}|$$

and

$$(C.3) \quad |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k]| = |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}|.$$

By (C.1), (C.2) and (C.3), it easily follows (3.10). Now we prove (3.11). For any $0 \leq k < n_W$, by (3.9) we have

$$\begin{aligned} Q_{k+1}^S &= |\mathcal{S}_S[k]| - N_S[k] + |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| \\ &\quad - |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| \\ &\geq |\mathcal{S}_S[k]| - N_S[k] - |\mathcal{S}_B[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| \\ &\geq |\mathcal{S}_S[k]| - N_S[k] - |\mathcal{V}_{S^c}[k]| \\ &= |\mathcal{S}_S[k]| - N_S[k] - N_{S^c}[k] = |\mathcal{S}_S[k]| - k, \end{aligned}$$

where we have applied the identities $|\mathcal{V}_S[k]| = N_S[k]$ and $k = N_S[k] + N_{S^c}[k]$. On the other hand

$$(C.4) \quad \begin{aligned} Q_{k+1}^S &= |\mathcal{S}_S[k]| - N_S[k] + |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| \\ &\quad - |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| \\ &\leq |\mathcal{S}_S[k]| - N_S[k] + |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k]| \leq |\mathcal{S}_S[k]|. \end{aligned}$$

The proof of relation (3.11) is completed.

APPENDIX D: PROOF OF LEMMA 3.9

We prove the first inequality. The second one can be proved in a similar way. Note that

$$|\mathcal{S}_R[k]| = \sum_{k_R, k_B: k_R + k_B = k} |\mathcal{S}_R[k]| \mathbf{1}_{\{N_R[k] = k_R, N_B[k] = k_B\}},$$

and by the definition of $\mathcal{N}_{k,h}$ we have

$$|\mathcal{S}_R[k]| \mathbf{1}_{\mathcal{N}_{k,h}} = \sum_{k_R, k_B: k_R + k_B = k, k_R \geq k-h, k_B \leq h} |\mathcal{S}_R[k]| \mathbf{1}_{\{N_R[k] = k_R, N_B[k] = k_B\}}.$$

For $a \geq 0$, we then have

$$\begin{aligned} &\mathbb{P}(|\mathcal{S}_R[k]| > a \mid \mathcal{N}_{k,h}) \mathbb{P}(\mathcal{N}_{k,h}) = \mathbb{P}(|\mathcal{S}_R[k]| \mathbf{1}_{\mathcal{N}_{k,h}} > a) \\ &= \mathbb{P} \left(\sum_{\substack{k_R, k_B \\ k_R + k_B = k, k_R \geq k-h, k_B \leq h}} |\mathcal{S}_R[k]| \mathbf{1}_{\{N_R[k] = k_R, N_B[k] = k_B\}} > a \right) \\ &= \sum_{\substack{k_R, k_B \\ k_R + k_B = k, k_R \geq k-h, k_B \leq h}} \mathbb{P}(|\mathcal{S}_R[k]| > a, N_R[k] = k_R, N_B[k] = k_B) \\ &\geq \sum_{\substack{k_R, k_B \\ k_R + k_B = k, k_R \geq k-h, k_B \leq h}} \mathbb{P}(\text{Bin}(n_W, \pi_R(k-h, h)) > a) \mathbb{P}(N_R[k] = k_R, N_B[k] = k_B) \\ &= \mathbb{P}(\text{Bin}(n_W, \pi_R(k-h, h)) > a) \mathbb{P}(\mathcal{N}_{k,h}), \end{aligned}$$

where the second equality is a consequence of the fact that for different values of (k_R, k_B) the events $\{N_R[k] = k_R, N_B[k] = k_B\}$ are disjoint and the inequality follows by (3.14) and the stochastic ordering properties of the binomial distribution which imply

$$|\mathcal{S}_R[k]| \mathbf{1}_{\{N_R[k] = k_R, N_B[k] = k_B\}} \stackrel{L}{=} \text{Bin}(n_W, \pi_R(k_R, k_B)) \geq_{st} \text{Bin}(n_W, \pi_R(k-h, h)).$$

□

APPENDIX E: PROOF OF PROPOSITION 7.1

Proposition 7.1 is an immediate consequence of the Borel-Cantelli lemma and the following propositions E.1 and E.2. Hereafter, when we write “for any $\kappa > 0$ ”, we implicitly assume that κ is arbitrarily chosen in $(0, z_R + z_B)$ in case $q = g$ and $\alpha_B < \alpha_R < 1$.

PROPOSITION E.1. Given η in (4.8), for any $\kappa > 0$ and $\delta > 0$, there exists a positive constant $c(\kappa, \delta) > 0$ such that

$$\max\{\mathbb{P}(\Upsilon_S(\kappa) > \delta), \mathbb{P}(\bar{\Upsilon}_S(\kappa) > \delta\eta q)\} \ll e^{-c(\kappa, \delta)\eta q}.$$

PROPOSITION E.2. For any $\kappa > 0$ and $\delta > 0$, there exists a positive constant $c(\kappa, \delta) > 0$ such that

$$(E.1) \quad \mathbb{P}(\mathfrak{N}_S(\kappa) > \delta) \ll e^{-c(\kappa, \delta)q}.$$

Proof of Proposition E.1: Auxiliary results. The proof of Proposition E.1 uses the following lemmas.

LEMMA E.3. There exists a positive constant $c(\kappa, \delta) > 0$ such that

$$\max\left\{\sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(\bar{Y}_S(\mathbf{k}) > \delta\eta q), \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(Y_S(\mathbf{k}) > \delta)\right\} \ll e^{-c(\kappa, \delta)\eta q}.$$

In turn the following lemmas E.4, E.5 and E.6 will be exploited to prove Lemma E.3. Hereafter, we set $\mathbf{k} := (k_R, k_B) \in (\mathbb{N} \cup \{0\})^2$. Recalling the definition of $\pi_S(\mathbf{k})$ given in Sect. 3.4.1, we have:

LEMMA E.4. (i) If $q = g$, then, for any $\kappa > 0$,

$$\sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \pi_S(\mathbf{k})}{(g\beta_S(k_S/g) + k_S/g)g} - 1 \right| \rightarrow 0.$$

(ii) If $g \ll q \ll n$, then, for any $\kappa > 0$,

$$\sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \pi_S(\mathbf{k})}{\eta q \beta_S(k_S/q)} - 1 \right| \rightarrow 0.$$

Hereafter, we set

$$\tilde{\pi}_S(\mathbf{k}) := \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) \geq 1), \quad \mathbf{k} := (k_R, k_B) \in (\mathbb{N} \cup \{0\})^2.$$

LEMMA E.5. Assume $q = g$. Then, for any $\kappa > 0$,

$$\sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \tilde{\pi}_S(\mathbf{k})}{r^{-1}[(1 - r^{-1})]^{r-1} (k_{S^c}/q + \alpha_{S^c})(k_S/q + \alpha_S) r q^2 p} - 1 \right| \rightarrow 0.$$

LEMMA E.6. Let $\{X_n\}_{n \in \mathbb{N}}$ and $\{X'_n\}_{n \in \mathbb{N}}$ be two sequences of non-negative random variables defined on the same probability space and such that $\mathbb{P}(X'_n \geq X_n) = 1$ for any $n \in \mathbb{N}$. Let $\mu_n \geq 0$ and $\mu'_n > 0$, $n \in \mathbb{N}$, be two deterministic sequences with $\inf \mu'_n = \mu > 0$. Then

$$\mathbb{P}\left(\left|\frac{X_n}{X'_n} - \frac{\mu_n}{\mu'_n}\right| > \varepsilon\right) \leq \mathbb{P}(|X_n - \mu_n| > \varepsilon\mu/4) + \mathbb{P}(|X'_n - \mu'_n| > \varepsilon\mu/4), \quad \forall \varepsilon \in (0, 1) \text{ and } n \in \mathbb{N}.$$

For the sake of clarity, we point out that, by the convention $0/0 := 1/2$ and the assumption $\mathbb{P}(X'_n \geq X_n) = 1$, for a Borel set $B \subseteq [0, \infty)$,

$$\left\{ \left| \frac{X_n}{X'_n} - \frac{\mu_n}{\mu'_n} \right| \in B \right\} = \left\{ \left| \frac{X_n}{X'_n} - \frac{\mu_n}{\mu'_n} \right| \in B, X'_n \neq 0 \right\} \cup \left\{ \left| \frac{1}{2} - \frac{\mu_n}{\mu'_n} \right| \in B, X'_n = 0 \right\}.$$

Here we prove Proposition E.1 and Lemma E.3. For the other lemmas, whose proof is rather elementary and tedious, we refer the reader to Appendix L.

Proof of Proposition E.1. We only prove Part (i). The proofs of Part (ii) and Part (iii) are (“mutatis mutandis”) analogous. By the union bound, for any $\kappa, \delta > 0$ we have

$$\mathbb{P}(\Upsilon_S(\kappa) > \delta) \leq \sum_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(Y_S(\mathbf{k}) > \delta) \leq |\mathbb{T}(\kappa)| \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(Y_S(\mathbf{k}) > \delta) \leq (\kappa q)^2 \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(Y_S(\mathbf{k}) > \delta)$$

$$\text{and} \quad \mathbb{P}(\bar{\Upsilon}_S(\kappa) > \delta q) \leq (\kappa q)^2 \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(\bar{Y}_S(\mathbf{k}) > \delta q).$$

The claim follows by Lemma E.3(i). \square

Proof of Lemma E.3. We prove only Part (i); Parts (ii), (iii) can be proved along similar lines and, for completeness, a sketch is reported in Appendix L. For $\kappa > 0$, define

$$(E.2) \quad \beta_{\min}(\kappa) := \min_{(x_R, x_B) \in \mathbb{T}'(\kappa)} (|\beta_R(x_R)| + |\beta_B(x_B)|),$$

where $\mathbb{T}'(\kappa)$ is defined in (7.4). Throughout this proof, for arbitrarily fixed $\kappa > 0$ and $\delta \in (0, 1)$ small enough, we denote with $n_{\kappa, \delta}$ a threshold value for n (depending on κ and δ) above which a given property holds. Note that $n_{\kappa, \delta}$ may vary from line to line.

We divide the proof of Part (i) in two sub-parts, where we show that there exists $c'(\kappa, \delta) > 0$ and $c''(\kappa, \delta) > 0$ (not depending on n) such that

$$(E.3) \quad (i) \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(\bar{Y}_S(\mathbf{k}) > \delta q) \ll e^{-c'(\kappa, \delta)q} \quad \text{and} \quad (ii) \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(Y_S(\mathbf{k}) > \delta) \ll e^{-c''(\kappa, \delta)q}.$$

The claim then follows by setting $c(\kappa, \delta) := \min\{c'(\kappa, \delta), c''(\kappa, \delta)\}$.

Proof of (E.3)-(i). By (3.9) we have

$$(E.4) \quad \begin{aligned} & \mathbb{P}(|Q_{k+1}^S - \beta_S(k_S/q)q| > \delta q \mid \mathbf{N}[k] = \mathbf{k}) \\ & \leq \mathbb{P}(|\mathcal{S}_S[k]| - k_S - \beta_S(k_S/q)q| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \\ & \quad + \mathbb{P}(|(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \\ & \quad + \mathbb{P}(|\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}). \end{aligned}$$

We proceed dividing the proof of (E.3)-(i) in two steps, where we give asymptotic exponential bounds (uniformly on $\mathbf{k} \in \mathbb{T}(\kappa)$) for the three terms in the right-hand side of (E.4).

Step 1: upper bound for the first addend in (E.4).

We prove that there exists $n_{\kappa, \delta}$ such that, for all $n \geq n_{\kappa, \delta}$,

$$(E.5) \quad \mathbb{P}(|\mathcal{S}_S[k]| - k_S - \beta_S(k_S/q)q| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \leq 2e^{-c_1(\kappa, \delta)q}, \quad \forall \mathbf{k} \in \mathbb{Z}(\kappa)$$

where $c_1(\kappa, \delta) > 0$ is a suitable positive constant (not depending on n). By (3.14) we have

$$\begin{aligned}
& \mathbb{P}(|\mathcal{S}_S[k]| - k_S - \beta_S(k_S/q)q| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \\
& \leq \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \leq (\beta_S(k_S/q) + k_S/q - \delta/3)q) \\
\text{(E.6)} \quad & + \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \geq (\beta_S(k_S/q) + k_S/q + \delta/3)q).
\end{aligned}$$

Taking $\delta' \in \left(0, \frac{\delta}{r^{-1}(1-r^{-1})^{r-1}(\kappa+\alpha_S)^r}\right)$, and using Lemma E.4, we can conclude that there exists $n_{\kappa, \delta} \geq 1$ such that for any $n \geq n_{\kappa, \delta}$ and for any $\mathbf{k} \in \mathbb{T}(\kappa)$:

$$\text{(E.7)} \quad (\beta_S(k_S/q) + k_S/q)q(1 - \delta'/3) < n_W \pi_S(\mathbf{k}) < (\beta_S(k_S/q) + k_S/q)q(1 + \delta'/3)$$

where by construction for any $n \geq n_{\kappa, \delta}$ and for any $k \in \mathbb{T}(\kappa)$

$$\begin{aligned}
\text{(E.8)} \quad & (\beta_S(k_S/q) + k_S/q)q(1 - \delta'/3) > (\beta_S(k_S/q) + k_S/q - \delta/3)q, \\
& \beta_S(k_S/q) + k_S/q)q(1 + \delta'/3) < (\beta_S(k_S/q) + k_S/q + \delta/3)q,
\end{aligned}$$

By (E.8), the usual concentration bound for the binomial distribution (see formula (I.2) in Appendix I) and the fact that the function H defined in (3.4) decreases on $[0, 1)$, for any $n \geq n_{\kappa, \delta}$, we have, uniformly in $\mathbf{k} \in \mathbb{T}(\kappa)$,

$$\begin{aligned}
\text{(E.9)} \quad & \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \leq (\beta_S(k_S/q) + k_S/q - \delta/3)q) \leq e^{-n_W \pi_S(\mathbf{k}) H\left(\frac{(\beta_S(k_S/q) + k_S/q - \delta/3)q}{n_W \pi_S(\mathbf{k})}\right)} \\
& \leq e^{-[r^{-1}(1-r^{-1})^{r-1} \alpha_S^r - \delta/3] H\left(\frac{1 - \delta/[3r^{-1}(1-r^{-1})^{r-1} \alpha_S^r]}{1 - \delta'/3}\right) q}.
\end{aligned}$$

Similarly, for any $n \geq n_{\kappa, \delta}$, uniformly on $\mathbf{k} \in \mathbb{T}(\kappa)$, we have:

$$\begin{aligned}
\text{(E.10)} \quad & \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \geq (\beta_S(k_S/q) + k_S/q + \delta/3)q) \\
& \leq e^{-[r^{-1}(1-r^{-1})^{r-1} \alpha_S^r - \delta/3] H\left(\frac{1 + \delta/[3r^{-1}(1-r^{-1})^{r-1}(\kappa+\alpha_S)^r]}{1 + \delta'/3}\right) q}.
\end{aligned}$$

Inequality (E.5) follows from (E.6), (E.9) and (E.10).

Step 2: upper bound for the second addend in (E.4). We prove that there exists $n_{\kappa, \delta}$ such that for all $n \geq n_{\kappa, \delta}$, and for all $\mathbf{k} \in \mathbb{T}(\kappa)$:

$$\text{(E.11)} \quad \mathbb{P}(|(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \leq e^{-c_2(\kappa, \delta)q},$$

where $c_2(\kappa, \delta) > 0$ is a suitable positive constant (not depending on n). Similarly, one can show that, for all $n \geq n_{\kappa, \delta}$ and for all $\mathbf{k} \in \mathbb{T}(\kappa)$:

$$\text{(E.12)} \quad \mathbb{P}(|\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \leq e^{-c_3(\kappa, \delta)q},$$

Inequality (E.3) easily follows from (E.4), (E.5), (E.11) and (E.12). To prove (E.11) note that

$$\begin{aligned}
& |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| \\
& \leq \sum_{v \in \mathcal{V}_W} \mathbf{1}\{D_S^{(v)}[k] - D_{S^c}^{(v)}[k] \leq r - 1, D_S^{(v)}[k] \geq r\} \leq \sum_{v \in \mathcal{V}_W} \mathbf{1}\{D_S^{(v)}[k] \geq r, D_{S^c}^{(v)}[k] \geq 1\}.
\end{aligned}$$

By construction it follows

$$\sum_{v \in \mathcal{V}_W} \mathbf{1}\{D_S^{(v)}[k] \geq r, D_{S^c}^{(v)}[k] \geq 1\} \Big| \{\mathbf{N}[k] = \mathbf{k}\} \stackrel{\text{L}}{=} \text{Bin}(n_W, \tilde{\pi}_S(\mathbf{k})), \quad \text{and}$$

$$\text{(E.13)} \quad \mathbb{P}(|\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| > (\delta q)/3 \mid \mathbf{N}[k] = \mathbf{k}) \leq \mathbb{P}(\text{Bin}(n_W, \tilde{\pi}_S(\mathbf{k})) > (\delta q)/3).$$

By Lemma E.5 there exists $n_{\kappa,\delta} \geq 1$ and constants $b_1, b_2 > 0$, such that, for any $n \geq n_{\kappa,\delta}$,

$$(1 - \delta)b_1q^2p < n_W \tilde{\pi}_S(\mathbf{k}) < (1 + \delta)b_2q^2p, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa).$$

By this relation, the concentration bound for the binomial distribution (see (I.1)) and the fact that function H increases on $(1, +\infty)$, for all $n \geq n_{\kappa,\delta}$, we have, uniformly on $\mathbf{k} \in \mathbb{T}(\kappa)$,

$$\begin{aligned} \mathbb{P}(\text{Bin}(n_W, \tilde{\pi}_S(\mathbf{k})) > (\delta q)/3) &\leq e^{-n_W \tilde{\pi}_S(\mathbf{k}) H\left(\frac{(\delta q)/3}{n_W \tilde{\pi}_S(\mathbf{k})}\right)} \\ (E.14) \quad &\leq e^{-(1-\delta)b_1qpH\left(\frac{\delta/3}{(1+\delta)b_2qp}\right)q} \leq e^{-c_2(\kappa,\delta)q}, \end{aligned}$$

for some positive constant $c_2(\kappa, \delta) > 0$ (not depending on n). Inequality (E.11) follows from (E.13) and (E.14).

Proof of (E.3)-(ii). By the previous part of the proof, we have, for all $n \geq n_{\kappa,\delta}$,

$$(E.15) \quad \mathbb{P}(|Q_{k+1}^S - \beta_S(k_S/q)q| > \delta q \mid \mathbf{N}[k] = \mathbf{k}) \leq \delta e^{-\tilde{c}(\kappa,\delta)q}, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa)$$

for some positive constant $\tilde{c}(\kappa, \delta) > 0$ (not depending on n). By the inequality $\|x - y\| \leq |x - y|$, $x, y \in \mathbb{R}$, it follows

$$(E.16) \quad \mathbb{P}(\|Q_{k+1}^S - |\beta_S(k_S/q)q|\| > \delta q \mid \mathbf{N}[k] = \mathbf{k}) \leq \delta e^{-\tilde{c}(\kappa,\delta)q}, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa).$$

By the triangular inequality and the union bound, we have

$$\begin{aligned} &\mathbb{P}(\|Q_{k+1}^S + |Q_{k+1}^{S^c} - (|\beta_S(k_S/q)q| + |\beta_{S^c}(k_{S^c}/q)q)\| > \delta q \mid \mathbf{N}[k] = \mathbf{k}) \\ &\leq \mathbb{P}(\|Q_{k+1}^S - |\beta_S(k_S/q)q|\| > (\delta q)/2 \mid \mathbf{N}[k] = \mathbf{k}) + \\ &\quad \mathbb{P}(\|Q_{k+1}^{S^c} - |\beta_{S^c}(k_{S^c}/q)q|\| > (\delta q)/2 \mid \mathbf{N}[k] = \mathbf{k}). \end{aligned}$$

Combining this relation with (E.16), for all $n \geq n_{\kappa,\delta}$, and for all $\mathbf{k} \in \mathbb{T}(\kappa)$ we have:

$$(E.17) \quad \mathbb{P}(\|Q_{k+1}^S + |Q_{k+1}^{S^c} - (|\beta_S(k_S/q)q| + |\beta_{S^c}(k_{S^c}/q)q)\| > \delta q \mid \mathbf{N}[k] = \mathbf{k}) \leq 2\delta e^{-c_4(\kappa,\delta)q},$$

for some positive constant $c_4(\kappa, \delta) > 0$ (not depending on n). By Lemma E.6, (E.15) and (E.17), for all $n \geq n_{\kappa,\delta}$, and for all $\mathbf{k} \in \mathbb{T}(\kappa)$ we have:

$$\mathbb{P}\left(\left|U_{k+1}^S - \frac{|\beta_S(k_S/q)q|}{|\beta_R(k_R/q)q| + |\beta_B(k_B/q)q|}\right| > \delta \mid \mathbf{N}[k] = \mathbf{k}\right) \leq c_5(\kappa, \delta, \beta_{\min})e^{-c_6(\kappa,\delta,\beta_{\min})q},$$

for suitable positive constants $c_5(\kappa, \delta, \beta_{\min})$ and $c_6(\kappa, \delta, \beta_{\min})$ (not depending on n), where constant $\beta_{\min} > 0$ is defined by (E.2). Claim (E.3)-(ii) easily follows by this last inequality.

Proof of Proposition E.2. We shall show later on that the process $\{\hat{N}_S[k]\}_{j \in \mathbb{N}}$ is an $\{\mathcal{H}_k\}_{k \in \mathbb{N}}$ -martingale with increments bounded by 1, i.e., $\hat{N}_S(1) \leq 1$ and $\hat{N}_S(k+1) - \hat{N}_S[k] \leq 1$, \mathbb{P} -a.s., for any $j \in \mathbb{N}$. Therefore, by the union bound and the Azuma inequality (see e.g. Theorem 2.8 p. 33 in [19]), it follows that, for every $\kappa, \delta > 0$,

$$\begin{aligned} \mathbb{P}(\mathfrak{N}_S(\kappa) > \delta) &\leq \sum_{k=1}^{\lfloor \kappa q \rfloor} \mathbb{P}(|\hat{N}_S[k]| > \delta q) \leq \sum_{k=1}^{\lfloor \kappa q \rfloor} \mathbb{P}\left(\left|\sum_{i=1}^k (\hat{N}_S[i] - \hat{N}_S[k-1])\right| > \delta q\right) \\ &\leq 2\kappa q \exp\left(-\frac{\delta^2 q^2}{2\lfloor \kappa q \rfloor}\right) \leq 2\kappa q \exp\left(-\frac{\delta^2 q}{2\kappa}\right). \end{aligned}$$

It remains to prove that the process $\{\hat{N}_S[k]\}_{k \in \mathbb{N}}$ is an $\{\mathcal{H}_k\}_{k \in \mathbb{N}}$ -martingale with increments bounded by 1. For any $j \in \mathbb{N}$, $\hat{N}_S[k]$ is clearly \mathcal{H}_k -measurable, moreover

$$\hat{N}_S[k+1] - \hat{N}_S[k] = N_S[k+1] - N_S[k] - U_{k+1}^S \mathbf{1}_{\{k < n_w\}} = M_{k+1}^S - U_{k+1}^S \mathbf{1}_{\{k < n_w\}} \quad \mathbb{P}\text{-a.s.}$$

Note that the second equality follows by (3.6), therefore, by Proposition 3.8, we have

$$(E.18) \quad \mathbb{E}[\hat{N}_S(k+1) | \mathcal{H}_k] - \hat{N}_S[k] = 0,$$

i.e., $\{\hat{N}_S[k]\}_{j \in \mathbb{N}}$ is an $\{\mathcal{H}_k\}$ -martingale. Moreover, $|M_{k+1}^S - U_{k+1}^S| < 1$.

APPENDIX F: PROOF OF PROPOSITION 7.2

Given $i \in \mathbb{N}$, $\mathbf{k} = (k_R, k_B) \in (\mathbb{N} \cup \{0\})^2$ with $k = k_R + k_B$, observe that by construction $\mathbf{N}[k+i] - \mathbf{N}[k]$ take values on \mathbb{I}_i (defined in (7.1)). Hence:

$$(F.1) \quad \sum_{\mathbf{i} \in \mathbb{I}_i} \mathbf{1}_{\mathcal{E}_i(k,i)} = 1 \quad \text{with} \quad \mathcal{E}_i(k,i) := \{\omega \in \Omega : \mathbf{N}[k+i] - \mathbf{N}[k] = \mathbf{i}\}, \quad \mathbf{i} \in \mathbb{I}_i \quad \text{and}$$

$$(F.2) \quad \mathbf{1}_{\mathcal{E}_i(k,i)} \mathbf{1}\{\mathbf{N}[k] = \mathbf{k}\} = \mathbf{1}_{\mathcal{E}_i(k,i)} \mathbf{1}\{\mathbf{N}[k+i] = \mathbf{k} + \mathbf{i}\}, \quad \text{for any } \mathbf{i} \in \mathbb{I}_i.$$

So, for any $z > 0$, recalling the definition of $J_S[k]$ in (7.3), we have

$$\begin{aligned} (J_S[k + \lfloor zq \rfloor] - J_S[k]) \mathbf{1}\{\mathbf{N}[k] = \mathbf{k}\} &= \left[\sum_{i=0}^{\lfloor zq \rfloor - 1} (J_S[k+i+1] - J_S[k+i]) \right] \mathbf{1}\{\mathbf{N}[k] = \mathbf{k}\} \\ &= \sum_{i=0}^{\lfloor zq \rfloor - 1} \sum_{\mathbf{i} \in \mathbb{I}_i} (J_S[k+i+1] - J_S[k+i]) \mathbf{1}_{\mathcal{E}_i(k,i)} \mathbf{1}\{\mathbf{N}[k] = \mathbf{k}\} \\ &= \sum_{i=0}^{\lfloor zq \rfloor - 1} \sum_{\mathbf{i} \in \mathbb{I}_i} U_{k+i+1}^S \mathbf{1}_{\mathcal{E}_i(k,i)} \mathbf{1}\{\mathbf{N}[k] = \mathbf{k}\} = \sum_{i=0}^{\lfloor zq \rfloor - 1} \sum_{\mathbf{i} \in \mathbb{I}_i} U_{k+i+1}^S \mathbf{1}_{\mathcal{E}_i(k,i)} \mathbf{1}\{\mathbf{N}[k+i] = \mathbf{k} + \mathbf{i}\}. \end{aligned}$$

Therefore, for any $y, z > 0$,

$$\begin{aligned} (F.3) \quad J_S[\lfloor yq \rfloor + \lfloor zq \rfloor] - J_S[\lfloor yq \rfloor] &= \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} (J_S[\lfloor yq \rfloor + \lfloor zq \rfloor] - J_S[\lfloor yq \rfloor]) \mathbf{1}\{\mathbf{N}[\lfloor yq \rfloor] = \mathbf{k}\} \\ &= \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \sum_{i=0}^{\lfloor zq \rfloor - 1} \sum_{\mathbf{i} \in \mathbb{I}_i} U_{\lfloor yq \rfloor + i + 1}^S \mathbf{1}_{\mathcal{E}_i(k,i)} \mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}. \end{aligned}$$

Now fix $\kappa < \kappa_f$ and assume $y + 2z \leq \kappa$, for any $\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}$ and $i \in \{1, \dots, \lfloor zq \rfloor - 1\}$, we have

$$k_R + k_B + i = \lfloor yq \rfloor + i \leq (y+z)q \leq (\kappa - z)q.$$

Therefore, for any $\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}$, $i = 1, \dots, \lfloor zq \rfloor - 1$ and $\mathbf{i} \in \mathbb{I}_i$, we have $\mathbf{k} + \mathbf{i} \in \mathbb{T}(\kappa)$. So, by the definition of Ω_κ in (7.10), for all $\omega \in \Omega_\kappa$ and any $\varepsilon > 0$ there exists $n(\omega, \varepsilon)$ such that for all $n \geq n(\omega, \varepsilon)$, we have

$$\mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}(\omega) \left| U_{\lfloor yq \rfloor + i + 1}^S(\omega) - \frac{|\beta_S((k_S + i_S)/q)|}{|\beta_R((k_R + i_R)/q)| + |\beta_B((k_B + i_B)/q)|} \right| < \varepsilon.$$

as long as $\mathbf{k} \in \mathbb{T}(\kappa)$. By this inequality, the fact that $q^{-1}(k_R + i_R, k_B + i_B) \in \mathbb{L}_{\mathbf{k}/q}(\kappa, z)$, (with $\mathbb{L}_{\mathbf{k}/q}(\kappa, z)$ defined in (7.5)) the definition of $\bar{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)}$ and $\underline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)}$ (in (7.6) or

(7.7) and (7.8)) and the fact that $0 \leq U_{\lfloor yq \rfloor + i + 1} \leq 1$, it follows that, for all $\omega \in \Omega_\kappa$ and any $\varepsilon > 0$, there exists $n(\omega, \varepsilon)$ such that, for all $n \geq n(\omega, \varepsilon)$,

$$\begin{aligned} \mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}(\omega)(\underline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} - \varepsilon) &\leq \mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}(\omega)U_{\lfloor yq \rfloor + i + 1}^S(\omega) \\ &\leq \mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}(\omega)(\overline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} + \varepsilon). \end{aligned}$$

Combining this with (F.3), we have that, for all $\omega \in \Omega_\kappa$ and any $\varepsilon > 0$, there exists $n(\omega, \varepsilon)$ such that for all $n \geq n(\omega, \varepsilon)$,

$$\begin{aligned} \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \sum_{i=0}^{\lfloor zq \rfloor - 1} \sum_{\mathbf{i} \in \mathbb{I}_i} \mathbf{1}_{\varepsilon_i(k, i)}(\omega) \mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}(\omega) (\underline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} - \varepsilon) \\ \leq J_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - J_S[\lfloor yq \rfloor](\omega) \\ \leq \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \sum_{i=0}^{\lfloor zq \rfloor - 1} \sum_{\mathbf{i} \in \mathbb{I}_i} \mathbf{1}_{\varepsilon_i(k, i)}(\omega) \mathbf{1}\{\mathbf{N}(\lfloor yq \rfloor + i) = \mathbf{k} + \mathbf{i}\}(\omega) (\overline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} + \varepsilon) \end{aligned}$$

i.e. (using (F.1) and (F.2)),

$$\begin{aligned} \lfloor zq \rfloor \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \mathbf{1}\{\mathbf{N}[\lfloor yq \rfloor] = \mathbf{k}\}(\omega) (\underline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} - \varepsilon) &\leq J_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - J_S[\lfloor yq \rfloor](\omega) \\ \text{(F.4)} \quad &\leq \lfloor zq \rfloor \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \mathbf{1}\{\mathbf{N}[\lfloor yq \rfloor] = \mathbf{k}\}(\omega) (\overline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} + \varepsilon). \end{aligned}$$

We note that, for any $\omega \in \Omega_\kappa$,

$$\begin{aligned} N_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - N_S[\lfloor yq \rfloor](\omega) \\ = J_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - J_S[\lfloor yq \rfloor](\omega) + \hat{N}_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - \hat{N}_S[\lfloor yq \rfloor](\omega). \end{aligned}$$

Since $\lfloor yq \rfloor \leq \lfloor yq \rfloor + 2\lfloor zq \rfloor \leq \kappa q$, by the definition of Ω_κ (in (7.10)), $\mathfrak{N}_S(\kappa)$ (in (7.2)) and $\hat{N}_S[k]$ (in (7.3)), we have that, for any $\omega \in \Omega_\kappa$ and any $\varepsilon > 0$, there exists $n'(\omega, \varepsilon)$ such that for any $n \geq n'(\omega, \varepsilon)$, we have

$$-\varepsilon q < \hat{N}_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - \hat{N}_S[\lfloor yq \rfloor](\omega) < \varepsilon q$$

and so

$$\begin{aligned} -\varepsilon q + J_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - J_S[\lfloor yq \rfloor](\omega) &< N_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - N_S[\lfloor yq \rfloor](\omega) \\ &< \varepsilon q + J_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - J_S[\lfloor yq \rfloor](\omega). \end{aligned}$$

Combining this inequality with (F.4), we have that, for all $\omega \in \Omega_\kappa$ and any $\varepsilon > 0$, there exists $n''(\omega, \varepsilon)$ such that, for all $n \geq n''(\omega, \varepsilon)$,

$$\begin{aligned} -\varepsilon q + \lfloor zq \rfloor \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \mathbf{1}\{\mathbf{N}[\lfloor yq \rfloor](\omega) = \mathbf{k}\} (\underline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} - \varepsilon) \\ < N_S[\lfloor yq \rfloor + \lfloor zq \rfloor](\omega) - N_S[\lfloor yq \rfloor](\omega) \\ < \varepsilon q + \lfloor zq \rfloor \sum_{\mathbf{k} \in \mathbb{I}_{\lfloor yq \rfloor}} \mathbf{1}\{\mathbf{N}[\lfloor yq \rfloor](\omega) = \mathbf{k}\} (\overline{\beta}_{S, \mathbb{L}_{\mathbf{k}/q}(\kappa, z)} + \varepsilon). \end{aligned}$$

The claim follows dividing this relation by q , then taking the limsup and the liminf as $n \rightarrow \infty$ and finally letting ε tend to zero. We recall that by construction $0 \leq \tilde{N}_S(xq) - N_S[\lfloor xq \rfloor] \leq 1$, for any $x > 0$ as consequence of definition (4.1). \square

APPENDIX G: PROOF OF LEMMA 7.3

We divide the proof in two steps. In the first step we prove the lemma assuming $a_{R,1} = a_{R,2}$. In the second step we consider the general case.

Step 1: the case $a_{R,1} = a_{R,2}$. Let $\mathcal{V}_{S,h}$, $S \in \{R, B\}$, $h \in \{1, 2\}$, denote the set of S -seeds for process h . Note that $|\mathcal{V}_{S,h}| = a_{S,h}$. Since $a_{R,1} = a_{R,2}$ and $a_{B,1} \geq a_{B,2}$, without loss of generality we assume that $\mathcal{V}_{R,1} \equiv \mathcal{V}_{R,2}$ and $\mathcal{V}_{B,1} \supseteq \mathcal{V}_{B,2}$. Note that $\mathcal{V}_{W,2} \supseteq \mathcal{V}_{W,1}$ and

$$(G.1) \quad \mathcal{V}_{W,2} \setminus \mathcal{V}_{W,1} = \mathcal{V}_{B,1} \setminus \mathcal{V}_{B,2}.$$

Let $\mathcal{V}_{S,h}(t)$ and $\mathcal{W}_{S,h}(t)$ be, respectively, the random subset of $\mathcal{V}_{W,1}$ and $\mathcal{V}_{W,2}$, defined on Ω , formed by S -active nodes at time t , for process h . We denote by $\mathcal{V}_{S,h}(\infty)$ and $\mathcal{W}_{S,h}(\infty)$, respectively, the random subset of $\mathcal{V}_{W,1}$ and $\mathcal{V}_{W,2}$, defined on Ω , formed by S -active nodes when process h stops. We shall show later on that

$$(G.2) \quad \mathcal{V}_{R,1}(\infty) \subseteq_{st} \mathcal{V}_{R,2}(\infty) \quad \text{and} \quad \mathcal{V}_{B,2}(\infty) \subseteq_{st} \mathcal{V}_{B,1}(\infty);$$

The claim then immediately follows from noticing that $|\mathcal{V}_{S,1}(\infty)| = N_{S,1}([0, \infty) \times \mathcal{V}_{W,1})$ and $|\mathcal{W}_{S,2}(\infty)| = N_{S,2}([0, \infty) \times \mathcal{V}_{W,2})$, $S \in \{R, B\}$, $h \in \{1, 2\}$. For instance, for what concerns the B -active nodes, by (G.2) (and the definition of \subseteq_{st}) we have that there exists a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ and two random subsets defined on it, say $\tilde{\mathcal{V}}_{B,h}(\infty)$, $h \in \{1, 2\}$, such that

$$\tilde{\mathcal{V}}_{B,h}(\infty) \stackrel{L}{=} \mathcal{V}_{B,h}(\infty) \quad \text{and} \quad \tilde{\mathcal{V}}_{B,2}(\infty) \subseteq \tilde{\mathcal{V}}_{B,1}(\infty), \quad \tilde{\mathbb{P}}\text{-a.s.}$$

From which, using an obvious notation,

$$\begin{aligned} \tilde{A}_{B,1}^* &= |\tilde{\mathcal{V}}_{B,1}(\infty)| + a_{B,1} \geq |\tilde{\mathcal{V}}_{B,2}(\infty)| + a_{B,1} \\ &= |\tilde{\mathcal{V}}_{B,2}(\infty)| + |\mathcal{V}_{W,2} \setminus \mathcal{V}_{W,1}| + a_{B,2} \geq |\tilde{\mathcal{W}}_{B,2}(\infty)| + a_{B,2} = \tilde{A}_{B,2}^*, \end{aligned}$$

where the second equation follows from (G.1). Last inequality instead descends from the fact that by construction $\tilde{\mathcal{W}}_{B,2}(\infty) \subseteq \tilde{\mathcal{V}}_{B,2}(\infty) \cup (\mathcal{V}_{W,2} \setminus \mathcal{V}_{W,1})$. Now to prove (G.2), first observe that $N^{(h)} := \sum_{v \in \mathcal{V}_{W,h}} N'_v$, and $\mathcal{V}_{W,1} \subseteq \mathcal{V}_{W,2}$ immediately imply that $N^{(1)} \subseteq_{st} N^{(2)}$.

Therefore, there exists a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ and two point processes $\tilde{N}^{(h)} = \{(\tilde{T}_k^{(h)}, \tilde{V}_k^{(h)})\}_{k \in \mathbb{N}}$, $h \in \{1, 2\}$, defined on this space, such that $\tilde{N}^{(h)} \stackrel{L}{=} N^{(h)}$, $h \in \{1, 2\}$, and $\tilde{N}^{(1)} \subseteq \tilde{N}^{(2)} = \tilde{N}^{(1)} \cup (\tilde{N}^{(2)} \setminus \tilde{N}^{(1)})$, $\tilde{\mathbb{P}}$ -almost surely. Hereon, we consider a copy of the competing bootstrap percolation process h defined on $\tilde{\Omega}$, and we denote by $\tilde{\mathcal{V}}_{S,h}(t)$ and $\tilde{\mathcal{W}}_{S,h}(t)$ the random subset of $\mathcal{V}_{W,1}$ and $\mathcal{V}_{W,2}$, defined on $\tilde{\Omega}$, formed by S -active nodes at time t , for the competing bootstrap percolation process h . Note that, by construction, for an arbitrarily fixed $k \in \mathbb{N}$, the set $\tilde{\mathcal{V}}_{S,h}(t)$ is constant for $\tilde{T}_k^{(1)} \leq t < \tilde{T}_{k+1}^{(1)}$, and may increase (with respect to the set inclusion), by the addition of a new S node, at time $t = \tilde{T}_{k+1}^{(1)}$. The relations (G.2) follow if we prove that, for any $k \in \mathbb{N}$,

$$(G.3) \quad \tilde{\mathcal{V}}_{R,1}(\tilde{T}_k^{(1)}) \subseteq \tilde{\mathcal{V}}_{R,2}(\tilde{T}_k^{(1)}), \quad \tilde{\mathcal{V}}_{B,2}(\tilde{T}_k^{(1)}) \subseteq \tilde{\mathcal{V}}_{B,1}(\tilde{T}_k^{(1)}), \quad \tilde{\mathbb{P}}\text{-a.s.}$$

Indeed by construction, for $S \in \{R, B\}$ and $h \in \{1, 2\}$,

$$(G.4) \quad \tilde{\mathcal{V}}_{S,h}(\infty) = \bigcup_{k \in \mathbb{N}} \tilde{\mathcal{V}}_{S,h}(\tilde{T}_k^{(1)}), \quad \tilde{\mathbb{P}}\text{-a.s.}$$

and $\tilde{\mathcal{V}}_{S,h}(\infty)$ has the same law of $\mathcal{V}_{S,h}(\infty)$.

We prove the relations (G.3) by induction on $k \geq 1$. Note that these relations are trivially true for $k = 1$, indeed $\tilde{\mathcal{V}}_{S,h}(\tilde{T}_1^{(1)}) = \emptyset$ for $h \in \{1, 2\}$ and $S \in \{R, B\}$. We assume (G.3) for

$k = j$ and prove these relations for $k = j + 1$, where j is arbitrarily chosen in \mathbb{N} . By the inductive hypothesis we have, $\tilde{\mathbb{P}}$ -almost surely,

$$\begin{aligned} \tilde{N}_{R,2}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,2}) &\geq \tilde{N}_{R,2}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}) \\ &= |\tilde{\mathcal{V}}_{R,2}(\tilde{T}_j^{(1)})| \geq |\tilde{\mathcal{V}}_{R,1}(\tilde{T}_j^{(1)})| = \tilde{N}_{R,1}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}) \end{aligned}$$

and

$$\tilde{N}_{B,2}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}) = |\tilde{\mathcal{V}}_{B,2}(\tilde{T}_j^{(1)})| \leq |\tilde{\mathcal{V}}_{B,1}(\tilde{T}_j^{(1)})| = \tilde{N}_{B,1}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}).$$

For $v \in \mathcal{V}_{W,2}$, let $\{\tilde{E}_i^{(v)}\}_{i \in \mathbb{N}}$ and $\{\tilde{E}'_i^{(v)}\}_{i \in \mathbb{N}}$ be independent sequences of independent random variables defined on $\tilde{\Omega}$ with the Bernoulli law of mean p , independent of $\tilde{N}^{(2)}$. By the above relations, we have that, for every $v \in \mathcal{V}_{W,1}$, $\tilde{\mathbb{P}}$ -almost surely,

$$\tilde{D}_{R,1}^{(v)}(\tilde{T}_j^{(1)}) := \sum_{i=1}^{\tilde{N}_{R,1}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}) + a_{R,1}} \tilde{E}_i^{(v)} \leq \sum_{i=1}^{\tilde{N}_{R,2}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,2}) + a_{R,2}} \tilde{E}_i^{(v)} =: \tilde{D}_{R,2}^{(v)}(\tilde{T}_j^{(1)})$$

and

$$\begin{aligned} \tilde{D}_{B,2}^{(v)}(\tilde{T}_j^{(1)}) &:= \sum_{i=1}^{\tilde{N}_{B,2}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,2}) + a_{B,2}} \tilde{E}'_i^{(v)} \leq \sum_{i=1}^{\tilde{N}_{B,2}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}) + a_{B,1}} \tilde{E}'_i^{(v)} \\ &\leq \sum_{i=1}^{\tilde{N}_{B,1}([0, \tilde{T}_j^{(1)}] \times \mathcal{V}_{W,1}) + a_{B,1}} \tilde{E}'_i^{(v)} =: \tilde{D}_{B,1}^{(v)}(\tilde{T}_j^{(1)}). \end{aligned}$$

note, indeed that $a_{B,1} = |\mathcal{V}_{W,2} \setminus \mathcal{V}_{W,1}| + a_{B,2}$. Therefore, $\tilde{\mathbb{P}}$ -almost surely, we have

$$\begin{aligned} \tilde{\mathcal{S}}_{R,1}((\tilde{T}_j^{(1)})) &:= \{v \in \mathcal{V}_{W,1} : \tilde{D}_{R,1}^{(v)}(\tilde{T}_j^{(1)}) - \tilde{D}_{B,1}^{(v)}(\tilde{T}_j^{(1)}) \geq r\} \\ &\subseteq \{v \in \mathcal{V}_{W,1} : \tilde{D}_{R,2}^{(v)}(\tilde{T}_j^{(1)}) - \tilde{D}_{B,2}^{(v)}(\tilde{T}_j^{(1)}) \geq r\} \\ &\subseteq \{v \in \mathcal{V}_{W,2} : \tilde{D}_{R,2}^{(v)}(\tilde{T}_j^{(1)}) - \tilde{D}_{B,2}^{(v)}(\tilde{T}_j^{(1)}) \geq r\} \\ \text{(G.5)} \quad &=: \tilde{\mathcal{S}}_{R,2}((\tilde{T}_j^{(1)})). \end{aligned}$$

Now note that, by construction, we have

$$v \in \tilde{\mathcal{V}}_{R,h}((\tilde{T}_{j+1}^{(1)})) \setminus \tilde{\mathcal{V}}_{R,h}((\tilde{T}_j^{(1)})), \quad h \in \{1, 2\}$$

if and only if

$$v \in \tilde{\mathcal{S}}_{R,h}((\tilde{T}_j^{(1)})) \setminus \tilde{\mathcal{V}}_{R,h}((\tilde{T}_j^{(1)})), \quad h \in \{1, 2\}.$$

Therefore, if $v \in \tilde{\mathcal{V}}_{R,1}((\tilde{T}_{j+1}^{(1)})) \setminus \tilde{\mathcal{V}}_{R,1}((\tilde{T}_j^{(1)}))$ then $v \in \tilde{\mathcal{S}}_{R,1}((\tilde{T}_j^{(1)}))$ and so by (G.5) it follows that $v \in \tilde{\mathcal{S}}_{R,2}((\tilde{T}_j^{(1)}))$, from which we have $v \in \tilde{\mathcal{V}}_{R,2}((\tilde{T}_{j+1}^{(1)}))$, and the proof of the first relation of (G.3) is completed (if $v \in \tilde{\mathcal{V}}_{R,1}((\tilde{T}_j^{(1)}))$ the claim immediately follows by the inductive hypothesis). The second relation (G.3) follows along similar lines noticing that

$$\begin{aligned} \tilde{\mathcal{S}}_{B,2}((\tilde{T}_j^{(1)})) \cap \mathcal{V}_{W,1} &= \{v \in \mathcal{V}_{W,1} : \tilde{D}_{B,2}^{(v)}(\tilde{T}_j^{(1)}) - D_{R,2}^{(v)}(\tilde{T}_j^{(1)}) \geq r\} \\ &\subseteq \{v \in \mathcal{V}_{W,1} : \tilde{D}_{B,1}^{(v)}(\tilde{T}_j^{(1)}) - D_{R,1}^{(v)}(\tilde{T}_j^{(1)}) \geq r\} \\ &= \tilde{\mathcal{S}}_{B,1}((\tilde{T}_j^{(1)})). \end{aligned}$$

Step 2: the case $a_{R,1} \leq a_{R,2}$. The general case can be easily obtained by introducing a third activation process with an initial configuration of the seeds given by $(a_{B,3}, a_{R,3}) = (a_{B,1}, a_{R,2})$. Indeed, since $a_{R,3} = a_{R,2}$ and $a_{B,3} = a_{B,1} \geq a_{B,2}$, by the previous step we have

$$(G.6) \quad A_{R,3}^* \leq_{st} A_{R,2}^* \quad \text{and} \quad A_{B,2}^* \leq_{st} A_{B,3}^*.$$

Again, since $a_{B,3} = a_{B,1}$ and $a_{R,3} = a_{R,2} \geq a_{R,1}$, by the previous step (noticing that in the proof the role of R and B can be interchanged) one has

$$(G.7) \quad A_{B,3}^* \leq_{st} A_{B,1}^* \quad \text{and} \quad A_{R,1}^* \leq_{st} A_{R,3}^*.$$

The claim follows combining (G.6) and (G.7). \square

APPENDIX H: PROOF OF (8.1)

By construction we have:

$$\{(T_k'^{\text{stop}}, V_k'^{\text{stop}})\}_k := \{(T_k', V_k')\}_k$$

and

$$\{E_i^{(v), \text{stop}}\}_{i \in \mathbb{N}} = \{E_i^{(v)}\}_{i \in \mathbb{N}}, \quad \{E_i'^{(v), \text{stop}}\}_{i \in \mathbb{N}} = \{E_i'^{(v)}\}_{i \in \mathbb{N}}.$$

Therefore

$$\mathcal{V}_S^{\text{stop}}(t) = \mathcal{V}_S(t), \quad S \in \{R, B\} \quad \text{on the event } \{t \leq Z_{\text{stop}}\}$$

In addition, on the event $\{t > Z_{\text{stop}}\}$, we have:

$$\mathcal{V}_R^{\text{stop}}(t) = \mathcal{V}_R^{\text{stop}}(Z_{\text{stop}}) = \mathcal{V}_R(Z_{\text{stop}}) \subseteq \mathcal{V}_R(t).$$

Therefore

$$(H.1) \quad D_R^{(v), \text{stop}}(T_k') \leq D_R^{(v)}(T_k') \quad \forall k \in \mathbb{N}, v \in \mathcal{V}_W.$$

We proceed proving by induction that

$$(H.2) \quad \mathcal{V}_B(T_k') \subseteq \mathcal{V}_B^{\text{stop}}(T_k') \quad \forall k \in \mathbb{N},$$

The relation (H.2) is clearly true for $k = 0$, indeed $\mathcal{V}_B(T_0') = \mathcal{V}_B^{\text{stop}}(T_0') = \emptyset$ \mathbb{P} -a.s.⁶ Assume that (H.2) is true for any $k \leq k_0$. Then

$$(H.3) \quad D_B^{(v), \text{stop}}(T'_{k_0}) \geq D_B^{(v)}(T'_{k_0}) \quad \forall v \in \mathcal{V}_W.$$

Combining (H.1) and (H.3) we have

$$\mathcal{S}_B(T'_{k_0}) \subseteq \mathcal{S}_B^{\text{stop}}(T'_{k_0})$$

which implies

$$\mathcal{V}_B(T'_{k_0+1}) \subseteq \mathcal{V}_B^{\text{stop}}(T'_{k_0+1}) \quad \forall k \in \mathbb{N}.$$

Indeed there are three cases:

$$i) V'_{k_0} \in \mathcal{V}_B(T'_{k_0}) \quad ii) V'_{k_0} \in \mathcal{V}_B^{\text{stop}}(T'_{k_0}) \setminus \mathcal{V}_B(T'_{k_0}) \quad iii) V'_{k_0} \notin \mathcal{V}_B^{\text{stop}}(T'_{k_0})$$

In the case *i*)

$$\mathcal{V}_B(T'_{k_0+1}) = \mathcal{V}_B(T'_{k_0}) \cup \{V'_{k_0}\} = \mathcal{V}_B(T'_{k_0}) \subseteq \mathcal{V}_B^{\text{stop}}(T'_{k_0}) = \mathcal{V}_B^{\text{stop}}(T'_{k_0}) \cup \{V'_{k_0}\} = \mathcal{V}_B^{\text{stop}}(T'_{k_0+1}).$$

⁶We recall that conventionally $T'_0 = 0$.

where the inclusion follows from the inductive hypothesis. In the case *ii*)

$$\mathcal{V}_B(T'_{k_0+1}) = \mathcal{V}_B(T'_{k_0}) \cup \{V'_{k_0}\} \subseteq \mathcal{V}_B^{\text{stop}}(T'_{k_0}) \cup \{V'_{k_0}\} = \mathcal{V}_B^{\text{stop}}(T'_{k_0}) = \mathcal{V}_B^{\text{stop}}(T'_{k_0+1}).$$

Finally, in the case *iii*)

$$\mathcal{V}_B(T'_{k_0+1}) = \mathcal{V}_B(T'_{k_0}) \cup \{V'_{k_0}\} \subseteq \mathcal{V}_B^{\text{stop}}(T'_{k_0}) \cup \{V'_{k_0}\} = \mathcal{V}_B^{\text{stop}}(T'_{k_0+1}).$$

Then (8.1) immediately follows since:

$$A_B^* = \left| \bigcup_k \mathcal{V}_B(T'_k) \right| + a_B \leq \left| \bigcup_k \mathcal{V}_B^{\text{stop}}(T'_k) \right| + a_B = A_B^{*,\text{stop}}. \quad \square$$

APPENDIX I: CONCENTRATION INEQUALITIES

Hereafter, H denotes the function defined in (3.4).

Let $\mu := mq$, $m \in \mathbb{N}$, $q \in (0, 1)$. For any integer $0 < k < m$, we have: if $k \geq \mu$, then:

$$(I.1) \quad \mathbb{P}(\text{Bin}(m, q) \geq k) \leq \begin{cases} e^{-\mu H\left(\frac{k}{\mu}\right)} & \text{for } k \geq \mu; \\ e^{-\left(\frac{k}{2}\right) \log\left(\frac{k}{\mu}\right)} & \text{for } k \geq e^2 \mu; \end{cases}$$

$$(I.2) \quad \mathbb{P}(\text{Bin}(m, q) \leq k) \leq e^{-\mu H\left(\frac{k}{\mu}\right)} \quad \text{for } k \geq e^2 \mu;$$

For any integer $0 \leq k \leq \lambda$, we have:

$$(I.3) \quad \mathbb{P}(\text{Po}(\lambda) \leq k) \leq e^{-\lambda H\left(\frac{k}{\lambda}\right)};$$

APPENDIX J: CAUCHY PROBLEM 4.2 - COMPLEMENTARY MATERIAL

Proof of Lemma A.1. Let \mathbf{g} be as in the statement and define \mathbf{f} on $[0, z(\kappa_{\mathbf{g}}))$, again as in the statement. Then $\mathbf{f}(0) = \mathbf{g}(0) = \mathbf{0}$ and, for any $x \in (0, z(\kappa_{\mathbf{g}}))$, we have

$$\begin{aligned} \mathbf{f}'(x) &= (z^{-1})'(x) \mathbf{g}'(z^{-1}(x)) = \frac{1}{z'(z^{-1}(x))} \boldsymbol{\beta}(\mathbf{g}(z^{-1}(x))) \\ &= \frac{\boldsymbol{\beta}(\mathbf{g}(z^{-1}(x)))}{\beta_R(g_R(z^{-1}(x))) + \beta_B(g_B(z^{-1}(x)))} \\ &= \frac{\boldsymbol{\beta}(\mathbf{f}(x))}{\beta_R(f_R(x)) + \beta_B(f_B(x))}, \end{aligned}$$

i.e., \mathbf{f} solves (4.2) with $\kappa := z(\kappa_{\mathbf{g}})$. □

Proof of Propositions A.2 and A.3 . Propositions A.2 and A.3 are immediate consequences of Lemmas J.1, J.2 and J.3, whose proof is given below. These lemmas consider one dimensional Cauchy's problems of the form

$$(J.1) \quad g'_S(x) = \beta_S(g_S(x)), \quad x \in (0, \kappa_{\mathbf{g}}), \quad g_S(0) = 0$$

where $\kappa_{\mathbf{g}} \in (0, +\infty]$ and $S \in \{R, B\}$.

LEMMA J.1. Let $q = g$ and assume $\alpha_S < 1$. Then the Cauchy problem (J.1) has a unique solution g_S on $(0, \infty)$ and $g_S(x) \uparrow z_S$, as $x \uparrow +\infty$.

LEMMA J.2. Let $q = g$ and assume $\alpha_S > 1$. Then the Cauchy problem (J.1) has a unique solution g_S on $(0, \kappa_{\mathbf{g}})$, with

$$\kappa_{\mathbf{g}} := \int_0^\infty \frac{dx}{-x + r^{-1}(1 - r^{-1})^{r-1}(\alpha_S + x)^r} < \infty$$

and $g_S(x) \uparrow +\infty$, as $x \uparrow \kappa_{\mathbf{g}}$.

LEMMA J.3. Let $g \ll q \ll p^{-1}$. Then the Cauchy problem (J.1) has a unique solution g_S on $(0, \kappa_{\mathbf{g}})$, with

$$\kappa_{\mathbf{g}} := \frac{r!}{(r-1)\alpha_S^{r-1}},$$

given by

$$g_S(x) = \frac{1}{(\alpha_S^{1-r} - \frac{r-1}{r!}x)^{1/(r-1)}} - \alpha_S.$$

J.1. Proof of Lemma J.2. By Remark 2.3 the function β_S is strictly positive. Moreover, $\lim_{x \rightarrow +\infty} \beta_S(x) = +\infty$, as it can be easily checked by a direct inspection. So equilibrium points do not exist, the unique solution g_S is strictly increasing, and $g'_S(x)$ is bounded away from zero for all x large enough. In particular, this latter property of the solution g_S guarantees that it has not horizontal asymptotes. Therefore there are only two possible cases: *i*) g_S is defined on the whole non-negative half-line $[0, \infty)$ and $g_S(x) \uparrow +\infty$, as $x \uparrow +\infty$; *ii*) g_S is defined on a finite interval of the form $[0, \kappa_{\mathbf{g}})$, for some $\kappa_{\mathbf{g}} \in (0, \infty)$ and $g_S(x) \uparrow +\infty$, as $x \uparrow \kappa_{\mathbf{g}}$. We are going to check that we fall in the case *ii*). Let \mathcal{D}_{g_S} be the domain of g_S . By the differential equation, we have

$$(J.2) \quad \frac{g'_S(x)}{\beta_S(g_S(x))} = 1, \quad \forall x \in \mathcal{D}_{g_S}$$

and so

$$(J.3) \quad \int_{g_S(0)}^{g_S(x)} \frac{1}{\beta_S(u)} du = \int_0^x \frac{g'_S(u)}{\beta_S(g_S(u))} du = \int_0^x du = x, \quad \forall x \in \mathcal{D}_{g_S}.$$

Note that

$$\int_0^\infty \frac{1}{\beta_S(x)} dx = \int_0^\infty \frac{dx}{-x + r^{-1}[(1 - r^{-1})]^{r-1}(\alpha_S + x)^r} = \kappa_{\mathbf{g}} < \infty.$$

Therefore by (J.3) we necessarily have $\mathcal{D}_{g_S} = [0, \kappa_{\mathbf{g}})$ and $g_S(x) \uparrow +\infty$, as $x \uparrow \kappa_{\mathbf{g}}$. \square

Proof of Lemma J.3. In this case we can compute explicitly the solution g_S . Indeed, letting \mathcal{D}_{g_S} denote the domain of g_S , we have

$$\int_0^{g_S(x)} \frac{1}{\beta_S(u)} du = r! \int_0^{g_S(x)} \frac{du}{(u + \alpha_S)^r} = x, \quad \forall x \in \mathcal{D}_{g_S}$$

and

$$\int_0^{g_S(x)} \frac{du}{(u + \alpha_S)^r} = \frac{\alpha_S^{1-r}}{r-1} - \frac{(g_S(x) + \alpha_S)^{1-r}}{r-1}.$$

This easily gives the claim. \square

Finally, we provide the proof of Lemma A.4.

Proof of Lemma A.4. We prove this Lemma by contradiction. First note that

$$f'_R(0) = \frac{\beta(0,0)}{\beta_R(0) + \beta_B(0)} > \frac{\tilde{\beta}(0,0)}{\tilde{\beta}_R(0) + \tilde{\beta}_B(0)} = \tilde{f}'_R(0)$$

and similarly $f'_B(0) < \tilde{f}'_B(0)$. Therefore $f_R(x) > \tilde{f}_R(x)$ and $f_B(x) < \tilde{f}_B(x)$ in a right-neighborhood of 0. Now, let $x_0 = \inf\{x : f_R(x) \leq \tilde{f}_R(x) \text{ or } f_B(x) \geq \tilde{f}_B(x)\}$, and assume $x_0 < \infty$, then

$$\begin{aligned} f_R(x_0) &= f_R(0) + \int_0^{x_0} \frac{\beta_R(\mathbf{f}(x))}{\beta_R(\mathbf{f}(x)) + \beta_B(\mathbf{f}(x))} dx \\ &> f_R(0) + \int_0^{x_0} \frac{\tilde{\beta}_R(f_R(x))}{\tilde{\beta}_R(f_R(x)) + \tilde{\beta}_B(f_B(x))} dx \\ &> \tilde{f}_R(0) + \int_0^{x_0} \frac{\tilde{\beta}_R(\tilde{f}_R(x))}{\tilde{\beta}_R(\tilde{f}_R(x)) + \tilde{\beta}_B(\tilde{f}_B(x))} dx = \tilde{f}_R(x_0) \end{aligned}$$

and similarly we obtain $f_B(x_0) < \tilde{f}_B(x_0)$, which contradict the assumption. Therefore we necessarily have $x_0 = \infty$. \square

APPENDIX K: PROOF OF THEOREM 4.7 - COMPLEMENTARY MATERIAL

Independence of $\{\underline{W}_k^{(\varepsilon)}\}_{1 \leq k \leq \lfloor xq \rfloor}$ and $\{\overline{W}_k^{(\varepsilon)}\}_{1 \leq k \leq \lfloor xq \rfloor}$. We only prove the independence of the r.v.'s $\{\underline{W}_k^{(\varepsilon)}\}_{1 \leq k \leq \lfloor xq \rfloor}$ as the independence of the r.v.'s $\{\overline{W}_k^{(\varepsilon)}\}_{1 \leq k \leq \lfloor xq \rfloor}$ can be shown similarly. For arbitrarily fixed $k, h \in \{1, \dots, \lfloor xq \rfloor\}$, $k \neq h$, and Borel sets $A, B \subseteq [0, \infty)$, we have

$$\begin{aligned} \mathbb{P}(\underline{W}_k^{(\varepsilon)} \in A, \underline{W}_h^{(\varepsilon)} \in B) &= \sum_{(q_s^R, q_s^B): 1 \leq s \leq \lfloor xq \rfloor} \mathbb{P}(\underline{W}_k^{(\varepsilon)} \in A, \underline{W}_h^{(\varepsilon)} \in B \mid \{(R_s^R, R_s^B) = (q_s^R, q_s^B)\}_{1 \leq s \leq \lfloor xq \rfloor}) \\ &\quad \times \mathbb{P}(\{(R_s^R, R_s^B) = (q_s^R, q_s^B)\}_{1 \leq s \leq \lfloor xq \rfloor}) \\ &= \sum_{(q_s^R, q_s^B)} \mathbb{P}\left(\frac{q_k^R + q_k^B}{\overline{R}_k^R(\varepsilon) + \overline{R}_k^B(\varepsilon)} \underline{W}_k \in A \mid (R_k^R, R_k^B) = (q_k^R, q_k^B)\right) \\ &\quad \times \mathbb{P}\left(\frac{q_h^R + q_h^B}{\overline{R}_h^R(\varepsilon) + \overline{R}_h^B(\varepsilon)} \underline{W}_h \in B \mid (R_h^R, R_h^B) = (q_h^R, q_h^B)\right) \mathbb{P}(\{(R_s^R, R_s^B) = (q_s^R, q_s^B)\}_{1 \leq s \leq \lfloor xq \rfloor}) \\ &= \sum_{(q_s^R, q_s^B)} \mathbb{P}(\underline{W}_k^{(\varepsilon)} \in A) \mathbb{P}(\underline{W}_h^{(\varepsilon)} \in B) \mathbb{P}(\{(R_s^R, R_s^B) = (q_s^R, q_s^B)\}_{1 \leq s \leq \lfloor xq \rfloor}) \\ &= \mathbb{P}(\underline{W}_k^{(\varepsilon)} \in A) \mathbb{P}(\underline{W}_h^{(\varepsilon)} \in B). \end{aligned}$$

Explicit tail bounds. Define the quantities

$$\overline{a}_*^{(\varepsilon)}(\kappa) := (1 + \varepsilon) \min_{0 \leq x \leq \kappa} \sum_S (\beta_S(x) + K^* \varepsilon), \quad \underline{a}_*^{(\varepsilon)}(\kappa) := (1 - \varepsilon) \min_{0 \leq x \leq \kappa} \sum_S (\beta_S(x) - K^* \varepsilon).$$

and $\underline{\mu}^{(\varepsilon)}(\kappa)$ and $\overline{\mu}^{(\varepsilon)}(\kappa)$ as in (7.30). We chose $\varepsilon > 0$ so small that $\overline{a}_*^{(\varepsilon)}(\kappa), \underline{a}_*^{(\varepsilon)}(\kappa) > 0$. By the bounds in [16], for any $\delta > 0$, we have

$$\mathbb{P}\left(\sum_{k=1}^{\lfloor \kappa q \rfloor} \overline{W}_k^{(\varepsilon)} > (1 + \delta) \overline{\mu}^{(\varepsilon)}(\kappa)\right) < \frac{1}{1 + \delta} \exp(-q \overline{a}_*^{(\varepsilon)}(\kappa) \overline{\mu}^{(\varepsilon)}(\kappa) (\delta - \log(1 + \delta)))$$

and

$$\mathbb{P}\left(\sum_{k=1}^{\lfloor \kappa q \rfloor} \underline{W}_k^{(\varepsilon)} > (1 + \delta) \underline{\mu}^{(\varepsilon)}(\kappa)\right) < \frac{1}{1 + \delta} \exp(-q \underline{a}_*^{(\varepsilon)}(\kappa) \underline{\mu}^{(\varepsilon)}(\kappa) (\delta - \log(1 + \delta))),$$

and, for $0 < \delta < 1$, we have

$$\mathbb{P}\left(\sum_{k=1}^{\lfloor \kappa q \rfloor} \overline{W}_k^{(\varepsilon)} < (1 - \delta) \overline{\mu}^{(\varepsilon)}(\kappa)\right) < \exp(-q \overline{a}_*^{(\varepsilon)}(\kappa) \overline{\mu}^{(\varepsilon)}(\kappa) (-\delta - \log(1 - \delta)))$$

and

$$\mathbb{P}\left(\sum_{k=1}^{\lfloor \kappa q \rfloor} \underline{W}_k^{(\varepsilon)} < (1 - \delta) \underline{\mu}^{(\varepsilon)}(\kappa)\right) < \exp(-q \underline{a}_*^{(\varepsilon)}(\kappa) \underline{\mu}^{(\varepsilon)}(\kappa) (-\delta - \log(1 - \delta))).$$

APPENDIX L: PROOF OF LEMMA E.3: PARTS (ii), (iii)

L.1. Proof of Part (ii). Although the proof is quite similar to the proof of Part (i), we provide some details. We divide the proof in two steps: for arbitrarily fixed $\kappa, \delta > 0$, we prove that there exist $c'(\kappa, \delta) > 0$ and $c''(\kappa, \delta) > 0$ (not depending on n) such that

$$(L.1) \quad \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(\tilde{Y}_S(\mathbf{k}) > \delta n (qp)^r) \ll e^{-c'(\kappa, \delta) n (qp)^r}$$

and

$$(L.2) \quad \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \mathbb{P}(Y_S(\mathbf{k}) > \delta) \ll e^{-c''(\kappa, \delta) n (qp)^r}.$$

Part (ii) then follows setting $c(\kappa, \delta) := \min\{c'(\kappa, \delta), c''(\kappa, \delta)\}$.

Step 1: Proof of (L.1). Arguing similarly to the derivation of (E.4), we have

$$\begin{aligned} & \mathbb{P}(|Q_{k+1}^S - \beta_S(k_S/q)n(qp)^r| > \delta n (qp)^r \mid \mathbf{N}[k] = \mathbf{k}) \\ & \leq \mathbb{P}(|\mathcal{S}_S[k]| - \beta_S(k_S/q)n(qp)^r| + N_S[k] \\ & \quad + |(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| \\ & \quad + |\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| \mid \mathbf{N}[k] = \mathbf{k}) \\ & \leq \mathbb{P}(|\mathcal{S}_S[k]| - \beta_S(k_S/q)n(qp)^r| > (\delta n (qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) \\ & \quad + \mathbb{P}(N_S[k] > (\delta n (qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) \\ & \quad + \mathbb{P}(|(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| > (\delta n (qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) \\ (L.3) \quad & \quad + \mathbb{P}(|\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| > (\delta n (qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}). \end{aligned}$$

Now, note that, for any $\mathbf{k} \in \mathbb{Z}(\kappa)$, we have

$$N_S[k] \leq \kappa q$$

$$|(\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}| \leq N_S[k] \leq \kappa q$$

and

$$|\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| > (\delta n (qp)^r)/4 \leq N_{S^c}(k) \leq \kappa q.$$

Since $q \ll n(qp)^r$ (which follows by (2.3) *ii*), we then have that there exists $n_{\kappa,\delta}$ such that, for all $n \geq n_{\kappa,\delta}$,

$$\mathbb{P}(N_S[k] > (\delta n(qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) = 0, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa)$$

$$\mathbb{P}((\mathcal{V}_W \setminus \mathcal{S}_S[k]) \cap \mathcal{V}_S[k] \cap \{v : D_S^v[k] \geq r\}) > (\delta n(qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) = 0, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa)$$

and

$$\mathbb{P}(|\mathcal{S}_S[k] \cap \mathcal{V}_{S^c}[k] \cap \{v : D_{S^c}^v[k] \geq r\}| > (\delta n(qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) = 0, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa).$$

Therefore, by (L.3), for any $n \geq n_{\kappa,\delta}$,

$$\begin{aligned} & \mathbb{P}(|Q_{k+1}^S - \beta_S(k_1/q)n(qp)^r| > \delta n(qp)^r \mid \mathbf{N}[k] = \mathbf{k}) \\ (L.4) \quad & \leq \mathbb{P}(|\mathcal{S}_S[k]| - \beta_S(k_1/q)n(qp)^r| > (\delta n(qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}), \quad \forall \mathbf{k} \in \mathbb{Z}(\kappa). \end{aligned}$$

We proceed distinguishing two sub-steps. In the first step we provide an exponential bound (uniformly in $\mathbf{k} \in \mathbb{T}(\kappa)$) for the probability in (L.4), in the second step we conclude the proof of (L.1).

Sub-step 1: Exponential bound for the probability (L.4). We prove that there exists $n_{\kappa,\delta} \geq 1$ such that, for all $n \geq n_{\kappa,\delta}$,

$$(L.5) \quad \mathbb{P}(|\mathcal{S}_S[k]| - n\beta_S(k_1/q)(qp)^r| > (\delta n(qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) \leq 2e^{-c_1(\kappa,\delta)n(qp)^r}, \quad \forall \mathbf{k} \in \mathbb{Z}(\kappa)$$

where $c_1(\kappa, \delta) > 0$ is a suitable positive constant (not depending on n). By (3.14) we have

$$\begin{aligned} & \mathbb{P}(|\mathcal{S}_S[k]| - n\beta_S(k_S/q)(qp)^r| > (n\delta(qp)^r)/4 \mid \mathbf{N}[k] = \mathbf{k}) \\ & \leq \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \leq n(qp)^r(\beta_S(k_S/q) - \delta/4)) \\ (L.6) \quad & + \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \geq n(qp)^r(\beta_S(k_S/q) + \delta/4)). \end{aligned}$$

Taking

$$\delta' \in \left(0, \frac{\delta(r!)}{(\kappa + \alpha_S)^r}\right),$$

and using Lemma E.4 we have that there exists $n_{\kappa,\delta} \geq 1$ such that, for any $n \geq n_{\kappa,\delta}$,

$$(L.7) \quad n_W \pi_S(\mathbf{k}) > n\beta_S(k_S/q)(qp)^r(1 - \delta'/4) > n(qp)^r(\beta_S(k_S/q) - \delta/4), \quad \forall \mathbf{k} \in \mathbb{T}(\kappa)$$

and

$$(L.8) \quad n_W \pi_S(\mathbf{k}) < n\beta_S(k_S/q)(qp)^r(1 + \delta'/4) < n(qp)^r(\beta_S(k_S/q) + \delta/4), \quad \forall \mathbf{k} \in \mathbb{T}(\kappa).$$

By (L.7), the usual concentration bound for the binomial distribution (see (I.2)) and the fact that the function H defined by (3.4) decreases on $[0, 1)$, for any $n \geq n_{\kappa,\delta}$, we have, uniformly in $\mathbf{k} \in \mathbb{T}(\kappa)$,

$$\begin{aligned} & \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \leq n(qp)^r(\beta_S(k_S/q) - \delta/4)) \\ & \leq \exp\left(-n_W \pi_S(\mathbf{k}) H\left(\frac{n(qp)^r(\beta_S(k_S/q) - \delta/4)}{n_W \pi_S(\mathbf{k})}\right)\right) \\ & \leq \exp\left(-n(qp)^r(\beta_S(k_S/q) - \delta/4) H\left(\frac{\beta_S(k_S/q) - \delta/4}{\beta_S(k_S/q)(1 - \delta'/4)}\right)\right) \\ (L.9) \quad & \leq \exp\left(-n(qp)^r(\alpha_S^r - \delta/4) H\left(\frac{1 - \delta/(4(\alpha_S)^r)}{1 - \delta'/4}\right)\right). \end{aligned}$$

By (L.7), (L.8), the usual concentration bound for the binomial distribution (see (I.1)) and the fact that the function H increases on $(1, \infty)$, for any $n \geq n_{\kappa, \delta}$, we have, uniformly in $\mathbf{k} \in \mathbb{T}(\kappa)$,

$$\begin{aligned}
& \mathbb{P}(\text{Bin}(n_W, \pi_S(\mathbf{k})) \geq n(qp)^r(\beta_S(k_S/q) + \delta/4)) \\
& \leq \exp\left(-n_W \pi_S(\mathbf{k}) H\left(\frac{n(qp)^r(\beta_S(k_S/q) + \delta/4)}{n_W \pi_S(\mathbf{k})}\right)\right) \\
& \leq \exp\left(-n(qp)^r(\beta_S(k_S/q) - \delta/4) H\left(\frac{\beta_S(k_S/q) + \delta/4}{\beta_S(k_S/q)(1 + \delta'/4)}\right)\right) \\
\text{(L.10)} \quad & \leq \exp\left(-n(qp)^r(\alpha_S^r - \delta/4) H\left(\frac{1 + \delta/(4(\alpha_S^r))}{1 + \delta'/4}\right)\right).
\end{aligned}$$

The inequality (L.5) follows by (L.6), (L.9) and (L.10).

Sub-step 2: Conclusion of the proof of (L.1). The claim easily follows by (L.4) and (L.5).

Step 2: Proof of (L.2). By the previous part of the proof, we have that, for all $n \geq n_{\kappa, \delta}$, (L.11)

$$\mathbb{P}(|Q_{k+1}^S - \beta_S(k_S/q)n(qp)^r| > \delta n(qp)^r \mid \mathbf{N}[k] = \mathbf{k}) \leq \delta e^{-\tilde{c}(\kappa, \delta)n(qp)^r}, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa)$$

for a suitable positive constant $\tilde{c}(\kappa, \delta) > 0$ (not depending on n). By the inequality $\|x\| - |y| \leq |x - y|$, $x, y \in \mathbb{R}$, it follows

$$\text{(L.12)} \quad \mathbb{P}(\|Q_{k+1}^S\| - |\beta_S(k_S/q)n(qp)^r| > \delta n(qp)^r \mid \mathbf{N}[k] = \mathbf{k}) \leq \delta e^{-\tilde{c}(\kappa, \delta)n(qp)^r}, \quad \forall \mathbf{k} \in \mathbb{T}(\kappa).$$

By the triangular inequality and the union bound, we have

$$\begin{aligned}
& \mathbb{P}(\|Q_{k+1}^S\| + |Q_{k+1}^{S^c}| - (|\beta_S(k_S/q)| + |\beta_{S^c}(k_{S^c}/q)|)n(qp)^r > \delta n(qp)^r \mid \mathbf{N}[k] = \mathbf{k}) \\
& \leq \mathbb{P}(\|Q_{k+1}^S\| - |\beta_S(k_S/q)n(qp)^r| > (\delta n(qp)^r)/2 \mid \mathbf{N}[k] = \mathbf{k}) + \\
& \quad \mathbb{P}(\|Q_{k+1}^{S^c}\| - |\beta_{S^c}(k_{S^c}/q)n(qp)^r| > (\delta n(qp)^r)/2 \mid \mathbf{N}[k] = \mathbf{k}).
\end{aligned}$$

Combining this relation with (L.12), for all $n \geq n_{\kappa, \delta}$, we have

$$\text{(L.13)} \quad \mathbb{P}(\|Q_{k+1}^S\| + |Q_{k+1}^{S^c}| - (|\beta_S(k_S/q)| + |\beta_{S^c}(k_{S^c}/q)|)n(qp)^r > \delta n(qp)^r \mid \mathbf{N}[k] = \mathbf{k}) \leq 2\delta e^{-c_2(\kappa, \delta)n(qp)^r},$$

$\forall \mathbf{k} \in \mathbb{T}(\kappa)$ and some positive constant $c_2(\kappa, \delta) > 0$ (not depending on n). By Lemma E.6, (L.11) and (L.13), for all $n \geq n_\delta$, we have

$$\mathbb{P}\left(\left|U_{k+1}^S - \frac{|\beta_S(k_S/q)|}{|\beta_S(k_S/q)| + |\beta_{S^c}(k_{S^c}/q)|}\right| > \delta \mid \mathbf{N}[k] = \mathbf{k}\right) \leq c_3(\kappa, \delta, \beta_{\min})e^{-c_4(\kappa, \delta, \beta_{\min})n(qp)^r},$$

$\forall \mathbf{k} \in \mathbb{T}(\kappa)$ and suitable positive constants $c_3(\kappa, \delta, \beta_{\min})$ and $c_4(\kappa, \delta, \beta_{\min})$ (not depending on n), where the constant $\beta_{\min} > 0$ is defined by (E.2). The claim (L.2) easily follows by this inequality.

Proof of Part (iii). The proof of Part (iii) follows the same lines as the proof of Part (ii). In particular, one first shows that, for any $\kappa, \delta > 0$, there exists $n_{\kappa, \delta} \geq 1$ such that, for any $n \geq n_{\kappa, \delta}$,

$$\text{(L.14)} \quad \begin{aligned} & \mathbb{P}(|Q_{k+1}^S - \beta_S(k_R/q, k_B/q)n| > \delta n \mid \mathbf{N}[k] = \mathbf{k}) \\ & \leq \mathbb{P}(\|\mathcal{S}_S[k]\| - \beta_S(k_R/q, k_B/q)n > (\delta n)/4 \mid \mathbf{N}[k] = \mathbf{k}), \quad \forall \mathbf{k} \in \mathbb{T}(\kappa). \end{aligned}$$

Then one provides an exponential bound (uniformly on $\mathbf{k} \in \mathbb{T}(\kappa)$) for the probability in (L.14), from which the claim follows. We omit the details.

Proof of Lemma E.4.

Proof of Part (i). We divide the proof of the Part (i) in two steps, where we prove that for every $\kappa > 0$

$$(L.15) \quad \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| 1 - \frac{[(k_S/q + \alpha_S)qp]^r / r!}{\pi_S(\mathbf{k})} \right| \rightarrow 0$$

and

$$(L.16) \quad \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_w((k_S/q + \alpha_S)qp)^r / r!}{(\beta_S(k_S/q) + k_S/q)q} - 1 \right| \rightarrow 0,$$

respectively. The claim then easily follows combining these two uniform convergences on $\mathbb{T}(\kappa)$.

Step 1: Proof of (L.15). We divide the proof of (L.15) in two further steps. In the first step, we show the non-uniform convergence, i.e., we prove that, for any sequence $\mathbf{k}_n = \mathbf{k} = (k_S, k_{S^c}) \in (\mathbb{N} \cup \{0\})^2$ such that $k_S/q \rightarrow x_S$, for some $x_S \in [0, \infty)^2$, it holds

$$(L.17) \quad \pi_S(\mathbf{k}) = \frac{[(k_S + a_S)p]^r}{r!} (1 + O((k_1 + a_S)p + (k_S + a_S)^{-1}))$$

$$(L.18) \quad \sim \frac{((x_S + \alpha_S)qp)^r}{r!}.$$

In the second step, we conclude the proof of (L.15) lifting the convergence (L.17) to a uniform convergence on $\mathbb{T}(\kappa)$. We warn the reader that in the proof of (L.17) and (L.18) we omit the dependence on n since no confusion arises in the computations. Such a dependence is instead made explicit in the second step.

Sub-Step 1: Proof of (L.17) and (L.18). We have

$$\pi_S(\mathbf{k}) = \sum_{m=0}^{k_S + a_S - r} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq m + r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = m).$$

By e.g. formula (8.1) in [8], we have, for any $j, \ell, m \in \mathbb{N}$,

$$\mathbb{P}(\text{Bin}(j + \ell, p) \geq m) = \frac{[(j + \ell)p]^m}{m!} (1 + O((j + \ell)p + (j + \ell)^{-1})).$$

Since $(1 - p)^{(k_{S^c} + a_{S^c})p} \rightarrow 1$, for n large enough we have

$$\begin{aligned} \pi_S(\mathbf{k}) &= \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = 0) \\ &+ \sum_{m=1}^{k_S + a_S - r} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq m + r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = m) \\ &= (1 - p)^{(k_{S^c} + a_{S^c})p} \frac{[(k_S + a_S)p]^r}{r!} (1 + O((k_1 + a_S)p + (k_S + a_S)^{-1})) \\ &+ \sum_{m=1}^{k_S + a_S - r} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq m + r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = m) \\ &= \frac{[(k_S + a_S)p]^r}{r!} \left((1 + O((k_1 + a_S)p + (k_S + a_S)^{-1})) \right) \end{aligned}$$

$$+ \frac{r!}{[(k_1 + a_S)p]^r} \times \sum_{m=1}^{k_S + a_S - r} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq m + r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = m) \Bigg).$$

The claim (L.17) follows if we check that (L.19)

$$\frac{r!}{[(k_S + a_S)p]^r} \sum_{m=1}^{k_S + a_S - r} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq m + r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = m) = O((k_S + a_S)p).$$

By the usual concentration bound for the binomial distribution (see (I.1)) letting H denote the function defined by (3.4), for n large enough we have

$$\begin{aligned} & \sum_{m=1}^{k_S + a_S - r} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq m + r) \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) = m) \\ & \leq \sum_{k \geq r+1} \mathbb{P}(\text{Bin}(k_S + a_S, p) \geq k) \\ & \leq \sum_{k \geq r+1} \exp\left(- (k_S + a_S)p H\left(\frac{k}{(k_S + a_S)p}\right)\right) \\ & \leq \sum_{k \geq r+1} \exp\left(-k \left(\log \frac{k}{(k_S + a_S)p} - 1\right)\right) \\ & \leq \sum_{k \geq r+1} \exp\left(- (r+1) \left(\log \frac{k}{(k_S + a_S)p} - 1\right)\right) \\ & = e^{r+1} \sum_{k \geq r+1} \left(\frac{(k_S + a_S)p}{k}\right)^{r+1} \\ & = e^{r+1} \left(\sum_{k \geq r+1} \frac{1}{k^{r+1}}\right) [(k_S + a_S)p]^{r+1}. \end{aligned}$$

The relation (L.19) follows by this inequality, and the proof of (L.17) is completed. As far as (L.18) is concerned, we note that by (2.2) and (2.3), we have

$$\begin{aligned} \frac{[(k_S + a_S)p]^r}{r!} (1 + O((k_S + a_S)p + (k_S + a_S)^{-1})) & \sim \frac{[(k_S + a_S)p]^r}{r!} \\ & \sim \frac{((x_S + \alpha_S)qp)^r}{r!}. \end{aligned}$$

Sub-Step 2: Conclusion of the proof of (L.15). Reasoning by contradiction, suppose that

$$\limsup_{n \rightarrow \infty} \sup_{\mathbf{k} \in \mathbb{T}_n(\kappa)} \left| 1 - \frac{[(k_S/q_n + \alpha_S)q_n p_n]^r / r!}{\pi_S(\mathbf{k}_{\mathbf{a}^{(n)}})} \right| = c > 0,$$

where $\mathbf{a}^{(n)} = (a_R^{(n)}, a_B^{(n)})$ and $c > 0$ is a positive constant. Letting $\{n'\}$ denote a subsequence that realizes the lim sup, we have

$$\lim_{n' \rightarrow \infty} \sup_{\mathbf{k} \in \mathbb{T}_{n'}(\kappa)} \left| 1 - \frac{[(k_S/q_{n'} + \alpha_S)q_{n'} p_{n'}]^r / r!}{\pi_S(\mathbf{k}_{\mathbf{a}^{(n')}})} \right| = \lim_{n' \rightarrow \infty} \max_{\mathbf{k} \in \mathbb{T}_{n'}(\kappa)} \left| 1 - \frac{[(k_S/q_{n'} + \alpha_S)q_{n'} p_{n'}]^r / r!}{\pi_S(\mathbf{k}_{\mathbf{a}^{(n')}})} \right| = c > 0.$$

Setting

$$\mathbf{k}_{n'}^*(\kappa) := \arg \max_{\mathbf{k} \in \mathbb{T}_{n'}} \left| 1 - \frac{[(k_S/q_{n'} + \alpha_S)q_{n'}p_{n'}]^r/r!}{\pi_S(\mathbf{k}_{\mathbf{a}(n')})} \right|,$$

we have (using an obvious notation)

$$(L.20) \quad \lim_{n' \rightarrow \infty} \left| 1 - \frac{[(k_{n'}^*(\kappa))_S/q_{n'} + \alpha_S]q_{n'}p_{n'}^r/r!}{\pi_S((\mathbf{k}_{n'}^*(\kappa))_{\mathbf{a}(n')})} \right| = c > 0.$$

Since the sequence $\mathbf{k}_{n'}^*(\kappa)/q_{n'}$ is contained in the compact $\mathbb{T}'(\kappa)$ defined as in (7.4), there exists a subsequence $\{n''\}$ such that $\mathbf{k}_{n''}^*(\kappa)/q_{n''} \rightarrow (z_S, z_{S^c}) \in \mathbb{K}(\kappa)$. So by (L.20) it follows

$$\lim_{n'' \rightarrow \infty} \left| 1 - \frac{[(k_{n''}^*(\kappa))_S/q_{n''} + \alpha_S]q_{n''}p_{n''}^r/r!}{\pi_S((\mathbf{k}_{n''}^*(\kappa))_{\mathbf{a}(n'')})} \right| = \lim_{n'' \rightarrow \infty} \left| 1 - \frac{[(z_S + \alpha_S)q_{n''}p_{n''}^r/r!]}{\pi_S((\mathbf{k}_{n''}^*(\kappa))_{\mathbf{a}(n'')})} \right| = c > 0,$$

which contradicts (L.18).

Step 2: Proof of (L.16). We have

$$(L.21) \quad n_W \frac{((k_S/q + \alpha_S)qp)^r}{r!} = (k_S/q + \alpha_S)^r qp n_W \frac{(qp)^{r-1}}{r!}.$$

So, by the definition of g and assumption $q = g$, it follows

$$(L.22) \quad \begin{aligned} n_W \frac{((k_S/q + \alpha_S)qp)^r}{r!} &\sim r^{-1}[1 - r^{-1}]^{r-1}(k_S/q + \alpha_S)^r q \\ &= (\beta_S(k_S/q) + k_S/q)q. \end{aligned}$$

Arguing as in the proof of Step 2 for (L.15) (i.e., reasoning by contradiction, considering a subsequence realizing the corresponding lim sup, using the compactness of $\mathbb{T}'(\kappa)$ and finally using (L.22)) one proves that the convergence (L.22) is indeed uniform on $\mathbb{T}(\kappa)$.

Proof of Part (ii). By (L.21), the current definition of the function β_S and the fact that $n \sim n_W$, it follows

$$n_W \frac{((k_S/q + \alpha_S)qp)^r}{r!} \sim n\beta_S(k_S/q)(qp)^r.$$

Arguing as in the proof of Step 2 for (L.15) one has

$$\sup_{\mathbf{k} \in \mathbb{Z}(\kappa)} \left| \frac{n_W(k_S/q + \alpha_S)^r (qp)^r/r!}{n\beta_S(k_S/q)(qp)^r} - 1 \right| \rightarrow 0.$$

The claim follows combining this uniform convergence with (L.15) (whose derivation does not depend on the assumptions on the asymptotic behavior of q and the particular definition of β_S).

Proof of Part (iii), for $q = p^{-1}$. We start noticing that

$$\begin{aligned} \pi_S(\mathbf{k}) &:= \mathbb{P}(\text{Bin}(k_S + a_S, p) - \text{Bin}(k_{S^c} + a_{S^c}, p) \geq r) \\ &= \sum_{r'=r}^{k_S+a_S} \mathbb{P}(\text{Bin}(k_S + a_S, p) = r') \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) \leq r' - r) \\ &= \sum_{r'=r}^{\infty} \mathbb{P}(\text{Bin}(k_S + a_S, p) = r') \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) \leq r' - r) \end{aligned}$$

and that

$$\begin{aligned}\widehat{\pi}_S(\mathbf{k}) &:= \mathbb{P}(\text{Po}((k_S + a_S)p) - \text{Po}((k_{S^c} + a_{S^c})p) \geq r) \\ &= \sum_{r'=r}^{\infty} \mathbb{P}(\text{Po}((k_S + a_S)p) = r') \mathbb{P}(\text{Po}((k_{S^c} + a_{S^c})p) \leq r' - r).\end{aligned}$$

This implies

$$|\pi_S(\mathbf{k}) - \widehat{\pi}_S(\mathbf{k})| \leq 2\kappa^2 p.$$

Indeed, letting d_{TV} denote the total variation distance and recalling that $d_{TV}(\text{Bin}(m, p), \text{Po}[mp]) \leq mp^2$, we have

$$\begin{aligned}|\pi_S(\mathbf{k}) - \widehat{\pi}_S(\mathbf{k})| &\leq \sum_{r'=r}^{\infty} \left| \mathbb{P}(\text{Bin}(k_S + a_S, p) = r') \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) \leq r' - r) \right. \\ &\quad \left. - \mathbb{P}(\text{Po}((k_S + a_S)p) = r') \mathbb{P}(\text{Po}((k_{S^c} + a_{S^c})p) \leq r' - r) \right| \\ &\leq \sum_{r'=r}^{\infty} \mathbb{P}(\text{Bin}(k_S + a_S, p) = r') \left| \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) \leq r' - r) - \mathbb{P}(\text{Po}((k_{S^c} + a_{S^c})p) \leq r' - r) \right| \\ &\quad + \sum_{r'=r}^{\infty} \left| \mathbb{P}(\text{Bin}((k_S + a_S)p) = r') - \mathbb{P}(\text{Po}((k_S + a_S)p) = r') \right| \mathbb{P}(\text{Po}((k_{S^c} + a_{S^c})p) \leq r' - r) \\ &\leq d_{TV}(\text{Bin}(k_{S^c} + a_{S^c}, p), \text{Po}((k_{S^c} + a_{S^c})p)) \sum_{r'=r}^{\infty} \mathbb{P}(\text{Bin}(k_S + a_S, p) = r') \\ &\quad + \sum_{r'=r}^{\infty} \left| \mathbb{P}(\text{Bin}((k_S + a_S)p) = r') - \mathbb{P}(\text{Po}((k_S + a_S)p) = r') \right| \\ &\leq d_{TV}(\text{Bin}(k_{S^c} + a_{S^c}, p), \text{Po}((k_{S^c} + a_{S^c})p)) + d_{TV}(\text{Bin}(k_S + a_S, p), \text{Po}((k_S + a_S)p)).\end{aligned}$$

Therefore, noticing that by (2.4) we have $\beta_S(k_R/q, k_B/q) = \widehat{\pi}_S(\mathbf{k})$, it follows

$$\begin{aligned}\sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \pi_S(\mathbf{k})}{n \beta_S(k_R/q, k_B/q)} - 1 \right| &= \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \pi_S(\mathbf{k})}{n \widehat{\pi}_S(\mathbf{k})} - 1 \right| \\ &= \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \pi_S(\mathbf{k}) - n \widehat{\pi}_S(\mathbf{k})}{n \widehat{\pi}_S(\mathbf{k})} \right| \\ &\leq \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{n_W \pi_S(\mathbf{k}) - n_W \widehat{\pi}_S(\mathbf{k})}{n \widehat{\pi}_S(\mathbf{k})} \right| + \frac{n - n_W}{n} \\ &\leq \sup_{\mathbf{k} \in \mathbb{T}(\kappa)} \left| \frac{2\kappa p}{\widehat{\pi}_S(\mathbf{k})} \right| + \frac{n - n_W}{n} \rightarrow 0,\end{aligned}$$

where the latter limit follows since $\inf_{\mathbf{k} \in \mathbb{T}(\kappa)} \widehat{\pi}_S(\mathbf{k})$ is bounded away from 0.

Proof of Part (iii), for $p^{-1} \ll q \ll n$. We start noticing that $\frac{k_S + a_S}{k_{S^c} + a_{S^c}} > 1$, and so, setting $\chi := \frac{(k_S + a_S + k_{S^c} + a_{S^c})qp}{2}$, we have

$$\begin{aligned}\pi_S(\mathbf{k}) &:= \mathbb{P}(\text{Bin}(k_S + a_S, p) - \text{Bin}(k_{S^c} + a_{S^c}, p) \geq r) = 1 - \mathbb{P}(\text{Bin}(k_S + a_S, p) - \text{Bin}(k_{S^c} + a_{S^c}, p) < r) \\ &\geq 1 - [\mathbb{P}(\text{Bin}(k_S + a_S, p) \leq \chi + r) + \mathbb{P}(\text{Bin}(k_{S^c} + a_{S^c}, p) \geq \chi)] \rightarrow 1,\end{aligned}$$

where the latter limit can be checked using the concentration inequalities reported in Appendix I to bound the last two terms. Similarly one can check that $\pi_{S^c}(\mathbf{k}) \rightarrow 0$.

Proof of Lemma E.5. By the definition of g , we have

$$(L.23) \quad n_W(k_{S^c}/q + \alpha_{S^c})(k_S/q + \alpha_S)^r \frac{(qp)^{r+1}}{r!} = (k_{S^c}/q + \alpha_{S^c})qp(k_S/q + \alpha_S)^r qp n_W \frac{(qp)^{r-1}}{r!} \\ \sim r^{-1}[1 - r^{-1}]^{r-1} (k_{S^c}/q + \alpha_{S^c})(k_S/q + \alpha_S)^r q^2 p.$$

Along similar lines as in the proof of Step 1 for (L.15) (see the proof of Lemma E.4), one has

$$(L.24) \quad n_W \hat{\pi}_S(\mathbf{k}) \sim n_W(k_2/q + \alpha_{S^c})(k_S/q + \alpha_S)^r \frac{(qp)^{r+1}}{r!}.$$

Arguing as in the proof of Step 2 for (L.15) (again, see the proof of Lemma E.4), one has that the convergences (L.23) and (L.24) are indeed uniform on $\mathbb{T}(\kappa)$, and the claim follows.

Proof of Lemma E.6. For $\varepsilon \in (0, 1)$, define the events

$$B_{\varepsilon\mu/4}^{(n)} := \left\{ |X_n - \mu_n| \leq \frac{\varepsilon\mu}{4} \right\}, \quad C_{\varepsilon\mu/4}^{(n)} := \left\{ |X'_n - \mu'_n| \leq \frac{\varepsilon\mu}{4} \right\}, \quad n \in \mathbb{N}.$$

Note that

$$\mu_n - \frac{\varepsilon\mu}{4} \leq X_n(\omega) \leq \mu_n + \frac{\varepsilon\mu}{4}, \quad \forall \omega \in B_{\varepsilon\mu/4}^{(n)}$$

and

$$0 < \mu'_n - \frac{\varepsilon\mu}{4} \leq X'_n(\omega) \leq \mu'_n + \frac{\varepsilon\mu}{4}, \quad \forall \omega \in C_{\varepsilon\mu/4}^{(n)}.$$

Therefore, $C_{\varepsilon\mu/4}^{(n)} \subseteq \{X'_n \neq 0\}$ and, for any $\omega \in B_{\varepsilon\mu/4}^{(n)} \cap C_{\varepsilon\mu/4}^{(n)}$, we have

$$(L.25) \quad \frac{4\mu_n - \varepsilon\mu}{4\mu'_n + \varepsilon\mu} \leq \frac{X_n(\omega)}{X'_n(\omega)} \leq \frac{4\mu_n + \varepsilon\mu}{4\mu'_n - \varepsilon\mu}.$$

We shall check later on that this inequality implies

$$(L.26) \quad \left| \frac{X_n(\omega)}{X'_n(\omega)} - \frac{\mu_n}{\mu'_n} \right| \leq \varepsilon.$$

Therefore,

$$B_{\varepsilon\mu/4}^{(n)} \cap C_{\varepsilon\mu/4}^{(n)} \subseteq \left\{ \left| \frac{X_n}{X'_n} - \frac{\mu_n}{\mu'_n} \right| \leq \varepsilon, X'_n \neq 0 \right\} \subseteq \left\{ \left| \frac{X_n}{X'_n} - \frac{\mu_n}{\mu'_n} \right| \leq \varepsilon \right\},$$

and so

$$\mathbb{P} \left(\left| \frac{X_n}{X'_n} - \frac{\mu_n}{\mu'_n} \right| > \varepsilon \right) \leq \mathbb{P} \left((B_{\varepsilon\mu/4}^{(n)})^c \cup (C_{\varepsilon\mu/4}^{(n)})^c \right) \leq \mathbb{P}(|X_n - \mu_n| > \varepsilon\mu/4) + \mathbb{P}(|X'_n - \mu'_n| > \varepsilon\mu/4).$$

It remains to check that (L.25) implies (L.26). Indeed

$$\frac{4\mu_n + \varepsilon\mu}{4\mu'_n - \varepsilon\mu} = \frac{4\mu_n + \varepsilon\mu}{4\mu'_n(1 - \frac{\varepsilon\mu}{4\mu'_n})} < \frac{4\mu_n + \varepsilon\mu}{4\mu'_n} \left(1 + \frac{2\varepsilon\mu}{4\mu'_n} \right) = \frac{\mu_n}{\mu'_n} + \frac{1}{4} \frac{\varepsilon\mu}{\mu'_n} + \frac{1}{2} \frac{\varepsilon\mu\mu_n}{(\mu'_n)^2} + \frac{1}{8} \frac{(\varepsilon\mu)^2}{(\mu'_n)^2} < \frac{\mu_n}{\mu'_n} + \varepsilon,$$

where the first inequality holds since $\frac{1}{1-x} < 1 + 2x$, $x \in (0, 1/2)$. Similarly,

$$\frac{4\mu_n - \varepsilon\mu}{4\mu'_n + \varepsilon\mu} = \frac{4\mu_n - \varepsilon\mu}{4\mu'_n(1 + \frac{\varepsilon\mu}{4\mu'_n})} > \frac{4\mu_n - \varepsilon\mu}{4\mu'_n} \left(1 - \frac{\varepsilon\mu}{4\mu'_n} \right) = \frac{\mu_n}{\mu'_n} - \frac{1}{4} \frac{\varepsilon\mu}{\mu'_n} - \frac{1}{4} \frac{\varepsilon\mu\mu_n}{(\mu'_n)^2} + \frac{1}{16} \frac{(\varepsilon\mu)^2}{(\mu'_n)^2} > \frac{\mu_n}{\mu'_n} - \varepsilon,$$

where the first inequality holds since $\frac{1}{1+x} > 1 - x$, $x \in (0, 1)$.

APPENDIX M: THE STOPPED PROCESS

Formal definition. N_S^{stop} is the point process on $[0, \infty) \times \mathcal{V}_W$ constructed by thinning $\{(T'_k, V'_k)\}_{k \in \mathbb{N}}$ in the following manner: for $k \in \mathbb{N}$ with $T'_k < Z_{\text{stop}}$, we retain those couples (T'_k, V'_k) for which, at time $(T'_k)^-$, the node V'_k is white and satisfies the “threshold condition with respect to S ”; for $k \in \mathbb{N}$ with $T'_k \geq Z_{\text{stop}}$, we keep, exclusively in the process N_B^{stop} , those couples (T'_k, V'_k) for which, at time $(T'_k)^-$, the node V'_k is still white and satisfies “threshold condition with respect to B ”. Note that the process N_S^{stop} is indistinguishable from N_S , up to time Z_{stop} . Furthermore, the process N_R^{stop} does not evolve after Z_{stop} , while the process N_B^{stop} evolves beyond Z_{stop} . Formally, for $A \in \mathcal{B}([0, \infty))$ and $L \subseteq \mathcal{V}_W$, we define the T_{stop} -stopped R activation process as

$$N_R^{\text{stop}}(A \times L) := \sum_{k \in \mathbb{N}: T'_k < Z_{\text{stop}}} \mathbf{1}_A(T'_k) \mathbf{1}_{\mathcal{V}_W(T'_k) \cap L}(V'_k) \mathbf{1}\{D_R^{(V'_k)}(T'_k) - D_B^{(V'_k)}(T'_k) \geq r\} \quad (\text{M.1})$$

and the t_{stop} -stopped B activation process as

$$N_B^{\text{stop}}(A \times L) := \sum_{k \in \mathbb{N}: T'_k < Z_{\text{stop}}} \mathbf{1}_A(T'_k) \mathbf{1}_{\mathcal{V}_W(T'_k) \cap L}(V'_k) \mathbf{1}\{D_B^{(V'_k)}(T'_k) - D_R^{(V'_k)}(T'_k) \geq r\} \\ + \sum_{k \in \mathbb{N}: T'_k \geq Z_{\text{stop}}} \mathbf{1}_A(T'_k) \mathbf{1}_{\mathcal{V}_{W, \text{stop}}(T'_k) \cap L}(V'_k) \mathbf{1}\{D_{B, \text{stop}}^{(V'_k)}(T'_k) - D_{R, \text{stop}}^{(V'_k)}(T'_k) \geq r\}.$$

We call the point process $N^{\text{stop}} := N_R^{\text{stop}} + N_B^{\text{stop}}$ R -stopped activation process, and denote by $\{(T_k^{\text{stop}}, V_k^{\text{stop}})\}_{k \in \mathbb{N}}$ its points. Setting

$$A^{*, \text{stop}} := A_R^{*, \text{stop}} + A_B^{*, \text{stop}}, \quad \text{where} \quad A_S^{*, \text{stop}} := N_S^{\text{stop}}([0, \infty) \times \mathcal{V}_W) + a_S,$$

and

$$K_{\text{stop}}^* := \min\{k \in \mathbb{N} : T_k^{\text{stop}}, \mathcal{S}_B(T_k^{\text{stop}}) \cap \mathcal{V}_W(T_k^{\text{stop}}) = \emptyset\},$$

it turns out

$$A^{*, \text{stop}} = K_{\text{stop}}^* + a_R + a_B - 1.$$

Note that by construction

$$A_R^{*, \text{stop}} = N_R^{\text{stop}}([0, Z_{\text{stop}}) \times \mathcal{V}_W) + a_R = N_R([0, Z_{\text{stop}}) \times \mathcal{V}_W) + a_R \leq A_R^* \quad \mathbb{P}\text{-a.s.}$$

Extending the stopped process beyond its natural termination. Note that the stopped process naturally terminates at $T_{K_{\text{stop}}^* - 1}^{\text{stop}}$. The above construction is used also to extend the R -stopped process beyond $T_{K_{\text{stop}}^* - 1}^{\text{stop}}$. In particular we define N_B beyond $T_{K_{\text{stop}}^* - 1}^{\text{stop}}$ as follows: the points (T_k, V_k) are defined by thinning the point process $\{(T'_k, V'_k)\}_{k: T'_k > T_{K_{\text{stop}}^* - 1}^{\text{stop}}}$ retaining only those points for which the corresponding node V'_k is still W , i.e. defined ℓ_k and ℓ_{k+1} as in (3.7) we set

$$(T_{k+1}, V_{k+1}) := (T'_{\ell_{k+1}}, V'_{\ell_{k+1}}).$$

Then we assign to V_{k+1} color B (regardless of the fact that V_{k+1} is B -susceptible or not). I.e., we define the processes N_B beyond $T_{K_{\text{stop}}^* - 1}^{\text{stop}}$ as

$$(T_{N_B[k]+1}^B, V_{N_B[k]+1}^B) := (T_{k+1}, V_{k+1}).$$

Of course the extension of the processes N_B and N described above has no effect on the evolution of the nodes' activation process up to time $T_{K_{\text{stop}}^* - 1}^{\text{stop}}$.

APPENDIX N: PROOFS OF THEOREMS 5.1, 5.2 AND 5.3- COMPLEMENTARY MATERIAL

N.1. Proof of Lemma 8.2. We prove the lemma reasoning by contradiction. Assume that there exists $\alpha > 0$ such that $\mathbb{P}(\limsup\{X_n > \alpha f_n\}) = \mathbb{P}(\bigcap_n \bigcup_{m \geq n} \{X_m > \alpha f_m\}) = \beta > 0$. Then

$$\liminf_{n \rightarrow \infty} \sum_{m \geq n} \mathbb{P}(X_m > \alpha f_m) \geq \lim_{n \rightarrow \infty} \mathbb{P}\left(\bigcup_{m \geq n} \{X_m > \alpha f_m\}\right) = \mathbb{P}\left(\bigcap_n \bigcup_{m \geq n} \{X_m > \alpha f_m\}\right) = \beta.$$

Therefore, we necessarily have

$$\sum_{n=0}^{\infty} \mathbb{P}(X_n > \alpha f_n) = \infty.$$

By the hypothesis about the stochastic ordering, it follows

$$\sum_{n=0}^{\infty} \mathbb{P}(Y_n > \alpha f_n) \geq \sum_{n=0}^{\infty} \mathbb{P}(X_n > \alpha f_n) = \infty.$$

By the Borel-Cantelli lemma, this implies $\mathbb{P}(\limsup\{Y_n > \alpha f_n\}) = 1$, which contradicts the assumption $Y_n/f_n \rightarrow 0$ a.s., as $n \rightarrow \infty$.

N.2. Proof of Theorem 5.1: Proof of $\psi > \tau$. Rewriting (J.2), we have:

$$\frac{g'_S(x)}{\beta_S(g_S(x))} = 1, \quad \forall x \in \mathcal{D}_{g_S}, S = \{R, B\}.$$

Therefore, for every $x > 0$, $x \in \mathcal{D}_{g_R} \cap \mathcal{D}_{g_B}$, we have

$$\int_0^x \frac{g'_R(y)}{\beta_R(g_R(y))} dy = \int_0^x \frac{g'_B(y)}{\beta_B(g_B(y))} dy = x.$$

By a change of variables it follows

$$\int_{g_R(0)}^{g_R(x)} \frac{1}{\beta_R(z)} dz = \int_0^{g_R(x)} \frac{1}{\beta_R(z)} dz = \int_{g_B(0)}^{g_B(x)} \frac{1}{\beta_B(z)} dz = \int_0^{g_B(x)} \frac{1}{\beta_B(z)} dz = x.$$

Recalling the definition of $\kappa_{\mathbf{g}}$ and its properties stated in Proposition A.2(ii) (or Proposition A.3), we have

$$\kappa_{\mathbf{g}} = \int_0^{\infty} \frac{dv}{\beta_R(z)} < \infty,$$

and $g_R(x) \uparrow \infty$ for $x \uparrow \kappa_{\mathbf{g}}$ and $g_B(\kappa_{\mathbf{g}}) < \infty$, which implies $\mathcal{D}_{g_R} \cap \mathcal{D}_{g_B} = [0, \kappa_{\mathbf{g}})$. Therefore for any $\kappa'_{\mathbf{g}} < \kappa_{\mathbf{g}}$

$$\int_{g_R(0)}^{g_R(\kappa'_{\mathbf{g}})} \frac{1}{\beta_R(z)} dz = \int_0^{g_B(\kappa'_{\mathbf{g}})} \frac{1}{\beta_B(z)} dz = \kappa'_{\mathbf{g}}$$

and so letting $\kappa'_{\mathbf{g}} \uparrow \kappa_{\mathbf{g}}$ we have

$$\int_0^{\infty} \frac{1}{\beta_R(z)} dz = \int_0^{g_B(\kappa_{\mathbf{g}})} \frac{1}{\beta_B(z)} dz = \kappa_{\mathbf{g}}.$$

The claim follows noticing that by the positiveness of $\beta_S(\cdot)$ we have

$$\tau := \int_0^{x_0} \frac{1}{\sum_S \beta_S(z)} dz < \int_0^{x_0} \frac{1}{\beta_R(z)} dz < \kappa_{\mathbf{g}} = \int_0^{g_B(\kappa_{\mathbf{g}})} \frac{1}{\beta_B(z)} dz < \int_0^{g_B(\kappa_{\mathbf{g}})+\varepsilon} \frac{1}{\beta_B(z)} dz = \psi.$$

N.3. Proof of (8.35). For $i < \bar{i}$, we have

$$2^{i+1}h_0 \leq 4^i k_0 < 8^{-1} \lfloor cn \rfloor, \text{ for all } n \text{ large enough.}$$

Therefore by (8.36) we have

$$N(Z_{i+1}) = N_R(Z_{i+1}) + N_B(Z_{i+1}) \leq 4^{i+1}k_0 < \lfloor cn \rfloor.$$

This proves the lower bound in (8.35). For the upper bound, note that for $i \geq \bar{i}$, we have $N(Z_{i+1}) = \lfloor cn \rfloor$, which follows by (8.33), and the fact that $4^{\bar{i}+1}k_0 > \lfloor cn \rfloor$ and $2^{\bar{i}+1}h_0 > \lfloor cn \rfloor$.

APPENDIX O: PROOF OF PROPOSITIONS 3.4 AND 3.5

O.1. Proof of Proposition 3.4. Note that by construction

$$(O.1) \quad \{U_k^s = u\} = \cup_{\mathbf{i} \in \mathcal{X}_u} \{\mathbf{X}_k = \mathbf{i}\} \rightarrow \mathbb{P}(U_k^s = u) = \sum_{\mathbf{i} \in \mathcal{X}_u} \mathbb{P}(\mathbf{X}_k = \mathbf{i})$$

Now, if $|\mathcal{S}_u| = 1$, the claim follows immediately from markovianity. Otherwise, first observe from Proposition 3.3 we immediately get:

$$(O.2) \quad \mathbb{P}(M_k^R = 1 \mid \mathbf{X}_k = \mathbf{i}) = u \quad \forall \mathbf{i} \in \mathcal{X}_u$$

Now, for $j \in \{0, 1\}$ we have:

$$\begin{aligned} \mathbb{P}(M_{k+1}^R = 1, M_h^R = j \mid U_k^R = u) &= \frac{\mathbb{P}(M_{k+1}^R = 1, M_h^R = j, U_k^R = u)}{\mathbb{P}(U_k^R = u)} \\ &= \sum_{\mathbf{i} \in \mathcal{X}_u} \frac{\mathbb{P}(M_{k+1}^R = 1, M_h^R = j, U_k^R = u, \mathbf{X}_k = \mathbf{i})}{\mathbb{P}(U_k^R = u)} \stackrel{(a)}{=} \sum_{\mathbf{i} \in \mathcal{X}_u} \frac{\mathbb{P}(M_{k+1}^R = 1, M_h^R = j, \mathbf{X}_k = \mathbf{i})}{\mathbb{P}(U_k^R = u)} \\ &= \sum_{\mathbf{i} \in \mathcal{X}_u} \frac{\mathbb{P}(M_{k+1}^R = 1, M_h^R = j \mid \mathbf{X}_k = \mathbf{i}) \mathbb{P}(\mathbf{X}_k = \mathbf{i})}{\mathbb{P}(U_k^R = u)} \\ &\stackrel{(b)}{=} \sum_{\mathbf{i} \in \mathcal{X}_u} \frac{\mathbb{P}(M_{k+1}^R = 1 \mid \mathbf{X}_k = \mathbf{i}) \mathbb{P}(M_h^R = j \mid \mathbf{X}_k = \mathbf{i}) \mathbb{P}(\mathbf{X}_k = \mathbf{i})}{\mathbb{P}(U_k^R = u)} \\ &\stackrel{(c)}{=} u \sum_{\mathbf{i} \in \mathcal{X}_u} \frac{\mathbb{P}(M_h^R = j \mid \mathbf{X}_k = \mathbf{i}) \mathbb{P}(\mathbf{X}_k = \mathbf{i})}{\mathbb{P}(U_k^R = u)} = u \sum_{\mathbf{i} \in \mathcal{X}_u} \frac{\mathbb{P}(M_h^R = j, \mathbf{X}_k = \mathbf{i})}{\mathbb{P}(U_k^R = u)} \\ &= u \frac{\mathbb{P}(M_h^R = j, U_k^R = u)}{\mathbb{P}(U_k^R = u)} \stackrel{(d)}{=} \mathbb{P}(M_{k+1} = 1 \mid U_k^R = u) \mathbb{P}(M_h = j \mid U_k^R = u) \end{aligned}$$

where equation (a) holds because from (O.1) we have $\{\mathbf{X}_k = \mathbf{i}\} \subseteq \{U_k^R = u\} \forall \mathbf{i} \in \mathcal{X}_u$, (b) from markovianity, while equations (c) and (d) from (O.2). \square

O.2. Proof of Proposition 3.5. The proof of Proposition 3.5 is omitted as it is a rather immediate consequence of (3.5) and the identity

$$\bigcap_{0 \leq h \leq k} \{\mathbf{X}_h \in \hat{\mathcal{X}}_{m_h}\} = \bigcup_{\mathbf{x}_1 \in \hat{\mathcal{X}}_{m_1}, \dots, \mathbf{x}_k \in \hat{\mathcal{X}}_{m_k}} \bigcap_{0 \leq h \leq k} \{\mathbf{X}_h = \mathbf{x}_h\}.$$