

SINGULAR VELOCITY OF THE STOKES AND NAVIER-STOKES EQUATIONS NEAR BOUNDARY IN THE HALF SPACE

Tongkeun Chang ¹ and Kyungkeun Kang ²

ABSTRACT. Local behaviors near boundary are analyzed for solutions of the Stokes and Navier-Stokes equations in the half space with localized non-smooth boundary data. We construct solutions of Stokes equations whose velocity field is not bounded near boundary away from the support of boundary data, although velocity and gradient velocity of solutions are locally square integrable. This is an improvement compared to known results in the sense that velocity field is unbounded itself, since previously constructed solutions were bounded near boundary, although their normal derivatives are singular. We also establish singular solutions and their derivatives that do not belong to L^q_{loc} near boundary with $q > 1$. For such examples, there corresponding pressures turn out not to be locally integrable. Similar construction via a perturbation argument is available to the Navier-Stokes equations near boundary as well.

2020 *Mathematics Subject Classification.* primary 35Q30, secondary 35B65.

Keywords and phrases: Singular velocity Stokes and Navier-Stokes equations, local regularity near boundary

1. INTRODUCTION

In this paper, we study local regularity of the Stokes and the Navier-Stokes equations near flat boundary. Let $B_r^+ := \{x = (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R} : |x| < r, x_n > 0\}$, $n \geq 2$. We consider the Stokes system near local boundaries.

$$(1.1) \quad u_t - \Delta u + \nabla \pi = 0 \quad \operatorname{div} u = 0 \quad \text{in } Q_2^+ := B_2^+ \times (0, 4),$$

where u indicates the velocity of the fluid and π is the associated pressure. We emphasize that no-slip boundary condition is given only on the flat boundary, i.e.

$$(1.2) \quad u = 0 \quad \text{on } \Sigma := (B_2 \cap \{x_n = 0\}) \times (0, 4).$$

Similar situation can be examined for the Navier-Stokes equations, namely

$$(1.3) \quad u_t - \Delta u + (u \cdot \nabla)u + \nabla \pi = 0, \quad \operatorname{div} u = 0 \quad \text{in } Q_2^+$$

together with the boundary condition (1.2).

Unlike the heat equation, due to non-local effect of incompressible fluid flow, we cannot expect local smoothing property for (1.1) or (1.3) near boundary. Indeed, the first author constructed a weak solution of (1.1) whose normal derivatives near boundary are unbounded. To be more precise, the singular solution, denoted by u , constructed in [11] is locally bounded weak solutions in Q_1^+ , namely,

$$(1.4) \quad \|u\|_{L^\infty(Q_1^+)} + \|u\|_{L^2(Q_1^+)} < \infty, \quad \|u\|_{L^\infty(Q_1^+)} = \infty.$$

In [12], such construction was extended to the Navier-Stokes equations as well and various types of blow-up rates were established near boundary. We remark that constructed singular solutions in [11] and [12] are due to nonlocal effect of solutions in the half-space caused by compactly supported singular boundary data.

Meanwhile, in [15], Seregin and Šverák constructed a Hölder continuous solution up to boundary in the form of shear flow whose normal derivatives are, however, unbounded near boundary in the half-space. It is remarkable that such a solution satisfies homogeneous initial and boundary conditions and the solution, however, is not of finite energy in the half space in contrast to ones constructed in [11] and [12].

¹Department of Mathematics, Yonsei University Seoul, 03722, South Korea, chang7357@yonsei.ac.kr

²Department of Mathematics, Yonsei University Seoul, 03722, South Korea, kkang@yonsei.ac.kr

On the other hand, a certain type of local maximal regularity was proved near boundary via a priori estimate. More precisely, in [14, Proposition 2] it was shown in three dimensions that for given $p, q \in (1, 2]$ the solution of the Stokes equations (1.1)-(1.2) satisfies the following a priori estimate: For any r with $p \leq r < \infty$

$$(1.5) \quad \|u_t\|_{L_t^q L_x^r(Q_{\frac{1}{4}}^+)} + \|\nabla^2 u\|_{L_t^q L_x^r(Q_{\frac{1}{4}}^+)} + \|\nabla \pi\|_{L_t^q L_x^r(Q_{\frac{1}{4}}^+)} \leq C \left(\|u\|_{L_t^q W_x^{1,p}(Q_{\frac{1}{4}}^+)} + \|\pi\|_{L_t^q L_x^p(Q_{\frac{1}{4}}^+)} \right).$$

Furthermore, due to parabolic embedding, it follows that

$$(1.6) \quad \|u\|_{C_t^{\frac{\alpha}{2}} C_x^\alpha(Q_{\frac{1}{4}}^+)} < \infty, \quad 0 < \alpha < 2 - \frac{2}{q}.$$

It is worth to mention that the estimates (1.5) is optimal, since the authors constructed in [8] an example that satisfies (1.5) but any higher integrability of the example fails (see also [12]).

We remark very recently that local boundary regularity of solutions was improved for the Stokes system, when boundary condition is given as Navier-boundary condition, not no-slip boundary condition (see [10]).

So far, singular solutions that has been constructed in above references are bounded, in fact, Hölder continuous but the normal derivatives of velocity are singular near boundary in the half space. One may wonder whether or not there exists a singular solution that is unbounded itself near boundary. Our main objective is to construct a weak solution of the Stokes and Navier-Stokes equations such that velocity field is singular near boundary in the half space (the definition of weak solutions of Stokes equations is given in section 3). Furthermore, we are also interested in establishing an example so that velocity field is bounded but it does not belong to C^α , class of Hölder continuous functions, with any exponent $\alpha > 0$ near boundary. We remark that since we are looking for solutions that are not Hölder continuous functions itself, due to above a priori estimate (1.5), the associated pressure is unlikely to belong to L_{loc}^p , $p > 1$, which turns out to be the case in examples of our construction. Our main results will be precisely stated in the next section.

2. MAIN RESULTS

For convenience, we introduce a tensor L_{ij} defined by

$$(2.1) \quad L_{ij}(x, t) = D_{x_j} \int_0^{x_n} \int_{\mathbb{R}^{n-1}} D_{z_n} \Gamma(z, t) D_{x_i} N(x - z) dz, \quad i, j = 1, 2, \dots, n,$$

where Γ and N are the n dimensional Gaussian kernel and Newtonian kernel defined by

$$(2.2) \quad \Gamma(x, t) = \begin{cases} \frac{1}{(4\pi t)^{\frac{n}{2}}} e^{-\frac{|x|^2}{4t}} & t > 0, \\ 0 & t < 0, \end{cases} \quad N(x) = \begin{cases} -\frac{1}{n(n-2)\omega_n} |x|^{-n+2} & n \geq 3, \\ \frac{1}{2\pi} \ln |x| & n = 2. \end{cases}$$

It is known that L_{ij} satisfies following relations:

$$(2.3) \quad \sum_{i=1}^n L_{ii} = \frac{1}{2} D_{x_n} \Gamma, \quad L_{ij} = L_{ji} \quad 1 \leq i, j \leq n-1, \quad L_{in} = L_{ni} + B_{in} \quad \text{if } i \neq n,$$

where

$$(2.4) \quad B_{in}(x, t) = \int_{\mathbb{R}^{n-1}} D_{x_n} \Gamma(x' - z', x_n, t) D_{z_i} N(z', 0) dz'.$$

Furthermore, we remind an estimate of L_{ij} defined in (2.1) (see (2.6) in [16])

$$(2.5) \quad |D_{x_n}^{l_0} D_{x'}^{k_0} L_{ij}(x, t)| \leq \frac{c}{t^{\frac{1}{2}} (|x|^2 + t)^{\frac{1}{2}n + \frac{1}{2}k_0} (x_n^2 + t)^{\frac{1}{2}l_0}},$$

where $1 \leq i, j \leq n$ except for the case that $i \neq n$ and $j = n$.

We recall the following Stokes equations in a half space:

$$(2.6) \quad \begin{aligned} w_t - \Delta w + \nabla p &= 0, & \operatorname{div} w &= 0 & \text{in } \mathbb{R}_+^n \times (0, \infty), \\ w|_{t=0} &= 0, & w|_{x_n=0} &= g. \end{aligned}$$

It is known that w is represented by

$$(2.7) \quad w_i(x, t) = \sum_{j=1}^n \int_0^t \int_{\mathbb{R}^{n-1}} K_{ij}(x' - y', x_n, t - s) g_j(y', s) dy' ds,$$

where the Poisson kernel K of the Stokes equations in $\mathbb{R}_+^n \times (0, \infty)$ is given as follows (see [16]):

$$(2.8) \quad \begin{aligned} K_{ij}(x' - y', x_n, t) &= -2\delta_{ij} D_{x_n} \Gamma(x' - y', x_n, t) + 4L_{ij}(x' - y', x_n, t) \\ &\quad + 2\delta_{jn} \delta(t) D_{x_i} N(x' - y', x_n). \end{aligned}$$

The formula of the associated pressure p is also given as

$$(2.9) \quad \begin{aligned} p(x, t) &= - \sum_{k=1}^{k=n} \int_{\mathbb{R}^{n-1}} D_{x_k} D_{x_n} N(x' - y', x_n) g_k(y', t) dy' \\ &\quad + 2 \sum_{k=1}^{k=n} (D_t - \Delta') \int_0^t \int_{\mathbb{R}^{n-1}} g_k(z', s) \int_{\mathbb{R}^{n-1}} D_{y_k} N(y', x_n) \Gamma(x' - y' - z', 0, t - s) dy' dz' ds \\ &\quad + N *' D_t g_n(x, t), \end{aligned}$$

We specify a boundary data $g : \mathbb{R}^{n-1} \times \mathbb{R}_+ \rightarrow \mathbb{R}^n$ with only non-zero n -th component defined as follows:

$$(2.10) \quad g(y', s) = \alpha(0, \dots, 0, g_n(y', s)) = \alpha(0, \dots, 0, g_n^S(y') g_n^T(s)),$$

where $\alpha > 0$ is a parameter, which will be taken sufficiently small for the case of the Navier-Stokes equations. Here $g_n^S \geq 0$ and $g_n^T \geq 0$ are assumed to satisfy

$$(2.11) \quad g_n^S \in C_c^\infty(A), \quad \text{supp } g_n^T \subset (0, 1), \quad g_n^T \in L^1(\mathbb{R}) \cap C^\infty([0, 1]),$$

where the set $A \subset \mathbb{R}^{n-1}$ is defined by

$$(2.12) \quad A = \{y' = (y_1, y_2, \dots, y_{n-1}) \in \mathbb{R}^{n-1} \mid 1 < |y'| < 2, -2 < y_i < -1, 1 \leq i \leq n-1\}.$$

Here we consider the case of non-smooth g_n^T , which is supposed to be three kinds of prototype types. More precisely, we assume the following for g_n^T :

Assumption 2.1. We suppose that $g_n^T(s) = \phi(1 - s)$, where ϕ satisfies

$$(2.13) \quad \phi \geq 0, \quad \text{supp } \phi \subset (0, 1), \quad \phi \in L^1(\mathbb{R}) \cap C^\infty((0, 1]), \quad \phi(1) = 0.$$

Assume further that either

$$(2.14) \quad \phi(s) = |\ln s| \quad \text{or} \quad \phi(s) = |\ln s|^{-1}, \quad s \in (0, \frac{1}{2}]$$

or ϕ satisfies

$$(2.15) \quad \phi^{(k)}(s) \approx \phi^{(k)}(2s), \quad s \in (0, \frac{1}{4}], \quad k = 0, 1.$$

Here, $A \approx B$ means that there exists a constant $c > 1$ such that $c^{-1}B \leq A \leq cB$. Furthermore, we also assume that

$$(2.16) \quad \int_\epsilon^1 s^{-\alpha} \phi(s) ds \approx \max(1, \epsilon^{1-\alpha} \phi(\epsilon)), \quad 0 < \alpha, \quad \alpha \neq 1.$$

$$(2.17) \quad \int_\epsilon^1 s^{-\alpha} \phi'(s) ds \approx \epsilon^{1-\alpha} \phi'(\epsilon), \quad 0 < \alpha, \quad \alpha \neq 1.$$

$$(2.18) \quad \int_0^\epsilon \phi(s) ds \approx \epsilon \phi(\epsilon).$$

Some comments are prepared for the Assumption 2.1.

(i) In case that $\frac{3}{4} < t < 1$, if ϕ satisfies (2.15), then one can see that

$$(2.19) \quad \phi^{(k)}(t-s) \approx \phi^{(k)}(1-s) \quad \text{for} \quad \frac{1}{2} < s < 2t-1 \quad k=0,1,$$

$$(2.20) \quad \phi^{(k)}(1-t) \approx \phi^{(k)}(1-s) \quad \text{for} \quad 2t-1 < s < t \quad k=0,1,$$

since $\frac{1}{2}(1-s) \leq t-s \leq (1-s)$ for $\frac{1}{2} < s < 2t-1$ and $\frac{1}{2}(1-s) \leq 1-t \leq (1-s)$ for $2t-1 < s < t$. Similarly, if $1 < t < \frac{5}{4}$, it is straightforward that

$$(2.21) \quad \phi^{(k)}(t-s) \approx \phi^{(k)}(1-s) \quad \text{for} \quad \frac{3}{4} < s < 2-t \quad k=0,1,$$

$$(2.22) \quad \phi^{(k)}(t-1) \approx \phi^{(k)}(t-s) \quad \text{for} \quad 2-t < s < 1 \quad k=0,1.$$

This is also due to facts that $(1-s) \leq t-s \leq 2(1-s)$ and $\frac{1}{2}(t-s) \leq t-1 \leq (t-s)$ for $0 < s < 2-t$ and $2-t < s < 1$, respectively.

(ii) One can check that $\phi(s) = |\ln s|$ or $\phi(s) = |\ln s|^{-1}$ satisfy (2.16)-(2.18). We note that $\phi(t) = t^{-a}$ with $a \in (0, 1)$, $a + \alpha \neq 1$ is another typical type that we will treat in this paper.

(iii) We note that $\phi(t) = t^{-a}\eta(t)$ is a prototype in Assumption 2.1, where η is cut off function with $\eta(t) = 1$ for $t \in (0, \frac{1}{2})$ and $\eta(t) = 0$ for $t \geq \frac{3}{4}$.

As a notational convention, we denote $\text{sgn}(a) = 1$ if $a > 0$ and $\text{sgn}(a) = -1$ if $a < 0$.

Next lemma, some elementary properties of Φ are stated and, for clarity, its proof will be given in section 3.

Lemma 2.2. *Let $0 < t < \frac{1}{4}$ and ϕ satisfy the Assumption 2.1. Then, for $0 < x_n < \frac{1}{4}$ and $\sqrt{t} < x_n$,*

$$(2.23) \quad \phi(t)e^{-\delta_1 \frac{x_n^2}{t}} \leq c\phi(x_n^2), \quad \text{sgn}(\phi')\phi'(t)e^{-\delta_1 \frac{x_n^2}{t}} \leq c\text{sgn}(\phi')\phi'(x_n^2).$$

In the case $\phi = t^{-a}$, there is a large constant $\epsilon_1 \gg 1$ such that if $\epsilon_1\sqrt{t} < x_n$, then there exists there a sufficiently small ϵ depending on ϵ_1 such that

$$(2.24) \quad t^{-\frac{1}{2}}e^{-\delta_1 \frac{x_n^2}{t}}\phi(t) \leq \epsilon x_n \text{sgn}(\phi')\phi'(x_n^2).$$

From (2.7), (2.8), and (2.3)₃, for $1 \leq i \leq n$, we have

$$(2.25) \quad \begin{aligned} w_i(x, t) &= 4 \int_0^t \int_{\mathbb{R}^{n-1}} L_{ni}(x' - y', x_n, t-s) g_n(y', s) dy' ds \\ &\quad + 4(1 - \delta_{in}) \int_0^t \int_{\mathbb{R}^{n-1}} B_{in}(x' - y', x_n, t-s) g_n(y', s) dy' ds \\ &\quad + 2 \int_{\mathbb{R}^{n-1}} D_{x_i} N(x' - y', x_n) g_n(y', t) dy' \\ &:= w_i^L(x, t) + w_i^B(x, t) + w_i^N(x, t). \end{aligned}$$

Form now on, for simplicity, we denote by Γ' the $n-1$ dimensional Gaussian kernel.

Definition 2.3. *As a notational convention, we denote $A \vee B := \max\{A, B\}$. Further, for a positive function g , we say that $f \approx \pm e^{-\delta g}$ if $c^{-1}e^{-g(x)} \leq |f(x)| \leq ce^{-\frac{1}{8}g(x)}$ for some constant $c > 0$.*

Lemma 2.4. *Let Γ' and N be $n-1$ dimensional Gaussian kernel and n dimensional Newtonian kernel defined in (2.2). Let $X' \in \mathbb{R}^{n-1}$ with $|X'| \geq 1$, $0 < x_n < \frac{1}{2}$ and $t > 0$. Then, there exists $c > 0$, independent of X' , x_n and t , such that*

$$(2.26) \quad \begin{aligned} \int_{\mathbb{R}^{n-1}} \Gamma'(X' - z', t) D_{x_n} N(z', x_n) dz' &= D_{x_n} N(X', x_n) + I(X', x_n, t) + J_1(X', x_n, t), \\ \int_{\mathbb{R}^{n-1}} \Gamma'(X' - z', t) D_{z_i} N(z', 0) dz' &= D_{X_i} N(X', 0) + J_2(X', t), \end{aligned}$$

where $i = 1, 2, \dots, n-1$ and I, J_1 and J_2 satisfy

$$(2.27) \quad I(x', x_n, t) \approx t^{-\frac{n-1}{2}} e^{-\delta \frac{|x'|^2}{t}}, \quad |J_1(X', x_n, t)| \leq c x_n t^{\frac{1}{2}}, \quad |J_2(X', t)| \leq c t^{\frac{1}{2}} \quad m \geq 1,$$

where " \approx " and δ are defined in Definition 2.3.

The proof of Lemma 2.4 will be provided in section 3.

Remark 2.5. Let $i = 1, 2, \dots, n-1$. Suppose that $x', y' \in \mathbb{R}^{n-1}$ with $|x'| < \frac{1}{2}$ and $y' \in A$ and, for convenience, we denote $X' = x' - y'$. Then, it is direct that $\frac{1}{2} < |X'| \leq \frac{5}{2}$, because $\frac{1}{2} \leq x_i - y_i$ and $|X'| \leq (|x'| + |y'|) \leq \frac{5}{2}$, which implies that there are positive constants c_1 and c_2 such that

$$c_1 \leq D_{x_i} N(x' - y', 0) \leq c_2, \quad c_1 x_n \leq D_{x_n} N(x' - y', x_n) \leq c_2 x_n.$$

Therefore, it follows from (2.26) and (2.27) that

$$(2.28) \quad \left| \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t) D_{z_i} N(z', 0) dz' \right| \leq c, \quad |x'| \leq \frac{1}{2}, \quad y' \in A,$$

$$(2.29) \quad \left| \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t) D_{x_n} N(z', x_n) dz' \right| \leq c \left(x_n + t^{-\frac{n-1}{2}} e^{-\frac{1}{8t}} \right), \quad |x'| \leq \frac{1}{2}, \quad y' \in A,$$

where $\frac{3}{4} < t < \frac{5}{4}$, $0 < s < t$ and $k \geq 0$. Furthermore, there exist $c_0 > 0$ and $t_0 \in (0, \frac{1}{4})$ such that if $0 < t - s < t_0$, then

$$(2.30) \quad c_0 \leq \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t - s) D_{z_i} N(z', 0) dz' \quad |x'| \leq \frac{1}{2}, \quad y' \in A,$$

$$(2.31) \quad c_0 (x_n + t^{-\frac{n-1}{2}} e^{-\frac{1}{t}}) \leq \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t - s) D_{x_n} N(z', x_n) dz', \quad |x'| \leq \frac{1}{2}, \quad y' \in A.$$

Indeed, the estimate (2.31) is due to the computations that

$$\begin{aligned} & \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t - s) D_{x_n} N(z', x_n) dz' \\ & \geq D_{x_n} N(X', x_n) + t^{-\frac{n-1}{2}} e^{-\frac{1}{t}} - |J_1(x', x_n, t - s)| \geq c (x_n + t^{-\frac{n-1}{2}} e^{-\frac{1}{t}}). \end{aligned}$$

In conclusion, due to (2.28)-(2.31), we obtain

$$(2.32) \quad \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t - s) D_{z_i} N(z', 0) dz' \approx c, \quad |x'| \leq \frac{1}{2}, \quad y' \in A,$$

$$(2.33) \quad \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t - s) D_{x_n} N(z', x_n) dz' \approx x_n + t^{-\frac{n-1}{2}} e^{-\frac{1}{t}}, \quad |x'| \leq \frac{1}{2}, \quad y' \in A.$$

□

We define for notational convention

$$\begin{aligned} \mathcal{D}_+ &= \left\{ (x, t) \in \mathbb{R}_+^n \times (0, \infty) : |t - 1|^{\frac{1}{2}} \geq x_n, |t - 1| < t_0, 0 < x_n < \frac{1}{2} \right\}, \\ \mathcal{D}_- &= \left\{ (x, t) \in \mathbb{R}_+^n \times (0, \infty) : |t - 1|^{\frac{1}{2}} < x_n, |t - 1| < t_0, 0 < x_n < \frac{1}{2} \right\}, \\ \mathcal{D}_0 &= \left\{ (x, t) \in \mathbb{R}_+^n \times (0, \infty) : \frac{1}{2} < \frac{x_n}{|t - 1|^{\frac{1}{2}}} < 1, |t - 1| < t_0, 0 < x_n < \frac{1}{2} \right\}. \end{aligned}$$

Here $t_0 > 0$ is given in Remark 2.5 and $t_1 < t_0$ will be determined later. We also introduce two different regions defined by

$$(2.34) \quad \begin{aligned} \tilde{\mathcal{D}}_+ &= \mathcal{D}_+ \cap \left\{ (x, t) : x_n \leq \epsilon_0 |t - 1|^{\frac{1}{2}} \right\}, \\ \tilde{\mathcal{D}}_- &= \mathcal{D}_- \cap \left\{ (x, t) : \epsilon_1 |t - 1|^{\frac{1}{2}} \leq x_n \right\}, \end{aligned}$$

where ϵ_0 and ϵ_1 be positive constants with $0 < \epsilon_0 \ll 1 \ll \epsilon_1 < \infty$, which will be specified later.

Theorem 2.6. *Let w be the solution of the Stokes equations (2.6) in the half-space with boundary data g given in (2.10) and (2.11). Suppose that ϕ satisfies Assumption 2.1. Then, there is $0 < t_1 < t_0$ such that if $|t - 1| < t_1$, $a \neq \frac{1}{2}$, $|x'| < \frac{1}{2}$ and $0 < x_n < \frac{1}{2}$ such that we prove the following: Let $i = 1, 2, \dots, n - 1$.*

$$(2.35) \quad |w_i(x, t)| \lesssim \begin{cases} 1 & \text{if } \phi(t) = |\ln t|^{-1} \\ \phi(|1 - t|) & \text{otherwise.} \end{cases}$$

In case that $\phi(t) \neq |\ln t|^{-1}$, there exist ϵ_0 and ϵ_1 , depending on t_1 , in (2.34) such that

$$(2.36) \quad w_i(x, t) \geq c\phi(1 - t) \quad \text{if } (x, t) \in \tilde{\mathcal{D}}_-, \quad t < 1.$$

$$(2.37) \quad w_i(x, t) \leq -c\phi(t - 1) \quad \text{if } (x, t) \in \tilde{\mathcal{D}}_- \cup \mathcal{D}_0, \quad 1 < t < 1 + t_0.$$

$$(2.38) \quad |D_{x_n} w_i(x, t)| \leq c \begin{cases} ((t - 1) \vee x_n^2)^{-\frac{1}{2}} \phi((t - 1) \vee x_n^2), & \text{if } 1 < t < \frac{5}{4}, \\ ((1 - t) \vee x_n^2)^{\frac{1}{2}} |\phi'((1 - t) \vee x_n^2)|, & \text{if } \frac{3}{4} < t < 1. \end{cases}$$

$$(2.39) \quad D_{x_n} w_i(x, t) \approx -(1 - t)^{\frac{1}{2}} \phi'(1 - t) \mathbf{I}_{\tilde{\mathcal{D}}_+} - x_n \phi'(x_n^2) \mathbf{I}_{\tilde{\mathcal{D}}_-}, \quad 1 - t_1 < t < 1.$$

$$(2.40) \quad D_{x_n} w_i(x, t) \approx -(t - 1)^{-\frac{1}{2}} \phi(t - 1) \quad \text{if } (x, t) \in \tilde{\mathcal{D}}_+, \quad 1 < t < 1 + t_0.$$

If $1 < t < 1 + t_1$ and $(x, t) \in \tilde{\mathcal{D}}_-$ then

$$(2.41) \quad D_{x_n} w_i(x, t) \approx \begin{cases} -x_n \phi'(x_n^2), & \text{if } \phi = t^{-a}, \\ -x_n \phi'(x_n^2) + (t - 1)^{-\frac{3}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}} \phi(t - 1), & \text{otherwise.} \end{cases}$$

$$(2.42) \quad p(x, t) \approx \begin{cases} -\phi'(1 - t), & \text{if } 1 - t_1 < t < 1, \\ x_n (t - 1)^{-\frac{1}{2}} \phi(t - 1) \pm 1, & \text{if } 1 < t < 1 + t_1. \end{cases}$$

Remark 2.7. *The results of Theorem 2.6 imply the following regularity of the solution for $|t - 1| < t_1$ and $|x| < \frac{1}{2}$ near boundary: Let $Q := B_{\frac{1}{2}}^+ \times (1 - t_1, 1 + t_1)$ and $i = 1, 2, \dots, n - 1$.*

(1) *(The case $\phi(t) = |\ln(t)|$) For any $p_1 < \infty$ and $p_2 < 3$*

$$(2.43) \quad \begin{aligned} w_i &\in L^{p_1}(Q), \quad D_{x_n} w_i \in L^{p_2}(Q), \quad p \in L_{weak}^1(Q) \quad \text{but} \\ w_i &\notin L^\infty(Q), \quad D_{x_n} w_i \notin L^3(Q), \quad p \notin L^1(Q). \end{aligned}$$

(2) *(The case $\phi(t) = |\ln(t)|^{-1}$) For any $p_3 > 3$, $p_4 > 1$ and $\alpha \in (0, 1)$*

$$(2.44) \quad \begin{aligned} w_i &\in L^\infty(Q), \quad D_{x_n} w_i \in L^3(Q), \quad p \in L^1(Q), \quad \text{but} \\ w_i &\notin C^\alpha(Q), \quad D_{x_n} w_i \notin L^{p_3}(Q), \quad p \notin L^{p_4}(Q). \end{aligned}$$

(3) *(The case $\phi(t) = t^{-a}$, $0 < a < 1$). For any $p_5 < \frac{3}{2a}$, $p_6 < \frac{3}{1+2a}$ and $p_7 < \frac{1}{1+a}$*

$$(2.45) \quad \begin{aligned} w_i &\in L^{p_5}(Q), \quad D_{x_n} w_i \in L^{p_6}(Q), \quad p \in L^{p_7}(Q), \quad \text{but} \\ w_i &\notin L^{\frac{3}{2a}}(Q), \quad D_{x_n} w_i \notin L^{\frac{3}{1+2a}}(Q), \quad p \notin L^{\frac{1}{1+a}}(Q). \end{aligned}$$

The case that $\phi(t) = t^{-a}$, $0 < a < 1$ is most interesting and, combining Theorem 2.6 and Remark 2.7, we make a table to show asymptotic behavior of the solution near boundary.

	Region	w_i	$D_{x_n} w_i$	p
$t < 1$	\tilde{D}_+		$\approx -(1-t)^{\frac{1}{2}} \phi'(1-t) \approx (1-t)^{-\frac{1}{2}-a}$	$\approx -\phi'(1-t) \approx (1-t)^{-1-a}$
	\tilde{D}_-	$\approx \phi(1-t)$	$\approx -x_n \phi'(x_n^2) \approx x_n^{-1-2a}$	
$t > 1$	\tilde{D}_+		$\approx -(t-1)^{-\frac{1}{2}} \phi(t-1) \approx -(t-1)^{-\frac{1}{2}-a}$	$\approx -x_n(t-1)^{-\frac{1}{2}} \phi(t-1) \approx (t-1)^{-a}$
	\tilde{D}_-	$\approx \phi(t-1)$	$\approx -x_n \phi'(x_n^2) \approx x_n^{-1-2a}$	
	\tilde{D}_0	$\approx \phi(t-1)$		
Integrable property		$\in L^p, p < \frac{3}{2a}$ but $\notin L^{\frac{3}{2a}}$	$\in L^p, p < \frac{3}{1+2a}$ but $\notin L^{\frac{3}{1+2a}}$	$\in L^p, p < \frac{1}{1+a}$ but $\notin L^{\frac{1}{1+a}}$

$$(\phi(t) = t^{-a}, 0 < a < 1)$$

Remark 2.8. When it comes to global estimate of the solution w in the half-space, it is worth mentioning that w belongs to the same spaces as shown in Remark 2.7. For example, in the case $\phi = |\ln t|^{-1}$, since the boundary data $g_n(t)$ in (2.10) is Dini-continuity for all $0 < t < \infty$, it follows due to [1] that w is bounded in the half space, i.e. $w \in L^\infty(\mathbb{R}_+^n \times (0, \infty))$ (see also [2] and [3]). Moreover, since $\phi \in L^\infty \cap \dot{B}_3^{\frac{1}{3}}$, using the results in [4] and [5], one can see that

$$(2.46) \quad \|\nabla w\|_{L^3(\mathbb{R}_+^n \times (0, \infty))} \leq c \left(\|\phi\|_{L^3(0, \infty)} + \|\phi\|_{\dot{B}_3^{\frac{1}{3}}(0, \infty)} \right) < \infty.$$

The other cases such as $\phi(t) = |\ln t|$ or t^{-a} can be treated similarly, and thus we skip their details. Our results can be understood that temporal singular behaviors caused by compactly supported boundary data are immediately transferred to all over the place and, in addition, unlike the interior local smoothing in spatial variables is not available near boundary. We remark that spatial singular behavior of compactly supported boundary data is not delivered to other places and local smoothing effect is available in any region away from the data (see [13]).

Remark 2.9. We remark that authors showed in [7] that there is a very weak solution of (1.1) or (1.3) so that ∇w is not square integrable, i. e., $\|\nabla w\|_{L^2(Q_1^+)} = \infty$, although $\|w\|_{L^2(Q_1^+)} < \infty$. It turns out that it is a special case almost close to $a = \frac{1}{4}$ and thus it is observable that w is not bounded and π is not integrable in Q_1^+ .

Next, we will establish existence of singular solutions of the Navier-Stokes equations as in the Stokes system. When it comes to the boundary data for temporal variable, we consider only a specific case in the Assumption 2.1, which satisfies the following:

$$(2.47) \quad \phi(t) = t^{-a}, \quad t \in (0, \frac{1}{2}], \quad a \in (0, 1),$$

since we think that it is most interesting. The other cases such as $\phi(t) = |\ln s|$ or $\phi(s) = |\ln s|^{-1}$ could be treated similarly, and thus we omit the details of those cases (see Remark 2.13).

Let $\varphi_1 \in C_c^\infty(\mathbb{R}^n)$ be a cut-off function satisfying $\varphi_1 \geq 0$, $\text{supp } \varphi_1 \subset B_{\frac{1}{\sqrt{2}}}$ and $\varphi_1 \equiv 1$ in $B_{\frac{3}{8}}$. Also, let $\varphi_2 \in C_c^\infty(-\infty, \infty)$ be a cut-off function satisfying $\varphi_2 \geq 0$, $\text{supp } \varphi_2 \subset (-3, 3)$ and $\varphi_2 \equiv 1$ in $(-2, 2)$. Let $\varphi(x, t) = \varphi_1(x)\varphi_2(t)$. Let $U = w\varphi$ and $\Pi = \pi\varphi$ such that $U|_{Q_{\frac{1}{4}}^+} = w$ and $\Pi|_{Q_{\frac{1}{4}}^+} = \pi$.

We consider the following perturbed Navier-Stokes equations in $\mathbb{R}_+^n \times (0, \infty)$:

$$(2.48) \quad v_t - \Delta v + \nabla q + \text{div}(v \otimes v + v \otimes U + U \otimes v) = -\text{div}(U \otimes U), \quad \text{div } v = 0$$

with homogeneous initial and boundary data, i.e.

$$(2.49) \quad v(x, 0) = 0 \text{ in } \mathbb{R}_+^n, \quad v(x, t) = 0 \text{ on } \{x_n = 0\} \times [0, \infty).$$

Proposition 2.10. *Suppose that ϕ satisfies Assumption 2.1 and (2.47). Let $0 < a < \frac{3}{2(n+2)}$ and $\frac{n+2}{2} < q < \frac{3}{4a}$. Then, we have $\|U\|_{L^{2q}} < c\alpha$, where $\alpha > 0$ is defined in (2.10). If α is taken to be sufficiently small, then there is a solution v of (2.48)-(2.49) such that*

$$(2.50) \quad \|\nabla v\|_{L^q(\mathbb{R}_+^n \times (0,1))} + \|v\|_{L^{2q}(\mathbb{R}_+^n \times (0,1))} \leq c(\|w\|_{L^q(Q_2^+)}, \|w\|_{L^{2q}(Q_2^+)}).$$

In case that $0 < a < \frac{3}{4(n+2)}$, we obtain $\|w\|_{L^{2q}} < c\alpha$ for $n+2 < q < \frac{3}{4a}$. If α is taken to be sufficiently small, then v satisfies

$$(2.51) \quad \|v\|_{L^\infty(\mathbb{R}_+^n \times (0,1))} \leq c(\|w\|_{L^q(Q_2^+)}, \|w\|_{L^{2q}(Q_2^+)}).$$

The proof of Proposition 2.10 will be given in section 5.

We then set $u := v + U$ and $p = \pi + q$, which becomes a weak solution of the Navier-Stokes equations in $\tilde{Q}_+ = B_{\frac{1}{2}}^+ \times (0, 2)$, namely

$$(2.52) \quad u_t - \Delta u + \nabla p = -\text{div}(u \otimes u), \quad \text{div } u = 0 \quad Q_1^+,$$

with boundary data $u(x, t) = 0$ on $\{x_n = 0\}$ and homogeneous initial data $u(x, 0) = 0$.

With the aid of Proposition 2.10, we can construct a solution of the Navier-Stokes equations such that its behaviors near boundary are more or less the same as those of the Stokes equations mentioned in Theorem 2.6. Since its verification is straightforward by combining Theorem 2.6 and Proposition 2.10, we just state it without providing its details.

Theorem 2.11. *Suppose that ϕ satisfies Assumption 2.1 and (2.47) with an additional restriction that $0 < a < \frac{3}{2(n+2)}$ for $\phi(t) = t^{-a}$. Then, there exists a solution of the Navier-Stokes equations, u , in (2.52) such that it satisfies the same regularity properties mentioned in Remark 2.7. Furthermore, if $0 < a < \frac{3}{4(n+2)}$, then u satisfies pointwise estimates (2.35), (2.36) and (2.37).*

Remark 2.12. *The pointwise estimate of u in Theorem 2.11 is the combination of (2.51) for v in Proposition 2.10 and (2.35)-(2.37) for w in Theorem 2.6. However, the pointwise estimate of ∇u is not clear because of the lack of that of ∇v , and instead, L^q estimate is only available up to $q < \frac{3}{4a}$ as shown in (2.50). Thus, we leave it as an open question on construction of solutions of the Navier-Stokes equations whose normal derivatives satisfies pointwise estimates near boundary given in Theorem 2.6.*

Remark 2.13. *In the case that $\phi(t) = |\ln t|$ or $\phi(s) = |\ln s|^{-1}$, the estimate (2.50) is available for any exponent $q < \infty$. Thus, it is also available to construct a singular solution of the Navier-Stokes equations with the same regularity properties as the Stokes system mentioned in Remark 2.7.*

This paper is organized as follows: In section 3, proofs of some lemmas are provided. Section 4 is devoted to proving Theorem 2.6. We present the proof of Proposition 2.10 in Section 5.

3. PRELIMINARIES

In this section, we mainly provide proofs of lemmas, which are useful for our analysis. Before we do that, we first define the notion of weak solutions of the Stokes equations in the half space.

Definition 3.1. *Let ϕ satisfy Assumption 2.1. We say that $w \in L^1_{\text{loc}}(0, \infty; W^{1,1}_{\text{loc}}(\mathbb{R}_+^n))$ is a weak solution of the Stokes equations (2.6) in the half-space with boundary data g given in (2.10) and (2.11), if the following equality holds:*

$$\int_0^T \int_{\mathbb{R}_+^n} w \cdot \Phi_t dx dt + \int_0^T \int_{\mathbb{R}_+^n} w \cdot \Delta \Phi dx dt + \int_0^T \int_{\partial \mathbb{R}_+^n} g_n \cdot D_{x_n} \Phi dx' dt = 0$$

for each $\Phi \in C_c^2(\overline{\mathbb{R}_+^n} \times [0, T])$ with

$$(3.1) \quad \operatorname{div} \Phi = 0, \quad \Phi|_{\partial \mathbb{R}_+^n \times (0, T)} = 0, \quad \Phi(\cdot, T) = 0.$$

In addition, for each $\Psi \in C_c^1(\overline{\mathbb{R}_+^n})$ with $\Psi|_{\partial \mathbb{R}_+^n} = 0$

$$(3.2) \quad \int_{\mathbb{R}_+^n} u(x, t) \cdot \nabla \Psi(x) dx = 0 \quad \text{for all } 0 < t < T.$$

For notational convention, we denote for fixed $\alpha \geq 0$ and $k = 0, 1$

$$(3.3) \quad \Psi_{\phi, k}(t) := \int_0^{\min(1, t)} (t-s)^{-\alpha} \phi^{(k)}(1-s) ds,$$

$$(3.4) \quad \tilde{\Psi}_{\phi, k}(t) := \int_0^{\min(1, t)} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi^{(k)}(1-s) ds.$$

The following is a key lemma to obtain main results.

Lemma 3.2. *Let $0 < \alpha$, $\alpha \neq 1$, $\alpha + a \neq 1$ and $\frac{3}{4} < t < \frac{5}{4}$. Suppose that ϕ satisfies Assumption 2.1. Then, $\Psi_{\phi, k}(t)$ and $\tilde{\Psi}_{\phi, k}(t)$ in (3.3)-(3.4) satisfy following:*

(i) *Let $1 < t < \frac{5}{4}$. Then,*

$$(3.5) \quad \Psi_{\phi, 0}(t) \approx \begin{cases} (t-1)^{1-\alpha} \phi(t-1) & \text{if } \alpha + a > 1, \\ 1 & \text{otherwise.} \end{cases}$$

$$(3.6) \quad \Psi_{\phi, 1}(t) \approx (t-1)^{1-\alpha} \phi'(t-1).$$

$$(3.7) \quad \tilde{\Psi}_{\phi, 0}(t) \approx \begin{cases} ((t-1) \vee x_n^2)^{1-\alpha} \phi((t-1) \vee x_n^2) & \text{if } \alpha + a > 1, \\ 1 & \text{otherwise.} \end{cases}$$

$$(3.8) \quad \tilde{\Psi}_{\phi, 1}(t) \approx ((t-1) \vee x_n^2)^{1-\alpha} \phi'((t-1) \vee x_n^2).$$

(ii) *Let $\frac{3}{4} < t < 1$. Suppose that $0 < \alpha < 1$. Then,*

$$(3.9) \quad \Psi_{\phi, 0}(t) \approx \begin{cases} (1-t)^{1-\alpha} \phi(1-t) & \text{if } \alpha + a > 1, \\ 1 & \text{otherwise.} \end{cases}$$

$$(3.10) \quad \Psi_{\phi, 1}(t) \approx (1-t)^{1-\alpha} \phi'(1-t).$$

$$(3.11) \quad \tilde{\Psi}_{\phi, 0}(t) \approx \begin{cases} \max(1, x_n^{-2\alpha+2} \phi(x_n^2)) + \phi(1-t)(1-t)^{2-\alpha} x_n^{-2} e^{-\delta \frac{x_n^2}{1-t}} & \text{if } 1-t < x_n^2, \quad \alpha > 0, \\ \phi(1-t) x_n^{2-2\alpha} & \text{if } x_n^2 < 1-t, \quad \alpha > 1, \\ \max(1, (1-t)^{-\alpha+1} \phi(1-t)) & \text{if } x_n^2 < 1-t, \quad \alpha < 1. \end{cases}$$

$$(3.12) \quad \tilde{\Psi}_{\phi,1}(t) \approx \begin{cases} x_n^{-2\alpha+2}\phi'(x_n^2) + \phi'(1-t)(1-t)^{2-\alpha}x_n^{-2}e^{-\delta\frac{x_n^2}{1-t}} & \text{if } 1-t < x_n^2, \quad \alpha > 0, \\ \phi'(1-t)x_n^{2-2\alpha} & \text{if } x_n^2 < 1-t, \quad \alpha > 1, \\ (1-t)^{-\alpha+1}\phi'(1-t) & \text{if } x_n^2 < 1-t, \quad \alpha < 1. \end{cases}$$

Proof. We first treat the cases $\Psi_{\phi,0}$ and $\tilde{\Psi}_{\phi,0}$. We first remind that

$$(3.13) \quad \begin{aligned} \frac{1}{2}(1-s) \leq t-s \leq (1-s), \quad \frac{3}{4} < t < 1 & \quad \text{for } 0 < s < 2t-1, \\ \frac{1}{2}(1-s) \leq 1-t \leq (1-s), \quad \frac{3}{4} < t < 1 & \quad \text{for } 2t-1 < s < t, \\ (1-s) \leq t-s \leq 2(1-s), \quad 1 < t < \frac{5}{4} & \quad \text{for } 0 < s < 2-t, \\ \frac{1}{2}(t-s) \leq t-1 \leq (t-s), \quad 1 < t < \frac{5}{4} & \quad \text{for } 2-t < s < 1. \end{aligned}$$

• (The case $1 < t < \frac{5}{4}$) We split the integral over $[0, 1]$ into $[0, 2-t]$ and $[2-t, 1]$. Then, from (3.13)_{3,4} and the change of variables, we have

$$(3.14) \quad \begin{aligned} \Psi_{\phi,0}(t) &\approx \int_0^{2-t} (1-s)^{-\alpha}\phi(1-s)ds + \int_{2-t}^1 (t-1)^{-\alpha}\phi(1-s)ds, \\ &= \int_{t-1}^1 s^{-\alpha}\phi(s)ds + (t-1)^{-\alpha} \int_0^{t-1} \phi(s)ds. \end{aligned}$$

Applying (2.16) and (2.18) in (3.14), we obtain (3.5).

Using (3.13)₄, we have $e^{-\frac{x_n^2}{2(t-1)}} \leq e^{-\frac{x_n^2}{2(t-s)}} \leq e^{-\frac{2x_n^2}{(t-1)}}$ for $2-t < s < 1$, namely $e^{-\frac{x_n^2}{4(t-1)}} \leq e^{-\frac{x_n^2}{4(t-s)}} \leq e^{-\frac{x_n^2}{(t-1)}}$. Thus, we note that $e^{-\frac{x_n^2}{4(t-s)}} \approx e^{-\delta\frac{x_n^2}{t-1}}$ for $2-t < s < 1$ (see Notation 2.3). Similarly, we split the following integral:

$$(3.15) \quad \tilde{\Psi}_{\phi,0}(t) \approx \int_{t-1}^1 s^{-\alpha}e^{-\frac{x_n^2}{4s}}\phi(s)ds + (t-1)^{-\alpha}e^{-\delta_1\frac{x_n^2}{t-1}} \int_0^{t-1} \phi(s)ds.$$

Due to (2.18), we have

$$(3.16) \quad (t-1)^{-\alpha}e^{-\delta_1\frac{x_n^2}{t-1}} \int_0^{t-1} \phi(s)ds \approx (t-1)^{-\alpha+1}\phi(t-1)e^{-\delta_1\frac{x_n^2}{t-1}}.$$

For the first term of the right-hand side of (3.15), we separate the cases that $x_n^2 \leq t-1$ and $x_n^2 > t-1$. In the case that $x_n^2 \leq t-1$, since $e^{-\frac{x_n^2}{4s}} \approx 1$ for $x_n^2 \leq t-1 \leq s < 1$, we can see via (2.16) that

$$(3.17) \quad \int_{t-1}^1 s^{-\alpha}e^{-\frac{x_n^2}{4s}}\phi(s)ds \approx \int_{t-1}^1 s^{-\alpha}\phi(s)ds \approx \max(1, (t-1)^{-\alpha+1}\phi(t-1)).$$

Thus, if $x_n^2 \leq t-1$, it follows from (3.15), (3.16) and (3.17) that

$$(3.18) \quad \tilde{\Psi}_{\phi,0}(t) \approx \max(1, (t-1)^{-\alpha+1}\phi(t-1)).$$

Secondly, we treat the case that $t-1 < x_n^2$. As in (3.17), we have

$$(3.19) \quad \int_{t-1}^1 s^{-\alpha}e^{-\frac{x_n^2}{4s}}\phi(s)ds = \int_{t-1}^{x_n^2} s^{-\alpha}e^{-\frac{x_n^2}{4s}}\phi(s)ds + \int_{x_n^2}^1 s^{-\alpha}e^{-\frac{x_n^2}{4s}}\phi(s)ds.$$

Since $e^{-\frac{x_n^2}{4s}} \approx 1$ for $x_n^2 < s < 1$, it follows from (2.16) that

$$(3.20) \quad \int_{x_n^2}^1 s^{-\alpha}e^{-\frac{x_n^2}{4s}}\phi(s)ds \approx \int_{x_n^2}^1 s^{-\alpha}\phi(s)ds \approx \max(1, x_n^{-2\alpha+2}\phi(x_n^2)).$$

This implies that

$$(3.21) \quad \max(1, x_n^{-2\alpha+2}\phi(x_n^2)) \lesssim \int_{t-1}^1 s^{-\alpha} e^{-\frac{x_n^2}{4s}} \phi(s) ds.$$

On the other hand, noting that $e^{-b} \leq c_m b^{-m}$ for any $b > 0$ and $m \in \mathbb{N}$, we have

$$(3.22) \quad \begin{aligned} \int_{t-1}^{x_n^2} s^{-\alpha} e^{-\frac{x_n^2}{4s}} \phi(s) ds &\leq c \int_{t-1}^{x_n^2} s^{-\alpha} \left(\frac{x_n^2}{4s}\right)^{-m} \phi(s) ds \leq c x_n^{-2m} \int_{t-1}^{x_n^2} s^{-\alpha+m} \phi(s) ds \\ &\leq c x_n^{-2m} x_n^{-2\alpha+2m+2} \phi(x_n^2) \leq c x_n^{-2\alpha+2} \phi(x_n^2). \end{aligned}$$

Estimates (3.19) -(3.22) implies that for $1 < t$ and $t-1 < x_n^2 < \frac{1}{2}$

$$(3.23) \quad \int_{t-1}^1 s^{-\alpha} e^{-\frac{x_n^2}{4s}} \phi(s) ds \approx \max(1, x_n^{-2\alpha+2} \phi(x_n^2)).$$

Adding up (3.15), (3.16), (3.18) and (3.23), we get (3.7).

• (The case $\frac{3}{4} < t < 1$) Noting that $\frac{1}{2} < 2t-1 < t$ and using (3.13)₁, (2.19), (2.20), (2.16) and change of variables, it follows that

$$(3.24) \quad \begin{aligned} \Psi_{\phi,0}(t) &\approx \int_0^{2t-1} (1-s)^{-\alpha} \phi(1-s) ds + \int_{2t-1}^t (t-s)^{-\alpha} \phi(1-t) ds \\ &= \int_{2-2t}^1 s^{-\alpha} \phi(s) ds + \phi(1-t) \int_0^{1-t} s^{-\alpha} ds \\ &\approx \max(1, (1-t)^{1-\alpha} \phi(1-t)). \end{aligned}$$

Hence, we obtain (3.9). It remains to show (3.11). We note first via (3.13)_{1,2}, (2.19) and (2.20) that

$$(3.25) \quad \begin{aligned} \tilde{\Psi}_{\phi,0}(t) &\approx \int_0^{2t-1} (1-s)^{-\alpha} e^{-\delta_1 \frac{x_n^2}{1-s}} \phi(1-s) ds + \int_{2t-1}^t (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi(1-t) ds \\ &:= I_1 + I_2. \end{aligned}$$

In case that $x_n^2 < 1-t$, since $e^{-\delta_1 \frac{x_n^2}{s}} \approx 1$ for $1-t < s < 1$, it follows due to (2.16) that

$$(3.26) \quad I_1 = \int_{2-2t}^1 s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi(s) ds \approx \int_{2-2t}^1 s^{-\alpha} \phi(s) ds \approx \max(1, (1-t)^{-\alpha+1} \phi(1-t)).$$

For I_2 , using the change of variables, we have

$$(3.27) \quad \begin{aligned} I_2 &= \phi(1-t) \int_0^{1-t} s^{-\alpha} e^{-\frac{x_n^2}{4s}} ds = \phi(1-t) x_n^{2-2\alpha} \int_{\frac{x_n^2}{1-t}}^{\infty} s^{-2+\alpha} e^{-\frac{s}{4}} ds \\ &\approx \begin{cases} \phi(1-t) x_n^{2-2\alpha} & \text{if } \alpha > 1, \\ \phi(1-t) (1-t)^{1-\alpha} & \text{if } \alpha < 1. \end{cases} \end{aligned}$$

Summing up (3.25), (3.26) and (3.27), we obtain

$$\begin{aligned} \tilde{\Psi}_{\phi,0}(t) &\approx \begin{cases} \phi(1-t) x_n^{2-2\alpha} + \max(1, (1-t)^{-\alpha+1} \phi(1-t)) & \text{if } \alpha > 1, \\ \max(1, (1-t)^{-\alpha+1} \phi(1-t)) & \text{if } \alpha < 1 \end{cases} \\ &\approx \begin{cases} \phi(1-t) x_n^{2-2\alpha} & \text{if } \alpha > 1, \\ \max(1, (1-t)^{-\alpha+1} \phi(1-t)) & \text{if } \alpha < 1. \end{cases} \end{aligned}$$

Thus, we have (3.11) for the case that $x_n^2 < 1-t$.

On the other hand, in case that $1-t < x_n^2$, we split I_1 as follows:

$$(3.28) \quad I_1 = \int_{2-2t}^{x_n^2} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi(s) ds + \int_{x_n^2}^1 s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi(s) ds := I_{11} + I_{12}.$$

It is straightforward from (2.16) that

$$(3.29) \quad I_{12} \approx \int_{x_n^2}^1 s^{-\alpha} \phi(s) ds \approx \max(1, x_n^{-2\alpha+2} \phi(x_n^2)).$$

With the same estimate of (3.22), we get

$$(3.30) \quad I_{11} = \int_{2-2t}^{x_n^2} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi(s) ds \leq c x_n^{-2\alpha+2} \phi(x_n^2).$$

Adding up (3.28), (3.29) and (3.30), we obtain for $1-t < x_n^2$

$$(3.31) \quad I_1 = \int_0^{2t-1} (1-s)^{-\alpha} e^{-\delta_1 \frac{x_n^2}{1-s}} \phi(1-s) ds \approx \max(1, x_n^{-2\alpha+2} \phi(x_n^2)).$$

Direct computations via the change of variables give

$$(3.32) \quad I_2 = \phi(1-t) x_n^{2-2\alpha} \int_{\frac{x_n^2}{1-t}}^{\infty} s^{-2+\alpha} e^{-\frac{s}{4}} ds \approx \phi(1-t) (1-t)^{2-\alpha} x_n^{-2} e^{-\delta \frac{x_n^2}{1-t}}.$$

Hence, from (3.25), (3.32), (3.31) and Lemma 2.2, for $1-t \leq x_n^2$,

$$\tilde{\Psi}_{\phi,0}(t) \approx \max(1, x_n^{-2\alpha+2} \phi(x_n^2)) + \phi(1-t) (1-t)^{2-\alpha} x_n^{-2} e^{-\delta \frac{x_n^2}{1-t}}.$$

We obtain (3.11).

Next, we prove the estimate (3.8) of $\tilde{\Psi}_{\phi,1}(t)$. Again we separately consider the cases that $t < 1$ and $t > 1$.

• (The case $1 < t < \frac{5}{4}$) For convenience, we define

$$(3.33) \quad \tilde{\Psi}_{\phi,k}(t; \tau) := \int_0^{\tau} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi^{(k)}(1-s) ds. \quad k = 0, 1.$$

We then split the integral, $\tilde{\Psi}_{\phi,1}(t; 2-t)$, as follows:

$$(3.34) \quad \tilde{\Psi}_{\phi,1}(t; 2-t) = \int_0^{\frac{3}{4}} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds + \int_{\frac{3}{4}}^{2-t} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds.$$

Since ϕ is smooth in $[\frac{1}{2}, 1]$, from direct calculation, we have for $0 < x_n < \frac{1}{2}$

$$(3.35) \quad \left| \int_0^{\frac{3}{4}} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds \right| \leq A_0 < \infty.$$

For the second term in (3.34), using (2.21) and (2.22), we have

$$(3.36) \quad \int_{\frac{1}{4}}^{2-t} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds \approx \int_{\frac{1}{4}}^{2-t} (1-s)^{-\alpha} e^{-\delta_1 \frac{x_n^2}{1-s}} \phi'(1-s) ds = \int_{t-1}^{\frac{1}{4}} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds.$$

In the case that $x_n^2 \leq t-1$, since $e^{-\delta_1 \frac{x_n^2}{s}} \approx 1$ for $x_n^2 \leq t-1 \leq s < 1$, we can see via (2.17) that

$$(3.37) \quad \int_{t-1}^{\frac{1}{4}} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds \approx \int_{t-1}^{\frac{1}{4}} s^{-\alpha} \phi'(s) ds \approx (t-1)^{-\alpha+1} \phi'(t-1).$$

Adding up (3.34), (3.35), (3.36) and (3.37), it follows that

$$(3.38) \quad \tilde{\Psi}_{\phi,1}(t; 2-t) \approx (t-1)^{-\alpha+1} \phi'(t-1),$$

and thus we obtain (3.8) for $x_n^2 < t-1$.

Secondly, if $t-1 < x_n^2$, we split the integral (3.37) as follows:

$$(3.39) \quad \int_{t-1}^{\frac{1}{4}} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds = \int_{t-1}^{x_n^2} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds + \int_{x_n^2}^{\frac{1}{4}} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds.$$

Since $e^{-\frac{x_n^2}{s}} \approx 1$ for $x_n^2 < s < 1$, it follows from (2.17) that

$$(3.40) \quad \int_{x_n^2}^{\frac{1}{4}} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds \approx \int_{x_n^2}^{\frac{1}{4}} s^{-\alpha} \phi'(s) ds \approx x_n^{-2\alpha+2} \phi'(x_n^2).$$

Recalling that $\text{sgn}(\phi')\phi'(s) > 0$ for $0 < s < \frac{1}{2}$ and $e^{-b} \leq c_m b^{-m}$ for any $b > 0$ and $m \in \mathbb{N}$, we have

$$(3.41) \quad \begin{aligned} \text{sgn}(\phi') \int_{t-1}^{x_n^2} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds &\leq c \text{sgn}(\phi') \int_{t-1}^{x_n^2} s^{-\alpha} \left(\frac{x_n^2}{\delta_1 s}\right)^{-m} \phi'(s) ds \\ &\leq c \text{sgn}(\phi') x_n^{-2m} \int_{t-1}^{x_n^2} s^{-\alpha+m} \phi'(s) ds \\ &\leq c x_n^{-2m} x_n^{-2\alpha+2m+2} \text{sgn}(\phi') \phi'(x_n^2) \\ &\leq c x_n^{-2\alpha+2} \text{sgn}(\phi') \phi'(x_n^2). \end{aligned}$$

Therefore, we deduce from estimates (3.39) -(3.41) that

$$(3.42) \quad \int_{t-1}^{\frac{1}{2}} s^{-\alpha} e^{-\delta_1 \frac{x_n^2}{s}} \phi'(s) ds \approx x_n^{-2\alpha+2} \phi'(x_n^2).$$

Adding up estimates (3.34), (3.35), (3.36) and (3.42), it follows in case $t-1 < x_n^2$ that

$$(3.43) \quad \tilde{\Psi}_{\phi,1}(t; 2-t) \approx x_n^{-2\alpha+2} \phi'(x_n^2),$$

which implies (3.8) for $t-1 < x_n^2$.

• (The case $\frac{3}{4} < t < 1$) Noting that $\frac{1}{2} < 2t-1 < t$. Hence, we have

$$(3.44) \quad \tilde{\Psi}_{\phi,1}(t) = \int_0^{\frac{1}{2}} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds + \int_{\frac{1}{2}}^t (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds.$$

Since ϕ is smooth in $[\frac{1}{2}, 1]$, it is direct for $0 < x_n < \frac{1}{2}$ that

$$(3.45) \quad \left| \int_0^{\frac{1}{2}} (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds \right| \leq B_0 < \infty.$$

Using (3.13)_{1,2}, it follows that

$$(3.46) \quad \begin{aligned} &\int_{\frac{1}{2}}^t (t-s)^{-\alpha} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds \\ &\approx \int_{\frac{1}{2}}^{2t-1} (1-s)^{-\alpha} e^{-\delta_1 \frac{x_n^2}{1-s}} \phi'(1-s) ds + \int_{2t-1}^t (t-s)^{-\alpha} e^{-\delta_1 \frac{x_n^2}{t-s}} \phi'(1-t) ds \\ &:= I_1 + I_2. \end{aligned}$$

In case that $x_n^2 \leq 1-t$, direct computations via the change of variables give

$$(3.47) \quad I_2 = \phi'(1-t) x_n^{2-2\alpha} \int_{\frac{x_n^2}{1-t}}^{\infty} s^{-2+\alpha} e^{-\delta_1 s} ds \approx \begin{cases} \phi'(1-t) x_n^{2-2\alpha} & \text{if } \alpha > 1, \\ \phi'(1-t) (1-t)^{1-\alpha} & \text{if } \alpha < 1. \end{cases}$$

Since $e^{-\delta_1 \frac{x_n^2}{s}} \approx 1$ for $1-t \leq s$, it follows from (2.16) that

$$(3.48) \quad I_1 \approx \int_{2-2t}^1 s^{-\alpha} \phi'(s) ds \approx (1-t)^{-\alpha+1} \phi'(1-t).$$

From (3.44) (3.45), (3.46), (3.47) and (3.48), we have

$$\tilde{\Psi}_{\phi,1}(t) \approx \begin{cases} \phi(1-t) x_n^{2-2\alpha} & \text{if } \alpha > 1, \\ (1-t)^{-\alpha+1} \phi(1-t) & \text{if } \alpha < 1. \end{cases}$$

We obtain (3.12) for $x_n^2 < 1 - t$.

On the other hand, in case that $1 - t < x_n^2$, we split I_1 as follows:

$$(3.49) \quad I_1 = \int_{2-2t}^{x_n^2} s^{-\alpha} e^{-\frac{x_n^2}{4s}} \phi'(s) ds + \int_{x_n^2}^{\frac{1}{2}} s^{-\alpha} e^{-\frac{x_n^2}{4s}} \phi'(s) ds := I_{11} + I_{12}.$$

It is straightforward from (3.40) that

$$(3.50) \quad I_{12} \approx \int_{x_n^2}^{\frac{1}{2}} s^{-\alpha} \phi'(s) ds \approx x_n^{-2\alpha+2} \phi'(x_n^2).$$

With the same estimate of (3.41), we have

$$(3.51) \quad \text{sgn}(\phi') \int_{2-2t}^{x_n^2} s^{-\alpha} e^{-\frac{x_n^2}{4s}} \phi'(s) ds \leq c \text{sgn}(\phi') x_n^{-2\alpha+2} \phi'(x_n^2).$$

Adding up (3.49) and (3.51), we obtain for $1 - t < x_n^2$

$$(3.52) \quad I_1 \approx \int_{\frac{1}{2}}^{2t-1} (1-s)^{-\alpha} e^{-\delta_1 \frac{x_n^2}{1-s}} \phi'(1-s) ds \approx x_n^{-2\alpha+2} \phi'(x_n^2).$$

Direct computations via the change of variables give

$$(3.53) \quad I_2 \approx \phi'(1-t) x_n^{2-2\alpha} \int_{\frac{x_n^2}{1-t}}^{\infty} s^{-2+\alpha} e^{-\delta_1 s} ds \approx \phi'(1-t) (1-t)^{2-\alpha} x_n^{-2} e^{-\delta \frac{x_n^2}{1-t}}.$$

Hence, from (3.44), (3.45), (3.46), (3.52), (3.53) and Lemma 2.2, we obtain (3.12) for $1 - t \leq x_n^2$. Following similar way of proving (3.5) and (3.9), one can also obtain that

$$(3.54) \quad \Psi_{\phi,1}(t) \approx |t-1|^{1-\alpha} \phi'(|t-1|), \quad \frac{3}{4} < t < \frac{5}{4}.$$

which implies (3.6) and (3.10). Since its proof is tedious repetition, the details are omitted. \square

Next, we provide the proofs of Lemma 2.2 and Lemma 2.4.

Proof of Lemma 2.2 Let $k \in \mathbb{N} \cup \{0\}$ be an integer with $2^k \leq \frac{x_n^2}{t} < 2^{k+1}$. Due to the Assumption 2.1, it follows that

$$(3.55) \quad c^{-k-1} \phi(t) \leq \phi(2^{k+1}t) \leq \phi(x_n^2).$$

We choose a positive integer m such that $c \leq 2^m$, where c is the constant in (3.55). Noting that $e^{-x} \leq c_m x^{-m}$, we obtain

$$e^{-\delta_1 \frac{x_n^2}{t}} \phi(t) \leq e^{-\delta_1 2^k} c^{k+1} \phi(x_n^2) \leq c_m (\delta_1 2^k)^{-m} c^{k+1} \phi(x_n^2) \leq c_m c \delta_1^{-m} \left(\frac{c}{2^m}\right)^k \phi(x_n^2) \leq c_m c \delta_1^{-m} \phi(x_n^2).$$

Since cc_m is an absolute constant, the first inequality in (2.23) is proved. For the second inequality, it follows from the following observation:

$$c^{-k-1} \text{sgn}(\phi'(t)) \phi'(t) \leq \text{sgn}(\phi'(2^{k+1}t)) \phi'(2^{k+1}t) \leq \text{sgn}(\phi'(x_n^2)) \phi(x_n^2).$$

Similar computations as above deduce the second inequality, and thus we skip its details. In the case that $\phi = t^{-\alpha}$, it is straightforward that $t^{-1} \phi(t) \approx \text{sgn}(\phi') \phi'(t)$ for $0 < t < \frac{1}{2}$. Again, the similar arguments as (2.23) implies (2.24), and the details are omitted. \square

Proof of Lemma 2.4 The second equality of (2.26) was shown in Lemma 3.2 in [8]. Thus it suffices to prove the first equality of (2.26). Let $x' \in \mathbb{R}^{n-1}$ with $|x'| \geq 1$. We divide \mathbb{R}^{n-1} by three disjoint sets D_1, D_2 and D_3 , which are defined by

$$D_1 = \left\{ z' \in \mathbb{R}^{n-1} : |x' - z'| \leq \frac{1}{10} |x'| \right\},$$

$$D_2 = \left\{ z' \in \mathbb{R}^{n-1} : |z'| \leq \frac{1}{10}|x'| \right\}, \quad D_3 = \mathbb{R}^{n-1} \setminus (D_1 \cup D_2).$$

We then split the following integral into three terms as follows:

$$(3.56) \quad \int_{\mathbb{R}^{n-1}} \Gamma'(x' - z', t) D_{x_n} N(z', x_n) dz' = \int_{D_1} \cdots + \int_{D_2} \cdots + \int_{D_3} \cdots := K_1 + I + K_3.$$

We first estimate I .

$$(3.57) \quad \begin{aligned} I &= \int_{|z'| \leq \frac{1}{10}|x'|} D_{x_n} N(z', x_n) \Gamma'(x' - z', t) dz' \\ &\approx ct^{-\frac{n-1}{2}} e^{-\delta \frac{|x'|^2}{t}} \int_{|z'| \leq \frac{1}{10}|x'|} \frac{x_n}{(|z'|^2 + x_n^2)^{\frac{n}{2}}} dz' \\ &\approx ct^{-\frac{n-1}{2}} e^{-\delta \frac{|x'|^2}{t}} \int_{|z'| \leq \frac{|x'|}{10x_n}} \frac{1}{(|z'|^2 + 1)^{\frac{n}{2}}} dz' \\ &\approx ct^{-\frac{n-1}{2}} e^{-\delta \frac{|x'|^2}{t}}. \end{aligned}$$

On the other hand, the term K_3 is controlled as follows:

$$(3.58) \quad \begin{aligned} |K_3| &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n}{2}}} t^{-\frac{n-1}{2}} \int_{\{|z'| \geq \frac{1}{10}|x'|\}} e^{-\frac{|z'|^2}{t}} dz' \\ &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n}{2}}} \int_{\{|z'| \geq \frac{|x'|}{10\sqrt{t}}\}} e^{-|z'|^2} dz' \\ &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n}{2}}} e^{-\frac{|x'|^2}{t}} \leq cx_n t^{\frac{1}{2}}, \end{aligned}$$

where we used that $e^{-\frac{|x'|^2}{2t}} \lesssim \frac{t^{\frac{m}{2}}}{|x'|^m}$ for $m \geq 0$ (in fact, we take $m = \frac{1}{2}$).

Meanwhile, we decompose K_1 in the following way.

$$\begin{aligned} K_1 &= \int_{D_1} \Gamma'(x' - z', t) D_{x_n} N(z', x_n) dz' \\ &= \int_{\{|z'| \leq \frac{1}{10} \frac{|x'|}{\sqrt{t}}\}} \Gamma'(z', 1) \left(D_{x_n} N(x' - \sqrt{t}z', x_n) - D_{x_n} N(x', x_n) \right) dz' \\ &\quad + D_{x_n} N(x', x_n) \int_{\mathbb{R}^{n-1}} \Gamma'(z', 1) dz' - D_{x_n} N(x', x_n) \int_{\{|z'| \geq \frac{1}{10} \frac{|x'|}{\sqrt{t}}\}} \Gamma'(z', 1) dz' \\ &= K_{11} + D_{x_n} N(x', x_n) + K_{12}, \end{aligned}$$

where we used $\int_{\mathbb{R}^{n-1}} \Gamma'(z', 1) dz' = 1$. Since $\frac{|x'|^2}{t} \geq 1$, we observe that

$$(3.59) \quad \begin{aligned} |K_{11}(x', t)| &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n+1}{2}}} t^{\frac{1}{2}} \int_{\{|z'| \leq \frac{1}{10} \frac{|x'|}{\sqrt{t}}\}} e^{-|z'|^2} |z'| dz' \\ &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n+1}{2}}} t^{\frac{1}{2}} \leq cx_n t^{\frac{1}{2}} \end{aligned}$$

and

$$(3.60) \quad \begin{aligned} |K_{12}(x', t)| &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n}{2}}} \int_{\{|z'| \geq \frac{1}{10} \frac{|x'|}{\sqrt{t}}\}} e^{-|z'|^2} dz' \\ &\leq c \frac{x_n}{(|x'|^2 + x_n^2)^{\frac{n}{2}}} e^{-\frac{|x'|^2}{2t}} \leq cx_n t^{\frac{1}{2}}. \end{aligned}$$

Adding up (3.57) -(3.60) and putting $J_1 := K_3 + K_{11} + K_{12}$, we deduce (2.26)-(2.27). \square

4. PROOF OF THEOREM 2.6

4.1. Estimate of w . We remind the decomposition of w , which given as $w = w^L + w^N + w^B$ in (2.25). Let $x' \in B_1' \subset \mathbb{R}^{n-1}$. Noting that $|x' - y'| \geq \frac{1}{2}$ for $y' \in A$ and recalling boundary conditions in (2.10)-(2.11) with Assumption 2.1, it follows for $\frac{3}{4} < t < \frac{5}{4}$ that

$$(4.1) \quad \begin{aligned} w_i^L(x, t) &= \int_0^t \int_A L_{ni}(x' - y', x_n, t - s) g_n(y', s) dy' ds \\ &\leq c \int_0^t (t - s)^{-\frac{1}{2}} \phi(1 - s) ds \\ &\approx \max(1, |t - 1|^{\frac{1}{2}} \phi(|t - 1|)), \end{aligned}$$

where we used (2.5).

On the other hand, we note that w_i^N vanishes for $t > 1$. Furthermore, it follows from (2.25) that

$$(4.2) \quad w_i^N(x, t) = 2 \int_A D_{x_i} N(x' - y', x_n) g_n(y', t) dy' = 2T_i g_n^S(x') \phi(1 - t),$$

where

$$T_i g_n^S(x') = \int_A D_{x_i} N(x' - y', x_n) g_n^S(y') dy'.$$

Since $T_i g_n^S(x')$ is positive and finite, we have $T_i g_n^S(x') \approx 1$ for $|x'| < \frac{1}{2}$ and $x_n \in (0, \frac{1}{2})$, which implies via (4.2) that

$$(4.3) \quad w_i^N(x, t) \approx \phi(1 - t) \mathbf{I}_{\{t < 1\}}.$$

It remains to estimate w_i^B . Since $w_i^B = 0$ for $i = n$, we note that $|x' - y'| \geq 1$ for $|x'| < \frac{1}{2}$ and $y' \in A$. Using (2.28) and (2.30), for $i \neq n$, we have for $\frac{3}{4} < t < \frac{5}{4}$

$$(4.4) \quad \begin{aligned} w_i^B(x, t) &= \int_0^t D_{x_n} \Gamma_1(x_n, t - s) g_n^T(s) \int_A \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t) D_{z_i} N(z', 0) dz' g_n^S(y') dy' ds \\ &\approx \int_0^t D_{x_n} \Gamma_1(x_n, t - s) g_n^T(s) ds. \end{aligned}$$

We separately treat (4.4) for the cases that $t < 1$ and for $t > 1$.

• **(The case $1 - t_0 < t < 1$)** Using (4.4) and (3.11), we observe that

$$(4.5) \quad \begin{aligned} w_i^B(x, t) &\approx - \int_0^t \frac{x_n}{(t - s)^{\frac{3}{2}}} e^{-\frac{x_n^2}{4(t-s)}} \phi(1 - s) ds \\ &\approx - \begin{cases} \phi(1 - t) & \text{if } x_n < (1 - t)^{\frac{1}{2}}, \\ \phi(x_n^2) + \phi(1 - t)(1 - t)^{\frac{1}{2}} x_n^{-1} e^{-\delta \frac{x_n^2}{1-t}} & \text{if } (1 - t)^{\frac{1}{2}} < x_n. \end{cases} \end{aligned}$$

Therefore, we have

$$(4.6) \quad |w_i^B(x, t)| \lesssim \begin{cases} 1 & \text{if } \phi(t) = |\ln t|^{-1} \\ \phi(1 - t) & \text{otherwise.} \end{cases}$$

Adding up (4.1), (4.3) and (4.6), we obtain

$$(4.7) \quad |w_i(x, t)| \lesssim \begin{cases} 1 & \text{if } \phi(t) = |\ln t|^{-1} \\ \phi(1 - t) & \text{otherwise.} \end{cases}$$

We complete the proof of (2.35) for $t < 1$.

Let $\phi \neq |\ln t|^{-1}$ and we choose a sufficiently large $\epsilon_1 \gg 1$. We then show that there exists a sufficiently small ϵ depending on ϵ_1 such that $w^B(x, t) > -\epsilon\phi(1-t)$ in $\tilde{\mathcal{D}}_- = \{(x, t) \in \mathcal{D}_- : \epsilon_1\sqrt{1-t} < x_n < \frac{1}{2}\}$. Indeed, recalling the second estimate in (4.5), it follows that

$$(4.8) \quad w_i^B(x, t) \gtrsim -\phi(\epsilon_1^2(1-t)) - \phi(1-t)\epsilon_1^{-1}e^{-\delta\epsilon_1^2} \geq -\epsilon\phi(1-t).$$

In a similar way, we can see via (4.1) that

$$(4.9) \quad |w^L(x, t)| < \epsilon\phi(1-t), \quad \text{for } (x, t) \in \tilde{\mathcal{D}}_-.$$

Summing up estimates (4.3), (4.8) and (4.9), we conclude in $\tilde{\mathcal{D}}_-$ that

$$(4.10) \quad w_i(x, t) \geq w_i^N(x, t) + w_i^B(x, t) - |w_i^L(x, t)| \geq (c - 2\epsilon)\phi(1-t) \geq \frac{c}{2}\phi(1-t).$$

We complete the proof of (2.36) for $t < 1$.

• **(The case $1 < t < 1 + t_0$)** With the aid of (3.7) and (4.4), we have

$$(4.11) \quad \begin{aligned} w_i^B(x, t) &\approx - \int_0^1 \frac{x_n}{(t-s)^{\frac{3}{2}}} e^{-\frac{x_n^2}{4(t-s)}} \phi(1-s) ds \\ &\approx - \begin{cases} x_n(t-1)^{-\frac{1}{2}}\phi(t-1) & \text{if } x_n^2 \leq t-1, \\ \max(1, \phi(x_n^2)) & \text{if } t-1 \leq x_n^2. \end{cases} \end{aligned}$$

Owing to (4.1), (4.3) and (4.11), it follows that

$$(4.12) \quad |w_i(x, t)| \lesssim \begin{cases} 1 & \text{if } \phi(t) = |\ln t|^{-1} \\ \phi(1-t) & \text{otherwise,} \end{cases}$$

which implies (2.35) for $t \in (1, \frac{5}{4})$ and thus together with (4.7), the proof of (2.35) is completed. Moreover, if $\phi \neq |\ln t|^{-1}$, by choosing sufficiently small t_0 such that for $\epsilon_1\sqrt{t-1} < x_n$ and $\sqrt{t-1} < t_0$, we have

$$(4.13) \quad w_i^B(x, t) + w_i^L(x, t) \leq (-c + c_1\sqrt{t-1})\phi(t-1) \leq -\frac{c}{2}\phi(t-1).$$

On the other hand, from (4.1) and (4.11), for $1 < t < 1 + t_0$ and $\frac{1}{2}\sqrt{t-1} < x_n < \sqrt{t-1}$, we have

$$w_i(x, t) \leq (-\frac{c}{2} + \epsilon)\phi(t-1) \leq -\frac{c}{4}\phi(t-1).$$

This completes the proof of (2.37). □

4.2. Estimate of ∇w . We note for $1 \leq i \leq n-1$ that

$$(4.14) \quad D_{x_n} w_i^L(x, t) = - \sum_{k=1}^{n-1} D_{x_k} w_{ik}^L(x, t) + \frac{1}{2} D_{x_i} f(x, t),$$

where

$$\begin{aligned} f(x, t) &= \int_0^t \int_{\mathbb{R}^{n-1}} D_{x_n} \Gamma(x' - z', x_n, t-s) g_n(z', s) dz' ds, \\ w_{ik}^L(x, t) &= \int_0^t \int_{\mathbb{R}^{n-1}} L_{ik}(x' - y', x_n, t-s) g_n(y', s) dy' ds, \quad k = 1, 2, \dots, n-1. \end{aligned}$$

Similar estimates as in (4.1) imply for $\frac{3}{4} < t < \frac{5}{4}$ that

$$(4.15) \quad D_{x_n} w_i^L(x, t) \leq c \max(1, |t-1|^{\frac{1}{2}} \phi(|t-1|)).$$

It is also straightforward as in (4.3) that

$$(4.16) \quad |D_x w_i^N(x, t)| \leq c\phi(|t-1|) \mathbf{I}_{\{t < 1\}}.$$

We can easily see that tangential derivatives of w_i^B satisfy essentially the same estimates as those of w_i^B , namely $|D_{x'} w_i^B(x, t)|$ satisfies (4.6). Therefore, it remains to estimate $D_{x_n} w_i^B$. As before, we treat the cases that $t < 1$

and $t > 1$ separately.

• **(The case $1 - t_0 < t < 1$)** We recall that

$$B_{in}(x', x_n, t) = D_{x_n} A_i(x', x_n, t),$$

where

$$(4.17) \quad A_i(x, t) = \int_{\mathbb{R}^{n-1}} \Gamma(x' - z', x_n, t) D_{z_i} N(z', 0) dz', \quad i = 1, 2, \dots, n-1.$$

Using that $(D_t - \Delta)A_i = 0$ for $(x, t) \in \mathbb{R}_+^n \times (0, \infty)$ and $g_n^T(y', 0) = 0$, we have

$$(4.18) \quad \begin{aligned} D_{x_n} w_i^B(x, t) &= \int_0^t \int_{\mathbb{R}^{n-1}} D_{x_n} B_{in}(x' - y', x_n, t - s) g_n^T(y', s) dy' ds \\ &= \int_0^t \int_{\mathbb{R}^{n-1}} (D_t - \Delta'_x) A_i(x' - y', x_n, t - s) g_n^T(y', s) dy' ds \\ &= \int_0^t \int_{\mathbb{R}^{n-1}} A_i(x' - y', x_n, t - s) (D_s - \Delta'_y) g_n^T(y', s) dy' ds. \end{aligned}$$

Due to the estimates (2.28), and (3.11), the second term is estimated as follows:

$$(4.19) \quad \begin{aligned} & \left| \int_0^t \int_{\mathbb{R}^{n-1}} A_i(x' - y', x_n, t - s) \Delta'_y g_n^T(y', s) dy' ds \right| \\ & \lesssim \int_0^t \frac{1}{(t-s)^{\frac{1}{2}}} e^{-\frac{x_n^2}{4(t-s)}} \phi(1-s) ds \\ & \approx \begin{cases} \max(1, x_n \phi(x_n^2)) + \phi(1-t)(1-t)^{\frac{3}{2}} x_n^{-2} e^{-\delta_1 \frac{x_n^2}{1-t}} & \text{if } 1-t < x_n^2, \\ \max(1, (1-t)^{\frac{1}{2}} \phi(1-t)) & \text{if } x_n^2 < 1-t. \end{cases} \end{aligned}$$

On the other hand, due to (2.28), (2.30) and (3.12), we compute for the first term as follows:

$$(4.20) \quad \begin{aligned} & \int_0^t \int_{\mathbb{R}^{n-1}} A_i(x' - y', x_n, t - s) D_s g_n^T(y', s) dy' ds \\ & \approx - \int_0^t \frac{1}{(t-s)^{\frac{1}{2}}} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds \\ & \approx - \begin{cases} x_n \phi'(x_n^2) + \phi'(1-t)(1-t)^{\frac{3}{2}} x_n^{-2} e^{-\delta_1 \frac{x_n^2}{1-t}} & \text{if } 1-t < x_n^2, \\ (1-t)^{\frac{1}{2}} \phi'(1-t) & \text{if } x_n^2 < 1-t. \end{cases} \end{aligned}$$

Thus, using Lemma 2.2, it follows that

$$(4.21) \quad |D_{x_n} w_i^B(x, t)| \leq -c(1-t)^{\frac{1}{2}} \phi'(1-t) \mathbf{I}_{\mathcal{D}_+} + x_n \phi'(x_n^2) \mathbf{I}_{\mathcal{D}_-}.$$

Hence, we obtain the second part in (2.38). In addition, we also note that

$$(4.22) \quad D_{x_n} w_i(x, t) \approx \begin{cases} -(1-t)^{\frac{1}{2}} \phi'(1-t) & \text{if } 0 < 1-t < \min\{t_0, t_1\} \text{ and } (x, t) \in \tilde{\mathcal{D}}_+ \\ -x_n \phi'(x_n^2) & \text{if } 0 < 1-t < \min\{t_0, t_1\} \text{ and } (x, t) \in \tilde{\mathcal{D}}_- \end{cases}$$

This completes the proof of (2.39).

• **(The case $1 < t < 1 + t_0$)**

We first recall that

$$(4.23) \quad D_{x_n} w_i^B(x, t) = \int_0^1 \int_{\mathbb{R}^{n-1}} D_{x_n}^2 A_i(x' - y', x_n, t - s) g_n(y', s) dy' ds,$$

where A_i is defined in (4.17). Since $|D_{x_n}^2 A_i(x' - y', x_n, t - s)| \leq c(t - s)^{-\frac{3}{2}} e^{-\frac{x_n^2}{4(t-s)}}$, it follows from (3.7) that

$$|D_{x_n} w_i^{\mathcal{B}}(x, t)| \leq c \int_0^1 (t - s)^{-\frac{3}{2}} e^{-\frac{x_n^2}{4(t-s)}} \phi(1 - s) ds \leq c ((t - 1) \vee x_n^2)^{-\frac{1}{2}} \phi((t - 1) \vee x_n^2).$$

We complete the proof of the first part in (2.38).

In case that $x_n \leq \epsilon_0 \sqrt{t - 1}$, it is direct that for $0 < s < 1$,

$$D_{x_n}^2 \Gamma_1(x_n, t - s) \approx -(t - s)^{-\frac{3}{2}} e^{-\delta_1 \frac{x_n^2}{t-s}} \approx -(t - s)^{-\frac{3}{2}}.$$

Hence, it follows from (2.28), (2.30) and (3.5) that

$$(4.24) \quad D_{x_n} w_i^{\mathcal{B}}(x, t) \approx -(t - 1)^{-\frac{1}{2}} \phi(t - 1).$$

Combining (4.15) and (4.24), for $x_n \leq \epsilon_0 \sqrt{t - 1}$, we have

$$D_{x_n} w_i(x, t) \approx -(t - 1)^{-\frac{1}{2}} \phi(t - 1),$$

which completes the proof of (2.40) for $x_n \leq \epsilon_0 \sqrt{t - 1}$.

Next, we consider the case that $\epsilon_1 \sqrt{t - 1} < x_n$. Dividing time interval into two parts such as $0 < s < 2 - t$ and $2 - t < s < 1$, we split the integral

$$(4.25) \quad \begin{aligned} D_{x_n} w_i^{\mathcal{B}}(x, t) &= \int_0^{2-t} \int_{\mathbb{R}^{n-1}} D_{x_n}^2 A_i(x' - y', x_n, t - s) g_n(y', s) dy' ds \\ &\quad + \int_{2-t}^1 \int_{\mathbb{R}^{n-1}} D_{x_n}^2 A_i(x' - y', x_n, t - s) g_n(y', s) dy' ds \\ &:= J_1 + J_2. \end{aligned}$$

Noting that $D_{x_n}^2 \Gamma_1(x_n, t - s) \approx (t - 1)^{-\frac{5}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}}$ for $2 - t < s$, we obtain

$$(4.26) \quad J_2 \approx (t - 1)^{-\frac{5}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}} \int_{2-t}^1 \phi(1 - s) ds \approx (t - 1)^{-\frac{3}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}} \phi(t - 1).$$

Using that $(D_t - \Delta)A_i = 0$ for $(x, t) \in \mathbb{R}_+^n \times (0, \infty)$. Since $\phi(1) = 0$ and integrating J_1 by parts, we get

$$(4.27) \quad \begin{aligned} J_1 &= \int_0^{2-t} \int_{\mathbb{R}^{n-1}} (D_t - \Delta'_x) A_i(x' - y', x_n, t - s) g_n(y', s) dy' ds \\ &= \int_0^{2-t} \int_{\mathbb{R}^{n-1}} A_i(x' - y', x_n, t - s) D_s g_n(y', s) dy' ds \\ &\quad - \int_0^{2-t} \int_{\mathbb{R}^{n-1}} A_i(x' - y', x_n, t - s) \Delta'_y g_n(y', s) dy' ds \\ &\quad - \int_{\mathbb{R}^{n-1}} A_i(x' - y', 2t - 2) \phi(t - 1) g_n^S(y') dy' \\ &:= J_{11} + J_{12} + J_{13}. \end{aligned}$$

It is straightforward from (2.28) and (2.30) that J_{13} satisfies

$$(4.28) \quad J_{13} = - \int_{\mathbb{R}^{n-1}} A_i(x' - y', 2t - 2) \phi(t - 1) g_n(y') dy' \approx - \frac{1}{(t - 1)^{\frac{1}{2}}} e^{-\frac{x_n^2}{4(t-1)}} \phi(t - 1).$$

From the proof of (3.7), the second term J_{12} is controlled as follows:

$$\begin{aligned}
|J_{12}| &\leq \left| \int_0^{2-t} \int_{\mathbb{R}^{n-1}} A_i(x' - y', x_n, t - s) \Delta'_y g_n(y', s) dy' ds \right| \\
&\leq c \int_0^{2-t} \frac{1}{(t-s)^{\frac{1}{2}}} e^{-\frac{x_n^2}{4(t-s)}} \phi(1-s) ds \\
(4.29) \quad &\leq c \int_0^{2-t} \frac{1}{(1-s)^{\frac{1}{2}}} e^{-\delta_1 \frac{x_n^2}{1-s}} \phi(1-s) ds \\
&= c \int_{t-1}^1 \frac{1}{s^{\frac{1}{2}}} e^{-\delta_1 \frac{x_n^2}{s}} \phi(s) ds \\
&\leq c \max(1, x_n \phi(x_n^2)).
\end{aligned}$$

Due to (3.8), we have

$$(4.30) \quad J_{11} \approx - \int_0^{2-t} \frac{1}{(t-s)^{\frac{1}{2}}} e^{-\frac{x_n^2}{4(t-s)}} \phi'(1-s) ds \approx -x_n \phi'(x_n^2).$$

We note for $\epsilon_1 \sqrt{t-1} < x_n$ with sufficiently large $\epsilon_1 > 0$ that

$$\begin{aligned}
(t-1)^{-\frac{3}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}} \phi(t-1) - (t-1)^{-\frac{1}{2}} e^{-\frac{x_n^2}{4(t-1)}} \phi(t-1) &\approx (t-1)^{-\frac{3}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}} \phi(t-1), \\
\max(1, x_n \phi(x_n^2)) - x_n \phi'(x_n^2) &\approx -x_n \phi'(x_n^2).
\end{aligned}$$

From (4.25) to (4.30), for $t-1 < t_0$ for small t_0 , we have

$$(4.31) \quad D_{x_n} w_i^{\mathcal{B}}(x, t) \approx -x_n \phi'(x_n^2) + (t-1)^{-\frac{3}{2}} x_n^2 e^{-\frac{x_n^2}{4(t-1)}} \phi(t-1).$$

With the aid of (4.15), (4.31) and Lemma 2.2, we complete the proofs of (2.41) for $\epsilon_1 \sqrt{t-1} \leq x_n$. \square

4.3. Estimate of pressure. For convenience, we denote

$$\mathcal{U}g_n(x', 0, t) = \int_0^t \int_A \Gamma(x' - y', 0, t-s) g_n(y', s) dy' ds.$$

Recalling that $g_n(x', 0) = 0$ and (2.9), we decompose $p = p_1 + p_2 + p_3$, where

$$\begin{aligned}
p_1(x, t) &= -D_{x_n} D_{x_n} N *' g_n(x, t), \\
p_2(x, t) &= 2(D_t - \Delta') D_{x_n} N *' \mathcal{U}g_n(x, t) \\
(4.32) \quad &= 2D_t D_{x_n} N *' \mathcal{U}g_n(x, t) - 2\Delta' D_{x_n} N *' \mathcal{U}g_n(x, t) \\
&:= p_{21} + p_{22}, \\
p_3(x, t) &= N *' D_t g_n(x, t).
\end{aligned}$$

Since $|x' - y'| \geq 1$ for $|x'| < \frac{1}{2}$ and $y' \in A$, we can estimate p_1 and p_3 as follows:

$$\begin{aligned}
(4.33) \quad |p_1(x, t)| &\leq c\phi(1-t) \mathbf{I}_{\{0 < t < 1\}}, \\
p_3(x, t) &\approx -\phi'(1-t) \mathbf{I}_{\{0 < t < 1\}}.
\end{aligned}$$

It remains to estimate p_2 . Firstly, for p_{22} we note that

$$\begin{aligned}
(4.34) \quad p_{22}(x, t) &= -2\Delta' D_{x_n} N *' \mathcal{U}g_n(x, t) = -2D_{x_n} N *' \mathcal{U}\Delta' g_n(x, t) \\
&= -2 \int_0^t \int_A (t-s)^{-\frac{1}{2}} \Delta' g_n(y', s) \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t-s) D_{x_n} N(z', x_n) dz' dy' ds.
\end{aligned}$$

From (4.34) and (2.29), we have

$$\begin{aligned}
(4.35) \quad |p_{22}(x, t)| &\leq c \int_0^t (t-s)^{-\frac{1}{2}} (x_n + (t-s)^{-\frac{n-1}{2}} e^{-\frac{1}{(t-s)}}) g_n^T(s) ds \\
&\leq c(1 + x_n \max(1, |t-1|^{\frac{1}{2}} \phi(|t-1|))) \\
&\leq c \max(1, x_n |t-1|^{\frac{1}{2}} \phi(|t-1|)).
\end{aligned}$$

• **(The case $\frac{3}{4} < t < 1$)**

Since $g_n(y', 0) = 0$, by intergration by parts and (2.31), we have

$$\begin{aligned}
(4.36) \quad p_{21}(x, t) &= \int_0^t \int_A (t-s)^{-\frac{1}{2}} \phi'(1-s) \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t-s) D_{x_n} N(z', x_n) dz' g_n(y') dy' ds \\
&\approx \int_0^t (t-s)^{-\frac{1}{2}} \phi'(1-s) (x_n + (t-s)^{-\frac{n-1}{2}} e^{-\delta \frac{1}{(t-s)}}) ds \\
&\approx \max(1, x_n (1-t)^{\frac{1}{2}} \phi'(1-t)).
\end{aligned}$$

Therefore, adding up (4.33), (4.35) and (4.36), we obtain for $x_n < \frac{1}{2}$ and $1-t < t_1$,

$$(4.37) \quad p(x, t) \approx -c\phi'(1-t).$$

We prove the first part of (2.42) for $t < 1$.

• **(The case $1 < t$)** Since $p_1(x, t) = p_3(x, t) = 0$ for $t > 1$, Lemma 2.4 implies that

$$\begin{aligned}
(4.38) \quad p_{22} &= -2 \int_0^1 \int_{\mathbb{R}^{n-1}} (t-s)^{-\frac{1}{2}} \Delta' g_n(y', s) (D_{x_n} N(x' - y', x_n) + (I + J_1)(x' - y', x_n, t)) dy' ds \\
&\leq c \int_0^1 (x_n (t-s)^{-\frac{1}{2}} + (t-s)^{-\frac{n}{2}} e^{-\frac{1}{8(t-s)}}) \phi(1-s) ds \\
&\leq c(1 + x_n \max(1, (t-1)^{\frac{1}{2}} \phi(t-1)))
\end{aligned}$$

and

$$\begin{aligned}
(4.39) \quad p_{21} &= \int_0^1 \int_{\mathbb{R}^{n-1}} (t-s)^{-\frac{3}{2}} \phi(1-s) \int_{\mathbb{R}^{n-1}} \Gamma'(x' - y' - z', t-s) D_{x_n} N(z', x_n) dz' g_n(y') dy' ds \\
&\approx \int_0^1 (x_n (t-s)^{-\frac{3}{2}} + (t-s)^{-\frac{n+1}{2}} e^{-\frac{1}{(t-s)}}) \phi(1-s) ds \\
&\approx x_n (t-1)^{-\frac{1}{2}} \phi(t-1) + 1.
\end{aligned}$$

Hence, we have

$$(4.40) \quad c_1 x_n (t-1)^{-\frac{1}{2}} \phi(t-1) - c_2 \leq p_{21}(x, t) \leq c_3 x_n (t-1)^{-\frac{1}{2}} \phi(t-1) + c_4,$$

which implies the second part of (2.42) for $t > 1$, and thus we complete the proof of (2.42). \square

5. PROOF OF PROPOSITION 2.10

To prove Proposition 2.10, we consider the following equations

$$(5.1) \quad \begin{cases} W_t - \Delta W + \nabla Q = f & \mathbb{R}_+^n \times (0, T), \\ \operatorname{div} W = 0 & \mathbb{R}_+^n \times (0, T), \\ W|_{t=0} = 0, \quad W|_{x_n=0} = 0, \end{cases}$$

where $0 < T \leq \infty$. We recall some known results for the system (5.1).

Proposition 5.1. ([7, Proposition 2.3]) *Let $1 < p, q < \infty$. Suppose that $f = \operatorname{div} \mathcal{F}$ with $F \in L^q(0, \infty; L^p(\mathbb{R}_+^n))$, $\mathcal{F}|_{x_n=0} = 0$. Then, the solution W of (5.1) satisfies*

$$(5.2) \quad \|\nabla W\|_{L^q(0, \infty; L^p(\mathbb{R}_+^n))} \leq c \|\mathcal{F}\|_{L^q(0, \infty; L^p(\mathbb{R}_+^n))}.$$

Proposition 5.2. ([6, Theorem 1.2]) *Let $1 < p, q < \infty$. Assume that $f = \operatorname{div} \mathcal{F}$ with $\mathcal{F} \in L^{q_1}(0, \infty; L^{p_1}(\mathbb{R}_+^n))$, where p_1, q_1 satisfy $n/p_1 + 2/q_1 = n/p + 2/q + 1$ with $q_1 \leq q$ and $p_1 \leq p$. Then, there is the unique very weak solution $W \in L^q(0, \infty; L^p(\mathbb{R}_+^n))$ of the Stokes equations (5.1) such that W satisfies*

$$\|W\|_{L^q(0, \infty; L^p(\mathbb{R}_+^n))} \leq c \|\mathcal{F}\|_{L^{q_1}(0, \infty; L^{p_1}(\mathbb{R}_+^n))}.$$

Proposition 5.3. [3, Theorem 2.2] *Let $n + 2 < p < \infty$. Let $f = \operatorname{div} \mathcal{F}$ with $\mathcal{F} \in L^p(\mathbb{R}_+^n \times (0, T))$. Then, there exists unique very weak solution, $w \in L^\infty(\mathbb{R}_+^n \times (0, T))$, of the Stokes equations (5.1) such that*

$$\|W\|_{L^\infty(\mathbb{R}_+^n \times (0, T))} \leq c T^{\frac{1}{2}(1 - \frac{n+2}{p})} \|\mathcal{F}\|_{L^p(\mathbb{R}_+^n \times (0, T))}.$$

Proposition 5.4. [9, Proposition 3.4] *Suppose that $f \in L^{q_1}(0, 1; L^{p_1}(\mathbb{R}_+^n))$ with $1 < p_1, q_1 < \infty$. If p and q satisfy $p > \frac{n}{n-1}p_1$ and $\frac{2}{q_1} + \frac{n}{p_1} - 1 \leq \frac{2}{q} + \frac{n}{p}$, then*

$$(5.3) \quad \|Q\|_{L^q(0, 1; L^p(\mathbb{R}_+^n))} < c \|f\|_{L^{q_1}(0, 1; L^{p_1}(\mathbb{R}_+^n))}.$$

Our aim is to establish the existence of solution v for (2.48) satisfying $v \in L^{2q}(\mathbb{R}_+^n \times (0, \infty))$ and $\nabla v \in L^q(\mathbb{R}_+^n \times (0, \infty))$ for fixed $q > \frac{n+2}{2}$. In order to do that, we consider the iterative scheme for (2.48), which is given as follows: For a positive integer $m \geq 1$

$$\begin{aligned} v_t^{m+1} - \Delta v^{m+1} + \nabla q^{m+1} &= -\operatorname{div} (v^m \otimes v^m + v^m \otimes U + U \otimes v^m + U \otimes U), \\ \operatorname{div} v^{m+1} &= 0 \end{aligned}$$

with homogeneous initial and boundary data, i.e. $v^{m+1}(x, 0) = 0$ and $v^{m+1}(x, t) = 0$ on $\{x_n = 0\}$.

Let $q > \frac{n+2}{2}$. We set $v^1 = 0$. We then have, due to Proposition 5.1 and Proposition 5.2,

$$(5.4) \quad \begin{aligned} \|\nabla v^2\|_{L^q(\mathbb{R}_+^n \times (0, T))} &\leq c \| |U|^2 \|_{L^q(\mathbb{R}_+^n \times (0, T))} \leq c \|U\|_{L^{2q}(\mathbb{R}_+^n \times (0, T))}^2 \leq c \|w\|_{L^{2q}}^2, \\ \|v^2\|_{L^{2q}(\mathbb{R}_+^n \times (0, T))} &\leq c \| |U|^2 \|_{L^{q_1}(0, T; L^q(\mathbb{R}_+^n))} \leq c T^{\frac{1}{2}(1 - \frac{n+2}{2q})} \| |U|^2 \|_{L^q(\mathbb{R}_+^n \times (0, T))} \\ &\leq c T^{\frac{1}{2}(1 - \frac{n+2}{2q})} \|w\|_{L^{2q}}^2, \end{aligned}$$

where $\frac{2}{q_1} + \frac{n}{q} = \frac{n+2}{2q} + 1$. We have $A := \|w\|_{L^{2q}} + \|\nabla w\|_{L^q} \leq c\alpha$, where $\alpha > 0$ is defined in (2.10). We take $T = 1$. Taking $\alpha > 0$ small such that $A < \frac{1}{8c}$, where c is the constant in (5.4) such that

$$\|\nabla v^2\|_{L^q(\mathbb{R}_+^n \times (0, 1))} + \|v^2\|_{L^{2q}(\mathbb{R}_+^n \times (0, 1))} \leq cA^2 < A.$$

Then, iterative arguments show that

$$(5.5) \quad \begin{aligned} \|v^{m+1}\|_{L^{2q}(\mathbb{R}_+^n \times (0, 1))} &\leq c (\|v^m\|^2 + |v^m U| + |U|^2)_{L^q(\mathbb{R}_+^n \times (0, 1))} \\ &\leq 2c \left(\|v^m\|_{L^{2q}(\mathbb{R}_+^n \times (0, 1))}^2 + \|w\|_{L^{2q}}^2 \right) \leq 4cA^2 < A, \\ \|\nabla v^{m+1}\|_{L^q(\mathbb{R}_+^n \times (0, 1))} &\leq c (\|v^m\|^2 + |v^m U| + |U|^2)_{L^q(\mathbb{R}_+^n \times (0, 1))} \\ &\leq 2c \left(\|v^m\|_{L^{2q}(\mathbb{R}_+^n \times (0, 1))}^2 + \|w\|_{L^{2q}(\mathbb{R}_+^n \times (0, 1))}^2 \right) \\ &\leq 4cA^2 < A. \end{aligned}$$

We denote $V^{m+1} := v^{m+1} - v^m$ and $Q^{m+1} := q^{m+1} - q^m$ for $m \geq 1$. We then see that (V^{m+1}, Q^{m+1}) solves

$$\begin{aligned} V_t^{m+1} - \Delta V^{m+1} + \nabla Q^{m+1} &= -\operatorname{div} (V^m \otimes v^m + v^{m-1} \otimes V^m + V^m \otimes U + U \otimes V^m), \\ \operatorname{div} V^{m+1} &= 0 \end{aligned}$$

with homogeneous initial and boundary data, i.e. $V^{m+1}(x, 0) = 0$ and $V^{m+1}(x, t) = 0$ on $\{x_n = 0\}$. It follows from (5.5) that

$$\begin{aligned} & \|\nabla V^{m+1}\|_{L^q(\mathbb{R}_+^n \times (0,1))} + \|V^{m+1}\|_{L^{2q}(\mathbb{R}_+^n \times (0,1))} \\ & \leq c \left(\|V^m v^m\| + |V^m v^{m-1}| + |V^m U| \right)_{L^q(\mathbb{R}_+^n \times (0,1))} \\ & \leq 3cA \|V^m\|_{L^{2q}(\mathbb{R}_+^n \times (0,1))} < \frac{1}{2} \|V^m\|_{L^{2q}(\mathbb{R}_+^n \times (0,1))}. \end{aligned}$$

Therefore, $(v^m, \nabla v^m)$ converges to $(v, \nabla v)$ in $L^{2q}(\mathbb{R}_+^n \times (0, 1)) \times L^q(\mathbb{R}_+^n \times (0, 1))$ such that v solves in the sense of distributions

$$\begin{aligned} v_t - \Delta v + \nabla q &= -\operatorname{div} (v \otimes v + v \otimes U + v \otimes U + U \otimes U), \\ \operatorname{div} v &= 0 \end{aligned}$$

with homogeneous initial and boundary data, i.e. $v(x, 0) = 0$ and $v(x, t) = 0$ on $\{x_n = 0\}$. In addition, via (5.5), we have

$$\|\nabla v\|_{L^q(\mathbb{R}_+^n \times (0,1))}, \quad \|v\|_{L^{2q}(\mathbb{R}_+^n \times (0,1))} \leq A.$$

Furthermore, if $q > n + 2$ and $\|w\|_{L^{2q}}^2 \leq c\alpha$, it follows from Proposition 5.3 that

$$\begin{aligned} (5.6) \quad \|v\|_{L^\infty(\mathbb{R}_+^n \times (0,2))} &\leq c \left(\|v\|^2 + |vU| + |U|^2 \right)_{L^q(\mathbb{R}_+^n \times (0,1))} \\ &\leq 2c \left(\|v\|_{L^{2q}(\mathbb{R}_+^n \times (0,1))}^2 + \|w\|_{L^{2q}}^2 \right) \leq 4cA^2 < A. \end{aligned}$$

This complete the proof of Proposition 2.10. \square

ACKNOWLEDGEMENT

T. Chang is supported by RS-2023-00244630. K. Kang is supported by RS-2024-00336346.

REFERENCES

- [1] T. Chang and H. Choe, *Maximum modulus estimate for the solution of the Stokes equations*, J. Differential Equations, **254** no. 7, 2682-2704 (2013).
- [2] T. Chang and H. Choe, *Quasi-Maximum Modulus Principle for the Stokes Equations*, J. Math. Fluid Mech. **19**, 135–149 (2017).
- [3] T. Chang, H. Choe and K. Kang, *On maximum modulus estimates of the Navier-Stokes equations with non-zero boundary data*, SIAM J. Math. Anal. **50**, no. 3, 3147-3171 (2018).
- [4] T. Chang and B. Jin, *Initial and boundary values for $L_\alpha^q(L_p)$ solution of the Navier-Stokes equations in the half-space*, J. Math. Anal. Appl. **439**, no. 1, 70-90 (2016).
- [5] T. Chang and B. Jin, *Initial-boundary value problem of the Navier–Stokes equations in the half space with nonhomogeneous data*, ANNALI DELL’UNIVERSITA’ DI FERRARA, **65**, 29–56, (2019).
- [6] T. Chang and B. Jin, *Initial and boundary values for $L_\alpha^q(L_p)$ solution of the Navier-Stokes equations in the half-space*, J. Math. Anal. Appl. **439**, no. 1, 70-90 (2016).
- [7] T. Chang and K. Kang, *On Caccioppoli’s inequalities of Stokes equations and Navier-Stokes equations near boundary*, J. Differential Equations, **269** no. 9, 6732–6757 (2020).
- [8] T. Chang and K. Kang, *Local regularity near boundary for the Stokes and Navier-Stokes equations*, Siam J. Math. Anal. **55** no. 5 5051-5085 (2023).
- [9] T. Chang and K. Kang, *Singular weak solutions near boundaries in a half space away from localized force for the Stokes and Navier-Stokes equations*, eprint arXiv:2303.05746.
- [10] H. Chen, S. Liang and T.-P. Tsai *Gradient estimates for the non-stationary Stokes system with the Navier boundary condition*, eprint arXiv:2306.16480.
- [11] K. Kang, *Unbounded normal derivative for the Stokes system near boundary*, Math. Ann., **331** no. 1, 87–109 (2005).
- [12] K. Kang, B. Lai, C.-C. Lai and T.-P. Tsai, *Finite energy Navier-Stokes flows with unbounded gradients induced by localized flux in the half-space*, Trans. Amer. Math. Soc., **375** no. 9, 6701–6746 (2022).
- [13] K. Kang and C. Min, *Local and global regularity for the Stokes and Navier-Stokes equations with boundary data in the half-space*, to appear in Commun. Pure Appl. Anal.
- [14] G. A. Seregin, *Some estimates near the boundary for solutions to the non-stationary linearized Navier-Stokes equations*, (English, Russian summary) Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **271** (2000), Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 31, 204–223, 317; translation in J. Math. Sci. (N. Y.) **115** no. 6, 2820-2831(2003).

- [15] G. A. Seregin and V. Šverák, *On a bounded shear flow in half-space*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) 385 (2010), Kraevye Zadachi Matematicheskoi Fiziki i Smezhnye Voprosy Teorii Funktsii. 41, 200–205, 236; J. Math. Sci. (New York) **178**, no. 3, 353–356 (2011).
- [16] V. A. Solonnikov, *On the theory of nonstationary hydrodynamic potentials. The Navier-Stokes equations: theory and numerical methods*, (Varenna, 2000), Lecture Notes in Pure and Appl. Math., 223, Dekker, New York, 113–129(2002).