

INFINITELY MANY HALF-VOLUME CONSTANT MEAN CURVATURE HYPERSURFACES VIA MIN-MAX THEORY

LIAM MAZUROWSKI AND XIN ZHOU

ABSTRACT. Let (M^{n+1}, g) be a closed Riemannian manifold of dimension $3 \leq n+1 \leq 5$. We show that, if the metric g is generic, then M contains infinitely many geometrically distinct constant mean curvature hypersurfaces, each enclosing half the volume of M . As an essential part of the proof, we develop an Almgren-Pitts type min-max theory for certain non-local functionals of the general form

$$\Omega \mapsto \text{Area}(\partial\Omega) - \int_{\Omega} h + f(\text{Vol}(\Omega)).$$

1. INTRODUCTION

The volume spectrum of a closed, Riemannian manifold M is a sequence of real numbers $\{\omega_p(M)\}_{p \in \mathbb{N}}$ associated to the geometry of M . The volume spectrum was introduced by Gromov [12] as a non-linear analog to the spectrum of the Laplacian. To define the volume spectrum, one first considers all p -sweepouts of M . Heuristically, a p -sweepout is a continuous family of hypersurfaces with the property that for any choice of points $x_1, \dots, x_p \in M$, there is a hypersurface Σ in the family with $x_1, \dots, x_p \in \Sigma$. Then the p -width $\omega_p(M)$ is defined as the min-max value

$$\omega_p(M) = \inf_{p\text{-sweepouts } \Phi} \left[\sup_{\text{hypersurfaces } \Sigma \text{ in } \Phi} \text{Area}(\Sigma) \right].$$

Liokumovich, Marques, and Neves [15] proved that the volume spectrum satisfies a Weyl law characterizing the asymptotic growth of $\omega_p(M)$ as a function of p .

In [22], the authors introduced the half-volume spectrum $\{\tilde{\omega}_p(M)\}_{p \in \mathbb{N}}$. This is defined analogously to the usual volume spectrum, except that one restricts to p -sweepouts by hypersurfaces that are each required to enclose half the volume of M . The half-volume spectrum also satisfies a Weyl law [22]. In this paper, we develop an Almgren-Pitts type min-max theory for finding hypersurfaces associated to the half-volume spectrum. Our main result is the following theorem.

Theorem 1.1. *Assume M^{n+1} is a closed manifold of dimension $3 \leq n+1 \leq 5$. Let g be a generic Riemannian metric on M . Then for each $p \in \mathbb{N}$ there exists an open set $\Omega_p \subset M$ with $\text{Vol}(\Omega_p) = \frac{1}{2} \text{Vol}(M)$ such that $\partial\Omega_p$ is smooth and almost embedded, has non-zero constant mean curvature, and satisfies $\text{Area}(\partial\Omega_p) = \tilde{\omega}_p(M)$.*

Combined with the Weyl law [22, Theorem 1] for the half-volume spectrum, this has the following immediate corollary.

Corollary 1.2. *Assume M^{n+1} is a closed manifold of dimension $3 \leq n+1 \leq 5$. Then, for a generic Riemannian metric g on M , there exist infinitely many geometrically distinct closed constant mean curvature hypersurfaces in M , each enclosing half the volume of M .*

Remark 1.3. In Theorem 1.1 and Corollary 1.2, the word generic is to be interpreted in the sense of Baire category. In other words, there is a subset Γ of the space of all smooth metrics on M

such that Γ is of second Baire category and the conclusions of Theorem 1.1 and Corollary 1.2 hold for all $g \in \Gamma$.

1.1. Background. Min-max theory is a powerful tool for finding critical points. In differential geometry, min-max theory has been used to great success to understand the existence of minimal surfaces, constant mean curvature surfaces, and more general prescribed mean curvature surfaces. In the early 1980s, Almgren [2], Pitts [24], and Schoen-Simon [25] developed a min-max theory for finding critical points of the area functional. Their combined work implies that every closed Riemannian manifold M^{n+1} with dimension $3 \leq n+1 \leq 7$ contains a smooth, closed, embedded minimal hypersurface. Around the same time, Yau [34] conjectured that every closed Riemannian 3-manifold should contain infinitely many closed minimal surfaces. Motivated by Yau's conjecture, Marques and Neves developed a program to understand the Morse theory of the area functional by further refining the Almgren-Pitts min-max theory. In [17], Marques and Neves proved the following theorem.

Theorem 1.4 (Marques-Neves [17]). *Let M^{n+1} be a closed Riemannian manifold of dimension $3 \leq n+1 \leq 7$ and fix an integer $p \in \mathbb{N}$. Then there exists a collection of disjoint closed minimal hypersurfaces $\Sigma_1, \dots, \Sigma_k$ in M together with multiplicities $m_1, \dots, m_k \in \mathbb{N}$ such that $\omega_p(M) = m_1 \text{Area}(\Sigma_1) + \dots + m_k \text{Area}(\Sigma_k)$.*

Theorem 1.4 alone does not suffice to prove Yau's conjecture because of the possible appearance of multiplicities. Indeed, different p -widths could be achieved by the same underlying collection of minimal surfaces, but with different multiplicities. Marques and Neves [19] conjectured that, for a generic choice of metric, all of the multiplicities in Theorem 1.4 can be taken to be 1. In other words, for a generic choice of metric, every ω_p is equal to the area of some (possibly disconnected) minimal hypersurface Σ_p . This *Multiplicity One Conjecture* was proven by the second named author [35], using the prescribed mean curvature regularization of the area functional developed by the second author and J. Zhu [38]. The combined work of Marques and Neves together with the resolution of the Multiplicity One Conjecture gives the following strong answer to Yau's conjecture for generic metrics.

Theorem 1.5 ([19],[35]). *Let M^{n+1} be a closed manifold with dimension $3 \leq n+1 \leq 7$. For a generic choice of metric g on M , there is a sequence of (possibly disconnected) closed minimal hypersurfaces Σ_p such that $\omega_p(M) = \text{Area}(\Sigma_p)$.*

We note that prior to Theorem 1.5, Irie, Marques, and Neves [13] had already established the existence of infinitely many closed minimal hypersurfaces for generic metrics. Moreover, Marques, Neves, and Song [20] proved that for a generic choice of metric on M , some sequence of minimal hypersurfaces becomes equidistributed in M . Finally, Song [29] proved that *every* metric g on M admits infinitely many minimal hypersurfaces, thus completely resolving Yau's conjecture. For a more detailed overview of this subject, we refer to the survey articles [16] and [36].

There is a parallel min-max procedure for constructing minimal hypersurfaces in Riemannian manifolds based on the theory of phase transitions and the Allen-Cahn equation. In this context, Gaspar and Guaraco [10] defined an Allen-Cahn analog to the volume spectrum $\{c_p(M)\}_{p \in \mathbb{N}}$ and proved a corresponding version of Theorem 1.4. They also showed that $c_p(M)$ satisfies a Weyl law [11]. In the Allen-Cahn setting, Chodosh and Mantoulidis [7] proved that the Multiplicity One Conjecture holds in ambient dimension $n+1 = 3$. In [22], the authors also defined an Allen-Cahn version of the half-volume spectrum $\{\tilde{c}_p(M)\}_{p \in \mathbb{N}}$.

In the context of constant mean curvature hypersurfaces, the A^c min-max theory of the second author and J. Zhu [37] can be used to find closed CMC hypersurfaces with prescribed mean curvature c . This min-max theory gives no control over the enclosed volume. It is natural to wonder if there is a corresponding min-max theory for producing *unstable* CMC hypersurfaces with prescribed volume (but with no control over the value of the mean curvature). Of course, by the solution of the isoperimetric problem, M admits a *stable* CMC enclosing volume v for every prescribed volume v . We are asking if there is a meaningful higher parameter version of this.

Intuitively, one expects that the half-volume spectrum should detect critical points of the area functional with a half-volume constraint. In [22], the authors used the Allen-Cahn min-max theory and the work of Bellettini-Wickramasekera [6] to show that each Allen-Cahn half-volume spectrum \tilde{c}_p is achieved by a constant mean curvature hypersurface enclosing half the volume of M , together with a collection of minimal hypersurfaces with even multiplicities. This is an analog to Theorem 1.4 in the half-volume Allen-Cahn setting.

In this paper, we prove a corresponding theorem for the Almgren-Pitts half-volume spectrum $\tilde{\omega}_p$. In fact, we are able to use a prescribed mean curvature regularization to obtain a multiplicity one type result. Thus we obtain Theorem 1.1, which is an analog to Theorem 1.5 in the half-volume Almgren-Pitts setting. As a corollary, we prove the existence of infinitely many half-volume CMCS for generic metrics. As part of the proof, we develop an Almgren-Pitts type min-max theory for certain non-local functionals. These arguments may be of interest in their own right, and we hope that they will prove useful in other contexts.

Remark 1.6. Most of our arguments work in the larger dimension range $3 \leq n+1 \leq 7$, but the extra restriction $3 \leq n+1 \leq 5$ is needed at exactly one point in the proof. This is related to the fact that stable CMCS are known to have diameter bounds in these dimensions [9].

1.2. Motivation and Conjectures. A well-known conjecture of Arnold [3] states that every Riemannian 2-sphere (S^2, g) admits at least two distinct closed curves with constant geodesic curvature κ for every $\kappa > 0$. The following “*Twin Bubble Conjecture*” [36] can be seen as a higher dimensional analog of Arnold’s conjecture; see also [8, 21] for some partial results.

Conjecture 1.7. *Every closed Riemannian manifold (M^{n+1}, g) with $3 \leq n+1 \leq 7$ admits at least two distinct closed hypersurfaces with constant mean curvature c for every constant $c > 0$.*

Our initial motivation for studying half-volume CMCS came from Conjecture 1.7. Naively, one expects that the two CMCS in Conjecture 1.7 come from “sweeping out M in opposite directions.” Of course, topologically all sweepouts of M are homotopic, and so it does not make sense to distinguish the direction of different sweepouts. Nevertheless, by considering the functional

$$\Omega \mapsto \max\{\text{Area}(\partial\Omega) - c \text{Vol}(\Omega), \text{Area}(\partial\Omega) - c \text{Vol}(M \setminus \Omega)\},$$

one can artificially attempt to see both a sweepout and “the sweepout in the opposite direction.” Performing min-max for this functional over all sweepouts of M , we expect to detect either a hypersurface with constant mean curvature c , or a half-volume CMC (with no control over the value of the mean curvature). The first case should correspond to the second c -CMC hypersurface in addition to [37]. Our current work provides a thorough understanding of the latter case via restriction of this functional to all half-volume hypersurfaces.

We also record the following counterpart to Conjecture 1.7, where prescribed mean curvature is replaced by a volume constraint.

Conjecture 1.8. *Every closed Riemannian manifold (M^{n+1}, g) with $3 \leq n+1 \leq 7$ admits at least two distinct closed CMC hypersurfaces enclosing volume v for every $v \in (0, \text{Vol}(M))$.*

1.3. Outline of Proof. In the remainder of the introduction, we will give a sketch of the proof of Theorem 1.1. This section is more technical than the rest of the introduction. We will assume that the reader is familiar with some of the notions from geometric measure theory introduced in Section 2, as well as the basic ideas in the Almgren-Pitts min-max theory.

Let M^{n+1} be a closed Riemannian manifold. The Almgren-Pitts min-max theory works with homotopy classes of maps into the space $\mathcal{Z}_n(M, \mathbb{Z}_2)$ of flat cycles mod 2. Because we work with a volume constraint, we will restrict to the space $\mathcal{B}(M, \mathbb{Z}_2)$ consisting of those $T \in \mathcal{Z}_n(M, \mathbb{Z}_2)$ such that $T = \partial\Omega$ for some Caccioppoli set Ω . Within $\mathcal{B}(M, \mathbb{Z}_2)$, we can further consider the space of half-volume cycles

$$\mathcal{H}(M, \mathbb{Z}_2) = \{T \in \mathcal{B}(M, \mathbb{Z}_2) : T = \partial\Omega \text{ and } \text{Vol}(\Omega) = \frac{1}{2} \text{Vol}(M)\}.$$

At first, one might hope to develop a min-max theory for the area functional directly on the space $\mathcal{H}(M, \mathbb{Z}_2)$.

However, a serious obstacle arises when attempting to design a pull-tight operation for the area functional restricted to $\mathcal{H}(M, \mathbb{Z}_2)$. In the unconstrained Almgren-Pitts min-max scheme, one first obtains a weak solution V in the space of varifolds $\mathcal{V}(M)$ as a limit of a min-max sequence. The pull-tight operation is certain pseudo-gradient flow that ensures that this weak solution V is stationary. The fact that V is stationary (or really just that V has bounded first variation) is then used in an essential way to prove the regularity of V .

In our case, the volume constraint does not make sense on the space of varifolds. Nevertheless, we can replace the space of varifolds by Almgren's VZ space, as in [30]. The space $\text{VZ}(M, \mathbb{Z}_2)$ consists of pairs $(V, T) \in \mathcal{V}(M) \times \mathcal{B}(M, \mathbb{Z}_2)$ such that there is a sequence $T_i \in \mathcal{B}(M, \mathbb{Z}_2)$ with $T_i \rightarrow T$ and $|T_i| \rightarrow V$. In particular, $\text{VZ}(M, \mathbb{Z}_2)$ is the natural space in which to consider the limit of a min-max sequence. The volume constraint makes sense in $\text{VZ}(M, \mathbb{Z}_2)$ and so we can look for a weak solution $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ with $T = \partial\Omega$ and $\text{Vol}(\Omega) = \frac{1}{2} \text{Vol}(M)$.

It is at this point the difficulty arises. One would like to design a pull-tight operation which ensures that the weak solution $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ with $T = \partial\Omega$ is stationary for the area functional with half-volume constraint, i.e.,

$$(1) \quad \delta V(X) = 0, \quad \text{for all } X \in \mathfrak{X}(M) \text{ with } \int_{\Omega} \text{div}(X) = 0.$$

Here $\mathfrak{X}(M)$ denotes the space of C^1 vector fields on M . It does not seem straightforward to achieve this. Moreover, even if such a pull-tight operation does exist, the condition (1) does not seem sufficient to prove regularity. Indeed, condition (1) gives no a priori bound on the first variation of V .

Thus we adopt a different approach. Rather than performing min-max theory for the area functional directly on $\mathcal{H}(M, \mathbb{Z}_2)$, we first perform min-max on all of $\mathcal{B}(M, \mathbb{Z}_2)$ with a functional E given by the area plus a term that penalizes the distance to the space of half-volume cycles. Let $\mathfrak{h} = \frac{1}{2} \text{Vol}(M)$. Then, given $k \in \mathbb{N}$, define

$$E_k(T) = \mathbf{M}(T) + k (\text{Vol}(\Omega) - \mathfrak{h})^2$$

where $\Omega \in \mathcal{C}(M)$ satisfies $\partial\Omega = T$. It is easy to see that $E_k(T)$ does not depend on the choice of Ω . Note that if $\Sigma = \partial\Omega$ is a smooth critical point of E_k then Σ has constant mean curvature

$$(2) \quad |H_{\Sigma}| = 2k |\text{Vol}(\Omega) - \mathfrak{h}| \leq k \text{Vol}(M).$$

Our idea to prove Theorem 1.1 is to first use min-max methods to construct a critical point of E_k for each $k \in \mathbb{N}$, and then to take a limit of these critical points as $k \rightarrow \infty$.

1.3.1. Min-Max Theory. Now assume $3 \leq n+1 \leq 7$. Fix some $k \in \mathbb{N}$. We would like to use min-max theory to find a critical point of E_k . We employ Almgren's VZ space $VZ(M, \mathbb{Z}_2)$ as discussed above. Note that the E_k functional extends naturally to $VZ(M, \mathbb{Z}_2)$ and that we can define the first variation $\delta E_k|_{(V,T)}$ at a point $(V, T) \in VZ(M, \mathbb{Z}_2)$. As a first step, we obtain a weak solution $(V, T) \in VZ(M, \mathbb{Z}_2)$ as a limit of a min-max sequence. We then design a suitable pull-tight argument which ensures that $\delta E_k|_{(V,T)} = 0$. As suggested by (2), one can then verify that V has $k \text{Vol}(M)$ -bounded first variation. Thus working with E_k has the major advantage that V has a priori bounded first variation.

The next step in the usual Almgren-Pitts construction is to use a combinatorial argument to show that some weak solution V is almost minimizing in annuli for the area functional. This almost minimizing property is defined in terms of (ε, δ) -deformations which, roughly speaking, are deformations which decrease the area of a cycle by at least ε without ever increasing it by more than δ . Because the combinatorial argument involves simultaneously applying many different localized (ε, δ) -deformations to a cycle T , it is essential at this point that area functional is *local*. In other words, given an open set $U \subset M$ and two cycles $S, T \in \mathcal{B}(M, \mathbb{Z}_2)$ which agree outside of U , one has

$$\mathbf{M}(T) - \mathbf{M}(S) = \mathbf{M}(T \llcorner U) - \mathbf{M}(S \llcorner U).$$

This property is used to ensure that if a cycle T is modified in several disjoint open sets, the total change in the mass of T is equal to the sum of the local changes.

The E_k functional, on the other hand, is *non-local*. If a cycle $T = \partial\Omega$ is modified in several disjoint open sets, the individual modifications may interact in a complicated way through the volume term. To circumvent this, we observe that the E_k functional still satisfies a certain quasi-locality property. Namely, if T is modified in several disjoint open sets, and each modification individually changes area by much more than it changes volume, then the net change in E_k resulting from performing all the modifications is approximately additive. Based on this, we define a new notion of almost-minimizing for the E_k functional. Heuristically, in our definition, we consider only those (ε, δ) -deformations which additionally change the area by much more than the volume at all times.

This new almost minimizing property still suffices to construct replacements. The first step is to solve a suitable constrained minimization problem. Because (ε, δ) -deformations which do not change the volume are always admissible, solutions to the constrained minimization problem are smooth and volume preserving stable for the area functional. By a result of Bellettini-Chodosh-Wickramasekera [5], the volume preserving stability implies a curvature estimate. This is then enough to construct a replacement as a limit of solutions to the constrained minimization problem. The smooth part of the replacement is either a multiplicity one CMC with non-zero mean curvature, or a collection of minimal hypersurfaces with multiplicities. Finally, one uses replacements in overlapping annuli to prove the regularity of (V, T) . Here the crucial point is to show that replacements in overlapping annuli will either both be minimal, or both have the same non-zero constant mean curvature. Using this, we are able to show that the support of T is a (possibly minimal) CMC bounding a region Ω , and that the remainder of $\text{supp } \|V\|$ consists of minimal hypersurfaces.

In fact, our min-max idea can handle more general functions of volume with very little extra work. See Section 2 for precise definitions.

Theorem 1.9. *Let (M^{n+1}, g) be a closed Riemannian manifold of dimension $3 \leq n+1 \leq 7$. Let $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ be a smooth function satisfying*

$$f(h-t) = f(h+t)$$

for all $t \in [0, \mathfrak{h}]$ where $\mathfrak{h} = \frac{1}{2} \text{Vol}(M)$. Define $E: \mathcal{B}(M, \mathbb{Z}_2) \rightarrow \mathbb{R}$ by

$$E(T) = \mathbf{M}(T) + f(\text{Vol}(\Omega))$$

where Ω satisfies $\partial\Omega = T$. Let Π be the X -homotopy class of a map $\Phi_0: X \rightarrow (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F})$ and assume that $L^E(\Pi) > 0$. Then there exists $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ with $E(V, T) = L^E(\Pi)$. Choose a set Ω with $\partial\Omega = T$ and let $H = -f'(\text{Vol}(\Omega))$.

- (i) If $H \neq 0$, then there exists a smooth, almost-embedded, (not necessarily connected) closed CMC hypersurface $\Lambda = \partial\Omega$, which has mean curvature H with respect to the normal pointing into Ω . Moreover, there exist a (possibly empty) collection of closed embedded minimal hypersurfaces $\Sigma_1, \dots, \Sigma_k$ and a collection of multiplicities $m_1, \dots, m_k \in \mathbb{N}$ such that

$$V = |\Lambda| + \sum_{i=1}^k m_i |\Sigma_i|.$$

The hypersurfaces $\Lambda, \Sigma_1, \dots, \Sigma_k$ are all disjoint.

- (ii) If $H = 0$, then there exists a collection of closed, embedded minimal hypersurfaces $\Lambda_1, \dots, \Lambda_q$ such that $\partial\Omega = \Lambda_1 \cup \dots \cup \Lambda_q$. Moreover, there exist a (possibly empty) collection of closed, embedded minimal hypersurfaces $\Sigma_1, \dots, \Sigma_k$ and a collection of multiplicities $\ell_1, \dots, \ell_q, m_1, \dots, m_k \in \mathbb{N}$ such that

$$V = \sum_{i=1}^q \ell_i |\Lambda_i| + \sum_{i=1}^k m_i |\Sigma_i|.$$

The hypersurfaces $\Lambda_1, \dots, \Lambda_q, \Sigma_1, \dots, \Sigma_k$ are all disjoint.

1.3.2. Regularization. At this point, we employ a prescribed mean curvature regularization to obtain improved regularity as in [35]. Again consider a fixed $k \in \mathbb{N}$. Given a smooth function $h: M \rightarrow \mathbb{R}$ and $\varepsilon > 0$, define $F_{k,\varepsilon}: \mathcal{C}(M) \rightarrow \mathbb{R}$ by

$$F_{k,\varepsilon}(\Omega) = \mathbf{M}(\partial\Omega) - \varepsilon \int_{\Omega} h + k (\text{Vol}(\Omega) - \mathfrak{h})^2.$$

For a good choice of h , we use similar arguments to find critical points of $F_{k,\varepsilon}$. Analogous to the PMC min-max theory [38], these critical points are smooth, almost-embedded multiplicity one hypersurfaces $\Sigma = \partial\Omega$ with mean curvature $H = \varepsilon h + h_0$, where $h_0 = -2k(\text{Vol}(\Omega) - \mathfrak{h})$ is a constant.

As above, we can actually handle more general functionals with very little extra effort. See Section 2 for precise definitions. In particular, see Definition 2.10 for the definition of property (T), which is a condition on the mean curvature of the level sets of h that ensures unique continuation. It is similar to, but stronger than, condition (\dagger) in [38].

Theorem 1.10. *Let (M^{n+1}, g) be a closed Riemannian manifold of dimension $3 \leq n+1 \leq 7$. Let $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ be an arbitrary smooth function, and let $h: M \rightarrow \mathbb{R}$ be a smooth Morse function satisfying property (T); see Definition 2.10. Define $F: \mathcal{C}(M) \rightarrow \mathbb{R}$ by*

$$F(\Omega) = \mathbf{M}(\partial\Omega) - \int_{\Omega} h + f(\text{Vol}(\Omega)).$$

Let Π be the (X, Z) -homotopy class of a map $\Phi_0: X \rightarrow (\mathcal{C}(M), \mathbf{F})$. Assume that

$$L^F(\Pi) > \sup_{z \in Z} F(\Phi_0(z)).$$

Then there exists a smooth, almost-embedded hypersurface $\Sigma = \partial\Omega$ satisfying

$$L^F(\Pi) = F(\Omega).$$

The mean curvature of Σ is given by $H = h - f'(\text{Vol}(\Omega))$, and the touching set of Σ is contained in a countable union of $(n-1)$ -dimensional manifolds.

1.3.3. Half-Volume CMCS. Fix some $p \in \mathbb{N}$. Consider a sequence of half-volume p -sweepouts $\Phi_k : X_k \rightarrow \mathcal{H}(M, \mathbb{Z}_2)$ with

$$\lim_{k \rightarrow \infty} \left[\sup_{x \in X_k} \mathbf{M}(\Phi_k(x)) \right] \rightarrow \tilde{\omega}_p.$$

Let Π_k be the homotopy class of Φ_k . Then

$$L^{E_k}(\Pi_k) \rightarrow \tilde{\omega}_p.$$

We can apply the min-max theory for E_k in the homotopy class of Π_k . We then follow the ideas in [35]. After regularizing to $F_{k,\varepsilon}$, we obtain smooth, multiplicity one hypersurfaces $\Sigma_{k,\varepsilon} = \partial\Omega_{k,\varepsilon}$ with

$$F_{k,\varepsilon}(\Omega_{k,\varepsilon}) \rightarrow L^{E_k}(\Pi_k), \quad \text{as } \varepsilon \rightarrow 0.$$

Moreover, the hypersurfaces $\Sigma_{k,\varepsilon}$ have mean curvature $H(\Sigma_{k,\varepsilon}) = \varepsilon h + h_{k,\varepsilon}$ where $h_{k,\varepsilon}$ is a constant.

Sending $\varepsilon \rightarrow 0$, we obtain weak convergence $\Sigma_{k,\varepsilon} \rightarrow V_k$ as varifolds and $\Omega_{k,\varepsilon} \rightarrow \Omega_k$ as Caccioppoli sets. In fact, for fixed k , the surfaces $\Sigma_{k,\varepsilon}$ all satisfy a certain almost minimizing property, and this is enough to obtain the regularity of V_k . This fact has already been observed in [14] and [30]. In our setting, there is a subtle extra challenge in this step which we address in Section 5. It follows that V_k is induced by either

- (i) an almost-embedded, multiplicity one CMC hypersurface $\Sigma_k = \partial\Omega_k$ with non-zero mean curvature, or
- (ii) a collection of smooth, embedded minimal hypersurfaces with multiplicities.

Moreover, in case (ii), some subcollection of the minimal hypersurfaces bounds Ω_k . Note that we do not a priori rule out case (ii). The primary benefit of the regularization is that it rules out the possibility of obtaining a mixture of CMC components with non-zero mean curvature and minimal components.

Next, we can send $k \rightarrow \infty$, and obtain convergence $V_k \rightarrow V$ as varifolds and $\Omega_k \rightarrow \Omega$ as Caccioppoli sets. It is straightforward to check that $\|V\|(M) \leq \tilde{\omega}_p$ and $\text{Vol}(\Omega) = \mathfrak{h}$. We would like to show that V is induced by a multiplicity one CMC. Again, each V_k satisfies a certain almost-minimizing property. However, unlike for $\Sigma_{k,\varepsilon}$, the varifolds V_k do not have a uniform a priori bound on their first variation. If we can show that the first variation stays bounded, then we can use the almost minimizing property as before to prove regularity of V . In this case, V is induced by either

- (i) an almost-embedded, multiplicity one CMC hypersurface $\Sigma = \partial\Omega$ with non-zero mean curvature, or
- (ii) a collection of smooth, embedded minimal hypersurfaces with multiplicities.

In the second case, some subcollection of the minimal hypersurfaces bounds Ω . We are able to show that, for a generic metric g , no collection of minimal hypersurfaces bounds a region with volume equal to \mathfrak{h} . Thus, for generic metrics, the second possibility cannot occur, and we obtain a multiplicity one CMC hypersurface $\Sigma = \partial\Omega$ enclosing half the volume of M and satisfying $\text{Area}(\Sigma) \leq \tilde{\omega}_p$.

To show that the mean curvature H_k of $\Sigma_k = \text{supp } \|V_k\|$ is uniformly bounded, we can argue as follows. Supposing instead that $H_k \rightarrow \infty$, we can rescale the surfaces Σ_k to obtain new surfaces Σ'_k with constant mean curvature 1. Moreover, we can use the almost minimizing property to ensure that the surfaces Σ'_k are volume preserving stable on a large set. It is at that this point that we need to further restrict to dimensions $3 \leq n+1 \leq 5$. In these dimensions, we can take advantage of the fact that stable 1-CMC surfaces are known to have diameter upper bounds by the work of Elbert-Nelli-Rosenberg [9]. This implies that Σ'_k resembles the union of many unit balls when k is large. At the original scale, this means that Σ_k is the union of many tiny components. However, this structure is incompatible with the half-volume constraint and the uniform area bound.

To complete the proof, it remains to verify the energy identity $\text{Area}(\Sigma) = \tilde{\omega}_p$. Since we know $E_k(V_k, \partial\Omega_k) \rightarrow \tilde{\omega}_p$, it is equivalent to show that

$$k(\text{Vol}(\Omega_k) - \mathfrak{h})^2 \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

In fact, this also follows from the fact that the mean curvature H_k is uniformly bounded. Indeed, since $(V_k, \partial\Omega_k)$ is stationary for E_k , we have $|H_k| = 2k|\text{Vol}(\Omega_k) - \mathfrak{h}|$. Hence

$$k(\text{Vol}(\Omega_k) - \mathfrak{h})^2 = \frac{|H_k|^2}{4k},$$

and this goes to 0 as $k \rightarrow \infty$, as needed.

1.4. Outline of the paper. We collect preliminary materials in Section 2. Section 3 and Section 4 are respectively devoted to the min-max theory for the F and E functionals, and the proof of Theorem 1.10 and Theorem 1.9. In Section 5, we prove several compactness results for min-max solutions associated with the E and F functionals. In Section 6, we apply the min-max theory for E and F and associated compactness results to prove our main Theorem 1.1 for half volume spectrum. In Appendix A, we prove that condition (T) is C^∞ generic among Morse functions, and in Appendix B, we prove a collection of results about closed CMC hypersurfaces in a generic metric.

Acknowledgement. L.M. acknowledges the support of an AMS Simons Travel Grant. X.Z. acknowledges the support by NSF grant DMS-1945178, an Alfred P. Sloan Research Fellowship, and a grant from the Simons Foundation (1026523, XZ). X.Z. also thanks the support of the math department at Princeton University and the Institute of Advanced Study, where part of the work was carried out.

2. PRELIMINARIES

In this section, we begin by reviewing some concepts from geometric measure theory needed in the paper. Then we introduce the functionals we will consider and related min-max notions.

We will need the following concepts from geometric measure theory; see [27]. Let (M^{n+1}, g) be a closed Riemannian manifold.

- Let $\mathcal{C}(M)$ denote the space of all Caccioppoli sets in M .
- Let $\mathcal{V}(M)$ denote the space of all n -dimensional varifolds on M .
- Let $\mathcal{Z}(M, \mathbb{Z}_2)$ denote the space of n -dimensional flat cycles in $M \bmod 2$.
- Given $\Omega \in \mathcal{C}(M)$, the notation $\partial\Omega$ denotes the element of $\mathcal{Z}(M, \mathbb{Z}_2)$ induced by the boundary of Ω .
- Given $T \in \mathcal{Z}(M, \mathbb{Z}_2)$, the notation $|T|$ stands for the varifold induced by T .
- Let $\mathcal{B}(M, \mathbb{Z}_2)$ denote the space of all $T \in \mathcal{Z}(M, \mathbb{Z}_2)$ such that $T = \partial\Omega$ for some $\Omega \in \mathcal{C}(M)$.

- Let $\mathcal{H}(M, \mathbb{Z}_2)$ denote the space of all $T \in \mathcal{Z}(M, \mathbb{Z}_2)$ such that $T = \partial\Omega$ for some $\Omega \in \mathcal{C}(M)$ with $\text{Vol}(\Omega) = \frac{1}{2} \text{Vol}(M)$.
- We use \mathcal{F} to denote the flat topology, \mathbf{F} to denote the \mathbf{F} -topology, and \mathbf{M} to denote the mass topology. By convention, the \mathbf{F} topology on $\mathcal{C}(M)$ is

$$\mathbf{F}(\Omega_1, \Omega_2) = \mathcal{F}(\Omega_1, \Omega_2) + \mathbf{F}(|\partial\Omega_1|, |\partial\Omega_2|).$$

- Let $\text{VZ}(M, \mathbb{Z}_2)$ denote Almgren's VZ space (see [2]).
- Let $\text{VC}(M)$ denote Almgren's VC space (see [2] and [30]).

Because the reader is likely less familiar with Almgren's VZ [2] and VC spaces [30, Definition 1.3], we include the relevant background below. The set $\text{VZ}(M, \mathbb{Z}_2)$ consists of all pairs $(V, T) \in \mathcal{V}(M) \times \mathcal{B}(M, \mathbb{Z}_2)$ such that there is a sequence $T_k \in \mathcal{B}(M, \mathbb{Z}_2)$ with $|T_k| \rightarrow V \in \mathcal{V}(M)$ and $T_k \rightarrow T \in \mathcal{B}(M, \mathbb{Z}_2)$. Note that it may or may not be true that $V = |T|$, but it is always true that $\| |T| \| \leq \|V\|$ as measures. We can endow $\text{VZ}(M, \mathbb{Z}_2)$ with the product metric, so that for any $(V, T), (V', T') \in \text{VZ}(M, \mathbb{Z}_2)$, the \mathcal{F} -distance between them is

$$\mathcal{F}((V, T), (V', T')) = \mathbf{F}(V, V') + \mathcal{F}(T, T').$$

We will use $\text{VZ}(M, \mathcal{F}, \mathbb{Z}_2)$ if we wish to emphasize the metric \mathcal{F} .

Given $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ and a C^1 diffeomorphism $\phi: M \rightarrow M$, define the pushforward $\phi_\#(V, T) = (\phi_\#V, \phi_\#T)$. It is easy to check that $\phi_\#(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ and the mapping $\phi_\#: \text{VZ}(M, \mathbb{Z}_2) \rightarrow \text{VZ}(M, \mathbb{Z}_2)$ is continuous.

Proposition 2.1. *Given any constant $L > 0$, the space*

$$(3) \quad Y_{z,L} = \{(V, T) \in \text{VZ}(M, \mathbb{Z}_2) : \|V\|(M) \leq L\}$$

is a compact metric space.

Proof. The flat topology on $\mathcal{B}(M, \mathbb{Z}_2)$ is metrizable. Moreover, the weak topology on the space $X = \{V \in \mathcal{V}(M) : \|V\|(M) \leq L\}$ is also metrizable. Therefore Y is a metric space. To see that Y is compact, assume that (V_i, T_i) is a sequence in Y . Since X is compact, by passing to a subsequence if necessary, we can assume that $V_i \rightarrow V$ for some $V \in X$. Since $\mathbf{M}(T_i) \leq L$ for all $i \in \mathbb{N}$, by passing to a further subsequence, we can assume that $T_i \rightarrow T$ for some $T \in \mathcal{B}(M, \mathbb{Z}_2)$. It remains to show that $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$. For each $i \in \mathbb{N}$, there is a sequence $T_{i,k} \in \mathcal{B}(M, \mathbb{Z}_2)$ such that $T_{i,k} \rightarrow T_i$ and $|T_{i,k}| \rightarrow V_i$ as $k \rightarrow \infty$. Since all spaces in question are metrizable, the diagonal sequence $T_{i,i}$ therefore satisfies $T_{i,i} \rightarrow T$ and $|\partial T_{i,i}| \rightarrow V$ as $i \rightarrow \infty$. Thus $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$, and it follows that Y is compact. \square

The VC space is entirely analogous but with $\mathcal{B}(M, \mathbb{Z}_2)$ replaced by $\mathcal{C}(M)$. The set $\text{VC}(M)$ consists of all pairs $(V, \Omega) \in \mathcal{V}(M) \times \mathcal{C}(M)$ such that there is a sequence $\Omega_k \in \mathcal{C}(M)$ with $|\partial\Omega_k| \rightarrow V \in \mathcal{V}(M)$ and $\Omega_k \rightarrow \Omega \in \mathcal{C}(M)$. We similarly endow $\text{VC}(M)$ with the product metric, so that for any $(V, \Omega), (V', \Omega') \in \text{VC}(M)$, the \mathcal{F} -distance between them is

$$\mathcal{F}((V, \Omega), (V', \Omega')) = \mathbf{F}(V, V') + \mathcal{F}(\Omega, \Omega').$$

Again, we will write $\text{VC}(M, \mathcal{F})$ if we wish to emphasize the metric \mathcal{F} . Given $(V, \Omega) \in \text{VC}(M)$ and a C^1 diffeomorphism $\phi: M \rightarrow M$, define the pushforward $\phi_\#(V, \Omega) = (\phi_\#V, \phi_\#\Omega)$. It is easy to check that $\phi_\#(V, \Omega) \in \text{VC}(M)$ and the mapping $\phi_\#: \text{VC}(M) \rightarrow \text{VC}(M)$ is continuous.

Proposition 2.2. *Given any constant $L > 0$, the space*

$$(4) \quad Y_{c,L} = \{(V, \Omega) \in \text{VC}(M) : \|V\|(M) \leq L\}$$

is a compact metric space.

2.1. The E Functional. Fix a closed, Riemannian manifold M .

Definition 2.3. Let $\mathfrak{h} = \frac{1}{2} \text{Vol}(M)$.

Let $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ be a non-constant smooth function and assume that f is “even” in the sense that

$$(5) \quad f(\mathfrak{h} - t) = f(\mathfrak{h} + t)$$

for all $t \in [0, \mathfrak{h}]$. Given $T \in \mathcal{B}(M, \mathbb{Z}_2)$, we write $f(\text{Vol}(T))$ to mean $f(\text{Vol}(\Omega))$ where $\Omega \in \mathcal{C}(M)$ satisfies $\partial\Omega = T$. This is well-defined since $|\text{Vol}(\Omega) - \mathfrak{h}| = |\text{Vol}(M \setminus \Omega) - \mathfrak{h}|$ and f satisfies (5).

Definition 2.4. Define the function $E^f: \mathcal{B}(M, \mathbb{Z}_2) \rightarrow \mathbb{R}$ by $E^f(T) = \mathbf{M}(T) + f(\text{Vol}(T))$. In the following, we will sometimes just write E instead of E^f if the choice of f is clear.

Assume $\Sigma = \partial\Omega$ is a smooth hypersurface in M . Let ν denote the inward pointing normal vector to Σ , and let H be the mean curvature of Ω with respect to ν . Let X be a C^1 vector field on M and let ϕ_t be the associated flow.

Proposition 2.5. *The first variation of E is given by*

$$\delta E|_{\Sigma}(X) = \frac{d}{dt} \Big|_{t=0} E(\phi_t(\Sigma)) = - \int_{\Sigma} H \langle X, \nu \rangle - f'(\text{Vol}(\Omega)) \int_{\Sigma} \langle X, \nu \rangle.$$

Thus Σ is critical for E if and only if Σ has constant mean curvature $H = -f'(\text{Vol}(\Omega))$.

Note that the E functional extends to a functional on $\text{VZ}(M, \mathbb{Z}_2)$ by

$$E(V, T) = \|V\|(M) + f(\text{Vol}(T)).$$

Moreover, if $T_i \in \mathcal{Z}(M, \mathbb{Z}_2)$ is a sequence with $T_i \rightarrow T$ and $|T_i| \rightarrow V$ then $E(T_i) \rightarrow E(V, T)$. It is also possible to define the first variation of E on $\text{VZ}(M, \mathbb{Z}_2)$.

Proposition 2.6. *The first variation of E on $\text{VZ}(M, \mathbb{Z}_2)$ is given by*

$$(6) \quad \begin{aligned} \delta E|_{(V, T)}(X) &= \frac{d}{dt} \Big|_{t=0} E((\phi_t)_{\#}(V, T)) \\ &= \delta V(X) + f'(\text{Vol}(\Omega)) \int_{\Omega} \text{div}(X) \end{aligned}$$

where $\Omega \in \mathcal{C}(M)$ satisfies $\partial\Omega = T$.

A pair $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ is called stationary for E if $\delta E|_{(V, T)}(X) = 0$ for all C^1 vector fields X on M .

Proposition 2.7. *Fix a C^1 vector field X on M . Assume that $(V_i, T_i) \rightarrow (V, T) \in \text{VZ}(M, \mathbb{Z}_2)$. Then $\delta E_{(V_i, T_i)}(X) \rightarrow \delta E_{(V, T)}(X)$.*

Proof. Let (V_i, T_i) and (V, T) be as in the statement of the theorem. It is possible to choose Caccioppoli sets Ω_i and Ω so that $\partial\Omega_i = T_i$ and $\partial\Omega = T$ and $\Omega_i \rightarrow \Omega$ as Caccioppoli sets. Since X is fixed, we have

$$\int_{G_n(M)} \text{div}_P X(x) dV_i(x, P) \rightarrow \int_{G_n(M)} \text{div}_P X(x) dV(x, P)$$

as $i \rightarrow \infty$. Again since X is fixed we have $\text{Vol}(\Omega_i) \rightarrow \text{Vol}(\Omega)$ and

$$\int_{\Omega_i} \text{div}(X) \rightarrow \int_{\Omega} \text{div}(X).$$

The result now follows from the first variation formula. \square

Corollary 2.8. *The set $\{(V, T) \in \text{VZ}(M, \mathbb{Z}_2) : (V, T) \text{ is stationary for } E\}$ is closed as a subset of $\text{VZ}(M, \mathbb{Z}_2)$.*

2.2. The F Functional. Fix a smooth, non-constant function $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$. Also fix a smooth Morse function $h: M \rightarrow \mathbb{R}$.

Definition 2.9. Given a regular point $x \in M$ for h , let $\Gamma(x)$ be the level set of h passing through x . Then define $v(h, x)$ to be the vanishing order at x of the mean curvature $H_{\Gamma(x)}$, regarded as a function on $\Gamma(x)$.

Definition 2.10. Let $h: M \rightarrow \mathbb{R}$ be a smooth Morse function. We say that h satisfies property (T) provided

(T) For every regular point x of h , we have $v(h, x) < \infty$.

In the following, fix a smooth Morse function $h: M \rightarrow \mathbb{R}$ satisfying property (T). In Appendix A, we show that the set of smooth Morse functions satisfying property (T) is dense in $C^\infty(M)$.

Definition 2.11. Define the functional $A^h: \mathcal{C}(M) \rightarrow \mathbb{R}$ by

$$A^h(\Omega) = \mathbf{M}(\partial\Omega) - \int_{\Omega} h.$$

Then define the functional $F^{h,f}: \mathcal{C}(M) \rightarrow \mathbb{R}$ by

$$F^{h,f}(\Omega) = A^h(\Omega) + f(\text{Vol}(\Omega)).$$

In the following, we will sometimes just write F instead of $F^{h,f}$ if the choice of h and f is clear.

Assume $\Sigma = \partial\Omega$ is a smooth hypersurface in M . Let ν denote the inward pointing normal vector to Σ , and let H be the mean curvature of Σ with respect to ν . Let X be a C^1 vector field on M and let ϕ_t be the associated flow.

Proposition 2.12. *The first variation of F is given by*

$$\delta F|_{\Omega}(X) = \left. \frac{d}{dt} \right|_{t=0} F(\phi_t(\Omega)) = \int_{\Sigma} (h - H - f'(\text{Vol}(\Omega))) \langle X, \nu \rangle.$$

Thus Ω is a critical point for F if and only if Σ has constant mean curvature given by $H = h - f'(\text{Vol}(\Omega))$.

Property (T) is used to control the touching set of almost-embedded hypersurfaces with mean curvature $h + h_0$ for some constant h_0 .

Proposition 2.13. *Assume that h satisfies property (T). Let Σ be an almost-embedded hypersurface with mean curvature $H = h + h_0$ for some constant h_0 . Then the touching set $S(\Sigma)$ is contained in a countable union of $(n - 1)$ -dimensional manifolds.*

Proof. For every regular point $x \in M$ for h , the proof of [38, Theorem 3.11] shows that there is some $r > 0$ for which $\mathcal{S}(\Sigma) \cap B_r(x)$ is contained in a countable union of $(n - 1)$ -dimensional manifolds. Let C be the finite set of all critical points of h . Let G_n be a sequence of closed subsets of $M \setminus C$ with $\cup_{n=1}^{\infty} G_n = M \setminus C$. Then each G_n is covered by finitely many balls B with the property that $\mathcal{S}(\Sigma) \cap B$ is contained in a countable union of $(n - 1)$ -dimensional submanifolds. Taking the union of these balls over all n , it follows that $\mathcal{S}(\Sigma) \cap (M \setminus C)$ is contained in a countable union of $(n - 1)$ -dimensional manifolds. Of course, the finite set C is also contained in a finite union of $(n - 1)$ -dimensional manifolds, and the result follows. \square

Note that the F functional extends to a functional on $\text{VC}(M)$ by

$$F(V, \Omega) = \|V\|(M) - \int_{\Omega} h + f(\text{Vol}(\Omega)).$$

Moreover, if $\Omega_i \in \mathcal{C}(M)$ is a sequence with $\Omega_i \rightarrow \Omega$ and $|\partial\Omega_i| \rightarrow V$ then $F(\Omega_i) \rightarrow F(V, \Omega)$. It is also possible to define the first variation of F on $\text{VC}(M)$.

Proposition 2.14. *The first variation of F on $\text{VC}(M)$ is given by*

$$\begin{aligned} \delta F|_{(V, \Omega)}(X) &= \left. \frac{d}{dt} \right|_{t=0} F((\phi_t)_{\#}(V, \Omega)) \\ (7) \quad &= \delta V(X) - \int_{\Omega} \text{div}(hX) + f'(\text{Vol}(\Omega)) \int_{\Omega} \text{div}(X). \end{aligned}$$

A pair $(V, \Omega) \in \text{VC}(M)$ is called stationary for F if $\delta F|_{(V, \Omega)}(X) = 0$ for all C^1 vector fields X on M .

Proposition 2.15. *Fix a C^1 vector field X on M . Assume that $(V_i, \Omega_i) \rightarrow (V, \Omega) \in \text{VC}(M)$. Then $\delta F|_{(V_i, \Omega_i)}(X) \rightarrow \delta F|_{(V, \Omega)}(X)$.*

Corollary 2.16. *The set $\{(V, \Omega) \in \text{VC}(M) : (V, \Omega) \text{ is stationary for } F\}$ is closed in $\text{VC}(M)$.*

2.3. Min-Max Notions. Let X be a cubical subcomplex of the cell complex $I(m, k_0)$ for some $m, k_0 \in \mathbb{N}$. Here $I(m, k) = I(1, k) \otimes \cdots \otimes I(1, k)$ (m -times), where $I(1, k)$ denotes the complex on $I = [0, 1]$ whose 1-cells and 0-cells are, respectively,

$$[1, 3^{-k}], [3^{-k}, 2 \cdot 3^{-k}], \dots, [1 - 3^{-k}, 1] \text{ and } [0], [3^{-k}], \dots, [1 - 3^{-k}], [1].$$

We refer to [35, Appendix A] for a summary of notions; (see also [18, Section 2.1]). We will define both absolute and relative homotopy classes.

2.3.1. Absolute Homotopy Classes. Given a continuous map $\Phi_0 : X \rightarrow (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F})$, we let Π be the collection of all sequences of continuous maps $\{\Phi_i : X \rightarrow (\mathcal{B}(M, \mathbb{Z}_2), \mathbf{F})\}$ such that, for each i , there exists a flat continuous homotopy map

$$\begin{aligned} H_i : X \times [0, 1] &\rightarrow (\mathcal{B}(M, \mathbb{Z}_2), \mathcal{F}), \\ H_i(x, 0) &= \Phi_0(x), \\ H_i(x, 1) &= \Phi_i(x). \end{aligned}$$

Definition 2.17. Such a sequence $\{\Phi_i\}_{i \in \mathbb{N}}$ is called an X -homotopy sequence of mappings into $\mathcal{B}(M, \mathbb{Z}_2)$, and Π is called the X -homotopy class of Φ_0 .

Definition 2.18. Fix a functional $E : \mathcal{B}(M, \mathbb{Z}_2) \rightarrow \mathbb{R}$. Define the min-max value of Π with respect to E by

$$L^E(\Pi) = \inf_{\{\Phi_i\} \in \Pi} \limsup_{i \rightarrow \infty} \left[\sup_{x \in X} E(\Phi_i(x)) \right].$$

A sequence $\{\Phi_i\} \in \Pi$ is called a critical sequence for E if

$$L^E(\{\Phi_i\}) := \limsup_{i \rightarrow \infty} \left(\sup_{x \in X} E(\Phi_i(x)) \right) = L^E(\Pi).$$

The critical set $\mathcal{K}(\{\Phi_i\})$ associated to a critical sequence $\{\Phi_i\}$ is the set of all $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ such that there exist $x_{i_j} \in X$ with $|\Phi_{i_j}(x_{i_j})| \rightarrow V$, and $\Phi_{i_j}(x_{i_j}) \rightarrow \Omega$, and $E(\Phi_{i_j}(x_{i_j})) \rightarrow L^E(\Pi)$. In the following, we will sometimes abbreviate L^E to L when the choice of E is clear.

Proposition 2.19. *The critical set $\mathcal{K}(\{\Phi_i\})$ is a compact subset of $\text{VZ}(M, \mathcal{F}, \mathbb{Z}_2)$.*

2.3.2. Relative Homotopy Classes. Let $Z \subset X$ be a cubical subcomplex. Given a continuous map $\Phi_0 : X \rightarrow (\mathcal{C}(M), \mathbf{F})$, we let Π be the collection of all sequences of continuous maps $\{\Phi_i : X \rightarrow (\mathcal{C}(M), \mathbf{F})\}$ such that, for each i , there exists a flat continuous homotopy map

$$\begin{aligned} H_i : X \times [0, 1] &\rightarrow (\mathcal{C}(M), \mathbf{F}), \\ H_i(x, 0) &= \Phi_0(x), \\ H_i(x, 1) &= \Phi_i(x), \end{aligned}$$

and, moreover,

$$\limsup_{i \rightarrow \infty} \left[\sup_{(z,t) \in Z \times [0,1]} \mathbf{F}(H_i(z,t), \Phi_0(z)) \right] \rightarrow 0$$

as $i \rightarrow \infty$.

Definition 2.20. Such a sequence $\{\Phi_i\}_{i \in \mathbb{N}}$ is called an (X, Z) -homotopy sequence of mappings into $\mathcal{C}(M)$, and Π is called the (X, Z) -homotopy class of Φ_0 .

Definition 2.21. Fix a functional $F : \mathcal{C}(M) \rightarrow \mathbb{R}$. Define the min-max value of Π with respect to F by

$$L^F(\Pi) = \inf_{\{\Phi_i\} \in \Pi} \limsup_{i \rightarrow \infty} \left[\sup_{x \in X} F(\Phi_i(x)) \right].$$

A sequence $\{\Phi_i\} \in \Pi$ is called a critical sequence for F if

$$L^F(\{\Phi_i\}) := \limsup_{i \rightarrow \infty} \left(\sup_{x \in X} F(\Phi_i(x)) \right) = L^F(\Pi).$$

The critical set $\mathcal{K}(\{\Phi_i\})$ associated to a critical sequence $\{\Phi_i\}$ is the set of all $(V, \Omega) \in \text{VC}(M)$ such that there exist $x_{i_j} \in X$ with $|\Phi_{i_j}(x_{i_j})| \rightarrow V$, $\Phi_{i_j}(x_{i_j}) \rightarrow \Omega$, and $F(\Phi_{i_j}(x_{i_j})) \rightarrow L^F(\Pi)$. In the following, we will sometimes abbreviate L^F to L when the choice of F is clear.

Proposition 2.22. *The critical set $\mathcal{K}(\{\Phi_i\})$ is a compact subset of $\text{VC}(M, \mathcal{F})$.*

2.4. Regularity and Curvature Estimates. When developing a min-max theory for the F functional, we will need to analyze volume constrained local minimizers of A^h .

Proposition 2.23. *Fix $3 \leq n+1 \leq 7$. Assume that Ω is a local minimizer for A^h in an open set U subject to a volume constraint. In other words, for any Θ with $\text{supp}(\Omega - \Theta) \subset U$ and $\text{Vol}(\Theta) = \text{Vol}(\Omega)$, we have $A^h(\Theta) \geq A^h(\Omega)$. Then $\partial\Omega$ is smooth and embedded in U .*

Proof. It suffices to verify condition (2) of [23, Proposition 3.1]. As in [23], we can find two points $p_1, p_2 \in U$ and deformations $\Omega_{i,t}$ such that

- (i) $\Omega_{i,0} = \Omega$, $\Omega_{i,t} \supset \Omega$ for $t > 0$, and $\Omega_{i,t} \subset \Omega$ for $t < 0$
- (ii) $\text{supp}(\Omega_{i,t} - \Omega)$ is contained in a small ball centered at p_i ,
- (iii) $|\mathcal{H}^n(\partial\Omega_{i,t}) - \mathcal{H}^n(\partial\Omega)| \leq C_1 |\text{Vol}(\Omega_{i,t}) - \text{Vol}(\Omega)|$

for $i = 1, 2$. Condition (i) implies that $|\text{Vol}(\Omega \Delta \Omega_{i,t})| = |\text{Vol}(\Omega) - \text{Vol}(\Omega_{i,t})|$. Therefore condition (iii) implies that

$$\begin{aligned} |A^h(\Omega_{i,t}) - A^h(\Omega)| &\leq |\mathcal{H}^n(\partial\Omega_{i,t}) - \mathcal{H}^n(\partial\Omega)| + \left| \int_{\Omega} h - \int_{\Omega_{i,t}} h \right| \\ &\leq C_1 |\text{Vol}(\Omega_{i,t}) - \text{Vol}(\Omega)| + (\sup |h|) \text{Vol}(\Omega \Delta \Omega_{i,t}) \\ &\leq C_2 |\text{Vol}(\Omega) - \text{Vol}(\Omega_{i,t})|. \end{aligned}$$

Now we can argue as in [38].

Let $p \in U$ and choose a small number $r > 0$ so that $B_r(p) \subset U$. We can suppose r is small enough that $B_r(p)$ is far away from a neighborhood of p_i for some choice of $i \in \{1, 2\}$. Consider any Θ with $\text{supp}(\Omega - \Theta) \subset B_r(p)$. Then it must be the case that

$$(8) \quad A^h(\Theta) - A^h(\Omega) \geq -C_2 |\text{Vol}(\Theta) - \text{Vol}(\Omega)|.$$

Otherwise, we could restore volume near p_i using the deformation above, and this would create a new competitor with the same volume as Ω but lower A^h . Let $A = \mathcal{H}^n(\partial\Omega)$ and $A' = \mathcal{H}^n(\partial\Theta)$. Then (8) implies that

$$A' - A \geq -C_2 |\text{Vol}(\Theta) - \text{Vol}(\Omega)| - \left| \int_{\Theta} h - \int_{\Omega} h \right| \geq -C_3 \text{Vol}(\Theta \Delta \Omega).$$

As in [38], this inequality implies condition (2) of [23, Proposition 3.1] provided r is sufficiently small. The regularity then follows. \square

Bellettini, Chodosh, and Wickramasekera proved curvature estimates for volume preserving stable CMCs. In particular, they proved the following Bernstein type theorem.

Theorem 2.24 ([5] Proposition 3). *Fix $3 \leq n+1 \leq 7$. Assume that $\Sigma \subset \mathbb{R}^{n+1}$ is a connected, embedded, volume preserving stable minimal hypersurface with no singularities. Further suppose that $\mathcal{H}^n(\Sigma \cap B_r) \leq \Lambda r^n$ for some constant $\Lambda \geq 1$ and all $r \geq 0$. Then Σ is a hyperplane.*

For the min-max theory, we need to extend the Bellettini-Chodosh-Wickramasekera curvature estimates to local minimizers of A^h with a volume constraint.

Proposition 2.25. *Fix an open set $U \subset M$. Assume that $\Sigma = \partial\Omega$ is smooth and properly embedded in U with mean curvature $H_{\Sigma} = h + h_0$ for some constant h_0 . Assume that Ω is volume preserving stable for A^{h+h_0} in U . Then there is a constant C depending only on U , $\|h\|_{C^3}$, an upper bound for $\text{Area}(\Sigma)$, and an upper bound for $\|H_{\Sigma}\|_{\infty}$ such that*

$$\|A_{\Sigma}\|^2(x) \leq \frac{C}{\text{dist}(x, \partial U)^2}$$

for all $x \in \Sigma \cap U$.

Proof. This follows from Theorem 2.24 by a standard argument. \square

Finally we record the following Simons type theorem for volume preserving stable cones.

Proposition 2.26. *Assume that Σ is a smooth, embedded, volume preserving stable minimal cone in $\mathbb{R}^{n+1} \setminus \{0\}$ with $3 \leq n+1 \leq 7$. Then Σ is a hyperplane.*

Proof. Note that Σ must be strongly stable outside some ball $B(0, R)$. Since $r\Sigma = \Sigma$ for every $r > 0$, it follows that Σ is strongly stable outside $B(0, rR)$ for every $r > 0$. Thus Σ is strongly stable in $\mathbb{R}^{n+1} \setminus \{0\}$ and the result follows from the usual Simons theorem. \square

3. RELATIVE MIN-MAX FOR F

The goal of this section is to develop a relative min-max theory for the F functional. Throughout this section, M is a fixed Riemannian manifold, $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ is a non-constant smooth function, $h: M \rightarrow \mathbb{R}$ is a smooth Morse function satisfying property (T), and we abbreviate $F = F^{h,f}$.

3.1. Pull Tight. We will employ a pull-tight operation to ensure that every point in the critical set is stationary for F . The construction is relatively standard, and so we shall be brief. The goal is to prove the following proposition.

Proposition 3.1. *Let Π be the (X, Z) -homotopy class of a continuous map $\Phi_0 : X \rightarrow (\mathcal{C}(M), \mathbf{F})$. Assume that*

$$L^F(\Pi) > \sup_{z \in Z} F(\Phi_0(z)).$$

Let $\{\Phi_i\}$ be a critical sequence for Π . There is another critical sequence $\{\Xi_i\}$ such that $\mathcal{K}(\{\Xi_i\}) \subset \mathcal{K}(\{\Phi_i\})$ and every $(V, \Omega) \in \mathcal{K}(\{\Xi_i\})$ is stationary for F .

Proof. Let $L = L^F(\{\Phi_i\}) + 1$. Recall that $Y = Y_{c,L}$ defined in (4) is the set of pairs in VC with mass bounded by L . Let

$$Y_0 = \{(V, \Omega) \in Y : (V, T) \text{ is stationary for } F\} \cup \Phi_0(Z).$$

Consider the concentric annuli of Y_0 under the \mathcal{F} -metric:

$$Y_1 = \{(V, \Omega) \in Y : \mathcal{F}((V, T), Y_0) \geq \frac{1}{2}\},$$

$$Y_j = \{(V, \Omega) \in Y : \frac{1}{2^j} \leq \mathcal{F}((V, \Omega), Y_0) \leq \frac{1}{2^{j-1}}\}, j \in \mathbb{N}, j \geq 2.$$

By compactness of Y_j under the \mathcal{F} -metric and the continuity of first variation of $(V, \Omega) \rightarrow \delta F_{V,\Omega}(\mathcal{X})$ for a fixed vector field \mathcal{X} (Proposition 2.15), for each $j \in \mathbb{N}$, we can find $c_j > 0$, such that for all $(V, \Omega) \in Y_j$, there exists a C^1 vector field $\mathcal{X}_{V,\Omega}$ on M with

$$\|\mathcal{X}_{V,\Omega}\|_{C^1} \leq 1, \quad \delta F_{V,\Omega}(\mathcal{X}_{V,T}) \leq -c_j < 0.$$

Next, we can follow the same procedure as in [30, Step 2 in Section 2.2] to construct a mapping

$$\mathcal{X} : Y \rightarrow \mathfrak{X}(M),$$

where $\mathfrak{X}(M)$ denotes the space of C^1 vector fields on M , such that the following lemma holds.

Lemma 3.2. *The map \mathcal{X} is continuous under the C^1 topology on $\mathfrak{X}(M)$. Moreover, there exist continuous functions $g : (0, \infty) \rightarrow (0, \infty)$ and $\rho : (0, \infty) \rightarrow (0, \infty)$ with $\lim_{t \rightarrow 0} g(t) = 0$ and $\lim_{t \rightarrow 0} \rho(t) = 0$ such that*

$$\delta F_{V',\Omega'}(\mathcal{X}(V, \Omega)) \leq -g(\mathcal{F}((V, \Omega), Y_0))$$

for all $(V', T'), (V, T) \in Y$ satisfying $\mathcal{F}((V', \Omega'), (V, \Omega)) \leq \rho(\mathcal{F}((V, \Omega), Y_0))$.

The next step is to construct a map into the space of isotopies. Again this can be done as in [30, Step 3 in Section 2.2]. Given $(V, \Omega) \in Y$ and $t > 0$, define $(V_t, \Omega_t) = \phi_{V,\Omega}(t)_\#(V, \Omega)$ where $\phi_{V,\Omega}(t)$ denotes the time t flow of the vector field $\mathcal{X}(V, \Omega)$.

Lemma 3.3. *There exist continuous functions $\mathcal{T} : (0, \infty) \rightarrow (0, \infty)$ and $\mathcal{L} : (0, \infty) \rightarrow (0, \infty)$ with $\lim_{t \rightarrow 0} \mathcal{T}(t) = 0$ and $\lim_{t \rightarrow 0} \mathcal{L}(t) = 0$ such that for all $(V, \Omega) \in Y$ it holds that*

$$F(V_{\mathcal{T}(\gamma)}, \Omega_{\mathcal{T}(\gamma)}) \leq F(V, \Omega) - \mathcal{L}(\gamma),$$

where we have abbreviated $\gamma = \mathcal{F}((V, \Omega), Y_0)$.

Define a map $\Psi : Y \times [0, 1] \rightarrow Y$ by

$$\Psi((V, \Omega), t) = (V_{\mathcal{T}(\gamma)t}, \Omega_{\mathcal{T}(\gamma)t}),$$

where again we have abbreviated $\gamma = \mathcal{F}((V, \Omega), Y_0)$. Then, for each i , define $\Xi_i : X \rightarrow \mathcal{C}(M)$ by $\Xi_i(x) = \pi \circ \Psi(|\Phi_i(x)|, \Phi_i(x), 1)$ where $\pi(V, \Omega) = \Omega$. It is straightforward to verify from the properties of Ψ that the sequence $\{\Xi_i\}$ is as required. \square

3.2. Replacements and the Almost Minimizing Property. In this section, we define a suitable almost-minimizing property and then use it to construct replacements. Note that previous Almgren-Pitts min-max schemes have always used for a local functional. For example, in the case of the A^h functional, if a competitor Ω is modified in a small open set U , it is possible to determine the change in A^h without knowledge of $\Omega \setminus (M - U)$. However, this is not the case for E . If a competitor Ω is modified in a small open set U , it is impossible to determine the change in $f(\text{Vol}(\Omega))$ without knowledge of $\Omega \setminus (M - U)$. The key observation is that the functional F still satisfies a quasi-locality type property if we restrict to modifications that change A^h faster than volume.

Definition 3.4. Define $a = \max_{v \in [0, \text{Vol}(M)]} |f'(v)|$. Note that $a > 0$ since we assume f is not constant.

Definition 3.5. Define $b = \sup_M |h|$.

Proposition 3.6. Assume U_1, U_2, \dots, U_N are disjoint open sets in M . Assume that $\Omega \in \mathcal{C}(M)$. Fix a small $\delta > 0$. Suppose there are sets Θ_k with $\text{supp}(\Omega - \Theta_k) \subset U_k$ and

$$2a |\text{Vol}(\Omega) - \text{Vol}(\Theta_i)| \leq A^h(\Omega) - A^h(\Theta_i) + \delta$$

for $k = 1, 2, \dots, N$. Let

$$\Omega^* = \Omega \setminus (U_1 \cup \dots \cup U_N) + \sum_{k=1}^N \Theta_k \setminus U_k.$$

Then there is an estimate

$$F(\Omega^*) \leq F(\Omega) + \frac{1}{2} \sum_{i=1}^N (A^h(\Theta_k) - A^h(\Omega)) + \frac{N\delta}{2}.$$

Proof. Note that

$$\text{Vol}(\Omega^*) = \text{Vol}(\Omega \setminus (U_1 \cup \dots \cup U_N)) + \sum_{k=1}^N \text{Vol}(\Theta_k \cap U_k).$$

Observe that

$$\text{Vol}(\Theta_k \cap U_k) - \text{Vol}(\Omega \cap U_k) = \text{Vol}(\Theta_k) - \text{Vol}(\Omega) \leq \frac{A^h(\Omega) - A^h(\Theta_k)}{2a} + \frac{\delta}{2a}.$$

for $k = 1, \dots, N$. It follows that

$$|\text{Vol}(\Omega^*) - \text{Vol}(\Omega)| \leq \frac{N\delta}{2a} + \sum_{k=1}^N \frac{A^h(\Omega) - A^h(\Theta_k)}{2a}.$$

Next observe that

$$\begin{aligned} f(\text{Vol}(\Omega^*)) &\leq f(\text{Vol}(\Omega)) + a |\text{Vol}(\Omega^*) - \text{Vol}(\Omega)| \\ &\leq f(\text{Vol}(\Omega)) + \frac{N\delta}{2} + \sum_{k=1}^N \frac{A^h(\Omega) - A^h(\Theta_k)}{2}. \end{aligned}$$

Thus we have

$$\begin{aligned}
F(\Omega^*) &= A^h(\Omega) + \sum_{k=1}^N (A^h(\Theta_k) - A^h(\Omega)) + f(\text{Vol}(\Omega^*)) \\
&\leq A^h(\Omega) + \frac{N\delta}{2} + \frac{1}{2} \sum_{k=1}^N (A^h(\Theta_k) - A^h(\Omega)) + f(\text{Vol}(\Omega)) \\
&= F(\Omega) + \frac{1}{2} \sum_{k=1}^N (A^h(\Theta_k) - A^h(\Omega)) + \frac{N\delta}{2}.
\end{aligned}$$

This proves the result. \square

3.2.1. The Almost-Minimizing Property. The above quasi-locality property is the motivation for the following definition of F almost-minimizing sets.

Definition 3.7. Let U be an open subset of M . Let ν denote either the \mathcal{F} , \mathbf{F} , or \mathbf{M} norm. Fix constants $\varepsilon, \delta > 0$ and $\gamma \geq 0$. Fix an element $\Omega \in \mathcal{C}(M)$. Let $\{\Omega_i\}_{i=1,\dots,k}$ be a sequence in $\mathcal{C}(M)$. Assume that

- (i) $\Omega_1 = \Omega$,
- (ii) $\text{supp}(\Omega - \Omega_i) \subset U$ for all i ,
- (iii) $2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_i)| \leq A^h(\Omega) - A^h(\Omega_i) + \gamma$ for all i ,
- (iv) $\nu(\partial\Omega_i, \partial\Omega_{i+1}) < \delta$ for all i ,
- (v) $A^h(\Omega_i) \leq A^h(\Omega) + \delta$ for all i .

We say Ω is $(F, \varepsilon, \delta, \gamma, \nu)$ -almost-minimizing in U if for all sequences $\{\Omega_i\}_{i=1,\dots,k}$ as above, we have $A^h(\Omega_k) \geq A^h(\Omega) - \varepsilon$.

Remark 3.8. Note that we allow $\gamma = 0$. When $\nu = \mathcal{F}$, we will always choose $\gamma = 0$. When $\nu = \mathbf{M}$, we will always choose $\gamma = \delta > 0$ to allow room for interpolation.

Definition 3.9. We define F -almost-minimizers. Let $(V, \Omega) \in \text{VC}(M)$ and let U be an open subset of M .

- (a) We say (V, Ω) is (F, \mathcal{F}) -almost-minimizing in U if there is a sequence $\Omega_i \in \mathcal{C}(M)$ such that Ω_i is $(F, \varepsilon_i, \delta_i, 0, \mathcal{F})$ -almost-minimizing in U , $|\partial\Omega_i| \rightarrow V$, $\Omega_i \rightarrow \Omega$, and $\varepsilon_i, \delta_i \rightarrow 0$.
- (b) We say (V, Ω) is (F, \mathbf{M}) -almost-minimizing in U if there is a sequence $\Omega_i \in \mathcal{C}(M)$ such that Ω_i is $(F, \varepsilon_i, \delta_i, \delta_i, \mathbf{M})$ almost-minimizing in U , $|\partial\Omega_i| \rightarrow V$, $\Omega_i \rightarrow \Omega$, and $\varepsilon_i, \delta_i \rightarrow 0$.

Proposition 3.10. Assume that Ω is $(F, \varepsilon, \rho, \rho, \mathbf{M})$ -almost-minimizing in U . If $\delta \leq \rho$ is small enough, then Ω is $(F, \varepsilon, \delta, 0, \mathcal{F})$ -almost-minimizing in any open set W compactly contained in U .

Proof. We rely on an interpolation process from [37] and [38]. Assume δ is small enough that we can apply [37, Lemma A.1] with $\eta = \min\{\rho/(2a+1), \rho/2\}$ and $c = \max\{1, b\}$. We can assume that $\delta \leq \eta$. Assume that Ω_1 and Ω_2 satisfy $\mathcal{F}(\partial\Omega_1, \partial\Omega_2) < \delta$ and $\text{supp}(\Omega - \Omega_i) \subset W$ and $A^h(\Omega_i) \leq A^h(\Omega) + \delta$ and

$$2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_i)| \leq A^h(\Omega) - A^h(\Omega_i)$$

for $i = 1, 2$. By [37, Lemma A.1] (also see [38, Lemma A.1]), there exists a sequence $\Omega_1 = \Lambda_0, \Lambda_1, \dots, \Lambda_m = \Omega_2 \in \mathcal{C}(M)$ such that for each $j = 1, \dots, m-1$ we have

- (a) $\text{supp}(\Lambda_j - \Omega) \subset U$,
- (b) $\mathbf{M}(\partial\Lambda_k - \partial\Lambda_{j+1}) \leq \eta$,
- (c) $A^h(\Lambda_j) \leq \max\{A^h(\Omega_1), A^h(\Omega_2)\} + \eta$,

(d) $|\text{Vol}(\Lambda_j) - \text{Vol}(\Omega_i)| \leq \eta$, for $i = 1, 2$.

One verifies immediately that (ii), (iv), and (v) hold along the sequence Λ_j with $\nu = \mathbf{M}$ and with δ replaced by ρ . Choose k so that $\max\{A^h(\Omega_1), A^h(\Omega_2)\} = A^h(\Omega_k)$. Observe that

$$\begin{aligned} 2a|\text{Vol}(\Omega) - \text{Vol}(\Lambda_j)| &= 2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_k) + \text{Vol}(\Omega_k) - \text{Vol}(\Lambda_j)| \\ &\leq 2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_k)| + 2a\eta, \end{aligned}$$

and therefore

$$\begin{aligned} A^h(\Omega) - A^h(\Lambda_j) &= A^h(\Omega) - A^h(\Omega_k) + A^h(\Omega_k) - A^h(\Lambda_j) \\ &\geq 2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_k)| - \eta \\ &\geq 2a|\text{Vol}(\Omega) - \text{Vol}(\Lambda_j)| - (2a + 1)\eta \\ &\geq 2a|\text{Vol}(\Omega) - \text{Vol}(\Lambda_j)| - \rho. \end{aligned}$$

Thus (iii) also holds along the sequence Λ_j with $\gamma = \rho$. It is easy to see that the above construction can be used to prove that if Ω is not $(F, \varepsilon, \delta, 0, \mathcal{F})$ almost-minimizing in W then Ω is not $(F, \varepsilon, \rho, \rho, \mathbf{M})$ almost-minimizing in U . \square

Proposition 3.11. *If $(V, \Omega) \in \text{VC}(M)$ is (F, \mathbf{M}) -almost-minimizing in U then (V, Ω) is (F, \mathcal{F}) -almost-minimizing in any open set W compactly contained in U .*

Proof. Assume that (V, Ω) is (F, \mathbf{M}) -almost-minimizing in U and let W be an open set compactly contained in U . According to the definition, there is a sequence $\Omega_i \in \mathcal{C}(M)$ such that Ω_i is $(F, \varepsilon_i, \rho_i, \rho_i, \mathbf{M})$ almost-minimizing in U and $|\partial\Omega_i| \rightarrow V$ and $\Omega_i \rightarrow \Omega$ and $\varepsilon_i, \rho_i \rightarrow 0$. By Proposition 3.10, for each i , there is a $\delta_i \leq \rho_i$ such that Ω_i is $(F, \varepsilon_i, \delta_i, 0, \mathcal{F})$ -almost-minimizing in W . Since $\varepsilon_i, \delta_i \rightarrow 0$, this witnesses that (V, Ω) is (F, \mathcal{F}) -almost-minimizing in W . \square

Recall that $a = \sup |f'|$ and $b = \sup |h|$.

Proposition 3.12. *Assume that $(V, \Omega) \in \text{VC}(M)$ is (F, \mathcal{F}) -almost-minimizing in U . Then V has $(2a + b)$ -bounded first variation in U .*

Proof. We will prove the contrapositive. Suppose that V does not have $(2a + b)$ -bounded first variation in U . Then there is a smooth vector field X compactly supported in U such that

$$\delta V(X) < -(2a + b + \varepsilon_0) \int_M |X| d\mu_V$$

for some $\varepsilon_0 > 0$. By continuity and the first variation formula, there is an $\varepsilon_1 > 0$ such that for all $\Omega \in \mathcal{C}(M)$ with $\mathbf{F}(|\partial\Omega|, V) < 2\varepsilon_1$ we have

$$\begin{aligned} \delta|\partial\Omega|(X) + 2a \int_M |X| d\mu_{\partial\Omega} &\leq 0, \\ \delta A^h|_{\Omega}(X) &\leq -\frac{\varepsilon_0}{2} \int_M |X| d\mu_{\partial\Omega}. \end{aligned}$$

Fix some Ω with $\mathbf{F}(|\partial\Omega|, V) < \varepsilon_1$. Deforming Ω along the flow of X for a uniform small time τ yields an \mathbf{F} -continuous family $(\Omega_t)_{t \in [0, \tau]}$ such that $\mathbf{F}(|\partial\Omega_t|, V) < 2\varepsilon_1$ for all $t \in [0, \tau]$. Observe that $t \mapsto A^h(\Omega_t)$ is decreasing for $t \in [0, \tau]$. We also have

$$A^h(\Omega_\tau) \leq A^h(\Omega) - \varepsilon_2$$

where $\varepsilon_2 > 0$ is a uniform constant that does not depend on the choice of Ω . Finally, note that for any $t \in [0, \tau]$ we have

$$\begin{aligned} 2a |\text{Vol}(\Omega) - \text{Vol}(\Omega_t)| &= 2a \left| \int_0^t \delta \text{Vol}|_{\Omega_t}(X) dt \right| \\ &\leq 2a \int_0^t \left(\int_{\partial\Omega_t} |X| \right) dt \leq - \int_0^t \delta |\partial\Omega_t|(X) dt. \end{aligned}$$

Therefore $\text{Area}(\partial\Omega) - \text{Area}(\partial\Omega_t) \geq 2a |\text{Vol}(\Omega) - \text{Vol}(\Omega_t)|$ for all $t \in [0, \tau]$. Discretizing the flow witnesses that Ω cannot be $(F, \varepsilon_2, \delta, 0, \mathcal{F})$ almost-minimizing in U for any $\delta > 0$. \square

Remark 3.13. Note that we cannot use a similar argument to prove that (V, Ω) is stable in a suitable sense for A^h in U as in [30], mainly due to the volume constraint in Definition 3.7(iii). This will bring in extra challenges to show compactness in Section 5.

Definition 3.14. An element $(V, \Omega) \in \text{VC}(M)$ is called F -almost-minimizing in annuli if for each $x \in M$ there is a number $\rho(x) > 0$ such that (V, Ω) is (F, \mathcal{F}) almost-minimizing in every annulus $\text{An}(x, s, r)$ with $s < r < \rho(x)$.

Proposition 3.15. Let Π be the (X, Z) -homotopy class of a continuous map $\Phi_0: X \rightarrow (\mathcal{C}(M), \mathbf{F})$, where X is a cubical subcomplex of $I(m, k)$ for some $m, k \in \mathbb{N}$. Assume that

$$L^F(\Pi) > \sup_{z \in Z} F(\Phi_0(z)).$$

Choose a pulled-tight critical sequence $\{\Phi_i\}$ and let $\mathcal{K} = \mathcal{K}(\{\Phi_i\})$ be the critical set. Then there is an element $(V, \Omega) \in \mathcal{K}$ which is F -almost minimizing in annuli. In fact, there exists some $(V, \Omega) \in \mathcal{K}$ satisfying the following stronger property:

- (R) There is a number $N = N(m)$ depending only on m such that for any collection of N concentric annuli $\text{An}(x, s_1, r_1), \dots, \text{An}(x, s_N, r_N)$ with $2r_j < s_{j+1}$, (V, Ω) is (F, \mathcal{F}) almost minimizing in at least one of the annuli.

Proof. Applying discretization [35, Theorem 1.11], we can find a homotopy sequence $\{\phi_i\}$ such that $L(\{\phi_i\}) = L(\Pi)$ and $\mathcal{K}(\{\phi_i\}) = \mathcal{K}(\{\Phi_i\})$. Assume for contradiction that no element in \mathcal{K} satisfies property (R). Then by Proposition 3.11, no element in \mathcal{K} satisfies property (R) with (E, \mathcal{F}) almost-minimizing replaced by (E, \mathbf{M}) almost-minimizing. Thus we can apply Parts 1-19 of the Almgren-Pitts combinatorial construction [24, Theorem 4.10] to $\{\phi_i\}$. This produces a homotopic sequence $\{\psi_i\}$.

Note that in Part 20, one can use Proposition 3.6 to verify that

$$L(\{\psi_i\}) < L(\{\phi_i\}) - \varepsilon$$

for some $\varepsilon > 0$. Let us be more precise. Consider disjoint annuli A_1, A_2, \dots, A_K . Fix $\Omega \in \mathcal{C}(M)$. As part of the combinatorial construction, one obtains small constants $\varepsilon, \delta > 0$ and Caccioppoli sets $\Omega_k(j)$, $k \in \{1, \dots, K\}$, $j \in \{1, \dots, J\}$ satisfying

- (i) $\Omega_k(1) = \Omega$,
- (ii) $\text{supp}(\Omega_k(j) - \Omega) \subset A_k$ for all j and k ,
- (iii) $2a |\text{Vol}(\Omega) - \text{Vol}(\Omega_k(j))| \leq A^h(\Omega) - A^h(\Omega_k(j)) + \delta$ for all j and k ,
- (iv) $\mathbf{M}(\partial\Omega_k(j) - \partial\Omega_k(j+1)) < \delta$ for all j and k ,
- (v) $A^h(\Omega_k(j)) \leq A^h(\Omega) + \delta$ for all j and k ,
- (vi) $A^h(\Omega_k(J)) \leq A^h(\Omega) - \varepsilon$.

For a choice of integers $j_1, \dots, j_N \in \{1, \dots, J\}$, one then considers a set of the form

$$\Omega^* = \Omega \setminus (A_1 \cup \dots \cup A_N) + \sum_{k=1}^K \Omega_k(j_k) \setminus A_k$$

and needs to estimate the difference $F(\Omega) - F(\Omega^*)$. This can be accomplished by applying Proposition 3.6 with $U_k = A_k$ and $\Theta_k = \Omega_k(j_k)$.

Applying interpolation [35, Theorem 1.12] to $\{\psi_i\}$ then gives a sequence

$$\{\Psi_i: X \rightarrow (\mathcal{C}(M), \mathbf{F})\}$$

in the (X, Z) -homotopy class Π with $\sup_{x \in X} F(\Psi_i(x)) < L(\Pi) - \varepsilon/2$ for all large i . This is a contradiction. \square

3.2.2. A Constrained Minimization Problem. Consider the following constrained minimization problem. Assume that Ω is $(F, \varepsilon, \delta, 0, \mathcal{F})$ almost minimizing in U . Let K be a compact subset of U . Let $\mathcal{A} = \mathcal{A}(\Omega, K, \delta)$ be the set of all $\Theta \in \mathcal{C}(M)$ such that there is a sequence $\{\Omega_i\}_{i=1, \dots, k}$ in $\mathcal{C}(M)$ satisfying

- (i') $\Omega_1 = \Omega$,
- (ii') $\text{supp}(\Omega - \Omega_i) \subset K$ for all i ,
- (iii') $2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_i)| \leq A^h(\Omega) - A^h(\Omega_i)$ for all i ,
- (iv') $\mathcal{F}(\partial\Omega_i, \partial\Omega_{i+1}) < \delta$ for all i ,
- (v') $A^h(\Omega_i) \leq A^h(\Omega) + \delta$ for all i ,

and $\Theta = \Omega_k$. We aim to minimize A^h among elements of \mathcal{A} .

Proposition 3.16. *There exists $\Theta \in \mathcal{A}$ such that $A^h(\Theta) = \inf_{\Xi \in \mathcal{A}} A^h(\Xi)$.*

Proof. Choose a minimizing sequence $\Theta_j \in \mathcal{A}$. Since the mass of $\partial\Theta_j$ is uniformly bounded, by passing to a subsequence if necessary, we can assume that $\Theta_j \rightarrow \Theta$ in $\mathcal{C}(M)$. Since A^h is lower-semicontinuous in the flat topology, one has

$$A^h(\Theta) \leq \inf_{\Xi \in \mathcal{A}} A^h(\Xi).$$

To complete the proof, it remains to show that $\Theta \in \mathcal{A}$. Select J large enough so that $\mathcal{F}(\partial\Theta_J, \partial\Theta) < \delta$. There is a sequence $\{\Omega_i\}_{i=1, \dots, k}$ satisfying (i')-(v') and such that $\Theta_J = \Omega_k$. Consider the extended sequence $\{\Omega_j\}_{j=1, \dots, k+1}$ where $\Omega_{k+1} = \Theta$. It is easy to see that (i'), (ii'), (iv'), (v') hold for the extended sequence. Note that

$$2a|\text{Vol}(\Omega) - \text{Vol}(\Theta_j)| \leq A^h(\Omega) - A^h(\Theta_j)$$

for all j . Passing to the limit as $j \rightarrow \infty$ and using the lower semi-continuity of A^h , it follows that

$$2a|\text{Vol}(\Omega) - \text{Vol}(\Theta)| \leq A^h(\Omega) - A^h(\Theta).$$

Thus (iii') holds as well, and it follows that $\Theta \in \mathcal{A}$, as needed. \square

Proposition 3.17. *Assume Ω is $(F, \varepsilon, \delta, 0, \mathcal{F})$ almost-minimizing in U and let K be a compact subset of U . Let Θ be a solution to the constrained minimization problem. Then Θ is $(F, \varepsilon, \delta, 0, \mathcal{F})$ almost-minimizing in U .*

Proof. Since $\Theta \in \mathcal{A}$, there is a sequence $\{\Omega_i\}_{i=1, \dots, k}$ satisfying (i')-(v') such that $\Theta = \Omega_k$. Now consider any sequence $\{\Theta_i\}_{i=1, \dots, j}$ such that

- (i'') $\Theta_1 = \Theta$,
- (ii'') $\text{supp}(\Theta - \Theta_i) \subset U$ for all i ,

- (iii'') $2a|\text{Vol}(\Theta) - \text{Vol}(\Theta_i)| \leq A^h(\Theta) - A^h(\Theta_i)$ for all i ,
- (iv'') $\mathcal{F}(\partial\Theta_i, \partial\Theta_{i+1}) < \delta$ for all i ,
- (v'') $A^h(\Theta_i) \leq A^h(\Theta) + \delta$ for all i .

Now consider the concatenated sequence $\Omega_1, \dots, \Omega_{k-1}, \Omega_k = \Theta_1, \Theta_2, \dots, \Theta_k$. It is clear that (i), (ii), (iv), and (v) hold along the extended sequence. Finally note that

$$\begin{aligned} 2a|\text{Vol}(\Omega) - \text{Vol}(\Theta_i)| &= 2a|\text{Vol}(\Omega) - \text{Vol}(\Theta) + \text{Vol}(\Theta) - \text{Vol}(\Theta_i)| \\ &\leq 2a|\text{Vol}(\Omega) - \text{Vol}(\Theta)| + 2a|\text{Vol}(\Theta) - \text{Vol}(\Theta_i)| \\ &\leq A^h(\Omega) - A^h(\Theta) + A^h(\Theta) - A^h(\Theta_i) \\ &= A^h(\Omega) - A^h(\Theta_i) \end{aligned}$$

for $i = 2, \dots, j$. Thus (iii) also holds along the extended sequence. Since Ω is almost-minimizing, it follows that $A^h(\Theta_k) \geq A^h(\Omega) - \varepsilon \geq A^h(\Theta) - \varepsilon$. \square

Proposition 3.18. *Let Θ be a solution to the constrained minimization problem. Then $\partial\Theta$ locally minimizes A^h with respect to volume preserving modifications in the interior of K . In the interior of K , $\partial\Theta$ is smooth with mean curvature $H = h + h_0$ for some constant h_0 and, moreover, $|H| \leq 2a + b$. The surface $\partial\Theta$ is a volume preserving stable critical point for A^{h+h_0} in the interior of K .*

Proof. There is a sequence $\{\Omega_i\}_{i=1, \dots, k}$ satisfying (i')-(v') such that $\Theta = \Omega_k$. We will show that if any of the claimed properties fail, it is possible to extend this sequence so as to contradict that Θ solved the constrained minimization problem.

Consider an open set $W \subset K$ with $\text{Vol}(W) < \delta$. Assume for contradiction that $\partial\Theta$ does not minimize area with respect to volume preserving modifications in W . Then there exists $\Xi \in \mathcal{C}(M)$ such that $\text{supp}(\Xi - \Theta) \subset W$ and $\text{Vol}(\Xi) = \text{Vol}(\Theta)$ and $A^h(\Xi) < A^h(\Theta)$. Consider the extended sequence $\{\Omega_i\}_{i=1, \dots, k+1}$ where $\Omega_{k+1} = \Xi$. It is easy to see that the extended sequence satisfies (i'), (ii'), and (v'). We have $\mathcal{F}(\partial\Omega_k, \partial\Omega_{k+1}) \leq \text{Vol}(W) < \delta$ and so (iv') holds. Finally observe that

$$\begin{aligned} A^h(\Omega) - A^h(\Xi) &\geq A^h(\Omega) - A^h(\Theta) \\ &\geq 2a|\text{Vol}(\Omega) - \text{Vol}(\Theta)| \\ &= 2a|\text{Vol}(\Omega) - \text{Vol}(\Xi)| \end{aligned}$$

and so (iii') holds. It follows that $\Xi \in \mathcal{A}$, and this is a contradiction.

Proposition 2.23 implies that $\partial\Theta$ is induced by a multiplicity one, smooth, embedded surface with mean curvature $H = h + h_0$ in the interior of K . Here h_0 is a constant. We claim that $|H| \leq 2a + b$. Suppose for contradiction that $|H| > 2a + b$ at some point $p \in \partial\Theta \cap \text{int}(K)$. Fix a small number $r > 0$. The assumption on the mean curvature gives the existence of $\Xi \in \mathcal{C}(M)$ with $\text{supp}(\Xi - \Theta) \subset B_r(p)$ and

$$\mathbf{M}(\partial\Theta) - \mathbf{M}(\partial\Xi) \geq (2a + b)|\text{Vol}(\Theta) - \text{Vol}(\Xi)| > 0.$$

Moreover, we can ensure that $\partial\Xi$ lies to one side of $\partial\Theta$ so that we have $\text{Vol}(\Xi \Delta \Theta) = |\text{Vol}(\Theta) - \text{Vol}(\Xi)|$. Note that this implies

$$\begin{aligned} A^h(\Theta) - A^h(\Xi) &\geq (2a + b)|\text{Vol}(\Theta) - \text{Vol}(\Xi)| - \int_{\Theta} h + \int_{\Xi} h \\ &\geq (2a + b)|\text{Vol}(\Theta) - \text{Vol}(\Xi)| - b\text{Vol}(\Theta \Delta \Xi) = 2a|\text{Vol}(\Theta) - \text{Vol}(\Xi)|. \end{aligned}$$

Consider the extended sequence $\{\Omega_i\}_{i=1,\dots,k+1}$ where $\Omega_{k+1} = \Xi$. It is easy to see that (i'), (ii'), (iv'), and (v') hold provided r is small enough. It remains to note that

$$\begin{aligned} A^h(\Omega) - A^h(\Xi) &= A^h(\partial\Omega) - A^h(\Theta) + A^h(\Theta) - A^h(\Xi) \\ &\geq 2a|\text{Vol}(\Omega) - \text{Vol}(\Theta)| + 2a|\text{Vol}(\Theta) - \text{Vol}(\Xi)| \\ &\geq 2a|\text{Vol}(\Omega) - \text{Vol}(\Xi)|. \end{aligned}$$

Thus (iii') holds so $\Xi \in \mathcal{A}$ and this is a contradiction.

It remains to show that Θ is a volume preserving stable critical point for A^{h+h_0} in the interior of K . Note that it is equivalent to show that Θ is a volume preserving stable critical point for A^h . Suppose this is not the case. Then there is a smooth function $\varphi: \partial\Theta \rightarrow \mathbb{R}$ with compact support in $\text{int}(K)$ such that

$$\int_{\partial\Theta \cap K} \varphi = 0, \text{ and } \int_{\partial\Theta \cap K} |\nabla \varphi|^2 - (|A|^2 + \text{Ric}(\nu, \nu) - \partial_\nu h) \varphi^2 < 0.$$

According to [4, Lemma 2.2], it is possible to find a smooth family $\partial\Theta_t$, $t \in [-\varepsilon_0, \varepsilon_0]$ such that

- (a) $\Theta_0 = \Theta$
- (b) $\text{supp}(\Theta_t - \Theta) \subset \text{int}(K)$ for all $t \in [-\varepsilon_0, \varepsilon_0]$
- (c) $\text{Vol}(\Theta_t) = \text{Vol}(\Theta)$ for all $t \in [-\varepsilon_0, \varepsilon_0]$,
- (d) one has

$$\left. \frac{d}{dt} \right|_{t=0} \partial\Theta_t = \varphi \cdot \nu$$

where ν is the unit normal pointing into Θ .

By the second variation of A^h and the volume preserving condition, there is an $\varepsilon_1 < \varepsilon_0$ such that $A^h(\Theta_t) < A^h(\Theta)$ for all $t \in (0, \varepsilon_1)$. Fix a very small $t_0 > 0$, and consider the extended sequence $\{\Omega_i\}_{i=1,\dots,k+1}$ where $\Omega_{k+1} = \Theta_{t_0}$. It is easy to verify that the extended sequence satisfies (i')-(v') provided t_0 is small enough, and this contradicts that Θ solved the constrained minimization problem. \square

3.2.3. Replacements. Next we turn to the construction of replacements.

Proposition 3.19. *Assume that V has $(2a+b)$ -bounded first variation, and also that (V, Ω) is (F, \mathcal{F}) -almost-minimizing in an open set U . Let K be a compact subset of U . There exists an element $(V^*, \Omega^*) \in \text{VC}(M)$ called a replacement for (V, Ω) such that*

- (i) $(V, \Omega) \llcorner (M - K) = (V^*, \Omega^*) \llcorner (M - K)$,
- (ii) $A^h(V, \Omega) = A^h(V^*, \Omega^*)$,
- (iii) (V^*, Ω^*) is (F, \mathcal{F}) almost-minimizing in U ,
- (iv) V^* has $(2a+b)$ -bounded first variation,
- (v) (V^*, Ω^*) , when restricted to the interior of K , is a limit of some solutions to the constrained minimization problems.

Proof. Let (V, Ω) and U and K be as in the assumptions of the theorem. By definition, there exists a sequence Ω_i such that $|\partial\Omega_i| \rightarrow |V|$ and $\Omega_i \rightarrow \Omega$ and Ω_i is $(F, \varepsilon_i, \delta_i, 0, \mathcal{F})$ -almost-minimizing in U and $\varepsilon_i, \delta_i \rightarrow 0$. Let Θ_i be the solution to a corresponding constrained minimization problem, i.e.,

$$A^h(\Theta_i) = \inf_{\Theta \in \mathcal{A}(\Omega_i, K, \delta_i)} A^h(\Theta).$$

By compactness, passing to a subsequence if necessary, we can assume that $|\partial\Theta_i| \rightarrow V^*$ and $\Theta_i \rightarrow \Omega^*$. We claim that (V^*, Ω^*) has the necessary properties.

Properties (i) and (v) are clear. To check property (iii), note that by Proposition 3.17, each Θ_i is $(F, \varepsilon_i, \delta_i, 0, \mathcal{F})$ -almost-minimizing in U . Since $|\partial\Theta_i| \rightarrow V^*$ and $\Theta_i \rightarrow \Omega^*$, it follows that (V^*, Ω^*) is (F, \mathcal{F}) -almost-minimizing in U . To see that (ii) holds, observe that

$$A^h(\Omega_i) - \varepsilon_i \leq A^h(\Theta_i) \leq A^h(\Omega_i)$$

for all i . We can let $i \rightarrow \infty$ in the above equation to deduce that

$$A^h(V, \Omega) = A^h(V^*, \Omega^*)$$

which is property (iii). It remains to check (iv). Since V has $(2a+b)$ -bounded first variation, it follows that V^* has $(2a+b)$ -bounded variation on $M-K$. Also V^* has $(2a+b)$ -bounded variation on U by Proposition 3.12. This implies that V^* has $(2a+b)$ -bounded variation on all of M . \square

Remark 3.20. Note that since the replacement (V^*, Ω^*) is still almost-minimizing in U , it is therefore possible to obtain a replacement (V^{**}, Ω^{**}) for (V^*, Ω^*) in any compact set $K' \subset U$, and so on.

To prove the regularity of replacements, we need to use the curvature estimates for volume preserving stable critical points of A^{h+h_0} .

Proposition 3.21. *Let (V^*, Ω^*) be a replacement for (V, Ω) in K . Then, in the interior of K , the varifold V^* is induced by a smooth, almost-embedded hypersurface Σ with multiplicity one. The surface Σ has mean curvature $H = h + h_0$ for some constant h_0 and its touching set is contained in a countable union of $(n-1)$ -dimensional submanifolds. The surface Σ coincides with $\partial\Omega^*$ in the interior of K . The surface Σ is volume preserving stable as an immersion for A^{h+h_0} in the interior of K . Finally, in the interior of K , there is a curvature estimate*

$$\|A_\Sigma\|^2(x) \leq \frac{C}{\text{dist}(x, \partial K)^2}$$

where C is a constant that depends only on M , a , b , and $\|V\|(M)$.

Proof. In the interior of K , the replacement V^* is a limit of smooth, embedded, volume preserving stable critical points Σ_k for A^{h+h_k} . Here $\{h_k\}$ is a sequence of constants. Moreover, the surfaces Σ_k have uniformly bounded mean curvature. Hence the regularity follows from the curvature estimates of Proposition 2.25. By Proposition 2.13, the touching set is contained in a countable union of $(n-1)$ -dimensional submanifolds since h satisfies property (T). \square

We conclude by investigating the tangent cones to almost-minimizers. We show that if (V, Ω) is (F, \mathcal{F}) -almost-minimizing in annuli, then every tangent cone to V is an integer multiple of a plane. In the following, $\eta_{p,r}$ denotes the map which rescales by a factor of $1/r$ centered at p .

Proposition 3.22. *Assume that (V, Ω) has $(2a+b)$ -bounded first variation and is (F, \mathcal{F}) -almost minimizing in a set U . Then for any sequence $p_i \rightarrow p \in U$ and any sequence of scales $r_i \rightarrow 0$, every varifold limit $\bar{V} = (\eta_{p_i, r_i})_\# V$ is an integer multiple of a complete, embedded minimal hypersurface. Moreover, every varifold tangent to V is an integer multiple of a hyperplane.*

Proof. Choose a sequence of points $p_i \rightarrow p$ and a sequence of scales $r_i \rightarrow 0$. Consider a varifold limit of the form

$$\bar{V} = \lim (\eta_{p_i, r_i})_\# V.$$

We can argue exactly as in [37, Lemma 5.10] to show that \bar{V} admits volume preserving stable minimal replacements in annuli. More precisely, for any annulus $\text{An} \subset T_p M$, there exists a varifold \bar{V}^* on $T_p M$ such that

- (i) $\overline{V} \llcorner (T_p M \setminus \text{An}) = \overline{V}^* \llcorner (T_p M \setminus \text{An})$,
- (ii) $\|\overline{V}\|(B) = \|\overline{V}^*\|(B)$ for a large ball B containing An ,
- (iii) the varifold \overline{V}^* is induced by a smooth, embedded, volume preserving stable minimal hypersurface with multiplicity in An .

In fact, this process can be iterated any number of times to construct replacements for \overline{V}^* , replacements for these replacements, and so on.

It is well known that the existence of replacements implies the regularity of \overline{V} when volume preserving stability is replaced by strong stability in (iii), c.f. [37, Appendix C]. However, strong stability is used only to get curvature estimates and to apply Simons theorem. Since volume preserving stable minimal hypersurfaces also have curvature estimates and satisfy Simons theorem (Proposition 2.26), the existence of volume preserving stable replacements is also enough to prove the regularity of \overline{V} .

Finally, once the regularity of all such \overline{V} is known, we can argue exactly as in [37, Proposition 5.11] to show that every varifold tangent to V is an integer multiple of a hyperplane. \square

3.3. Regularity. The goal of this subsection is to prove the regularity of the min-max pair (V, Ω) . Given a smooth almost-embedded hypersurface Σ with mean curvature $h + h_0$, let $\mathcal{R}(\Sigma)$ be the set of embedded points of Σ and let $\mathcal{S}(\Sigma)$ denote the touching set of Σ .

Next we construct replacements for (V, Ω) on overlapping annuli. To prove regularity, it is essential that the consecutive replacements can be selected to have matching mean curvature. The next proposition shows that this is always possible.

Proposition 3.23. *Assume (V, Ω) is F -almost-minimizing in annuli and fix a point $p \in \text{supp } \|V\|$. Fix sufficiently small numbers $s_1 < r_1$ and let (V^*, Ω^*) be a replacement for (V, Ω) in $\text{An}(p, s_1, r_1)$. Let Σ^* be the smooth, almost-embedded hypersurface inducing V^* in $\text{An}(p, s_1, r_1)$ and suppose Σ^* has mean curvature $h + h_0$. Choose $s_1 < r_2 < r_1$ so that $\partial B_{r_2}(p)$ intersects Σ^* and the countable union of manifolds containing $\mathcal{S}(\Sigma^*)$ transversally. Fix any $s < s_1$ and let $(V_s^{**}, \Omega_s^{**})$ be a replacement for (V^*, Ω^*) in $\text{An}(p, s, r_2)$. Let Σ_s^{**} be the smooth, almost-embedded hypersurface inducing V_s^{**} in $\text{An}(p, s, r_2)$. Then Σ_s^{**} has mean curvature $H = h + h_0$ for the same constant h_0 .*

Proof. Choose $s < s_1 < r_2 < r_1$ as in the statement of the proposition. Observe that the maximum principle ensures that Σ^* and Σ_s^{**} are both non-empty provided r_1 is sufficiently small. Note that Σ_s^{**} has mean curvature $H = h + h_1$ for some constant h_1 , and we need to show that $h_0 = h_1$.

Suppose to the contrary that $h_0 \neq h_1$. Pick a point $p \in \mathcal{R}(\Sigma^*) \cap \text{An}(p, r_2, r_1)$ and pick a point $q \in \mathcal{R}(\Sigma_s^{**}) \cap \text{An}(p, s, r_2)$. Let ν be the unit normal pointing into Ω_s^{**} . Let X be a vector field supported in a small neighborhood of $\{p, q\}$ such that

$$\int_{\Sigma^*} \langle X, \nu \rangle = - \int_{\Sigma_s^{**}} \langle X, \nu \rangle \neq 0.$$

Note that this implies $\delta \text{Vol}|_{\Omega_s^{**}}(X) = 0$. Since $h_0 \neq h_1$, after replacing X by $-X$ if necessary, we obtain that

$$\delta A^h|_{V_s^{**}, \Omega_s^{**}}(X) < 0.$$

Hence, by continuity, there are $\varepsilon_1, \varepsilon_2 > 0$ such that

$$\delta A^h|_{\Theta}(X) < -\varepsilon_2, \quad |\delta \text{Vol}|_{\Theta}(X)| \leq \frac{\varepsilon_2}{4a}$$

for all $\Theta \in \mathcal{C}(M)$ with $\mathcal{F}(|\partial\Theta|, \Theta), (V_s^{**}, \Omega_s^{**})) < 2\varepsilon_1$. It follows that for any $\Theta \in \mathcal{C}(M)$ with $\mathcal{F}(|\partial\Theta|, \Theta), (V_s^{**}, \Omega_s^{**})) < \varepsilon_1$, we can flow Θ along the flow of X for a uniform short time τ to obtain a family Θ_t such that

- (i) $A^h(\Theta_t) \leq A^h(\Theta)$ for all $t \in [0, \tau]$,
- (ii) $2a|\text{Vol}(\Theta) - \text{Vol}(\Theta_t)| \leq A^h(\Theta) - A^h(\Theta_t)$ for all $t \in [0, \tau]$,
- (iii) $A^h(\Theta_\tau) \leq A^h(\Theta) - \varepsilon_3$.

Here $\varepsilon_3 > 0$ is a uniform constant that does not depend on Θ . Discretizing this family shows that Θ is not $(F, \varepsilon_3, \delta, 0, \mathcal{F})$ -almost-minimizing for any $\delta > 0$. Since this is true for all Θ which are \mathcal{F} -close to $(V_s^{**}, \Omega_s^{**})$, this contradicts that $(V_s^{**}, \Omega_s^{**})$ is (F, \mathcal{F}) -almost-minimizing. \square

We can now prove the regularity.

Proposition 3.24. *Assume that (V, Ω) is stationary for F . Further suppose that (V, Ω) is F -almost-minimizing in annuli. Then V is induced by a smooth, closed, almost-embedded hypersurface Σ with multiplicity one. The touching set of Σ is contained in a countable union of $(n-1)$ -dimensional submanifolds. Moreover, Σ coincides with the boundary of Ω . Finally, the mean curvature of Σ satisfies $H = h - f'(\text{Vol}(\Omega))$.*

Proof. Once we know that successive replacements on overlapping annuli must have the same mean curvature, then we can proceed exactly as in [38] to deduce local regularity: for each $p \in \text{supp } \|V\|$ there is an $r > 0$ such that V is induced by a smooth, almost-embedded hypersurface $\Sigma(p, r)$ with multiplicity one in $B_r(p)$. Moreover, the touching set of $\Sigma(p, r)$ is contained in a countable union of $(n-1)$ -dimensional manifolds. Finally, $\Sigma(p, r)$ has mean curvature $h + h_0(p, r)$ for some constant $h_0(p, r)$ that a priori depends on p and r .

The local regularity implies that V is induced by finitely many smooth, almost-embedded, multiplicity one components $(\Gamma_i)_{i=1}^k$ with mean curvature $H_i = h + h_i$, where h_i is a constant that depends a priori on the component. Since $\|\partial\Omega\| \leq \|V\|$, the constancy theorem implies that

$$|\partial\Omega| = \sum_{i=1}^k a_i |\Gamma_i|,$$

where each a_i is either 0 or 1. We claim that in fact $a_i = 1$ for all i . Indeed, this follows from the fact that (V, Ω) is stationary for F . If some a_i was equal to 0, then since the mean curvature of Γ_i is not identically 0, we could construct a local deformation near Γ_i decreasing the area to first order, and leaving the region Ω unchanged. But such a deformation would also decrease F to first order. Thus a_i must equal 1, as claimed.

Finally, the fact that (V, Ω) is stationary for F implies that the mean curvature of Γ_i with respect to the normal vector pointing into Ω must equal $h - f'(\text{Vol}(\Omega))$ for every component Γ_i . Otherwise it would again be possible to construct a deformation decreasing the F functional to first order. This completes the proof. \square

Theorem 1.10 now follows by combining Proposition 3.1, Proposition 3.15, and Proposition 3.24.

4. ABSOLUTE MIN-MAX FOR E

The goal of this section is to develop a min-max theory for the E functional. Throughout this section, we fix a closed manifold M , and a smooth function $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ satisfying (5), and we abbreviate $E = E^f$.

4.1. Pull Tight. We will employ a pull-tight operation to ensure that every point in the critical set is stationary for E .

Proposition 4.1. *Let Π be the homotopy class of an \mathbf{F} continuous map $\Phi_0 : X \rightarrow \mathcal{B}(M, \mathbb{Z}_2)$. Let $\{\Phi_i\}$ be a critical sequence for Π . There is another critical sequence $\{\Xi_i\}$ such that $\mathcal{K}(\{\Xi_i\}) \subset \mathcal{K}(\{\Phi_i\})$ and every $(V, T) \in \mathcal{K}(\{\Xi_i\})$ is stationary for E .*

Proof. The proof is almost identical to Proposition 3.1, except that one replaces the VC space by the VZ space and lets $Z = \emptyset$. \square

4.2. Replacements and the Almost Minimizing Property. In this section, we define a suitable almost-minimizing property and then use it to construct replacements. Most of the constructions are very similar to those for the F functional, although at some points we need to exercise care to ensure that nothing depends on whether we consider $T = \partial\Omega$ or $T = \partial(M \setminus \Omega)$. We will omit proofs that are essentially identical to those for F -almost minimizers.

As in the previous section, the key point is that the functional E still satisfies a quasi-locality type property if we restrict to modifications that change area faster than volume. As before, let $a = \sup |f'|$.

Proposition 4.2. *Assume U_1, U_2, \dots, U_N are disjoint open sets in M . Assume that $T \in \mathcal{B}(M, \mathbb{Z}_2)$ and choose $\Omega \in \mathcal{C}(M)$ with $\partial\Omega = T$. Let $\delta > 0$ be given. Suppose there are $S_k = \partial\Theta_k \in \mathcal{B}(M, \mathbb{Z}_2)$ with $\text{supp}(\Omega - \Theta_k) \subset U_k$ and*

$$2a|\text{Vol}(\Omega) - \text{Vol}(\Theta_i)| \leq \mathbf{M}(\partial\Omega) - \mathbf{M}(\partial\Theta_i) + \delta$$

for $k = 1, 2, \dots, N$. Let

$$T^* = T \llcorner (M \setminus (U_1 \cup \dots \cup U_k)) + \sum_{k=1}^N S_k \llcorner U_k.$$

Then there is an estimate

$$E(T^*) \leq E(T) + \frac{1}{2} \sum_{i=1}^N (\mathbf{M}(S_k) - \mathbf{M}(T)) + \frac{N\delta}{2}.$$

4.2.1. The Almost-Minimizing Property. The above quasi-locality property is the motivation for the following definition of E almost-minimizing cycles.

Definition 4.3. Let U be an open subset of M . Let ν denote either the \mathcal{F} , \mathbf{F} , or \mathbf{M} norm. Fix constants $\varepsilon, \delta > 0$ and $\gamma \geq 0$. Fix an element $T \in \mathcal{B}(M, \mathbb{Z}_2)$. Choose $\Omega \in \mathcal{C}(M)$ with $\partial\Omega = T$. Let $(\Omega_i)_{i=1, \dots, k}$ be a sequence in $\mathcal{C}(M)$. Assume that

- (i) $\Omega_1 = \Omega$,
- (ii) $\text{supp}(\Omega - \Omega_i) \subset U$ for all i ,
- (iii) $2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_i)| \leq \mathbf{M}(\partial\Omega) - \mathbf{M}(\partial\Omega_i) + \gamma$ for all i ,
- (iv) $\nu(\partial\Omega_i, \partial\Omega_{i+1}) < \delta$ for all i ,
- (v) $\mathbf{M}(\partial\Omega_i) \leq \mathbf{M}(\partial\Omega) + \delta$ for all i .

We say T is $(E, \varepsilon, \delta, \gamma, \nu)$ -almost-minimizing in U if for all sequences $(\Omega_i)_{i=1, \dots, k}$ as above, we have $\mathbf{M}(\partial\Omega_k) \geq \mathbf{M}(T) - \varepsilon$.

Remark 4.4. The above definition does not depend on the choice of Ω . Indeed if (Ω_i) is a sequence satisfying (i)-(v) with respect to Ω then $M - \Omega_i$ is a sequence satisfying (i)-(v) with respect to $M - \Omega$.

Definition 4.5. We define E almost-minimizers. Let $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ and let U be an open subset of M .

- (a) We say (V, T) is (E, \mathcal{F}) almost-minimizing in U if there is a sequence $T_i \in \mathcal{B}(M, \mathbb{Z}_2)$ such that T_i is $(E, \varepsilon_i, \delta_i, 0, \mathcal{F})$ -almost-minimizing in U and $|T_i| \rightarrow V$ and $T_i \rightarrow T$ and $\varepsilon_i, \delta_i \rightarrow 0$.
- (b) We say (V, T) is (E, \mathbf{M}) almost-minimizing in U if there is a sequence $T_i \in \mathcal{B}(M, \mathbb{Z}_2)$ such that T_i is $(E, \varepsilon_i, \delta_i, \delta_i, \mathbf{M})$ almost-minimizing in U and $|T_i| \rightarrow V$ and $T_i \rightarrow T$ and $\varepsilon_i, \delta_i \rightarrow 0$.

Proposition 4.6. Assume that T is $(E, \varepsilon, \rho, \rho, \mathbf{M})$ almost-minimizing in U . If $\delta \leq \rho$ is small enough, then T is $(E, \varepsilon, \delta, 0, \mathcal{F})$ almost-minimizing in any open set W compactly contained in U .

Proposition 4.7. If $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ is (E, \mathbf{M}) almost-minimizing in U then (V, T) is (E, \mathcal{F}) almost-minimizing in any open set W compactly contained in U .

Proposition 4.8. Assume that $(V, T) \in \text{VZ}(M)$ is (E, \mathcal{F}) almost-minimizing in U . Then V has $2a$ -bounded first variation in U .

Definition 4.9. An element $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ is called E -almost-minimizing in annuli if for each $x \in M$ there is a number $\rho(x) > 0$ such that (V, T) is (E, \mathcal{F}) almost-minimizing in every annulus $\text{An}(x, s, r)$ with $s < r < \rho(x)$.

Proposition 4.10. Let Π be the X -homotopy class of a map $\Phi_0: X \rightarrow (B(M, \mathbb{Z}_2), \mathbf{F})$, where X is a cubical subcomplex of $I(m, k)$ for some $m, k \in \mathbb{N}$. Choose a pulled-tight critical sequence $\{\Phi_i\}$ and let $\mathcal{K} = \mathcal{K}(\{\Phi_i\})$ be the critical set. Then there is an element $(V, T) \in \mathcal{K}$ which is (E, \mathcal{F}) almost minimizing in annuli. In fact, there exists some $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ satisfying the following stronger property:

- (R) There is a number $N = N(m)$ depending only on m such that for any collection of N concentric annuli $\text{An}(x, s_1, r_1), \dots, \text{An}(x, s_N, r_N)$ with $2r_j < s_{j+1}$, (V, T) is (E, \mathcal{F}) almost minimizing in at least one of the annuli.

Proof. Applying discretization [35, Theorem 1.11], we can find a homotopy sequence $\{\phi_i\}$ such that $L(\{\phi_i\}) = L(\Pi)$ and $\mathcal{K}(\{\phi_i\}) = \mathcal{K}(\{\Phi_i\})$. Assume for contradiction that no element in \mathcal{K} satisfies property (R). Then by Proposition 4.6, no element in \mathcal{K} satisfies property (R) with (E, \mathcal{F}) almost-minimizing replaced by (E, \mathbf{M}) almost-minimizing. Thus we can apply Parts 1-19 of the Almgren-Pitts combinatorial construction [24, Theorem 4.10] to $\{\phi_i\}$. This produces a homotopic sequence $\{\psi_i\}$.

Let us give more details. In Part 9, given σ and a vertex of σ , one defines $T(j, 1)$ and $T(j, 2)$ in the same way as Pitts. Note that

$$\mathbf{M}(T(j, 1) - T(j, 2)) < \delta_i.$$

Choose Ω_1 so that $\partial\Omega_1 = T(j, 1)$. Then by the isoperimetric theorem, there is a *unique* choice of Ω_2 so that $\partial\Omega_2 = T(j, 2)$ and $|\text{Vol}(\Omega_1) - \text{Vol}(\Omega_2)| < \frac{\delta_i}{2a}$. Moreover, $\text{supp}(\Omega_2 - \Omega_1) \subset A_k$. Then, using the fact that $T(j, 2)$ is not $(E, \varepsilon, \delta_i, \delta_i, \mathcal{M})$ almost minimizing in an annulus $a_k \subset A_k$, we obtain a sequence $(\Omega_q)_{q=2}^{3^{N_1}}$ such that

- (i) $\text{supp}(\Omega_2 - \Omega_q) \subset a_k$ for all q ,
- (ii) $2a|\text{Vol}(\Omega_2) - \text{Vol}(\Omega_q)| \leq \mathbf{M}(\partial\Omega_2) - \mathbf{M}(\partial\Omega_q) + \delta_i$ for all i ,
- (iii) $\mathbf{M}(\partial\Omega_2, \partial\Omega_q) < \delta_i$ for all q ,
- (iv) $\mathbf{M}(\partial\Omega_q) \leq \mathbf{M}(\partial\Omega_2) + \delta_i$ for all q ,

$$(v) \mathbf{M}(\partial\Omega_{3^{N_1}}) \leq \mathbf{M}(\partial\Omega_2) - \varepsilon.$$

Note then that

- (i) $\text{supp}(\Omega_1 - \Omega_q) \subset A_k$ for all q ,
- (ii) $2a|\text{Vol}(\Omega_1) - \text{Vol}(\Omega_q)| \leq \mathbf{M}(\partial\Omega_1) - \mathbf{M}(\partial\Omega_q) + 3\delta_i$ for all i ,
- (iii) $\mathbf{M}(\partial\Omega_1, \partial\Omega_q) < 3\delta_i$ for all q ,
- (iv) $\mathbf{M}(\partial\Omega_q) \leq \mathbf{M}(\partial\Omega_1) + 3\delta_i$ for all q ,
- (v) $\mathbf{M}(\partial\Omega_{3^{N_1}}) \leq \mathbf{M}(\partial\Omega_1) - \varepsilon/2$.

We set $T(j, q) = \partial\Omega_q$. Again, we emphasize that all properties in the above list continue to hold if we replace Ω_q by $M \setminus \Omega_q$ for $q = 1, \dots, 3^{N_1}$. One then continues through Part 19 as in [24].

Now in Part 20, one can use Proposition 4.2 to verify that

$$L(\{\psi_i\}) < L(\{\phi_i\}) - \varepsilon_1$$

for some $\varepsilon_1 > 0$. Again let us be more precise. Consider disjoint annuli A_1, A_2, \dots, A_I . Fix $T \in \mathcal{B}(M, \mathbb{Z}_2)$. For each annulus, consider a sequence $T_k(q) = T_k(j, q)$ as above. For a choice of integers $q_1, \dots, q_N \in \{1, \dots, 3^{N_1}\}$, one then considers a cycle of the form

$$T^* = T \llcorner (M \setminus (A_1 \cup \dots \cup A_I)) + \sum_{k=1}^I T_k(q_k) \llcorner A_k$$

and needs to estimate the difference $E(T) - E(T^*)$. Choose Ω so that $T = \partial\Omega$. Then, for each k and q , let $\Omega_k(q)$ be the sets that were selected above which satisfy $\partial\Omega_k(q) = T_k(q)$ and were used to define $T_k(q)$. If $\Omega_k(1) = \Omega$, then set $\Theta_k(q) = \Omega_k(q)$ for all q . Otherwise, if $\Omega_k(1) = M \setminus \Omega$, then set $\Theta_k(q) = M \setminus \Omega_k(q)$ for all q . Then we can apply Proposition 4.2 with $S_k = T_k(q_k)$ and $\Theta_k = \Omega_k(q_k)$ to estimate $E(T) - E(T^*)$, as needed.

Applying interpolation [35, Theorem 1.12] to $\{\psi_i\}$ then gives a sequence

$$\{\Psi_i: X \rightarrow (\mathcal{C}(M), \mathbf{F})\}$$

in the (X, Z) -homotopy class Π with $\sup_{x \in X} F(\Psi_i(x)) < L(\Pi) - \varepsilon_2$ for all large i . This is a contradiction. \square

4.2.2. A Constrained Minimization Problem. Consider the following constrained minimization problem. Assume that T is $(E, \varepsilon, \delta, 0, \mathcal{F})$ almost minimizing in U . Let K be a compact subset of U . Choose a set $\Omega \in \mathcal{C}(M)$ such that $\partial\Omega = T$. Let $\mathcal{A} = \mathcal{A}(T, K, \delta)$ be the set of all $S \in \mathcal{B}(M, \mathbb{Z}_2)$ such that there is a sequence $(\Omega_i)_{i=1, \dots, k}$ in $\mathcal{C}(M)$ satisfying

- (i') $\Omega_1 = \Omega$,
- (ii') $\text{supp}(\Omega - \Omega_i) \subset K$ for all i ,
- (iii') $2a|\text{Vol}(\Omega) - \text{Vol}(\Omega_i)| \leq \mathbf{M}(\partial\Omega) - \mathbf{M}(\partial\Omega_i)$ for all i ,
- (iv') $\mathcal{F}(\partial\Omega_i, \partial\Omega_{i+1}) < \delta$ for all i ,
- (v') $\mathbf{M}(\partial\Omega_i) \leq \mathbf{M}(\partial\Omega) + \delta$ for all i ,

and $S = \partial\Omega_k$. We aim to minimize area among elements of \mathcal{A} .

Proposition 4.11. *There exists $S \in \mathcal{A}$ such that $\mathbf{M}(S) = \inf_{R \in \mathcal{A}} \mathbf{M}(R)$.*

Proposition 4.12. *Assume $T = \partial\Omega$ is $(E, \varepsilon, \delta, 0, \mathcal{F})$ almost-minimizing in U and let K be a compact subset of U . Let S be an area minimizer in \mathcal{A} . Then S is $(E, \varepsilon, \delta, 0, \mathcal{F})$ almost-minimizing in U .*

Proposition 4.13. *Let S be an area minimizer in \mathcal{A} . Then S locally minimizes area with respect to volume preserving modifications in the interior of K . Moreover, S is smooth with*

constant mean curvature $|H| \leq 2a$ in K . The surface S is volume preserving stable for the area functional in the interior of K .

4.2.3. *Replacements.* Next we turn to the construction of replacements.

Proposition 4.14. *Assume that V has $2a$ -bounded first variation, and also that (V, T) is (E, \mathcal{F}) almost-minimizing in an open set U . Let K be a compact subset of U . There exists an element $(V^*, T^*) \in \text{VZ}(M)$ called a replacement for (V, T) such that*

- (i) $V \llcorner (M - K) = V^* \llcorner (M - K)$,
- (ii) $\|V\|(M) = \|V^*\|(M)$,
- (iii) (V^*, T^*) is (E, \mathcal{F}) almost-minimizing in U ,
- (iv) V^* has $2a$ -bounded first variation,
- (v) (V^*, T^*) , when restricted to the interior of K , is a limit of some solutions to the constrained minimization problems.

Proposition 4.15. *Let (V^*, T^*) be a replacement for (V, T) in K . Then in the interior of K , either*

- (i) *the varifold V^* is induced by a smooth, almost-embedded CMC hypersurface Σ with $|H| \leq 2a$ and multiplicity one; or*
- (ii) *the support of V^* is a smooth, embedded (not necessarily connected) minimal hypersurface Σ and V^* is induced by the connected components of Σ equipped with some integer multiplicities.*

Moreover, if $\Omega^* \in \mathcal{C}(M)$ satisfies $\partial\Omega^* = T^*$, then $\partial\Omega^*$ coincides with the odd multiplicity components of V^* in the interior of K . Finally, in the interior of K , there is a curvature estimate

$$\|A_\Sigma\|^2(x) \leq \frac{C}{\text{dist}(x, \partial K)^2}$$

where C is a constant that depends only on M , a , and an upper bound for $\|V\|(M)$.

Proof. In the interior of K , the replacement V^* is a limit of smooth, embedded, volume preserving stable CMC hypersurfaces with uniformly bounded mean curvature. Hence the regularity follows from the curvature estimates for volume preserving stable CMCs. \square

We conclude by investigating the tangent cones to E almost-minimizers. We show that if (V, T) is (E, \mathcal{F}) almost-minimizing in annuli, then every tangent cone to V is an integer multiple of a plane.

Proposition 4.16. *Assume that (V, T) has $2a$ -bounded first variation and is (E, \mathcal{F}) -almost minimizing in a set U . Then for any sequence $p_i \rightarrow p \in U$ and any sequence of scales $r_i \rightarrow 0$, every varifold limit $\overline{V} = (\eta_{p_i, r_i})_\# V$ is an integer multiple of a complete, embedded minimal hypersurface. Moreover, every varifold tangent to V is an integer multiple of a hyperplane.*

4.3. Regularity. The goal of this subsection is to prove the regularity of the min-max pair (V, T) . As in the case of the F functional, the key point is to ensure that successive replacements on overlapping annuli have the same mean curvature.

Given a smooth almost-embedded surface Σ with constant mean curvature, recall that $\mathcal{R}(\Sigma)$ and $\mathcal{S}(\Sigma)$ denote respectively the set of embedded points and the touching set of Σ .

The next proposition shows that, once a replacement has non-zero mean curvature, all subsequent replacements must also have matching non-zero mean curvature. It does not seem straightforward to show that if the initial replacement is minimal with multiplicity, then subsequent replacements will also be minimal with multiplicity. Fortunately, the following is already enough to prove regularity.

Proposition 4.17. *Assume (V, Ω) is E almost-minimizing in annuli and fix a point $p \in \text{supp } \|V\|$. Fix sufficiently small numbers $s_1 < r_1$ and let (V^*, Ω^*) be a replacement for (V, Ω) in $\text{An}(p, s_1, r_1)$. Let Σ^* be the smooth, almost-embedded hypersurface inducing V^* in $\text{An}(p, s_1, r_1)$ and assume that Σ^* has non-zero mean curvature h_0 . Choose $s_1 < r_2 < r_1$ so that $\partial B_{r_2}(p)$ intersects Σ^* and the countable union of manifolds containing $\mathcal{S}(\Sigma^*)$ transversally and so that $\partial B_{r_2}(p) \cap \mathcal{R}(\Sigma^*)$ is non-empty. Fix any $s < s_1$ and let $(V_s^{**}, \Omega_s^{**})$ be a replacement for (V^*, Ω^*) in $\text{An}(p, s, r_2)$. Let Σ_s^{**} be the smooth hypersurface inducing V_s^{**} in $\text{An}(p, s, r_2)$. Then Σ_s^{**} also has mean curvature h_0 .*

Proof. There are several other possibilities to rule out. First, suppose for contradiction that Σ_s^{**} is a multiplicity one CMC with constant mean curvature $h_1 \neq h_0$. Then we can argue as in Proposition 3.23 to construct a volume preserving deformation which decreases the area. This contradicts the E -almost-minimizing property.

Second, suppose for contradiction that Σ_s^{**} is minimal. If the multiplicity of Σ_s^{**} is one, then Σ_s^{**} coincides with the boundary of Ω_s^{**} and we can again construct a volume preserving deformation which decreases the area. As before, this violates the almost-minimizing property. Finally suppose that Σ_s^{**} has higher multiplicity. Then the multiplicity function θ for V_s^{**} satisfies $\theta(x) \geq 2$ for all $x \in \Sigma_s^{**}$. By assumption, there is a point $y \in \partial B_{r_2} \cap \mathcal{R}(\Sigma^*)$. We know any varifold tangent to V_s^{**} at y is an integer multiple of a plane. By considering Σ^* , we see that in fact the only possible varifold tangent is $T_y \Sigma^*$ with multiplicity one. Thus, by transversality, there must be a sequence of points $x_i \in \Sigma_s^{**}$ with $x_i \rightarrow y$. But, by upper semi-continuity of the density, this implies that $\theta_{V_s^{**}}(y) \geq 2$, contradicting that the varifold tangent $T_y \Sigma^*$ has multiplicity one. Therefore this also cannot occur. \square

We can now complete the regularity argument.

Proposition 4.18. *Assume that (V, T) satisfies $\delta E(V, T) = 0$. Further suppose that (V, T) is (E, \mathcal{F}) almost-minimizing in annuli. Choose a set Ω with $\partial\Omega = T$ and let $H = -f'(\text{Vol}(\Omega))$.*

- (i) *If $H \neq 0$, then there exists a smooth, almost-embedded (not necessarily connected) CMC hypersurface $\Lambda = \partial\Omega$, which has mean curvature H with respect to the normal pointing into Ω . Moreover, there exist a (possibly empty) collection of minimal hypersurfaces $\Sigma_1, \dots, \Sigma_k$ and a collection of multiplicities $m_1, \dots, m_k \in \mathbb{N}$ such that*

$$V = |\Lambda| + \sum_{i=1}^k m_i |\Sigma_i|.$$

The hypersurfaces $\Lambda, \Sigma_1, \dots, \Sigma_k$ are all disjoint.

- (ii) *If $H = 0$, then there exists a collection of smooth, embedded minimal hypersurfaces $\Lambda_1, \dots, \Lambda_q$ such that $\partial\Omega = \Lambda_1 \cup \dots \cup \Lambda_q$. Moreover, there exist a (possibly empty) collection of minimal hypersurfaces $\Sigma_1, \dots, \Sigma_k$ and a collection of multiplicities $\ell_1, \dots, \ell_q, m_1, \dots, m_k \in \mathbb{N}$ such that*

$$V = \sum_{i=1}^q \ell_i |\Lambda_i| + \sum_{i=1}^k m_i |\Sigma_i|.$$

The hypersurfaces $\Lambda_1, \dots, \Lambda_q, \Sigma_1, \dots, \Sigma_k$ are all disjoint.

Proof. The argument is slightly more complicated than for the F functional, because Proposition 4.17 is weaker than Proposition 3.23. To start, choose a point $p \in \text{supp } \|V\|$. Choose very small numbers $s_1 < r_1$ and let (V^*, Ω^*) be a replacement for (V, Ω) in $\text{An}(p, s_1, r_1)$. There are now several cases to consider.

First, suppose that (V^*, Ω^*) is induced by a smooth, almost-embedded CMC hypersurface Σ^* with non-zero mean curvature in $\text{An}(p, s_1, r_1)$. Then choose $s_1 < r_2 < r_1$ so that $\partial B_{r_2}(p)$ is transverse to Σ^* and the union of manifolds containing $\mathcal{S}(\Sigma^*)$ and so that $\partial B_{r_2}(p) \cap \mathcal{R}(\Sigma^*)$ is non-empty. Then for $s < s_1$ let $(V_s^{**}, \Omega_s^{**})$ be a replacement for (V^*, Ω^*) in $\text{An}(p, s, r_2)$. According to Proposition 4.17, V_s^{**} must be induced by a multiplicity one hypersurface Σ_s^{**} whose mean curvature matches that of Σ^* . One can now proceed exactly as in [37] to deduce that V is induced by a smooth, almost-embedded CMC hypersurface with multiplicity one in a small neighborhood of p .

Second, suppose that (V^*, Ω^*) is induced by a smooth, embedded, minimal hypersurface Σ^* , possibly with multiplicity. Choose $s_1 < r_2 < r_1$ so that $\partial B_{r_2}(p)$ intersects Σ^* transversally. Then for $s < s_1$ let $(V_s^{**}, \Omega_s^{**})$ be a replacement for (V^*, Ω^*) in $\text{An}(p, s, r_2)$. If for every $s < s_1$, V_s^{**} is induced by a smooth minimal surface with multiplicity, then we can argue exactly as in the usual Almgren-Pitts construction to deduce that V is induced by a smooth, embedded minimal hypersurface with multiplicity in a small neighborhood of p .

The remaining possibility is that for some choice of $s_2 < s_1$, the varifold $V_{s_2}^{**}$ is induced by a smooth, almost-embedded CMC hypersurface $\Sigma_{s_2}^{**}$ with non-zero mean curvature. In this case, we will take a *third* replacement. Choose $s_2 < r_3 < r_2$ so that $\partial B_{r_3}(p)$ intersects $\Sigma_{s_2}^{**}$ and the countable union of manifolds containing $\mathcal{S}(\Sigma_{s_2}^{**})$ transversally and so that $\partial B_{r_3}(p) \cap \mathcal{R}(\Sigma_{s_2}^{**})$ is non-empty. Then for any $s < s_2$, let $(V_s^{***}, \Omega_s^{***})$ be a replacement for $(V_{s_2}^{**}, \Omega_{s_2}^{**})$ in $\text{An}(p, s, r_3)$. According to Proposition 4.17 (applied with $(V, \Omega) = (V_{s_2}^{**}, \Omega_{s_2}^{**})$), the varifold V_s^{***} is induced by a smooth, almost-embedded hypersurface Σ_s^{***} in $\text{An}(p, s, r_3)$ with non-zero mean curvature matching that of $\Sigma_{s_2}^{**}$. Arguing exactly as in [37], one can show that $\Sigma_{s_2}^{**}$ and Σ_s^{***} glue smoothly along $\partial B_{r_3}(p)$. One can then let $s \rightarrow 0$ and continue to argue exactly as in [37] to deduce that V is induced by a smooth, almost-embedded CMC hypersurface with multiplicity one in a small neighborhood of p .

We have now proven the following local regularity: for every $p \in \text{supp } \|V\|$ there is an $r > 0$ such that in $B_r(p)$ the varifold V is induced by either

- (i) a smooth, almost-embedded, multiplicity one CMC surface Λ with non-zero mean curvature, or
- (ii) a collection of smooth, embedded minimal hypersurfaces with integer multiplicities.

The local regularity then implies that V is induced by a collection of multiplicity one, almost-embedded CMCs together with a collection of smooth, embedded minimal hypersurfaces with multiplicities.

Next note that since $\| |T| \| \leq \|V\|$, the constancy theorem implies that $|T|$ is induced by some subcollection of the surfaces inducing V . Hence we can list the components of $\text{supp } \|V\|$ as $\Lambda_1, \dots, \Lambda_\ell, \Sigma_1, \dots, \Sigma_k$ so that

$$|T| = \sum_{i=1}^{\ell} |\Lambda_i|,$$

$$V = \sum_{i=1}^{\ell} \ell_i |\Lambda_i| + \sum_{j=1}^k m_j |\Sigma_j|.$$

Here $\ell_i, m_j \in \mathbb{N}$ are some multiplicities.

Now we use the fact that (V, T) is stationary for E to deduce the remaining claims of the proposition. First note that every surface Σ_j must be minimal. Indeed, if some surface Σ_j was not minimal, then we could construct a local deformation near Σ_j decreasing the area and

leaving T unchanged. Such a deformation would decrease E to first order. Therefore each Σ_j must be minimal.

Choose a set Ω with $\partial\Omega = T$. It remains to verify that all the surfaces Λ_i must have mean curvature $-f'(\text{Vol}(\Omega))$, computed with respect to the normal pointing into Ω . If, to the contrary, Λ_i had a different mean curvature, then we could construct a local deformation near Λ_i decreasing the E functional to first order. This completes the proof of the proposition. \square

Theorem 1.9 now follows by combining Proposition 4.1, Proposition 4.10, and Proposition 4.18.

5. COMPACTNESS

In this section, we record some compactness properties satisfied by the surfaces produced by the min-max theorems.

5.1. Compactness for \mathbf{F} . Assume that $f: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ satisfies (5) and that $h: M \rightarrow \mathbb{R}$ is a smooth Morse function satisfying property (T) 2.10. Choose a sequence $\varepsilon_k \rightarrow 0$ and let

$$F_k(\Omega) = \mathbf{M}(\partial\Omega) - \varepsilon_k \int_{\Omega} h + f(\text{Vol}(\Omega)).$$

Also define

$$E(T) = \mathbf{M}(\partial\Omega) + f(\text{Vol}(T)).$$

We would like to prove certain compactness for a sequence (V_k, Ω_k) of critical points for F_k as $k \rightarrow \infty$.

The basic strategy is to try and show that the almost-minimizing property implies volume preserving stability. As noted in Remark 3.13, it seems we cannot adapt the argument in [30, Lemma 3.3] because of the extra volume constraint in the definition of almost-minimizers. Thus we will instead argue that the min-max hypersurfaces coincide with their replacements, which are known to be volume preserving stable.

Proposition 5.1. *Assume that $\Sigma = \partial\Omega$ is a smooth, almost-embedded hypersurface with mean curvature $\varepsilon_k h + h_0$ for some constant h_0 . Assume that $(|\Sigma|, \Omega)$ is (F_k, \mathcal{F}) almost-minimizing in an open set U and that $\text{An}(p, s, r) \subset U$ is a closed annulus. Let (V^*, Ω^*) be a replacement for $(|\Sigma|, \Omega)$ in $\text{An}(p, s, r)$. Let Σ^* be the smooth, almost-embedded hypersurface inducing V in $\text{An}(p, s, r)$. Then $\Sigma \cap (M \setminus \text{An}(p, s, r))$ and Σ^* have the same mean curvature, and glue smoothly along the boundary of $\text{An}(p, s, r)$.*

Proof. Consider any point $q \in \partial\text{An}(p, s, r)$. Note that (V^*, Ω^*) is (F_k, \mathcal{F}) -almost-minimizing in small annuli centered at q . Therefore, by the same local regularity argument as in Proposition 3.24, it follows that V^* is induced by a smooth, almost-embedded hypersurface in a neighborhood of q . Therefore, $\text{supp } \|V^*\|$ is smooth in a neighborhood of $\partial\text{An}(p, s, r)$. This implies the conclusion of the proposition. \square

Remark 5.2. An almost-embedded hypersurface in M can be viewed as an immersion $\Sigma \rightarrow M$. In the following, by a component of an almost-embedded hypersurface, we mean the image of a connected component of Σ under this immersion. Thus two spheres touching tangentially at a point define two components.

Corollary 5.3. *Assume that $\Sigma = \partial\Omega$ is a smooth, almost-embedded hypersurface with mean curvature $\varepsilon_k h + h_0$ for some constant h_0 . Assume that $(|\Sigma|, \Omega)$ is (F_k, \mathcal{F}) almost-minimizing in an open set U and that $\text{An}(p, s, r) \subset U$ is a closed annulus. Assume that a local embedded sheet*

Γ of Σ crosses the boundary of $An(p, s, r)$. Then the component of Σ containing this sheet is volume preserving stable for $A^{\varepsilon_k h + h_0}$ in the interior of $An(p, s, r)$.

Proof. Consider a replacement (V^*, Ω^*) for $(|\Sigma|, \Omega)$ in $An(p, s, r)$. Then V^* is induced by a smooth, almost-embedded hypersurface Σ^* in the interior of $An(p, s, r)$. According to Proposition 5.1, $\Sigma \cap (M \setminus An(p, s, r))$ glues smoothly to Σ^* along the boundary of $An(p, s, r)$. Now let Σ_1 be the component of Σ containing Γ . Then by unique continuation, $\Sigma_1 \cap \text{int}(An(p, s, r))$ is contained in Σ^* and the conclusion of the proposition follows since replacements are volume preserving stable. \square

Corollary 5.4. *Assume that $\Sigma = \partial\Omega$ is a smooth, almost-embedded hypersurface with mean curvature $\varepsilon_k h + h_0$ for some constant h_0 . Assume that $(|\Sigma|, \Omega)$ is (F_k, \mathcal{F}) almost-minimizing in an open set U . There is a radius $\rho > 0$ depending only on M and an upper bound c for $|\varepsilon_k h + h_0|$ such that if $s < r < \rho$ and $An(p, s, r) \subset U$, then Σ is volume preserving stable for $A^{\varepsilon_k h + h_0}$ in the interior of $An(p, s, r)$.*

Proof. Let (V^*, Ω^*) be a replacement for $(|\Sigma|, \Omega)$ in $An(p, s, r)$. Then V^* is induced by a smooth, almost-embedded hypersurface Σ^* in the interior of $An(p, s, r)$. Again Proposition 5.1 implies that Σ^* glues smoothly to $\Sigma \cap (M \setminus An(p, s, r))$ along the boundary of $An(p, s, r)$. By the maximum principle, if ρ is sufficiently small depending on M and an upper bound for the mean curvature of Σ , then Σ and Σ^* can have no closed components contained entirely in $An(p, s, r)$. Therefore, by unique continuation, $\Sigma = \Sigma^*$ in $An(p, s, r)$ and the result follows. \square

Proposition 5.5. *Consider a sequence $\{(V_k, \Omega_k)\} \in \text{VC}(M)$. Assume that (V_k, Ω_k) is stationary for F_k and that (V_k, Ω_k) satisfies property (R) (3.15) for F_k with an integer m that does not depend on k . Finally, suppose that $\|V_k\|(M)$ is bounded uniformly above, and that the varifolds V_k have c -bounded first variation for a uniform constant $c > 0$. Then, up to a subsequence, $(V_k, \partial\Omega_k) \rightarrow (V, T) \in \text{VZ}(M, \mathbb{Z}_2)$. Moreover, (V, T) is stationary for E and we have the following alternative. Either:*

- (i) *The varifold V is induced by a smooth, almost-embedded, multiplicity one constant mean curvature surface Λ with non-zero mean curvature; or*
- (ii) *The varifold V is induced by a collection of minimal hypersurfaces with multiplicities.*

The following property also holds:

- (R') *There is a number $N = N(m)$ depending only on m and a number $\rho > 0$ depending only on c , such that for any collection of N concentric annuli $An(x, s_1, r_1), \dots, An(x, s_N, r_N)$ with $2r_j < s_{j+1}$ and $r_N < \rho$, $\text{supp } \|V\|$ is volume preserving stable for the area as an immersion in at least one of the annuli.*

Proof. Choose a sequence $\{(V_k, \Omega_k)\}$ as in the statement of the theorem and let $T_k = \partial\Omega_k$. Then $(V_k, T_k) \in \text{VZ}(M, \mathbb{Z}_2)$ and so, up to a subsequence, $(V_k, T_k) \rightarrow (V, T)$ in $\text{VZ}(M, \mathbb{Z}_2)$ since $\|V_k\|(M)$ is bounded uniformly from above. It follows immediately from the first variation formulas (6)(7) that (V, T) is stationary for E .

Note that Proposition 3.24 implies that V_k is induced by a smooth, almost-embedded hypersurface $\Sigma_k = \partial\Omega_k$ with mean curvature $H_k = \varepsilon_k h + h_k$, where h_k is a constant. Moreover, the constants h_k are uniformly bounded. Passing to a further subsequence, we can suppose that $h_k \rightarrow h_\infty$.

To prove property (R'), let ρ be the constant from Corollary 5.4, which depends only on c . Consider a collection of N concentric annuli $An(x, s_1, r_1), \dots, An(x, s_N, r_N)$ with $2r_j < s_{j+1}$ and $r_N < \rho$. Since (V_k, Ω_k) satisfies property (R), there is a choice of $j \in \{1, \dots, N\}$ and a subsequence (V_{k_i}, Ω_{k_i}) such that (V_{k_i}, Ω_{k_i}) is (F_{k_i}, \mathcal{F}) -almost minimizing in $An = An(x, s_j, r_j)$

for every i . According to Proposition 5.4, Σ_{k_i} is volume preserving stable for $A^{\varepsilon_k h + h_{k_i}}$ as an immersion in An .

This stability implies a curvature estimate for Σ_{k_i} in An . If $h_\infty \neq 0$, then passing to a further subsequence, Σ_{k_i} converges smoothly with multiplicity one to an almost-embedded limit Σ with mean curvature h_∞ in An . If $h_\infty = 0$, then passing to a further subsequence, Σ_{k_i} converges smoothly, possibly with multiplicity, to a collection of minimal hypersurfaces in An . In both cases, this implies that $\text{An} \cap \text{supp } \|V\|$ is smooth and volume preserving stable for the area as an immersion. This proves that property (R') holds.

Finally, we need to show that either (i) or (ii) holds. First suppose that $h_\infty \neq 0$. Then the above argument shows that for any collection of N concentric annuli $\text{An}(x, s_1, r_1), \dots, \text{An}(x, s_N, r_N)$ with $2r_j < s_{j+1}$ and $r_N < \rho$, V is induced by a smooth, almost-embedded, volume preserving stable CMC with mean curvature h_∞ in one of the annuli. This implies that there is a radius $\sigma(x) > 0$ such that for any annulus $\text{An}(x, s, r)$ with $s < r < \sigma(x)$, the varifold V is induced by a smooth, almost-embedded, volume preserving stable CMC with mean curvature h_∞ in $\text{An}(x, s, r)$. This property implies that alternative (i) holds by the removable singularity part in [37, Section 6, Step 4].

If instead $h_\infty = 0$, then the above argument shows that for any collection of N concentric annuli $\text{An}(x, s_1, r_1), \dots, \text{An}(x, s_N, r_N)$ with $2r_j < s_{j+1}$ and $r_N < \rho$, V is induced by a collection of smooth, volume preserving stable minimal surfaces with multiplicity in one of the annuli. This implies that there is a radius $\sigma(x) > 0$ such that for any annulus $\text{An}(x, s, r)$ with $s < r < \sigma(x)$, the varifold V is induced by a collection of smooth, volume preserving stable minimal surfaces with multiplicity. This property implies that alternative (ii) holds by the removable singularity part of the Almgren-Pitts regularity argument [24]. \square

5.2. Compactness for E. Next we record another compactness property for the E functional. Consider a sequence of smooth functions $f_k: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ satisfying (5) and set

$$E_k(T) = \mathbf{M}(T) + f_k(\text{Vol}(T)).$$

Let $\{(V_k, T_k)\}$ be a sequence such that (V_k, T_k) is a critical point for E_k for each $k \in \mathbb{N}$. We show that if the mean curvature of the CMC portion of (V_k, T_k) stays bounded then there is a convergence subsequence. We will omit the proofs in this subsection, since they are very similar to the previous subsection.

Proposition 5.6. *Assume that (V, T) has the regularity described in Proposition 4.18. Assume that (V, T) is (E_k, \mathcal{F}) almost-minimizing in an open set U and that $\text{An}(p, s, r) \subset U$ is a closed annulus. Let (V^*, T^*) be a replacement for (V, T) in $\text{An}(p, s, r)$. Then $\text{supp } \|V^*\| \cap \text{An}(p, s, r)$ and $\text{supp } \|V\| \cap (M \setminus \text{An}(p, s, r))$ glue smoothly along the boundary of $\text{An}(p, s, r)$.*

Corollary 5.7. *Assume that (V, T) has the regularity described in Proposition 4.18. Assume that (V, T) is (E_k, \mathcal{F}) almost-minimizing in an open set U and that $\text{An}(p, s, r) \subset U$ is a closed annulus. Assume that a local embedded sheet of $\text{supp } \|V\|$ crosses the boundary of $\text{An}(p, s, r)$. Then the component of $\text{supp } \|V\|$ containing this sheet is volume preserving stable for the area in the interior of $\text{An}(p, s, r)$.*

Corollary 5.8. *Assume that (V, T) has the regularity described in Proposition 4.18. Assume that (V, T) is (E_k, \mathcal{F}) almost-minimizing in an open set U . There is a radius $\rho > 0$ depending only on M and an upper bound c for the first variation of V such that if $s < r < \rho$ and $\text{An}(p, s, r) \subset U$, then every component of $\text{supp } \|V\|$ intersecting $\text{An}(p, s, r)$ has the same mean curvature. Moreover, $\text{supp } \|V\|$ is volume preserving stable for the area in $\text{An}(p, s, r)$.*

Proposition 5.9. *Consider a sequence $(V_k, T_k) \in \text{VZ}(M, \mathbb{Z}_2)$. Assume that (V_k, T_k) is stationary for E_k and that (V_k, T_k) satisfies property (R) (4.10) for E_k with an integer m that does not depend on k . Suppose that $\|V_k\|(M)$ is uniformly bounded, and that all the varifolds V_k have c -bounded first variation for a uniform constant c . Then, up to a subsequence, $(V_k, T_k) \rightarrow (V, T) \in \text{VZ}(M, \mathbb{Z}_2)$. Moreover, (V, T) has the regularity described in Proposition 4.18 and the following property holds:*

(R') *There is a number $N = N(m)$ depending only on m and a number $\rho > 0$ depending only on c , such that for any collection of N concentric annuli $An(x, s_1, r_1), \dots, An(x, s_N, r_N)$ with $2r_j < s_{j+1}$ and $r_N < \rho$, there is an annulus An in the collection such that every component of $\text{supp } \|V\|$ intersecting An has the same mean curvature and is volume preserving stable for the area in An .*

6. CONSTRUCTING HALF-VOLUME CMCS

The goal of this section is to prove Theorem 1.1 and Corollary 1.2. To begin, we recall the notion of the half-volume spectrum.

6.1. The Half-Volume Spectrum. In [22], the authors introduced the half-volume spectrum of a manifold. This is similar to the usual volume spectrum introduced by Gromov, except that all hypersurfaces in the construction are now additionally required to enclose half the volume of M .

The Almgren isomorphism theorem [1] implies that the space $\mathcal{B}(M, \mathbb{Z}_2)$ is weakly homotopy equivalent to $\mathbb{R}P^\infty$. Marques and Neves [19] later gave a substantially simpler proof of this fact. In [22], it is shown that $\mathcal{H}(M, \mathbb{Z}_2)$ is homotopy equivalent to $\mathcal{B}(M, \mathbb{Z}_2)$. In particular, this means that the cohomology ring $H^*(\mathcal{H}(M, \mathbb{Z}_2), \mathbb{Z}_2)$ is isomorphic to $\mathbb{Z}_2[\lambda]$ where the generator λ is in degree 1.

Definition 6.1. Fix an integer $p \in \mathbb{N}$. Let X be a cubical complex. A flat continuous map $\Phi: X \rightarrow \mathcal{H}(M, \mathbb{Z}_2)$ is called a half-volume p -sweepout if $\Phi^*(\lambda^p) \neq 0$ in $H^*(X, \mathbb{Z}_2)$.

Definition 6.2. A flat continuous map $\Phi: X \rightarrow \mathcal{H}(M, \mathbb{Z}_2)$ is said to have *no concentration of mass* provided

$$\lim_{r \rightarrow 0} \left[\sup_{x \in X} \sup_{y \in M} \mathbf{M}(\Phi(x) \llcorner B(y, r)) \right] = 0.$$

Definition 6.3. Let \mathcal{Q}_p be the set of all half-volume p -sweepouts of M with no concentration of mass. Note that different half-volume p -sweepouts are allowed to have different domains.

Definition 6.4. For an integer $p \in \mathbb{N}$, the half-volume p -width of M is

$$\tilde{\omega}_p = \inf_{\Phi \in \mathcal{Q}_p} \left[\sup_{x \in \text{dom}(\Phi)} \mathbf{M}(\Phi(x)) \right].$$

The sequence $\{\tilde{\omega}_p\}_{p \in \mathbb{N}}$ is called the half-volume spectrum of M .

The next proposition says that the definition of the half-volume p -width is unchanged if we restrict to only those half-volume p -sweepouts whose domain is a subset of $I(2p+1, k)$ for some $k \in \mathbb{N}$. This was proved for (ordinary) p -sweepouts by Y. Li [14].

Proposition 6.5. *For every $\varepsilon > 0$, there is half-volume p -sweepout $\Psi: Y \rightarrow \mathcal{H}(M, \mathbb{Z}_2)$ whose domain Y is a cubical subcomplex of $I(2p+1, k)$ for some $k \in \mathbb{N}$ and which satisfies*

$$\sup_{y \in Y} \mathbf{M}(\Psi(y)) \leq \tilde{\omega}_p + \varepsilon.$$

Moreover, Ψ has no concentration of mass.

Proof. This follows from the proof of [14, Proposition 3.2]. Indeed, given $\varepsilon > 0$, we can find a half-volume p -sweepout $\Phi: X \rightarrow \mathcal{H}(M, \mathbb{Z}_2) \subset \mathcal{B}(M, \mathbb{Z}_2)$ with no concentration of mass and with

$$\sup_{x \in X} \mathbf{M}(\Phi(x)) \leq \tilde{\omega}_p + \varepsilon.$$

Applying the construction in [14] verbatim to Φ produces a cubical subcomplex Y of $I(2p+1, k)$ and a flat continuous p -sweepout $\Psi: Y \rightarrow \mathcal{B}(M, \mathbb{Z}_2)$ with the property that $\Psi(Y) \subset \Phi(X)$. In particular, the inclusion $\Psi(Y) \subset \Phi(X)$ implies that Ψ takes values in $\mathcal{H}(M, \mathbb{Z}_2)$ and that

$$\sup_{y \in Y} \mathbf{M}(\Psi(y)) \leq \tilde{\omega}_p + \varepsilon,$$

and that Ψ has no concentration of mass. \square

6.2. The Penalized Functionals. Intuitively, we expect that each $\tilde{\omega}_p$ is associated with a critical point of the area functional restricted to the space of half-volume cycles. Therefore, $\tilde{\omega}_p$ should be achieved by a half-volume CMC. Rather than directly developing a min-max theory for the area on $\mathcal{H}(M, \mathbb{Z}_2)$, we first apply min-max on all of $\mathcal{B}(M, \mathbb{Z}_2)$ with a functional that consists of the area plus a term that penalizes the distance to the space of half-volume cycles.

Fix a closed Riemannian manifold (M^{n+1}, g) with dimension $3 \leq n+1 \leq 7$. Let $\mathfrak{h} = \frac{1}{2} \text{Vol}(M)$. For each $k \in \mathbb{N}$ define $f_k: [0, \text{Vol}(M)] \rightarrow \mathbb{R}$ by

$$f_k(v) = k(v - \mathfrak{h})^2.$$

Then define $E_k: \mathcal{B}(M, \mathbb{Z}_2) \rightarrow \mathbb{R}$ by

$$E_k(T) = \mathbf{M}(T) + f_k(\text{Vol}(\Omega)) = \mathbf{M}(T) + k(\text{Vol}(\Omega) - \mathfrak{h})^2$$

where $\Omega \in \mathcal{C}(M)$ satisfies $\partial\Omega = T$. Since $|\text{Vol}(\Omega) - \mathfrak{h}| = |\text{Vol}(M \setminus \Omega) - \mathfrak{h}|$, it is easy to see that the definition does not depend on the choice of Ω .

Fix $p \in \mathbb{N}$. For each $k \in \mathbb{N}$, select a half-volume p -sweepout $\Phi_k^*: X_k \rightarrow \mathcal{H}(M, \mathcal{F}, \mathbb{Z}_2)$ with no concentration of mass for which

$$\sup_{x \in X_k} \mathbf{M}(\Phi_k^*(x)) \leq \tilde{\omega}_p + \frac{1}{k}.$$

By Proposition 6.5, we can further ensure that X_k is a cubical subcomplex of $I(2p+1, \ell)$ for some $\ell \in \mathbb{N}$. By applying discretization [35, Theorem 1.11] followed by interpolation [35, Theorem 1.12], we can replace Φ_k^* with a new \mathbf{F} -continuous map $\Phi_k: X_k \rightarrow \mathcal{B}(M, \mathbf{F}, \mathbb{Z}_2)$ such that

$$\sup_{x \in X_k} \mathbf{M}(\Phi_k(x)) \leq \tilde{\omega}_p + \frac{2}{k}, \quad \sup_{x \in X_k} |\text{Vol}(\Phi_k(x)) - \mathfrak{h}| \leq \frac{1}{k}.$$

Recall here that $\text{Vol}(\Phi_k(x))$ stands for $\text{Vol}(\Omega)$ for any set Ω with $\partial\Omega = \Phi_k(x)$.

Let Π_k be the X_k -homotopy class of the map Φ_k . Observe that

$$L^{E_k}(\Pi_k) \leq \sup_{x \in X_k} E_k(\Phi_k(x)) = \sup_{x \in X_k} \mathbf{M}(\Phi_k(x)) + k \sup_{x \in X_k} |\text{Vol}(\Phi_k(x)) - \mathfrak{h}|^2 \leq \tilde{\omega}_p + \frac{3}{k}.$$

Now suppose for contradiction that

$$\liminf_{k \rightarrow \infty} L^{E_k}(\Pi_k) < \tilde{\omega}_p.$$

Then, after passing to a subsequence, we can find an $\eta > 0$ and maps $\Psi_k: X_k \rightarrow \mathcal{B}(M, \mathbb{Z}_2)$ homotopic to Φ_k such that

$$\sup_{x \in X_k} E_k(\Psi_k(x)) \leq \tilde{\omega}_p - \eta$$

for all k . Note in particular this implies that for any $x \in X_k$ we have

$$\mathbf{M}(\Psi_k(x)) \leq \tilde{\omega}_p - \eta, \quad \text{and} \quad |\text{Vol}(\Psi_k(x)) - \mathfrak{h}| \leq \sqrt{\frac{\tilde{\omega}_p}{k}}$$

Now recall the deformation retraction $\theta: \mathcal{B}(M, \mathbb{Z}_2) \times [0, 1] \rightarrow \mathcal{H}(M, \mathbb{Z}_2)$ constructed in [22].

Lemma 6.6. *There is a continuous function $w: [0, \infty) \rightarrow [0, \infty)$ with $w(0) = 0$ such that*

$$\mathbf{M}(\theta(T, 1)) \leq \mathbf{M}(T) + w(|\text{Vol}(T) - \mathfrak{h}|)$$

for any $T \in \mathcal{B}(M, \mathbb{Z}_2)$.

Proof. Let $f: M \rightarrow [0, 1]$ be a Morse function and let $U_t = \{f < t\}$ for $t \in [0, 1]$. We first claim that for every $\varepsilon > 0$ there is a $\delta > 0$ such that for all $t \in [0, 1]$ and all $W \subset U_t$ we have

$$\text{Vol}(W) \geq \text{Vol}(U_t) - \delta \implies \mathbf{M}(\partial W) \geq \mathbf{M}(\partial U_t) - \varepsilon.$$

Suppose to the contrary that for some $\varepsilon > 0$ there is no $\delta > 0$ which makes the assertion true. Then for $\delta = \frac{1}{n}$ there exist $t_n \in [0, 1]$ and $W_n \subset U_{t_n}$ with $\text{Vol}(W_n) > \text{Vol}(U_{t_n}) - \frac{1}{n}$ but $\mathbf{M}(\partial W_n) < \mathbf{M}(\partial U_{t_n}) - \varepsilon$. After passing to a subsequence, we can suppose that $t_n \rightarrow t_\infty \in [0, 1]$ and that $W_n \rightarrow W_\infty \in \mathcal{C}(M)$. Note that $W_\infty \subset U_{t_\infty}$ and $\text{Vol}(W_\infty) = \text{Vol}(U_{t_\infty})$ and so $W_\infty = U_{t_\infty}$. Since $t \mapsto \mathbf{M}(\partial U_t)$ is continuous, this implies that

$$\begin{aligned} \mathbf{M}(\partial U_{t_\infty}) &\leq \liminf \mathbf{M}(W_n) \\ &\leq \liminf \mathbf{M}(\partial U_{t_n}) - \varepsilon = \mathbf{M}(\partial U_{t_\infty}) - \varepsilon. \end{aligned}$$

This is a contradiction.

Now, to prove the lemma, it suffices to show that for each $\varepsilon > 0$ there is a $\delta > 0$ such that

$$|\text{Vol}(T) - \mathfrak{h}| < \delta \implies \mathbf{M}(\theta(T, 1)) \leq \mathbf{M}(T) + \varepsilon.$$

So let $\varepsilon > 0$ be given and choose δ according to the previous claim. Assume that T satisfies $|\text{Vol}(T) - \mathfrak{h}| < \delta$. Choose a set Ω with $\text{Vol}(\Omega) \leq \mathfrak{h}$ and $\partial\Omega = T$ and note that $\mathfrak{h} - \text{Vol}(\Omega) < \delta$. The set $\theta(T, 1)$ is defined as $\partial(\Omega \cup U_t)$ where t is chosen so that $\text{Vol}(\Omega \cup U_t) = \mathfrak{h}$. Note that $\Omega \cap U_t \subset U_t$ and that $\text{Vol}(\Omega \cap U_t) = \text{Vol}(\Omega) + \text{Vol}(U_t) - \text{Vol}(\Omega \cup U_t) \geq \text{Vol}(U_t) - \delta$. According to the previous claim, this implies that

$$\mathbf{M}(\partial(\Omega \cap U_t)) \geq \mathbf{M}(\partial U_t) - \varepsilon.$$

Finally note that

$$\begin{aligned} \mathbf{M}(\partial(\Omega \cup U_t)) &\leq \mathbf{M}(\partial\Omega) + \mathbf{M}(\partial U_t) - \mathbf{M}(\partial(\Omega \cap U_t)) \\ &\leq \mathbf{M}(\partial\Omega) + \varepsilon, \end{aligned}$$

as needed. \square

By the lemma, for k large enough, the map $\Xi_k: X_k \rightarrow \mathcal{H}(M, \mathbb{Z}_2)$ given by $\Xi_k(x) = \theta(\Psi_k(x), 1)$ is a half-volume p -sweepout with no concentration of mass (guaranteed by properties of θ [22]) which satisfies

$$\sup_{x \in X_k} \mathbf{M}(\Xi_k(x)) \leq \tilde{\omega}_p - \eta + w\left(\sqrt{\frac{\tilde{\omega}_p}{k}}\right) < \tilde{\omega}_p.$$

This contradicts the definition of $\tilde{\omega}_p$. It follows that

$$L^{E_k}(\Pi_k) \rightarrow \tilde{\omega}_p$$

as $k \rightarrow \infty$.

6.3. Bounding the Mean Curvature. We assume that $3 \leq \dim(M) \leq 5$ from this point onward. Again $p \in \mathbb{N}$ is fixed. Applying the E_k -min-max theorem in the homotopy class Π_k gives the existence of critical points for E_k . We would like to show that these critical points converge to a regular limit as $k \rightarrow \infty$. The key point is to show that the mean curvature of the critical points stays uniformly bounded as $k \rightarrow \infty$. Then we can appeal to the compactness results of the previous section.

The following diameter bound for stable CMCs is due to Elbert-Nelli-Rosenberg [9].

Theorem 6.7 ([9]). *Assume M^{n+1} is a Riemannian manifold of dimension $3 \leq n+1 \leq 5$. Assume the sectional curvatures of M are bounded below by $-\kappa$ for some $\kappa \geq 0$. Let Σ be a stable, immersed H -CMC hypersurface in M . There is a constant $c = c(n, H, \kappa)$ so that $\text{dist}_\Sigma(q, \partial\Sigma) \leq c$ for all $q \in \Sigma$ provided $|H| > 2\sqrt{\kappa}$.*

The next proposition will be used to show that half-volume CMCs with bounded area cannot consist entirely of tiny components.

Proposition 6.8. *Let M^{n+1} be a closed Riemannian manifold. Let $\Lambda = \partial\Omega$ be a smooth, almost-embedded hypersurface with non-vanishing mean curvature. Assume that the mean curvature vector of Λ points consistently into or consistently out of Ω . Assume that $\text{Area}(\Lambda) \leq A$ and that $\frac{1}{3} \text{Vol}(M) \leq \text{Vol}(\Omega) \leq \frac{2}{3} \text{Vol}(M)$. There is a positive constant $\delta > 0$, depending only on M and A , such that Λ has a component with extrinsic diameter at least δ .*

Proof. Let I_M denote the isoperimetric profile of M . By the asymptotics of the isoperimetric profile for small volumes, there are constants $c > 0$ and $V_0 > 0$ such that $I_M(v) \geq cv^{n/(n+1)}$ for all $v \in (0, V_0)$. Let $\alpha = \frac{1}{3} \text{Vol}(M)$. Shrinking V_0 if necessary, we can suppose that

$$v^{\frac{n}{n+1}} \geq \frac{2Av}{c\alpha}$$

for all $v \in (0, V_0)$. Now choose $\delta > 0$ so that any closed hypersurface in M with extrinsic diameter less than δ encloses a region with volume less than V_0 .

Suppose for contradiction that every component of Λ has extrinsic diameter smaller than δ . Replacing Ω by $M \setminus \Omega$ if necessary, we can suppose that Ω is the union of the small volume regions enclosed by these components. Let J be the number of connected components of Λ , and list the volumes of these components as v_1, \dots, v_J . Note that

$$\sum_{j=1}^J v_j \geq \alpha$$

and that $v_j \leq V_0$ for all $j = 1, \dots, J$. Thus the area of Λ satisfies

$$\text{Area}(\Lambda) \geq c \sum_{j=1}^J v_j^{n/(n+1)} \geq \frac{2A}{\alpha} \sum_{j=1}^J v_j \geq 2A.$$

This contradicts the definition of A . □

Now we can prove that the mean curvature does not blow up. Let \mathcal{S}_k be the set of all $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ such that

- (i) (V, T) is stationary for E_k ,
- (ii) (V, T) satisfies property (R) (4.10) for E_k with $m = 2p + 1$,
- (iii) $\|V\|(M) \leq \tilde{\omega}_p + 3/k$.
- (iv) $|\text{Vol}(T) - \mathfrak{h}| \leq \sqrt{2\tilde{\omega}_p/k}$.

Note that every (V, T) in \mathcal{S}_k is regular in the sense of Proposition 4.18. Also note that the critical points produced by the E_k min-max theory in the homotopy class Π_k belong to \mathcal{S}_k when k is sufficiently large. Define

$$H_k = \inf\{c \geq 0 : \text{every } (V, T) \in \mathcal{S}_k \text{ has } c\text{-bounded first variation}\}.$$

and note that $H_k \leq k \text{Vol}(M)$.

Proposition 6.9. *Assume that $3 \leq n+1 \leq 5$. Then $H = \sup\{H_k : k \in \mathbb{N}\}$ is finite.*

Proof. Suppose to the contrary that $H_k \rightarrow \infty$. Then we can find $(V_k, T_k) \in \mathcal{S}_k$ such that the mean curvature of the CMC portion Λ_k of $\text{supp } \|V_k\|$ goes to infinity as $k \rightarrow \infty$. By assumption, each (V_k, T_k) has the following property:

- (R) For any collection of $N = N(2p+1)$ concentric annuli $\text{An}(x, s_1, r_1), \dots, \text{An}(x, s_N, r_N)$ with $2r_j < s_{j+1}$ for all j , (V_k, T_k) is (E_k, \mathcal{F}) -almost-minimizing in at least one of the annuli.

Let $\eta_{x,\rho}$ denote the map which rescales by a factor of $1/\rho$ centered at x . Choose $\rho_k \rightarrow 0$ so that $\eta_{x,\rho_k}(\Lambda_k)$ has mean curvature 1.

Choose $\delta > 0$ according to Proposition 6.8. Note that Λ_k has a uniform upper bound on area, and $\Lambda_k = \partial\Omega_k$ with $\text{Vol}(\Omega_k) \rightarrow \mathfrak{h}$. Therefore, to get a contradiction, it suffices to show that, for some large k , every connected component of Λ_k has extrinsic diameter less than δ . If this is not the case, then (passing to a subsequence) we can find points $x_k \in \Lambda_k$ such that the connected component Γ_k of Λ_k containing x_k has extrinsic diameter at least δ . Fix some positive numbers $s_1 < r_1 < s_2 < r_2 < \dots < s_N < r_N$ satisfying $2r_j < s_{j+1}$ for all j . Choose a sequence $\sigma_k \rightarrow \infty$ such that $\rho_k \sigma_k \rightarrow 0$.

For each k , consider the collection of $N(2p+1)$ -concentric annuli $\text{An}(x_k, s_1/\sigma_k, r_1/\sigma_k), \dots, \text{An}(x_k, s_N/\sigma_k, r_N/\sigma_k)$ and note that this collection is admissible for property (R). Therefore (passing to a subsequence) there is a $j \in \{1, \dots, N\}$ such that (V_k, T_k) is (E_k, \mathcal{F}) -almost-minimizing in

$$\text{An}_k = \text{An}(x_k, s_j/\sigma_k, r_j/\sigma_k)$$

for all k . For notational convenience, let $s = s_j$ and $r = r_j$. Then for k large enough, since Γ_k has extrinsic diameter at least δ , it follows that Γ_k crosses the boundary of An_k . Hence, by Proposition 5.7, it follows that Γ_k is volume preserving stable in the interior of An_k .

Define

$$C_k = \partial B\left(x_k, \frac{s+r}{2\sigma_k}\right) \subset \text{An}_k.$$

Choose a point $y_k \in \Gamma_k \cap C_k$, which must exist since Γ_k has diameter at least δ . Now consider the balls

$$B_k = B\left(y_k, \frac{r-s}{2\sigma_k}\right) \subset \text{An}_k, \quad D_k = B\left(y_k, \frac{r-s}{4\sigma_k}\right) \subset B_k.$$

Note that Γ_k must intersect $\partial B(y_k, t)$ for every $0 < t < (r-s)/(2\sigma_k)$ since Γ_k is connected and contains both x_k and y_k .

Define $\Lambda'_k = \eta_{y_k, \rho_k}(\Lambda_k)$, $\Gamma'_k = \eta_{y_k, \rho_k}(\Gamma_k)$, $B'_k = \eta_{y_k, \rho_k}(B_k)$, and $D'_k = \eta_{y_k, \rho_k}(D_k)$. Note that Λ'_k has constant mean curvature 1 and is volume preserving stable in B'_k . Because $\rho_k \sigma_k \rightarrow 0$, the balls B'_k resemble large, nearly Euclidean balls when k is large. Note that Γ'_k must be strongly stable as a 1-CMC immersion in either D'_k or $B'_k \setminus D'_k$, but this violates the diameter estimate of Theorem 6.7. Indeed, there are points $q_k \in \Gamma'_k \cap D'_k$ with $\text{dist}(q_k, \partial D'_k) \rightarrow \infty$ and points $z_k \in \Gamma'_k \cap [B'_k \setminus D'_k]$ with $\text{dist}(z_k, \partial B'_k \cup \partial D'_k) \rightarrow \infty$. \square

This implies the following compactness property.

Proposition 6.10. *Assume that $3 \leq n+1 \leq 5$. Consider a sequence $(V_k, T_k) \in \mathcal{S}_k$. Then, up to a subsequence, $(V_k, T_k) \rightarrow (V, T)$ in $\text{VZ}(M, \mathbb{Z}_2)$. Moreover, (V, T) has the regularity described in Proposition 4.18 and the mean curvature of the CMC part of (V, T) is at most H . We have $\text{Vol}(T) = \mathfrak{h}$. Finally, (V, T) satisfies property (R') (5.5) with $m = 2p+1$ and ρ depending only on H .*

Proof. The mean curvature of (V_k, T_k) is uniformly bounded by H . Choosing Ω_k so that $\partial\Omega_k = T_k$, one has $\text{Vol}(\Omega_k) \rightarrow \mathfrak{h}$. Therefore the result follows from Proposition 5.9. \square

6.4. The Lifting Construction. At this point, we assume the metric g on M is generic so that, by the results of Appendix B, we can assume that M is bumpy, that M has no half-volume minimal hypersurfaces, that half-volume CMCs are isolated in M , and that every closed almost-embedded half-volume CMC in M is actually embedded.

Define $H = \sup\{H_k : k \in \mathbb{N}\}$ as in the previous subsection. Choose a scale $\rho > 0$ depending on H according to Proposition 6.9. Let \mathcal{S} be the set of all pairs $(V, T) \in \text{VZ}(M, \mathbb{Z}_2)$ such that

- (i) $\|V\|(M) \leq \tilde{\omega}_p$,
- (i) (V, T) satisfies property (R') (5.5) with $m = 2p+1$ and scale ρ ,
- (ii) $\text{Vol}(T) = \mathfrak{h}$,
- (iii) There is a smooth, embedded CMC hypersurface Λ with non-zero mean curvature $|H_\Lambda| \leq H$. Moreover, there there exist a (possibly empty) collection of closed minimal hypersurfaces $\Sigma_1, \dots, \Sigma_k$ and a collection of multiplicities $m_1, \dots, m_k \in \mathbb{N}$ such that

$$V = |\Lambda| + \sum_{i=1}^k m_i |\Sigma_i|.$$

Here the hypersurfaces $\Lambda, \Sigma_1, \dots, \Sigma_k$ are all disjoint.

Proposition 6.11. *Assume the metric g on M is generic and that $3 \leq n+1 \leq 5$. Then the set \mathcal{S} is finite.*

Proof. Suppose to the contrary that \mathcal{S} is not finite. Then there is a sequence of distinct elements $(V_k, T_k) \in \mathcal{S}$. Let Λ_k be the CMC portion of (V_k, T_k) . Since the mean curvature of Λ_k stays uniformly bounded, we can argue as in Proposition 5.9 to see that that, up to a subsequence, $(V_k, T_k) \rightarrow (V, T) \in \mathcal{S}$. Let Λ denote the CMC portion of (V, T) . Note, in particular, Λ is guaranteed to be embedded with non-zero mean curvature, since the metric g admits no half-volume minimal hypersurfaces, and every almost-embedded half-volume CMC is embedded.

Since Λ is genuinely embedded and the convergence of the CMC portion occurs with multiplicity one, Allard's regularity theorem implies that the convergence $\Lambda_k \rightarrow \Lambda$ is actually smooth. This contradicts that half-volume CMCs are isolated in M ; see Corollary B.4. Thus $\Lambda_k = \Lambda$ for sufficiently large k .

Now let V'_k be the minimal portion of V_k and let V' be the minimal portion of V . Then Property (R') implies that for each $x \in \text{supp}\|V'\|$ there is a $\rho(x) > 0$ such that if $s < r < \rho(x)$ then the convergence of V'_k to V' is smooth in $\text{An}(x, s, r)$. This implies that the convergence $V'_k \rightarrow V'$ is actually smooth away from finitely many points. Therefore, $V'_k = V'$ for sufficiently large k , as otherwise it would be possible to extract a Jacobi field on V' . This shows that in fact $(V_k, T_k) = (V, T)$ for sufficiently large k and this is a contradiction. \square

The following lemma is due to Marques and Neves.

Lemma 6.12 ([18] Corollary 3.6). *Let \mathcal{T} be a finite subset of $\mathcal{B}(M, \mathbb{Z}_2)$. If $\eta > 0$ is sufficiently small, then every map $\Phi: S^1 \rightarrow B_\eta^{\mathcal{F}}(\mathcal{T})$ is homotopically trivial.*

Enumerate the set $\mathcal{S} = \{(W_1, S_1), \dots, (W_Q, S_Q)\}$ and let $\mathcal{T} = \{S_1, \dots, S_Q\}$. Choose $\eta > 0$ according to the previous lemma. Recall the set \mathcal{S}_k defined in the previous subsection. By Proposition 6.9 and Proposition 5.9, it follows that for any $\eta > 0$ there is a $K \in \mathbb{N}$ such that $\mathcal{F}((V, T), \mathcal{S}) \leq \eta/4$ for all $(V, T) \in \mathcal{S}_k$ provided $k \geq K$. Note in particular that if $\mathcal{F}((V, T), \mathcal{S}) \leq \eta/4$ then $\mathcal{F}(T, \mathcal{T}) \leq \eta/4$.

Now fix some $k \geq K$. Consider a pulled-tight E_k -min-max sequence $\Psi_{k,j}: X_k \rightarrow \mathcal{B}(M, \mathcal{F}, \mathbb{Z}_2)$ for the homotopy class Π_k . Choose a sequence $\ell_j \rightarrow \infty$ so that

$$\sup\{\mathbf{F}(\Psi_{k,j}(x), \Psi_{k,j}(y)) : \alpha \in X_k(\ell_j), x, y \in \alpha\} < \frac{\eta}{2}.$$

Let $\tilde{Z}_{k,j} \subset X_k$ be the union of all the cells $\alpha \in X_k(\ell_j)$ such that

$$\mathcal{F}(\Psi_{k,j}(x), \mathcal{S}) \geq \eta/2$$

for all vertices $x \in \alpha$. Here and in the following, we use $\Psi_{k,j}(x)$ to denote its image under the natural inclusion $\mathcal{B}(M, \mathbb{Z}_2) \rightarrow \text{VZ}(M, \mathbb{Z}_2)$. Then

$$\mathcal{F}(\Psi_{k,j}(x), \mathcal{S}) \geq \eta/2$$

for all $x \in \tilde{Z}_{k,j}$. Let $\tilde{Y}_{k,j} = \overline{X_k \setminus \tilde{Z}_{k,j}}$. Consider the restricted sequence $\Psi_{k,j}|_{\tilde{Z}_{k,j}}$. Then either

- (i) $L^{E_k}(\{\Psi_{k,j}|_{\tilde{Z}_{k,j}}\}) < L^{E_k}(\Pi_k)$, or
- (ii) no element in $\mathcal{K}(\{\Psi_{k,j}|_{\tilde{Z}_{k,j}}\})$ satisfies property (R) for E_k with $m = 2p + 1$.

Indeed, if both (i) and (ii) failed then there would be some element

$$(W, S) \in \mathcal{K}(\{\Psi_{k,j}|_{\tilde{Z}_{k,j}}\}) \cap \mathcal{S}_k.$$

But then $\mathcal{F}((W, S), \mathcal{S}) \leq \eta/4$ contradicting the choice of $\tilde{Z}_{k,j}$.

We can now argue exactly as in [35, Section 5, Step 2] to obtain a new min-max sequence $\Psi'_{k,j}: X_k \rightarrow \mathcal{B}(M, \mathbb{Z}_2)$ such that

- (i) X_k can be decomposed into $Y_{k,j}$ and $Z_{k,j}$ where $Y_{k,j} = \overline{X_k \setminus Z_{k,j}}$. Moreover, we have

$$(\Psi'_{k,j}|_{Y_{k,j}})^* \lambda = 0 \text{ in } H^1(Y_i, \mathbb{Z}_2), \quad (\Psi'_{k,j}|_{Z_{k,j}})^* (\lambda^{p-1}) \neq 0 \text{ in } H^{p-1}(Z_{k,j}, \mathbb{Z}_2)$$

provided j is large enough,

- (ii) $L^{E_k}(\{\Psi'_{k,j}\}) = L^{E_k}(\Pi_k)$,
- (iii) $\limsup_{j \rightarrow \infty} \sup\{E_k(\Psi'_{k,j}(z)) : z \in Z_{k,j}\} < L^{E_k}(\Pi_k)$.

Define $\tilde{E}_k: \mathcal{C}(M) \rightarrow \mathbb{R}$ by

$$\tilde{E}_k(\Omega) = \mathbf{M}(\partial\Omega) + f_k(\text{Vol}(\Omega)).$$

Let \tilde{X}_k be the double cover of X_k associated to the cohomology class $(\Psi'_{k,j})^* \lambda$. Let $\tilde{\Pi}_{k,j}$ be the $(\tilde{X}_k, \tilde{Z}_{k,j})$ relative homotopy class of $\Psi'_{k,j}$. By the same argument as [35, Section 5, Step 3], if j is large enough then

$$L^{\tilde{E}_k}(\tilde{\Pi}_{k,j}) > \sup_{z \in \tilde{Z}_{k,j}} \tilde{E}_k(\Psi'_{k,j}(z)).$$

Moreover, we have $L^{\tilde{E}_k}(\tilde{\Pi}_{k,j}) \rightarrow L^{E_k}(\Pi_k)$ as $j \rightarrow \infty$.

6.5. Conclusion of the Proof. Let $h: M \rightarrow \mathbb{R}$ be a smooth Morse function satisfying property (T) (2.10). Choose a sequence $\varepsilon_j \rightarrow 0$. Define $F_{k,j}: \mathcal{C}(M) \rightarrow \mathbb{R}$ by

$$F_{k,j}(\Omega) = \mathbf{M}(\partial\Omega) - \varepsilon_j \int_{\Omega} h + f_k(\text{Vol}(\Omega)).$$

Then $F_{k,j}(\Omega) = \tilde{E}_k(\Omega) - \varepsilon_j \int_{\Omega} h$, and we have

$$L^{F_{k,j}}(\tilde{\Pi}_{k,j}) > \sup_{z \in \tilde{Z}_{k,j}} F_{k,j}(\Psi'_{k,j}(z))$$

for large enough j , and $L^{F_{k,j}}(\tilde{\Pi}_{k,j}) \rightarrow L^{E_k}(\Pi_k)$ as $j \rightarrow \infty$. Therefore, by applying min-max theory for $F_{k,j}$ (Theorem 1.10) in the relative homotopy class $\tilde{\Pi}_{k,j}$, there exist critical points $(V_{k,j}, \Omega_{k,j})$ for $F_{k,j}$ with $F_{k,j}(V_{k,j}, \Omega_{k,j}) \rightarrow L^{E_k}(\Pi_k)$. By Proposition 5.5, after passing to a subsequence, $(V_{k,j}, \partial\Omega_{k,j}) \rightarrow (V_k, T_k) \in \text{VZ}(M, \mathbb{Z}_2)$, (V_k, T_k) is stationary for E_k , $E_k(V_k, T_k) = L^{E_k}(\Pi_k)$, and V_k is induced by either

- (i) a smooth, almost-embedded, multiplicity one constant mean curvature surface with non-zero mean curvature; or
- (ii) a collection of minimal hypersurfaces with multiplicities.

It remains to take a limit as $k \rightarrow \infty$. Again the key step is to show that the mean curvature doesn't blow up.

Proposition 6.13. *Consider the pairs $(V_k, T_k) \in \text{VZ}(M, \mathbb{Z}_2)$ defined above. Then the mean curvature of $\Lambda_k = \text{supp } \|V_k\|$ stays uniformly bounded as $k \rightarrow \infty$.*

Proof. Suppose to the contrary that the mean curvature H_k of Λ_k goes to ∞ . Choose $\rho_k \rightarrow 0$ so that $\eta_{x, \rho_k}(\Lambda_k)$ has mean curvature 1.

Let $\Lambda_{k,j}$ denote $\text{supp } \|V_{k,j}\|$ and let $H_{k,j}$ be the mean curvature of $\Lambda_{k,j}$. Then for fixed k , we have $H_{k,j} \rightarrow H_k$ as $j \rightarrow \infty$. Choose $\delta > 0$ according to Proposition 6.8. Then, if j is large enough depending on k , $\Lambda_{k,j}$ has a connected component $\Gamma_{k,j}$ with extrinsic diameter at least δ . Pick a point $x_{k,j} \in \Gamma_{k,j}$.

Let $N = N(2p+1)$ and fix $s_1 < r_1 < s_2 < r_2 < \dots < s_N < r_N$ with $2r_i < s_{i+1}$. Choose a sequence $\sigma_k \rightarrow \infty$ such that $\rho_k \sigma_k \rightarrow 0$. Now fix an integer $k \in \mathbb{N}$. For each j , consider the collection of concentric annuli $\text{An}(x_{k,j}, s_1/\sigma_k, r_1/\sigma_k), \dots, \text{An}(x_{k,j}, s_N/\sigma_k, r_N/\sigma_k)$. Since $(V_{k,j}, \Omega_{k,j})$ satisfies property (R) (3.15), there is an $i(k) \in \{1, \dots, N\}$ and a (non-relabeled) subsequence of j 's such that $(V_{k,j}, \Omega_{k,j})$ is $(F_{k,j}, \mathcal{F})$ -almost minimizing in the annulus $\text{An}(x_{k,j}, s_{i(k)}/\sigma_k, r_{i(k)}/\sigma_k)$ for all j . Since $\Gamma_{k,j}$ has extrinsic diameter at least δ , it follows that $\Gamma_{k,j}$ must cross the boundary of $\text{An}(x_{k,j}, s_{i(k)}/\sigma_k, r_{i(k)}/\sigma_k)$. By Corollary 5.3, it follows that $\Gamma_{k,j}$ is volume preserving stable for $A^{\varepsilon_j h}$ in $\text{An}(x_{k,j}, s_{i(k)}/\sigma_k, r_{i(k)}/\sigma_k)$.

Now consider the rescaled surfaces $\Gamma'_{k,j} = \eta_{x_{k,j}, \rho_k}(\Gamma_{k,j})$. Passing to a further subsequence in j , we can suppose that $x_{k,j} \rightarrow x_k$ as $j \rightarrow \infty$. For a fixed k , the surfaces $\Gamma'_{k,j}$ have a uniform upper bound on mean curvature, a uniform upper bound on area, and are volume preserving stable for $A^{\varepsilon_j \rho_k h}$ in $\text{An}'_{k,j} = \eta_{x_{k,j}, \rho_k}(\text{An}(x_{k,j}, s_{i(k)}/\sigma_k, r_{i(k)}/\sigma_k))$. Therefore, these surfaces have a uniform curvature estimate, and converge smoothly to a volume preserving stable CMC hypersurface Γ'_k in $\text{An}'_k = \eta_{x_k, \rho_k}(\text{An}(x_k, s_{i(k)}/\sigma_k, r_{i(k)}/\sigma_k))$ with mean curvature 1. Note that the surface Γ'_k is connected and intersects both the inner and outer boundary of An'_k . Therefore, we can now argue exactly as in Proposition 6.9 to get a contradiction for large enough k . \square

Once we know that the mean curvature doesn't blow up, it follows that (V_k, T_k) satisfies property (R') (5.5) at a uniform scale ρ that does not depend on k . This implies that $(V_k, T_k) \rightarrow (V, T)$, $\|V\|(M) \leq \tilde{\omega}_p$, $\text{Vol}(T) = \mathfrak{h}$, and V is induced by either

- (i) a smooth, almost-embedded, multiplicity one constant mean curvature surface with non-zero mean curvature; or
- (ii) a collection of minimal hypersurfaces with multiplicities.

Choose Ω so that $\partial\Omega = T$. Note that case (ii) cannot occur because then some collection of minimal hypersurfaces in M would bound Ω . But we have assumed the metric g is generic so that this cannot happen by Proposition B.1. Therefore case (i) occurs and $\partial\Omega$ is an almost-embedded constant mean curvature hypersurface.

It remains to show that $\|V\|(M) = \tilde{\omega}_p$. Since $E_k(V_k, T_k) \rightarrow \tilde{\omega}_p$, it is equivalent to show that

$$k(\text{Vol}(\Omega_k) - \mathfrak{h})^2 \rightarrow 0$$

as $k \rightarrow \infty$, where $\partial\Omega_k = T_k$. But this follows from the fact that (V_k, T_k) is stationary for E_k and the fact that the mean curvature does not blow up. Indeed, let H_k denote the mean curvature of $\text{supp } \|V_k\|$. Then we know that $|H_k| = 2k|\text{Vol}(\Omega_k) - \mathfrak{h}|$ since (V_k, T_k) is stationary for E_k . It follows that

$$k(\text{Vol}(\Omega_k) - \mathfrak{h})^2 = \frac{|H_k|^2}{4k},$$

and this goes to 0 as $k \rightarrow \infty$ since H_k is uniformly bounded. Therefore, $\|V\|(M) = \tilde{\omega}_p$, as needed. This proves Theorem 1.1, and Corollary 1.2 follows immediately.

APPENDIX A. CONDITION (T)

Let $h: M \rightarrow \mathbb{R}$ be a smooth Morse function. Given a regular point $x \in M$ for h , let $\Gamma(x)$ be the level set of h passing through x . Then define $v(h, x)$ to be the vanishing order at x of the mean curvature $H_{\Gamma(x)}$, regarded as a function on $\Gamma(x)$. Recall that our min-max theory requires h to satisfy the following property (Definition 2.10):

(T) For every regular point x of h , we have $v(h, x) < \infty$.

This property is used to show that the touching set of an almost-embedded $(h + h_0)$ -PMC is contained in a countable union of $(n - 1)$ -dimensional manifolds. The goal of this appendix is to show that an arbitrary Morse function h can be perturbed slightly in the smooth topology to a nearby Morse function that satisfies condition (T).

Note the following lemmas.

Lemma A.1. *Assume that $u: M \rightarrow \mathbb{R}$ is a smooth Morse function. Let C be the set of all critical points of u . Let $G \subset M \setminus C$ be a closed set such that for each $x \in G$ we have $v(u, x) < \infty$. Then there is a finite $k \in \mathbb{N}$ such that $v(u, x) \leq k$ for all $x \in G$. Moreover, if w is a sufficiently small smooth perturbation of u then $v(w, x) \leq k$ for all $x \in G$ as well.*

Proof. The function $v(u, x)$ is upper semicontinuous, i.e.,

$$\limsup_{w \rightarrow u, y \rightarrow x} v(w, y) \leq v(u, x).$$

If there were points $x_n \in G$ with $v(u, x_n) \rightarrow \infty$, then there would be a convergent subsequence $x_{n_k} \rightarrow x \in G$. But then the semi-continuity implies $v(u, x) = \infty$, contrary to assumption. Likewise, if there is a sequence $u_n \rightarrow u$ such that there are points $x_n \in G$ with $v(u_n, x_n) \geq k + 1$, then we can find a convergent subsequence $x_{n_k} \rightarrow x \in G$. The semi-continuity implies that $v(u, x) \geq k + 1$, contrary to assumption. \square

Lemma A.2. *Assume that $u: M \rightarrow \mathbb{R}$ is a smooth Morse function. Let C be the set of all critical points of u . Let $G \subset M \setminus C$ be a closed set such that for each $x \in G$ we have $v(u, x) < \infty$. Then there is an $\varepsilon > 0$ such that $v(u, x) < \infty$ for every $x \in M$ with $\text{dist}(x, G) \leq \varepsilon$.*

Proof. Again this follows from the upper semicontinuity of the vanishing order. Suppose to the contrary that there is no such ε . Then there is a sequence of points $x_n \rightarrow x \in G$ with $v(u, x_n) \rightarrow \infty$. But this implies that $v(u, x) = \infty$. \square

Now we can prove the main result of this appendix.

Proposition A.3. *Let $h: M \rightarrow \mathbb{R}$ be a smooth Morse function. Then there exist Morse functions satisfying property (T) which are arbitrarily close to h in $C^\infty(M)$.*

Proof. The topology on $C^\infty(M)$ is complete and metrizable. Fix a complete metric ρ on $C^\infty(M)$ inducing the topology of $C^\infty(M)$. Consider the following two constructions.

Construction 1: Let $u: M \rightarrow \mathbb{R}$ be a Morse function. Let $B = \{b_1, \dots, b_k\}$ be the set of all critical values of u and let C denote the set of all critical points of u . Fix a small radius $r > 0$. Let Σ_i be a smooth, closed, embedded surface which coincides with $\{u = b_i\}$ outside $N_r(C) = \{x \in M : \text{dist}(x, C) < r\}$. Choose a short time s , and let $\Sigma_{i,t}$, $t \in [0, s]$ be the surface obtained by running mean curvature flow for time t starting from Σ_i . There exists a smooth, time-dependent vector field X_t on M which coincides with the mean curvature vector of $\Sigma_{i,t}$ on $\Sigma_{i,t}$ at time t . We can suppose that X_t is supported in $\{x \in M : \text{dist}(u(x), B) < r\}$ for all t . Let ζ be a smooth function on M which is identically 1 outside $N_{2r}(C)$ and identically 0 inside $N_r(C)$. Let ϕ_t be the flow of ζX_t and define

$$u_t = u \circ \phi_t^{-1}.$$

If t is small enough, the following properties hold:

- (i) u_t is Morse and has exactly the same critical points and critical values as u ,
- (ii) if $x \in (u_t)^{-1}(B) \setminus N_{2r}(C)$ then $v(u_t, x) < \infty$ (c.f. [38, Proposition 3.9]),
- (iii) if $\text{dist}(b, B) \geq r$ then $\{u_t = b\} = \{u = b\}$.

Moreover, $u_t \rightarrow u$ in $C^\infty(M)$ as $t \rightarrow 0$. We select a small time τ and then set $\tilde{u} = u_\tau$. \blacksquare

Construction 2: Let $u: M \rightarrow \mathbb{R}$ be a Morse function. Let $B = \{b_1, \dots, b_k\}$ be the set of critical values of u . Let $\mathcal{N}_\rho(B) = \{b \in \mathbb{R} : \text{dist}(b, B) < \rho\}$. Fix a small number $r > 0$. By the argument in [38, Proposition 3.9], there exists a small positive number s and smooth family of smooth functions $w_t: M \rightarrow \mathbb{R}$, $t \in [0, s]$ such that $\Sigma_{b,t} = \{w_t = b\}$ is the surface obtained from running mean curvature flow for time t starting from $\{u = b\}$ for all $b \in \mathbb{R} \setminus \mathcal{N}_{r/2}(B)$ and all $t \in [0, s]$. Choose a smooth cut-off function ζ which is identically 1 outside $u^{-1}(\mathcal{N}_r(B))$ and identically 0 inside $u^{-1}(\mathcal{N}_{r/2}(B))$. Then set

$$u_t = \zeta w_t + (1 - \zeta)u.$$

If t is small enough then the following properties hold:

- (i) u_t is Morse and has exactly the same critical points and critical values as u ,
- (ii) if $x \in M \setminus (u_t)^{-1}(\mathcal{N}_{2r}(B))$ then $v(u_t, x) < \infty$ (c.f. [38, Proposition 3.9]),
- (iii) $(u_t)^{-1}(B) = u^{-1}(B)$.

Moreover, $u_t \rightarrow u$ in $C^\infty(M)$ as $t \rightarrow 0$. We select a small time τ and then set $\tilde{u} = u_\tau$. \blacksquare

Now choose a smooth Morse function $h: M \rightarrow \mathbb{R}$. Let C be the set of critical points of h and let $B = \{b_1, \dots, b_k\}$ be the set of critical values. Fix some $\varepsilon_0 > 0$. To begin, apply Construction 1 with $u = h$ and $r = r_0 = 1$ and τ small enough to ensure that

$$\rho(u, \tilde{u}) < \frac{\varepsilon_0}{2}.$$

Then define $h_1 = \tilde{u}$ and let G_1 be a closed set containing a neighborhood of the set $(h_1)^{-1}(B) \setminus N_2(C)$ with the property that $v(h_1, x) < \infty$ for all $x \in G_1$. Note that such a G_1 exists by Lemma A.2. We further select $\varepsilon_1 > 0$ small enough that if $\rho(w, h_1) < \varepsilon_1$ then

- (i) $v(w, x) < \infty$ for all $x \in G_1$,
- (ii) $w^{-1}(B) \setminus N_2(C) \subset G_1$.

This is possible by Lemma A.1.

Next we apply Construction 2 with $u = h_1$ and $r = r_1 = 2^{-1}$ and τ small enough to ensure that

$$\rho(u, \tilde{u}) < \min \left\{ \frac{\varepsilon_1}{2}, \frac{\varepsilon_0}{4} \right\}.$$

Then we set $h_2 = \tilde{u}$. We let G_2 be a closed set containing G_1 and a neighborhood of $M \setminus (h_2)^{-1}(\mathcal{N}_1(B))$ with the property that $v(h_2, x) < \infty$ for all $x \in G_2$. Such a set G_2 exists by Lemma A.2. Note that G_2 also contains a neighborhood of $(h_2)^{-1}(B) \setminus N_2(C) = (h_1)^{-1}(B) \setminus N_2(C)$. We select $\varepsilon_2 > 0$ small enough that if $\rho(w, h_2) < \varepsilon_2$ then

- (i) $v(w, x) < \infty$ for all $x \in G_2$,
- (ii) $w^{-1}(B) \setminus N_2(C) \subset G_2$.
- (iii) $M \setminus w^{-1}(\mathcal{N}_1(B)) \subset G_2$.

This is possible by Lemma A.1.

Now, for $n \geq 2$, suppose inductively that $h_n: M \rightarrow \mathbb{R}$ is a smooth Morse function with the same critical points and critical values as h . Let $r_n = 2^{-n}$. Assume that G_n is a closed subset of $M \setminus C$ such that $v(h_n, x) < \infty$ for all $x \in G_n$. Suppose that G_n contains a neighborhood of $(h_n)^{-1}(B) \setminus N_{4r_n}(C)$ when n is odd, and that G_n contains a neighborhood of $(h_n)^{-1}(B) \setminus N_{8r_n}(C)$ when n is even. Further suppose that G_n contains a neighborhood of $M \setminus (h_n)^{-1}(\mathcal{N}_{4r_n}(B))$ when n is even, and that G_n contains a neighborhood of $M \setminus (h_n)^{-1}(\mathcal{N}_{8r_n}(B))$ when n is odd. Choose ε_n small enough that if $\rho(w, h_n) < \varepsilon_n$ then

- (i) $v(w, x) < \infty$ for all $x \in G_n$,
- (ii) $w^{-1}(B) \setminus N_{8r_n}(C) \subset G_n$.
- (iii) $M \setminus w^{-1}(\mathcal{N}_{8r_n}(B)) \subset G_n$.

This is possible by Lemma A.1.

Now if n is even, apply Construction 1 with $u = h_n$ and $r = r_n$ and τ small enough to ensure that

$$\rho(u, \tilde{u}) < \min \left\{ \frac{\varepsilon_n}{2}, \frac{\varepsilon_{n-1}}{2^2}, \dots, \frac{\varepsilon_0}{2^{n+1}} \right\}.$$

Then set $h_{n+1} = \tilde{u}$. Let G_{n+1} be a closed set containing G_n together with a neighborhood of $(h_{n+1})^{-1}(B) \setminus N_{4r_{n+1}}(C)$ with the property that $v(h_{n+1}, x) < \infty$ for all $x \in G_{n+1}$. Again such a choice is possible by Lemma A.2. Further note that G_{n+1} contains a neighborhood of $M \setminus (h_{n+1})^{-1}(\mathcal{N}_{8r_{n+1}}(B))$ since $M \setminus (h_{n+1})^{-1}(\mathcal{N}_{8r_{n+1}}(B)) = M \setminus (h_n)^{-1}(\mathcal{N}_{4r_n}(B))$ and G_n contains a neighborhood of $M \setminus (h_n)^{-1}(\mathcal{N}_{4r_n}(B))$.

If n is odd, apply Construction 2 with $u = h_n$ and $r = r_n$ and τ small enough to ensure that

$$\rho(u, \tilde{u}) < \min \left\{ \frac{\varepsilon_n}{2}, \frac{\varepsilon_{n-1}}{2^2}, \dots, \frac{\varepsilon_0}{2^{n+1}} \right\}.$$

Then set $h_{n+1} = \tilde{u}$. Let G_{n+1} be a closed set containing G_n and a neighborhood of $M \setminus h_{n+1}^{-1}(\mathcal{N}_{4r_{n+1}})$ with the property that $v(h_{n+1}, x) < \infty$ for all $x \in G_{n+1}$. Again such a choice is possible by Lemma A.2. Note that G_{n+1} contains a neighborhood of $(h_{n+1})^{-1}(B) \setminus N_{8r_{n+1}}(C)$ since $(h_{n+1})^{-1}(B) \setminus N_{8r_{n+1}}(C) = (h_n)^{-1}(B) \setminus N_{4r_n}(C)$ and G_n contains a neighborhood of $(h_n)^{-1}(B) \setminus N_{4r_n}(C)$.

Now observe that (h_n) is a Cauchy sequence with respect to ρ and so it converges to some smooth function h^* with $\rho(h, h^*) < \varepsilon_0$. If ε_0 is chosen sufficiently small, then h^* is also Morse. Since each critical point y of h is also a critical point for h^* with $h(y) = h^*(y)$, it follows that

h^* has the same critical points and critical values as h . Moreover, we know that $v(h^*, x) < \infty$ for all $x \in G = \cup_{n=1}^{\infty} G_n$. To complete the proof, it remains to show that $G = M \setminus C$.

First, we claim that G contains $(h^*)^{-1}(B) \setminus C$. It suffices to show that G contains $(h^*)^{-1}(B) \setminus N_{8r_n}(C)$ for all n . To see this, observe that by construction one has

$$\rho(h^*, h_n) < \sum_{j=1}^{\infty} \frac{\varepsilon_n}{2^j} = \varepsilon_n.$$

By the choice of ε_n , this implies that $(h^*)^{-1}(B) \setminus N_{8r_n}(C) \subset G_n \subset G$. This proves the claim.

Second, we claim that G contains $M \setminus (h^*)^{-1}(B)$. It suffices to show that G contains $M \setminus (h^*)^{-1}(\mathcal{N}_{8r_n}(B))$ for all n . To see this, again observe that

$$\rho(h^*, h_n) < \sum_{j=1}^{\infty} \frac{\varepsilon_n}{2^j} = \varepsilon_n.$$

By the choice of ε_n , this implies that $M \setminus (h^*)^{-1}(\mathcal{N}_{8r_n}(B)) \subset G_n \subset G$. This completes the proof. \square

APPENDIX B. GENERIC METRICS

In this appendix, we prove three theorems about generic metrics. The first result says that, generically, half-volume minimal hypersurfaces do not exist. The second result says that, generically, half-volume CMCs are isolated. Finally, the third result says that, generically, every almost-embedded half volume CMC is actually embedded.

Proposition B.1. *Let M be a closed manifold. Then, for a generic smooth metric g on M , there is no smooth, closed, (not necessarily connected) minimal hypersurface which encloses half the volume of M .*

Proof. Given $C > 0$ and $I \in \mathbb{N}$ and a smooth metric g on M , let $\mathcal{M}_{C,I}(g)$ be the set of all smooth, closed, embedded (not necessarily connected) minimal hypersurfaces in (M, g) with area at most C and index at most I . Let $\mathcal{G}_{C,I}$ be the set of all smooth metrics g on M such that the set $\mathcal{M}_{C,I}(g) = \{\Sigma_1, \dots, \Sigma_n\}$ is finite, each surface Σ_i is non-degenerate, and none of the surfaces Σ_i encloses half the volume of M .

Let \mathcal{G} be the set of all smooth metrics on M . It suffices to show that for each fixed $C > 0$ and $I \in \mathbb{N}$, the set $\mathcal{G}_{C,I}$ is open and dense in \mathcal{G} . First, we claim that $\mathcal{G}_{C,I}$ is open. Indeed, choose some $g \in \mathcal{G}_{C,I}$ and write $\mathcal{M}_{C,I} = \{\Sigma_1, \dots, \Sigma_n\}$. Then, since each Σ_i is non-degenerate, there is a neighborhood U_1 of g in \mathcal{G} such that for every $\tilde{g} \in U_1$ and every $i = 1, \dots, n$ there is a unique small perturbation of Σ_i to a hypersurface $\Sigma_i(\tilde{g})$ which is minimal and non-degenerate in (M, \tilde{g}) . Then, by Sharp's compactness theorem [26], it follows that for \tilde{g} in a possibly smaller neighborhood U_2 of g we have

$$\mathcal{M}_{C,I}(\tilde{g}) \subset \{\Sigma_1(\tilde{g}), \dots, \Sigma_n(\tilde{g})\}.$$

Finally, since none of the hypersurfaces Σ_i enclose half the volume of (M, g) , it follows that for \tilde{g} in a potentially smaller neighborhood U_3 of g , none of the hypersurfaces $\Sigma_i(\tilde{g})$ enclose half the volume of (M, \tilde{g}) . This proves that $\mathcal{G}_{C,I}$ is open.

It remains to show that $\mathcal{G}_{C,I}$ is dense in \mathcal{G} . Choose an arbitrary metric $g \in \mathcal{G}$. Then there exists a small perturbation g_1 of g which is bumpy. By Sharp's compactness theorem, it follows that the set $\mathcal{M}_{C,I}(g_1) = \{\Sigma_1, \dots, \Sigma_n\}$ is finite and consists entirely of non-degenerate minimal hypersurfaces. Again there is a small neighborhood U_1 of g_1 in \mathcal{G} such that for every $\tilde{g} \in U_1$ there is a small perturbation of Σ_i to a hypersurface $\Sigma_i(\tilde{g})$ which is minimal and non-degenerate

in (M, \tilde{g}) . As above, by Sharp's compactness theorem, for \tilde{g} in a possibly smaller neighborhood U_2 of g_1 we have

$$\mathcal{M}_{C,I}(\tilde{g}) \subset \{\Sigma_1(\tilde{g}), \dots, \Sigma_n(\tilde{g})\}.$$

We now inductively perturb g_1 to ensure that no surfaces in this collection encloses half the volume.

First consider $\Sigma_1(g_1)$. If $\Sigma_1(g_1)$ does not enclose half the volume of (M, g_1) then we set $g^1 = g_1$. Suppose instead that $\Sigma_1 = \partial\Omega_1$ does enclose half the volume of M . Then we set $g^1 = e^{2\phi}g_1$ where ϕ is non-negative, and vanishes on the closure of Ω_1 , and is positive at some point in $M \setminus \Omega_1$. By selecting ϕ close enough to 0, we can ensure that $g^1 \in U_2$. Then $\Sigma_1(g^1) = \Sigma_1$ as sets but $\Sigma_1(g^1)$ no longer encloses half the volume of (M, g^1) .

Next consider $\Sigma_2(g^1)$. If $\Sigma_2(g^1)$ does not enclose half the volume of (M, g^1) then set $g^2 = g^1$. Suppose instead that $\Sigma_2(g^1) = \partial\Omega_2$ does enclose half the volume of (M, g^1) . Then we set $g^2 = e^{2\phi}g^1$ where ϕ is non-negative, and vanishes on the closure of Ω_2 together with the support of $\Sigma_1(g^1)$, and is positive at some point in $M \setminus \Omega_2$. By selecting ϕ close enough to 0, we can ensure that $g^2 \in U_2$ and that $\Sigma_1(g^2)$ does not enclose half the volume of (M, g^2) . Note that $\Sigma_2(g^2) = \Sigma_2(g^1)$ as sets but $\Sigma_2(g^2)$ no longer encloses half the volume of (M, g^2) .

Continuing this process inductively, one arrives at a metric $g^n \in U_2$ with the property that none of the hypersurfaces $\Sigma_1(g^n), \dots, \Sigma_n(g^n)$ enclose half the volume of (M, g^n) . Thus $g^n \in \mathcal{G}_{C,I}$ and so g^n is the required perturbation of g . \square

Let g be a smooth Riemannian metric on M . Assume that $u: \Sigma \rightarrow M$ is a two-sided, null-homologous embedding with a preferred choice of normal vector ν with respect to g . Then there is a unique region Ω whose boundary is $u(\Sigma)$ and such that ν points into Ω . We define the volume enclosed by u to be the volume of Ω . Note that the enclosed volume depends only on the image $u(\Sigma)$ and the preferred choice of normal vector, and not the particular immersion u .

Now suppose that $u: \Sigma \rightarrow M$ is a two-sided, null-homologous CMC almost-embedding with non-zero mean curvature and a preferred choice of normal vector ν with respect to g . Consider another immersion $w: \Sigma \rightarrow M$ such that

$$w(x) = \exp_{u(x)}^g(f(x)\nu)$$

for some $f \in C^{j,\alpha}(\Sigma)$ with small norm. Regard $w(\Sigma)$ as an integer multiplicity current (oriented by the preferred choice of normal vector). Then exactly one of the following is true (depending on whether the normal vector and the mean curvature vector point in the same direction or opposite directions):

- (i) There exists a unique $\Theta = \theta(x)\llcorner \mathcal{H}^{n+1}$ with $\theta(x) \in \{0, 1, 2\}$ such that $\partial\Theta = w(\Sigma)$.
- (ii) There exists a unique $\Theta = \theta(x)\llcorner \mathcal{H}^{n+1}$ with $\theta(x) \in \{-1, 0, 1\}$ such that $\partial\Theta = w(\Sigma)$.

We define the volume enclosed by w to be

$$\int_M \theta(x) d\mathcal{H}^{n+1}(x).$$

Observe that the enclosed volume of w actually depends only on $w(\Sigma)$ and the preferred choice of normal vector, and not on the particular immersion w .

Let M^{n+1} and Σ^n be smooth, closed manifolds. Fix $q, j \in \mathbb{N}$ and $\alpha \in (0, 1)$ with $q \geq j + 3$. Let \mathcal{P} be the set of all triples $(g, [u], H)$ where g is a C^q Riemannian metric on M , and $[u]$ is the equivalence class of a simple, two-sided, null-homologous, $C^{j,\alpha}$ immersion $u: \Sigma \rightarrow M$ with constant mean curvature H and a preferred choice of normal vector. Let $\mathcal{P}' \subset \mathcal{P}$ be a neighborhood of the triples $(g, [u], H)$ where u is an almost-embedding. We can suppose that \mathcal{P}' is small enough that the enclosed volume function is defined on \mathcal{P}' . Then let $\mathcal{M} \subset \mathcal{P}'$ be

the set of all triples $(g, [u], H)$ where the volume enclosed by u is equal to $\frac{1}{2} \text{Vol}(M, g)$. Define $\Pi: \mathcal{M} \rightarrow \Gamma$ by $\Pi(g, [u], H) = g$. Let \mathcal{M}_∞ be the subset of \mathcal{M} consisting of triples $(g, [u], H)$ where g and u are smooth.

Proposition B.2. *There is an open subset $\mathcal{U} \subset \mathcal{M}$ containing \mathcal{M}_∞ such that \mathcal{U} has the structure of a separable C^{q-j} Banach manifold. Moreover, the differential of Π restricted to \mathcal{U} is Fredholm with index 0.*

Proof. The proof is almost identical to White's manifold structure theorem [31]. We will provide a sketch of the argument. Let Γ be the space of all C^q Riemannian metrics on M . Fix an arbitrary smooth metric $g \in \Gamma$. Fix a smooth, simple, two-sided, null-homologous, almost-embedding $u: \Sigma \rightarrow M$ with a preferred choice of normal vector ν , constant mean curvature H with respect to g , and enclosed volume $\frac{1}{2} \text{Vol}(M, g)$ with respect to g . Here we compute the mean curvature of H with respect to ν and the enclosed volume is defined as above.

Define a map $\mathcal{H}: \Gamma \times C^{j,\alpha}(\Sigma) \rightarrow C^{j-2,\alpha}(\Sigma)$ by letting $\mathcal{H}(h, f)$ be the mean curvature of the immersion

$$x \mapsto \exp_{u(x)}^g(f(x)\nu),$$

computed with respect to h . The differential of the function \mathcal{H} at the point $(g, 0)$ is well-known. If φ is a smooth function on M and $h(t) = e^{2t\varphi}g$ then

$$D_1\mathcal{H}(h'(0)) = -2\varphi H - n\langle \nabla\varphi, \nu \rangle.$$

If $w \in C^{j,\alpha}(\Sigma)$ then

$$D_2\mathcal{H}(w) = Jw = \Delta w + (|A|^2 + \text{Ric}(\nu, \nu))w,$$

where the Laplacian and second fundamental form are those of $u(\Sigma)$ with respect to g .

Now consider the volume functional. Define $\mathcal{V}: \Gamma \times C^{j,\alpha}(\Sigma) \rightarrow \mathbb{R}$ by letting $\mathcal{V}(h, f)$ be the volume enclosed by the immersion

$$x \mapsto \exp_{u(x)}^g(f(x)\nu),$$

computed with respect to h . As explained above, this is well-defined if $\|f\|_{j,\alpha}$ is small-enough, and it depends only on the image of the immersion and the preferred choice of normal vector. The differential of \mathcal{V} is also well-known. Let Ω be the region enclosed by $u(\Sigma)$. If ψ is a smooth function on M supported in Ω and $h(t) = e^{2t\psi}g$ then

$$D_1\mathcal{V}(h'(0)) = (n+1) \int_{\Omega} \psi dV_g.$$

Also one has

$$D_2\mathcal{V}(w) = - \int_{\Sigma} w dA_{u^*g}.$$

for $w \in C^{j,\alpha}(\Sigma)$.

Now define

$$\begin{aligned} \mathcal{F}: \Gamma \times C^{j,\alpha}(\Sigma) \times \mathbb{R} &\rightarrow C^{j-2,\alpha}(\Sigma) \times \mathbb{R}, \\ \mathcal{F}(h, f, a) &= \left(\mathcal{H}(h, f) - a, \mathcal{V}(h, f) - \frac{\text{Vol}(M, h)}{2} \right). \end{aligned}$$

Also define

$$\begin{aligned} L: C^{j,\alpha}(\Sigma) \times \mathbb{R} &\rightarrow C^{j-2,\alpha}(\Sigma) \times \mathbb{R}, \\ L(w, b) &= D\mathcal{F}(0, w, b) = \left(Jw - b, - \int_{\Sigma} w dA_g \right). \end{aligned}$$

The key point is that L is formally self-adjoint, with finite dimensional kernel K , and image $(K^\perp) \cap (C^{j-2,\alpha}(\Sigma) \times \mathbb{R})$. Moreover, it is easy to see from the explicit formulas for $D_1\mathcal{H}$ and $D_1\mathcal{V}$ along conformal paths of metrics that no non-zero element of K is orthogonal to the image of $D_1\mathcal{F}$. As in [31], these facts imply that $D\mathcal{F}$ is onto, and that the kernel of $D\mathcal{F}$ is complemented.

It now follows by the implicit function theorem that $\mathcal{N} = \mathcal{F}^{-1}(0,0)$ has the structure of a C^{q-j} Banach manifold in a neighborhood of $(g,0,H)$. Now consider the map

$$\begin{aligned}\Pi: \Gamma \times C^{j,\alpha}(\Sigma) \times \mathbb{R} &\rightarrow \Gamma, \\ \Pi(h, f, a) &= h.\end{aligned}$$

Again, the same argument as in [31] shows that the differential $D(\Pi|_{\mathcal{N}})$, evaluated at $(g,0,H)$, is Fredholm with index 0.

The above construction gives a chart for \mathcal{M} in a neighborhood of every $(g,[u],H)$ in which g is smooth and in which u is a smooth almost-embedding. Let \mathcal{U} be the open subset of \mathcal{M} covered by these charts. As in [31], one can show that the transition maps between charts are C^{q-j} and therefore that this gives \mathcal{U} the claimed manifold structure. \square

Corollary B.3. *Let M^{n+1} be a smooth, closed manifold. Then for a (Baire) generic set of smooth Riemannian metrics on M , there are no almost-embedded half-volume CMC hypersurfaces Σ in (M,g) which carry a non-zero function φ such that $J_\Sigma\varphi$ is constant and $\int_\Sigma\varphi = 0$.*

Proof. Fix q, j , and α as before. Let Γ_q be the space of C^q metrics on M . Let Γ_∞ be the space of smooth Riemannian metrics on M . Fix a smooth, closed n -dimensional manifold Σ and consider the manifold $\mathcal{U} \subset \mathcal{M}$ containing \mathcal{M}_∞ from the previous proposition. Since $\Pi: \mathcal{U} \rightarrow \Gamma_q$ is Fredholm with index 0 and \mathcal{U} is separable, it follows from Smale's infinite dimensional Sard theorem [28] that the set \mathcal{R} of regular values of Π is of second category in Γ_q . According to [32, Theorem 2.10], the set $\mathcal{R} \cap \Gamma_\infty$ is therefore of second category in Γ_∞ . Now observe that if $g \in \mathcal{R} \cap \Gamma_\infty$ and $u: \Sigma \rightarrow M$ is a smooth half-volume almost-embedded CMC then there is no non-zero function φ on Σ such that $J_\Sigma\varphi$ is constant and $\int_\Sigma\varphi = 0$. Since there are only countably many possible diffeomorphism classes for Σ , the result follows. \square

Corollary B.4. *Let M^{n+1} be a smooth, closed manifold. Then for a (Baire) generic set of smooth Riemannian metrics on M , every almost-embedded half-volume CMC hypersurface Σ in (M,g) is isolated.*

Proof. Let Σ be an almost-embedded half-volume CMC hypersurface in (M,g) . Consider the operator $\mathcal{F}: C^{j,\alpha}(\Sigma) \times \mathbb{R} \rightarrow C^{j-2,\alpha}(\Sigma) \times \mathbb{R}$ given by

$$F(f, a) = \left(\mathcal{H}(g, f) - a, \mathcal{V}(g, f) - \frac{\text{Vol}(M, g)}{2} \right).$$

Then the linearization of \mathcal{F} at $(0,0)$ is

$$L(w, b) = \left(J_\Sigma w - b, - \int_\Sigma w \right).$$

Therefore the previous corollary implies that, for generic g , the map L is injective and hence an isomorphism. The inverse function theorem now implies that Σ is isolated. \square

Proposition B.5. *Assume that M^{n+1} is a smooth, closed manifold. Then for a (Baire) generic set of smooth Riemannian metrics on M , every almost-embedded half-volume CMC hypersurface in (M,g) is actually embedded.*

Proof. Again we can follow White's self-transversality argument [33] almost identically. Let $\Pi: \mathcal{M}_\infty \rightarrow \Gamma_\infty$ be the projection to the metric, and let \mathcal{M}_{reg} be the set of points $(g, [u], H)$ in \mathcal{M}_∞ at which the differential $D\Pi$ has trivial kernel. Pick a point $(g, [u], H) \in \mathcal{M}_{\text{reg}}$ and assume that $u: \Sigma \rightarrow M$ is an almost-embedding. Pick a small open set $W \subset M$ such that $u(\Sigma)$ is embedded in W . Let

$$V_0 = \{f \in C^\infty(\Sigma) : Jf \text{ is supported in } W\}.$$

Let $\Omega^2\Sigma = \{(x_1, x_2) \in \Sigma \times \Sigma : x_1 \neq x_2\}$ and then let

$$C = \{(x_1, x_2) \in \Omega^2\Sigma : u(x_1) = u(x_2)\}$$

which is a compact subset of $\Omega^2(\Sigma)$. According to [33, Theorem 29], there is a finite dimensional subspace $V \subset V_0$ such that given any $(x_1, x_2) \in C$ and $a_1, a_2 \in \mathbb{R}$, there is an $f \in V$ with $f(x_i) = a_i$ for $i = 1, 2$.

Let f_1, \dots, f_d be a basis for V . Next, we can argue as in [33, Proposition 8] to find maps

$$\gamma: B^d(0, \varepsilon) \rightarrow C^\infty(M),$$

$$\mathcal{W}: B^d(0, \varepsilon) \times \Sigma \rightarrow M$$

such that

- (i) $\gamma(0) = 0$,
- (ii) $\mathcal{W}(0, \cdot) = u(\cdot)$,
- (iii) for each $\tau \in B^d(0, \varepsilon)$, the map $\mathcal{W}(\tau, \cdot)$ is a half-volume CMC immersion with respect to the metric $e^{2\gamma(\tau)}g$,
- (iv) For each $i = 1, \dots, d$ we have

$$\left. \frac{d}{dt} \right|_{t=0} \mathcal{W}(te_i, x) = f_i(x).$$

Indeed, since Jf_i is supported in W , it is clear from the formula for $D_1\mathcal{F}$ that there exists $\varphi_i \in C^\infty(M)$ such that $D_1\mathcal{F}(v_i) = (-Jf_i, \int_\Sigma f_i)$, where $h_i(t) = e^{2t\varphi_i}g$ and $v_i = h'_i(0)$. Then define

$$\begin{aligned} \gamma: \mathbb{R}^d \times M &\rightarrow \mathbb{R}, \\ \gamma(\tau, x) &= \sum_{i=1}^d 2\tau_i \varphi_i(x). \end{aligned}$$

Since $(g, [u], H) \in \mathcal{M}_{\text{reg}}$, it follows that for a sufficiently small $\varepsilon > 0$, for each $\tau \in B^d(0, \varepsilon)$ there is a unique function $w(\tau)$ with small norm on Σ such that

$$\mathcal{W}(\tau, x) = \exp_{u(x)}^g(w(\tau)\nu)$$

is a half-volume CMC immersion with respect to $e^{2\gamma(\tau)}g$ with mean curvature $H(\tau)$. Moreover, the map

$$\mathcal{W}: B^d(0, \varepsilon) \times \Sigma \rightarrow M$$

is smooth. Finally, observe that $\mathcal{F}(e^{2\gamma(\tau)}g, w(\tau), H(\tau)) \equiv 0$, and hence

$$\begin{aligned} (0, 0) &= \frac{d}{d\tau_i} \mathcal{F}(e^{2\gamma(\tau)}g, w(\tau), H(\tau)) \\ &= D_1\mathcal{F}(v_i) + L\left(\frac{d}{d\tau_i}w, \frac{d}{d\tau_i}H\right), \end{aligned}$$

where L is defined as above. This implies that

$$L\left(\frac{d}{d\tau_i}w, \frac{d}{d\tau_i}H\right) = L(f_i, 0)$$

Since $(g, [u], H) \in \mathcal{M}_{\text{reg}}$ the operator L is invertible, and therefore this implies that

$$\frac{d}{dt}\Big|_{t=0} \mathcal{W}(te_i, x) = \frac{d}{d\tau_i}w = f_i,$$

as needed.

Let $\Delta_2 M = \{(x, x) \in M^2 : x \in M\}$. The same argument as [33, Theorem 14] now shows that the map

$$\begin{aligned} \tilde{\mathcal{U}}: B^d(0, \varepsilon) \times \Omega^2 \Sigma &\rightarrow M \times M, \\ \tilde{\mathcal{U}}(\tau, x_1, x_2) &= (\mathcal{U}(\tau, x_1), \mathcal{U}(\tau, x_2)) \end{aligned}$$

is a submersion at every point in $\tilde{\mathcal{U}}^{-1}(\Delta^2 M)$. From this construction, it now follows as in [33, Theorem 15] that, for a generic smooth metric g on M , every half-volume CMC immersion is self-transverse. In particular, for such metrics, every almost-embedded half-volume CMC is embedded. \square

REFERENCES

- [1] Frederick Almgren. The homotopy groups of the integral cycle groups. *Topology*, 1(4):257–299, 1962.
- [2] Frederick Almgren. The theory of varifolds. *Mimeographed notes*, 1965.
- [3] Vladimir I Arnold. *Arnold’s problems*. Springer, 2004.
- [4] J Lucas Barbosa, Manfredo do Carmo, and Jost Eschenburg. Stability of hypersurfaces of constant mean curvature in riemannian manifolds. In *Manfredo P. do Carmo–Selected Papers*, pages 291–306. Springer, 2012.
- [5] Costante Bellettini, Otis Chodosh, and Neshan Wickramasekera. Curvature estimates and sheeting theorems for weakly stable cmc hypersurfaces. *Advances in Mathematics*, 352:133–157, 2019.
- [6] Costante Bellettini and Neshan Wickramasekera. The inhomogeneous allen–cahn equation and the existence of prescribed-mean-curvature hypersurfaces. *arXiv preprint arXiv:2010.05847*, 2020.
- [7] Otis Chodosh and Christos Mantoulidis. Minimal surfaces and the allen–cahn equation on 3-manifolds: Index, multiplicity, and curvature estimates. *Annals of Mathematics*, 191(1):213–328, 2020.
- [8] Akashdeep Dey. Existence of multiple closed CMC hypersurfaces with small mean curvature. *J. Differential Geom.*, 125(2):379–403, 2023.
- [9] Maria Elbert, Barbara Nelli, and Harold Rosenberg. Stable constant mean curvature hypersurfaces. *Proceedings of the American Mathematical Society*, 135(10):3359–3366, 2007.
- [10] Pedro Gaspar and Marco AM Guaraco. The allen–cahn equation on closed manifolds. *Calculus of Variations and Partial Differential Equations*, 57:1–42, 2018.
- [11] Pedro Gaspar and Marco AM Guaraco. The weyl law for the phase transition spectrum and density of limit interfaces. *Geometric and Functional Analysis*, 29:382–410, 2019.
- [12] Mikhael Gromov. Dimension, non-linear spectra and width. In *Geometric Aspects of Functional Analysis: Israel Seminar (GAFA) 1986–87*, pages 132–184. Springer, 2006.
- [13] Kei Irie, Fernando Marques, and André Neves. Density of minimal hypersurfaces for generic metrics. *Annals of Mathematics*, 187(3):963–972, 2018.
- [14] Yangyang Li. Existence of infinitely many minimal hypersurfaces in higher-dimensional closed manifolds with generic metrics. *Journal of Differential Geometry*, 124(2):381–395, 2023.
- [15] Yevgeny Liokumovich, Fernando Marques, and André Neves. Weyl law for the volume spectrum. *Annals of Mathematics*, 187(3):933–961, 2018.
- [16] F. C. Marques and A. Neves. Morse theory for the area functional. *São Paulo J. Math. Sci.*, 15(1):268–279, 2021.
- [17] Fernando C Marques and André Neves. Morse index and multiplicity of min-max minimal hypersurfaces. *Cambridge Journal of Mathematics*, 4(4):463–511, 2016.

- [18] Fernando C Marques and André Neves. Existence of infinitely many minimal hypersurfaces in positive ricci curvature. *Inventiones mathematicae*, 209(2):577–616, 2017.
- [19] Fernando C Marques and André Neves. Morse index of multiplicity one min-max minimal hypersurfaces. *Advances in Mathematics*, 378:107527, 2021.
- [20] Fernando C Marques, André Neves, and Antoine Song. Equidistribution of minimal hypersurfaces for generic metrics. *Inventiones mathematicae*, 216(2):421–443, 2019.
- [21] Liam Mazurowski. CMC doublings of minimal surfaces via min-max. *J. Geom. Anal.*, 32(3):Paper No. 104, 28, 2022.
- [22] Liam Mazurowski and Xin Zhou. The half-volume spectrum of a manifold. *arXiv preprint arXiv:2302.07722*, 2023.
- [23] Frank Morgan. Regularity of isoperimetric hypersurfaces in riemannian manifolds. *Transactions of the American Mathematical Society*, 355(12):5041–5052, 2003.
- [24] Jon T Pitts. *Existence and regularity of minimal surfaces on Riemannian manifolds.(MN-27)*, volume 27. Princeton University Press, 2014.
- [25] Richard Schoen and Leon Simon. Regularity of stable minimal hypersurfaces. *Communications on Pure and Applied Mathematics*, 34(6):741–797, 1981.
- [26] Ben Sharp. Compactness of minimal hypersurfaces with bounded index. *Journal of Differential Geometry*, 106(2):317–339, 2017.
- [27] Leon Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis, Australian National University*. Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [28] S. Smale. An infinite dimensional version of sard’s theorem. *American Journal of Mathematics*, 87(4):861, 1965.
- [29] Antoine Song. Existence of infinitely many minimal hypersurfaces in closed manifolds. *Annals of Mathematics*, 197(3):859 – 895, 2023.
- [30] Zhichao Wang and Xin Zhou. Existence of four minimal spheres in S^3 with a bumpy metric. *arXiv preprint arXiv:2305.08755*, 2023.
- [31] Brian White. The space of minimal submanifolds for varying riemannian metrics. *Indiana University Mathematics Journal*, pages 161–200, 1991.
- [32] Brian White. On the bumpy metrics theorem for minimal submanifolds. *American Journal of Mathematics*, 139(4):1149–1155, 2017.
- [33] Brian White. Generic transversality of minimal submanifolds and generic regularity of two-dimensional area-minimizing integral currents. *arXiv preprint arXiv:1901.05148*, 2019.
- [34] Shing-Tung Yau. *Seminar on differential geometry*. Princeton University Press, 1982.
- [35] Xin Zhou. On the multiplicity one conjecture in min-max theory. *Annals of Mathematics*, 192(3):767–820, 2020.
- [36] Xin Zhou. Mean curvature and variational theory. In *ICM—International Congress of Mathematicians. Vol. IV. Sections 5–8*, pages 2696–2717. EMS Press, Berlin, [2023] ©2023.
- [37] Xin Zhou and Jonathan J Zhu. Min-max theory for constant mean curvature hypersurfaces. *Inventiones mathematicae*, 218:441–490, 2019.
- [38] Xin Zhou and Jonathan J Zhu. Existence of hypersurfaces with prescribed mean curvature i-generic min-max. *Cambridge Journal of Mathematics*, 8(2):311–362, 2020.

CORNELL UNIVERSITY, DEPARTMENT OF MATHEMATICS, ITHACA, NEW YORK 14850

Email address: `lmm334@cornell.edu`

CORNELL UNIVERSITY, DEPARTMENT OF MATHEMATICS, ITHACA, NEW YORK 14850

Email address: `xinzhou@cornell.edu`