

SOME REMARKS ON PERIODIC GRADINGS

OKSANA S. YAKIMOVA

ABSTRACT. Let \mathfrak{q} be a finite-dimensional Lie algebra, $\vartheta \in \text{Aut}(\mathfrak{q})$ a finite order automorphism, and \mathfrak{q}_0 the subalgebra of fixed points of ϑ . Using ϑ one can construct a pencil \mathcal{P} of compatible Poisson brackets on $\mathcal{S}(\mathfrak{q})$, and thereby a ‘large’ Poisson-commutative subalgebra $\mathcal{Z}(\mathfrak{q}, \vartheta)$ of $\mathcal{S}(\mathfrak{q})^{\mathfrak{q}_0}$. In this article, we study one particular bracket $\{ , \}_{\infty} \in \mathcal{P}$ and the related Poisson centre \mathcal{Z}_{∞} . It is shown that \mathcal{Z}_{∞} is a polynomial ring, if \mathfrak{q} is reductive.

CONTENTS

Introduction	1
1. Preliminaries on Poisson brackets and polynomial contractions	3
2. Properties of $\mathfrak{q}_{(\infty)}$ and of the Poisson centre $\mathcal{Z}_{\infty} \subset \mathcal{S}(\mathfrak{q}_{(\infty)})$	8
3. The reductive case	9
4. Properties of \mathcal{Z}_{∞} in the reductive case	14
References	20

INTRODUCTION

The ground field \mathbb{k} is algebraically closed and $\text{char}(\mathbb{k}) = 0$. Let $\mathfrak{q} = (\mathfrak{q}, [,], \cdot)$ be a finite-dimensional algebraic Lie algebra, i.e., $\mathfrak{q} = \text{Lie } Q$, where Q is a connected affine algebraic group. The dual space \mathfrak{q}^* is a Poisson variety, i.e., the algebra of polynomial functions on \mathfrak{q}^* , $\mathbb{k}[\mathfrak{q}^*] \simeq \mathcal{S}(\mathfrak{q})$, is equipped with the Lie–Poisson bracket $\{ , \}$. Here $\{x, y\} = [x, y]$ for $x, y \in \mathfrak{q}$. Poisson-commutative subalgebras of $\mathbb{k}[\mathfrak{q}^*]$ are important tools for the study of geometry of the coadjoint action of Q and representation theory of \mathfrak{q} .

0.1. There is a well-known method, *the Lenard–Magri scheme*, for constructing “large” Poisson-commutative subalgebras of $\mathbb{k}[\mathfrak{q}^*]$, which is related to *compatible Poisson brackets*, see e.g. [GZ00]. Two Poisson brackets $\{ , \}'$ and $\{ , \}''$ are said to be *compatible*, if any linear combination $\{ , \}_{a,b} := a\{ , \}' + b\{ , \}''$ with $a, b \in \mathbb{k}$ is a Poisson bracket. Then one defines a certain dense open subset $\Omega_{\text{reg}} \subset \mathbb{k}^2$ that corresponds to the *regular* brackets in the pencil $\mathcal{P} = \{ \{ , \}_{a,b} \mid (a, b) \in \mathbb{k}^2 \}$. Let $\mathcal{Z}_{a,b} \subset \mathcal{S}(\mathfrak{q})$ denote the Poisson centre of $(\mathcal{S}(\mathfrak{q}), \{ , \}_{a,b})$. Then the subalgebra $\mathcal{Z} \subset \mathcal{S}(\mathfrak{q})$ generated by $\mathcal{Z}_{a,b}$ with $(a, b) \in \Omega_{\text{reg}}$ is Poisson-commutative

2010 *Mathematics Subject Classification*. 17B63, 14L30, 17B08, 17B20, 22E46.

Key words and phrases. index of Lie algebra, contraction, commutative subalgebra, symmetric invariants.

This work is funded by the DFG (German Research Foundation) — project number 404144169.

w.r.t. $\{ , \}'$ and $\{ , \}''$. An obvious first step is to take the initial Lie–Poisson bracket $\{ , \}$ as $\{ , \}'$. The rest depends on a clever choice of $\{ , \}''$.

0.2. Let ϑ be an automorphism of \mathfrak{q} of finite order m . Then \mathfrak{q} is equipped with a \mathbb{Z}_m -grading $\mathfrak{q} = \bigoplus_{i=0}^{m-1} \mathfrak{q}_i$, where each \mathfrak{q}_i is an eigenspace of ϑ . One can naturally construct a compatible Poisson bracket $\{ , \}''$ associated with the grading [PY, PY21]. In this case, all Poisson brackets in \mathcal{P} are linear and there are two lines $l_1, l_2 \subset \mathbb{k}^2$ such that $\Omega = \mathbb{k}^2 \setminus (l_1 \cup l_2) \subset \Omega_{\text{reg}}$ and the Lie algebras corresponding to $(a, b) \in \Omega$ are isomorphic to \mathfrak{q} . The lines l_1 and l_2 give rise to new Lie algebras, denoted $\mathfrak{q}_{(0)}$ and $\mathfrak{q}_{(\infty)}$. These new algebras are different *contractions* of \mathfrak{q} . A definition and basic properties of contractions are discussed in Section 1.2. Let $\text{ind } \mathfrak{q}$ denote the *index* of \mathfrak{q} (see Section 1.1). Then we have $\text{ind } \mathfrak{q} \leq \text{ind } \mathfrak{q}_{(t)}$ for $t \in \{0, \infty\}$. Let $\mathcal{S}(\mathfrak{q})^{\mathfrak{q}} \simeq \mathbb{k}[\mathfrak{q}^*]^{\mathfrak{q}}$ be the ring of *symmetric invariants* of \mathfrak{q} , i.e., the Poisson centre of $\mathcal{S}(\mathfrak{q})$. Let further $\mathcal{Z} = \mathcal{Z}(\mathfrak{q}, \vartheta)$ be the Poisson-commutative subalgebra associated with \mathcal{P} . Many features of \mathcal{Z} depend on the properties of $\mathfrak{q}_{(0)}$ and $\mathfrak{q}_{(\infty)}$.

In [PY21, PY23], we have studied the ring $\mathcal{Z}_0 = \mathcal{S}(\mathfrak{q}_{(0)})^{\mathfrak{q}_{(0)}}$ in case $\mathfrak{q} = \mathfrak{g}$ is reductive. For some \mathfrak{g} such that $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$, this is a polynomial ring with $\text{rk } \mathfrak{g}$ generators. However, there are exceptions even if $m = 2$ [Y17]. Partial results were obtained for $\mathcal{Z}_{\infty} = \mathcal{S}(\mathfrak{q}_{(\infty)})^{\mathfrak{q}_{(\infty)}}$. Namely, if \mathfrak{g} is reductive and ϑ is an inner automorphism, then $\mathcal{Z}_{\infty} = \mathcal{S}(\mathfrak{g}_0)$ [PY21].

In Section 1, we collect basic facts on the coadjoint action and symmetric invariants. Explicit descriptions of the algebras $\mathfrak{q}_{(0)}$ and $\mathfrak{q}_{(\infty)}$ are presented in Section 1.3.

0.3. If $\text{ind } \mathfrak{q}_{(\infty)} > \text{ind } \mathfrak{q}$, then \mathcal{Z}_{∞} does not have to be Poisson-commutative. Our first result states that $\{\mathcal{Z}_{\infty}^{\mathfrak{q}_0}, \mathcal{Z}_{\infty}^{\mathfrak{q}_0}\} = 0$, if \mathfrak{q}_0^* contains a regular in \mathfrak{q}^* element. Furthermore, under that assumption, the algebra $\text{alg}\langle \mathcal{Z}, \mathcal{Z}_{\infty}^{\mathfrak{q}_0} \rangle$, generated by \mathcal{Z} and $\mathcal{Z}_{\infty}^{\mathfrak{q}_0}$, is still Poisson-commutative, see Theorem 2.3. Both statements have applications related to the current algebra $\mathfrak{q}[t]$. Namely, one can construct a large Poisson-commutative subalgebra of $\mathcal{S}(\mathfrak{q}[t]^{\vartheta})$ following the ideas of [PY21, Sect. 8]. However, our new approach works for several non-reductive Lie algebras and does not require the assumption that $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$, which is imposed in [PY21]. The construction will appear in a forthcoming paper.

Section 3 contains a brief summary of Kac’s classification of finite order automorphisms for a semisimple \mathfrak{g} [Ka69]. In particular, we describe a relation between the roots of \mathfrak{g} and of \mathfrak{g}^{σ} , where σ is a diagram automorphism of \mathfrak{g} . Then in Section 3.3, we state main results of [PY21] on generators of $\mathcal{Z}(\mathfrak{g}, \vartheta)$. Many crucial properties of $\mathcal{Z}(\mathfrak{g}, \vartheta)$, including its transcendence degree, depend on the equality $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$, see e.g. [PY21, Thm 3.10] or Theorem 3.6 here. Conjecture 3.1 in [PY23] states that $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$ for all \mathfrak{g} and all ϑ . In Section 3.4, we recollect several instances, where the equality holds, and provide a few new positive examples, see Theorem 3.10.

In the more favourable reductive case, we prove that \mathcal{Z}_∞ and $\mathcal{Z}_\infty^{\mathfrak{g}_0}$ are always polynomial rings and describe their generators explicitly, see Section 4. We consider also the non-reductive Lie algebra $\tilde{\mathfrak{g}} = \mathfrak{g}_0 \ltimes \mathfrak{g}_{(\infty)}$ and its coadjoint representation. It is shown that $\text{ind } \tilde{\mathfrak{g}} = \text{rk } \mathfrak{g} + \text{rk } \mathfrak{g}_0$ and that $\tilde{\mathfrak{g}}$ has the *codim-2* property, see Theorem 4.4. Then by Theorem 4.5, $\mathcal{S}(\tilde{\mathfrak{g}})^{\tilde{\mathfrak{g}}}$ is a polynomial ring with $\text{ind } \tilde{\mathfrak{g}}$ generators. These generators are described explicitly.

Non-reductive Lie algebras \mathfrak{q} such that $\mathcal{S}(\mathfrak{q})^{\mathfrak{q}}$ is a polynomial ring with $\text{ind } \mathfrak{q}$ generators attract a lot of attention, see e.g. [J06, PPY, P07, P09, Y14, CM16, FL18]. A quest for this type of algebras continues. Many examples that are found so far are related to particular simple Lie algebras. For instance, assertions of [J06] and [PPY] hold in full generality only for \mathfrak{sl}_n and \mathfrak{sp}_{2n} . Note that our results on $\mathfrak{g}_{(\infty)}$ and $\tilde{\mathfrak{g}}$ are independent of the type of \mathfrak{g} and apply to all finite order automorphisms. We prove also that $\tilde{\mathfrak{g}}$ is a Lie algebra of *Kostant type* in the terminology of [Y14].

The Lie algebra $\tilde{\mathfrak{g}}$ is *quadratic*, i.e., there is a $\tilde{\mathfrak{g}}$ -invariant non-degenerate symmetric bilinear form on $\tilde{\mathfrak{g}}$. This implies that the adjoint and coadjoint representations of $\tilde{\mathfrak{g}}$ are isomorphic, see e.g. [P09, Sect. 1.1]. In [P09, Sect. 4], invariants of the adjoint action of $\tilde{\mathfrak{g}}$ are studied. In the notation of that paper, $\tilde{\mathfrak{g}} = \mathfrak{g}\langle m+1 \rangle_0$. A more general object $\mathfrak{g}\langle nm+1 \rangle_0$ of [P09] can be also interpreted in our context.

Suppose that $\mathfrak{g} = \mathfrak{h}^{\oplus n}$ is a sum of n copies of a reductive Lie algebra \mathfrak{h} and ϑ is a composition of a finite order automorphism $\tilde{\vartheta} \in \text{Aut}(\mathfrak{h})$ and a cyclic permutation of the summands. Formally we have

$$(0.1) \quad \vartheta((x_1, x_2, \dots, x_n)) = (x_n, \tilde{\vartheta}(x_1), x_2, \dots, x_{n-1}) \quad \text{for } (x_1, x_2, \dots, x_n) \in \mathfrak{h} \oplus \mathfrak{h} \oplus \dots \oplus \mathfrak{h}.$$

Then $\tilde{\mathfrak{g}}$ associated with ϑ is equal to $\mathfrak{h}\langle nm+1 \rangle_0$. In [P09, Thm 4.1(ii)], it is shown that $\mathbb{k}[\mathfrak{g}\langle nm+1 \rangle_0]^{\mathfrak{g}\langle nm+1 \rangle_0}$ is a polynomial algebra of Krull dimension $n \text{rk } \mathfrak{g} + \text{rk } \mathfrak{g}_0$, if \mathfrak{g}_0 contains a regular nilpotent element of \mathfrak{g} . Here we have no assumptions on \mathfrak{g}_0 , i.e., we show that [P09, Thm 4.1(ii)] holds for all \mathfrak{g} and ϑ .

Our general reference for semisimple Lie groups and algebras is [Lie3].

1. PRELIMINARIES ON POISSON BRACKETS AND POLYNOMIAL CONTRACTIONS

Let Q be a connected affine algebraic group with Lie algebra \mathfrak{q} . The symmetric algebra of \mathfrak{q} over \mathbb{k} is \mathbb{N}_0 -graded, i.e., $\mathcal{S}(\mathfrak{q}) = \bigoplus_{i \geq 0} \mathcal{S}^i(\mathfrak{q})$. It is identified with the algebra of polynomial functions on the dual space \mathfrak{q}^* .

1.1. The coadjoint representation. The group Q acts on \mathfrak{q}^* via the coadjoint representation and then $\text{ad}^* : \mathfrak{q} \rightarrow \text{GL}(\mathfrak{q}^*)$ is the *coadjoint representation* of \mathfrak{q} . The algebra of Q -invariant polynomial functions on \mathfrak{q}^* is denoted by $\mathcal{S}(\mathfrak{q})^Q$ or $\mathbb{k}[\mathfrak{q}^*]^Q$. Write $\mathbb{k}(\mathfrak{q}^*)^Q$ for the field of Q -invariant rational functions on \mathfrak{q}^* .

Let $\mathfrak{q}^\xi = \{x \in \mathfrak{q} \mid \text{ad}^*(x) \cdot \xi = 0\}$ be the *stabiliser* in \mathfrak{q} of $\xi \in \mathfrak{q}^*$. The *index* of \mathfrak{q} ,

ind \mathfrak{q} , is the minimal codimension of Q -orbits in \mathfrak{q}^* . Equivalently, $\text{ind } \mathfrak{q} = \min_{\xi \in \mathfrak{q}^*} \dim \mathfrak{q}^\xi$. By the Rosenlicht theorem (see [Sp89, IV.2]), one also has $\text{ind } \mathfrak{q} = \text{tr.deg } \mathbb{k}(\mathfrak{q}^*)^Q$. Set $\mathbf{b}(\mathfrak{q}) = (\dim \mathfrak{q} + \text{ind } \mathfrak{q})/2$. Since the Q -orbits in \mathfrak{q}^* are even-dimensional, $\mathbf{b}(\mathfrak{q})$ is an integer. If \mathfrak{q} is reductive, then $\text{ind } \mathfrak{q} = \text{rk } \mathfrak{q}$ and $\mathbf{b}(\mathfrak{q})$ equals the dimension of a Borel subalgebra.

The Lie–Poisson bracket on $\mathcal{S}(\mathfrak{q})$ is defined on $\mathcal{S}^1(\mathfrak{q}) = \mathfrak{q}$ by $\{x, y\} := [x, y]$. It is then extended to higher degrees via the Leibniz rule. Hence $\mathcal{S}(\mathfrak{q})$ has the usual associative-commutative structure and additional Poisson structure. Whenever we refer to *subalgebras* of $\mathcal{S}(\mathfrak{q})$, we always mean the associative-commutative structure. Then a subalgebra $\mathcal{A} \subset \mathcal{S}(\mathfrak{q})$ is said to be *Poisson-commutative*, if $\{H, F\} = 0$ for all $H, F \in \mathcal{A}$. It is well known that if \mathcal{A} is Poisson-commutative, then $\text{tr.deg } \mathcal{A} \leq \mathbf{b}(\mathfrak{q})$, see e.g. [Vi90, 0.2]. More generally, suppose that $\mathfrak{h} \subset \mathfrak{q}$ is a Lie subalgebra and $\mathcal{A} \subset \mathcal{S}(\mathfrak{q})^{\mathfrak{h}}$ is Poisson-commutative. Then $\text{tr.deg } \mathcal{A} \leq \mathbf{b}(\mathfrak{q}) - \mathbf{b}(\mathfrak{h}) + \text{ind } \mathfrak{h}$, see [MY19, Prop. 1.1].

The *centre* of the Poisson algebra $(\mathcal{S}(\mathfrak{q}), \{, \})$ is

$$\mathcal{Z}(\mathfrak{q}) := \{H \in \mathcal{S}(\mathfrak{q}) \mid \{H, F\} = 0 \ \forall F \in \mathcal{S}(\mathfrak{q})\} = \mathcal{S}(\mathfrak{q})^{\mathfrak{q}} = \mathbb{k}[\mathfrak{q}^*]^{\mathfrak{q}} = \mathbb{k}[\mathfrak{q}^*]^Q.$$

Since the quotient field of $\mathbb{k}[\mathfrak{q}^*]^Q$ is contained in $\mathbb{k}(\mathfrak{q}^*)^Q$, we deduce from the Rosenlicht theorem that

$$(1.1) \quad \text{tr.deg } (\mathcal{S}(\mathfrak{q})^{\mathfrak{q}}) \leq \text{ind } \mathfrak{q}.$$

The set of *regular* elements of \mathfrak{q}^* is

$$(1.2) \quad \mathfrak{q}_{\text{reg}}^* = \{\eta \in \mathfrak{q}^* \mid \dim \mathfrak{q}^\eta = \text{ind } \mathfrak{q}\} = \{\eta \in \mathfrak{q}^* \mid \dim Q \cdot \eta \text{ is maximal}\}.$$

It is a dense open subset of \mathfrak{q}^* . Set $\mathfrak{q}_{\text{sing}}^* = \mathfrak{q}^* \setminus \mathfrak{q}_{\text{reg}}^*$. We say that \mathfrak{q} has the *codim– n* property if $\text{codim } \mathfrak{q}_{\text{sing}}^* \geq n$. The *codim–2* property is going to be most important for us.

For $\gamma \in \mathfrak{q}^*$, let $\hat{\gamma}$ be the skew-symmetric bilinear form on \mathfrak{q} defined by $\hat{\gamma}(\xi, \eta) = \gamma([\xi, \eta])$ for $\xi, \eta \in \mathfrak{q}$. It follows that $\ker \hat{\gamma} = \mathfrak{q}^\gamma$. The 2-form $\hat{\gamma}$ is related to the *Poisson tensor (bivector)* π of the Lie–Poisson bracket $\{, \}$ as follows.

Let dH denote the differential of $H \in \mathcal{S}(\mathfrak{q}) = \mathbb{k}[\mathfrak{q}^*]$. Then π is defined by the formula $\pi(dH \wedge dF) = \{H, F\}$ for $H, F \in \mathcal{S}(\mathfrak{q})$. Then $\pi(\gamma)(d_\gamma H \wedge d_\gamma F) = \{H, F\}(\gamma)$ and therefore $\hat{\gamma} = \pi(\gamma)$. In this terms, $\text{ind } \mathfrak{q} = \dim \mathfrak{q} - \text{rk } \pi$, where $\text{rk } \pi = \max_{\gamma \in \mathfrak{q}^*} \text{rk } \pi(\gamma)$.

For a subalgebra $A \subset \mathcal{S}(\mathfrak{q})$ and $\gamma \in \mathfrak{q}^*$, set $d_\gamma A = \langle d_\gamma F \mid F \in A \rangle_{\mathbb{k}}$. By the definition of $\mathcal{S}(\mathfrak{q})^{\mathfrak{q}}$, we have

$$(1.3) \quad d_\gamma \mathcal{S}(\mathfrak{q})^{\mathfrak{q}} \subset \ker \pi(\gamma)$$

for each $\gamma \in \mathfrak{q}^*$.

1.2. Contractions and invariants. We refer to [Lie3, Ch. 7, §2] for basic facts on contractions of Lie algebras. In this article, we consider contractions of the following form. Let $\mathbb{k}^* = \mathbb{k} \setminus \{0\}$ be the multiplicative group of \mathbb{k} and $\varphi : \mathbb{k}^* \rightarrow \text{GL}(\mathfrak{q})$, $s \mapsto \varphi_s$, a polynomial representation. That is, the matrix entries of $\varphi_s : \mathfrak{q} \rightarrow \mathfrak{q}$ are polynomials in s w.r.t. some

(any) basis of \mathfrak{q} . Define a new Lie algebra structure on the vector space \mathfrak{q} and associated Lie–Poisson bracket by

$$(1.4) \quad [x, y]_{(s)} = \{x, y\}_{(s)} := \varphi_s^{-1}[\varphi_s(x), \varphi_s(y)], \quad x, y \in \mathfrak{q}, \quad s \in \mathbb{k}^*.$$

All the algebras $(\mathfrak{q}, [,]_{(s)})$ are isomorphic and $(\mathfrak{q}, [,]_{(1)})$ is the initial Lie algebra \mathfrak{q} . The induced \mathbb{k}^* -action on the variety of structure constants is not necessarily polynomial, i.e., $\lim_{s \rightarrow 0} [x, y]_{(s)}$ may not exist for all $x, y \in \mathfrak{q}$. Whenever such a limit exists, we obtain a new linear Poisson bracket, denoted $\{ , \}_0$, and thereby a new Lie algebra $\mathfrak{q}_{(0)}$, which is said to be a *contraction* of \mathfrak{q} . If we wish to stress that this construction is determined by φ , then we write $\{x, y\}_{(\varphi, s)}$ for the bracket in (1.4) and say that $\mathfrak{q}_{(0)} = \mathfrak{q}_{(0, \varphi)}$ is the φ -contraction of \mathfrak{q} or is the *zero limit of \mathfrak{q} w.r.t. φ* . A criterion for the existence of $\mathfrak{q}_{(0)}$ can be given in terms of Lie brackets of the φ -eigenspaces in \mathfrak{q} , see [Y17, Sect. 4]. We identify all algebras $\mathfrak{q}_{(s)}$ and $\mathfrak{q}_{(0)}$ as vector spaces. The semi-continuity of index implies that $\text{ind } \mathfrak{q}_{(0)} \geq \text{ind } \mathfrak{q}$.

The map φ_s , $s \in \mathbb{k}^*$, is naturally extended to an invertible transformation of $\mathcal{S}^j(\mathfrak{q})$, which we also denote by φ_s . The resulting graded map $\varphi_s : \mathcal{S}(\mathfrak{q}) \rightarrow \mathcal{S}(\mathfrak{q})$ is nothing but the comorphism associated with $s \in \mathbb{k}^*$ and the dual representation $\varphi^* : \mathbb{k}^* \rightarrow \text{GL}(\mathfrak{q}^*)$. Since $\mathcal{S}^j(\mathfrak{q})$ has a basis that consists of $\varphi(\mathbb{k}^*)$ -eigenvectors, any $F \in \mathcal{S}^j(\mathfrak{q})$ can be written as $F = \sum_{i \geq 0} F_i$, where the sum is finite and $\varphi_s(F_i) = s^i F_i \in \mathcal{S}^j(\mathfrak{q})$. Let F^\bullet denote the non-zero component F_i with maximal i .

Proposition 1.1 ([Y14, Lemma 3.3]). *If $F \in \mathcal{Z}(\mathfrak{q})$ and $\mathfrak{q}_{(0)}$ exists, then $F^\bullet \in \mathcal{Z}(\mathfrak{q}_{(0)})$.*

1.3. Periodic gradings of Lie algebras and related compatible brackets. Let $\vartheta \in \text{Aut}(\mathfrak{q})$ be a Lie algebra automorphism of finite order $m \geq 2$ and $\zeta = \sqrt[m]{1}$ a primitive root of unity. Write also $\text{ord}(\vartheta)$ for the order of ϑ . If \mathfrak{q}_i is the ζ^i -eigenspace of ϑ , $i \in \mathbb{Z}_m$, then the direct sum $\mathfrak{q} = \bigoplus_{i \in \mathbb{Z}_m} \mathfrak{q}_i$ is a *periodic grading* or \mathbb{Z}_m -grading of \mathfrak{q} . The latter means that $[\mathfrak{q}_i, \mathfrak{q}_j] \subset \mathfrak{q}_{i+j}$ for all $i, j \in \mathbb{Z}_m$. Here $\mathfrak{q}_0 = \mathfrak{q}^\vartheta$ is the fixed-point subalgebra for ϑ and each \mathfrak{q}_i is a \mathfrak{q}_0 -module.

We choose $\{0, 1, \dots, m-1\} \subset \mathbb{Z}$ as a fixed set of representatives for $\mathbb{Z}_m = \mathbb{Z}/m\mathbb{Z}$. Under this convention, we have $\mathfrak{q} = \mathfrak{q}_0 \oplus \mathfrak{q}_1 \oplus \dots \oplus \mathfrak{q}_{m-1}$ and

$$(1.5) \quad [\mathfrak{q}_i, \mathfrak{q}_j] \subset \begin{cases} \mathfrak{q}_{i+j}, & \text{if } i+j \leq m-1, \\ \mathfrak{q}_{i+j-m}, & \text{if } i+j \geq m. \end{cases}$$

This is needed below, when we consider \mathbb{Z} -graded contractions of \mathfrak{q} associated with ϑ .

The presence of ϑ allows us to split the Lie–Poisson bracket on \mathfrak{q}^* into a sum of two compatible Poisson brackets. Consider the polynomial representation $\varphi : \mathbb{k}^* \rightarrow \text{GL}(\mathfrak{q})$ such that $\varphi_s(x) = s^j x$ for $x \in \mathfrak{q}_j$. As in Section 1.2, this defines a family of linear Poisson brackets on $\mathcal{S}(\mathfrak{q})$ parametrised by $s \in \mathbb{k}^*$, see (1.4).

Below we outline some results of [PY21]:

- (i) There is a limit $\lim_{s \rightarrow 0} \{x, y\}_{(s)} =: \{x, y\}_0$, which is a linear Poisson bracket on $\mathcal{S}(\mathfrak{q})$.
- (ii) The difference $\{ , \} - \{ , \}_0 =: \{ , \}_\infty$ is a linear Poisson bracket on $\mathcal{S}(\mathfrak{q})$, which is obtained as the zero limit w.r.t. the polynomial representation $\psi: \mathbb{k}^* \rightarrow \mathrm{GL}(\mathfrak{q})$ such that $\psi_s = s^m \cdot \varphi_{s^{-1}} = s^m \cdot \varphi_s^{-1}$, $s \in \mathbb{k}^*$. In other words, $\{ , \}_\infty = \lim_{s \rightarrow 0} \{ , \}_{(\psi, s)}$.
- (iii) For any $\vartheta \in \mathrm{Aut}(\mathfrak{q})$ of finite order, the Poisson brackets $\{ , \}_0$ and $\{ , \}_\infty$ are compatible, and the corresponding pencil contains the initial Lie–Poisson bracket.

Set

$$\{ , \}_t = \{ , \}_0 + t \{ , \}_\infty,$$

where $t \in \mathbb{P} := \mathbb{k} \cup \{\infty\}$ and the value $t = \infty$ corresponds to the bracket $\{ , \}_\infty$. Let $\mathfrak{q}_{(t)}$ stand for the Lie algebra corresponding to $\{ , \}_t$. All these Lie algebras have the same underlying vector space.

Proposition 1.2 ([PY21, Prop. 2.3]). *The Lie algebras $\mathfrak{q}_{(0)}$ and $\mathfrak{q}_{(\infty)}$ are \mathbb{N}_0 -graded. More precisely, if $\mathfrak{r}[i]$ stands for the component of grade $i \in \mathbb{N}_0$ in an \mathbb{N}_0 -graded Lie algebra \mathfrak{r} , then*

$$\mathfrak{q}_{(0)}[i] = \begin{cases} \mathfrak{q}_i & \text{for } i = 0, 1, \dots, m-1 \\ 0 & \text{otherwise} \end{cases}, \quad \mathfrak{q}_{(\infty)}[i] = \begin{cases} \mathfrak{q}_{m-i} & \text{for } i = 1, 2, \dots, m \\ 0 & \text{otherwise} \end{cases}.$$

In particular, $\mathfrak{q}_{(\infty)}$ is nilpotent and the subspace \mathfrak{q}_0 , which is the highest grade component of $\mathfrak{q}_{(\infty)}$, belongs to the centre of $\mathfrak{q}_{(\infty)}$. \square

Suppose that $t \in \mathbb{k}^*$. Then $\{ , \}_t = \{ , \}_{(s)}$, where $s^m = t$. The algebras $\mathfrak{q}_{(t)}$ with $t \in \mathbb{k}^*$ are isomorphic and they have one and the same index. We say that $t \in \mathbb{P}$ is *regular* if $\mathrm{ind} \mathfrak{q}_{(t)} = \mathrm{ind} \mathfrak{q}$ and write $\mathbb{P}_{\mathrm{reg}}$ for the set of regular values. Then $\mathbb{P}_{\mathrm{sing}} := \mathbb{P} \setminus \mathbb{P}_{\mathrm{reg}} \subset \{0, \infty\}$ is the set of singular values.

Let $Q_0 \subset Q$ be the connected subgroup of Q with $\mathrm{Lie} Q_0 = \mathfrak{q}_0$. It is easy to see that Q_0 is an algebraic group. Hence there are connected algebraic groups $Q_{(t)}$ such that $\mathfrak{q}_{(t)} = \mathrm{Lie} Q_{(t)}$ for each $t \in \mathbb{P}$.

Let \mathcal{Z}_t be the centre of the Poisson algebra $(\mathcal{S}(\mathfrak{q}), \{ , \}_t)$. In particular, $\mathcal{Z}_1 = \mathcal{S}(\mathfrak{q})^{\mathfrak{q}}$. If $t = s^m \in \mathbb{k}^*$, then $\mathcal{Z}_t = \varphi_s^{-1}(\mathcal{Z}_1)$. By Eq. (1.1), we have $\mathrm{tr.deg} \mathcal{Z}_t \leq \mathrm{ind} \mathfrak{q}_{(t)}$. In [PY21], we have studied the subalgebra $\mathcal{Z} \subset \mathcal{S}(\mathfrak{q})$ generated by the centres \mathcal{Z}_t with $t \in \mathbb{P}_{\mathrm{reg}}$, i.e.,

$$\mathcal{Z} = \mathcal{Z}(\mathfrak{q}, \vartheta) = \mathrm{alg} \langle \mathcal{Z}_t \mid t \in \mathbb{P}_{\mathrm{reg}} \rangle.$$

By a general property of compatible brackets, the algebra \mathcal{Z} is Poisson-commutative w.r.t. **all** brackets $\{ , \}_t$ with $t \in \mathbb{P}$, cf. [PY, Sect. 2]. Note that the Lie subalgebra $\mathfrak{q}_0 \subset \mathfrak{q} = \mathfrak{q}_{(1)}$ is also the **same** Lie subalgebra in any $\mathfrak{q}_{(t)}$ with $t \neq \infty$ (cf. Proposition 1.2 for $\mathfrak{q}_{(0)}$). Therefore,

$$(1.6) \quad \mathcal{Z}_t \subset \mathcal{S}(\mathfrak{q})^{\mathfrak{q}_0} \text{ for } t \neq \infty.$$

Convention. We think of \mathfrak{q}^* as the dual space for any Lie algebra $\mathfrak{q}_{(t)}$ and sometimes omit the subscript $'(t)'$ in $\mathfrak{q}_{(t)}^*$. However, if $\xi \in \mathfrak{q}^*$, then the stabiliser of ξ with respect to the coadjoint representation of $\mathfrak{q}_{(t)}$ is denoted by $\mathfrak{q}_{(t)}^{\xi}$. Set $\mathfrak{q}_{\infty, \mathrm{reg}}^* := (\mathfrak{q}_{(\infty)}^*)_{\mathrm{reg}}$.

1.4. Good generating systems. Let \mathfrak{g} be a reductive Lie algebra. Consider a contraction $\mathfrak{g}_{(0)}$ of \mathfrak{g} given by $\{\varphi_s \mid s \in \mathbb{k}^\times\}$. Let $\{x_1, \dots, x_{\dim \mathfrak{g}}\}$ be basis of \mathfrak{g} and $\omega = x_1 \wedge \dots \wedge x_{\dim \mathfrak{g}}$ a volume form. Then $\varphi(\omega) = s^D \omega$, where D is a non-negative integer. We set $D_\varphi := D$.

The Poisson centre $\mathcal{S}(\mathfrak{g})^\mathfrak{g}$ is a polynomial algebra of Krull dimension $l = \text{rk } \mathfrak{g}$ and $\text{ind } \mathfrak{g} = l$. Hence one has now the equality in Eq. (1.1). Let $\{H_1, \dots, H_l\}$ be a set of homogeneous algebraically independent generators of $\mathcal{S}(\mathfrak{g})^\mathfrak{g}$ and $d_i = \deg H_i$. Then $\sum_{i=1}^l d_i = \mathbf{b}(\mathfrak{g})$. As above, each H_j decomposes as $H_j = \sum_{i \geq 0} H_{j,i}$, where $\varphi_s(H_j) = \sum_{i \geq 0} s^i H_{j,i}$. The polynomials $H_{j,i}$ are called bi-homogeneous components of H_j . By definition, the φ -degree of $H_{j,i}$ is i , also denoted by $\deg_\varphi H_{j,i}$. Then H_j^\bullet is the non-zero bi-homogeneous component of H_j with maximal φ -degree. We set $\deg_{\varphi} H_j = \deg_{\varphi} H_j^\bullet$ and $d_j^\bullet = \deg_{\varphi} H_j^\bullet$.

Definition 1.3. Let us say that H_1, \dots, H_l is a *good generating system* in $\mathcal{S}(\mathfrak{g})^\mathfrak{g}$ (g.g.s. for short) for φ , if $H_1^\bullet, \dots, H_l^\bullet$ are algebraically independent. Then we also say that φ admits a g.g.s.

The property of being ‘good’ really depends on a generating system. The importance of g.g.s. is manifestly seen in the following result.

Theorem 1.4 ([Y14, Theorem 3.8]). *Let H_1, \dots, H_l be an arbitrary set of homogeneous algebraically independent generators of $\mathcal{S}(\mathfrak{g})^\mathfrak{g}$. Then*

- (i) $\sum_{j=1}^l \deg_\varphi H_j \geq D_\varphi$;
- (ii) H_1, \dots, H_l is a g.g.s. if and only if $\sum_{j=1}^l \deg_\varphi H_j = D_\varphi$;
- (iii) if $\mathfrak{g}_{(0)}$ has the *codim–2 property*, $\text{ind } \mathfrak{g}_{(0)} = l$, and H_1, \dots, H_l is a g.g.s., then $\mathcal{Z}_0 = \mathcal{S}(\mathfrak{g}_{(0)})^{\mathfrak{g}_{(0)}}$ is a polynomial algebra freely generated by $H_1^\bullet, \dots, H_l^\bullet$ and

$$\{\xi \in \mathfrak{g}^* \mid d_\xi H_1^\bullet \wedge \dots \wedge d_\xi H_l^\bullet = 0\} = (\mathfrak{g}_{(0)})_{\text{sing}}^*. \quad \square$$

Let $F_1, \dots, F_N \in \mathbb{k}[x_i \mid 1 \leq i \leq n] = \mathbb{k}[\mathbb{A}^n]$ be algebraically independent homogeneous polynomials. Set

$$\mathcal{J}(F_1, \dots, F_N) := \{x \in \mathbb{A}^n \mid d_x F_1 \wedge \dots \wedge d_x F_N = 0\}.$$

Then $\mathcal{J}(F_1, \dots, F_N)$ is a proper closed subset of \mathbb{A}^n . An open subset of \mathbb{A}^n is said to be *big*, if its complement does not contain divisors. Thereby the differentials dF_i with $1 \leq i \leq N$ are linearly independent on a big open subset if and only if $\dim \mathcal{J}(F_1, \dots, F_N) \leq n - 2$.

By the *Kostant regularity criterion* for \mathfrak{g} , $\mathcal{J}(H_1, \dots, H_l) = \mathfrak{g}_{\text{sing}}^*$, see [K63, Theorem 9]. By [K63], \mathfrak{g} has the *codim–3 property*, i.e., $\dim \mathfrak{g}_{\text{sing}}^* \leq \dim \mathfrak{g} - 3$.

Part (iii) of Theorem 1.4 states that the Kostant regularity criterion holds for $\mathfrak{g}_{(0)}$. One of the ingredients of the proof, which will be used in this paper as well, is the following statement.

Theorem 1.5 ([PPY, Theorem 1.1]). *Let F_1, \dots, F_N be as above. If $\dim \mathcal{J}(F_1, \dots, F_N) \leq n - 2$, then $\mathcal{F} = \mathbb{k}[F_j \mid 1 \leq j \leq N]$ is an algebraically closed subalgebra of $\mathbb{k}[\mathbb{A}^n]$, i.e., if $H \in \mathbb{k}[\mathbb{A}^n]$ is algebraic over the field $\mathbb{k}(F_1, \dots, F_N)$, then $H \in \mathcal{F}$.*

2. PROPERTIES OF $\mathfrak{q}_{(\infty)}$ AND OF THE POISSON CENTRE $\mathcal{Z}_{\infty} \subset \mathcal{S}(\mathfrak{q}_{(\infty)})$

By Proposition 1.2, $\mathfrak{q}_{(\infty)}$ is a nilpotent \mathbb{N} -graded Lie algebra. Recall also that the subspace \mathfrak{q}_0 belongs to the centre of $\mathfrak{q}_{(\infty)}$. Let π_t be the Poisson tensor of $\mathfrak{q}_{(t)}$. We identify \mathfrak{q}_0^* with the annihilator $\text{Ann}(\bigoplus_{i=1}^{m-1} \mathfrak{q}_i) \subset \mathfrak{q}^*$ and regard it as a subspace of \mathfrak{q}^* .

For any $\xi \in \mathfrak{q}_0^* \subset \mathfrak{q}^*$, we have

$$(2.1) \quad \mathfrak{q}_{(\infty)}^{\xi} = \mathfrak{q}_0 \oplus \bigoplus_{j \geq 1} \mathfrak{q}_j^{\xi}, \quad \text{where } \mathfrak{q}_j^{\xi} = \{y \in \mathfrak{q}_j \mid \xi([y, \mathfrak{q}_{m-j}]) = 0\}.$$

Furthermore, $\mathfrak{q}^{\xi} = \mathfrak{q}_0^{\xi} \oplus \bigoplus_{j \geq 1} \mathfrak{q}_j^{\xi}$. If this ξ is regular in \mathfrak{q}^* , then $\xi \in (\mathfrak{q}_0^*)_{\text{reg}}$.

Theorem 2.1. *Suppose that $\mathfrak{q}_0^* \cap \mathfrak{q}_{\text{reg}}^* \neq \emptyset$. Then $\text{ind } \mathfrak{q}_{(\infty)} = \dim \mathfrak{q}_0 + \text{ind } \mathfrak{q} - \text{ind } \mathfrak{q}_0$.*

Proof. (1) Take any $\xi \in \mathfrak{q}_0^* \cap \mathfrak{q}_{\text{reg}}^*$. Then $\mathfrak{q}_{(\infty)}^{\xi} = \mathfrak{q}^{\xi} + \mathfrak{q}_0$. Furthermore $\mathfrak{q}^{\xi} \cap \mathfrak{q}_0 = \mathfrak{q}_0^{\xi}$. As we have explained above, $\xi \in (\mathfrak{q}_0^*)_{\text{reg}}$. Thereby $\dim \mathfrak{q}_0^{\xi} = \text{ind } \mathfrak{q}_0$ and hence $\dim \mathfrak{q}_{(\infty)}^{\xi} = \text{ind } \mathfrak{q} + \dim \mathfrak{q}_0 - \text{ind } \mathfrak{q}_0$. This leads to $\text{ind } \mathfrak{q}_{(\infty)} \leq \text{ind } \mathfrak{q} + \dim \mathfrak{q}_0 - \text{ind } \mathfrak{q}_0$.

(2) Let us prove the opposite inequality. Take any $\xi \in \mathfrak{q}_{\text{reg}}^*$ such that $\bar{\xi} = \xi|_{\mathfrak{q}_0} \in (\mathfrak{q}_0^*)_{\text{reg}}$. Note that there is a non-empty open subset consisting of suitable elements. For all but finitely many t , we have $\dim \mathfrak{q}_{(t)}^{\xi} = \text{ind } \mathfrak{q}$. Hence $\mathfrak{v} = \lim_{t \rightarrow \infty} \mathfrak{q}_{(t)}^{\xi}$ is a well-defined subspace of \mathfrak{q} of dimension $\text{ind } \mathfrak{q}$. If $t \neq \infty$, then $\pi_t(\xi)|_{\mathfrak{q}_0 \times \mathfrak{q}} = \pi_1(\xi)|_{\mathfrak{q}_0 \times \mathfrak{q}}$ and $\pi_1(\xi)(\mathfrak{q}_0, \mathfrak{q}_{(t)}^{\xi}) = \pi_t(\xi)(\mathfrak{q}_0, \mathfrak{q}_{(t)}^{\xi}) = 0$. Therefore

$$(2.2) \quad \pi_1(\xi)(\mathfrak{q}_0, \mathfrak{v}) = 0$$

and $\mathfrak{v} \cap \mathfrak{q}_0 \subset \mathfrak{q}_0^{\bar{\xi}}$. By the construction, $\mathfrak{v} \subset \mathfrak{q}_{(\infty)}^{\xi}$. Thus

$$(2.3) \quad \dim \mathfrak{q}_{(\infty)}^{\xi} \geq \dim(\mathfrak{v} + \mathfrak{q}_0) = \dim \mathfrak{v} + \dim \mathfrak{q}_0 - \dim(\mathfrak{q}_0 \cap \mathfrak{v}) \geq \text{ind } \mathfrak{q} + \dim \mathfrak{q}_0 - \text{ind } \mathfrak{q}_0.$$

This finishes the proof, since the inequality holds on a non-empty open subset. \square

The assumption $\mathfrak{q}_0^* \cap \mathfrak{q}_{\text{reg}}^* \neq \emptyset$ is satisfied in the reductive case, since the reductive subalgebra $\mathfrak{g}_0 = \mathfrak{g}^{\vartheta}$ contains regular semisimple elements of \mathfrak{g} , see e.g. [Ka83, §8.8]. Thus, we obtain a new proof of [PY21, Theorem 3.2].

Corollary 2.2 ([PY21]). *Suppose that $\mathfrak{q} = \mathfrak{g}$ is reductive. Then one has $\infty \in \mathbb{P}_{\text{reg}}$ if and only if $\dim \mathfrak{g}_0 = \text{rk } \mathfrak{g}_0$, i.e., \mathfrak{g}_0 is an abelian subalgebra of \mathfrak{g} .*

Recall that $\mathfrak{q}_0 \subset \mathcal{Z}_{\infty}$. Thereby \mathcal{Z}_{∞} is not Poisson-commutative, unless \mathfrak{q}_0 is commutative. However, if $[\mathfrak{q}_0, \mathfrak{q}_0] = 0$ and $\mathfrak{q}_0^* \cap \mathfrak{q}_{\text{reg}}^* \neq \emptyset$, then $\text{ind } \mathfrak{q}_0 = \dim \mathfrak{q}_0$ and $\{ , \}_{\infty}$ is a regular structure in the pencil spanned by $\{ , \}$ and $\{ , \}_0$, see Theorem 2.1. In this case, $\mathcal{Z}_{\infty} \subset \mathcal{Z}(\mathfrak{q}, \vartheta)$. Thereby a description of \mathcal{Z}_{∞} is desirable.

The group Q_0 acts on $\mathfrak{q}_{(\infty)}$. Hence one may consider $\mathcal{Z}_{\infty}^{\mathfrak{q}_0} \subset \mathcal{Z}_{\infty}$.

Theorem 2.3. *Assume that $\mathfrak{q}_0^* \cap \mathfrak{q}_{\text{reg}}^* \neq \emptyset$.*

(i) *We have $\{\mathcal{Z}_{\infty}^{\mathfrak{q}_0}, \mathcal{Z}_{\infty}^{\mathfrak{q}_0}\} = 0$.*

(ii) *The algebra $\text{alg}\langle \mathcal{Z}, \mathcal{Z}_\infty^{\mathfrak{q}_0} \rangle$ is still Poisson-commutative.*

(iii) *We have also $\text{tr.deg } \mathcal{Z}_\infty^{\mathfrak{q}_0} \leq \text{ind } \mathfrak{q}$.*

Proof. Take $\xi \in \mathfrak{q}_{\text{reg}}^* \cap \mathfrak{q}_{\infty, \text{reg}}^*$ such that $\bar{\xi} = \xi|_{\mathfrak{q}_0} \in (\mathfrak{q}_0^*)_{\text{reg}}$. Note that there is a non-empty open subset consisting of suitable elements. Following the proof of Theorem 2.1, set $\mathfrak{v} = \lim_{t \rightarrow \infty} \mathfrak{q}_{(t)}^\xi$. Recall that $\mathfrak{v} + \mathfrak{q}_0 \subset \mathfrak{q}_{(\infty)}^\xi$ and that $\dim(\mathfrak{v} + \mathfrak{q}_0) \geq \text{ind } \mathfrak{q}_{(\infty)}$, see (2.3). Since $\xi \in \mathfrak{q}_{\infty, \text{reg}}^*$, there is the equality $\mathfrak{q}_{(\infty)}^\xi = \mathfrak{v} + \mathfrak{q}_0$.

Next $d_\xi \mathcal{Z}_\infty^{\mathfrak{q}_0} \subset \mathfrak{q}_{(\infty)}^\xi$ by (1.3) and $\pi_1(\xi)(d_\xi \mathcal{Z}_\infty^{\mathfrak{q}_0}, \mathfrak{q}_0) = 0$, since $\mathcal{Z}_\infty^{\mathfrak{q}_0}$ consists of \mathfrak{q}_0 -invariants. In the proof of Theorem 2.1, we have established that $\pi_1(\xi)(\mathfrak{v}, \mathfrak{q}_0) = 0$, see (2.2). Suppose that $y \in \mathfrak{q}_0$ and $\pi_1(\xi)(y, \mathfrak{q}_0) = 0$. Then $y \in \mathfrak{q}_0^{\bar{\xi}}$. In particular, $d_\xi \mathcal{Z}_\infty^{\mathfrak{q}_0} \subset \mathfrak{v} + \mathfrak{q}_0^{\bar{\xi}}$ and $\mathfrak{v} \cap \mathfrak{q}_0 \subset \mathfrak{q}_0^{\bar{\xi}}$. By the dimension reasons, $\mathfrak{v} \cap \mathfrak{q}_0 = \mathfrak{q}_0^{\bar{\xi}}$. Thus $d_\xi \mathcal{Z}_\infty^{\mathfrak{q}_0} \subset \mathfrak{v} + \mathfrak{q}_0^{\bar{\xi}} \subset \mathfrak{v}$. The inclusion proves the inequality $\text{tr.deg } \mathcal{Z}_\infty^{\mathfrak{q}_0} \leq \dim \mathfrak{v} = \text{ind } \mathfrak{q}$.

For almost all $t \in \mathbb{P}$, we have $\dim \mathfrak{q}_{(t)}^\xi = \text{ind } \mathfrak{q}$. Hence \mathfrak{v} is a subspace of

$$L(\xi) := \sum_{t: \text{rk } \pi_t(\xi) = \text{dim } \mathfrak{q} - \text{ind } \mathfrak{q}} \mathfrak{q}_{(t)}^\xi$$

and the latter is known to be isotropic w.r.t. $\pi_t(\xi)$ for any t , see e.g. [PY08, Appendix]. Then, in particular, any $F \in \{\mathcal{Z}_\infty^{\mathfrak{q}_0}, \mathcal{Z}_\infty^{\mathfrak{q}_0}\}$ vanishes at ξ , and, since ξ is generic, the first claim is settled.

If $\infty \in \mathbb{P}_{\text{reg}}$, then $\mathcal{Z}_\infty^{\mathfrak{q}_0} \subset \mathcal{Z}_\infty \subset \mathcal{Z}$ and part (ii) is clear. Suppose that $\infty \in \mathbb{P}_{\text{sing}}$. Then $\mathcal{Z} \subset \text{alg}\langle \mathcal{Z}_t \mid t \neq \infty \rangle$. For any $t \neq \infty$, the brackets $\{ , \}_\infty$ and $\{ , \}_t$ span

$$\mathcal{P} = \{a\{ , \}_0 + b\{ , \}_\infty \mid (a, b) \in \mathbb{k}^2\}$$

and $\{\mathcal{Z}_\infty, \mathcal{Z}_t\}_\infty = \{\mathcal{Z}_\infty, \mathcal{Z}_t\}_t = 0$. Thereby $\{\mathcal{Z}_\infty, \mathcal{Z}_t\} = 0$ for each $t \neq \infty$. This finishes the proof. \square

3. THE REDUCTIVE CASE

In most of this section, we recollect known results about automorphisms of reductive Lie algebras and properties of $\mathcal{Z}(\mathfrak{g}, \vartheta)$. Statements of Section 3.2 are crucial for the proof of Theorem 4.4. Theorem 3.10 on the index of $\mathfrak{g}_{(0)}$ is a new result.

3.1. The Kac diagram of a finite order automorphism. We describe briefly Kac's classification of finite order automorphisms in the semisimple case [Ka69].

A pair $(\mathfrak{g}, \vartheta)$ is *decomposable*, if \mathfrak{g} is a direct sum of two non-trivial ϑ -stable ideals. Otherwise $(\mathfrak{g}, \vartheta)$ is said to be *indecomposable*. Classification of finite order automorphism readily reduces to the indecomposable case. The centre of \mathfrak{g} is always a ϑ -stable ideal and automorphisms of an abelian Lie algebra have no particular significance (in our context). Therefore assume that \mathfrak{g} is semisimple.

Suppose that \mathfrak{g} is not simple and $(\mathfrak{g}, \vartheta)$ is indecomposable. Then $\mathfrak{g} = \mathfrak{h}^{\oplus n}$ is a sum of n copies of a simple Lie algebra \mathfrak{h} and ϑ is a composition of an automorphism of \mathfrak{h} and a cyclic permutation of the summands.

Below we assume that \mathfrak{g} is simple. By a result of R. Steinberg [St68, Theorem 7.5], every semisimple automorphism of \mathfrak{g} fixes a Borel subalgebra of \mathfrak{g} and a Cartan subalgebra thereof. Let \mathfrak{b} be a ϑ -stable Borel subalgebra and $\mathfrak{t} \subset \mathfrak{b}$ a ϑ -stable Cartan subalgebra. This yields a ϑ -stable triangular decomposition $\mathfrak{g} = \mathfrak{u}^- \oplus \mathfrak{t} \oplus \mathfrak{u}$, where $\mathfrak{u} = [\mathfrak{b}, \mathfrak{b}]$. Let $\Delta = \Delta(\mathfrak{g})$ be the set of roots of \mathfrak{g} related to \mathfrak{t} , Δ^+ the set of positive roots corresponding to \mathfrak{u} , and $\Pi \subset \Delta^+$ the set of simple roots. Let \mathfrak{g}_γ be the root space for $\gamma \in \Delta$. Hence $\mathfrak{u} = \bigoplus_{\gamma \in \Delta^+} \mathfrak{g}_\gamma$. Let $e_\gamma \in \mathfrak{g}_\gamma$ be a non-zero root vector.

Clearly, ϑ induces a permutation of Π , which is an automorphism of the Dynkin diagram, and ϑ is inner if and only if this permutation is trivial. Accordingly, ϑ can be written as a product $\sigma \circ \vartheta'$, where ϑ' is inner and σ is the so-called *diagram automorphism* of \mathfrak{g} . We refer to [Ka83, §8.2] for an explicit construction and properties of σ . In particular, σ depends only on the connected component of $\text{Aut}(\mathfrak{g})$ that contains ϑ and $\text{ord}(\sigma)$ equals the order of the corresponding permutation of Π .

— The case of an inner ϑ :

Set $\Pi = \{\alpha_1, \dots, \alpha_l\}$ and let $\delta = \sum_{i=1}^l n_i \alpha_i$ be the highest root in Δ^+ . An inner periodic automorphism with $\mathfrak{t} \subset \mathfrak{g}_0$ is determined by an $(l+1)$ -tuple of non-negative integers, *Kac labels*, $\mathbf{p} = (p_0, p_1, \dots, p_l)$ such that $\gcd(p_0, \dots, p_l) = 1$ and $\mathbf{p} \neq (0, \dots, 0)$. Set $m := p_0 + \sum_{i=1}^l n_i p_i$ and let \overline{p}_i denote the unique representative of $\{0, 1, \dots, m-1\}$ such that $p_i \equiv \overline{p}_i \pmod{m}$. The \mathbb{Z}_m -grading $\mathfrak{g} = \bigoplus_{i=0}^{m-1} \mathfrak{g}_i$ corresponding to $\vartheta = \vartheta(\mathbf{p})$ is defined by the conditions that

$$\mathfrak{g}_{\alpha_i} \subset \mathfrak{g}_{\overline{p}_i} \text{ for } i = 1, \dots, l, \quad \mathfrak{g}_{-\delta} \subset \mathfrak{g}_{\overline{p}_0}, \text{ and } \mathfrak{t} \subset \mathfrak{g}_0.$$

The *Kac diagram* $\mathcal{K}(\vartheta)$ of $\vartheta = \vartheta(\mathbf{p})$ is the **affine** (= extended) Dynkin diagram of \mathfrak{g} , $\tilde{\mathcal{D}}(\mathfrak{g})$, equipped with the labels p_0, p_1, \dots, p_l . In $\mathcal{K}(\vartheta)$, the i -th node of the usual Dynkin diagram $\mathcal{D}(\mathfrak{g})$ represents α_i and the extra node represents $-\delta$. It is convenient to assume that $\alpha_0 = -\delta$ and $n_0 = 1$. Then $(l+1)$ -tuple (n_0, n_1, \dots, n_l) yields the coefficients of a linear dependence for $\alpha_0, \alpha_1, \dots, \alpha_l$.

— The case of an outer ϑ :

Let σ be the diagram automorphism of \mathfrak{g} related to ϑ . The order of a nontrivial diagram automorphism is either 2 or 3, there 3 is possible only for \mathfrak{g} of type D_4 . Therefore, σ defines either a \mathbb{Z}_2 - or \mathbb{Z}_3 -grading of \mathfrak{g} , say $\mathfrak{g} = \mathfrak{g}^\sigma \oplus \mathfrak{g}_1^{(\sigma)}$ or $\mathfrak{g} = \mathfrak{g}^\sigma \oplus \mathfrak{g}_1^{(\sigma)} \oplus \mathfrak{g}_2^{(\sigma)}$.

The Kac diagrams of outer periodic automorphisms are supported on the twisted affine Dynkin diagrams of index 2 and 3, see [Vi76, §8] and [Lie3, Table 3]. Such a diagram has $r+1$ nodes, where $r = \text{rk } \mathfrak{g}^\sigma$, certain r nodes comprise the Dynkin diagram of the simple Lie algebra \mathfrak{g}^σ , and the additional node represents the lowest weight $-\delta_1$ of the

\mathfrak{g}^σ -module $\mathfrak{g}_1^{(\sigma)}$. Write $\delta_1 = \sum_{i=1}^r a'_i \nu_i$, where the elements ν_i are the simple roots of \mathfrak{g}^σ , and set $a'_0 = 1$. Then the $(r+1)$ -tuple $(a'_0, a'_1, \dots, a'_r)$ yields coefficients of linear dependence for $-\delta_1, \nu_1, \dots, \nu_r$.

Let $\mathbf{p} = (p_0, p_1, \dots, p_r)$ be an $(r+1)$ -tuple of non-negative integers, *Kac labels*, such that $\mathbf{p} \neq (0, 0, \dots, 0)$ and $\gcd(p_0, p_1, \dots, p_r) = 1$. The Kac diagram of $\vartheta = \vartheta(\mathbf{p})$ is the required twisted affine diagram equipped with the labels (p_0, p_1, \dots, p_r) over the nodes. Then $m = \text{ord}(\vartheta(\mathbf{p})) = \text{ord}(\sigma) \cdot \sum_{i=0}^r a'_i p_i$.

3.2. Relation between roots of \mathfrak{g} and \mathfrak{g}^σ . Let σ be a diagram automorphism of \mathfrak{g} associated with \mathfrak{b} and $\mathfrak{t} \subset \mathfrak{b}$. Then $\sigma(\mathfrak{b}) = \mathfrak{b}$ and $\sigma(\mathfrak{t}) = \mathfrak{t}$ by the construction. Let $\Delta_{\mathfrak{g}^\sigma}^+$ be the set of positive roots of \mathfrak{g}^σ associated with $(\mathfrak{b}^\sigma, \mathfrak{t}_0)$, where $\mathfrak{t}_0 = \mathfrak{t}^\sigma$.

Take any $\alpha \in \Delta^+$. If $\sigma(\alpha) \neq \alpha$, then the restriction $\bar{\alpha} = \alpha|_{\mathfrak{t}_0}$ is a positive root of \mathfrak{g}^σ . In any case, $\bar{\alpha} = \sigma(\alpha)|_{\mathfrak{t}_0}$. Suppose $\beta \in \Delta_{\mathfrak{g}^\sigma}$. Then there is a non-zero root vector $x_\beta \in \mathfrak{g}^\sigma$. We can write $x_\beta = \sum_{\alpha \in \Delta} b_\alpha e_\alpha$ with $b_\alpha \in \mathbb{k}$. Then $\alpha|_{\mathfrak{t}_0} = \beta$, whenever $b_\alpha \neq 0$. In particular, $\Delta_{\mathfrak{g}^\sigma}$ is contained in $\Delta|_{\mathfrak{t}_0}$.

Let $\langle \sigma \rangle \subset \text{Aut}(\mathfrak{g})$ be a subgroup generated by σ . In all types except A_{2n} , the restriction of roots from \mathfrak{t} to \mathfrak{t}_0 produces bijections between $\langle \sigma \rangle$ -orbits on Δ^+ and $\Delta_{\mathfrak{g}^\sigma}^+$. If $\mathfrak{g} = \mathfrak{sl}_{2n+1}$, then the situation is slightly different. These are well-known facts, nevertheless we give a brief explanation below. Set $\mathfrak{s}_\alpha = \mathfrak{g}_\alpha \oplus \mathfrak{g}_{-\alpha} \oplus [\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}]$.

Lemma 3.1 (cf. [Ka83, § 8.2]). *Suppose $\text{ord}(\sigma) = 3$. Then there is a bijection between $\langle \sigma \rangle$ -orbits on Δ^+ and $\Delta_{\mathfrak{g}^\sigma}^+$.*

Proof. Here \mathfrak{g} is of type D_4 . There are six $\langle \sigma \rangle$ -orbits on Δ^+ . Three of these orbits have three elements, namely

$$\{\alpha_1, \alpha_3, \alpha_4\}, \{\alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_2 + \alpha_4\}, \{\alpha_1 + \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_4, \alpha_2 + \alpha_3 + \alpha_4\}.$$

The other three consist of fixed points: $\{\alpha_2\}, \{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4\}, \{\alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4\}$. The Lie algebra \mathfrak{g}^σ is of type G_2 and it has 6 positive roots. Therefore there is nothing more to show. \square

Lemma 3.2. *Suppose $\text{ord}(\sigma) = 2$. Let $x_\mu = \sum_{\alpha \in \Delta} b_\alpha e_\alpha \in (\mathfrak{b} \cap \mathfrak{g}^\sigma)$ with $b_\alpha \in \mathbb{k}$ be a non-zero root vector of \mathfrak{g}^σ . Fix one $\alpha \in \Delta^+$ such that $b_\alpha \neq 0$. Suppose further that $\bar{\beta} = \beta|_{\mathfrak{t}_0} \in \mathbb{k}\bar{\alpha}$ for some $\beta \in \Delta^+$ that does not belong to the $\langle \sigma \rangle$ -orbit of α . Then $\beta = \alpha + \sigma(\alpha)$.*

Proof. Since $\sigma(x_\mu) = x_\mu$, we have $(\mathfrak{s}_\alpha + \sigma(\mathfrak{s}_\alpha))^\sigma \simeq \mathfrak{sl}_2$, and if $\sigma(\alpha) = \alpha$, then $\sigma|_{\mathfrak{g}_\alpha} = \text{id}$.

Let $h \in \mathfrak{t}_0$ be a subregular element of \mathfrak{g}^σ such that $\bar{\alpha}(h) = 0$ and $\gamma(h) \neq 0$, whenever $\gamma \in \Delta$ and $\gamma|_{\mathfrak{t}_0} \notin \mathbb{k}\bar{\alpha}$. Consider \mathfrak{g}^h . On the one hand, we have $(\mathfrak{g}^h)^\sigma = (\mathfrak{g}^\sigma)^h = \mathfrak{sl}_2 + \mathfrak{t}_0$. On the other hand, if $\sigma(\beta) \neq \beta$, then

$$(\mathfrak{g}^h)^\sigma \supset (\mathfrak{s}_\alpha + \mathfrak{s}_{\sigma(\alpha)})^\sigma \oplus (\mathfrak{s}_\beta + \mathfrak{s}_{\sigma(\beta)})^\sigma \simeq \mathfrak{sl}_2 \oplus \mathfrak{sl}_2,$$

which is a contradiction. This shows that $\sigma(\beta) = \beta$.

By the same reason as above, $\sigma|_{\mathfrak{g}_\beta} \neq \text{id}$. It follows that $\sigma|_{\mathfrak{g}_\beta} = \sigma|_{\mathfrak{g}_{-\beta}} = -\text{id}$, since $\text{ord}(\sigma) = 2$. Then $[\mathfrak{g}_\beta, \mathfrak{g}_{-\beta}] \subset \mathfrak{t}_0$. Let $h_\beta \in [\mathfrak{g}_\beta, \mathfrak{g}_{-\beta}]$ be such that $\beta(h_\beta) = 2$. Since $h_\beta \in \mathfrak{t}_0$ and $\bar{\beta} \in \mathbb{k}\bar{\alpha}$, we have $\alpha(h_\beta) \neq 0$. In view of this, $[\mathfrak{s}_\beta, \mathfrak{s}_\alpha] \neq 0$.

Set $\mathfrak{f} = [\mathfrak{g}^h, \mathfrak{g}^h]$. Then $\mathfrak{s}_\alpha + \mathfrak{s}_\beta \subset \mathfrak{f}$. Furthermore, if $\gamma \in \Delta$ is a root of \mathfrak{f} , then $\bar{\gamma} = \gamma|_{\mathfrak{t}_0}$ belongs to $\mathbb{k}\bar{\alpha}$. Hence, $\text{rk } \mathfrak{f} \geq 2$ and \mathfrak{f} is simple. By the construction, $\mathfrak{f}^\sigma = \mathfrak{sl}_2 + \tilde{\mathfrak{t}}$, where $\tilde{\mathfrak{t}} \subset \mathfrak{t}_0$ is a maximal torus of \mathfrak{f}^σ . The involution σ induces an automorphism of \mathfrak{f} of order 2. It cannot be inner, because the restrictions of α and β to $\tilde{\mathfrak{t}}$ coincide. From the description of outer involutions, we deduce that $\text{rk } \mathfrak{f} \leq 2$. Therefore the only possibility for $(\mathfrak{f}, \mathfrak{f}^\sigma)$ is the pair $(\mathfrak{sl}_3, \mathfrak{so}_3)$. Here $\beta = \alpha_1 + \alpha_2$ with α_1, α_2 being simple roots of \mathfrak{sl}_3 and $\sigma(\alpha_1) = \alpha_2$. Since $e_\beta = [e_{\alpha_1}, e_{\alpha_2}]$ in \mathfrak{sl}_3 , up to a suitable normalisation, we have also $e_\beta = [e_\alpha, e_{\sigma(\alpha)}]$ in \mathfrak{g} and $\beta = \alpha + \sigma(\alpha)$ in Δ . \square

3.3. Properties and generators of algebras $\mathcal{Z}(\mathfrak{g}, \vartheta)$. From now on, G is a connected semisimple algebraic group and $\mathfrak{g} = \text{Lie } G$. We consider $\vartheta \in \text{Aut}(\mathfrak{g})$ of order $m \geq 2$ and freely use the previous notation and results, with \mathfrak{q} being replaced by \mathfrak{g} . In particular,

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_{m-1},$$

where $\{0, 1, \dots, m-1\}$ is the fixed set of representatives for \mathbb{Z}_m , and G_0 is the connected subgroup of G with $\text{Lie } G_0 = \mathfrak{g}_0$. Then $\mathfrak{g}_{(t)}$ is a family of Lie algebras parameterised by $t \in \mathbb{P} = \mathbb{k} \cup \{\infty\}$, where the algebras $\mathfrak{g}_{(t)}$ with $t \in \mathbb{k}^*$ are isomorphic to $\mathfrak{g} = \mathfrak{g}_{(1)}$, while $\mathfrak{g}_{(0)}$ and $\mathfrak{g}_{(\infty)}$ are different \mathbb{N}_0 -graded contractions of \mathfrak{g} .

Note that \mathfrak{g}_0 is a reductive Lie algebra. Let κ be the Killing form on \mathfrak{g} . We identify \mathfrak{g} and \mathfrak{g}_0 with their duals via κ . Moreover, since $\kappa(\mathfrak{g}_i, \mathfrak{g}_j) = 0$ if $i + j \notin \{0, m\}$, the dual space of $\mathfrak{g}_j, \mathfrak{g}_j^*$, can be identified with \mathfrak{g}_{m-j} . We identify also \mathfrak{t} with \mathfrak{t}^* and \mathfrak{t}_0 with \mathfrak{t}_0^* . Set $\mathcal{Z} = \mathcal{Z}(\mathfrak{g}, \vartheta)$.

Theorem 3.3 ([PY21]). *Suppose that $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$. Then $\text{tr.deg } \mathcal{Z} = \mathbf{b}(\mathfrak{g}, \vartheta) := \mathbf{b}(\mathfrak{g}) - \mathbf{b}(\mathfrak{g}_0) + \text{rk } \mathfrak{g}_0$.* \square

Note that $\mathcal{Z} \subset \mathcal{S}(\mathfrak{g})^{\mathfrak{g}_0}$ by [PY21, Eq. (3.6)]. Thereby $\text{tr.deg } \mathcal{Z} \leq \mathbf{b}(\mathfrak{g}, \vartheta)$, see [MY19, Prop. 1.1]. Thus, if $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$, then $\text{tr.deg } \mathcal{Z}$ takes the maximal possible value. Note also that $\mathbf{b}(\mathfrak{g}, \vartheta) = \mathbf{b}(\mathfrak{g})$ if and only if $[\mathfrak{g}_0, \mathfrak{g}_0] = 0$.

Since ϑ acts on $\mathcal{S}(\mathfrak{g})^{\mathfrak{g}}$, there is a generating set $\{H_1, \dots, H_l\} \subset \mathcal{S}(\mathfrak{g})^{\mathfrak{g}}$ consisting of ϑ -eigenvectors. Then $\vartheta(H_i) = \zeta^{r_i} H_i$ with $0 \leq r_i < m$. The integers r_i depend only on the connected component of $\text{Aut}(\mathfrak{g})$ that contains ϑ , and if a is the order of ϑ in $\text{Aut}(\mathfrak{g})/\text{Int}(\mathfrak{g})$, then $\zeta^{ar_i} = 1$. Therefore, if \mathfrak{g} is simple, then $\zeta^{r_i} = \pm 1$ for all types but D_4 .

Recall from Section 1.3 that $\mathfrak{g}_{(0)} = \mathfrak{g}_{(0, \varphi)} =: \mathfrak{g}_{(0, \vartheta)}$ is a contraction of \mathfrak{g} defined by φ . Below we freely use notation of Sections 1.2, 1.4.

Lemma 3.4 ([PY21]). *For any $\vartheta \in \text{Aut}(\mathfrak{g})$ of order m , we have*

- (1) $\vartheta(H_j) = H_j$ if and only if $d_j^\bullet \in m\mathbb{Z}$;
- (2) $\sum_{j=1}^l r_j = \frac{1}{2}m(\text{rk } \mathfrak{g} - \text{rk } \mathfrak{g}_0)$;

$$(3) \operatorname{rk} \mathfrak{g}_0 = \#\{j \mid \vartheta(H_j) = H_j\}. \quad \square$$

Lemma 3.5. *For each j , the restriction of H_j to \mathfrak{g}_0^* is non-zero if and only if $\vartheta(H_j) = H_j$.*

Proof. If $H_j|_{\mathfrak{g}_0^*} \neq 0$, then the lowest φ -component $H_{j,0} \in \mathcal{S}(\mathfrak{g}_0)$ is non-zero and hence we have $\vartheta(H_j) = H_j$.

Consider some $x \in \mathfrak{t}_0^* \cap \mathfrak{g}_{\text{reg}}^*$, which exists by [Ka83, §8.8], and apply Kostant's regularity criterion [K63, Theorem 9] to x . According to this criterion, $\langle d_x H_i \mid 1 \leq i \leq l \rangle_{\mathbb{k}} = \mathfrak{g}^x = \mathfrak{t}$. Here $d_x H_i \in \mathfrak{g}_0^x = \mathfrak{t}_0$ if and only if $\vartheta(H_i) = H_i$. In view of Lemma 3.4(3), we have $d_x H_i \neq 0$ for each i such that $r_i = 0$. Then also $H_{i,0} \neq 0$, whenever $r_i = 0$. \square

Theorem 3.6 ([PY21]). *Suppose that $\vartheta \in \operatorname{Aut}(\mathfrak{g})$ admits a g.g.s. and $\operatorname{ind} \mathfrak{g}_{(0)} = \operatorname{rk} \mathfrak{g}$. Then*

(i) $\mathcal{Z}_\times := \operatorname{alg}\langle H_{j,i} \mid 1 \leq j \leq l, 0 \leq i \leq d_j^* \rangle \subset \mathcal{Z}$ is a polynomial Poisson-commutative subalgebra of $\mathcal{S}(\mathfrak{g})^{\mathfrak{g}_0}$ having the maximal transcendence degree.

(ii) More precisely, if H_1, \dots, H_l is a g.g.s. that consists of ϑ -eigenvectors, then \mathcal{Z}_\times is freely generated by the non-zero bi-homogeneous components of all H_j . \square

A precise relationship between \mathcal{Z} and \mathcal{Z}_\times depends on further properties of ϑ . Two complementary assertions are given below.

Corollary 3.7 ([PY21]). *In addition to the hypotheses of Theorem 3.6, suppose that $\mathfrak{g}_{(0)}$ has the codim-2 property and $\mathfrak{g}_0 = \mathfrak{g}^\vartheta$ is not abelian. Then $\mathcal{Z} = \mathcal{Z}_\times$ is the polynomial algebra freely generated by all non-zero bi-homogeneous components $H_{j,i}$.*

Corollary 3.8 ([PY21]). *In addition to the hypotheses of Theorem 3.6, suppose that $\mathfrak{g}_{(0)}$ has the codim-2 property, ϑ is inner, and $\mathfrak{g}_0 = \mathfrak{g}^\vartheta$ is abelian. Then $\mathcal{Z}_\infty = \mathcal{S}(\mathfrak{g}_0)$ and $\mathcal{Z} = \operatorname{alg}\langle \mathcal{Z}_\times, \mathfrak{g}_0 \rangle$ is a polynomial algebra.*

3.4. The equality for the index of $\mathfrak{g}_{(0)}$. The equality $\operatorname{ind} \mathfrak{g}_{(0)} = \operatorname{rk} \mathfrak{g}$ is very important in our context. If it holds, then \mathcal{Z} can be extended to a Poisson-commutative subalgebra of the maximal possible transcendence degree $\mathfrak{b}(\mathfrak{g})$ in the same way as in [PY, Sect. 6.2].

- It holds if $\operatorname{ord}(\vartheta)$ is two or three [P07, PY23].
- It holds if \mathfrak{g} is either \mathfrak{so}_n or of type G_2 [PY23].
- It holds if $\mathfrak{g}_1 \cap \mathfrak{g}_{\text{reg}} \neq \emptyset$ by [P09, Prop. 5.3].

Below we will see a few more positive examples.

Let $(\mathfrak{g}, \vartheta)$ be an indecomposable pair, where $\mathfrak{g} = \mathfrak{h}^{\oplus n}$, the algebra \mathfrak{h} is simple, and ϑ corresponds to $\tilde{\vartheta} \in \operatorname{Aut}(\mathfrak{h})$. Note that $\tilde{\vartheta}$ may be trivial. We remark also that $\mathfrak{g}_0 \simeq \mathfrak{h}^{\tilde{\vartheta}}$.

Let $\mathfrak{h}_{(0)}$ be the contraction of \mathfrak{h} associated with $\tilde{\vartheta}$.

Lemma 3.9 ([PY21, Lemma 8.1]). *If $\operatorname{ind} \mathfrak{h}_{(0)} = \operatorname{rk} \mathfrak{h}$, then $\operatorname{ind} \mathfrak{g}_{(0)} = \operatorname{rk} \mathfrak{g}$. If there is a g.g.s. for $\tilde{\vartheta}$ in $\mathcal{S}(\mathfrak{h})$, then there is a g.g.s. for ϑ in $\mathcal{S}(\mathfrak{g})$. \square*

Theorem 3.10. *Suppose that either $\mathfrak{g}_0 = \mathfrak{t}_0 \subset \mathfrak{t}$ or \mathfrak{g} is simple and among the Kac labels of ϑ , see Section 3.1 for the definition, only p_0 is zero. Then $\operatorname{ind} \mathfrak{g}_{(0)} = \operatorname{rk} \mathfrak{g}$.*

Proof. Making use of Lemma 3.9, we may assume that \mathfrak{g} is simple. If $\mathfrak{g}_0 = \mathfrak{t}_0$, then each Kac label of ϑ is non-zero, see [Vi76, Prop. 17] or e.g. [PY23, Sect. 2.4]. Let ϑ_1 be the automorphism obtained from ϑ by changing each non-zero Kac label of ϑ to 1. Then ϑ_1 may be different from ϑ and it may define a different \mathbb{Z}_m -grading of \mathfrak{g} . However, the Lie brackets $[\cdot, \cdot]_{(0)}$ defined by ϑ and ϑ_1 coincide [PY23]. Thus we may suppose that each non-zero Kac label of ϑ is 1. By our assumptions on ϑ , only p_0 may not be equal to 1.

Let Π be a set of the simple roots of \mathfrak{g} chosen in the same way as in Section 3.1. Let $e_i \in \mathfrak{g}_{\alpha_i}$ be a non-zero root vector corresponding to a simple root $\alpha_i \in \Pi$. If ϑ is inner, then $e = \sum_{i=1}^l e_i \in \mathfrak{g}_1$ is a regular nilpotent element in \mathfrak{g} . Thereby $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$ by [P09, Prop. 5.3].

Suppose that ϑ is outer. Let σ be the diagram automorphism of \mathfrak{g} associated with ϑ and $\Pi' = \{\nu_1, \dots, \nu_r\}$ the set of simple roots of \mathfrak{g}^σ used in Section 3.1. Then there is a bijection between Π' and the set of $\langle \sigma \rangle$ -orbits on Π . Namely, ν_j corresponds to the $\langle \sigma \rangle$ -orbit $\langle \sigma \rangle \cdot \alpha_i$ of $\alpha_i \in \Pi$, if $\nu_j = \alpha_i|_{\mathfrak{t}_0}$. There is a normalisation of the root vectors $e_i = e_{\alpha_i}$ such that $e'_j = \sum_{\alpha \in \langle \sigma \rangle \cdot \alpha_i} e_\alpha$ is a simple root vector of \mathfrak{g}^σ of the weight $\nu_j = \alpha_i|_{\mathfrak{t}_0}$ for each j [Ka83, §8.2]. If the Kac label p_j is 1, then $e'_j \in \mathfrak{g}_1$ according to the description of \mathfrak{g}_1 given in [Vi76, Prop. 17]. Since $p_j = 1$ for any $j \geq 1$, we obtain $e = \sum_{i=1}^l e_i \in \mathfrak{g}_1$. Then again by [P09, Prop. 5.3], we have $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$. \square

The case $\mathfrak{g}_0 = \mathfrak{t}_0$ is of particular importance, because $\text{tr.deg } \mathcal{Z}(\mathfrak{g}, \vartheta) = \mathfrak{b}(\mathfrak{g})$ here. If $\mathfrak{g}_0 = \mathfrak{t}$, then $\mathcal{Z}(\mathfrak{g}, \vartheta)$ coincides with the Poisson-commutative subalgebra $\mathcal{Z}(\mathfrak{b}, u^-)$ constructed in [PY'], see [PY23, Example 4.5]. In particular, this algebra is known to be maximal [PY', Theorem 5.5]. If $\mathfrak{g}_0 = \mathfrak{t}_0$ is a proper subspace of \mathfrak{t} , then we obtain a less studied subalgebra. We conjecture that it is still maximal.

4. PROPERTIES OF \mathcal{Z}_∞ IN THE REDUCTIVE CASE

In order to understand $\mathcal{Z}_\infty^{\mathfrak{g}_0}$, we study symmetric invariants of $\tilde{\mathfrak{g}} = \mathfrak{g}_0 \times \mathfrak{g}_{(\infty)}$. The Lie algebra $\tilde{\mathfrak{g}}$ is a contraction of $\mathfrak{g}_0 \oplus \mathfrak{g}$ associated with a map $\tilde{\varphi} : \mathbb{k}^* \rightarrow \text{GL}(\mathfrak{g}_0 \oplus \mathfrak{g})$, where $\tilde{\varphi}(s) = \tilde{\varphi}_s$. In order to define $\tilde{\varphi}_s$, we consider a vector space decomposition

$$\mathfrak{g}_0 \oplus \mathfrak{g} = \mathfrak{g}_0^{\text{d}} \oplus \mathfrak{g}_{m-1} \oplus \dots \oplus \mathfrak{g}_2 \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_0^{\text{ab}},$$

where the first summand $\mathfrak{g}_0^{\text{d}}$ is embedded diagonally into $\mathfrak{g}_0 \oplus \mathfrak{g}$ and the last summand $\mathfrak{g}_0^{\text{ab}}$ is embedded anti-diagonally. Then set $\tilde{\varphi}_s|_{\mathfrak{g}_0^{\text{d}}} = \text{id}$, $\tilde{\varphi}_s|_{\mathfrak{g}_{m-j}} = s^j \text{id}$, and finally $\tilde{\varphi}_s|_{\mathfrak{g}_0^{\text{ab}}} = s^m \text{id}$. If we consider $\xi \in (\mathfrak{g}_0^{\text{ab}})^*$ as an element of \mathfrak{g}^* and of $\tilde{\mathfrak{g}}^*$, then $\tilde{\mathfrak{g}}^\xi = \mathfrak{g}_0^\xi \oplus \mathfrak{g}^\xi$ as a vector space. If ξ is regular in \mathfrak{g}^* , then $\dim \tilde{\mathfrak{g}}^\xi = \text{ind } \mathfrak{g}_0 + \text{ind } \mathfrak{g}$, cf. (2.1). Then, in view of [Ka83, §8.8], which states that $\mathfrak{g}_0^* \cap \mathfrak{g}_{\text{reg}}^* \neq \emptyset$, we have $\text{ind } \tilde{\mathfrak{g}} \leq \text{ind } \mathfrak{g}_0 + \text{ind } \mathfrak{g}$. Since $\tilde{\mathfrak{g}}$ is a contraction of $\mathfrak{g}_0 \oplus \mathfrak{g}$, there is the equality, cf. [P09, Sect. 6.2],

$$(4.1) \quad \text{ind } \tilde{\mathfrak{g}} = \text{ind}(\mathfrak{g}_0 \oplus \mathfrak{g}) = \text{rk } \mathfrak{g}_0 + \text{rk } \mathfrak{g}.$$

Note that ϑ is also an automorphism of $\mathfrak{g}_0 \oplus \mathfrak{g}$, acting on $\mathfrak{g}_0 \oplus \mathfrak{g}_0$ trivially, as well as of $\tilde{\mathfrak{g}}$. However, the contraction defined by $\tilde{\varphi}$ is not directly related to ϑ .

4.1. Small rank examples. Let ϑ be the outer automorphism of \mathfrak{sl}_3 of order 4 with the Kac labels $\mathbf{p} = (0, 1)$. Then the corresponding \mathbb{Z}_4 -grading of $\mathfrak{g} = \mathfrak{sl}_3$ looks as follows

$$\mathfrak{g} = \mathfrak{sl}_2 \oplus \mathbb{k}_I^2 \oplus \mathfrak{t}_1 \oplus \mathbb{k}_{II}^2,$$

where $\dim \mathbb{k}_I^2 = \dim \mathbb{k}_{II}^2 = 2$ and $\mathfrak{t}_1 \subset \mathfrak{t}$ is one-dimensional.

Let $E_{ij} \in \mathfrak{gl}_n$ be elementary matrices (matrix units).

Lemma 4.1. *Suppose $\tilde{\mathfrak{g}} = \mathfrak{g}_0 \times \mathfrak{g}_{(\infty)}$ corresponds to the pair $(\mathfrak{g}, \vartheta)$ described above. Let $e \in (\mathfrak{g}_0^{\text{ab}})^*$ be a non-zero nilpotent element. Then $e + y \in \tilde{\mathfrak{g}}_{\text{reg}}^*$ for a generic $y \in \mathfrak{g}_1^*$.*

Proof. We may suppose that $\mathfrak{g}_0 = \mathfrak{sl}_2$ is embedded into \mathfrak{g} as $\langle E_{11} - E_{33}, E_{13}, E_{31} \rangle_{\mathbb{k}}$. Then $\mathbb{k}_I^2 = \langle E_{12} - E_{23}, E_{21} + E_{32} \rangle_{\mathbb{k}}$, $\mathbb{k}_{II}^2 = \langle E_{12} + E_{23}, E_{21} - E_{32} \rangle_{\mathbb{k}}$, $\mathfrak{t} = \mathbb{k}(E_{11} - 2E_{22} + E_{33})$, and e is identified with $\kappa(E_{13}, \cdot)$. As y we choose $\kappa(E_{21} - E_{32}, \cdot)$. Set $\gamma = y + e$ and let π be the Poisson tensor of $\tilde{\mathfrak{g}}$. We identify $\mathfrak{g}_0^{\text{d}}$ and $\mathfrak{g}_0^{\text{ab}}$ with \mathfrak{sl}_2 in a natural way.

Observe that $\pi(\gamma)(\mathfrak{t}_1, \mathbb{k}(E_{12} + E_{23})) \neq 0$ and that

$$(4.2) \quad \pi(\gamma)(\mathfrak{t}_1 + \mathbb{k}(E_{12} + E_{23}), \mathfrak{g}_0^{\text{d}} + \mathbb{k}_I^2 + \mathbb{k}(E_{21} - E_{32}) + \mathfrak{g}_0^{\text{ab}}) = 0.$$

Therefore $\tilde{\mathfrak{g}}^\gamma \subset \mathfrak{g}_0^{\text{d}} + \mathbb{k}_I^2 + \mathbb{k}(E_{21} - E_{32}) + \mathfrak{g}_0^{\text{ab}}$. Clearly $E_{13} \in \mathfrak{g}_0^{\text{ab}}$ belongs to $\tilde{\mathfrak{g}}^\gamma$. Now suppose that $x = x_0 + x_1 + x_3 + x_4$ with $x_0 \in \mathfrak{g}_0^{\text{d}}$, $x_1 \in \mathbb{k}(E_{21} - E_{32})$, $x_3 \in \mathbb{k}_I^2$, $x_4 \in \mathfrak{g}_0^{\text{ab}}$ belongs to $\tilde{\mathfrak{g}}^\gamma$. We may assume that $x_4 \in \mathbb{k}E_{31} + \mathbb{k}(E_{11} - E_{33})$. Then

$$(4.3) \quad \pi(\gamma)(x, \mathbb{k}_I^2) = \pi(\gamma)(x_0 + x_1, \mathbb{k}_I^2) = 0, \quad \pi(\gamma)(x, \mathfrak{g}_0^{\text{ab}}) = \pi(\gamma)(x_0, \mathfrak{g}_0^{\text{ab}}) = 0, \quad \text{and}$$

$$(4.4) \quad \pi(\gamma)(x, \mathfrak{g}_0^{\text{d}}) = \pi(\gamma)(x_3 + x_4, \mathfrak{g}_0^{\text{d}}) = 0.$$

Thereby $x_0 \in \mathbb{k}E_{13}$, which implies $x_0 + x_1 \in \mathbb{k}(E_{13} + E_{21} - E_{32})$. Since $\pi(\gamma)(x_4, E_{13}) = 0$ and $\pi(\gamma)(E_{21} + E_{32}, E_{13}) \neq 0$ for $E_{13} \in \mathfrak{g}_0^{\text{d}}$, we obtain $x_3 \in \mathbb{k}(E_{12} - E_{23})$. Looking at the commutators with $E_{11} - E_{33} \in \mathfrak{g}_0^{\text{d}}$, we obtain $x_3 + x_4 \in \mathbb{k}(E_{12} - E_{23} + E_{31})$.

We conclude that $\dim \tilde{\mathfrak{g}}^\gamma \leq 3$. It cannot be smaller than $3 = \text{ind } \tilde{\mathfrak{g}}$, see (4.1). Thus γ is a regular point and the same holds for the elements of a non-empty open subset of \mathfrak{g}_1^* . \square

Next we generalise Lemma 4.1 to periodic automorphisms of order $4n$ of direct sums $\mathfrak{sl}_3^{\oplus n}$. Let now $\tilde{\vartheta}$ be the outer automorphism of $\mathfrak{h} = \mathfrak{sl}_3$ of order 4 with the Kac labels $(1, 0)$. Set $\mathfrak{g} = \mathfrak{h}^{\oplus n}$ and let $\vartheta \in \text{Aut}(\mathfrak{g})$ be obtained from $\tilde{\vartheta}$ and a cyclic permutation of the summands of \mathfrak{g} , see (0.1).

Lemma 4.2. *Suppose $\tilde{\mathfrak{g}} = \mathfrak{g}_0 \times \mathfrak{g}_{(\infty)}$ corresponds to the pair $(\mathfrak{g}, \vartheta)$ described above. Let $e \in (\mathfrak{g}_0^{\text{ab}})^*$ be a non-zero nilpotent element. Then $e + y \in \tilde{\mathfrak{g}}_{\text{reg}}^*$ for a generic $y \in \mathfrak{g}_1^*$.*

Proof. It is convenient to identify \mathfrak{g}_0^d with \mathfrak{h}_0 , \mathfrak{g}_0^{ab} with $\mathfrak{h}_0 \bar{t}^{4n}$, each \mathfrak{g}_{4k+i} , where $0 \leq i \leq 3$, with $\mathfrak{h}_i \bar{t}^{4n-4k-i}$, assuming that $\bar{t}^{4n+1} = 0$. Then $[x\bar{t}^j, y\bar{t}^u] = [x, y]\bar{t}^{j+u}$ in $\tilde{\mathfrak{g}}$ for $\tilde{\vartheta}$ -eigenvectors $x, y \in \mathfrak{h}$.

We understand e as a linear function such that $e(E_{13}\bar{t}^{4n}) = e((E_{11} - E_{33})\bar{t}^{4n}) = 0$ and $e(E_{31}\bar{t}^{4n}) = 1$. Modelling the proof of Lemma 4.1, take y such that $y((E_{12} - E_{23})\bar{t}^{4n-1}) = 2$ and $y((E_{21} + E_{32})\bar{t}^{4n-1}) = 0$. Set $\gamma = e + y$. Note that $\gamma([x\bar{t}^j, y\bar{t}^u]) = 0$, if $j + u \notin \{4n-1, 4n\}$. Iterating computations in the spirit of (4.2), (4.3), (4.4), we obtain

$$\tilde{\mathfrak{g}}^\gamma = \langle E_{13}\bar{t}^{4n}, E_{13}\bar{t}^{4k} + (E_{21} - E_{32})\bar{t}^{4k+1}, (E_{12} - E_{23})\bar{t}^{4k+3} + E_{31}\bar{t}^{4k+4} \mid 0 \leq k < n \rangle_{\mathbb{k}}.$$

In particular, $\dim \tilde{\mathfrak{g}}^\gamma = 1 + 2n = \text{ind } \tilde{\mathfrak{g}}$ and $\gamma \in \tilde{\mathfrak{g}}_{\text{reg}}^*$. The same holds for the elements of a non-empty open subset of \mathfrak{g}_1^* . \square

4.2. General computations. Let us begin with a technical lemma.

Lemma 4.3. *Let \mathfrak{q} be a finite-dimensional Lie algebra. Suppose we have $\eta \in \mathfrak{q}^*$ and $\bar{y} \in (\mathfrak{q}^\eta)^*$ such that $\dim(\mathfrak{q}^\eta)^{\bar{y}} = \text{ind } \mathfrak{q}$. Then for any extension y of \bar{y} to a linear function on \mathfrak{q} , there is $s \in \mathbb{k}^*$ such that $s\eta + y \in \mathfrak{q}_{\text{reg}}^*$.*

Proof. Write $\mathfrak{q} = \mathfrak{m} \oplus \mathfrak{q}^\eta$, where $\hat{\eta}$ is non-degenerate on \mathfrak{m} . Let $\mathfrak{m}_1 \subset \mathfrak{q}^\eta$ be a subspace of dimension $\dim \mathfrak{q}^\eta - \text{ind } \mathfrak{q}$ such that \hat{y} is non-degenerate on \mathfrak{m}_1 . By the construction $\dim(\mathfrak{m} \oplus \mathfrak{m}_1) = \dim \mathfrak{q} - \text{ind } \mathfrak{q}$. We extend \bar{y} to a linear function y on \mathfrak{q} in some way, which is of no importance. Choose bases in \mathfrak{m} , \mathfrak{m}_1 , take their union, and let $M = \begin{pmatrix} sA + C & B \\ -B^t & A_1 \end{pmatrix}$ be the matrix of $s\hat{\eta} + \hat{y}$, where $s \in \mathbb{k}$, in this basis; here A is the matrix of $\hat{\eta}|_{\mathfrak{m}}$ and A_1 is the matrix of $\hat{y}|_{\mathfrak{m}_1}$. Then

$$\det(M) = s^{\dim \mathfrak{m}} \det(A) \det(A_1) + (\text{terms of smaller degree in } s).$$

Since $\det(A) \det(A_1) \neq 0$, for almost all $s \in \mathbb{k}$, we have $\det(M) \neq 0$. Whenever this happens, $\text{rk}(s\hat{\eta} + \hat{y}) \geq \dim \mathfrak{q} - \text{ind } \mathfrak{q}$ and $s\eta + y \in \mathfrak{q}_{\text{reg}}^*$. \square

Theorem 4.4. *The algebra $\tilde{\mathfrak{g}}$ has the codim-2 property.*

Proof. The subalgebra \mathfrak{g}_0 is reductive and it contains a semisimple element x that is regular in \mathfrak{g} , see e.g. [Ka83, §8.8]. Thus, there is $\eta \in (\mathfrak{g}_0^{\text{ab}})^* \subset \tilde{\mathfrak{g}}^*$ that corresponds to a regular element of \mathfrak{g} and here $\dim \tilde{\mathfrak{g}}^\eta = \text{rk } \mathfrak{g} + \text{rk } \mathfrak{g}_0 = \text{ind } \tilde{\mathfrak{g}}$, cf. (2.1).

Take now $\xi \in \tilde{\mathfrak{g}}^*$ such that $\bar{\xi} = \xi|_{\mathfrak{g}_0^{\text{ab}}}$ corresponds to a regular element of \mathfrak{g} . Note that $\lim_{s \rightarrow 0} s^m \tilde{\varphi}_s(\xi) = \bar{\xi}$. Since $\bar{\xi} \in \tilde{\mathfrak{g}}_{\text{reg}}^*$, we have $\varphi_s(\xi) \in \tilde{\mathfrak{g}}_{\text{reg}}^*$ for almost all s . Each map $\tilde{\varphi}_s: \tilde{\mathfrak{g}} \rightarrow \tilde{\mathfrak{g}}$ is an automorphism of the Lie algebra $\tilde{\mathfrak{g}}$. Thus $\xi \in \tilde{\mathfrak{g}}_{\text{reg}}^*$.

Assume that $D \subset \tilde{\mathfrak{g}}_{\text{sing}}^*$ is a divisor in $\tilde{\mathfrak{g}}^*$. Then $\{\xi \mid \xi \in D\} \subset (\mathfrak{g}_0^{\text{ab}})^*$ lies in a proper closed subset of $(\mathfrak{g}_0^{\text{ab}})^*$. Thereby $D = D_0 \times \text{Ann}(\mathfrak{g}_0^{\text{ab}})$ for some divisor $D_0 \subset (\mathfrak{g}_0^{\text{ab}})^*$. Since $\tilde{\mathfrak{g}}_{\text{sing}}^*$ is a G_0 -stable subset of $\tilde{\mathfrak{g}}^*$, the divisor D_0 is G_0 -stable as well. Since $\tilde{\mathfrak{g}}_{\text{sing}}^*$ is a conical subset, D_0 is the zero set of a homogeneous polynomial.

A generic fibre of the categorical quotient $\varpi: (\mathfrak{g}_0^{\text{ab}})^* \rightarrow (\mathfrak{g}_0^{\text{ab}})^* // G_0$ consists of a single G_0 -orbit. Since D_0 is G_0 -stable, it follows that $\dim \overline{\varpi(D_0)} = \text{rk } \mathfrak{g}_0 - 1$. Then also $\varpi^{-1}(y) \subset D_0$ for any $y \in \varpi(D_0)$, because each fibre of ϖ is irreducible.

The intersection $D_0 \cap \mathfrak{t}_0^*$ does not contain regular in \mathfrak{g} elements, thereby it is a subspace of codimension 1, the zero set of $\bar{\alpha} = \alpha|_{\mathfrak{t}_0}$ for some $\alpha \in \Delta^+$. Let $\eta \in D_0 \cap \mathfrak{t}_0^*$ be a generic point. Then η is either regular or subregular in \mathfrak{g}_0 . Furthermore, if $\beta \in \Delta$ and $\beta(\eta) = 0$, then $\bar{\beta} = \beta|_{\mathfrak{t}_0} \in \mathbb{k}\bar{\alpha}$. Set $\mathfrak{f} = [\mathfrak{g}^\eta, \mathfrak{g}^\eta]$. Note that \mathfrak{f} is semisimple and that the pair $(\mathfrak{f}, \mathfrak{f}^\vartheta)$ is indecomposable.

(\diamond) Case 1. Suppose that η is not regular in \mathfrak{g}_0 . Then necessarily $[\mathfrak{g}_0^\eta, \mathfrak{g}_0^\eta] = \mathfrak{sl}_2$. Let $e \in [\mathfrak{g}_0^\eta, \mathfrak{g}_0^\eta]$ be a non-zero nilpotent element, which we regard also as a point in $(\mathfrak{g}_0^{\text{ab}})^*$. Then $\eta + e \in D_0$, since $\varpi(\eta + e) = \varpi(\eta)$. Suppose $\{\beta \in \Delta^+ \mid \beta(\eta) = 0\} = \{\vartheta^k(\alpha) \mid k \geq 0\}$. Then $\vartheta^k(\alpha) + \vartheta^{k'}(\alpha)$ with $k, k' \geq 0$ is never a root of \mathfrak{g} , the subalgebra \mathfrak{f} is a direct sum of copies of \mathfrak{sl}_2 , the nilpotent element e is regular in \mathfrak{f} , and $\eta + e \in \mathfrak{g}_{\text{reg}}^*$. This is a contradiction.

Suppose that there is no equality for the sets above. Then $\vartheta|_{\mathfrak{f}}$ is constructed from an outer automorphism $\tilde{\vartheta}$ of a simple Lie algebra \mathfrak{h} . From Section 3.2 we know that $\mathfrak{h} = \mathfrak{sl}_3$. Next $\mathfrak{h}^{\tilde{\vartheta}} \simeq \mathfrak{sl}_2$. The fixed points subalgebra can be embedded in two different ways, as \mathfrak{so}_3 or as $\langle E_{11} - E_{33}, E_{13}, E_{31} \rangle_{\mathbb{k}}$. For the first embedding, e is still regular in \mathfrak{f} , a contradiction.

Suppose that $\mathfrak{h}^{\tilde{\vartheta}} = \langle E_{11} - E_{33}, E_{13}, E_{31} \rangle_{\mathbb{k}}$. The twisted affine Dynkin digram of type A_2 has two nodes. Our choice of $\mathfrak{h}^{\tilde{\vartheta}}$ implies that $\tilde{\vartheta}$ is the automorphism considered in Lemma 4.1. Then by Lemma 4.2, there is $y \in \mathfrak{f}_1^* \simeq \mathfrak{f}_i \subset \mathfrak{g}_j$, where $0 < j < m$ and i depends on $\vartheta|_{\mathfrak{f}}$, such that $e + y \in \tilde{\mathfrak{f}}_{\text{reg}}^*$. Note that $s\eta + e + y \in D$ for any $s \in \mathbb{k}^*$.

The vector spaces $\tilde{\mathfrak{g}}^\eta$ and $(\mathfrak{g}_0 \oplus \mathfrak{g})^\eta$ coincide. We have $\tilde{\mathfrak{g}}^\eta = \tilde{\mathfrak{f}} + \mathfrak{t}_0 + \mathfrak{t}$, where $\tilde{\mathfrak{g}}^\eta$ and $\tilde{\mathfrak{f}}$ are contractions of \mathfrak{g}^η and $\mathfrak{f} \oplus \mathfrak{f}^\vartheta$, respectively, given by the restrictions of $\tilde{\varphi}$. We extend e and y to linear functions on $\tilde{\mathfrak{g}}^\eta$ and $\tilde{\mathfrak{g}}$ keeping the same symbols for the extensions. Then

$$(4.5) \quad \dim(\tilde{\mathfrak{g}}^\eta)^{e+y} = (\text{rk } \mathfrak{g} + \text{rk } \mathfrak{g}_0) - (\text{rk } \mathfrak{f} + \text{rk } \mathfrak{f}_0) + \text{ind } \tilde{\mathfrak{f}} = \text{ind } \tilde{\mathfrak{g}}.$$

By Lemma 4.3, $s\eta + e + y \in \tilde{\mathfrak{g}}_{\text{reg}}^*$ for some $s \in \mathbb{k}^*$, a contradiction.

(\diamond) Case 2. Suppose now that η is regular in \mathfrak{g}_0 . Then $\mathfrak{f}_0 \subset \mathfrak{t}_0$. The quotient of $\tilde{\mathfrak{f}}$ by $(\mathfrak{t}_0^{\text{ab}} \cap \tilde{\mathfrak{f}}) \subset \mathfrak{g}_0^{\text{ab}}$ is the contraction $\mathfrak{f}_{(0), \vartheta^{-1}}$ associated with the restriction of ϑ^{-1} to \mathfrak{f} . For $y \in \tilde{\mathfrak{f}}^*$ such that $y(\mathfrak{t}_0^{\text{ab}}) = 0$, we have $\tilde{\mathfrak{f}}^y = \mathfrak{f}_{(0), \vartheta^{-1}}^y \oplus (\mathfrak{t}_0^{\text{ab}} \cap \tilde{\mathfrak{f}})$. By Theorem 3.10, $\text{ind } \mathfrak{f}_{(0), \vartheta^{-1}} = \text{rk } \mathfrak{f}$. Therefore there is $y \in \tilde{\mathfrak{g}}^*$ such that $y(\mathfrak{t}_0^{\text{ab}}) = 0$ and $\dim \tilde{\mathfrak{f}}^y = \text{rk } \mathfrak{f} + \text{rk } \mathfrak{f}_0$ for $\bar{y} = y|_{\tilde{\mathfrak{f}}}$. Repeating the argument of (4.5) and using again Lemma 4.3, we conclude that $s\eta + y \in \tilde{\mathfrak{g}}_{\text{reg}}^*$ for some $s \in \mathbb{k}^*$. This final contradiction shows that there is no divisor D . \square

4.3. Symmetric invariants. Set $r = \text{rk } \mathfrak{g}_0$ and choose a set of homogeneous generators $F_1, \dots, F_r \in \mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0}$. As above, let $\mathfrak{g}_0^{\text{ab}}$ stand for the abelian ideal of $\tilde{\mathfrak{g}}$, which is isomorphic to \mathfrak{g}_0 as a \mathfrak{g}_0 -module. Let further $\{H_j \mid 1 \leq j \leq l\} \subset \mathcal{S}(\mathfrak{g})^{\mathfrak{g}}$ be a generating set consisting of homogeneous polynomials that are ϑ -eigenvectors. Assume that $\vartheta(H_j) = H_j$ if and only

if $j \leq r$, cf. Lemma 3.4. Unless stated otherwise, we use upper bullets for the highest $\tilde{\varphi}$ -components of $H \in \mathcal{S}(\mathfrak{g}_0 \oplus \mathfrak{g})$.

Theorem 4.5. *There is a g.g.s. F_i, \tilde{H}_i, H_j with $1 \leq i \leq r < j \leq l$ for the contraction $\mathfrak{g}_0 \oplus \mathfrak{g} \rightsquigarrow \tilde{\mathfrak{g}}$ defined by $\tilde{\varphi}$. Furthermore, for $i \leq r$, $\tilde{H}_i \in H_i + \mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0}$, where \mathfrak{g}_0 is a direct summand of $\mathfrak{g}_0 \oplus \mathfrak{g}$, and the ring $\mathcal{S}(\tilde{\mathfrak{g}})^{\tilde{\mathfrak{g}}}$ is freely generated by $\{F_i^\bullet, \tilde{H}_i^\bullet, H_j^\bullet \mid 1 \leq i \leq r < j \leq l\}$.*

Proof. First we compute the number $D_{\tilde{\varphi}}$ using the properties of the grading on $\tilde{\mathfrak{g}}$:

$$D_{\tilde{\varphi}} = m \dim \mathfrak{g}_0 + \sum_{j=1}^{m-1} (m-j) \dim \mathfrak{g}_j = m \dim \mathfrak{g}_0 + \frac{m}{2} (\dim \mathfrak{g} - \dim \mathfrak{g}_0) = \frac{m}{2} (\dim \mathfrak{g} + \dim \mathfrak{g}_0).$$

Since $\mathfrak{g}_0^{\text{ab}}$ is embedded anti-diagonally in $\mathfrak{g}_0 \oplus \mathfrak{g}$, we have $\deg_{\tilde{\varphi}} F_i^\bullet = m \deg F_i$.

Suppose that $\vartheta(H_j) = H_j$. Then the restriction of H_j from \mathfrak{g}^* to $\mathfrak{g}_0^* \subset \mathfrak{g}^*$ is non-zero by Lemma 3.5. Thus $H_j^\bullet \in \mathcal{S}(\mathfrak{g}_0^{\text{ab}})^{\mathfrak{g}_0}$. Hence there is a homogeneous polynomial \tilde{H}_j in $H_j + \mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0}$, where \mathfrak{g}_0 is a direct summand of $\mathfrak{g}_0 \oplus \mathfrak{g}$, such that $\tilde{d}_j^\bullet = \deg_{\tilde{\varphi}} \tilde{H}_j^\bullet < m \deg H_j$. We have $\vartheta(\tilde{H}_j^\bullet) = \zeta^{-\tilde{d}_j^\bullet} \tilde{H}_j^\bullet$ and at the same time $\vartheta(\tilde{H}_j^\bullet) = \tilde{H}_j^\bullet$. Thereby $\tilde{d}_j^\bullet \leq m \deg H_j - m$.

Suppose now that $\vartheta(H_j) \neq H_j$, i.e., $\vartheta(H_j) = \zeta^{r_j} H_j$ with $0 < r_j < m$. Here $H_j^\bullet \notin \mathcal{S}(\mathfrak{g}_0^{\text{ab}})$ and $\deg_{\tilde{\varphi}} \tilde{H}_j^\bullet \leq m \deg H_j - r_j$. According to Lemma 3.4, $\sum_{j=1}^l r_j = \frac{1}{2}m(\text{rk } \mathfrak{g} - \text{rk } \mathfrak{g}_0)$. Thus

$$(4.6) \quad \sum_{i=1}^r \deg_{\tilde{\varphi}} F_i^\bullet + \sum_{j=1}^r \deg_{\tilde{\varphi}} \tilde{H}_j^\bullet + \sum_{j=r+1}^l \deg_{\tilde{\varphi}} H_j^\bullet \leq \\ \leq m\mathbf{b}(\mathfrak{g}_0) + m\mathbf{b}(\mathfrak{g}) - m\text{rk } \mathfrak{g}_0 - \frac{1}{2}m(\text{rk } \mathfrak{g} - \text{rk } \mathfrak{g}_0) = \frac{m}{2}(\dim \mathfrak{g} + \dim \mathfrak{g}_0) = D_{\tilde{\varphi}}.$$

By Theorem 1.4(i), we have the equality in (4.6) and the polynomials F_i, \tilde{H}_i, H_j form a good generating system for $\tilde{\varphi}$. We obtain also

$$(4.7) \quad \deg_{\tilde{\varphi}} H_j^\bullet = m \deg H_j - r_j \quad \text{if } 0 < r_j < m.$$

By Theorem 4.4, the Lie algebra $\tilde{\mathfrak{g}}$ has the *codim*-2 property. Then Theorem 1.4(iii) states that $\mathcal{S}(\tilde{\mathfrak{g}})^{\tilde{\mathfrak{g}}}$ is freely generated by $F_i^\bullet, \tilde{H}_i^\bullet, H_j^\bullet$ with $1 \leq i \leq r < j \leq l$. \square

Recall that $H_j = \sum_{i \geq 0} H_{j,i}$, where $\varphi_s(H_j) = \sum_{i \geq 0} s^i H_{j,i}$. Let $H_{j,\bullet}$ be the non-zero bi-homogeneous component of H_j with the minimal φ -degree. We regard each $H_{j,i}$ as an element of $\mathcal{S}(\mathfrak{g}_{(\infty)}) \subset \mathcal{S}(\tilde{\mathfrak{g}})$ identifying \mathfrak{g} with the subspace

$$\mathfrak{g}_{(\infty)} = \mathfrak{g}_{m-1} \oplus \dots \oplus \mathfrak{g}_1 \oplus \mathfrak{g}_0^{\text{ab}} \subset \tilde{\mathfrak{g}}.$$

Note that $\deg_{\tilde{\varphi}} H_{j,i} = m \deg H_j - i$.

Theorem 4.6. *The algebra $\mathcal{Z}_\infty^{\mathfrak{g}_0}$ is freely generated by $F_i^\bullet \in \mathcal{S}(\mathfrak{g}_0^{\text{ab}})$ with $1 \leq i \leq r$ and the set $\{H_{j,\bullet} \mid r < j \leq l\}$, where $H_{j,\bullet} = H_j^\bullet$ for each j .*

Proof. For any $H_j \in \mathcal{S}(\mathfrak{g}) \subset \mathcal{S}(\mathfrak{g}_0 \oplus \mathfrak{g})$, the highest $\tilde{\varphi}$ -component $H_j^\bullet \in \mathcal{S}(\tilde{\mathfrak{g}})$ is obtained by taking first $H_{j,\bullet} \in \mathcal{S}(\mathfrak{g})$ and then replacing in it each element of \mathfrak{g}_0 by its copy in $\mathfrak{g}_0^{\text{ab}}$. This is exactly how we understand $H_{j,\bullet} \in \mathcal{S}(\tilde{\mathfrak{g}})$.

By the construction and Proposition 1.1,

$$\mathcal{H} := \{F_i^\bullet, H_j^\bullet \mid 1 \leq i \leq r, r < j \leq l\} \subset \mathcal{Z}_\infty^{\mathfrak{g}_0}.$$

These homogeneous polynomials are algebraically independent, see Theorem 4.5. In view of Theorem 2.3(iii), we have $\text{tr.deg } \mathcal{Z}_\infty^{\mathfrak{g}_0} = \text{rk } \mathfrak{g} = l$. By Theorems 1.4(iii), 4.4, 4.5, $\tilde{\mathfrak{g}}$ satisfies the Kostant regularity criterion and the differentials of the generators $F_i^\bullet, \tilde{H}_i^\bullet, H_j^\bullet$ with $1 \leq i \leq r < j \leq l$ are linearly independent on a big open subset. Thus also

$$(4.8) \quad \dim \mathcal{J}(\mathcal{H}) \leq \dim \mathfrak{g} - 2,$$

where $\mathcal{J}(\mathcal{H}) \subset \mathfrak{g}_{(\infty)}^*$. By Theorem 1.5, the subalgebra generated by \mathcal{H} is algebraically closed in $\mathcal{S}(\mathfrak{g}_{(\infty)})$ and hence it coincides with $\mathcal{Z}_\infty^{\mathfrak{g}_0}$. \square

If ϑ is inner, then $\mathcal{Z}_\infty = \mathcal{S}(\mathfrak{g}_0)$ [PY21] and $\mathcal{Z}_\infty^{\mathfrak{g}_0} = \mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0}$. One does not need any heavy machinery in that case. If ϑ is outer, the situation is different. We have seen this in Theorem 4.6 and are going to obtain a description of \mathcal{Z}_∞ next.

Proposition 4.7. *If ϑ is outer, then a basis $\{y_1, \dots, y_R\}$ of \mathfrak{g}_0 together with the lowest φ -components $H_{j,\bullet}$, where $r < j \leq l$, freely generate \mathcal{Z}_∞ .*

Proof. We check that the differentials dy_i and $dH_{j,\bullet}$ of the proposed generators are linearly independent on a big open subset of \mathfrak{g}^* . For $\xi \in \mathfrak{g}^*$, set $\bar{\xi} = \xi|_{\mathfrak{g}_0}$. Note that $\bar{\xi} \in (\mathfrak{g}_0^*)_{\text{reg}}$ for all ξ in a big open subset of \mathfrak{g}^* . Let $\varpi_\xi: T_\xi^* \mathfrak{g}^* \rightarrow T_\xi^*(G_0 \cdot \xi)$ be the restriction map. Since $H_{j,\bullet}$ is a G_0 -invariant, we have $\varpi_\xi(d_\xi H_{j,\bullet}) = 0$ for each j . Note that $\ker(\varpi_\xi|_{\mathfrak{g}_0}) = \mathfrak{g}_0^{\bar{\xi}}$.

Suppose that $\bar{\xi} \in (\mathfrak{g}_0^*)_{\text{reg}}$. Then $\dim \varpi_\xi(\mathfrak{g}_0) = \dim \mathfrak{g}_0 - \text{rk } \mathfrak{g}_0$. By the Kostant regularity criterion [K63, Theorem 9] applied to \mathfrak{g}_0 , we have

$$\langle d_\xi F_i^\bullet \mid 1 \leq i \leq r \rangle_{\mathbb{k}} = \mathfrak{g}_0^{\bar{\xi}}.$$

Suppose now that $\xi \notin \mathcal{J}(\mathcal{H})$. This additional restriction still leaves us a big open subset of suitable elements, see (4.8). Then

$$\dim(\mathfrak{g}_0^{\bar{\xi}} + \langle d_\xi H_{j,\bullet} \mid r < j \leq l \rangle_{\mathbb{k}}) = l.$$

Thus, on a big open subset, the dimension of $V_\xi = \langle d_\xi y_i, d_\xi H_{j,\bullet} \mid 1 \leq i \leq R, r < j \leq l \rangle_{\mathbb{k}}$ is equal to the sum of $\dim \varpi_\xi(V_\xi) = \dim \mathfrak{g}_0 - \text{rk } \mathfrak{g}_0$ and $l = \dim \ker(\varpi_\xi|_{V_\xi})$, i.e., $\dim V_\xi = R + (l - r)$. This number is equal to the number of generators. By Theorem 1.5, the subalgebra

$$\text{alg}\langle y_i, H_{j,\bullet} \mid 1 \leq i \leq R, r < j \leq l \rangle$$

is algebraically closed in $\mathcal{S}(\mathfrak{g}_{(\infty)})$. Furthermore,

$$\text{alg}\langle y_i, H_{j,\bullet} \mid 1 \leq i \leq R, r < j \leq l \rangle \subset \mathcal{Z}_\infty \quad \text{and} \quad \text{tr.deg } \mathcal{Z}_\infty \leq \text{ind } \mathfrak{g}_{(\infty)} = R + l - r,$$

see Theorem 2.1 for the last equality. Thereby $\text{alg}\langle y_i, H_{j,\bullet} \mid 1 \leq i \leq R, r < j \leq l \rangle = \mathcal{Z}_\infty$. \square

In [PY21, Sect. 5], we have considered a larger Poisson-commutative subalgebra $\tilde{\mathcal{Z}} := \text{alg}\langle \mathcal{Z}, \mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0} \rangle$.

Corollary 4.8. *Suppose that $\text{ind } \mathfrak{g}_{(0)} = \text{rk } \mathfrak{g}$ and that $\mathcal{S}(\mathfrak{g}_{(0)})^{\mathfrak{g}_{(0)}} = \mathbb{k}[H_1^\bullet, \dots, H_l^\bullet]$, where each H_j is homogeneous, $\vartheta(H_j) \in \mathbb{k}H_j$, and the polynomials H_j^\bullet are highest φ -components.*

- *If $\infty \in \mathbb{P}_{\text{reg}}$, then $\text{alg}\langle \mathcal{Z}, \mathcal{Z}_\infty^{\mathfrak{g}_0} \rangle = \text{alg}\langle \mathcal{Z}, \mathcal{Z}_\infty \rangle = \tilde{\mathcal{Z}} = \mathcal{Z} = \text{alg}\langle \mathcal{Z}_\times, \mathfrak{g}_0 \rangle$;*
- *if $\infty \notin \mathbb{P}_{\text{reg}}$, then $\text{alg}\langle \mathcal{Z}, \mathcal{Z}_\infty^{\mathfrak{g}_0} \rangle = \tilde{\mathcal{Z}} = \text{alg}\langle \mathcal{Z}_\times, \mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0} \rangle$.*

In both cases, $\tilde{\mathcal{Z}}$ is a free (polynomial) algebra.

Proof. Our assumptions on $\mathcal{S}(\mathfrak{g}_{(0)})^{\mathfrak{g}_{(0)}}$ imply that $\mathcal{Z}_0 \subset \mathcal{Z}_\times$. Furthermore, $\{H_1, \dots, H_l\}$ is a g.g.s. for ϑ . Then \mathcal{Z}_\times is a polynomial algebra by Theorem 3.6. It has $\mathfrak{b}(\mathfrak{g}, \vartheta)$ algebraically independent generators $H_{j,i}$. Exactly $r = \text{rk } \mathfrak{g}_0$ of these generators belong to $\mathcal{S}(\mathfrak{g}_0)$, see Lemmas 3.4, 3.5. If we replace them with algebraically independent generators of $\mathcal{S}(\mathfrak{g}_0)^{\mathfrak{g}_0}$, the new algebra is still polynomial. This finishes the proof in view of Theorem 4.6. \square

The first statement of Corollary 4.8 generalises Corollary 3.8 to outer automorphisms.

REFERENCES

- [CM16] J.-Y. CHARBONNEL and A. MOREAU. The symmetric invariants of centralizers and Slodowy grading, *Math. Z.*, **282** (2016), no. 1-2, 273–339.
- [FL18] F. FAUQUANT-MILLET and P. LAMPROU. Polynomiality for the Poisson centre of truncated maximal parabolic subalgebras, *J. Lie Theory*, **28** (2018), 1063–1094.
- [GZ00] I.M. GELFAND and I.S. ZAKHAREVICH. Webs, Lenard schemes, and the local geometry of bi-Hamiltonian Toda and Lax structures, *Selecta Math., New Ser.*, **6** (2000), 131–183.
- [Lie3] V.V. GORBATSEVICH, A.L. ONISHCHIK and E.B. VINBERG. “Lie Groups and Lie Algebras III” (Encyclopaedia Math. Sci., vol. 41) Berlin: Springer 1994.
- [J06] A. JOSEPH. On semi-invariants and index for biparabolic (seaweed) algebras, I, *J. Algebra*, **305** (2006), 487–515.
- [Ka69] V.G. KAC. Automorphisms of finite order of semisimple Lie algebras, *Funct. Anal. Appl.* **3** (1969), 252–254.
- [Ka83] V.G. KAC. “Infinite-dimensional Lie algebras”. Birkhäuser, 1983.
- [K63] B. KOSTANT. Lie group representations on polynomial rings, *Amer. J. Math.*, **85** (1963), 327–404.
- [MY19] A. MOLEV and O. YAKIMOVA. Quantisation and nilpotent limits of Mishchenko–Fomenko subalgebras, *Represent. Theory*, **23** (2019), 350–378.
- [P07] D. PANYUSHEV. On the coadjoint representation of \mathbb{Z}_2 -contractions of reductive Lie algebras, *Adv. Math.*, **213** (2007), 380–404.
- [P09] D. PANYUSHEV. Periodic automorphisms of Takiff algebras, contractions, and θ -groups, *Transformation Groups*, **14**, no. 2 (2009), 463–482.
- [PPY] D. PANYUSHEV, A. PREMÉT and O. YAKIMOVA. On symmetric invariants of centralisers in reductive Lie algebras, *J. Algebra*, **313** (2007), 343–391.
- [PY08] D. PANYUSHEV and O. YAKIMOVA. The argument shift method and maximal commutative subalgebras of Poisson algebras, *Math. Res. Letters*, **15** (2008), no. 2, 239–249.

- [PY] D. PANYUSHEV and O. YAKIMOVA. Poisson-commutative subalgebras of $\mathcal{S}(\mathfrak{g})$ associated with involutions, *Int. Math. Res. Notices*, **2021** no. 23 (2021), 18367–18406.
- [PY'] D. PANYUSHEV and O. YAKIMOVA. Compatible Poisson brackets associated with 2-splittings and Poisson commutative subalgebras of $\mathcal{S}(\mathfrak{g})$, *J. LMS*, **103** (2021), no. 4, 1577–1595.
- [PY21] D. PANYUSHEV and O. YAKIMOVA. Periodic automorphisms, compatible Poisson brackets, and Gaudin subalgebras, *Transformation groups*, **26** (2021), 641–670.
- [PY23] D. PANYUSHEV and O. YAKIMOVA. Automorphisms of finite order, periodic contractions, and Poisson-commutative subalgebras of $\mathcal{S}(\mathfrak{g})$, *Math. Z.* **303**, article 51 (2023), (DOI).
- [Sp89] T.A. SPRINGER. Aktionen reductiver Gruppen auf Varietäten, in: “Algebraische Transformationsgruppen und Invariantentheorie”, DMV-Seminar, Bd. **13**, Basel–Boston–Berlin: Birkhäuser 1989, pp. 3–39.
- [St68] R. STEINBERG. Endomorphisms of algebraic groups, *Memoirs Amer. Math. Soc.*, № **80**, AMS, Providence, R.I. (1968), 108 pp.
- [Vi76] E.B. VINBERG. The Weyl group of a graded Lie algebra, *Math. USSR-Izv.* **10** (1976), 463–495.
- [Vi90] E.B. VINBERG. On certain commutative subalgebras of a universal enveloping algebra, *Math. USSR-Izv.* **36** (1991), 1–22.
- [Y14] O. YAKIMOVA. One-parameter contractions of Lie–Poisson brackets, *J. Eur. Math. Soc.*, **16** (2014), 387–407.
- [Y17] O. YAKIMOVA. Symmetric invariants of \mathbb{Z}_2 -contractions and other semi-direct products, *Intern. Math. Res. Notices*, (2017) 2017 (6): 1674–1716.

(O.Y.) INSTITUT FÜR MATHEMATIK, FRIEDRICH-SCHILLER-UNIVERSITÄT JENA, 07737 JENA, DEUTSCHLAND

Email address: oksana.yakimova@uni-jena.de