

INVERSE IMAGES OF A POSITIVE CLOSED CURRENT FOR A HOLOMORPHIC ENDOMORPHISM OF A COMPACT KÄHLER MANIFOLD

TAEYONG AHN

ABSTRACT. In this paper, we prove that for a given surjective holomorphic endomorphism f of a compact Kähler manifold X and for some integer p with $1 \leq p \leq k$, there exists a proper invariant analytic subset E for f such that if S is smooth in a neighborhood of E , the sequence $d_p^{-n}(f^n)^*(S - \alpha_S)$ converges to 0 exponentially fast in the sense of currents where d_p denotes the dynamical degree of order p and α_S is a closed smooth form in the de Rham cohomology class of S .

1. INTRODUCTION

Let (X, ω) be a compact Kähler manifold of complex dimension $k \geq 2$ such that $\int_X \omega^k = 1$. Let $f : X \rightarrow X$ be a surjective holomorphic map. For $0 \leq s \leq k$, the dynamical degree d_s of order s of f is the spectral radius of the pull-back operator f^* acting on the Hodge cohomology group $H^{s,s}(X, \mathbb{C})$. It is known that d_s itself is an eigenvalue of f^* on $H^{s,s}(X, \mathbb{C})$. An inequality due to Khovanskii, Teissier and Gromov ([13], [32]) implies that the function $s \rightarrow \log d_s$ is concave on $0 \leq s \leq k$. In particular, there are integers p and p' with $0 \leq p \leq p' \leq k$ such that

$$d_0 < \dots < d_p = \dots = d_{p'} > \dots > d_k.$$

We always have $d_0 = 1$. The last dynamical degree d_k is also called the topological degree of f because it is equal to the cardinality of $f^{-1}(x)$ for a generic point x in X . We call d_p the main dynamical degree of f .

The aim of this paper is to study the dynamics of a holomorphic endomorphism on a compact Kähler manifold by proving the following theorem:

Theorem 1.1. *Let (X, ω) be a compact Kähler manifold of dimension $k \geq 2$ and $f : X \rightarrow X$ a surjective holomorphic endomorphism. Let $1 \leq p \leq k$ be an integer such that $d_{p-1} < d_p$. Then, there exists a proper analytic subset E invariant under f such that for a positive closed (p, p) -current S smooth in a neighborhood of E , we have*

$$d_p^{-n}(f^n)^*(S - \alpha_S) \rightarrow 0$$

exponentially fast in the sense of currents where α_S is a smooth closed form of bidegree (p, p) in the de Rham cohomology class of S .

The equidistribution of inverse images of an analytic subset or a positive closed current under holomorphic/meromorphic endomorphisms has been much studied on complex projective space \mathbb{P}^k . To list a few, [34],[30] for $k = 1$; for general $k \geq 2$ and $p = k$, [28], [8], [21]; for $k = 2$ and $p = 1$, [26],[27] (see also [33]); for general $k \geq 2$, [29],[36],[37],[33],[17],[38] (see also [35]); for $1 \leq p \leq k$, see [1], [2], [4], [3]

However, it has not been much studied on compact Kähler manifolds. For a meromorphic map and for $p = k$, see [14]. For a holomorphic endomorphism, in [5], non-pluripolar products were considered. For a holomorphic automorphism, see [22], [9]. Notice that our theorem works for general bidegrees and also for holomorphic automorphisms as well.

In [1], the following was proved:

Date: May 2, 2024.

Theorem 1.2. *Let $f : \mathbb{P}^k \rightarrow \mathbb{P}^k$ be a holomorphic endomorphism of degree $d \geq 2$. Let T^p denote the Green (p, p) -current associated with f on \mathbb{P}^k . Then there is a proper invariant analytic subset E for f such that $d^{-pn}(f^n)^*(S)$ converges to T^p exponentially fast in the sense of currents for every positive closed (p, p) -current S of mass 1 which is smooth near E .*

One important property of a holomorphic endomorphism f of \mathbb{P}^k is that the graph of f^{-1} is a holomorphic correspondence. Intuitively, if a point is not in the Julia set, its forward images accumulate near the attracting set under iteration. So, if a point is not trapped in the attracting set, iterates push its inverse images away from the attracting set to the boundary of the Fatou set. Since f^{-1} is a holomorphic correspondence, once the initial current has no mass on the attracting set, then its inverse images never have mass on the set due to the work in [18]. The condition in Theorem 1.2 is a sufficient condition that if the initial current has no mass on the set, then there is no mass charged on the attracting set and therefore, all the mass move towards the Julia set. Theorem 1.2 is a quantitative explanation about it. Technically, this property appears in the availability of the Lojasiewicz type inequality as in Lemma 3.3 in [1]. Then, using the Lojasiewicz type inequality, we can control the influence from the set E , which appears in the Hölder continuity of the quasi-potential of $f_*(\omega)$.

On a general compact Kähler manifold, Lemma 4.7 in [15] guarantees that the same is true and motivates us to generalize the work in [1] to general compact Kähler manifolds.

Lemma 1.3 (Lemma 4.7 in [15]). *Let f be a surjective holomorphic map from X to X . Then f^{-1} is a holomorphic correspondence.*

However, there are many differences between projective space and compact Kähler manifold. Firstly, general compact Kähler manifolds may not have many automorphisms and therefore, it is not clear that positive closed currents can be approximated by smooth positive closed currents. Secondly, the Green potential may not be negative. In addition, the cohomology groups of a general compact Kähler manifold are not simple compared to those of a complex projective space. Lastly, the existence of the Green current is not clear. These are the main difficulties for the work to be done on general compact Kähler manifolds.

To resolve these obstacles, the following strategy is used. For the lack of symmetry, we quantify the approximation theorem of positive closed currents introduced by Dinh-Sibony in [16] and [22] so as to control the C^1 -norm of approximating smooth closed currents. Compared to the case of \mathbb{P}^k , approximating smooth closed currents may not be positive. So, we need additional estimates in the form of (semi-)regular transforms. In this work, we use the Green potential kernel induced from the work by Bost-Gillet-Soulé in [7]. In general, the Green potential is not negative and therefore, we had to extend the estimates for the Green quasi-potential in [20] and [1] to (semi-)regular transforms of positive closed currents. Since the cohomology groups may be complicated, the mass of a positive closed current may not behave as nicely as in the case of \mathbb{P}^k and so, we normalize the inverse image of a positive closed current with the dynamical degree. Since the existence of the Green current is not clear, we rather observe the current $S - \alpha_S$ instead of S and later in Section 9, we consider sufficient conditions for the Green current to exist. Summarizing the discussion in Section 9, we obtain the following:

Theorem 1.4. *Let (X, ω) be a compact Kähler manifold of dimension $k \geq 2$ and $f : X \rightarrow X$ a surjective holomorphic endomorphism such that that $d_{p-1} < d_p$ and d_p is a simple eigenvalue of $f^* : H^{p,p}(X, \mathbb{C}) \rightarrow H^{p,p}(X, \mathbb{C})$ and that other eigenvalues of f^* on $H^{p,p}(X, \mathbb{C})$ have modulus strictly less than d_p . Then, the limit $T_p^+ := \lim_{n \rightarrow \infty} d_p^{-n}(f^n)^*\omega^p$ exists and there exists a proper analytic subset E invariant under f such that for every positive closed (p, p) -current S smooth in a neighborhood of E , we have*

$$d_p^{-n}(f^n)^*S \rightarrow c_S T_p^+$$

exponentially fast in the sense of currents where $T_p^+ = \lim_{n \rightarrow \infty} d_p^{-n}(f^n)^* \omega^p$, $\{\cdot\}$ denotes the de Rham cohomology class, and $c_S = \lim_{n \rightarrow \infty} \frac{(f^n)^* \{S\} \sim \{\omega^{k-p}\}}{d_p^n \{T_p^+\} \sim \{\omega^{k-p}\}}$. In particular, if the action of f is simple on cohomology and p is the order of the main dynamical degree, $c_S = \frac{\langle S, T^- \rangle}{\langle \omega^p, T^- \rangle}$.

Corollary 1.5. *Assume f and E as in Theorem 1.4. Then, if an analytic subset H of pure codimension (p, p) does not meet E , then*

$$d_p^{-n}(f^n)^*[H] \rightarrow c_H T_p^+$$

exponentially fast in the sense of currents where $[H]$ denotes the current of integration on H and $c_H = \lim_{n \rightarrow \infty} \frac{(f^n)^* \{[H]\} \sim \{\omega^{k-p}\}}{d_p^n \{T_p^+\} \sim \{\omega^{k-p}\}}$.

The condition in Theorem 1.4 means a version of hyperbolicity so that there is only one maximally increasing direction in cohomology and all the others are decreasing directions. If $X = \mathbb{P}^k$, then $d_s = d^s$ and Theorem 1.4 implies Theorem 1.2.

One remark is that while our primary interest is on holomorphic endomorphisms, Theorem 1.1 applies to holomorphic automorphisms as well. Based on the proof, if f is a holomorphic automorphism of X , then $E = \emptyset$, which means that every positive closed current satisfies Theorem 1.1. One representative example of this is a holomorphic automorphism of a compact Kähler manifold with positive entropy. Concerning Theorem 1.4, in the case of holomorphic automorphisms, [9] obtained finer results without speed of convergence. See also [22].

Another remark is that in this context, it is reasonable to consider a version of the Dinh-Sibony conjecture in the following sense and our work partially answers the non-intersecting case.

Question 1.6 (See Conjecture 1.4 in [19]). *Let (X, ω) be a compact Kähler manifold of dimension $k \geq 2$ and $f : X \rightarrow X$ a surjective holomorphic endomorphism with simple action on cohomology. Let p be the order of the main dynamical degree and T^\pm the associated Green currents. Then $d_p^{-n}(f^n)^*[H]$ converges to $c_H T^+$ for every analytic subset H of X of pure dimension $k - p$ which is generic, where $c_H = \frac{\langle [H], T^- \rangle}{\langle \omega^p, T^- \rangle}$. Here, H is generic if either $H \cap \mathcal{E} = \emptyset$ or $\text{codim } H \cap \mathcal{E} = p + \text{codim } \mathcal{E}$ for any irreducible component \mathcal{E} of every totally invariant proper analytic subset of X .*

Notation. We denote by Φ_n and Ψ_n the hypersurface of the critical points and that of the critical values of f^n , respectively for $n = 1, 2, \dots$. For an analytic subset V of X , $[V]$ means the current of integration over V and $\text{multi}_x V$ means the multiplicity of V at $x \in V$ as an analytic subset. For a positive closed current R , $\nu(x, R)$ means the Lelong number R at $x \in X$.

The distance $\text{dist}(\cdot, \cdot)$ on a compact Kähler manifold means the distance with respect to the natural metric associated with the Kähler form. For a set $A \subset X$ and a constant $\theta > 0$, the θ -neighborhood A_θ of A denotes the set of points $x \in X$ such that $\text{dist}(x, A) < \theta$. The norms $\|\cdot\|_\infty$, $\|\cdot\|_{C^\alpha}$ and $\|\cdot\|_{L^\alpha}$ (or $\|\cdot\|_{\infty, U}$, $\|\cdot\|_{C^\alpha, U}$ and $\|\cdot\|_{L^\alpha, U}$ for a subset U of X) of a function or a form are the norms of the function or the sum of the usual corresponding norms of its coefficients (on U) with respect to a fixed finite atlas of X , respectively.

Acknowledgments. The research of the author was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. RS-2023-00250685).

2. PRELIMINARIES

2.1. Currents. For the basics of currents, we refer the reader to [10]. In this section, we introduce some notations that we will use.

Let s be an integer such that $1 \leq s \leq k$. If S is a positive or negative (s, s) -current on X , we define the mass of S by

$$\|S\| := |\langle S, \omega^{k-s} \rangle|.$$

Let \mathcal{C}_s denote the cone of positive closed (s, s) -currents on X , \mathcal{D}_s the real vector space generated by \mathcal{C}_s and \mathcal{D}_s^0 the space of currents $S \in \mathcal{D}_s$ such that $\{S\} = 0$ in $H^{s,s}(X, \mathbb{R})$. The duality between the cohomology groups implies that if S is a current in \mathcal{C}_s , its mass depends only on the class $\{S\}$ in $H^{s,s}(X, \mathbb{R})$. We define the $*$ -norm $\|\cdot\|_*$ on \mathcal{D}_s by

$$\|S\|_* := \min(\|S^+\| + \|S^-\|),$$

where the minimum is taken over $S^\pm \in \mathcal{C}_s$ such that $S = S^+ - S^-$. A subset in \mathcal{D}_s is said to be $*$ -bounded if it is bounded with respect to the $*$ -norm. We will consider the following $*$ -topology on \mathcal{D}_s and \mathcal{D}_s^0 . We say that S_n converges to S in \mathcal{D}_s if $S_n \rightarrow S$ weakly as currents and if the set $\{\|S_n\|_*\}$ is bounded. Note that if we restrict the $*$ -topology to a $*$ -bounded subset of \mathcal{D}_s , then it coincides with the usual weak topology on the space of currents. According to [22], smooth forms are dense in \mathcal{D}_s and \mathcal{D}_s^0 with respect to the $*$ -topology. We denote by $\widetilde{\mathcal{D}}_s$ and $\widetilde{\mathcal{D}}_s^0$ the subsets of smooth forms in \mathcal{D}_s and \mathcal{D}_s^0 , respectively.

We will also work on the space of L^1 functions which can be written as a difference of two quasi-plurisubharmonic functions on X . On that space, we will use the DSH-norm which will be denoted by $\|\cdot\|_{\text{DSH}}$ and defined by

$$\|f\|_{\text{DSH}} := \|f\|_{L^1} + \|dd^c f\|_*.$$

There are also some other natural norms and distances on \mathcal{D}_s which are closely related to the weak topology. For $\alpha > 0$, if S and S' are currents in \mathcal{D}_s , we define

$$\|S\|_{C^{-\alpha}} := \sup_{\|\varphi\|_{C^\alpha} \leq 1} |\langle S, \varphi \rangle| \quad \text{and} \quad \text{dist}_\alpha(S, S') := \|S - S'\|_{C^{-\alpha}}$$

where the supremum is taken over the set of the smooth test $(k-s, k-s)$ -forms φ on X with $\|\varphi\|_{C^\alpha} \leq 1$. Observe that $\|\cdot\|_{C^{-\alpha}} \lesssim \|\cdot\|_*$ for every $\alpha > 0$.

Thanks to the standard theory of interpolation between Banach spaces, we have

Proposition 2.1 (Section 2.1 in [20]). *Let α and α' be strictly positive real numbers with $\alpha < \alpha'$. Then on any $*$ -bounded subset of \mathcal{D}_s , the topology induced from dist_α or from $\text{dist}_{\alpha'}$ coincides with the weak topology. Moreover, on any $*$ -bounded subset of \mathcal{D}_s , there is a constant $c_{\alpha, \alpha'} > 0$ such that*

$$\text{dist}_{\alpha'} \leq \text{dist}_\alpha \leq c_{\alpha, \alpha'} [\text{dist}_{\alpha'}]^{\alpha/\alpha'}.$$

In particular, a function on a $$ -bounded subset of \mathcal{D}_s is Hölder continuous with respect to dist_α if and only if it is Hölder continuous with respect to $\text{dist}_{\alpha'}$.*

2.2. (semi-)regular transforms. We recall (semi-)regular transforms of currents in [22].

Consider the compact Kähler manifold $\mathfrak{X} := X \times X$. Let $\pi_i : \mathfrak{X} \rightarrow X$ denote the canonical projection on its factor for $i = 1, 2$. Then, $\omega_{\mathfrak{X}} := \pi_1^* \omega + \pi_2^* \omega$ is a natural Kähler form on \mathfrak{X} . Let $\pi : \widehat{\mathfrak{X}} \rightarrow \mathfrak{X}$ be the blow-up of \mathfrak{X} along the diagonal submanifold Δ in $\mathfrak{X} = X \times X$ and let $\widehat{\Delta} := \pi^{-1}(\Delta)$ denote the exceptional hypersurface. We define $\Pi_i := \pi_i \circ \pi$ for $i = 1, 2$. Then, Π_i and its restriction to $\widehat{\Delta}$ are both submersions for $i = 1, 2$. By a theorem of Blanchard in [6], $\widehat{\mathfrak{X}}$ is a compact Kähler manifold. We fix a Kähler form $\omega_{\widehat{\mathfrak{X}}}$ on $\widehat{X} \times \widehat{X}$ throughout the article. For later use in Section 3, we assume that $\omega_{\widehat{\mathfrak{X}}}$ is normalized as follows. The current $\pi_*([\widehat{\Delta}] \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1})$ has support in Δ and is a positive closed current of bidimension (k, k) . Hence, by the support theorem, $\pi_*([\widehat{\Delta}] \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1})$ is a positive constant multiple of $[\Delta]$. By multiplying a proper positive constant to $\omega_{\widehat{\mathfrak{X}}}$, we may assume that $\pi_*([\widehat{\Delta}] \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}) = [\Delta]$.

Definition 2.2. Let $0 \leq q \leq k$ be an integer. Let \mathcal{Q} be a form of bidegree (q, q) on $\widehat{\mathcal{X}}$ which is smooth outside $\widehat{\Delta}$ and such that

$$|\mathcal{Q}| \lesssim -\log \text{dist}(\cdot, \widehat{\Delta}) \quad \text{and} \quad |\nabla \mathcal{Q}| \lesssim \text{dist}(\cdot, \widehat{\Delta})^{-1}$$

near $\widehat{\Delta}$. Let s be an integer such that $k - q \leq s \leq k$. A linear mapping $\mathcal{L}^{\mathcal{Q}}$ on the space of currents of bidegree (s, s) on X to the space of currents $(s + q - k, s + q - k)$ on X defined by

$$\mathcal{L}^{\mathcal{Q}}(S) := (\Pi_2)_*(\Pi_1^*(S) \wedge \mathcal{Q})$$

is called a semi-regular transform of bidegree $(q - k, q - k)$ associated with the form \mathcal{Q} .

If \mathcal{Q} is smooth, then the transform $\mathcal{L}^{\mathcal{Q}}$ is said to be regular. If \mathcal{Q} is positive, then the transform $\mathcal{L}^{\mathcal{Q}}$ is said to be positive. If \mathcal{Q} is closed, then the transform $\mathcal{L}^{\mathcal{Q}}$ is said to be closed.

Here, $|\mathcal{Q}|$ and $|\nabla \mathcal{Q}|$ mean the sum of the coefficients of \mathcal{Q} and the sum of the estimates of their gradients with respect to a fixed finite atlas of $\widehat{\mathcal{X}}$, respectively.

Remark 2.3. A positive semi-regular transform maps positive currents to positive currents. A closed semi-regular transform $\mathcal{L}^{\mathcal{Q}}$ maps closed currents to closed currents and satisfies $\mathcal{L}^{\mathcal{Q}}(dd^c S) = dd^c \mathcal{L}^{\mathcal{Q}}(S)$ for every current S .

Young's inequality (or the Hölder inequality) gives the following proposition.

Proposition 2.4. [Lemma 2.1 in [16] or Proposition 2.3.2 in [21]] Any semi-regular transform can be extended to a linear continuous operator from the space of currents of order 0 to the space of $L^{1+1/k}$ -forms. It defines a linear continuous operator from the space of L^α -forms, $\alpha \geq 1$, to the space of $L^{\alpha'}$ -forms where α' is given by $(\alpha')^{-1} + 1 = \alpha^{-1} + (1 + 1/k)^{-1}$ if $\alpha < k + 1$ and $\alpha' = \infty$ if $\alpha \geq k + 1$. It also defines a linear continuous operator from the space of L^∞ -forms to the space of C^1 -forms.

2.3. Superpotentials. For the superpotential on a compact Kähler manifold, we refer the reader to [22]. See also [20] for the theory on complex projective space.

Let $\beta = \{\beta_1, \dots, \beta_{h_s}\}$ with $h_s = \dim H^{s,s}(X, \mathbb{R})$ be a fixed family of real smooth closed (s, s) -forms such that the family of classes $\{\beta\} := \{\{\beta_1\}, \dots, \{\beta_{h_s}\}\}$ is a basis of $H^{s,s}(X, \mathbb{R})$. We can also find another family $\beta^* = \{\beta_1^*, \dots, \beta_{h_s}^*\}$ of real smooth closed $(k - s, k - s)$ -forms so that $\{\beta^*\} := \{\{\beta_1^*\}, \dots, \{\beta_{h_s}^*\}\}$ is a basis of $H^{k-s, k-s}(X, \mathbb{R})$ and is the dual basis of $\{\beta\}$ with respect to the cup-product \smile .

Let R be a current in \mathcal{D}_{k-s+1}^0 and U_R a potential of R , that is, a $(k - s, k - s)$ -current such that $dd^c U_R = R$. Adding to U_R a suitable combination of β_i^* 's allows us to assume that $\langle U_R, \beta_i \rangle = 0$ for $i = 1, \dots, h_s$. We say that U_R is β -normalized.

Definition 2.5. Let β and β^* be given families of real smooth closed forms of bidegree (s, s) and $(k - s, k - s)$ as above, respectively. Let S be a current in \mathcal{D}_s . The β -normalized superpotential \mathcal{U}_S of S is the following function defined on smooth forms R in \mathcal{D}_{k-s+1}^0 by

$$\mathcal{U}_S(R) := \langle S, U_R \rangle,$$

where U_R is an β -normalized smooth potential of R . We say that S has a bounded superpotential if \mathcal{U}_S is bounded on each $*$ -bounded subset of $\widehat{\mathcal{D}}_{k-s+1}^0$. We say that S has a continuous superpotential if \mathcal{U}_S can be extended to a continuous function on \mathcal{D}_{k-s+1}^0 . Here, the topology is with respect to the norm $\|\cdot\|_{C^{-\alpha}}$ for some $\alpha > 0$. In this case, the extension is also denoted by \mathcal{U}_S and is also called a superpotential of S . We say that S has a Hölder continuous superpotential if \mathcal{U}_S is continuous and Hölder continuous on \mathcal{D}_{k-s+1}^0 with respect to the norm $\|\cdot\|_{C^{-\alpha}}$ for some $\alpha > 0$.

Remark 2.6. *The definition of the β -normalized superpotential \mathcal{U}_S of S is independent of the choice of β^* . If S belongs to \mathcal{D}_s^0 , the notion of superpotential is independent of the choice of the family β . So, when we are dealing with superpotentials of currents $S \in \mathcal{D}_s^0$, we will not specify the families β and β^* .*

We recall the Green potential kernel and the Green potential of S for $S \in \mathcal{D}_s^0$ in [22], which are useful in computing superpotentials.

The integration on the diagonal submanifold Δ of \mathfrak{X} defines a positive closed (k, k) -current $[\Delta]$. By the Künneth formula, we have a canonical isomorphism

$$H^{k,k}(\mathfrak{X}, \mathbb{C}) \simeq \sum_{0 \leq i \leq k} H^{i, k-i}(X, \mathbb{C}) \otimes H^{k-i, i}(X, \mathbb{C}).$$

Then, $[\Delta]$ is cohomologous to a smooth real closed (k, k) -form α_Δ which is a finite combination of forms of type $\pi_1^*(\psi) \wedge \pi_2^*(\psi')$ where ψ and ψ' are closed forms on X of bidegree $(i, k-i)$ and $(k-i, i)$, respectively. So, α_Δ satisfies $d_x \alpha_\Delta = d_y \alpha_\Delta = 0$. Replacing $\alpha_\Delta(x, y)$ by $[\alpha_\Delta(x, y) + \alpha_\Delta(y, x)]/2$, we may assume that α_Δ is symmetric, i.e. invariant under the involution $(x, y) \rightarrow (y, x)$ where (x, y) denotes the coordinates of \mathfrak{X} .

According to [7], there is a real smooth closed $(k-1, k-1)$ -form η on $\widehat{\mathfrak{X}}$ such that $\pi^*(\alpha_\Delta)$ is cohomologous to $[\widehat{\Delta}] \wedge \eta$, where $[\widehat{\Delta}]$ is the positive closed $(1, 1)$ -current of integration on $\widehat{\Delta}$. Hence, $\pi_*([\widehat{\Delta}] \wedge \eta)$ is cohomologous to α_Δ and therefore to $[\Delta]$. Since the map $(x, y) \rightarrow (y, x)$ induces an involution on $\widehat{\mathfrak{X}}$, we can also choose η symmetric with respect to this involution.

Let $\alpha_{\widehat{\Delta}}$ be a real closed $(1, 1)$ -form on $\widehat{\mathfrak{X}}$, which is cohomologous to $[\widehat{\Delta}]$. We can choose $\alpha_{\widehat{\Delta}}$ symmetric. There is a quasi-plurisubharmonic (q-psh for short) function u on $\widehat{\mathfrak{X}}$ such that $dd^c u = [\widehat{\Delta}] - \alpha_{\widehat{\Delta}}$. This function is necessarily symmetric. Subtracting from u a constant, we may assume that $u < -2$.

Choose a real smooth $(k-1, k-1)$ -form $\delta_{\widehat{\mathfrak{X}}}$ on $\widehat{\mathfrak{X}}$ such that $dd^c \delta_{\widehat{\mathfrak{X}}} = \pi^*(\alpha_\Delta) - \alpha_{\widehat{\Delta}} \wedge \eta$ where α_Δ , $\alpha_{\widehat{\Delta}}$ and η are as above. We can choose $\delta_{\widehat{\mathfrak{X}}}$ to be symmetric. Let \mathcal{L}^K be the semi-regular transform associated with the form $K := u\eta - \delta_{\widehat{\mathfrak{X}}}$. The following result was obtained in Proposition 2.1 in [22].

Proposition 2.7. *Let S be a current in \mathcal{D}_s^0 with $s \geq 1$. Then $U_S := \mathcal{L}^K(S)$ is a potential of S , that is, $dd^c U_S = S$ in the sense of currents. Moreover, we have*

$$\|U_S\|_{L^{1+1/k}} \leq c \|S\|_*$$

for some constant $c > 0$ independent of S .

We will call $U_S := \mathcal{L}^K(S)$ the Green potential of S . Notice that K is not necessarily positive. There exists a constant $m_K > 0$ such that $\eta + m_K \omega_{\widehat{\mathfrak{X}}}^{k-1}$ and $\delta_{\widehat{\mathfrak{X}}} + m_K \omega_{\widehat{\mathfrak{X}}}^{k-1}$ are both strictly positive. We define the forms K_\pm to be

$$K_+ := u(\eta + m_K \omega_{\widehat{\mathfrak{X}}}^{k-1}) - (m_K \omega_{\widehat{\mathfrak{X}}}^{k-1}) \quad \text{and} \quad K_- := u(m_K \omega_{\widehat{\mathfrak{X}}}^{k-1}) - (\delta_{\widehat{\mathfrak{X}}} + m_K \omega_{\widehat{\mathfrak{X}}}^{k-1}).$$

Both K_\pm are strictly negative and $\mathcal{L}^K = \mathcal{L}^{K_+} - \mathcal{L}^{K_-}$. In particular, the transforms \mathcal{L}^{K_\pm} associated with the forms K_\pm are negative.

3. REGULARIZATION OF POSITIVE CLOSED CURRENTS

The regularization of positive closed currents in this section was introduced in [16] and [22]. The main purpose of this section is to study its quantitative aspects.

Let u be the q-psh function on X in Subsection 2.3. Let χ be a smooth convex increasing function on $\mathbb{R} \cup \{-\infty\}$ such that $\chi(t) = t$ for $t \geq 0$, $\chi(t) = -1$ for $t \leq -2$ and $0 \leq \chi' \leq 1$. For $\theta \in \mathbb{C}$, we define

$$\chi_\theta(t) := \chi(t - \log |\theta|) + \log |\theta| \quad \text{and} \quad u_\theta := \chi_\theta(u).$$

Then, $u_\theta = u_{|\theta|}$ and u_θ decreasingly converges to u as $|\theta| \rightarrow 0$. Let $m_{\widehat{\Delta}} > 0$ be sufficiently large so that $m_{\widehat{\Delta}}\omega_{\widehat{\mathfrak{X}}} - \alpha_{\widehat{\Delta}}$ is positive. Then, we have

$$dd^c u_\theta = (\chi''_\theta \circ u)du \wedge d^c u + (\chi'_\theta \circ u)dd^c u = -(\chi'_\theta \circ u)\alpha_{\widehat{\Delta}} \geq -m_{\widehat{\Delta}}\omega_{\widehat{\mathfrak{X}}}.$$

So, for $\theta \in \mathbb{C}$, the smooth closed $(1, 1)$ -current $\alpha_{\widehat{\Delta}} + dd^c u_\theta$ can be written as a difference of two smooth positive closed $(1, 1)$ -currents as $\alpha_{\widehat{\Delta}} + dd^c u_\theta = (m_{\widehat{\Delta}}\omega_{\widehat{\mathfrak{X}}} + dd^c u_\theta) - (m_{\widehat{\Delta}}\omega_{\widehat{\mathfrak{X}}} - \alpha_{\widehat{\Delta}})$.

Recall that $\pi_*([\widehat{\Delta}] \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}) = [\Delta]$. Let $\theta \in \mathbb{C}$ be such that $|\theta| < 1$. For notational convenience, we write

$$\mathcal{L}_\theta^+ := \mathcal{L}^{(m_{\widehat{\Delta}}\omega_{\widehat{\mathfrak{X}}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}}, \quad \mathcal{L}^- := \mathcal{L}^{(m_{\widehat{\Delta}}\omega_{\widehat{\mathfrak{X}}} - \alpha_{\widehat{\Delta}}) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}} \quad \text{and} \quad \mathcal{L}_\theta := \mathcal{L}^{(\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}}.$$

Then, we obviously have $\mathcal{L}_\theta = \mathcal{L}_\theta^+ - \mathcal{L}^-$ and $\mathcal{L}_0 = \text{id}$ on smooth forms (see [16, p.486]). The transforms \mathcal{L}_θ^+ , \mathcal{L}^- and \mathcal{L}_θ are all positive closed and of bidegree $(0, 0)$.

Our regularizing kernel $(\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}$ is slightly different from the one in [22] but it works in the same way in [22, Lemma 2.4.6]. Also, in [22, Lemma 2.4.6], we do not actually need the closedness of a smooth current. So, we get the following lemma.

Lemma 3.1. *Let φ be a smooth form of bidegree $(k-s, k-s)$. Then, for every $\theta \in \mathbb{C}$, $\mathcal{L}_\theta(\varphi)$ is smooth and*

$$\|\mathcal{L}_\theta(\varphi) - \varphi\|_\infty \leq c|\theta|\|\varphi\|_{C^1}$$

where $c > 0$ is a constant independent of φ and θ .

From [22, Lemma 2.4.1], we can deduce the following lemma. The constant $c > 0$ below comes from the description of the support of $dd^c u_\theta$ in terms of θ in the lemma.

Lemma 3.2. *Let $\theta \in \mathbb{C}^*$ be sufficiently small. Then, there exist constants $c_1, c_2 > 0$ independent of θ such that for any subset U of X and for any form S with C^1 -coefficients and $\text{supp} S \Subset U$, $\mathcal{L}_\theta(S)$ is C^1 with compact support in $U_{c_1\theta}$ such that*

$$\|\mathcal{L}_\theta(S)\|_{C^1, U_{c_1\theta}} = \|\mathcal{L}^{(\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}}(S)\|_{C^1, U_{c_1\theta}} \leq c_2\|S\|_{C^1, U}.$$

Proof. The uniform norm estimate is quite straightforward. So, we omit it and focus on the estimate of its gradient. Since S has C^1 coefficients, we have

$$\begin{aligned} \mathcal{L}^{(\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}}(S) &= (\Pi_2)_*(\Pi_1^*(S) \wedge (\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}) \\ &= (\pi_2)_*(\pi_1^*(S) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1})). \end{aligned}$$

According to [22, Lemma 2.4.1], there exists a constant $c > 0$ such that the support of $\pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1})$ is in $(\Delta)_{c|\theta|}$ for all θ 's with sufficiently small $|\theta|$. Also, notice that the support of $\pi_*(dd^c u_\theta)$ is away from Δ . By use of a partition of unity, we may assume that the support of S and the support of $\mathcal{L}_\theta(S)$ belongs to the same coordinate neighborhood. Let $P \in X$ be point. Let $(x, y) = (x_1, \dots, x_k, y_1, \dots, y_k) \in X \times X$ be local coordinates in a neighborhood of a point $(P, P) \in X \times X$ such that $(0, \dots, 0, 0, \dots, 0)$ corresponds to (P, P) . Let P' be a point in the support of $\mathcal{L}_\theta(S)$ and $y' = (y'_1, \dots, y'_k)$ its coordinates.

Let $v = (v_1, \dots, v_k) \in \mathbb{C}^k$ be such that $|v_1|^2 + \dots + |v_k|^2 = 1$. Then, we consider the form of

$$\begin{aligned} & \frac{1}{h} [(\pi_2)_*(\pi_1^*(S) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(y' + hv) \\ & \quad - (\pi_2)_*(\pi_1^*(S) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(y')] \\ & = \frac{1}{h} \int_x [(\pi_1^*(S)(x) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y' + hv) \\ & \quad - \pi_1^*(S)(x) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y')] \end{aligned}$$

Note that since S has C^1 coefficients,

$$\begin{aligned} & \lim_{h \rightarrow \infty} \frac{1}{h} \int_x [(\pi_1^*(S)(x + hv) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y') \\ & \quad - \pi_1^*(S)(x) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y')] \end{aligned}$$

is a continuous form whose uniform norm is bounded by $\|S\|_{C^1}$ up to a constant multiple independent of θ . Indeed, this is just the transform of the derivative of S in the direction of v , which is a continuous form. So, it is enough to estimate

$$\begin{aligned} & \frac{1}{h} \int_x [(\pi_1^*(S)(x + hv) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y') \\ & \quad - \pi_1^*(S)(x) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y' + hv)] \\ & = \frac{1}{h} \int_x [(\pi_1^*(S)(x + hv) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y') \\ & \quad - \pi_1^*(S)(x + hv) \wedge \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x + hv, y' + hv)] \\ (3.1) \quad & = \frac{1}{h} \int_x [(\pi_1^*(S)(x + hv) \wedge [\pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y') \\ & \quad - \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x + hv, y' + hv)]] \end{aligned}$$

for all h with sufficiently small $|h|$ and show that this quantity is bounded by $\|S\|_\infty$ up to a constant independently of θ . The first equality comes from the change of the variable $x \rightarrow x - hv$ inside the integral. Since S is bounded, it suffices to consider the estimate of

$$\frac{1}{h} [\pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x, y') - \pi_*((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1}))(x + hv, y' + hv)]$$

After shrinking the neighborhood of (P, P) if necessary, we change variables $(x, y) \rightarrow (x, y - x)$ and denote by $z = y - x$. Then, in the neighborhood of $(P, 0)$, $\Delta = \{z = 0\}$ and the operator $(\pi_2)_*$ is the integration with respect to the x -variable. Since the support of $\pi_*(dd^c u_\theta)$ is away from Δ , it is enough to consider the case where one of z_i 's is not equal to 0. Without loss of generality, we only consider the open set $z_1 \neq 0$. Then, once again, we change variables $(x, z) \rightarrow (x, z_1, z_2/z_1, \dots, z_k/z_1)$ in that neighborhood. We denote by $w_1 = z_1$ and $w_i = z_i/z_1$ for $i = 2, \dots, k$. With respect to the coordinates (x, w) , the above quantity becomes

$$(3.2) \quad \frac{1}{h} [((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1})(x, w') - ((\alpha_{\widehat{\Delta}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathfrak{X}}}^{k-1})(x + hv, w')]$$

where $w'_1 = y'_1 - x_1$ and $w'_i = (y'_i - x_i)/(y'_1 - x_1)$ for $i = 2, \dots, k$. With respect to the coordinates (x, w) , the function u can be written as $u = \log|w_1| + \psi$ for a smooth function ψ of z and w . Then, we have

$$dd^c u_\theta = dd^c[\chi_\theta(u)] = \chi''_\theta(u) d(\log|w_1| + \psi) \wedge d^c(\log|w_1| + \psi) + \chi'_\theta(u) dd^c(\log|w_1| + \psi).$$

Hence, as $h \rightarrow 0$, we get some extra derivatives of ψ and $\chi_\theta(u)$ with respect to x , which are bounded independently of θ . So, by the argument used in [22, Lemma 2.4.1], |(3.2)| is bounded by $1/|\theta|^2$ up to a constant multiple independent of θ . Given y' , due to the smallness of the support of $dd^c(\alpha_{\widehat{\Delta}} + dd^c u_\theta)$, we integrate the bounded integrand on $|x_1 - y'_1| < c'|\theta|$ for some $c' > 0$

independent of θ , which gives us a value of order $|\theta|^2$ and cancels the factor $1/|\theta|^2$. So, the value |(3.1)| is bounded by $\|S\|_\infty$ up to a constant multiple independent of θ and S . \square

The above two lemmas imply the following:

Proposition 3.3. *Let φ be a smooth test form of bidegree $(k-s, k-s)$. Then, $\|\mathcal{L}_\theta^{k+2}(\varphi) - \varphi\|_\infty$ converges to 0 as $\theta \rightarrow 0$.*

Proof. It is enough to observe that

$$\mathcal{L}_\theta^{k+2}(\varphi) - \varphi = \sum_{i=0}^{k+1} \mathcal{L}_\theta(\mathcal{L}_\theta^i(\varphi)) - \mathcal{L}_\theta^i(\varphi)$$

and that $\|\mathcal{L}_\theta^i(\varphi)\|_{C^1}$ is uniformly bounded by $\|\varphi\|_{C^1}$ up to a constant independent of θ and $i = 0, \dots, k+1$. Then, a previous lemma completes the proof. \square

For $S \in \mathcal{D}_s$, we define

$$S_\theta := \mathcal{L}_\theta \circ \dots \circ \mathcal{L}_\theta(S) = \mathcal{L}_\theta^{k+2}(S).$$

Proposition 2.4 implies that S_θ is a current with C^1 -coefficients. According to [16], S_θ belongs to the same cohomology class as S does. This regularization is essentially the same as the one in [16] but slightly different from the one in [22]. Since currents in \mathcal{D}_s are of order 0, Proposition 3.3 implies the convergence $S_\theta \rightarrow S$ in the sense of currents

Corollary 3.4. *For each current $S \in \mathcal{D}_s$, the sequence (S_θ) of C^1 -smooth currents converges to S in the sense of currents.*

Note that S_θ is not positive in general. However, the regularization S_θ can be written as $S_\theta = \sum_i (-1)^{\text{sgn}(\mathcal{L}_{i,1}, \dots, \mathcal{L}_{i,k+2})} \mathcal{L}_{i,1} \circ \dots \circ \mathcal{L}_{i,k+2}(S)$ where $\mathcal{L}_{i,j}$ is either \mathcal{L}_θ^+ or \mathcal{L}_θ^- and $\text{sgn}(\mathcal{L}_{i,1}, \dots, \mathcal{L}_{i,k+2})$ is the number of indices $\mathcal{L}_{i,j}$ such that $\mathcal{L}_{i,j} = \mathcal{L}_\theta^-$. Then, S_θ can be written as

$$S_\theta = S_\theta^+ - S_\theta^-$$

where S_θ^\pm are positive closed currents of bidegree (s, s) . Indeed, S_θ^+ (or S_θ^-) is just the sum of terms $(-1)^{\text{sgn}(\mathcal{L}_{i,1}, \dots, \mathcal{L}_{i,k+2})} \mathcal{L}_{i,1} \circ \dots \circ \mathcal{L}_{i,k+2}(S)$ with even (or odd) sgn . Proposition 2.4 implies that S_θ^\pm are of C^1 -coefficients. We give C^1 -estimates of S_θ^\pm in terms of θ .

Proposition 3.5. *There exists a constant $c_{\text{reg}} > 0$ such that for all θ with sufficiently small $|\theta|$ and for all $S \in \mathcal{C}_s$, we have*

$$\|S_\theta^\pm\| \leq c_{\text{reg}} \|S\| \quad \text{and} \quad \|S_\theta^\pm\|_{C^1} \leq c_{\text{reg}} |\theta|^{-(2k+2)(k+2)} \|S\|.$$

Lemma 3.6. *Let S be a current in \mathcal{C}_s . Then for all $\theta \in \mathbb{C}^*$ with sufficiently small $|\theta|$, the current $\mathcal{L}^{dd^c u_\theta \wedge \omega_{\widehat{\mathbb{X}}}^{k-1}}(S)$ is smooth and we have*

$$\|\mathcal{L}^{dd^c u_\theta \wedge \omega_{\widehat{\mathbb{X}}}^{k-1}}(S)\|_{C^1} \leq c |\theta|^{-(2k+2)} \|S\|$$

where $c > 0$ is a constant independent of S and θ .

Proof. The first assertion is from the observation that the support of $dd^c u_\theta$ is away from $\widehat{\Delta}$. So, we consider the second assertion. We have

$$\begin{aligned} \mathcal{L}^{dd^c u_\theta \wedge \omega_{\widehat{\mathbb{X}}}^{k-1}}(S) &= (\Pi_2)_*(\Pi_1^*(S) \wedge dd^c u_\theta \wedge \omega_{\widehat{\mathbb{X}}}^{k-1}) \\ &= (\pi_2)_*(\pi_1^*(S) \wedge \pi_*(dd^c u_\theta) \wedge \pi_*(\omega_{\widehat{\mathbb{X}}})^{k-1}). \end{aligned}$$

For the support of $dd^c u_\theta$ does not intersect $\widehat{\Delta}$. We estimate $\pi_*(dd^c u_\theta) \wedge \pi_*(\omega_{\widehat{\mathbb{X}}})^{k-1}$.

Since X is compact and all the derivatives of u_θ are uniformly bounded outside a neighborhood of Δ , it suffices to estimate in a neighborhood of a point $(P, P) \in \Delta \subset X \times X$ for a point $P \in X$. Let

$(x, y) = (x_1, \dots, x_k, y_1, \dots, y_k) \in X \times X$ be local coordinates in a neighborhood of a point $(P, P) \in X \times X$. After shrinking the neighborhood of (P, P) if necessary, we make a change of variables $(x, y) \rightarrow (x, y - x)$ and denote by $z = y - x$. Then, in the neighborhood, $\Delta = \{z = 0\}$ and the operator $(\pi_2)_*$ is the integration with respect to the x -variable.

Since the support of $\pi_*(dd^c u_\theta \wedge \omega_{\mathbb{X}}^{k-1})$ is away from Δ , it is enough to estimate the C^1 -norm of $\pi_*(dd^c u_\theta \wedge \omega_{\mathbb{X}}^{k-1})$ in a sector \mathfrak{S} of the form $|z_1| < \varepsilon$ and $|z_j| < 2|z_1|$ where $z = (z_1, \dots, z_k)$. Here, for simplicity, we just take $\varepsilon = 1$. We apply the same arguments to other sectors.

The following is a slight modification of [22, Lemma 2.4.1]. We again make a change of variables $w_1 = z_1$ and $w_j = z_j/z_1$ for $j \geq 2$ so that $(x, w_1, w_2, \dots, w_k)$ becomes the natural coordinates of $\pi^{-1}(\mathfrak{S}) \subset \bar{X} \times \bar{X}$ and $\bar{\Delta} = \{w_1 = 0\}$.

As in the proof of [22, Lemma 2.4.1], the support of $dd^c u_\theta$ sits inside $\{c_1|\theta| \leq |w_1|\}$ for some constant $c_1 > 0$ independent of θ . As previously, with respect to the coordinates (x, w) , the function u can be written as $u = \log|w_1| + \psi$ for a smooth function ψ of x and w . Then, we have

$$dd^c u_\theta = dd^c[\chi_\theta(u)] = \chi_\theta''(u)d(\log|w_1| + \psi) \wedge d^c(\log|w_1| + \psi) + \chi_\theta'(u)dd^c(\log|w_1| + \psi).$$

Observe that $\pi_*(dd^c u_\theta)$ is obtained from replacing w_1, \dots, w_k by $z_1, z_2/z_1, \dots, z_k/z_1$. So, we have

$$\begin{aligned} \pi_*(dd^c u_\theta) &= \chi_\theta''(u)d(\log|z_1| + \psi(x, z_1, z_2/z_1, \dots, z_k/z_1)) \wedge d^c(\log|z_1| + \psi(x, z_1, z_2/z_1, \dots, z_k/z_1)) \\ &\quad + \chi_\theta'(u)dd^c(\log|z_1| + \psi(x, z_1, z_2/z_1, \dots, z_k/z_1)). \end{aligned}$$

Also, each component of $\omega_{\mathbb{X}}^{k-1}$ is of the form $C(x, w)dx_I \wedge d\bar{x}_J \wedge dw_{I'} \wedge d\bar{w}_{J'}$ where $C(x, w)$ is a smooth function of x and w , and $dx_I \wedge d\bar{x}_J \wedge dw_{I'} \wedge d\bar{w}_{J'}$ is the wedge product of dx_1, \dots, dx_k and dw_1, \dots, dw_k of bidegree $(k-1, k-1)$. Again, $\pi_*(\omega_{\mathbb{X}}^{k-1})$ is obtained from replacing w_1, \dots, w_k by $z_1, z_2/z_1, \dots, z_k/z_1$. By direct computations, we have

$$d\left(\frac{y_i - x_i}{y_1 - x_1}\right) = \frac{1}{y_1 - x_1}d(y_i - x_i) - \frac{y_i - x_i}{(y_1 - x_1)^2}d(y_1 - x_1)$$

for each $i = 2, \dots, k$. After changing coordinates back to (x, y) from (x, z) , each coefficient of $\pi_*(\omega_{\mathbb{X}}^{k-1})$ is of the form $(y_1 - x_1)^{-l}(\bar{y}_1 - \bar{x}_1)^{-l'}C'(x, y_1 - x_1, (y_2 - x_2)/(y_1 - x_1), \dots, (y_k - x_k)/(y_1 - x_1))$ where $0 \leq l, l' \leq k-1$ and C' is a smooth function of $x, y_1 - x_1, (y_2 - x_2)/(y_1 - x_1), \dots, (y_k - x_k)/(y_1 - x_1)$. The support of $dd^c u_\theta$ lies in $\{c_1|\theta| \leq |z_1|\}$. The C^1 -norm of each coefficient of $\pi_*(\omega_{\mathbb{X}}^{k-1})$ with respect to y is bounded by $|\theta|^{-2k+1}$ up to a positive constant multiple independent of θ . The proof of [22, Lemma 2.4.1] implies that the C^1 -norm of the coefficients of $dd^c u_\theta$ on its support is bounded by $|\theta|^{-3}$ up to a positive constant multiple independent of θ . So, we get

$$\|(\pi_2)_*(\pi_1^*(S) \wedge \pi_*(dd^c u_\theta) \wedge \pi_*(\omega_{\mathbb{X}}^{k-1}))\|_{C^1} \lesssim |\theta|^{-(2k+2)} \|S\|$$

where the inequality \lesssim means \leq up to a constant multiple independent of S and θ . \square

Proof of Proposition 3.5. The first estimate is from the same argument used in [16, Theorem 1.1]. We consider the second inequality. Without loss of generality, we may assume that S is a positive closed (p, p) -current on X . Notice that $\mathcal{L}_\theta^+ = \mathcal{L}^{m_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^k} + \mathcal{L}^{dd^c u_\theta \wedge \omega_{\mathbb{X}}^{k-1}}$ and $\mathcal{L}^- = \mathcal{L}^{m_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^k} - \mathcal{L}^{\alpha_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^{k-1}}$. Hence, S_θ^{\pm} 's are the sum of $(k+2)$ -times compositions of $\mathcal{L}^{m_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^k}$, $\mathcal{L}^{\alpha_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^{k-1}}$ or $\mathcal{L}^{dd^c u_\theta \wedge \omega_{\mathbb{X}}^{k-1}}$. The terms containing only either $\mathcal{L}^{m_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^k}$ or $\mathcal{L}^{\alpha_{\mathbb{X}} \wedge \omega_{\mathbb{X}}^{k-1}}$ are independent of θ . So, the dominating part of the operator is $(\mathcal{L}^{dd^c u_\theta \wedge \omega_{\mathbb{X}}^{k-1}})^{\circ(k+2)}$ and the previous lemma implies the desired estimate. \square

Lemma 3.7. *Let S be a positive current of bidegree (p, p) . There exists a constant $c > 0$ such that for all θ with sufficiently small $|\theta|$, we have*

$$\|\mathcal{L}_\theta^+(S)\| \leq c(1 + |\theta|)\|S\|, \quad \|\mathcal{L}^-(S)\| \leq c\|S\| \quad \text{and} \quad \|\mathcal{L}_\theta(S)\|_* \leq c(2 + |\theta|)\|S\|$$

where the constant $c > 0$ is independent of S and θ .

Proof. From Lemma 3.1, we get

$$\langle \mathcal{L}_\theta^+(S), \omega^{k-p} \rangle = \langle \mathcal{L}_\theta(S) + \mathcal{L}^-(S), \omega^{k-p} \rangle = \langle S, \mathcal{L}_\theta(\omega^{k-p}) + \mathcal{L}^-(\omega^{k-p}) \rangle \leq c'(1 + |\theta|) \|S\|$$

and

$$\langle \mathcal{L}^-(S), \omega^{k-p} \rangle = \langle S, \mathcal{L}^-(\omega^{k-p}) \rangle \leq c'' \|S\|.$$

for some $c' > 0$ and $c'' > 0$. Using the definition of the $*$ -norm, we get the last inequality. \square

Lemma 3.8. *Let S_1 and S_2 be positive currents of bidegree (s_1, s_1) and (s_2, s_2) such that $s_1 + s_2 = k$ and $\text{supp}S_1 \cap \text{supp}S_2 = \emptyset$. Then, for all θ with sufficiently small $|\theta|$, we have*

$$\langle S_1, \mathcal{L}^\theta(S_2) \rangle = 0.$$

Proof. [22, Lemma 2.4.1] implies that the support of $\pi_*((\alpha_{\widehat{x}} + dd^c u_\theta) \wedge \omega_{\widehat{x}}^{k-1})$ uniformly shrinks to Δ as $\theta \rightarrow 0$. \square

Lemma 3.9. *Let S_1 and S_2 be positive currents of bidegree (s_1, s_1) and (s_2, s_2) such that $s_1 + s_2 = k$ and $\text{supp}S_1 \cap \text{supp}S_2 = \emptyset$. Then, we have*

$$|\langle S_1, \mathcal{L}^-(S_2) \rangle| \lesssim \text{dist}(\text{supp}S_1, \text{supp}S_2)^{2-2k} \|S_1\| \cdot \|S_2\|.$$

The inequality \lesssim means \leq up to a constant multiple independent of S_1 and S_2 .

Proof. The support of $\pi_1^*(S_1) \wedge \pi_2^*(S_2)$ is away from Δ . On its support, Lemma 3.1 in [16] implies that

$$\|\pi_* \omega_{\widehat{x}}^k\|_{\infty, \text{supp}\pi_1^*(S_1) \wedge \pi_2^*(S_2)} \lesssim \text{dist}(\text{supp}S_1, \text{supp}S_2)^{2-2k}.$$

Since S_1 and S_2 are positive, the mass of $\pi_1^*(S_1) \wedge \pi_2^*(S_2)$ is bounded by $\|S_1\| \cdot \|S_2\|$. \square

4. ANALYTIC (SUB)MULTIPLICATIVE COCYCLES

The notion of analytic (sub)multiplicative cocycles was first introduced by Favre [24], [25] and further studied by Dinh [12] and Gignac [31].

Let Z be an irreducible compact complex space of dimension l , not necessarily smooth. Let $g : Z \rightarrow Z$ be an open holomorphic map.

Definition 4.1. [Definition 1.1 in [12]] *A sequence $\{\kappa_n\}$ of functions $\kappa_n : Z \rightarrow (0, \infty)$ for $n \geq 0$ is said to be an analytic submultiplicative (resp., multiplicative) cocycle (with respect to g) if for all $m, n \geq 0$ and for all $z \in Z$,*

- (1) κ_n is upper-semicontinuous (usc for short) with respect to the Zariski topology on Z and $\kappa_n \geq c_\kappa^n$ for some constant $c_\kappa > 0$, and
- (2) $\kappa_{m+n}(z) \leq \kappa_n(z) \cdot \kappa_m(g^n(z))$ (resp., =).

Definition 4.2 (Introduction in [12]).

$$\kappa_{-n}(z) := \max_{w \in g^{-n}(z)} \kappa_n(w).$$

Observe that κ_{-n} is usc in the Zariski sense. The following theorem is the key in this section.

Theorem 4.3. [Theorem 1.2 in [12]] *The sequence $\{(\kappa_{-n})^{1/n}\}$ converges to a function κ_- defined over Z with the following properties: for all $\delta > \inf_Z \kappa_-$, the set $\{\kappa_- \geq \delta\}$ is a proper analytic subset of Z , invariant under g and contained in the orbit of $\{\kappa_n \geq \delta^n\}$ for all $n \geq 0$. In particular, κ_- is usc in the Zariski sense.*

The above notions and related properties work well on compact Kähler manifolds and so we will adopt the settings used in [1]. From now on, we assume the hypotheses of Theorem 1.1. Then, f is a finite-to-one map and by the open mapping theorem between complex spaces, f is open.

4.1. Local multiplicity of f^n . For each $n \in \mathbb{N}$, define $\mu_n(x)$ to be the local multiplicity of f^n at $x \in X$. Then, $\{\mu_n\}$ is an analytic multiplicative cocycle with respect to f . By Theorem 4.3, the limit function μ_- with $\min_X \mu_- = 1$ exists for the sequence $\{\mu_n\}$. Since X is compact and μ_- is usc in the Zariski sense, there exists a constant $d_f > 0$ such that $d_f = \max_X \mu_-$. Since the topological degree of the map f is d_k , $\mu_1(x) \leq d_k$ for all $x \in X$ and therefore, by (2) of Definition 4.1, $(\mu_{-n}(x))^{1/n} \leq d_k$ and $d_f \leq d_k$.

If $d_f > 1$, then we define $E_\lambda := \{\mu_- \geq d_f \lambda^{-1}\}$ for $1 \leq \lambda < d_f$.

4.2. Multiplicity of the analytic subset defined by the set of critical values of f^n . For notational convenience, we denote $\xi(x, \varphi) := \nu(x, dd^c \log |\varphi|)$ for a point $x \in X$ and a holomorphic function $\varphi : X \rightarrow \mathbb{C}$ where $\nu(x, dd^c \log |\varphi|)$ denotes the Lelong number of the current $dd^c \log |\varphi|$ at $x \in X$. Then, by the chain rule, we have $\xi(x, J_{f^{m+n}}) = \xi(x, J_{f^n}) + \xi(x, J_{f^m} \circ f^n)$ for any $x \in X$ and for any $m, n \in \mathbb{N}$ and also the following proposition:

Proposition 4.4 (See Remark 3 in [23]; for a sharper version, see also [35]). *For any $x \in X$ and for any $m, n \geq 0$, the following inequality holds:*

$$\xi(x, J_{f^m} \circ f^n) \leq (2k - 1 + 2\xi(x, J_{f^n})) \cdot \xi(f^n(x), J_{f^m}).$$

For each $n \in \mathbb{N}$, we define $\mu'_n(x) := 2k - 1 + 2\xi(x, J_{f^n})$ on X , where J_{f^n} denotes the Jacobian determinant of f^n . Then, the above proposition implies that $\{\mu'_n\}$ is an analytic submultiplicative cocycle with respect to f (see Section 3 in [26]). Hence, by Theorem 4.3, the limit function μ'_- exists for $\{\mu'_n\}$. We have $\min_X \mu'_- = 1$. Since μ'_- is usc in the Zariski sense, there exists a constant $d'_f > 0$ such that $d'_f = \max_X \mu'_-$.

If $d'_f > 1$, we define $E'_{\lambda'} := \{\mu'_- \geq d'_f (\lambda')^{-1}\}$ for $1 \leq \lambda' < d'_f$. The sets E_λ and $E'_{\lambda'}$ are proper analytic subsets of X invariant under f .

Lemma 4.5 (Lemma 2.6 in [1]). *Assume that $d_f, d'_f > 1$. Let E_λ and $E'_{\lambda'}$ be defined for λ and λ' with $1 < \lambda < d_f$ and $1 < \lambda' < d'_f$, respectively. Let $E := E_\lambda \cup E'_{\lambda'}$. Then, E is invariant under f and there exists $n_E \in \mathbb{N}$ such that for every $m \in \mathbb{N}$,*

- (1) $\mu_{-n_E m}(x) < \left(\frac{d_f}{\lambda}\right)^{n_E m}$ for $x \in \Psi_{n_E m} \setminus E$;
- (2) $\text{multi}_x \Psi_{n_E m} = \nu(x, [\Psi_{n_E m}]) < c_\Psi n_E m \left(\left(\frac{d_f}{\lambda}\right)^k \frac{d'_f}{\lambda'}\right)^{n_E m}$ for $x \in \Psi_{n_E m} \setminus E$,

where c_Ψ denotes the number of the irreducible components in the hypersurface Ψ_1 of the critical values of f .

5. HÖLDER CONTINUITY

In this section, we collect the Hölder continuity properties of some functions and some superpotentials which will be used to prove Theorem 1.1.

Definition 5.1. *Let K and α be positive constants. Let U be an open subset of X . A continuous map $g : X \rightarrow \mathbb{R}$ is said to be (K, α) -Hölder continuous on U if for every $x, y \in U$, we have*

$$|g(x) - g(y)| \leq K \text{dist}(x, y)^\alpha.$$

Let X and $f : X \rightarrow X$ be as in Theorem 1.1. Let E be an analytic subset invariant under f and $\delta > 1$ a real number such that $\mu_{-1} < \delta$ on $X \setminus E$. As a corollary to [21, Proposition 4.2], we obtain

Corollary 5.2 (Corollary 4.4 in [21]). *There are an integer N_E and a constant $c_E \geq 1$ such that if $0 < t < 1$ is a constant and if x, y are two points in X with $\text{dist}(x, E) > t$ and $\text{dist}(y, E) > t$, then we can write*

$$f^{-1}(x) = \{x_1, \dots, x_{d_k}\} \quad \text{and} \quad f^{-1}(y) = \{y_1, \dots, y_{d_k}\}$$

with $\text{dist}(x_i, y_i) \leq c_E t^{-N_E} \text{dist}(x, y)^{1/\delta}$.

Let ω_f be a smooth closed $(1, 1)$ -form in $\{f_*(\omega)\}$. Then, there exists a q-psh function $u_f : X \rightarrow \mathbb{R}$ such that $f_*(\omega) - \omega_f = dd^c u_f$. The function u_f is unique up to a constant. By adding a proper constant, we may assume that $u_f < -1$. We can investigate the Hölder continuity property of u_f outside the set E .

Lemma 5.3. *Assume that $s > 0$ be a sufficiently small real number. Then, u_f is $(c_f s^{-N_E}, \delta^{-1})$ -Hölder continuous on $X \setminus E_s$ for some $c_f > 0$ independent of s where N_E is a constant in Corollary 5.2.*

Lemma 5.4. *Let $g : X \rightarrow \mathbb{R}$ be a continuous function. Let K, α and θ be positive real numbers. Assume that $|\theta|$ be sufficiently small. Let W_1 and W_2 be two neighborhoods of E such that $W_2 \Subset W_1$ and $(W_1^c)_\theta \subset W_2^c$. Assume that g is (K, α) -Hölder continuous on W_2^c . Then, there exists a smooth function \tilde{g} defined in a neighborhood of W_1^c to \mathbb{R} such that*

$$\|\tilde{g}\|_{C^2, W_1^c} \leq c\theta^{-2(k+1)} \quad \text{and} \quad \|g - \tilde{g}\|_{\infty, W_1^c} \leq cK\theta^\alpha$$

where the constant $c > 0$ is independent of K, α, θ, W_1 and W_2 .

Proof. We first consider the following local case. Denote by B and B' two balls in a coordinate chart with center at 0 and of radii 1 and 2, respectively. Let K, α and θ be positive real numbers. Let $E \subset \mathbb{C}^k$ be an analytic subset, and V and W two open neighborhoods of E such that $W^c \subset (W^c)_\theta \subset V^c$ and that $B \setminus W \neq \emptyset$ in order to avoid triviality. Let $g : B' \rightarrow \mathbb{R}$ be a continuous function which is (K, α) -Hölder continuous on $B' \setminus V$.

Let g_{reg} denote the restriction to the set $B \setminus W$ of the standard regularization by convolution of g . Then, $g_{\text{reg}} : B \setminus W \rightarrow \mathbb{R}$ becomes a desired smooth function. More precisely, let $\psi : \mathbb{C}^k \rightarrow \mathbb{R}$ be a smooth function such that $\text{supp } \psi \Subset \{|z| < 1\}$, $\psi(z) \geq 0$ for all $z \in \mathbb{C}^k$, $\psi(z) = \psi(|z|)$, ψ is decreasing in $|z|$ and $\int_{\mathbb{C}^k} \psi(z) d\lambda(z) = 1$ where λ denotes the standard Lebesgue measure on \mathbb{C}^k . Then, g_{reg} is defined by

$$g_{\text{reg}}(x) := \int_{y \in \mathbb{C}^k} g(x - y) |\theta|^{-2k} \psi(y/\theta) d\lambda(y)$$

Since the support of $\psi(y/\theta)$ sits inside $\{|y| < |\theta|\}$, for any $x \in B \setminus W$,

$$\begin{aligned} |g(x) - g_{\text{reg}}(x)| &= \left| \int_{y \in \mathbb{C}^k} (g(x) - g(x - y)) |\theta|^{-2k} \psi(y/\theta) d\lambda(y) \right| \\ &\leq \int_{y \in \mathbb{C}^k} |g(x) - g(x - y)| |\theta|^{-2k} \psi(y/\theta) d\lambda(y) \\ &\leq \int_{y \in \mathbb{C}^k} K |y|^\alpha |\theta|^{-2k} \psi(y/\theta) d\lambda(y) \leq \int_{y \in \mathbb{C}^k} K |\theta|^\alpha |\theta|^{-2k} \psi(y/\theta) d\lambda(y) \leq K |\theta|^\alpha \end{aligned}$$

For the second argument, we use the change of variables. This is also standard.

$$g_{\text{reg}}(x) = \int_{y \in \mathbb{C}^k} g(x - y) |\theta|^{-2k} \psi(y/\theta) d\lambda(y) = \int_{z \in \mathbb{C}^k} g(z) |\theta|^{-2k} \psi((x - z)/\theta) d\lambda(z).$$

Since ψ is smooth, we have

$$\|g_{\text{reg}}\|_{C^2} \lesssim \sup_{B'} |g| \cdot |\theta|^{-2(k+1)} \|\psi\|_{C^2}.$$

Observe that the constants involved in the inequalities other than K, α, θ and $\sup_{B'} |g|$ are some constants related to ψ . So, for a fixed $g : B' \rightarrow \mathbb{R}$, we get the same inequalities for different K, α, θ, V and W as long as $W^c \subset (W^c)_\theta \subset V^c$ holds and g is (K, α) -Hölder continuous on $B' \setminus V$.

Now, we consider the general case. At each point $x_0 \in X$, we take an open neighborhood U_{x_0} and a coordinate chart $\phi_{x_0} : U_{x_0} \rightarrow \mathbb{C}^k$ such that $B' \subset \phi_{x_0}(U_{x_0})$. This can be always achieved by scaling ϕ_{x_0} . Then, we let $B_{x_0} := \phi_{x_0}^{-1}(B)$ and $B'_{x_0} := \phi_{x_0}^{-1}(B')$. Since X is compact, we can find a finite cover $(B_j)_{j=1, \dots, N}$ out of $(B_{x_0})_{x_0 \in X}$. Let $(\chi_j)_{j=1, \dots, N}$ be a partition of unity subordinated to the cover $(B_j)_{j=1, \dots, N}$.

Notice that on each B'_j , the Euclidean metric and the metric induced from X are equivalent. So, there exists a constant $c_j > 1$ such that $c_j^{-1} \text{dist}(x, y) \leq |x - y| \leq c_j \text{dist}(x, y)$ for $x, y \in B'_j$. So, we apply the above model case with θ replace by $c_j^{-1} \theta_i$ and with $V = W_2 \cap B'_j$ and $W = W_1 \cap B'_j$ and denote by $g_{i,j}$ the resulting function g_{reg} on B_j . Then, the desired function g_i is obtained by

$$g_i = \sum_{j=1}^N \chi_j g_{i,j}.$$

□

Now, we consider the Hölder continuity of some superpotentials. Since f is not just a holomorphic correspondence but a holomorphic map, $f^*(\alpha)$ is smooth whenever α is smooth. Hence, we can apply the same argument in the proof Lemma 5.4.3 in [20] and we get the desired Hölder continuity.

Lemma 5.5. *Let $1 \leq s \leq k$ be an integer. The superpotential of $f_*(\omega^s) - \alpha_{f_*(\omega^s)}$ admits Hölder continuous superpotentials, where $\alpha_{f_*(\omega^s)}$ is a real closed smooth (s, s) -form in $\{f_*(\omega^s)\}$.*

Proof. We know that $f_*(\omega)$ admits Hölder continuous quasi-potentials. Then, by [22, Proposition 3.4.2], we have the Hölder continuity of superpotentials of $[f_*(\omega)]^l$. Then, since $f_*(\omega^l) \leq [f_*(\omega)]^l$, the domination principle in [15, Theorem 1.1] implies that the superpotential of $f_*(\omega^l)$ is Hölder continuous. □

6. PROOF OF THEOREM 1.1

Since E is invariant under f , without loss of generality, it suffices to prove the theorem with f replaced by some power f^{N_f} of it. We first find E and then a good iterate f^{N_f} of f . For notational convenience, we will write operators $L := d_p^{-1} f^*$, $\Lambda := d_{p-1}^{-1} f_*$ on \mathcal{D}_p , \mathcal{D}_{k-p+1} , respectively. Note that since the set of indeterminacy is empty, $(f^n)^* = (f^*)^n$. So, $L^n(\cdot) = d_p^{-n} (f^*)^n(\cdot)$ on \mathcal{D}_p .

Recall the two constants d_f and d'_f associated with f introduced in Section 4. We first consider the case of $d_f > 1$ and $d'_f > 1$ and then other cases later.

6.1. $d_f > 1$ and $d'_f > 1$. We define a non-negative integer n_{dyn} such that

$$d_f \leq \left(\frac{d_s}{d_{s-1}} \right)^{n_{\text{dyn}}} \quad \text{for all } 1 \leq s \leq p.$$

Let $1 \leq p \leq k$ be an integer such that $d_{p-1} < d_p$. We denote $d := d_p/d_{p-1}$.

We can find a sufficiently large $N_1 \in \mathbb{N}$ so that for every $n \geq N_1$,

$$(40k^2 c_\Psi n)^{8n_{\text{dyn}} k} < d_f^n,$$

where c_Ψ is the number of the irreducible components in the hypersurface Ψ_1 of the critical values of f as in Section 4.

Recall the definition of the dynamical degree. There exists a $N_2 \in \mathbb{N}$ such that for every $n \geq N_2$,

$$\langle (f^n)^* \omega^p, \omega^{k-p} \rangle \leq \left(d_p + \frac{1}{4} \right)^n \quad \text{and} \quad \langle (f^n)_* \omega^{k-p+1}, \omega^{p-1} \rangle \leq \left(d_{p-1} + \frac{1}{4} \right)^n.$$

Let $c_m > 0$ be a constant such that for every integer $1 \leq l \leq k$, $-c_m \|\varphi\|_\infty \omega^l \leq \varphi \leq c_m \|\varphi\|_\infty \omega^l$ holds for every smooth (l, l) -form φ . Then, if $n \geq N_2$, we have

$$\langle L^n(S), \omega^{k-p} \rangle = \langle L^n(\alpha_S), \omega^{k-p} \rangle \leq c_m \left(\frac{5}{4}\right)^n \|\alpha_S\|_\infty$$

for every current $S \in \mathcal{C}_p$ where α_S is a smooth closed form in $\{S\}$. In the same way, if $n \geq N_2$

$$\langle \Lambda^n(R), \omega^{p-1} \rangle = \langle \Lambda^n(\alpha_R), \omega^{p-1} \rangle \leq c_m \left(\frac{5}{4}\right)^n \|\alpha_R\|_\infty$$

for every current $R \in \mathcal{C}_{k-p+1}$ where α_R is a smooth closed form in $\{R\}$.

Then, take $N = N_1 + N_2$ and we can choose $1 < \lambda < d_f$ such that

$$(40k^2 c_\Psi N)^{(Nk)^{-1}} d_f^{1-(8n_{\text{dyn}}k^2)^{-1}} < \lambda < d_f.$$

Further, we can also choose $1 < \lambda' < d'_f$ such that

$$(6.1) \quad 1 < \left(\frac{d_f}{\lambda}\right)^k \left(\frac{d'_f}{\lambda'}\right) < (40k^2 c_\Psi N)^{-N^{-1}} d_f^{(8n_{\text{dyn}}k)^{-1}} < (40k^2 c_\Psi N)^{-N^{-1}} d^{(8k)^{-1}}.$$

For such $1 < \lambda < d_f$ and $1 < \lambda' < d'_f$, let $E := E_\lambda \cup E'_{\lambda'}$ where $E_\lambda := \{\mu_- \geq d_f \lambda^{-1}\}$ and $E'_{\lambda'} := \{\mu'_- \geq d'_f (\lambda')^{-1}\}$ as in Section 4. For each $j \in \mathbb{N}$ and for $\delta = c_\Psi n_E N j \left(\left(\frac{d_f}{\lambda}\right)^k \frac{d'_f}{\lambda'}\right)^{n_E N j}$, Lemma 4.5 implies

- (1) $\mu_{-n_E N j}(x) < \left(\frac{d_f}{\lambda}\right)^{n_E N j} < \delta$ for $x \in \Psi_{n_E N j} \setminus E$;
- (2) $\text{multi}_x \Psi_{n_E N j} = \nu(x, [\Psi_{n_E N j}]) < c_\Psi n_E N j \left(\left(\frac{d_f}{\lambda}\right)^k \frac{d'_f}{\lambda'}\right)^{n_E N j} = \delta$ for $x \in \Psi_{n_E N j} \setminus E$,

where c_Ψ denotes the number of the irreducible components in the hypersurface Ψ_1 of the critical values of f .

We look into the relationship between δ and d which is a crucial condition for the proof. From (6.1) and $40k^2 c_\Psi N > 3$, we get

$$40k^2 \delta = (40k^2 c_\Psi n_E N j) \left(\left(\frac{d_f}{\lambda}\right)^k \left(\frac{d'_f}{\lambda'}\right)\right)^{n_E N j} < (40k^2 c_\Psi N)^{n_E j} \left(\left(\frac{d_f}{\lambda}\right)^k \left(\frac{d'_f}{\lambda'}\right)\right)^{n_E N j} < d^{n_E N j (8k)^{-1}}.$$

Hence, we take $N_f = n_{\text{dyn}} n_E N$. Since E is invariant under f , it is invariant under f^{N_f} . We replace f by f^{N_f} , λ by λ^{N_f} and λ' by $(\lambda')^{N_f}$. Also Ψ_1 is replaced by Ψ_{N_f} , which will be denoted by V . By the definition, the dynamical degree d_s is replaced by $d_s^{N_f}$ for $0 \leq s \leq k$, and d_f and d'_f are replaced by $d_f^{N_f}$ and $d'_f{}^{N_f}$. Then, we may assume that $f : X \rightarrow X$ satisfy **Condition (M)**

- (1) $\mu_{-1}(x) < \delta$ for $x \in V \setminus E$ and
- (2) $\text{multi}_x V < \delta$ for $X \in V \setminus E$

where $\delta = c_\Psi n_{\text{dyn}} n_E N \left(\frac{d_f}{\lambda}\right)^k \left(\frac{d'_f}{\lambda'}\right)$. But note that we keep c_Ψ from the replacement, and that n_{dyn} , n_E and N are unchanged. The number $c_\Psi n_{\text{dyn}} n_E N$ will play the role of an upper bound of the number of irreducible components in Ψ_{N_f} . After the replacement, we have

$$(6.2) \quad (40k^2 \delta)^{6k} < d^{3/4}$$

and the mass estimate as follows:

$$(6.3) \quad \|L^n(S)\| \leq c_m \left(\frac{5}{4}\right)^{N_f n} \|\alpha_S\|_\infty \quad \text{and} \quad \|\Lambda^n(R)\| \leq c_m \left(\frac{5}{4}\right)^{N_f n} \|\alpha_R\|_\infty$$

for $S \in \mathcal{C}_p$ and for $R \in \mathcal{C}_{k-p+1}$.

Now, we prove the statement of Theorem 1.1. Let $S \in \mathcal{C}_p$ be a current such that S has a smooth representation in a neighborhood of E . Let φ be a smooth test form of bidegree $(k-p, k-p)$. Our goal is to show

$$|\langle L^n(S) - L^n(\alpha_S), \varphi \rangle| = |\mathcal{U}_{L^n(S-\alpha_S)}(dd^c \varphi)| \leq c \|\varphi\|_{C^2} \rho^n \quad \text{for all } n \in \mathbb{N}$$

for some constants $c > 0$ and $0 < \rho < 1$ independent of φ . Similarly to c_m , we can find a constant $c'_m > 0$ such that for any φ , $c'_m \|\varphi\|_{C^2} \omega^{k-p+1} + dd^c \varphi$ is a positive form. We can write

$$dd^c \varphi = (c'_m \|\varphi\|_{C^2} \omega^{k-p+1} + dd^c \varphi) - c'_m \|\varphi\|_{C^2} \omega^{k-p+1}.$$

So, we will estimate $\mathcal{U}_{L^n(S-\alpha_S)}(R)$ for smooth $R \in \mathcal{D}_{k-p+1}^0$ such that $R = R_+ - R_-$ where R_\pm are smooth currents in \mathcal{C}_{k-p+1} with $\|R_\pm\|_\infty \leq 1$.

To this end, we define some sequences $(s_{n,i}, \varepsilon_{n,i}, t_{n,i})$ below) for computational purposes which will be used in splitting X into three regions. Roughly speaking, the first region is a neighborhood of a subset of $V \setminus E$, the second one a neighborhood of E , and the last one the complement of the two sets. The following two lemmas will be used to find an appropriate size of the neighborhood of a subset of $V \setminus E$ and E .

Below was induced from a Lojasiewicz type inequality in [1]. See also [29] and [21]. It is of local nature, and so it is valid in our case as well. Let α_V be a closed smooth $(1,1)$ -form cohomologous to the current of integration $[V]$. Then, we can find a unique q-psh function φ_V over X such that $\sup_X \varphi_V = 0$ and $dd^c \varphi_V = [V] - \alpha_V$. Let $E_V = V \cap E$. Then, from our construction, for all $x \in V \setminus E_V$, $\text{multi}_x V < \delta$ from (2) of **Condition (M)**.

Lemma 6.1 (See Lemma 3.3 in [1]). *There are constants $C, A > 0$ such that for $x \in X$,*

$$\delta \log \text{dist}(x, V) + C \log \text{dist}(x, E_V) - A \leq \varphi_V(x) \leq \log \text{dist}(x, V) + A.$$

Lemma 6.2 (See Lemma 3.1 in [21]). *There is a constant $m_f \geq 1$ such that for every subset A and B of X ,*

$$\text{dist}(f^{-j}(A), f^{-j}(B)) \geq m_f^{-j} \text{dist}(A, B) \quad \text{for every } j \geq 0.$$

In particular, if $f(B) \subseteq B$,

$$\text{dist}(f^{-j}(A), B) \geq m_f^{-j} \text{dist}(A, B) \quad \text{for every } j \geq 0.$$

For the moment, let $\varepsilon > 0$ be a sufficiently small positive real number so that $\varepsilon < m_f^{-1}$. Later, this number will be precisely defined in terms of n . We define the following sequences of real numbers and currents for $n \in \mathbb{N}$ and $1 \leq i \leq n$.

- $s_{n,i} = \varepsilon^{n C_i}$,
- $\varepsilon_{n,i} = \varepsilon^{nC(4k+N_E)} (40k^2 \delta)^{6ki}$, and
- $t_{n,i} = \varepsilon_{n,i} (10k)^{-1}$,
- $R_{n,0} = R$ and $R_{n,i} := (\Lambda(R_{n,i-1}))_{\varepsilon_{n,i}}$.

where the subscript means the regularization in Section 3. The constant $C > 0$ is from Lemma 6.1 and the constant N_E is from Corollary 5.2. Then, following [20] and [1], we can expand

$$\begin{aligned}
\langle U_{L^n(S-\alpha_S)}, R \rangle &= \langle LU_{L^{n-1}(S-\alpha_S)}, R \rangle = d^{-1} \langle U_{L^{n-1}(S-\alpha_S)}, \Lambda(R) \rangle \\
&= d^{-1} \langle U_{L^{n-1}(S-\alpha_S)}, \Lambda(R_{n,0}) - R_{n,1} \rangle + d^{-1} \langle U_{L^{n-1}(S-\alpha_S)}, R_{n,1} \rangle \\
&= d^{-1} \langle U_{L^{n-1}(S-\alpha_S)}, \Lambda(R_{n,0}) - R_{n,1} \rangle + d^{-2} \langle U_{L^{n-2}(S-\alpha_S)}, \Lambda(R_{n,1}) \rangle \\
&= d^{-1} \langle U_{L^{n-1}(S-\alpha_S)}, \Lambda(R_{n,0}) - R_{n,1} \rangle \\
&\quad + \cdots + d^{-i} \langle U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \\
&\quad + \cdots + d^{-n} \langle U_{S-\alpha_S}, \Lambda(R_{n,n-1}) - R_{n,n} \rangle + d^{-n} \langle U_{S-\alpha_S}, R_{n,n} \rangle
\end{aligned}$$

where the potentials are assumed to be the Green potentials. Here, the pairings in the above make sense since the Green potentials are currents of order 0 and $\Lambda(R_{n,i})$'s admit continuous superpotentials for all $n \in \mathbb{N}$ and for all $1 \leq i \leq n$ as $R_{n,i}$'s have C^1 -coefficients. Since the operator Λ and the regularizing operators are linear, we need to estimate

- (1) $d^{-i} \langle U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$ for smooth $R \in \mathcal{C}_{k-p+1}$ with $\|R\|_\infty \leq 1$ and for $i = 1, \dots, n$, and
- (2) $d^{-n} \langle U_{S-\alpha_S}, R_{n,n} \rangle$ for smooth $R \in \mathcal{D}_{k-p+1}^0$ such that $R = R_+ - R_-$ where R_\pm are smooth currents in \mathcal{C}_{k-p+1} with $\|R_\pm\|_\infty \leq 1$.

Suppose that whenever $\varepsilon > 0$ is sufficiently small, then (1) $\lesssim \varepsilon$ and (2) $\lesssim nd^{n/4}(-\log \varepsilon)$ for all integers n and i with $1 \leq i \leq n$ where the inequality is independent of n and i . (These estimates of (1) and (2) will be proved in Section 7 and Section 8.) Then, from the expansion, we have

$$|\langle U_{L^n(S-\alpha_S)}, R \rangle| \lesssim 6n\varepsilon + nd^{n/4}(-\log \varepsilon)$$

for all sufficiently small $\varepsilon > 0$. We take $\varepsilon = d^{-n}$ and let $n \rightarrow \infty$. Then, $\rho = d^{1/8}$ proves the statement for the case of $d_f > 1$ and $d'_f > 1$.

6.2. $d_f = 1$ or $d'_f = 1$. In these cases, we get either $E = \emptyset$ or $E' = \emptyset$. We will consider only the case of $d_f = 1$ and $d'_f = 1$. For the other two cases can be easily deduced. It is enough to find a good iterate f^{N_f} of f so that we can apply the previous argument. Let $d := d_p/d_{p-1}$. Choose a sufficiently large $N_1 \in \mathbb{N}$ so that for every $n \geq N_1$, we have

$$(40k^2 c_\psi n)^{8k} < d^n.$$

Then, as in the case of $d_f > 1$ and $d'_f > 1$, we can find $1 < \lambda < d$ and $0 < \lambda' < 1$ so that

$$1 < \left(\frac{d}{\lambda}\right)^k \left(\frac{1}{\lambda'}\right) < (40k^2 c_\psi N_1)^{-N_1^{-1}} d^{(8k)^{-1}}.$$

As in the case of $d_f > 1$ and $d'_f > 1$, there exist constants $c_m > 0$ and $N_2 \in \mathbb{N}$ such that

$$\langle L^n(S), \omega^{k-p} \rangle \leq c_m \left(\frac{5}{4}\right)^n \|\alpha_S\|_\infty \quad \text{and} \quad \langle \Lambda^n(R), \omega^{p-1} \rangle \leq c_m \left(\frac{5}{4}\right)^n \|\alpha_R\|_\infty$$

for every $n \geq N_2$ where $S \in \mathcal{C}_p$ and $R \in \mathcal{C}_{k-p+1}$ are currents, and α_S and α_R are smooth closed forms in the same cohomology class as S and R , respectively. Let $N = N_1 + N_2$.

For such $1 < \lambda < d$ and $0 < \lambda' < 1$, as remarked in the above, we have $E := \emptyset$ and the arguments in Lemma 4.5 imply that for some large enough $n_E \in \mathbb{N}$, we have

- (1) $\mu_{-n_EN}(x) < \left(\frac{d}{\lambda}\right)^{n_EN} < \delta$ for $x \in \Psi_{n_EN}$;
- (2) $\text{multi}_x \Psi_{n_EN} < c_\Psi n_EN \left(\left(\frac{d}{\lambda}\right)^k \frac{1}{\lambda'}\right)^{n_EN} = \delta$ for $x \in \Psi_{n_EN}$,

where $\delta = c_\Psi n_E N \left(\left(\frac{d}{\lambda} \right)^k \frac{1}{\lambda'} \right)^{n_E N}$ and c_Ψ denotes the number of the irreducible components in the hypersurface Ψ_1 of the critical values of f . We replace f by f^{N_f} where $N_f := n_E N$. Then, with these multiplicity conditions, our arguments in the case of $d_f > 1$ and $d'_f > 1$ work in the same way and give us the desired conclusion. Note that in this case, we have $E = \emptyset$, we only need to compute on $W_{n,i,1}$ and $W_{n,i,3}$ and that every positive closed current of bidegree (p, p) satisfies Theorem 1.1. \square

We end this section with a lemma about some estimates related to the currents $R_{n,i}$, which were not needed in the case of \mathbb{P}^k due to the simplicity of its cohomology.

Lemma 6.3. *Let $C_m = 2c_{\text{reg}} c_m \left(\frac{5}{4} \right)^{N_f}$. Let $R \in \mathcal{C}_{k-p+1}$ be a smooth current. Then, for all positive integers n and i such that $1 \leq i \leq n$, the current $R_{n,i}$ can be expressed $R_{n,i} = R_{n,i}^+ - R_{n,i}^-$ where $R_{n,i}^\pm$ are smooth positive closed $(k-p+1, k-p+1)$ -currents such that*

$$(1) \quad \|R_{n,i}^+\|_{C^1} + \|R_{n,i}^-\|_{C^1} \leq \|R\|_{C^1} (C_m)^i \left(\prod_{j=1}^i \varepsilon_{n,j} \right)^{-6k^2} \quad \text{and}$$

$$(2) \quad \|R_{n,i}^+\| + \|R_{n,i}^-\| \leq \|R\|_{C^1} (C_m)^i \left(\prod_{j=1}^{i-1} \varepsilon_{n,j} \right)^{-6k^2}.$$

Proof. We prove by induction. When $i = 1$, Then, $R_{n,1} = (\Lambda(R))_{\varepsilon_{n,1}}^+ - (\Lambda(R))_{\varepsilon_{n,1}}^-$. Proposition 3.5 and the estimates (6.3) prove it. So, we may assume that it is true for $i - 1$ with $i \geq 2$. Denote by $r_{n,i} = \|R_{n,i}^+\|_{C^1} + \|R_{n,i}^-\|_{C^1}$. Then, from Section 3, we have

$$R_{n,i} = (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^+ + (\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^- - (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^- - (\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^+$$

and define

$$R_{n,i}^+ := (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^+ + (\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^- \quad \text{and} \quad R_{n,i}^- := (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^- + (\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^+.$$

From Proposition 3.5 and the estimates (6.3), we have

$$\|(\Lambda(R_{n,i-1}^\pm))_{\varepsilon_{n,i}}^\pm\|_{C^1} \leq c_{\text{reg}} \varepsilon_{n,i}^{-6k^2} \|\Lambda(R_{n,i-1}^\pm)\| \leq c_{\text{reg}} c_m \left(\frac{5}{4} \right)^{N_f} \varepsilon_{n,i}^{-6k^2} \|R_{n,i-1}^\pm\|_{C^1},$$

respectively and thus by the triangle inequality, we get

$$r_{n,i} \leq 2c_{\text{reg}} c_m \left(\frac{5}{4} \right)^{N_f} \varepsilon_{n,i}^{-6k^2} r_{n,i-1}.$$

For the mass estimate, we have

$$\begin{aligned} \|R_{n,i}^+\| &\leq \|(\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^+\| + \|(\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^-\| \leq c_{\text{reg}} (\|\Lambda(R_{n,i-1}^+)\| + \|\Lambda(R_{n,i-1}^-)\|) \\ &\leq c_{\text{reg}} c_m \left(\frac{5}{4} \right)^{N_f} r_{n,i-1}. \end{aligned}$$

The same is true for $R_{n,i}^-$. So, we proved for i as desired. \square

7. ESTIMATES OF $d^{-i} \langle U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$

We start by constructing two sequences of cut-off functions playing the role of smoothing out the boundary of the neighborhoods of E and a subset of $V \setminus E$, on each of which the currents $L^{n-i}(S - \alpha_S)$ and $f_*(\omega^{k-p+1})$ have different analytic properties.

Lemma 7.1 (See Lemma 2.2.6 in [20]). *Let $\chi : \mathbb{R} \cup \{-\infty\} \rightarrow \mathbb{R}$ be a convex increasing function such that χ' is bounded. Then, for every DSH function φ , $\chi(\varphi)$ is DSH and*

$$\|\chi(\varphi)\|_{\text{DSH}} \lesssim 1 + \|\varphi\|_{\text{DSH}}.$$

In particular, $\inf \|T_\chi^-\|$ is bounded by $\|\chi'\|_\infty \inf \|T^-\|$, where T_χ^\pm and T^\pm are positive closed currents such that $dd^c \chi(\varphi) = T_\chi^+ - T_\chi^-$ and $dd^c \varphi = T^+ - T^-$ and \inf is taken over such T_χ^- 's and T^- 's, respectively.

The following is a basic regularization lemma. We can get it through patching the locally regularized functions on each coordinate chart by use of a partition of unity.

Lemma 7.2. *Let E be the analytic subset as above. There are constants $s_0 > 0$ and $0 < r_E < 1$ such that for all $0 < s \leq s_0$, there exists a smooth function $\chi_s^E : X \rightarrow \mathbb{R}$ such that $\chi_s^E \equiv 1$ on $E_{r_E s}$, $\text{supp} \chi_s^E \subseteq E_s$ and $\|\chi_s^E\|_{C^2} \lesssim 1/s^2$ where the inequality \lesssim means \leq up to a constant multiple independent of s .*

The following crucial lemma is induced from Lemma 6.1. The argument in [1, Lemma 7.3] is of local nature and so it works in our case as well. This is one of the two places where the multiplicity assumption comes to play. The other is the Hölder continuity of u_f where u_f is a function such that $f_*(\omega) - \omega_f = dd^c u_f$.

Lemma 7.3 (Lemma 7.3 in [1]). *Let n and i be positive integers such that $1 \leq i \leq n$. Let s, t be two positive real numbers such that $\frac{1}{2} r_E^{k+2} s_{n,i} \leq s \leq 2s_{n,i}$ and $\frac{1}{2} t_{n,i} \leq t \leq 2t_{n,i} \left(\frac{2}{2\delta+1}\right)^{k+2}$ for $s_{n,i}, t_{n,i}$ with sufficiently small $\varepsilon > 0$. Then, there is a function $\chi_{s,t} : X \rightarrow \mathbb{R}$ with $0 \leq \chi_{s,t} \leq 1$, such that $\chi_{s,t} \equiv 1$ on $V_t \setminus E_s$, $\text{supp}(\chi_{s,t}) \subseteq V_{t(\delta+1/2)^{-1}} \setminus E_{r_E s}$, and $\|\chi_{s,t}\|_{\text{DSH}} \leq c_\chi \max\{1, 9s^{-2}\}$, where $c_\chi > 0$ is a constant independent of s, t , and ε, i, n in the definition of $s_{n,i}, t_{n,i}$.*

We define

$$\begin{aligned} \chi_{n,i,1} &:= \left(1 - \chi_{s_{n,i}/r_E}^E\right) \chi_{s_{n,i}, t_{n,i}}, & W_{n,i,1} &:= V_{t_{n,i}(\delta+1/2)^{-1}} \setminus E_{s_{n,i}}, \\ \chi_{n,i,2} &:= \chi_{s_{n,i}/r_E}^E, & W_{n,i,2} &:= E_{s_{n,i}/r_E} \quad \text{and} \\ \chi_{n,i,3} &:= 1 - \chi_{n,i,1} - \chi_{n,i,2} = \left(1 - \chi_{s_{n,i}/r_E}^E\right) \left(1 - \chi_{s_{n,i}, t_{n,i}}\right), & W_{n,i,3} &:= X \setminus (V_{t_{n,i}} \cup E_s). \end{aligned}$$

Then, we have

$$\begin{aligned} \chi_{n,i,1} &\equiv 1 \text{ on } V_{t_{n,i}} \setminus E_{s_{n,i}/r_E}, & \text{supp}(\chi_{n,i,1}) &\subseteq W_{n,i,1}, \\ \chi_{n,i,2} &\equiv 1 \text{ on } E_{s_{n,i}}, & \text{supp}(\chi_{n,i,2}) &\subseteq W_{n,i,2} \text{ and} \\ \chi_{n,i,3} &\equiv 1 \text{ on } X \setminus \left(V_{t_{n,i}(\delta+1/2)^{-1}} \cup E_s\right), & \text{supp}(\chi_{n,i,3}) &\subseteq W_{n,i,3} \end{aligned}$$

and we can write

$$d^{-i} \langle U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle = \sum_{j=1}^3 d^{-i} \langle \chi_{n,i,j} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$$

each of which will be estimated below.

7.1. Estimate of $d^{-i} \langle \chi_{n,i,1} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$ in $W_{n,i,1}$. In this region, we use the Hölder continuity of quasi-potentials of $f_*(\omega)$.

For notational convenience, we will use the following notation throughout this subsection. Let $\theta \in \mathbb{C}$ be a constant. Later, θ will be chosen to be $\varepsilon_{n,i}$ with sufficiently small ε . Let Θ_j be a smooth positive closed (k, k) -form on $\widehat{\mathbb{X}}$, which equals either $(m_{\widehat{\Delta}} \omega_{\widehat{\mathbb{X}}} + dd^c u_\theta) \wedge \omega_{\widehat{\mathbb{X}}}^{k-1}$ or $(m_{\widehat{\Delta}} \omega_{\widehat{\mathbb{X}}} - \alpha_{\widehat{\Delta}}) \wedge \omega_{\widehat{\mathbb{X}}}^{k-1}$ for $j = 1, \dots, k+2$ so that its associated positive closed semi-regular transform \mathcal{L}^{Θ_j} of bidegree $(0, 0)$ coincides with either \mathcal{L}_θ^+ or \mathcal{L}_θ^- for $j = 1, \dots, k+2$. Let $\chi^{E,j} := \chi_{\left(\frac{1}{2}r_E\right)^j}^E$ in Lemma 7.2 for

$j = 1, \dots, k+2$. We also denote by $\mathcal{L}^m = \mathcal{L}^{\Theta_1} \circ \dots \circ \mathcal{L}^{\Theta_m}$ for $0 \leq m \leq k+2$. Notice that \mathcal{L}^m is positive and closed, and that it may depend on θ .

Differently from [1], we cannot use the negativity of $U_{L^{n-i}(S-\alpha_S)}$. See [7]. So, not only do we need to estimate integrals of the form $\langle \chi_{n,i,1} \mathcal{L}^\Phi(S), f_*(\omega^{k-p+1}) \rangle$ and $\langle \chi_{n,i,1} \mathcal{L}^{u\Phi}(S), f_*(\omega^{k-p+1}) \rangle$, but also those of the form $\langle \chi_{n,i,1} \mathcal{L}^\Phi(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle$ and $\langle \chi_{n,i,1} \mathcal{L}^{u\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle$. By the continuity of the (semi-)regular transformation and the superpotential of $f_*(\omega^{k-p+1})$, we may assume that S is smooth positive and closed.

Since the proof of [20, Lemma 2.3.7] is of local nature, it gives the following.

Lemma 7.4. *Let $S \in \mathcal{C}_p$ be a smooth current. For all sufficiently small $t > 0$, we have*

$$\int_{V_t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(\omega^{k-p+1}) \lesssim \|\Phi\|_\infty \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|S\| t^{3/2}.$$

The inequality \lesssim means \leq up to a constant multiple independent of t , θ , Φ and S .

The proof is similar to the proof of Lemma 2.3.7. But X is different from \mathbb{P}^k in that X is not homogeneous.

Proof. Here since S , ω^{k-p+1} and Φ are smooth and positive, we can write

$$\begin{aligned} \int_{V_t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(\omega^{k-p+1}) &\leq \|\Phi\|_\infty \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \int_{V_t} \mathcal{L}^{\omega_{\tilde{x}}^{k-1}}(S) \wedge \omega^{k-p+1} \\ &= \|\Phi\|_\infty \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \int_{y \in V_t} \left[\int_{x \in X \setminus \{y\}} \pi_* \left(\omega_{\tilde{x}}^{k-1} \right) (x, y) \wedge S(x) \right] \wedge \omega^{k-p+1}(y) \end{aligned}$$

Since X is compact, it is enough to prove the estimate in a coordinate neighborhood $B \subset X$. So, we prove that

$$\int_{y \in V_t \cap B} \left[\int_{x \in X \setminus \{y\}} \pi_* \left(\omega_{\tilde{x}}^{k-1} \right) (x, y) \wedge S(x) \right] \wedge \omega^{k-p+1}(y) \lesssim t.$$

Let $B_z(4t)$ denote the ball of radius $4t$ centered at some $z \in V \cap B$. By shrinking B , we may assume that $B_z(4t)$ lies in the coordinate chart when $t > 0$ is sufficiently small. We estimate

$$\int_{y \in B_z(4t)} \left[\int_{x \in X \setminus \{y\}} \pi_* \left(\omega_{\tilde{x}}^{k-1} \right) (x, y) \wedge S(x) \right] \wedge \omega^{k-p+1}(y).$$

By the representation of a flat current in terms of a vector field and a Radon measure (or one might think of it as a version of the Radon-Nikodym theorem), we may write it as

$$\int_{x \in X} \left(\int_{y \in B_z(4t) \setminus \{y\}} \left[\Psi_S(x) \lrcorner \pi_* \left(\omega_{\tilde{x}}^{k-1} \right) (x, y) \right] \wedge \omega^{k-p+1}(y) \right) d\mu(x).$$

Here, $\Psi_S(y)$ is actually a $(k-p)$ -vector field with $\|\Psi_S(y)\|_\infty \leq 1$ and $\left[\Psi_S(y) \lrcorner \pi_* \left(\omega_{\tilde{x}}^{k-1} \right) (x, y) \right]$ is actually a form in x smooth except y and the singularity is at most $\text{dist}(x, y)^{2-2k}$ and one might think of μ as the trace measure of S . Hence, $\int_X d\mu$ can be bounded in terms of $\|S\|$. Since $\|\Psi_S(y)\|_\infty \leq 1$ and $\omega_{\tilde{x}}^q$ is smooth, the inner integral is bounded by a constant multiple of $\left[\left(\sum_{|I|=k-p} e_I(y) \right) \lrcorner \pi_* \left(\omega_{\tilde{x}}^{k-1} \right) (x, y) \right]$ (in Federer's notation) and further bounded by the following uniformly with respect to y .

$$\int_{B_z(4t)} |x|^{2-2k} (dd^c|x|^2)^k \lesssim t^2.$$

So, the arguments using the Lelong number and the Abel transform in [20, Lemma 2.3.7] complete the proof. \square

Next, we modify [1, Proposition 7.6] to prove the following:

Lemma 7.5. *Let n and i be positive integers such that $1 \leq i \leq n$. Let l be an integer such that $0 \leq l \leq k - p + 1$. Let s, t be two positive real numbers such that $\frac{1}{2}r_E^{k-p-l+2}s_{n,i} \leq s \leq 2s_{n,i}$ and $\frac{1}{2}t_{n,i} \leq t \leq 2(t_{n,i})^{\frac{2}{2\delta+1}}^{k-p-l+2}$ with sufficiently small $\varepsilon > 0$. Then, we have*

$$\left| \int_{V_i \setminus E_s} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^l \wedge \omega^{k-p-l+1}) \right| \lesssim \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty s^{-2(k+l)-N_E} t^{\beta_l},$$

where $\beta_l = (2(k+1)(2\delta+1))^{-l}\delta^{-l}$. The inequality \lesssim means \leq up to a constant multiple independent of $n, i, s, t, \theta, \Phi, S$ and α_S .

For this estimate, we need an auxiliary lemma. Lemma 5.3 and (1) of **Condition (M)** imply that u_f is Hölder continuous on X and that u_f is $(c_f s^{-N_E}, \delta^{-1})$ -Hölder continuous in $X \setminus E_s$.

Lemma 7.6. *Under the assumptions of Lemma 7.5, for any $1 \leq j \leq m$, we have*

$$\left| \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^{\Theta_1}(\dots \mathcal{L}^{\Theta_{j-2}}(\mathcal{L}^{\Theta_{j-1}}((1 - \chi^{E,j-1}) \mathcal{L}^{\Theta_j}(\chi^{E,j} \mathcal{L}^{\Theta_{j+1}}(\chi^{E,j+1} \dots \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2}u_f) \wedge \omega^{k-p-l+1}))) \dots))) \dots) \right| \lesssim \|\Phi\|_\infty \|S\| s^{-2k} t^{3/(2\delta+1)}$$

where $\chi_{s,t}$ is the cut-off function in Lemma 7.3. The inequality \lesssim means \leq up to a constant multiple independent of n, i, s, t, θ, Φ and S .

Proof.

$$\begin{aligned} & \int_X \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^{\Theta_1}(\dots \mathcal{L}^{\Theta_{j-2}}(\mathcal{L}^{\Theta_{j-1}}((1 - \chi^{E,j-1}) \mathcal{L}^{\Theta_j}(\chi^{E,j} \mathcal{L}^{\Theta_{j+1}}(\dots \\ & \quad \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2}u_f) \wedge \omega^{k-p-l+1}))) \dots))) \dots) \\ &= \int_X (1 - \chi^{E,j-1}) \mathcal{L}^{\Theta_{j-1}}(\mathcal{L}^{\Theta_{j-2}}(\dots \mathcal{L}^{\Theta_1}(\chi_{s,t} \mathcal{L}^\Phi(S)) \dots)) \wedge \mathcal{L}^{\Theta_j}(\chi^{E,j} \mathcal{L}^{\Theta_{j+1}}(\dots \\ & \quad \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2}u_f) \wedge \omega^{k-p-l+1}))) \dots) \end{aligned}$$

The currents $(1 - \chi^{E,j-1}) \mathcal{L}^{\Theta_{j-1}}(\mathcal{L}^{\Theta_{j-2}}(\dots \mathcal{L}^{\Theta_1}(\chi_{s,t} \mathcal{L}^\Phi(S)) \dots))$ and $\chi^{E,j} \mathcal{L}^{\Theta_{j+1}}(\dots \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2}u_f) \wedge \omega^{k-p-l+1}))) \dots)$ have disjoint support and the distance between them is bounded below by $(\frac{1}{2}r_E)^{k+2} s$. Further, from Lemma 3.7 and Lemma 7.4, we see that the mass of the current $(1 - \chi^{E,j-1}) \mathcal{L}^{\Theta_{j-1}}(\mathcal{L}^{\Theta_{j-2}}(\dots \mathcal{L}^{\Theta_1}(\chi_{s,t} \mathcal{L}^\Phi(S)) \dots))$ is bounded by $\|\Phi\|_\infty \|S\| t^{3/(2\delta+1)}$ up to a constant multiple independent of n, i, s, t, θ, Φ and S . Note that u_f is Hölder continuous and $\chi^{E,k+2}$ is smooth on X . So, in the sense of currents, we may write

$$\begin{aligned} dd^c(\chi^{E,k+2}u_f) &= d(\chi^{E,k+2} + u_f) \wedge d^c(\chi^{E,k+2} + u_f) - d\chi^{E,k+2} \wedge d^c\chi^{E,k+2} - du_f \wedge d^c u_f \\ &\quad + u_f dd^c\chi^{E,k+2} + \chi^{E,k+2} dd^c u_f. \end{aligned}$$

The first three terms are positive or negative currents and the last two terms can be written as a difference of positive currents of mass under control. Using Stokes' theorem, we have

$$\begin{aligned} & \left| \int_X d(\chi^{E,k+2} + u_f) \wedge d^c(\chi^{E,k+2} + u_f) \wedge \omega^{k-1} \right| = \left| \int_X (\chi^{E,k+2} + u_f) \wedge dd^c(\chi^{E,k+2} + u_f) \wedge \omega^{k-1} \right| \\ & \leq c \left(\sup_X |u_f| + 1 \right) (\|\chi^{E,k+2}\|_{C^2} + \|f_*(\omega)\| + 1) \end{aligned}$$

where $c > 0$ is a constant independent of n, i, s, t, θ, Φ and S . The second and third terms can be treated in the same way. By Lemma 3.7, the current $\chi^{E,j} \mathcal{L}^{\Theta_{j+1}}(\dots \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2}u_f) \wedge \omega^{k-p-l+1}))) \dots)$ can be written as a finite sum of positive or negative currents, the mass of each of which is bounded by s^{-2} up to a constant multiple independent of n, i, s, t, θ, Φ

and S . By Lemma 3.8 and Lemma 3.9,

$$\left| \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^{\Theta_1}(\dots \mathcal{L}^{\Theta_{j-2}}(\mathcal{L}^{\Theta_{j-1}} \underline{ig}((1 - \chi^{E,j-1}) \mathcal{L}^j(\chi^{E,j} \mathcal{L}^{j+1}(\dots \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2} u_f) \wedge \omega^{k-p-l+1})) \dots)))))) \right| \lesssim \|\Phi\|_\infty \|S\| s^{-2k} t^{3/(2\delta+1)}$$

The inequality \lesssim means \leq up to a constant multiple independent of n, i, s, t, θ, Φ and S . \square

Proof of Lemma 7.5. Our proof is by induction. Observe that $\|S\| = |\langle S, \omega^{k-p} \rangle| = |\langle \alpha_S, \omega^{k-p} \rangle| \leq c_m \|\alpha_S\|_\infty$ and therefore, Lemma 7.4 proves the case $l = 0$. So, we assume that the statement is true for $l - 1$ with $l \geq 1$. Let $\chi_{s,t}$ is the cut-off function in Lemma 7.3. Then, from the positivity of the integrand, we have

$$(7.1) \quad \int_{V_t \setminus E_s} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^l \wedge \omega^{k-p-l+1}) \leq \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^l \wedge \omega^{k-p-l+1})$$

$$= \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge \omega_f \wedge \omega^{k-p-l+1})$$

$$(7.2) \quad + \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c u_f \wedge \omega^{k-p-l+1}).$$

Since ω is a Kähler form, the bound of the integral (7.1) is obtained from the induction hypothesis.

$$|(7.1)| \lesssim 2 \|\omega_f\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty (r_E s)^{-2(k+l-1)-N_E} (t^{(\delta+1/2)^{-1}})^{\beta_{l-1}}.$$

We only need to estimate the integral (7.2). This is the part that we should be careful when compared to the works in [20] and [1]. We know that u_f is $(c_f (r_E (\frac{1}{2} r_E)^{k+3} s)^{-N_E}, \delta^{-1})$ -Hölder continuous on $X \setminus E_{r_E (\frac{1}{2} r_E)^{k+3} s}$ from Lemma 5.3. For a sufficiently small $\gamma > 0$ so that

$\overline{(X \setminus E_{(\frac{1}{2} r_E)^{k+3} s})_\gamma} \cap E_{r_E (\frac{1}{2} r_E)^{k+3} s} = \emptyset$ with $W_1 = E_{(\frac{1}{2} r_E)^{k+3} s}$ and $W_2 = E_{r_E (\frac{1}{2} r_E)^{k+3} s}$, Lemma 5.4 implies that we can find a smooth function \tilde{v}_γ defined in a neighborhood $X \setminus E_{(\frac{1}{2} r_E)^{k+3} s}$ such that

$$\|\tilde{v}_\gamma\|_{C^2, X \setminus E_{(\frac{1}{2} r_E)^{k+3} s}} \leq c_H \gamma^{-2(k+1)} \quad \text{and} \quad \|u_f - \tilde{v}_\gamma\|_{\infty, X \setminus E_{(\frac{1}{2} r_E)^{k+3} s}} \leq c_H c_f \left(r_E \left(\frac{1}{2} r_E \right)^{k+3} s \right)^{-N_E} \gamma^{\delta-1}$$

where $c_H > 0$ is the constant in Lemma 5.4 and is independent of s . We define

$$v_\gamma = \chi^{E,k+2} u_f + (1 - \chi^{E,k+2}) \tilde{v}_\gamma.$$

Then, (7.2) is equal to the sum

$$(7.3) \quad \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c v_\gamma \wedge \omega^{k-p-l+1})$$

$$(7.4) \quad + \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c (u_f - v_\gamma) \wedge \omega^{k-p-l+1}).$$

The integral (7.3) is equal to the sum

$$(7.5) \quad \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c (\chi^{E,k+2} u_f) \wedge \omega^{k-p-l+1})$$

$$(7.6) \quad + \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c ((1 - \chi^{E,k+2}) \tilde{v}_\gamma) \wedge \omega^{k-p-l+1})$$

We can write the integral (7.5) as below:

$$(7.5) = \sum_{j=1}^m \int \chi_{s,t} \mathcal{L}^\Phi(S) \wedge \mathcal{L}^{\Theta_1}(\dots \mathcal{L}^{\Theta_{j-2}}(\mathcal{L}^{\Theta_{j-1}}((1 - \chi^{E,j-1}) \mathcal{L}^{\Theta_j}(\chi^{E,j} \mathcal{L}^{\Theta_{j+1}}(\dots \chi^{E,m-1}(\mathcal{L}^{\Theta_m}(f_*(\omega)^{l-1} \wedge dd^c(\chi^{E,k+2} u_f) \wedge \omega^{k-p-l+1})) \dots)))))) \dots).$$

Notice that $\chi^{E,m} dd^c (\chi^{E,k+2} u_f) = dd^c (\chi^{E,k+2} u_f)$ is used in the very last line. Then, Lemma 7.6 implies that

$$|(7.5)| \lesssim \|\Phi\|_\infty \|S\| s^{-2k} t^{3/(2\delta+1)}.$$

Since $\|\tilde{v}_\gamma\|_{C^2, X \setminus E} (\frac{1}{2} r_E)^{k+3} \leq c_H \gamma^{-2(k+1)}$ and $\|\chi^{E,k+2}\|_{C^2} \lesssim s^{-2}$, the induction hypothesis implies

$$|(7.6)| \lesssim \gamma^{-2(k+1)} s^{-2} \cdot \|\Phi\|_\infty \|\alpha_S\|_\infty (r_E s)^{-2(k+l-1)-N_E} \left(t^{(\delta+1/2)^{-1}}\right)^{\beta_{l-1}}$$

and therefore, we get

$$|(7.3)| \lesssim \|\Phi\|_\infty \|\alpha_S\|_\infty \gamma^{-2(k+1)} s^{-2(k+l)-N_E} \left(t^{(\delta+1/2)^{-1}}\right)^{\beta_{l-1}}$$

Since \mathcal{L}^m is closed, we have

$$(7.4) = \int dd^c \chi_{s,t} \wedge \mathcal{L}^\Phi(S) \wedge \mathcal{L}^m((u_f - v_\gamma) f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1}).$$

Observe that

$$u_f - v_\gamma = (1 - \chi^{E,k+2})(u_f - \tilde{v}_\gamma) \text{ and } \|u_f - \tilde{v}_\gamma\|_{\infty, X \setminus E} (\frac{1}{2} r_E)^{k+3} \lesssim s^{-N_E} \gamma^{\delta^{-1}}.$$

Since $\|\chi_{s,t}\|_{\text{DSH}} \lesssim s^{-2}$, we can find two positive closed $(1,1)$ -currents $\vartheta_{s,t}^\pm$ such that $\vartheta_{s,t}^+ - \vartheta_{s,t}^- = dd^c \chi_{s,t}$ and $\|\vartheta_{s,t}^\pm\| \lesssim s^{-2}$. Hence, since $\vartheta_{s,t}^\pm$, Φ , S , $f_*(\omega)$, ω are all positive and closed, a cohomological argument gives us

$$\begin{aligned} |(7.4)| &\leq 2 \|u_f - v_\gamma\|_\infty \|\Phi\|_\infty \left(\int \vartheta_{s,t}^+ \wedge \mathcal{L}^{\omega_{\tilde{x}}^{k-1}}(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1}) \right) \\ &= 2 \|u_f - v_\gamma\|_\infty \|\Phi\|_\infty \left(\int \vartheta_{s,t}^+ \wedge \mathcal{L}^{\omega_{\tilde{x}}^{k-1}}(\alpha_S) \wedge \mathcal{L}^m(\omega_f^{l-1} \wedge \omega^{k-p-l+1}) \right) \\ &= 2 \|u_f - v_\gamma\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty \|\omega_f\|^{l-1} \left(\int \vartheta_{s,t}^+ \wedge \mathcal{L}^{\omega_{\tilde{x}}^{k-1}}(\omega^p) \wedge \mathcal{L}^m(\omega^{k-p}) \right) \\ &\lesssim 2 s^{-N_E} \gamma^{\delta^{-1}} \|\Phi\|_\infty \|\alpha_S\|_\infty \|\omega_f\|^{l-1} s^{-2} \lesssim \|\Phi\|_\infty \|\alpha_S\|_\infty \gamma^{\delta^{-1}} s^{-2-N_E}. \end{aligned}$$

We take $\gamma = t^{(2(k+1)(2\delta+1))^{-l} \delta^{-(l-1)}}$. Then, from the above, we have

$$\begin{aligned} |(7.3)| &\lesssim \|\Phi\|_\infty \|\alpha_S\|_\infty \gamma^{-2(k+1)} s^{-2(k+l)-N_E} \left(t^{(\delta+1/2)^{-1}}\right)^{\beta_{l-1}} \\ &\lesssim \|\Phi\|_\infty \|\alpha_S\|_\infty s^{-2(k+l)-N_E} t^{\frac{-2(k+1)}{(2(k+1)(2\delta+1))^l \delta^{l-1}} + \frac{1}{(2(k+1)(2\delta+1))^{l-1} \delta^{l-1} (\delta+1/2)}} \leq \|\Phi\|_\infty \|\alpha_S\|_\infty s^{-2(k+l)-N_E} t^{\beta_l} \end{aligned}$$

and

$$|(7.4)| \lesssim \|\Phi\|_\infty \|\alpha_S\|_\infty s^{-N_E} \gamma^{\delta^{-1}} s^{-2} \lesssim \|\Phi\|_\infty \|\alpha_S\|_\infty s^{-2-N_E} t^{\frac{1}{(2(k+1)(2\delta+1))^l \delta^l}} \leq \|\Phi\|_\infty \|\alpha_S\|_\infty s^{-2-N_E} t^{\beta_l}.$$

Combining these two gives the estimate of (7.2), which proves the desired statement for l . \square

Remark 7.7. In the above proof, from the relationship between $s_{n,i}$ and $t_{n,i}$, our choice of γ satisfies the desired smallness for the argument if $\varepsilon > 0$ is sufficiently small. So, our proof works.

We consider $\langle \chi_{n,i,1} \mathcal{L}^{u_\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle$. Recall the definition of u_τ in the beginning of Section 3. For all $\tau \in \mathbb{C} \setminus \{0\}$ with sufficiently small $|\tau| < 1$, we have

$$(\log |\tau| - 1) + (u - u_\tau) \leq u \leq 0$$

and therefore,

$$(7.7) \quad \begin{aligned} & (\log |\tau| - 1) \mathcal{L}^\Phi(S) + \mathcal{L}^{(u-u_\tau)\Phi}(S) \leq \mathcal{L}^{u\Phi}(S) \leq 0 \quad \text{and} \\ & (\log |\tau| - 1) \langle \chi_{s_n, i, t_n, i} \mathcal{L}^\Phi(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle + \langle \chi_{s_n, i, t_n, i} \mathcal{L}^{(u-u_\tau)\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle \\ & \leq \langle \chi_{s_n, i, t_n, i} \mathcal{L}^{u\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle \leq 0 \end{aligned}$$

The first term containing $(\log |\tau| - 1) \mathcal{L}^\Phi(S)$ can be deduced from the previous estimate and $u - u_\tau$ is also negative and has support in a small neighborhood of $\widehat{\Delta}$. The estimate about the second term is essentially the same as Lemma 2.3.9 and Lemma 2.3.10 in [20].

Lemma 7.8. *We have*

$$\left| \langle \mathcal{L}^{(u-u_\tau)\Phi}(S), \mathcal{L}^m(\omega^{k-p+1}) \rangle \right| \lesssim \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|S\| |\tau|^{1/2}$$

The inequality \lesssim means \leq up to a positive constant multiple independent of θ , S and τ .

Proof. From the negativity of $\mathcal{L}^{(u-u_\tau)\Phi}(S)$, we have

$$\left| \langle \mathcal{L}^{(u-u_\tau)\Phi}(S), \mathcal{L}^m(\omega^{k-p+1}) \rangle \right| \lesssim \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \left| \langle \mathcal{L}^{(u-u_\tau)\Phi}(S), \omega^{k-p+1} \rangle \right|$$

Since ω^{k-p+1} is smooth, so we have

$$\langle \mathcal{L}^{(u-u_\tau)\Phi}(S), \omega^{k-p+1} \rangle = \int_{X \times X \setminus \Delta} \pi_* [(u - u_\tau)\Phi] \wedge \pi_1^*(S) \wedge \pi_2^*(\omega^{k-p+1}).$$

Since we will not use the closedness, by disintegration as in Lemma 7.4, we can write it as

$$\int_{y \in X} \left[\int_{x \in X \setminus \{y\}} \Psi_S(y) \lrcorner [(u - u_\tau)\Phi](x, y) \wedge \omega^{k-p+1}(x) \right] d\mu(y)$$

in terms of Federer's notation. Note that the form $\Psi_S(y) \lrcorner [\pi_* [(u - u_\tau)\Phi]]$ has a singularity of $-\|\Phi\|_\infty \text{dist}(x, y)^{2-2k} \log \text{dist}(x, y)$ and its support lies inside a ball with center at y and of radius $c\tau$ where $c > 0$ is a uniform constant independent of τ and one might think of μ as the trace measure of S . Hence, $\int_X d\mu$ can be bounded in terms of $\|S\|$. Since $\|\Psi_S\| \leq 1$, the integrand $\int_{x \in X \setminus \{y\}} \Psi_S(y) \lrcorner [\pi_* [(u - u_\tau)\Phi]] \wedge \pi_2^*(\omega^{k-p+1})$ is bounded by a uniform constant multiple of $\tau^{1/2} \|\Phi\|_\infty$. So, it is proved. \square

Lemma 7.9. *For every integer l with $0 \leq l \leq k - p + 1$, we have*

$$\left| \int \mathcal{L}^{(u-u_\tau)\Phi}(S) \wedge \mathcal{L}^m(f_*(\omega)^l \wedge \omega^{k-p-l+1}) \right| \lesssim \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty |\tau|^{\frac{1}{2(4kd_k)^l}}$$

The inequality \lesssim means \leq up to a positive constant multiple independent of θ , S , α_S and τ .

Proof. We prove it by induction. We have $\|S\| = |\langle S, \omega^{k-p} \rangle| = |\langle \alpha_S, \omega^{k-p} \rangle| \leq c_m \|\alpha_S\|_\infty$. The previous lemma corresponds to the case of $l = 0$. Assume that it is true for $l - 1$.

$$(7.8) \quad \begin{aligned} & \int \mathcal{L}^{(u-u_\tau)\Phi}(S) \wedge \mathcal{L}^m(f_*(\omega)^l \wedge \omega^{k-p-l+1}) \\ & = \int \mathcal{L}^{(u-u_\tau)\Phi}(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge \omega_f \wedge \omega^{k-p-l+1}) \end{aligned}$$

$$(7.9) \quad + \int \mathcal{L}^{(u-u_\tau)\Phi}(S) \wedge \mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c u_f \wedge \omega^{k-p-l+1}).$$

The estimate of (7.8) comes from the induction hypothesis. Since u_f is (K, d_k^{-1}) -Hölder continuous on X for some $K > 0$, we can use Lemma 5.4 to find a smooth function v_γ such that

$\|u_f - v_\gamma\|_\infty \leq K\gamma^{1/d_k}$ and $\|v_\gamma\|_{C^\infty} \lesssim K|\gamma|^{-2(k+1)}$ where $\gamma = |\tau|^{\frac{1}{2(4k)d_k^{l-1}}}$. Then, since \mathcal{L}^m is closed,

$$(7.9) = \int_{\overline{X \times X}} (u - u_\tau) \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c u_f \wedge \omega^{k-p-l+1}))$$

$$(7.10) = \int_{\overline{X \times X}} (u - u_\tau) \Phi \wedge \Pi_1^*(S) \wedge dd^c \Pi_2^*(\mathcal{L}^m((u_f - v_\gamma) f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1}))$$

$$(7.11) + \int_{\overline{X \times X}} (u - u_\tau) \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m(f_*(\omega)^{l-1} \wedge dd^c(v_\gamma) \wedge \omega^{k-p-l+1}))$$

Since the $*$ -norm of $dd^c(u - u_\tau)$ is uniformly bounded independently of τ , we use cohomological arguments to get an estimate of (7.10) as follows:

$$\begin{aligned} |(7.10)| &= \left| \int_{\overline{X \times X}} dd^c(u - u_\tau) \wedge \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m((u_f - v_\gamma) f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1})) \right| \\ &\leq \left| \int_{\overline{X \times X}} (m_{\overline{\Delta}} \omega_{\overline{\mathfrak{X}}} + dd^c u) \wedge \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m((u_f - v_\gamma) f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1})) \right| \\ &\quad + \left| \int_{\overline{X \times X}} (m_{\overline{\Delta}} \omega_{\overline{\mathfrak{X}}} + dd^c u_\tau) \wedge \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m((u_f - v_\gamma) f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1})) \right| \\ &\leq 2\|u_f - v_\gamma\|_\infty \left| \int_{\overline{X \times X}} m_{\overline{\Delta}} \omega_{\overline{\mathfrak{X}}} \wedge \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m(f_*(\omega)^{l-1} \wedge \omega^{k-p-l+1})) \right| \\ &= 2\|u_f - v_\gamma\|_\infty \left| \int_{\overline{X \times X}} m_{\overline{\Delta}} \omega_{\overline{\mathfrak{X}}} \wedge \Phi \wedge \Pi_1^*(\alpha_S) \wedge \Pi_2^*(\mathcal{L}^m(\omega_f^{l-1} \wedge \omega^{k-p-l+1})) \right| \\ &\lesssim 2\|u_f - v_\gamma\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty \|\omega_f\|_\infty^{l-1} \left| \int_{\overline{X \times X}} m_{\overline{\Delta}} \omega_{\overline{\mathfrak{X}}} \wedge \omega_{\overline{\mathfrak{X}}}^{k-1} \wedge \Pi_1^*(\omega^p) \wedge \Pi_2^*(\mathcal{L}^m(\omega^{k-p})) \right| \\ &\lesssim \|u_f - v_\gamma\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty \|\mathcal{L}^m(\omega^{k-p+1})\| \lesssim \gamma^{1/d_k} \|\Phi\|_\infty \|\alpha_S\|_\infty = \|\Phi\|_\infty \|\alpha_S\|_\infty |\tau|^{\frac{1}{2(4kd_k)^l}}. \end{aligned}$$

The currents S , $f_*(\omega)$ and ω are all positive and $u - u_\tau \leq 0$. So, the induction hypothesis gives

$$\begin{aligned} |(7.11)| &\lesssim \|v_\gamma\|_{C^2} \left| \int_{\overline{X \times X}} (u - u_\tau) \Phi \wedge \Pi_1^*(S) \wedge \Pi_2^*(\mathcal{L}^m(f_*(\omega)^{l-1} \wedge \omega^{k-p-l+2})) \right| \\ &\lesssim |\gamma|^{-2(k+1)} \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty |\tau|^{\frac{1}{2(4kd_k)^{l-1}}} \\ &= \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty |\tau|^{\frac{-2(k+1)}{2(4k)d_k^{l-1}} + \frac{1}{2(4kd_k)^{l-1}}} \\ &= \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty |\tau|^{\frac{2(k-1)}{2(4k)d_k^{l-1}}} < \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty |\tau|^{\frac{1}{2(4kd_k)^l}} \end{aligned}$$

□

The current $\mathcal{L}^{u\Phi}(S)$ is negative and the current $\mathcal{L}^m(f_*(\omega^{k-p+1}))$ is positive. If we take $l = k - p + 1$ and $\tau = e^{1-t_{n,i}^{-\beta_{k-p+2/2}}}$, then the inequality (7.7), Lemma 7.5 and Lemma 7.9 give us

$$\begin{aligned} 0 &\geq \langle \chi_{n,i,1} \mathcal{L}^{u\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle \geq \langle \chi_{s_{n,i}, t_{n,i}} \mathcal{L}^{u\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle \\ &\geq - \left(t_{n,i}^{-\beta_{k-p+2/2}} \cdot s^{-4k-N_E} \left(t_{n,i}^{(\delta+1/2)^{-1}} \right)^{\beta_{k-p+1}} + e^{\frac{1-t_{n,i}^{-\beta_{k-p+2/2}}}{2(4kd_k)^{k-p+1}}} \right) \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty \\ &\gtrsim - \left(s^{-4k-N_E} t_{n,i}^{\beta_{k-p+2}} + \frac{2(4kd_k)^{k-p+1}}{t_{n,i}^{-\beta_{k-p+2/2}} - 1} \right) \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty \\ &\gtrsim -s_{n,i}^{-4k-N_E} t_{n,i}^{\beta_{k+1/2}} \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty \end{aligned}$$

if $\varepsilon > 0$ is sufficiently small. So, we get

Lemma 7.10. *Let n and i be positive integers such that $1 \leq i \leq n$. Then, for all sufficiently small $\varepsilon > 0$, we have*

$$\left| \langle \chi_{n,i,1} \mathcal{L}^{u\Phi}(S), \mathcal{L}^m(f_*(\omega^{k-p+1})) \rangle \right| \lesssim \|\mathcal{L}^m(\omega^{k-p+1})\|_\infty \|\Phi\|_\infty \|\alpha_S\|_\infty s_{n,i}^{-4k-N_E} t_{n,i}^{\beta_{k+1}/2}$$

where $\beta_{k+1} = (2(k+1)(2\delta+1))^{-(k+1)} \delta^{-(k+1)}$ and the inequality \lesssim means \leq up to a positive constant multiple independent of n, i, θ, S and α_S .

Now, we are ready to estimate $d^{-i} \langle \chi_{n,i,1} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$. Recall Lemma 6.3. We have $R_{n,i} = (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^+ + (\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^- - (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^- - (\Lambda(R_{n,i-1}^-))_{\varepsilon_{n,i}}^+$ for $i \geq 2$. When $i = 1$, we simply remove terms containing $R_{n,0}^-$. Due to Proposition 2.7, the Green potential kernel can be written as $\mathcal{L}^K = \mathcal{L}^{K^+} - \mathcal{L}^{K^-}$. Therefore, $|\langle \chi_{n,i,1} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle|$ is bounded by the sum of $|\langle \chi_{n,i,1} \mathcal{L}^{K^\pm}(L^{n-i}(S)), \Lambda(R_{n,i-1}^\pm) \rangle|$ and $|\langle \chi_{n,i,1} \mathcal{L}^{K^\pm}(L^{n-i}(S)), (\Lambda(R_{n,i-1}^\pm))_{\varepsilon_{n,i}}^\pm \rangle|$. The forms K_\pm are negative and the current $L^{n-i}(S)$ is positive. When $i \geq 2$, each term can be estimated by Lemma 7.5, Lemma 7.10 and Lemma 6.3 as below:

$$\begin{aligned} & \left| \langle \chi_{n,i,1} \mathcal{L}^{K^+}(L^{n-i}(S)), \Lambda(R_{n,i-1}^+) \rangle \right| = \left| \langle \chi_{n,i,1} \mathcal{L}^{u(\eta+m_K \omega_{\tilde{x}}^{k-1}) - m_K \omega_{\tilde{x}}^{k-1}}(L^{n-i}(S)), \Lambda(R_{n,i-1}^+) \rangle \right| \\ & \lesssim \|R_{n,i-1}^+\|_\infty (\|\eta\|_\infty + m_K) \left| \langle \chi_{n,i,1} (\mathcal{L}^{u\omega_{\tilde{x}}^{k-1}}(L^{n-i}(S)) + \mathcal{L}^{\omega_{\tilde{x}}^{k-1}}(L^{n-i}(S))), \Lambda(\omega^{k-p+1}) \rangle \right| \\ & \lesssim \|R_{n,i-1}^+\|_\infty \left| \langle \chi_{n,i,1} (\mathcal{L}^{u\omega_{\tilde{x}}^{k-1}}(L^{n-i}(S)) + \mathcal{L}^{\omega_{\tilde{x}}^{k-1}}(L^{n-i}(S))), f_*(\omega)^{k-p+1} \rangle \right| \\ & \lesssim \|R_{n,i-1}^+\|_\infty \|L^{n-i}(\alpha_S)\|_\infty s_{n,i}^{-4k-N_E} t_{n,i}^{\beta_{k+1}/2} \\ & \lesssim \|R\|_{C^1}(C_m)^{i-1} \left(\prod_{j=1}^{i-1} \varepsilon_{n,j} \right)^{-6k^2} (1 + \|f\|_{C^1})^{kn} \|\alpha_S\|_\infty s_{n,i}^{-4k-N_E} t_{n,i}^{\beta_{k+1}/2} \\ & \lesssim (C_m(1 + \|f\|_{C^1})^k)^n \varepsilon_{n,i-1}^{-12k^2} s_{n,i}^{-4k-N_E} t_{n,i}^{\beta_{k+1}/2} \\ & \lesssim (C_m(1 + \|f\|_{C^1})^k)^n \varepsilon^{nC(4k+N_E)} \left[(40k^2\delta)^{6k(i-1)} \left(\frac{(40k^2\delta)^{6k}}{20k(2(k+1)(2\delta+1)\delta)^{k+1}} - 12k^2 \right) - i \right] \\ & \leq (C_m(1 + \|f\|_{C^1})^k)^n \varepsilon^{nC(4k+N_E)} \left[(40k^2\delta)^{6k(i-1)} ((40k^2\delta)^{3k} - 12k^2) - i \right] < \varepsilon \end{aligned}$$

for all sufficiently small $\varepsilon > 0$. For $i = 1$, it can be done in the same way. Since the operation $(\cdot)_{\varepsilon_{n,i}}^+$ is the $(k+2)$ -times compositions of either \mathcal{L}_θ^+ or \mathcal{L}^- , and the real numbers $\varepsilon_{n,i}$ and $t_{n,i}$ are sufficiently small compared to $s_{n,i}$ when $\varepsilon > 0$ is small enough, we get

$$\begin{aligned} & \left| \langle \chi_{n,i,1} \mathcal{L}^{K^+}(L^{n-i}(S)), (\Lambda(R_{n,i-1}^+))_{\varepsilon_{n,i}}^+ \rangle \right| \lesssim \|R_{n,i-1}^+\|_\infty \left| \langle \chi_{n,i,1} \mathcal{L}^{K^+}(L^{n-i}(S)), (\Lambda(\omega^{k-p+1}))_{\varepsilon_{n,i}}^+ \rangle \right| \\ & \lesssim \|R_{n,i-1}^+\|_\infty \left| \langle \chi_{n,i,1} \mathcal{L}^{K^+}(L^{n-i}(S)), (f_*(\omega)^{k-p+1})_{\varepsilon_{n,i}}^+ \rangle \right| < \varepsilon. \end{aligned}$$

in the same way as previously. All other terms can be computed in the same way. For the terms containing α_S , we can apply the same argument as well. Hence, we have

$$\left| \langle \chi_{n,i,1} \mathcal{L}^{K^\pm}(L^{n-i}(S)), \Lambda(R_{n,i-1}^\pm) \rangle \right|, \left| \langle \chi_{n,i,1} \mathcal{L}^{K^\pm}(L^{n-i}(S)), (\Lambda(R_{n,i-1}^\pm))_{\varepsilon_{n,i}}^\pm \rangle \right| \lesssim \varepsilon$$

for all positive integers n and i such that $1 \leq i \leq n$ and for all sufficiently small $\varepsilon > 0$. Here, the inequality \lesssim means \leq up to a positive constant multiple independent of n, i and S .

Lemma 7.11. *For all n and i be positive integers such that $1 \leq i \leq n$ and for all sufficiently small $\varepsilon > 0$, we have*

$$\left| \langle \chi_{n,i,1} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \right| \lesssim \varepsilon.$$

Here, the inequality \lesssim means \leq up to a positive constant multiple independent of n, i and S .

7.2. Estimate of $d^{-i}\langle \chi_{n,i,2} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$ in $W_{n,i,2}$. We will use the property that in $W_{n,i,2}$, $L^{n-i}(S)$ is smooth.

Lemma 7.12. *Assume that $S \in \mathcal{D}_p$ is smooth in E_{s/r_E^3} for $s > 0$. Let Φ be a smooth positive closed current of bidegree $(k-1, k-1)$ on \mathfrak{X} . The currents $\chi_s^E \mathcal{L}^{u\Phi}(S)$ and $\chi_s^E \mathcal{L}^\Phi(S)$ is C^1 . Furthermore,*

$$\begin{aligned} \|\chi_s^E \mathcal{L}^{u\Phi}(S)\|_{C^1} &\lesssim s^{1-2k} (\|S\|_{\infty, E_{s/r_E^2}} + \|S\|_*) \|\Phi\|_{C^1} \quad \text{and} \\ \|\chi_s^E \mathcal{L}^\Phi(S)\|_{C^1} &\lesssim s^{1-2k} (\|S\|_{\infty, E_{s/r_E^2}} + \|S\|_*) \|\Phi\|_{C^1}. \end{aligned}$$

Here, the inequality \lesssim means \leq up to a constant multiple independent of s , S and Φ .

Proof. The same proof works for both. So, we only consider $\chi_s^E \mathcal{L}^{u\Phi}(S)$. We have

$$\chi_s^E \mathcal{L}^{u\Phi}(S) = \chi_s^E \mathcal{L}^{u\Phi}(\chi_{s/r_E^2}^E S) + \chi_s^E \mathcal{L}^{u\Phi}\left(\left(1 - \chi_{s/r_E^2}^E\right) S\right)$$

As in [22, Section 2.3], $\pi_*(u\Phi)$ is smooth outside Δ and its gradient satisfies $|\nabla \pi_*(u\Phi)| \lesssim \|\Phi\|_{C^1} (\text{dist}(\cdot, \Delta))^{1-2k}$. Hence, we have

$$\begin{aligned} \left\| \chi_s^E \mathcal{L}^{u\Phi}(\chi_{s/r_E^2}^E S) \right\|_{C^1} &\leq 2 \|\chi_s^E\|_{C^1} \left\| \mathcal{L}^{u\Phi}(\chi_{s/r_E^2}^E S) \right\|_{C^1} \\ &\lesssim \|\chi_s^E\|_{C^2} \left\| \chi_{s/r_E^2}^E S \right\|_{\infty} \|\Phi\|_{C^1} \leq s^{-2} \|S\|_{\infty, E_{s/r_E^2}} \|\Phi\|_{C^1} \end{aligned}$$

Let φ be a smooth test form on X . Then, we have

$$\begin{aligned} \left\langle \chi_s^E \mathcal{L}^{u\Phi}\left(\left(1 - \chi_{s/r_E^2}^E\right) S\right), \varphi \right\rangle &= \left\langle \mathcal{L}^{u\Phi}\left(\left(1 - \chi_{s/r_E^2}^E\right) S\right), \chi_s^E \varphi \right\rangle \\ &= \int_X \chi_s^E \left[\int_{X \setminus \{z\}} \pi_*(u\Phi)(z, \xi) \wedge \left(\left(1 - \chi_{s/r_E^2}^E\right) S\right)(\xi) \right] \wedge \varphi. \end{aligned}$$

Since $\text{dist}\left(\text{supp} \chi_s^E, \text{supp}\left(1 - \chi_{s/r_E^2}^E\right)\right) \geq (r_E^{-1} - 1)s$, again according to [22, Section 2.3], we see that the current $\chi_s^E \left[\int_{X \setminus \{z\}} \pi_*(u\Phi)(z, \xi) \wedge \left(\left(1 - \chi_{s/r_E^2}^E\right) S\right)(\xi) \right]$ is smooth and

$$\left\| \chi_s^E \left[\int_{X \setminus \{z\}} \pi_*(u\Phi)(z, \xi) \wedge \left(\left(1 - \chi_{s/r_E^2}^E\right) S\right)(\xi) \right] \right\|_{C^1} \lesssim s^{1-2k} \|S\|_* \|\Phi\|_{C^1}.$$

□

Lemma 7.13. *Let $\tilde{R} \in \mathcal{D}_{k-p+1}$. Under the hypotheses in Lemma 7.12, for a sufficiently small $\theta > 0$,*

$$\begin{aligned} \left| \langle \chi_s^E \mathcal{L}^{u\Phi}(S), \tilde{R} - (\tilde{R})_\theta \rangle \right| &\lesssim s^{1-2k} \theta \|S\|_{\infty, E_{s/r_E^2}} \|\tilde{R}\|_* \|\Phi\|_{C^1} \quad \text{and} \\ \left| \langle \chi_s^E \mathcal{L}^\Phi(S), \tilde{R} - (\tilde{R})_\theta \rangle \right| &\lesssim s^{1-2k} \theta \|S\|_{\infty, E_{s/r_E^2}} \|\tilde{R}\|_* \|\Phi\|_{C^1}. \end{aligned}$$

Here, the inequality \lesssim means \leq up to a constant multiple independent of s , S , θ and Φ .

Proof. As previously, we consider the first estimate. The same applies to the second. We have

$$\langle \chi_s^E \mathcal{L}^{u\Phi}(S), \tilde{R} - (\tilde{R})_\theta \rangle = \langle \chi_s^E \mathcal{L}^{u\Phi}(S) - (\chi_s^E \mathcal{L}^{u\Phi}(S))_\theta, \tilde{R} \rangle.$$

Lemma 7.12 implies that $\chi_s^E \mathcal{L}^{u\Phi}(S)$ is a form with C^1 coefficients and Lemma 3.1 implies

$$\left\| \chi_s^E \mathcal{L}^{u\Phi}(S) - (\chi_s^E \mathcal{L}^{u\Phi}(S))_\theta \right\|_{\infty} \lesssim s^{1-2k} \theta (\|S\|_{\infty, E_{s/r_E^2}} + \|S\|_*) \|\Phi\|_{C^1}.$$

Hence, we get the estimate. □

Corollary 7.14. *Assume that S be as in Theorem 1.1. For all sufficiently small $\varepsilon > 0$, we have*

$$\left| \langle \chi_{n,i,2} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \right| < (\|S\|_{\infty, E_{s_0}} + \|\alpha_S\|_{\infty}) \varepsilon.$$

Here, the inequality \lesssim means \leq up to a positive constant multiple independent of n , i and S .

Proof. Notice that $U_{L^{n-i}(S-\alpha_S)}$ is the difference of $\mathcal{L}^{K_+}(L^{n-i}(S-\alpha_S)) - \mathcal{L}^{K_-}(L^{n-i}(S-\alpha_S))$ where $K_+ = u\left(\eta + m_K \omega_{\frac{\mathbb{X}}{2}}^{k-1}\right) - \left(m_K \omega_{\frac{\mathbb{X}}{2}}^{k-1}\right)$ and $K_- = u\left(m_K \omega_{\frac{\mathbb{X}}{2}}^{k-1}\right) - \left(\delta_{\frac{\mathbb{X}}{2}} + m_K \omega_{\frac{\mathbb{X}}{2}}^{k-1}\right)$. From Lemma 7.13 and Lemma 6.3, we get

$$\begin{aligned} & \left| \langle \chi_{n,i,2} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \right| = \left| \langle \chi_{n,i,2} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - (\Lambda(R_{n,i-1}))_{\varepsilon_{n,i}} \rangle \right| \\ & \lesssim s_{n,i}^{1-2k} \varepsilon_{n,i} \left(\|L^{n-i}(S-\alpha_S)\|_{\infty, E_{s/r_E^2}} + \|L^{n-i}(S)\| + \|L^{n-i}(\alpha_S)\| \right) \|\Lambda(R_{n,i-1})\|_* . \end{aligned}$$

By Lemma 6.2, for all sufficiently small $\varepsilon > 0$, S is smooth in E_{ε/r_E^3} and $L^{n-i}(S-\alpha)$ is smooth in $E_{s_{n,i}/r_E^3}$. From direct computations, we get

$$\|L^{n-i}(S-\alpha_S)\|_{\infty, E_{s_{n,i}/r_E^3}} \leq (1 + \|f\|_{C^1})^{k(n-i)} \|S-\alpha_S\|_{\infty, E_{s_0}}$$

where S is smooth in E_{s_0} for some fixed $s_0 > 0$. Hence, together with (6.3) and Lemma 6.3,

$$\begin{aligned} & \left| \langle \chi_{n,i,2} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \right| \\ & \lesssim ((1 + \|f\|_{C^1})^{kn} \|S-\alpha_S\|_{\infty, E_{s_0}} + 2c_m \left(\frac{5}{4}\right)^{N_f n} \|\alpha_S\|_{\infty}) (\|\Lambda(R_{n,i-1}^+)\| + \|\Lambda(R_{n,i-1}^-)\|) s_{n,i}^{1-2k} \varepsilon_{n,i} \\ & \lesssim \left(\left(\left(\frac{5}{4}\right)^{N_f} + \|f\|_{C^1} \right)^{kn} (\|S\|_{\infty, E_{s_0}} + \|\alpha_S\|_{\infty}) \right) c_m \left(\frac{5}{4}\right)^{N_f} C_m^i \left(\prod_{j=1}^{i-1} \varepsilon_{n,j} \right)^{-6k^2} s_{n,i}^{1-2k} \varepsilon_{n,i} \\ & \lesssim (\|S\|_{\infty, E_{s_0}} + \|\alpha_S\|_{\infty}) \left(\left(\frac{5}{4}\right)^{N_f} + \|f\|_{C^1} \right)^{kn} C_m^i \varepsilon_{n,i-1}^{-12k^2} s_{n,i}^{1-2k} \varepsilon_{n,i} \lesssim (\|S\|_{\infty, E_{s_0}} + \|\alpha_S\|_{\infty}) \varepsilon \end{aligned}$$

for all sufficiently small $\varepsilon > 0$ where $R_{n,i-1}^{\pm}$ are smooth positive closed currents in Lemma 6.3. By plugging-in $s_{n,i}$ and $\varepsilon_{n,i}$, we see the desired estimate. \square

7.3. Estimate of $d^{-i}\langle \chi_{n,i,3} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle$ in $W_{n,i,3}$. In this region, $\Lambda(R_{n,i})$ is smooth.

Lemma 7.15 (Lemma 5.4.7 in [20]). *Let $\alpha \geq 0$. For all sufficiently small $t > 0$, we have*

$$\|\Lambda(R)\|_{\mathcal{C}^\alpha(V_i^c)} \lesssim \|R\|_{\mathcal{C}^\alpha} t^{-(4+\alpha)k},$$

for any smooth form R of bidegree (p, p) with $0 \leq p \leq k$. The inequality is up to a constant multiple independent of t and R .

Corollary 7.16. *For all sufficiently small $\varepsilon > 0$,*

$$d^{-i}\langle \chi_{n,i,3} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \lesssim \varepsilon.$$

Here, the inequality \lesssim means \leq up to a positive constant multiple independent of n , i and S .

Proof. Note that the $*$ -norm of the Green potential is uniformly bounded if the mass of S is bounded. Indeed, for $S \in \mathcal{C}_p$, we have

$$\left| \langle \mathcal{L}^{K_{\pm}}(S), \omega^{k-p+1} \rangle \right| = \left| \langle S, \mathcal{L}^{K_{\pm}}(\omega^{k-p+1}) \rangle \right| \lesssim \max \left\{ \|\mathcal{L}^{K_{\pm}}(\omega^{k-p+1})\|_{\infty} \right\} \|S\|$$

and $U_S = \mathcal{L}^{K_+}(S) - \mathcal{L}^{K_-}(S)$. So, from (6.3), Lemma 3.2 and Lemma 6.3, we get

$$\begin{aligned} & \left| d^{-i}\langle \chi_{n,i,3} U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \right| \lesssim \|\Lambda(R_{n,i-1}) - R_{n,i}\|_{\infty, W_{n,i,3}} \|L^{n-i}(S-\alpha_S)\|_* \\ & \lesssim \|\Lambda(R_{n,i-1})\|_{C^1, W_{n,i,3}} \varepsilon_{n,i} \|L^{n-i}(S-\alpha_S)\|_* \lesssim \|R_{n,i-1}\|_{C^1} t_{n,i}^{-5k} \varepsilon_{n,i} \|L^{n-i}(S-\alpha_S)\|_* \\ & \lesssim \|R\|_{C^1} (C_m)^{i-1} \left(\prod_{j=1}^{i-1} \varepsilon_{n,j} \right)^{-6k^2} \varepsilon_{n,i}^{1/2} \left(3c_m \left(\frac{5}{4}\right)^{N_f n} \|\alpha_S\|_{\infty} \right) \lesssim C_m^n \left(\frac{5}{4}\right)^{N_f n} \varepsilon_{n,i-1}^{-12k^2} \varepsilon_{n,i}^{1/2} \leq \varepsilon \end{aligned}$$

for all sufficiently small $\varepsilon > 0$. \square

From Lemma 7.11, Corollary 7.14, Corollary 7.16, we have

Lemma 7.17. For all sufficiently small $\varepsilon > 0$,

$$d^{-i}\langle U_{L^{n-i}(S-\alpha_S)}, \Lambda(R_{n,i-1}) - R_{n,i} \rangle \lesssim \varepsilon.$$

Here, the inequality \lesssim means \leq up to a positive constant multiple independent of n , i and S .

8. ESTIMATES OF $d^{-n}\langle U_{S-\alpha_S}, R_{n,n} \rangle$

For this estimate, we use a version of exponential estimate [22, Theorem 3.2.6].

Theorem 8.1 (Theorem 3.2.6 in [22]). Let S be a current in \mathcal{D}_p and \mathcal{U}_S be the β normalized superpotential of S . Then we have for R smooth in \mathcal{D}_{k-p+1}^0 with $\|R\|_* \leq 1$,

$$|\mathcal{U}_S(R)| \leq c\|S\|_*(1 + \log^+ \|R\|_{C^1}),$$

where $\log^+ := \max\{\log, 0\}$ and $c > 0$ is a constant independent of S and R .

Corollary 8.2. For all sufficiently small $\varepsilon > 0$,

$$|d^{-n}\langle U_{S-\alpha_S}, R_{n,n} \rangle| \lesssim \|S\|(1 + \log^+ \|R_{n,n}\|_{C^1}) \lesssim \log(\varepsilon_{n,n}^{-6k^2}).$$

Here, the inequality \lesssim means \leq up to a positive constant multiple independent of n and S .

Proof. Let R_{\pm} be smooth positive closed currents such that $R \in \mathcal{D}_{k-p+1}^0$, $R = R_+ - R_-$ and $\|R_{\pm}\|_{C^1} \leq 1$. Then, from Proposition 3.5 and Lemma 6.3, we get

$$\begin{aligned} |d^{-n}\langle U_{S-\alpha_S}, R_{n,n} \rangle| &\lesssim d^{-n}\|S\| |1 + \log^+ \|R_{n,n}\|_{C^1}| \leq d^{-n}\|S\| |1 + \log^+ (\|(R_+)_{n,n}\|_{C^1} + \|(R_-)_{n,n}\|_{C^1})| \\ &\lesssim d^{-n} \left| 1 + 2 \log \left(2\|R\|_{C^1} C_m^n \left(\prod_{j=1}^n \varepsilon_{n,j} \right)^{-6k^2} \right) \right| \leq d^{-n} \left| 1 + 2 \log \left(2\|R\|_{C^1} C_m^n \varepsilon_{n,n}^{-12k^2} \right) \right| \\ &\lesssim \frac{12k^2 n C (4k + N_E) (40k^2 \delta)^{6kn}}{d^n} (-\log \varepsilon) \lesssim n \left(\frac{(40k^2 \delta)^{6k}}{d} \right)^n (-\log \varepsilon) \lesssim n d^{n/4} (-\log \varepsilon) \end{aligned}$$

for all sufficiently small $\varepsilon > 0$. □

9. GREEN CURRENTS AND (ALMOST) SIMPLE ACTION

On a general compact Kähler manifold, the convergence of the sequence $d_p^{-n}(f^n)^* \omega^p$ is not clear. For the existence and construction of the Green current, see [17], [22], [9], [15]. In this section, we summarize some related results for our purpose.

We first summarize some results from [15]. Let X and f be as in Theorem 1.1. We say that the action of f on cohomology is simple if there is an integer $0 \leq p \leq k$ such that

- (1) the dynamical degree d_p of order p of f is strictly larger than the other dynamical degrees,
- (2) d_p is a simple eigenvalue of f^* on $H^{p,p}(X, \mathbb{R})$, and
- (3) the other (real or complex) eigenvalues of this operator have modulus strictly smaller than d_p .

Under this condition, we have the existence of the Green current T^+ .

Proposition 9.1 ([15, Proposition 5.5]). The sequence $d_p^{-n}(f^n)^*(\omega^p)$ converges weakly to a positive closed (p, p) -current T^+ as n tends to infinity. The sequence $d_p^{-n}(f^n)_*(\omega^{k-p})$ converges weakly to a positive closed $(k-p, k-p)$ -current T^- as n tends to infinity. Moreover, T^+ and T^- have continuous superpotentials.

Lemma 4.1 in [5] implies the following convergence.

Corollary 9.2. We have $d_p^{-n}(f^n)^*(\alpha_S) \rightarrow c_{\alpha_S} T^+$ exponentially fast where $c_{\alpha_S} = \frac{\langle \alpha_S, T^- \rangle}{\langle \omega^p, T^- \rangle}$.

Together with Theorem 1.1, we get

Theorem 9.3. *Let X and f be as in Theorem 1.1. Assume further that the action of f is simple. Then, there exists a proper analytic subset E invariant under f such that if $S \in \mathcal{C}_p$ is smooth near E , then*

$$d_p^{-n}(f^n)^* S \rightarrow c_S T^+$$

exponentially fast where $c_S = \frac{\langle S, T^- \rangle}{\langle \omega^p, T^- \rangle}$. In particular, if H is an analytic subset of pure dimension $k - p$ such that $H \cap E = \emptyset$, then the sequence $d_p^{-n}(f^n)^*[H]$ converges to $c_H T^+$ where $c_H = \frac{\langle [H], T^- \rangle}{\langle \omega^p, T^- \rangle}$.

If we do not require the existence of the limit $d_p^{-n}(f^n)_*(\omega^{k-p})$, the convergence of $d_p^{-n}(f^n)^*(\omega^p)$ is true under slightly more relaxed conditions.

- (1) $d_{p-1} < d_p$,
- (2) d_p is a simple eigenvalue of f^* on $H^{p,p}(X, \mathbb{R})$, and
- (3) the other (real or complex) eigenvalues of this operator on $H^{p,p}(X, \mathbb{R})$ have modulus strictly smaller than d_p .

The proofs of lemmas and propositions in [15] work in the same way. So, we have Theorem 1.4.

REFERENCES

- [1] AHN, T.: Equidistribution in higher codimension for holomorphic endomorphisms of \mathbb{P}^k , Trans. Am. Math. Soc. 368 (2016) 3359 – 3388.
- [2] AHN, T.: Local regularity of super-potentials and equidistribution of positive closed currents on \mathbb{P}^k , Math. Ann. 371 (2018), no. 3–4, 1163–1190.
- [3] AHN, T.: An equidistribution theorem for certain birational maps of \mathbb{P}^k , Internat. J. Math. 32 (2021), no. 3, Paper No. 2150017, 19 pp.
- [4] AHN, T.; NGUYEN, N. C.: Equidistribution of non-pluripolar products associated with quasi-plurisubharmonic functions of finite energy, Proc. Amer. Math. Soc. 148 (2020), no. 2, 719–729.
- [5] AHN, T.; VU, D.-V.: Equidistribution for non-pluripolar currents on compact Kähler manifolds, arXiv:2309.12099, preprint
- [6] BLANCHARD, A.: Sur les variétés analytiques complexes, Ann. Sci. Ecole Norm. Sup. (3) 73 (1956) 157–202.
- [7] BOST, J. -B.; GILLET, H.; SOULÉ, C.: Heights of projective varieties and positive Green forms, J. Amer. Math. Soc. 7 (1994) no. 4, 903–1027.
- [8] BRIEND, J. -Y.; DUVAL, J.: Deux caractérisations de la mesure d'équilibre d'un endomorphisme de $\mathbb{P}^k(C)$ (French, with English and French summaries), Publ. Math. Inst. Hautes Études Sci. 93 (2001), 145–159
- [9] DE THÉLIN, H.; DINH, T.-C.: Dynamics of automorphisms on compact Kähler manifolds, Adv. Math. 229 (2012), no. 5, 2640–2655.
- [10] DEMAILLY, J.-P.: Complex Analytic and Differential Geometry.
- [11] DINH, T.-C.: Suites d'applications méromorphes multivaluées et courants laminaires, J. Geom. Anal. 15 (2005) 207 – 227.
- [12] DINH, T.-C.: Analytic multiplicative cocycles over holomorphic dynamical systems, Complex Var. Elliptic Equ. 54 (2009) no. 3-4, 243–251.
- [13] DINH, T.-C.; NGUYÊN, V.-A.: The mixed Hodge-Riemann bilinear relations for compact Kähler manifolds, Geom. Fuct. Anal. 16 (2006) 838–849.
- [14] DINH, T.-C.; NGUYÊN, V.-A.; Truong, T. T.: Equidistribution for meromorphic maps with dominant topological degree, Indiana Univ. Math. J. 64 (2015) no.6, 1805–1828.
- [15] DINH, T.-C.; NGUYÊN, V.-A.; VU, D.-V.: Super-potentials, densities of currents and number of periodic points for holomorphic maps
- [16] DINH, T.-C.; SIBONY, N.: Regularization of currents and entropy, Ann. Sci. Éc. Norm. Supér. 37 (2004), 959–971.
- [17] DINH, T.-C.; SIBONY, N.: Green currents for holomorphic automorphisms of compact Kähler manifolds, J. Amer. Math. Soc., 18 (2005), no.2, 291–312.
- [18] DINH, T.-C.; SIBONY, N.: Pull-back of currents by holomorphic maps, Manuscripta Math. 123 (2007), no. 3, 357–371.
- [19] DINH, T.-C.; SIBONY, N.: Equidistribution towards the Green current for holomorphic maps, Ann. Sci. Éc. Norm. Supér. (4) 41 (2008), no. 2, 307–336.

- [20] DINH, T.-C.; SIBONY, N.: Super-potentials of positive closed currents, intersection theory and dynamics, *Acta Math.*, 203 (2009), 1–82.
- [21] DINH, T.-C.; SIBONY, N.: Equidistribution speed for endomorphisms of projective spaces, *Math. Ann.* 347 (2010), no.3, 613–626.
- [22] DINH, T.-C.; SIBONY, N.: Super-potentials for currents on compact Kähler manifolds and dynamics of automorphisms, *J. Algebraic Geom.* 19 (2010), no. 3, 473–529.
- [23] FAVRE, C.: Note on pull-back and Lelong number of currents (English, with English and French summaries), *Bull. Soc. Math. France* 127 (1999), no.3, 445–458.
- [24] FAVRE, C.: Dynamique des applications rationnelles, PhD thesis, Université de Paris-Sud, Orsay, 2000.
- [25] FAVRE, C.: Multiplicity of holomorphic functions, *Math. Ann.* 316 (2000), no. 2, 355–378.
- [26] FAVRE, C.; JONSSON, M.: Brolin’s theorem for curves in two complex dimensions (English, with English and French summaries), *Ann. Inst. Fourier (Grenoble)* 53 (2003), no.5, 1461–1501.
- [27] FAVRE, C.; JONSSON, M.: Eigenvaluations (English, with English and French summaries), *Ann. Sci. École Norm. Sup. (4)* 40 (2007), no. 2, 309–349
- [28] FORNÆSS, J. E.; SIBONY, N.: Complex dynamics in higher dimensions, *Complex potential theory (Montreal, PQ, 1993)*, NATO Adv. Sci. Inst. Ser. CMath. Phys. Sci., vol. 439, Kluwer Acad. Publ., Dordrecht, 1994, pp. 131–186. Notes partially written by Estela A. Gavosto.
- [29] FORNÆSS, J. E.; SIBONY, N.: Complex dynamics in higher dimension II. In *Modern methods in complex analysis (Princeton, NJ, 1992)*, volume 137 of *Ann. Math. Stud.*, pages 135–182. Princeton University Press, 1995.
- [30] FREIRE, A.; LOPES, A.; MÁÑÉ, M.: An invariant measure for rational maps, *Bol. Soc. Brasil. Mat.* 14 (1983), no. 1, 45–62.
- [31] GIGNAC, William: Measures and dynamics on Noetherian spaces, *J. Geom. Anal.* 24 (2014), no. 4, 1770–179.
- [32] GROMOV, M.: *Convex sets and Kähler manifolds*, World Sci. Publishing, Teaneck, NJ, 1990.
- [33] GUEDJ, V.: Equidistribution towards the Green current (English, with English and French summaries), *Bull. Soc. Math. France* 131 (2003), no. 3, 359–372.
- [34] LJUBICH, M.: Entropy properties of rational endomorphisms of the Riemann sphere, *Ergodic Theory Dynam. Systems* 3 (1983), no. 3, 351–385
- [35] PARRA, Rodrigo: The Jacobian cocycle and equidistribution towards the Green current, arXiv:1103.4633.
- [36] RUSSAKOVSKII, A.; SHIFFMAN, B.: Value distribution for sequences of rational mappings and complex dynamics, *Indiana Univ. Math. J.* 46 (1997), no. 3, 897–932
- [37] SIBONY, N.: Dynamique des applications rationnelles de P^k (French, with English and French summaries), *Dynamique et géométrie complexes (Lyon, 1997)*, Panor. Synthèses, vol. 8, Soc. Math. France, Paris, 1999, pp. ix–x, xi–xii, 97–185.
- [38] TAFLIN, J.: Equidistribution speed towards the Green current for endomorphisms of P^k , *Adv. Math.* 227 (2011), no. 5, 2059–2081

(AHN) DEPARTMENT OF MATHEMATICS EDUCATION, INHA UNIVERSITY, 100 INHA-RO, MICHUHOL-GU, INCHEON 22212, REPUBLIC OF KOREA
Email address: t.ahn@inha.ac.kr