

Higher-order Asymptotic Profiles of Solutions to the Cauchy Problem for the Convection-Diffusion Equation with Variable Diffusion

Ikki Fukuda and Shinya Sato

Dedicated to Professor Takayoshi Ogawa on the occasion of his 60th birthday

Abstract

We consider the asymptotic behavior of solutions to the convection-diffusion equation:

$$\partial_t u - \operatorname{div}(a(x)\nabla u) = d \cdot \nabla (|u|^{q-1} u), \quad x \in \mathbb{R}^n, \quad t > 0$$

with an integrable initial data $u_0(x)$, where $n \geq 1$, $q > 1 + \frac{1}{n}$ and $d \in \mathbb{R}^n$. Moreover, we take $a(x) = 1 + b(x) > 0$, where $b(x)$ is smooth and decays fast enough at spatial infinity. It is known that the asymptotic profile of the solution to this problem can be given by the heat kernel. Moreover, the second asymptotic profile of the solution have already been studied. In particular, the following three cases are distinguished: $1 + \frac{1}{n} < q < 1 + \frac{2}{n}$; $q = 1 + \frac{2}{n}$; $q > 1 + \frac{2}{n}$. More precisely, the second asymptotic profile has different properties in each of these three cases. In this paper, we focus on the critical case of $q = 1 + \frac{2}{n}$. By analyzing the corresponding integral equation in details, we have succeeded to give the more higher-order asymptotic expansion of the solution, which generalizes the previous works.

Keywords: Convection-diffusion equation; Variable diffusion; Higher-order asymptotic profiles.

1 Introduction

We consider the Cauchy problem for the convection-diffusion equation of the following form:

$$\begin{aligned} \partial_t u - \operatorname{div}(a(x)\nabla u) &= d \cdot \nabla (|u|^{q-1} u), \quad x \in \mathbb{R}^n, \quad t > 0, \\ u(x, 0) &= u_0(x), \quad x \in \mathbb{R}^n, \end{aligned} \tag{1.1}$$

where $u = u(x, t)$ is a real-valued unknown function, $u_0(x)$ is a given initial data, $n \geq 1$, $q > 1$ and $d \in \mathbb{R}^n$. For the variable diffusion coefficient $a(x)$, we take asymptotically constant diffusion of the form $a(x) = 1 + b(x) > 0$ with $b(x)$ satisfying $b \in L^1(\mathbb{R}^n) \cap C^{1,\alpha}(\mathbb{R}^n)$ ($0 < \alpha \leq 1$) and $\|b\|_{L^\infty} < 1$ such that

$$|b(x)| + (1 + |x|^2)^{\frac{1}{2}} |\nabla b(x)| \leq C (1 + |x|^2)^{-\frac{\delta}{2}}, \quad x \in \mathbb{R}^n \tag{1.2}$$

for some positive constant $C > 0$ and $\delta > 0$. This kind of equation arises from the study in the context of the diffusion of pollution in fluids (see, e.g. [2]). In this paper, we would like to analyze the asymptotic behavior of the solutions to (1.1). In particular, we are interested in finding out how the variable diffusion and the nonlinearity affect the solution at $t \rightarrow \infty$.

First of all, let us recall some known results related to the Cauchy problem (1.1). In [5], Escobedo–Zuazua studied (1.1) when $a(x) \equiv 1$ (i.e. $b(x) \equiv 0$). They showed that if $q > 1$ and $u_0 \in L^1(\mathbb{R}^n)$, then there exists a unique solution $u \in C([0, \infty); L^1(\mathbb{R}^n))$ satisfying

$$u \in C((0, \infty); W^{2,p}(\mathbb{R}^n)) \cap C^1((0, \infty); L^p(\mathbb{R}^n)).$$

for any $p \in (1, \infty)$. Moreover, if we additionally assume $u_0 \in L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$, then the solution $u(x, t)$ also satisfies the following property:

$$u \in C([0, \infty); L^1(\mathbb{R}^n)) \cap C((0, \infty); L^\infty(\mathbb{R}^n)) \cap L^\infty(\mathbb{R}^n \times [0, \infty)). \tag{1.3}$$

2020 Mathematics Subject Classification: 35B40, 35K15, 35K55.

We note that the above results are also summarized very well, in the lecture note written by Zuazua [11]. Next, let us introduce the additional properties of the solution $u(x, t)$ to (1.1). Now, integrating the equation (1.1) over \mathbb{R}^n , we deduce that the mass is conserved,

$$\text{i.e. } \int_{\mathbb{R}^n} u(x, t) dx = \int_{\mathbb{R}^n} u_0(x) dx, \quad t > 0.$$

On the other hand, the following L^p -decay estimates hold for any $p \in [1, \infty]$ (cf. [5, 11]):

$$\|u(t)\|_{L^p} \leq C t^{-\frac{n}{2}\left(1-\frac{1}{p}\right)}, \quad t > 0, \quad (1.4)$$

$$\|\nabla u(t)\|_{L^p} \leq C t^{-\frac{n}{2}\left(1-\frac{1}{p}\right)-\frac{1}{2}}, \quad t > 1. \quad (1.5)$$

Furthermore, Duro–Zuazua [4] studied the more general cases of (1.1) with the variable diffusion satisfying (1.2) and generalized the results given in [5]. Actually, the all properties mentioned above also can be established for our target problem (1.1). We note that the constants appeared in (1.4) and (1.5) do not depend on $b(x)$, by virtue of (1.2). Also, we emphasize that these decay rates are the same as for the solution to the corresponding linear problem.

In addition to the decay estimates (1.4) and (1.5), the more detailed asymptotic behavior of the solution has been studied in [4, 5]. The decay rates given in (1.4) and (1.5) do not depend on the nonlinear exponent q . However, the asymptotic profile of the solution deeply depends on the nonlinearity. Actually, when $a(x) \equiv 1$, the existence of a critical value of the exponent q . Roughly speaking, the following results have been shown:

- If $q > 1 + \frac{1}{n}$, the solution behaves like a self-similar solution of the linear heat equation.
- If $q = 1 + \frac{1}{n}$, the solution behaves like a self-similar solution of the full equation.
- If $1 < q < 1 + \frac{1}{n}$, the solution behaves like a self-similar solution of a reduced equation.

For the more precise statements, see [5]. In the present paper, we focus on the super critical case $q > 1 + \frac{1}{n}$. Under this situation, the asymptotic profile of the solution can be given by the heat kernel, even if the more variable diffusion case (cf. [4]). That is, neither the variable diffusion, nor the nonlinearity appear in the first approximation. To state such a result, we set

$$G(x, t) := \frac{1}{(4\pi t)^{\frac{n}{2}}} e^{-\frac{|x|^2}{4t}}, \quad M := \int_{\mathbb{R}^n} u_0(x) dx. \quad (1.6)$$

Moreover, for $p \in [1, \infty)$, let us introduce the following weighted Lebesgue spaces:

$$L^p(\mathbb{R}^n; 1 + |x|) := \left\{ f \in L^p(\mathbb{R}^n); \int_{\mathbb{R}^n} \{|f(x)|(1 + |x|)\}^p dx < \infty \right\}.$$

Then, the following asymptotic formula has been established in [4]:

Proposition 1.1 ([4]). *Let $n \geq 1$ and $q > 1 + \frac{1}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$ and $t \geq 1$, the solution $u(x, t)$ to (1.1) satisfies the following asymptotic formula:*

$$\|u(t) - MG(t)\|_{L^p} \leq C \begin{cases} t^{-\frac{n}{2}\left(q-\frac{1}{p}\right)+\frac{1}{2}}, & 1 + \frac{1}{n} < q < 1 + \frac{2}{n}, \\ t^{-\frac{n}{2}\left(1-\frac{1}{p}\right)-\frac{1}{2}} \log(2 + t), & q = 1 + \frac{2}{n}, \\ t^{-\frac{n}{2}\left(1-\frac{1}{p}\right)-\frac{1}{2}}, & q > 1 + \frac{2}{n}, \end{cases} \quad (1.7)$$

where $G(x, t)$ and M are defined by (1.6).

As we can see from the above result, the asymptotic rates given in (1.7) depend on the exponent q . In addition, the following three cases are distinguished:

$$1 + \frac{1}{n} < q < 1 + \frac{2}{n}; \quad q = 1 + \frac{2}{n}; \quad q > 1 + \frac{2}{n}.$$

Roughly speaking, we note that the above critical exponent $q = 1 + \frac{2}{n}$ appears due to the integrability of the following integral:

$$\int_1^t \int_{\mathbb{R}^n} |u(x, \tau)|^q dx d\tau = \int_1^t \|u^q(\cdot, \tau)\|_{L^1} d\tau \leq C \int_1^t \tau^{-\frac{n}{2}(q-1)} d\tau, \quad (1.8)$$

where we used the decay estimate (1.4). Based on the above result, we can expect that the nonlinearity strongly affects on the second asymptotic profile of the solution. Actually, such a result has already obtained in [3, 12]. First, Zuazua [12] succeeded to derive the three kinds of second asymptotic profiles of the solution $u(x, t)$ depending on q , when $a(x) \equiv 1$. After that, these results were generalized by Duro-Carpio [3], including the variable diffusion cases. We will see here that in view of the second asymptotic profiles, the asymptotic rates to $MG(x, t)$ given in (1.7) are optimal, with respect to the time decaying order. On the other hand, the importance of the perturbation $b(x)$ is greater in lower dimensions, particularly in one-dimension $n = 1$. Indeed in [3], they proved that the variable diffusion appears in the second asymptotic profiles only when $q > 1 + \frac{2}{n}$ or when $n = 1$ and $q = 3$. More precisely, the following three propositions have been shown in [3]:

Proposition 1.2 ([3]). *Let $n \geq 1$ and $1 + \frac{1}{n} < q < 1 + \frac{2}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^{n(q-1)}(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following asymptotic formula:*

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}(q-\frac{1}{p})-\frac{1}{2}} \left\| u(t) - MG(t) - |M|^{q-1} MZ(t) \right\|_{L^p} = 0, \quad (1.9)$$

where $G(x, t)$ and M are defined by (1.6), while $Z(x, t)$ is the solution to the following problem:

$$\begin{aligned} \partial_t Z - \Delta Z &= d \cdot \nabla (G^q), \quad x \in \mathbb{R}^n, \quad t > 0, \\ Z(x, 0) &= 0, \quad x \in \mathbb{R}^n. \end{aligned}$$

Proposition 1.3 ([3]). *Let $q = 1 + \frac{2}{n}$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following asymptotic formulas:*

(i) *If $n \geq 2$ and $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^q(\mathbb{R}^n; 1 + |x|) \cap L^2(\mathbb{R}^n)$, then*

$$\lim_{t \rightarrow \infty} \frac{t^{\frac{n}{2}(1-\frac{1}{p})+\frac{1}{2}}}{\log t} \left\| u(t) - MG(t) - \alpha_n |M|^{\frac{2}{n}} M(\log t) d \cdot \nabla G(t) \right\|_{L^p} = 0. \quad (1.10)$$

(ii) *If $n = 1$ and $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^3(\mathbb{R}^n; 1 + |x|)$, then*

$$\lim_{t \rightarrow \infty} \frac{t^{\frac{1}{2}(1-\frac{1}{p})+\frac{1}{2}}}{\log t} \left\| u(t) - MG(t) - (K(t) + \alpha_1 d M^3) (\log t) \partial_x G(t) \right\|_{L^p} = 0, \quad (1.11)$$

where $G(x, t)$ and M are defined by (1.6), while α_n and $K(t)$ are defined by

$$\alpha_n := \frac{1}{4\pi} \left(1 + \frac{2}{n}\right)^{-\frac{n}{2}}, \quad K(t) := \frac{1}{\log t} \int_0^t \int_{\mathbb{R}} b(y) \partial_x u(y, \tau) dy d\tau. \quad (1.12)$$

Remark 1.4. In the above proposition, we must distinguish the cases $n \geq 2$ and $n = 1$, due to the integrability for the integral $\int_1^\infty \int_{\mathbb{R}^n} |b(y)\nabla u(y, \tau)| dy d\tau$. For details, see Remark 3.8 below. However, in the case of $n = 1$, if we further assume that the initial data can be expressed as $u_0(x) = \partial_x v_0(x)$ with some $v_0 \in L^1(\mathbb{R})$, then we have the following fact (for the proof, see [3]):

$$\lim_{t \rightarrow \infty} K(t) = \lim_{t \rightarrow \infty} \left\{ \frac{1}{\log t} \int_0^t \int_{\mathbb{R}} b(y) \partial_x u(y, \tau) dy d\tau \right\} = 0.$$

Therefore, under this situation, we can see that (1.10) holds even if the case of $n = 1$.

Proposition 1.5 ([3]). Let $q > 1 + \frac{2}{n}$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following results:

(i) If $n \geq 2$ and $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^{n(q-1)}(\mathbb{R}^n)$, then

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}(1-\frac{1}{p})+\frac{1}{2}} \|u(t) - MG(t) - \beta \cdot \nabla G(t)\|_{L^p} = 0, \quad (1.13)$$

where $G(x, t)$ and M are defined by (1.6), while β is defined by

$$\beta := d \int_0^\infty \int_{\mathbb{R}^n} (|u|^{q-1} u)(y, \tau) dy d\tau + \int_0^\infty \int_{\mathbb{R}^n} b(y) \nabla u(y, \tau) dy d\tau - m, \quad (1.14)$$

$$m := (m_1, m_2, \dots, m_n), \quad m_i := \int_{\mathbb{R}^n} x_i u_0(x) dx. \quad (1.15)$$

(ii) If $n = 1$, $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^q(\mathbb{R}^n; 1 + |x|)$ and the initial data can be expressed as $u_0(x) = \partial_x v_0(x)$ for some $v_0 \in L^1(\mathbb{R})$, then (1.13) holds with β as in (1.14).

Remark 1.6. We note that the term $\int_0^\infty \int_{\mathbb{R}^n} b(y) \nabla u(y, \tau) dy d\tau = - \int_0^\infty \int_{\mathbb{R}^n} \nabla b(y) u(y, \tau) dy d\tau$ does not appear if $b(x)$ is an identically constant. This fact agrees with the result given in [12].

Comparing the asymptotic rates $o(t^{-(n/2)(q-1/p)+1/2})$ in (1.9), $o(t^{-(n/2)(1-1/p)-1/2} \log t)$ in (1.10) and (1.11), and $o(t^{-(n/2)(1-1/p)-1/2})$ in (1.13) obtained from the three propositions above, we can find that $o(t^{-(n/2)(1-1/p)-1/2})$ in (1.13) gives the fastest asymptotic rate. Note that this asymptotic rate is the same as the asymptotic rate that appears in the second order asymptotic expansion for the solution to the linear heat equation (see, (3.3) below). Compared with this result and the other two cases, Propositions 1.2 and 1.3 appear to be less accurate approximations to the solution. Moreover, it should also be noted that a very interesting phenomenon occurs in the critical case of $q = 1 + \frac{2}{n}$. Actually, the logarithmic term $\log t$ appears not only in the asymptotic rate $o(t^{-(n/2)(1-1/p)-1/2} \log t)$ but also in the shape of the second asymptotic profiles:

$$\alpha_n |M|^{\frac{2}{n}} M(\log t) d \cdot \nabla G(x, t) \quad \text{in (1.10),} \quad (K(t) + \alpha_1 d M^3) (\log t) \partial_x G(x, t) \quad \text{in (1.11).} \quad (1.16)$$

Also, we remark that the logarithmic term $\log t$ came from both of the effects of the nonlinearity and the variable diffusion. We note that similar phenomenon were also observed for other nonlinear parabolic equations. For example, let us introduce the following parabolic system of chemotaxis of the form

$$\begin{aligned} u_t &= \Delta u - \nabla \cdot (u \nabla v), \quad x \in \mathbb{R}^n, \quad t > 0, \\ v_t &= \Delta v - v + u, \quad x \in \mathbb{R}^n, \quad t > 0. \end{aligned}$$

Indeed, in Nagai–Yamada [10], the related result to (1.10) and (1.11) has been derived for the above system, in the case of $n = 1$ (we can also refer to [9]).

On the other hand, Ishige–Kawakami [8] succeeded to derive the different asymptotic formula for any $q > 1 + \frac{1}{n}$, which has the same asymptotic rate $o(t^{-(n/2)(1-1/p)-1/2})$. Actually, the following proposition has been established:

Proposition 1.7 ([8]). *Let $n \geq 1$ and $q > 1 + \frac{1}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$ and $a(x) \equiv 1$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following asymptotic formula:*

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}(1-\frac{1}{p})+\frac{1}{2}} \|u(t) - \hat{u}(t)\|_{L^p} = 0, \quad (1.17)$$

where $\hat{u}(x, t)$ is defined by

$$\begin{aligned} \hat{u}(x, t) &:= MG(x, 1+t) - c(t) \cdot \nabla G(x, 1+t) \\ &\quad + |M|^{q-1} M \int_0^t (a \cdot \nabla G(t-\tau) * G(1+\tau)^q)(x) d\tau, \\ c_i(t) &:= \int_{\mathbb{R}^n} x_i u(x, t) dx - |M|^{q-1} M \int_0^t \int_{\mathbb{R}^n} x_i \{a \cdot \nabla (G(x, 1+\tau)^q)\} dx d\tau, \\ c(t) &:= (c_1(t), c_2(t), \dots, c_n(t)), \end{aligned} \quad (1.18)$$

while $G(x, t)$ and M are defined by (1.6).

Here, we remark that actually in [8], not only the convection-diffusion equation (1.1) but also the asymptotic behavior of solutions to the more general parabolic partial differential equations of the form

$$\begin{aligned} \partial_t u - \Delta u &= F(x, t, u, \nabla u), \quad x \in \mathbb{R}^n, \quad t > 0, \\ u(x, 0) &= u_0(x), \quad x \in \mathbb{R}^n \end{aligned} \quad (1.19)$$

with some nonlinear function $F \in C(\mathbb{R}^n \times (0, \infty) \times \mathbb{R} \times \mathbb{R}^n)$ were studied. The above Proposition 1.7 is a direct consequence of its general theory. For details, see the original paper [8].

By virtue of Proposition 1.7, an asymptotic expansion of the solution has been accomplished up to the order $o(t^{-(n/2)(1-1/p)-1/2})$. However, we note that these asymptotic profiles obtained in Proposition 1.7 and Propositions 1.2, 1.3, 1.5 are different each other. The above formula (1.17) has the advantage of being able to express the asymptotic behavior of the solution to (1.1) in a unified manner for any $q > 1 + \frac{1}{n}$. However, compared with the results obtained in [3, 12], the effects of the nonlinearity and the variable diffusion on the asymptotic profiles are a little bit difficult to see, especially in the critical case $q = 1 + \frac{2}{n}$ (logarithmic term does not appear in (1.17)). On the other hand, in [3, 12], the second asymptotic profiles are decomposed and expressed in detail depending on the nonlinear exponent q , but the given asymptotic rate is slower than $o(t^{-(n/2)(1-1/p)-1/2})$ by $o(\log t)$. Taking into account this situation, we believe that it might be effective to derive a new asymptotic formula which is better in terms of both the shape of the asymptotic profile and the asymptotic rate.

In this study, we focus on the critical case $q = 1 + \frac{2}{n}$ and study the asymptotic behavior of the solution to (1.1), based on the above perspectives. Especially, we investigate whether the asymptotic rate $o(t^{-(n/2)(1-1/p)-1/2})$ can be realized when the second asymptotic profiles are fixed to the form in (1.16). Namely, we consider that whether the asymptotic rate $o(t^{-(n/2)(1-1/p)-1/2} \log t)$ given in (1.10) and (1.11) is optimal or not. As a result, by analyzing the corresponding integral equation (3.1) below in details, we have succeeded to give the more higher-order asymptotic expansion of the solution, which generalizes Proposition 1.3 given by Duro-Carpio [3].

The rest of this paper is organized as follows. In Section 2, we shall state our main results Theorems 2.1 and 2.5 below. Next, we would like to give the proofs of these results in Section 3. This section is divided into two parts. Subsection 3.1 is devoted to prove Theorem 2.1. In this part, we mainly study the effect of the nonlinearity. On the other hand, Theorem 2.5 will be proven in Subsection 3.2. The effect of the variable diffusion is investigated in this part. The main novelty of this paper is to derive the higher-order asymptotic profiles which take into account the effects of both the nonlinearity and the variable diffusion. The key estimates are in Propositions 3.4 and 3.11 below, and the main techniques of the proofs are based on the idea used in [6] for the asymptotic analysis of a dispersive-dissipative equation.

2 Main Results

In this section, we would like to state our main results. First, let us mention about the result in the case of $n \geq 2$. In order to doing that, we shall introduce the following new function $\Psi(x, t)$:

$$\Psi(x, t) := t^{-\frac{n}{2}-\frac{1}{2}}\Psi_*\left(\frac{x}{\sqrt{t}}\right), \quad \Psi_*(x) := d \cdot \nabla \left(\int_0^1 (G(1-s) * F(s))(x) ds \right), \quad (2.1)$$

$$F(y, s) := s^{-\frac{n}{2}-1}F_*\left(\frac{y}{\sqrt{s}}\right), \quad F_*(y) := \frac{1}{2^{n+2}\pi^{\frac{n}{2}+1}} \left\{ e^{-\frac{|y|^2}{4}\left(1+\frac{2}{n}\right)} - \left(1+\frac{2}{n}\right)^{-\frac{n}{2}} e^{-\frac{|y|^2}{4}} \right\}. \quad (2.2)$$

In addition, we shall define the constants \mathcal{M} and \mathcal{N} by

$$\mathcal{M} := d \int_0^1 \int_{\mathbb{R}^n} \left(|u|^{\frac{2}{n}} u \right)(y, \tau) dy d\tau + d \int_1^\infty \int_{\mathbb{R}^n} \left(|u|^{\frac{2}{n}} u - |MG|^{\frac{2}{n}} MG \right)(y, \tau) dy d\tau. \quad (2.3)$$

$$\mathcal{N} := \int_0^\infty \int_{\mathbb{R}^n} b(y) \nabla u(y, \tau) dy d\tau, \quad n \geq 2. \quad (2.4)$$

Then, we are able to obtain the following higher-order asymptotic expansion of the solution:

Theorem 2.1. *Let $n \geq 2$ and $q = 1 + \frac{2}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following asymptotic formula:*

$$\begin{aligned} \lim_{t \rightarrow \infty} t^{\frac{n}{2}\left(1-\frac{1}{p}\right)+\frac{1}{2}} \left\| u(t) - MG(t) - \alpha_n |M|^{\frac{2}{n}} M(\log t) d \cdot \nabla G(t) \right. \\ \left. - (\mathcal{M} + \mathcal{N} - m) \cdot \nabla G(t) - |M|^{\frac{2}{n}} M \Psi(t) \right\|_{L^p} = 0, \end{aligned} \quad (2.5)$$

where $G(x, t)$ and M, m and α_n are defined by (1.6), (1.15) and (1.12), respectively. In addition, $\Psi(x, t)$, \mathcal{M} and \mathcal{N} are defined by (2.1), (2.3) and (2.4), respectively.

Remark 2.2. *In view of the higher-order asymptotic expansion, we can obtain the optimal asymptotic rate to the second asymptotic profile. More precisely, under the same assumptions in Theorem 2.1, the solution $u(x, t)$ to (1.1) satisfies the following estimate:*

$$\left\| u(t) - MG(t) - \alpha_n |M|^{\frac{2}{n}} M(\log t) d \cdot \nabla G(t) \right\|_{L^p} = (C_* + o(1)) t^{-\frac{n}{2}\left(1-\frac{1}{p}\right)-\frac{1}{2}} \quad (2.6)$$

as $t \rightarrow \infty$, for any $p \in [1, \infty]$, where the constant C_* is defined by

$$C_* := \left\| (\mathcal{M} + \mathcal{N} - m) \cdot \nabla G(1) + |M|^{\frac{2}{n}} M \Psi_* \right\|_{L^p}. \quad (2.7)$$

From this result, for the asymptotic rate to the second asymptotic profile, we can conclude that the optimal asymptotic rate is $O(t^{-(n/2)(1-1/p)-1/2})$ if and only if $C_* \neq 0$. This fact implies that the asymptotic rate given in (1.10) can be improved, i.e. our result (2.6) generalizes the result (1.10) given in [3].

Remark 2.3. *Compared with Proposition 1.7 given by Ishige–Kawakami [8], our formula is a different from their result with respect to a structure of the asymptotic profile. More precisely, we emphasize that our new obtained asymptotic profile $\Psi(x, t)$ has the following parabolic self-similarity:*

$$\Psi(x, t) = \lambda^{n+1} \Psi(\lambda x, \lambda^2 t), \quad \text{for } \lambda > 0.$$

Remark 2.4. *Similarly as Remark 1.6, if $b(x)$ is an identically constant, then the constant \mathcal{N} defined by (2.4) satisfies $\mathcal{N} = 0$. Therefore, in this case, the above results (2.5) and (2.6) with $\mathcal{N} = 0$ are also valid for the case of $n = 1$.*

Next, we would like to introduce the result in the case of $n = 1$ and $q = 3$. As we have already seen in (1.11), for deriving the asymptotic profile in this case, we need to introduce the function $K(t)$ defined by (1.12). In order to lead the higher-order asymptotic expansion, we shall slightly modify this function. Now, let us define the following new functions $\mathcal{K}(t)$ and $\Phi(x, t)$, and the constant \mathcal{L} :

$$\mathcal{K}(t) := \frac{1}{\log t} \int_1^t \int_{\mathbb{R}} b(y) \partial_x u(y, \tau) dy, \quad (2.8)$$

$$\Phi(x, t) := t^{-1} \Phi_* \left(\frac{x}{\sqrt{t}}, t \right), \quad \Phi_*(x, t) := t \partial_x \left(\int_0^1 \left(G(1-s) * \phi(\sqrt{t} \cdot, ts) G(s) \right) (x) ds \right), \quad (2.9)$$

$$\phi(y, \tau) := b(y) \partial_x u(y, \tau) - \left(\int_{\mathbb{R}} b(z) \partial_x u(z, \tau) dz \right), \quad \mathcal{L} := \int_0^1 \int_{\mathbb{R}} b(y) \partial_x u(y, \tau) dy d\tau. \quad (2.10)$$

Then, we can state the following higher-order asymptotic expansion of the solution for $n = 1$:

Theorem 2.5. *Let $n = 1$ and $q = 3$. Suppose $u_0 \in L^1(\mathbb{R}; 1 + |x|) \cap L^\infty(\mathbb{R})$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following asymptotic formula:*

$$\begin{aligned} \lim_{t \rightarrow \infty} t^{\frac{1}{2} \left(1 - \frac{1}{p}\right) + \frac{1}{2}} \left\| u(t) - MG(t) - (\alpha_1 dM^3 + \mathcal{K}(t)) (\log t) \partial_x G(t) \right. \\ \left. - (\mathcal{M} + \mathcal{L} - m) \partial_x G(t) - M^3 \Psi(t) - \Phi(t) \right\|_{L^p} = 0, \end{aligned} \quad (2.11)$$

where $G(x, t)$ and M, m and α_1 are defined by (1.6), (1.15) and (1.12), respectively. In addition, $\Psi(x, t)$ and \mathcal{M} are defined by (2.1) and (2.3), respectively. Moreover, $\Phi(x, t)$, $\mathcal{K}(t)$ and \mathcal{L} are defined by (2.9), (2.8) and (2.10), respectively.

Remark 2.6. *Since the functions $\mathcal{K}(t)$ and $\Phi_*(x, t)$ are depend on the original solution $u(x, t)$, $\mathcal{K}(t)(\log t) \partial_x G(x, t)$ and $\Phi(x, t)$ in (2.11) perhaps should be called the ‘‘correction terms’’ rather than the asymptotic profiles. However, we can find their good properties at $t \rightarrow \infty$. Actually, for any $p \in [1, \infty]$, the following uniform boundedness hold:*

$$|\mathcal{K}(t)| \leq C, \quad \|\Phi_*(\cdot, t)\|_{L^p} \leq C, \quad t > 1.$$

For the proof, see (3.35) and (3.44) below. Therefore, taking into account this facts, we can conclude that $\mathcal{K}(t)(\log t) \partial_x G(x, t)$ and $\Phi(x, t)$ decay at least on the order $O(t^{-(n/2)(1-1/p)-1/2} \log t)$ and $O(t^{-(n/2)(1-1/p)-1/2})$, respectively.

Remark 2.7. *Similarly as Remarks 1.6 and 2.4, if $b(x)$ is an identically constant, then all of $\mathcal{K}(t)$, $\Phi(x, t)$ and \mathcal{L} always vanish. Therefore, in this case, the corresponding result of (2.6) is also valid for the case of $n = 1$.*

3 Proof of the Main Results

In this section, we would like to prove our main results Theorems 2.1 and 2.5, i.e. we shall derive the asymptotic formulas (2.5) and (2.11). In what follows, let us consider the case of $q = 1 + \frac{2}{n}$. In order to show these results, we need to analyze the corresponding integral equation for (1.1):

$$\begin{aligned} u(x, t) &= (G(t) * u_0)(x) + \int_0^t \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u \right) (\tau) \right) (x) d\tau \\ &\quad + \sum_{i=1}^n \int_0^t \left(\frac{\partial G}{\partial x_i}(t - \tau) * b \frac{\partial u}{\partial x_i}(\tau) \right) (x) d\tau \\ &=: (G(t) * u_0)(x) + D_1(x, t) + D_2(x, t). \end{aligned} \quad (3.1)$$

Then, to complete the proof of Theorems 2.1 and 2.5, we shall derive the leading term for the all terms in the right hand side of (3.1). In order to doing that, we start with introducing the basic properties for the heat kernel $G(x, t)$. First, let us mention about the L^p -decay estimate. Let $\alpha \in \mathbb{Z}_+^n$ and $p \in [1, \infty]$. Then, the following estimate holds (for the proof, see e.g. [7]):

$$\|\partial_x^\alpha G(t)\|_{L^p} \leq Ct^{-\frac{n}{2}\left(1-\frac{1}{p}\right)-\frac{|\alpha|}{2}}, \quad t > 0. \quad (3.2)$$

Moreover, we would like to introduce the following asymptotic formula for $(G(t) * u_0)(x)$. For the proof, let us refer to [1]. This formula plays an important role in the proofs of main results.

Proposition 3.1 ([1]). *Let $u_0 \in L^1(\mathbb{R}^n; 1 + |x|)$ and $p \in [1, \infty]$. Then, we have*

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}\left(1-\frac{1}{p}\right)+\frac{1}{2}} \|G(t) * u_0 - MG(t) + m \cdot \nabla G(t)\|_{L^p} = 0, \quad (3.3)$$

where $G(x, t)$ and M are defined by (1.6), while m is defined by (1.15).

By virtue of this proposition, to completes the proofs of the main results, we only need to consider the asymptotic behavior of the Duhamel term in (3.1). In what follows, we shall derive the asymptotic profiles for $D_1(x, t)$ and $D_2(x, t)$ in Subsections 3.1 and 3.2, respectively.

3.1 Proof of Theorem 2.1

In this subsection, we would like to prove Theorem 2.1. First, let us derive the asymptotic profiles for $D_1(x, t)$ in the right hand side of (3.1). As we can see from the results obtained in Zuazua [12] and Duro-Carpio [3], the asymptotic profile of $D_1(x, t)$ is given by $\alpha_n |M|^{\frac{2}{n}} M(\log t) d \cdot \nabla G(x, t)$. In the present paper, we construct the second asymptotic profile of $D_1(x, t)$ and generalize the results given in [3, 12]. More precisely, the following asymptotic formula can be established:

Theorem 3.2. *Let $n \geq 1$ and $q = 1 + \frac{2}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}\left(1-\frac{1}{p}\right)+\frac{1}{2}} \left\| D_1(t) - \alpha_n |M|^{\frac{2}{n}} M(\log t) d \cdot \nabla G(t) - |M|^{\frac{2}{n}} M \Psi(t) - \mathcal{M} \cdot \nabla G(t) \right\|_{L^p} = 0, \quad (3.4)$$

where $D_1(x, t)$ is defined by (3.1), while $G(x, t)$ and M , and α_n are defined by (1.6) and (1.12), respectively. In addition, $\Psi(x, t)$, \mathcal{M} and \mathcal{N} are defined by (2.1), (2.3) and (2.4), respectively.

Remark 3.3. *For the proof of this theorem, we note that the assumption $n \geq 2$ is unnecessary. This restriction of the dimension $n \geq 2$ will be used in Proposition 3.7 below.*

To prove this theorem, let us introduce the following new functions $v(x, t)$ and $w(x, t)$:

$$v(x, t) := \int_1^t \left(d \cdot \nabla G(t - \tau) * \left(|G|^{\frac{2}{n}} G \right) (\tau) \right) (x) d\tau, \quad (3.5)$$

$$\begin{aligned} w(x, t) &:= \int_0^1 \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u \right) (\tau) \right) (x) d\tau \\ &\quad + \int_1^t \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u - |MG|^{\frac{2}{n}} MG \right) (\tau) \right) (x) d\tau. \end{aligned} \quad (3.6)$$

By, using the functions $v(x, t)$ and $w(x, t)$ above, $D_1(x, t)$ in (3.1) can be split as follows:

$$\begin{aligned} D_1(x, t) &= \int_0^t \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u \right) (\tau) \right) (x) d\tau \\ &= \int_1^t \left(d \cdot \nabla G(t - \tau) * \left(|MG|^{\frac{2}{n}} MG \right) (\tau) \right) (x) d\tau \end{aligned}$$

$$\begin{aligned}
& + \int_0^1 \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u \right) (\tau) \right) (x) d\tau \\
& + \int_1^t \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u - |MG|^{\frac{2}{n}} MG \right) (\tau) \right) (x) d\tau = |M|^{\frac{2}{n}} Mv(x, t) + w(x, t).
\end{aligned}$$

Then, from the above expression, we can see that the following relation holds:

$$\begin{aligned}
D_1(x, t) - \alpha_n |M|^{\frac{2}{n}} M (\log t) d \cdot \nabla G(x, t) - \mathcal{M} \cdot \nabla G(x, t) - |M|^{\frac{2}{n}} M \Psi(x, t) \\
= |M|^{\frac{2}{n}} M \{v(x, t) - \alpha_n (\log t) d \cdot \nabla G(x, t) - \Psi(x, t)\} + \{w(x, t) - \mathcal{M} \cdot \nabla G(x, t)\}. \quad (3.7)
\end{aligned}$$

Therefore, in order to complete the proof of Theorem 3.2, we need to evaluate the first and second terms of the above equation. First, let us derive the following estimate for the first term:

Proposition 3.4. *Let $n \geq 1$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\|v(t) - \alpha_n (\log t) d \cdot \nabla G(t) - \Psi(t)\|_{L^p} \leq C t^{-\frac{n}{2} \left(1 - \frac{1}{p}\right) - 1}, \quad t > 1, \quad (3.8)$$

where $v(x, t)$, $G(x, t)$, $\Psi(x, t)$ and α_n are defined by (3.5), (1.6), (2.1) and (1.12), respectively.

Proof. First, we set $V(x, t) := \alpha_n (\log t) d \cdot \nabla G(x, t)$. Then, it follows that

$$\begin{aligned}
\partial_t V(x, t) &= \frac{\alpha_n}{t} (d \cdot \nabla G(x, t)) + \alpha_n (\log t) \partial_t (d \cdot \nabla G(x, t)) \\
&= \frac{\alpha_n}{t} (d \cdot \nabla G(x, t)) + \alpha_n (\log t) \Delta (d \cdot \nabla G(x, t)).
\end{aligned}$$

Thus, we can see that $V(x, t)$ is the solution to the following Cauchy problem:

$$\begin{aligned}
\partial_t V - \Delta V &= \frac{\alpha_n}{t} (d \cdot \nabla G(x, t)), \quad x \in \mathbb{R}^n, \quad t > 1, \\
V(x, 1) &= 0, \quad x \in \mathbb{R}^n.
\end{aligned} \quad (3.9)$$

By virtue of the Duhamel principle, $V(x, t)$ can be rewritten by

$$V(x, t) = \int_1^t (d \cdot \nabla G(t - \tau) * \alpha_n \tau^{-1} G(\tau)) (x) d\tau.$$

Now, it follows from (1.6) and (2.2) that $(|G|^{\frac{2}{n}} G - \alpha_n \tau^{-1} G)(y, \tau) = F(y, \tau)$. Therefore, using the above expression and (3.5), we are able to obtain

$$\begin{aligned}
v(x, t) - V(x, t) &= \int_1^t \left(d \cdot \nabla G(t - \tau) * \left(|G|^{\frac{2}{n}} G - \alpha_n \tau^{-1} G \right) (\tau) \right) (x) d\tau \\
&= d \cdot \nabla \int_1^t \int_{\mathbb{R}^n} G(x - y, t - \tau) F(y, \tau) dy d\tau =: d \cdot \nabla I(x, t). \quad (3.10)
\end{aligned}$$

Next, let us transform $I(x, t)$ in the above equation (3.10), by using the scaling argument. Actually, from (1.6) and (2.2), we note that the following self-similar structures of $G(x, t)$ and $F(y, \tau)$ hold true:

$$G(x, t) = \lambda^n G(\lambda x, \lambda^2 t), \quad F(y, \tau) = \lambda^{n+2} F(\lambda y, \lambda^2 \tau), \quad \text{for } \lambda > 0.$$

By virtue of the above properties, we can easily obtain

$$G(x - y, t - \tau) = t^{-\frac{n}{2}} G\left(\frac{x - y}{\sqrt{t}}, 1 - \frac{\tau}{t}\right), \quad F(y, \tau) = t^{-\frac{n}{2} - 1} F\left(\frac{y}{\sqrt{t}}, \frac{\tau}{t}\right). \quad (3.11)$$

Therefore, using the above equations and the change of variables, we obtain

$$\begin{aligned}
I(x, t) &= t^{-n-1} \int_1^t \int_{\mathbb{R}^n} G\left(\frac{x-y}{\sqrt{t}}, 1-\frac{\tau}{t}\right) F\left(\frac{y}{\sqrt{t}}, \frac{\tau}{t}\right) dy d\tau \\
&= t^{-\frac{n}{2}} \int_{\frac{1}{t}}^1 \int_{\mathbb{R}^n} G\left(\frac{x}{\sqrt{t}}-z, 1-s\right) F(z, s) dz ds \\
&= t^{-\frac{n}{2}} \int_{\frac{1}{t}}^1 (G(1-s) * F(s))\left(\frac{x}{\sqrt{t}}\right) ds.
\end{aligned} \tag{3.12}$$

Now, noting the definition (2.1) of the function $\Psi(x, t)$, it can be rewritten as follows:

$$\begin{aligned}
\Psi(x, t) &= t^{-\frac{n}{2}-\frac{1}{2}} \Psi_*\left(\frac{x}{\sqrt{t}}\right) = t^{-\frac{n}{2}-\frac{1}{2}} d \cdot \nabla \left(\int_0^1 (G(1-s) * F(s))(x) ds \right) \Big|_{x=\frac{x}{\sqrt{t}}} \\
&= t^{-\frac{n}{2}} d \cdot \nabla \int_0^1 (G(1-s) * F(s))\left(\frac{x}{\sqrt{t}}\right) ds.
\end{aligned} \tag{3.13}$$

Thus, for any $p \in [1, \infty]$, it follows from (3.10) and (3.12) that

$$\begin{aligned}
\|v(t) - V(t) - \Psi(t)\|_{L^p} &= t^{-\frac{n}{2}} \left\| d \cdot \nabla \int_0^{\frac{1}{t}} (G(1-s) * F(s))\left(\frac{\cdot}{\sqrt{t}}\right) ds \right\|_{L^p} \\
&= t^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}} \left\| d \cdot \nabla \int_0^{\frac{1}{t}} G(1-s) * F(s) ds \right\|_{L^p}.
\end{aligned} \tag{3.14}$$

In what follows, we shall evaluate the right hand side of the above (3.14). Now, noticing $\int_{\mathbb{R}^n} F_*(z) dz = 0$. Therefore, using the mean value theorem, (3.1), the Schwarz inequality, the Young inequality and (3.2), we obtain

$$\begin{aligned}
&\left\| d \cdot \nabla \int_0^{\frac{1}{t}} G(1-s) * F(s) ds \right\|_{L^p} \\
&\leq \int_0^{\frac{1}{t}} \left\| d \cdot \nabla \int_{\mathbb{R}^n} G(\cdot - y, 1-s) s^{-\frac{n}{2}-1} F_*\left(\frac{y}{\sqrt{s}}\right) dy \right\|_{L^p} ds \\
&= \int_0^{\frac{1}{t}} s^{-\frac{n}{2}-1} \left\| d \cdot \nabla \int_{\mathbb{R}^n} \left(\int_0^1 y \cdot \nabla G(\cdot - \theta y, 1-s) d\theta \right) F_*\left(\frac{y}{\sqrt{s}}\right) dy \right\|_{L^p} ds \\
&= \int_0^{\frac{1}{t}} s^{-\frac{n}{2}-1} \left\| \int_{\mathbb{R}^n} \left(\int_0^1 d \cdot (D^2 G(\cdot - \theta y, 1-s) y) d\theta \right) F_*\left(\frac{y}{\sqrt{s}}\right) dy \right\|_{L^p} ds \\
&\leq |d| \int_0^{\frac{1}{t}} s^{-\frac{n}{2}-\frac{1}{2}} \int_0^1 \left\| \frac{1}{\theta^n} \int_{\mathbb{R}^n} \left| D^2 G\left(\frac{\cdot}{\theta} - y, \frac{1-s}{\theta^2}\right) \right| \frac{|y|}{\sqrt{s}} \left| F_*\left(\frac{y}{\sqrt{s}}\right) \right| dy \right\|_{L^p} d\theta ds \\
&\leq C \int_0^{\frac{1}{t}} s^{-\frac{n}{2}-\frac{1}{2}} \int_0^1 \theta^{-n(1-\frac{1}{p})-2} (1-s)^{-\frac{n}{2}(1-\frac{1}{p})-1} \theta^{n(1-\frac{1}{p})+2} s^{\frac{n}{2}} \| \cdot \|_{L^1} |F_*|_{L^1} d\theta ds \\
&\leq C \| \cdot \|_{L^1} \int_0^{\frac{1}{t}} s^{-\frac{1}{2}} (1-s)^{-\frac{n}{2}(1-\frac{1}{p})-1} ds \leq C \| \cdot \|_{L^1} t^{-\frac{1}{2}}, \quad t > 1,
\end{aligned} \tag{3.15}$$

for any $p \in [1, \infty]$. Finally, combining (3.14) and (3.15), we can conclude that the desired estimate (3.8) is true. This completes the proof. \square

Next, we would like to treat the second term of (3.7). To doing that, we prepare a lemma:

Lemma 3.5. *Let $n \geq 1$ and $q = 1 + \frac{2}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the solution $u(x, t)$ to (1.1) satisfies the following estimate:*

$$\left\| \left(|u|^{\frac{2}{n}} u \right) (t) - \left(|MG|^{\frac{2}{n}} MG \right) (t) \right\|_{L^p} \leq C t^{-\frac{n}{2} \left(1 - \frac{1}{p}\right) - \frac{3}{2}} \log t, \quad t > 1, \quad (3.16)$$

where $G(x, t)$ and M are defined by (1.6).

Proof. First, we recall that the following inequality holds:

$$\left| |\alpha|^{q-1} \alpha - |\beta|^{q-1} \beta \right| \leq C \left(|\alpha|^{q-1} + |\beta|^{q-1} \right) |\alpha - \beta|, \quad \alpha, \beta \in \mathbb{R}, \quad q > 1.$$

Therefore, it follows from (1.4), (1.7) and (3.2) that

$$\begin{aligned} & \left\| \left(|u|^{\frac{2}{n}} u \right) (t) - \left(|MG|^{\frac{2}{n}} MG \right) (t) \right\|_{L^p} \\ & \leq C \left(\|u(t)\|_{L^\infty}^{\frac{2}{n}} + \|MG(t)\|_{L^\infty}^{\frac{2}{n}} \right) \|u(t) - MG(t)\|_{L^p} \\ & \leq C \left(\left(t^{-\frac{n}{2}} \right)^{\frac{2}{n}} + \left(|M| t^{-\frac{n}{2}} \right)^{\frac{2}{n}} \right) t^{-\frac{n}{2} \left(1 - \frac{1}{p}\right) - \frac{1}{2}} \log t \leq C t^{-\frac{n}{2} \left(1 - \frac{1}{p}\right) - \frac{3}{2}} \log t, \quad t > 1. \end{aligned}$$

This completes the proof. \square

In what follows, let us deal with the second term in (3.7), i.e. we would like to introduce the asymptotic profile for $w(x, t)$. Actually, the following formula can be established:

Proposition 3.6. *Let $n \geq 1$ and $q = 1 + \frac{2}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2} \left(1 - \frac{1}{p}\right) + \frac{1}{2}} \|w(t) - \mathcal{M} \cdot \nabla G(t)\|_{L^p} = 0, \quad (3.17)$$

where $w(x, t)$, $G(x, t)$ and \mathcal{M} are defined by (3.6), (1.6) and (2.3), respectively.

Proof. First of all, for simplicity, let us define $\rho(x, t)$, \mathcal{M}_0 and \mathcal{M}_1 as follows:

$$\rho(x, t) := \left(|u|^{\frac{2}{n}} u - |MG|^{\frac{2}{n}} MG \right) (x, t), \quad (3.18)$$

$$\mathcal{M}_0 := d \int_0^1 \int_{\mathbb{R}^n} \left(|u|^{\frac{2}{n}} u \right) (y, \tau) dy d\tau, \quad \mathcal{M}_1 := d \int_1^\infty \int_{\mathbb{R}^n} \rho(y, \tau) dy d\tau. \quad (3.19)$$

Using the definition of \mathcal{M} in (2.3), \mathcal{M}_0 and \mathcal{M}_1 in (3.19), we have

$$\mathcal{M} = d \int_0^1 \int_{\mathbb{R}^n} \left(|u|^{\frac{2}{n}} u \right) (y, \tau) dy d\tau + d \int_1^\infty \int_{\mathbb{R}^n} \rho(y, \tau) dy d\tau = \mathcal{M}_0 + \mathcal{M}_1.$$

Then, it follows from the definition of $w(x, t)$ in (3.6) that

$$\begin{aligned} & w(x, t) - \mathcal{M} \cdot \nabla G(x, t) \\ & = w(x, t) - \mathcal{M}_0 \cdot \nabla G(x, t) - \mathcal{M}_1 \cdot \nabla G(x, t) \\ & = \left\{ \int_1^t \left(d \cdot \nabla G(t - \tau) * \rho(\tau) \right) (x) d\tau - \mathcal{M}_1 \cdot \nabla G(x, t) \right\} \\ & \quad + \left\{ \int_0^1 \left(d \cdot \nabla G(t - \tau) * \left(|u|^{\frac{2}{n}} u \right) (\tau) \right) (x) d\tau - \mathcal{M}_0 \cdot \nabla G(x, t) \right\}. \end{aligned} \quad (3.20)$$

Therefore, in order to prove (3.34), it is sufficient to derive the following asymptotic formulas:

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2} \left(1 - \frac{1}{p}\right) + \frac{1}{2}} \left\| \int_1^t d \cdot \nabla G(t - \tau) * \rho(\tau) d\tau - \mathcal{M}_1 \cdot \nabla G(t) \right\|_{L^p} = 0, \quad (3.21)$$

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}(1-\frac{1}{p})+\frac{1}{2}} \left\| \int_0^1 d \cdot \nabla G(t-\tau) * \left(|u|^{\frac{2}{n}} u \right) (\tau) d\tau - \mathcal{M}_0 \cdot \nabla G(t) \right\|_{L^p} = 0, \quad (3.22)$$

for any $p \in [1, \infty]$. First, we shall prove (3.21). From the definition of \mathcal{M}_1 by (3.19), we have

$$\begin{aligned} & \int_1^t (d \cdot \nabla G(t-\tau) * \rho(\tau))(x) d\tau - \mathcal{M}_1 \cdot \nabla G(x, t) \\ &= \int_1^t \int_{\mathbb{R}^n} d \cdot \nabla G(x-y, t-\tau) \rho(y, \tau) dy d\tau - d \cdot \nabla G(x, t) \int_1^\infty \int_{\mathbb{R}^n} \rho(y, \tau) dy d\tau \\ &= \int_1^t \int_{\mathbb{R}^n} d \cdot \nabla (G(x-y, t-\tau) - G(x, t)) \rho(y, \tau) dy d\tau - d \cdot \nabla G(x, t) \int_t^\infty \int_{\mathbb{R}^n} \rho(y, \tau) dy d\tau \\ &=: X(x, t) + Y(x, t). \end{aligned} \quad (3.23)$$

In addition, let us take $\varepsilon \in (0, 1)$ and split the integral of $X(x, t)$ as follows:

$$\begin{aligned} X(x, t) &= \int_{\frac{\varepsilon t}{2}}^t \int_{\mathbb{R}^n} d \cdot \nabla (G(x-y, t-\tau) - G(x, t)) \rho(y, \tau) dy d\tau \\ &\quad + \int_1^{\frac{\varepsilon t}{2}} \left(\int_{|y| \geq \varepsilon \sqrt{t}} + \int_{|y| \leq \varepsilon \sqrt{t}} \right) d \cdot \nabla (G(x-y, t-\tau) - G(x, t)) \rho(y, \tau) dy d\tau \\ &=: X_1(x, t) + X_2(x, t) + X_3(x, t). \end{aligned} \quad (3.24)$$

In what follows, we would like to evaluate $X_1(x, t)$, $X_2(x, t)$ and $X_3(x, t)$. Here and later, let $p \in [1, \infty]$. First, from the definition of $\rho(y, \tau)$ in (3.18) and Lemma 3.5, we obtain

$$\|\rho(\tau)\|_{L^p} = \left\| \left(|u|^{\frac{2}{n}} u \right) (\tau) - \left(|MG|^{\frac{2}{n}} MG \right) (\tau) \right\|_{L^p} \leq C \tau^{-\frac{n}{2}(1-\frac{1}{p})-\frac{3}{2}} \log \tau, \quad \tau > 1. \quad (3.25)$$

Now, we shall treat $X_1(x, t)$. From the Young inequality, (3.2) and (3.25), we have

$$\begin{aligned} \|X_1(t)\|_{L^p} &\leq |d| \int_{\frac{\varepsilon t}{2}}^t \|\nabla G(t-\tau)\|_{L^1} \|\rho(\tau)\|_{L^p} d\tau + |d| \|\nabla G(t)\|_{L^p} \int_{\frac{\varepsilon t}{2}}^t \|\rho(\tau)\|_{L^1} d\tau \\ &\leq C \int_{\frac{\varepsilon t}{2}}^t (t-\tau)^{-\frac{1}{2}} \tau^{-\frac{n}{2}(1-\frac{1}{p})-\frac{3}{2}} \log \tau d\tau + C t^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}} \int_{\frac{\varepsilon t}{2}}^t \tau^{-\frac{3}{2}} \log \tau d\tau \\ &\leq C \left(\varepsilon^{-\frac{n}{2}(1-\frac{1}{p})-\frac{3}{2}} + \varepsilon^{-\frac{3}{2}} \right) t^{-\frac{n}{2}(1-\frac{1}{p})-1} \log t, \quad t > \frac{2}{\varepsilon}. \end{aligned} \quad (3.26)$$

Next, let us deal with $X_2(x, t)$. From (3.2), we can easily obtain

$$\begin{aligned} \|X_2(t)\|_{L^p} &\leq |d| \int_1^{\frac{\varepsilon t}{2}} (\|\nabla G(\cdot, t-\tau)\|_{L^p} + \|\nabla G(t)\|_{L^p}) \int_{|y| \geq \varepsilon \sqrt{t}} |\rho(y, \tau)| dy d\tau \\ &\leq C \int_1^{\frac{\varepsilon t}{2}} \left((t-\tau)^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}} + t^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}} \right) \int_{|y| \geq \varepsilon \sqrt{t}} |\rho(y, \tau)| dy d\tau \\ &\leq C t^{-\frac{n}{2}(1-\frac{1}{p})-\frac{1}{2}} z(t), \quad t > \frac{2}{\varepsilon}, \end{aligned} \quad (3.27)$$

where the function $z(t)$ is defined by

$$z(t) := \int_1^{\frac{\varepsilon t}{2}} \int_{|y| \geq \varepsilon \sqrt{t}} |\rho(y, \tau)| dy d\tau = \int_1^\infty \chi_{[1, \frac{\varepsilon t}{2}]}(\tau) \int_{|y| \geq \varepsilon \sqrt{t}} |\rho(y, \tau)| dy d\tau.$$

Moreover, it follows from (3.25) that $\int_1^\infty \int_{\mathbb{R}^n} |\rho(y, \tau)| dy d\tau < \infty$. Therefore, from the Lebesgue dominated convergence theorem, we obtain

$$\lim_{t \rightarrow \infty} z(t) = \int_1^\infty \lim_{t \rightarrow \infty} \left(\chi_{[1, \frac{\varepsilon t}{2}]}(\tau) \int_{|y| \geq \varepsilon \sqrt{t}} |\rho(y, \tau)| dy \right) d\tau = 0. \quad (3.28)$$

Finally, we would like to evaluate $X_3(x, t)$. First, recalling $\partial_t G = \Delta G$ and using (3.2), the following estimate holds for $1 \leq \tau \leq \frac{\varepsilon t}{2}$:

$$\begin{aligned} & \|d \cdot \nabla (G(t - \tau) - G(t))\|_{L^p} \\ &= \tau \left\| d \cdot \nabla \int_0^1 \partial_t G(t - \theta\tau) d\theta \right\|_{L^p} \leq \tau \int_0^1 \|d \cdot \nabla \Delta G(t - \theta\tau)\|_{L^p} d\theta \\ &\leq C\tau \int_0^1 (t - \theta\tau)^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{3}{2}} d\theta \leq C\varepsilon t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}}. \end{aligned} \quad (3.29)$$

Moreover, under this situation, if we also assume $|y| \leq \varepsilon\sqrt{t}$, the following estimate holds too:

$$\begin{aligned} & \|d \cdot \nabla (G(\cdot - y, t - \tau) - G(\cdot, t - \tau))\|_{L^p} \leq |d||y| \int_0^1 \|D^2 G(\cdot - \theta y, t - \tau)\|_{L^p} d\theta \\ &\leq C\varepsilon\sqrt{t} (t - \tau)^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-1} \leq C\varepsilon t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}}. \end{aligned} \quad (3.30)$$

Thus, from (3.29) and (3.30), we obtain

$$\begin{aligned} & \|d \cdot \nabla (G(\cdot - y, t - \tau) - G(\cdot, t))\|_{L^p} \\ &\leq \|d \cdot \nabla (G(\cdot - y, t - \tau) - G(\cdot, t - \tau))\|_{L^p} + \|d \cdot \nabla (G(\cdot, t - \tau) - G(\cdot, t))\|_{L^p} \\ &\leq C\varepsilon t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}}, \quad |y| \leq \varepsilon\sqrt{t}, \quad 1 \leq \tau \leq \frac{\varepsilon t}{2}. \end{aligned} \quad (3.31)$$

Therefore, combining the fact $\int_1^\infty \int_{\mathbb{R}^n} |\rho(y, \tau)| dy d\tau < \infty$ and (3.31), we are able to see that

$$\begin{aligned} \|X_3(t)\|_{L^p} &\leq \int_1^{\frac{\varepsilon t}{2}} \int_{|y| \leq \varepsilon\sqrt{t}} \|d \cdot \nabla (G(\cdot - y, t - \tau) - G(\cdot, t))\|_{L^p} |\rho(y, \tau)| dy d\tau \\ &\leq C\varepsilon t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}} \int_1^{\frac{\varepsilon t}{2}} \int_{|y| \leq \varepsilon\sqrt{t}} |\rho(y, \tau)| dy d\tau \leq C\varepsilon t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}}, \quad t > \frac{2}{\varepsilon}. \end{aligned} \quad (3.32)$$

Next, let us evaluate $Y(x, t)$. It follows from (3.2) and (3.25) that

$$\begin{aligned} \|Y(t)\|_{L^p} &\leq |d| \|\nabla G(t)\|_{L^p} \int_t^\infty \|\rho(\tau)\|_{L^1} d\tau \leq C t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}} \int_t^\infty \tau^{-\frac{3}{2}} \log \tau d\tau \\ &= C t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}} \left(-2 \left[\tau^{-\frac{1}{2}} \log \tau \right]_t^\infty + 2 \int_t^\infty \tau^{-\frac{3}{2}} d\tau \right) \\ &\leq C t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-\frac{1}{2}} \cdot C t^{-\frac{1}{2}} (\log t + 1) \leq C t^{-\frac{n}{2}} \left(1 - \frac{1}{p}\right)^{-1} (\log t + 1), \quad t > 1. \end{aligned} \quad (3.33)$$

Summarizing up (3.23), (3.24), (3.26), (3.27), (3.28), (3.32) and (3.33), we eventually obtain the following estimate:

$$\limsup_{t \rightarrow \infty} t^{\frac{n}{2} \left(1 - \frac{1}{p}\right) + \frac{1}{2}} \left\| \int_1^t d \cdot \nabla G(t - \tau) * \rho(\tau) d\tau - \mathcal{M}_1 \cdot \nabla G(t) \right\|_{L^p} \leq C\varepsilon.$$

Thus, we obtain (3.21) because $\varepsilon > 0$ can be chosen arbitrarily small. Here, we remark that (3.22) can be shown by the similar way. Actually, by virtue of the property (1.3), we have

$$\int_0^1 \int_{\mathbb{R}^n} |u(y, \tau)|^{1 + \frac{2}{n}} dy d\tau \leq \int_0^1 \|u(\tau)\|_{L^\infty}^{\frac{2}{n}} \|u(\tau)\|_{L^1} d\tau \leq C.$$

Therefore, we can prove (3.22) more easily than (3.21). As a result, we are able to see that the asymptotic formula (3.34) is true. This completes the proof. \square

Proof of Theorem 3.2. Combining Propositions 3.4, 3.6 and (3.7), we can immediately see that the desired asymptotic formula (3.4) is true. This completes the proof. \square

Finally, in the rest of this subsection, let us mention about the asymptotic profile for $D_2(x, t)$ in (3.1) when $n \geq 2$. Actually, this is already known by the previous works. More precisely, the following formula is given by Duro–Carpio [3] (for the proof, see Proposition 1 in [3]).

Theorem 3.7 ([3]). *Let $n \geq 2$ and $q = 1 + \frac{2}{n}$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\lim_{t \rightarrow \infty} t^{\frac{n}{2}(1-\frac{1}{p})+\frac{1}{2}} \|D_2(t) - \mathcal{N} \cdot \nabla G(t)\|_{L^p} = 0, \quad (3.34)$$

where $D_2(x, t)$, $G(x, t)$ and \mathcal{N} are defined by (3.1), (1.6) and (2.4), respectively.

Remark 3.8. *In order to prove this proposition, the assumption $n \geq 2$ is necessary. Actually, if $n = 1$, we cannot guarantee $|\mathcal{N}| < \infty$. Now, let us consider the following integral:*

$$\begin{aligned} \int_0^t \int_{\mathbb{R}^n} |b(y) \nabla u(y, \tau)| dy d\tau &\leq \int_0^1 \|\nabla u(\tau)\|_{L^1} \|b\|_{L^\infty} d\tau + \int_1^t \|\nabla u(\tau)\|_{L^\infty} \|b\|_{L^1} d\tau \\ &\leq C \|b\|_{L^\infty} \int_0^1 \tau^{-\frac{1}{2}} d\tau + C \|b\|_{L^1} \int_1^t \tau^{-\frac{n}{2}-\frac{1}{2}} d\tau, \quad t > 1. \end{aligned} \quad (3.35)$$

Here, we used (1.5) and the following estimate (for the proof, see Lemma 1 in [3]):

$$\|\nabla u(t)\|_{L^1} \leq C t^{-\frac{1}{2}}, \quad t > 0. \quad (3.36)$$

Therefore, if $n \geq 2$, we are able to see that $|\mathcal{N}| \leq C \int_0^\infty \int_{\mathbb{R}^n} |b(y) \nabla u(y, \tau)| dy d\tau < \infty$. On the other hand, the right hand side of (3.35) grows with the order $O(\log t)$ when $n = 1$.

End of the proof of Theorem 2.1. Applying Proposition 3.6, Theorems 3.2 and 3.7 for (3.1), we can conclude that the desired asymptotic formula (2.5) holds. This completes the proof. \square

3.2 Proof of Theorem 2.5

In this subsection, we would like to prove Theorem 2.5. In what follows, let us consider the case of $n = 1$ and $q = 3$. By virtue of Theorem 3.2, we have already obtained the asymptotic formula for $D_1(x, t)$ even if $n = 1$. Therefore, it is sufficient to derive the leading term of $D_2(x, t)$ when $n = 1$. In [3], Duro–Carpio derived the asymptotic profile for $D_2(x, t)$. Here, we shall slightly modify their obtained asymptotic profile and derive the second asymptotic profile for $D_2(x, t)$. Indeed, we have succeeded to show the following asymptotic formula:

Theorem 3.9. *Let $n = 1$ and $q = 3$. Suppose $u_0 \in L^1(\mathbb{R}; 1 + |x|) \cap L^\infty(\mathbb{R})$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\lim_{t \rightarrow \infty} t^{\frac{1}{2}(1-\frac{1}{p})+\frac{1}{2}} \|D_2(t) - \mathcal{K}(t)(\log t) \partial_x G(t) - \Phi(t) - \mathcal{L} \partial_x G(t)\|_{L^p} = 0, \quad (3.37)$$

where $D_2(x, t)$ is defined by (3.1), while $G(x, t)$, $\Phi(x, t)$, $\mathcal{K}(t)$ and \mathcal{L} are defined by (1.6), (2.9), (2.8) and (2.10), respectively.

In order to prove this theorem, let us decompose $D_2(x, t)$ as follows:

$$\begin{aligned} D_2(x, t) &= \int_0^1 (\partial_x G(t - \tau) * b \partial_x u(\tau))(x) d\tau + \int_1^t (\partial_x G(t - \tau) * b \partial_x u(\tau))(x) d\tau \\ &=: \eta(x, t) + h(x, t). \end{aligned} \quad (3.38)$$

Since $|\mathcal{L}| < \infty$ from (3.35), for the first term $\eta(x, t)$, we can easily show the following asymptotic formula. We omit the proof this proposition because it can be given in the completely same way to prove Proposition 3.6.

Proposition 3.10. *Let $n = 1$ and $q = 3$. Suppose $u_0 \in L^1(\mathbb{R}^n; 1 + |x|) \cap L^\infty(\mathbb{R}^n)$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\lim_{t \rightarrow \infty} t^{\frac{1}{2}\left(1 - \frac{1}{p}\right) + \frac{1}{2}} \|\eta(t) - \mathcal{L}\partial_x G(t)\|_{L^p} = 0, \quad (3.39)$$

where $\eta(x, t)$, $G(x, t)$ and \mathcal{L} are defined by (3.38), (1.6) and (2.10), respectively.

Finally, let us consider the asymptotic behavior of $h(x, t)$ in (3.38). We have slightly modified the proof of Proposition 3.4 and succeeded to derive the following asymptotic formula:

Proposition 3.11. *Let $n = 1$ and $q = 3$. Suppose $u_0 \in L^1(\mathbb{R}; 1 + |x|) \cap L^\infty(\mathbb{R})$. Then, for any $p \in [1, \infty]$, the following asymptotic formula holds:*

$$\|h(t) - \mathcal{K}(t)(\log t)\partial_x G(t) - \Phi(t)\|_{L^p} \leq Ct^{-\frac{1}{2}\left(1 - \frac{1}{p}\right) - 1}, \quad t > 1, \quad (3.40)$$

where $h(x, t)$, $G(x, t)$, $\Phi(x, t)$ and $\mathcal{K}(t)$ are defined by (3.38), (1.6), (2.9) and (2.8), respectively.

Proof. First, we set $H(x, t)$ as follows:

$$H(x, t) := \mathcal{K}(t)(\log t)\partial_x G(x, t) = \left(\int_1^t \int_{\mathbb{R}} b(y)\partial_x u(y, \tau) dy d\tau \right) \partial_x G(x, t).$$

Then, in the same way to get (3.9), we can see that $H(x, t)$ satisfies

$$\begin{aligned} \partial_t H - \partial_x^2 H &= \left(\int_{\mathbb{R}} b(y)\partial_x u(y, t) dy \right) \partial_x G(x, t), \quad x \in \mathbb{R}^n, \quad t > 1, \\ H(x, 1) &= 0, \quad x \in \mathbb{R}^n. \end{aligned}$$

By virtue of the Duhamel principle, $H(x, t)$ can be transformed as follows:

$$H(x, t) = \int_1^t \int_{\mathbb{R}} \partial_x G(x - y, t - \tau) \left(\int_{\mathbb{R}} b(z)\partial_x u(z, \tau) dz \right) \partial_x G(y, \tau) dy d\tau.$$

Therefore, it follows from (3.38), (2.10) and the above expression that

$$\begin{aligned} &h(x, t) - H(x, t) \\ &= \int_1^t \int_{\mathbb{R}} \partial_x G(x - y, t - \tau) \left\{ b(y)\partial_x u(y, \tau) - \left(\int_{\mathbb{R}} b(z)\partial_x u(z, \tau) dz \right) \right\} G(y, \tau) dy d\tau \\ &= \int_1^t \int_{\mathbb{R}} \partial_x G(x - y, t - \tau) \phi(y, \tau) G(y, \tau) dy d\tau =: \partial_x E(x, t). \end{aligned} \quad (3.41)$$

Next, let us transform $E(x, t)$ in the above equation (3.10), by using the scaling argument again. Now, recalling (3.1) and (3.11), it follows from the change of variables that

$$\begin{aligned} E(x, t) &= t^{-1} \int_1^t \int_{\mathbb{R}^n} G\left(\frac{x-y}{\sqrt{t}}, 1 - \frac{\tau}{t}\right) \phi(y, \tau) G\left(\frac{y}{\sqrt{t}}, \frac{\tau}{t}\right) dy d\tau \\ &= \sqrt{t} \int_{\frac{1}{t}}^1 \int_{\mathbb{R}^n} G\left(\frac{x}{\sqrt{t}} - z, 1 - s\right) \phi(\sqrt{t}z, ts) G(z, s) dz ds \\ &= \sqrt{t} \int_{\frac{1}{t}}^1 \left(G(1-s) * \phi(\sqrt{t}\cdot, ts) G(s) \right) \left(\frac{x}{\sqrt{t}} \right) ds. \end{aligned} \quad (3.42)$$

Here, similarly as (3.13), from the definition (2.9) of the function $\Phi(x, t)$, we get

$$\Phi(x, t) = \sqrt{t} \partial_x \left(\int_0^1 \left(G(1-s) * \phi(\sqrt{t}\cdot, ts) G(s) \right) \left(\frac{x}{\sqrt{t}} \right) ds \right)$$

Hence, for any $p \in [1, \infty]$, we have from (3.41) and (3.42) that

$$\begin{aligned} \|h(t) - H(t) - \Phi(t)\|_{L^p} &= \sqrt{t} \left\| \partial_x \left(\int_0^{\frac{1}{t}} (G(1-s) * \phi(\sqrt{t}\cdot, ts) G(s)) \left(\frac{\cdot}{\sqrt{t}} \right) ds \right) \right\|_{L^p} \\ &= t^{\frac{1}{2p}} \left\| \partial_x \int_0^{\frac{1}{t}} G(1-s) * \phi(\sqrt{t}\cdot, ts) G(s) ds \right\|_{L^p}. \end{aligned} \quad (3.43)$$

In what follows, let us evaluate the right hand side of the above (3.43). Since $\int_{\mathbb{R}} \phi(y, \tau) dy = 0$ for all $\tau > 0$, analogously as (3.15), we are able to obtain the following estimate:

$$\begin{aligned} & \left\| \partial_x \left(\int_0^{\frac{1}{t}} G(1-s) * \phi(\sqrt{t}\cdot, ts) G(s) ds \right) \right\|_{L^p} \\ &= \left\| \int_0^{\frac{1}{t}} \int_{\mathbb{R}} \left(\int_0^1 \partial_x^2 G(\cdot - \theta y, 1-s) d\theta \right) \phi(\sqrt{t}y, ts) \frac{y}{\sqrt{s}} G\left(\frac{y}{\sqrt{s}}, 1\right) dy ds \right\|_{L^p} \\ &= \int_0^{\frac{1}{t}} \int_0^1 \left\| \int_{\mathbb{R}} \partial_x^2 \left(\frac{1}{\theta} G\left(\frac{\cdot}{\theta} - y, \frac{1-s}{\theta^2}\right) \right) \phi(\sqrt{t}y, ts) \frac{y}{\sqrt{s}} G\left(\frac{y}{\sqrt{s}}, 1\right) dy \right\|_{L^p} d\theta ds \\ &\leq C \int_0^{\frac{1}{t}} \left(\int_0^1 \eta^{-1} \eta^{-2} \eta^{\frac{1}{p}} \eta^{1-\frac{1}{p}+2} d\eta \right) (1-s)^{-\frac{1}{2}(1-\frac{1}{p})-1} s^{\frac{1}{2}} \left\| \cdot |G(\cdot, 1) \phi(\sqrt{ts}\cdot, ts) \right\|_{L^1} ds \\ &\leq C \left\| \cdot |G(\cdot, 1) \right\|_{L^1} \int_0^{\frac{1}{t}} s^{\frac{1}{2}} (1-s)^{-\frac{1}{2}(1-\frac{1}{p})-1} \left\| \phi(\sqrt{ts}\cdot, ts) \right\|_{L^\infty} ds \\ &\leq C \int_0^{\frac{1}{t}} s^{\frac{1}{2}} (1-s)^{-\frac{1}{2}(1-\frac{1}{p})-1} (\|b\|_{L^\infty} + \|b\|_{L^1}) \|\partial_x u(ts)\|_{L^\infty} ds \\ &\leq C (\|b\|_{L^\infty} + \|b\|_{L^1}) \int_0^{\frac{1}{t}} s^{\frac{1}{2}} (1-s)^{-\frac{1}{2}(1-\frac{1}{p})-1} (ts)^{-1} ds \\ &\leq C (\|b\|_{L^\infty} + \|b\|_{L^1}) t^{-\frac{3}{2}}, \quad t > 1, \end{aligned} \quad (3.44)$$

for any $p \in [1, \infty]$. Here, we used the following estimate:

$$\|\partial_x u(t)\|_{L^\infty} \leq Ct^{-1}, \quad t > 0.$$

We omit the proof of this estimate because it can be given by the same argument used for the proof of (3.36) (for details, see e.g. [3, 11]).

Summarizing up (3.43) and (3.44), we can eventually conclude that the desired result (3.40) is true. This completes the proof. \square

Proof of Theorem 3.9. Combining Propositions 3.10, 3.11 and (3.38), we can immediately see that the desired asymptotic formula (3.37) is true. This completes the proof. \square

End of the proof of Theorem 2.5. Applying Proposition 3.6, Theorems 3.2 and 3.9 for (3.1), we can conclude that the desired asymptotic formula (2.11) holds. This completes the proof. \square

Acknowledgment

This study is supported by Grant-in-Aid for Young Scientists Research No.22K13939, Japan Society for the Promotion of Science. The authors would like to express their appreciation to Professor Ryosuke Nakasato for his useful comments and stimulating discussions.

References

- [1] J. Duoandikoetxea and E. Zuazua: *Moments, masses de Dirac et décomposition de fonctions*, C.R. Acad. Sci. Paris, t. 315, Serie I (1992) 693–698.
- [2] C.N. Dawson, C.J. van Duijn and R.E. Grundy: *Long time asymptotics in contaminant transport in porous media*, SIAM J. Appl. Maths. **56** (1996) 965–993.
- [3] G. Duro and A. Carpio: *Asymptotic profiles for convection-diffusion equations with variable diffusion*, Nonlinear Anal. **45** (2001) 407–433.
- [4] G. Duro and E. Zuazua: *Large time behavior for convection-diffusion equations in \mathbb{R}^n with asymptotically constant diffusion*, Comm. P.D.E. **24** (1999) 1283–1240.
- [5] M. Escobedo and E. Zuaua: *Large time behavior for convection-diffusion equations in \mathbb{R}^n* , J. Funct. Anal. **100** (1991) 119–161.
- [6] I. Fukuda and Y. Irino: *Higher-order asymptotic profiles for solutions to the Cauchy problem for a dispersive-dissipative equation with a cubic nonlinearity*, arXiv.2211.04667.
- [7] M.-H. Giga, Y. Giga and J. Saal: *Nonlinear partial differential equations—Asymptotic behavior of solutions and self-similar solutions—*, Birkhäuser, Boston, Basel, Berlin, 2010.
- [8] K. Ishige and T. Kawakami: *Asymptotic expansions of solutions of the Cauchy problem for nonlinear parabolic equations*, J. Anal. Math. **121** (2013) 317–351.
- [9] M. Kato: *Sharp asymptotics for a parabolic system of chemotaxis in one space dimension*, Differ. Integral Equ. **22** (2009) 35–51.
- [10] T. Nagai and T. Yamada: *Large time behavior of bounded solutions to a parabolic system of chemotaxis in the whole space*, J. Math. Anal. Appl. **336** (2007) 704–726.
- [11] E. Zuazua: *Asymptotic behavior of scalar convection-diffusion equations*, arXiv.2003.11834.
- [12] E. Zuazua: *Weakly nonlinear large time behavior for scalar convection-diffusion equations*, Differ. Integral Equ. **6** (1993) 1481–1492.

Ikki Fukuda

Division of Mathematics and Physics,
Faculty of Engineering, Shinshu University
4-17-1, Wakasato, Nagano, 380-8553, JAPAN
E-mail: i_fukuda@shinshu-u.ac.jp

Shinya Sato

Department of Engineering,
Graduate School of Science and Technology, Shinshu University
4-17-1, Wakasato, Nagano, 380-8553, JAPAN
E-mail: sato.shinya@qri.jp