

CONTINUOUS EXTENSION OF THE DISCRETE SHIFT TRANSLATIONS ON ONE-DIMENSIONAL QUANTUM LATTICE SYSTEMS

HAJIME MORIYA AND HEIDE NARNHOFER

ABSTRACT. We study the continuous extension of discrete shift translations on one-dimensional quantum lattice systems. We explore a specific example provided by a quasi-free C^* -flow on the one-dimensional fermion lattice system, comparing it with its original discrete shift-translations. Specifically, we demonstrate its explicit dynamical formula, which reveals violations of causality and locality. Furthermore, we prove that this quasi-free C^* -flow on the one-dimensional fermion lattice system cannot be extended to the one-dimensional quantum spin lattice system via the Jordan-Wigner transformation.

1. INTRODUCTION

We investigate the continuous extension of discrete shift translations on the one-dimensional fermion lattice system and those on the one-dimensional quantum spin lattice system to C^* -flows. To provide context for our investigation, we review some relevant previous results. In [24], it has been shown that the continuous extension of the shift-translation automorphism group on the one-dimensional quantum spin lattice system cannot be achieved by local Hamiltonians satisfying certain locality condition. This non-existence statement (Corollary 6.1 of [24]) is a generalization of the non-implementability of the shift translation by finite quantum circuits [12]. We also refer to [29], where the implementation of shift translations by Hamiltonians on finite quantum-spin chains is studied in detail.

Our ultimate objective is to demonstrate all C^* -flows (strongly continuous one-parameter groups of automorphisms) that implement continuous shift translations on one-dimensional quantum systems, or to disprove the existence of such C^* -flows. In this article, we narrow our focus to a quasi-free extension of discrete shift translations on the one-dimensional fermion lattice system. Due to the aforementioned no-go statements, this quasi-free C^* -flow lacks both finite propagation speed and the sufficient locality of interactions that implies the Lieb-Robinson bound. We examine some properties of this quasi-free C^* -flow in comparison with its original discrete shift-translation automorphism group on the fermion system. Additionally, we demonstrate that the quasi-free C^* -flow on the fermion system is not

2020 *Mathematics Subject Classification.* Primary 82B10, 46N55.

Key words and phrases. C^* -flows, Shift-translations, discrete versus continuous dynamics, quantum spin lattice systems versus fermion lattice systems, Hamiltonian versus momentum operators.

extendable to the one-dimensional quantum spin lattice system; this constitutes a form of no-go theorem.

From the mathematical perspective, the pursuit of continuous extensions of a discrete automorphism group (or a single automorphism) finds its origins in Kuipier's seminal work [16], which explores the topological structure of unitaries on infinite-dimensional Hilbert spaces. Previous studies [12] [24] mentioned before have examined the topological structure of the discrete shift translations on one-dimensional quantum spin systems using Index theory, and recent research [14] has further developed the topological aspects shown in [12]. According to [22], which generalizes Kuipier's work, the σ -weakly continuous extension of the discrete shift translations on von Neumann algebras (generated by the CAR algebra) obviously exists.

Finally, let us mention the work [13], which deals with a somewhat similar extension problem for the shift translations on the quantum spin chain. The work [13] explores the continuous limit of shift translations on the one-dimensional quantum-spin lattice system by letting the spacing of the unit lattice approach zero, whereas we always consider the given quantum lattice system as fixed. Therefore, our work and [13] tackle different issues.

2. MATHEMATICAL FORMULATION

2.1. Quantum systems. We introduce the spinless fermion lattice system and the spin-one-half quantum spin lattice system on \mathbb{Z} as quasi-local C^* -systems. We refer to [5] for the reference of C^* -algebraic quantum statistical mechanics.

For a subset $I \subset \mathbb{Z}$, $|I|$ denotes the number of sites in I . If $I \subset \mathbb{Z}$ has a finite $|I| < \infty$, we shall denote $I \Subset \mathbb{Z}$. Let c_i and c_i^* denote the annihilation operator and the creation operator of a spinless fermion sitting at $i \in \mathbb{Z}$, respectively. These obey the canonical anti-commutation relations (CARs):

$$(2.1) \quad \{c_i^*, c_j^*\} = \{c_i, c_j\} = 0, \quad \{c_i^*, c_j\} = \delta_{i,j} 1, \quad i, j \in \mathbb{Z}.$$

For each $I \Subset \mathbb{Z}$, define the finite system $\mathfrak{A}^F(I)$ by the $*$ -algebra generated by $\{c_i^*, c_i; i \in I\}$. It is isomorphic to $M_{2^{|I|}}(\mathbb{C})$, the algebra of all $2^{|I|} \times 2^{|I|}$ complex matrices. For $I \subset J \Subset \mathbb{Z}$, $\mathfrak{A}^F(I)$ is naturally embedded into $\mathfrak{A}^F(J)$ as a subalgebra. Let us take

$$(2.2) \quad \mathfrak{A}_\circ^F := \bigcup_{I \Subset \mathbb{Z}} \mathfrak{A}^F(I).$$

By taking the norm (operator norm) completion of \mathfrak{A}_\circ^F we obtain a C^* -algebra \mathfrak{A}^F called the CAR algebra. The dense $*$ -subalgebra \mathfrak{A}_\circ^F in \mathfrak{A}^F is called the local algebra of \mathfrak{A}^F .

Let Θ denote the automorphism on the C^* -algebra \mathfrak{A}^F determined by

$$(2.3) \quad \Theta(c_i) = -c_i, \quad \Theta(c_i^*) = -c_i^*, \quad i \in \mathbb{Z}.$$

As the automorphism $\Theta \in \text{Aut}(\mathfrak{A}^F)$ satisfies $\Theta \circ \Theta = \text{id} \in \text{Aut}(\mathfrak{A}^F)$, it provides the graded structure with \mathfrak{A}^F :

$$(2.4) \quad \begin{aligned} \mathfrak{A}_+^F &:= \{A \in \mathfrak{A}^F \mid \Theta(A) = A\}, & \mathfrak{A}_-^F &:= \{A \in \mathfrak{A}^F \mid \Theta(A) = -A\}, \\ \mathfrak{A}^F &= \mathfrak{A}_+^F + \mathfrak{A}_-^F. \end{aligned}$$

By the graded structure of \mathfrak{A}^F , any element $A \in \mathfrak{A}^F$ has its unique even-odd decomposition:

$$(2.5) \quad A = A_+ + A_-, \quad A_+ := \frac{1}{2}(A + \Theta(A)) \in \mathfrak{A}_+^F, \quad A_- := \frac{1}{2}(A - \Theta(A)) \in \mathfrak{A}_-^F.$$

Next, we will introduce the quantum spin lattice system \mathfrak{A}^S via the fermion lattice system \mathfrak{A}^F using the C^* -algebraic formalism given in [2]. This somewhat roundabout method provides a rigorous frame to deal with the Jordan-Wigner transformation connecting the fermion lattice system to the quantum spin lattice system. Let

$$(2.6) \quad \Theta_-(c_i^\sharp) = \begin{cases} c_i^\sharp & i \geq 1 \\ -c_i^\sharp & i \leq 0 \end{cases},$$

where c_i^\sharp denotes c_i or c_i^* . Let $\hat{\mathfrak{A}}$ denote the C^* -crossed product of \mathfrak{A}^F by the \mathbb{Z}_2 -action $\Theta_- \in \text{Aut}(\mathfrak{A}^F)$. Concretely, $\hat{\mathfrak{A}}$ is generated by the algebra \mathfrak{A}^F and the additional element T satisfying the following algebraic relations:

$$(2.7) \quad T = T^*, \quad T^2 = 1, \quad TAT = \Theta_-(A) \text{ for } A \in \mathfrak{A}^F.$$

The grading Θ on \mathfrak{A}^F is uniquely extended to $\hat{\mathfrak{A}}$ by determining

$$(2.8) \quad \Theta(T) = T.$$

Obviously, $\hat{\mathfrak{A}}$ is decomposed into the following four linear subspaces:

$$(2.9) \quad \hat{\mathfrak{A}} = \mathfrak{A}_+^F + T\mathfrak{A}_+^F + \mathfrak{A}_-^F + T\mathfrak{A}_-^F.$$

By definition, $\hat{\mathfrak{A}}$ includes both the fermion lattice system $\mathfrak{A}^F (= \mathfrak{A}_+^F + \mathfrak{A}_-^F)$ and the quantum spin lattice system \mathfrak{A}^S defined by

$$(2.10) \quad \mathfrak{A}^S \equiv \mathfrak{A}_+^F + T\mathfrak{A}_-^F.$$

The spin operators $\{\sigma_j^x, \sigma_j^y, \sigma_j^z \in M_2(\mathbb{C})\}$ at each $j \in \mathbb{Z}$ are explicitly given by

$$(2.11) \quad \sigma_j^z := 2c_j^*c_j - 1, \quad \sigma_j^x := TS^{(j)}(c_j + c_j^*), \quad \sigma_j^y := iTS^{(j)}(c_j - c_j^*),$$

with

$$(2.12) \quad S^{(j)} := \begin{cases} \sigma_1^z \cdots \sigma_{j-1}^z & j \geq 2 \\ 1 & j = 1 \\ \sigma_0^z \cdots \sigma_j^z & j \leq 0 \end{cases}.$$

By the above definition, using the CARs (2.1), we see that the set $\{\sigma_j^x, \sigma_j^y, \sigma_j^z \in \hat{\mathfrak{A}}; j \in \mathbb{Z}\}$ generates the quantum spin algebra \mathfrak{A}^S on \mathbb{Z} given in (2.10) as

a subalgebra of $\hat{\mathfrak{A}}$. Let Θ^S denote the automorphism on the C^* -algebra \mathfrak{A}^S determined by

$$(2.13) \quad \Theta^S(\sigma_j^x) = -\sigma_j^x, \quad \Theta^S(\sigma_j^y) = -\sigma_j^y, \quad \Theta^S(\sigma_j^z) = \sigma_j^z, \quad i \in \mathbb{Z}.$$

As $\Theta^S \circ \Theta^S = \text{id} \in \text{Aut}(\mathfrak{A}^S)$, Θ^S provides the graded structure with \mathfrak{A}^S as

$$(2.14) \quad \begin{aligned} \mathfrak{A}_+^S &:= \{A \in \mathfrak{A}^S \mid \Theta^S(A) = A\}, \quad \mathfrak{A}_-^S := \{A \in \mathfrak{A}^S \mid \Theta^S(A) = -A\}, \\ \mathfrak{A}^S &= \mathfrak{A}_+^S + \mathfrak{A}_-^S. \end{aligned}$$

We easily see that \mathfrak{A}_+^S is generated by the set $\{\sigma_j^z \in \mathfrak{A}^S; j \in \mathbb{Z}\}$ and even monomials of $\{\sigma_j^x, \sigma_j^y; j \in \mathbb{Z}\}$, and \mathfrak{A}_-^S is generated by odd monomials of $\{\sigma_j^x, \sigma_j^y \in \mathfrak{A}^S; j \in \mathbb{Z}\}$. The following identities give the location of the fermion algebra and the quantum spin algebra in $\hat{\mathfrak{A}}$:

$$(2.15) \quad \mathfrak{A}_+^S = \mathfrak{A}_+^F, \quad \mathfrak{A}_-^S = T\mathfrak{A}_-^F.$$

For each $I \in \mathbb{Z}$, define the subalgebra $\mathfrak{A}^S(I)$ of \mathfrak{A}^S by the finite-dimensional algebra generated by $\{\sigma_j^x, \sigma_j^y, \sigma_j^z; j \in I\}$. Let

$$(2.16) \quad \mathfrak{A}_\circ^S := \bigcup_{I \in \mathbb{Z}} \mathfrak{A}^S(I).$$

This norm dense $*$ -subalgebra in \mathfrak{A}^S is called the local algebra of \mathfrak{A}^S .

2.2. Discrete and continuous shift translations. We first define the discrete shift translations on \mathfrak{A}^F and then those on \mathfrak{A}^S . For each $i \in \mathbb{Z}$, let

$$(2.17) \quad \tau_i^F(c_j) = c_{j+i}, \quad \tau_i^F(c_j^*) = c_{j+i}^*, \quad j \in \mathbb{Z}.$$

The above formulas determine the shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ on \mathfrak{A}^F . For each $i \in \mathbb{Z}$, let

$$(2.18) \quad \tau_i^S(\sigma_j^a) = \sigma_{j+i}^a, \quad a \in \{x, y, z\}, \quad j \in \mathbb{Z}.$$

Then $\{\tau_i^S \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$ gives the shift-translation automorphism group on \mathfrak{A}^S . Our purpose is to extend these \mathbb{Z} -shift-translation automorphism groups to strongly continuous one-parameter group of automorphisms, i.e. C^* -flows.

In the following, we construct a continuous shift-translation group on \mathfrak{A}^F . First, note that our fermion lattice system \mathfrak{A}^F is identical to the CAR algebra over the complex Hilbert space $l^2(\mathbb{Z})$. This identification is done by setting $c_j = c(\chi_j)$ for $j \in \mathbb{Z}$, where $\chi_j \in l^2(\mathbb{Z})$ denotes the indicator determined by

$$(2.19) \quad \chi_j(k) := \delta_{jk}, \quad (k \in \mathbb{Z}).$$

For each $f = (f_j)_{j \in \mathbb{Z}} \in l^2(\mathbb{Z})$, define

$$(2.20) \quad c(f) := \sum_{j \in \mathbb{Z}} f_j c_j \in \mathfrak{A}^F, \quad c^*(f) := \sum_{j \in \mathbb{Z}} f_j c_j^* \in \mathfrak{A}^F.$$

\mathfrak{A}^F is also isomorphic to the CAR algebra over $L^2[-\pi, \pi]$ as C^* -algebras by the isometry of $l^2(\mathbb{Z})$ and $L^2[-\pi, \pi]$ as Hilbert spaces. Here, the closed interval $[-\pi, \pi]$ can be interpreted as the range of the possible momentum of each fermion particle. The connection between $f = (f_j)_{j \in \mathbb{Z}} \in l^2(\mathbb{Z})$ and $\tilde{f}(k) \in L^2[-\pi, \pi]$ is given by the Fourier transformation as follows:

$$(2.21) \quad \begin{aligned} \tilde{f}(k) &:= \sum_{j \in \mathbb{Z}} f_j e^{ijk}, \quad k \in [-\pi, \pi], \\ f_j &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \tilde{f}(k) e^{-ijk} dk \quad j \in \mathbb{Z}. \end{aligned}$$

Hence for each $j \in \mathbb{Z}$

$$(2.22) \quad \widetilde{\chi}_j(k) = e^{ijk}, \quad k \in [-\pi, \pi].$$

The one-step (right) shift translation U_1 on the Hilbert space $l^2(\mathbb{Z})$ is given by

$$(2.23) \quad (U_1 f)_j := f_{j-1}, \quad j \in \mathbb{Z}, \quad \text{for } f = (f_j)_{j \in \mathbb{Z}} \in l^2(\mathbb{Z}).$$

The \mathbb{Z} -group of unitaries $\{U_i \in \mathcal{U}(L^2[-\pi, \pi]); i \in \mathbb{Z}\}$ is given by

$$U_0 = I, \quad U_k := \underbrace{U_1 \cdots U_1}_{k\text{-times}}, \quad U_{-k} := U_k^{-1} \quad (k \in \mathbb{N}).$$

By the Fourier transformation, $U_1 \in \mathcal{U}(l^2(\mathbb{Z}))$ has the following expression on $L^2[-\pi, \pi]$

$$(2.24) \quad \widetilde{U_1} f(k) = e^{ik} \tilde{f}(k), \quad k \in [-\pi, \pi].$$

By the second quantization procedure, the \mathbb{Z} -group of unitaries $\{U_i \in \mathcal{U}(L^2[-\pi, \pi]); i \in \mathbb{Z}\}$ generates the discrete-shift translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ given in (2.17). To interpolate $\{U_i \in \mathcal{U}(L^2[-\pi, \pi]); i \in \mathbb{Z}\}$, we define unitaries U_t ($t \in \mathbb{R}$) on $L^2[-\pi, \pi]$ as

$$(2.25) \quad \widetilde{U_t} f(k) := e^{itk} \tilde{f}(k), \quad k \in [-\pi, \pi].$$

Now we obtain a strongly continuous one-parameter group of unitaries:

$$(2.26) \quad \{U_t \in \mathcal{U}(L^2[-\pi, \pi]); t \in \mathbb{R}\},$$

equivalently,

$$(2.27) \quad \{U_t \in \mathcal{U}(l^2(\mathbb{Z})); t \in \mathbb{R}\}.$$

By Stone's theorem [26], there exists a unique self-adjoint operator \bar{h} on $L^2[-\pi, \pi]$ such that

$$(2.28) \quad U_t = e^{it\bar{h}}, \quad t \in \mathbb{R}.$$

This one-parameter group of unitaries generates a strongly continuous one parameter group of quasi-free (Bogoliubov) automorphisms on \mathfrak{A}^F :

$$(2.29) \quad \{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}.$$

By construction, it is a desired C^* -flow that extends the discrete shift-automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ from the discrete parameter \mathbb{Z} to the continuous parameter \mathbb{R} . We see that the continuous shift-translation C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ given above preserves the grading Θ :

$$(2.30) \quad \Theta(\tau_t^F(A)) = \tau_t^F(\Theta(A)) \quad \text{for all } A \in \mathfrak{A}^F \text{ for each } t \in \mathbb{R}.$$

Hence it induces a C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}_+^F), t \in \mathbb{R}\}$ on the even algebra \mathfrak{A}_+^F by restriction.

The time evolution $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ (2.29) on \mathfrak{A}^F is associated to the second quantized Hamiltonian $\tilde{H}_\infty = \Gamma(\bar{h})$ of the one-particle Hamiltonian \bar{h} . In the following, we shall derive a concrete expression of \tilde{H}_∞ by a formal (non rigorous) method. Let

$$(2.31) \quad \hat{c}_k := \sum_{j \in \mathbb{Z}} c_j e^{ijk}, \quad \hat{c}_k^* := \sum_{j \in \mathbb{Z}} c_j^* e^{ijk}, \quad k \in [-\pi, \pi]$$

Let $\hat{N}_k := \hat{c}_k^* \hat{c}_k$, namely the number operator for the momentum $k \in [-\pi, \pi]$. In terms of \hat{N}_k , \tilde{H}_∞ has the following expression in the momentum space:

$$(2.32) \quad \tilde{H}_\infty = \int_{-\pi}^{\pi} k \hat{N}_k dk.$$

Using

$$[\hat{N}_k, \hat{c}_k] = -i\hat{c}_k, \quad [\hat{N}_k, \hat{c}_k^*] = i\hat{c}_k^*$$

we rewrite \tilde{H}_∞ in the lattice space as

$$\tilde{H}_\infty = \int_{-\pi}^{\pi} \sum_{n \in \mathbb{Z}} \sum_{m \in \mathbb{Z}} k e^{ik(n-m)} c_n^* c_m dk = \sum_{n, m \in \mathbb{Z}} h_{n, m} c_n^* c_m,$$

where for $n, m \in \mathbb{Z}$

$$h_{n, m} = G(n - m) \in \mathbb{C}, \quad G(x) := \int_{-\pi}^{\pi} k e^{ikx} dk.$$

By integration by parts

$$\begin{aligned} G(x) &= \left[k \frac{e^{ikx}}{ix} \right]_{-\pi}^{\pi} - \int_{-\pi}^{\pi} \frac{e^{ikx}}{ix} dk = \pi \frac{e^{i\pi x} + e^{-i\pi x}}{ix} + \frac{e^{i\pi x} - e^{-i\pi x}}{x^2} \\ &= -2\pi i \frac{\cos(\pi x)}{x} + 2i \frac{\sin(\pi x)}{x^2} \end{aligned}$$

By plugging $n - m$ to x , we obtain

$$(2.33) \quad \tilde{H}_\infty = \sum_{n, m \in \mathbb{Z}} h_{n, m} c_n^* c_m, \quad h_{n, m} = \frac{-2\pi i (-1)^{n-m}}{n - m}.$$

In fact, the construction of continuous shift translations described above has been given in §2.1 of [12], §6 of [24], Appendix E of [28], and §III of [30]. We, however, emphasize that the second-quantization method for the fermion system seems not valid for the *infinite* quantum spin lattice system \mathfrak{A}^S . This assertion will be rigorously validated later in Proposition 3.

As the shift-translation C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ is a quasi-free dynamics, we can track its dynamical behavior as follows:

Proposition 1. *The discrete shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ on the one-dimensional fermion lattice system \mathfrak{A}^F can be extended to a quasi-free C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ on \mathfrak{A}^F . It has the exact dynamical formulas in terms of the sinc function $\text{sinc } x := \frac{\sin x}{x}$ as follows. For $j \in \mathbb{Z}$ and $t \in \mathbb{R} \setminus \mathbb{Z}$,*

$$(2.34) \quad \begin{aligned} \tau_t^F(c_j) &= \sum_{l \in \mathbb{Z}} \text{sinc}(\pi(j+t-l))c_l = \sum_{l \in \mathbb{Z}} \frac{(-1)^{j-l} \sin(\pi t)}{\pi(j+t-l)} c_l \in \mathfrak{A}^F, \\ \tau_t^F(c_j^*) &= \sum_{l \in \mathbb{Z}} \text{sinc}(\pi(j+t-l))c_l^* = \sum_{l \in \mathbb{Z}} \frac{(-1)^{j-l} \sin(\pi t)}{\pi(j+t-l)} c_l^* \in \mathfrak{A}^F. \end{aligned}$$

For $j \in \mathbb{Z}$ and $k \in \mathbb{Z}$,

$$(2.35) \quad \tau_k^F(c_j) = c_{j+k} \in \mathfrak{A}_\circ^F, \quad \tau_k^F(c_j^*) = c_{j+k}^* \in \mathfrak{A}_\circ^F.$$

The time-shift translation $\tau_t^F \in \text{Aut}(\mathfrak{A}^F)$ ($\forall t \in \mathbb{R} \setminus \mathbb{Z}$) violates the locality in the sense of Definition 2.2.1 of [10]: There exist a strict local operator whose time-translation is not a local operator.

Proof. The existence of such C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ is given in the text, and (2.35) is obvious. As $\widetilde{U_t \chi_j}(k) = e^{i(j+t)k}$ ($k \in [-\pi, \pi]$) by (2.22) (2.25), we have

$$(2.36) \quad U_t \chi_j(l) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(j+t)k} e^{-ilk} dk = \frac{1}{\pi(j+t-l)} \sin(j+t-l)\pi, \quad (l \in \mathbb{Z}).$$

This yields the formula (2.34). The formula (2.34) implies that for any $j \in \mathbb{Z}$, neither $\tau_t^F(c_j)$ nor $\tau_t^F(c_j^*)$ belongs to \mathfrak{A}_\circ^F for $t \in \mathbb{R} \setminus \mathbb{Z}$. Actually, for each $t \in \mathbb{R} \setminus \mathbb{Z}$, there are many more $A \in \mathfrak{A}_\circ^F$ such that $\tau_t^F(A) \notin \mathfrak{A}_\circ^F$. Thus the locality as stated is violated for all $t \in \mathbb{R} \setminus \mathbb{Z}$. \square

We shall provide several consequences of Proposition 1. The first one is related to the following question posed in § 6.1 of [10]: Is there a meaningful notion of a ground state for QCA(quantum cellular automata)? A similar question naturally arises: How about equilibrium states for discrete automorphism groups? The following corollary provides a positive answer to this question for the shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ on the one-dimensional fermion lattice system \mathfrak{A}^F . (However, we do not know the answer for $\{\tau_i^S \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$.)

Corollary 2. *The continuous shift-translation automorphism group $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ on \mathfrak{A}^F has a unique KMS state φ_β at any inverse temperature $\beta \in \mathbb{R}$. It has a unique ground state φ_∞ at $\beta = \infty$ and a unique ceiling state $\varphi_{-\infty}$ at $\beta = -\infty$.*

Proof. Due to Theorem 1 of [25], there is a unique KMS state φ_β on \mathfrak{A}^F for each $\beta \in \mathbb{R}$ with respect to the time evolution $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$.

There is no invariant vector in $L^2[-\pi, \pi]$ under the action of $\{U_t \in \mathcal{U}(L^2[-\pi, \pi]); t \in \mathbb{R}\}$. Thus by the discussion given in Example 5.3.20 [5], we see that a unique ground state φ_∞ with respect to $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ exists. Similarly, there is a unique ceiling state $\varphi_{-\infty}$ with respect to $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$. \square

Remark 1. For each $\beta \in \mathbb{R}$, the quasi-free KMS state φ_β is not a Gibbs state, since its one-particle density operator $\exp(-\beta\bar{h})$ on $L^2[-\pi, \pi]$ is not of trace-class, see Prop. 5.2.23 [5]. By Theorem 4.3 [23] the KMS state φ_β induces a III_1 factor, since the assumption on the time evolution of this theorem is obviously satisfied by the quasi-free time evolution $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$.

Remark 2. As the second-quantized Hamiltonian $\tilde{H}_\infty = \int_{-\pi}^{\pi} k \hat{N}_k dk$ given in (2.32) generates the spatial translations on \mathfrak{A}^F , it may be naturally identified with the momentum operator “ P ”. Although the existence of a ground state for P on the infinite homogeneous quantum system \mathfrak{A}^F seems to be unexpected, it should be noted that the ground state φ_∞ and the ceiling state $\varphi_{-\infty}$ correspond to the fermi sea occupied by all left-(right-) moving particles, respectively. Of course the identification $\tilde{H}_\infty = P$ is formal; we refer to §4.5 of [21] which compares Hamiltonians and momentum operators in terms of spatial and time crystals.

Remark 3. In addition to the violation of the locality described in Proposition 1, causality, which denotes the one-directional nature of dynamics, is not perfectly satisfied by the continuous extension of shift translations $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$. This is because backward dynamics with respect to the time parameter emerges, as illustrated in (2.34). The violation of the causality is also observed in the continuous shift translations on the finite-quantum spin chains, as depicted in Fig. 1 of [12]. On the infinitely extended system \mathfrak{A}^F , however, the violation of causality for any shift-translation C^* -flow becomes negligible in a large time scale, as we will see later in Proposition 4.

Remark 4. The Lieb-Robinson bound [18] is a key assumption of the work [24], whereas we do not require it. Lieb-Robinson bounds have been shown for power-law interactions with their decay rate $\frac{1}{r^\alpha}$ ($\alpha > d$), where d is the spatial dimension of the quantum lattice system, see [8] [17] [20] [28], particularly [11] for free fermion models on lattices. On the other hand, the formal Hamiltonian \tilde{H}_∞ of the continuous shift translations on the one-dimensional fermion lattice system consists of the two-body translation invariant interactions with $\frac{1}{r}$ -decay ($\alpha = d = 1$) rate, as given in (2.33). Therefore, the quasi-free fermion model associated to the continuous-shift translations has such a long range that it is outside the scope of these previous works.

3. NON EXTENDABILITY OF CONTINUOUS QUASI-FREE SHIFTS TO THE QUANTUM SPIN SYSTEM

In the preceding section, we provided a C^* -flow that extends the discrete shift-translation automorphism group on the fermion lattice system \mathfrak{A}^F . The following proposition demonstrates that it is not possible to extend this quasi-free C^* -flow on \mathfrak{A}^F given in Proposition 1 to the quantum spin lattice system \mathfrak{A}^S .

Proposition 3. *There is no C^* -flow on \mathfrak{A}^S which extends the discrete shift-translation automorphism group $\{\tau_i^S \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$ on \mathfrak{A}^S and coincides with the quasi-free C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ on the even subalgebra $\mathfrak{A}_+^S = \mathfrak{A}_+^F$:*

$$(3.1) \quad \tau_t^S(A) = \tau_t^F(A) \quad \forall A \in \mathfrak{A}_+^S = \mathfrak{A}_+^F \quad \forall t \in \mathbb{R}.$$

Proof. Let $\{\tau_t^S \in \text{Aut}(\mathfrak{A}^S), t \in \mathbb{R}\}$ denote a C^* -flow extension of the discrete shift-translation automorphism group $\{\tau_i^S \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$ on \mathfrak{A}^S . Suppose that it satisfies the assumption (3.1). In the following, we will prove non-existence of such C^* -flow. To this end, we recall Lemma 4.2 of [19], see also [9]. Let $B_2 := c_2^* + c_2$. For each $t \in \mathbb{R}$, we define

$$(3.2) \quad V_t := \tau_t^S(B_2 T) \tau_t^F(B_2) T \in \mathfrak{A}_+^F.$$

The inclusion $V_t \in \mathfrak{A}_+^F$ above is verified as follows. By (3.1) τ_t^S and τ_t^F are the same shift-translation group on $\mathfrak{A}_+^S = \mathfrak{A}_+^F$, and hence they automatically preserve the grading Θ^S, Θ , respectively. So $\tau_t^S(B_2 T) \in \mathfrak{A}_-^S$ for $B_2 T \in \mathfrak{A}_-^S$, and $\tau_t^F(B_2) \in \mathfrak{A}_-^F$ for $B_2 \in \mathfrak{A}_-^F$. Thus we have $\tau_t^S(B_2 T) \in T\mathfrak{A}_-^F$ and $\tau_t^F(B_2) T \in T\mathfrak{A}_-^F$. From the identity $T^2 = 1$ and the inclusion $\mathfrak{A}_-^F \cdot \mathfrak{A}_-^F \subset \mathfrak{A}_+^F$, it follows that the product $\tau_t^S(B_2 T) \cdot \tau_t^F(B_2) T \equiv V_t$ belongs to \mathfrak{A}_+^F . Let

$$(3.3) \quad (\theta_- f)_j := \begin{cases} f_j & j \geq 1 \\ -f_j & j \leq 0 \end{cases}$$

for $f = (f_j)_{j \in \mathbb{Z}} \in l^2(\mathbb{Z})$. The unitary operator θ_- on $l^2(\mathbb{Z})$ generates Θ_- on \mathfrak{A}^F given in (2.6) by the second-quantization procedure. For each $t \in \mathbb{R}$, we define

$$(3.4) \quad \omega_t := U_t \theta_- U_{-t} \theta_- \in \mathcal{U}(l^2(\mathbb{Z})),$$

where $U_t \in \mathcal{U}(l^2(\mathbb{Z}))$ is given in (2.27). In Lemma 4.2 of [19] the following relation is given:

$$(3.5) \quad \text{Ad}(V_t) = \tau_t^S \circ \Theta_- \circ \tau_{-t}^S \circ \Theta_- \text{ on } \mathfrak{A}_+^F.$$

From (3.4) (3.5) it follows that

$$(3.6) \quad V_t c(f) V_t^* = c(\omega_t f), \quad V_t c^*(f) V_t^* = c^*(\omega_t f).$$

We now consider the continuous shift operator U_t in the place of the NESS-dynamics u_t of the original paper [19]. Due to $V_t \in \mathfrak{A}_+^F$, each $\beta_t := \text{Ad}(V_t)$ is an inner automorphism of \mathfrak{A}^F . Accordingly, $1 - \omega_t$ is a Hilbert

Schmidt operator on $l^2(\mathbb{Z})$, see [1]. This entails the following boundedness condition.

$$(3.7) \quad I_t \equiv \sum_{m \in \mathbb{N}} \sum_{n \in \mathbb{N}} |(e_{-m}, U_t e_n)|^2 < \infty, \quad t \in \mathbb{R}.$$

Next, we will show that the required boundedness (3.7) is not satisfied. For each $t \in \mathbb{R}$, we compute

$$(3.8) \quad \begin{aligned} (e_{-m}, U_t e_n) &= \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{-imk} e^{-itk} e^{-ink} dk = \frac{1}{2\pi} \left[\frac{e^{i(-m-t-n)k}}{i(-m-t-n)} \right]_{-\pi}^{\pi} \\ &= \frac{1}{2\pi i(-m-t-n)} \left(e^{i\pi(-m-n)} e^{-it\pi} - e^{i\pi(m+n)} e^{it\pi} \right) = \frac{(-1)^{-m-n-1} \sin(\pi t)}{\pi(-m-t-n)}. \end{aligned}$$

Thus, we obtain

$$(3.9) \quad I_t = \frac{1}{\pi^2} \sum_{m \in \mathbb{N}} \sum_{n \in \mathbb{N}} \frac{|\sin(\pi t)|^2}{|(m+t+n)|^2}.$$

Hence, unless $\sin(\pi t) = 0$, this double infinite sum diverges, namely,

$$(3.10) \quad I_t = \infty \quad \forall t \in \mathbb{R} \setminus \mathbb{Z}.$$

Due to the inconsistency between (3.7) and (3.10), the existence of the inner automorphism $\text{Ad}(V_t)$ is negated. Therefore, the assumed $V_t \in \mathfrak{A}^F$ actually does not exist for all non-integer real numbers; of course, integers $t \in \mathbb{Z}$ are excluded due to the existence of the discrete shift-translation automorphism group $\{\tau_i^S \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$. We conclude that the C^* -flow $\{\tau_t^S \in \text{Aut}(\mathfrak{A}^S), t \in \mathbb{R}\}$ satisfying the assumption does not exist. \square

4. ASYMPTOTICALLY ABELIAN CONDITION

Proposition 1 (together with Remark 3) demonstrates that the locality and causality (one-sided direction) satisfied by the discrete shift-translation automorphism group of \mathfrak{A}^F are lost by its continuous extension. We ask what dynamical properties are shared by both the discrete shift-translation automorphism group and its continuous extensions. We will show that the asymptotically abelian condition serves as an example of such shared properties.

The asymptotically abelian condition is a general characterization of chaotic quantum dynamics that ensures delocalization [5]. It implies that every local operator will escape from its original position, and never come back there after some time.

We shall recall the precise definition of the asymptotically abelian condition [7] in the following. First, we give the definition for the case \mathfrak{A}^S . Consider a C^* -flow $\{\alpha_t \in \text{Aut}(\mathfrak{A}^S), t \in \mathbb{R}\}$ on \mathfrak{A}^S . We say that it satisfies the (norm) asymptotically abelian condition, if the following asymptotic

formula in C^* -norm holds:

$$(4.1) \quad \lim_{|t| \rightarrow \infty} [A, \alpha_t(B)] = 0 \quad \text{for every } A, B \in \mathfrak{A}^S,$$

where the norm convergence is used. Similarly, consider a \mathbb{Z} -automorphism group $\{\tilde{\alpha}_i \in \mathfrak{A}^S, i \in \mathbb{Z}\}$. It satisfies the (norm) asymptotically abelian condition if

$$(4.2) \quad \lim_{|i| \rightarrow \infty} [A, \tilde{\alpha}_i(B)] = 0 \quad \text{for every } A, B \in \mathfrak{A}^S,$$

where the norm convergence is used. We then move to the case \mathfrak{A}^F , which has the graded structure (2.4) with the grading Θ . The graded commutator on \mathfrak{A}^F is defined by the following mixture of the commutator and the anti-commutator:

$$(4.3) \quad \begin{aligned} [A_+, B]_{\Theta} &:= [A_+, B] \quad \text{for } A_+ \in \mathfrak{A}_+, B \in \mathfrak{A}, \\ [A_-, B_-]_{\Theta} &:= \{A_-, B_-\} \quad \text{for } A_-, B_- \in \mathfrak{A}_-. \end{aligned}$$

A C^* -flow $\{\alpha_t \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ on the fermion system \mathfrak{A}^F satisfies the Θ -graded (norm) asymptotically abelian condition, if

$$(4.4) \quad \lim_{|t| \rightarrow \infty} [A, \alpha_t(B)]_{\Theta} = 0 \quad \text{for every } A, B \in \mathfrak{A}^F,$$

where the norm convergence is used. Similarly, $\{\tilde{\alpha}_i \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ satisfies the (norm) Θ -graded asymptotically abelian condition, if

$$(4.5) \quad \lim_{|i| \rightarrow \infty} [A, \tilde{\alpha}_i(B)]_{\Theta} = 0 \quad \text{for every } A, B \in \mathfrak{A}^F,$$

where the norm convergence is used. In what follows, we shortly call the asymptotically abelian condition ‘AA-condition’ and the Θ -graded asymptotically abelian condition ‘ Θ -AA-condition’.

As a typical example, the discrete shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$ on \mathfrak{A}^S satisfies AA-condition, whereas the discrete shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ on \mathfrak{A}^F satisfies Θ -AA-condition. If a C^* -flow $\{\alpha_t \in \text{Aut}(\mathfrak{A}^S)$ or $\in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ satisfies AA condition or Θ -AA-condition, then so does its restricted \mathbb{Z} -automorphism group $\{\alpha_i \in \text{Aut}(\mathfrak{A}^S)$ or $\in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$. The following proposition demonstrates that the converse implication holds in general.

Proposition 4. *Let $\{\alpha_t \in \text{Aut}(\mathfrak{A}^S), t \in \mathbb{R}\}$ be any C^* -flow on \mathfrak{A}^S that extends the discrete shift-translation automorphism group $\{\tau_i^S \in \text{Aut}(\mathfrak{A}^S), i \in \mathbb{Z}\}$. Then $\{\alpha_t \in \text{Aut}(\mathfrak{A}^S), t \in \mathbb{R}\}$ satisfies AA-condition. Let $\{\alpha_t \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ be any C^* -flow on \mathfrak{A}^F that extends the discrete shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$. Then $\{\alpha_t \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ satisfies Θ -AA-condition.*

Proof. We consider the case \mathfrak{A}^S . We introduce some convenient notation. Each $t \in \mathbb{R}_+$ is uniquely written as the sum of its non-negative integer part $t_{\text{int}} \in \mathbb{N} \cup 0$ and its decimal part $t_{\text{dec}} \in [0, 1)$. For example, we have $t_{\text{int}} = 3$

and $t_{\text{dec}} = 0.14159$ for $t = 3.14159$. Let \mathcal{K} be some subset of \mathfrak{A}^S . Let $\varepsilon > 0$ and $A \in \mathfrak{A}^S$. If $\inf_{X \in \mathcal{K}} \|A - X\| < \varepsilon$, then we write $A \in^\varepsilon \mathcal{K}$. For $I \in \mathbb{Z}$, its translation by $x \in \mathbb{Z}$ is denoted as $I + x := \{y + x \in \mathbb{Z}; y \in I\}$. For $N \in \mathbb{N} \cup \{0\}$ define the following discrete segment centered at $0 \in \mathbb{Z}$

$$[N] \equiv \{-N, -N + 1, \dots, -1, 0, 1, \dots, N - 1, N\}$$

We denote

$$\mathfrak{A}_{[N]}^S \equiv \mathfrak{A}^S([N]).$$

For $x \in \mathbb{Z}$, let

$$\mathfrak{A}_{[N]+x}^S \equiv \tau_x^S(\mathfrak{A}_{[N]}^S)$$

To show the formula (4.1) it is enough to derive

$$(4.6) \quad \lim_{t \rightarrow +\infty} [A, \alpha_t(B)] = 0 \quad \text{for each pair of } A, B \in \mathfrak{A}_o^S.$$

For any given $A, B \in \mathfrak{A}_o^S$, there is a minimum $N \in \mathbb{N}$ such that $A, B \in \mathfrak{A}_{[N]}^S$. Let $B(t) := \alpha_t(B) \in \mathfrak{A}^S$, which is an \mathfrak{A}^S -valued norm continuous function for $t \in \mathbb{R}$. We consider $B(t)$ on the closed segment $[0, 1] \equiv \{t \in \mathbb{R}; 0 \leq t \leq 1\}$. For $p \in \mathbb{N}$, take the following finite subset of \mathfrak{A}^S :

$$(4.7) \quad \mathcal{B}_p := \{B(0), B(1/p), B(2/p), \dots, B(k/p), \dots, B(1-1/p), B(1)\}, \quad (0 \leq k \leq p, k \in \mathbb{N}).$$

For any fixed $\varepsilon > 0$, there exists a sufficiently large $p \in \mathbb{N}$ such that

$$(4.8) \quad B(t) \in^\varepsilon \mathcal{B}_p \quad \text{for every } t \in [0, 1],$$

Since \mathcal{B}_p is a finite subset of \mathfrak{A}^S , there exists a sufficiently large $M (> N) \in \mathbb{N}$ such that

$$(4.9) \quad \mathcal{B}_p \in^\varepsilon \mathfrak{A}_{[M]}^S.$$

From (4.8) (4.9) we have

$$(4.10) \quad B(t) \in^{2\varepsilon} \mathfrak{A}_{[M]}^S \quad \text{for every } t \in [0, 1].$$

We will crudely estimate the location of $\alpha_t(B)$ for large $t \in \mathbb{R}$. By definition, we have

$$(4.11) \quad B(t) = B(t_{\text{int}} + t_{\text{dec}}) = \alpha_{t_{\text{int}}}(B(t_{\text{dec}})) = \tau_{t_{\text{int}}}^S(B(t_{\text{dec}})) \quad \text{for each } t (= t_{\text{int}} + t_{\text{dec}}) \in \mathbb{R}.$$

From (4.10) and (4.11) it follows that

$$(4.12) \quad B(t) \in^{2\varepsilon} \mathfrak{A}_{[M]+t_{\text{int}}}^S \quad \text{for each } t \in \mathbb{R}.$$

Hence, the intersection $[N] \cap [M] + t_{\text{int}}$ will eventually become empty as $t \rightarrow +\infty$ ($t_{\text{int}} \rightarrow +\infty$). Thus, due to the local commutativity of \mathfrak{A}^S , we have shown (4.6).

The proof for (4.6) provided above demonstrates that the location of $\alpha_t(B)$ is near to the shift-translation $\tau_{t_{\text{int}}}^S(B)$ and becomes far away from

the original location of the local operator B as the time $t \in \mathbb{R}$ proceeds. This characteristic also holds true in the case of \mathfrak{A}^F . Thus, we can show

$$(4.13) \quad \lim_{t \rightarrow \infty} [A, \alpha_t(B)]_{\Theta} = 0 \quad \text{for each pair of } A, B \in \mathfrak{A}_0^F,$$

repeating the same argument for (4.6). We need not consider distinctions arising from the graded structure of the fermion system \mathfrak{A}^F . \square

Remark 5. This is a side remark. The Rohlin property [15] is another property which becomes invalid by the continuous extension of discrete shift translations. The shift-translation automorphism group $\{\tau_i^F \in \text{Aut}(\mathfrak{A}^F), i \in \mathbb{Z}\}$ is known to have the Rohlin property [4], whereas its quasi-free extension $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ fails to satisfy it because of the existence of KMS states Corollary 2.

5. DISCUSSION

We specified the formal Hamiltonian \tilde{H}_{∞} for the shift-translation quasi-free C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ given in Proposition 1. It consists of the two-body translation invariant interactions with $\frac{1}{r}$ -decay. Let us recapitulate its formula given in (2.33): $\tilde{H}_{\infty} = \sum_{n,m \in \mathbb{Z}} \frac{-2\pi i (-1)^{n-m}}{n-m} c_n^* c_m$.

We note that its derivation is not rigorous; the meaning of \tilde{H}_{∞} in C^* -theory [5] remains unclear. More precisely, we do not know whether and how it is associated to a pre-generator of the C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$. Although the quasi-free C^* -flow $\{\tau_t^F \in \text{Aut}(\mathfrak{A}^F), t \in \mathbb{R}\}$ has KMS states as shown in Corollary 2, the one-site energy, which is required for the description of equilibrium states (attaining the minimum free-energy), seems not obvious. At least, its existence does not follow from the general theory of fermion lattice systems [3] [5]. We conjecture that the alternative \pm sign of each creation-annihilation pair interaction may provide a clue for the cancellation of long-range effects, yielding a well-defined C^* -flow on the infinite fermion lattice system. (There are some long-range quantum lattice models which do not have C^* -flows, see [27] [6]. However, these are mean-field models, and therefore have no relevance to our quasi-free fermion lattice model.)

The above quasi-free C^* -flow on \mathfrak{A}^F is a special example of continuous extensions of discrete shift translations. Proposition 3 merely excludes one specific extension method from \mathfrak{A}^F to \mathfrak{A}^S ; it is not a thorough no-go statement. There may be other continuous extensions of discrete shift translations on \mathfrak{A}^F and \mathfrak{A}^S . We shall raise the following questions.

Question A: Is there another (non-quasi-free) C^* -flow on \mathfrak{A}^F that extends the discrete shift translations?

Question B: Is there a C^* -flow on \mathfrak{A}^S that extends the discrete shift translations?

Question C: Is there a C^* -flow on \mathfrak{A}^S satisfying AA condition?

If a positive example for **Question A** exists on \mathfrak{A}^F , then we will argue whether its extension to \mathfrak{A}^S is possible or not. Of course, by Proposition 4, if a positive example of **Question B** is found, then it gives a positive example for **Question C** as well.

ACKNOWLEDGMENTS

HM would like to thank Jean-Bernard Bru, Toshihiko Masuda and Itaru Sasaki for correspondence. HM acknowledges Kakenhi (grant no. 21K03290) for the financial support.

REFERENCES

- [1] Araki, H., 1970: On quasifree states of CAR and Bogoliubov automorphisms. *Publ. Res. Inst. Math. Sci.* **6**, 385-442.
- [2] Araki, H., 1984: On the XY-model on two-sided infinite chain. *Publ. Res. Inst. Math. Sci.* **20**, 277-296.
- [3] Araki, H., and Moriya, H., 2003: Equilibrium statistical mechanics of Fermion lattice systems. *Rev. Math. Phys.* **15**, 93-198.
- [4] Bratteli, O., Kishimoto, A., Rørdam, M., and Størmer, E., 1993: The crossed product of a UHF algebra by a shift. *Ergodic Theory and Dynamical Systems.* **13**, 615-626.
- [5] Bratteli, O., and Robinson, D. W., 1997: *Operator Algebras and Quantum Statistical Mechanics 2. 2nd Edition.* Springer-Verlag, 517pages.
- [6] Bru J.-B., and de Siqueira Pedra W., 2020: Quantum dynamics generated by long-range interactions for lattice fermions and quantum spins. *J. Math. Anal. Appl.* **493** 124517.
- [7] Doplicher, S., Kadison, R. V., Kastler, D., and Robinson, D. W., 1967: Asymptotically Abelian systems. *Commun. Math. Phys.* **6**, 101-120.
- [8] Else, D. V., Machado, F., Nayak, C., and Yao, N. Y., 2020: An improved Lieb-Robinson bound for many-body Hamiltonians with power-law interactions. *Phys. Rev. A.* **101** 022333.
- [9] Evans, D. E., and Lewis, J. T., 1986: On a C^* -algebra approach to phase transition in the two-dimensional Ising Model. II. *Commun. Math. Phys.* **102**, 521-535.
- [10] Farrelly, T., 2020: A review of Quantum Cellular Automata. *Quantum* **4**, 368.
- [11] Gong, Z., Guaita, T., and J. I. Cirac., 2023: Long-range free fermions: Lieb-Robinson bound, clustering properties, and topological phases. *Phys. Rev. Lett.* **130** (2023).
- [12] Gross, D., Nesme, V., Vogts, H., and Werner, R. F., 2012: Index theory of one dimensional quantum walks and cellular automata. *Commun. Math. Phys.* **310**, 419-454.
- [13] Jones, V. F. R., 2018: A no-go theorem for the continuum limit of a periodic quantum spin chain. *Commun. Math. Phys.* **357**, 295-317.
- [14] Kato, K., Kishimoto, D., and Mitsunobu, T. 2023: Homotopy type of the space of finite propagation unitary operators on \mathbb{Z} . *Homology, Homotopy and Applications.* **25**, 375-400.
- [15] Kishimoto, A., 1996: A Rohlin property for one-parameter automorphism groups. *Commun. Math. Phys.* **179**, 599-622
- [16] Kuiper, N. H., 1965: The homotopy type of the unitary group of Hilbert space. *Topology.* **3**, 19-30.
- [17] Kuwahara, T., and Saito, K., 2020: Strictly linear light cones in long-range interacting systems of arbitrary dimensions. *Phys. Rev. X.* **10** 031010.
- [18] Lieb, E., and Robinson, D. W., 1972: The finite group velocity of quantum spin systems. *Commun. Math. Phys.* **28** 313-321.
- [19] Matsui, T., and Ogata, Y., 2003: Variational principle for non-equilibrium steady states of the XX Model. *Rev. Math. Phys.* **15**, 905-923.
- [20] Matsuta, T., Koma, T., Nakamura, S., 2017: Improving the Lieb-Robinson bound for long-range interactions. *Ann. Henri. Poincaré* **18**, 519-528.
- [21] Moriya, H., 2024: Non-existence of spontaneous symmetry breakdown of time-translation symmetry on general quantum systems: Any macroscopic order parameter moves not! *Rev. Math. Phys.* **36**, 2330008.
- [22] Popa, S., Takesaki, M., 1993: The topological structure of the unitary and automorphism groups of a factor. *Commun. Math. Phys.* **155**, 93-101.

- [23] Pusz, W., and Woronowicz, S. L., 1978: Passive states and KMS states for general quantum systems, *Commun. Math. Phys.* **58**, 273-290.
- [24] Ranard, D., Walter, M., and Witteveen, F., 2022: A converse to Lieb-Robinson bounds in one dimension using Index theory. *Ann. Henri. Poincaré* **23**, 3905-3979.
- [25] Rocca, F., Sirugue, M., and Testard, D., 1969: On a class of equilibrium states under the Kubo-Martin-Schwinger boundary condition. I. Fermions *Commun. Math. Phys.* **13**, 317-334.
- [26] Stone, M. H., 1932: On one-parameter unitary groups in Hilbert Space. *Ann. of Math.* **33(3)** 643-648.
- [27] Thirring, W., and Wehrl A., 1967: On the mathematical structure of the B.C.S.-model. *Commun. Math. Phys.* **4**, 303-314.
- [28] Wilming, H., and Werner, A. H., 2022: Lieb-Robinson bounds imply locality of interactions. *Phys. Rev. B.* **105**, 125101.
- [29] Yin, C., Lucas, A., and Stephen, D. T., 2024: Frobenius light cone and the shift unitary. arXiv:2402.07990
- [30] Zimborás, Z., Farrelly, T., Farkas, S., Masanes, L., 2022: Does causal dynamics imply local interactions? *Quantum* **6**, 748.

FACULTY OF MECHANICAL ENGINEERING, INSTITUTE OF SCIENCE AND ENGINEERING, KANAZAWA UNIVERSITY, KAKUMA, KANAZAWA 920-1192, JAPAN

Email address: hmoriya4@se.kanazawa-u.ac.jp

FACULTY OF PHYSICS, UNIVERSITY OF VIENNA, BOLTZMANNGASSE 5, A-1090 VIENNA, AUSTRIA

Email address: heide.narnhofer@univie.ac.at