MATRIX INVERTIBLE EXTENSIONS OVER COMMUTATIVE RINGS. PART III: HERMITE RINGS

GRIGORE CĂLUGĂREANU, HORIA F. POP, ADRIAN VASIU

ABSTRACT. We reobtain and often refine prior criteria due to Kaplansky, Mc-Govern, Roitman, Shchedryk, Wiegand, and Zabavsky–Bilavska and obtain new criteria for a Hermite ring to be an EDR. We mention three criteria: (1) a Hermite ring R is an EDR iff for all pairs $(a, c) \in R^2$, the product homomorphism $U(R/Rac) \times U(R/Rc(1-a)) \rightarrow U(R/Rc)$ between groups of units is surjective; (2) a reduced Hermite ring is an EDR iff it is a pre-Schreier ring and for each $a \in R$, every zero determinant unimodular 2×2 matrix with entries in R/Ra lifts to a zero determinant matrix with entries in R; (3) a Bézout domain R is an EDD iff for all triples $(a, b, c) \in R^3$ there exists a unimodular pair $(e, f) \in R^2$ such that (a, e) and (be + af, 1 - a - bc) are unimodular pairs. We use these criteria to show that each Bézout ring R that is an $(SU)_2$ ring (as introduced by Lorenzini) such that for each nonzero $a \in R$ there exists no nontrivial self-dual projective R/Ra-module of rank 1 generated by 2 elements (e.g., all its elements are squares), is an EDR.

1. INTRODUCTION

Let R be a commutative ring with identity. Let U(R), N(R), J(R), Z(R) and Pic(R) be its group of units, its nilpotent radical, its Jacobson radical, its set of zero divisors, and its Picard group (respectively). For $a \in R$, let $Ann_R(a)$ be its annihilator in R. For $n \in \mathbb{N} = \{1, 2, ...\}$, let $\mathbb{M}_n(R)$ be the R-algebra of $n \times n$ matrices with entries in R. Let $GL_n(R)$ be the general linear group of units of $\mathbb{M}_n(R)$. Let $SL_n(R) := \{M \in GL_n(R) | \det(M) = 1\}$ be the special linear subgroup of $GL_n(R)$. Let Um(F) be the set of unimodular elements of a free R-module F. We say that $A, B \in \mathbb{M}_2(R)$ are congruent modulo an ideal I of R if $A - B \in \mathbb{M}_2(I)$.

In this paper we study Bézout rings, i.e., rings whose finitely generated ideals are principal. Each Bézout ring is an arithmetical ring, i.e., its lattice of ideals is distributive and all its localizations at prime ideals are valuation rings (see [10], Thms. 1 and 2). The ring R is a Bézout ring iff each diagonal matrix with entries in R admits diagonal reduction (see [11], Thm. (3.1)). Each Hermite ring R in the sense of Kaplansky, i.e., defined by the identity $R^2 = RUm(R^2)$, is a Bézout ring but the converse does not hold (see [6], Ex. 3.4, [20], Ex. 3.3, or [2], Prop. 8). However, a Hermite domain is the same as a Bézout domain. Bézout domains are GCD (greatest common divisors exist) domains and hence pre-Schreier domains with trivial Picard groups. As R is a Hermite ring iff each 1×2 matrix with entries in R admits diagonal reduction, each EDR is a Hermite ring. We will use the tools

²⁰²⁰ Mathematics Subject Classification. Primary: 15A83, 13G05, 19B10. Secondary: 13A05, 13F05, 13F25, 15B33, 16U10.

Key words and phrases. ring, matrix, projective module, unimodular.

of Parts I and II in order to obtain necessary and sufficient criteria for a Hermite ring (or domain) to be an *EDR* (or an *EDD*, i.e., an elementary divisor domain).

Let $A \in Um(\mathbb{M}_2(R))$. Recall that A is called extendable if it is obtained from a matrix $A^+ \in SL_3(R)$ by removing its third row and its third column (see [3], Def. 1.1). If we can choose A^+ such that its (3,3) entry is 0, then A is called simply extendable. Recall that A is called determinant liftable (resp. weakly determinant liftable) if there exists $B \in Um(\mathbb{M}_2(R))$ (resp. $B \in \mathbb{M}_2(R)$) congruent to A modulo $R \det(A)$ and with $\det(B) = 0$ (see [3], Def. 1.1). The *only* general implications between these 4 notions on A are the ones in the diagram

(I) simply extendable
$$\implies$$
 extendable \Downarrow determinant liftable \implies weakly determinant liftable

(see [4], Thm. 1.3 for the vertical ones; for the fact that their converses do not hold and that there exist no implications between extendable and determinant liftable see [3], Exs. 5.3 and 6.2 and [4], paragraph after Thm. 1.3 and Ex. 1.9). Recall that a matrix $B \in \mathbb{M}_2(R)$ is called non-full if it is the product of a 2 × 1 and a 1 × 2 matrix with entries in R.

Recall from [3], Def. 1.2 that R is called a Π_2 ring if each matrix in $Um(\mathfrak{M}_2(R))$ of zero determinant is extendable, equivalently it is simply extendable by [3], Lem. 4.1(1) and that R is called an E_2 (resp. an SE_2) ring if each matrix in $Um(\mathfrak{M}_2(R))$ is extendable (resp. simply extendable). If Pic(R) is trivial (say, R is a Bézout domain), then R is a Π_2 ring (see [3], paragraph after Thm. 1.4).

See [3], Def. 1.5 for stable ranges notation. If R is a Hermite ring, then $sr(R) \leq 2$ (see [14], Prop. 8(i) or [21], Cor. 2.1.1). In fact, a Bézout ring R is a Hermite ring iff $sr(R) \leq 2$ (see [21], Thm. 2.1.2). Bézout domains that have stable range 1 or 1.5 were also studied in [16] and respectively [18], [19] and [1]. For Hermite rings, Parts I and II get summarized in an example and a theorem as follows.

Example 1.1. For Hermite rings R, simply extendable and extendable properties on a matrix in $Um(\mathbb{M}_2(R))$ are equivalent (see [3], Thm. 1.6). For Hermite rings Rthat are Π_2 rings, simply extendable, extendable and determinant liftable properties on a matrix in $Um(\mathbb{M}_2(R))$ are equivalent (see [4], Cor. 1.5). For Hermite rings Rsuch that zero determinant matrices in $Um(\mathbb{M}_2(R))$ are non-full, the 4 notions on a matrix in $Um(\mathbb{M}_2(R))$ are equivalent (see [4], Cor. 1.8).

Theorem 1.2. For a Hermite ring R the following statements are equivalent:

(1) The ring R is an EDR.

(2) The ring R is an SE_2 ring.

(3) The ring R is an E_2 ring.

(4) For each $a \in R$, R/Ra is a Π_2 ring (equivalently, every projective R/Ra-module of rank 1 generated by 2 elements is free).

(5) Each matrix in $Um(\mathbb{M}_2(R))$ is determinant liftable and R is a Π_2 ring.

If all zero determinant matrices in $\mathbb{M}_2(R)$ are non-full, then these five statements are also equivalent to:

(6) Each matrix in $Um(\mathbb{M}_2(R))$ is weakly determinant liftable.

See [3], Cor. 1.8 for $(1) \Leftrightarrow (2) \Leftrightarrow (3)$, see [3], Cors. 2.5 and 4.2 for $(3) \Leftrightarrow (4)$, see [3], Thm. 1.4 for the equivalence of (4), and see [4], Thm. 1.4 for (2) \Leftrightarrow (5). See [4], Thm. 1.7 for (3) \Leftrightarrow (6). The equivalence (1) \Leftrightarrow (4) can be viewed as a more practical way to check that a Hermite ring is an *EDR* than [20], Thm. 2.1 (which does not restrict to only 2 generators) or [23], Thms. 3 and 5 (which work with finitely generated projective modules over all quotients of R).

Recall that R is a Hermite ring iff for all $p, q \in \mathbb{N}$ and each $p \times q$ matrix B with entries in R, there exist $M \in GL_p(R)$ and $N \in GL_q(R)$ such that MB and BN are both lower (equivalently, upper) triangular (see [7], Thm. 3). Thus, in Theorem 1.2(4) or (5) it suffices to consider only upper triangular matrices, and it is part of the goals of this paper to show that in fact one can restrict to specific upper triangular matrices (e.g., see Propositions 2.2 and 2.3).

Example 1.3. For a ring R (resp. a ring R with N(R) = 0) and $(a, b, c) \in Um(R^3)$, $A := \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \in Um(\mathbb{M}_2(R))$ is determinant (resp. weakly determinant) liftable iff there exists $(x, y, z, w) \in R^4$ such that ax + by + cw = 1 and xw = yz (resp. such that 1 - ax - by - cw + ac(xw - yz) = 0), see [4], Thm. 1.2 (resp. Thm. 5.1(2) and (3)). Note that 1 - ax - by - cw + ac(xw - yz) = (1 - ax)(1 - cw) - (b + acz)y (cf. [4], Eq. (II)), hence A is weakly determinant liftable iff there exists $(x, z, w) \in R^3$ such that b + acz divides (1 - ax)(1 - cw).

Based on Example 1.3 and $(1) \Leftrightarrow (5)$ of Theorem 1.2 it follows directly:

Corollary 1.4. Assume R is a Hermite ring and a Π_2 ring. Then R is an EDR iff for each $(a, b, c) \in Um(R^3)$ there exists $(x, y, z, w) \in R^4$ such that ax + by + cw = 1and xw = yz (i.e., there exists $(x, y, w) \in R^3$ such that ax + by + cw = 1 and $xw \in Ry$).

For Hermite rings, to get necessary and/or sufficient conditions to be *EDRs* that involve units, one would like to get unit interpretations that would characterize the inequality $sr(R) \leq 2$ in some way similar to [3], Prop. 2.4. As a progress in this direction we introduce using units the following class of rings.

Definition 1.5. We say that R is a U_2 ring if for each $(a, b) \in Um(R^2)$ and $c \in R$, the natural product homomorphism

$$\psi_{a,b,c}: U(R/Rac) \times U(R/Rbc) \rightarrow U(R/Rc)$$

is surjective, i.e., the functorial commutative diagram

(II)
$$U(R/(Rac \cap Rbc)) \longrightarrow U(R/Rac)$$

$$\downarrow \qquad \qquad \downarrow$$

$$U(R/Rbc) \longrightarrow U(R/Rc)$$

whose arrows are natural reductions, is not only a pullback but is also a pushout.

For $(a, b) \in Um(R^2)$, if $(a', b') \in R^2$ is such that aa' + bb' = 1, then $v_{(aa', 1-aa', c)}$ factors through $v_{a,b,c}$. Hence R is a U_2 ring iff $v_{a,1-a,c}$ is surjective for all $(a, c) \in R^2$. The fact that a unit in U(R/Rc) does or does not belong to $Im(v_{a,1-a,c})$ relates to a certain zero determinant matrix being or not being non-full (see Example 3.1).

Example 1.6. Assume asr(R) = 1, i.e., R has almost stable range 1. We show that R is a U_2 ring. Let $(a, b) \in Um(R^2)$ and $c \in R$. If $c \in J(R)$, then U(R) surjects

onto U(R/Rc). If $c \notin J(R)$, then, as c is a linear combination of ac and bc, either $ac \notin J(R)$ or $bc \notin J(R)$. Hence either $U(R/Rac) \to U(R/(Rac + Rc)) = U(R/Rc)$ or $U(R/Rbc) \to U(R/(Rbc + Rc)) = U(R/Rc)$ is a surjective homomorphisms (see [3], Prop. 2.4(3)). We incur that $v_{a,b,c}$ is surjective.

Recall that a ring R is called pre-Schreier if each $x \in R$ is primal, i.e., if x divides yz, with $(y, z) \in R^2$, then there exists $(u, v) \in R^2$ such that x = uv, u divides y and v divides z. The next two examples underline the relevance of this class of rings.

Example 1.7. Let R be such that each zero determinant matrix $B \in M_2(R)$ is non-full. We show that R is a pre-Schreier ring. Let $(x, y, z) \in R^3$ be such that x divides yz. Let $w \in R$ be such that xw = yz. Let $C := \begin{bmatrix} x & y \\ z & w \end{bmatrix} \in M_2(R)$. As $\det(C) = 0$, there exists $(l, m, o, q) \in R^4$ such that $C = \begin{bmatrix} l \\ m \end{bmatrix} \begin{bmatrix} o & q \end{bmatrix}$. Hence x = lo with l dividing y = lq and o dividing z = mo. Thus R is a pre-Schreier ring. **Example 1.8.** Assume N(R) = 0. Let $B \in M_2(R)$ with $\det(B) = 0$. We check that if B admits diagonal reduction, then B is non-full. Let $M, N \in GL_2(R)$ be such that $MBN = \begin{bmatrix} a & 0 \\ 0 & ab \end{bmatrix}$ with $(a, b) \in R^2$. As $\det(BN) = \det(M) \det(B) \det(N) = 0 = a^2b$, it follows that $(ab)^2 = 0$ and thus, as N(R) = 0, we have ab = 0. So, $B = M^{-1} \begin{bmatrix} a & 0 \\ 0 & 0 \end{bmatrix} N^{-1} = M^{-1} \begin{bmatrix} a \\ 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \end{bmatrix} N^{-1}$ is non-full.

From Examples 1.7 and 1.8 we get directly:

Corollary 1.9. If R is an EDR with N(R) = 0, then R is a pre-Schreier ring.

Section 3 proves the following theorem.

Theorem 1.10. We assume that one of the following conditions holds:

(1) for each $(a, b, c) \in Um(R^3)$ there exists $(x, y, z, w) \in R^4$ such that 1 - ax - by - cw + ac(xw - yz) = 0 and R is a pre-Schreier ring;

(2) each upper triangular matrix in $Um(\mathbb{M}_2(R))$ is simply extendable and R is either a Π_2 ring or an integral domain (e.g., R is an SE₂ ring).

Then R is a U_2 ring.

Example 1.11. Assume each upper triangular matrix in $Um(\mathbb{M}_2(R))$ is weakly determinant liftable (e.g., this holds if R is a $WJ_{2,1}$ ring in the sense of [4], Def. 1.10(1), see [4], Thm. 1.11(1)). If R is also a pre-Schreier ring with N(R) = 0, then condition (1) of Theorem 1.10 holds by Example 1.3, hence R is a U_2 ring.

Each Dedekind domain is a U_2 ring by Example 1.6. Hence Dedekind domains which are not *PIDs* are U_2 rings which are not Π_2 rings (see [3], Thm. 1.7(4)). In Section 4 we prove the following 'units supplement' to Theorem 1.2.

Theorem 1.12. For a Hermite ring R the following statements are equivalent:

(1) The ring R is an EDR.

(2) The ring R/N(R) is a pre-Schreier ring and each matrix in $Um(\mathbb{M}_2(R))$ is weakly determinant liftable.

(3) The ring R is a U_2 ring.

(4) Given two unimodular pairs $(a, b), (c, d) \in Um(R^2)$, there exists $t \in R$ such that we can factor $d + ct = d_1d_2$ with $(a, d_1), (b, d_2) \in Um(R^2)$.

(5) Given $(a, d) \in \mathbb{R}^2$ and $c \in 1 + \mathbb{R}d$, there exists $t \in \mathbb{R}$ such that we can factor $d + ct = d_1d_2$ with $(a, d_1), (1 - a, d_2) \in Um(\mathbb{R}^2)$.

Example 1.13. Let R be a Hermite ring which is not an EDR (see [6], Sect. 4 and Ex. 4.11 and [2], Prop. 8). From Theorems 1.2 and 1.12 it follows that R is neither an E_2 ring nor a U_2 ring and there exists $a \in R$ such that the Hermite ring R/Ra is not a Π_2 ring.

Theorem 1.12 implies directly the following result.¹

Corollary 1.14. Assume R is a Hermite ring and a pre-Schreier ring with N(R) = 0. Then R is an EDR iff each matrix in $Um(\mathbb{M}_2(R))$ is weakly determinant liftable and iff for each $(a, b, c) \in Um(R^3)$ there exists $(x, y, z, w) \in R^4$ such that

(III)
$$(1 - ax)(1 - cw) = (b + acz)y.$$

If asr(R) = 1, then R is an U_2 ring (see Example 1.6), thus McGovern's theorem reproved as [3], Cor. 1.9(2) follows from the equivalence (1) \Leftrightarrow (3) of Theorem 1.12. In Section 5 we prove the following ' Π_2 supplement' to Theorem 1.2.

Theorem 1.15. Assume R is a Hermite ring with N(R) = 0. Then R is an EDR iff for each $a \in R$ the quotient ring $R/\operatorname{Ann}_R(a)$ is a Π_2 ring and each matrix in $Um(\mathbb{M}_2(R))$ is weakly determinant liftable (equivalently, and for every $(a, b, c) \in$ $Um(R^3)$ there exists $(x, y, z, w) \in R^4$ such that Equation (III) holds).

Lorenzini introduced 3 classes of rings that are 'between' Hermite rings and EDRs (see [12], Prop. 4.11). For the first class, called $J_{2,1}$ (see [12], Def. 4.6), it was proved in [4], Thm. 1.11(2) that each $J_{2,1}$ ring which is a Π_2 ring is an EDR. We recall the last 2 classes of rings (see [12], Def. 2.1) in a slightly different way.

Definition 1.16. For $n \ge 2$ we say that R is:

(1) a $(WSU')_n$ (resp. $(WSU)_n$) ring if for each $A \in Um(\mathbb{M}_n(R))$ there exists $N \in SL_n(R)$ (resp. $N \in GL_n(R)$) such that AN is symmetric;

(2) an $(SU')_n$ (resp. $(SU)_n$) ring if it a Hermite ring and a $(WSU')_n$ (resp. $(WSU)_n$) ring.

In Definition 2.1(1), the existence of N is equivalent to the existence of $M \in SL_n(R)$ (resp. $M \in GL_n(R)$) such that MA is symmetric and equivalent to the existence of $M, N \in SL_n(R)$ (resp. $M, N \in GL_n(R)$) such that MAN is symmetric. This follows via conjugation from the fact that for each symmetric matrix $O \in \mathbb{M}_n(R)$ and every $M \in GL_n(R)$, MOM^T is symmetric.

Each $(WSU')_n$ ring is a $(WSU)_n$ ring. A $(WSU)_2$ ring is an E_2 (resp. SE_2 ring) iff each symmetric matrix in $Um(\mathbb{M}_2(R))$ is extendable (resp. simply extendable). Section 6 proves basic properties of $(WSU')_n$ and $(WSU)_n$ rings; e.g., our definition of $(SU')_n$ or $(SU)_n$ rings is equivalent to the one in [12] by Proposition 6.1.

In Section 7 we prove the following theorem.

¹Similar to Corollary 1.4, Corollary 1.14 can be also deduced from Example 1.3 and Theorem 1.2 via the result proved in [8] that a Bézout ring with N(S) = 0 is pre-Schreier iff all zero determinant matrices in $\mathbb{M}_2(S)$ are non-full.

Theorem 1.17. Let R be a $(WSU)_2$ ring. Then the following properties hold:

(1) For each $(a,b) \in Um(R^2)$ and $c \in R$, $Coker(v_{a,b,c})$ is a Boolean group, i.e., we have an inclusion $\{x^2 | x \in U(R/Rc)\} \subset Im(v_{a,b,c})$.

(2) If $sr(R) \leq 4$, then for each $d \in R$, every projective R/Rd-module of rank 1 generated by 2 elements is self-dual.

(3) Assume that each element of R is the square of an element of R (e.g., this holds if R is an integrally closed domain with an algebraically closed field of fractions or is a perfect ring of characteristic 2). Then R is a U_2 ring. If moreover R is a Hermite ring, then R is an EDR.

In Section 8 we prove the following theorem.

Theorem 1.18. Let R be a Hermite ring with the property that for each $(a, b) \in Um(R^2)$ and $c \in R$, $Coker(v_{a,b,c})$ is a Boolean group. Then for each $d \in R$, every projective R/Rd-module of rank 1 generated by 2 elements is self-dual.

To connect with Pell-type equations and provide more examples of $(WSU)_2$ rings that are *EDRs*, we first prove in Section 9 the following non-full Pell-type criterion.

Criterion 1.19. Let $A = \begin{bmatrix} a & b \\ b & c \end{bmatrix} \in Um(\mathbb{M}_2(R))$ be symmetric and with zero determinant. Then the following properties hold:

(1) The matrix A is simply extendable if there exists $(e, f) \in \mathbb{R}^2$ such that $ae^2 - cf^2 \in U(\mathbb{R})$, and the converse holds if R has characteristic 2.

(2) Assume that R is a Hermite ring of characteristic 2 with N(R) = 0. If b is not a zero divisor, then A is simply extendable.

By combining Theorem 1.17(1) with Criterion 1.19, we obtain the following Pelltype criterion proved in Section 9.

Criterion 1.20. Let R be a $(SU)_2$ ring. Then R is an EDR if for all $(a, b, c) \in Um(R^3)$ there exists $(e, f) \in R^2$ such that $(ae^2 - cf^2, ac - b^2) \in Um(R^2)$, and the converse holds if R has characteristic 2.

We recall that, if the characteristic of R if a prime, then its perfection R_{perf} is the inductive limit of the inductive system indexed by $n \in \mathbb{N}$ whose all transition homomorphisms are the Frobenius endomorphism of R. If R is Hermite (or a Bézout) ring, then so is $R_{perf} \cong (R/N(R))_{perf}$.

Example 1.21. Assume R has characteristic 2. If for all $(a, c) \in R^2$, $Coker(v_{a,1-a,c})$ is a Boolean group, then R_{perf} is a U_2 ring. From this and Theorem 1.17(3) it follows that perfections of $(WSU)_2$ rings of characteristic 2 are U_2 rings. Thus perfections of $(SU)_2$ rings of characteristic 2 are EDRs by Theorem 1.12.

As an application of Corollary 1.14 in Section 10 we prove:

Criterion 1.22. For a Bézout domain R the following statements are equivalent:

(1) The ring R is an EDD.

(2) For each $(a, b, s) \in \mathbb{R}^3$, there exists $(q, r) \in \mathbb{R}^2$ such that by defining y := r+s-asq-bqr and t := 1+q-aq-br we have $t \in \mathbb{R}y+\mathbb{R}at$ (equivalently, there exists a product decomposition $y = y_1y_2$ such that y_1 divides t and $(a, y_2) \in Um(\mathbb{R}^2)$).

(3) For each $(a, b, s) \in \mathbb{R}^3$ there exists $(e, f) \in Um(\mathbb{R}^2)$ such that (a, e), $(be + af, 1 - bs - a) \in Um(\mathbb{R}^2)$.

E.g., statement (3) holds if $(a, s) \in Um(R^2)$ as we can take (e, f) := (s, 1), if $(1-a, b) \in Um(R^2)$ as we can take (e, f) := (1, 0), or if there exists $q \in R$ such that $(b+aq, 1-bs-a) \in Um(R^2)$ as we can take (e, f) := (1-a, q+b). So, each Bézout domain R with the property that for all $(a, b, s) \in R^3$ with $(a, s), (b, 1-a) \notin Um(R^2)$ there exists $q \in R$ such that $(b+aq, 1-bs-a) \in Um(R^2)$, is an EDD.

The implicit and explicit questions raised in the literature, such as, "Is a Bézout domain of finite Krull dimension [at least 2] an *EDD*?" (see [5], Ch. III, Probl. 5, p. 122), and, 'What classes of Bézout domains which are not *EDDs* exist?', remain unanswered. However, the above results reobtain or can be easily used to reobtain multiple other criteria existing in the literature of when a Hermite ring is an *EDR*. The equivalence $(1) \Leftrightarrow (4)$ of Theorem 1.12 was proved for Bézout domains in [17], Thm. 5 and a variant of it was proved for Hermite rings in [15], Prop. 2.9. Equation (III) generalizes and refines the equation one would get based on [15], Rm. 2.8. The last two references were reinterpreted in terms of neat stable range (see [22], Def. 21) 1 in [22], Thms. 31 and 33; see [13] for clean and neat rings.

2. Test matrices

Definition 2.1. Let $A \in Um(\mathbb{M}^2(R))$. An upper triangular matrix $B \in Um(\mathbb{M}^2(R))$ will be called a companion test matrix for A if there exists $(a, b, c, a', c') \in R^5$ such that A is equivalent to $\begin{bmatrix} a & b \\ 0 & c \end{bmatrix}$ and $B = \begin{bmatrix} aa' & b \\ 0 & cc' \end{bmatrix}$.

Let \mathcal{P} be one of the 4 notions extendable, simply extendable, determinant liftable and weakly determinant liftable. Definition 2.1 is justified by the next proposition.

Proposition 2.2. If \mathcal{P} is weakly determinant liftable we assume that N(R) = 0. If $A \in Um(\mathbb{M}^2(R))$ has a companion test matrix B which is \mathcal{P} , then A is \mathcal{P} .

Proof. As \mathcal{P} property depends only on equivalence classes (see [3], Lem. 4.3(1) for the case of extendable and simply extendable notions), it suffices to show that if $B = \begin{bmatrix} aa' & b \\ 0 & cc' \end{bmatrix}$ is \mathcal{P} , then so is $C = \begin{bmatrix} a & b \\ 0 & c \end{bmatrix}$. First we assume that \mathcal{P} is extendable (resp. simply extendable). From [3], Cor.

First we assume that \mathcal{P} is extendable (resp. simply extendable). From [3], Cor. 4.7 (resp. [3], Thm. 4.3) it follows that there exists $(e,g) \in \mathbb{R}^2$ (resp. $(e,g) \in Um(\mathbb{R}^2)$) such that $(aa'e, be + cc'g, aa'cc') \in Um(\mathbb{R}^3)$ (resp. $(aa'e, be + cc'g) \in Um(\mathbb{R}^2)$). Denoting $f := c'g \in \mathbb{R}$, as $(aa'e, be + cf, aa'cc') \in Um(\mathbb{R}^3)$ (resp. $(aa'e, be + cf) \in Um(\mathbb{R}^2)$), it follows that $(ae, be + cf, ac) \in Um(\mathbb{R}^3)$ (resp. $(ae, be + cf) \in Um(\mathbb{R}^2)$). Thus C is \mathcal{P} by [3], Cor. 4.7 (resp. [3], Thm. 4.3).

Next we assume that \mathcal{P} is determinant liftable (resp. weakly determinant liftable). From [4], Thm. 1.2 (resp. [4], Thm. 5.1(2) and 3) it follows that there exists $(x, y, z, w) \in \mathbb{R}^4$ such that 1 - aa'x - by - cc'w = 0 and xw = yz (resp. such that 1 - aa'x - by - cc'w = 0 and xw = yz (resp. such that 1 - aa'x - by - cc'w + aa'cc'(xw - yz) = 0). If $(x', y', z', w') := (a'x, y, a'c'z, c'w) \in \mathbb{R}^4$, then 1 - ax' - by' - cw' = 0 and x'w' = y'z' (resp. 1 - ax' - by' - cw' + ac(x'w' - y'z') = 0), so C is \mathcal{P} by [4], Thm. 1.2 (resp. Thm. 5.1(2) and 3).

Until Proposition 2.3 below we assume that R is a Hermite ring, i.e., for every pair $(p,q) \in R^2$ there exists $(r,s,t) \in R^3$ such that p = rs, q = rt and $(s,t) \in Um(R^2)$. If moreover R is an integral domain (i.e., if R is a Bézout domain), then r is unique up to a multiplication with a unit of R and is called the greatest common divisor of p and q and one writes $r = \gcd(x, y)$. Our convention is $\gcd(0, 0) = 0$ (so that we can still write $0 = 0 \cdot 1$ with $\gcd(1, 1) = 1$).

Let $A \in Um(\mathbb{M}_2(R))$ and $M \in SL_2(R)$ be such that $B := MA = \begin{bmatrix} g & u \\ 0 & h \end{bmatrix}$ is upper triangular. We write g = ac and h = bc with $(a, b) \in Um(R^2)$ and $c \in R$. As $B \in Um(\mathbb{M}_2(R))$ and Rg + Rh = Rc, we have $(c, u) \in Um(R^2)$. Let $a', b', c', u' \in R$ be such that aa' + bb' = cc' + uu' = 1. For $d \in R$, A and B are equivalent to

$$C := \begin{bmatrix} 1 & db' \\ 0 & 1 \end{bmatrix} B \begin{bmatrix} 1 & da' \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} ac & u + cd(aa' + bb') \\ 0 & bc \end{bmatrix} = \begin{bmatrix} ac & u + cd \\ 0 & bc \end{bmatrix}.$$

Here the role of u + cd is that of an arbitrary element of R whose reduction modulo Rc is the 'fixed' unit $u + Rc \in U(R/Rc)$. Note that each matrix $D_{a',b',d} := \begin{bmatrix} aa'cc' & u + cd \\ 0 & bb'cc' \end{bmatrix} \in Um(\mathbb{M}_2(R))$ is a companion test matrix for A and that $D_{a',b',0}$ is the image of the 'first universal test matrix for Hermite rings'

$$\mathcal{D} := \begin{bmatrix} x(1-yz) & y \\ 0 & (1-x)(1-yz) \end{bmatrix} \in Um\big(\mathbb{M}_2(\mathbb{Z}[x,y,z])\big)$$

via the ring homomorphism $\mathbb{Z}[x, y, z] \to R$ that maps x, y, and z to aa', u, and u' (respectively); so 1 - x maps to bb' = 1 - aa' and 1 - yz maps to 1 - uu' = cc'. We also note that \mathcal{D} is equivalent to the matrix

$$\mathcal{E} := \begin{bmatrix} 1 & 0 \\ z(x-1)(1-yz) & 1 \end{bmatrix} \mathcal{D} \begin{bmatrix} 1 & 0 \\ xz & 1 \end{bmatrix} = \begin{bmatrix} x & y \\ 0 & (1-x)(1-yz)^2 \end{bmatrix}$$

which is the image of the 'second universal test matrix for Hermite rings'

$$\mathcal{F} := \begin{bmatrix} x & y \\ 0 & (1-x)(1-yz) \end{bmatrix} \in Um\big(\mathbb{M}_2(\mathbb{Z}[x,y,z])\big)$$

via the endomorphism of $\mathbb{Z}[x, y, z]$ that fixes x and y and maps z to $2z - yz^2$. See Corollary 4.2 for the usage of universal test matrix in this paragraph.

The 'universal test upper triangular matrix for all rings' is

$$\mathcal{G} := \left[egin{array}{cc} x & y \ 0 & 1-x-yz \end{array}
ight] \in Umig(\mathbb{M}_2(\mathbb{Z}[x,y,z])ig).$$

Proposition 2.3. If \mathcal{P} is weakly determinant liftable we assume that N(R) = 0. Then the following properties hold:

(1) Each upper triangular matrix in $Um(\mathbb{M}_2(R))$ is \mathcal{P} iff for each homomorphism $\phi : \mathbb{Z}[x, y, z] \to R$, the image of $\mathcal{G} \in Um(\mathbb{M}_2(\mathbb{Z}[x, y, z]))$ in $Um(\mathbb{M}_2(R))$ via ϕ is \mathcal{P} .

(2) Each zero determinant upper triangular matrix in $Um(\mathbb{M}_2(R))$ is \mathcal{P} iff for each homomorphism $\phi : \mathbb{Z}[x, y, z] \to R$ with $x(1 - x - yz) \in \text{Ker}(\phi)$, the image of $\mathcal{G} \in Um(\mathbb{M}_2(\mathbb{Z}[x, y, z]))$ in $Um(\mathbb{M}_2(R))$ via ϕ is \mathcal{P} .

Proof. The 'only if' parts are clear. For the 'if' part of (1), let $A = \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \in Um(\mathbb{M}_2(R))$. Let $(a', b', c') \in R^3$ be such that aa' + bb' + cc' = 1. Let $\phi : \mathbb{Z}[x, y, z] \to R$ be the homomorphism that maps x, y and z to aa', b and b' (respectively). The image $\begin{bmatrix} aa' & b \\ 0 & cc' \end{bmatrix}$ of \mathcal{G} via ϕ is \mathcal{P} , so A is \mathcal{P} by Proposition 2.2. The 'if' part of (2) is proved similarly as ac = 0 implies aa'cc' = 0 and $x(1 - x - yz) \in \operatorname{Ker}(\phi)$. \Box

3. Proof of Theorem 1.10

Let $(a, b) \in Um(\mathbb{R}^2)$ and $c \in \mathbb{R}$. For a fixed unit $\bar{u} \in U(\mathbb{R}/\mathbb{R}c)$, let $u \in \mathbb{R}$ be such that $\bar{u} = u + (c)$ and consider the matrix $A = \begin{bmatrix} ac & u \\ 0 & bc \end{bmatrix} \in Um(\mathbb{M}_2(\mathbb{R})).$

Assume R is a pre-Schreier ring and for $(ac, u, bc) \in Um(R^3)$ there exists $(x, y, z, w) \in R^4$ such that $1 - acx - uy - bcw + abc^2(xw - yz) = 0$; so $(1 - acx)(1 - bcw) = y(u+abc^2z)$ (cf. Example 1.3). As $u+abc^2z$ divides (1-acx)(1-bcw) and R is a pre-Schreier ring, we will only use the existence of a nonuple $(x, y, z_1, z_2, w, q, r, s, t) \in R^9$, e.g., $\nu := (x, y, bcz, 0, w, 1, 1, 1, 1) \in R^9$, such that $(ac, sq), (bc, rt) \in Um(R^2)$ and we can decompose $qru + acz_1 + bcz_2 = u_a u_b$, with $u_a \in R$ dividing s - acx and $u_b \in R$ dividing t - bcw. Hence $(q + Rac)^{-1} \cdot (u_a + Rac) \in U(R/Rac)$ and $(r + Rbc)^{-1} \cdot (u_b + Rbc) \in U(R/Rbc)$ are such that the product of their images in U(R/Rc) is \bar{u} . Thus R is a U_2 ring.

Assume R is a Π_2 ring and A is simply extendable. So A is determinant liftable. Let $(x, y, z, w) \in \mathbb{R}^4$ be such that $(1 - acx)(1 - bcw) = (u + abc^2z)y$ and xw = yz (see Example 1.3). Clearly, $(1 - acx, y, 1 - bcw) \in Um(\mathbb{R}^3)$. We will only use the existence of a nonuple $(x, y, z_1, z_2, w, q, r, s, t) \in \mathbb{R}^9$, e.g., ν above, such that

$$D := \begin{bmatrix} s - acx & qru + acz_1 + bcz_2 \\ y & t - bcw \end{bmatrix} \in Um(\mathbb{M}_2(R))$$

has zero determinant, $(ac, sq), (bc, rt) \in Um(R^2)$ and $(s-acx, y, t-bcw) \in Um(R^3)$. As R is a Π_2 ring, D is non-full (see [3], Thm. 1.4) and thus we can decompose $qru + acz_1 + bcz_2 = u_a u_b$, with $u_a \in R$ dividing s - acx and $u_b \in R$ dividing t - bcw. As in the previous paragraph we argue that R is a U_2 ring.

In this paragraph we assume that R is an integral domain and A is simply extendable. To prove that $v_{a,b,c}$ is surjective we can assume that $abc \neq 0$. As A is simply extendable, from [3], Prop. 5.1(1) it follows that its reduction modulo $Rabc^2$ is non-full and hence the system of congruences

 $xy \equiv ac \mod abc^2, \ xw \equiv u \mod abc^2, \ yz \equiv 0 \mod abc^2, \ zw \equiv bc \mod abc^2$

has a solution $(x, y, z, w) \in \mathbb{R}^4$. From the second congruence it follows that $(c, xw) \in Um(\mathbb{R}^2)$. From this and the first and fourth congruences, it follows that there exists $(y', z') \in \mathbb{R}^2$ such that y = cy' and z = cz'. Thus, as \mathbb{R} is an integral domain, the above first system of congruences is equivalent to a second one

 $xy' \equiv a \mod abc, \ xw \equiv u \mod abc^2, \ y'z' \equiv 0 \mod ab, \ z'w \equiv b \mod abc.$

As $(a, b) \in Um(\mathbb{R}^2)$, from the last two congruences it follows firstly that $(a, z') \in Um(\mathbb{R}^2)$ and secondly that there exists $y'' \in \mathbb{R}$ such that y' = ay''. Thus the second system of congruences is equivalent to a third one

 $xy'' \equiv 1 \mod bc, \ xw \equiv u \mod abc^2, \ y''z' \equiv 0 \mod b, \ z'w \equiv b \mod abc.$

From the first and third congruences it follows firstly that $(b, y'') \in Um(\mathbb{R}^2)$ and secondly that there exists $z'' \in \mathbb{R}$ such that z' = bz''. Thus the third system of congruences is equivalent to a fourth one that has only three congruences

$$xy'' \equiv 1 \mod bc, \ xw \equiv u \mod abc^2, \ z''w \equiv 1 \mod ac.$$

Thus $x \in U(R/Rbc)$ and $w + Rac \in U(R/Rac)$ are such that the product of their images in U(R/Rc) is \bar{u} . Hence R is a U_2 ring. Thus Theorem 1.10 holds.

Example 3.1. Let $(a,c) \in R^2$ be such that $Rac \cap R(1-a)c = 0$ and let $u \in R$ be such that $(c,u) \in Um(R^2)$; so a(1-a)c = 0 and $A = \begin{bmatrix} ac & u \\ 0 & (1-a)c \end{bmatrix} \in Um(\mathbb{M}_2(R))$ has zero determinant. The reduction of A modulo Rac is $A_1 := \begin{bmatrix} 0 & u \\ 0 & c \end{bmatrix}$ and modulo R(1-a)c is $A_2 := \begin{bmatrix} c & u \\ 0 & 0 \end{bmatrix}$. Both A_1 and A_2 are non-full. For $i \in \{1, 2\}$, to solve the matrix equation $A_i = \begin{bmatrix} l_i \\ m_i \end{bmatrix} \begin{bmatrix} o_i & q_i \end{bmatrix}$ is the same as solving the system of equations $l_io_i = c\delta_{i1}$, $l_iq_i = u$, $m_io_i = 0$, $m_iq_i = c\delta_{i2}$, where δ_{i1} and δ_{i2} are Kronecker deltas. As $(c, u) \in Um(R^2)$, we have $l_1 \in U(R/Rac)$ and $q_2 \in U(R/R(1-a)c)$. So the solution sets are $\{(l_1, 0, cl_1^{-1}, ul_1^{-1})|l_1 \in U(R/Rac)\}$ for i = 1 and $\{(uq_2^{-1}, cq_2^{-1}, 0, q_2)|q_2 \in U(R/R(1-a)c)\}$ for i = 2. As Rc = Rac + R(a-a)c, the matrix A is non-full iff there exists $(o_1, q_2) \in U(R/Rac) \times U(R/R(1-a)c)$ such that $l_2o_1 + Rc = u + Rc$ and iff $u + Rf \in Im(v_{a,1-a,c})$.

4. Proof of Theorem 1.12 and applications

To prove Theorem 1.12, let S := R/N(R). We show that $(1) \Rightarrow (2)$. As R is an EDR, each matrix in $Um(\mathbb{M}_2(R))$ is determinant liftable by Theorem 1.2 and so it is weakly determinant liftable. From [9], Thm. 3 it follows that S is an EDR and hence it is a pre-Schreier ring by Example 1.8. So $(1) \Rightarrow (2)$ holds.

We show that $(2) \Rightarrow (3)$ if N(R) = 0. As N(R) = 0, from Example 1.3 it follows that condition (1) of Theorem 1.12 holds, hence R is a U_2 ring. We have $(3) \Leftrightarrow (4)$, as for $(a, b), (c, d) \in Um(R^2)$, (4) only translates what means that the unit $d + Rc \in U(R/Rc)$ is in the image of $v_{a,b,c}$. Clearly, (4) \Rightarrow (5) (with b = 1 - a).

To check that $(5) \Rightarrow (4)$, let $(a, b), (c, d) \in Um(R^2)$. If $(e, f, s, t) \in R^4$ is such that ae + bf = cs + dt = 1, then by applying (5) to (ae, d) and $sc \in 1 + Rd$ it follows that there exists $t_1 \in R$ such that we can decompose $d + t_1sc = d_1d_2$ with $(ae, d_1), (1 - ae, d_2) = (b, d_2) \in Um(R_2)$ and thus $(a, d_1), (b, d_2) \in Um(R_2)$ and statement (4) holds by taking $t := t_1s$.

We show that $(3) \Rightarrow (1)$. As R is an EDR iff it is an E_2 ring (see Theorem 1.2), it suffices to show that if R is a U_2 ring, then each $A \in Um(\mathbb{M}_2(R))$ is extendable. Based on [3], Lem. 4.1(3), we can assume that A is upper triangular, and hence we can write $A = \begin{bmatrix} ac & u \\ 0 & bc \end{bmatrix}$ where $(a, b, c, u) \in R^4$ is such that $\det(A) = abc^2$ and $(a, b) \in Um(R^2)$ and u in R is (up to equivalence, see Section 2) an arbitrary representative of a fixed unit $u + Rc \in U(R/Rc)$. As R is a U_2 ring, based on the arbitrariness part we can assume there exists $(d, e) \in R^2$ such that u = de, $d + Rac \in U(R/Rac)$ and $e + Rbc \in U(R/Rbc)$. Let $d', e' \in R$ be such that d' + Racis the inverse of d + Rac and e' + Rac is the inverse of e + Rbc. Let $f, g \in R$ be such that dd' = 1 + acf and ee' = 1 + bcq. The non-full matrix

$$B := \begin{bmatrix} ac(1+bcg) & u \\ abc^2d'e' & bc(1+acf) \end{bmatrix} = \begin{bmatrix} acee' & de \\ abc^2d'e' & bcdd' \end{bmatrix} \in \mathbb{M}_2(R)$$

is congruent to A modulo $R \det(A)$. So A modulo $R \det(A)$ is non-full, thus simply extendable (see [3], Prop. 5.1(2)). Hence A is extendable by [3], Lem. 4.1(1).

We are left to show that $(2) \Rightarrow (3)$ in general (i.e., without assuming that N(R) = 0). For $A \in Um(\mathbb{M}_2(R))$, let $\overline{A} \in Um(\mathbb{M}_2(S))$ be its reduction modulo N(R). If $B \in Um(\mathbb{M}_2(S))$, we choose A such that $B = \overline{A}$. As A is weakly

determinant liftable, there exists $C \in \mathbb{M}_2(R)$ congruent to A modulo $R \det(A)$ and $\det(C) = 0$. Hence $\overline{C} \in \mathbb{M}_2(S)$ is congruent to B modulo $S \det(B)$ and $\det(\overline{C}) = 0$. So S has the same properties as R (note that N(S) = 0 and S = S/N(S) = R/N(R) is a pre-Schreier ring). As we proved that $(2) \Rightarrow (3)$ if N(R) = 0, it follows that S is a U_2 ring. As $(3) \Rightarrow (1)$, we incur that S is an *EDR*. It is well-known that this implies that R is an *EDR*: e.g., see [9], Thm. 3; this also follows from Theorem 1.2 via the fact that A is (simply) extendable iff \overline{A} is so as one can easily check based on [3], Cor. 4.7. As R is an *EDD*, it is an SE_2 ring (see Theorem 1.2) and hence a U_2 ring (see Theorem 1.12(2)). Thus $(2) \Rightarrow (3)$. Hence Theorem 1.12 holds.

Corollary 4.1. Assume R is a Bézout domain such that for each $(a, b) \in Um(R^2)$ and $c \in R$, the image of the functorial homomorphism $U(R/Rabc) \rightarrow U(R/Rab) \cong$ $U(R/Ra) \times U(R/Rb)$ is the product of the images of the functorial homomorphisms $U(R/Rabc) \rightarrow U(R/Ra)$ and $U(R/Rabc) \rightarrow U(R/Rb)$. Then R is an EDR.

Proof. Based on Theorem 1.12, it suffices to show that R is a U_2 ring. Given a unit $u + Rc \in U(R/Rc)$ and $(a, b) \in Um(R^2)$, let $a' := \gcd(a, c)$ and $b' := \gcd(b, c)$. As gcd(a, b) = 1, there exists $c' \in R$ such that c = a'b'c'. From our hypothesis applied to $(a',b') \in Um(R^2)$ and $c' \in R$ it follows that there exist units $e + Rc, f + Rc \in C$ U(R/Rc) such that their images in $U(R/Ra') \times U(R/Rb')$ are (u+Ra', 1+Rb') and (1+Ra', u+Rb') (respectively). Then $(u+Ra, 1+Rb) \in R/Rab \cong R/Ra \times R/Rb$ and $e + Rc \in R/Rc$ (resp. $(1 + Ra, u + Rb) \in R/Rab \cong R/Ra \times R/Rb$ and $f + Rc \in R/Rc$) map to the same element in $R/Ra'b' \cong R/Ra' \times R/Rb'$ and hence, as abc' is the least common multiple of c and ab, there exists $(g,h) \in \mathbb{R}^2$ such that g + Rabc'(resp. h + Rabc') reduces to both of them. As $g + Rbc \in U(R/Rbc)$ and $h + Rac \in U(R/Rbc)$ U(R/Rac), it follows that $(e+Rc)(f+Rc) \in \text{Im}(v_{a,b,c})$. Hence to show that $v_{a,b,c}$ is surjective, by replacing u + Rc with $(u + Rc)[(e + Rc)(f + Rc)]^{-1}$ we can assume that $u-1 \in Ra'b'$, i.e., the images of u+Rc and 1+Rab in Ra'b' are equal, which implies that u + Rc belongs to $Im(U(R/Rabc') \rightarrow U(R/Rc))$. As $abc' + Rabc \in R/Rabc$ has square 0, the homomorphism $U(R/Rabc) \rightarrow U(R/Rabc')$ is surjective and thus u + Rc belongs also to $Im(U(R/Rabc) \rightarrow U(R/Rc))$ and hence to $Im(v_{a,b,c})$.

Corollary 4.2. Let R be a Hermite ring. Then R is an EDR iff for all homomorphisms $\phi : \mathbb{Z}[x, y, z] \to R$, the image of $\mathcal{D} \in Um(\mathbb{M}_2(\mathbb{Z}[x, y, z]))$ (equivalently, $\mathcal{F} \in Um(\mathbb{M}_2(\mathbb{Z}[x, y, z]))$, see Section 2) in $Um(\mathbb{M}_2(R))$ via ϕ is extendable.

Proof. The 'only if' part is obvious. To prove the 'if' part, based on Theorem 1.12, it suffices to prove that R is a U_2 ring, i.e., for all $a, c \in R$ and every $(u, c) \in Um(R^2)$, we have $u + Rc \in Im(v_{a,1-a,c})$. Let $s, t \in R$ be such that cs + ut = 1. As $u + Rcs \in U(R/Rcs)$, it suffices to prove that $u + Rcs \in Im(v_{a,1-a,cs})$. Thus by replacing cwith cs we can assume that c = 1 - ut. Hence $A = \begin{bmatrix} a(1 - ut) & u \\ 0 & (1 - a)(1 - ut) \end{bmatrix} \in Um(\mathbb{M}_2(R))$ is the image of \mathcal{D} via the homomorphism $\phi : \mathbb{Z}[x, y, z] \to R$ that maps x, y, and z to a, u, and t (respectively) and so A is extendable. As \mathcal{D} is equivalent to \mathcal{E} (see Section 2), it follows that A is equivalent to the matrix $B := \begin{bmatrix} a & u \\ 0 & (1 - a)(1 - uw) \end{bmatrix} \in Um(\mathbb{M}_2(R))$, where $w := 2t - ut^2$. From [3], Lem. 4.1(3) it follows that B is extendable and hence its reduction modulo det $(B) = Ra(1 - a)(1 - uw) = Ra(1 - a)c^2$ is non-full (see [3], Prop. 5.1(2)). We will denote *R*. We have a product decomposition $\bar{u} = \bar{u_1}\bar{u_2}$ with $\bar{u_1} = u_1 + Ra(1-a)c^2 \in \bar{R}$ dividing \bar{a} and $\bar{u_2} = u_2 + Ra(1-a)c^2 \in \bar{R}$ dividing $(1-\bar{a})(1-\bar{w}\bar{u})$ and hence also $1-\bar{a}$; here $(u_1, u_2) \in R^2$. Thus there exists a triple $(d, e, f) \in R^3$ such that $u_1u_2 = u + a(1-a)c^2d$, u_1 divides $a + a(1-a)c^2e$ and u_2 divides $1-a+a(1-a)c^2f$. It follows that $(u_1, (1-a)c), (u_2, ac) \in Um(R^2)$, and therefore we have an identity $u + Rc = u_1u_2 + Rc = v_{a,1-a,c}(u_2 + Rac, u_1 + R(1-a)c)$.

Corollary 4.3. Let R be a Bézout domain. Then R is an EDD iff for all $(a, u, t) \in R^3$ with $u \neq 0$ there exists $(s, l, z) \in R^3$ such that

(IV)
$$(1 - us - al)^2 + l - usl - al^2 - (s + t - ust)z = 0$$

Proof. The Hermite ring R is an EDR iff for all $(a, u, t) \in R^3$ the matrix $A = \begin{bmatrix} a(1-ut) & u \\ 0 & (1-a)(1-ut) \end{bmatrix} \in Um(\mathbb{M}_2(R))$ is extendable (see Corollary 4.2) iff it is determinant liftable (see Example 1.1(1)) and hence (see [4], Thm. 1.2) iff there exists $(x, y, z, w) \in R^4$ such that

$$1 - a(1 - ut)x - uy - (1 - a)(1 - ut)w = 0 = xw - yz.$$

If u = 0, then A is diagonal and hence simply extendable (see [3], Ex. 4.9(3)). Thus we can assume $u \neq 0$. The equation 1 - a(1 - ut)x - uy - (1 - a)(1 - ut)w = 0 can be rewritten as 1 - uy = (1 - ut)(ax - aw + w) and working modulo Ru with $u \neq 0$ it follows that its general solution is ax + (1 - a)w = 1 - us and y = s + t - ust with $s \in R$. As R is an integral domain, the general solution of ax + (1 - a)w = 1 - us is x = 1 - us + (1 - a)l and w = 1 - us - al and the equation xw - yz = 0 becomes [1 - us + (1 - a)l](1 - us - al) - (s + t - ust)z = 0 and the corollary follows. \Box

5. Proof of Theorem 1.15

For $(a, b) \in \mathbb{R}^2$, let Diag(a, b) be the diagonal matrix whose (1, 1) and (2, 2) entries are a and b (respectively). We first prove the following general lemmas.

Lemma 5.1. Let $(d, e) \in R^2$ be such that Rd = Re. Then there exists $N \in SL_2(R)$ such that NDiag(d, 0) = Diag(e, 0) (so Diag(d, 0) and Diag(e, 0) are equivalent). Proof. Let $(u, v) \in R^2$ be such that (d, e) = (eu, dv). As $1 - uv \in Ann_R(d)$, for $N := \begin{bmatrix} v & -1 \\ 1 - uv & u \end{bmatrix} \in SL_2(R)$ one computes NDiag(d, 0) = Diag(e, 0). \Box

Lemma 5.2. Let $e \in R$ be such that for $m \in \{2, 4\}$ the reduction map $Um(R^m) \to Um((R/\operatorname{Ann}_R(e))^m)$ is surjective². Assume N(R) = 0. Then all zero determinant matrices in $eUm(\mathbb{M}_2(R))$ admit diagonal reduction iff $R/\operatorname{Ann}_R(e)$ is a Π_2 ring. *Proof.* For $* \in R \cup \mathbb{M}_2(R)$, let $\bar{*}$ be the reduction of * modulo $\operatorname{Ann}_R(e)$. We compute

(V) $\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix} Diag(e,0) \begin{bmatrix} l_{11} & l_{12} \\ l_{21} & l_{22} \end{bmatrix} = \begin{bmatrix} em_{11}l_{11} & em_{11}l_{12} \\ em_{21}l_{11} & em_{21}l_{12} \end{bmatrix}.$

The right hand side of Equation (V) depends only on the pairs $(m_{11}, m_{21}), (l_{11}, l_{12})$ in \mathbb{R}^2 . Given such two pairs, we have $(m_{11}, m_{21}), (l_{11}, l_{12}) \in Um(\mathbb{R}^2)$ iff there exists $(m_{12}, m_{22}, l_{21}, l_{22}) \in \mathbb{R}^4$ such that $\begin{bmatrix} m_{11} & m_{12} \\ m_{21} & m_{22} \end{bmatrix}, \begin{bmatrix} l_{11} & l_{12} \\ l_{21} & l_{22} \end{bmatrix} \in SL_2(\mathbb{R}).$

²Each element of $Um((R/Ann_R(e))^m)$ is the image of an element of $Um((R/Rf)^4)$ for some $f \in Ann_R(e)$ and hence, if $sr(R) \leq m$, of an element in $Um(R^4)$ by [3], Prop. 2.4(1).

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in Um(\mathbb{M}_2(R))$ be such that det(eA) = 0; so $e^2 \det(A) = 0$ and as N(R) = 0 it follows that $e \det(A) = 0$. Thus $\det(A) \in \operatorname{Ann}_R(e)$ and hence $\det(\bar{A}) = 0$. The matrix eA admits diagonal reduction iff it is equivalent to (f,0) (see Example 1.8) with Re = Rf and hence iff it is equivalent to (e,0) (see Lemma 5.1). From this and the previous paragraph it follows that eA admits diagonal reduction iff there exists $(m_{11}, m_{21}), (l_{11}, l_{12}) \in Um(R^2)$ such that (ea, eb, ec, ed) is equal to $(em_{11}l_{11}, em_{11}l_{12}, em_{21}l_{11}, em_{21}l_{12})$ and, as the map $Um(R^2) \to Um((R/\operatorname{Ann}_R(e))^2)$ is surjective, iff $\bar{A} = \begin{bmatrix} \overline{m_{11}} \\ \overline{m_{21}} \end{bmatrix} \begin{bmatrix} \overline{l_{11}} & \overline{l_{12}} \end{bmatrix}$ is non-full. Each $C \in Um(\mathbb{M}_2(R/\operatorname{Ann}_R(e)))$ is \bar{B} for some $B \in Um(\mathbb{M}_2(R))$; if $\det(C) = 0$, then $\det(B) \in \operatorname{Ann}_R(e)$ and hence $\det(eB) = 0$. Thus we conclude that all zero determinant matrices in $aUm(\mathbb{M}_2(R))$ admit diagonal reduction iff all zero determinant matrices in $Um(\mathbb{M}_2(R/\operatorname{Ann}_R(e)))$ are non-full (equivalently, are simply extendable by [3], Prop. 5.1(1)) and iff $R/\operatorname{Ann}_R(e)$ is a Π_2 ring.

We prove Theorem 1.15. If R is an EDR then all zero determinant matrices in $\mathbb{M}_2(R)$ admit diagonal reduction. If for each $a \in R$, $R/Ann_R(a)$ is a Π_2 ring, then all zero determinant matrices in $RUm(\mathbb{M}_2(R)) = \mathbb{M}_2(R)$ admit diagonal reduction by Lemma 5.2. In the last two sentences we can replace 'admit diagonal reduction' by 'are non-full' by Example 1.8, hence Theorem 1.15 follows from Theorem 1.2.

6. On $(WSU')_n$ AND $(WSU)_n$ RINGS

For $F \in \mathbb{M}_n(R)$, let F^T be its transpose and let Ker_F and Im_F be the kernel and the image (respectively) of the *R*-linear map $L_F : \mathbb{R}^n \to \mathbb{R}^n$ defined by it.

Proposition 6.1. A ring R is an $(SU')_n$ (resp. $(SU)_n$) ring in the sense of Definition 2.1(2) iff it is so in the sense of [12], Def. 2.1.

Proof. For the 'only if' part, let $B \in \mathbb{M}_n(R)$. As R is a Hermite ring, there exist $e \in R$ and $C \in Um(\mathbb{M}_n(R))$ such that B = eC. As R is an $(SU')_n$ (resp. $(SU)_n$) ring, there exists $N \in SL_n(R)$ (resp. $N \in GL_n(R)$) such that AN is symmetric. Hence BN = e(AC) is symmetric, thus R is an $(SU')_n$ (resp. $(SU)_n$) ring in the sense of [12], Def. 2.1. For the 'if part, clearly R is a $(WSU')_n$ (resp. $(WSU)_n$) ring and it is a Hermite ring by [12], Prop. 3.1.

As in [12], Sect. 2, if U(R) is a 2-divisible abelian group, R is a $(WSU')_n$ ring iff it is a $(WSU)_n$ ring. For $(WSU)_n$ rings we have the following general result.

Theorem 6.2. Assume R is a $(WSU)_n$ ring. Then the following properties hold:

(1) Each unimodular matrix $A \in Um(\mathbb{M}_n(R))$ is equivalent to its transpose A^T .

(2) If n = 2, then every projective R-module of rank 1 and generated by 2 elements is self-dual.

(3) If $sr(R) \leq n^2$, then for each $d \in R$, R/Rd is a $(WSU)_n$ ring.

Proof. If $N \in GL_n(R)$ is such that AN is symmetric, then $(N^T)^{-1}AN = A^T$ and hence part (1) holds. To check part (2), let P be a projective R-module of rank 1 generated by 2 elements. We consider an isomorphism $P \oplus Q \cong R^2$ to be viewed as an identification. Taking determinants it follows that Q is the dual of P. Let $B \in Um(\mathbb{M}_2(R))$ be such that $\operatorname{Ker}_B = Q$ and $\operatorname{Im}_B = P$. As B^T and B are equivalent by part (1), the R-modules P and Q are isomorphic, so P is selfdual. Thus part (2) holds. To check part (3) for $d \in R$, let $\overline{A} \in Um(\mathbb{M}_n(R/Rd))$. As $sr(R) \leq n^2$, there exists $A \in Um(\mathbb{M}_n(R))$ such that its reduction modulo Rdis \overline{A} (see [3], Prop. 2.4(1)). Let $N \in GL_n(R)$ be such that AN is symmetric. If $\overline{N} \in GL_n(R/Ra)$ is the reduction of N modulo Rd, then \overline{AN} , being the reduction of AN modulo Rd, is symmetric. Thus R/Rd is a $(WSU)_n$ ring; so part (3) holds. \Box

Remark 6.3. If R is an integral domain such that for each $d \in R$, every projective R/Rd-module of rank 1 and generated by 2 elements is self-dual, then each $A \in Um(\mathbb{M}_n(R))$ is equivalent to its transpose A^T .

7. Proof of Theorem 1.17 and applications

Let Spec R be the spectrum of R and let Max R be its subset of maximal ideals. To prove part (1), it suffices to show that for all $(u, c) \in Um(R^2)$ and $a \in R$, we have $u^2 + Rc \in Im(v_{a,1-a,c})$. To check this we can assume that $c \neq 0$. As in the proof of Corollary 4.2, we can assume that there exists $t \in R$ such that c = 1-ut. As we are assuming that R is a $(WSU)_2$ ring, there exists $M = \begin{bmatrix} x & y \\ z & w \end{bmatrix} \in GL_2(R)$ such that $M \begin{bmatrix} ac & u \\ 0 & (1-a)c \end{bmatrix}$ is a symmetric matrix, i.e., we have an identity (VI) acz = ux + (1-a)cy.

As c divides ux and $(u, c) \in Um(R^2)$, c divides x. Let $s \in R$ be such that x = cs. Let $\mu := \det(M) = csw - yz \in U(R)$. Thus $(cs, yz) \in Um(R^2)$.

We first assume that $c \notin Z(R)$. Dividing Equation (VI) by c it follows that

(VII)
$$az + (a-1)y = us$$

We check that the assumption $(a, s) \notin Um(R^2)$ leads to a contradiction. This assumption implies that there exists $\mathfrak{m} \in Max R$ such that $Ra + Rs \subseteq \mathfrak{m}$. From this and Equation (VII) it follows that $(a-1)y \in \mathfrak{m}$. As $a-1 \notin \mathfrak{m}$, it follows that $y \in \mathfrak{m}$, hence $(s, yz) \notin Um(R^2)$, a contradiction. As $(a, s) \in Um(R^2)$, $s + Ra \in U(R/Ra)$.

From Equation (VII) it follows that the images of $(-y + Rc)(u + Rc)^{-1} \in U(R/Rc)$ and $s + Ra \in U(R/Ra)$ in U(R/(Ra + Rc)) are equal. Thus there exists a unit of the quotient ring $R/(Ra \cap Rc)$ that maps to $(-y+Rc)(u+Rc)^{-1} \in U(R/Rc)$. As $(Ra \cap Rc)/Rac$ is a square 0 ideal of R/Rac, the homomorphism $U(R/Rac) \rightarrow U(R/(Ra \cap Rc))$ is surjective, hence $(-y+Rc)(u+Rc)^{-1} \in Im(v_{a,1-a,c})$.

Due to the symmetry of Equation (VII) in az and (a-1)y, similar arguments give $s + R(1-a) \in U(R/R(1-a))$ and $(z+Rc)(u+Rc)^{-1} \in Im(v_{a,1-a,c})$. Clearly $\mu + Rc \in Im(v_{a,1-a,c})$. As $\mu + Rc = (-y+Rc)(z+Rc)$, it follows that

$$(u+Rc)^{2} = [(-y+Rc)(u+Rc)^{-1}]^{-1}[(z+Rc)(u+Rc)^{-1}]^{-1}(\mu+Rc) \in Im(v_{a,1-a,c}).$$

We use a trick with annihilators to show that an analogue of Equation (VII) holds even if $c \in Z(R)$. As $(c, u), (c, yz) \in Um(R^2)$, it follows that $(c, uyz) \in Um(R^2)$. This implies that $\operatorname{Ann}_R(c) \subset Ruyz$. From this and the identity c[az+(a-1)y-us] =0 it follows that there exists $q \in R$ such that az + (a - 1)y - su = uyzq. If $s_1 := s + yzq$, then $(s_1, yz) \in Um(R^2)$ and the identity $az + (a - 1)y = us_1$ holds. Thus, using s_1 and the last identity instead of s and Equation (VII), as above we argue that $(u + Rc)^2 \in Im(v_{a,1-a,c})$. So part (1) holds. Part (2) follows from Theorem 6.2(2) and (3). The first part of part (3) follows from part (1) and definitions. Based on it, the second part of part (3) follows from Theorem 1.12. Thus Theorem 1.17 holds.

Remark 7.1. If $(a, y) \in Um(R^2)$ (resp. $(1 - a, z) \in Um(R^2)$), then $-y + Rac \in U(R/Rac)$ (resp. $z + R/R(1-a)c \in U(R/R(1-a)c)$), hence $u + Rc \in Im(v_{a,1-a,c})$.

Corollary 7.2. For a Bézout domain the following statements are equivalent:

(1) The ring R is an $(WSU)_2$ (resp. $(WSU')_2$) ring.

(2) For all $(a,b), (c,u) \in Um(R^2)$ with $abc \neq 0$, if $(a',b') \in R^2$ such that aa' + bb' = 1 is given, then there exists a triple $(s,l,w) \in R^3$ with the property that $csw + (a'us - bl)(b'us + al) \in U(R)$ (resp. csw + (a'us - bl)(b'us + al) = 1).

Proof. If $A_1, A_2 \in \mathbb{M}_2(R)$ are equivalent, then there exists $M_1 \in GL_2(R)$ such that M_1A_1 is symmetric iff there exists $M_2 \in GL_2(R)$ such that M_2A_2 is symmetric. This is so, as for $M, N, M_1 \in GL_2(R)$ such that $A_2 = MA_1N$ and M_1A_1 is symmetric, by denoting $M_2 := N^T M_1 M^{-1} \in GL_2(R)$, $M_2A_2 = N^T (M_1A_1)N$ is symmetric. Also, if $M, N, M_1 \in SL_2(R)$, then $M_2 \in SL_2(R)$.

Based on the previous paragraph and Section 2, the ring R is a $(WSU)_2$ (resp. $(WSU')_2$) ring iff for all $(a, b), (c, u) \in Um(R^2)$ there exists $M = \begin{bmatrix} x & y \\ z & w \end{bmatrix}$ in $GL_2(R)$ (resp. in $SL_2(R)$) such that the matrix $M \begin{bmatrix} ac & u \\ 0 & bc \end{bmatrix}$ is symmetric, i.e., the equation acz = ux + bcy holds. If abc = 0, then $\begin{bmatrix} ac & u \\ 0 & bc \end{bmatrix}$ is simply extendable (see [3], Ex. 4.9(3)) and hence equivalent to a diagonal matrix (see [3], Thm. 4.3) and from the previous paragraph we get that M exists. Thus we can assume that

and from the previous paragraph we get that M exists. Thus we can assume that $abc \neq 0$. So $c \notin Z(R)$ and hence as in the proof of Theorem 1.17, we write x = cs with $s \in R$ and the identity az = us + by holds. The general solution of the equation az - by = us in the indeterminates y, z is z = a'us - bl and y = -b'us - al with $l \in R$. As det(M) = xw - yz = csw + (a'us - bl)(b'us + al), the corollary hods. \Box

Remark 7.3. Referring to Corollary 7.2(2) with b = 1 - a, we can take a' = 1 = b', and one is searching for $(s, l, w) \in \mathbb{R}^3$ such that $csw + (us + al - l)(us + al) \in U(\mathbb{R})$ (resp. and csw + (us + al - l)(us + al) = 1, which for c = 1 - ut with $t \in \mathbb{R}$ is closely related to Equation (IV) via a substitution of the form z := -w).

Remark 7.4. A symmetric matrix $A = \begin{bmatrix} a & b \\ b & d \end{bmatrix} \in Um(\mathbb{M}_2(R))$ is determinant liftable iff there exists $(x, w)^2 \in R$ such that b divides 1 - ax - dw and for a suitable $s \in R$ such that bs = 1 - ax - cw, the equation $X^2 + sX + xw = 0$ has two solutions $y, z \in R$ (so xw - yz = 0 and 1 - ax - by - bz - dw = 0).

Recall from [24], Sect. 1 that R is said to have square stable range 1, and one writes ssr(R) = 1, if for each $(a, b) \in Um(R^2)$, there exists $r \in R$ such that $a^2 + br \in U(R)$. Clearly, if sr(R) = 1, then ssr(R) = 1.

Example 7.5. Assume ssr(R) = 1. Then for each $c \in R$, the cokernel of the reduction homomorphism $U(R) \to U(R/Rc)$ is a Boolean group and hence for all $(a,b) \in Um(R^2)$, $Coker(v_{a,b,c})$ is a Boolean group. If $(c,u) \in Um(R^2)$, let $w \in R$ be such that $u^2 + cw \in U(R)$. Then $M := \begin{bmatrix} c & -u \\ u & w \end{bmatrix} \in GL_2(R)$ and the

product matrix $M\begin{bmatrix} ac & u \\ 0 & (1-a)c \end{bmatrix}$ is symmetric. Hence, if R is also a Hermite ring, then each upper triangular matrix in $Um(\mathbb{M}_2(R))$ has a companion test matrix equivalent to a symmetric matrix; thus, from this, Theorem 1.2 and Proposition 2.2 it follows that R is an EDR iff each symmetric matrix in $Um(\mathbb{M}_2(R))$ is extendable.

8. Proof of Theorem 1.18

Let P be a projective R/Rd-module of rank 1 generated by 2 elements. We consider an isomorphism $P \oplus Q \cong (R/Rd)^2$. Let $D \in Um(\mathbb{M}_2(R/Rd))$ be such that $\operatorname{Ker}_D = Q$ and $\operatorname{Im}_D = P$. As $sr(R) \leq 2$, there exists $A \in Um(\mathbb{M}_2(R))$ such that its reduction modulo Rd is D (see [3], Prop. 2.4(1)). Let $\overline{R} := R/R \det(A)$ and $\operatorname{let} \overline{A} \in Um(\mathbb{M}_2(\overline{R}))$ be the reduction of A modulo $R \det(A)$. We have $\det(A) \in Rd$. We know that $\operatorname{Ker}_{\overline{A}}$ and $\operatorname{Im}_{\overline{A}}$ are dual projective \overline{R} -modules of rank 1 generated by 2 elements (see [3], Sect. 3) and their reductions modulo the ideal $Rd/R \det(A)$ of \overline{R} are Q and P. Thus, to end the proof, it suffices to show that the classes of $\operatorname{Ker}_{\overline{A}}$ and $\operatorname{Im}_{\overline{A}}$ in $Pic(\overline{R})$ are equal (equivalently, have orders 1 or 2 or equivalently, $\operatorname{Ker}_{\overline{A}}^2$ and $\operatorname{Im}_{\overline{A}}^2$ are free R-modules of rank 2). So, d and D will not be mentioned again.

and $\operatorname{Im}_{\overline{A}}^2$ are free *R*-modules of rank 2). So, *d* and *D* will not be mentioned again. Based on Section 2, we can assume that $A = \begin{bmatrix} ac & u \\ 0 & bc \end{bmatrix} \in \mathbb{M}_2(R)$ with $(a, b, c) \in \mathbb{R}^3$ such that $(a, b), (c, u) \in Um(\mathbb{R}^2)$ and we will only use that $\operatorname{Coker}(v_{a,b,c})$ is a Boolean group. As $\operatorname{Coker}(v_{a,b,c})$ makes sense for R/Rabc and as two finitely generated projective R/Rabc^2 -modules are isomorphic if and only if their reductions modulo the square 0 ideal $\operatorname{Rabc}/\operatorname{Rabc}^2$ are isomorphic, by reducing modulo Rabc we can assume that abc = 0. Hence $(R, A) = (\overline{R}, \overline{A})$ and we will only use (R, A).

Let $(s,t) \in R^2$ be such that cs + ut = 1. By [3], Ex. 5.2 we have $\operatorname{Ker}_A = Rv_1 + Rv_2$, where $v_1 := \begin{bmatrix} -u \\ ac \end{bmatrix}$ and $v_2 := \begin{bmatrix} -bc \\ 0 \end{bmatrix}$. As $(a,b) \in Um(R^2)$, we have $Spec R = Spec R_a \cup Spec R_b$. Note $(\operatorname{Ker}_A)_a = R_a v_1$ and $(\operatorname{Ker}_A)_b = R_b(tv_1 + \frac{s}{b}v_2)$. Over R_{ab} we have c = 0 and hence $v_1 = u(tv_1 + \frac{s}{b}v_2)$. Thus $(\operatorname{Ker}_A^{\otimes 2})_a = R(v_1 \otimes v_1)$, $(\operatorname{Ker}_A^{\otimes 2})_b = R((tv_1 + \frac{s}{b}v_2) \otimes (tv_1 + \frac{s}{b}v_2))$, and over R_{ab} we have an identity $v_1 \otimes v_1 = u^2((tv_1 + \frac{s}{b}v_2) \otimes (tv_1 + \frac{s}{b}v_2))$. As $u^2 + Rc \in \operatorname{Im}(v_{a,b,c})$, there exists $u_a \in U(R/Rbc)$ and $u_b \in U(R/Rac)$ such that the product of their images in U(R/Rc) is $u^2 + Rc$. As R_a is a localization of R/Rbc and R_b is a localization of R/Rac, we will denote also by u_a and u_b their images in R_a and R_b (respectively). Thus $u_a^{-1}(v_1 \otimes v_1)$ and $u_b^{-1}((tv_1 + \frac{s}{b}v_2) \otimes (tv_1 + \frac{s}{b}v_2))$ coincide over R_{ab} and hence there exists an element $v \in \operatorname{Ker}_A^{\otimes 2}$ whose images in $(\operatorname{Ker}_A^{\otimes 2})_a$ and $(\operatorname{Ker}_A^{\otimes 2})_b$ are equal to $u_a^{-1}(v_1 \otimes v_1)$ and $u_b^{-1}((tv_1 + \frac{s}{b}v_2) \otimes (tv_1 + \frac{s}{b}v_2))$ (respectively). Thus $Rv = \operatorname{Ker}_A^{\otimes 2}$, hence $\operatorname{Ker}_A^{\otimes 2} \cong R$.

Corollary 8.1. Let R be a Hermite ring such that for each $a \in R$, every selfdual projective R/Ra-module of rank 1 generated by 2 elements is free. Then the following properties hold:

(1) The ring R is an EDR iff for all $(a,b) \in Um(R^2)$ and $c \in R$, $Coker(v_{a,b,c})$ is a Boolean group.

(2) If R is a $(WSU)_2$ ring, then R is an EDR.

Proof. For part (1), its 'only if' part follows from Theorem 1.12. To check the 'if' part, from Theorem 1.18 it follows that for each $A \in Um(\mathbb{M}_2(R))$, the reduction \overline{A} of A modulo $R \det(A)$ is such that $\operatorname{Im}_{\overline{A}} \cong R/R \det(A)$, hence \overline{A} is non-full and thus extendable (see [3], Prop. 5.1(2)). Hence R is an E_2 ring and thus an *EDR* by Theorem 1.2. Thus Corollary 8.1(1) holds. Corollary 8.1(2) follows from Corollary 8.1(1) and Theorem 1.17(1) (it also follows from Theorems 6.2(2) and 1.2).

9. Proof of Criteria 1.19 and 1.20

We first prove Criterion 1.19(1). If there exists $(e, f) \in R^2$ such that $ae^2 - cf^2 \in U(R)$, then from the identity $ae^2 - cf^2 = e(ae + bf) - f(be + cf)$ it follows that $(ae + bf, be + cf) \in Um(R^2)$, so A is simply extendable by [3], Thm. 4.3. For the converse, assume char(R) = 2 and A is simply extendable. Let $(e, f) \in Um(R^2)$ be such that $(ae + bf, be + cf) \in Um(R^2)$ by loc. cit. To prove that $ae^2 - cf^2 \in U(R)$ it suffices to show that the assumption that there exist $\mathfrak{m} \in MaxR$ such that $ae^2 - cf^2 \in \mathfrak{m}$, leads to a contradiction. By replacing R with R/\mathfrak{m} , we can assume that R is a field and we know that either $a \neq 0$ or $c \neq 0$, either $ae + bf \neq 0$ or $be + cf \neq 0$, $ac = b^2$ and $ae^2 = cf^2$; hence either $e \neq 0$ or $f \neq 0$. If acef = 0, then by the symmetry in (a, c) and (e, f) we can assume that ae = 0; if a = 0, then b = 0, $c \neq 0$, and hence f = 0, thus ae + bf = be + cf = 0, a contradiction, and if e = 0, then $f \neq 0$, hence c = 0, and by the symmetry in the pair (a, c), we similarly reach a contradiction. Thus we can assume that $acef \neq 0$, hence $b^2 = ac = e^{-2}c^2f^2 = f^{-2}a^2e^2$; as char(R) = 2 and R is a field, it follows that $b = e^{-1}cf = f^{-1}ae$, thus ae + bf = be + cf = 0, a contradiction. So part (1) holds.

To prove part (2), as $b \notin Z(R)$, for each $\mathfrak{m} \in MaxR$, b is a nonzero element of $R_{\mathfrak{m}}$. As R is a Hermite ring and N(R) = 0, each $R_{\mathfrak{m}}$ is a valuation domain (e.g., see [4], Ex. 2.4). Let $d \in R$ and $(a', b') \in Um(R^2)$ be such that a = da' and b = db'. As d divides a and $(a, c) \in Um(R^2)$, we have $(d, c) \in Um(R^2)$. Thus also $(d^2, c) \in Um(R^2)$. From this, as d^2 divides ac, it follows that d^2 divides a; let $u \in R$ be such that $a = d^2u$. As $b \in R_{\mathfrak{m}} \setminus \{0\}$, it is easy to see that the image of u in $R_{\mathfrak{m}}$ is a unit for all $\mathfrak{m} \in MaxR$. Hence $u \in U(R)$. By symmetry, there exist $g \in R$ and $v \in U(R)$ such that $c = g^2v$. As $b \notin Z(R)$, we have $d, g \notin Z(R)$.

To complete the proof that the matrix A is extendable, based on part (1) it suffices to show that there exists $(e, f) \in R^2$ such that $ae^2 - cf^2 = ae^2 + cf^2 \in U(R)$. Replacing A by $u^{-1}A$ (see [3], Lem. 4.1(3)), we can assume that u = 1. Thus $a = d^2$. From this and the identities $ac = b^2$ and $c = g^2v$, the element $w := \frac{b}{gd}$ of the total ring of fractions of R has the property that $w^2 = v$ in each valuation domain $R_{\mathfrak{p}}$ with $\mathfrak{p} \in Spec R$, hence $w \in R_{\mathfrak{p}}$, and we conclude that $w \in R$ and $v = w^2$. By replacing (v,g) with (1,gw), we can assume that v = 1 and $c = g^2$. As $(a,c) = (d^2,g^2) \in Um(R^2)$, it follows that $(d,g) \in Um(R^2)$, hence there exists $(e,f) \in Um(R^2)$ such that de + gf = 1, and so $ae^2 + cf^2 = 1 \in U(R)$. Thus part (2) holds and so Criterion 1.19 holds.

We prove Criterion 1.20. As R is an $(SU)_2$ ring, it is a Hermite ring, and, as the extendability is preserved under equivalence (see [3], Lem. 4.1(3)), it is an E_2 ring iff each symmetric matrix in $Um(\mathbb{M}_2(R))$ is extendable. Thus R is an EDR iff each symmetric matrix in $Um(\mathbb{M}_2(R))$ is extendable by Theorem 1.2. Let $A = \begin{bmatrix} a & b \\ b & c \end{bmatrix} \in Um(\mathbb{M}_2(R))$. Based on [3], Lem. 4.1(1) and Criterion 1.19(1) applied to A modulo $R(ac - b^2)$, A is extendable if there exists $(e, f) \in R^2$ such that $ae^2 - cf^2 + R(ac - b^2) \in U(R/[R(ac - b^2)])$, equivalently, if $(ae^2 - cf^2, ac - b^2) \in Um(R^2)$, and the converse holds if char(R) = 2; hence Criterion 1.20 holds.

10. Proof of Criterion 1.22

As R is a Bézout domain, R is an EDD iff for all $(a, b, s) \in \mathbb{R}^3$, the equation

(VIII)
$$[1 - (1 - a - bs)w - by](1 - ax) = ay[(1 - a - bs)z + bx]$$

in the indeterminates x, y, z and w has a solution in \mathbb{R}^4 (see Corollary 1.14 and Proposition 2.3(1)). To solve this equation, we note that, as $(a, 1 - ax) \in Um(\mathbb{R}^2)$, a divides 1 - (1 - a - bs)w - by. Thus, as R is an integral domain, Equation (IX) has a solution in \mathbb{R}^4 iff the system of two equations

(IX)
$$1 - (1 - a - bs)w - by = ta \text{ and } t(1 - ax) = y[(1 - a - bs)z + bx]$$

in the indeterminates t, x, y, z and w has a solution in \mathbb{R}^5 (this holds even if a = 0!).

The first equation of System (IX), rewritten as w = 1 + a(w - t) + b(sw - y), has the general solution given by w = 1 - aq - br, t = q + w = 1 + q - aq - br and y = r + sw = r + s(1 - aq - br), where $(q, r) \in \mathbb{R}^2$ can be arbitrary.

The second equation of System (IX), rewritten as t = (at+by)x + (1-a-bs)yz, has a solution iff $t \in R(at+by) + R(1-a-bs)y$. As 1 = (1-a-bs)w + (at+by), we have $t \in R(at+by) + R(1-a-bs)y$ iff $t \in R(at+by) + Ry = Rat + Ry$.

From the last two paragraphs it follows that System (IX) has a solution in \mathbb{R}^5 iff there exists $(q, r) \in \mathbb{R}^2$ as mentioned in statement (2). Thus (1) \Leftrightarrow (2).

If $(1-a,b) \in Um(R^2)$, then we can choose $(q,r) \in Um(R^2)$ such that t = 1 + q(1-a) + rb = 0, hence $t = 0 \in Rat + Ry = Ry$, and for e := 1 and f := 0, we have $(a, e), (be + af, 1 - bs - a) \in Um(R^2)$. Thus to prove that $(2) \Leftrightarrow (3)$ we can assume that $(1-a,b) \notin Um(R^2)$, and hence always $t \neq 0$.

For $t_1 := \gcd(r-qs, t)$ we write $t = t_1t_2$ and $r-qs = \alpha t_1$ with $(\alpha, t_2) \in Um(R^2)$; thus $y = r + s(t-q) = st + \alpha t_1$. We would like to find $(q, r) \in R^2$ such that there exists $(g, h) \in R^2$ with the property that gy+aht = t, equivalently, $gst+g\alpha t_1+aht = t$, and thus we must have $g \in Rt_2$. Writing $g = t_2\beta$ with $\beta \in R$, we want to find $(q, r) \in R^2$ such that there exists $(\beta, h) \in R^2$ satisfying $1 = \beta(st_2 + \alpha) + ah$, equivalently, such that $(a, st_2 + \alpha) \in Um(R^2)$. Replacing $r = qs + \alpha t_1$ in t, it follows that $t = 1 + q - bqs - b\alpha t_1 - aq$, hence there exists $\gamma \in R$ such that $1 + q(1 - bs - a) = t_1\gamma$; we have $t_2 = \gamma - b\alpha$ and thus $e := st_2 + \alpha = s\gamma + (1 - bs)\alpha$. As $(\alpha, t_2) \in Um(R^2)$, it follows that $(\alpha, \gamma) \in Um(R^2)$. There exists a unique $f \in R$ such that be + f(1 - bs) divides 1 + q(1 - bs - a) iff $(be + f(1 - bs), 1 - bs - a) \in Um(R^2)$ and hence iff $(be + af, 1 - bs - a) \in Um(R^2)$. Thus $(2) \Leftrightarrow (3)$. So Criterion 1.22 holds.

Acknowledgement. The third author would like to thank SUNY Binghamton for good working conditions. Competing interests: the authors declare none.

References

- V. A. Bovdi, V. P. Shchedryk Commutative Bézout domains of stable range 1.5. Linear Algebra Appl. 568 (2019), 127–134.
- [2] P. Carbonne Anneaux de Bézout, Hermite et Kaplansky "universels". Canad. Math. Bull. 30 (1987), no. 4, 461–470.

- [3] G. Călugăreanu, H. F. Pop, and A. Vasiu Matrix invertible extensions over commutative rings. Part I: general theory. 17 pages manuscript dated April 2, 2024, https://arxiv.org/abs/2404.05780.
- [4] G. Călugăreanu, H. F. Pop, and A. Vasiu Matrix invertible extensions over commutative rings. Part II: determinant liftability. 13 pages manuscript dated April 25, 2024.
- [5] L. Fuchs, L. Salce Modules over non-Noetherian domains. Math. Surveys Monogr. 84, American Mathematical Society, Providence, RI, 2001.
- [6] L. Gillman, M. Henriksen Rings of continuous functions in which every finitely generated ideal is principal. Trans. Amer. Math. Soc. 82 (1956), 366–391.
- [7] L. Gillman, M. Henriksen Some remarks about elementary divisor rings. Trans. Amer. Math. Soc. 82 (1956), 362–365.
- [8] O. Gabber, A. Vasiu Pathologies for commutative Bézout rings. Work in progress.
- [9] M. Henriksen Some remarks on elementary divisor rings. II. Michigan Math. J. 3 (1955/56), 159–163.
- [10] C. U. Jensen Arithmetical rings. Acta Math. Acad. Sei. Hungar. 17 (1966), 115–123.
- [11] M. D. Larsen, W. J. Lewis, T. S. Shores Elementary divisor rings and finitely presented modules. Trans. Amer. Math. Soc. 187 (1974), 231–248.
- [12] D. Lorenzini Elementary divisor domains and Bézout domains. J. Algebra 371 (2012), 609– 619.
- [13] W. W. McGovern Neat rings. J. Pure Appl. Algebra 205 (2006), no. 2, 243–265.
- [14] P. Menal, J. Moncasi On regular rings with stable range 2. J. Pure Appl. Algebra 24 (1982), no. 1, 25–40.
- [15] M. Roitman The Kaplansky condition and rings of almost stable range 1. Proc. Amer. Math. Soc. 141 (2013), no. 9, 3013–3018.
- [16] D. E. Rush Bézout domains with stable range 1. J. Pure Appl. Algebra 158 (2001), no. 2–3, 309–324.
- [17] V. P. Shchedryk Commutative domains of elementary divisors and some properties of their elements. Ukrainian Math. J. 64 (2012), no. 1, 140–155.
- [18] V. P. Shchedryk Bézout rings of stable range 1.5. Translation of Ukrainian Math. J. 67 (2015), no. 6, 849–860.
- [19] V. P. Shchedryk Bézout rings of stable rank 1.5 and the decomposition of a complete linear group into products of its subgroups. Ukrainian Math. J. 69 (2017), no. 1, 113–120; translation in Ukrainian Math. J. 69 (2017), no. 1, 138–147.
- [20] R. Wiegand, S. Wiegand Finitely generated modules over Bézout rings. Pacific J. Math. 58 (1975), no. 2, 655–664.
- [21] B. V. Zabavsky Diagonal reduction of matrices over rings. Math. Stud. Monog. Ser., Vol. 16, VNTL Publishers, L'viv, 2012. 251 pp.
- [22] B. V. Zabavsky Diagonal reduction of matrices over finite stable range rings. Mat. Stud. 41 (2014), no. 1, 101–108.
- [23] B. V. Zabavsky, S. I. Bilavska Decomposition of finitely generated projective modules over Bézout ring. Mat. Stud. 40 (2013), no. 1, 104–107.
- [24] B. V. Zabavsky, O. Romaniv A Bézout ring of stable range 2 which has square stable range 1. Comm. Algebra 47 (2019), no. 12, 5392–5397.

Grigore Călugăreanu E-mail: calu@math.ubbcluj.ro

Address: Department of Mathematics, Babeş-Bolyai University,

1 Mihail Kogălniceanu Street, Cluj-Napoca 400084, Romania.

Horia F. Pop E-mail: horia.pop@ubbcluj.ro

Address: Department of Computer Science, Babeş-Bolyai University, 1 Mihail Kogălniceanu Street, Cluj-Napoca 400084, Romania.

Adrian Vasiu, E-mail: avasiu@binghamton.edu

Address: Department of Mathematics and Statistics, Binghamton University,

P. O. Box 6000, Binghamton, New York 13902-6000, U.S.A.