

A MODEL PROBLEM FOR FIRST ORDER MEAN FIELD GAMES WITH DISCRETE INITIAL DATA

P. JAMESON GRABER AND BRADY ZIMMERMAN

ABSTRACT. In this article, we study a simplified version of a density-dependent first-order mean field game, in which the players face a penalization equal to the population density at their final position. We consider the problem of finding an equilibrium when the initial distribution is a discrete measure. We show that the problem becomes finite-dimensional: the final piecewise smooth density is completely determined by the weights and positions of the initial measure. We establish existence and uniqueness of a solution using classical fixed point theorems. Finally, we show that Newton's method provides an effective way to compute the solution. Our numerical simulations provide an illustration of how density penalization in a mean field game tends to smooth the initial distribution.

1. INTRODUCTION

Mean field games were introduced by Lasry and Lions [LL06a, LL06b, LL07] and Caines, Huang, and Malhamé [HMC06] in order to model the interactions between large numbers of agents playing a differential game; see for instance [BFY13, CD17, G⁺14, ACD⁺20] for an overview of the field. In this article, we are interested in first-order, or deterministic, mean field games. Typically, these are modeled using a first-order coupled system of partial differential equations, as follows:

$$(1.1) \quad \begin{cases} -\partial_t u + H(x, \nabla_x u) = F(x, m), & x \in \mathbb{R}^d, t \in (0, T), \\ \partial_t m - \nabla_x \cdot (D_p H(x, \nabla_x u) m) = 0, & x \in \mathbb{R}^d, t \in (0, T), \\ m(x, 0) = m_0(x), & x \in \mathbb{R}^n, \\ u(x, T) = G(x, m(x, T)), & x \in \mathbb{R}^n. \end{cases}$$

Here H is the Hamiltonian for an optimal control problem, which a representative agent solves for a given population density $m(x, t)$. The functions F and G model the cost this player must pay as a function of the density m . In Nash equilibrium, the density will evolve according to the continuity equation driven by the optimal feedback vector field $-D_p H(x, \nabla_x u)$.

There are now many results on the existence and uniqueness of suitably defined *weak* solutions to System (1.1) [Car15, Gra14, CG15, CGPT15] as well as regularity of solutions [PS17, GM18, Mun22]. As far as we know, all of these results apply only when the initial distribution m_0 is a density, i.e. it has no singular part with respect to Lebesgue measure. It remains an open question how to analyze weak solutions in the case where m_0 is replaced with a more general measure, which could have singular parts.

In this article, we contribute to the theory of mean field games by analyzing an example in which the initial measure is entirely discrete, i.e. it is a weighted sum of Dirac measures. We simplify the problem by choosing $F = 0$, $H(x, p) = \frac{1}{2}p^2$, and $G(x, m(x, T)) = m(x, T)$. We will also set the

Date: July 26, 2024.

2020 Mathematics Subject Classification. 49N80, 35Q89.

Key words and phrases. mean field games, optimal control.

This research was supported in part by National Science Foundation through NSF Grant DMS-2045027.

space dimension $d = 1$. In this case the problem can be viewed as a static, “one-shot” game in which a player starting at a point x only has to choose a point y to minimize the cost

$$\frac{(x - y)^2}{2T} + m(y, T).$$

There is nothing particularly important about the value of T , so we will set $T = 1/2$, and we will denote by $f(y)$ the final density $m(y, T)$. Hence the cost to a player starting at point x and choosing to move to point y is given by

$$(1.2) \quad J(x, y, f) = (x - y)^2 + f(y),$$

where f can in principle be any probability density function, i.e. a non-negative measurable function such that $\int_{-\infty}^{\infty} f(x) dx = 1$. To avoid dealing with too many measure-theoretic issues, we will impose the restriction that f must be continuous.

The initial distribution will be given by $m = \sum_{j=1}^n a_j \delta_{x_j}$ with $a_1 + \dots + a_n = 1, a_j \geq 0$ and x_1, \dots, x_n distinct points in \mathbb{R} . Thus m is an empirical measure in which a_j is the proportion of players initially located at each x_j . Without loss of generality, we will assume $x_1 < \dots < x_n$, i.e. we put the points in the initial measure in order from left to right.

In this context, a *Nash equilibrium* is a measure π on $\mathbb{R} \times \mathbb{R}$ such that:

- (1) $\pi(A \times \mathbb{R}) = m(A) = \sum_{j=1}^n a_j \delta_{x_j}(A)$, where $\delta_{x_j}(A) = \begin{cases} 1 & \text{if } x_j \notin A \\ 0 & \text{if } x_j \in A \end{cases}$, i.e. first marginal of π is m ,
- (2) $\pi(\mathbb{R} \times B) = \int_B f(x) dx$ for a continuous density f , i.e. the second marginal of π is the density f , and
- (3) for π - a.e. (x, y) , $y \in \operatorname{argmin} J(x, \cdot, f)$, i.e. π couples points x and y such that y is an optimal move starting from x .

In other words, given the cost function J , a density function f is chosen to possibly be an equilibrium measure. With the knowledge of this function f , the players choose the optimal strategy for themselves, resulting in some final distribution. If this final distribution matches f , then f is in fact an equilibrium. The idea of a “Self-Fulfilling Prophecy” can be helpful to understand this: f is “prophesied” to be an equilibrium measure, and, if it turns out to result in an equilibrium, then f in a sense “fulfilled its prophecy.” The below diagram demonstrates this phenomenon.




FIGURE 1. Mean Field Games Diagram

In this article, our purpose is two-fold:

- (1) prove there exists a unique Nash equilibrium, and
- (2) compute solutions using a numerical method.

As far as we know, this constitutes the first result on existence and uniqueness of solutions to first-order mean field game with singular measures. Our approach relies on the fact that we take an entirely discrete measure, which allows us to rewrite the problem in an equivalent finite-dimensional formulation. Intuitively, all we need to compute is how each Dirac mass $a_j \delta_{x_j}$ in the initial distribution will “fan out” into a density of total mass a_j centered around point x_j . As explained below, the precise shape of this density is known through a priori considerations. This approach makes the problem amenable to classical methods. We believe it also provides geometric insight for what is going on more generally in density-penalized mean field games of first order.

Remark 1.1. *The main results of this article were first announced in the second author’s undergraduate thesis [Zim24].*

2. REFORMULATION OF THE EQUILIBRIUM PROBLEM

Let π be an equilibrium, and let f be the corresponding final density.

Definition 2.1. *The support of π , $\text{supp } \pi$, is the set of all (x, y) such that, for all $\epsilon > 0$, $\pi(B_\epsilon(x, y)) > 0$, where $B_\epsilon(x, y) = \{(z, w) \mid \sqrt{(x - z)^2 + (y - w)^2} < \epsilon\}$, or, equivalently, $\pi(U) > 0$ for every open set containing (x, y) .*

The support of f , $\text{supp } f$, is the closure of the set of all y such that $f(y) > 0$.

Lemma 2.2. *If $(x, y) \in \text{supp } \pi$, then $x = x_j$ for some j and $y \in \text{supp } f$. Conversely, if $y \in \text{supp } f$, then there exists x_j such that $(x_j, y) \in \text{supp } \pi$.*

Proof. If $(x, y) \in \text{supp } \pi$, then for all $\epsilon > 0$ we have $m(B_\epsilon(x)) = \pi(B_\epsilon(x) \times \mathbb{R}) > 0$ and $\int_{B_\epsilon(y)} f(z) dz = \pi(\mathbb{R} \times B_\epsilon(y)) > 0$. It follows that $x = x_j$ for some j , and $y \in \text{supp } f$ since for every $\epsilon > 0$ there exists $z \in B_\epsilon(y)$ such that $f(z) > 0$.

Conversely, suppose $y \in \text{supp } f$. For every $\epsilon > 0$, $\pi(\mathbb{R} \times B_\epsilon(y)) = \int_{B_\epsilon(y)} f > 0$. Since $\pi(\mathbb{R} \times B_\epsilon(y)) = \sum_{j=1}^n \pi(\{x_j\} \times B_\epsilon(y))$, for each ϵ there is a j such that $\pi(\{x_j\} \times B_\epsilon(y)) > 0$. We can then find a sequence $\epsilon_k \downarrow 0$ and a fixed j such that $\pi(\{x_j\} \times B_{\epsilon_k}(y)) > 0$ for all k , which implies $(x_j, y) \in \text{supp } \pi$. \square

Now if $(x_j, y) \in \text{supp } \pi$, the definition of equilibrium implies $y \in E_j$ where

$$E_j = \operatorname{argmin} \left\{ y : (x_j - y)^2 + f(y) \right\},$$

i.e. E_j is the set of minimizers y for the cost $J(x_j, y, f)$. By Lemma 2.2, $\text{supp } f \subset \bigcup_{j=1}^n E_j$; the intuitive meaning is that every player must move into one of the sets E_j . On the other hand, for $y \in E_j$ we have $f(y) = C_j - (x_j - y)^2$, where

$$C_j = \min \left\{ y : (x_j - y)^2 + f(y) \right\}.$$

Thus, f is completely determined by C_j and E_j , and these in turn are coupled by the definition of E_j . So how do we determine C_j and E_j ? A first clue is the following proposition, whose proof is elementary:

Proposition 2.3. $E_j = \left\{ y : C_j - (x_j - y)^2 \geq (C_k - (x_k - y)^2)_+ \quad \forall k \right\}$, where $x_+ := \max\{x, 0\}$.

Proof. If $y \in E_j$, then $(x_j - y)^2 + f(y) = C_j$, and at the same time $(x_k - y)^2 + f(y) \geq C_k$ for all k by definition of C_k . Recalling that $f \geq 0$, we see that $C_j - (x_j - y)^2 = f(y) \geq (C_k - (x_k - y)^2)_+$ for all k .

Conversely, suppose $y \notin E_j$. If $y \in E_k$ for some other k , then $C_j - (x_j - y)^2 < f(y) = C_k - (x_k - y)^2$. If $y \notin \bigcup_1^n E_k$, then $f(y) = 0$, hence $C_j - (x_j - y)^2 < 0$. It follows that $C_j - (x_j - y)^2 < (C_k - (x_k - y)^2)_+$ for at least one k . \square

Corollary 2.4. Let $f_j(y) = C_j - (x_j - y)^2$. Then $f(y) = \max\{0, f_1(y), \dots, f_n(y)\}$ and $E_j = \{y : f(y) = f_j(y)\}$.

Intuitively, all the players initially concentrated at x_j should spread out according to the density function f_j over the set E_j . We now make this intuition rigorous.

Proposition 2.5. $E_j = \{y : (x_j, y) \in \text{supp } \pi\}$, and $\int_{E_j} f = \int_{E_j} f_j = a_j$.

Proof. If $(x_j, y) \in \text{supp } \pi$, then by definition of equilibrium, it follows that $y \in E_j$. Conversely, suppose $y \in E_j$. As explained below in Lemma 3.1, E_j is an interval, and if y is in the interior of that interval, then $y \notin \bigcup_{k \neq j} E_k$. By Corollary 2.4 and Lemma 2.2 we deduce that $(x_j, y) \in \text{supp } \pi$. If y is on the boundary of E_j , there is a sequence $y_k \rightarrow y$ with y_k in the interior of E_j , so $(x_j, y_k) \in \text{supp } \pi$ and thus $(x_j, y) \in \text{supp } \pi$.

To prove the remaining statements, start with the following inequality:

$$(2.1) \quad a_j = m(\{x_j\}) = \pi(\{x_j\} \times \mathbb{R}) = \pi(\{x_j\} \times E_j) \leq \pi(\mathbb{R} \times E_j) = \int_{E_j} f.$$

Summing over j , we get

$$(2.2) \quad 1 = \sum_{j=1}^n a_j \leq \sum_{j=1}^n \int_{E_j} f = \int f = 1.$$

It follows that $\int_{E_j} f = a_j$, as desired. \square

With these results, we can now reformulate the problem using only the parameters C_1, \dots, C_n .

Definition 2.6. For a given vector $C = (C_1, \dots, C_n) \in \mathbb{R}^n$,

- let $f_j^{(C)}(x) = C_j - (x - x_j)^2$ for $j = 1, \dots, n$;
- let $f^{(C)}(x) = \max\{f_1^{(C)}(x), \dots, f_n^{(C)}(x), 0\}$;
- let $E_j^{(C)} = \left\{ x : f_j^{(C)}(x) = f^{(C)}(x) \right\}$ for $j = 1, \dots, n$;
- let $F_j(C) = \int_{E_j^{(C)}} f^{(C)}(x) dx$ when $E_j^{(C)}$ is nonempty, and let $F_j(C) = 0$ if $E_j^{(C)}$ is empty, for $j = 1, \dots, n$; and
- let $F(C) = (F_1(C), \dots, F_n(C))$.

We remark that, in what follows, $f_j^{(C)}$, $f^{(C)}$, and $E_j^{(C)}$ will often be written more cleanly as f_j , f , and E_j , when it is clear from context that C is fixed.

To better understand these definitions, consider the following image: The graph of $f(x)$ is the




FIGURE 2. Visualization of the density $f(x)$

solid-color part (i.e. the maximum). Each individual curve $f_j(x)$ can be thought of the density after the population sitting at point x_j “fans out” and forms a “bubble” surrounding that point.

With these definitions, we now state the problem we wish to solve as follows:

Problem 2.7. *Given a vector $a = (a_1, \dots, a_n)$, whose components are the weights in the discrete measure $m = \sum_{j=1}^n a_j \delta_{x_j}$, we want to find a vector C such that $F(C) = a$.*

By the arguments we have just given, if $f(x)$ is an equilibrium density, then $f(x) = f^{(C)}(x)$ where C is the solution to Problem 2.7. Conversely, suppose C is the solution to Problem 2.7 and let $f(x) = f^{(C)}(x)$. To see that $f(x)$ is an equilibrium density, first define a function $T : \mathbb{R} \rightarrow \mathbb{R}$ that sends every point in the interior of $E_j = E_j^{(C)}$ to the point x_j . Notice that m is now the *push-forward* of the density through T . (We say a measure ν is the push-forward of a measure μ through a function T if $\nu(B) = T\#\mu(B) := \mu(T^{-1}(B))$; in this case $\nu = m$ and $d\mu = f(x) dx$.) Set π to be the push-forward of the density f by the function $y \mapsto (T(y), y)$. It follows that π is an equilibrium.

We now state our main result:

Theorem 2.8 (Existence and uniqueness). *For each a such that $a_j > 0$ for $j = 1, \dots, n$, there exists a unique C such that $F(C) = a$.*

Theorem 2.8 immediately implies the existence and uniqueness of a Nash equilibrium for any game where the initial distribution is discrete. We will prove Theorem 2.8 in two sections: Section 4 shows existence, and Section 5 proves uniqueness. These proofs rely on our analysis of $F(C)$ in Section 3, in which we establish certain formulas for $F_j(C)$ and its derivatives, prove continuity and coercivity, and the convexity of the effective domain of F (namely, the set of all C for which $F_j(C) > 0$ for all j).

3. STRUCTURE OF $F(C)$

The proof of Theorem 2.8 is based on various properties of $F(C)$, as given in Definition 2.6. Whenever C is fixed, we will usually suppress the argument C and simply write $f_j(x)$, $f(x)$, and E_j instead of $f_j^{(C)}(x)$, $f^{(C)}(x)$, and $E_j^{(C)}$. In Section 3.1, we study the set E_j , namely the set over which $f(x) = f_j(x)$. We also study the effective domain of $F(C)$, over which we look for solutions to Problem 2.7, and we prove it is convex. In Section 3.2 we seek more or less explicit formulas for $F_j(C)$ and its derivatives. Section 3.3 shows that F is a *coercive* function on certain domains, while Section 3.4 shows that it is continuous everywhere. In reading these technical results, it will be useful to recall that we have ordered the points x_1, \dots, x_n from left to right, i.e. $x_1 < \dots < x_n$ without loss of generality.

3.1. Properties of E_j . We begin with the following claim:

Lemma 3.1. *E_j is convex and compact, hence either empty, a singleton, or a closed, bounded interval.*

Proof. First note that f_j is concave and continuous. Then observe that for every i , $f_j - f_i$ is affine, hence also concave and continuous. Indeed,

$$(3.1) \quad f_i(x) - f_j(x) = C_i - C_j + 2(x_i - x_j) \left(x - \frac{x_i + x_j}{2} \right).$$

We can write E_j is the intersection of super-level sets

$$(3.2) \quad E_j = \cap_i \{x : f_j(x) - f_i(x) \geq 0\} \cap \{x : f_j(x) \geq 0\},$$

which is therefore convex and closed by the first three observations. Moreover, it is compact because $\{x : f_j(x) \geq 0\} = \{x : (x - x_j)^2 \leq C_j\}$ is compact. The proof is complete. \square

In light of this, we introduce the following notation:

Definition 3.2. *In the case that E_j is an interval, we define α_j and β_j as the left and right end points of E_j , i.e. $[\alpha_j, \beta_j] = E_j$. We denote the length of E_j by $\|E_j\| = \beta_j - \alpha_j$.*

Before proceeding, we observe that Definition 2.6 leaves open the possibility that $F_j(C) = 0$. Since we are trying to solve $F(C) = a$ where all the components of a are positive, the true domain of interest is given as follows:

Definition 3.3. $\mathbb{G} = \{C_j : F(C_j) > 0 \text{ for } j = 1, \dots, n\}$.

Corollary 3.4. *If $C \in \mathbb{G}$, then E_j is an interval for all $j = 1, \dots, n$.*

Proof. First, by definition of \mathbb{G} , if $C \in \mathbb{G}$, then $F_j(C) > 0$ for all $j = 1, \dots, n$. Further, by Definition 2.6, it is clear that, in order for $F_j(C)$ to be greater than 0, then E_j cannot be empty or a singleton. Thus, by 3.1, if $F(C) \in \mathbb{G}$, then E_j must be an interval for all $j = 1, \dots, n$. \square

The most crucial points of interest will be the intersections between parabolas, namely where $f_i(x) = f_j(x)$, or $C_i - (x - x_i)^2 = C_j - (x - x_j)^2$. We define the unique solution by

$$(3.3) \quad \gamma_{ij} = \frac{C_i - C_j}{2(x_j - x_i)} + \frac{x_j + x_i}{2}, i \neq j.$$

By Equation (3.1), $f_i - f_j$ is increasing when $i > j$ and decreasing when $i < j$. It follows that if $x > \gamma_{ij}$ for some $i > j$ or if $x < \gamma_{ij}$ for some $i < j$, then $f_i(x) > f_j(x)$ and thus $x \notin E_j$. We deduce that $\beta_j \leq \min\{\gamma_{ij} : i > j\}$ and $\alpha_j \geq \max\{\gamma_{ij} : i < j\}$. Conversely, if $\max\{\gamma_{ij} : i < j\} \leq x \leq \min\{\gamma_{ij} : i > j\}$, then we deduce $f_j(x) \geq f_i(x)$ for all i . It follows that E_j is the intersection of the interval $\max\{\gamma_{ij} : i < j\} \leq x \leq \min\{\gamma_{ij} : i > j\}$ with the set where $f_j(x) \geq 0$, i.e. the interval $x_j - \sqrt{C_j} \leq x \leq x_j + \sqrt{C_j}$. We are especially interested in the case when E_j is not empty.

The content of the following lemma is essentially to say that when $C \in \mathbb{G}$ then Figure 2 is exactly the right visualization of the function $f^{(C)}(x)$, in the sense that the points of intersections between parabolas can be ordered from left to right. In particular, in order to locate $E_j^{(C)}$, it is sufficient to identify where $f_j^{(C)}(x)$ and its nearest neighbors $f_{j+1}^{(C)}(x)$ and $f_{j-1}^{(C)}(x)$ intersect.

Lemma 3.5. *For $C \in \mathbb{G}$, we have $\min\{\gamma_{ij} : i > j\} = \gamma_{j(j+1)}$ and $\max\{\gamma_{ij} : i < j\} = \gamma_{(j-1)j}$. Thus α_j and β_j are continuous functions of $C \in \mathbb{G}$ given by $\alpha_j = \max\{\gamma_{(j-1)j}, x_j - \sqrt{C_j}\}$ and $\beta_j = \min\{\gamma_{j(j+1)}, x_j + \sqrt{C_j}\}$, which results in the following formulas:*

$$(3.4) \quad \alpha_j = \begin{cases} \frac{C_{j-1} - C_j}{2(x_j - x_{j-1})} + \frac{x_j + x_{j-1}}{2} & \text{if } f_j(\alpha_j) > 0, \\ x_j - \sqrt{C_j}, & \text{if } f_j(\alpha_j) = 0, \end{cases} \quad \beta_j = \begin{cases} \frac{C_j - C_{j+1}}{2(x_{j+1} - x_j)} + \frac{x_{j+1} + x_j}{2}, & \text{if } f_j(\beta_j) > 0, \\ x_j + \sqrt{C_j}, & \text{if } f_j(\beta_j) = 0. \end{cases}$$

Moreover, they satisfy the ordering property $\beta_{j-1} \leq \alpha_j < \beta_j$ for all j (where $\beta_0 := -\infty$ for convenience), and $\beta_{j-1} = \alpha_j$ if and only if $f_j(\gamma_{(j-1)j}) \geq 0$.

Proof. We claim that $\min\{\gamma_{ij} : i > j\} = \gamma_{j(j+1)}$. Suppose to the contrary that there exists some $i > j + 1$ such that $\gamma_{ij} < \gamma_{j(j+1)}$. We will now show that E_{j+1} is empty. Indeed, if $x < \gamma_{j(j+1)}$, then we have $f_j(x) > f_{j+1}(x)$, so $x \notin E_{j+1}$. On the other hand, since $\gamma_{j(j+1)} > \gamma_{ij}$, it follows that $f_i(\gamma_{j(j+1)}) > f_j(\gamma_{j(j+1)}) = f_{j+1}(\gamma_{j(j+1)})$. Then since $f_i - f_{j+1}$ is increasing, we see that $f_i(x) > f_{j+1}(x)$ for all $x \geq \gamma_{j(j+1)}$, so E_{j+1} is empty. This contradicts the assumption that $C \in \mathbb{G}$, so the claim follows. The mirror image of this argument shows that $\max\{\gamma_{ij} : i < j\} = \gamma_{(j-1)j}$.

We have already established that E_j is the intersection of the interval $\max\{\gamma_{ij} : i < j\} \leq x \leq \min\{\gamma_{ij} : i > j\}$ with the set where $f_j(x) \geq 0$, i.e. the interval $x_j - \sqrt{C_j} \leq x \leq x_j + \sqrt{C_j}$. From here it is straightforward to deduce the formulas in Equation (3.4). The ordering property $\beta_{j-1} \leq \alpha_j$ follows from $\beta_{j-1} \leq \gamma_{(j-1)j} \leq \alpha_j$, and $\alpha_j < \beta_j$ because E_j is not empty. If $\beta_{j-1} = \alpha_j$, then they both must equal $\gamma_{(j-1)j}$, which implies that $f_j(\gamma_{(j-1)j}) \geq 0$. Conversely, if $f_j(\gamma_{(j-1)j}) = f_{j-1}(\gamma_{(j-1)j}) \geq 0$, then we deduce that $x_j - \sqrt{C_j} \leq \gamma_{(j-1)j} \leq x_{j-1} + \sqrt{C_{j-1}}$, and this implies $\beta_{j-1} = \alpha_j = \gamma_{(j-1)j}$. \square

Corollary 3.6. *If $C \in \mathbb{G}$, then the length of $E_j(C)$ is given by*

$$(3.5) \quad \|E_j\| = k_j(C) = \min\{\gamma_{j,j+1}(C), x_j + \sqrt{C_j}\} - \max\{\gamma_{j-1,j}(C), x_j - \sqrt{C_j}\}.$$

In fact, $C \in \mathbb{G}$ if and only if $k_j(C) > 0$ for all j . In particular, \mathbb{G} is a convex set.

Proof. If $C \in \mathbb{G}$, use Lemma 3.5 to see that $\|E_j\| = \beta_j - \alpha_j = k_j(C) > 0$. If $C \notin \mathbb{G}$, then at least one E_j must be a singleton or empty, i.e. $f_j(x)$ is never the maximum in $f(x) = \max\{f_1(x), \dots, f_n(x), 0\}$. We deduce that one of the following inequalities must be true:

- (1) $\gamma_{j(j+1)} < \gamma_{(j-1)j}$
- (2) $\gamma_{(j-1)j} > x_j + \sqrt{C_j}$

$$(3) \quad \gamma_{j(j+1)} < x_j - \sqrt{C_j}$$

hence $k_j(C) \leq 0$.

Since $k_j(C)$ is a concave function and $\mathbb{G} = \{C : k_j(C) > 0 \forall j\}$, it follows that \mathbb{G} is convex. \square

We make one more simple observation about \mathbb{G} .

Lemma 3.7. \mathbb{G} is non-empty.

Proof. To show \mathbb{G} is non-empty, by 3.3, we need a C such that $F_j(C) > 0$ for all $j = 1, \dots, n$. So, let $\epsilon > 0$. Choose ϵ small enough so that, if $C_j = \epsilon$ for all $j = 1, \dots, n$, then $f_j(\alpha_j) = 0 = f_j(\beta_j)$ for all $j = 1, \dots, n$, i.e. no f_j intersects any $f_i, i \neq j$ above the x -axis. Then $E_j = [x_j - \sqrt{\epsilon}, x_j + \sqrt{\epsilon}]$ for all $j = 1, \dots, n$. Thus, we have

$$(3.6) \quad F_j(C) = \frac{4}{3}\epsilon^{3/2},$$

which is greater than 0. Thus, \mathbb{G} is non-empty. \square

Thus, the set \mathbb{G} is both non-empty and convex, which will be useful in the proof of uniqueness of equilibria.

3.2. Computing $F_j(C)$ and its Derivatives. In this section, we want to calculate $F_j(C)$ and all of its derivatives for $C \in \mathbb{G}$. So, we have the following:

Proposition 3.8. $F_j(C) = C_j(\beta_j - \alpha_j) - \frac{1}{3}(\beta_j - x_j)^3 + \frac{1}{3}(\alpha_j - x_j)^3$ for every $C \in \mathbb{G}$.

Proof. We can compute this explicitly:

$$(3.7) \quad F_j(C) = \int_{\alpha_j}^{\beta_j} C_j - (x - x_j)^2 dx = C_j(\beta_j - \alpha_j) - \frac{1}{3}(\beta_j - x_j)^3 + \frac{1}{3}(\alpha_j - x_j)^3.$$

In particular, taking into account the different possibilities for α_j and β_j (see 3.2):

$$F_j(C) = \int_{\alpha_j}^{\beta_j} C_j - (x - x_j)^2 dx = C_j(\beta_j - \alpha_j) - \frac{1}{3}(\beta_j - x_j)^3 + \frac{1}{3}(\alpha_j - x_j)^3$$

If $f_j(\alpha_j), f_j(\beta_j) > 0$,

$$F_j(C) = C_j \left(\frac{C_j - C_{j+1}}{2(x_{j+1} - x_j)} + \frac{x_{j+1} + x_j}{2} \right) - \frac{1}{3} \left(\frac{C_j - C_{j+1}}{2(x_{j+1} - x_j)} + \frac{x_{j+1} + x_j}{2} - x_j \right)^3 \\ - C_j \left(\frac{C_{j-1} - C_j}{2(x_j - x_{j-1})} + \frac{x_j + x_{j-1}}{2} \right) + \frac{1}{3} \left(\frac{C_{j-1} - C_j}{2(x_j - x_{j-1})} + \frac{x_j + x_{j-1}}{2} - x_j \right)^3,$$

or if $f_j(\alpha_j), f_j(\beta_j) = 0$,

$$F_j(C) = \left(C_j(x_j + \sqrt{C_j}) - \frac{1}{3}(x_j + \sqrt{C_j} - x_j)^3 \right) - \left(C_j(x_j - \sqrt{C_j}) - \frac{1}{3}(x_j - \sqrt{C_j} - x_j)^3 \right) = \frac{4}{3}C_j^{3/2}.$$

The other two cases, when $f_j(\alpha_j) > 0, f_j(\beta_j) = 0$ and when $f_j(\alpha_j) = 0, f_j(\beta_j) > 0$, are straightforward variations of these two cases. \square

Remark 3.9. These calculations show that, in particular, $0 \leq F_j(C) \leq \frac{4}{3}C_j^{3/2}$ for any C . This will be useful later.

Proposition 3.10. $\frac{\partial F_j(C)}{\partial C_j}(C) = \|E_j\| + \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} + \frac{f_j(\alpha_j)}{2(x_j - x_{j-1})}$ for every $C \in \mathbb{G}$.

Proof. With $F(C)$ now calculated, we can easily calculate $\frac{\partial F_j(C)}{\partial C_j}$, taking into account the different possibilities for α_j and β_j as above:

(3.8)

If $f_j(\alpha_j), f_j(\beta_j) > 0$,

$$\begin{aligned} \frac{\partial F_j(C)}{\partial C_j} &= \left(\frac{C_j - C_{j+1}}{2(x_{j+1} - x_j)} + \frac{x_{j+1} + x_j}{2} \right) + \left(\frac{C_j}{2(x_{j+1} - x_j)} \right) - \frac{1}{2(x_{j+1} - x_j)} \left(\frac{C_j - C_{j+1}}{2(x_{j+1} - x_j)} + \frac{x_{j+1} - x_j}{2} \right)^2 \\ &\quad - \left(\frac{C_{j-1} - C_j}{2(x_j - x_{j-1})} + \frac{x_j + x_{j-1}}{2} \right) + \left(\frac{C_i}{2(x_j - x_{j-1})} \right) - \frac{1}{2(x_j - x_{j-1})} \left(\frac{C_{j-1} - C_j}{2(x_j - x_{j-1})} - \frac{x_j - x_{j-1}}{2} \right)^2 \\ &= \|E_j\| + \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} + \frac{f_j(\alpha_j)}{2(x_j - x_{j-1})}, \end{aligned}$$

or if $f_j(\alpha_j), f_j(\beta_j) = 0$,

$$\frac{\partial F_j(C)}{\partial C_j} = 2C_j.$$

Again, the other two cases are straightforward adaptations. In any case, we have

$$(3.9) \quad \frac{\partial F_j(C)}{\partial C_j} = \|E_j\| + \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} + \frac{f_j(\alpha_j)}{2(x_j - x_{j-1})},$$

which is a continuous function of C . Notice that this holds for every case; for example, if $f_j(\alpha_j), f_j(\beta_j) = 0$, then those components of the partial derivative would be 0, leaving $\|E_j\|$, which is in fact $2C_j$ in that case. In particular, the partial derivative is seen to be continuous everywhere, since as we approach the boundary cases where $f_j(\alpha_j)$ or $f_j(\beta_j)$ goes to zero, the formula still holds in the limit. \square

Theorem 3.11. For every $C \in \mathbb{G}$, $\frac{\partial F_j(C)}{\partial C_{j-1}} = \frac{f_j(\alpha_j)}{2(x_j - x_{j-1})}$, $\frac{\partial F_j(C)}{\partial C_{j+1}} = \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)}$, and $\frac{\partial F_j(C)}{\partial C_i} = 0$ if $i \notin \{j-1, j, j+1\}$.

Proof. Suppose $\beta_{j-1} < \alpha_j$. From Lemma 3.5, we see that $f_j(\gamma_{(j-1)j}) < 0$ and thus E_j does not depend on C_{j-1} for small perturbations of C . Similarly, if $\beta_j < \alpha_{j+1}$, then E_j does not depend on C_{j+1} for small perturbations of C . More generally, we deduce that E_j never depends on C_i for $C \in \mathbb{G}$ if $i \notin \{j-1, j, j+1\}$.

So we only consider the case when $f_j(\gamma_{(j-1)j}), f_j(\gamma_{j(j+1)}) \geq 0$. In this case Lemma 3.5 implies that $\alpha_j = \gamma_{(j-1)j}$ and $\beta_j = \gamma_{j(j+1)}$. The case $f_j(\gamma_{(j-1)j}) = f_j(\alpha_j) > 0$ is stable under small perturbations of C , and we differentiate the formula from Proposition 3.8 to get

(3.10)

$$\frac{\partial F_j(C)}{\partial C_{j-1}} = -\frac{1}{2(x_j - x_{j-1})} \left(\left(\frac{C_j - C_{j-1}}{2(x_{j-1} - x_j)} + \frac{x_{j-1} - x_j}{2} \right)^2 - C_j \right) = \frac{f_j(\gamma_{(j-1)j})}{2(x_j - x_{j-1})} = \frac{f_j(\alpha_j)}{2(x_j - x_{j-1})}.$$

As $f_j(\gamma_{(j-1)j}) \rightarrow 0$, this formula converges to 0, which agrees with the case $f_j(\gamma_{(j-1)j}) < 0$. Similarly, the case $f_j(\gamma_{j(j+1)}) = f_j(\beta_j) > 0$ is stable under small perturbations of C , and we

differentiate the formula from Proposition 3.8 to get

$$(3.11) \quad \frac{\partial F_j(C)}{\partial C_{j+1}} = \frac{1}{2(x_j - x_{j+1})} \left(\left(\frac{C_{j+1} - C_j}{2(x_j - x_{j+1})} - \frac{x_j - x_{j+1}}{2} \right)^2 - C_j \right) = \frac{f_j(\gamma_{j(j+1)})}{2(x_{j+1} - x_j)} = \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)}.$$

As $f_j(\gamma_{j(j+1)}) \rightarrow 0$, this formula converges to 0, which agrees with the case $f_j(\gamma_{j(j+1)}) < 0$. This establishes the existence and continuity of the partial derivatives. \square

We notice the following, deduced from the formulas in Lemma 3.5 and the result of Theorem 3.11:

Remark 3.12.

$$\frac{\partial F_i(C)}{\partial C_j} = \frac{\partial F_j(C)}{\partial C_i}$$

for every $C \in \mathbb{G}$ and for every $i, j \in \{1, \dots, n\}$.

This implies that F is in fact a gradient, i.e. there exists a smooth function $\Phi : \mathbb{G} \rightarrow \mathbb{R}$ such that $F(C) = \nabla \Phi(C)$ for all $C \in \mathbb{G}$. This is a remarkable coincidence, as it is already well-known that the mean field game (1.1) is a potential game, meaning that there exists a potential whose critical points are Nash equilibria. However, it must be emphasized that usually this potential is defined on the space of (paths defined on) probability measures, and its derivative is supposed to be the cost paid to individual players. See e.g. [BCS17, CH17, Gra24], [LL07, Section 2.6], [CD17] and references therein. Here the context is quite different, as we have significantly transformed the problem, so the “potential” in this case does not have precisely the same meaning as in these references.

3.3. Coercivity. We say that the vector field $F(C)$ is coercive provided that $\lim_{\|C\| \rightarrow \infty} \frac{F(C) \cdot C}{\|C\|} = \infty$,

where $\|C\| = (C_1^2 + \dots + C_n^2)^{1/2}$ is the norm of the vector C . This is the usual condition for a vector field to be surjective, as in the Minty-Browder Theorem [Bro67]. There is no chance for $F(C)$ to be coercive on all of \mathbb{R}^n , since $F_j(C) = 0$ whenever $C_j \leq 0$. However, this will not matter, as we are always aiming at restricting to vectors C with non-negative components. Instead, we will use the following substitute for coercivity to prove existence of solutions.

Lemma 3.13. *There exist constants $\delta \in (0, 1)$ and $M > 0$ such that if C is any vector in \mathbb{R}^n with $C_j = \max\{C_1, \dots, C_n\} \geq M$, then $F_j(C) \geq \delta C_j$.*

Proof. Choose a C_j such that $C_j = \max\{C_1, \dots, C_n\} \geq M$. We now seek to estimate $F_j(C)$. Recall that, from the discussion before Lemma 3.5, E_j is the intersection of the interval $\max\{\gamma_{ij} : i < j\} \leq x \leq \min\{\gamma_{ij} : i > j\}$ with the interval $x_j - \sqrt{C_j} \leq x \leq x_j + \sqrt{C_j}$. By the maximality of C_j ,

$$\begin{aligned} \max\{\gamma_{ij} : i < j\} &\leq \max \left\{ \frac{x_j + x_i}{2}, i < j \right\} = \frac{x_j + x_{j-1}}{2}, \\ \min\{\gamma_{ij} : i > j\} &\leq \min \left\{ \frac{x_j + x_i}{2}, i > j \right\} = \frac{x_j + x_{j+1}}{2}. \end{aligned}$$

We can assume that

$$M \geq \max \left\{ \left(\frac{x_{i+1} - x_i}{2} \right)^2 : i = 1, \dots, n-1 \right\}.$$

Then it follows that

$$x_j - \sqrt{C_j} \leq \frac{x_j + x_{j-1}}{2}, \quad x_j + \sqrt{C_j} \geq \frac{x_j + x_{j+1}}{2}.$$

We can now conclude that $E_j = [\alpha_j, \beta_j]$ with

$$\alpha_j \leq \frac{x_{j-1} + x_j}{2}, \quad \beta_j \geq \frac{x_j + x_{j+1}}{2}.$$

This gives a lower bound on how close α_j and β_j can get to x_j . Thus,

$$F_j(C) \geq \int_{\frac{x_{j-1} + x_j}{2}}^{\frac{x_j + x_{j+1}}{2}} C_j - (x - x_j)^2 dx = C_j \frac{x_{j+1} - x_{j-1}}{2} - \frac{1}{3} \left(\left(\frac{x_{j+1} - x_j}{2} \right)^3 - \left(\frac{x_j - x_{j-1}}{2} \right)^3 \right).$$

Define

$$\bar{\delta} := \frac{\min_j \{x_{j+1} - x_{j-1}\}}{2}, \quad \bar{M} := \frac{1}{3} \max_j \left(\left(\frac{x_{j+1} - x_j}{2} \right)^3 - \left(\frac{x_j - x_{j-1}}{2} \right)^3 \right)_+,$$

where we recall $x_+ := \max\{x, 0\}$. Then

$$F_j(C) \geq \bar{\delta} C_j - \bar{M}.$$

Pick any $0 < \delta < \min\{\bar{\delta}, 1\}$ and assume $M \geq (\bar{\delta} - \delta)^{-1} \bar{M}$. If $C_j \geq M$, we get $F_j(C) \geq \delta C_j$, as desired. \square

We remark that Lemma 3.13 can be used to prove that F is coercive on $\mathbb{R}_{\geq 0}^n$, which is defined to be the set of all (x_1, \dots, x_n) such that $x_j \geq 0$ for every j . The reason is that if $C \in \mathbb{R}_{\geq 0}^n$ and $C_j = \max\{C_1, \dots, C_n\}$, then

$$F(C) \cdot C \geq F_j(C) C_j.$$

For the remaining details, let us first define the norm

$$\|x\|_\infty = \max\{|x_1|, \dots, |x_n|\}$$

and note that $\|x\|_\infty = \max\{x_1, \dots, x_n\}$ for $x \in \mathbb{R}_{\geq 0}^n$. Recall that

$$\|x\|_\infty \leq \|x\| \leq \sqrt{n} \|x\|_\infty.$$

Let M, δ be as in Lemma 3.13. If $C \in \mathbb{R}_{\geq 0}^n$ and $\|C\| \geq M\sqrt{n}$, then $\|C\|_\infty \geq M$ and so

$$F(C) \cdot C \geq F_j(C) C_j \geq \delta C_j^2 = \delta \|C\|_\infty^2 \geq \frac{\delta}{\sqrt{n}} \|C\|^2.$$

Thus F is coercive on $\mathbb{R}_{\geq 0}^n$. However, we also note that only Lemma 3.13, and not coercivity per se, will be used directly to prove the existence of solutions.

3.4. Continuity. With coercivity proved, we move to prove that $F(C)$ is continuous. Before doing so, we need to explore what happens in $E_j = \emptyset$.

Lemma 3.14. *Let $I \subseteq \{1, \dots, n\}$ and \mathcal{C}_I be the set of all C such that $E_j(C) = \emptyset$ for all $j \in I$. Then \mathcal{C}_I is open, i.e. for every $C \in \mathcal{C}_I$, there exists $\epsilon > 0$ such that, if $\|\tilde{C} - C\| < \epsilon$, then $\tilde{C} \in \mathcal{C}_I$.*

Proof. Let $C \in \mathcal{C}_I$. Define $\tilde{f}^{(C)}(x) = \max\{f_j^{(C)}(x) : j \in I\}$. Then $f^{(C)}(x) > \tilde{f}^{(C)}(x)$ for all x , because all of the sets $E_j(C)$, $j \in I$, are empty. Since all of the $f_j^{(C)}(x)$ are parabolas that go to $-\infty$ as $|x| \rightarrow \infty$, there exists $M > 0$ such that if $|x| > M$, then $f^{(C)}(x) - \tilde{f}^{(C)}(x) \geq 1$. Now $f^{(C)}(x) - \tilde{f}^{(C)}(x)$ is a continuous and positive function, hence it has a minimum $\delta > 0$ on the compact set $[-M, M]$. Without loss of generality, we can assume $\delta \leq 1$; in particular, we

have $f^{(C)}(x) - \tilde{f}^{(C)}(x) \geq \delta$ for every x . Now assume $|\tilde{C}_j - C_j| < \delta/2$ for $j = 1, 2, \dots, n$. Then $\left| f_j^{(\tilde{C})}(x) - f_j^{(C)}(x) \right| < \delta/2$ for every j and every x . It follows that $\left| f^{(\tilde{C})}(x) - f^{(C)}(x) \right| < \delta/2$ and $\left| \tilde{f}^{(\tilde{C})}(x) - \tilde{f}^{(C)}(x) \right| < \delta/2$, so

$$(3.12) \quad \begin{aligned} f^{(\tilde{C})}(x) - \tilde{f}^{(\tilde{C})}(x) &= f^{(\tilde{C})}(x) - f^{(C)}(x) + f^{(C)}(x) - \tilde{f}^{(C)}(x) + \tilde{f}^{(C)}(x) - \tilde{f}^{(\tilde{C})}(x) \\ &> -\delta/2 + \delta - \delta/2 = 0. \end{aligned}$$

It follows that all the sets $E_j(\tilde{C})$, $j \in I$, are empty, and so \tilde{C} must also be in \mathcal{C}_I . Thus, \mathcal{C}_I is open. \square

With this in mind, we can now prove continuity:

Theorem 3.15. *$F(C)$ is continuous on \mathbb{R}^n .*

Proof. It is sufficient to prove that each component function $F_j(C)$ is continuous. First, let us point out that we can remove certain components from consideration if the corresponding sets $E_j(C)$ are empty. Let I be the set of all indices j such that $E_j(C) = \emptyset$, so that $C \in \mathcal{C}_I$, the set defined in Lemma 3.14. Take a sequence of vectors $C^{(k)} \rightarrow C$. Then, for k sufficiently large, $C^{(k)} \in \mathcal{C}_I$ also. So $F_j(C^{(k)}) = F_j(C) = 0$ for every $j \in I$. In light of this, we can remove the indices in I and relabel the sequence $\{1, \dots, n\} \setminus I$, calling it $\{1, \dots, m\}$. Then $f(x)$ will still be the $\max\{f_1(x), f_2(x), \dots, f_m(x), 0\}$, as those indices which may have been removed will not affect the maximum. Thus, we can assume (without loss of generality) for the remainder of this proof that $E_j(C) \neq \emptyset$ for every j , and thus we will be able to appeal to previously derived explicit formulas for the component functions $F_j(C)$.

Let j be fixed but arbitrary. Since $E_j(C)$ is not empty, by Lemma 3.1 it is either an interval or a single point. Suppose that $E_j(C)$ is an interval. Choose a sequence of vectors $C^{(k)} \rightarrow C$. By Proposition 3.8, $F_j(C)$ is simply a polynomial in terms of C , and thus $F_j(C^{(k)}) \rightarrow F_j(C)$.

Lastly, suppose $E_j(C)$ is a single point. Then, by the contra-positive of Corollary 3.4, $F_j(C) = 0$. Again, we choose a sequence of vectors $C^{(k)} \rightarrow C$. So, we need to show $F_j(C^{(k)}) \rightarrow 0$. If $F_j(C^{(k)}) > 0$ then $E_j(C^{(k)})$ is an interval whose length is given by $k_j(C^{(k)})$ defined in Corollary 3.6. Further, since $k_j(C)$ is a continuous function of C and $E_j(C)$ is a single point, then $k_j(C^{(k)}) \rightarrow 0$ as $k \rightarrow \infty$. Therefore, whenever $E_j(C^{(k)})$ is an interval, its length shrinks to zero as $k \rightarrow \infty$, and since in any other case we have $F_j(C^{(k)}) = 0$, we deduce that $F_j(C^{(k)}) \rightarrow 0$ as $k \rightarrow \infty$. \square

We have now sufficiently explored various properties of $f(x)$ and $F(C)$, so we may now move onto our first goal:

4. PROVING EXISTENCE OF A SOLUTION

In this section, we seek to show that a solution C to the equation $F(C) = a$ exists for all a with positive components. Before we begin the proof, we state Brouwer's Fixed Point Theorem:

Lemma 4.1. *Every continuous function from a nonempty convex compact subset K of a Euclidean space to K itself has a fixed point.*

See e.g. [Flo03]. We now begin the main result:

Theorem 4.2. *Let $a = (a_1, \dots, a_n)$ with $a_j > 0$ for each j . Then there exists a solution C to the equation $F(C) = a$.*

Proof. Let a be given. Define $L : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}^n$ as $L_i(C) := (C_i + a_i - F_i(C))_+$ (recall $x_+ := \max\{x, 0\}$). We will use the norm

$$\|C\|_1 := |C_1| + \dots + |C_n| = C_1 + \dots + C_n.$$

It is useful to recall that

$$\|C\|_\infty \leq \|C\|_1 \leq n\|C\|_\infty.$$

By Lemma 3.13, there exist constants $\delta \in (0, 1)$ and $M > 0$ such that if $C_j = \|C\|_\infty \geq M$, then $F_j(C) \geq \delta C_j$. In this case,

$$L_j(C) \leq ((1 - \delta)C_j + a_j)_+ = (1 - \delta)C_j + a_j.$$

In general,

$$L_i(C) \leq C_i + a_i.$$

Summing over all i , we deduce that

$$\|L(C)\|_1 \leq \|C\|_1 + \|a\|_1 - \delta C_j.$$

Recall that $C_j = \|C\|_\infty \geq n^{-1}\|C\|_1$. Let $r > \|a\|_1 n/\delta$. Then we have

$$(4.1) \quad \|L(C)\|_1 \leq \|C\|_1 + \|a\|_1 - \delta n^{-1}\|C\|_1 < \|C\|_1 \quad \text{if } \|C\|_1 \geq r.$$

We now restrict to a compact, convex set. Let M_r be the set of all $x \in \mathbb{R}_{\geq 0}^n$ such that $\|x\|_1 \leq r$. Then M_r is closed and bounded, so M_r is compact, and it is clearly nonempty. Lastly, M_r is convex, as it is the intersection of two convex sets, namely, $\mathbb{R}_{\geq 0}^n$ and the closed unit ball for the 1-norm. Thus, by Lemma 4.1, any continuous function $g : M_r \rightarrow M_r$ has a fixed point.

Define a function

$$(4.2) \quad P(D) := \begin{cases} D, & \|D\|_1 \leq r \\ \frac{rD}{\|D\|_1}, & \|D\|_1 > r. \end{cases}$$

Note that P maps $\mathbb{R}_{\geq 0}^n$ into M_r . Finally, let $Q(C) := P(L(C))$ for $C \in M_r$. Then Q is continuous, as it is the composition of continuous functions, and it maps M_r to itself. Thus, by Lemma 4.1, Q has a fixed point $C \in M_r$. If $\|C\|_1 = r$, then

$$(4.3) \quad r = \|C\|_1 = \|Q(C)\|_1 = \|P(L(C))\|_1.$$

By construction of P , this implies

$$(4.4) \quad \|L(C)\|_1 \geq r.$$

But this contradicts (4.1). It follows that $\|C\|_1 < r$. Thus,

$$(4.5) \quad r > \|C\|_1 = \|Q(C)\|_1 = \|P(L(C))\|_1.$$

Hence, $P(L(C)) = L(C)$, so finally we have

$$(4.6) \quad C = Q(C) = P(L(C)) = L(C).$$

This implies that for each i ,

$$C_i = (C_i + a_i - F_i(C))_+.$$

If $C_i > 0$, this implies

$$C_i = C_i + a_i - F_i(C) \Rightarrow F_i(C) = a_i.$$

If $C_i = 0$, then $F_i(C) = 0$ and we get

$$0 = (a_i)_+ = a_i$$

which contradicts the assumption $a_i > 0$. Hence $C_i > 0$ for all i and $F(C) = a$, as desired. \square

5. PROVING UNIQUENESS OF A SOLUTION

By Theorem 4.2, we know that for any vector a with positive entries there exists a solution $C \in \mathbb{G}$ to $F(C) = a$. In this section, we prove that the solution is unique. To do so, we show that $F(C)$ is strictly monotone in a classical sense. It is interesting to note that monotonicity is often used to prove uniqueness in mean field games, the most commonly used notion being the ‘‘Lasry-Lions monotonicity condition’’ (cf. [LL07]). (For a discussion on alternative monotonicity conditions, see [GM23].) Here, the coupling is indeed precisely through the density of players, and therefore the Lasry-Lions monotonicity condition holds. However, it is interesting to note that the monotonicity of $F(C)$, a function of the parameters C_1, \dots, C_n , is not a direct consequence the Lasry-Lions monotonicity condition in any obvious way.

Lemma 5.1. *On \mathbb{G} , $F(C)$ is strictly monotone, i.e. for any $C, \tilde{C} \in \mathbb{G}$, $(F(C) - F(\tilde{C}))(C - \tilde{C}) > 0$.*

Proof. Recall that \mathbb{G} is convex by Corollary 3.6. To show F is strictly monotone, then, it suffices to show that

$$\sum_{i=1}^n \sum_{j=1}^n \frac{\partial F_i(C)}{\partial C_j} \cdot (v_i v_j) > 0 \quad \forall C, v \in \mathbb{R}^n \text{ with } v \neq 0.$$

Assume for the sake of notation that $v_0 = 0 = v_{n+1}$. By Lemma 2.11,

$$S := \sum_{i=1}^n \sum_{j=1}^n \frac{\partial F_i(C)}{\partial C_j} v_i v_j = \sum_{j=1}^n \left(\frac{\partial F_j(C)}{\partial C_j} (v_j)^2 + \frac{\partial F_j(C)}{\partial C_{j-1}} (v_{j-1} v_j) + \frac{\partial F_j(C)}{\partial C_{j+1}} (v_j v_{j+1}) \right).$$

Using the derivative formulas from (3.10),

$$\begin{aligned} S = \sum_{j=1}^n \left(\left\| E_j \right\| + \frac{f_j(\alpha_j)}{2(x_j - x_{j-1})} + \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} \right) (v_j)^2 \\ - \left(\frac{f_j(\alpha_j)}{2(x_j - x_{j-1})} \right) (v_{j-1} v_j) - \left(\frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} \right) (v_j v_{j+1}). \end{aligned}$$

Since $f_1(\alpha_1) = 0 = f_n(\beta_n)$, we can rewrite this as

$$\begin{aligned} S = \sum_{j=1}^n \left\| E_j \right\| v_j^2 + \sum_{j=2}^n \left(\frac{f_j(\alpha_j)}{2(x_j - x_{j-1})} (v_j)^2 - \left(\frac{f_j(\alpha_j)}{2(x_j - x_{j-1})} \right) (v_{j-1} v_j) \right) \\ + \sum_{j=1}^{n-1} \left(\frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} (v_j)^2 - \left(\frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} \right) (v_j v_{j+1}) \right). \end{aligned}$$

Since $f_{j+1}(\alpha_{j+1})$ is either $f_j(\beta_j)$ or 0, we can combine to get

$$\begin{aligned} S &= \sum_{j=1}^n \|E_j\| v_j^2 + \sum_{j=1}^{n-1} \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} (v_{j+1})^2 - \frac{2f_j(\beta_j)}{2(x_{j+1} - x_j)} (v_j v_{j+1}) + \frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} (v_j)^2 \\ &= \sum_{j=1}^n \|E_j\| v_j^2 + \sum_{j=1}^{n-1} \left(\frac{f_j(\beta_j)}{2(x_{j+1} - x_j)} \right) (v_{j+1} - v_j)^2 > 0. \end{aligned}$$

Thus, $F(C)$ is strictly monotone on \mathbb{G} . □

We now have everything we need to prove uniqueness.

Theorem 5.2. *The solution to the equation $F(C) = a$ is unique.*

Proof. Suppose, for the sake of contradiction, that $C_1, C_2 \in \mathbb{G}$ with $C_1 \neq C_2$ are both solutions to $F(C) = a$. Then $F(C_1) = F(C_2)$, so $(F(C_1) - F(C_2))(C_1 - C_2) = 0$. But this contradicts 5.1. Thus, the solution to $F(C)$ is unique. □

Remark 5.3. *As a result of Lemma 5.1, the potential Φ defined in Remark 3.12 is strictly convex on \mathbb{G} . The unique solution of $F(C) = a$ can be characterized as the unique minimizer of $\Phi(C) - C \cdot a$ over $C \in \mathbb{G}$.*

We have now shown that a unique solution to $F(C) = a$ exists and is unique. We now move on to numerical approximations.

6. NUMERICAL SIMULATIONS

We used Newton's Method to find approximate solutions to $F(C) = a$. By algebraically solving $\frac{4}{3}C_j^{3/2} = a_j$, we obtain an initial guess $C = C^{(0)}$ such that $F_j(C) \leq a_j$ for each j (see Remark 3.9). Then we define an approximating sequence $C^{(n)}$ by

$$(6.1) \quad C^{(n+1)} = C^{(n)} - DF \left(C^{(n)} \right)^{-1} \left(F \left(C^{(n)} \right) - a \right),$$

where $DF(C)$ is the Jacobian matrix computed in Section 3.2. The residual $\left\| F \left(C^{(n)} \right) - a \right\|$ can be computed easily; in practice it becomes vanishingly small after only a few iterations. Below we report on three simulations. In each case, the initial distribution $m = \sum_{j=1}^n a_j \delta_{x_j}$ is illustrated a set of n bars located at position x_j with height a_j . Then the final density $f(x)$ is then graphed over these bars. The function $f(x)$ is piecewise quadratic with the j th parabola centered at x_j and representing the spreading of the population that starts at x_j ; the area under this parabola above the x -axis equals a_j . This accords with our geometric intuition about the game: players spread out as much as possible so as to avoid areas of high population density, and the shape of the final density matches the quadratic cost $(x - y)^2$ of traveling from x to y .




FIGURE 3. Example 1: $x = (1, 2.25, 3, 3.75)$, $a = (0.25, 0.25, 0.25, 0.25)$

In the first simulation (Figure 3), we took an initial population uniformly distributed over four points: $m = \frac{1}{4}(\delta_1 + \delta_{2.25} + \delta_3 + \delta_{3.75})$. In this example, the leftmost quarter of the population is far enough away from rest that it can spread out without colliding with the others, whereas the other three are close enough to each other that players have less room to move, resulting in a narrower probability density.




FIGURE 4. Example 2: $x = (1, 2, \dots, 10)$, $a = \text{random}$




FIGURE 5. Example 3: $x = \text{random}$, $a = \text{random}$

In the second simulation (Figure 4), we let $x_j - x_{j-1}$ be constant and let a be random. (Note that we did not normalize the components of a to sum to 1, but this does not change the shape of the solution.) Notice how in this sample the probability density has a very interesting contour. For example, the “bubble” (i.e. piece of parabola) corresponding to $x = 9$ barely appears because it is crowded by neighboring “bubbles.” This demonstrates how the different weights locally change the shape of the density.

In the last simulation (Figure 5), the vectors x and a are random (and a is normalized to be a probability vector). Although the initial measure is highly irregular, the final density will still be smooth, though it appears less so as the number of random points are chosen. Indeed, as the proof of uniqueness in Section 5 suggests, the problem becomes more and more ill-conditioned as the points x_j get closer together, since the lengths of the intervals E_j necessarily get smaller.

As a final remark, we recall that $F = \nabla\Phi$ for some potential Φ , and that solving $F(C) = a$ is equivalent to minimizing $\Phi(C) - C \cdot a$ (see Remarks 3.12 and 5.3). Since Φ is strictly convex, one might hope that Newton’s method converges globally. However, it is not true in general that strict convexity implies global convergence for Newton’s method, as explained in this Stack Exchange discussion [new]; see also [BV04]. The proof of Lemma 5.1 suggests that the lower bound on the Hessian $\nabla^2\Phi(C)$ goes to zero as any of the lengths $\|E_j\|$ vanishes. One can infer that the problem is better conditioned if the points x_j are sufficiently spread out and/or the weights a_j are such that the equilibrium density has a uniform lower bound on the lengths of the intervals E_j . We see this in numerical experiments: if the distance between points x_j becomes too small, the algorithm appears not to converge.

7. CONCLUSION

In this paper we have studied a mean field game with final cost equal to the population density of players. We considered the case where the initial measure is discrete, which seems to be new in the literature. We showed that the problem reduces to a finite dimensional problem, which can prove is well-posed by classical methods. Our numerical simulations illustrate the “smoothing effect” that mean field games with density penalization are expected to have. For future research, it would be interesting to see if this approach could be extended to more general mean field games with density-dependent costs and discrete initial measures.

REFERENCES

- [ACD⁺20] Yves Achdou, Pierre Cardaliaguet, François Delarue, Alessio Porretta, Filippo Santambrogio, Pierre Cardaliaguet, and Alessio Porretta. An introduction to mean field game theory. *Mean Field Games: Cetraro, Italy 2019*, pages 1–158, 2020.
- [BCS17] Jean-David Benamou, Guillaume Carlier, and Filippo Santambrogio. Variational mean field games. In *Active Particles, Volume 1*, pages 141–171. Springer, 2017.
- [BFY13] Alain Bensoussan, Jens Frehse, and Phillip Yam. *Mean field games and mean field type control theory*. Springer, 2013.
- [Bro67] F.E. Browder. Existence and perturbation theorems for nonlinear maximal monotone operators in Banach spaces. *Bull. Amer. Math. Soc.*, 73:322–327, 1967.
- [BV04] Stephen Boyd and Lieven Vandenberghe. *Convex optimization*. Cambridge university press, 2004.
- [Car15] Pierre Cardaliaguet. Weak solutions for first order mean field games with local coupling. In *Analysis and Geometry in Control Theory and its Applications*, volume 11 of *Springer INdAM Series*, pages 111–158. Springer, 2015.
- [CD17] R Carmona and F Delarue. Probabilistic Theory of Mean Field Games: vol. I, Mean Field FBSDEs, Control, and Games. *Stochastic Analysis and Applications, Springer Verlag*, 2017.
- [CG15] Pierre Cardaliaguet and P. Jameson Graber. Mean field games systems of first order. *ESAIM: COCV*, 21(3):690–722, 2015.

- [CGPT15] Pierre Cardaliaguet, P. Jameson Graber, Alessio Porretta, and Daniela Tonon. Second order mean field games with degenerate diffusion and local coupling. *Nonlinear Differential Equations and Applications NoDEA*, 22(5):1287–1317, 2015.
- [CH17] Pierre Cardaliaguet and Saeed Hadikhanloo. Learning in mean field games: The fictitious play. *ESAIM: Control, Optimisation and Calculus of Variations*, 23(2):569–591, 2017.
- [Flo03] Monique Florenzano. *General equilibrium analysis: existence and optimality properties of equilibria*. Springer Science & Business Media, 2003.
- [G⁺14] Diogo A Gomes et al. Mean field games models—a brief survey. *Dynamic Games and Applications*, 4(2):110–154, 2014.
- [GM18] P. Jameson Graber and Alpár R. Mészáros. Sobolev regularity for first order mean field games. *Annales de l’Institut Henri Poincaré C, Analyse Non Linéaire*, 35(6):1557–1576, September 2018.
- [GM23] P. Jameson Graber and Alpár R. Mészáros. On monotonicity conditions for mean field games. *Journal of Functional Analysis*, 285(9):110095, 2023.
- [Gra14] P Jameson Graber. Optimal control of first-order Hamilton–Jacobi equations with linearly bounded Hamiltonian. *Applied Mathematics & Optimization*, 70(2):185–224, 2014.
- [Gra24] P Jameson Graber. Remarks on potential mean field games. *arXiv preprint arXiv:2405.15921*, 2024.
- [HMC06] Minyi Huang, Roland P Malhamé, and Peter E Caines. Large population stochastic dynamic games: closed-loop McKean-Vlasov systems and the nash certainty equivalence principle. *Communications in Information & Systems*, 6(3):221–252, 2006.
- [LL06a] Jean-Michel Lasry and Pierre-Louis Lions. Jeux à champ moyen. I—Le cas stationnaire. *Comptes Rendus Mathématique*, 343(9):619–625, 2006.
- [LL06b] Jean-Michel Lasry and Pierre-Louis Lions. Jeux à champ moyen. II—Horizon fini et contrôle optimal. *Comptes Rendus Mathématique*, 343(10):679–684, 2006.
- [LL07] Jean-Michel Lasry and Pierre-Louis Lions. Mean field games. *Japanese Journal of Mathematics*, 2(1):229–260, 2007.
- [Mun22] Sebastian Munoz. Classical and weak solutions to local first-order mean field games through elliptic regularity. *Annales de l’Institut Henri Poincaré C*, 39(1):1–39, 2022.
- [new] Newton-rhapon method on general strongly convex functions. <https://math.stackexchange.com/questions/3408436/newton-rhapon-method-on-general-strongly-convex-functions>.
- [PS17] Adam Proskinski and Filippo Santambrogio. Global-in-time regularity via duality for congestion-penalized mean field games. *Stochastics*, 89(6-7):923–942, 2017.
- [Zim24] Brady Zimmerman. A finite dimensional approximation of a density dependent mean field game, 2024. Bachelor’s Thesis.

J. GRABER: BAYLOR UNIVERSITY, DEPARTMENT OF MATHEMATICS;, SID RICHARDSON BUILDING, 1410 S. 4TH STREET, WACO, TX 76706, TEL.: +1-254-710-, FAX: +1-254-710-3569

Email address: Jameson_Graber@baylor.edu

B. ZIMMERMAN: BAYLOR UNIVERSITY,, SID RICHARDSON BUILDING, 1410 S. 4TH STREET, WACO, TX 76706

Email address: Brady_Zimmerman1@baylor.edu