A Formulation of Quantum Fluid Mechanics and Trajectories

James P. Finley:

Department of Physical Sciences, Eastern New Mexico University, Portales, NM 88130

E-mail: james.finley@enmu.edu

Abstract.

A formalism of classical mechanics is given for time-dependent many-body states of quantum mechanics, describing both fluid flow and point mass trajectories. The familiar equations of energy, motion, and those of Lagrangian mechanics are obtained. An energy and continuity equation is demonstrated to be equivalent to the real and imaginary parts of the time dependent Schrödinger equation, respectively, where the Schrödinger equation is in density matrix form. For certain stationary states, using Lagrangian mechanics and a Hamiltonian function for quantum mechanics, equations for point-mass trajectories are obtained. For 1-body states and fluid flows, the energy equation and equations of motion are the Bernoulli and Euler equations of fluid mechanics, respectively. Generalizations of the energy and Euler equations are derived to obtain equations that are in the same form as they are in classical mechanics. The fluid flow type is compressible, inviscid, irrotational, with the nonclassical element of *local* variable mass. Over all space mass is conserved. The variable mass is a necessary condition for the fluid flow to agree with the zero orbital angular momentum for s states of hydrogen. Cross flows are examined, where velocity directions are changed without changing the kinetic energy. For one-electron atoms, the velocity modification gives closed orbits for trajectories, and mass conservation, vortexes, and density stratification for fluid flows. For many body states, Under certain conditions, and by hypotheses, Euler equations of orbital-flows are obtained. One-body Schrödinger equations that are a generalization of the Hartree-Fock equations are also obtained. These equations contain a quantum Coulomb's law, involving the 2-body pair function of reduced density matrix theory that replace the charge densities.

1. Introduction

A formalism of classical mechanics is presented for time-dependent many-body states of quantum mechanics, describing both fluid flows and point mass trajectories. An energy and continuity equation is demonstrated to be equivalent to the real and imaginary parts of the time dependent Schrödinger equation, respectively, where the Schrödinger equation is in density matrix form. For a class at stationary states, the energy equation implies a Hamiltonian that is used in Lagrangian mechanics to obtain equations of motion.

For fluid flow of 1-body states, the equation of motion is the Euler equation, and the energy equation is the Bernoulli equation. The flow type is compressible, inviscid, irrotational, with the nonclassical element of *local* variable mass. Over all space mass is conserved. For many body states, under certain conditions, and by hypotheses, Euler equations of orbital-flows are obtained. Under additional conditions, 1-body Schrödinger equations that are a generalization of the Hartree-Fock equations are also obtained. These equations contain a quantum Coulomb's law, involving the 2-body pair function of reduced density matrix theory, replacing charge densities. In order to obtain local conservation of mass for fluids flows and trajectories with continuous velocity vectors, cross flows are examined, where velocity directions are changed without changing the kinetic energy. These cross flows provide vortexes for one-electron atoms and closed orbits for trajectories. Conditions are given for continuity satisfaction and incompressible fluid flow, giving density stratification, where the mass density is only constant of each streamline. As in the relationship between Madelung and Bohminan mechanics for one-body states, the streamlines of the fluid flow are also the paths of point mass trajectories. Because of the this close relationship, we move freely back and forth between the two descriptions.

Needed wavefunction identities are obtained Sec. 4. Sec. 5 defines variables for quantum states that correspond to linear self-adjoint operators, and explicit forms of these variables are given with the wavefunction in polar form. Sec. 6 uncovers much mathematical structure, with many relationships, involving the terms that originate in the imaginary part of the Schrödinger equation in density matrix form.

The energy equation of Sec. 7 is a combination and generalization of the energy equations of the Bernoulli equation of fluid mechanics, used to describes 1-body states with a real valued wavefunctions [1], and the one from Bohmian Mechanics [2–13], used to describe states with complex value wavefunctions. The energy equation can be represented with a certain pressure, for fluids, or a certain body potential, for trajectories. There are two linearly independent velocity components: one velocity component of Bohmian mechanics, denoted \mathbf{v} , and one from the Bernoullian formalism mentioned above, denoted \mathbf{u} . Because the energy equation has a nonclassical feature of kinetic energy when $\mathbf{u} \cdot \mathbf{v}$ does not vanish, and we wish to remove such elements, a generalized energy equation is derived without this feature.

In Sec. 8, for 1-body states, an Euler equation of fluid dynamics and a continuity equation is proved to be equivalent to the time dependent Schrödinger equation. Because terms of the momentum time-derivative of classical fluids are missing in the Euler equation, a range of generalizations are obtained. The most generalized one has all classical terms, and the most number of additional forces, including bulk viscosity. The additional forces involve coupling between the two velocity field components. Interpretations of the various equations, and the identifications of variables are given in Secs. 8.2 and 8.3. Sec. 9 covers cross flows. Sec. 10 provides Lagrangian mechanics for quantum states. Sec. 11 obtains orbital flows, and mixed flows, involving a single 1-body wavefunction describing all bodies. These equations contain a quantum Coulomb's law, involving the 2-body pair function of reduced density matrix theory that replace the charge densities.

There are a number of overlapping formalism that go by various names for the description of quantum states: the De Broglie–Bohm theory [2–13], also called Bohmian mechanics and quantum hydrodynamics, and Madelung fluid mechanics [14, 15]. All of these methods have the same Madelung velocity \mathbf{v} , and, except for the earliest ones, explicitly contain the quantum potential Q [2, 3, 8]. For many body states, the theories based on a Madelung fluid differ from the Bohmian particle trajectories [13]. Heifetz and coworkers [16, 17] explores the thermodynamics of Madelung fluids. There are many modifications and generalizations of the Madelung equations [18–23]. The generalization by Broadbridge [19] and Jamali [22] use a complex velocity. Tsekov [24] also uses a complex velocity to derives a complex Navier–Stokes equation. Vadasz [25] derived an extension of the Schrödinger equation from the Navier–Stokes equation. Quantum hydrodynamic theory has been employed to treat systems with single particle wave functions [26–41]. The method also also been generalized to treat many particle systems [42, 43]. Application of this formalism include the investigation of spin effects [44, 45], Bose–Einstein condensates [46], graphene [47] and plasmas [48–50].

The quantum potential of Bohmian mechanics suffers from kinetic energy Because of the missing kinetic energy, Bohmian mechanics has a contamination. difficulty in explaining the nonvanishing expectation value of kinetic energy for states with real valued wavefunction, given that these states are described as being static. Bohmian mechanics theory provides no information about these states, nor explain what the Bohmian potential does for static states. It is possible that spin angular momentum might be accounted for by an orbiting electron, and there is a formula for the electron motion derived by Holland [51-53]. However, the kinetic energy responsible for the creation of the orbital momentum must be present in nonrelativistic quantum mechanics, and the only place it could be is in the Bohm potential. Bohmian mechanics also cannot explain the x and y portions of orbital angular momentum. Furthermore, the Bohm potential, besides being difficult to interpret, gives a stress tensor for force for equations of motion [8, 43], even for stationary states, where the forces should be conservative and be defined by a potential. The addition of the velocity vector component \mathbf{u} of the Bernoullian formulation, and the elimination of the Bohm potential, removes these problems [1].

Adding the velocity vector \mathbf{u} and a pressure P has other advantages [1]. The force from the pressure removes the necessity of tunneling, so that nothing special happens when a particle moves into a region that is, otherwise, classically forbidden. The velocity vector \mathbf{u} explains the zero angular momentum for the *s* electronic states of the hydrogen atom, since the velocity field is in the radial direction. It is also easily demonstrated that the pressure force $-\nabla P$ helps stabilize one-electron atoms, giving, for the hydrogen 1s state, a strong repulsive force close to the nucleus and a weak attractive force beyond a distance of about 1.37 a.u. For energy, a portion of the pressure cancels the energy from the proton nucleus, giving a finite energy at the nucleus. Equations for the speed of sound have also been obtained, where the extrema of the momentum per volume occur at points that are Mach 1 speed [54].

2. Runner Notation System

In this section, a notational system is introduced, applicable to the sections that follow, where equations for the general *n*-body case have the same appearance as in the special case where (n = 1). This notation permits the omission of the summation sign along with the corresponding subscripts that run over an index set. Consider the two equations: $(-\nabla^2 \Psi/2 + U\Psi = E)$ and $(K \doteq -\nabla^2 \Psi/2)$. It is understood, by the rules given below, that the first equation is a single equation, such that ∇^2 denotes the series $(\nabla_1^2 + \cdots + \nabla_n^2)$. The second one is a sequence of *n* definitions, the *i*th being $(K_i = -\nabla^2 \Psi_i \div 2)$.

Definition 1 The core family $(\Psi \doteq \{Re\Psi, Im\Psi, \rho, S\})$ is a set of real-valued functions from the Cartesian product $\mathbf{S} \times (\mathbb{R} \times \mathbb{R}^{3n})$. The spin space is defined by $\mathbf{S} \doteq \{(\omega_1, \omega_2, \cdots, \omega_n) | \omega_i \in \{-1, +1\}\}, and \mathbb{R} \times \mathbb{R}^{3n} is a Euclidean space.$ The members of Ψ are two times continuously differentiable almost everywhere on the space $\mathbb{R} \times \mathbb{R}^{3n}$ for all $\mathbf{s} \in \mathbf{S}$. We write $\mathbb{R} \times \mathbb{R}^{3n}$ to emphasize that the variable of \mathbb{R} is time. The elements of Ψ are related via the polar form of the wavefunction: $(\Psi = \sqrt{\rho}e^{iS/\hbar})$.

If two functions are equal almost everywhere (a.e.), they are considered physically indistinguishable. The symbol "=" means equal a.e. The same idea is used for equivalence of equations: A function satisfies one of the equations a.e, if and only if it satisfies the other one a.e. Hence, there will be on reason to write "a.e," since the condition is worked into the formalism. For the most part, we do not use, nor do we need, the special structure from the $L^3(\mathbb{R}^{3n})$ Hilbert space, that is not already in the underlining vector space.

Definition 2 (Runners) Let $(\Sigma, +)$ be a vector space of n-tuples fitted with the sequence to series map $(X) \mapsto [X]$, where $(X) = (X_1, X_2, \dots X_n) \in \Sigma$ and $([X] \doteq X_1 + X_2 \dots + X_n)$. The runner notation system uses special shorthand notation: If $(X), (Y) \in \Sigma$, then the n-equations

$$X_i = Y_i, \qquad i = 1, \cdots, n,$$

are denoted by (X = Y). However, if $U \notin \Sigma$ appears as a term in an equation with one or more elements from Σ , then, for example, (U = X + Y) denotes the series equality: (U = [X] + [Y]), that is

$$U = X_1 + X_2 + \dots + X_n + Y_1 + Y_2 + \dots + Y_n.$$

The elements (X) and (Y) are said to run in the equation series (X = Y + U) and step in the equation sequence (X = Y). The members of $(\Sigma, +)$ are called runners.

Let $\rho, S \in \Psi$ and $U \notin \Sigma$. For runners $(\nabla \rho)$, $(\nabla S) \in \Sigma$ of an n-body state, here are two examples of the notation:

$$\nabla^2 Y \doteq \nabla \rho \cdot \nabla S \quad means \quad \nabla_i^2 Y = \nabla_i \rho \cdot \nabla_i S, \quad i \in \{1, 2, \dots n\}$$
$$\nabla^2 Y = \nabla \rho \cdot \nabla S + U \quad means \quad \sum_{i=1}^n \nabla_i^2 Y = \sum_{i=1}^n \nabla_i \rho \cdot \nabla_i S + U,$$

where it is understood that the collection Σ is extended to include Y, and the symbol " \doteq " is used for definitions. If $f \in \Psi$ and $(\hat{O} = \hat{O}_1 + \cdots + \hat{O}_n)$ is a differential operator, where \hat{O}_i acts on the *i*th component of \mathbb{R}^{3n} , then we require that $((\hat{O}f) \in \Sigma)$.

Other notation used includes the following items. 1) A semicolon in an equation means "such that" or "where." For example, " $(A > B^{-1}; B \neq 0)$ " means "A is greater than B such that $(B \neq 0)$." 2) "Eq. $(N) \times c$ " means Eq. (N) that is multiplied by c. 3) The symbol ∂ is the partial time derivative operator, that is $([\partial \Psi](t) \doteq \partial \Psi/\partial t)$. 4) The following notation is used for spin summations and spatial integrations:

 \sum_{m} and \int_{m} are the sum and integration over the mth component of elements

from **S** and \mathbb{R}^{3n} , respectively, For example, for (n = 3) and (m = 2), we have

$$\left[\int_{2}\sum_{2}\Psi\right](\mathbf{r}_{1},\omega_{1};\mathbf{r}_{3},\omega_{3}) = \int\sum_{\omega_{2}\in\{-1,+1\}}\left[\Psi(\mathbf{r}_{1},\omega_{1};\mathbf{r}_{2},\omega_{2};\mathbf{r}_{3},\omega_{3})\right]d\mathbf{r}_{2}$$

The following notation is used:

$$\int_{m+} = \int_{m+1} \cdots \int_{n}, \quad \sum_{m+} = \sum_{m+1} \cdots \sum_{n}, \quad \hat{\int}_{m} = \sum_{m} \int_{m}, \quad \hat{\int}_{m+} = \sum_{m+} \int_{m+},$$

and $\hat{\int} = \hat{\int}_{1} \cdots \hat{\int}_{n}.$ (1)

It is common in mathematical proofs to prove that a statement, involving a given function, is true for only a value f(x) of the function f. It is then pointed out that, since the point x is arbitrary, the proof holds for the function f. We use this approach for spin, where the spin variables are treated as parameters. For density matrices, this differs from the usual convention, where spin is an explicit variable or the spin variables are summed over, giving spinless density matrices.

3. Correspondence Variables

In quantum mechanics, physical properties are replaced by observables. Linear selfadjoint operators represent observables that, together with rules, provide predictions for the statistical outcome of a sufficiently large collection of measurements of an observable. There are special probabilistic predictions given for the order of measurements and the simultaneous measurements of related pairs of observables, position and momentum being an example. There is a special axiom declaring the physical meaning of the probability distribution ($\rho \doteq \Psi^* \Psi$), called the Born rule.

In this paper, variables are defined that correspond to observables of quantum mechanics. These correspondence variables are needed to give a description of quantum states based on classical mechanics. These variables are generalized real-valued fields, with domain $\hat{\mathbb{R}} \times \mathbb{R}^{3n}$, that are defined using the wavefunction Ψ and operators of quantum mechanics. This approach yields two variables in some cases, one from the

real part, and one from the imaginary part, of a defining complex valued-function. There are two classes of correspondence variables: *particle and fluid variables*. The two classes being related by the map: $(Y \mapsto Y\rho)$, where Y is a particle variable, $Y\rho$ is a fluid variable, and $(\rho \doteq \Psi^*\Psi)$ is the probability distribution. These definitions are given to avoid saying "per mass" and "per volume."

It is a requirement of a wavefunction for all times that the subset of \mathbb{R}^{3n} where the probability distribution ρ vanish has measure zero, and that a wavefunction and its first derivatives vanish at infinity. These conditions can be satisfied by requiring the external potentials to be members of the $(L^{3/2} + L^{\infty})$ Hilbert space [55]. Fluid variables have the same domain as the wavefunction they are determined by. For particle variables, the domain is the support of $\Psi^*\Psi$, defined as the region of \mathbb{R}^{3n} space where $\Psi^*\Psi$ does not vanish, for a given point of time.

Next we define the classes of correspondence variables. Let \hat{L} be an operator that represents an observable that is given by a sum over 1-body operators. For example, $(-L = i\hbar\nabla_1 + \cdots + i\hbar\nabla_n).$

1a) (Weak variable from operator) Let $((\Psi^* \hat{L} \Psi) \in \Sigma)$. A weak fluid variable $((Y) \in \Sigma)$, corresponding to operator \hat{L} , is a real-valued generalized field, with domain \mathbb{R}^{3n} , that satisfies the following equation sequence:

$$\left\langle \hat{L} \right\rangle \doteq \hat{\int} \Psi^* \hat{L} \Psi = \hat{\int} Y,$$
 (2)

where $\langle \hat{L} \rangle$ is the expectation value of an observable that corresponds to \hat{L} , and $\hat{\int}$ is the *n*-body spin-sum, spatial-integration operator (1).

1b) (Strong variable by operator) The two strong fluid variables, corresponding to linear self adjoint operator \hat{L} , are the real and imaginary parts of the function $(\Psi^*\hat{L}\Psi)$. The strong particle variables are the real or imaginary parts of $((\Psi^*L\Psi/\rho)) \in \Sigma)$. It follows from the definitions that the strong fluid variable $(\text{Re}\Psi^*\hat{L}\Psi)$ is weak, since the expectation value of a Hermitian operator must be real. The strong-particle variables by composition are defined by the real or imaginary parts of the composite $((g \circ (\Psi^*L\Psi/\rho)) \in \Sigma)$, involving a given map g that acts on each component of the sequence $(\Psi^*\hat{L}\Psi/\rho)$. We also take the external potential U and $(i\hbar\Psi^*\partial\Psi/\rho)$ to be strong particle variables of potential and total energy. These variables are not runners.

1c) (First order reduced variables by operator) Let f and g be the strong particle variables of \hat{L} :

$$f\rho + ig\rho \doteq \Psi^* \hat{L} \Psi; \qquad f, g \in \Sigma.$$
 (3)

The first order, reduced, strong particle-variables, \hat{f} and \hat{g} , corresponding to operator \hat{L} , are

$$\hat{f}\hat{\rho} + i\hat{g}\hat{\rho} \doteq \int_{1+}^{\infty} \Psi^* \hat{L}_1 \Psi, \qquad (4)$$

where $\hat{\int}_{1+}$ is the spin-sum, spatial-integration operation (1) with (m = 1); the probability density $\hat{\rho}$, defined by

$$\hat{\rho} \doteq \hat{\int}_{1+} \rho; \qquad \rho = \Psi^* \Psi, \quad \hat{\int}_1 \hat{\rho} = 1,$$
(5)

is normalized to unity, instead of the usual n normalization for a wavefunction normalized to unity. Defs. (3) an (4) give

$$\hat{f}\hat{\rho} = \int_{1+}^{\hat{c}} f_1\rho \quad \text{and} \quad \hat{g}\hat{\rho} = \int_{1+}^{\hat{c}} g_1\rho.$$
 (6)

2)(Variable from equation) For the special case of 1-body (n = 1), the field is a variable of an equation of classical mechanics, and the equation is equivalent to, or implied by, a 1-body Schrödinger equation.

3) (A free, or distributive, variable) is a fluid variable whose integration over all space \mathbb{R}^{3n} vanish at all times. If $(Y = W\rho)$ and (Y) is free, (W) is a weak particle variable. Note that if (Y) is a weak fluid variable, such that

$$Y = Z + X,\tag{7}$$

where X is free, then (Z) is also a weak variable of operator. The strong fluid variable $(\text{Im}\Psi^*\hat{L}\Psi)$ is free, since the expectation value of a Hermitian operator must be real.

4. Wavefunction Identities

The lemmas in this section are needed to obtain formulae for certain correspondence variables. Note that displayed equations in Lemmas 1 and 3 below, are sequences of n equations. Let

$$(\Psi \nabla^2 \Psi), \Psi \nabla \Psi, |\nabla \Psi|^2, \nabla \cdot (\Psi^* \nabla \Psi) \in \mathbf{\Sigma}, \quad \text{and} \quad \hat{\mathbf{P}}_i \doteq -i\hbar \nabla_i.$$

Lemma 1 (Momentum Operator Form of the Laplacian)

$$\Psi^*\left(-\hbar^2\nabla^2\Psi\right) = |\hat{\mathbf{P}}\Psi|^2 - i\hbar\nabla\cdot\left(\Psi^*\hat{\mathbf{P}}\Psi\right), \quad \hat{\mathbf{P}} \doteq -i\hbar\nabla.$$
(8)

$$|\hat{\mathbf{P}}\Psi|^2 \rho = |\Psi^* \hat{\mathbf{P}}\Psi|^2. \tag{9}$$

$$\Psi^* \left(-\frac{\hbar^2}{2m} \nabla^2 \Psi \right) \rho = \frac{1}{2m} |\Psi^* \hat{\mathcal{P}} \Psi|^2 - i \frac{\hbar}{2m} \nabla \cdot \left(\Psi^* \hat{\mathcal{P}} \Psi \right) \rho.$$
(10)

Proof Substituting the following two identities into the third one gives Eq. (8):

$$\begin{split} |\hat{\mathbf{P}}\Psi|^2 &= (-i\hbar\nabla\Psi) \cdot (i\hbar\nabla\Psi^*) = \hbar^2 (\nabla\Psi^* \cdot \nabla\Psi), \\ i\hbar\nabla \cdot \left(\Psi^* \hat{\mathbf{P}}\Psi\right) &= \hbar^2 \nabla \cdot (\Psi^* \nabla\Psi) \\ \Psi^* \left(-\hbar^2 \nabla^2 \Psi\right) &= \hbar^2 \nabla \Psi^* \cdot \nabla \Psi - \hbar^2 \nabla \cdot (\Psi^* \nabla\Psi) \,. \end{split}$$

For Eq. (9), $|\hat{\mathbf{P}}\Psi|^2 \rho = \rho(-i\hbar\nabla\Psi) \cdot (i\hbar\nabla\Psi^*) = (-i\hbar\Psi^*\nabla\Psi) \cdot (i\hbar\Psi\nabla\Psi^*) = |\Psi^*\hat{\mathbf{P}}\Psi|^2$.

The multiplication of identity (8) by $\rho/2m$ and using identity (9) gives Eq. (10).

Lemma 2 (Schrödinger Eq. and Vanish of Divergence Momentae Integral) The n-body Schrödinger equation, in the form

$$i\hbar\Psi^*\partial\Psi = \Psi^*\hat{H}\Psi; \qquad \hat{H} \doteq = -\frac{\hbar^2}{2m}\nabla^2 + U, \quad and \quad \nabla^2\Psi \in \Sigma,$$
 (11)

can be written

$$i\Psi^*\hbar\partial\Psi = \frac{1}{2m}|\hat{\mathbf{P}}\Psi|^2 - i\frac{\hbar}{2m}\nabla\cdot\left(\Psi^*\hat{\mathbf{P}}\Psi\right) + |\Psi|^2U.$$
(12)

The real and imaginary parts are

$$-\hbar Im\left(\Psi^*\partial\Psi\right) = \frac{1}{2m}|\hat{\mathbf{P}}\Psi|^2 + \frac{\hbar}{2m}Im\left(\nabla\cdot\left(\Psi^*\hat{\mathbf{P}}\Psi\right)\right) + |\Psi|^2U, \quad and \tag{13}$$

$$-\hbar Re\left(\Psi^*\partial\Psi\right) = \frac{\hbar}{2m} Re\left(\nabla\cdot\left(\Psi^*\hat{\mathbf{P}}\Psi\right)\right),\tag{14}$$

respectively, where U is the external potential. For each component $(\nabla_i \cdot (\Psi^* \hat{\mathbf{P}}_i \Psi))$ of the sequence $(\nabla \cdot (\Psi^* \hat{\mathbf{P}} \Psi) \in \Sigma)$,

if
$$\Psi$$
 vanish at infinity then $\int_{\mathbb{R}^3} \nabla_i \cdot \left(\Psi^* \hat{\mathbf{P}}_i \Psi \right)$ vanish.

Proof Eq. (12) follows from the substitution of Eq. (8)÷2*m* into Eq. (11). Eqs. (13) and (14) follow by separating Eq. (12) into real and imaginary parts and using the identities ($\operatorname{Re}(iX) = -\operatorname{Im}X$) and ($\operatorname{Im}(iX) = \operatorname{Re}X$). For the second part,

let
$$\bar{\mathbf{P}}_{\alpha} = -i\hbar \frac{\partial}{\partial \alpha}, \quad \alpha \in \{x, y, z\} = \mathbf{r} \in (\mathbf{r}_1, \mathbf{r}_2, \cdots, \mathbf{r}_n).$$

$$\int_{\mathbb{R}} \frac{d}{d\alpha} \left(\Psi^* \bar{\mathbf{P}}_{\alpha} \Psi \right) = \Psi^* \bar{\mathbf{P}}_{\alpha} \Psi \Big|_{-\infty}^{\infty} = 0.$$

Definition 3

We replace the wavefunction Ψ as the fundamental variable, with the variables ρ and S, such that

$$\Psi \doteq Re^{iS/\hbar} \doteq e^{(iS-\theta)/\hbar}, \quad \rho \doteq \Psi^* \Psi = R^2, \quad \int \rho = 1, \tag{15}$$

where Ψ is defined to within a phase factor. Hence, we have the composite $\Psi = \Psi(\rho, S)$. Also, θ is defined under the condition $(R > \mathbf{0})$, giving $(\rho = e^{-2\theta/\hbar})$. (For convenience, this definition is given in a different form with Defs. (22).)

Lemma 3 (Gradient, Divergence and Laplacian Identities in Polar Form)

$$\Psi^* \hat{\mathbf{P}} \Psi = \rho \nabla S - i \frac{\hbar}{2} \nabla \rho.$$
(16)

$$\frac{1}{2m}|\Psi^*\hat{\mathbf{P}}\Psi|^2 = \frac{\hbar^2}{8m}|\nabla\rho|^2 + \frac{1}{2m}\rho^2|\nabla S|^2.$$
(17)

A Formulation of Quantum Fluid Mechanics and Trajectories

$$-i\frac{\hbar}{2m}\nabla\cdot\left(\Psi^{*}\hat{\mathbf{P}}\Psi\right) = -\frac{\hbar^{2}}{4m}\nabla^{2}\rho - i\frac{\hbar}{2m}\nabla\cdot\left(\rho\nabla S\right).$$
(18)

$$\Psi^*\left(-\frac{\hbar^2}{2m}\nabla^2\Psi\right)\rho = \frac{\hbar^2}{8m}|\nabla\rho|^2 + \frac{1}{2m}\rho^2|\nabla S|^2 - \frac{\hbar^2}{4m}\rho\nabla^2\rho - i\frac{\hbar}{2m}\rho\nabla\cdot(\rho\nabla S).$$
(19)

Proof Let $(S' \doteq S/\hbar)$. Consider the development

$$Re^{-iS'}(-\nabla)Re^{iS'} = -iR^2\nabla S' - R\nabla R = -i\rho\nabla S' - \frac{1}{2}\nabla\rho.$$

Multiplying the equation by $i\hbar$, setting $(S' = S/\hbar)$, and using the definition of the momentum operator $(\hat{P}\Psi) = (-i\hbar\nabla\Psi)$, we obtain Eq. (16). Eqs. (17) and (18) follow immediately. Substituting Eqs. (17) and (18) into Eq. (10)÷(2m) we obtain Eq. (19).

For later convenience, we note that Eq. (16) holds with the replacement $(-\nabla) \rightarrow \partial$:

$$i\hbar\Psi^*\partial\Psi = -\rho\partial S + i\frac{\hbar}{2}\partial\rho.$$
⁽²⁰⁾

Only the derivatives of S are needed for the correspondence variables. It follows from (16) and (20) that ∇S and ∂S can be computed from a wavefunction, as the real parts of $\Psi^* \hat{P} \Psi / \rho$ and $-i\hbar \Psi^* \partial \Psi / \rho$, respectively, even if the wavefunction is not written in polar form, as in a linear combination.

5. Polar Forms of Correspondence Variables

In this section, polar forms of the correspondence are derived.

5.1. Momentae Strong and Kinetic Energy Strong by a Composition

In this subsection, polar forms of the correspondence strong variables for momentum are given, where the ordered momentum pair is called a momentae. The kinetic energy variable by composition is defined and obtained. A discussion of one of the velocity fields is given.

Definition 4 (Momentae Runners $\{(\rho_m \mathbf{v}), (\rho_m \mathbf{u})\} \subset \Sigma$)

$$\Psi^* \dot{\mathbf{P}} \Psi \doteq \rho_m \mathbf{v} + i\rho_m \mathbf{u}, \quad \rho_m \doteq m\rho, \tag{21}$$

$$\zeta_0 \doteq \frac{\hbar}{2}, \qquad \zeta = \frac{1}{m}\zeta_0, \qquad \eta = \zeta_0\rho, \quad \theta \doteq -\zeta_0 \ln \rho.$$
(22)

The parameter m is the mass of each and every identical particle of a quantummechanical system. The functions \mathbf{v} and \mathbf{u} are the first and second velocities of the order pair (\mathbf{v}, \mathbf{u}) . (To define $\ln \rho$, ρ is unitless in $\ln \rho$.)

Comparing (16) and (21), and using (22), we find that

$$\rho_m \mathbf{v} = \rho \nabla S \doteq \zeta_0 \nabla \phi + \zeta_0 \nabla \times \mathbf{J} \doteq \zeta_0 \mathbf{Q}, \qquad \rho_m \mathbf{u} = \rho \nabla \theta = -\zeta_0 \nabla \rho, \tag{23}$$

A Formulation of Quantum Fluid Mechanics and Trajectories

where the first equations string defines the scalar (ϕ) and vector (**J**) fields by Helmholtz decomposition. By definition (21) and definition **1b** of Sec 3, (($\rho_m \mathbf{v}$) = ($\rho \nabla S$)) and (($\rho_m \mathbf{u}$) = ($\rho \nabla \theta$)) are the strong-fluid momentae variables. Similarly, ($m\mathbf{v}$) and ($m\mathbf{u}$) are the strong-particle momentae variables. These variables are irrotational fields with S and θ particle-momentae potentials. For a fluid interpretation of 1-body states, ρ_m is the mass density, and the real and imaginary parts of the following are the mass fluxes:

$$\nabla \cdot \left(\Psi^* \hat{\mathbf{P}} \Psi \right) = \nabla \cdot (\rho_m \mathbf{v}) + i \nabla \cdot (\rho_m \mathbf{u}).$$

Using the composite $(g(\rho^{-1}\Psi^*\hat{P}\Psi))$ with $(g(x) = |x|^2/2m)$, where x is complex valued, and noting that $(\rho^{-1}\Psi^*\hat{P}\Psi = m\mathbf{v} + im\mathbf{u})$, we obtain a strong-particle kinetic-energy variable by composition:

$$\frac{1}{2m} \left| \frac{\Psi^* \hat{P} \Psi}{\Psi^* \Psi} \right|^2 = \frac{1}{2m} |\nabla \theta|^2 + \frac{1}{2m} |\nabla S|^2 = \frac{1}{2} m u^2 + \frac{1}{2} m v^2.$$
(24)

For later convenience in Corollary (18), we note that starting from the second equation from (23), we have

$$m\mathbf{u} = -\zeta_0 \frac{\nabla \rho}{\rho} \Rightarrow \frac{1}{2}mu^2 = \frac{1}{2}\zeta_0^2 m |\nabla \rho|^2 \rho^{-2}.$$
 (25)

As in Def. (5) for probability density $\hat{\rho}$, let

$$\hat{\gamma} = \int_{1+}^{\hat{\gamma}} \gamma, \qquad \gamma = \phi, \mathbf{J}.$$
 (26)

As in Eq. (23), let $\hat{\mathbf{v}}$ and $\hat{\mathbf{u}}$ be defined by

$$\hat{\rho}_m \hat{\mathbf{u}} \doteq -\zeta_0 \nabla_1 \hat{\rho}, \quad \hat{\rho}_m \hat{\mathbf{v}} \doteq \zeta_0 \nabla_1 \hat{\phi} + \zeta_0 \nabla \times \hat{\mathbf{J}}.$$
(27)

Next we show that these definitions for $\hat{\mathbf{u}}$ and $\hat{\mathbf{v}}$ are equivalent to the strong, first order reduced variables given by Def. (6):

$$\hat{\rho}\hat{\mathbf{u}} = \int_{1+}^{\hat{\rho}} \rho \mathbf{u}_1, \qquad \hat{\rho}\hat{\mathbf{v}} = \int_{1+}^{\hat{\rho}} \rho \mathbf{v}_1.$$
(28)

Def. (23.2) and (26) give

$$\hat{\rho}_m \hat{\mathbf{u}} = \int_{1+}^{\hat{\rho}} \rho_m \mathbf{u}_1 = -\zeta_0 \int_{1+}^{\hat{\rho}} \nabla_1 \rho \, d\mathbf{r}_1' = -\zeta_0 \nabla_1 \hat{\rho}.$$
(29)

Similarly, Def. (23.1) and (26) give

$$\hat{\rho}_m \hat{\mathbf{v}} = \sum_{1+} \int \rho_m \mathbf{v}_1 \, d\mathbf{r}_1' = \zeta_0 \sum_{1+} \int \left(\nabla_1 \phi + \zeta_0 \nabla_1 \times \mathbf{J} \right) \, d\mathbf{r}_1' = \zeta_0 \nabla_1 \hat{\phi} + \zeta_0 \nabla_1 \times \hat{\mathbf{J}}. \tag{30}$$

Hence, the two definitions for vector fields $\hat{\mathbf{u}}$ and $\hat{\mathbf{v}}$ are equivalent. This same result holds for any vector field-variable that can be expressed as a Helmholtz decomposition, where $\hat{\rho}_m \hat{\mathbf{u}}$ is special, since it is irrotational.

(Velocity singularities) The velocities (\mathbf{v}) and (\mathbf{u}) have singularities when ρ vanish at nodes, and this causes some equations obtained below not to be defined for these subspaces [1, 8, 54]. Since these subspaces have measure zero, these singularities pose no problems if one works in a Hilbert space. These singularities can be removed by using the fluid-variable momentum to replace the particle-variable momentum, since such a modification give finite limits as $\rho \to 0$, at least for a few systems that have been studied [1]. However, such treatments seem to be counter production, since singularities in variables of classical mechanics can be useful. For example, the electric field on the surface of a conductor can be discontinuous, giving singularities in the electric-field gradient. Since singularities involving ($m\mathbf{u}$) and ($m\mathbf{v}$) might have some usefulness, and these singularities arise naturally out of the formalism, in some equations and definitions we choose to use the particle momentum-variables ($m\mathbf{v}$) and ($m\mathbf{u}$), instead of the corresponding fluid ones ($\rho_m \mathbf{v}$) and ($\rho_m \mathbf{u}$).

(Other work with velocity u) The second velocity u of Eq. (25) for one-body systems, appears in other investigations. The function $2\rho_m |\mathbf{u}|^2$ is a term of the Hamiltonian functional of the generalized fluid-dynamics formalism by Broer [56], where the Hamiltonian functional is derived from the time-dependent Schrödinger equation. Salesi [57] obtains a Lagrangian function that is equivalent to the Madelung equations that contains the term $\rho_m |\mathbf{u}|^2/2$. He interprets $|\mathbf{u}|^2/2$ as the internal energy of the relative motion in the center of mass coordinate frame from the Zitterbewegung (ZWB) model of spin. The velocity magnitude $|\mathbf{u}_{\pm}|$ follows as a non-relativistic approximation of a velocity expression of Hestenes [58] of Schrödinger–Pauli theory. Tsekov [24] obtains the same velocity choice **u** of Eq. (25), as the imaginary component of a complex velocity, where the formalism involves diffusion. Furthermore, the velocity choice **u** is used in the Bernoullian equation, an equation equivalent to the real-valued one-body Schrödinger equation, where the flow is compressible with variable-mass [1,54].

5.2. Kinetic energy strong and weak

In this subsection, polar forms of strong and weak correspondence variables for kinetic energy are given. The following lemma contains definitions and a proof that are needed below.

Lemma 4 (The Free Pressure Variables) The definition

$$-i\frac{\hbar}{2m}\nabla\cdot\left(\Psi^{*}\hat{\mathbf{P}}\Psi\right) \doteq P + ip, \qquad \hat{\mathbf{P}}\Psi = -i\hbar\nabla\Psi, \tag{31}$$

is equivalent to the following two alternative definitions:

1)
$$P \doteq \zeta_0 \nabla \cdot (\rho \mathbf{u}), \quad P \doteq -\zeta \zeta_0 \nabla^2 \rho, \quad P \in \Sigma.$$
 (32)

2)
$$p \doteq -\zeta_0 \nabla \cdot (\rho \mathbf{v}), \quad p \doteq -\zeta \zeta_0 \nabla^2 \phi, \qquad p \in \Sigma.$$
 (33)

Also, for each component of the sequences (P) and (p),

$$\int P_i, \quad and \quad \int p_i \quad vanish. \tag{34}$$

Proof Substituting Eq. (21) into (31), followed by the substituting of Eq. (23) and using vector calculus identity $(\nabla \cdot \nabla \times \mathbf{J} = \mathbf{0})$, gives Eqs. (32) and (33). Statement (34) follows from lemma 2 and definition (31).

Lemma 5 (Correspondence Variables from the Kinetic Energy.) Let (P) and (p) be defined by (31), and let $(u) \doteq (|\mathbf{u}|)$ and $(v) \doteq (|\mathbf{v}|)$, where (\mathbf{u}) and (\mathbf{v}) are defined by (21), in other words

$$v\rho_m \doteq \left| \operatorname{Re}(\Psi^* \hat{\mathrm{P}} \Psi) \right|, \quad u\rho_m \doteq \left| \operatorname{Im}(\Psi^* \hat{\mathrm{P}} \Psi) \right|.$$
 (35)

$$\Psi^*\left(-\frac{\hbar^2}{2m}\nabla^2\Psi\right) = \frac{1}{2}\rho_m u^2 + \frac{1}{2}\rho_m v^2 + P + ip.$$
(36)

Proof From definition (21), and the ones for (u) and (v) above, we have

$$\frac{1}{2m}|\Psi^*\hat{\mathbf{P}}\Psi|^2\rho^{-1} = \frac{1}{2}\rho_m u^2 + \frac{1}{2}\rho_m v^2.$$
(37)

Substituting this result, and also the definitions for (P) and (p), Eq. (31), into identity $(10) \times \rho^{-1}$,

$$\Psi^*\left(-\frac{\hbar^2}{2m}\nabla^2\Psi\right) = \frac{1}{2m}|\Psi^*\hat{\mathbf{P}}\Psi|^2\rho^{-1} - i\frac{\hbar}{2m}\nabla\cdot\left(\Psi^*\hat{\mathbf{P}}\Psi\right),$$

we obtain Eq. (36).

The real and imaginary parts of Eq. (36) gives formulae for the two strong-fluid kineticenergy variables, and Eq.(19)× ρ^{-1} gives the polar form representation of these variables. Combining identities (36) and (10), we discover, after using Eq. (34) and the last result from Lemma 2, that the expectation value of the kinetic energy can be written

$$\left\langle -\frac{\hbar^2}{2m} \nabla^2 \right\rangle = \int \frac{1}{2m} |\Psi^* \hat{\mathbf{P}} \Psi|^2 \rho^{-1} = \int \frac{1}{2} \rho_m u^2 + \frac{1}{2} \rho_m v^2.$$
 (38)

From this result and Eq. (24), we find that $\left(|\Psi^*\hat{P}\Psi|^2\rho^{-1}/(2m)\right)$ is a fluid kinetic-energy variable that is strong by composition and weak by equality (38).

Since the probability distribution is symmetric with respect to the interchange of two electron coordinates, e.g., $\rho(\mathbf{r}_1, \mathbf{r}_2, \cdots) = \rho(\mathbf{r}_2, \mathbf{r}_1, \cdots)$, each of the *n* terms of the integrand in Eq. (38)—the terms involving a runner—are equal, so we can write

$$\left\langle -\frac{\hbar^2}{2m} \nabla^2 \right\rangle = n \int_1^2 \left(\frac{1}{2} \rho_m u_1^2 + \frac{1}{2} \rho_m v_1^2 \right) = \int_1 K(\mathbf{r}), \tag{39}$$

where the second integration is over \mathbb{R}^3 , and

$$K(\mathbf{r}) \doteq n \sum_{1} \hat{\int}_{1+} \left(\frac{1}{2} \rho_m u_1^2 + \frac{1}{2} \rho_m v_1^2 \right).$$

Equation (39) is an alternative way to express the kinetic energy functional used in reduced density matrix theory [59,60], replacing the integrand $(-\hbar^2/2m)\nabla_{\mathbf{r}}\rho_1(\mathbf{r},\mathbf{r}')|_{\mathbf{r'=r}}$ of the kinetic energy functional with the kinetic energy density K, where ρ_1 is the reduced, one-body, density matrix. Comparing the two, it seems that the extra variable of $\rho_1(\mathbf{r},\mathbf{r'})$ has no physical meaning. For ground states, the kinetic energy functional (39) is equal to the kinetic energy functional of the Hohenberg–Kohn theorem [59, 61], a functional that is also used in Kohn–Sham density functional theory as part of the kinetic energy correction.

5.3. Total energy strong and weak

In this subsection, polar forms of correspondence variables for the total energy are given.

Definition 5

$$i\hbar\Psi^*\partial\Psi \doteq E\rho + iF\rho, \quad E, F \notin \Sigma.$$
 (40)

The potential U of the Hamiltonian operator \hat{H} is a multiplicative operator such that for each eigenvector Ψ of \hat{H} , the subspace $\mathbf{Z} \doteq \{\Psi(\mathbf{r},t) = 0 | \mathbf{r} \in \mathbb{R}^{3n}\}$ has measure zero for all $t \in \mathbb{R}$. (A condition on U that is satisfied if $U \subset (L^{3/2} + L^{\infty})$, where $(L^{3/2} + L^{\infty})$ is a Hilbert space [55].)

Comparing Eq. (20) and (40) and using Eq. (22) for θ , we discover that

$$E = -\partial S, \quad F\rho = \zeta_0 \partial \rho, \quad \text{and} \quad F = -\partial \theta.$$
 (41)

Taking the gradient and then using the particle momentae Eq. (23), we have

$$-\nabla E = m\partial \mathbf{v}, \qquad -\nabla F = m\partial \mathbf{u},\tag{42}$$

the local balance of particle momentae. The fields $\partial \mathbf{v}$ and $\partial \mathbf{u}$ of 1-body states are identified as local accelerations of classical fluid dynamics [62]. $\partial \mathbf{v}(\mathbf{r}, t)$ and $\partial \mathbf{u}(\mathbf{r}, t)$ are the rate of change of the velocity fields \mathbf{v} and \mathbf{u} at point $\mathbf{r} \in \mathbb{R}^3$ and time $t \in \mathbb{R}$. For steady flow, these partial derivatives vanish, and the acceleration of a fluid particle is obtained by the time derivative of the composite, $((\mathbf{u} + \mathbf{v} = [\mathbf{u} + \mathbf{v}](\mathbf{r}))$, such that $\mathbf{r} = \mathbf{r}(t)$. We identify $-(\nabla E)$ and $-(\nabla F)$ of Eqs. (42) as correspondence variables by equation of local components of force, as defined in Sec. (3). Sec. 6.6 demonstrates that the energae (E, F) have the important property of being conserved over all space for all times.

Adding the potential-energy term $U\rho$ from the Schrödinger equation to both sides of Eq. (36), and then using the definition of the Hamiltonian operator \hat{H} , Eq. (11.2), we find the pair of strong-fluid energy variables from the Hamiltonian operator \hat{H} :

$$\Psi^* \hat{H} \Psi = \frac{1}{2} \rho_m u^2 + P + \frac{1}{2} \rho_m v^2 + U\rho + ip.$$
(43)

It follows from equalities (34) and (38), and the definition of the Hamiltonian operator \hat{H} , that

$$\left\langle \hat{H} \right\rangle = \int \hat{I} \frac{1}{2m} |\Psi^* \hat{P} \Psi|^2 \rho^{-1} + U\rho = \int \hat{I} \frac{1}{2} \rho_m u^2 + \frac{1}{2} \rho_m v^2 + U\rho.$$
 (44)

Hence, the integrand on the rhs is a weak fluid-variable of the energy.

5.4. Energy equations

In this subsection we obtain an energy equation that is a generalization of the Bernoulli equation of fluid mechanics. In Sec. 7, this equation, together with a continuity equation, is proved to be equivalent to the space time Schrödinger equation $(i\hbar\partial\Psi = \hat{H}\Psi)$. A discussion about the energy equation is given, where correspondence variables by equation are identified.

So far in this paper, all equalities are either definitions or they are implied by definitions. Therefore, their satisfaction does not required the wavefunction determining the variables to satisfy a Schrödinger equation. For this paragraph, we impose the condition that the wavefunction Ψ is a solution of the Schrödinger equation $(i\hbar\partial\Psi = \hat{H}\Psi)$ with potential U. This requirement and definition (40) implies that $(E\rho + iF\rho = \Psi^*\hat{H}\Psi)$. Substituting Eqs. (43) into this implication, we discover two energy equations:

$$E\rho = \frac{1}{2}\rho_m u^2 + \frac{1}{2}\rho_m v^2 + P + U\rho, \qquad F\rho = p.$$
(45)

It immediately follows Eqs (45), (44), and (34) that the expectation value of the energy from the Hamiltonian operator is

$$\langle \hat{H} \rangle = \langle E \rangle; \qquad \langle E \rangle \doteq \int \hat{E} \rho.$$
 (46)

A reasonable generalization involving F and some operator \hat{G} is $\langle \hat{G} \rangle$, such that $(\Psi^*\hat{G}\Psi \doteq F\rho)$, and from Eqs. (45) and (34), $\langle \hat{G} \rangle$ vanish. From Sec. 3, definition **3**, F is a free particle-variable of the energy.

6. Time derivative relations

In this section, a number of relationships are obtain in the case of continuity equation $(\partial \rho = -\nabla \cdot (\rho \mathbf{v}))$ satisfaction involving the first velocity (generalized) field \mathbf{v} .

6.1. Equivalent forms of the continuity equation

From the following theorem, six equations are shown to be equivalent to the continuity equation, including the imaginary part of the Schödinger Eq. (11). A set of scalar fields containing the element $\partial \rho$ is identified, such that if one member vanish then all members vanish.

Theorem 1 (The Continuity Six and One for All Vanish)

I. The following equations are equivalent:

1)
$$p = \zeta_0 \partial \rho.$$
 2) $\partial \rho + \nabla \cdot (\rho \mathbf{v}) = \mathbf{0}.$ 3) (45): $F \rho = p.$ (47)
4) $Im(i\hbar\Psi^*\partial\Psi) = Im(\Psi^*\hat{H}\Psi).$

5)
$$\partial \rho = -\nabla \cdot \mathbf{j}; \quad \mathbf{j} = \frac{\hbar}{2mi} \left(\Psi^* \nabla \Psi - \Psi \nabla \Psi^* \right).$$
 (48)

A Formulation of Quantum Fluid Mechanics and Trajectories

6)
$$Eq. (14): \hbar Re(\Psi^* \partial \Psi) = -\frac{\hbar}{2m} Re\left(\nabla \cdot \left(\Psi^* \hat{P}\Psi\right)\right).$$

II. Let the continuity set \mathbb{C} be

$$\mathbb{C} = \{\partial\rho, p, F, \, \nabla \cdot (\rho \mathbf{v}), \nabla \cdot \mathbf{j}, \, \operatorname{Re} \nabla \cdot (\Psi^* \hat{\mathbf{P}} \Psi), \operatorname{Im}(i\hbar \Psi^* \partial \Psi)\},\$$

where it is understood that the runner members run, e.g., p above means $p_1 + \cdots + p_n$. Let $\mathbb{S} \in \mathbb{R}^{3n}$ have non zero measure. Let Eq. #4 be true on \mathbb{S} . If there exist an element $X \in \mathbb{C}$ that vanish on \mathbb{S} , that is $(X|_S = \mathbf{0})$, then each and every element of \mathbb{C} vanish on \mathbb{S} .

IIIa Let $(\Psi = \Psi(\rho, S))$ satisfy the Schrödinger Eq. (11). Let E be a constant. $(-\partial S = E)$ implies all members of the continuity set \mathbb{C} vanish.

IIIb Let $(\Psi = \Psi(\rho, S))$ satisfy the continuity Eq. (1.1). $(\nabla S = \mathbf{0})$ implies all members of the continuity set \mathbb{C} vanish.

IIIc Stationary state implies $S(\mathbf{r},t) = Et + \beta(\mathbf{r})$; $t \in \mathbb{R}$ and $\beta \in \mathbb{R}^{3n}$ and E = const.

Proof

I. by inspection of Def. (32) for p, we find: $\boxed{\mathbf{1} \Leftrightarrow \mathbf{2}}$. By inspection of Eq. (41), $(F\rho = \zeta_0 \partial \rho)$, we find: $\boxed{\mathbf{1} \Leftrightarrow \mathbf{3}}$. Consider Eq. **6**. From Def. (20), we find: $(\hbar \operatorname{Re}(\Psi^* \partial \Psi) = \hbar \partial \rho/2)$, and from Def. (31), we find: $-\hbar \operatorname{Re}\left(\nabla \cdot \left(\Psi^* \hat{P}\Psi\right)\right) \div 2m = p$. Hence, $\boxed{\mathbf{1} \Leftrightarrow \mathbf{6}}$. From Lemma 2, $\boxed{\mathbf{6} \Leftrightarrow \mathbf{4}}$. Consider Eq. **5**. From Def. (31) for p, we have

$$p = -\frac{1}{2m} \operatorname{Im} \left(i\hbar \nabla \cdot \left(\Psi^* \hat{\mathbf{P}} \Psi \right) \right) = -\frac{1}{2m} \operatorname{Im} \left(\hbar^2 \nabla \cdot \left(\Psi^* \nabla \Psi \right) \right),$$

$$p = -\frac{\hbar^2}{4mi} \nabla \cdot \left(\Psi^* \nabla \Psi - \Psi \nabla \Psi^* \right).$$
(49)

Using Eq. 1, in the form $(\partial \rho = \zeta_0^{-1} p)$, we find: $\mathbf{1} \Leftrightarrow \mathbf{5}$.

So far we have **1** is equivalent to **2,3,5** and **6**. We also have: **1** \Leftrightarrow **6** and **6** \Leftrightarrow **4** \Rightarrow **1** \Leftrightarrow **4**. Hence, **1** is equivalent to each and every one. By the transitive property of equivalence relations, each and every pair from the set of six statements is equivalent.

II. Let the implication $(A|_S = \mathbf{0}) \Rightarrow (B|_S = \mathbf{0})$, a relation between $A, B \in \mathbb{C}$, be denoted $A \to B$. Let the equivalence relation $A \leftrightarrow B$ be $A \to B$ and $B \to A$. We need to prove that for all $C, D \in \mathbb{C}, C \leftrightarrow D$. Let $(Q = \text{Re}\nabla \cdot (\Psi^* \hat{P}\Psi))$.

First we prove that for all $C, D \in \mathbb{C}$, except C = Q or $D = Q, C \leftrightarrow D$. From 1 and 2, $p \to \partial \rho \to \nabla \cdot (\rho \mathbf{v})$. From 3 and Def. (32) for $p, \nabla \cdot (\rho \mathbf{v}) \to F$ a.e. From Def. (40) for $F, F \to \operatorname{Im}(i\hbar\Psi^*\partial\Psi)$. From condition 4, we also have $\operatorname{Im}(i\hbar\Psi^*\partial\Psi) \to \operatorname{Im}(\Psi^*\hat{H}\Psi) \notin \mathbb{C}$.

The implication $\operatorname{Im}(\Psi^*\hat{H}\Psi) \to \nabla \cdot \mathbf{j}$ requires three steps. From identity (10), and because $U\rho$ is real valued, \mathbf{i}) $\operatorname{Im}(\Psi^*\hat{H}\Psi) \leftrightarrow Q$ a.e. By noting that \mathbf{ii}) $Q \to \operatorname{Im}(i\hbar\nabla \cdot (\Psi^*\hat{P}\Psi))$, and examination Eqs. (49) with the beginning part 'p =" omitted, \mathbf{iii}) $\operatorname{Im}(i\hbar\nabla \cdot (\Psi^*\hat{P}\Psi)) \to \nabla \cdot \mathbf{j}$. Hence, $\operatorname{Im}(\Psi^*\hat{H}\Psi) \to \nabla \cdot \mathbf{j}$. From $\mathbf{5}$, $\nabla \cdot \mathbf{j} \to \partial \rho$. From $\mathbf{1}$, $\overline{\partial \rho \to p}$. Hence, for all $C, D \in \mathbb{C}$, except C = Q or $D = Q, C \leftrightarrow D$. Since, as mentioned above, $\operatorname{Im}(\Psi^*\hat{H}\Psi) \leftrightarrow Q$, if follows from the transitive property of equivalence relations that each every element $(X|_S = \mathbf{0})$, or equation, from on the set $\{(X|_S = \mathbf{0}) | X \in \mathbb{C}\}$ is equivalent.

IIIa. $(-\partial S = E)$ implies that $\Psi = \Psi(\rho, S)$ satisfies the time independent Schrödinger equation, implying that $(\partial \rho = \mathbf{0})$.

IIIb. The definition $(m\mathbf{v} \doteq \nabla S)$ continuity Eq. (47.2) gives

$$\nabla S = \mathbf{0} \Rightarrow \mathbf{v} = \mathbf{0} \Rightarrow \nabla \cdot (\rho \mathbf{v}) = \mathbf{0} \Rightarrow \partial \rho = \mathbf{0}; \quad (\partial \rho \in \mathbb{C}).$$

IIIc. With no loss of generality, let $(S(\mathbf{r}, t) = E't + \epsilon(\mathbf{r}, t)t + \beta(\mathbf{r}))$. The only solution of $(i\hbar\Psi^*\partial\Psi = E\rho)$, with E being a constant, is $(\partial\epsilon = 0)$, where $(E = E' + \epsilon)$.

The converse of part **IIIa** is false: The vanishing of $\partial \rho$ does not imply a stationary state. For such a state to be time dependent, from Eq. (47.2), $(\nabla \cdot \rho \mathbf{v}) = 0$) must hold with ρ fixed and \mathbf{v} changing with time. In this special case, the real part of the Schrödinger is a relation between ∇S , ∂S and U with fixed ρ . Note that Eq. (48) is an equation of probability conservation with \mathbf{j} being the probability current density [63].

The equalities obtained for the remaining subsections of this section hold with continuity equation satisfaction: $(\partial \rho + \nabla \cdot (\rho \mathbf{v}) = \mathbf{0})$.

6.2. Equations implied by the orthogonal condition $(\nabla \rho \cdot \nabla S = \mathbf{0})$.

In this subsection, the special relationships are found when the orthogonal condition, defined by $(\nabla \rho \cdot \nabla S = \mathbf{0})$; $\nabla \rho \cdot \nabla S \in \Sigma$, is satisfied. The interpretations given are for one-body states. Note that from Defs. (23), $(\nabla \rho \cdot \nabla S = \mathbf{0})$ and $(\mathbf{u} \cdot \mathbf{v} = \mathbf{0})$ are equivalent.

The logic:

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{0} \Leftrightarrow \ m\nabla \cdot \mathbf{v} = \nabla^2 S = -\frac{\partial \rho_m}{\rho}; \qquad \nabla^2 S \in \mathbf{\Sigma}, \tag{50}$$

follows from the continuity equation, written $(\partial \rho + \nabla \rho \cdot \mathbf{v} + \rho \nabla \cdot \mathbf{v} = \mathbf{0})$, $(m\mathbf{v} = \nabla S)$, and the observation that \mathbf{u} and $\nabla \rho$ are perpendicular, if $(\mathbf{u} \cdot \mathbf{v} = \mathbf{0})$. Applying the operator ∂ to Eq. (50), and using Eqs. (41), we have

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{0} \Rightarrow -\nabla^2 \partial S = \partial \left(\frac{\partial \rho_m}{\rho}\right) \Rightarrow \nabla^2 E = \frac{1}{\zeta_0} \partial F.$$

The Poisson equation (50) reduces to the Laplace equation for the time-independent case.

$$\mathbf{u} \cdot \mathbf{v} = \mathbf{0} \text{ and } \partial \rho = \mathbf{0} \quad \Leftrightarrow \quad \nabla \cdot \mathbf{v} = \nabla^2 S = \mathbf{0}; \qquad \nabla \cdot \mathbf{v} \in \mathbf{\Sigma}.$$
 (51)

For 1-body, steady fluid flow with mass density ρ_m , we have $(\partial \rho_m = \mathbf{0})$, implying that $(\nabla \cdot \mathbf{v} = \mathbf{0})$, so that \mathbf{v} is both irrotational and solenoidal. A mass conserving steady flow of classical mechanics that is both irrotational and solenoidal, and, therefore, incompressible, is called potential flow, where the velocity potential satisfies Laplace's equation.

6.3. Local balance of the fluid momentum $(\rho_m \mathbf{u})$

Applying the gradient sequence $((\nabla) \doteq \nabla_1 + \cdots + \nabla_n)$ to the first continuity Eq. (47), gives the equation sequence $(\nabla[p] = \zeta_0 \partial \nabla \rho)$, where, in each of the *n* equations, $([p] = p_1 + \cdots + p_n)$. Using Eq. (23)× ρ , $(\zeta_0 \nabla \rho = -\rho_m \mathbf{u})$, we obtain the objective:

$$-\nabla[p] = \partial(\rho_m \mathbf{u}),\tag{52}$$

the local balance of fluid momentum $\rho_m(\mathbf{u})$, where (∇) and (\mathbf{u}) step. For a one-body system, if the above equation is for a fluid with mass density ρ_m and velocity field \mathbf{u} , then p is the portion of the total pressure responsible for the local momentum time-derivative $[\partial \rho_m \mathbf{u}](\mathbf{r}, t)$, at the fixed point $(\mathbf{r}, t) \in \mathbb{R}^3 \times \mathbb{R}$. According to Sec. (3), this identifies [p]as a correspondence pressure component from equation. For later convenience, we note, from the derivation of Eq. (52), that

$$m\mathbf{u} = -\zeta_0 \frac{\nabla \rho}{\rho}$$
 and $p = \zeta_0 \partial \rho \Rightarrow Eq. (52).$ (53)

The remaining material from this section are not used later.

6.4. Law of the pressures

Applying the sequence operation $\zeta(\nabla \cdot)$ to Eq. (52) with switched order, we have

$$\zeta_0 \partial \left(\nabla \cdot (\rho \mathbf{u}) \right) = -\zeta \nabla^2 [p].$$

Substituting Eq. (32), we discover

$$\partial P = -\zeta \nabla^2[p],\tag{54}$$

the law of the pressures: A sequence of Poisson equations.

6.5. Laws of the momentae

Multiplying the second continuity equation (47) by ζ_0 , and using $(m\mathbf{v} = \nabla S)$, we have

$$-\zeta_0 \partial \rho = \zeta_0 \nabla \cdot (\rho \nabla S), \quad \text{and} \quad$$

then using the definition: $\theta \doteq -\zeta_0 \ln \rho \Rightarrow -\zeta_0 \partial \rho = \rho \partial \theta$, from (22), gives

$$\rho \partial \theta = \zeta \nabla \cdot (\rho \nabla S), \tag{55}$$

the law of the particle-momentae potentials.

Applying the sequence operator $\zeta_0 \nabla$ to the to the second continuity Eq. (47), written $(\partial \rho = -\nabla \cdot (\rho \mathbf{v}))$, where $[\nabla \cdot \rho \mathbf{v}]$ runs, we obtain

$$\zeta_0 \partial \nabla \rho = -\zeta_0 \nabla \left[\nabla \cdot \rho \mathbf{v} \right].$$

Substituting the second definition (23), written $(-\zeta_0 \nabla \rho = \rho_m \mathbf{u}, \text{ gives})$

$$\partial(\rho_m \mathbf{u}) = \zeta_0 \nabla \left[\nabla \cdot \rho \mathbf{v} \right],\tag{56}$$

the law of the fluid momentae.

6.6. All space momentae, energae and probability density conservation

The proof below gives a relationship between the two energy functions E and F, involving two nondegenerate eigenstates of the Hamiltonian operator. Such states satisfy and do not satisfy the time dependent and the time independent Schrödinger equation, respectively. Both energy quantities have the important property of being conserved over all space for all times. A discussion is then give on how it can be generalized. The same idea holds for $\partial \rho$ and $\rho \partial S$, given Eq. (41).

Theorem 2 (Time dependence of $\partial \rho$ and $\rho \partial S$) Let

$$\begin{bmatrix} e^{-i\varepsilon\circ} \end{bmatrix}(t) - e^{-i\varepsilon t}, \quad [\cos\varepsilon\circ](t) = \cos(\varepsilon t), \quad [\sin\varepsilon\circ](t) = \sin(\varepsilon t)$$

$$\psi_i \doteq e^{-i\varepsilon_i\circ}, \quad |C_1|^2 + |C_2|^2 = 1, \ C_i \in \mathbb{R}, \ \rho_i \doteq \phi_i\phi_i; \ and \ i \in \{1, 2\}$$

Let $\phi_1(\mathbf{r})\psi_1(t)$ and $\phi_2(\mathbf{r})\psi_2(t)$ be values of nondegenerate eigenfunctions of a Hamiltonian operator, at time $t \in \mathbb{R}$ and position $\mathbf{r} \in \mathbb{R}^{3n}$. Let Ψ be defined by the following:

$$\Psi(\mathbf{r},t) \doteq C_1 \phi_1(\mathbf{r}) \psi_1(t) + C_2 \phi_2(\mathbf{r}) \psi_2(t),$$

$$F\rho = C_1 C_2 \phi_1 \phi_2(\varepsilon_2 - \varepsilon_1) \sin[(\varepsilon_1 - \varepsilon_2) \circ] = \frac{\hbar}{2} \partial \rho, \quad and \tag{57}$$

$$E\rho = C_1 C_2 \phi_1 \phi_2(\varepsilon_1 + \varepsilon_2) \cos[(\varepsilon_2 - \varepsilon_1)\circ] + C_1^2 \varepsilon_1 \rho_1 + C_2^2 \varepsilon_2 \rho_2 = -\rho \partial S.$$
(58)

Both of the fluid energy fields have nonlocal conservation in space at all points of time:

$$\hat{\int} F\rho = \hat{\int} \partial\rho = 0 \quad and \quad \hat{\int} E\rho = \hat{\int} \rho \partial S = C_1^2 \varepsilon_1 + C_2^2 \varepsilon_2.$$
(59)

Proof In the following equation sequence, we start with the linear combination wavefunction Ψ , and calculate the left-hand side of Eq. (40), for an *n*-body system:

$$\begin{split} \Psi^* i \partial \Psi &= (C_1 (\phi_1 \psi_1)^* + C_2 (\phi_2 \psi_2)^*) (C_1 \varepsilon_1 \phi_1 \psi_1 + C_2 \varepsilon_2 \phi_2 \psi_2), \\ \Psi^* i \partial \Psi &= C_1^2 \varepsilon_1 \rho_1 + C_2^2 \varepsilon_2 \rho_2 + C_1 C_2 (\varepsilon_1 \phi_1 \phi_2^* \psi_2^* \psi_1 + \varepsilon_2 \phi_2 \phi_1^* \psi_1^* \psi_2), \\ \Psi^* i \partial \Psi &= C_1^2 \varepsilon_1 \rho_1 + C_2^2 \varepsilon_2 \rho_2 + C_1 C_2 \phi_1 \phi_2 \left(\varepsilon_1 e^{i(\varepsilon_2 - \varepsilon_1)^\circ} + \varepsilon_2 e^{-i(\varepsilon_2 - \varepsilon_1)^\circ} \right). \end{split}$$

Using (40) and Defs. (41), we obtain energy Eq. (57) and (58). Since the functions ϕ_1 and ϕ_2 are orthonormal, the spacial energy averages, obtained by integrating (57) and (58) over all space \mathbb{R}^{3n} and summing over all spin variables, are given by Eqs. (59).

If we repeat the derivation in the proof with the following change $C_2 \rightarrow -C_2$, then the sign of the C_1C_2 terms in Eq. (57) and (58) change. If, instead, we make the following change $C_2 \rightarrow iC_2$, nothing dramatic happens: The β terms in Eq. (58), for E, and Eq. (57), for F, switch places. For the general case, using a set of m orthonormal spatial wavefunctions $\{\phi_1, \dots, \phi_m\}$ that are eigenfunctions of a single Hamiltonian operator, the same result is obtained: E and F are conserved over space. This result is easily seen to follow because, with no loss of generality, the members of the set $\{\phi_1, \dots, \phi_m\}$ are mutually orthogonal. The same result is obtained for a wavefunction defined by an infinite sequence of orthonormal eigenfunctions, since each term of the corresponding sequences, for both E and F, are conserved, and such sets are complete.

7. Energy Equation and Schrödinger Equation Equivalence

7.1. Eqs. (45) and Schrödinger Eq. (11) equivalence

Theorem 1 proved that the imaginary part of the Schrödinger Eq. (11) is equivalent to five other equations, including the second energy Eq. (45): $(F\rho = p)$. The following proof demonstrates that the real part of the Schrödinger Eq. (11) is equivalent to Eq. (45.1). Since it is convenient to do so, an alternate proof is given for the equivalence of the imaginary parts. The theorem from Appendix A proves that the two equations of Bohmian mechanics are equivalent to the Schrödinger equation.

Theorem 3 (Energy Eq. and Schrödinger Eq. Equivalence) In the L^2 Hilbert space, the real and imaginary parts of the Schrödinger Eq. (11) are equivalent to first and second equations of the 2-set Eqs. (45), respectively, where the wavefunction Ψ satisfies Def. (15.1), the scalar fields (ρ , S) are such that the velocities **u** and **v** satisfy Eqs. (23), the pressure P satisfies Eq. (32.1), and E and F satisfy Eqs. (41).

Proof Since the two equation $(A_1 = B_1)$ and $(A_2 = B_2)$ imply $(A_1 = A_2)$ and $(B_1 = B_2)$ are equivalent, Eqs. (20) and (43) imply that the Schrödinger equation, in the form $(i\hbar\Psi^*\partial\Psi = \Psi^*\hat{H}\Psi)$, is equivalent to

$$-\rho\partial S + i\zeta_0\partial\rho = \frac{1}{2}\rho_m u^2 + P + \frac{1}{2}\rho_m v^2 + U\rho + ip.$$

Substituting Eqs. (41) gives the 2-set Eqs. (45), from the real and imaginary parts.

(Discussion about the energy Eq. (45)) For 1-body states, Eq. (45) is the Bernoulli equation of fluid dynamics with ρ_m being the mass density for one-body states. However, this equations is being applied in a more general way, since quantum flows are compressible and have variable mass. Using (45.1), flows for one-body real valued wavefunctions have been investigated [1,54], where the generalized Bernoulli equation (45) describes compressible, irrotational, steady flow with *local* variable mass and ($\mathbf{v} = \mathbf{0}$). Over all space, mass is conserved, because the rate of mass creation from the sources are equal to the rate of mass annihilation from the sinks. Also, each fluid element has a constant energy per mass E/m. Having local variable mass is required for the flow of the 1s state of the hydrogen atom to be steady with zero angular momentum, since such flows must be radially directed, and a radially directed flow with conserved mass requires infinite mass.

As given above in Eq. (46), $\langle E \rangle$ is the expectation value $\langle \hat{H} \rangle$ of the Hamiltonian operator. By considering the following equation for a one-body system with a fluid interpretation, it is easily seen that the proper interpretation of the vector field E for these systems is that E/m is the energy per mass:

$$\langle E \rangle = \int (E/m)\rho_m = \int E\rho.$$

Similarly, E/q can be interpreted as the energy per charge for a system with identical particles, each with charge q. Hence, E, despite having an energy unit, can be interpreted to be the energy per mass and energy per charge that is multiplied by the total mass and total charge, respectively, and these are intensive variables. However, for steady states, where ρ is normalized to unity, the field E is uniform, giving $(E = \langle E \rangle)$, where $\langle E \rangle$ is an extensive variable. Hence, E can be considered intensive or extensive, depending on the context. For *n*-body systems, for proper scaling with respect to the number of particles, ρ should be normalized to n instead of unity, and E/m becomes E/(nm). If this is done, a similar interpretation for the energy can be given as in the one-body case. With the identification of Eq. (45) as a generalization of the Bernoulli equation of fluid dynamics, the function (P) is a correspondence variables from equation for the pressure, with an energy per volume unit. Since, from Eq. (32), it integrates to zero, from definition 3 of Sec, 3, it a free, or distributive, variable, and a system has a total of zero [P] energy. The presence of [P] as an energy does not require the input of energy, only the separation of zero energy into positive and negative parts, where the energy is distributed over space. For any change a state undergoes, the total free energy from [P] is conserved as zero.

We also identify two velocity vectors (**u**) and (**v**) and their kinetic energy as correspondence variables from equation (45). However, this kinetic energy identification is less than perfect, since the formula for the kinetic energy, Eq. (24), involves the square of the complex valued vector-field ($\mathbf{v} + i\mathbf{u}$), instead of the square of a real valued vectorfield, as is done is classical mechanics. Hence, the kinetic energy definition has a non classical element for certain complex valued wavefunctions. While some non classical elements can be tolerated, we wish to minimize such elements. In the next subsection an energy equation is derived where the kinetic energy is calculated from the singe velocity vector ($\mathbf{w} = \mathbf{u} + \mathbf{v}$), instead of a sum of two kinetic energies from the 2-set (\mathbf{v}, \mathbf{u}).

7.2. Single velocity vector energy equation

A way to remove the nonclassical kinetic energy problem, mentioned above, is given below by Theorem 4, where a single velocity vector \mathbf{w} is defined to replace the 2–set (\mathbf{v}, \mathbf{u}) . The alternate energy equation also introduces a new term that is proportional to the volumetric dilatation rate $(\nabla \cdot \mathbf{v})$. The 2–set of energies (E, F) are also combined into one energy \overline{E} . A discussion about the energy equation is also given. The following definition and lemma are used for the theorem below.

Definition 6

$$\bar{E} \doteq E + F = -\partial(S + \theta), \quad \mathbf{w} \doteq \mathbf{u} + \mathbf{v}.$$
 (60)

Lemma 6

Definition (32.1) for P and (23) for $\mathbf{u} \Rightarrow P = -\rho_m \mathbf{u} \cdot \mathbf{u} + \eta \nabla \cdot \mathbf{u}$ (61)

Definition (32.2) for p and (23) for $\mathbf{u} \Rightarrow p = -\rho_m \mathbf{u} \cdot \mathbf{v} - \eta \nabla \cdot \mathbf{v}$. (62)

Proof Using definition (32) for P and p, respectively, definition above for η , and (23) for **u**, we have

$$P = \zeta_0 \nabla \cdot (\rho \mathbf{u}) = \zeta_0 \rho \nabla \cdot \mathbf{u} + \zeta_0 \nabla \rho \cdot \mathbf{u} = \eta \nabla \cdot \mathbf{u} - \rho_m \mathbf{u} \cdot \mathbf{u}, \text{ and}$$
$$p = -\zeta_0 \nabla \cdot (\rho \mathbf{v}) = -\zeta_0 \rho \nabla \cdot \mathbf{v} - \zeta_0 \nabla \rho \cdot \mathbf{v} = -\eta \nabla \cdot \mathbf{v} + \rho_m \mathbf{u} \cdot \mathbf{v}.$$

Theorem 4 (Single Velocity Total Energy Representation) The following equations are equivalent to the equation 2-set (45), and, therefore, also the Schödinger equation from theorem 3:

$$\bar{E}\rho = \frac{1}{2}\rho_m w^2 + P - \eta(\nabla \cdot \mathbf{v}) + U\rho, \quad F\rho = p.$$
(63)

Proof Using the definition for (\mathbf{w}) from Eqs. (60) and Def. (62), we discover that

$$\frac{1}{2}\rho_m w^2 - \eta(\nabla \cdot \mathbf{v}) = \frac{1}{2}\rho_m u^2 + \frac{1}{2}\rho_m v^2 + \rho_m \mathbf{u} \cdot \mathbf{v} - \eta(\nabla \cdot \mathbf{v}) = \frac{1}{2}\rho_m u^2 + \frac{1}{2}\rho_m v^2 + p.$$

Substituting this result into Eq. (63) and using $(F\rho = p)$, we find that the resulting equation is the first equation of (45) with $F\rho$ added to both sides. Hence, this resulting equation is equivalent the first equation of the 2-set (63). (The first equations of (45) and (63) are not equivalent.)

(Discussion about the energy Eq. (63)) Besides combing the 2-set of velocities (\mathbf{u}, \mathbf{v}) vectors into one, Eq. (63) combines the two energies into one \overline{E} . From Eq. (63.2) and Eq. (34), F is a free particle energy. Hence, while F can change the energy locally, it has no effect on the total energy. Note that because the proof uses $(F\rho = p)$, Eq. (63.1), by itself, is not equivalent to the real part of the Schödinger equation.

8. Euler fluid equations for one-body states

Definition 7 (Quantum flow of one-body states) A flow $(\mathcal{F} \doteq (p, \phi, \mathbf{x}))$ is a set containing a pressure p, mass density ϕ and velocity \mathbf{x} , with a given body force. A quantum flow $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$, of a 1-body state, is a flow such that there exists a pair of scalar fields (ρ, S) , where the velocities \mathbf{u} and \mathbf{v} satisfy Eqs. (23), and the pressure Psatisfies Eq. (32.1). Also, E, F and p use the definitions above. If there is no chance of confusion, some or all of the variables ρ_m , P, \mathbf{u} , and \mathbf{v} in the notation $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$ can be suppressed, and a body force can be added.

An improper steady flow is a flow where $(\partial \rho = \mathbf{0})$ and the corresponding wavefunction is not a stationary state. Since, we will not consider such flows, given a pair of scalar fields (ρ, S) determining a wavefunction Ψ and flow \mathcal{F} , the following statements are considered equivalent: $(\partial \rho = \mathbf{0})$, the flow \mathcal{F} is steady, and the wavefunction Ψ is a stationary state. For such a case, the members of the continuity set \mathbb{C} of Theorem 1 vanish.

A smooth flow is a flow $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$ such that the sequence $(\nabla S \cdot \nabla \rho)$, or, equivalently $(\mathbf{u} \cdot \mathbf{v} = \mathbf{0})$, vanish identically. As in Sec. (6.2), the equation $(\nabla S \cdot \nabla \rho = \mathbf{0})$ is called the orthogonal condition.

For smooth flow stationary states, including one-electron atoms, giving $(\nabla \rho \cdot \mathbf{v} = \mathbf{0})$, it follows from continuity Eq. (47.2) that \mathbf{v} is solenoidal, also F vanish from Theorem 1. Hence, in this case, Eqs. (63) reduces to (45).

8.1. Euler fluid equations of motion for one-body states

In this subsection, for one-body states, four Euler equations are derived that are equivalent to energy equations (45.1). The flow type identified is compressible, irrotational flow with a velocity field \mathbf{w} that is a sum of two velocity fields, \mathbf{u} and \mathbf{v} . The velocity fields \mathbf{u} and \mathbf{v} are not, in general, perpendicular. The \mathbf{u} velocity does not satisfy a continuity equation. For stationary states, the flow is inviscid. Of the four derived time dependent equations, only one has the force $-\nabla p$, and one has bulk a viscosity $\nabla[\eta(\nabla \cdot \mathbf{v}])$ involving velocity \mathbf{v} . Three out of the four Euler equations are missing one or more terms involving both velocity fields, $\mathbf{v}\nabla \cdot (\rho_m \mathbf{u})$ being an example. All terms of the Euler equation are present for stationary states. Interpretations of the results is given at the end of the section.

The next lemma, that can be proved by inspection, is needed for theorems that follow. The lemma uses definitions (23), (32), and (33).

Lemma 7 (Mass flux equations)

$$\nabla \cdot (\rho_m \mathbf{u}) = \frac{1}{\zeta} P \quad and \quad \mathbf{u} = -\zeta \frac{\nabla \rho}{\rho} \quad \Rightarrow \quad \nabla \cdot (\rho_m \mathbf{u}) \mathbf{u} = -\frac{P}{\rho} \nabla \rho = \frac{1}{\zeta} P \mathbf{u}, \tag{64}$$

$$\nabla \cdot (\rho_m \mathbf{v}) = -\frac{1}{\zeta} p \quad and \quad \mathbf{v} = \zeta \frac{\mathbf{Q}}{\rho} \quad \Rightarrow \quad \nabla \cdot (\rho_m \mathbf{v}) \mathbf{v} = -\frac{p}{\rho} \mathbf{Q} = -\frac{1}{\zeta} p \mathbf{v}, \tag{65}$$

$$\nabla \cdot (\rho_m \mathbf{u}) = \frac{1}{\zeta} P \quad and \quad \mathbf{v} = \zeta \frac{\mathbf{Q}}{\rho} \quad \Rightarrow \quad \nabla \cdot (\rho_m \mathbf{u}) \mathbf{v} = \frac{P}{\rho} \mathbf{Q} = \frac{1}{\zeta} P \mathbf{v}, \tag{66}$$

$$\nabla \cdot (\rho_m \mathbf{v}) = -\frac{1}{\zeta} p \quad and \quad \mathbf{u} = -\zeta \frac{\nabla \rho}{\rho} \Rightarrow \quad \nabla \cdot (\rho_m \mathbf{v}) \mathbf{u} = -\frac{p}{\rho} \nabla \rho = -\frac{1}{\zeta} p \mathbf{u}.$$
(67)

In the derivation of differential forms of the equations of motion of fluids, the special case of conserved mass is assumed [62, 64, 65]. Since we have variable mass flows, we need to make sure we have the correct momentum time-derivative. This is disposed of in the next Lemma. The Lemma demonstrates that if mass is not conserved for a state with velocity \mathbf{w} , then the total momentum time-derivative has the additional term $\mathbf{w}(\partial \rho_m + \nabla \cdot (\rho \mathbf{w}))$. If \mathbf{w} is given by ($\mathbf{w} = \mathbf{u} + \mathbf{v}$), and $\mathbf{w}(\partial \rho_m + \nabla \cdot (\rho \mathbf{v}))$ vanish, then the additional term reduces to $\mathbf{w}[\partial \rho_m + \nabla \cdot (\rho \mathbf{u})]$.

Lemma 8 (Momentum time derivatives)

Let \mathbb{V} be a time dependent subset of space \mathbb{R}^3 with surface S, and occupied by a fluid with variable mass. Let the velocity field of the fluid be $(\mathbf{w} : \mathbb{R}^3 \times \hat{\mathbb{R}} \to \mathbb{R}^3)$, where $\mathbf{r} \in \mathbb{R}^3$ and $t \in \hat{\mathbb{R}}$ are spatial and temporal variables, respectively. Let the path of a fluid particle be $(\mathbf{r} = \mathbf{r}(t))$, giving the composition $(\mathbf{w} = \mathbf{w}(\mathbf{r}(t)))$, thereby, permitting total time derivatives. Let \mathcal{D} be the total time derivative, that is expressed as the material derivative for steady flow when the chain rule is used. Let the tensor $\nabla \mathbf{A} \cdot \mathbf{B}$ be

$$(\nabla \mathbf{A} \cdot \mathbf{B})_i \doteq \nabla A_i \cdot \mathbf{B} = \mathbf{B} \cdot \nabla A_i, \quad i = 1, 2, 3, \qquad \mathbf{A} : \mathbb{R}^3 \to \mathbb{R}, \quad \mathbf{B} : \mathbb{R}^3 \to \mathbb{R},$$

giving the spacial part of the material derivative $\nabla \mathbf{w} \cdot \mathbf{w}$ as a special case.

$$\mathcal{D}\!\!\int_{\mathbb{V}} \mathbf{Y} = \int_{\mathbb{V}} \mathcal{D}\mathbf{Y} + \mathbf{Y}(\nabla \cdot \mathbf{w}); \quad \mathbf{Y} : \hat{\mathbb{R}} \longrightarrow \mathbb{R}^3, \tag{68}$$

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho = \int_{\mathbb{V}} \partial\rho + \nabla \cdot (\rho \mathbf{w}), \tag{69}$$

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho \mathbf{w} = \int_{\mathbb{V}} \mathcal{D}(\rho \mathbf{w}) + \rho \mathbf{w} (\nabla \cdot \mathbf{w}), \tag{70}$$

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho \mathbf{w} = \int_{\mathbb{V}} \rho \mathcal{D}\mathbf{w}, \quad if \ \mathcal{D}\!\!\int_{\mathbb{V}} \rho = \mathbf{0}, \quad and$$
(71)

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho \mathbf{w} = \int_{\mathbb{V}} \partial(\rho \mathbf{w}) + \rho \nabla \mathbf{w} \cdot \mathbf{w} + \nabla \cdot (\rho \mathbf{w}) \mathbf{w} = \int_{\mathbb{V}} \partial(\rho \mathbf{w}) + \int_{\mathcal{S}} \rho \mathbf{w} (\mathbf{w} \cdot \hat{\mathbf{n}}).$$
(72)

If \mathbf{w} is irrotational, then

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho \mathbf{w} = \int_{\mathbb{V}} \partial(\rho \mathbf{w}) + \frac{1}{2} \rho \nabla w^2 + \nabla \cdot (\rho \mathbf{w}) \mathbf{w}.$$
(73)

Let $(\mathbf{w} \doteq \mathbf{u} + \mathbf{v})$. If \mathbf{w} is irrotational and $(\partial \rho + \nabla \cdot (\rho \mathbf{v}) = \mathbf{0})$, then

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho \mathbf{w} = \int_{\mathbb{V}} \rho \partial \mathbf{w} + \frac{1}{2} \rho \nabla w^2 + \nabla \cdot (\rho \mathbf{u}) \mathbf{w}.$$
(74)

Proof Let $\mathbb{V} : \hat{\mathbb{R}} \longrightarrow \mathcal{P}(\mathbb{R}^3)$, where $\hat{\mathbb{R}}$ is temporal and $\mathcal{P}(\mathbb{R}^3)$ is the power set of \mathbb{R}^3 . In other words, the value $\mathbb{V}(t)$ is the Cartesian subspace $\mathbb{V} \in \mathbb{R}^3$ at time $t \in \hat{\mathbb{R}}$. Let \mathbf{w} be the velocity field of a compressible fluid that occupies $\mathbb{V}(t)$ at time t. Let $\mathbf{Y} : \mathbb{R}^3 \to \mathbb{R}$. The time derivative of the *j*th Cauchy sum of the integral $\int_{\mathbb{V}} \mathbf{Y}$, involving a partitioning into m_j subsets: $(\mathbb{V} = \mathbb{V}_1 \cup \mathbb{V}_2 \cdots \mathbb{V}_{m_j})$ with corresponding volumes $(V = V_1 \cup V_2 \cdots V_{m_j})$, is

$$\mathcal{D}\sum_{i=1}^{m_j} \mathbf{Y}(\mathbf{r}_i) V_i = \sum_{i=1}^{m_j} \left[\mathcal{D}\mathbf{Y}(\mathbf{r}_i) \right] V_i + \sum_{i=1}^{m_j} \mathbf{Y}(\mathbf{r}_i) \dot{V}_i; \qquad \dot{V}_i \doteq \mathcal{D}V_i, \tag{75}$$

where **Y** and V_i are time dependent composites involving $(\mathbf{r} = \mathbf{r}(t))$. Also, $\mathbf{r}_i \in \mathbb{V}_i$, and $m_j \longrightarrow \infty$ as $j \longrightarrow \infty$. By the definition of the divergence, in the limit of $j \longrightarrow \infty$, we have

$$\lim_{m_j \to \infty} \sum_{i=1}^{m_j} [\mathbf{Y} \dot{V}_i](\mathbf{r}_i) = \lim_{m_j \to \infty} \sum_{i=1}^{m_j} [\mathbf{Y} (\nabla \cdot \mathbf{w}) V_i](\mathbf{r}_i),$$

where $\nabla \cdot \mathbf{w}$ is the fluid volumetric dilation. Because both functions have the same limit, the rhs of this equation, without the limit, can replace the term on the rhs of Eq. (75), giving the same result for the limit. Hence

$$\mathcal{D}\sum_{i=1}^{m_j} \mathbf{Y} V_i = \sum_{i=1}^{m_j} (\mathcal{D}(\mathbf{Y}) + \mathbf{Y}(\nabla \cdot \mathbf{w})) V_i.$$

By definition of Riemmann integration, we have Eq. (68).

Substituting $(\rho \mathbf{w} = \mathbf{Y})$ into Eq. (68), we have

(70):
$$\mathcal{D}_{\int_{\mathbb{V}}} \rho \mathbf{w} = \int_{\mathbb{V}} \mathcal{D}(\rho \mathbf{w}) + \rho \mathbf{w} (\nabla \cdot \mathbf{w}).$$

Eq. (71) follows from the given mass conservation condition Eq. (69), and

$$\mathcal{D}(\rho \mathbf{w}) + \rho \mathbf{w} (\nabla \cdot \mathbf{w}) = \rho \mathcal{D} \mathbf{w} + (\mathcal{D}\rho + \rho (\nabla \cdot \mathbf{w})) \mathbf{w}.$$

Let **w** be given by the composite ($\mathbf{w} = \mathbf{w}(t, \mathbf{r}(t))$; ($\mathbf{r} \in \mathbb{R}^3$). Using the chain rule, the total derivative of $\rho \mathbf{w}$ is, by definition, the material derivative of $\rho \mathbf{w}$:

$$\mathcal{D}(\rho \mathbf{w}) = \rho \mathcal{D} \mathbf{w} + \mathbf{w} \mathcal{D} \rho = \rho \partial \mathbf{w} + \rho \nabla \mathbf{w} \cdot \mathbf{w} + \mathbf{w} \partial \rho + \mathbf{w} \nabla \rho \cdot \mathbf{w}.$$

Hence, $\mathcal{D}(\rho \mathbf{w}) + \rho \mathbf{w}(\nabla \cdot \mathbf{w}) = \partial(\rho \mathbf{w}) + \rho \nabla \mathbf{w} \cdot \mathbf{w} + \nabla \cdot (\rho \mathbf{w}) \mathbf{w}.$

Substituting into Eq. (70), we have (72.1). Equation. (73) follows from the vector calculus equality $(2\nabla \mathbf{w} \cdot \mathbf{w} = \nabla w^2)$, for irrotational field \mathbf{w} . Equation. (69), follows from Eq. (68) with \mathbf{Y} replaced by ρ . Subtracting the continuity equation, multiplied by \mathbf{w} : $(\mathbf{w}\partial\rho_m + \nabla \cdot (\rho_m \mathbf{v})\mathbf{w} = \mathbf{0})$, from Eq. (73), gives Eq. (74).

Let $(\mathbf{p} \doteq \rho \mathbf{w})$. For Eq. (72.2) we only need to to prove that each of three components of the vector equation, given by

$$\int_{\mathcal{S}} \rho \mathbf{w}(\mathbf{w} \cdot \hat{\mathbf{n}}) = \int_{\mathbb{V}} \rho \nabla \mathbf{w} \cdot \mathbf{w} + \nabla \cdot (\rho \mathbf{w}) \mathbf{w},$$

are equal:

$$\left(\int_{S} \rho \mathbf{w}(\mathbf{w} \cdot \hat{\mathbf{n}}) \right)_{i} = \int_{S} p_{i} \mathbf{w} \cdot \hat{\mathbf{n}} = \int_{\mathbb{V}} \nabla \cdot (p_{i} \mathbf{w}) = \int_{\mathbb{V}} \nabla p_{i} \cdot \mathbf{w} + p_{i} \nabla \cdot \mathbf{w}$$
$$= \int_{\mathbb{V}} \nabla (\rho w_{i}) \cdot \mathbf{w} + \rho w_{i} \nabla \cdot \mathbf{w} = \int_{\mathbb{V}} \rho \nabla w_{i} \cdot \mathbf{w} + w_{i} (\nabla \rho \cdot \mathbf{w}) + w_{i} (\rho \nabla \cdot \mathbf{w})$$
$$= \int_{\mathbb{V}} \rho \nabla w_{i} \cdot \mathbf{w} + \nabla \cdot (\rho_{m} \mathbf{w}) w_{i} = \left(\int_{\mathbb{V}} \rho \nabla \mathbf{w} \cdot \mathbf{w} + \nabla \cdot (\rho \mathbf{w}) \mathbf{w} \right)_{i}.$$

Next we obtain the first Euler equation of the set of four. This one is the simplest one. In the special case of steady flow, the equation reduces to the complete Euler equation with variable mass.

Theorem 5 (Equation of motion for one-body states part 1)

For quantum flows $\mathcal{F}(\rho_m, P, \mathbf{u}, \mathbf{v})$ of one-body states, we have the following:

1) The first energy equation of (45) is equivalent to the Euler equation:

$$\rho_m \partial \mathbf{v} + \frac{1}{2} \rho_m \nabla \left(u^2 + v^2 \right) + \nabla P + \nabla \cdot \left(\rho_m \mathbf{u} \right) \mathbf{u} + \rho \nabla U = \mathbf{0}, \tag{76}$$

that is, (45.1) \Leftrightarrow (76). 2) (76) $\Leftrightarrow Re(i\hbar\Psi^*\partial\Psi = \Psi^*\hat{H}\Psi).$

Proof Applying the gradient-sequence operation (∇) to the energy equation $(45) \times \rho^{-1}$, we have the equation sequence:

$$\nabla E = \frac{1}{2}m\nabla\left(u^2 + v^2\right) + \nabla\left(P\rho^{-1}\right) + \nabla U.$$
(77)

Since this equation with a constant of integration is equivalent to Eq. (45.1) for E, we only need to prove that $(77) \Leftrightarrow (76)$ to prove part 1 of the theorem. Multiplying Eq. (77) by ρ , followed by substituting of Eq. (42).1× ρ : ($\rho \nabla E = \rho_m \partial \mathbf{v}$), we have

$$\rho_m \partial \mathbf{v} + \frac{1}{2} \rho_m \nabla \left(u^2 + v^2 \right) + \rho \nabla \left(P \rho^{-1} \right) + \rho \nabla U = \mathbf{0}.$$
(78)

Since this equation is equivalent to Eq. (77), we only need to prove that this equation is equivalent to Eq. (76), that is, (78) \Leftrightarrow (76). Substituting implication (64) into the identity

$$\rho \nabla \left(\frac{P}{\rho}\right) = \nabla P - P \frac{\nabla \rho}{\rho},\tag{79}$$

gives the following equality:

$$\rho \nabla \left(P \rho^{-1} \right) = \nabla P + \nabla \cdot \left(\rho_m \mathbf{u} \right) \mathbf{u}.$$
(80)

Substituting this equation into the Eq. (78), we obtain Eq. (76). Hence $(78) \Leftrightarrow (76)$. For part **2**, from Theorem 3 we have $\left(\operatorname{Re}(i\hbar\Psi^*\partial\Psi) = \operatorname{Re}(\Psi^*\hat{H}\Psi)\right) \Leftrightarrow$ (45.1) combining this with part $1: (45.1) \Leftrightarrow (76)$, along with the transitive and symmetric property of equivalence relations, gives statement $2:(76) \Leftrightarrow (\operatorname{Re}(i\hbar\Psi^*\partial\Psi))$ $= \operatorname{Re}(\Psi^* \hat{H} \Psi) \Big).$

It follows from Lemma 8, Eq. (73), that the following terms are missing in Euler Eq. (76): $\rho_m \partial \mathbf{u}, \rho_m \nabla \mathbf{u} \cdot \mathbf{v}$ and $\nabla \cdot (\rho_m \mathbf{u}) \mathbf{v}$. The next theorem obtains an equation of motion that improves the situation by obtaining the first two. A second derived equation of motion includes the force $-\nabla p$, but at the cost of adding the term $\nabla \cdot (\rho_m \mathbf{v}) \mathbf{u}$.

Theorem 6 (Equation of motion for one-body states part 2) Consider the following equations

$$\rho_m \partial \mathbf{w} + \frac{1}{2} \rho_m \nabla w^2 + \nabla P + \rho \nabla U + \nabla \cdot (\rho_m \mathbf{u}) \mathbf{u} = \eta \nabla (\nabla \cdot \mathbf{v}).$$
(81)

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho_m \mathbf{w} - \int_{\mathbb{V}} \nabla \cdot (\rho_m \mathbf{u}) \mathbf{v} = \int_{\mathbb{V}} -\nabla P - \rho \nabla U + \eta \nabla (\nabla \cdot \mathbf{v}) \,. \tag{82}$$

$$\partial(\rho_m \mathbf{u}) + \rho_m \partial \mathbf{v} + \frac{1}{2} \rho_m \nabla \left(u^2 + v^2 \right) + \nabla (P + p) + \nabla \cdot (\rho_m \mathbf{u}) \mathbf{u} + \rho \nabla U = \mathbf{0}.$$
(83)

$$-\int_{\mathbb{V}} \nabla \cdot (\rho_m \mathbf{v}) \mathbf{u} + \mathcal{D} \int_{\mathbb{V}} \rho_m \mathbf{w} - \int_{\mathbb{V}} \nabla \cdot (\rho_m \mathbf{u}) \mathbf{v} + \rho_m \nabla (\mathbf{u} \cdot \mathbf{v})$$
$$= \int_{\mathbb{V}} -\nabla P - \nabla P - \nabla P - \rho \nabla U + \eta \nabla (\nabla \cdot \mathbf{v}).$$
(84)

For one-body quantum flows $\mathcal{F}(\rho_m, P, \mathbf{u}, \mathbf{v})$, under the condition of the satisfaction of continuity Eq. (47.2), the following three statements are equivalent: **1)** Eq. (76), **2)** Eq. (81), and **3)** Eq. (83). Also, Eqs. (81) and (82) are equivalent; Eqs. (83) and (84) are equivalent.

Proof First we show that $(76) \Leftrightarrow (81)$, giving $\mathbf{1} \Leftrightarrow \mathbf{2}$. Since, from Theorem 1, Eq. (45).1: $(F\rho = p)$ is equivalent to the continuity Eq. (47) displayed above, implication $(62) \times \rho^{-1}$ holds with $p\rho^{-1}$ replaced by F. Thus, continuity satisfaction implies

$$\nabla F = m\nabla(\mathbf{u} \cdot \mathbf{v}) - \zeta_0 \nabla(\nabla \cdot \mathbf{v}).$$

Using the Eq. (42.2), $(-\nabla F = m\partial \mathbf{u})$, and definition $(\eta = \zeta_0 \rho)$, we can write this equation, after multiplication by ρ , as

$$\rho_m \partial \mathbf{u} + \rho_m \nabla (\mathbf{u} \cdot \mathbf{v}) = \eta \nabla (\nabla \cdot \mathbf{v}). \tag{85}$$

Adding this result to Eq. (76) gives (81). Hence $(\underline{76}) \Leftrightarrow (\underline{81})$, giving $\underline{1} \Leftrightarrow \underline{2}$. Next we show that $(\underline{76}) \Leftrightarrow (\underline{83})$, giving $\underline{1} \Leftrightarrow \underline{3}$. from Theorem 1, the continuity Eq. (47) displayed above is equivalent to $(p = \zeta_0 \partial \rho)$. From logic (53), we have Eq. (52): $(\partial(\rho_m \mathbf{u})) + \nabla p = \mathbf{0})$. Adding this equation to Eq. (76), gives Eq. (83). Hence $\underline{1} \Leftrightarrow \underline{3}$. By the transitive and symmetric property of equivalence relations, we have $\underline{1} \Leftrightarrow \underline{2} \Rightarrow$ $\underline{2} \Leftrightarrow \underline{1}$ and $\underline{1} \Leftrightarrow \underline{3} \Rightarrow \underline{2} \Leftrightarrow \underline{3}$

It follows from Eq. (74) that Eqs. (81) and (82) are equivalent. For the same reason, Eqs. (83) and (84) are equivalent, after making the substitution $(\mathbf{u}\partial\rho_m = -\nabla(\rho_m \mathbf{v})\mathbf{u})$.

For equation of motion (81), the only term missing is $\nabla \cdot (\rho_m \mathbf{u}) \mathbf{v}$. The extra term $\eta \nabla (\nabla \cdot \mathbf{v})$ can be interpreted as a sort of bulk viscosity, discussed below.

The next theorem, obtains all terms of the time derivative of the momentum. This is achieved with additional forces.

Theorem 7 (Equation of motion for one-body states part 3) Let the subspace $(\mathbb{V} \in \mathbb{R}^3)$ be occupied by a fluid particle that satisfies the Euler Eq. (83). Let $(\mathcal{P} \doteq -P)$.

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho_m \mathbf{w} = \int_{\mathbb{V}} -\nabla P - \rho \nabla U + \nabla [\eta \left(\nabla \cdot \mathbf{v}\right)] + \frac{1}{\zeta} \Big(P \mathbf{v} + \mathcal{P} \mathbf{u} + (\mathbf{v} \cdot \mathbf{u}) \rho_m \mathbf{u} \Big). \tag{86}$$

Proof It follow from theorem (7) that we only need to prove that Eq. (86) is equivalent to Euler Eq. (81):

$$\rho_m \partial \mathbf{w} + \frac{1}{2} \rho_m \nabla w^2 + \nabla \cdot (\rho_m \mathbf{u}) \mathbf{u} = -\nabla P - \rho \nabla U + \eta \nabla (\nabla \cdot \mathbf{v}).$$

For treating the term with factor η , consider the development:

$$\nabla \eta (\nabla \cdot \mathbf{v}) = \zeta_0 \nabla \rho \nabla \cdot \mathbf{v} = -\rho_m \mathbf{u} \nabla \cdot \mathbf{v} = m \mathbf{u} (-\rho \nabla \cdot \mathbf{v}) = m \mathbf{u} \nabla \rho \cdot \mathbf{v} - m \mathbf{u} \nabla \cdot (\rho \mathbf{v}),$$

iving

giving

$$\nabla \eta (\nabla \cdot \mathbf{v}) - m \mathbf{u} \nabla \rho \cdot \mathbf{v} + \nabla \cdot (\rho_m \mathbf{v}) \mathbf{u} = \mathbf{0}.$$
(87)

The second term on the lbs satisfies:

$$-m\mathbf{u}\nabla\rho\cdot\mathbf{v} = (m\mathbf{u}\rho)\frac{1}{\zeta}\left(\frac{-\zeta\nabla\rho}{\rho}\right)\cdot\mathbf{v} = (\rho_m\mathbf{u})\frac{1}{\zeta}(\mathbf{u}\cdot\mathbf{v}),$$

giving

$$\nabla \eta (\nabla \cdot \mathbf{v}) + \frac{1}{\zeta} (\mathbf{u} \cdot \mathbf{v}) \rho_m \mathbf{u} + \nabla \cdot (\rho_m \mathbf{v}) \mathbf{u} = \mathbf{0}.$$

Substituting equation (67): $(\nabla \cdot (\rho_m \mathbf{v})\mathbf{u} = -\zeta^{-1}p\mathbf{u})$ and then adding equation. (66): $(\nabla \cdot (\rho_m \mathbf{u})\mathbf{v} = \zeta^{-1}P\mathbf{v})$, we have

$$\nabla \cdot (\rho_m \mathbf{u}) \mathbf{v} = \nabla \eta (\nabla \cdot \mathbf{v}) + \frac{1}{\zeta} \Big((\mathbf{u} \cdot \mathbf{v}) \rho_m \mathbf{u} + P \mathbf{v} - p \mathbf{u} \Big).$$
(88)

Substituting into Euler Eq. (81) and using the definition $(\mathcal{P} \doteq -P)$, we have

$$\rho_m \partial \mathbf{w} + \frac{1}{2} \rho_m \nabla w^2 + \nabla \cdot (\rho_m \mathbf{u}) \mathbf{w} = -\nabla P - \rho \nabla U + \nabla (\eta \nabla \cdot \mathbf{v}) + \Gamma_{\rho}[\mathbf{u}, \mathbf{v}];$$
$$\Gamma_{\rho}[\mathbf{u}, \mathbf{v}] = \frac{1}{\zeta} (\mathbf{u} \cdot \mathbf{v}) \rho_m \mathbf{u} + \zeta^{-1} P \mathbf{v} + \zeta^{-1} \mathcal{P} \mathbf{u}$$

Eq. (86) follows from Eq. (74).

8.2. Correspondence variables by Euler Equations

We identify, by definition **2** of Sec 3, the mass density ρ_m , pressure P, and velocity $(\mathbf{u}+\mathbf{v})$ as corresponding variables from Eq. (86). The pressure P is not a thermodynamic pressure, because, being a free energy, it is not positive definite. Euler Eq. (86) is interpreted as a generalization of the other three Euler equations, with Euler Eq. (76) being the most specialized. Consider equality (88), if this equality is not satisfied, there can be solutions of Eq. (86) that do not satisfy the other Euler equations. This specialization, in the form of relationships between variables, explains why certain terms, especially $\nabla \cdot (\rho_m \mathbf{u}) \mathbf{v}$, are missing in the specialized Euler equations: The missing terms cancel.

For equation of motion (84), the force $-\nabla p$ is included, but at the cost of introducing the term $-\nabla \cdot (\rho_m \mathbf{u})\mathbf{v}$. Because of this shortcoming, we lack sufficient evidence to interpret P as a corresponding pressure. However, Eq. (52) does identify

the series [p] as a force. The first definitions from Eqs. (32) and (33) suggests that the sign of p is wrong, indicating that the definition $(\mathcal{P} = -p)$ is more satisfying. If we interpret \mathcal{P} as a pressure, we can substitute $(p = -\mathcal{P})$ into the gradient of Eq. (62), giving the possibility of finding an Euler equation with $-\nabla \mathcal{P}$ as a force.

The forces given by the last term on the rhs of Eq. (86) are unknown. These velocity dependent forces are coupling forces involving the two velocity components. They might be associated with irreversible, unsteady flow; a sort of "friction" caused by nonsmooth flow. However, states of wavefunctions that satisfy the space time Schrödinger equation are closed, so if the flow satisfies Euler Eq. (76), the total energy is conserved. Since satisfaction of Eq. (86) does not imply the satisfaction of the Schrödinger equation, it might be applicable for mixed states, where the system of interest is part of a larger closed system.

In order to interpret the term $\nabla(\eta \nabla \cdot \mathbf{v})$, consider the stress tensor of the Navier– Stokes equation:

$$\sigma = -p\mathbf{I} + \lambda(\nabla \cdot \mathbf{u})\mathbf{I} + 2\mu(\operatorname{\mathbf{grad}}\mathbf{u})_{+}; \quad (\operatorname{\mathbf{grad}})_{ij} = \frac{\partial u_i}{\partial x_j}, \quad (\operatorname{\mathbf{grad}}\mathbf{u})_{+} = \operatorname{\mathbf{sym}}(\operatorname{\mathbf{grad}}\mathbf{u}),$$

giving $\sigma_{ij} = -p\delta_{ij} + \lambda \frac{\partial u_k}{\partial x_k}\delta_{ij} + \mu \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i}\right),$

where $\mu : \mathbb{R}^3 \to \mathbb{R}$ and $\lambda : \mathbb{R}^3 \to \mathbb{R}$ are the dynamic- and second-viscosity coefficients, respectively. Since the corresponding force is

$$\operatorname{\mathbf{div}} \sigma = -\nabla p + \nabla [\lambda \nabla \cdot \mathbf{u}] + 2(\operatorname{\mathbf{grad}} \mathbf{u}) \nabla \mu + \mu \nabla (\nabla \cdot \mathbf{u}),$$

where $((\mathbf{div} \ \sigma)_i = \partial_1 \sigma_{i1} / \partial x_1 + \partial_1 \sigma_{i2} / \partial x_2 + \partial_1 \sigma_{i3} / \partial x_3; \mathbf{r} = (x_1, x_2, x_3))$, and the implication $(\nabla \times \mathbf{u} = \mathbf{0} \Rightarrow \mu \nabla^2 \mathbf{u} = \mu \nabla (\nabla \cdot \mathbf{u}))$ is substituted, $\nabla (\eta \nabla \cdot \mathbf{v})$ can only come from the diagonal elements of the stress tensor. We now give two possible interpretations of $\nabla (\eta \nabla \cdot \mathbf{v})$. One is obtained by choosing $(\mu = \mathbf{0})$, giving $(\eta = \lambda)$. The second one is to take η as the nonpressure part of the average normal stress:

$$\frac{1}{3}\mathrm{tr}\sigma = \frac{1}{3}\sigma_{ii} = -p + \lambda \frac{\partial u_k}{\partial x_k} + \frac{2}{3}\mu \left(\frac{\partial u_i}{\partial x_i}\right) = -p + \left(\lambda + \frac{2}{3}\mu\right)\nabla\cdot\mathbf{u},$$

giving $(\eta = \lambda + 2\mu \div 3)$, where $(\lambda + 2\mu)$ is the bulk viscosity. Since this assignment is an approximation, where $2(\mathbf{grad u})\nabla\mu$ is neglected, the first one seems like a better interpretation.

Since η is not a constant, the interpretation of $\eta \nabla (\nabla \cdot \mathbf{u})$ from Eq. (81) is more difficult. Because of the conditions imposed on the stress tensor—giving the forces of the Navier-Stokes equation—may not be applicable to quantum flows, it seems reasonable to interpreted the function $\eta \nabla (\nabla \cdot \mathbf{u})$ as a sort of bulk viscosity. So we do that: Both $\nabla (\eta \nabla \cdot \mathbf{u})$ and $\eta \nabla (\nabla \cdot \mathbf{u})$ are interpreted as corresponding variable of bulk viscosity.

8.3. Interpretations of energy and Euler equations

We have demonstrated that states of quantum mechanical satisfy energy equations and equations of motion. We also have identified corresponding variables of velocity, energy, and force. This is the easy part. The hard part is giving a complete description of states of quantum mechanics that is consistent with these equations and variables.

While the equations derived have been identified as equations of fluid dynamics with variable mass, one can imagine other possible descriptions that are consistent with these equations, fluid pulsation and particle oscillation, being examples. Also, given the independence of the two velocity vectors \mathbf{u} and \mathbf{v} , and the the couplings between them, the two velocities could be associated with different modes, or forms, of energy. For example, Salesi [57], using approximations of relativistic quantum-mechanics, assigns the kinetic energy from the Madelung velocity \mathbf{v} to the center of mass motion, and the velocity \mathbf{u} to the internal energy of the electron.

For a particle trajectory description, we take equation of motion $(78) \div \rho$, where, for one-body states, the streamlines of the fluid are the possible paths for the particle. This approach is Bohmmian mechanics with velocity **u** and body force $P\rho^{-1}$ replacing the quantum potential. The energy equation with the addition of pressure removes tunneling and gives a constraint between the variables. The addition of \mathbf{u} gives states with kinetic energies that agrees with the corresponding expectation value. It also provides the necessary additional orbital angular momentum, since, for example, velocity \mathbf{v} alone can only provide the correct z component of angular momentum of the electronic states of the hydrogen atom. However, the trajectories with velocity \mathbf{u} are unstable, since they go off into infinity with a velocity **u** direction of $-\nabla \rho$. Therefore, if the velocity directions is not changed, as considered in the next section, the electron must frequently jump from one path to another, or go back and forth, and such behavior, involving velocity discontinuity, has been predicted to exist [66]. It is worth mentioning that such jumps might be needed in standard quantum mechanics, if two systems in the same state are to be identical over time, otherwise an electron would be stuck in a single orbit. Also, for the hydrogen s states with standard quantum mechanics, in order to satisfy the vanishing angular momentum-requirement, the electron must stay in a straight radial line, going back and forth, frequently changing to the opposite direction, and perhaps, go through the nucleus.

Since the *n*s wavefunctions of hydrogen are eigenfunctions of \hat{L}^2 —the operator for the square of the magnitude of the angular momentum—and the eigenvalues are zero, they must have radially directed momentum, so that the angular momentum vector-field is the zero function. For a fluid description of the *n*s state of hydrogen, the vanishing angular momentum condition is satisfied by velocity **u** of Eq. (23). Since the flows are steady, the flows must have local variable mass, since a steady radial flow requires infinite mass if mass is conserved. Over all space, the flows conserve mass, because the sources cancel the sinks. Anyone who works with quantum mechanics has to swallow "quantum pills," tunneling and the uncertainty principle being two examples. As quantum pills go, the local variable mass pill is, perhaps, not too large too swallow. (Fluid flows with velocity **u** is explored elsewhere, including the investigation of the hydrogen atom 1s and 2s states [1,54].)

While local variable mass and a jumping electron are possibilities, these are

30

nonclassical elements, and we wish investigate if these elements can be eliminated. The next section investigates a way to modify the direction of quantum flows, and two of the applications in doing so is for mass conservation for fluid flow and stable orbitals for trajectories.

9. One-Body Cross Flows

In this section, for one-body states, we investigate ways to change velocity directions of steady, quantum flows, without changing the kinetic or total energy, and conditions are given for these flows to conserve mass and also to be incompressible along streamlines. A special type of flow, called cross flow, is explored, where a velocity field μ is introduced that replaces \mathbf{u} , and μ has the same speed as \mathbf{u} . In order to get an idea what we have in mind, consider the familiar orbital plot of the $3d_z$ electronic state of the hydrogen atom, containing one level surface of the probability density. Imagine the torus filled with electron fluid undergoing steady flow. From symmetry, the mass density and speed are constant, mass is conserved, and the flow is incompressible, and, therefore, also solenoidal. Such incompressible flows have density stratification, where the mass density is only constant of each streamline. Since velocity \mathbf{v} is irrotational, if μ is solenoidal, the two velocities can be considered components of a velocity vector that can be expressed by a Helmolholtz decomposition ($\boldsymbol{\omega} = \nabla S \div m + \nabla \times \mathbf{K}$), such that ($\boldsymbol{\mu} = \nabla \times \mathbf{K}$).

In Sec. 10, equations of motion are obtained for many-body states. Under certain conditions, these equations are reduced to 1-body in Sec. 11, leading to orbital equations. However, a difficulty arises in the integrations. This problem removed in Sec. 11.5 by a change in velocity direction to give incompressible flow.

Diverting velocity direction away from $-\nabla\rho$ might provide a way to describe electron spin, given that electron spin could be from electron motion. The following velocity field formula was obtained for spin one-half particles from approximations of the Pauli equations by Holland [51], in spherical coordinates:

$$\mathbf{x}_{\pm} \times \hat{\mathbf{z}} = \frac{\nabla \rho}{m\rho} \times \mathbf{s}_{\mp} = \mathbf{u} \sin \theta \,\hat{\phi}, \qquad \mathbf{s}_{\mp} = \mp \frac{\hbar}{2} \hat{\mathbf{z}}. \tag{89}$$

This formula was applied to states of the hydrogen atom by Colijn and Vrscay [52, 53]. For fluid flow, the formula provides incompressible flow with mass conservation and density stratification for the electronic states of the hydrogen atom. Because of the sin θ factor, for hydrogen states, formula (89) accounts for all the kinetic energy for either a fluid particles or an orbiting electron when they are on the x-y plane containing the nucleus. In the assignment of spin angular momentum to translation, a difficulty arises in how to describe both spin and orbital momentum with translation in such a way that the two forms of momentum can be distinguished, and such that the correct total kinetic energy is obtained. For example, if all the kinetic energy is used to describe the y and

z components of orbital angular momentum for the hydrogen atom, since, as mentioned above, this portion of the angular momentum cannot be described by velocity \mathbf{v} alone.

To get some idea about the special conditions that cross flows might satisfy, consider Euler Eq. (76) for a steady, mass conserving-flow of classical mechanics, where u^2 is replaced by μ^2 , $\mathbf{v} \cdot \nabla \rho_m$ vanish, and, using the definition $(\mathbf{u}(\rho, \nabla \rho) = -\zeta \nabla \ln \rho)$, the mass flux-term $\nabla \cdot (\rho_m \mathbf{u})\mathbf{u}$ is considered a functional of the density. Also, the velocity vector is not necessarily irrotational, giving

$$\rho_m \mathcal{D}\boldsymbol{\mu} + \nabla P + [\nabla \cdot (\rho_m \mathbf{u})\mathbf{u}](\rho, \nabla \rho) + \rho \nabla U = \mathbf{0},$$

where $\mathcal{D}\boldsymbol{\mu}$ is a material derivative. Furthermore, the only requirement we place on $\boldsymbol{\mu}$ is $(\mu^2 = u^2)$. Using the standard approach to integrate this equation of motion [62], the dot product of the unit vector in the direction of $\boldsymbol{\mu}$ is taken. In order to obtain the resulting energy Eq. (45), where \mathbf{u} is replaced by $\boldsymbol{\mu}$, the density functional $\nabla \cdot (\rho_m \mathbf{u})\mathbf{u}$ must vanish. Hence, the direction of $\boldsymbol{\mu}$ must satisfy $(\boldsymbol{\mu} \cdot \mathbf{u} = \mathbf{0})$, giving constant density flow: $(\boldsymbol{\mu} \cdot \nabla \rho_m = \mathbf{0})$. With this condition and mass conservation, we have $(\mathbf{0} = \nabla \cdot (\rho_m \boldsymbol{\mu}) = \rho_m \nabla \cdot \boldsymbol{\mu})$, giving $(\nabla \cdot \boldsymbol{\mu} = \mathbf{0})$. Hence, the flow is incompressible on each streamline. With u^2 replaced by $\boldsymbol{\mu}^2$, and $(u^2 = \boldsymbol{\mu}^2)$, energy Eq. (45) is the Bernoulli equation, for the general case of density stratification, and the special case, where the energy is uniform for the entire flow field. This analysis suggests that the correct quantum flow, at least in this special case, is incompressible, density stratified, mass conserving flow. Next cross flow is defined. Under special conditions these flows are mass conserving and solenoidal.

Definition 8 (Cross Flow for one-body states) Let (S, ρ) determine the quantum flow $\mathcal{F}(\mathbf{u}, \mathbf{v})$ and the 1-body wavefunction Ψ . Let $\boldsymbol{\mu}$ be a vector field that satisfies $(\boldsymbol{\mu} \cdot \mathbf{u} = \mathbf{0})$ and $|\boldsymbol{\mu}| = |\mathbf{u}|$. Let $(\boldsymbol{\omega} \doteq \boldsymbol{\mu} + \mathbf{v})$. The flow $\mathcal{F}_{\times}(\boldsymbol{\mu}, \mathbf{v})$ is called a cross flow of \mathcal{F} , where both flows have the same quantum parameters P and ρ_m . If $\boldsymbol{\mu}$ is solenoidal: $(\nabla \cdot \boldsymbol{\mu} = \mathbf{0})$, \mathcal{F}_{\times} is said to be cross velocity solenoidal. If \mathcal{F} is smooth $(\nabla \rho \cdot \nabla S = \mathbf{0})$, \mathcal{F}_{\times} is also said to be smooth. The cross flow Euler equation for steady, smooth flow is Eq. (76) without the mass flux term $(\nabla \cdot (\rho_m \mathbf{u})\mathbf{u}, but$ they can also have additional nowork forces. Since energy is invariant to velocity direction and the presence of nowork forces, a cross flow $\mathcal{F}_{\times}(\boldsymbol{\mu}, \mathbf{v})$ that satisfies Euler Eq. (76), also satisfy energy Eq. (45), with the same energy uniform E as its corresponding flow $\mathcal{F}(\boldsymbol{\mu}, \mathbf{v})$.

Given an ensemble of a quantum state with a distribution of initial configurations, distributed according to the Born rule, the Born rule is preserved over time if the velocity vector is \mathbf{v} : The distribution of members of the ensemble satisfy the Born rule at all times [67]. For particle trajectories of one-body states, a cross flow velocity vector $\boldsymbol{\omega}$ also satisfies this condition, because the direction of the velocity vector component $\boldsymbol{\mu}$ is always on a constant level surface of the probability density ρ , since $\boldsymbol{\mu} \cdot \nabla \rho$ vanish.

For cross flows with vortexes, or orbiting electrons, there is an additional force. For real valued wavefunctions, in order to achieve the necessary radius of curvature, such a force must act perpendicular to the velocity vector $\boldsymbol{\mu}$. The force also must do no work, or, at least, be defined by a free energy-potential. Otherwise, such a force would

contribute to the energy. With the addition of a nowork force, the Euler equations above are only satisfied along the velocity directions, but they still integrate to the same energy equation. For the hydrogen s states, in atomic units, normal to the velocity direction, the force \hat{F} must satisfy

$$-\nabla U \cdot \hat{\mathbf{r}} - \rho^{-1} \nabla P \cdot \hat{\mathbf{r}} + \hat{F} \cdot \hat{\mathbf{r}} = -\frac{\mu^2}{r^2}; \qquad \mu^2 = u^2, \quad -\nabla U = -\frac{1}{r^2} \hat{\mathbf{r}},$$

where $-\mu^2/r^2$ is the centrifugal acceleration. For hydrogen 1s fluid flow, the speed has a constant value of 1 a.u, the same as in the Bohr model, and the pressure satisfies $(-\nabla P \cdot \hat{\mathbf{r}}/\rho = r^{-2} + 2r^{-1} - 2)$ [1]. An orbiting electron-treatment of the 1s electron, where the force \hat{F} is left out, gives a modified Bohr model, where the radius of the sole stable orbit is $3/2 a_0$, compared to 1 a_0 for the Bohr atom. For excited states, we expect that this model will give more then one orbit. With the force \hat{F} included, in order to have a stable orbit at the Bohr radius, the force \hat{F} must be $-\hat{\mathbf{r}}$ a.u. For fluid flow, the force must be such that the Euler equations are satisfied perpendicular to the streamlines over the entire flow field.

The next lemma proves that if the velocity \mathbf{u} of a flow $\mathcal{F}(\rho_m, P, \mathbf{u}, \mathbf{v})$ is replaced by velocity $\boldsymbol{\mu}$, such that the direction of $\boldsymbol{\mu}$ is $\mathbf{u} \times \mathbf{v}$, then $\boldsymbol{\mu}$ is solenoidal. The theorem is applicable to any two irrotational vector fields.

Lemma 9 (The divergence of the cross product of irrotational fields) Let $(\mu \doteq \mathbf{x} \times \mathbf{u})$, where \mathbf{x} and \mathbf{u} are irrotational vector fields. The vector field μ is solenoidal.

Proof Let the scalar field θ and θ_{\times} be such that $(\mathbf{u} = \nabla \theta)$ and $(\mathbf{x} = \nabla \theta_{\times})$, respectively. Using the identity

$$\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B}),$$

and the vanishing of the curl of the gradient, we have,

$$abla \cdot oldsymbol{\mu} =
abla \cdot (\mathbf{x} imes \mathbf{u}) =
abla \cdot (
abla heta_{ imes} imes
abla heta) = \mathbf{0}.$$

The following theorem demonstrates that smooth cross flows have constant density: $(\boldsymbol{\omega} \cdot \nabla \rho = \mathbf{0})$; $(\boldsymbol{\omega} \doteq \boldsymbol{\mu} + \mathbf{v})$. Also, $(\nabla \cdot \boldsymbol{\mu} = \mathbf{0})$ implies mass conservation, and the previous lemma demonstrates one way to define the vector field $\boldsymbol{\mu}$ so that it is solenoidal. Also, conditions for incompressible flow are also given. Since it is easy to overlook, it is worth mentioning that steady, constant density-flow does not imply mass conservation.

Theorem 8 Let \mathcal{F}_{\times} be a cross flow of \mathcal{F} .

1) If \mathcal{F} is smooth, each streamline of \mathcal{F}_{\times} has a constant density.

2) The following statements are equivalent: \mathcal{F}_{\times} is cross velocity solenoidal, $\rho \mu$ is solenoidal, \mathcal{F}_{\times} is mass conserving.

If \mathcal{F}_{\times} is cross velocity solenoidal, then vector field $\boldsymbol{\omega}$ can be represented by the Helmholtz decomposition:

$$m\boldsymbol{\omega} = \nabla S + \nabla \times \mathbf{K}, \qquad \boldsymbol{\mu} = \nabla \times \mathbf{K}.$$
 (90)

3) If \mathcal{F}_{\times} is cross velocity solenoidal, smooth and steady, $\boldsymbol{\omega}$ is incompressible. Here is a summary:

1) \mathcal{F}_{\times} is smooth $\Rightarrow (\nabla \rho_m \cdot \boldsymbol{\omega} = \mathbf{0}).$ 2) $(\nabla \cdot \boldsymbol{\mu} = 0) \Leftrightarrow (\nabla \cdot (\rho_m \boldsymbol{\mu}) = \mathbf{0}) \Leftrightarrow (\partial \rho + \nabla \cdot (\rho_m \boldsymbol{\omega}) = \mathbf{0}) \Rightarrow Eq. (90).$ 3) $(\nabla \cdot \boldsymbol{\mu} = 0)$ and smooth and steady $\mathcal{F}_{\times} \Rightarrow (\nabla \cdot \boldsymbol{\omega} = \mathbf{0}).$

Proof

1) For a cross flow \mathcal{F}_{\times} from a corresponding smooth flow \mathcal{F} , we have $(\mathbf{u} \cdot \boldsymbol{\mu} = \mathbf{0} \Rightarrow \nabla \rho_m \cdot \boldsymbol{\mu} = \mathbf{0})$ and $(\nabla \rho \cdot \nabla S = \mathbf{0} \Rightarrow \nabla \rho \cdot \mathbf{v} = \mathbf{0})$, respectively, giving a constant density cross flow: $(\nabla \rho \cdot \boldsymbol{\omega} = \mathbf{0})$.

2) Combining the solenoidal requirement $(\nabla \cdot \boldsymbol{\mu} = \mathbf{0})$ with the cross flow requirement $(\nabla \rho_m \cdot \boldsymbol{\mu} = \mathbf{0})$, we have $(\nabla \cdot (\rho_m \boldsymbol{\mu}) = \mathbf{0})$. Hence, $\rho \boldsymbol{\mu}$ is solenoidal. The logic:

$$\nabla \cdot (\rho_m \boldsymbol{\mu}) = \boldsymbol{0} + \text{Continuity Eq. (47.2)} \Rightarrow \partial \rho + \nabla \cdot (\rho_m \boldsymbol{\omega}) = \boldsymbol{0}, \quad (91)$$

then indicates that mass is conserved. Equation (90) holds because a solenoidal vector field $\boldsymbol{\mu}$ can be written as the curl of vector potential **K**. By reversing the above two implications, \mathcal{F}_{\times} being mass conserving implies it is cross velocity solenoidal. Equation (90) holds by the Helmholtz decomposition theorem.

3) For smooth, steady flow the continuity Eq. (47.2) reduces to $(\nabla \cdot \mathbf{v} = \mathbf{0})$. Combining this result with the cross velocity solenoidal property $(\nabla \cdot \boldsymbol{\mu} = \mathbf{0})$, we have $(\nabla \cdot \boldsymbol{\omega} = \mathbf{0})$.

The following corollary uses information from lemma 9: $(\nabla \cdot \boldsymbol{\mu} = \mathbf{0})$, Eq. (51) for steady smooth flow: $(\nabla \cdot \mathbf{v} = \mathbf{0})$, and Theorem 8.

Corollary 1 Steady ($\partial \rho = \mathbf{0}$) smooth ($\nabla \rho \cdot \nabla S = 0$) cross flows $\mathcal{F}_{\times}(\boldsymbol{\mu}, \mathbf{v})$ with $(\boldsymbol{\mu} = \nabla S/|\nabla S| \times \mathbf{u})$ are solenoidal ($\nabla \cdot (\boldsymbol{\mu} + \mathbf{v}) = \mathbf{0}$), conserves mass ($\nabla \cdot \rho_m(\boldsymbol{\mu} + \mathbf{v}) = \mathbf{0}$), incompressible ($\nabla \cdot (\boldsymbol{\mu} + \mathbf{v}) = \mathbf{0}$), and have a constant density ($\rho|_s = \text{const.}$) on each streamline s, if ∇S does not vanish on positive measure.

The next theorem provides conditions for smooth, cross flows to satisfy that imply that they satisfy energy equation (45.1). From Theorem 8, these flows have a constant density on each streamline, but the density depends on the streamline. Let $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$ be a smooth flow, whose wavefunction satisfies the spatial Schrödinger equation. For the next theorem, it is demonstrated that if there exists a cross flow $\mathcal{F}_{\times}(P, \rho_m, \boldsymbol{\mu}, \mathbf{v})$, corresponding to smooth flow $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$, that satisfies the equation of motion

$$\rho_m \mathcal{D}\boldsymbol{\omega} + \nabla P + \rho \nabla U = \rho \vec{F}; \qquad \boldsymbol{\omega} = \boldsymbol{\mu} + \mathbf{v}, \tag{92}$$

with any nowork force \vec{F} , the cross flow \mathcal{F}_{\times} satisfies the same energy Eq. (45.1) as the smooth flow \mathcal{F} . However, the theorem given is slightly more general, where the wavefunction Ψ of the smooth flow \mathcal{F} is not necessarily an eigenfunction of the Schrödinger equation. Section 11.5 demonstrates that such cross flows for many body states can yield reduced 1-body Euler equations.

Theorem 9 (Smooth Cross Flow Implications) Let S and ρ be time-independent 1-body fields, giving the wavefunction Ψ and smooth, steady flow $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$. However, Ψ is not necessarily an eigenvector of Hamiltonian H with external potential U. Let \mathcal{C} be the set of all smooth, cross flows $\mathcal{F}_{\times}(P, \rho_m, \boldsymbol{\mu}, \mathbf{v})$ of \mathcal{F} that satisfy Euler Eq. (92) with a nonwork force \vec{F} , where \vec{F} can depend on the cross flow.

The following hold if C in nonempty.

1) The flow \mathcal{F} and each and every $\mathcal{F}_{\times} \in \mathcal{C}$ satisfy the energy Eq. (45.1), where the energy scalar field E may depend on the streamline.

2) Under the condition of continuity Eq. (47.2) satisfaction for flow \mathcal{F} , the following two statements are equivalent:

a) Ψ is an eigenvector of the TISE. **b**) The energy scalar field E is uniform.

3) Eq. (92) is an Euler equation for the cross flows $\mathcal{F}_{\times}(P, \rho_m, \boldsymbol{\mu}, \mathbf{v}) \in \mathcal{C}$, having the proper time derivative, if and only if mass is conserved.

Proof

1) Let \mathcal{F}_{\times} satisfy Eq. (92). Note that

$$\rho_m \mathcal{D}\boldsymbol{\omega} = \frac{1}{2} \rho_m \nabla \omega^2 - \boldsymbol{\omega} \times (\nabla \times \boldsymbol{\omega}), \tag{93}$$

where, by definition of the cross product \times , the second term on the rhs, a vector field, is perpendicular to the streamline—the direction of $\boldsymbol{\omega}$. Let $\hat{\mathbf{e}}$ be a unit vector tangent to the streamline, that is, in the direction of $\pm \boldsymbol{\omega}$. Hence, $(\boldsymbol{\omega} \times (\nabla \times \boldsymbol{\omega}) \cdot \hat{\mathbf{e}} = \mathbf{0})$. Next we perform the following usual procedure for the fields restricted to a streamline:

- 1) Substituting Eq. (93) into Eq. (92).
- 2) Take the dot product of the resulting equation and $\hat{\mathbf{e}}$.
- 3) Use $(\vec{F} \cdot \hat{\mathbf{e}} = \mathbf{0})$ —that holds because \vec{F} does no work.

4) Use $(\boldsymbol{\omega} \times (\nabla \times \boldsymbol{\omega}) \cdot \hat{\mathbf{e}} = \mathbf{0})$ —that holds by definition of $\hat{\mathbf{e}}$.

The procedure gives

$$\frac{1}{2}\rho_m \nabla \mu^2 \cdot \hat{\mathbf{e}} + \frac{1}{2}\rho_m \nabla v^2 \cdot \hat{\mathbf{e}} + \nabla P \cdot \hat{\mathbf{e}} + \rho_m \nabla U \cdot \hat{\mathbf{e}} = \mathbf{0}.$$
(94)

From Theorem 8, for smooth, cross flows, the density is constant on each streamline, giving $(d(\rho_m \omega)/dt = \rho_m d\omega/dt)$. Using this result, the differential form of the above equation is

$$\frac{1}{2}d(\rho_m\mu^2) + \frac{1}{2}d(\rho_mv^2) + dP + \rho \, dU = \mathbf{0}.$$
(95)

Since the density ρ_m is constant on each streamline, the integration of Eq. (95) gives energy Eq. (45.1) with velocity vector field $\boldsymbol{\mu}$ replacing \mathbf{u} . Hence, \mathcal{F}_{\times} satisfies Eq. (45.1). Since the path integration of the above equation is restricted to a streamline, the energy is only necessarily constant on each streamline. Since $(u^2 = \mu^2)$, and both flows share the same density ρ and pressure P, \mathcal{F} also satisfies energy Eq. (45.1). **2 part 1)** If *E* is a constant, then \mathcal{F} satisfies energy Eq. (45.1), as is, with a constant energy *E*. According to Theorem (3), the wavefunction ($\Psi = \Psi(\rho, S)$) satisfies the real part of the TISE. Since, by hypothesis, \mathcal{F} also satisfies continuity Eq. (47.2), and from Theorem 1, continuity Eq. (47.2) is equivalent to the imaginary part of the TISE (11), the wavefunction Ψ also satisfies the imaginary part of the TISE (11). Hence, a constant energy *E* implies that Ψ is an eigenvector of the TISE.

2 part 2) If the wavefunction ($\Psi = \Psi(\rho, S)$) is an eigenvector of Hamiltonian H with external potential U, then, according the Theorem 3, \mathcal{F} satisfies the energy Eq. (45.1), as is, with a constant energy E. (Since, both \mathcal{F} and \mathcal{F}_{\times} satisfy the energy Eq. (45.1) with the same energy E, the flow \mathcal{F}_{\times} also satisfies the Euler Eq. (76), as is, with a constant energy E.)

3) If mass is not conserved, Eq. (92) is not an Euler equation, only because it is missing the mass flux term $(\nabla \cdot (\rho_m \boldsymbol{\omega}) \boldsymbol{\omega})$. If mass is conserved, this term vanish.

Next we make an argument that, because the direction of \mathbf{u} is $-\nabla \rho$, and \mathbf{u} is irrotational, there is always a combination of a nowork force and prescribed cross flow velocity direction that gives incompressible flow with mass conservation. Given that the nowork force is at our disposal, we can "steer" a fluid particle with speed $|\mathbf{u}|$ in any continuous path, including with a streamline with tangent vector $\nabla S/|\nabla S| \times \mathbf{u}/|\mathbf{u}|$. According to Corollary (1), such a flow is incompressible and mass conserving. In cases where ∇S is the zero vector, we have more options for the velocity direction, so there could be more possibilities. We simply need a potential ϕ , such that $\nabla \phi$ is not in the same direction as $\nabla \rho$, and then take velocity $\boldsymbol{\mu}$ to be $\nabla \phi/|\nabla \phi| \times \mathbf{u}$. For time dependence, if the flow is laminar and changing in a continuous manner, it seems like a reasonable assumption that you can find such a flow with a time dependent nowork force.

Next we consider if a variable mass flow $\mathcal{F}_{vm}(\rho_m, P, \mathbf{u})$ can have meaning in a flow of classical mechanics, where \mathbf{u} is not necessarily given by Def. (23.2). Let the flow $\mathcal{F}(\rho_m, P, \boldsymbol{\mu})$ be an actual flow of a classical fluid. Let the variable mass-flow $\mathcal{F}_{vm}(\rho_m, P, \mathbf{u})$ have a velocity direction perpendicular the $\boldsymbol{\mu}$ and satisfy $(u^2 = \mu^2)$. Consider \mathcal{F}_{vm} as a fictitious, or imaginary flow, making \mathcal{F}_{vm} a sort of cross flow of $\mathcal{F}(\rho_m, P, \boldsymbol{\mu})$. The imaginary flow \mathcal{F}_{vm} can be used to obtain information about flow \mathcal{F} along lines with directions that are normal to the streamlines of \mathcal{F} . A fluid particle in the imaginary flow is like a probe, giving information about the mass density, speed, pressure, and kinetic and potential energies of flow \mathcal{F} along its streamlines. The imaginary flow carries all the information about the actual flow, except velocity direction.

Next we try to give meaning to formula (32.1) for pressure P. Consider again the imaginary $\mathcal{F}(\rho_m, P, \mathbf{u})$ flow of flow $\mathcal{F}_{\times}(\rho_m, P, \boldsymbol{\mu})$, where both flows satisfy Bernoulli Eq. (45), and \mathcal{F} satisfies the Euler equation:

$$\frac{1}{2}\nabla u^2 + \rho^{-1} \left(\nabla \cdot (\rho_m \mathbf{u})\mathbf{u} + \nabla P\right) + \nabla U = \vec{G},\tag{96}$$

with nowork force \vec{G} . The integration of this equation must give the Bernoulli Eq. (45). Hence,

$$\nabla \cdot (\rho_m \mathbf{u}) \mathbf{u} + \nabla P = \rho_m \nabla \left(P \rho_m^{-1} \right) = \nabla P - \rho_m^{-1} P \nabla \rho_m, \quad \text{giving}$$
$$P \nabla \rho_m = -\nabla \cdot (\rho_m \mathbf{u}) \rho_m \mathbf{u}. \tag{97}$$

Taking the dot product with \mathbf{u} , we have

$$-P = \frac{\nabla \cdot (\rho_m \mathbf{u})}{\nabla \rho_m \cdot \mathbf{u}} \rho_m u^2 = \left(1 + \frac{\nabla \cdot \mathbf{u}}{\nabla \rho_m \cdot \mathbf{u}}\right) \rho_m u^2; \qquad \nabla \rho \neq 0,$$

and these equations are satisfied in the special case with Defs. (32.1) and (23) for Pand \mathbf{u} , respectively. (We also have $(\nabla \rho_m \cdot \mathbf{u} = \pm |\nabla \rho_m| |\mathbf{u}|$.) Hence, formula (32.1) for the pressure P is a special case of the more general relation (97). Note that if $\nabla \cdot \mathbf{u}$ vanish but $\nabla \rho_m \cdot \mathbf{u}$ does not vanish, then the pressure is -2 times the fluid kinetic energy. For relation (97) to hold for a compressible, inviscid fluid of classical mechanics, there must be a nowork force \vec{G} , such that Eq. (96) is satisfied, with the restrictions mentioned above, that is, $(\nabla \rho_m \cdot \boldsymbol{\mu} = \mathbf{0})$ and $(u^2 = \mu^2)$. Also, the pressure P must be nonnegative, and the rhs of Eq. (97) must vanish if $\nabla \rho_m$ vanish. However, the pressure P can be replaced by pressure (P + const.), since Eq. (96) is invariant to P with an additive constant. Note that P is invariant to a change in sign of \mathbf{u} .

10. Lagrange's equations of motion for Many Bodies

Consider the classical Hamiltonian for n identical point masses of mass m with external V and interacting W potentials:

$$\bar{W} \doteq \sum_{i=1}^{n} W_{i}, \qquad W_{i}(\mathbf{r}_{1}, \cdots \mathbf{r}_{n}) \doteq \sum_{j \neq i}^{n} W_{12}(\mathbf{r}_{j}, \mathbf{r}_{i})$$
$$E = \sum_{i=1}^{n} \frac{1}{2} m \mathbf{s}_{i} \cdot \mathbf{s}_{i} + V(\mathbf{r}_{i}) + \frac{1}{2} \sum_{i=1}^{n} \sum_{j \neq i} W(\mathbf{r}_{i}, \mathbf{r}_{j}), \quad E = \text{const},$$
(98)

where velocities \mathbf{s}_i depends on time. As is well known, it follows from Lagrangian mechanics, that the *n* equations of motion are

$$m\frac{d\mathbf{s}_i}{dt} = -\nabla_i V + \sum_{j \neq i}^n \nabla_i W(\mathbf{r}_i, \mathbf{r}_j), \qquad i = 1, \cdots n.$$
(99)

Let $[(\mathbf{r}_1, \dots, \mathbf{r}_n)](t)$ be the particle configuration at time t. If this particle configurationfunction is invertible, giving $(t = t(\mathbf{r}_1, \dots, \mathbf{r}_n))$, then we can define a generalized velocity fields: $(\mathbf{v}_i(\mathbf{r}_1, \dots, \mathbf{r}_n) = \mathbf{s}_i [t(\mathbf{r}_1, \dots, \mathbf{r}_n)])$ with domain \mathbb{R}^{3n} . The Hamiltonian above can then be written

$$E = \sum_{i=1}^{n} \frac{1}{2} m \mathbf{v}_{i}^{2} + V(\mathbf{r}_{i}) + \frac{1}{2} \sum_{j \neq i} W(\mathbf{r}_{i}, \mathbf{r}_{j}).$$
(100)

Furthermore, if there exists a time range $\tau \in (t_0, t_1)$, where $(\mathbf{r}_i(t_a) = \mathbf{r}_i(t_b)) \Rightarrow (t_a = t_b)$, for all $i \in \{1, \dots, n\}$, then, during the time interval we have the maps $\mathbf{r}_i \mapsto t \in \tau$, and we can define the velocity fields $(\mathbf{y}_i(\mathbf{r}_i) = \mathbf{s}_i[t(\mathbf{r}_i)])$ with domain \mathbb{R}^3 . Using the chain rule for the acceleration, the equation of motion for velocity field $\mathbf{y}_{\alpha} \in \{\mathbf{y}_1, \dots, \mathbf{y}_n\}$ can be written

$$m\nabla \mathbf{y}_{\alpha} \cdot \mathbf{y}_{\alpha} = -\nabla_i V(\mathbf{r}_{\alpha}) - \sum_{i \neq \alpha}^n \nabla_{\alpha} W(\mathbf{r}_{\alpha}, \mathbf{r}_i); \qquad (\nabla \mathbf{y}_{\alpha} \cdot \mathbf{y}_{\alpha})_i = (\nabla y_{\alpha i}) \cdot \mathbf{y}_{\alpha}.$$

Given the similarities of Hamiltonians $(45) \times \rho^{-1}$ and (100), it should be possible to obtain *n* equations of motion, for *n* interacting point masses involving Hamiltonian (45.1). This section does that using Lagrangian dynamics for stationary states under the orthogonal condition. The Lagrangian equations obtained are in particle, or per mass units, where the pressure *P* appears as a factor in the product $P\rho^{-1}$. For trajectories of point masses, $P\rho^{-1}$ is a body force. Equation of motion forms applicable to fluid flow are also given. An equation of motion form involving velocity fields \mathbf{u}_1 and \mathbf{v}_1 , parameterized by the other $\mathbf{r}_2, \cdots \mathbf{r}_m$ position variables, is also obtained. This equation is used in the section that follows to obtain reduced 1-body equations. Except for difficulties that must be overcome, the reduced equation, for a particle with position variable \mathbf{r}_1 , is obtained by integrating an equation of motion over the variables $\mathbf{r}_2, \cdots, \mathbf{r}_n$.

For later use we note that, since the forces are conservative in the equations of motion sequence (99), via integration, there exist constants $\epsilon_1, \dots \epsilon_n$, such that

$$\epsilon_i = \frac{1}{2}m\mathbf{s}_i \cdot \mathbf{s}_i - V(\mathbf{r}_i) - \sum_{j \neq i}^n W(\mathbf{r}_i, \mathbf{r}_j).$$

Summing over the particle energies we see that the potential energy of 2-body interactions is double counted when compared to energy Eq. (98):

$$\sum_{i=1}^{n} \epsilon_i = \sum_{i=1}^{n} \frac{1}{2} m \mathbf{s}_i \cdot \mathbf{s}_i + V(\mathbf{r}_i) + \sum_{i=1}^{n} \sum_{j \neq i} W(\mathbf{r}_i, \mathbf{r}_j).$$
(101)

10.1. Many Body Trajectories and Stream Neighborhoods

In this subsection, using the generalized velocity fields $\mathbf{u}_i, \mathbf{v}_i \colon \mathbb{R}^{3n} \to \mathbb{R}^3$, time and field dependent velocities are defined for stationary quantum states under the orthogonal condition. For trajectories of particles, the velocities functions \mathbf{u}_i and \mathbf{v}_i are replaced by time dependent ones, denoted below by $\dot{\mathbf{r}}_i \colon \Omega \to \mathbb{R}^3$ and $\dot{\mathbf{s}}_i \colon \Omega \to \mathbb{R}^3$, where $\Omega \subset \hat{\mathbb{R}}$, and $\hat{\mathbb{R}}$ is the real number line with time as the variable. In classical mechanics, if the displacement as a function of time, for a given time range, is invertible, velocities as a function of position can be defined. These velocity fields, denoted by $\mathbf{x}_i, \mathbf{y}_i \colon \mathbb{R}^3 \to \mathbb{R}^3$, are also defined.

Let $\mathbf{u}_i : \mathbb{R}^{3n} \to \mathbb{R}$ and $\mathbf{v}_i : \mathbb{R}^{3n} \to \mathbb{R}$. Let

$$\vec{\mathbf{q}} \doteq (\mathbf{q}_1, \cdots, \mathbf{q}_n) \in \mathbb{R}^{3n}$$
 and $\vec{\mathbf{q}}_0 \doteq (\mathbf{q}_{0,1}, \cdots, \mathbf{q}_{0,n}) \in \mathbb{R}^{3n}$

where $\vec{\mathbf{q}}_0$ and $(\Omega \doteq (t_0, t) \in \hat{\mathbb{R}})$ are, respectively, the initial configuration and time range, of an *n*-particle trajectory. Also, $\vec{\mathbf{q}} : \Omega \to \mathbb{R}^{3n}$, meaning that $\vec{\mathbf{q}}(\tau)$ is the configuration at time $\tau \in \Omega$. The unique path *P* of motion, denoted $(P[\vec{\mathbf{q}}_0] : \Omega \to \mathbb{R}^{3n})$, for initial configuration $\vec{\mathbf{q}}_0$, is the solutions of the first order differential equations:

$$\dot{\mathbf{r}}_i = \mathbf{u}_i, \quad \dot{\mathbf{s}}_i \doteq \mathbf{v}_i; \quad \dot{\mathbf{q}}_i \doteq \dot{\mathbf{r}}_i + \dot{\mathbf{s}}_i, \quad i = 1, 2, \cdots n,$$
(102)

where $\dot{\mathbf{z}}_i : \Omega \to \mathbb{R}^3$; $(\dot{\mathbf{z}}_i = \dot{\mathbf{r}}_i, \dot{\mathbf{s}}_i, \dot{\mathbf{q}}_i)$, are the *i*th time-dependent velocities. The path of motion $P[\vec{\mathbf{q}}_0]$ is given as an *n*-tuple of curves, the *i*th one being $(\mathbf{q}_i : \Omega \to \mathbb{R}^3)$. If $(\vec{\mathbf{q}}(\tau) = P[\vec{\mathbf{q}}_0](\tau))$, then

$$\dot{\mathbf{q}}_{i}(\tau) = [\mathbf{u}_{i} + \mathbf{v}_{i}] \Big(\vec{\mathbf{q}}(\tau) \Big) \doteq \mathbf{w}_{i} \Big(\vec{\mathbf{q}}(\tau) \Big); \quad \tau \in \Omega.$$
(103)

The path of motion $P[\vec{\mathbf{q}}_0]$ is said to be component invertible if $(\mathbf{q}_i(\tau_1) = \mathbf{q}_i(\tau_2)) \Rightarrow$ $(\tau_1 = \tau_2))$ for each $i \in \{1, \dots, n\}$. For component invertible paths, denoted $P_I[\vec{\mathbf{q}}_0]$, since each value $\mathbf{q}_i(t)$ from $(t \in \Omega)$ is unique, we have the following bijective map: $(\mathbf{q}_i \mapsto \vec{\mathbf{q}})$. For component invertible path $P_I[\vec{\mathbf{q}}_0]$, using this bijection, let the velocity fields \mathbf{x}_i : Range $(\mathbf{q}_i) \to \mathbb{R}^3$ and \mathbf{y}_i : Range $(\mathbf{q}_i) \to \mathbb{R}^3$ be

$$\mathbf{x}_{i}(\mathbf{q}_{i}) \doteq \mathbf{u}_{i}(\vec{\mathbf{q}}), \quad \mathbf{y}_{i}(\mathbf{q}_{i}) \doteq \mathbf{v}_{i}(\vec{\mathbf{q}}); \quad \vec{\mathbf{q}} = \vec{\mathbf{q}}(\mathbf{q}_{i}), \quad (\vec{\mathbf{q}}) \in \text{Range } P_{I}[\vec{\mathbf{q}}_{0}]. \tag{104}$$

Comparing Eq. (102) and (104), restricted to $P_I[\vec{\mathbf{q}}_0]$, we find the following logic

$$\left(\mathbf{q}_{1}(\tau),\cdots,\mathbf{q}_{n}(\tau)\right) = P_{I}[\vec{\mathbf{q}}_{0}](\tau) \quad \Rightarrow \quad \dot{\mathbf{r}}_{i}(\tau) = \mathbf{x}_{i}(\mathbf{q}_{i}), \quad \dot{\mathbf{s}}_{i}(\tau) \doteq \mathbf{y}_{i}(\mathbf{q}_{i}), \tag{105}$$

and the sum of the two implications gives

$$\left(\mathbf{q}_{1}(\tau),\cdots,\mathbf{q}_{n}(\tau)\right) = P[\vec{\mathbf{q}}_{0}](\tau) \Rightarrow \dot{\mathbf{q}}_{i}(\tau) = \mathbf{x}_{i}(\mathbf{q}_{i}) + \mathbf{y}_{i}(\mathbf{q}_{i}).$$
 (106)

10.2. Lagrangian equations of motion and the Hamiltonian

In this subsection, using the relationships defined in the previous subsection, a Hamiltonian, Lagrangian and equations of motion are obtained for n-body stationary quantum states that satisfy the orthogonal condition and a local mass-conservation law. We begin with the Hamiltonian.

Theorem 10 (The Hamiltonian Function of Quantum Mechanics) Let $P_I[\vec{\mathbf{q}}_0]$ be a component invertible path. The n-body spatial Schrödinger equation:

$$-\frac{\hbar}{2m}\nabla^2\Psi + U\Psi = E\Psi; \quad \nabla^2\Psi \in \mathbf{\Sigma},\tag{107}$$

and the orthogonal condition: $(\nabla \rho \cdot \nabla S = \mathbf{0})$, is equivalent to the Hamiltonian

$$H(\dot{\vec{\mathbf{q}}},\vec{\mathbf{q}}) \doteq T(\dot{\vec{\mathbf{q}}}) + V(\vec{\mathbf{q}}); \quad H = E = const., \tag{108}$$

and the continuity Eq. (47.2), where

$$T(\dot{\vec{\mathbf{q}}}) \doteq \frac{1}{2}m\dot{\mathbf{q}} \cdot \dot{\mathbf{q}}, \qquad V(\vec{\mathbf{q}}) \doteq [P\rho^{-1}](\mathbf{q}) + U(\vec{\mathbf{q}}), \tag{109}$$

and the wavefunction Ψ is restricted to Range $(P_I[\vec{\mathbf{q}_0}])$. The *i*th components of

 $(\dot{\mathbf{q}}\cdot\dot{\mathbf{q}}), \ [P\rho^{-1}](\mathbf{q}), \in \mathbf{\Sigma},$

are, respectively, the values $\dot{\mathbf{q}}_i \cdot \dot{\mathbf{q}}_i$ and

$$[P\rho^{-1}]_i(\mathbf{q}_i) \doteq [-m\mathbf{x}_i \cdot \mathbf{x}_i + \zeta_0 \nabla \cdot \mathbf{x}_i](\mathbf{q}_i).$$
(110)

Proof From Theorem 3, the time dependent Schrödinger equation $\Leftrightarrow Eqs.$ (45), and from Theorem 1, the imaginary part of the TISE is equivalent to continuity Eq. (47.2). For a stationary state, E is constant; (S(t) = -Et), and the time independent Schrödinger equation (TISE) is equivalent to Eqs. (45) with constant E. The orthogonal condition $(\mathbf{u} \cdot \mathbf{v} = \mathbf{0})$ and $(\mathbf{w} = \mathbf{u} + \mathbf{v})$, give $(\mathbf{w} \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{v})$. Hence, the real part of the TISE and the orthogonal condition is equivalent the following special case of Eq. (45.1) $\div \rho$:

$$E = \frac{1}{2}m\mathbf{w} \cdot \mathbf{w} + [P\rho^{-1}](\mathbf{u}) + U(\vec{\mathbf{q}}); \quad E = \text{const},$$
(111)

where, in accordance with Eqs. (61),

$$[P\rho^{-1}](\mathbf{u}) = -m\mathbf{u} \cdot \mathbf{u} + \zeta_0 \nabla \cdot \mathbf{u}.$$

Let the functions of Eq. (111) be restricted to Range $(P_I[\vec{\mathbf{q}_0}]) \in \mathbb{R}^{3n}$. With this restriction, using (104), we make the following replacement in Eq. (111): $(\mathbf{u}) \to (\mathbf{x})$. Also, using Eq. (103), we can replace (\mathbf{w}) by $(\dot{\mathbf{q}})$. Hence, $(\mathbf{107}) \Leftrightarrow (\mathbf{108})$ with the given restriction to $P_I[\vec{\mathbf{q}_0}]$.

Using the Hamiltonian from the previous theorem, we annow define a Lagrangian that corresponds to this Hamiltonian. The theorem that follows obtains the relationship between the two functions, and the Lagrangian equations of motion for quantum states.

Definition 9 (The Lagrangian of Quantum Mechanics) The Lagrangian L of quantum mechanics for stationary states, defined on Range $(P[\vec{\mathbf{q}_0}]) \in \mathbb{R}^{3n}$, is

$$L(\vec{\mathbf{q}}, \vec{\mathbf{q}}) = T(\vec{\mathbf{q}}) + Q\mathbf{A} \cdot \dot{\mathbf{q}} - V(\vec{\mathbf{q}}), \qquad (\vec{\mathbf{q}}) \in Range(P[\vec{\mathbf{q}}_0]), \qquad (112)$$

where the kinetic T and potential V energies are defined by Eq. (109). Also, the scalar function Q and vectors $(\mathbf{A}) \in \Sigma$ are arbitrary. Furthermore, $\mathbf{A}_i \cdot \dot{\mathbf{q}}_i$ is the *i*th component of the sequence $(\mathbf{A} \cdot \dot{\mathbf{q}})$, and the *i*th component of $(P\rho^{-1})$ is given by Eq. (110).

Theorem 11 (The Lagrange's equation of motion and Hamiltonian part 1) 1) The Lagrangian equations of motion, given by the vector equation sequence

$$\frac{\partial L}{\partial \mathbf{q}} = \frac{d}{dt} \frac{\partial L}{\partial \dot{\mathbf{q}}}; \quad \partial L / \partial \mathbf{q}, \ \partial L / \partial \dot{\mathbf{q}} \in \mathbf{\Sigma}, \tag{113}$$

is equivalent to the Hamiltonian H being constant in time:

$$H \doteq \dot{\mathbf{q}} \cdot \frac{\partial L}{\partial \dot{\mathbf{q}}} - L,\tag{114}$$

for $\partial L/\partial \mathbf{q}$ restricted to Range $P_I[\vec{\mathbf{q}}_0]$, and where the Hamiltonian is given by Eq. (108).

2) The Lagrange's equations of motion for quantum states are the following form of Newton's second law:

$$\frac{d}{dt}(m\dot{\mathbf{q}}) + Q\frac{d\mathbf{A}}{dt} = \left[-\nabla(P\rho^{-1}) - \nabla U\right](\mathbf{q}) + Q\nabla\left(\mathbf{A}\cdot\dot{\mathbf{q}}\right),\tag{115}$$

a sequence of n vector equations.

Proof For part 1, using the chain rule, the time derivative of $(L = L(\mathbf{q}, \dot{\mathbf{q}}))$ is

$$\frac{dL}{dt} = \frac{\partial L}{\partial \mathbf{q}} \cdot \dot{\mathbf{q}} + \frac{\partial L}{\partial \dot{\mathbf{q}}} \cdot \ddot{\mathbf{q}}, \qquad (\partial L/\partial \mathbf{q}) \cdot \dot{\mathbf{q}}, (\partial L/\partial \dot{\mathbf{q}}) \cdot \ddot{\mathbf{q}} \in \mathbf{\Sigma}, \quad \partial L/\partial t \notin \mathbf{\Sigma}, \quad (116)$$

where, according to Eq. (112), L does not explicitly depend on time t. Suppose the Lagrangian equations of motion (113) is true. The substitution of the Lagrangian equations into the above one gives

$$\frac{dL}{dt} = \left(\frac{d}{dt}\frac{\partial L}{\partial \dot{\mathbf{q}}}\right) \cdot \dot{\mathbf{q}} + \frac{\partial L}{\partial \dot{\mathbf{q}}} \cdot \ddot{\mathbf{q}} \Rightarrow \frac{d}{dt}\left(\dot{\mathbf{q}} \cdot \frac{\partial L}{\partial \dot{\mathbf{q}}} - L\right) = \mathbf{0}.$$

Comparison with definition (114), we find that (H = const.). Suppose (H = const.). Comparing the last equality of the logic

$$H = \dot{\mathbf{q}} \cdot \frac{\partial L}{\partial \dot{\mathbf{q}}} - L = \text{const.} \Rightarrow \frac{d}{dt} \left(\dot{\mathbf{q}} \cdot \frac{\partial L}{\partial \dot{\mathbf{q}}} - L \right) = \mathbf{0} \Rightarrow \frac{dL}{dt} = \left(\frac{d}{dt} \frac{\partial L}{\partial \dot{\mathbf{q}}} \right) \cdot \dot{\mathbf{q}} + \frac{\partial L}{\partial \dot{\mathbf{q}}} \cdot \ddot{\mathbf{q}},$$

with Eq. (116), gives the Lagrangian equations of motion (113).

For part $\mathbf{2}$, equations of motion (115) follows by substitution of Eq. (112) into (113) and using Eq. (109).

The next theorem obtains explicit forms of the Hamiltonian and equations of motion for stationary quantum states that satisfy the orthogonal condition and a local mass conservation restriction. The following lemma is needed for the theorem that follows.

Lemma 10 The *n*-body time independent Schrödinger equation with orthogonal condition ($\nabla \rho \cdot \nabla S = \mathbf{0}$) implies Lagrange's equations of motion (115) with the conservative forces restricted to component invertible path $P_I[\vec{\mathbf{q}}_0]$, within definitions of **A** and *Q*.

Proof From Theorem (10), the *n*-body TISE and $(\nabla \rho \cdot \nabla S = \mathbf{0})$ imply both the kinetic T and potential V energy functions of Hamiltonian (108), restricted to $P_I[\vec{\mathbf{q}}_0]$. Via the Lagrange's Eq. (113), we have (108) \Rightarrow (115).

Theorem 12 (The Lagrange's equation of motion and Hamiltonian part 2) The *n*-body time independent Schrödinger equation imply

$$H = \frac{1}{2}m\dot{\mathbf{r}}\cdot\dot{\mathbf{r}} + \frac{1}{2}m\dot{\mathbf{s}}\cdot\dot{\mathbf{s}} + [P\rho^{-1}](\mathbf{q}) + U(\vec{\mathbf{q}}), \qquad (117)$$

and the following two sequences of n equations:

$$\frac{1}{2}\rho_m \nabla x^2 + \frac{1}{2}\rho_m \nabla y^2 + \nabla \cdot (\rho_m \mathbf{x})\mathbf{x} = -\nabla P - \rho \nabla U, \qquad (118)$$

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho_m \dot{\mathbf{r}} + \rho_m \dot{\mathbf{s}} - \int_{\mathbb{V}} \nabla \cdot (\rho_m \mathbf{x}) \mathbf{y} = \int_{\mathbb{V}} -\nabla P - \rho \nabla U; \quad \mathbb{V} \subset \mathbb{R}^3,$$
(119)

under the orthogonal condition $(\nabla \rho \cdot \nabla S = \mathbf{0})$, the condition that the potential energy functions are restricted to Range $(P_I[\vec{\mathbf{q_0}}]) \in \mathbb{R}^{3n}$, and $(\mathbf{A} \cdot \dot{\mathbf{q}})$ is not present. In

addition, for implication (118) and (119), body mass conservation: $(\nabla_i \cdot (\rho_m \mathbf{v}_i) = \mathbf{0})$, for $(i = 1, 2, \dots n)$, and this condition replaces, and implies, the weaker condition $(\sum_i \nabla_i \cdot (\rho_m \mathbf{v}_i) = \mathbf{0})$. (For one-body states with steady flow, Eq. (117) correspond to energy Eq. (45); Eq. (118) correspond to the Euler Eq. (76); Eq. (119) corresponds to Euler Eq. (81), using Eq. (51): $(\nabla \cdot \mathbf{v} = \mathbf{0})$.)

Proof For implication (117), since, from Theorem (10), the *n*-body time independent Schrödinger equation with orthogonal condition implies Eq. (108), we only need to prove that (108) \Rightarrow (117). From the orthogonal condition ($\mathbf{u} \cdot \mathbf{v} = \mathbf{0}$), we have $(\dot{\mathbf{q}} \cdot \dot{\mathbf{q}} = \dot{\mathbf{r}} \cdot \dot{\mathbf{r}} + \dot{\mathbf{s}} \cdot \dot{\mathbf{s}})$. Substituting this condition into Eq. (108) via Eq. (109), we obtain Eq. (117).

For implications (118) and (119), since, from Lemma (10), the n-body time independent Schrödinger equation and the orthogonal condition imply Eq. (115), we only need to prove the two implications below.

 $(\underline{115}) \Rightarrow (\underline{118})$. Consider a velocity function of time $\mathbf{z} : \Omega \to \mathbb{R}^3$. If the path is invertible, then, in Cartesian coordinates, we can write the velocity as a composite $(\mathbf{z}(t) = \mathbf{z}(x(t), y(t), z(t))$, as is done in the usual procedure of fluid dynamics for treating acceleration in the derivation of the Euler equation. This also gives the velocity field: $(\mathbf{z} = \mathbf{z}(x, y, z))$. Hence, the same symbol \mathbf{z} is used for two different functions, both representing the same physical quantity. If \mathbf{z} is irrotational and the flow is steady, the two functions satisfy

$$\frac{d\mathbf{z}}{dt} = \frac{1}{2}\nabla\mathbf{z}\cdot\mathbf{z},$$

and this follows by taking the material derivative and utilizing an equality for irrotational vectors. From the development in subsection 10.1, since different symbol for the two functions are used, we have

$$\frac{d}{dt}(m\dot{\mathbf{q}}) = \frac{1}{2}m\nabla[(\mathbf{x} + \mathbf{y}) \cdot (\mathbf{x} + \mathbf{y})] = \frac{1}{2}m\nabla\mathbf{x} \cdot \mathbf{x} + \frac{1}{2}m\nabla\mathbf{y} \cdot \mathbf{y},$$

where the orthogonal condition is used, and also that \mathbf{u}_i and \mathbf{v}_i being irrotational imply that \mathbf{x}_i and \mathbf{y}_i are also irrotational. It follows from logic (106) that this equation is satisfied on path of motion $P_I[\vec{\mathbf{q}}_0]$, connecting the time dependent trajectory velocities and the field velocities. Substituting into Eq. (115) and setting ((\mathbf{A}) = $\mathbf{0}$), we have

$$\frac{1}{2}m\nabla x^2 + \frac{1}{2}m\nabla y^2 = -\nabla(P\rho^{-1}) - \nabla U.$$

Multiplying by ρ , and using equality (80), generalized to a sequence, we obtain

Eq. (118):
$$\frac{1}{2}\rho_m \nabla x^2 + \nabla \cdot (\rho_m \mathbf{x})\mathbf{x} + \frac{1}{2}\rho_m \nabla y^2 = -\nabla P - \rho \nabla U.$$

 $(118) \Rightarrow (119)$. Consider Eq. (73) that holds for any irrotational velocity field. Using our explicit notation for the velocity as a function of time $(\dot{\mathbf{r}} + \dot{\mathbf{s}})$ and the velocity as a field \mathbf{w} , for steady flow, and under the orthogonal condition, this equation becomes

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho_m \dot{\mathbf{r}} + \rho_m \dot{\mathbf{s}} = \int_{\mathbb{V}} \frac{1}{2} \rho_m \nabla \mathbf{x} \cdot \mathbf{x} + \frac{1}{2} \rho_m \nabla \mathbf{y} \cdot \mathbf{y} + \nabla \cdot (\rho_m \mathbf{w}) \mathbf{w}.$$
(120)

By hypothesis, $((\nabla \cdot (\rho_m \mathbf{v})))$ vanish identically, giving $((\nabla \cdot (\rho_m \mathbf{y})) = (\mathbf{0})$, with \mathbf{y} given by Def. (104). Hence,

$$\mathcal{D}\!\!\int_{\mathbb{V}} \rho_m \dot{\mathbf{r}} + \rho_m \dot{\mathbf{s}} = \int_{\mathbb{V}} \frac{1}{2} \rho_m \nabla \mathbf{x} \cdot \mathbf{x} + \frac{1}{2} \rho_m \nabla \mathbf{y} \cdot \mathbf{y} + \nabla \cdot (\rho_m \mathbf{x}) \mathbf{x} + \nabla \cdot (\rho_m \mathbf{x}) \mathbf{y}.$$

Integrating Eq. (118) over $(V \in P_I[\vec{\mathbf{q}}_0])$, followed by substituting into Eq. (118), we obtain Eq. (119).

10.3. Parameterized one body Euler and Schrödinger equations

Next we specialize the potential U to a forms that is applicable to electronic systems.

Definition 10 (External Potential from 1- and 2-body potentials)

The potentials $\overline{V} : \mathbb{R}^{3n} \to \mathbb{R}$ and $\overline{W} : \mathbb{R}^{3n} \to \mathbb{R}$ are sums over one-body $V_1 : \mathbb{R}^3 \to \mathbb{R}$ and two-body $W_i : \mathbb{R}^3 \times \mathbb{R}^3 \to \mathbb{R}$ functions:

$$\bar{V}(\mathbf{r}_1, \mathbf{r}_2, \cdots \mathbf{r}_n) \doteq V(\mathbf{r}_1) + V(\mathbf{r}_2) + \cdots + V(\mathbf{r}_n), \quad and \tag{121}$$

$$\bar{W} \doteq \sum_{i=1}^{n} W_i, \qquad W_i(\mathbf{r}_1, \cdots \mathbf{r}_n) \doteq \sum_{j \neq i}^{n} W_{12}(\mathbf{r}_j, \mathbf{r}_i).$$
(122)

The body potential U of Eq. (45) of the TISE is the following sum of one-body and two-body potentials:

$$U \doteq \overline{V} + \frac{1}{2}\overline{W} \Rightarrow \nabla_1 U \doteq \nabla_1 V + \nabla_1 W_1; \qquad W_{12}(\mathbf{r}, \mathbf{r}') = W_{12}(\mathbf{r}', \mathbf{r}), \qquad (123)$$

with the condition that \bar{V} and \bar{W}_i vanish at infinity.

Consider the first component \mathbf{u}_1 of the *n*-tuple (\mathbf{u}). For such first component functions, the next theorem treats the (n-1) variables from $(\mathbf{q}'_1 = (\mathbf{r}_2, \dots, \mathbf{r}_n) \in \mathbb{R}^{3(n-1)})$ as parameters. A 1-body Euler equation, Newton's 2nd law, an energy equation, and a spatial Schrödingerödinger equation is obtained. The equation of motion is used in the next section to obtain 1-body reduced equations of motion and a reduced spatial Schrödinger equation, under certain conditions.

Theorem 13 (From n-body to parameterized 1-body Schrödger Equations)

Let $((\mathbf{q}_1, \mathbf{q}'_1) \in \mathbb{R} \times \mathbb{R}^{n-1} = \mathbb{R}^{3n})$. The n-body time-independent Schrödger equation

$$-\frac{\hbar}{2m}\nabla^2\Psi + \left(\bar{V} + \frac{1}{2}\bar{W}\right)\Psi = E\Psi; \quad \nabla^2\Psi \in \mathbf{\Sigma}, \quad \Psi: \mathbb{R}^{3n} \to \mathbb{C},$$
(124)

under the orthogonal condition, and the sequence of equations

$$Re\left(\nabla_{i}\cdot\left(\Psi^{*}\hat{\mathbf{P}}_{i}\Psi\right)\right) = \mathbf{0}, \quad \hat{\mathbf{P}}_{i} \doteq -i\hbar\nabla_{i}, \quad i = 1, \cdots n,$$
(125)

imply the 1-body Euler equation, Newton's 2nd law, the energy equation, and the TISE:

$$\frac{1}{2}\rho_m \nabla_1 (u_1^2 + v_1^2) + \nabla_1 \cdot (\rho_m \mathbf{u}_1) \mathbf{u}_1 = -\nabla_1 P_1 - \rho \nabla_1 V - \rho \nabla_1 W_1, \qquad (126)$$

$$\mathcal{D}(\rho_m \mathbf{u}_1 + \rho_m \mathbf{v}_1) + \rho_m \mathbf{u}_1 \nabla_1 \cdot \mathbf{u}_1 = -\nabla_1 P_1 - \rho \nabla_1 V - \rho \nabla_1 W_1, \qquad (127)$$

A Formulation of Quantum Fluid Mechanics and Trajectories

$$E_1 \rho = \frac{1}{2} \rho_m u_1^2 + \frac{1}{2} \rho_m v_1^2 + P_1 + (V + W_1)\rho, \qquad (128)$$

$$-\frac{\hbar}{2m}\nabla_1^2\Psi + (V+W_1)\Psi = E_1\Psi; \qquad \Psi: \mathbb{R}^3 \to \mathbb{C},$$
(129)

where the points $(\mathbf{q}'_1) \in \mathbb{R}^{n-1}$ are taken as parameters for Ψ in Eq. (129), and, therefore, the symbol Ψ represents different functions in Eqs. (124) and (129).

Proof Let the subspace $\mathbb{R}_1^{3n}[\mathbf{q}_1'] \subset \mathbb{R}^{3n}$ parameterized by $((\mathbf{q}_1') \doteq (\mathbf{q}_2, \cdots, \mathbf{q}_n) \in \mathbb{R}^{3(n-1)})$ and isomorphic with \mathbb{R}^3 , be

$$\mathbb{R}_1^{3n}[\mathbf{q}_1'] \doteq \{ (\mathbf{q}_1, \mathbf{q}_1') \in \mathbb{R}^{3n} | (\mathbf{q}_1') \in \mathbb{R}_1^{3(n-1)} \text{ is fixed.} \}.$$

Consider Theorem 1 part I result $4 \Leftrightarrow 6$. For a 1-body stationary state, this logic reduces to the special case

$$\operatorname{Re}\left(\nabla_{1}\cdot\left(\Psi^{*}\hat{\mathbf{P}}_{1}\Psi\right)\right)=\mathbf{0}\Leftrightarrow\operatorname{Im}\left(\Psi^{*}\hat{H}\Psi\right)=\mathbf{0}.$$
(130)

Since $\operatorname{Im}(\Psi^*E_1\Psi)$ vanish identically, the second equation of this logic is the imaginary part of the 1-body TISE (11). We identify the first equation of *n*-body sequence (125) with domain $\mathbb{R}_1^{3n}[\mathbf{q}_1']$ as the lhs equation of logic (130), parameterized with (\mathbf{q}_1') . Hence, we have the following: The *n*-body equation sequence (125), restricted to subspace $\mathbb{R}_1^{3n}[\mathbf{q}_1'] \subset \mathbb{R}^{3n}$, implies the satisfaction of the imaginary part of the 1-body TISE (129), that is multiplied by Ψ^* and parameterized with (\mathbf{q}_1') .

Substituting Def. (21) into condition (125), we recognize that condition (125) is equivalent to body mass conservation: $(\nabla_i \cdot (\rho_m \mathbf{v}_i) = \mathbf{0})$. From Theorem 12, under the condition $(\nabla_i \cdot (\rho_m \mathbf{v}_i) = \mathbf{0})$, the *n*-body TISE (124) \Rightarrow (118) with the functions restricted to Range $P[\vec{\mathbf{q}}_0]$. Consider the first equation from equation sequence (118) with initial position $(\mathbf{q}_1, \mathbf{q}'_1) \in \mathbb{R}^{3n}$ at time t_0 of range $(t_0, t) \in \mathbb{R}$ of path $P[(\mathbf{q}_1, \mathbf{q}'_1)]$. At this initial point of space $(\mathbf{q}, \mathbf{q}'_1) \in \mathbb{R}^{3n}$ and initial time $t_0 \in \mathbb{R}$, using Eq. (104) with $((\vec{\mathbf{q}}) = (\vec{\mathbf{q}}_0) = (\mathbf{q}_1, \mathbf{q}'_1))$, this equation becomes Eq. (126), where the body force $-\nabla_1 U$ of Def. (123) is substituted. Since \mathbf{q}_1 of the point $(\mathbf{q}_1, \mathbf{q}'_1)$ is arbitrary, Eq. (126) can be extended from the point $(\mathbf{q}_1, \mathbf{q}'_1)$ to $\mathbb{R}_1^{3n}[\mathbf{q}'_1]$ a.e. Hence, by this extension, $(124) \Rightarrow (126)$,

Almost everywhere, by inspection, Eq. (126) is Euler Eq. (76) for one-body stationary states, under the following special conditions: 1) \mathbf{q}' are parameters, 2) ∇U is defined by definition (123), 3) the *n*-body state is stationary, and 4) constraint (125), where the flow is a 1-body quantum flow. From statement 2 of Theorem 5, we have the following: The 1-body Euler Eq. (76) \Rightarrow the real part of the 1-body TISE (129), multiplied by Ψ^* . Combining this with the results above, we have Eq. (124) and Eq. (125)) \Rightarrow Eq. (129). Implication Eq. (128) follows, because from Theorem 5, we have Eq. (126)) \Rightarrow Eq. (128).

For the implication of Euler Eq. (127), we begin by combining the following equalities for steady, irrotation, smooth flow:

$$\frac{1}{2}\rho_m \nabla_1 (u_1^2 + v_1^2) + \nabla_1 \cdot (\rho_m \mathbf{u}_1) \mathbf{u}_1 = \rho_m \mathcal{D}(\mathbf{u}_1 + \mathbf{v}_1) + \mathbf{u}_1 \nabla_1 \rho_m \cdot \mathbf{u}_1 + \rho_m \mathbf{u}_1 \nabla_1 \cdot \mathbf{u}_1,$$

A Formulation of Quantum Fluid Mechanics and Trajectories

$$\mathcal{D}(\rho_m \mathbf{u}_1) = \rho_m \mathcal{D} \mathbf{u}_1 + \mathbf{u}_1 \nabla_1 \rho_m \cdot \mathbf{u}_1, \quad \mathcal{D}(\rho_m \mathbf{v}_1) = \rho_m \mathcal{D} \mathbf{v}_1 + \mathbf{v}_1 \nabla_1 \rho_m \cdot \mathbf{v}_1 = \rho_m \mathcal{D} \mathbf{v}_1,$$

to obtain

$$\frac{1}{2}\rho_m \nabla_1 (u_1^2 + v_1^2) + \nabla_1 \cdot (\rho_m \mathbf{u}_1) \mathbf{u}_1 = \mathcal{D}(\rho_m \mathbf{u}_1 + \rho_m \mathbf{v}_1) + \rho_m \mathbf{u}_1 \nabla_1 \cdot \mathbf{u}_1$$

Substituting this result in Eq. (126) we obtain Eq. (127).

Renziehausen and Barth [13] obtain an equation of motion for *n*-body trajectories of Bohmian mechanics. The equation has a single total momentum derivative involving all *n* bodies. Also, an Euler equation for *n*-body states that generalizes Euler Eq (83) has also been obtained [68]. For, say particle #1, the equation contains velocities and pressures of the other (n - 1) bodies that act like forces. These equations could be derived for classical systems, but they would be more difficult to solve, then the standard collection of *n* equations of motion for *n* bodies.

11. Quantum Flow Mixture and Orbital Flow

In this section, the forces and fluid momentae of Eq. (127) are reduced to 1-body. A generalization of Coulomb's law is identified. Inferences and hypotheses are employed to move forward and obtain 1-body quantum orbital flow, with motion, energy and Schrödinger equations for n orbital states. Before that, a flow mixture approach is also obtained, where many-body states are treated as 1-body mixtures.

11.1. The quantum reduced forces

In this subsection, the forces of Euler Eq. (127) are reduced to 1-body. The next theorem obtains a modified Coulomb's law with the product of charge densities are replaced by the pair function ρ_2 , a two-body function of reduced density matrix theory [59,60]. We begin with notation.

Definition 11 (Scalar V_e and vector J_e potentials)

$$\vec{\mathcal{E}}(\mathbf{r}_1) \doteq - \left[\nabla_1 V_e + \nabla_1 \times \mathbf{J}_e\right](\mathbf{r}_1) \doteq -2 \int_2 Q(\mathbf{r}_1, \mathbf{r}_2) \nabla_1 W_{12}(\mathbf{r}_2, \mathbf{r}_1) \, d\mathbf{r}_2, \tag{131}$$

$$Q(\mathbf{r}_{1},\mathbf{r}_{2}) \doteq \frac{\rho_{2}(\mathbf{r}_{1},\mathbf{r}_{2})}{\hat{\rho}(\mathbf{r}_{1})}, \quad \rho_{2} \doteq \frac{(n-1)}{2} \sum_{2} \hat{\int}_{2+}^{\hat{\rho}} \rho, \qquad (132)$$

$$Q(\mathbf{r}_1, \mathbf{r}_2) \doteq \frac{1}{2}\hat{\rho}(\mathbf{r}_2) + \frac{1}{2}\rho_{xc}(\mathbf{r}_1, \mathbf{r}_2),$$
(133)

$$1/|\mathbf{r}_1 \cdot \mathbf{r}_2| \doteq \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|}, \quad and \quad 1/|\overrightarrow{|\mathbf{r}_1 \cdot \mathbf{r}_2|^2} \doteq \frac{\mathbf{r}_1 - \mathbf{r}_2}{|\mathbf{r}_1 - \mathbf{r}_2|^3}, \tag{134}$$

where the map $(W_{12}, Q) \mapsto (V_e, \mathbf{J}_e)$, defined by Eq. (131) is such that $|\nabla_1 \times \mathbf{J}_e|$ is minimized. For electronic systems, $(W_{12}(\mathbf{r}_1, \mathbf{r}_2) \doteq |\mathbf{r}_1 \setminus \mathbf{r}_2|^{-1})$. Reduced m-body density

matrices of the wavefunction Ψ are called m-body dentrices. The function Q is the quantum charge density, and ρ_2 is the pair- or 2-density, normalized to (n-1), instead of the usual number of pairs $n(n-1) \div 2$. We have $(\rho_2(\mathbf{r}, \mathbf{r}') = \gamma_2(\mathbf{r}, \mathbf{r}'; \mathbf{r}, \mathbf{r}'))$, where γ_2 is the reduced 2-body dentrix, normalized to $(n-1) \div 2$. As indicated in Def. (26), the probability density function $(\hat{\rho} : \mathbb{R}^3 \to \mathbb{R})$ is parameterized by one spin variable. As indicated in Def. (132.2), the second spin variable ω_2 of ρ_2 , corresponding the second spatial component of $(\mathbf{r}_1, \cdots \mathbf{r}_n) \in \mathbb{R}^3$, is summed over. However, the spatial variable \mathbf{r}_2 is not integrated over. Therefore, ρ_2 has one spin parameter that corresponds to \mathbf{r}_1 .

The requirement that $|\nabla_1 \times \mathbf{J}_e|$ is minimized is given because a Helmholtz decomposition is not unique. This is easily verified by recognizing that a vector field can be both irrotational and solenoidal; such vectors have potentials that satisfy Laplace's equation. An example being the velocity vector \mathbf{v} for hydrogen electron states, except for the s states.

In this section, we make a hypothesis when we want to know if a statement is true, and we think it might be true. Such propositions are considered useful even if they are false, if they provide good approximations for a range of states.

Hypothesis 1 (Force $\vec{\mathcal{E}}$ conservation.) $(\nabla_1 \times \mathbf{J}_e = \mathbf{0})$.

Theorem 14 (Quantum Coulomb's law) Let $W_{12}(\mathbf{r}_1, \mathbf{r}_2) \doteq |\mathbf{r}_1 \setminus \mathbf{r}_2|^{-1}$.

$$\vec{\mathcal{E}}(\mathbf{r}_1) = \frac{1}{4\pi\epsilon_0} \int_2 2Q(\mathbf{r}_1, \mathbf{r}_2) / |\vec{\mathbf{r}}_1 \setminus \mathbf{r}_2|^2; \qquad 4\pi\epsilon_0 = 1.$$
(135)

Under hypothesis 1: $(\vec{\mathcal{E}} \cdot d\hat{\mathbf{s}} = -\nabla V_e \cdot d\hat{\mathbf{s}})$, we have

$$V_e(\mathbf{r}_1) = \frac{1}{4\pi\epsilon_0} \int_{\infty}^{\mathbf{r}_1} \left(-\int_2 2Q(\mathbf{s}, \mathbf{r}_2) / |\mathbf{s}_{\backslash} \mathbf{r}_2|^2 \right) \cdot d\mathbf{s},$$
(136)

that can be written

$$V_e(\mathbf{r}_1) = \frac{1}{2\pi\epsilon_0} \int_2 Q(\mathbf{r}_1, \mathbf{r}_2) / |\mathbf{r}_1 \cdot \mathbf{r}_2| - \frac{1}{2\pi\epsilon_0} \int_{\infty}^{\mathbf{r}_1} \left(\int_2 \nabla_\mathbf{s} Q(\mathbf{s}, \mathbf{r}_2) / |\mathbf{s}_1 \cdot \mathbf{r}_2| \right) \cdot d\mathbf{s}.$$
(137)

Using the definition for the exchange correlation hole ρ_{xc} , Def. (133), we have

$$\vec{\mathcal{E}}(\mathbf{r}_1) = \frac{1}{4\pi\epsilon_0} \int d\mathbf{r}_2 \, \hat{\rho}(\mathbf{r}_2) / |\vec{\mathbf{r}_1 \cdot \mathbf{r}_2}|^2 + \frac{1}{4\pi\epsilon_0} \rho_{xc}(\mathbf{r}_2, \mathbf{r}_1) / |\vec{\mathbf{r}_1 \cdot \mathbf{r}_2}|^2, \tag{138}$$

Proof Using Def. (131) and the identity

$$\nabla_1 \left(1/|\mathbf{r}_1 \cdot \mathbf{r}_2| \right) = -1/|\overrightarrow{\mathbf{r}_1 \cdot \mathbf{r}_2}|^2, \tag{139}$$

we obtain Eq. (135), and from path integration and by Hypothesis 1, Eq. (136). Equations (138) follows immediately by Def. (133). Similarly, for Eq. (137), we have

$$\nabla_{\mathbf{s}} \left(Q(\mathbf{s}, \mathbf{r}_2) / |\mathbf{s}_{\backslash} \mathbf{r}_2| \right) = \nabla_{\mathbf{s}} Q(\mathbf{s}, \mathbf{r}_2) / |\mathbf{s}_{\backslash} \mathbf{r}_2| - Q(\mathbf{s}, \mathbf{r}_2) / |\overrightarrow{\mathbf{s}_{\backslash} \mathbf{r}_2}|^2.$$

Solving for the last term on the rhs and substituting the result into Eq. (136), we obtain Eq. (137).

If we use the 2-density of a closed shell determinantal-state, given by, for our normalizations,

$$\rho_2(\mathbf{r}_1, \mathbf{r}_2) = \rho(\mathbf{r}_1)\rho(\mathbf{r}_2) - \rho_1(\mathbf{r}_1, \mathbf{r}_2)\rho_1(\mathbf{r}_2, \mathbf{r}_1),$$

for the first term on the rhs of Eq. (137), and neglect the second term of $V_{\rm e}$, we have

$$\int d\mathbf{r}_1 V_{\mathbf{e}} \rho(\mathbf{r}_1) = \frac{1}{4\pi\epsilon_0} \int \int d\mathbf{r}_1 d\mathbf{r}_2 \rho(\mathbf{r}_1) \rho(\mathbf{r}_1) / |\mathbf{r}_1 \cdot \mathbf{r}_2| - \rho_1(\mathbf{r}_1, \mathbf{r}_2) \rho_1(\mathbf{r}_2, \mathbf{r}_1) / |\mathbf{r}_1 \cdot \mathbf{r}_2|.$$

This is two times the Hartree Fock Coulomb and exchange energies, in agreement with their contributions to the sum of the occupied orbital energies of Hartree Fock theory, and this double counting is also in agreement with the classical energy Eq. (101) for n point masses.

Theorem 15 (The quantum reduced forces and momentae derivatives) Let

$$\hat{P}_{1} \doteq \hat{\int}_{1+} P = -\zeta \zeta_{0} \nabla^{2} \hat{\int}_{1+} \rho = -\zeta \zeta_{0} \nabla^{2} \hat{\rho}.$$
(140)

$$\int_{1+} -\nabla_1 P_1 - \rho \nabla_1 V - \rho \nabla_1 W_1 = -\nabla_1 \hat{P} - \hat{\rho} \nabla_1 V + \hat{\rho} \vec{\mathcal{E}}.$$
(141)

$$\int_{1+} -\rho \nabla_1 W_1 = \hat{\rho} \vec{\mathcal{E}}; \qquad \vec{\mathcal{E}} \doteq -\nabla_1 V_e - \nabla_1 \times \mathbf{J}_e.$$
(142)

$$\nabla_1 \cdot (\rho \mathbf{v}_1) = \mathbf{0} \; \Rightarrow \; \nabla \cdot (\hat{\rho} \hat{\mathbf{v}}) = \mathbf{0}. \tag{143}$$

$$\hat{\int}_{1+} \frac{d}{dt} (\rho_m \mathbf{u}_1 + \rho_m \mathbf{v}_1) = \frac{d}{dt} (\hat{\rho}_m \hat{\mathbf{u}} + \hat{\rho}_m \hat{\mathbf{v}}).$$
(144)

Steady flow and $\nabla_1 \cdot (\rho \mathbf{v}_1)$ vanish imply

$$\frac{d}{dt}(\hat{\rho}_m\hat{\mathbf{u}} + \hat{\rho}_m\hat{\mathbf{v}}) + \int_{1+}^{\hat{\rho}} \rho_m \mathbf{u}_1 \nabla_1 \cdot \mathbf{u}_1 = -\nabla_1 \hat{P} - \hat{\rho} \nabla_1 V + \hat{\rho} \vec{\mathcal{E}}.$$
(145)

Proof The reduction of $-\nabla P_1$ follows from Def. (140). Recognizing from Def. (121) that V only depends on \mathbf{r}_1 , we have

$$\hat{\int}_{1+}^{\hat{\rho}} \rho \nabla_1 V = \hat{\rho} \nabla_1 V. \tag{146}$$

For the remaining term, using Def. (122), we have

$$\left[\hat{\int}_{1+}^{n} \rho \nabla_1 W_1\right] (\mathbf{r}_1) = \sum_{i>1}^{n} \hat{\int}_{1+}^{n} \rho(\mathbf{r}_1 \cdots \mathbf{r}_n) \nabla_1 W_{12}(\mathbf{r}_i, \mathbf{r}_1) \, d\mathbf{r}_1'.$$

As is done in density matrix theory, consider the term involving $W_{12}(\mathbf{r}_3, \mathbf{r}_1)$. By interchanging the \mathbf{r}_2 and \mathbf{r}_3 dummy variables in the integrand, and using the fact that ρ is symmetric with respect to interchange of coordinates, we recognize that the terms with $W_{12}(\mathbf{r}_2, \mathbf{r}_1)$ and $W_{12}(\mathbf{r}_3, \mathbf{r}_1)$ are equal. Hence, by a generalization, each of the (n-1)terms of the sum are equal under integration, giving

$$\left[\int_{1+}^{\hat{\rho}} \nabla_1 W_1\right](\mathbf{r}_1) = 2 \int_2 \rho_2(\mathbf{r}_1, \mathbf{r}_2) \nabla_1 W_{12}(\mathbf{r}_2, \mathbf{r}_1),$$

where Def. (132.2) for the 2-density ρ_2 is used. Def. (131) for $\vec{\mathcal{E}}$ and Def. (132) for Q then give Eq. (142). Eq. (143) follows from Def. (28.2).

For Eq. (144), first we recognize that the time derivative acts on the first component n-tuple. (It is understood that this first variable is a composition time and position, and the derivative is computed using the chain rule.) Under the assumption of the interchange of the order of differentiation and integration, Eq. (144) follows from Eqs. (28). Eq. (145) follows immediately from Eqs. (127) and (141).

The reduced pressure \hat{P} also appears as a scalar pressure in the many-body hydrodynamics equations of motion of Renziehausen and Barth [13], a formalism involving a pressure tensor without fluid velocity $\hat{\mathbf{u}}$.

We now define a type of flow with reduced variables.

Definition 12 (Reduced quantum flows) The reduced flow $\mathcal{F}(\hat{\rho}, \hat{P}, \hat{\mathbf{u}}, \hat{\mathbf{v}})$, of the quantum n-body flow $\mathcal{F}(\rho, P, \mathbf{u}, \mathbf{v})$, is the flow that has the reduced variables of $\hat{\rho}$, \hat{P} , $\hat{\mathbf{u}}$, $\hat{\mathbf{v}}$ of $\mathcal{F}(\rho, P, \mathbf{u}, \mathbf{v})$, where $\hat{\mathbf{u}}$ and $\hat{\mathbf{v}}$ are given by Def. (28), and the reduced pressure is $(\hat{P} = -\zeta\zeta_0\nabla^2\hat{\rho})$.

Unfortunately we do not know how to reduce the spoiler integral in Eq. (145) involving the volumetric dilation, given the nonlinear velocity \mathbf{u} dependence, thwarting a complete reduced equation of motion. Ideally, we would like to replace Eq. (145) with

$$\mathcal{D}\!\!\int_{\mathbb{V}} \hat{\rho}_m \hat{\mathbf{u}} + \hat{\rho}_m \hat{\mathbf{v}} = \int_{\mathbb{V}} -\nabla_1 \hat{P} - \hat{\rho} \nabla_1 V + \hat{\rho} \vec{\mathcal{E}}.$$
(147)

To try to find a way around this problem, we make an argument that the spoiler integral can be neglected. Consider a generalization of equation of motion (127), where we have a moving control volume with velocity \mathbf{x}_c . Letting ($\mathbf{x} = \mathbf{u}_1 - \mathbf{x}_c$) be the relative velocity of an observer that moves with the control, and taking ($\mathbf{v} = \mathbf{0}$) and steady flow, we have

$$\int_{\mathcal{S}} \rho_m \mathbf{u}_1(\mathbf{x} \cdot \hat{\mathbf{n}}) = \int_{\mathbb{V}} -\nabla_1 P_1 - \rho \nabla_1 V - \rho \nabla_1 W_1 + \rho_m G(\mathbf{x}_c), \qquad (148)$$

where \hat{G} is the noninertial force caused by the accelerating control volume. Let $\mathbf{x} : \mathbb{R}^3 \to \mathbb{R}^3$ be unparameterized: velocity \mathbf{x} is required to be the same velocity field for@all parameters from $\mathbb{R}^{3(n-1)}$. Eqs. (127) and (148) are two different coordinate systems observing the same event, and, therefore, carry the same information. In the special case where ($\mathbf{x}_c = \mathbf{0}$) is taken, equation of motion (148) becomes Eq. (127). (The momentum time derivatives of the two are related by identities (70) and (72).) Applying the operation $\hat{\int}_{1+}^{1+}$, we have

$$\int_{\mathcal{S}} \hat{\rho}_m \hat{\mathbf{u}}(\mathbf{x} \cdot \hat{\mathbf{n}}) = \int_{\mathbb{V}} -\nabla_1 \hat{P} - \hat{\rho} \nabla_1 V + \hat{\rho} \vec{\mathcal{E}} + \hat{\rho}_m \hat{G}(\mathbf{x}_c).$$

Since **x** is at our disposal, we choose $(\mathbf{x} = \hat{\mathbf{u}})$, giving

$$\int_{\mathcal{S}} \hat{\rho}_m \hat{\mathbf{u}}(\hat{\mathbf{u}} \cdot \hat{\mathbf{n}}) = \int_{\mathbb{V}} -\nabla_1 \hat{P} - \hat{\rho} \nabla_1 V + \hat{\rho} \vec{\mathcal{E}} + \int_{1+}^{\hat{\rho}} \rho_m G(\mathbf{u}_1 - \hat{\mathbf{u}}).$$
(149)

Consider the generalized mean value theorem for an integrand that is a product of two functions $f, g : \mathbb{R} \to \mathbb{R}$ that are continuous on \mathbb{R} : For some $\alpha \in \mathbb{R}$, we have

$$\lim_{a \to \infty} \int_{-a}^{+a} fg = g^* \int_{-\infty}^{+\infty} f; \qquad g^* = g(\alpha).$$

Similarly, in our case, we have

$$\int_{1+} \rho_m G(\mathbf{u}_1 - \hat{\mathbf{u}}) = \hat{\rho}_m G(\mathbf{u}_1^* - \hat{\mathbf{u}}).$$

Since $\hat{\mathbf{u}}$ is an average of \mathbf{u}_1 over space $\mathbb{R}^{3(n-1)}$ with weight ρ_m , $(\mathbf{u}_1^* = \hat{\mathbf{u}})$ is, at least, a reasonable approximation. Taking this approximation, the relative velocity \mathbf{x} vanish, and the above equation of motion becomes

$$\mathcal{D}\!\!\int_{\mathbb{V}}\hat{\rho}\hat{\mathbf{w}} = \int_{\mathbb{V}} -\nabla_1\hat{P} - \hat{\rho}\nabla_1V + \hat{\rho}\vec{\mathcal{E}} + \int_{1+}^{\hat{\rho}}\rho_m G(\mathbf{0}),$$

where time derivative identity (72) is substituted. We interpret this equation as the equation of motion for a person at rest relative to the fluid velocity, such that $G(\mathbf{0})$ is the inertial force per mass. This equation and Eq. (147) are, therefore, equations of motions for the same fluid particle of the same flow, but with different observers: Eq. (147) is for an observer at rest, and the one above is for an observer that moves with the fluid particle. Therefore, the two equations are equivalent.

For the general case, where \mathbf{v}_1 does not vanish, we note that for a bases set such that one of the basis vectors is in the direction of \mathbf{v}_1 , for smooth flow, velocities \mathbf{u}_1 and \mathbf{v}_1 do not appear in the same equations of motion (127), and the equation involving \mathbf{v}_1 can be reduced to 1-body.

Another approach is to use the mean value for the momentum flux directly:

$$\hat{\int}_{1+} \int_{\mathcal{S}} \rho_m \mathbf{u}_1(\mathbf{u}_1 \cdot \hat{\mathbf{n}}) = \int_{\mathcal{S}} \hat{\rho}_m \hat{\mathbf{u}}(\mathbf{u}_1^* \cdot \hat{\mathbf{n}}),$$

followed by the same approximation $(\mathbf{u}_1^* = \hat{\mathbf{u}})$.

Renziehausen and Barth [13], in their many-body formalism of Madelung fluid mechanics, are able to reduce the *n*-body momentum time derivative involving velocity (**v**) to 1-body, and obtain reduced equations of motion. Their formalism does not include the second velocity (**u**) and contains a pressure tensor. They are also successful in the reduction of the continuity equation to 1-body: $(\partial \hat{\rho} + \nabla \cdot (\hat{\rho} \hat{\mathbf{v}}) = \mathbf{0})$. This suggests that it might be possible to remove the body mass conservation condition: $(\nabla \cdot (\rho_i \mathbf{v}_i = \mathbf{0})$ from the formalism presented here, but note that it is used in Theorem 13.

In subsection 11.5, a special type of incompressible, cross flows are considered, where the volumetric dilation is absent, permitting, under certain conditions, the complete reduction of the equation of motion to 1-body. In this section, we move forward by inferences and hypotheses.

11.2. Quantum Flow Mixture

In this subsection, we describe quantum flow mixes, obtained by taking force conservation Hypothesis 1, and treating the system as, effectively, having one body. We consider electronic systems with the Bohn–Oppenheimer approximation. In quantum mechanics, the external potential and the number of bodies determines the spatial Schrödinger equation and, thus, all eigenfunctions. Taking Hypothesis 1 and 1-body, we identity, from the quantum reduced forces (141), the 1-body external potential $(V+V_e)$, giving the spatial 1-body Schrödinger equation:

$$-\frac{\hbar^2}{2m}\nabla^2 \mathcal{I} + (V + V_{\rm e})\mathcal{I} = \hat{E}\mathcal{I}.$$
(150)

Let ρ_2 be the pair density of an *n*-body wavefunction with external potential V. Let V_e be determined by ρ_2 via Eq. (136) and (132). Let $\hat{\rho}$ be the 1-density determined by ρ_2 by integration. We want to know if, given such maps: $\rho_2 \mapsto V_e$ and $\rho_2 \mapsto \hat{\rho}$, is there an eigenfunction \mathcal{I} of Eq. (150), such that $(\rho_2 \mapsto \hat{\rho} = \mathcal{II}^*)$.

In classical mechanics, once the forces are known, the momentum is determined by Newton's second law. Comparing Euler Eq. (76) with quantum forces (141), we can infer the equations of motion for the first order, reduced, fluid momentum variables $\hat{\rho}_m \hat{\mathbf{u}}$ and $\hat{\rho}_m \hat{\mathbf{v}}$. We do this now by hypothesis, where the hypothesis is for the most general case, without necessarily the orthogonal condition. This is followed by a definition for quantum flow mixture.

Hypothesis 2 (Quantum flow mixture) Let $\mathcal{F}_R(\hat{\rho}, \hat{P}, \hat{\mathbf{u}}, \hat{\mathbf{v}})$ be the reduced flow of quantum flow $\mathcal{F}_Q(\rho, P, \mathbf{u}, \mathbf{v})$ of an electronic system. Let the wavefunction of \mathcal{F}_Q satisfy the space time Schrödinger equation, and let ρ_2 be the pair density of \mathcal{F}_Q . Let $(V_e = V_e(Q))$ be given by Eq. (136), where, it follows from

$$Q(\mathbf{r}_1, \mathbf{r}_2) \doteq \frac{\rho_2(\mathbf{r}_1, \mathbf{r}_2)}{\hat{\rho}(\mathbf{r}_1)} \quad and \quad \hat{\rho}(\mathbf{r}_1) = \frac{2}{n-1} \int_2 \rho_2(\mathbf{r}_1, \mathbf{r}_2), \tag{151}$$

that $(Q = Q(\rho_2))$. The reduced flow $\hat{\mathcal{F}}_R$ satisfies the first order, reduced Euler-equation:

$$\hat{\rho}_m \partial \hat{\mathbf{v}} + \frac{1}{2} \hat{\rho}_m \nabla \left(\hat{u}^2 + \hat{v}^2 \right) + \nabla \cdot \left(\hat{\rho}_m \hat{\mathbf{u}} \right) \hat{\mathbf{u}} + \nabla \hat{P} + \hat{\rho} \nabla (V + V_e) = \hat{\rho} \vec{\mathcal{E}}, \qquad (152)$$

such that $(\vec{\mathcal{E}} = \vec{\mathcal{E}}(\rho_2))$ and $(\hat{\rho} = \hat{\rho}(\rho_2))$. Also the velocity vector $\hat{\mathbf{v}}$ is irrotational, permitting the following definition for momentum potential \hat{S} and 1-body wavefunction \mathcal{I} :

$$\nabla \hat{S} \doteq m \hat{\mathbf{v}} \Rightarrow \mathcal{I}(\hat{\rho}, \hat{S}) \doteq \sqrt{\hat{\rho}} e^{i\hat{S}/\hbar}, \qquad (153)$$

where \mathcal{I} , normalized to n, is called the orbital of state.

Taking Hypothesis 1, it follows from Theorems (3) and (5) that the reduced continuity Eq. (143) and Euler equation (152) imply

$$\hat{E}\hat{\rho} = \frac{1}{2}\hat{\rho}_m\hat{u}^2 + \frac{1}{2}\hat{\rho}_m\hat{v}^2 + \hat{P} + (V+V_e)\hat{\rho},$$
(154)

and the space time Schrödinger equation:

$$i\hbar\partial\mathcal{I} = -\frac{\hbar^2}{2m}\nabla^2\mathcal{I} + (V+V_e)\mathcal{I},\tag{155}$$

becoming Eq. (150) for stationary states.

Definition 13 (Quantum flow mixture) The flow mixture $\mathcal{F}_M(\mathcal{I}, \hat{P}, \hat{\mathbf{u}}, \hat{\mathbf{v}})$ is the reduced quantum flow that satisfies Eq. (152) and (153) with orbital of state ($\mathcal{I} = \mathcal{I}(\hat{\rho}, \hat{S})$), such that the pair density ρ_2 , from the corresponding quantum flow, determines the 1-density $\hat{\rho}$ and the quantum Coulomb's force $\hat{\mathcal{E}}$ of Euler Eq. (152).

The kinetic energy from Eq. (154) is calculated using the strong, first order reduced velocity variables given by Def. (6). This kinetic energy differs from the sum of the strong, first order reduced kinetic energy variables:

$$\int_{1+}^{\hat{}} \rho_m u^2 + \rho_m v^2.$$

Given a choice between the two kinetic energies, the strong variables of the velocity is a better fit for Euler Eq. (152), because equations of motion involves the velocity, not kinetic energy.

The next definition gives a modification of the Dirac notation that explicitly distinguishes between operators and their defining kernel, useful for denoting density matrices, that we call dentrices.

Definition 14 (Explicit Dirac Notation) Let $\alpha \rangle \doteq \alpha : \mathbb{R}^{3n} \to \mathbb{C}$ and $\langle \beta \doteq \beta^* : \mathbb{R}^{3n} \to \mathbb{C}$. The kernel of the integral operator $|\alpha\rangle\langle\beta|$, written in Dirac notation, and involving given functions α and β , is denoted $\alpha\rangle\langle\beta$, in other words, we have $(\alpha\rangle\langle\beta(\mathbf{r},\mathbf{r}') = \alpha(\mathbf{r})\beta^*(\mathbf{r}'))$. The maps $\langle \alpha|$ and $\beta\rangle\langle\alpha|$ are the functional and operator with values $\langle \alpha|\gamma\rangle$ and $\alpha\rangle\langle\alpha|\gamma\rangle$ at the point $\gamma \in L^2$, respectively.

Since the function $(\hat{\rho}_1 \doteq \mathcal{I})\langle \mathcal{I}\rangle$ is given as a complex product of a wavefunction \mathcal{I} of a noninteracting state, it cannot be the 1-dentrix ρ_1 of a wavefunction Ψ of an interacting *n*-body state [59,61]. However, this difference is not relevant in the evaluation of the truth value of Schrödinger Eq. (155). To investigate this kinetic energy issue further, consider the well known energy functional of the 2-dentrix for stationary states [59,60]:

$$E = n \int_{1}^{\hat{\mu}} -\frac{\hbar^{2}}{2m} \hat{\nabla}^{2} \rho_{1} + V \hat{\rho} + n \int_{1}^{\hat{\mu}} \int_{2}^{\hat{\mu}} \frac{1}{\mathbf{r}_{12}} \rho_{2}; \qquad \mathbf{r}_{12}(\mathbf{r}, \mathbf{r}') = |\mathbf{r} - \mathbf{r}'|, \qquad (156)$$

$$\left[\hat{\nabla}^2 \rho_1\right](\mathbf{r}) = \nabla^2_{\mathbf{r}} \rho_1(\mathbf{r}, \mathbf{r}') \big|_{\mathbf{r}'=\mathbf{r}}, \quad \rho_1(\mathbf{r}, \mathbf{r}') = \int_{1+}^{2} \Psi(\mathbf{r}, \mathbf{r}_2, \cdots, \mathbf{r}_n) \Psi^*(\mathbf{r}', \mathbf{r}_2, \cdots, \mathbf{r}_n).$$

where the *n* factors appear because of our normalization choice (132) for ρ_2 . This equation is derived from the expectation value of the Hamiltonian \hat{H} . In order to express the kinetic energy in the integral form above, it is necessary to introduce an additional variable that endows the 1-dentrix ρ_1 with domain $\mathbb{R}^{3n} \times \mathbb{R}^{3(n+1)}$. Such an extended domain, giving an unphysical function, does not appear in the kinetic energy identity (36). Therefore, we cannot end up with $\mathcal{I} \rangle \langle \mathcal{I}$ being the 1-dentrix ρ_1 of the wavefunction Ψ of the corresponding quantum flow. (Note that \mathbf{r}_{12} , defined above, is a function, not the value of a function at $(\mathbf{r}_1, \mathbf{r}_2)$.)

11.3. Quantum Orbital Flow

In this subsection, we describe quantum orbital flow, obtained by taking force conservation Hypothesis 1, where *n*-body systems are treated, explicitly, as having *n* bodies. We take the *n* body external potential as a sum over a one body potentials, with value $(W(\mathbf{r}_1) + \cdots + W(\mathbf{r}_n))$, where $(W = V + V_e)$ is the potential of the reduced quantum forces (141).

Suppose we replace the Coulomb potential $(W_{12}(\mathbf{r}, \mathbf{r}') = 1 \div |\mathbf{r} - \mathbf{r}'|)$ by a modification that is body dependent $(W_i(\mathbf{r}_i, \mathbf{r}') = \alpha_i |\mathbf{r}_i - \mathbf{r}'|^{-1})$; $(\alpha_i \in \mathbb{R})$ and $(i = 1, \dots, n)$. This symmetry breakage gives n reduced equations of motions, replacing the single Eq. (145), where the *i*th equation is obtained by integration over $\mathbf{r}_1, \dots, \mathbf{r}_{i-1}, \mathbf{r}_{i+1}, \dots, \mathbf{r}_n$, and summation over the corresponding spin variables. Under a hypothesis that is similar to Hypothesis 2, we obtain n Schrödinger equations, replacing Eq. (155), where we normalize each orbital \mathcal{I}_i to unity. It is reasonable to expect that a solution set of n linearly independent orbitals exist for the n Schrödinger equations. If we take the limit $(\alpha_i \to 1)$, we would then expect to have n orthonormal orbitals satisfying the n Schrödinger equations. Hence, encoded in the orbital of state \mathcal{I} must be n linearly independent 1-body orbitals that can be extracted.

In Hartree–Fock theory for fermions, antisymmetry of the wavefunction with respect to interchange of bodies is imposed on the wavefunction by using a Slater determinant, giving n orthogonal orbitals for a solution set, instead of one occupied orbital for all bodies. Similarly, we aim to replace Eq. (155) with n orbital equations. However, the variational approach used by the Hartree–Fock method does not work for our situation. Therefore, we must satisfy this need by hypothesis. Even though the hypothesis is as general as possible, without modifications, it might only be suitable for states that can be represented by a single Slater determinant: closed shell singlet and spin polarized open shells on top of a closed shell.

Hypothesis 3 (Quantum Orbital Flow) Let the density $\hat{\rho}$ be normalized to the usual n, given by Def. (5.3) with the replacement $1 \rightarrow n$. There exists a set of linearly independent orbitals $\{\psi_1, \psi_2, \dots, \psi_n\}$; $(\psi_i = \rho_i e^{iS_i/\hbar})$, that satisfy the following sequence of n Euler equations:

$$\bar{\rho}_i \partial \mathbf{v}_i + \frac{1}{2} \bar{\rho}_i \nabla \left(u_i^2 + v_i^2 \right) + \nabla P_i + \left(\nabla \cdot \bar{\rho}_i \mathbf{u}_i \right) \mathbf{u}_i + \rho_i \nabla (V + V_e) = \rho_i \vec{\mathcal{E}}; \quad \bar{\rho}_i = m \rho_i, \quad (157)$$

and

$$\partial \rho_i + \nabla \cdot (\rho_i \mathbf{v}_i) = \mathbf{0} \quad (\Rightarrow \ \partial \hat{\rho} + \nabla \cdot (\hat{\rho} \hat{\mathbf{v}})),$$
(158)

involving orbital densities $(\rho_i \doteq \psi_i \psi_i^*)$, such that $(\vec{\mathcal{E}} = \vec{\mathcal{E}}(\rho_2))$, $(\hat{\rho} = \hat{\rho}(\rho_2))$, $(\hat{\rho} = \rho_1 + \cdots + \rho_n)$, and the set $\{S_1, \cdots, S_n\}$ satisfies the weighted average equation for \hat{S} :

$$\sum_{i}^{n} \hat{\rho}_{i} \nabla S_{i} \doteq \nabla \hat{S}; \quad \hat{\rho}_{i} \doteq \frac{\rho_{i}}{\hat{\rho}} \doteq \frac{\psi_{i} \psi_{i}^{*}}{\hat{\rho}} \Rightarrow \hat{\rho} \hat{\mathbf{v}} = \sum_{i} \rho_{i} \mathbf{v}_{i}.$$
(159)

Also under hypothesis (1), giving $(\vec{\mathcal{E}} = -\nabla V_e)$, it follows from Theorems (3) and (5) that the Euler equation (157) and continuity Eq. (158) imply corresponding sequences of Schrödinger and energy equations.

not appear.

Definition 15 (Quantum orbital flow) Consider the following sequence of n orbital field equations:

$$i\hbar\partial\psi = -\frac{\hbar^2}{2m}\nabla^2\psi + (V+V_e)\psi; \quad \hat{\rho} = \psi\psi^*, \quad where \quad \psi \in \Sigma,$$
 (160)

reducing to the following equation sequence for stationary states:

$$-\frac{\hbar^2}{2m}\nabla^2\psi + (V+V_e)\psi = \varepsilon\psi, \qquad \varepsilon\psi \in \Sigma.$$
(161)

Let \mathcal{F}_Q be a quantum flow, whose wavefunction satisfies the space time Schrödinger equation. We take the hypothesis that $\hat{\mathbf{v}}$ of \mathcal{F}_Q is irrotational, giving definition $(m\hat{\mathbf{v}} \doteq \nabla \hat{S})$. The corresponding n component orbital flow $\hat{\mathcal{F}}_O(\{\psi_i, P_i, \mathbf{u}_i, \mathbf{v}_i\})$ is defined by a set of linearly independent orbitals of state $\{\psi_1, \dots, \psi_n\}$ that satisfy the sequence of n Schrödinger Eqs (160), and the elements of the set $\{S_1, \dots, S_n\}$ satisfy Eq. (159), where

$$\psi_i \doteq \pm \sqrt{\rho_i} e^{iS_i/\hbar}, \quad \psi_i : \mathbb{R}^3 \to \mathbb{C}, \quad \rho_i \doteq \psi_i \psi_i^*, \quad \int_1 \psi_i \psi_i^* = 1.$$
 (162)

Also, the pair density ρ_2 of \mathcal{F}_Q determines both the density $\hat{\rho}$ and potential V_e . Furthermore, as in the above definitions, we have

$$\bar{\rho}_i \mathbf{u}_i \doteq -\zeta_0 \nabla \rho_i, \quad m \mathbf{v}_i \doteq \nabla S_i, \qquad P_i \doteq \zeta_0 \nabla \cdot (\rho_i \mathbf{u}_i), \quad p_i \doteq -\zeta_0 \nabla \cdot (\rho_i \mathbf{v}_i),$$
$$\varepsilon_i = -\partial S_i, \quad and \quad \varphi_i = -\partial \theta_i.$$

In addition, according to Theorems (3) and (5), the orbital flow $\hat{\mathcal{F}}_O$ satisfy the following two n equations sequences, where all terms run:

$$\frac{1}{2}\rho_m u^2 + \frac{1}{2}\rho_m v^2 + P + (V + V_e)\rho = \varepsilon\rho, \quad \partial\rho + \nabla \cdot (\rho \mathbf{v}) = \mathbf{0}, \quad and \tag{163}$$

$$\rho_m \partial \mathbf{v} + \frac{1}{2} \rho_m \nabla \left(u^2 + v^2 \right) + \nabla P + (\nabla \cdot \rho \mathbf{u}) \mathbf{u} + \rho \nabla (V + V_e) = \mathbf{0}.$$
(164)

Next we show that the orbital and mixed quantum flows are related.

Theorem 16 Consider the following equations:

1) The n-body Schrödinger equation

$$i\hbar\partial\hat{\mathcal{I}} = \hat{H}\hat{\mathcal{I}}; \qquad \hat{H} = -\frac{\hbar^2}{2m}\sum_{i=1}^n \nabla_i^2 + \sum_{i=1}^n \left[V + V_e\right](\mathbf{r}_i),$$
(165)

where $(\hat{\mathcal{I}}: \mathbb{R}^{3n} \to \mathbb{C})$ is the Slater determinant $\hat{\mathcal{I}}$ constructed from $\{\psi_1, \dots, \psi_n\}$. 2) The sequence of n Schrödinger Eqs. (160) of the orbitals of state $\{\psi_1, \dots, \psi_n\}$, in the form

$$i\hbar\partial\psi\rangle\langle\psi=-\frac{\hbar^2}{2m}\nabla^2\psi\rangle\langle\psi+(V+V_e)\psi\rangle\langle\psi.$$

A Formulation of Quantum Fluid Mechanics and Trajectories

3) The single Schrödinger Eq. (155), in the form

$$i\hbar\partial\mathcal{I}\rangle\langle\mathcal{I}=-\frac{\hbar^2}{2m}\nabla^2\mathcal{I}\rangle\langle\mathcal{I}+(V+V_e)\mathcal{I}\rangle\langle\mathcal{I},$$

for the special case, such that

$$\mathcal{I} \langle \mathcal{I} = \psi_1 \rangle \langle \psi_1 + \dots + \psi_n \rangle \langle \psi_n; \qquad \int \psi_i \psi_j^* = \delta_{ij}.$$
(166)

 $1 \Rightarrow 2 \Rightarrow 3$. Also, the function $\mathcal{I} \setminus \langle \mathcal{I} : \mathbb{R}^3 \times \mathbb{R}^3 \longrightarrow \mathbb{C}$ is the 1-dentrix from the Slater determinant $\hat{\mathcal{I}}$ constructed from the orthonormal orbital set $\{\psi_1, \dots, \psi_n\}$.

Proof 1 Since Eq. (165) is separable, there must be a set of n orthonormal orbitals giving Slater determinant $\hat{\mathcal{I}}$ that satisfies Eq. (165), such that the orbital set satisfies the equation sequence (160). Summing Eq. (160)× ψ^* over the n equations of the sequence, we obtain Eqs. (155), in the form above, where the 1-dentrix satisfies Eq. (166). Since there is a one-to-one correspondence between determinantal states and 1-dentrixes given by the sum from Eq. (166), the function $\mathcal{I} \setminus \langle \mathcal{I} \rangle$ is the 1-dentrix from the Slater determinant $\hat{\mathcal{I}}$ formed from $\{\psi_1, \dots, \psi_n\}$.

11.4. Obtaining solutions of Schrödinger Eq. (161) and energy Eq. (156)

In the discussions that follows, we restrict the discussion to stationary states. Furthermore, we assume that Hypothesis 3 holds, or provides a good approximation, for a range of states, especially states that can be described by a single Slater determinant.

One approach to obtain an approximate solution sets $\{\psi_1, \dots, \psi_n\}$ for the spatial Schrödinger Eq. (161), where V_e is determined by the 2-density, is to use an approximate map $\hat{\rho} \mapsto \rho_2$ of the 1-density, where $\hat{\rho}$ is equal to the orbital density sum $|\psi|^2$; $|\psi|^2 \in \Sigma$. Given such a map as a composite function or functional, the Schrödinger Eqs. (161) can be solved using the self consistent field method. The energy can then be calculated using Eq. (156), where, as in the approximation of the kinetic energy in Kohm–Sham density functional theory, the kinetic energy is approximated by the one obtained from the 1-dentrix $\mathcal{I} \setminus \langle \mathcal{I} \rangle$ of Eq. (166) of the noninteracting state \mathcal{I} .

Approximations of ρ_2 can come from functionals of the electron interaction energy $E_{\rm e}$ —the last term in the rhs of energy Eq. (156.1)—given as a functional of the density, where the \mathbf{r}_{12} function is replaced by variable W_{12} :

$$[E_{\mathbf{e}}(W_{12})](\hat{\rho}) \doteq \left[\int_{1}^{\hat{\rho}} \int_{2}^{\hat{\rho}} W_{12} \rho_{2}\right](\hat{\rho}).$$

The 2-density ρ_2 is then given by a functional derivative:

$$\rho_{2} = \frac{\delta E_{e}}{\delta \mathbf{r}_{12}^{-1}}, \qquad \frac{\delta E_{e}}{\delta \mathbf{r}_{12}^{-1}} \doteq \left[\frac{\delta E_{e}}{\delta W_{12}}\right] \left(\mathbf{r}_{12}^{-1}\right); \quad E_{e}[\rho] = \frac{1}{2} \int_{1}^{2} \int_{2}^{2} \frac{1}{\mathbf{r}_{12}} \hat{\rho}(\mathbf{r}_{1}) \hat{\rho}(\mathbf{r}_{2}) + E_{\mathrm{XC}}[\rho].$$

There is a disconnect between most of the functional used in Kohm–Sham density functional theory, since the theory does not provide for the construction of a useful

way to obtain approximation. The same functionals are applicable other approaches. For example, the functionals can be used for reference state one-particle density-matrix theory, involving Brueckner orbitals [69, 70]. Similarly, the same exchange-correlation energy functionals $E_{\rm XC}$ used in Kohm–Sham density functional theory can be used to obtain an approximate for $(\rho_2 = \rho_2[\hat{\rho}])$.

The 2-body function $\mathcal{I} \subset \mathcal{I}$ from a determinantal state $\hat{\mathcal{I}}$, given by Eq. (166), cannot be the 1-dentrix of an *n*-body interacting state. We now consider if it is at all possible that the formalism can be modified so that the n orbital expression (166) can deliver an interacting state 1-dentrix. First note that n linearly independent orbitals can give the 1-dentrix of an interacting state, if the orbitals are nonorthogonal. This can be seen by partitioning natural orbitals into n separate sums of orbitals, where some orbitals are divided into two by using $(\psi = c\psi + (1-c)\psi)$, such that each of the *n* orbitals are normalized. Since mixed states of quantum mechanics are, in general, nonorthogonal, the proper description for the embedded states considered here might be nonorthogonal orbitals. Unfortunately, nonorthogonal orbitals cannot provide a solution to equation sequence (161), because the eigenstates of a Hermitian operator are orthogonal. Therefore, the only option is to generalize the Hamiltonian to be nonhermitian, or use a generalized Euler equation. The use of nonhermitian operators provide opportunities for interpretations arising from the possibility of linearly dependent-eigenfunctions and complex valued-eigenvalues, given that the orbital states can be considered part of a mixed state.

11.5. Reduced Cross Flows

In this subsection, we investigate an approach, based on smooth, cross flows, that permit a way to get around the problem of a complete reduction of equation of motion (145). Such flows do not have the spoiler integral in their equation of motion. From Theorem 8, these flows have a constant density on each streamline, but the density depends on the streamline. The next theorem demonstrates that if there exists a constant density cross flow $\mathcal{F}_{\times}(P_1, \rho_m, \boldsymbol{\mu}_1, \mathbf{v}_1)$, corresponding to smooth flow $\mathcal{F}(P_1, \rho_m, \mathbf{u}_1, \mathbf{v}_1)$, that satisfies the equation of motion

$$\rho_m \mathcal{D}\boldsymbol{\omega} + \nabla P_1 + \rho \nabla V_1 + \rho \nabla W_1 = \rho \vec{F}, \qquad (167)$$

for any non work force \vec{F} , then flows $\mathcal{F}_{\times}(P, \rho_m, \boldsymbol{\mu}, \mathbf{v})$ and $\mathcal{F}(P, \rho_m, \mathbf{u}, \mathbf{v})$ satisfy the same Euler equation, except for nowork forces. Conditions for the energy to be uniform and the satisfaction of the spatial Schrödinger equation are also given. Also, if the cross flow is solenoidal, mass is conserved.

Theorem 17 (Cross Flow Euler Equation Reduction) Let C_1 be the set of all cross flows $\mathcal{F}_{\times}(P_1, \rho_m, \boldsymbol{\mu}_1, \mathbf{v}_1)$ of parameterized, smooth flow $\mathcal{F}(P_1, \rho_m, \mathbf{u}_1, \mathbf{v}_1)$ that satisfy Euler cross flow Eq. (167), corresponding to Euler Eq. (127), with an additional nowork force \vec{F} , where ($\boldsymbol{\omega} = \boldsymbol{\mu}_1 + \mathbf{v}_1$), and \vec{F} can depend on the cross flow. In addition, the n-body wavefunction ($\Psi = \Psi(S, \rho)$) of \mathcal{F} is parameterized by the point set (\mathbf{q}'_1) $\in \mathbb{R}^{n-1}$, and Ψ satisfies the TISE (129).

A Formulation of Quantum Fluid Mechanics and Trajectories

1) If C_1 is nonempty, each $\mathcal{F}_{\times}(P_1, \rho_m, \boldsymbol{\mu}_1, \mathbf{v}_1) \in C_1$ satisfies energy equation (128), and their corresponding reduced flows $\mathcal{F}_{\times}(\hat{P}, \hat{\rho}_m, \hat{\boldsymbol{\mu}}, \hat{\mathbf{v}})$ satisfy the following reduced cross flow Euler equation:

$$\mathcal{D}(\hat{\rho}_m \hat{\boldsymbol{\omega}}) + \nabla \hat{P} + \hat{\rho} \nabla V_1 = \hat{\rho} \vec{\mathcal{E}} + \hat{\rho} \vec{f}; \qquad \hat{\rho} \vec{f} = \int_{1+} \rho \vec{F}, \qquad (168)$$

where $(\hat{\boldsymbol{\omega}} = \hat{\boldsymbol{\mu}} + \hat{\mathbf{v}}).$

Parts 2 and 3 below hold under hypothesis 1: $(\vec{\mathcal{E}} \cdot d\hat{\mathbf{s}} = -\nabla V_{\mathbf{e}} \cdot d\hat{\mathbf{s}})$, giving from Eq. (168),

$$\mathcal{D}(\hat{\rho}_m \hat{\boldsymbol{\omega}}) + \nabla \hat{P}_1 + \hat{\rho} \nabla V_1 + \hat{\rho} \nabla V_e = \hat{\rho} \vec{f}.$$
(169)

2) If C_1 is nonempty, then each $\mathcal{F}_{\times}(\hat{\boldsymbol{\mu}}, \hat{\mathbf{v}})$ of $\mathcal{F}_{\times}(\boldsymbol{\mu}_1, \mathbf{v}_1) \in C_1$ satisfies the following energy equation:

$$\hat{E}\hat{\rho} = \frac{1}{2}\hat{\rho}_m\hat{\mu}^2 + \frac{1}{2}\hat{\rho}_m\hat{v}^2 + \hat{P} + (V_1 + V_e)\hat{\rho}.$$
(170)

The continuity equation $(\nabla \cdot \hat{\rho}_m \hat{\mathbf{v}}) = \mathbf{0}$ is also satisfied.

3) If $(\hat{u}^2 = \hat{\mu}^2)$ then reduced flow $\mathcal{F}(\hat{\mathbf{u}}, \hat{\mathbf{v}})$ also satisfies energy Eq. (170). If, in addition, energy \hat{E} of Eq. (170) is uniform, and $\hat{\mathbf{v}}$ is irrotational, giving $(\nabla \mathcal{S} \doteq m \hat{\mathbf{v}})$, then the wavefunction $(\mathcal{I} = \sqrt{\hat{\rho}} e^{i\mathcal{S}/\hbar})$ of reduced from $\mathcal{F}(\hat{\mathbf{u}}, \hat{\mathbf{v}})$ satisfies the 1-body Schrödinger Eq. (150).

Proof

1) Each $\mathcal{F}_{\times}(P_1, \rho_m, \boldsymbol{\mu}_1, \mathbf{v}_1) \in \mathcal{C}_1$ satisfies energy equation (128) follows from Theorem 9. Each $\mathcal{F}_{\times}(\hat{P}, \hat{\rho}_m, \hat{\boldsymbol{\mu}}, \hat{\mathbf{v}})$ satisfies the reduced cross flow Euler Eq. (168), follows from Theorem (15), equality (141), for the forces, and that $(\rho_m \mathcal{D}\boldsymbol{\omega} = \mathcal{D}(\rho_m \boldsymbol{\omega}))$, because, according to Theorem 8, smooth cross flows have constant density, and $\rho_m \boldsymbol{\omega}$ can be reduced.

2 Each $\mathcal{F}_{\times}(\hat{\boldsymbol{\mu}}, \hat{\mathbf{v}})$ satisfies reduced energy equation (170) follows from the direct integration of Euler Eq. (169) along the streamline, as is done in Theorem (9). Since the wavefunction Ψ satisfies the TISE, the continuity equation, given by $(\nabla_1 \cdot (\rho_m \mathbf{v}_1) = \mathbf{0})$, is satisfied, implying that $(\nabla \cdot (\hat{\rho}_m \hat{\mathbf{v}}) = \mathbf{0})$.

3 If $(\hat{u}^2 = \hat{\mu}^2)$ then reduced flow $\mathcal{F}(\hat{\mathbf{u}}, \hat{\mathbf{v}})$ also satisfies energy Eq. (170) because the other flow parameters are the same for both reduced flows. The satisfaction of the Schrödinger equation follows from Theorem 3.

12. Summary and Conclusion

The work presented here provides a foundation of a formalism for quantum states based on the cause and effect of Newton's second law. While the energy and equations of motion derived here are applicable to models involving fluid flow and particle trajectories, the best possible description of the nature of states of quantum systems,

consistent with these equations, might be something different. Since a trajectory model has limitations set by the uncertainty principle involving position and momentum, a model based on a continuum is preferred over the trajectory of point masses. According to Broer [56], at least during the 1970s, there was considerable interests in fluid models for this reason. At this stage of development, it is useful to develop many interpretations of the equations, for use in a process of elimination.

Theorem 1 connects different definitions associated with the imaginary part of the Schrödinger equation in density matrix form. Theorem 3 demonstrates that n-body Bernoullian Energy Eq. (45.1) and continuity Eq. (45.2) are equivalent to the Schrödinger equation. Theorem 4 provides a generalized equation that removes a nonclassical feature of the kinetic energy formula. Section 8.2 gives a complete identification of the terms in the spectrum of the Euler equations of Theorem 5, 6, and 7, and identifies the flow type as compressible, inviscid, irrotational, and variable mass. The cross flows of Sec. 9 demonstrate that it is possible to assign kinetic energy to a velocity vector that have properties that are closer to ones seen in systems of classical mechanics, including orbit trajectories with continuous velocities and conserved mass fluid flow. The standard Lagrangian formulation of classical mechanics is described in Sec. 10.

Via theorems, and reasonable hypotheses, for orbital flows of *n*-body states, Sec. 11 obtains sets of *n* Energy (163), Euler (164) and 1-body Schrödinger Eqs. (160). The Schrödinger equations are generalizations of the Hartree–Fock equations with the Coulomb and exchange potentials replaced by potential $V_{\rm e}$, giving a quantum Coulomb's law of force. The same set of three equations are obtained for flows, called mixed flows, that are defined by Quantum Flow Defs. (15). For mixed flows, instead of *n* 1-body equations, there is only one reduced Euler, energy and Schrödinger equation, to represent an *n*-body state. Section 11.5 demonstrates that certain cross flows, that are investigated in Sec. 9, can be reduced to the same Euler equation of mixed flows, but without the need for hypothesis.

Appendix A. Bohmian mechanics Eqs. and Schrödinger Eq. equivalence

Theorem 18 (Bohmian Mechanics Equations) The Bohmian Mechanics equations

$$-\partial S = \frac{1}{2}mv^2 + Q + U, \quad \partial \rho + \frac{1}{m}\nabla \cdot (\rho \nabla S) = \mathbf{0}, \tag{A.1}$$

and the definition of the quantum potential Q:

$$Q \doteq -\frac{\hbar^2}{2m} \rho^{-1} R \nabla^2 R, \quad Q \in \mathbf{\Sigma}, \tag{A.2}$$

are equivalent to equation set $(45) \times \rho^{-1}$, definitions (41) for E and F, definition (32) for P and p, and the second equation of (25) for u^2 . The Bohmian mechanics Eqs. (A.1) and (A.2) are also equivalent to the Schrödinger Eq. (11). A Formulation of Quantum Fluid Mechanics and Trajectories

Note that both approaches use

$$((m\mathbf{v}) \doteq (\nabla S))$$
 giving $\partial \rho + \nabla \cdot (\rho \mathbf{v}) = \mathbf{0},$ (A.3)

but this is not needed for the proof that follows.

Proof First we prove that the first equations of $(45) \times \rho^{-1}$ and (A.1) are equivalent. Comparing these two equations, and using the first definition from (41) $(E = -\partial S)$, the definition above for (Q), and noting that the equations have the same definition for (v^2) , it follows that the two equations are equivalent, if and only if

$$-\frac{\hbar^2}{2m}\rho^{-1}R\nabla^2 R = \frac{1}{2}mu^2 + P\rho^{-1}.$$
 (A.4)

The identity:

$$-\frac{\hbar^2}{2m}\rho^{-1}R\nabla^2 R = \frac{\hbar^2}{8m}\rho^{-2}\nabla\rho\cdot\nabla\rho - \frac{\hbar^2}{4m}\rho^{-1}\nabla^2\rho,$$

is easily proved by expanding out $(\nabla^2 \rho)$ or by setting S to a constant in (19). (A proof is also give elsewhere [1].) Hence, we only need to prove that the rhs of this identity is equivalent to the rhs of (A.4). Substituting definition (25) for u^2 and (33) for P, the above identity becomes (A.4), completing the first part of the proof.

To complete the proof for the first statement, we show that the second equations of (45) and (A.1) are equivalent:

Substituting definitions (41): $F\rho \doteq \frac{\hbar}{2}\partial\rho$, and (33): $p \doteq -\frac{\hbar}{2m}\nabla \cdot (\rho\nabla S)$, into definition (45): $(F\rho = p)$, gives the second equation of (A.1), if the equation is multiplied by $2/\hbar$ and solved for the zero function.

For the second statement, the equivalence of the Bohmian mechanics Eqs. (A.1) and (A.2) and the Schrödinger Eq. (11) follows immediately from Theorem 3.

- [1] J. P. Finley. J. Phys. Commun., 6:04002, 2022.
- [2] D. Bohm. *Phys. Rev.*, 85:166, 1952.
- [3] D. Bohm. Phys. Rev., 85:180, 1952.
- [4] T. Takabayasi. Prog. Theor. Phys., 11:341, 1954.
- [5] A. S. Sanz. Front. Phys, 14:11301, 2019.
- [6] A. S. Sanz and S. Miret-Arteś. A Trajectory Description of Quantum Processes. II. Applications, vol. 831 of Lecture Notes in Physics. Springer, Berlin, 2014.
- [7] D. Dürr and S. Teufel. Bohmian Mechanics. Springer, Heidelberg, 2009.
- [8] R. E. Wyatt. Quantum Dynamics with Trajectories. Springer, New York, 2005.
- [9] P. K. Chattaraj, ed. Quantum Trajectories. CRC Taylor and Francis, New York, 2010.
- [10] K. H. Hughes and G. Parlant, Eds. Quantum Trajectories. CCP6, Daresbury, 2011.
- [11] X. Oriols and J. Mompart, Eds. Applied Bohmian Mechanics: From Nanoscale Systems to Cosmology. Pan Standford Publishing, Singapore, 2012.
- [12] K. Jung. J. Phys.: Conf. Ser., page 442, 2013.
- [13] K. Renziehausen and I. Barth. Found. Phys., 50:772, 2020.
- [14] E. Madelung. Naturwissenschaften, 14:1004, 1926.
- [15] E. Madelung. Z. Phys., 40:322, 1927.
- [16] E. Heifetz and E. Cohen. Found. Phys., 45:1514, 2015.

57

- [17] E. Heifetz et al. On entropy production in the Madelung fluid and the role of Bohm's potential in classical diffusion, the manuscript is available from http://arXiv.org/abs/1509.01265v2, 2015.
- [18] A. L. Sorokin. Dokl. Phys, 46:576, 2001.
- [19] P. Broadbridge. Symmetry, 7:1803, 2015.
- [20] M. Schönberg. Il Nuovo Cimento, 12:103, 1954.
- [21] M Caliari et al. Il Nuovo Cimento, 6:69, 2004.
- [22] A. Jamali. Nonlinear Generalisation of Quantum Mechanics, the manuscript is available from http://preprints.org/manuscript/202108.0525/v1, 2021.
- [23] M. Waegell. Quantum Stud.: Math. Found., pages 1–18, 2024.
- [24] R. Tsekov. Ann. Univ. Sofia, Fac. Phys. Special Edition, [arXiv 0904.0723], page 112, 2012.
- [25] P. Vadasz. Fluids, 1:18, 2016.
- [26] T. Takabayasi. Prog. Theor. Phys., 8:143, 1952.
- [27] T. Takabayasi. Prog. Theor. Phys., 14:283, 1955.
- [28] D. Bohm, R. Schiller, and J. Tiomno. Nuovo Cimento 1, 1:48, 1955.
- [29] R. J. Harvey. Phys. Rev., 152:1115, 1966.
- [30] I. Bialynicki-Birula and Z. Bialynicka-Birula. Phys. Rev. D, 3:2410, 1971.
- [31] N. Rosen. Nuovo Cimento B, 19:90, 1974.
- [32] B. M. Deb and A. S. Bamzai. Mol. Phys, 35:1349, 1978.
- [33] T. C. Wallstrom. Phys. Lett. A, 184:229, 1994.
- [34] E. Recami and G. Salesi. Phys. Rev. A, 57:98, 1998.
- [35] R. E. Wyatt. J. Chem. Phys., 117:9568, 2002.
- [36] C. L. Lopreore and R. E. Wyatt. Phys. Rev. Lett., 82:5190, 1999.
- [37] C. L. Lopreore and R. E. Wyatt. Chem. Phys. Lett., 325:73, 2000.
- [38] C. L. Lopreore and R. E. Wyatt. J. Chem. Phys., 116:1228, 2002.
- [39] T. Koide. Phys. Rev. C, 87:034902, 2013.
- [40] R. E. Wyatt, D. J. Kouri, and D. K. Hoffman. J. Chem. Phys., 112:10730, 2000.
- [41] R. E. Wyatt and E. R. Bittner. J. Chem. Phys., 113:8898, 2000.
- [42] L.S. Kuzmenkov and S.G.Maksimov. Theor. Math. Phys., 118:227, 1999.
- [43] K. Renziehausen and I. Barth. Prog. Theor. Exp. Phys, page 013A05, 2018.
- [44] L.S. Kuzmenkov, S.G.Maksimov, and V. V. Fedoseev. Theor. Math. Phys., 126:110, 2001.
- [45] P.A. Andreev and L. S. Kuzmenkov. Russ. Phys. J., 50:1251, 2007.
- [46] P. A. Andreev and L. S. Kuzmenkov. Phys. Rev. A, 78:053624, 2008.
- [47] P. A. Andreev. In: PIERS Proceedings, Moscow, p. 154, page 19, 2012.
- [48] P. A. Andreev. Int. J. Mod. Phys. B, 26:1250186, 2012.
- [49] M.I. Trukhanova and P. A. Andreev. Phys. Plasmas, 22:022128, 2015.
- [50] M. I. Trukhanova. rog. Theor. Exp. Phys, 2013:111101, 2013.
- [51] P. Holland. Phys. Rev. A, 60:4326, 1999.
- [52] C. Colijn and E.R. Vrscay. *Phys. Lett. A*, 300:334, 2002.
- [53] C. Colijn and E.R. Vrscay. J. Phys. A, 36:4689, 2003.
- [54] J. P. Finley. The Fluid Dynamics of the One-Body Stationary States of Quantum Mechanics with Real Valued Wavefunctions, the manuscript is available from http://arXiv.org/abs/2107.10315, 2021.
- [55] E. H. Lieb. Int. J. Quantum Chem., 24:243, 1983.
- [56] L. J. F. Broer. Physica, 76:364, 1974.
- [57] G. Salesi. Mod. Phys. Lett. A, 11:1815, 1996.
- [58] D. Hestenes. Found. Phys., 23:365, 1993.
- [59] W. Yang R. G. Parr. Density Functional Theory of Atoms and Molecules. Oxford University Press, Oxford, 1989.
- [60] J. Cioslowski. Many-Electron Densities and Reduced Density Matrices. Kluwer, New York, Boston, London, 2000.
- [61] R. M. Dreizler and E.K.U. Gross. *Density Functional Theory*. Springer-Verlag, Berlin, New York,

1990.

- [62] B. R. Munson, D. F. Young, and T. H. Okiishi. Fundamentals of Fluid Dynamics. Wiley, Hoboken, 5th edition, 2006.
- [63] B. H. Bransden and C. J. Joachain. Physics of Atoms and Molecules. Longman, London and New York, 1989.
- [64] A. H. Shapiro. The Dynamics and Thermodynamics of Compressible Fluid Flow, volume 1. Ronald, New York, 1953.
- [65] P. A. Kelly. Foundations of continuum mechanics. Foundations of Continuum Mechanics, Available from http://pkel015.connect.amazon.auckland.ac.nz, 2020.
- [66] A. Oldofredi. Latosensu, 5(1):77–85, 2018.
- [67] M. D.Towler. Proc. R. Soc. A, 468:990-1013, 2011.
- [68] J. P. Finley. Field Equations of Classical Mechanics for Quantum Mechanics, the manuscript is available from http://arXiv.org/abs/2207.04349, 2022.
- [69] J. P. Finley. Phys. Rev. A, 69:042514, 2004.
- [70] J. P. Finley. J. Mol. Phys., 102:627–639, 2004.