

Some Winnability Results for the Neighborhood and Group Labeling Lights Out Games

Brittany Doherty* Christian J. Miller† Darren B. Parker‡

MR Subject Classifications: 05C20, 06B99

Keywords: Lights out, light-switching game, winnability, graph game

Abstract

We look at both the *group labeling lights out game* and the *neighborhood lights out game*. Our main focus is to determine necessary and sufficient conditions for when the group labeling lights out game on path graphs, cycle graphs, and complete bipartite graphs can be won for every possible initial labeling. In the process of solving this problem, we demonstrate a new proof for when the neighborhood lights out game on complete bipartite graphs can be won for every possible initial labeling.

1 Introduction

The lights out game on graphs or directed graphs is an example of a light-switching game. In any light-switching game, there is a collection of lights that can be on, off, or perhaps have multiple on-states (which can be interpreted as different colors or different intensities of the same color). There is also a collection of switches, where each switch can change the states of one or more of the lights when toggled. The object of the game is usually to get all the lights into the “off” state. Examples of light-switching games are the Berlekamp (or Gale-Berlekamp) light-switching game (see [BM15], [CS04], and [Sch11]) and Merlin’s Magic Square (see [Pel87] and [Sto89]).

Lights out is a commonly studied light-switching game that was originally an electronic game created by Tiger Electronics in 1995. The idea behind this game has since been extended to several light-switching games on graphs. Some of these extensions are direct generalizations of the original game, like the σ^+ -game in [Sut89], the neighborhood lights out game developed independently in [Ara12] and [GP13], and a matrix-generated version of the game in [KP]. Other versions are explored in [Pel87], [Ara00], [CMP09]) and [DP].

*brittanycastle@gmail.com

†Department of Mathematics, Michigan State University, East Lansing, MI 48824, USA, mill3706@msu.edu,

‡Department of Mathematics, Grand Valley State University, Allendale, Michigan 49401-6495, parkerda@gvsu.edu

Each version of the game begins with some labeling of the vertices, usually by elements of \mathbb{Z}_m for some $m \geq 2$. The switches in this game are the vertices. The game is won when we achieve some desired labeling, usually where each vertex has label 0 (i.e. 0 is considered the “off” label). We call this labeling the *zero labeling*.

The most direct generalization of the original lights out game is called the *neighborhood m -lights out game*, where $m \geq 2$ is an integer. In this game, the labels come from \mathbb{Z}_m . When a vertex v is toggled, the labels of v and every vertex adjacent to v are increased by 1 modulo m . As noted above, the game is won when the zero labeling is achieved. This game is closely linked to the *neighborhood matrix* of G , which we denote by $N(G)$, or simply N if it is clear what the graph is. Thus, we also call the neighborhood m -lights out game the (N, m) -lights out game.

We also study a version of the lights out game defined in [PZ21], where the vertex labels come from a group H . The game begins with a graph G and an initial *labeling* of the vertices with elements of H , which we primarily express as a function $\lambda_0 : V(G) \rightarrow H$. The game is played by toggling vertices. Each time a vertex is toggled, it changes the labeling of G . If at any point in the game we have a labeling $\lambda : V(G) \rightarrow H$, then when a vertex v is toggled, this changes the labeling to λ' , where for each $w \in V(G)$, we have

$$\lambda'(w) = \begin{cases} \lambda(v) * \lambda(w), & w = v \text{ or } vw \in E(G) \\ \lambda(w), & \text{otherwise.} \end{cases}$$

The game is won when we achieve the *identity labeling* where the label of each vertex is the identity element of H . If H is abelian and the binary operation is addition, we also call this the *zero labeling* as in the neighborhood lights out game. We call this game the *H -labeling lights out game* (or, more briefly, the *H -labeling game*).

In both of the above games, if we begin the game with an unfortunate labeling, it may be impossible to win the game. If it is possible to win the game when we begin with the labeling λ_0 , we call λ_0 an (N, m) -winnable labeling in the neighborhood lights out game and an H -winnable labeling in the H -labeling lights out game. Depending on the game, we say that G is (N, m) -Always Winnable or H -Always Winnable (abbreviated (N, m) -AW or H -AW) if every labeling of $V(G)$ is (N, m) -winnable or H -winnable, respectively.

For the group labeling lights out game, we focus here on games where the group is a cyclic group \mathbb{Z}_m . Our main goal is to determine which path graphs, cycle graphs, and complete bipartite graphs are \mathbb{Z}_m -AW. In Section 2, we narrow down the values of m that can possibly support \mathbb{Z}_m -AW graphs. We also narrow down the groups that are necessary to consider when proving a graph is \mathbb{Z}_m -AW. In Section 3, we show how we can use the neighborhood lights out game to help us find necessary conditions for our graphs to be \mathbb{Z}_m -AW. Along the way, we give a new, simpler proof of (N, m) -AW complete bipartite graphs. Finally, in Section 4, we prove that the necessary conditions from Section 3 are sufficient as well.

2 Winnability with the Group \mathbb{Z}_m

We begin with a result from [PZ21] that narrows down the possible cyclic groups that lead to always winnable graphs.

Theorem 2.1. [PZ21, Thm. 2.4] If $m = 2^k d$, where d is odd, then there is a one-to-one correspondence between \mathbb{Z}_m -winnable labelings of G and \mathbb{Z}_{2^k} -winnable labelings of G .

This implies that if $d > 1$, then it is impossible for all labelings to be winnable. We thus get the following.

Corollary 2.2. If G is \mathbb{Z}_m -AW, then $m = 2^k$ for some $k \geq 1$.

Thus, for the remainder of the paper, we assume the vertex labels come from \mathbb{Z}_{2^k} for some $k \geq 1$.

For our next results, it is helpful to recall that every element of \mathbb{Z}_{2^k} is a congruence class (i.e. a set of integers that are congruent to one another modulo 2^k). Moreover, for each $S \in \mathbb{Z}_{2^k}$, since 2^k is even, either every element of S is odd or every element is even. Thus, it makes sense to say that S is even if every integer in S is even, and S is odd if every integer in S is odd. For $q \in \mathbb{Q}$, we define $qS = \{x \in \mathbb{Q} : x = qs \text{ for some } s \in S\}$. Note that if S is even, then $\frac{1}{2}S \in \mathbb{Z}_{2^{k-1}}$ (see [PZ21] for details).

Thinking of the labels this way helps us view a \mathbb{Z}_{2^k} -labeling lights out game as having a simultaneous \mathbb{Z}_2 -labeling lights out game. For a graph G and labeling $\lambda : V(G) \rightarrow \mathbb{Z}_{2^k}$, define $\lambda_2 : V(G) \rightarrow \mathbb{Z}_2$ by

$$\lambda_2(v) = \begin{cases} 1, & \lambda(v) \text{ is odd} \\ 0, & \lambda(v) \text{ is even} \end{cases}$$

Lemma 2.3. Let G be a graph, let $\lambda : V(G) \rightarrow \mathbb{Z}_{2^k}$ be a labeling, and let $\lambda_2 : V(G) \rightarrow \mathbb{Z}_2$ be as defined above.

1. If we toggle $v \in V(G)$ under the rules of the \mathbb{Z}_{2^k} -labeling game with labeling λ to get the labeling π , then toggling v under the rules of the \mathbb{Z}_2 -labeling game with labeling λ_2 results in the labeling π_2 .
2. Suppose λ_2 is \mathbb{Z}_2 -winnable. If we begin the \mathbb{Z}_{2^k} -labeling game with labeling λ , then it is possible to toggle the vertices of G so that every vertex has an even label.

Proof. For (1), we consider the parity of $\lambda(v)$. If $\lambda(v)$ is even, then when we toggle v in the \mathbb{Z}_{2^k} -labeling game, each label of an adjacent vertex is increased by an even number, and labels of vertices not adjacent to v are unchanged. Thus, the parities of all vertices are unchanged and so $\pi_2 = \lambda_2$. Also, $\lambda_2(v) = 0$, and so toggling v in the \mathbb{Z}_2 -labeling game leaves the labels of all vertices unchanged. Thus, the resulting labeling is λ_2 , which we determined above to be π_2 . In the case of $\lambda(v)$ odd, toggling v in the \mathbb{Z}_{2^k} -labeling game gives us π , which is the same as λ on all vertices not equal or adjacent to v . The label on v and each vertex adjacent to v changes parity, since we add the odd number $\lambda(v)$ to

each of these labels. In the \mathbb{Z}_2 -labeling game, $\lambda_2(v) = 1$, and so toggling v changes the parity of all adjacent vertices and leaves all other labels unchanged (just like with π). We end up with the labeling π_2 .

For (2), an easy induction using (1) gives us that for any sequence of toggles in the \mathbb{Z}_{2^k} -labeling game giving us a labeling π , the same toggles used in the corresponding \mathbb{Z}_2 -labeling game gives us the labeling π_2 . Now suppose we begin the \mathbb{Z}_{2^k} -labeling game with labeling λ . Since λ_2 is \mathbb{Z}_2 -winnable, if we begin with labeling λ_2 , we can toggle the vertices of G in the \mathbb{Z}_2 -labeling game to achieve the zero labeling. We now apply these same toggles in the \mathbb{Z}_{2^k} -labeling game. The resulting labeling π in the \mathbb{Z}_{2^k} -labeling game has π_2 as the zero labeling, which means that $\pi(v)$ is even for all $v \in V(G)$. This completes the proof. \square

In the case that the range of λ consists only of even labels, we get a similar relationship as above between the \mathbb{Z}_{2^k} -labeling game and the $\mathbb{Z}_{2^{k-1}}$ -labeling game.

Lemma 2.4. Let G be a graph, and let $k \geq 2$. Suppose $\lambda : V(G) \rightarrow \mathbb{Z}_{2^k}$ is a labeling where each vertex has an even label. Define $\pi : V(G) \rightarrow \mathbb{Z}_{2^{k-1}}$ by $\pi(w) = \frac{1}{2}\lambda(w)$ for all $w \in V(G)$. If we toggle the vertex v in the \mathbb{Z}_{2^k} -labeling game with labeling λ to get the labeling λ' , then toggling v in the $\mathbb{Z}_{2^{k-1}}$ -labeling game with labeling π results in the labeling π' , where $\pi'(w) = \frac{1}{2}\lambda'(w)$ for all $w \in V(G)$.

Proof. When we toggle v in the \mathbb{Z}_{2^k} -labeling game with labeling λ , each adjacent vertex w gets the label $\lambda'(w) = \lambda(v) + \lambda(w)$. Toggling v in the $\mathbb{Z}_{2^{k-1}}$ -labeling game with labeling π gives us $\pi'(w) = \pi(v) + \pi(w) = \frac{1}{2}\lambda(v) + \frac{1}{2}\lambda(w)$. It is easy to show that (as sets) $\frac{1}{2}\lambda(v) + \frac{1}{2}\lambda(w) = \frac{1}{2}(\lambda(v) + \lambda(w)) = \frac{1}{2}\lambda'(w)$. Thus, $\pi'(w) = \frac{1}{2}\lambda'(w)$. Since vertices not adjacent to v are unchanged in both games, we get $\pi'(w) = \frac{1}{2}\lambda'(w)$ for them as well. \square

We use Lemma 2.3 and Lemma 2.4 to prove a result that allows us to assume our vertex labelings come from \mathbb{Z}_2 .

Theorem 2.5. Let G be a graph and let $k \geq 1$. Then G is \mathbb{Z}_{2^k} -AW if and only if G is \mathbb{Z}_2 -AW.

Proof. First we assume G is \mathbb{Z}_{2^k} -AW. To prove G is \mathbb{Z}_2 -AW, we let $\lambda : V(G) \rightarrow \mathbb{Z}_2$ be a labeling and prove λ is \mathbb{Z}_2 -winnable. We define $\pi : V(G) \rightarrow \mathbb{Z}_{2^k}$ by $\pi(v) = \lambda(v) \bmod 2^k$. It is clear from the definition that $\pi_2 = \lambda$. Since G is \mathbb{Z}_{2^k} -AW, π is \mathbb{Z}_{2^k} -winnable. Thus, we can toggle the vertices in the \mathbb{Z}_{2^k} -labeling game to get the zero labeling, which we will call π' . By Lemma 2.3(1), if we do this same toggling in the \mathbb{Z}_2 -labeling game with initial labeling $\lambda = \pi_2$, we end up with the labeling π'_2 . Since π' is the zero labeling, all labels are even, and so π'_2 is the zero labeling as well. This proves that λ is \mathbb{Z}_2 -winnable, and so G is \mathbb{Z}_2 -AW.

For the other direction, assume G is \mathbb{Z}_2 -AW and let $\lambda : V(G) \rightarrow \mathbb{Z}_{2^k}$. We prove λ is \mathbb{Z}_{2^k} -winnable by induction. The case $k = 1$ follows from the assumption that G is \mathbb{Z}_2 -AW. For $k > 1$, by Lemma 2.3(2), since G is \mathbb{Z}_2 -AW, we can toggle the vertices in the \mathbb{Z}_{2^k} -lights out game so that all vertices have even labels. Thus, we can assume $\lambda(v)$ is

even for all $v \in V(G)$. Since all labels of λ are even, we can define $\pi : V(G) \rightarrow \mathbb{Z}_{2^{k-1}}$ by $\pi(w) = \frac{1}{2}\lambda(w)$.

By the induction hypothesis, π is $\mathbb{Z}_{2^{k-1}}$ -winnable, so we can toggle the vertices in the $\mathbb{Z}_{2^{k-1}}$ -labeling game so that we obtain the zero labeling π_0 . If we toggle the vertices identically in the \mathbb{Z}_{2^k} -labeling game, an easy induction using Lemma 2.4 implies that this results in a labeling λ' such that $\pi_0(w) = \frac{1}{2}\lambda'(w)$ for all $w \in V(G)$. Since $\pi_0(w) = 0$ for all $w \in V(G)$, it follows that $\lambda'(w) = 0$ as well, making λ' the zero labeling. Thus, λ is \mathbb{Z}_{2^k} -winnable. Since λ is arbitrary, that makes G \mathbb{Z}_{2^k} -AW, proving the theorem. \square

3 Neighborhood Lights Out Game and \mathbb{Z}_{2^k} -Winnability

Our main results for the \mathbb{Z}_{2^k} -labeling game will be characterizations of \mathbb{Z}_{2^k} -AW path graphs, cycle graphs, and complete bipartite graphs. This requires us to prove our conditions on the graphs are both necessary and sufficient for the graphs to be \mathbb{Z}_{2^k} -AW. We have already simplified our task a great deal. Theorem 2.5 implies that we need only find necessary and sufficient conditions for our graphs to be \mathbb{Z}_2 -AW. In this section, we use the neighborhood lights out game to help us find necessary conditions for winnability.

The neighborhood lights out game is advantageous to our study of the group labeling game for two reasons. First, the $(N, 2)$ -lights out game plays very similarly to the \mathbb{Z}_2 -labeling game. In the \mathbb{Z}_2 -labeling game, when we toggle a vertex whose label is 1, that changes its own label to 0 and the label of each adjacent vertex from 0 to 1 or from 1 to 0. This is identical to the $(N, 2)$ -lights out game. Also, if a vertex has label 0, then toggling it in the \mathbb{Z}_2 -labeling game has no effect on the labeling of the graph. Thus, the \mathbb{Z}_2 -labeling game is equivalent to playing the $(N, 2)$ -lights out game where we only toggle vertices that have label 1. This means that if we want even a chance to win the \mathbb{Z}_2 -labeling game, we have to be able to win the $(N, 2)$ -lights out game as well. This gives us the following.

Lemma 3.1. If a graph G is \mathbb{Z}_2 -AW, then G is $(N, 2)$ -AW.

The second advantage of the $(N, 2)$ -lights out game is that winnability is generally easier to determine than in the \mathbb{Z}_2 -labeling game. We illustrate this on a general result for complete bipartite graphs. We should note that (N, m) -AW complete bipartite graphs were characterized in [GP13]. However, here we give a much more elegant proof.

Recall that a complete bipartite graph is a graph whose vertices can be partitioned into two sets, P_1 and P_2 , such that $E(G) = \{vw : v \in P_1 \text{ and } w \in P_2\}$. Before proving our main result, we must prove the following.

Lemma 3.2. Let $k, n, p \geq 1$ and $m \geq 2$.

1. Suppose we are playing either the (N, m) -lights out game or the \mathbb{Z}_{2^k} -labeling game on $K_{n,p}$. If P_1 and P_2 are the parts from the partition of $V(K_{n,p})$ (with $|P_1| = n$ and $|P_2| = p$), then for any initial labeling of $V(G)$, we can toggle the vertices in such a way that all vertices in P_1 have label 0 and all vertices in P_2 have the same label.

2. Suppose that an (N, m) -winnable labeling has all vertices in P_1 (respectively P_2) having the same label. In a winning toggling, each vertex in P_1 (resp. P_2) is toggled the same number of times.

Proof. For (1), in both games we obtain the desired labeling by first toggling each vertex in P_2 until it has label 0. This can be done in the neighborhood game because each toggle increases the label of the toggled vertex by 1 modulo m , so we get to label 0 within $k - 1$ toggles. This can be done in the \mathbb{Z}_{2^k} -labeling game since each toggle doubles the label of the toggled vertex, so we get to label 0 within k toggles. Since no two vertices in P_2 are adjacent, this results in all vertices in P_2 having label 0. Then we toggle each vertex in P_1 until it has label 0. As before, this results in each vertex in P_1 having label 0. Also, since every vertex in P_1 is adjacent to every vertex in P_2 , every vertex in P_2 will have the same label, since each toggle of a vertex in P_1 increases every vertex in P_2 by the same number.

For (2), assume that all vertices in P_1 have the same label. For contradiction, assume two vertices $v, w \in P_1$ are toggled a different number of times for a winning toggling. Once this is done, v and w have different labels. This will still hold when all vertices in P_1 are toggled, since none of them are adjacent to v or w . Each time we toggle a vertex in P_2 , this increases the labels of v and w by 1 modulo m . Thus, once all vertices in P_2 are toggled, v and w still have different labels. But now all vertices have been toggled, and it is impossible for both v and w to have label 0. This contradicts our assumption of having a winning toggling, and so v and w must be toggled the same number of times. The proof for all vertices of P_2 having the same label is similar. \square

We are now ready to characterize (N, m) -AW complete bipartite graphs.

Theorem 3.3. The graph $K_{n,p}$ is (N, m) -AW if and only if $\gcd(m, np - 1) = 1$.

Proof. We begin by assuming $K_{n,p}$ (with parts P_1 and P_2) is (N, m) -AW and proving that $\gcd(m, np - 1) = 1$. Let $\lambda : V(G) \rightarrow \mathbb{Z}_m$ be any labeling. By Lemma 3.2(1), we can assume that there is some $a \in \mathbb{Z}_m$ such that $\lambda(v) = 0$ for all $v \in P_1$ and $\lambda(v) = a$ for all $v \in P_2$. By Lemma 3.2(2) and the fact that $K_{n,p}$ is (N, m) -winnable, there exist $x, y \in \mathbb{Z}$ such that toggling every vertex in P_1 x times and toggling every vertex in P_2 y times results in the zero labeling. Considering each vertex individually, each vertex in P_1 gets its label increased by each vertex in P_2 by y . Each such vertex is also increased by its own toggling, which increases its label by x . Since there are p vertices in P_2 , and since each vertex in P_1 begins with label 0, the terminal label for each vertex in P_1 is $x + py$. Similarly, the terminal label for each vertex in P_2 is $nx + y + a$. Note that the a appears in this last expression because the initial label of each vertex of P_2 is a .

Putting together the expressions we get for the terminal labels of the vertices, along with the fact that each terminal label is 0, we get the equations $x + py = 0$ and $nx + y + a = 0$. Since λ is (N, m) -winnable, this system of equations must always have a solution. Conversely, if a solution of this system of equations exists, then if each vertex of P_1 is toggled x times and each vertex of P_2 is toggled y times, this is a winning toggling for the game. Thus, $K_{n,p}$ is (N, m) -AW if and only if the system $x + py = 0$ and $nx + y + a = 0$

has a solution. By solving the system directly or computing the determinant of $\begin{bmatrix} 1 & p \\ n & 1 \end{bmatrix}$, a solution always exists if and only if $np - 1$ is a unit in \mathbb{Z}_m . This occurs precisely when $\gcd(m, np - 1) = 1$. \square

If we let $m = 2^k$, the condition $\gcd(2^k, np - 1) = 1$ is equivalent to one or both of n and p being even. Putting this together with Lemma 3.1 and Lemma 2.5, we get the following.

Corollary 3.4. If $K_{m,n}$ is \mathbb{Z}_{2^k} -AW, then one or both of m and n is even.

In [GP13], winnability in the neighborhood lights out game was also determined for paths and cycles for all (N, m) -lights out games. If we state these results in the case that $m = 2$, we get that P_n is $(N, 2)$ -AW precisely when $n \equiv 0$ or $1 \pmod{3}$ and C_n is $(N, 2)$ -AW precisely when $n \equiv 1$ or $2 \pmod{3}$. Putting these results together with Lemma 3.1 gives us the following.

Lemma 3.5. Let $n, k \in \mathbb{N}$.

1. If $n \geq 1$ and P_n is \mathbb{Z}_{2^k} -AW, then $n \equiv 0$ or $1 \pmod{3}$.
2. If $n \geq 3$ and C_n is \mathbb{Z}_{2^k} -AW, then $n \equiv 1$ or $2 \pmod{3}$.

4 Sufficient Conditions for \mathbb{Z}_{2^k} -Winnability

Corollary 3.4 and Lemma 3.5 give necessary conditions for P_n , C_n , and $K_{n,p}$ to be \mathbb{Z}_{2^k} -AW. In this section, we show that these conditions are sufficient as well. The notation we use for P_n and C_n is $V(P_n) = V(C_n) = \{v_1, v_2, \dots, v_n\}$, $E(P_n) = \{v_i v_{i+1} : 1 \leq i \leq n-1\}$, and $E(C_n) = E(P_n) \cup \{v_1 v_n\}$.

We prove a given condition is sufficient for a graph to be \mathbb{Z}_2 -AW by proving each possible labeling is \mathbb{Z}_2 -winnable. The first step in doing this is to prove we can convert each labeling into a labeling that is easy to work with, much like Lemma 3.2(1). We do this in the following lemma.

Lemma 4.1. Let $n \in \mathbb{N}$, and suppose we are playing the \mathbb{Z}_2 -labeling game on P_n or C_n .

1. For any initial labeling of P_n , we can toggle the vertices to achieve a labeling λ with $\lambda(v_i) = 0$ for all $i \geq 2$.
2. If $n \geq 3$, then for any initial labeling of C_n , we can toggle the vertices to achieve a labeling λ with $\lambda(v_i) = 0$ for all $i \geq 3$.

Proof. For (1), let π be a labeling on P_n . We prove our result by induction on the maximum number k such that $\pi(v_k) = 1$ (i.e. the “last” vertex to have a nonzero label). If $k = 1$, then $\pi(v_i) = 0$ for all $i \geq 2$, and so π is the desired labeling. For $k > 1$, we have two cases. If $k = n$, then we toggle $v_k = v_n$. This causes v_k to have label 0, which means

the maximum i with v_i having a nonzero label is at most $k - 1$. By induction, we can toggle the vertices to achieve labeling λ with $\lambda(v_i) = 0$ for all $j \geq 2$. In the case $k < n$, we consider the subcases $\pi(v_{k-1}) = 1$ and $\pi(v_{k-1}) = 0$. If $\pi(v_{k-1}) = 1$, we toggle v_{k-1} . This changes the labels of v_{k-1} and v_k to 0, which makes the maximum i with v_i having a nonzero label at most $k - 2$. We then apply induction as in the $k = n$ case. In the case $\pi(v_{k-1}) = 0$, we first toggle v_k . This changes the labels of v_k to 0 and of v_{k-1} and v_{k+1} to 1. We then toggle v_{k-1} . This changes the labels of v_{k-1} to 0 and v_k to 1. Finally, we toggle v_k again, which changes the labels of v_k and v_{k+1} to 0, and v_{k-1} to 1. This gives us a labeling where the maximum i with v_i having a nonzero label is $k - 1$. We then apply induction as in the previous cases, and the lemma is proved. The proof for (2) is almost identical. The only change is that the base cases are $k = 0$ and $k = 1$. \square

Now we look to prove the conditions sufficient for the \mathbb{Z}_2 -labeling game. We begin with path graphs.

Lemma 4.2. Let $n \in \mathbb{N}$.

1. If $n \geq 1$ and $n \equiv 0$ or $1 \pmod{3}$, then P_n is \mathbb{Z}_2 -AW.
2. If $n \geq 3$ and $n \equiv 1$ or $2 \pmod{3}$, then C_n is \mathbb{Z}_2 -AW.

Proof. For (1), we assume λ is a labeling of P_n and prove λ is \mathbb{Z}_2 -winnable. By Lemma 4.1(1), we can assume $\lambda(v_i) = 0$ for $i \geq 2$. If $\lambda(v_1) = 0$, we have the zero labeling, making λ \mathbb{Z}_2 -winnable, so we can assume $\lambda(v_1) = 1$. We begin our toggling strategy by toggling each v_i in order from v_1 to v_n . An easy induction implies that for all $1 \leq i \leq n - 1$, when we toggle v_i , every v_j with $j < i$ has label 1, v_{i+1} has label 1, and all other vertices have label 0. When we toggle v_n all vertices have label 1, except v_n , which has label 0.

Now we consider the cases $n \equiv 0 \pmod{3}$ and $n \equiv 1 \pmod{3}$ separately. Suppose $n \equiv 0 \pmod{3}$. For each $0 \leq s \leq r - 1$, we then toggle v_{3s+1} . For $s = 0$, this changes the labels of v_1 and v_2 to 0. For each $1 \leq s \leq r - 1$, this changes the labels of v_{3s} , v_{3s+1} , and v_{3s+2} to 0. This results in the zero labeling, which makes λ \mathbb{Z}_2 -winnable.

If $n \equiv 1 \pmod{3}$, then we toggle each v_{3s+2} for all $0 \leq s \leq r - 1$. This changes the labels of each v_{3s+1} , v_{3s+2} , and v_{3s+3} to 0, which again results in the zero labeling. Thus, λ is \mathbb{Z}_2 -winnable, and so P_n is \mathbb{Z}_2 -AW.

The proof for (2) is similar. We assume λ is a labeling of C_n and prove λ is \mathbb{Z}_2 -winnable. By Lemma 4.1(2), we can assume $\lambda(v_i) = 0$ for all $i \geq 3$. If $\lambda(v_1) = \lambda(v_2) = 0$, then λ is the zero labeling and is thus \mathbb{Z}_2 -winnable. If $\lambda(v_1) = \lambda(v_2) = 1$, then we can toggle v_2 to get $\lambda(v_i) = 0$ for all $i \neq 3$. By relabeling the vertices starting at v_3 , we can thus assume $\lambda(v_1) = 1$ and $\lambda(v_i) = 0$ for all $i \neq 1$.

In the case $n \equiv 1 \pmod{3}$, we have $n = 3r + 1$ for some $r \geq 1$. Similarly as the proof for P_n , we toggle each v_i in order from v_1 to $v_{n-2} = v_{3r-1}$. As in the proof for P_n , for each $i < n - 2$ and $i = n - 1$, v_i has label 1. Since v_n is adjacent to v_1 , then v_n also has label 1. Thus, v_i has label 1 for all $i \neq n - 2$ and v_{n-2} has label 0. We toggle each v_{3s} for $1 \leq s \leq r$. Using similar reasoning as in the proof above for P_n , $n \equiv 1 \pmod{3}$, this results in the zero labeling.

In the case $n \equiv 2 \pmod{3}$, we have $n = 3r + 2$ for some $r \geq 1$. We toggle each v_i in order from v_1 to $v_{n-1} = v_{3r+1}$. Using similar reasoning as before, every vertex has label 1 except v_{n-1} and v_n , which both have label 0. By toggling all v_{3s+2} for $0 \leq s \leq r - 1$, we get the zero labeling. Thus, λ is \mathbb{Z}_2 -winnable, and so C_n is \mathbb{Z}_2 -AW. \square

Our next result is for complete bipartite graphs.

Lemma 4.3. If $n, p \in \mathbb{N}$ and one or both of n and p is even, then $K_{n,p}$ is \mathbb{Z}_2 -AW.

Proof. Let P_1 and P_2 be the parts of $K_{n,p}$ with $|P_1| = n$ and $|P_2| = p$. We assume λ is a labeling of $K_{n,p}$ and prove that λ is \mathbb{Z}_2 -winnable. By Lemma 3.2(1), we can assume that all vertices in P_1 have label 0 and all vertices in P_2 have the same label. If the vertices of P_2 have label 0, that makes λ the zero labeling, which is obviously \mathbb{Z}_2 -winnable, so we can assume each vertex in P_2 has label 1. Without loss of generality, we can assume p is even (if not, we can toggle each vertex in P_2 to make every vertex in P_2 have label 0 and every vertex in P_1 have label 1).

We then achieve the zero labeling by toggling each vertex in P_2 once. Each toggling increases the labels of each vertex in P_1 by 1 mod 2. Since there is an even number of vertices in P_2 , each vertex in P_1 has its label increased by 0 mod 2. Thus, each vertex in P_1 ends up with label 0. Since each vertex in P_2 changes its own label to 0, this results in the zero labeling. Thus, λ is \mathbb{Z}_2 -winnable, which makes $K_{n,p}$ \mathbb{Z}_2 -AW. \square

We can now state our main result, which follows directly from Theorem 2.5, Lemma 3.5, Lemma 4.2, and Lemma 4.3.

Theorem 4.4. Let $k, n, p \in \mathbb{N}$.

1. If $n \geq 1$, then P_n is \mathbb{Z}_{2^k} -AW if and only if $n \equiv 0$ or $1 \pmod{3}$.
2. If $n \geq 3$, then C_n is \mathbb{Z}_{2^k} -AW if and only if $n \equiv 1$ or $2 \pmod{3}$.
3. If $n, p \geq 1$, then $K_{n,p}$ is \mathbb{Z}_{2^k} -AW if and only if one or both of n and p is even.

5 Open Problems

We close with three possible directions for further research.

- When we set out to determine whether or not our graphs were \mathbb{Z}_{2^k} -AW, we used a two-step process. We first looked at the $(N, 2)$ -lights out game to find necessary conditions for a graph to be \mathbb{Z}_{2^k} -AW. Then we proved that these conditions are sufficient as well. However, this would not work if we had a graph whose winnability in the \mathbb{Z}_2 -labeling game was different than in the $(N, 2)$ -lights out game. Are there any graphs that are $(N, 2)$ -AW but not \mathbb{Z}_2 -AW?

- If the answer to the above question is yes, that complicates our determination of necessary conditions for a graph to be \mathbb{Z}_{2^k} -AW. What makes studying the neighborhood lights out game so nice is that winning or losing the game depends only on how many times we toggle each vertex, not the order in which we toggle the vertices. Along with some other nice properties of the game, this makes it possible to determine winnability using systems of linear equations. In the H -labeling lights out game, this is not the case (see [PZ21] for a detailed discussion). If we cannot depend on the neighborhood lights out game to help us prove group labeling lights out theorems, what other techniques can we use?
- While there seems to be much to learn from the group labeling games with cyclic groups, it would be nice to see how the game works with non-cyclic groups. By the same reasoning that led us to restricting our attention to \mathbb{Z}_{2^k} , we can only have H -winnable graphs when $|H| = 2^k$ for some $k \in \mathbb{N}$. But there are no other obvious restrictions. Non-abelian groups are especially tempting to look at.

References

- [Ara00] P.V. Araujo, *How to turn all the lights out*, Elem. Math. **55** (2000), 135–141.
- [Ara12] C. Arangala, *The $4 \times n$ multistate lights out game*, Math. Sci. Int. Res. J. **1** (2012), 10–13.
- [BM15] R.A. Brualdi and S.A. Meyer, *A Gale-Berlekamp permutation switching problem*, European J. Combin. **44** (2015), 43–56.
- [CMP09] D. Craft, Z. Miller, and D. Pritikin, *A solitaire game played on 2-colored graphs*, Discrete Math. **309** (2009), 188–201.
- [CS04] J. Carlson and D. Stolarski, *The correct solution to Berlekamp’s switching game*, Discrete Math. **287** (2004), 145–150.
- [DP] T.E. Dettling and D.B. Parker, *The Lights Out Game on Directed Graphs*, Submitted to Involve.
- [GP13] A. Giffen and D.B. Parker, *On Generalizing the “Lights Out” Game and a Generalization of Parity Domination*, Ars Combin. **111** (2013), 273–288.
- [KP] L. Keough and D.B. Parker, *An Extremal Problem for the Neighborhood Lights Out Game*, To appear in Discuss. Math. Graph Theory.
- [Pel87] D. Pelletier, *Merlin’s Magic Square*, Amer. Math. Monthly **94** (1987), 143–150.
- [PZ21] D.B. Parker and V. Zadorozhnyy, *A Group Labeling Version of the Lights Out Game*, Involve **14** (2021), 541–554.

- [Sch11] U. Schauz, *Colorings and Nowhere-Zero Flows of Graphs in Terms of Berlekamp's Switching Game*, *Electronic J. Combin.* **18** (2011), #P65.
- [Sto89] D.L. Stock, *Merlin's Magic Square Revisited*, *Amer. Math. Monthly* **96** (1989), 608–610.
- [Sut89] K. Sutner, *The σ -game and cellular automata*, *Amer. Math. Monthly* **97** (1989), 24–34.