

STABILITY IN THE BANACH ISOMETRIC CONJECTURE AND NEARLY MONOCHROMATIC FINSLER SURFACES

GAUTAM AISHWARYA AND DMITRY FAIFMAN

ABSTRACT. The Banach isometric conjecture asserts that a normed space with all of its k -dimensional subspaces isometric, where $k \geq 2$, is Euclidean. The first case of $k = 2$ is classical, established by Auerbach, Mazur and Ulam using an elegant topological argument. We refine their method to arrive at a stable version of their result: if all 2-dimensional subspaces are almost isometric, then the space is almost Euclidean. Furthermore, we show that a 2-dimensional surface, which is not a torus or a Klein bottle, equipped with a near-monochromatic Finsler metric, is approximately Riemannian. The stability is quantified explicitly using the Banach–Mazur distance.

1. INTRODUCTION AND RESULTS

In 1932, Banach made the following conjecture [2].

Conjecture 1.1. *Let V be a normed space over the reals, and $k \geq 2$ an integer smaller than $n = \dim(V)$. Suppose all linear k -dimensional subspaces of V are isometric to each other. Then, V must necessarily be a Hilbert space.*

A seemingly more general conjecture, known to be equivalent to Conjecture 1.1 [18], is the following.

Conjecture 1.2. *Let V be a normed space over the reals, and $2 \leq k < \dim V$. If $K \subset V$ is a convex body, and all its k -dimensional sections through an interior point of K are affinely equivalent, then K is an ellipsoid.*

Here two subsets of two respective affine spaces are *affinely equivalent* if one is the image of the other under an affine map.

This conjecture has been proven in various cases, starting with the work of Auerbach–Mazur–Ulam in 1935 [1], who proved it for $k = 2$. Their proof is based on a topological obstruction - an idea later pushed further by Gromov [8] to settle the conjecture for even k , as well as for $\dim V \geq k + 2$. Recently, Bor–Hernandez Lamonedá–Jimenez Desantiago–Montejano Peimbert [3] combined topological and convex-geometric ideas to prove the conjecture for $k = 4m + 1$, with the possible exception of $n = 133$; while for $k = 3$, the conjecture has been confirmed by Ivanov, Mamaev, and Nordskova [11]. A local version of Conjecture 1.1 for $k = 2, 3$ was established in [10, 12]. Some results were also obtained for the Banach conjecture in complex normed spaces [8, 4].

MSC classification: 52A21, 46C15, 57R15.

GA was supported by ISF grant 1468/19 and NSF-BSF grant DMS-2247834. DF was supported by ISF grant No. 1750/20.

1.1. Results. In this note, we initiate the study of stability in Banach’s conjecture. Stability results in convex geometry concern themselves either with geometric inequalities, showing near-extremizers to be close to the extremizers; or with the geometric characterization of a certain type of convex body, showing e.g. that a convex body approximately satisfying a property exhibited only by ellipsoids, must be close to an ellipsoid. This work studies the latter type of stability. Previous stability results of this type appeared in several works, such as Burger and Schneider [5], Groemer [7], Gruber [9] (see also references therein).

Loosely speaking, we wish to know if, when the linear k -dimensional sections of a convex body K with 0 in its interior are approximately linearly/affinely equivalent, K must be approximately an ellipsoid. To quantify the deviation from linear/affine equivalence, we use the Banach–Mazur distance.

Definition 1.3. *Let V, W be isomorphic Banach spaces (with possibly non-symmetric norms). The Banach–Mazur distance between them is defined by*

$$d_{BM}(V, W) = \inf\{\|T\|\|T^{-1}\| : T \in \text{GL}(V, W)\}.$$

One can identify (non-symmetric) normed spaces with their unit balls, which are convex bodies with the origin a fixed point in their interior. The Banach–Mazur distance yields a corresponding notion of distance on such pointed bodies.

Definition 1.4. *Let $K \subset V, L \subset W$ be convex bodies with the origin in their interior. The linear Banach–Mazur distance is*

$$d_{BM}^{\text{lin}}(K, L) = \inf\{\lambda > 0 : K \subset TL \subset \lambda K, \text{ for some } T \in \text{GL}(W, V)\}.$$

When affine maps are considered, we need to allow additional freedom in the choice of origin. Denote by $\overline{\text{GL}}(W, V)$ the set of affine maps from W to V .

Definition 1.5. *Let $K \subset V, L \subset W$ be convex bodies with non-empty interior.*

$$d_{BM}(K, L) = \inf\{\lambda > 0 : K \subset TL \subset \lambda K + z, \text{ for some } T \in \overline{\text{GL}}(W, V), z \in V\}.$$

Note that if the convex bodies $K, L \subset \mathbb{R}^n$ are the unit balls of norms $\|\bullet\|_K$, resp $\|\bullet\|_L$ on \mathbb{R}^n , we have $d_{BM}(K, L) \leq d_{BM}^{\text{lin}}(K, L) = d_{BM}(\|\bullet\|_K, \|\bullet\|_L)$ in general. Equality holds e.g. when both K, L are centrally symmetric.

The Banach–Mazur distance is usually left in multiplicative form. Thus K, L are affinely equivalent if and only if $d_{BM}(K, L) = 1$, and $d_{BM}(K, L) \geq 1$ in general.

We can now formulate a quantitative version of the Banach conjecture. Denote by \mathbb{E}^n the Euclidean n -dimensional space, and by B^n the unit ball therein.

Conjecture 1.6. *For any $\epsilon > 0$ and $n > k \geq 2$ there is $\delta = \delta(n, k, \epsilon) > 0$ such that the following holds. For any n -dimensional normed space V , possibly non-symmetric, such that $d_{BM}(E, F) < 1 + \delta$ for all k -dimensional linear subspaces $E, F \subset V$, it must hold that $d_{BM}(V, \mathbb{E}^n) < 1 + \epsilon$.*

As before, this is the linear case of the general affine conjecture.

Conjecture 1.7. *For any $\epsilon > 0$ and $n > k \geq 2$ there is $\delta = \delta(n, k, \epsilon) > 0$ such that the following holds. For any convex body $K \subset \mathbb{R}^n$ with 0 in its interior, such that $d_{BM}(K \cap E, K \cap F) < 1 + \delta$ for all k -dimensional linear subspaces $E, F \subset \mathbb{R}^n$, we must have $d_{BM}(K, B^n) < 1 + \epsilon$.*

We prove Conjecture 1.7 for $k = 2$ with an explicit stability estimate. In the following, $c > 0$ denotes various universal constants that can be computed explicitly.

Theorem 1.8. *Let $K \subset \mathbb{R}^n$ be a convex body with 0 in its interior. Assume that $d_{BM}(K \cap E, K \cap F) < 1 + \delta$ for all linear 2-dimensional subspaces $E, F \subset \mathbb{R}^n$. Then $d_{BM}(K, B^n) < 1 + cn^{2n^2}\delta^{1/6}$.*

For Conjecture 1.6 with $k = 2$ we obtain better stability.

Theorem 1.9. *Let $V = \mathbb{R}^n$ be a normed space, not necessarily symmetric. Assume that $d_{BM}(E, F) < 1 + \delta$ for all 2-dimensional subspaces $E, F \subset V$. Then $d_{BM}(V, \mathbb{E}^n) < 1 + cn^2\delta^{1/3}$.*

In the special case of symmetric norms this can be stated as follows.

Corollary 1.10. *Let $K \subset \mathbb{R}^n$ be a centrally-symmetric convex body. Assume that $d_{BM}(K \cap E, K \cap F) < 1 + \delta$ for all linear 2-dimensional subspaces $E, F \subset \mathbb{R}^n$. Then $d_{BM}(K, B^n) < 1 + cn^2\delta^{1/3}$.*

We deduce these results as a corollary of the following general theorem on fields of convex bodies. A *continuous field of convex bodies* over a manifold Σ is a family of convex bodies with non-empty interior $K_x \subset T_x\Sigma$, $x \in \Sigma$, continuous with respect to the Hausdorff distance within each chart.

Theorem 1.11. *Let Σ be a closed surface which is not a torus or a Klein bottle. If a continuous field of convex bodies $(K_x)_{x \in \Sigma}$ satisfies $d_{BM}(K_x, K_y) < 1 + \delta$ for all $x, y \in \Sigma$, then for all $x \in \Sigma$, $d_{BM}(K_x, B^2) < 1 + c\delta^{1/3}$.*

Thus a near-monochromatic Finsler metric on Σ is approximately Riemannian:

Corollary 1.12. *Let Σ be as in Theorem 1.11, and assume it is equipped with a C^0 Finsler structure such that $d_{BM}(T_x\Sigma, T_y\Sigma) < 1 + \delta$ for all $x, y \in \Sigma$. Then $d_{BM}(T_x\Sigma, \mathbb{E}^2) < 1 + c\delta^{1/3}$ for all $x \in \Sigma$.*

The restriction on the topology of Σ is necessary as one can easily construct monochromatic Finsler structures on the torus and the Klein bottle which are not Riemannian. It will be seen from the proof that the key required property is the non-existence of a field of tangent lines on the surface. We conjecture that Theorem 1.11, with some stability bounds, holds in greater generality, in particular for even dimensional spheres.

1.2. Ideas and plan of the proof. The proof is inspired by the proof of Auerbach–Mazur–Ulam of Conjecture 1.7 for $k = 2$, $\dim V = 3$, which we now recall using modern language. Letting $K_x = K \cap x^\perp$, we get a continuous field of convex bodies in the tangent planes of S^2 , $x \mapsto K_x \subset T_x S^2$. A convex body in \mathbb{R}^2 is either an ellipse, or has a finite group of affine symmetries. Fixing $x_0 \in S^2$ and assuming that none of K_x are ellipses, the affine maps $\text{Aff}(K_{x_0}, K_x)$ mapping K_{x_0} to K_x define a covering space over S^2 , which must have a section, $x \mapsto g_x$, as S^2 is simply connected. Fixing a line $L \subset T_{x_0} S^2$, and letting L_x be the linear line in $T_x S^2$ parallel to $g_x(L)$, we obtain a field $x \mapsto L_x$ of tangent lines on S^2 , which is impossible. Thus all sections K_x are ellipses. A separate geometric argument, showing that a convex body all of whose 2-dimensional sections through an interior point are ellipses must be an ellipsoid, completes the proof.

As the proof of Auerbach–Mazur–Ulam, as do most other known cases of the Banach conjecture, has a topological obstruction at its heart, it is somewhat surprising that a stability result for $k = 2$ can be obtained by following the scheme of proof outlined above.

The proof of Theorem 1.11 appears in section 3. We make use of an associated Riemannian structure on Σ given by the Binet–Legendre ellipsoids $\mathcal{E}(K_x)$. We consider sets of approximate isometries $G_\alpha(K_x, K_y)$, consisting of affine maps $g : T_x\Sigma \rightarrow T_y\Sigma$ that preserve the Binet–Legendre ellipsoids, and map K_x to K_y up to an error α with respect to the Hausdorff distance. We observe that if the elements of $G_\alpha(K_x)$ form a fine net in the special orthogonal group, then K_x must be close to an ellipse. Provided this is not the case, the key step is then to find a range of values α for which $G_\alpha(K_{x_0})$, or more precisely a low-resolution version thereof, is fixed. This then allows to define a covering space consisting of approximate isometries in $G_{\alpha'}(K_{x_0}, K_x)$, and proceed as before.

The corollaries of Theorem 1.11 to stability in the Banach conjecture are proved in section 4. To pass from Theorem 1.11 to Theorem 1.8, we prove a stable version of independent interest of the characterization of ellipsoids through planar sections mentioned above. For Theorem 1.9, we get better stability by utilizing instead a stable version of the von Neumann–Jordan theorem due to Passer.

1.3. Acknowledgements. The authors are thankful to Juan Carlos Alvarez Paiva for his inspiring talk at the Integral and Metric Geometry 2022 BIRS-CMO workshop. The second-named author is grateful to Vitali D. Milman for introducing him to the Banach conjecture, and to Károly J. Böröczky for a helpful discussion.

2. PRELIMINARIES

For an ellipsoid \mathcal{E} , we denote by $d_{\mathcal{E}}$ the Hausdorff distance with respect to the Euclidean metric for which \mathcal{E} is a unit ball. Denote by $\text{cent}(K)$ the centroid (center of mass) of K . We say K is *centered* if $\text{cent}(K) = 0$. A basic property of centered convex bodies in \mathbb{R}^n , which we will use repeatedly, is that $-K \subset nK$ [20, Lemma 2.3.3]. For a centered convex body K , the *Binet–Legendre ellipsoid* $\mathcal{E}(K) \subset \mathbb{R}^n$ of $K \subset \mathbb{R}^n$, also called the ellipsoid of inertia, is the dual ellipsoid of the unit ball in $(\mathbb{R}^n)^*$ of the inner product

$$\langle \phi, \psi \rangle_K = \frac{n+2}{\text{vol}(K)} \int_K \phi(x)\psi(x) d\text{vol}(x),$$

where vol is an arbitrary Lebesgue measure on V . For general K , we set $\mathcal{E}(K) := \mathcal{E}(K - \text{cent}(K)) + \text{cent}(K)$.

Evidently, if $g \in \overline{\text{GL}}(\mathbb{R}^n)$ then $\mathcal{E}(gK) = g\mathcal{E}(K)$. If K is an ellipsoid, $\mathcal{E}(K) = K$. Note that unlike the definition in [15], which uses a fixed origin, we use the centroid of K as the center of the Binet–Legendre ellipsoid.

Lemma 2.1. *Let $K, L \subset \mathbb{R}^n$ be centered convex bodies. If $\frac{1}{\lambda_1}K \subset L \subset \lambda_2 K$, then*

$$\mathcal{E}(K) \subset \lambda_1^{\frac{n+2}{2}} \lambda_2^{\frac{n}{2}} \mathcal{E}(L), \quad \mathcal{E}(L) \subset \lambda_1^{\frac{n}{2}} \lambda_2^{\frac{n+2}{2}} \mathcal{E}(K).$$

Furthermore, there are positive constants a_n, b_n such that $a_n\mathcal{E}(K) \subset K \subset b_n\mathcal{E}(K)$. One can take explicitly $a_n = 2^{-\frac{n}{2}-1}n^{-\frac{3n}{4}-\frac{3}{2}}$, $b_n = 2^{\frac{n}{2}+1}n^{\frac{3n}{4}+\frac{3}{2}}$.

Proof. We follow [16]. We have

$$\int_L \phi(x)^2 dx \leq \int_{\lambda_2 K} \phi(x)^2 dx = \lambda_2^{n+2} \int_K \phi(x)^2 dx.$$

As $\text{vol}(L) \geq \lambda_1^{-n} \text{vol}(K)$, we conclude that

$$\langle \phi, \phi \rangle_L \leq \lambda_1^n \lambda_2^{n+2} \langle \phi, \phi \rangle_K \Rightarrow \mathcal{E}(L) \subset \lambda_1^{\frac{n}{2}} \lambda_2^{\frac{n+2}{2}} \mathcal{E}(K).$$

The second inclusion follows by symmetry.

For the second part, define $K_s = \frac{1}{2}(K + (-K))$. As $-K \subset nK$, we have

$$\frac{1}{n}K_s \subset \frac{2}{n+1}K_s \subset K \subset 2K_s,$$

and by the first part,

$$\mathcal{E}(K_s) \subset 2^{\frac{n}{2}} n^{\frac{n}{2}+1} \mathcal{E}(K), \quad \mathcal{E}(K) \subset 2^{\frac{n}{2}+1} n^{\frac{n}{2}} \mathcal{E}(K_s).$$

We make use of John's maximal volume ellipsoid J of K_s , which satisfies $J \subset K_s \subset \sqrt{n}J$ by John's theorem. Observe that $\mathcal{E}(J) = J$. By the first part, it holds that

$$J \subset n^{\frac{n}{4}} \mathcal{E}(K_s), \quad \mathcal{E}(K_s) \subset n^{\frac{n+2}{4}} J.$$

As

$$\frac{1}{n}J \subset \frac{1}{n}K_s \subset K \subset 2K_s \subset 2\sqrt{n}J,$$

we conclude that

$$K \subset 2^{\frac{n}{2}+1} n^{\frac{3n}{4}+\frac{3}{2}} \mathcal{E}(K), \quad \mathcal{E}(K) \subset 2^{\frac{n}{2}+1} n^{\frac{3n}{4}+\frac{3}{2}} K.$$

□

Definition 2.2. *The Binet–Legendre distance between $K, L \subset \mathbb{R}^n$ is*

$$d_{BL}(K, L) = \min\{d_{\mathcal{E}(g_1K)}(g_1K, g_2L) : g_1, g_2 \in \overline{\text{GL}}(\mathbb{R}^n), \mathcal{E}(g_1K) = \mathcal{E}(g_2L)\}.$$

Lemma 2.3. *d_{BM} and d_{BL} are equivalent distances on $\mathcal{K}(\mathbb{R}^n)$. Explicitly, there is $\epsilon_n > 0$ such that for $d_{BM}(K, L) \leq 1 + \epsilon_n$ we have*

$$a_n(d_{BM}(K, L) - 1) \leq d_{BL}(K, L) \leq b'_n(d_{BM}(K, L) - 1),$$

where $b'_n = (3n + 2)b_n(1 + 6n\frac{b_n}{a_n})$.

Furthermore, for $\epsilon < \epsilon_n$, if $d_{BM}(K, L) = 1 + \epsilon$ and $\mathcal{E}(K) = \mathcal{E}(L) = B^n$, then there is $T \in \text{O}(\mathbb{R}^n)$ such that

$$(1 + d_n\epsilon)^{-1}K \subset TL \subset (1 + d_n\epsilon)K,$$

where $d_n = 4n\frac{b_n}{a_n}(1 + 6n\frac{b_n}{a_n})$.

Later we will use the larger value $d_n = c4^n n^{3n+8}$ for simplicity.

Proof. Write $a = a_n, b = b_n$. Assume $d_{BL}(K, L) = \epsilon$. We may assume that a Euclidean structure is fixed such that $\mathcal{E}(K) = \mathcal{E}(L) = B^n$, and $d_{B^n}(K, L) = \epsilon$. That is, $K \subset L + \epsilon B^n, L \subset K + \epsilon B^n$. Then by Lemma 2.1, $K \subset L + \frac{\epsilon}{a}L = (1 + \frac{\epsilon}{a})L$, and similarly $L \subset (1 + \frac{\epsilon}{a})K$. Thus $d_{BM}(K, L) \leq 1 + \frac{\epsilon}{a} = 1 + \frac{d_{BL}(K, L)}{a}$.

In the other direction, assume $d_{BM}(K, L) = 1 + \epsilon$. We may assume $\mathcal{E}(K) = B^n$, and $K \subset L \subset (1 + \epsilon)K + z$ for some $z \in \mathbb{R}^n$. It holds that

$$q := \text{cent}(L) = \frac{1}{\text{vol}(L)} \int_L x d \text{vol}(x) = \frac{1}{\text{vol}(L)} \int_{L \setminus K} x d \text{vol}(x).$$

Since $K \subset (1 + \epsilon)K + z$, we deduce that

$$-z \in \epsilon K \Rightarrow z \in n\epsilon K.$$

In particular, $|z| \leq n\epsilon b$, and $\max_L |x| \leq |z| + (1 + \epsilon) \max_K |x| \leq b(1 + (n + 1)\epsilon)$.

Consequently by Lemma 2.1, and since for $\epsilon < \frac{1}{n}$

$$(1 + \epsilon)^n - 1 = \sum_{j=1}^n \frac{n(n-1) \cdots (n-j+1)}{j!} \epsilon^j \leq \sum_{j=1}^{\infty} \frac{(n\epsilon)^j}{j!} = e^{n\epsilon} - 1 < 2n\epsilon,$$

it follows that for $\epsilon < \frac{1}{n+1}$,

$$|q| \leq \frac{\text{vol}(L \setminus K)}{\text{vol}(L)} \max_{L \setminus K} |x| \leq \frac{((1 + \epsilon)^n - 1) \text{vol}(K)}{\text{vol}(K)} \max_L |x| \leq 4nb\epsilon$$

and so $\pm q \in \frac{4nb}{a} \epsilon a B^n \subset \frac{4nb}{a} \epsilon K$. Thus

$$(1 - 4\frac{nb}{a}\epsilon)K \subset L - q \subset (1 + \epsilon)K + \frac{nb\epsilon}{a} a B^n + \frac{4nb\epsilon}{a} a B^n \subset \left(1 + (1 + \frac{5nb}{a})\epsilon\right) K.$$

Denote $b'_n = 1 + 5\frac{nb}{a}$. Then (for ϵ sufficiently small)

$$\frac{1}{1 + b'_n \epsilon} K \subset L - q \subset (1 + b'_n \epsilon) K,$$

and it follows again by Lemma 2.1 that $(1 + b'_n \epsilon)^{-n-1} B^n \subset \mathcal{E}(L - q) \subset (1 + b'_n \epsilon)^{n+1} B^n$, and so we can choose a positive definite map T with $T(\mathcal{E}(L - q)) = B^n$ with the eigenvalues of T satisfying $(1 + b'_n \epsilon)^{-n-1} \leq \lambda_1, \dots, \lambda_n \leq (1 + b'_n \epsilon)^{n+1}$. Writing $T = \text{Id} + S$, it follows that the Euclidean operator norm $\|S\| \leq 2nb'_n \epsilon$ (for ϵ small enough). We similarly have $T^{-1} = \text{Id} + S'$ with $\|S'\| \leq 2nb'_n \epsilon$.

We deduce that $L' = T(L - q)$ has $\mathcal{E}(L') = B^n$, and

$$\begin{aligned} L' &\subset T(1 + b'_n \epsilon)K = K + b'_n \epsilon K + S(1 + b'_n \epsilon)K \\ &\subset K + b'_n \epsilon b B^n + b(1 + b'_n \epsilon)S(B^n) \\ &\subset K + (b'_n \epsilon b + 2nb'_n \epsilon b(1 + b'_n \epsilon))B^n. \end{aligned}$$

Similarly

$$\begin{aligned} K &\subset (1 + b'_n \epsilon)T^{-1}L' \\ &\subset L' + (b'_n \epsilon b + 2nb'_n \epsilon b(1 + b'_n \epsilon))B^n. \end{aligned}$$

that is, $d_{BL}(K, L) \leq d_{B^n}(K, L') \leq (3n + 1)b'_n b \epsilon$.

In particular, since

$$L' \subset (1 + \frac{b}{a}(b'_n \epsilon + 2nb'_n \epsilon(1 + b'_n \epsilon)))K, \quad K \subset (1 + \frac{b}{a}(b'_n \epsilon + 2nb'_n \epsilon(1 + b'_n \epsilon)))L',$$

setting $d_n := 4nb'_n \frac{b}{a}$ establishes the last claim of the lemma. \square

Corollary 2.4. *If $d_{BM}(K, B^n) < 1 + \epsilon$ and $\text{cent}(K) = 0$ then $(1 + d_n \epsilon)^{-1} \mathcal{E}(K) \subset K \subset (1 + d_n \epsilon) \mathcal{E}(K)$, and consequently $d_{BM}^{\text{Lin}}(K, B^n) < 1 + d_n \epsilon$.*

Given an ellipse $\mathcal{E} \subset \mathbb{R}^2$, we equip $\partial \mathcal{E}$ with the angular distance function $d_{\mathcal{E}}$ induced by the Euclidean structure for which it is a unit ball. Given two ellipses $\mathcal{E}_i \subset E_i$, $i = 1, 2$, where E_i are 2-dimensional linear spaces, denote by $\text{O}(\mathcal{E}_1, \mathcal{E}_2)$ the set of affine maps $E_1 \rightarrow E_2$ mapping \mathcal{E}_1 to \mathcal{E}_2 . If $E_1 = E_2 = E$, $\mathcal{E}_1 = \mathcal{E}_2 = \mathcal{E}$, we write $\text{O}(\mathcal{E}) = \text{O}(\mathcal{E}, \mathcal{E})$, and define $\text{SO}(\mathcal{E}) \subset \text{O}(\mathcal{E})$ to be the orientation-preserving subgroup. Note that $\text{SO}(\mathcal{E})$ is a circle with distance function $d(g, h) = d_{\mathcal{E}}(gx, hx)$,

which is independent of the choice of $x \in \partial\mathcal{E}$. For general $\mathcal{E}_1, \mathcal{E}_2$, $O(\mathcal{E}_1, \mathcal{E}_2)$ is the disjoint union of two circles, and we define a metric on it by

$$d(g_1, g_2) = \begin{cases} d_{\mathcal{E}_1}(g_1^{-1}g_2, \text{Id}), & g_1^{-1}g_2 \in \text{SO}(\mathcal{E}_1), \\ \infty, & g_1^{-1}g_2 \notin \text{SO}(\mathcal{E}_1). \end{cases}$$

Our main technical tool will be the subset of λ -approximate isometries of convex bodies, defined as follows.

Definition 2.5. *Let $\lambda > 0$, and let E, F be linear spaces of equal dimension. For two convex bodies $K \subset E$, $L \subset F$ we set $G_\lambda(K, L) = \{g \in O(\mathcal{E}(K), \mathcal{E}(L)) : d_{\mathcal{E}(L)}(L, gK) < \lambda\}$.*

We write $G_\lambda(K) := G_\lambda(K, K) \subset O(\mathcal{E}(K))$. We can then also define the subset $G_\lambda^+(K) = G_\lambda(K) \cap \text{SO}(\mathcal{E}(K))$.

The following is immediate from definitions.

Lemma 2.6. *If $g \in G_\lambda(K_1, K_2)$ then $g^{-1} \in G_\lambda(K_2, K_1)$. If $g_j \in G_{\lambda_j}(K_j, K_{j+1})$ for $j = 1, 2$ then $g_2g_1 \in G_{\lambda_1+\lambda_2}(K_1, K_3)$.*

We say that a set $A \subset S^1$ is an ϵ -net if any closed ϵ -interval intersects A .

The following technical lemma is a stable version of the basic fact that a convex body in \mathbb{R}^2 is either an ellipse, or has a finite group of symmetries.

Proposition 2.7. *For all $\epsilon > 0$, if $\alpha, \beta > 0$ satisfy $\frac{1}{a_2}\alpha + \frac{b_2}{a_2}\beta \leq \epsilon$ and $G_\alpha^+(K)$ is a β -net in $\text{SO}(\mathcal{E}(K))$ then $d_{BM}(K, B^2) < 1 + \epsilon$.*

We will later use

$$\alpha(\epsilon) = \frac{a_2}{2}\epsilon, \quad \beta(\epsilon) = \frac{a_2}{2b_2}\epsilon, \tag{1}$$

so that $\alpha = \alpha(\epsilon)$, $\beta = \beta(\epsilon)$ satisfy the assumption of the proposition.

Proof. Fix a Euclidean structure $|\bullet|$ on \mathbb{R}^2 whose unit ball is $\mathcal{E}(K)$. Assume $p \in K$ satisfies $|p| = \max\{|x| : x \in K\}$. By Lemma 2.1, $|p| \leq b_2$. Denoting $D = |p|\mathcal{E}(K)$, we have $K \subset D$. On the other hand, since $G_\alpha^+(K)$ is a β -net, it holds that

$$D \subset K + 2|p|\sin\left(\frac{\beta}{2}\right)\mathcal{E}(K) + \alpha\mathcal{E}(K) \subset \left(1 + \frac{2b_2}{a_2}\sin\frac{\beta}{2} + \frac{1}{a_2}\alpha\right)K,$$

so that $d_{BM}(K, B^2) = d_{BM}(K, D) < 1 + \frac{1}{a_2}\alpha + \frac{b_2}{a_2}\beta$. \square

To pass from Theorem 1.11 to Theorem 1.9, we make use of the von Neumann–Jordan constant, which we now recall.

Definition 2.8. *The von Neumann–Jordan constant $C_{NJ}(V)$ of a normed space V is the least $M \geq 1$ such that for all non-zero $x, y \in V$,*

$$\frac{1}{M} \leq \frac{\|x+y\|^2 + \|x-y\|^2}{2(\|x\|^2 + \|y\|^2)} \leq M.$$

The celebrated von Neumann–Jordan theorem [13] asserts that $C_{NJ}(V) = 1$ implies V is Euclidean. The opposite direction is trivial, asserting simply the validity of the parallelogram identity in Euclidean space.

We will need a stable version of these facts, comparing the von Neumann–Jordan constant with the Banach–Mazur distance to Euclidean space. The fact that near-Euclidean normed spaces satisfy an approximate parallelogram identity appeared in [14].

Lemma 2.9 (Kato–Maligranda–Takahashi [14], Theorem 5). *For any normed space V it holds that $C_{NJ}(V) \leq d_{BM}(V, \mathbb{E}^n)^2$.*

A stable version of the von Neumann–Jordan theorem was established by Passer.

Theorem 2.10 (Passer [19], Theorem 1.2). *For an n -dimensional, symmetric normed space V it holds that $d_{BM}(V, \mathbb{E}^n) \leq 1 + cn^2(C_{NJ}(V) - 1)$.*

Here as before $c > 0$ is an explicit universal constant.

3. FIELDS OF CONVEX BODIES ON A SURFACE

We now prove Theorem 1.11. It will be more convenient to work with the following equivalent formulation.

Theorem 3.1. *Let Σ be a smooth closed surface which is not a torus or a Klein bottle. For any $\delta > 0$ and any continuous field of convex bodies $(K_x)_{x \in \Sigma}$, $K_x \subset T_x \Sigma$, such that $d_{BM}(K_x, K_y) < 1 + \delta$ for all $x, y \in \Sigma$, there must exist $x \in \Sigma$ such that $d_{BM}(K_x, B^2) < 1 + c\delta^{1/3}$.*

Here and in the proof, c, C are explicit universal constants which assume different values in distinct appearances, whose values we omit for simplicity.

Proof. We denote by d the distance function on various circles appearing in the proof.

Let (K_x) be as above, and assume that $d_{BM}(K_x, B^2) \geq 1 + \epsilon$, for all x . We may assume that $\text{cent}(K_x) = 0$ for all x .

Step 1: choose parameters α_3, β_0 such that $G_{\alpha_3}(K_x, K_y)$ does not contain a pair of elements g_1, g_2 with $d(g_1, g_2) \in [\beta_0/2, \beta_0]$.

Denote $\alpha_0 = \alpha(\epsilon)$, $\beta'_0 = \beta(\epsilon)$ as in eq. (1). Let $\beta_0 \leq \beta'_0$ be the largest real of the form $\beta_0 = \frac{2\pi}{B_0}$ with $B_0 \geq 3$ an integer. By Proposition 2.7, $G_{\alpha_0}^+(K_x)$ is not a β_0 -net. Note that $\alpha_0, \beta_0 \in [c\epsilon, C\epsilon]$.

Denote $\alpha_1 = \alpha_0/(2B_0)$, so that $c\epsilon^2 \leq \alpha_1 \leq C\epsilon^2$. It follows that $G_{\alpha_1}^+(K_x)$ does not contain elements g with $d(g, \text{Id}) \in [\beta_0/2, \beta_0]$: if g were such an element, then the iterates $g^j \in G_{\alpha_0}^+(K_x)$ for $j = 0, 1, \dots, 2B_0$ form a β_0 -net, in contradiction.

Denote $\alpha_2 = \alpha_1/2$, $\beta_1 = \beta_0/2$. It follows that the distance between $g_1, g_2 \in G_{\alpha_2}(K_x)$ cannot lie in $[\beta_1, \beta_0]$, or else $(g_1^{-1}g_2)^{\pm 1} \in G_{\alpha_1}^+(K_x) \cap [\beta_1, \beta_0]$.

Using Lemma 2.3, we let $\delta' = b'_2\delta$ be the upper bound on $d_{BL}(K, L)$ when $d_{BM}(K, L) \leq 1 + \delta$. Assume $\alpha_3 := \alpha_2 - \delta' > 8B_0\delta' \iff \delta' < \frac{1}{8B_0+1}\alpha_2$, which holds when $\delta < C\epsilon^3$, as we will henceforth assume to arrive at a contradiction. It follows in particular that $\alpha_3 > \delta'$, and so $G_{\alpha_3}(K_x, K_y)$ is non-empty. Furthermore, $G_{\alpha_3}(K_x, K_y)$ does not contain a pair of elements g_1, g_2 with $d(g_1, g_2) \in [\beta_1, \beta_0]$: fixing $h \in \mathcal{O}(\mathcal{E}(K_x), \mathcal{E}(K_y))$ such that $d_{BL}(K_x, K_y) = d_{\mathcal{E}(K_y)}(hK_x, K_y)$, we would otherwise have elements $\hat{g}_1 := g_1h^{-1}, \hat{g}_2 := g_2h^{-1} \in G_{\alpha_3+\delta'}(K_y) = G_{\alpha_2}(K_y)$, with $d(\hat{g}_1, \hat{g}_2) \in [\beta_1, \beta_0]$, in contradiction.

Step 2: find a range of parameters α for a fixed point $x_0 \in \Sigma$ for which $G_\alpha(K_{x_0})$ has the same β_1 -clusters.

By a β -cluster in $A \subset S^1$ we understand a maximal subset of A of diameter at most β .

Recall that $G_{\alpha_3}(K_x, K_y) \subset \mathcal{O}(\mathcal{E}(K_x), \mathcal{E}(K_y))$, and the latter disjoint union of two circles is equipped with a metric.

It follows by construction that any two distinct β_1 -clusters in $G_{\alpha_3}(K_x, K_y)$ are disjoint, and moreover separated by a distance of at least β_0 . Thus given $g \in G_{\alpha_3}(K_x, K_y)$, we define $[g]$ to be the unique β_1 -cluster containing g . Furthermore, for $\alpha \leq \alpha_3$ and $0 < \eta < \alpha$, if $Z \subset G_{\alpha-\eta}(K_x, K_y)$ is a β_1 -cluster, and $h \in G_\eta(K_y, K_z)$, write $h[Z] := [h \circ g] \subset G_\alpha(K_x, K_z)$, where $g \in Z$ can be arbitrary.

The number of β_1 -clusters in $G_\alpha(K_x, K_y)$ is a non-decreasing function of $\alpha \in (0, \alpha_3]$, which is bounded from above by $2B_0$. Given $0 < \alpha < \alpha' \leq \alpha_3$, there is a natural injective map from the β_1 -clusters of $G_\alpha(K_x, K_y)$ into the β_1 -clusters of $G_{\alpha'}(K_x, K_y)$, given by set inclusion.

Fix $x_0 \in \Sigma$. Since $\alpha_3 > 8B_0\delta'$, we can find $\gamma_0 \in [\delta', \alpha_3 - 3\delta']$ such that the number of β_1 -clusters in $G_\alpha(K_{x_0})$ is constant for $\alpha \in [\gamma_0, \gamma_0 + 3\delta']$. Putting $\gamma_1 = \gamma_0 + 3\delta'$, the β_1 -clusters of $G_\alpha(K_{x_0})$ are therefore naturally identified across the range $\alpha \in [\gamma_0, \gamma_1]$.

Step 3: construct a covering space of Σ out of the β_1 -clusters for K_{x_0} .

For all $x \in \Sigma$, choose $h_x \in \mathcal{O}(\mathcal{E}(K_{x_0}), \mathcal{E}(K_x))$ such that $d_{\mathcal{E}(K_x)}(h_x K_{x_0}, K_x) < \frac{3}{2}\delta'$. The map $Z \mapsto h_x[Z]$ gives an embedding of the set of β_1 -clusters in $G_{\gamma_0}(K_{x_0})$ into the set of β_1 -clusters in $G_{\gamma_1}(K_x, K_x)$. Furthermore, the image C_x of this embedding is independent of the choice of h_x : if \hat{h}_x is another such map, and $g \in G_{\gamma_0}(K_{x_0})$, then $g' = \hat{h}_x^{-1} h_x g \in G_{\gamma_1}(K_{x_0})$. Recall by construction that we may find $g'' \in G_{\gamma_0}(K_{x_0})$ with $d(g', g'') < \beta_1$. It then follows that $d(h_x g, \hat{h}_x g'') = d(\hat{h}_x g', \hat{h}_x g'') < \beta_1$, and so $\hat{h}_x g''$ and $h_x g$ define the same β_1 -cluster in $G_{\gamma_1}(K_x, K_x)$, proving the claim. Let m be the number of elements (clusters) in C_x , which is the same for all x by construction.

Denote by $p : \widehat{C} \rightarrow \Sigma$ the bundle with fiber C_x over $x \in \Sigma$. Our next goal will be to topologize \widehat{C} so as to make it a covering space of Σ . We will next describe a natural identification of the fibers of p over small subsets of Σ .

Fix any Riemannian metric on Σ , and denote by $B_r(z)$ an open ball of radius r centered at z in this metric. Take r_0 small enough so that $B_{r_0}(z)$ can be identified with an open subset of $T_z \Sigma$ through the exponential map, for all $z \in \Sigma$. For $x, y \in B_{r_0}(z)$, $x = \exp_z(\xi)$, $y = \exp_z(\eta)$, we let $\bar{u}_{xy} : T_z \Sigma \rightarrow T_z \Sigma$ be the unique positive definite map, with respect to the inner product determined by $d_\xi \exp^{-1}(\mathcal{E}(K_x))$, mapping $d_\xi \exp^{-1}(\mathcal{E}(K_x))$ to $d_\eta \exp^{-1}(\mathcal{E}(K_y))$.

Thus $u_{xy} := d_\eta \exp \circ \bar{u}_{xy} \circ d_\xi \exp^{-1} \in \text{SO}(\mathcal{E}(K_x), \mathcal{E}(K_y))$. Decreasing r_0 further, we can achieve $u_{xy} \in G_{\delta'/4}^+(K_x, K_y)$ for all $x, y \in B_{r_0}(z)$ by the continuity of $x \mapsto K_x$.

Choose $h_x \in G_{\delta'}(K_{x_0}, K_x)$. Now if Z is a β_1 -cluster in $G_{\gamma_0}(K_{x_0})$, then $h_x[Z]$ is a β_1 -cluster in C_x , while $u_{xy} \circ h_x[Z]$ is a β_1 -cluster in C_y , and we define the bijection $b_{xy} : p^{-1}(x) \rightarrow p^{-1}(y)$ by setting $b_{xy}(h_x[Z]) = u_{xy} \circ h_x[Z]$, for all Z . One readily checks that this correspondence is independent of the choice of h_x , and that $b_{yx'} \circ b_{xy} = b_{xx'}$ for all $x, y, x' \in B_{r_0}(z)$.

This now allows us to topologize \widehat{C} : Taking $U \subset B_{r_0}(z)$ open, $p^{-1}(U)$ is, by the above, the disjoint union of m copies U_1, \dots, U_m of U . The collection of all such U_j is a basis of the topology of \widehat{C} . With this topology, $p : \widehat{C} \rightarrow \Sigma$ is an m -sheeted covering map. Moreover, \widehat{C} naturally inherits a smooth structure from Σ such that p is a local diffeomorphism.

The bundle $p^* \widehat{C}$ over \widehat{C} trivially admits a section $y \mapsto J_y$. Recall that $J_y \subset G_{\gamma_1}(K_{x_0}, K_{p(y)})$.

Step 4: find a continuous section. We next choose a continuous field of closed intervals $\widehat{C} \ni y \mapsto I_y \subset \mathcal{O}(\mathcal{E}(K_{x_0}), \mathcal{E}(K_{p(y)}))$ of length $3\beta_1$ such that $J_y \subset I_y$. This can be done as follows. Choose a finite cover of \widehat{C} by r_0 -balls $U_j = B_{r_0}(z_j)$. Let ρ_j be a subordinate partition of unity. In each U_j , one can easily choose a field of $3\beta_1$ -intervals \widehat{I}_y^j for $y \in U_j$ such that $J_y \subset \widehat{I}_y^j$, and $u_{p(y)p(z)}(\widehat{I}_y^j) \subset \mathcal{O}(\mathcal{E}(K_{x_0}), \mathcal{E}(K_{p(z)}))$ is independent of y , if r_0 is sufficiently small. Then define $I_y := \sum_j \rho_j(y) \widehat{I}_y^j$, namely the weighted mean of the intervals \widehat{I}_y^j with weights $\rho_j(y)$, which is well defined since all of the intervals \widehat{I}_y^j are contained in a fixed interval I'_y of length $2\beta_1 + \beta_1 + 2\beta_1 = 5\beta_1 < 2\pi$ containing J_y , and so the averaging is carried out inside I'_y . Define P_y as the center of I_y .

Thus P_y is a continuous global section over \widehat{C} of the principal $\mathrm{O}(2)$ -bundle with fiber $\mathcal{O}(\mathcal{E}(K_{x_0}), \mathcal{E}(K_{p(y)})) \subset \overline{\mathrm{GL}}(T_{x_0}\Sigma, T_y\widehat{C})$ over $y \in \widehat{C}$, which readily implies that \widehat{C} admits a continuous field of tangent lines: taking any line $L \subset T_{x_0}\Sigma$, $P_y L$ defines such a field by parallel-shifting the lines to pass through the origin. However $\chi(\widehat{C}) = m\chi(\Sigma) \neq 0$, a contradiction.

This contradiction means that one cannot have $\delta < C\epsilon^3$, concluding the proof. \square

Proof of Corollary 1.12. Denote by $K_x \subset T_x\Sigma$ the unit ball of the Finsler structure. By Theorem 1.11, $d_{BM}(K_x, B^2) < 1 + c\delta^{1/3}$ for all $x \in \Sigma$.

The assignment $x \mapsto \mathrm{cent}(K_x)$ defines a vector field over Σ , which must vanish somewhere by the Poincaré-Hopf theorem since by assumption $\chi(\Sigma) \neq 0$. Taking z a point where $\mathrm{cent}(K_z) = 0$, we deduce by Corollary 2.4 that $d_{BM}^{\mathrm{Lin}}(K_z, B^2) < 1 + c\delta^{1/3}$. As $d_{BM}^{\mathrm{Lin}}(K_x, K_z) < 1 + \delta$ for all x , the conclusion follows. \square

4. THE STABLE BANACH CONJECTURE FOR PLANAR SECTIONS

Proof of Theorem 1.9. Let K be the unit ball of V . Let $F \subset V$ be a 3-dimensional subspace. By Theorem 1.11, $d_{BM}(K \cap E, B^2) \leq 1 + c\delta^{1/3}$ for all planes $E \subset F$.

Fix a Euclidean structure on F , and denote by $S(F)$ the unit sphere. For a 2-dimensional subspace $\theta^\perp \subset F$ with $\theta \in S(F)$, define $p(\theta) := \mathrm{cent}(K \cap \theta^\perp) \in \theta^\perp$. As $p(\theta)$ defines a continuous vector field over $S(F)$, it must vanish somewhere. Let $E_0 \subset F$ be a plane such that $\mathrm{cent}(K \cap E_0) = 0$.

By Corollary 2.4, there is an absolute constant $c > 0$ such that

$$(1 + c\delta^{1/3})^{-1} \mathcal{E}(K \cap E_0) \subset K \cap E_0 \subset (1 + c\delta^{1/3}) \mathcal{E}(K \cap E_0).$$

In particular, $K \cap E_0$ is nearly centrally-symmetric:

$$-(K \cap E_0) \subset (1 + c\delta^{1/3})(K \cap E_0).$$

By assumption, it holds for any other linear plane $E \subset V$ that $d_{BM}^{\mathrm{Lin}}(K \cap E, K \cap E_0) < 1 + \delta$, and it easily follows that $-(K \cap E) \subset (1 + c\delta^{1/3})(K \cap E)$ for all E . Therefore, $-K \subset (1 + c\delta^{1/3})K$.

Let $K_s = \frac{1}{2}(K + (-K))$ be the Minkowski symmetrization of K . It follows readily that

$$(1 + c\delta^{1/3})^{-1} K_s \subset K \subset (1 + c\delta^{1/3}) K_s. \quad (2)$$

It follows by the above that $d_{BM}(K_s \cap E, B^2) < 1 + c\delta^{1/3}$ for all linear planes E . By Lemma 2.9, the von Neumann–Jordan constant of each section $K_s \cap E$ satisfies

$C_{NJ}(K_s \cap E) < 1 + c\delta^{1/3}$. Consequently, $C_{NJ}(K_s) < 1 + c\delta^{1/3}$. By Theorem 2.10,

$$d_{BM}(K_s, B^n) = d_{BM}^{\text{Lin}}(K_s, B^n) < 1 + cn^2\delta^{1/3},$$

and by eq. (2), $d_{BM}^{\text{Lin}}(K, B^n) < 1 + cn^2\delta^{1/3}$. \square

For the proof of Theorem 1.8 we will need a stable version of the simple geometric fact proved in [1] (see also [6, Lemma 16.12]), asserting that if the 2-dimensional sections of a convex body through a fixed interior point are all ellipses, then the body is an ellipsoid.

Proposition 4.1. *Let $K \subset \mathbb{R}^n$ be a convex body with $0 \in \text{int}(K)$ such that for all 2-dimensional linear planes E , $d_{BM}(K \cap E, B^2) < 1 + \epsilon$. Then $d_{BM}(K, B^n) < 1 + C_n\sqrt{\epsilon}$, where $C_n = cn^{2n^2}$.*

Proof. We prove the statement with $C'_n = c^n d_{n-1} d_{n-2} \cdots d_2$ replacing C_n , as $C'_n < C_n$. We proceed by induction on n , with the base $n = 2$ being trivial.

First consider the map $\text{Gr}_{n-1}(\mathbb{R}^n) \ni H \mapsto \text{cent}(K \cap H) \in H$. This is a global section of the vector bundle γ^\perp with fiber H over $\mathbb{R}\mathbb{P}^{n-1} = \mathbb{P}((\mathbb{R}^n)^*)$, which has full Stiefel-Whitney class $w(\gamma^\perp) = 1 + a + \cdots + a^{n-1}$, where $a \in H^1(\mathbb{R}\mathbb{P}^{n-1}, \mathbb{Z}_2)$ is the generator (see e.g. [17, chapter 4]). It follows that any global section must vanish somewhere, and we fix a hyperplane H_0 such that $\text{cent}(K \cap H_0) = 0$. Denote $\mathcal{E}_0 = \mathcal{E}(K \cap H_0)$.

By the induction hypothesis, $d_{BM}(K \cap H_0, B^{n-1}) < 1 + C'_{n-1}\sqrt{\epsilon}$. By Corollary 2.4,

$$(1 + d_{n-1}C'_{n-1}\sqrt{\epsilon})^{-1}\mathcal{E}_0 \subset K \cap H_0 \subset (1 + d_{n-1}C'_{n-1}\sqrt{\epsilon})\mathcal{E}_0. \quad (3)$$

Since any chord of \mathcal{E}_0 through the origin is bisected by it, it follows that if $[A, A']$ is a chord of $K \cap H_0$ through the origin, then for any Euclidean norm one has

$$(1 + d_{n-1}C'_{n-1}\sqrt{\epsilon})^{-1} \leq \frac{|A|}{|A'|} \leq 1 + d_{n-1}C'_{n-1}\sqrt{\epsilon} \quad (4)$$

Let H_1, H_2 be the affine hyperplanes supporting K and parallel to H_0 . Assume H_1 is at least as far from the origin as H_2 (with respect to any norm on \mathbb{R}^n/H_0). Fix $z \in \partial K \cap H_1$.

Now let E be any linear 2-dimensional plane through z , and define $L = E \cap H_0$. Denote $\mathcal{E}_E := \mathcal{E}(K \cap E)$, $p_E := \text{cent}(\mathcal{E}_E) = \text{cent}(K \cap E)$. We will use the Euclidean structure on E with unit ball $\mathcal{E}_E - p_E$. Note that by Corollary 2.4,

$$(1 + c\epsilon)^{-1}(\mathcal{E}_E - p_E) \subset K \cap E - p_E \subset (1 + c\epsilon)(\mathcal{E}_E - p_E). \quad (5)$$

Let $L_i = H_i \cap E$, $i = 1, 2$ be the lines in E which are supporting lines of $K \cap E$ and are parallel to L . As the distance between L_1 and L_2 is at least the diameter of $(1 + c\epsilon)^{-1}(\mathcal{E}_E - p_E)$, it follows that

$$\text{dist}(L_1, 0) \geq 1 - c\epsilon. \quad (6)$$

We will repeatedly make use of the following simple fact: If two concentric discs B_1, B_2 of radii $1 - c\epsilon$, resp. $1 + C\epsilon$ are given, and L is a line, then for $\epsilon > 0$,

$$\text{Length}(L \cap (B_2 \setminus B_1)) < c\sqrt{\epsilon}. \quad (7)$$

Orient L arbitrarily. Denote $[A, A'] = K \cap L$, $[A_E, A'_E] = \mathcal{E}_E \cap L$. By eq. (4),

$$\left| \frac{A + A'}{2} - 0 \right| \leq cd_{n-1}C'_{n-1}\sqrt{\epsilon}.$$

By eq. (7), $|A - A_E| \leq c\sqrt{\epsilon}$, and $|A' - A'_E| \leq c\sqrt{\epsilon}$. Consequently, setting $0_E := (A_E + A'_E)/2$, we find

$$|0_E - 0| \leq cd_{n-1}C'_{n-1}\sqrt{\epsilon}. \quad (8)$$

Let z_E be the touching point on $\partial\mathcal{E}_E$ of a translate of L that is on the same side of L as L_1 . By eq. (7),

$$|z - z_E| \leq c\sqrt{\epsilon}. \quad (9)$$

Let $[z, w]$ be the chord in $K \cap E$ defined by the line through $z, 0$. Let $[z_E, w_E]$ be the chord in \mathcal{E}_E defined by the line through $z_E, 0_E$. Let p be the midpoint of $[z, w]$, and note that the midpoint of $[z_E, w_E]$ is p_E .

We next claim that

$$|w - w_E| \leq cd_{n-1}C'_{n-1}\sqrt{\epsilon}. \quad (10)$$

Denoting by w' the intersection of the ray R from z to w with $\partial\mathcal{E}_E$, it holds by eq. (7) that $|w - w'| \leq c\sqrt{\epsilon}$. Next, from eq. (6) we conclude that the angle formed by the ray R_E from z_E to 0_E with $\partial\mathcal{E}_E$ lies in $[\frac{\pi}{4}, \frac{\pi}{2}]$. From eqs. (8), (9), and (6) we deduce that the angle between R and R_E is at most $cd_{n-1}C'_{n-1}\sqrt{\epsilon}$. As $\text{dist}(R \cap \mathcal{E}_E, R_E \cap \mathcal{E}_E) \leq c\sqrt{\epsilon}$, we conclude that $|w_E - w'| \leq cd_{n-1}C'_{n-1}\sqrt{\epsilon}$, and so $|w - w_E| \leq |w - w'| + |w' - w_E| \leq cd_{n-1}C'_{n-1}\sqrt{\epsilon}$, showing eq. (10).

We deduce from eqs. (9), (10) that also $|p - p_E| \leq cd_{n-1}C'_{n-1}\sqrt{\epsilon}$.

From (5) it follows that $K \cap E$ is close to an ellipse with a center at p :

$$(1 + cd_{n-1}C'_{n-1}\sqrt{\epsilon})^{-1}(\mathcal{E}_E - p) \subset K \cap E - p \subset (1 + cd_{n-1}C'_{n-1}\sqrt{\epsilon})(\mathcal{E}_E - p). \quad (11)$$

We now let \mathcal{E} be the unique ellipsoid with center at p such that both z and $\partial\mathcal{E}_0$ lie on $\partial\mathcal{E}$. It follows from eqs. (11), (3), (9), (6) that

$$(1 + cd_{n-1}C'_{n-1}\sqrt{\epsilon})^{-1}(\mathcal{E}_E - p) \subset (\mathcal{E} - p) \cap E \subset (1 + cd_{n-1}C'_{n-1}\sqrt{\epsilon})(\mathcal{E}_E - p),$$

and thus also

$$(1 + cd_{n-1}C'_{n-1}\sqrt{\epsilon})^{-1}(\mathcal{E} - p) \subset K - p \subset (1 + cd_{n-1}C'_{n-1}\sqrt{\epsilon})(\mathcal{E} - p).$$

Taking $C'_n = cd_{n-1}C'_{n-1}$ completes the induction and the proof. \square

Proof of Theorem 1.8. From Theorem 1.11 it follows that $d_{BM}(K \cap E, B^2) < 1 + c\delta^{1/3}$ for all linear planes E . We then deduce from Proposition 4.1 that $d_{BM}(K, B^n) < 1 + cn^{2n^2}\delta^{1/6}$, concluding the proof. \square

REFERENCES

1. H Auerbach, S Mazur, and S Ulam, *On a characteristic property of the ellipsoïde*, Monatshefte für Mathematik und Physik **42** (1935), 45–48.
2. S Banach, *Théorie des opérations linéaires. monografie matematyczne, 1, polskie towarzystwo matematyczne, warszawa, 1932.*[5, Théorème sur les ensembles de première catégorie. Fund. Math **16** (1930), 395–398.
3. Gil Bor, Luis Hernández Lamonedá, Valentín Jiménez-Desantiago, and Luis Montejano, *On the isometric conjecture of Banach*, Geometry & Topology **25** (2021), no. 5, 2621–2642.
4. Javier Bracho and Luis Montejano, *On the complex Banach conjecture*, J. Convex Anal. **28** (2021), no. 4, 1211–1222 (English).
5. Thomas Burger and Rolf Schneider, *On convex bodies close to ellipsoids*, Journal of Geometry **47** (1993), 16–22.
6. Herbert Busemann, *The geometry of geodesics*, reprint of the 1955 original ed., Mineola, NY: Dover Publications, 2005 (English).
7. H Groemer, *Stability theorems for ellipsoids and spheres*, Journal of the London Mathematical Society **49** (1994), no. 2, 357–370.
8. ML Gromov, *On a geometric hypothesis of Banach*, Izv. Akad. Nauk SSSR Ser. Mat **31** (1967), 1105–1114.

9. Peter M Gruber, *Stability of Blaschke's characterization of ellipsoids and Radon norms*, Discrete & Computational Geometry **17** (1997), 411–427.
10. Sergei Ivanov, *Monochromatic Finsler surfaces and a local ellipsoid characterization*, Proceedings of the American Mathematical Society **146** (2018), no. 4, 1741–1755.
11. Sergei Ivanov, Daniil Mamaev, and Anya Nordskova, *Banach's isometric subspace problem in dimension four*, Inventiones mathematicae **233** (2023), no. 3, 1393–1425.
12. Sergei Ivanov, Daniil Mamaev, and Anya Nordskova, *Local Kakutani's ellipsoid characterization*, 2023.
13. P. Jordan and J. von Neumann, *On inner products in linear, metric spaces*, Ann. Math. (2) **36** (1935), 719–723 (English).
14. Mikio Kato, Lech Maligranda, and Yasuji Takahashi, *On James and Jordan-von Neumann constants and the normal structure coefficient of Banach spaces*, Studia Mathematica **144** (2001), no. 3, 275–295.
15. Vladimir S Matveev and Marc Troyanov, *The Binet–Legendre metric in Finsler geometry*, Geometry & Topology **16** (2012), no. 4, 2135–2170.
16. ———, *Completeness and incompleteness of the Binet–Legendre metric*, European Journal of Mathematics **1** (2015), no. 3, 483–502.
17. John W. Milnor and James D. Stasheff, *Characteristic classes*, Texts Read. Math., vol. 32, New Delhi: Hindustan Book Agency, 2005 (English).
18. Luis Montejano, *Convex bodies all whose sections (projections) are equal*, European congress of mathematics. Proceedings of the 8th congress, 8ECM, Portorož, Slovenia, June 20–26, 2021, Berlin: European Mathematical Society (EMS), 2023, pp. 857–883 (English).
19. Benjamin Passer, *An approximate version of the Jordan von Neumann theorem for finite-dimensional real normed spaces*, Linear and Multilinear Algebra **63** (2015), no. 1, 68–77.
20. Rolf Schneider, *Convex bodies: the Brunn–Minkowski theory*, no. 151, Cambridge university press, 2014.

Email address: gautamaish@gmail.com

FACULTY OF MATHEMATICS, TECHNION - ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA 3200003, ISRAEL

Email address: faifmand@tauex.tau.ac.il

SCHOOL OF MATHEMATICAL SCIENCES, TEL AVIV UNIVERSITY, TEL AVIV 6997801, ISRAEL