ON EXTREME VALUES OF QUADRATIC TWISTS OF DIRICHLET-TYPE L-FUNCTIONS

SANOLI GUN AND RASHI LUNIA

ABSTRACT. In a recent work [2], it has been shown that *L*-functions associated with arbitrary non-zero cusp forms take large values at the central critical point. The goal of this note is to derive analogous results for twists of Dirichlet-type functions. More precisely, for an odd integer q > 1, let *F* be a non-zero \mathbb{C} -linear combination of primitive, complex, even Dirichlet characters of conductor *q*. We show that for any $\epsilon > 0$ and sufficiently large *X*, there are $\gg X^{1-\epsilon}$ fundamental discriminants 8d with $X < d \leq 2X$ and (d, 2q) = 1 such that $|L(1/2, F \otimes \chi_{8d})|$ is large.

1. INTRODUCTION

For a Dirichlet character χ modulo q and a complex number s with $\Re(s) > 1$, the *L*-function associated with it is defined by

$$L(s,\chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}.$$

This *L*-function has a meromorphic continuation to the entire complex plane and satisfies a functional equation relating its value at *s* with its value at 1 - s. Consequently, values of these *L*-functions at the central point s = 1/2 are interesting as well as mysterious.

Jutila [6] obtained asymptotic formulae for first and second moments of Dirichlet *L*-functions attached to real characters. Application of Cauchy-Schwartz inequality ensures that there are $\gg \frac{X}{\log X}$ real characters χ of conductor at most X for which $L(1/2, \chi) \neq 0$. This result was improved by Soundararajan in [12] to show that $L(1/2, \chi) \neq 0$ for at least 87.5% of the real characters χ .

In his thesis, Soundararajan [11] showed that for an even, non-quadratic, primitive Dirichlet character ψ of odd conductor q, $L(1/2, \psi \otimes \chi_{8d}) \neq 0$ for at least 20% of the square-free integers $d \geq 0$ with (d, 2q) = 1 and indicated that it can be improved to 33%. In a novel article [13], Soundararajan introduced the so-called resonance method to study large values of Dirichlet *L*-functions at 1/2. This method was used by the first author along with Kohnen and Soundararajan [2] to obtain large values *L*-functions associated with arbitrary non-zero cusp forms at central critical point. The goal of this article is to extend this investigation to the study of large values

²⁰¹⁰ Mathematics Subject Classification. 11M06.

Key words and phrases. Extremal values, Dirichlet-type functions.

of quadratic twists of \mathbb{C} -linear combination of Dirichlet *L*-functions. More precisely, we prove the following theorem.

Theorem 1. Let q > 1 be an odd integer and let

$$F = \sum_{\psi \pmod{q}}^{*} c_{\psi} \psi$$

be a non-zero linear combination of even, non-quadratic, primitive Dirichlet characters of conductor qwith coefficients $c_{\psi} \in \mathbb{C}$. For any $\epsilon > 0$ and sufficiently large X, there exists $\gg X^{1-\epsilon}$ fundamental discriminants 8d with $X < d \leq 2X$ and (d, 2q) = 1 such that

$$|L(1/2, F \otimes \chi_{8d})| > \exp\left(\frac{1}{81}\sqrt{\frac{\log X}{\log\log X}}\right)$$

Remark 1.1. Theorem 1 also holds when F is a non-zero \mathbb{C} -linear combination of odd, non-quadratic, primitive Dirichlet characters of odd conductor q > 1.

This article is arranged as follows. In §2, we recall some preliminaries and state the results leading to the proof of the main theorem. In §3, §4 and §5, we prove the auxiliary results. Finally in §6, we prove Theorem 1.

2. PRELIMINARIES

Throughout the article, p and ℓ will denote primes. For a fundamental discriminant D, let $\chi_D(\cdot) = \left(\frac{D}{\cdot}\right)$ be the real Dirichlet character with conductor |D|. Let q > 1 be an odd integer and ψ a non-quadratic, primitive Dirichlet character modulo q. For simplicity, we shall assume $\psi(-1) = 1$. If (q, d) = 1, the character $\psi \otimes \chi_{8d}$ defined as $\psi \otimes \chi_{8d}(\cdot) = \psi(\cdot)\chi_{8d}(\cdot)$, is primitive with conductor 8dq. The Dirichlet *L*-function attached to $\psi \otimes \chi_{8d}$ is defined as

$$L(s,\psi\otimes\chi_{8d})=\sum_{n=1}^{\infty}\frac{\psi(n)\chi_{8d}(n)}{n^s},\qquad\Re(s)>1.$$

Since ψ is non-quadratic, $\psi \otimes \chi_{8d}$ is non-principal and hence $L(s, \psi \otimes \chi_{8d})$ has an analytic continuation to the entire complex plane. It satisfies the functional equation

$$\Lambda(s,\psi\otimes\chi_{8d})=\epsilon(d)\Lambda(1-s,\psi\otimes\chi_{8d}),$$

where

$$\Lambda(s,\psi\otimes\chi_{8d}) = \pi^{-s/2}(8qd)^{s/2}\Gamma(s/2)L(s,\psi\otimes\chi_{8d}), \quad \epsilon(d) = \psi(8d)\left(\frac{8d}{q}\right)\frac{\tau(\psi)}{\sqrt{q}} \tag{1}$$

and $\tau(\psi) = \sum_{a=1}^{q} \psi(a) e^{2\pi i a/q}$ is the Gauss sum associated to ψ (see [1] and also eq 1.4 and page 13 of [11]). For an odd positive square-free integer *d*, we have (see Lemma 2.2 and eq 1.7 of

[11])

$$|L(1/2,\psi\otimes\chi_{8d})|^2 = 2\sum_{n=1}^{\infty} \left(\frac{8d}{n}\right) \frac{d_{\psi}(n)}{\sqrt{n}} V\left(\frac{\pi n}{8dq}\right)$$
(2)

where $d_{\psi}(n) = \sum_{ab=n} \psi(a) \overline{\psi(b)}$ and for real numbers c > 1/2 and x > 0,

$$V(x) = \frac{1}{2\pi i} \int_{(c)} \left(\frac{\Gamma(s/2 + 1/4)}{\Gamma(1/4)}\right)^2 x^{-s} \frac{ds}{s} \text{ and } V(0) = 1.$$
(3)

We note that (see Lemma 2.1 of [11]) V(x) is a real-valued smooth function on $[0, \infty)$ and for any $\epsilon > 0$, it satisfies

$$V(x) = 1 + O(x^{1/2 - \epsilon})$$
 and $V(x) \ll e^{-x/2}$. (4)

From now onwards, \sum' denotes that the sum is over square-free integers. We record the following lemma on sums of quadratic Dirichlet characters.

Lemma 2. Let $u \leq X$ be an odd natural number. If u is a square, then

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(u) = \frac{X}{\zeta(2)} \prod_{p|2uq} \left(\frac{p}{p+1}\right) + O(u^{\frac{1}{4}}X^{\frac{1}{2}}),$$

while if *u* is not a square then as *X* tends to infinity,

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(u) = O(u^{\frac{1}{4}} X^{\frac{1}{2}} (\log X)^{\frac{3}{4}}).$$

Here the constants in O are dependent on q.

Proof. Since
$$\mu^{2}(d) = \sum_{a^{2}|d} \mu(a)$$
, we get

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(u) = \sum_{\substack{X < d \le 2X \\ (d,2q)=1}} \left(\frac{8d}{u}\right) \sum_{a^{2}|d} \mu(a) = \sum_{\substack{a \le \sqrt{2X} \\ (a,2q)=1}} \mu(a) \left(\frac{8a^{2}}{u}\right) \sum_{\substack{X < b \le \frac{2X}{a^{2}} \\ (b,2q)=1}} \left(\frac{b}{u}\right).$$
(5)

When u is a square, we get

$$\sum_{\substack{a \le \sqrt{2X} \\ (a,2uq)=1}} \mu(a) \sum_{\substack{X \ge b \le \frac{2X}{a^2} \\ (b,2uq)=1}} 1 = \sum_{\substack{a \le \sqrt{2X} \\ (a,2uq)=1}} \mu(a) \left[\frac{X}{a^2} \prod_{p|2uq} \left(1 - \frac{1}{p} \right) + O(\sigma_0(u)) \right]$$
$$= \frac{X}{\zeta(2)} \prod_{p|2uq} \left(\frac{p}{p+1} \right) + O(\sigma_0(u)X^{\frac{1}{2}}),$$

where $\sigma_0(n)$ denotes the number of divisors of n and the O constant is dependent on q. Let χ_0 be the principal character modulo 2q. Then from (5), we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(u) = \left(\frac{8}{u}\right) \sum_{\substack{a \le \sqrt{2X} \\ (a,2uq)=1}} \mu(a) \sum_{\substack{X \\ a^2} < b \le \frac{2X}{a^2}} \chi_0(b) \left(\frac{b}{u}\right).$$
(6)

If *u* is not a square, then $\chi_0(\cdot)\left(\frac{\cdot}{u}\right)$ is a non-principal character and using Pólya-Vinogradov inequality, we get

$$\sum_{\substack{X\\a^2} < b \le \frac{2X}{a^2}} \chi_0(b) \left(\frac{b}{u}\right) \ll \min\left(\sqrt{qu}\log(8qu), \frac{X}{a^2}\right) \ll \left(\sqrt{qu}\log(8qu)\frac{X}{a^2}\right)^{\frac{1}{2}}.$$

Therefore

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(u) \ll u^{\frac{1}{4}} (\log(qu))^{\frac{1}{2}} X^{\frac{1}{2}} \sum_{\substack{a \le \sqrt{2X} \\ (a,2uq)=1}} \frac{1}{a} \ll u^{\frac{1}{4}} (\log(8qu))^{\frac{1}{2}} X^{\frac{1}{2}} \log(X) \ll u^{\frac{1}{4}} X^{\frac{1}{2}} (\log X)^{\frac{3}{4}},$$

where the constant in \ll depends on q. This completes the proof of Lemma 2.

From now on, fix a primitive, non-quadratic Dirichlet character ψ_0 of odd conductor q > 1 such that $\psi_0(-1) = 1$. We define the resonator function as

$$R(d) = \sum_{n \le N} r(n)\chi_{8d}(n)\psi_0(n)$$

where $N = X^{\frac{1}{24}}$ and r(n) is a multiplicative function defined as follows. For a prime p, set

$$r(p) = \begin{cases} \frac{L}{\sqrt{p \log p}} & \text{if } L^2 \le p \le L^4\\ 0 & \text{otherwise,} \end{cases}$$

where $L = \frac{1}{8}\sqrt{\log N \log \log N}$ and set $r(p^n) = 0$ for n > 1. We note that $|r(n)| \le 1$ for all n. In §3, we will prove the following proposition.

Proposition 3. Let the notations be as above. We have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' |R(d)|^2 \le \frac{X}{\zeta(2)} \prod_{L^2 \le p \le L^4} (1 + r(p)^2) + O(X^{\frac{5}{8}} (\log X)^{\frac{3}{4}}),$$

where O constant depends on q. Further

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |R(d)|^6 \ll X \exp\left(O\left(\frac{\log X}{\log\log X}\right)\right),$$

where the constants depend on q.

Using random matrix theory, Keating and Snaith [7] conjectured that

$$\sum_{|d| \le X}^{\flat} L(1/2, \chi_d)^k \sim C_k X(\log X)^{\frac{k(k+1)}{2}},$$

where C_k is a positive constant depending on k. Throughout $\sum_{i=1}^{b}$ implies that the sum is over fundamental discriminants d. In [4], Heath-Brown showed that for any $\epsilon > 0$ and $\sigma \ge 1/2$,

$$\sum_{|d| \le X}^{\flat} |L(\sigma + it, \chi_d)|^4 \ll X^{1+\epsilon} (1+|t|)^{1+\epsilon}.$$

In [9], Rudnick and Soundararajan showed that the predicted lower bounds of Keating and Snaith are true for the *k*-th moments of $|L(1/2, \chi_d)|$. From the work of Harper [3] (see also Soundararajan [14]), conditionally on the Generalised Riemann Hypothesis, upper bounds of correct order are also known. A recent result of Chen [10] proves an asymptotic formula for the fourth moment of $L(1/2, \chi_d)$, conditionally on the Generalised Riemann hypothesis. To the best of our knowledge *k*-th moments of $|L(1/2, \psi \otimes \chi_{8d})|$ for $k \ge 3$ have not been studied in the literature. However, for our purposes, we need a much weaker upper bound on the fourth moment of $|L(1/2, \psi \otimes \chi_{8d})|$ which is analogous to the bound of Heath-Brown. More precisely, in §4, we also prove the following proposition.

Proposition 4. *For any* $\epsilon > 0$ *, we have*

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(1/2, \psi \otimes \chi_{8d})|^4 \ll X^{1+\epsilon},$$

where the constant depends on q.

Finally, in §5, we prove the following proposition which plays a crucial role in proving Theorem 1.

Proposition 5. Let ψ be a primitive, non-quadratic, even Dirichlet character with conductor q. When $\psi = \psi_0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(1/2,\psi_0 \otimes \chi_{8d})|^2 |R(d)|^2 \gg X \log X \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(\left(\frac{1}{8\sqrt{24}} + o(1)\right)\sqrt{\frac{\log X}{\log\log X}}\right),$$

while for $\psi \neq \psi_0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(1/2, \psi \otimes \chi_{8d})|^2 |R(d)|^2 \ll X \log X \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(o\left(\frac{L}{\log L}\right)\right).$$

Remark 2.1. *Results analogous to Proposition 3 and Proposition 5 for modular L-functions are obtained in* [2].

3. PROOF OF PROPOSITION 3

By the definition of resonator function, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' |R(d)|^2 = \sum_{m,n \le N} r(m)r(n)\psi_0(m)\overline{\psi_0(n)} \sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(mn).$$
(7)

Since r(n) is non-zero only when n is square free, we note that m, n are square-free in the above sum. We now apply Lemma 2 to the inner sum of the right hand side of (7). Contribution of

the main term in $\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \chi_{8d}(mn)$ occurs when mn is a square. Since m, n are square-free, mn is a square only when m = n. Therefore, the main term of (7) is

$$\leq \frac{X}{\zeta(2)} \sum_{m \leq N} r(m)^2 \prod_{p \mid 2qm} \left(\frac{p}{p+1}\right) + O(X^{\frac{5}{8}}) \leq \frac{X}{\zeta(2)} \sum_{m \leq N} r(m)^2 + O(X^{\frac{5}{8}}) \\ \leq \frac{X}{\zeta(2)} \prod_{L^2 \leq p \leq L^4} \left(1 + r(p)^2\right) + O(X^{\frac{5}{8}}).$$

We bound the remaining terms of (7) by

$$\ll X^{\frac{1}{2}} (\log X)^{\frac{3}{4}} \sum_{m,n \le N} (mn)^{\frac{1}{4}} \ll X^{\frac{1}{2}} (\log X)^{\frac{3}{4}} N^{\frac{5}{2}} \ll X^{\frac{5}{8}} (\log X)^{\frac{3}{4}}.$$

This completes the proof of the first part of Proposition 3. Again we apply Lemma 2 to calculate

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |R(d)|^6 = \sum_{n_1, \dots, n_6 \le N} r(n_1) \dots r(n_6) \psi_0(\prod_{i=1}^3 n_i) \overline{\psi}_0(\prod_{i=4}^6 n_i) \sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} \chi_{8d}(\prod_{i=1}^6 n_i).$$

As before, the main term occurs when $\prod_{i=1}^{6} n_i$ is a square. Since n_i 's are square-free, each prime dividing the product in this case divides an even number of n_i . Therefore, we get

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |R(d)|^6 \ll X \sum_{\substack{n_1, \dots, n_6 \le N \\ n_1 \dots n_6 = \Box}} r(n_1) \dots r(n_6)$$
$$\ll X \prod_{L^2 \le p \le L^4} \left(1 + \binom{6}{2} r(p)^2 + \binom{6}{4} r(p)^4 + \binom{6}{6} r(p)^6 \right)$$
$$\ll X \exp\left(45 \sum_{L^2 \le p \le L^4} \frac{L^2}{p(\log p)^2} \right) \ll X \exp\left(O\left(\frac{\log X}{\log \log X}\right) \right).$$

This completes the proof of Proposition 3.

4. PROOF OF PROPOSITION 4

In order to prove Proposition 4, we first prove the following proposition.

Proposition 6. Let ψ , χ_{8d} be as before and $s_0 = \sigma_0 + it_0 \in \mathbb{C}$ with $1/2 \le \sigma_0 \le 1$, $T_0 = 1 + |t_0|$. Then for any sufficiently large real number X and any $\epsilon > 0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' |L(s_0, \psi \otimes \chi_{8d})|^4 \ll_{\epsilon} (X + (XT_0)^{2-2\sigma_0})(XT_0)^{\epsilon}.$$

We need the following lemma to prove Proposition 6.

Lemma 7. (Heath-Brown, Corollary 3 of [4]) Let M, Q be positive integers and let a_1, \ldots, a_M be arbitrary complex numbers. Let S(Q) denote the set of all real primitive characters of conductor at most Q. Then for any $\epsilon > 0$, we have

$$\sum_{\chi \in S(Q)} \Big| \sum_{n \le M} a_n \chi(n) \Big|^2 \ll_{\epsilon} Q^{\epsilon} M^{1+\epsilon} (Q+M) \max_{n \le M} |a_n|^2.$$

Proof of Proposition 6. We follow the proof of Theorem 2 of [4]. Consider the integral

$$I = \frac{1}{2\pi i} \int_{(2)} |L(s, \psi \otimes \chi_{8d})|^2 \Gamma(s - s_0) U^{s - s_0} ds$$

for a real parameter U. By Mellin inversion (see Page 99, Exercise 6.6.3 of [8]), we have

$$I = \sum_{n=1}^{\infty} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}}$$
(8)

Moving the line of integration to (α) with $0 \le \alpha < \sigma_0 \le 1$, we get

$$I = |L(s_0, \psi \otimes \chi_{8d})|^2 + \frac{1}{2\pi i} \int_{(\alpha)} |L(s, \psi \otimes \chi_{8d})|^2 \Gamma(s - s_0) U^{s - s_0} ds.$$
(9)

Let T = 1 + |t| and let $\nu(\beta)$ for $0 \le \beta \le 1$ be the infimum of all ν such that

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' |L(\beta + it, \psi \otimes \chi_{8d})|^4 \ll (X + (XT)^{2-2\beta})(XT)^{\nu}$$
(10)

uniformly in X and t. Applying Cauchy-Schwartz inequality, using (8) and (9), we get

$$|L(s_0,\psi\otimes\chi_{8d})|^4 \ll \left|\sum_{n=1}^{\infty}\frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}}e^{-\frac{n}{U}}\right|^2 + \left|\int_{(\alpha)}|L(s,\psi\otimes\chi_{8d})|^2\Gamma(s-s_0)U^{s-s_0}ds\right|^2$$

Again using Cauchy-Schwartz inequality and Stirling's formula, we get

$$\left|\int_{(\alpha)} |L(s,\psi\otimes\chi_{8d})|^2 \Gamma(s-s_0) U^{s-s_0} ds\right|^2 \ll U^{2(\alpha-\sigma_0)} \int_{-\infty}^{\infty} |L(\alpha+it,\psi\otimes\chi_{8d})|^4 e^{-|t-t_0|} dt.$$

Therefore for $0 \le \alpha < \sigma_0 \le 1$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(s_0, \psi \otimes \chi_{8d})|^4 \ll \sum_{\substack{X < d \le 2X \\ (d,2q)=1}} \left| \sum_{n=1}^{\infty} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}} \right|^2$$

$$+ U^{2(\alpha - \sigma_0)} \int_{-\infty}^{\infty} \sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(\alpha + it, \psi \otimes \chi_{8d})|^4 e^{-|t-t_0|} dt.$$
(11)

Applying Stirling's formula and functional equation (1), we get

$$|L(\alpha+it,\psi\otimes\chi_{8d})| \ll d^{1/2-\alpha}T^{1/2-\alpha}|L(1-\alpha+it,\psi\otimes\chi_{8d})|.$$
(12)

Applying (10), (11) and (12), for any $\epsilon > 0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(s_0, \psi \otimes \chi_{8d})|^4 \ll \sum_{\substack{X < d \le 2X \\ (d,2q)=1}} \left| \sum_{n=1}^{\infty} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}} \right|^2 + U^{2(\alpha-\sigma_0)} (XT_0)^{2-4\alpha} (X + (XT_0)^{2\alpha}) (XT_0)^{\nu(1-\alpha)+\epsilon}.$$

Choosing $N_0 = U \log^2(XT_0)$, we note that

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \Big| \sum_{n > N_0} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}} \Big|^2 \ll \sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' \left(\int_{N_0}^{\infty} e^{-n/U} \right)^2 \ll XU^2 e^{-2N_0/U}.$$

Let k be the integer such that $2^k < N_0 \le 2^{k+1}$. We split the sum

$$\sum_{n \le N_0} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}}$$

in $O(\log N_0)$ intervals of the form $M/2 < n \le M$ with $M \ll N_0$. Now, using Lemma 7, we see that

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} \Big| \sum_{M/2 < n \le M} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}} \Big|^2 \ll X^{\epsilon} M^{1-2\sigma_0+\epsilon}(X+M).$$

Therefore, applying Cauchy-Schwartz inequality, for any $1/2 \le \sigma_0 \le 1$, we get

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} \left| \sum_{n=1}^{\infty} \frac{d_{\psi}(n)\chi_{8d}(n)}{n^{s_0}} e^{-\frac{n}{U}} \right|^2 \ll X^{\epsilon} N_0^{1-2\sigma_0+\epsilon}(X+N_0) + XU^2 e^{-2N_0/U} \\ \ll (XT_0U)^{\epsilon}(X+U^{2-2\sigma_0}) + XU^2 e^{-2\log^2 XT_0},$$

where the constants in \ll depend on α and σ_0 . Thus for $1/2 \le \sigma_0 \le 1$ and $0 \le \alpha < \sigma_0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(s_0, \psi \otimes \chi_{8d})|^4 \ll (XT_0U)^{\epsilon} (X + U^{2-2\sigma_0}) + XU^2 e^{-2\log^2 XT_0} + U^{2(\alpha - \sigma_0)} (XT_0)^{2-4\alpha} (X + (XT_0)^{2\alpha}) (XT_0)^{\nu(1-\alpha) + \epsilon},$$
(13)

where the constant in \ll depends on α and σ_0 . For $\sigma_0 > 1/2$, choosing $\alpha = 1 - \sigma_0$ and $U = (XT_0)^{1+\delta}$ for any $0 < \delta < 1$ in (13), we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(s_0, \psi \otimes \chi_{8d})|^4 \ll (XT_0)^{\epsilon} (X + (XT_0)^{(2-2\sigma_0)})((XT_0)^{\delta} + (XT_0)^{(2-4\sigma_0)\delta + \nu(\sigma_0)})$$

uniformly in t_0 and the constant in \ll depends on σ_0 . Since $\epsilon > 0$ is arbitrary, by (10), we have

$$\nu(\sigma_0) \le \max\{\delta, \ \nu(\sigma_0) - (4\sigma_0 - 2)\delta\}.$$

Since $1/2 < \sigma_0 \le 1$, we get $\nu(\sigma_0) \le \delta$. Since δ can be chosen arbitrarily small, we get $\nu(\sigma_0) \le 0$. Now let $\sigma_0 = 1/2$. Choosing $\alpha = 1/2 - \epsilon$ in (13) and using the fact that $\nu(1/2 + \epsilon) \leq 0$ from above, we get

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(s_0, \psi \otimes \chi_{8d})|^4 \ll (XT_0U)^{\epsilon} (X+U) + XU^2 e^{-2\log^2 XT_0} + U^{-2\epsilon} (X+(XT_0)^{1-2\epsilon}) (XT_0)^{5\epsilon}.$$
 (14)

Finally, choosing $U = XT_0$ and noting that $\epsilon > 0$ can be chosen arbitrarily small, we get $\nu(1/2) \leq 0$. This completes the proof of Proposition 6.

As an immediate consequence of Proposition 6, we get

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(1/2, \psi \otimes \chi_{8d})|^4 \ll_{\epsilon} X^{1+\epsilon}.$$

This completes the proof of Proposition 4.

5. PROOF OF PROPOSITION 5

Let ψ, ψ' be arbitrary primitive, non-quadratic even Dirichlet characters of conductor q and $d_{\psi}(n) = \sum_{i} \psi(a) \overline{\psi}(b)$. Applying prime number theorem and eq 5.79 on page 124 of [5], we

have

$$\sum_{p \le x} d_{\psi}(p)^2 = (1 + o(1)) \frac{2X}{\log X} \text{ and for } \psi \ne \psi', \quad \sum_{p \le x} d_{\psi}(p) d_{\psi'}(p) = o\left(\frac{x}{\log x}\right), \tag{15}$$

where the *o* constants depend on *q*. Let $\Phi(t)$ be a smooth function supported on [1/2, 5/2] and $\hat{\Phi}(0) = \int_0^\infty \Phi(t) dt$. In this set-up, we have the following Lemma.

Lemma 8. Let ψ be a primitive, non-quadratic even Dirichlet character with odd conductor q > 1, $u \leq X$ be an integer with (u, 2q) = 1 and h be a multiplicative function defined by

$$h(p^k) = 1 + \frac{1}{p} + \frac{1}{p^2} - \frac{d_{\psi}(p)^2}{p(p+1)}$$

for primes p and integers $k \ge 1$. Write $u = u_1 u_2^2$, where u_1 is square-free. Then for any $\epsilon > 0$, we have

$$\sum_{\substack{d \ (d,2q)=1}}^{d'} \chi_{8d}(u) |L(1/2,\psi \otimes \chi_{8d})|^2 \Phi\left(\frac{d}{X}\right)$$

$$= A_{\psi} \hat{\Phi}(0) \frac{d_{\psi}(u_1)}{h(u)} \frac{\sqrt{u_1}}{\sigma(u_1)} X\left(\log \frac{D_1 X}{u_1} + \sum_{p|u_1} \frac{D_2(p)}{p} \log p + \sum_{\substack{p|u_1\\p \nmid u_1}} \frac{D_3(p)}{p} \log p\right)$$

$$+ O\left(u^{1/2+\epsilon} X^{3/4+\epsilon} + X^{7/8+\epsilon}\right),$$
(16)

where $\sigma(n)$ denotes the sum of positive divisors of n and

$$A_{\psi} = \frac{\phi(2q)}{2q\zeta(2)} |L(1,\psi^2)|^2 \left| 1 - \frac{\psi(2)^2}{2} \right|^2 \prod_{p|2q} \left(1 - \frac{1}{p} \right) h(p) \prod_{p|2q} \left(1 + \frac{1}{p} \right)^{-1}.$$

Here D_1 *is a positive constant which depends on* Φ *,* ψ *and* $D_2(p)$ *,* $D_3(p)$ *are* $\ll 1$ *.*

Proof. Note that

$$\mu^{2}(d) = \sum_{v^{2}|d} \mu(v) = \sum_{\substack{v^{2}|d\\v \leq X^{1/8}}} \mu(v) + \sum_{\substack{v^{2}|d\\v > X^{1/8}}} \mu(v).$$

Therefore, we can split the sum in (16) as

$$\sum_{\substack{d\\(d,2q)=1}}^{\prime} \chi_{8d}(u) |L(1/2,\psi\otimes\chi_{8d})|^2 \Phi\left(\frac{d}{X}\right) = \sum_{\substack{d\\(d,2q)=1}}^{\prime} \sum_{\substack{v^2|d\\v\leq X^{1/8}}} \mu(v)\chi_{8d}(u) |L(1/2,\psi\otimes\chi_{8d})|^2 \Phi\left(\frac{d}{X}\right) + \sum_{\substack{d\\(d,2q)=1}}^{\prime} \sum_{\substack{v^2|d\\v>X^{1/8}}} \mu(v)\chi_{8d}(u) |L(1/2,\psi\otimes\chi_{8d})|^2 \Phi\left(\frac{d}{X}\right).$$
(17)

Using Proposition 1.4 of [11], we see that the first term on the right hand side of (17) is

$$\begin{split} &\sum_{\substack{d\\(d,2q)=1}}\sum_{\substack{v^{2}\mid d\\v\leq X^{1/8}}}\mu(v)\chi_{8d}(u)|L(1/2,\psi\otimes\chi_{8d})|^{2}\Phi\left(\frac{d}{X}\right)\\ &=A_{\psi}\hat{\Phi}(0)\frac{d_{\psi}(u_{1})}{h(u)\sqrt{u_{1}}}\frac{u_{1}}{\sigma(u_{1})}X\Big(\log\frac{D_{1}X}{u_{1}}+\sum_{p\mid u_{1}}\frac{D_{2}(p)}{p}\log p +\sum_{p\mid u_{1}}\frac{D_{3}(p)}{p}\log p\Big)\\ &+O\Big(X^{7/8}\log X \ +\ u^{1/2+\epsilon}X^{3/4+\epsilon}\Big),\end{split}$$

where $h, A_{\psi}, D_1, D_2(p)$ and $D_3(p)$ are before.

Now we shall show that

$$\sum_{\substack{d \ (d,2q)=1}} \sum_{\substack{v^2 \mid d \\ v > X^{1/8}}} \mu(v) \chi_{8d}(u) |L(1/2, \psi \otimes \chi_{8d})|^2 \Phi\left(\frac{d}{X}\right) = O(X^{7/8+\epsilon}).$$

Note that there is a one-to-one correspondence between the sets

$$\left\{ X/2 < d \le 5X/2 \mid (d, 2q) = 1, \ v^2 | d \implies v > X^{1/8} \right\}$$

and

$$\Big\{X/2 < v^2 m \le 5X/2 \mid m \text{ square-free }, (vm, 2q) = 1, v > X^{1/8}\Big\}.$$

Since
$$\sum_{\substack{v>X^{1/8}\\v>X^{1/8}}} \mu(v) \ll d^{\epsilon}$$
, we get

$$\sum_{\substack{d\\(d,2q)=1\\v>X^{1/8}}} \sum_{\substack{v^{2}|d\\v>X^{1/8}}} \mu(v)\chi_{8d}(u)|L(1/2,\psi\otimes\chi_{8d})|^{2}\Phi\left(\frac{d}{X}\right)$$

$$\ll \sum_{\substack{d\\(d,2q)=1\\v^{2}|d\Rightarrowv>X^{1/8}}} d^{\epsilon}|L(1/2,\psi\otimes\chi_{8d})|^{2}\Phi\left(\frac{d}{X}\right)$$

$$\ll X^{\epsilon} \sum_{\substack{X^{1/8}< v\leq\sqrt{5X/2}\\(v,2q)=1}} \sum_{\substack{x^{2}\leq m\leq\frac{5X}{2v^{2}}\\(m,2q)=1}} |L(1/2,\psi\otimes\chi_{8v^{2}m})|^{2}.$$
(18)

Now for $\Re(s) > 1$, we have

$$L(s,\psi\otimes\chi_{8v^2m}) = L(s,\psi\otimes\chi_{8m})\prod_{p\mid v} \left(1-\frac{\psi(p)\chi_{8m}(p)}{p^s}\right).$$
(19)

Since ψ is a non-quadratic character, both $\psi \otimes \chi_{8v^2m}$ and $\psi \otimes \chi_{8m}$ are non-principal characters and hence $L(s, \psi \otimes \chi_{8v^2m})$ and $L(s, \psi \otimes \chi_{8m})$ are entire functions. Therefore by identity theorem (19) holds on \mathbb{C} . Applying Cauchy-Schwartz inequality to the inner sum in (18) and using Proposition 4, we have

$$\sum_{\substack{\frac{X}{2v^2} \le m \le \frac{5X}{2v^2} \\ (m,2q)=1}}' |L(1/2,\psi \otimes \chi_{8v^2m})|^2 \ll \frac{\sqrt{X}}{v} \left(v^{\epsilon} \sum_{\substack{\frac{X}{2v^2} \le m \le \frac{5X}{2v^2} \\ (m,2q)=1}}' |L(1/2,\psi \otimes \chi_{8m})|^4 \right)^{1/2} \ll \frac{X^{1+\epsilon}}{v^2}.$$

Finally taking sum over v, we get

$$\sum_{\substack{d\\(d,2q)=1\\v>X^{1/8}}} \sum_{\substack{v^2|d\\v>X^{1/8}}} \mu(v) \left(\frac{8d}{u}\right) |L(1/2,\psi\otimes\chi_{8d})|^2 \Phi\left(\frac{d}{X}\right) \ll X^{1+\epsilon} \sum_{\substack{X^{1/8} < v \le \sqrt{5X/2}\\(v,2q)=1}} \frac{1}{v^2} \ll X^{7/8+\epsilon}.$$
 (20)

This completes the proof of Lemma 8.

We now use Lemma 8 to complete the proof of Proposition 5.

Proof of Proposition 5. We have

$$\sum_{\substack{d \ (d,2q)=1}}^{d'} |L(1/2,\psi \otimes \chi_{8d})|^2 |R(d)|^2 \Phi\left(\frac{d}{X}\right)$$

=
$$\sum_{m,n \le N} r(m)r(n)\psi_0(m)\overline{\psi_0}(n) \sum_{\substack{d \ (d,2q)=1}}^{d'} \chi_{8d}(mn)|L(1/2,\psi \otimes \chi_{8d})|^2 \Phi\left(\frac{d}{X}\right).$$
(21)

Using Lemma 8, we see that the contribution of the error term from (16) is

$$\ll \sum_{m,n \le N} (X^{7/8+\epsilon} + (mn)^{1/2+\epsilon} X^{3/4+\epsilon}) \ll N^2 X^{7/8+\epsilon} + N^{3+\epsilon} X^{3/4+\epsilon} \ll X^{23/24+\epsilon}.$$

Applying (16) of Lemma 8 and taking sum over all m, n, we see that the main term of (21) is

$$A_{\psi}\hat{\Phi}(0)X\log X\sum_{m,n}r(m)r(n)\psi_{0}(m)\overline{\psi_{0}}(n) \frac{d_{\psi}\left(\frac{mn}{(m,n)^{2}}\right)}{h(mn)}\frac{\sqrt{\frac{mn}{(m,n)^{2}}}}{\sigma\left(\frac{mn}{(m,n)^{2}}\right)}$$

= $A_{\psi}\hat{\Phi}(0)X\log X\prod_{p}\left(1 + \frac{r(p)^{2}}{h(p^{2})}\right)\left(1 + \frac{r(p)}{h(p)}d_{\psi_{0}}(p)d_{\psi}(p)\frac{\sqrt{p}}{p+1}\right).$

Now we show that the contribution from the terms with $\max\{m,n\} > N$ is negligible. Let $\alpha = \frac{1}{8 \log L}$. Then we have

$$\sum_{\max\{m,n\}>N} r(m)r(n)\psi_0(m)\overline{\psi_0}(n) \frac{d_{\psi}\left(\frac{mn}{(m,n)^2}\right)}{h(mn)} \frac{\sqrt{\frac{mn}{(m,n)^2}}}{\sigma\left(\frac{mn}{(m,n)^2}\right)} \\ \ll \sum_{\max\{m,n\}>N} r(m)r(n) \left| d_{\psi}\left(\frac{mn}{(m,n)^2}\right) \right| \frac{\sqrt{\frac{mn}{(m,n)^2}}}{\sigma\left(\frac{mn}{(m,n)^2}\right)} \left(\frac{mn}{N}\right)^{\alpha} \\ \ll \exp\left(-\frac{\log X}{400\log\log X}\right).$$

Finally, the contribution of the remaining terms of (16) in the sum (21) is

$$\ll \sum_{m,n \le N} \frac{r(m)r(n)}{h(mn)} \frac{\left| d_{\psi}\left(\frac{mn}{(m,n)^{2}}\right) \right|}{\sqrt{\frac{mn}{(m,n)^{2}}}} \Big(\log D_{1} + \sum_{p \mid \frac{mn}{(m,n)^{2}}} \left(\frac{1}{p} + 1 \right) \log p + \sum_{p \mid (m,n)} \frac{\log p}{p} \Big)$$

$$\ll \sum_{d=1}^{\infty} \sum_{\substack{m,n=1\\(m,n)=1}}^{\infty} \frac{r(d)^{2}r(m)r(n)}{\sqrt{mn}} \frac{|d_{\psi}(mn)|}{h(d^{2})h(mn)} \Big(\log D_{1} + \sum_{p \mid mn} \left(\frac{1}{p} + 1 \right) \log p + \sum_{p \mid d} \frac{\log p}{p} \Big)$$

$$\ll \prod_{p} \Big(1 + \frac{r(p)^{2}}{h(p^{2})} \Big) \Big(1 + \frac{4r(p)}{h(p)\sqrt{p}} \Big) \Big(\log D_{1} + \sum_{\ell} \frac{r(\ell)}{\sqrt{\ell}} \log \ell + \sum_{\ell} r(\ell)^{2} \frac{\log \ell}{\ell} \Big)$$

$$\ll L \prod_{L^{2} \le p \le L^{4}} \Big(1 + \frac{r(p)^{2}}{h(p^{2})} \Big) \Big(1 + \frac{4r(p)}{h(p)\sqrt{p}} \Big).$$

Therefore, we get

$$\sum_{\substack{d \ (d,2q)=1}}^{d'} |L(1/2,\psi \otimes \chi_{8d})|^2 |R(d)|^2 \Phi\left(\frac{d}{X}\right)$$

$$= A_{\psi} \hat{\Phi}(0) X \log X \prod_{L^2 \le p \le L^4} \left(1 + \frac{r(p)^2}{h(p^2)}\right) \left(1 + \frac{r(p)}{h(p)} d_{\psi_0}(p) d_{\psi}(p) \frac{\sqrt{p}}{p+1}\right)$$

$$+ O\left(X \sqrt{\log X \log \log X} \prod_{L^2 \le p \le L^4} \left(1 + \frac{r(p)^2}{h(p^2)}\right) \left(1 + \frac{4r(p)}{\sqrt{p}h(p)}\right)\right). \tag{22}$$

Using the fact that $\frac{1}{h(p^k)} = 1 + O(\frac{1}{p})$ for $k \ge 1$, we get

$$\left(1 + \frac{r(p)^2}{h(p^2)}\right) \left(1 + \frac{r(p)}{h(p)} d_{\psi_0}(p) d_{\psi}(p) \frac{\sqrt{p}}{p+1}\right) = (1 + r(p)^2) \left(1 + \frac{r(p)}{\sqrt{p}} d_{\psi_0}(p) d_{\psi}(p) + O\left(\frac{r(p)^3}{\sqrt{p}}\right)\right)$$
$$= (1 + r(p)^2) \exp\left(\frac{r(p)}{\sqrt{p}} d_{\psi_0}(p) d_{\psi}(p) + O\left(\frac{r(p)^3}{\sqrt{p}}\right)\right).$$

Therefore the product in the main term of (22) becomes

$$\prod_{p} (1+r(p)^{2}) \exp\left(\sum_{p} \left(\frac{r(p)}{\sqrt{p}} d_{\psi_{0}}(p) d_{\psi}(p) + O\left(\frac{r(p)^{3}}{\sqrt{p}}\right)\right)\right)$$

=
$$\prod_{p} (1+r(p)^{2}) \exp\left(\sum_{p} \frac{r(p)}{\sqrt{p}} d_{\psi_{0}}(p) d_{\psi}(p) + O\left(\frac{L}{\log^{3} L}\right)\right).$$

Similarly, the product in the error term of (22) becomes

$$\prod_{p} (1+r(p)^2) \exp\left(\sum_{p} \frac{4r(p)}{\sqrt{p}} + O\left(\frac{L}{\log^3 L}\right)\right)$$

When $\psi = \psi_0$, we choose $\Phi(t)$ to be a smooth function supported on [1,2] with $0 \le \Phi(t) \le 1$ for all t and $\Phi(t) = 1$ on [1.1, 1.9]. Then we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}' |L(1/2,\psi_0 \otimes \chi_{8d})|^2 |R(d)|^2$$

$$\geq \sum_{\substack{d \\ (d,2q)=1}}' |L(1/2,\psi \otimes \chi_{8d})|^2 |R(d)|^2 \Phi\left(\frac{d}{X}\right)$$

$$\geq \frac{4}{5} A_{\psi} X \log X \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(\sum_{L^2 \le p \le L^4} \frac{r(p)}{\sqrt{p}} d_{\psi_0}(p) d_{\psi}(p) + O\left(\frac{L}{\log^3 L}\right)\right)$$

$$+ O\left(X \sqrt{\log X \log \log X} \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(\sum_{L^2 \le p \le L^4} \frac{4r(p)}{\sqrt{p}} + O\left(\frac{L}{\log^3 L}\right)\right)\right).$$

Using (15), we note that

$$\sum_{L^2 \le p \le L^4} \frac{r(p)}{\sqrt{p}} d_{\psi_0}(p)^2 = \left(\frac{1}{2} + o(1)\right) \frac{L}{\log L}.$$

This implies that

$$\begin{split} \sum_{\substack{X < d \le 2X \\ (d, 2q) = 1}}^{\prime} |L(1/2, \psi_0 \otimes \chi_{8d})|^2 |R(d)|^2 \gg X \log X \prod_{L^2 \le p \le L^4} (1 + r(p)^2) \exp\left(\left(\frac{1}{2} + o(1)\right) \frac{L}{\log L}\right) \\ \gg X \log X \prod_{L^2 \le p \le L^4} (1 + r(p)^2) \exp\left(\left(\frac{1}{8\sqrt{24}} + o(1)\right) \sqrt{\frac{\log X}{\log \log X}}\right). \end{split}$$

When $\psi \neq \psi_0$, we choose $\Phi(t)$ to be a smooth function supported on [1/2, 5/2] with $0 \leq \Phi(t) \leq 1$ for all t and $\Phi(t) = 1$ on [1, 2]. Then

$$\begin{split} &\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(1/2, \psi \otimes \chi_{8d})|^2 |R(d)|^2 \\ \le &\sum_{\substack{d \\ (d,2q)=1}} |L(1/2, \psi \otimes \chi_{8d})|^2 |R(d)|^2 \Phi\left(\frac{d}{X}\right) \\ \le & 2A_{\psi} X \log X \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(\sum_{L^2 \le p \le L^4} \frac{r(p)}{\sqrt{p}} d_{\psi_0}(p) d_{\psi}(p) + O\left(\frac{L}{\log^3 L}\right)\right) \\ &+ O\left(X \sqrt{\log X \log \log X} \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(\sum_{L^2 \le p \le L^4} \frac{4r(p)}{\sqrt{p}} + O\left(\frac{L}{\log^3 L}\right)\right)\right). \end{split}$$

When $\psi \neq \psi_0$, applying (15), we get

$$\sum_{p} \frac{r(p)}{\sqrt{p}} d_{\psi_0}(p) d_{\psi}(p) = o\left(\frac{L}{\log L}\right).$$

This implies for $\psi \neq \psi_0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(1/2, \psi \otimes \chi_{8d})|^2 |R(d)|^2 \ll X \log X \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(o\left(\frac{L}{\log L}\right)\right).$$

This completes the proof of Proposition 5.

14

6. PROOF OF THEOREM 1

Since *F* is non-zero, $c_{\psi_0} \neq 0$ for some ψ_0 . Without loss of generality, we can assume that $c_{\psi_0} = 1$. Applying triangle inequality and Cauchy-Schwartz inequality, we have

$$2|L(1/2, F \otimes \chi_{8d})|^{2} = 2\Big|\sum_{\substack{\psi \pmod{q} \\ \psi \pmod{q}}}^{*} c_{\psi}L(1/2, \psi \otimes \chi_{8d})\Big|^{2} \\ \ge |L(1/2, \psi_{0} \otimes \chi_{8d})|^{2} - A\left(\sum_{\substack{\psi \pmod{q} \\ \psi \neq \psi_{0}}}^{*} |L(1/2, \psi \otimes \chi_{8d})|^{2}\right)$$

for some positive constant A > 0. Let S denote the set of square free integers d co-prime to 2q with $X < d \le 2X$ such that

Г

_

$$|L(1/2,\psi_0 \otimes \chi_{8d})|^2 > A \sum_{\substack{\psi \pmod{q} \\ \psi \neq \psi_0}}^* |L(1/2,\psi \otimes \chi_{8d})|^2 + \exp\left(\frac{1}{40}\sqrt{\frac{\log X}{\log\log X}}\right)$$

Therefore

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}} |L(1/2, \psi_0 \otimes \chi_{8d})|^2 |R(d)|^2 \le \sum_{d \in \mathcal{S}} |L(1/2, \psi_0 \otimes \chi_{8d})|^2 |R(d)|^2$$

$$+ \sum_{\substack{X < d \le 2X \\ (d,2q)=1}} \left(A \sum_{\substack{\psi \pmod{q} \\ \psi \neq \psi_0}}^* |L(1/2, \psi \otimes \chi_{8d})|^2 + \exp\left(\frac{1}{40}\sqrt{\frac{\log X}{\log \log X}}\right) \right) |R(d)|^2.$$
(23)

Using Proposition 3 and Proposition 5, we see that the contribution of the second term on the right hand side of the above equation is negligible. This is because for any $\psi \neq \psi_0$, we have

$$\sum_{\substack{X < d \le 2X \\ (d,2q)=1}}^{\prime} |L(1/2, \psi \otimes \chi_{8d})|^2 |R(d)|^2 \ll X \log X \prod_{L^2 \le p \le L^4} (1+r(p)^2) \exp\left(o\left(\sqrt{\frac{\log X}{\log\log X}}\right)\right).$$

Again applying Proposition 5, we see that the lower bound of the left hand side of (23) is

$$X \log X \prod_{L^2 \le p \le L^4} (1 + r(p)^2) \exp\left(\left(\frac{1}{8\sqrt{24}} + o(1)\right) \sqrt{\frac{\log X}{\log\log X}}\right) \ll \sum_{d \in \mathcal{S}} |L(1/2, \psi_0 \otimes \chi_{8d})|^2 |R(d)|^2.$$
(24)

Using Cauchy-Schwartz inequality, Hölder's inequality, Proposition 3, and Proposition 4, for any $\epsilon > 0$, we see that

$$\sum_{d\in\mathcal{S}} |L(1/2,\psi_0\otimes\chi_{8d})|^2 |R(d)|^2 \ll \left(\sum_{\substack{X
(25)$$

Using (24) and (25), we conclude that $|S| \gg X^{1-\epsilon}$. This completes the proof of Theorem 1. \Box

SANOLI GUN AND RASHI LUNIA

Acknowledgments. We would like to thank K. Soundararajan for sharing his PhD thesis with us.

REFERENCES

- [1] H. Davenport, Multiplicative number theory, Grad. Texts in Math. 74, Third edition, *Springer-Verlag, New York*, (2000).
- [2] S. Gun, W. Kohnen and K. Soundararajan, *Large Fourier coefficients of Half-Integer weight modular forms*, to appear in Amer J. Math.
- [3] A. Harper, Sharp conditional bounds for moments of the Riemann zeta function, preprint, arXiv:1305.4618 (2013).
- [4] D. Heath-Brown, A mean value estimate for real character sums, Acta Arith. 72 (1995), no.3, 235–275.
- [5] H. Iwaniec and E. Kowalski, Analytic Number Theory, American Math. Soc. Colloq. Publ. 53, American Mathematical Society, Providence, 2004.
- [6] M. Jutila, On the mean value of $L(1/2, \chi)$ for real characters, Analysis 1 (1981), no.2, 149–161.
- [7] J. Keating and N. Snaith, *Random matrix theory and L-functions at* s = 1/2, Comm. Math. Phys. **214** (2000), no.1, 91–110.
- [8] M. R. Murty, Problems in analytic number theory, Graduate Texts in Mathematics 206, Second Edition, Springer, (2008).
- [9] Z. Rudnick and K. Soundararajan, *Lower bounds for moments of L-functions: symplectic and orthogonal examples*, Multiple Dirichlet series, automorphic forms, and analytic number theory, Proc. Sympos. Pure Math. 75 (2006) American Mathematical Society, Providence, RI, 293–303.
- [10] Q. Shen, The fourth moment of quadratic Dirichlet L-functions, Math. Z. 298 (2021), no.1–2, 713–745.
- [11] K. Soundararajan, Quadratic twists of Dirichlet L-functions, Ph.D. Thesis, Princeton University ProQuest LLC, Ann Arbor, MI, (1998).
- [12] K. Soundararajan, Nonvanishing of quadratic Dirichlet L-functions at s = 1/2, Ann. of Math. (2) 152 (2000), no.2, 447–488.
- [13] K. Soundararajan, Extreme values of zeta and L-functions, Math. Ann. 342 (2008), no. 2, 467–486.
- [14] K. Soundararajan, Moments of the Riemann zeta function, Ann. of Math. (2) 170 (2009), no.2, 981–993.

SANOLI GUN AND RASHI LUNIA,

THE INSTITUTE OF MATHEMATICAL SCIENCES, A CI OF HOMI BHABHA NATIONAL INSTITUTE, CIT CAMPUS, TARAMANI, CHENNAI 600 113, INDIA.

Email address: sanoli@imsc.res.in
Email address: rashisl@imsc.res.in