

TETRAHEDRON DUALITY

JUNYA YAGI

ABSTRACT. A certain two-dimensional supersymmetric gauge theory is argued to satisfy a relation that promotes the Zamolodchikov tetrahedron equation to an infrared duality between two quantum field theories. Solutions of the tetrahedron equation with continuous spin variables are obtained from partition functions of this theory and its variants.

1. INTRODUCTION

Dualities of supersymmetric gauge theories yield solutions of the Yang–Baxter equation [3, 4, 27, 31]. In this paper we discuss solutions of the Zamolodchikov tetrahedron equation [32] which similarly arise from gauge theory dualities.

1.1. The tetrahedron equation for IRC models. The tetrahedron equation is a three-dimensional analog of the Yang–Baxter equation and appears as a condition for integrability in the context of classical spin models on three-dimensional lattices.

In an interaction-round-a-cube (IRC) model, spin variables are located at the vertices of a simple cubic lattice and their interactions take place inside the cubes of the lattice. The Boltzmann weight for the interaction in a single cube is a function

$$(1) \quad W(a|e, f, g|b, c, d|h; s, t, u)$$

of eight spin variables a, b, \dots, h at the vertices of the cube. It may also depend on parameters s, t, u , called the *spectral parameters*, which we think of as assigned to three planes bisecting the cube. See Figure 1(a).




FIGURE 1. (a) The configuration of spin variables for the Boltzmann weight (1). (b) The tetrahedron equation in the IRC form.




FIGURE 2. The presentations of the Boltzmann weight (1) and the tetrahedron equation (2) using curves in a plane. Spin variables are assigned to the regions separated by the curves.

The tetrahedron equation for IRC models reads [1]

$$\begin{aligned}
 (2) \quad & \int_d \mu(d) W(a_4|c_2, c_1, c_3|b_1, b_3, b_2|d; s_1, s_3, s_2) W(c_1|b_2, a_3, b_1|c_4, d, c_6|b_4; s_1, s_4, s_2) \\
 & \quad \times W(b_1|d, c_4, c_3|a_2, b_3, b_4|c_5; s_1, s_4, s_3) W(d|b_2, b_4, b_3|c_5, c_2, c_6|a_1; s_2, s_4, s_3) \\
 & = \int_d \mu(d) W(b_1|c_1, c_4, c_3|a_2, a_4, a_3|d; s_2, s_4, s_3) W(c_1|b_2, a_3, a_4|d, c_2, c_6|a_1; s_1, s_4, s_3) \\
 & \quad \times W(a_4|c_2, d, c_3|a_2, b_3, a_1|c_5; s_1, s_4, s_2) W(d|a_1, a_3, a_2|c_4, c_5, c_6|b_4; s_1, s_3, s_2),
 \end{aligned}$$

where the integrals are performed for the value of the spin variable d with respect to a measure $\mu(d)$. Geometrically, the two sides of the tetrahedron equation represent two ways of decomposing a rhombic dodecahedron into four cubes, as depicted in Figure 1(b).

The Boltzmann weight (1) may also be presented as a *Yang–Baxter move* (or *braid move*) on a configuration of three curves in a plane. In Figure 1(a), imagine a fourth plane that is perpendicular to the line through a and h . Its intersections with the other three planes make three straight lines. As the fourth plane moves from the front to the back side of the cube, the three lines undergo a Yang–Baxter move. The tetrahedron equation amounts to an equality between two sequences of four Yang–Baxter moves applied to a configuration of four curves. See Figure 2.

1.2. Solution by the brane box model. The solutions of the tetrahedron equation (2) discussed in this paper originate from a two-dimensional gauge theory with $\mathcal{N} = (0, 2)$ supersymmetry as well as its variants. This gauge theory, known as the




FIGURE 3. (a) Part of the brane box model that contains $2 \times 2 \times 2$ cubes. (b) The natural unit cell of the brane box model. (c) The oblique unit cell which satisfies the tetrahedron duality.

brane box model [18, 24], is realized by intersecting branes that form a cubic lattice in a three-dimensional subspace of the ten-dimensional spacetime of string theory.

The gauge group and the matter content of the brane box model may be presented as if they constitute a spin model analogous to the Ising model. The “spins” are located at the vertices of a simple cubic lattice, and each corresponds to a $U(N)$ factor in the gauge group. There are two types of pairwise “interactions” (or “bonds”) between spins, produced by chiral multiplets and Fermi multiplets in bifundamental representations.¹ Part of this spin model is illustrated in Figure 3(a), where spins are drawn as circles and interactions are represented by solid and dashed arrows.

Given a spin model on a lattice, the choice of unit cell from which the model is constructed is far from unique. In the case of the brane box model, however, there is a unique unit cell of smallest size whose faces are parallel to branes in the underlying string theory picture.² This unit cell is shown in Figure 3(b). It is a cube of unit volume with eight spins located at its vertices and represents a theory with eight $U(N)$ flavor symmetry³ groups. The brane box model is obtained from infinite copies of this theory glued together. The gluing is done by gauging: to glue two copies of the unit cell, one identifies the $U(N)$ symmetries at the overlapping vertices of the two cubes and changes them to gauge symmetries.

In [29], the author argued that the above natural choice of unit cell should lead to a solution of the tetrahedron equation. The argument adapted a beautiful idea of Kevin Costello [8, 9] and involves surface defects in a partially topological quantum field theory. Although the reasoning was seemingly solid, there was one problem: the natural unit cell did not appear to satisfy the tetrahedron equation!

In the present work we partially resolve this problem. Consider another unit cell, which is *slanted* compared to the natural unit cell by one unit along one of the three axes of the cubic lattice, as shown in Figure 3(c). We will call it the *oblique unit cell* of the brane box model. Under a reasonable assumption about the

¹The actual interactions in the brane box model as a field theory are due to coupling to a vector multiplet and a cubic superpotential.

²There is still a choice to be made regarding the locations of seven arrows in the unit cell.

³A *flavor symmetry* of a supersymmetric field theory is a global symmetry that commutes with the supersymmetry of the theory.

cancellation of gauge anomaly, we will show that the oblique unit cell does satisfy the tetrahedron equation.

1.3. The tetrahedron duality. The statement “the oblique unit cell satisfies the tetrahedron equation” needs to be made more precise.

Let

$$(3) \quad (a|e, f, g|b, c, d|h; s, t, u)$$

denote the theory represented by the oblique unit cell. This is an $\mathcal{N} = (0, 2)$ supersymmetric field theory with eight $U(N)$ flavor symmetry groups, $U(N)_a, U(N)_b, \dots, U(N)_h$, and the R-charge assignment for its matter content depends on spectral parameters s, t, u .

The claim is that the oblique unit cell (3) satisfies the *tetrahedron duality*

$$(4) \quad \text{Gauge}_d \left(\begin{array}{l} (a_4|c_2, c_1, c_3|b_1, b_3, b_2|d; s_1, s_3, s_2)(c_1|b_2, a_3, b_1|c_4, d, c_6|b_4; s_1, s_4, s_2) \\ (b_1|d, c_4, c_3|a_2, b_3, b_4|c_5; s_1, s_4, s_3)(d|b_2, b_4, b_3|c_5, c_2, c_6|a_1; s_2, s_4, s_3) \end{array} \right) \\ \simeq \text{Gauge}_d \left(\begin{array}{l} (b_1|c_1, c_4, c_3|a_2, a_4, a_3|d; s_2, s_4, s_3)(c_1|b_2, a_3, a_4|d, c_2, c_6|a_1; s_1, s_4, s_3) \\ (a_4|c_2, d, c_3|a_2, b_3, a_1|c_5; s_1, s_4, s_2)(d|a_1, a_3, a_2|c_4, c_5, c_6|b_4; s_1, s_3, s_2) \end{array} \right),$$

where Gauge_d is the operation of gauging $U(N)_d$ and \simeq means equivalence of theories at low energies. Each side of this relation is an $\mathcal{N} = (0, 2)$ supersymmetric gauge theory constructed from four copies of the oblique unit cell coupled by gauging $U(N)_d$. The two sides have the same gauge group and global symmetry groups, but their matter contents are different.

In fact, the tetrahedron duality (4) is a consequence of the $(0, 2)$ *trinality*, an infrared equivalence of a triplet of $\mathcal{N} = (0, 2)$ supersymmetric gauge theories with unitary gauge groups [16], again under a reasonable assumption about anomaly cancellation.

1.4. Elliptic genus solutions. Since the tetrahedron duality (4) holds at the level of quantum field theory, any physical quantity computed at low energies in the theories represented by the two sides match. If a quantity W is compatible with the gauging operation in the sense that it satisfies the relation

$$(5) \quad W \text{Gauge}_d \simeq \int_d \mu(d)W,$$

then W computed in the oblique unit cell (3) (for which we will use the notation (1)) is a solution of the tetrahedron equation (2). The partition function of the lattice model defined by this solution equals W computed in the brane box model.

Elliptic genera are excellent examples of such quantities. There are different versions of elliptic genera, but all of them are partition functions on a torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$, $\text{Im } \tau > 0$, and take the form

$$(6) \quad \text{Tr}_{\mathcal{H}_{\text{BPS}}} \left((-1)^F q^{H_L} \prod_{\alpha} u_{\alpha}^{K_{\alpha}} \right).$$

Here \mathcal{H}_{BPS} is a space of states preserving half of $\mathcal{N} = (0, 2)$ supersymmetry, $(-1)^F$ is a fermion parity operator, H_L is a left-moving Hamiltonian and $q = e^{2\pi i\tau}$; the theory has $U(1)$ flavor symmetries $U(1)_{\alpha}$, $\alpha = 1, \dots, N_f$, K_{α} is a generator of $U(1)_{\alpha}$ and u_{α} is an associated parameter, called the *fugacity* for K_{α} . Being a partition function on an elliptic curve that can be written as a trace, an elliptic

genus is a power series in q and u_α with integer coefficients and has a nice modular property.

Not only are elliptic genera compatible with gauging, but they are also invariant under continuous deformations of the theory (barring wall-crossing phenomena). As such, they can be computed exactly in the weak coupling limit, and integral formulas are known for theories with standard Lagrangian descriptions [5, 6].

1.5. Abelian gauge anomalies. The elliptic genera of the brane box model thus provide solutions of the tetrahedron equation (2). There is a caveat, however.

For the brane box model to define a consistent quantum field theory, the gauge anomaly in the abelian part of the gauge group, generated by chiral fermions, must be canceled. The theory represented by the oblique unit cell is also plagued by an abelian gauge anomaly. A version of the Green–Schwarz mechanism is expected to cancel these anomalies [24].

The $(0, 2)$ triality, on the other hand, has been mostly supported by calculations using another method of cancelling abelian gauge anomalies. In this method, additional Fermi multiplets in determinant representations are introduced.

In the claim made above that the oblique unit cell satisfies the tetrahedron duality, we have assumed that *the cancellation of abelian gauge anomaly in the brane box model does not compromise the validity of the $(0, 2)$ triality*. There is a good reason to believe that this is the case based on the study of brane brick models [10–13], of which the brane box model is one example.

That said, the precise mechanism of anomaly cancellation for the brane box model is yet to be understood. This fact prevents us from writing down the elliptic genus solutions.

1.6. Solutions by the 1D brane box model and the $(0, 1)$ brane box model.

The problem stemming from the lack of understanding of anomaly cancellation can be circumvented in two ways:⁴

- (1) Dimensionally reduce the brane box model to one dimension. We will call the resulting theory the *1D brane box model*. Since chiral anomalies are absent in odd dimensions, the abelian gauge anomaly vanishes after the dimensional reduction. The oblique unit cell for the 1D brane box model satisfies the tetrahedron duality and its Witten genus (which is a partition function on a circle) is a solution of the tetrahedron equation.
- (2) Replace the brane box model with an $\mathcal{N} = (0, 1)$ supersymmetric gauge theory with special orthogonal gauge group. This theory, which we will refer to as the $(0, 1)$ *brane box model*, is free of gauge anomaly and enjoys a triality almost identical to the $(0, 2)$ triality [19]. Consequently, the oblique unit cell for the $(0, 1)$ brane box model satisfies the tetrahedron duality and its elliptic genera solve the tetrahedron equation, though they lack spectral parameters since there is no continuous R-symmetry for $\mathcal{N} = (0, 1)$ supersymmetry.

⁴Here are some of the failed attempts that the author has made to cancel the abelian gauge anomaly and write down the elliptic genus solutions: (1) Replace $U(N)$ gauge groups with $SU(N)$. (There is no triality for $SU(N)$ gauge groups.) (2) Introduce extra Fermi multiplets. (It seems hard to cancel the anomaly while preserving the locality of the lattice model, let alone the $(0, 2)$ triality.) (3) Add a determinant Fermi multiplet for every bifundamental chiral multiplet and a determinant chiral multiplet for every bifundamental Fermi multiplet. (The additional multiplets are not compatible with the $(0, 2)$ triality.)

We will write down two solutions of the tetrahedron equations produced by the 1D brane box model and the $(0, 1)$ brane box model. The lattice models defined by these solutions have continuous spin variables, coming from the fugacities for the flavor symmetry groups of the oblique unit cells.

1.7. Open questions. While we obtain solutions of the tetrahedron equation from the brane box model and related theories, we leave many questions unanswered in this paper. Some important ones are:

- Why does the unit cell need to be oblique in order for it to satisfy the tetrahedron duality?
- How is the abelian gauge anomaly canceled in the brane box model?
- What is the significance of the third theory related by the triality to the two sides of the tetrahedron duality?

Seeking connections with other works, we can further ask interesting questions. Here are two:

- String theory realizations of the trialities of brane brick models and their $\mathcal{N} = (0, 1)$ supersymmetric projections [14] were given in [12] and [15]. Do brane brick models provide more examples of theories satisfying the tetrahedron duality?
- Solutions of the tetrahedron equation were constructed in [20–22, 28] by a method using quantum cluster algebras. They are closely related to the solution [2, 23] realized by a brane configuration in M-theory [30]. What are the relations, if any, of these solutions to the ones obtained in this paper?

In connection with the former question, it should be mentioned that in [12] a rhombic dodecahedron structure was observed in brane brick models and a possibility of its relevance to the tetrahedron equation was suggested.

2. THE BRANE BOX MODEL AND THE TETRAHEDRON DUALITY

In this section we introduce the brane box model and explain how the $(0, 2)$ triality implies that its oblique unit cell satisfies the tetrahedron duality. We also discuss the cancellation of abelian gauge anomaly and the relation of spectral parameters to boundary conditions on the lattice.

2.1. The brane box model. Consider a three-dimensional Euclidean subspace \mathbb{R}^3 of the ten-dimensional Minkowski spacetime of type IIA string theory. Let (x, y, z) be the standard coordinates of \mathbb{R}^3 and

$$(7) \quad X_l = \{x = l + \frac{1}{2}\}, \quad Y_m = \{y = m + \frac{1}{2}\}, \quad Z_n = \{z = n + \frac{1}{2}\},$$

be planes forming a cubic lattice in \mathbb{R}^3 , where l, m, n run over all integers. Introduce a stack of D4-branes filling \mathbb{R}^3 and place an NS5-brane on each of the planes (7). These branes also extend in additional directions in ten dimensions, especially a two-dimensional Minkowski subspace (on which the brane box model emerges).

The brane box model is a two-dimensional $\mathcal{N} = (0, 2)$ supersymmetric gauge theory that describes the low-energy dynamics of D4-branes diced into cubes by NS5-branes as above. The gauge group of the brane box model is the product

$$(8) \quad \prod_{(l,m,n) \in \mathbb{Z}^3} \mathrm{U}(N)_{(l,m,n)},$$

where N is the number of D4-branes. The $U(N)$ factor $U(N)_{(l,m,n)}$ arises from the piece of the stack of N D4-branes that contains the point (l, m, n) .

The brane box model has a $U(1)$ global symmetry $U(1)_R$, called an *R-symmetry*, under which the two supercharges of $\mathcal{N} = (0, 2)$ supersymmetry have charge $+1$ and -1 . The definition of $U(1)_R$ is not unique since the R-charge R generating $U(1)_R$ can be shifted by a linear combination of generators of $U(1)$ flavor symmetries. For example, there are $U(1)$ flavor symmetries $U(1)_{X_l}$, $U(1)_{Y_m}$, $U(1)_{Z_n}$ associated with X_l , Y_m , Z_n , respectively, so we can assign real numbers ξ_l , η_m , ζ_n to X_l , Y_m , Z_n and set

$$(9) \quad R = R_0 + \sum_l \xi_l X_l + \sum_m \eta_m Y_m + \sum_n \zeta_n Z_n.$$

Here R_0 is the old R-charge and X_l , Y_m , Z_n are generators of $U(1)_{X_l}$, $U(1)_{Y_m}$, $U(1)_{Z_n}$. The new R-charge generates a $U(1)$ global symmetry that transforms the supercharges in the same way as before and hence still deserves to be called an R-symmetry. We will see that parameters assigned to planes, similar to those just introduced, make an appearance as spectral parameters in the tetrahedron duality.

In order to describe the matter content of the brane box model, let us introduce a couple of notations. For $a, b \in \mathbb{Z}^3$, we write

$$(10) \quad a \xrightarrow{r} b$$

and

$$(11) \quad a \xrightarrow{r} b$$

to denote, respectively, a chiral multiplet and a Fermi multiplet with $R = r$ in the bifundamental representation $(\overline{\mathbf{N}}, \mathbf{N})$ of $U(N)_a \times U(N)_b$. There is an equivalence

$$(12) \quad a \xrightarrow{r} b \cong b \xrightarrow{-r} a,$$

which is a general property of $\mathcal{N} = (0, 2)$ supersymmetric field theories.

For the choice (9) of the R-charge, with $R_0 = \frac{1}{2}$ for chiral multiplets and 0 for Fermi multiplets, the list of matter multiplets of the brane box model is the following:

$$(13) \quad (l, m, n) \xrightarrow{\frac{1}{2} + \xi_l} (l + 1, m, n),$$

$$(14) \quad (l, m, n) \xrightarrow{\frac{1}{2} + \eta_m} (l, m + 1, n),$$

$$(15) \quad (l, m, n) \xrightarrow{\frac{1}{2} + \zeta_n} (l, m, n + 1),$$

$$(16) \quad (l, m, n) \xleftarrow{\frac{1}{2} - \xi_l - \eta_m - \zeta_n} (l + 1, m + 1, n + 1),$$

$$(17) \quad (l, m, n) \xrightarrow{\eta_m + \zeta_n} (l, m + 1, n + 1),$$

$$(18) \quad (l, m, n) \xrightarrow{\xi_l + \zeta_n} (l + 1, m, n + 1),$$

$$(19) \quad (l, m, n) \xrightarrow{\xi_l + \eta_m} (l + 1, m + 1, n).$$

A matter multiplet has charge ± 1 under $U(1)_W$, $W = X_l, Y_m, Z_n$, if the arrow intersects the corresponding plane W . The sign of the charge is correlated with the direction of intersection.

The matter fields interact with one another via coupling with a vector multiplet for the gauge group and through a superpotential. There is a cubic term in the superpotential for each triangle consisting of two solid arrows and one dashed arrow with the sum of R-charges equal to one. The superpotential is important data of the brane box model, but we will only be interested in the infrared effective theory in which all superpotential terms allowed by the symmetries of the model are expected to be generated. Thus, in our discussions the ultraviolet superpotential merely plays the role of a constraint for the possible charge assignments for $U(1)_R$ and flavor symmetries, and we will suppress it in what follows.

Reflecting the lattice structure of the string theory construction, the brane box model has an interpretation as a spin model on a cubic lattice, as described in section 1. For a fixed $(l, m, n) \in \mathbb{Z}^3$, the matter multiplets (13)–(19) and the eight $U(N)$ groups connected by them comprise the natural unit cell shown in Figure 3(b).

2.2. Cancellation of abelian gauge anomaly. The matter content of the brane box model described above leads to anomaly in the abelian part of the gauge symmetry. Let us explain one way of cancelling this anomaly. See e.g. [7, 25, 26] for detailed discussions.

Let Σ be the spacetime of the brane box model and $U(1)_a$ be the overall $U(1)$ subgroup of $U(N)_a$. Under the gauge transformation by

$$(20) \quad (e^{i\Lambda_a})_{a \in \mathbb{Z}^3} : \Sigma \rightarrow \prod_{a \in \mathbb{Z}^3} U(1)_a,$$

the path integral measure is multiplied by the phase factor

$$(21) \quad \exp\left(i \sum_{a, b \in \mathbb{Z}^3} \frac{\mathcal{A}^{ab}}{4\pi} \int_{\Sigma} \Lambda_a F_b\right),$$

where F_b is the field strength of the gauge field for $U(1)_b$ and

$$(22) \quad \mathcal{A}^{ab} = \begin{cases} 2 & a = b, \\ -1 & a \rightarrow b \text{ or } b \rightarrow a, \\ 1 & a \dashrightarrow b \text{ or } b \dashrightarrow a, \\ 0 & \text{otherwise.} \end{cases}$$

To cancel this phase factor, we introduce a chiral multiplet Φ_a that is valued in $\mathbb{C}/2\pi i\mathbb{Z}$ and transforms under the gauge transformation (20) as

$$(23) \quad \Phi_a \rightarrow \Phi_a + i\Lambda_a.$$

Then, the gauge transformation of

$$(24) \quad - \sum_{a, b \in \mathbb{Z}^3} \frac{\mathcal{A}^{ab}}{4\pi} \int \text{Im } \phi_a F_b$$

cancels the anomalous variation of the path integral measure, where ϕ_a is the scalar component of Φ_a . We add the supersymmetric completion of this term to the action functional of the brane box model. Since each nonzero summand is associated with a spin site (if $a = b$) or an arrow (if $a \neq b$), the added term can be given a lattice model interpretation.

We expect that the fields $\text{Im } \phi_a$ descend from the Ramond–Ramond 3-form field $C^{(3)}$ in type IIA string theory. For a stack of D4-branes carrying a gauge field with field strength F , the worldvolume action is given by $\int_{\mathbb{R}^{4,1}} C^{(3)} \wedge \text{Tr } F$.

2.3. The oblique unit cell and the tetrahedron duality. The choice of R-charge (9) is adapted to the set of planes $\{X_l, Y_m, Z_n\}_{l,m,n \in \mathbb{Z}}$ in the sense that its parametrization is generated by the flavor symmetries associated with these planes. The R-charge assignment relevant for the tetrahedron duality is adapted to another set of planes $\{X'_l, Y'_m, Z'_n\}_{l,m,n \in \mathbb{Z}}$.

Let (x', y', z') be the coordinates of \mathbb{R}^3 related to (x, y, z) by

$$(25) \quad x' = x, \quad y' = -y + z, \quad z' = z.$$

Then, this set of planes is defined by

$$(26) \quad X'_l = \{x' = l + \frac{1}{2}\}, \quad Y'_m = \{y' = m + \frac{1}{2}\}, \quad Z'_n = \{z' = n + \frac{1}{2}\}.$$

Associated with X'_l, Y'_m, Z'_n there are $U(1)$ flavor symmetries $U(1)_{X'_l}, U(1)_{Y'_m}, U(1)_{Z'_n}$, generated by charges X'_l, Y'_m, Z'_n . We choose the R-charge to be

$$(27) \quad R = R_0 + \sum_l \left(s_l + \frac{1}{2}\right) X'_l + \sum_m \left(t_m - \frac{1}{2}\right) Y'_m - \sum_n u_n Z'_n.$$

In terms of the new coordinates and R-charge, the matter content of the brane box model is

$$(28) \quad (l, m, n) \xrightarrow{1+s_l} (l+1, m, n),$$

$$(29) \quad (l+1, m+1, n+1) \xrightarrow{1-t_m} (l+1, m, n+1),$$

$$(30) \quad (l, m, n) \xrightarrow{t_m - u_n} (l, m+1, n+1),$$

$$(31) \quad (l, m, n) \xleftarrow{-s_l + u_n} (l+1, m, n+1),$$

$$(32) \quad (l+1, m, n) \xrightarrow{-u_n} (l+1, m, n+1),$$

$$(33) \quad (l, m, n) \xrightarrow{s_l + t_m - u_n} (l+1, m+1, n+1),$$

$$(34) \quad (l, m+1, n+1) \xrightarrow{1+s_l - t_m} (l+1, m, n+1).$$

The above list of matter multiplets defines the oblique unit cell of the brane box model:⁵

$$(35) \quad (a|e, f, g|b, c, d|h; s, t, u) = \left(\begin{array}{c} (a \xrightarrow{t-u} b)(b \xrightarrow{1+s-t} c)(c \xrightarrow{-s+u} a) \\ (a \xrightarrow{s+t-u} h)(h \xrightarrow{1-t} c)(a \xrightarrow{1+s} e)(e \xrightarrow{-u} c) \end{array} \right).$$

See Figure 4. By the right-hand side we mean an $\mathcal{N} = (0, 2)$ supersymmetric field theory consisting of four chiral multiplets and three Fermi multiplets, each of which transforms in a bifundamental representation under two of eight $U(N)$ flavor symmetry groups $U(N)_a, U(N)_b, \dots, U(N)_h$. We emphasize that these $U(N)$ groups are not gauged; only when multiple copies of the oblique unit cell are combined, the $U(N)$ groups at the overlapping vertices are gauged. The oblique unit cell also has a cubic superpotential, but otherwise its matter fields do not interact with one another.

Now we argue that the oblique unit cell satisfies the tetrahedron duality.

⁵The theory defined by the first row of the oblique unit cell (35) also satisfies the tetrahedron duality, but the corresponding lattice is the disjoint union of infinitely many copies of a two-dimensional lattice.




FIGURE 4. The oblique unit cell of the brane box model.

Our argument relies on two infrared equivalences. One is the (0,2) triality, as already highlighted in section 1. The other is

$$(36) \quad (a \xrightarrow{r} b)(b \xrightarrow{1-r} a) \simeq 1.$$

Here 1 means the trivial multiplet (that is to say, no fields at all). This relation holds because in any theory that contains the two multiplets on the left-hand side, a quadratic term given by their product can be generated in the superpotential. Such a superpotential term makes the pair of multiplets massive and disappear at low energies.

Plugging the definition (35) of the oblique unit cell into the tetrahedron duality (4), we find that the left-hand side is given by

$$(37) \quad \text{Gauge}_d \left(\begin{array}{l} (a_4 \xrightarrow{s_3-s_2} b_1)_1 (b_1 \xrightarrow{1+s_1-s_3} b_3)_2 (b_3 \xrightarrow{-s_1+s_2} a_4)_3 \\ (a_4 \xrightarrow{s_1+s_3-s_2} d)(d \xrightarrow{1-s_3} b_3)_4 (a_4 \xrightarrow{1+s_1} c_2)_5 (c_2 \xrightarrow{-s_2} b_3)_6 \\ (c_1 \xrightarrow{s_4-s_2} c_4)(c_4 \xrightarrow{1+s_1-s_4} d)(d \xrightarrow{-s_1+s_2} c_1) \\ (c_1 \xrightarrow{s_1+s_4-s_2} b_4)(b_4 \xrightarrow{1-s_4} d)(c_1 \xrightarrow{1+s_1} b_2)_7 (b_2 \xrightarrow{-s_2} d)_8 \\ (b_1 \xrightarrow{s_4-s_3} a_2)_9 (a_2 \xrightarrow{1+s_1-s_4} b_3)_{10} (b_3 \xrightarrow{-s_1+s_3} b_1)_2 \\ (b_1 \xrightarrow{s_1+s_4-s_3} c_5)(c_5 \xrightarrow{1-s_4} b_3)_{11} (b_1 \xrightarrow{1+s_1} d)(d \xrightarrow{-s_3} b_3)_4 \\ (d \xrightarrow{s_4-s_3} c_5)(c_5 \xrightarrow{1+s_2-s_4} c_2)(c_2 \xrightarrow{-s_2+s_3} d) \\ (d \xrightarrow{s_2+s_4-s_3} a_1)(a_1 \xrightarrow{1-s_4} c_2)_{12} (d \xrightarrow{1+s_2} b_2)_8 (b_2 \xrightarrow{-s_3} c_2)_{13} \end{array} \right),$$

whereas the right-hand side is

$$(38) \quad \text{Gauge}_d \left(\begin{array}{l} (b_1 \xrightarrow{s_4-s_3} a_2)_9 (a_2 \xrightarrow{1+s_2-s_4} a_4)_{14} (a_4 \xrightarrow{-s_2+s_3} b_1)_1 \\ (b_1 \xrightarrow{s_2+s_4-s_3} d)(d \xrightarrow{1-s_4} a_4)(b_1 \xrightarrow{1+s_2} c_1)(c_1 \xrightarrow{-s_3} a_4) \\ (c_1 \xrightarrow{s_4-s_3} d)(d \xrightarrow{1+s_1-s_4} c_2)(c_2 \xrightarrow{-s_1+s_3} c_1) \\ (c_1 \xrightarrow{s_1+s_4-s_3} a_1)(a_1 \xrightarrow{1-s_4} c_2)_{12} (c_1 \xrightarrow{1+s_1} b_2)_7 (b_2 \xrightarrow{-s_3} c_2)_{13} \\ (a_4 \xrightarrow{s_4-s_2} a_2)_{14} (a_2 \xrightarrow{1+s_1-s_4} b_3)_{10} (b_3 \xrightarrow{-s_1+s_2} a_4)_3 \\ (a_4 \xrightarrow{s_1+s_4-s_2} c_5)(c_5 \xrightarrow{1-s_4} b_3)_{11} (a_4 \xrightarrow{1+s_1} c_2)_5 (c_2 \xrightarrow{-s_2} b_3)_6 \\ (d \xrightarrow{s_3-s_2} c_4)(c_4 \xrightarrow{1+s_1-s_3} c_5)(c_5 \xrightarrow{-s_1+s_2} d) \\ (d \xrightarrow{s_1+s_3-s_2} b_4)(b_4 \xrightarrow{1-s_3} c_5)(d \xrightarrow{1+s_1} a_1)(a_1 \xrightarrow{-s_2} c_5) \end{array} \right).$$

Among the 56 arrows that appear in the tetrahedron duality, 14 pairs (underlined and numbered above) cancel against each other by virtue of the equivalences (12) and (36). After the cancellations we have

$$(39) \quad \text{Gauge}_d \left(\begin{array}{l} (d \xrightarrow{-s_1+s_2} c_1)(d \xrightarrow{s_4-s_3} c_5) \\ (d \xrightarrow{s_2+s_4-s_3} a_1)(a_4 \xrightarrow{s_1+s_3-s_2} d)(c_4 \xrightarrow{1+s_1-s_4} d) \\ (b_1 \xrightarrow{1+s_1} d)(b_4 \xrightarrow{1-s_4} d)(c_2 \xrightarrow{-s_2+s_3} d) \\ (c_1 \xrightarrow{s_1+s_4-s_2} b_4)(c_1 \xrightarrow{s_4-s_2} c_4)(b_1 \xrightarrow{s_1+s_4-s_3} c_5)(c_5 \xrightarrow{1+s_2-s_4} c_2) \end{array} \right)$$

and

$$(40) \quad \text{Gauge}_d \left(\begin{array}{l} (c_1 \xrightarrow{s_4-s_3} d)(c_5 \xrightarrow{-s_1+s_2} d) \\ (d \xrightarrow{1+s_1} a_1)(d \xrightarrow{1-s_4} a_4)(d \xrightarrow{s_3-s_2} c_4) \\ (b_1 \xrightarrow{s_2+s_4-s_3} d)(d \xrightarrow{s_1+s_3-s_2} b_4)(d \xrightarrow{1+s_1-s_4} c_2) \\ (b_1 \xrightarrow{1+s_2} c_1)(c_1 \xrightarrow{-s_3} a_4)(c_2 \xrightarrow{-s_1+s_3} c_1)(c_1 \xrightarrow{s_1+s_4-s_3} a_1) \\ (a_4 \xrightarrow{s_1+s_4-s_2} c_5)(c_4 \xrightarrow{1+s_1-s_3} c_5)(b_4 \xrightarrow{1-s_3} c_5)(a_1 \xrightarrow{-s_2} c_5) \end{array} \right)$$

for the left-hand side and the right-hand side. In this step we have used the equivalence (12) once (when cancelling pair #4).

Further applying the equivalence (12) to the six underlined arrows and moving some arrows across the two sides using the equivalence (36), we arrive at the infrared duality

$$(41) \quad \text{Gauge}_d \left(\begin{array}{l} (d \xrightarrow{-s_1+s_2} c_1)(d \xrightarrow{s_4-s_3} c_5) \\ (d \xrightarrow{s_2+s_4-s_3} a_1)(d \xrightarrow{-s_1-s_3+s_2} a_4)(d \xrightarrow{-1-s_1+s_4} c_4) \\ (b_1 \xrightarrow{1+s_1} d)(b_4 \xrightarrow{1-s_4} d)(c_2 \xrightarrow{-s_2+s_3} d) \\ (c_1 \xrightarrow{-s_2} b_1)(c_1 \xrightarrow{s_1+s_4-s_2} b_4)(c_1 \xrightarrow{1+s_1-s_3} c_2) \\ (c_5 \xrightarrow{-s_1-s_4+s_3} b_1)(c_5 \xrightarrow{s_3} b_4)(c_5 \xrightarrow{1+s_2-s_4} c_2) \end{array} \right) \\ \simeq \text{Gauge}_d \left(\begin{array}{l} (c_1 \xrightarrow{s_4-s_3} d)(c_5 \xrightarrow{-s_1+s_2} d) \\ (d \xrightarrow{1+s_1} a_1)(d \xrightarrow{1-s_4} a_4)(d \xrightarrow{s_3-s_2} c_4) \\ (d \xrightarrow{-s_2-s_4+s_3} b_1)(d \xrightarrow{s_1+s_3-s_2} b_4)(d \xrightarrow{1+s_1-s_4} c_2) \\ (a_1 \xrightarrow{-s_1-s_4+s_3} c_1)(a_4 \xrightarrow{s_3} c_1)(c_4 \xrightarrow{1-s_4+s_2} c_1) \\ (a_1 \xrightarrow{-s_2} c_5)(a_4 \xrightarrow{s_1+s_4-s_2} c_5)(c_4 \xrightarrow{1+s_1-s_3} c_5) \end{array} \right).$$

The two sides of this relation are two of a triplet of $\mathcal{N} = (0, 2)$ supersymmetric gauge theories related by the $(0, 2)$ triality. Indeed, they are the theories $T_{3N, 2N, 3N}$ and $T_{2N, 3N, 3N}$ in the notation of Appendix A, except that a different mechanism is employed for the cancellation of abelian gauge anomaly. Therefore, the tetrahedron duality holds under the assumption that the $(0, 2)$ triality is not broken by the anomaly cancellation mechanism of the brane box model.

2.4. Spectral parameters and boundary conditions. An elliptic genus of the oblique unit cell (or any unit cell for that matter) defines a Boltzmann weight for a lattice model whose partition function is equal to the elliptic genus of the brane box model. In order to make the partition function finite, one usually makes the lattice finite by some means. For example, we can impose periodic boundary conditions on the lattice, placing it in a 3-torus \mathbb{T}^3 rather than \mathbb{R}^3 . Let us take the size of the resulting lattice to be $L \times M \times N$.

The partition function of the lattice model is a function of the spectral parameters, which are assigned to the planes (26) and appear in the definition of the R-charge (27). Although the total number of the spectral parameters is $L + M + N$, the partition function actually depends only on three linear combinations of them because of gauge symmetry.

For any oriented closed surface S in \mathbb{T}^3 that does not intersect with the integral points, there is an associated $U(1)$ symmetry $U(1)_S$ in the brane box model, defined in a similar manner to $U(1)_{X'_l}$, $U(1)_{Y'_m}$, $U(1)_{Z'_n}$. If S is the boundary of a three-dimensional region V of \mathbb{T}^3 , then $U(1)_S$ is a gauge symmetry: for $S = \partial V$ with orientation given by the normal vectors pointing outward,

$$(42) \quad U(1)_S = \prod_{(l,m,n) \in V} U(1)_{(l,m,n)}.$$

Thus, the number of independent $U(1)$ flavor symmetries is given by the rank of the second homology group of \mathbb{T}^3 , which is 3. Without loss of generality, we can set all spectral parameters but those assigned to X'_0 , Y'_0 , Z'_0 to zero. The three remaining parameters s_0 , t_0 , u_0 may be interpreted as specifying twisted boundary conditions in the x' , y' , z' directions, respectively.

3. A SOLUTION BY THE 1D BRANE BOX MODEL

Dimensional reduction of the brane box model yields the 1D brane box model, an $\mathcal{N} = 2$ supersymmetric gauge theory that has the same gauge group and matter content as its parent theory (except that the chiral and Fermi multiplets are, of course, those for $\mathcal{N} = 2$ supersymmetry in one dimension). If the dimensional reduction commutes with the triality, then the oblique unit cell for the 1D brane box model also satisfies the tetrahedron duality and its Witten index, computed by a partition function on a circle, is a solution of the tetrahedron equation.

Crucially, the abelian gauge anomaly disappears after the dimensional reduction since chiral anomalies are absent in one dimension. Thus, we can write down the Witten index by the dimensional reduction of an integral formula for the relevant elliptic genus [5, 6]. The formula shows that the Witten index of the 1D brane box model is given by the partition function of an Ising-type spin model with continuous spin variables and pairwise interaction. Below we describe this lattice model and verify that its Boltzmann weight indeed solves the tetrahedron equation.

3.1. The lattice model. The spin model in question is defined on a simple cubic lattice. At each point $a \in \mathbb{Z}^3$ a set of N periodic spin variables

$$(43) \quad a = (a_i)_{i=1}^N \in \mathbb{T}^N$$

is placed, where $\mathbb{T} \simeq \mathbb{R}/2\pi\mathbb{Z}$ is the unit circle in the complex plane. We will use the same symbol to refer to both a spin site and the set of spin variables located there.

The interactions between spin variables at neighboring sites come in two types and are represented by solid arrows and dashed arrows in the list (28)–(34). The Boltzmann weights for these interactions are given by⁶

$$(44) \quad W(a \xrightarrow{r} b) = \prod_{i,j} -\frac{1}{\phi(e^{4\pi i r} b_i/a_j)},$$

$$(45) \quad W(a \xrightarrow{-r} b) = \prod_{i,j} \phi(e^{4\pi i r} b_i/a_j),$$

where

$$(46) \quad \phi(z) = z^{1/2} - z^{-1/2}.$$

The number r above an arrow is a coupling constant controlling the interaction. We take $r \in \mathbb{R}$.

Since $\phi(z) = -\phi(1/z)$, the equivalence (12) becomes an equality up to a sign for the Boltzmann weights:

$$(47) \quad W(a \xrightarrow{-r} b) = (-1)^N W(b \xrightarrow{-r} a).$$

The cancellation relation (36) holds without a sign:

$$(48) \quad W(a \xrightarrow{r} b) W(b \xrightarrow{1-r} a) = 1.$$

The partition function of the lattice model is the integral over all spin variables of the product of all local Boltzmann weights. The integration over spin variables $a \in \mathbb{T}^N$ is performed with respect to the measure

$$(49) \quad \mu(a) = \frac{1}{N!} \prod_i \frac{da_i}{2\pi i a_i} \prod_{i \neq j} \phi\left(\frac{a_i}{a_j}\right).$$

The factor $\prod_{i \neq j} \phi(a_i/a_j)$ may be thought as the Boltzmann weight for self-interaction among $(a_i)_{i=1}^N$.

In performing this integration we encounter poles coming from factors of the form $W(b \xrightarrow{r} a)$ and factors of the form $W(a \xrightarrow{r} b)$. We agree to deform the domain of integration in such a way that the compact region bounded by the domain contains all poles from the former but excludes any poles from the latter. Alternatively, we may modify the definition of the Boltzmann weight (44) by multiplying the argument of the function ϕ by e^ϵ and later take the limit $\epsilon \rightarrow +0$.

3.2. The tetrahedron equation. The tetrahedron equation for the oblique unit cell of this lattice model simplifies to the following equality, obtained by evaluating

⁶Note that only the non-integer part of r appears in the Boltzmann weights. In the language of the Witten index, $e^{4\pi i r}$ is the product of the flavor fugacities for $U(1)_{X'_l}$, $U(1)_{Y'_m}$, $U(1)_{Z'_n}$.

the Witten indices of the two sides of the equivalence relation (41):

$$(50) \quad \int_d \mu(d) \left(\begin{array}{l} W(d \xrightarrow{-s_1+s_2} c_1) W(d \xrightarrow{s_4-s_3} c_5) \\ \times W(d \xrightarrow{s_2+s_4-s_3} a_1) W(d \xrightarrow{-s_1-s_3+s_2} a_4) W(d \xrightarrow{-1-s_1+s_4} c_4) \\ \times W(b_1 \xrightarrow{1+s_1} d) W(b_4 \xrightarrow{1-s_4} d) W(c_2 \xrightarrow{-s_2+s_3} d) \\ \times W(c_1 \xrightarrow{-s_2} b_1) W(c_1 \xrightarrow{s_1+s_4-s_2} b_4) W(c_1 \xrightarrow{1+s_1-s_3} c_2) \\ \times W(c_5 \xrightarrow{-s_1-s_4+s_3} b_1) W(c_5 \xrightarrow{s_3} b_4) W(c_5 \xrightarrow{1+s_2-s_4} c_2) \end{array} \right) \\ = (-1)^N \int_d \mu(d) \left(\begin{array}{l} W(c_1 \xrightarrow{s_4-s_3} d) W(c_5 \xrightarrow{-s_1+s_2} d) \\ \times W(d \xrightarrow{1+s_1} a_1) W(d \xrightarrow{1-s_4} a_4) W(d \xrightarrow{s_3-s_2} c_4) \\ \times W(d \xrightarrow{-s_2-s_4+s_3} b_1) W(d \xrightarrow{s_1+s_3-s_2} b_4) W(d \xrightarrow{1+s_1-s_4} c_2) \\ \times W(a_1 \xrightarrow{-s_1-s_4+s_3} c_1) W(a_4 \xrightarrow{s_3} c_1) W(c_4 \xrightarrow{1-s_4+s_2} c_1) \\ \times W(a_1 \xrightarrow{-s_2} c_5) W(a_4 \xrightarrow{s_1+s_4-s_2} c_5) W(c_4 \xrightarrow{1+s_1-s_3} c_5) \end{array} \right).$$

The sign $(-1)^N$ comes from the use of the relation (12) seven times in the simplification of the tetrahedron duality.

Let N_1, N_2, N_3, n_2, n_3 be nonnegative integers such that

$$(51) \quad n_2 = \frac{1}{2}(N_1 - N_2 + N_3), \quad n_3 = \frac{1}{2}(N_1 + N_2 - N_3).$$

Then, we have the following equality

$$(52) \quad \frac{(-1)^{n_3 N_2}}{n_3!} \int_{\mathcal{D}} \prod_i \frac{dz_i}{2\pi i z_i} \prod_{i \neq j} \phi\left(\frac{z_i}{z_j}\right) \frac{\prod_{i,\gamma} \phi(\tau_\gamma w_\gamma / z_i) \prod_{\alpha,\beta} \phi(u_\alpha / \rho_\alpha \sigma_\beta v_\beta)}{\prod_{i,\alpha} \phi(\rho_\alpha z_i / u_\alpha) \prod_{i,\beta} \phi(\sigma_\beta v_\beta / z_i)} \\ = \frac{(-1)^{n_2(N_2+N_3)}}{n_2!} \int_{\mathcal{D}'} \prod_{\bar{i}} \frac{dz'_{\bar{i}}}{2\pi i z'_{\bar{i}}} \prod_{\bar{i} \neq \bar{j}} \phi\left(\frac{z'_{\bar{i}}}{z'_{\bar{j}}}\right) \frac{\prod_{\bar{i},\beta} \phi(\sigma'_\beta v_\beta / z'_{\bar{i}}) \prod_{\gamma,\alpha} \phi(w_\gamma / \tau'_\gamma \rho'_\alpha u_\alpha)}{\prod_{\bar{i},\gamma} \phi(\tau'_\gamma z'_{\bar{i}} / w_\gamma) \prod_{\bar{i},\alpha} \phi(\rho'_\alpha u_\alpha / z'_{\bar{i}})}.$$

Here

$$(53) \quad (u_\alpha, \rho_\alpha)_{\alpha=1}^{N_1} \in \mathbb{T}^{2N_1}, \quad (v_\beta, \sigma_\beta)_{\beta=1}^{N_2} \in \mathbb{T}^{2N_2}, \quad (w_\gamma, \tau_\gamma)_{\gamma=1}^{N_3} \in \mathbb{T}^{2N_3},$$

are arbitrary parameters and

$$(54) \quad \rho'_\alpha = e^{4\pi i \Delta} \rho_\alpha^{-1}, \quad \sigma'_\beta = e^{4\pi i(-1+\Delta)} \sigma_\beta, \quad \tau'_\gamma = e^{4\pi i(1-\Delta)} \tau_\gamma^{-1},$$

with $\Delta \in \mathbb{R}$ arbitrary and fixed. For each variable $z_i, i = 1, \dots, n_3$, the integrand of the left-hand side is a meromorphic differential on \mathbb{CP}^1 which have no poles at $z_i = 0$ and ∞ ; all poles come from the factors $\prod_{i,\alpha} \phi(\rho_\alpha z_i / u_\alpha)^{-1}$ and $\prod_{i,\beta} \phi(\sigma_\beta v_\beta / z_i)^{-1}$. The domain \mathcal{D} bounds a region that contains 0 and all poles from the former factor but no poles from the latter. Similarly, for each variable $z'_{\bar{i}}, \bar{i} = 1, \dots, n_2$, the integrand of the right-hand side is a meromorphic differential on \mathbb{CP}^1 with no poles at $z'_{\bar{i}} = 0$ and ∞ , and \mathcal{D}' bounds a region that contains all poles from the factor $\prod_{\bar{i},\gamma} \phi(\tau'_\gamma z'_{\bar{i}} / w_\gamma)^{-1}$ but no poles from $\prod_{\bar{i},\alpha} \phi(\rho'_\alpha u_\alpha / z'_{\bar{i}})^{-1}$.

Equation (50) is a special case of (52), hence the oblique unit cell is a solution of the tetrahedron equation. Indeed, if we take $N_1 = 2N, N_2 = N_3 = 3N$ (for which

$n_2 = n_3 = N$),

$$(55) \quad (u_\alpha)_{\alpha=1}^{N_1} = (c_1, c_5), \quad \rho_\alpha = \begin{cases} e^{4\pi i(s_4 - s_3)} & (1 \leq \alpha \leq N), \\ e^{4\pi i(-s_1 + s_2)} & (N + 1 \leq \alpha \leq 2N), \end{cases}$$

$$(56) \quad (v_\beta)_{\beta=1}^{N_2} = (a_1, a_4, c_4), \quad \sigma_\beta = \begin{cases} e^{4\pi i(1 + s_1)} & (1 \leq \alpha \leq N), \\ e^{4\pi i(1 - s_4)} & (N + 1 \leq \alpha \leq 2N), \\ e^{4\pi i(s_3 - s_2)} & (2N + 1 \leq \alpha \leq 3N), \end{cases}$$

$$(57) \quad (w_\gamma)_{\gamma=1}^{N_3} = (b_1, b_4, c_2), \quad \tau_\gamma = \begin{cases} e^{4\pi i(-s_2 - s_4 + s_3)} & (1 \leq \alpha \leq N), \\ e^{4\pi i(s_1 + s_3 - s_2)} & (N + 1 \leq \alpha \leq 2N), \\ e^{4\pi i(1 + s_1 - s_4)} & (2N + 1 \leq \alpha \leq 3N), \end{cases}$$

and

$$(58) \quad \Delta = s_4 - s_3 - s_1 + s_2,$$

then the left-hand side and the right-hand of (52) become the right-hand side and the left-hand side of (50), respectively.

Equation (52) can be demonstrated in a way similar to the proof of the equality between the elliptic genera of theories related by the (0, 2) triality, treated in detail in Appendix A. Using

$$(59) \quad \operatorname{Res}_{z=w} \frac{1}{z\phi(z/w)} = 1,$$

we find that the left-hand side evaluates to

$$(60) \quad \sum_{\substack{\mathfrak{S} \subset \{1, \dots, N_1\} \\ |\mathfrak{S}| = n_3}} \frac{\prod_{i \in \mathfrak{S}} \prod_\gamma \phi(\tau_\gamma \rho_i w_\gamma / u_i) \prod_{\bar{i} \in \bar{\mathfrak{S}}} \prod_\beta \phi(u_{\bar{i}} / \rho_{\bar{i}} \sigma_\beta v_\beta)}{\prod_{i \in \mathfrak{S}} \prod_{\bar{i} \in \bar{\mathfrak{S}}} \phi(\rho_{\bar{i}} u_i / \rho_i u_{\bar{i}})},$$

where $\bar{\mathfrak{S}} = \{1, \dots, N_1\} \setminus \mathfrak{S}$. For the right-hand side, we can multiply it by $(-1)^{n_2}$ and sum over the residues at the poles from the factor $\prod_{\bar{i}, \alpha} \phi(\rho'_\alpha u_\alpha / z'_i)^{-1}$ since the sum of the residues of a meromorphic differential on any compact Riemann surface is zero. The sign factor $(-1)^{n_2}$ just introduced is cancelled by the same factor coming from the residues, and we get

$$(61) \quad \sum_{\substack{\bar{\mathfrak{S}} \subset \{1, \dots, N_1\} \\ |\bar{\mathfrak{S}}| = n_2}} \frac{\prod_{\bar{i} \in \bar{\mathfrak{S}}} \prod_\beta \phi(\rho'_{\bar{i}} u_{\bar{i}} / \sigma'_\beta v_\beta) \prod_{i \in \mathfrak{S}} \prod_\gamma \phi(w_\gamma / \tau'_\gamma \rho'_i u_i)}{\prod_{i \in \mathfrak{S}} \prod_{\bar{i} \in \bar{\mathfrak{S}}} \phi(\rho'_i u_i / \rho_i u_{\bar{i}})},$$

with $\mathfrak{S} = \{1, \dots, N_1\} \setminus \bar{\mathfrak{S}}$. Looking at the relation (54) we see that the above two sums are equal.

4. A SOLUTION BY THE (0, 1) BRANE BOX MODEL

$\mathcal{N} = (0, 1)$ supersymmetric gauge theories with special orthogonal gauge groups are also known to exhibit triality [19]. Since this triality has the same combinatorial structure as its (0, 2) counterpart, we obtain another solution of the tetrahedron duality by mere reinterpretation of notations that turns the brane box model into the relevant $\mathcal{N} = (0, 1)$ supersymmetric gauge theory, which we call the (0, 1) brane box model.

For the (0, 1) brane box model, each point $a \in \mathbb{Z}^3$ represents an $\mathrm{SO}(2N)$ gauge group $\mathrm{SO}(2N)_a$, accompanied by a Fermi multiplet in the symmetric representation

of $\mathrm{SO}(2N)_a$. A solid arrow $a \rightarrow b$ and a dashed arrow $a \dashrightarrow b$ are a chiral multiplet and a Fermi multiplet in the bivector representation of $\mathrm{SO}(2N)_a \times \mathrm{SO}(2N)_b$, respectively.⁷ All of these multiplets are those for $\mathcal{N} = (0, 1)$ supersymmetry. In particular, there is no R-symmetry and hence no number (R-charge) assigned to arrows. The $(0, 1)$ brane box model is free of gauge anomaly without the need of introducing additional matter multiplets.

Although no formula is known for the elliptic genera of $\mathcal{N} = (0, 1)$ supersymmetric gauge theories in general, in the case relevant to the $(0, 1)$ triality the elliptic genus in the Ramond–Ramond sector has been computed in the low-energy effective sigma model description [19]. It turns out that the result can be expressed as an integral akin to integral formulas for the elliptic genera of $\mathcal{N} = (0, 2)$ supersymmetric gauge theories. Based on this observation we propose that the elliptic genus of the $(0, 1)$ brane box model is equal to the partition function of a lattice model defined by a solution of the tetrahedron equation, which we now describe.

4.1. The lattice model. Fix a complex number q with $|q| < 1$. Let

$$(62) \quad \eta = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$$

be the Dedekind eta function and

$$(63) \quad \hat{\theta}(z) = i \frac{\theta_1(z)}{\eta} = q^{1/12} (z^{1/2} - z^{-1/2}) \prod_{n=1}^{\infty} (1 - zq^n)(1 - z^{-1}q^n),$$

where θ_1 is the Jacobi theta function. Note that $q^{-1/12} \hat{\theta}(z) \rightarrow \phi(z)$ as $q \rightarrow 0$; this limit corresponds to dimensional reduction. We have $\hat{\theta}(z) = -\hat{\theta}(1/z)$ and

$$(64) \quad \hat{\theta}(qz) = -q^{-1/2} z^{-1} \hat{\theta}(z).$$

The lattice model arising from the elliptic genus of the $(0, 1)$ brane box model is much like the one for the 1D brane box model described in section 3.1. A set of continuous spin variables $a = (a_i)_{i=1}^N$ resides at $a \in \mathbb{Z}^3$. They interact according to the list (28)–(34) (with the numbers above the arrows ignored). The Boltzmann weights for these interactions are

$$(65) \quad W(a \rightarrow b) = \prod_{i,j} -\frac{1}{\hat{\theta}(b_i a_j^{\pm 1})},$$

$$(66) \quad W(a \dashrightarrow b) = \prod_{i,j} \hat{\theta}(b_i a_j^{\pm 1})$$

and satisfy the relations (47) and (48). Here we have introduced the shorthand notation $\hat{\theta}(b_i a_j^{\pm 1}) = \hat{\theta}(b_i a_j) \hat{\theta}(b_i a_j^{-1})$. The integration over spin variables a is performed with respect to the measure

$$(67) \quad \mu(a) = \frac{1}{2^{N-1} N!} \prod_i \frac{da_i}{2\pi i a_i} \eta^2 \hat{\theta}(a_i^2) \prod_{i \neq j} \hat{\theta}(a_i a_j^{\pm 1}).$$

The numerical factor $2^{N-1} N!$ is the order of the Weyl group of $\mathrm{SO}(2N)$.

⁷From the point of view of field theory there is nothing that distinguishes $a \rightarrow b$ from $b \rightarrow a$ or $a \dashrightarrow b$ from $b \dashrightarrow a$. Nevertheless, we will keep the orientations of arrows because somehow they carry essential information with regard to the integral formula for the elliptic genus of the $(0, 1)$ brane box model.

The choice of the domain of integration is slightly more complicated here than the case of the 1D brane box model. As a function of a_i , $W(b \rightarrow a)$ has poles at $a_i = q^n b_j^{\pm 1}$, $n \in \mathbb{Z}$. The integration contour for a_i is such that it encircles all poles of the form $a_i = b_j^{\pm 1}$ in $W(b \rightarrow a)$ and excludes all other poles. In the 1D limit $q \rightarrow 0$, the poles outside the contour are all sent to 0 or ∞ .

4.2. The tetrahedron equation. The proof of the tetrahedron equation for the oblique unit cell of this lattice model is entirely analogous to the case of the 1D brane box model in section 3.2.

Let N_1, N_2, N_3, n_2, n_3 be positive integers satisfying (51). The tetrahedron equation is a special case of the following equation:

$$(68) \quad \frac{(-1)^{n_3 N_2}}{2^{n_3-1} n_3!} \int_{\mathcal{D}} \prod_i \frac{dz_i}{2\pi i z_i} \eta^2 \hat{\theta}(z_i^2) \prod_{i \neq j} \hat{\theta}(z_i z_j^{\pm 1}) \frac{\prod_{i,\gamma} \hat{\theta}(w_\gamma z_i^{\pm 1}) \prod_{\alpha,\beta} \hat{\theta}(u_\alpha v_\beta^{\pm 1})}{\prod_{i,\alpha} \hat{\theta}(z_i u_\alpha^{\pm 1}) \prod_{i,\beta} \hat{\theta}(v_\beta z_i^{\pm 1})}$$

$$= \frac{(-1)^{n_2(N_2+N_3)}}{2^{n_2-1} n_2!} \int_{\mathcal{D}'} \prod_{\bar{i}} \frac{dz'_{\bar{i}}}{2\pi i z'_{\bar{i}}} \eta^2 \hat{\theta}(z'_{\bar{i}}{}^2) \prod_{\bar{i} \neq \bar{j}} \hat{\theta}(z'_{\bar{i}} z'_{\bar{j}}^{\pm 1}) \frac{\prod_{\bar{i},\beta} \hat{\theta}(v_\beta z'_{\bar{i}}{}^{\pm 1}) \prod_{\gamma,\alpha} \hat{\theta}(w_\gamma u_\alpha^{\pm 1})}{\prod_{\bar{i},\gamma} \hat{\theta}(z'_{\bar{i}} w_\gamma^{\pm 1}) \prod_{\bar{i},\alpha} \hat{\theta}(u_\alpha z'_{\bar{i}}{}^{\pm 1})}.$$

The domains \mathcal{D} and \mathcal{D}' enclose all poles where $\{z_i\}_{i=1}^{n_3} \subset \{u_\alpha^{\pm 1}\}_{\alpha=1}^{N_1}$ and $\{z'_{\bar{i}}\}_{\bar{i}=1}^{n_2} \subset \{w_\gamma^{\pm 1}\}_{\gamma=1}^{N_3}$, respectively, but no other poles. The integrand on each side is a meromorphic differential on the elliptic curve $\mathbb{C}^\times/q\mathbb{Z}$ for each integration variable.

Let \mathfrak{S} be a subset of $\{1, \dots, N_1\}$ of order n_3 and consider the set of poles where $\{z_i\}_{i=1}^{n_3} \subset \{u_i^{\pm 1}\}_{i \in \mathfrak{S}}$ in the integrand on the left-hand side. The poles where $z_i = z_j^{\pm 1}$ for some i, j with $i \neq j$ do not contribute to the integral because of the factor $\hat{\theta}(z_i z_j^{\pm 1})$. There are $2^{n_3} n_3!$ remaining poles from this set. Using

$$(69) \quad \text{Res}_{z=w} \frac{\eta^2}{z \hat{\theta}(z/w)} = 1,$$

we find that the sum of their residues is

$$(70) \quad 2 \frac{\prod_{i \in \mathfrak{S}} \prod_\gamma \hat{\theta}(w_\gamma u_i^{\pm 1}) \prod_{\bar{i} \in \bar{\mathfrak{S}}} \prod_\beta \hat{\theta}(u_{\bar{i}} v_\beta^{\pm 1})}{\prod_{i \in \mathfrak{S}} \prod_{\bar{i} \in \bar{\mathfrak{S}}} \hat{\theta}(u_i u_{\bar{i}}^{\pm 1})}.$$

For the right-hand side, we can trade the sum over the residues at the poles where $\{z'_{\bar{i}}\}_{\bar{i}=1}^{n_2} \subset \{w_\gamma^{\pm 1}\}_{\gamma=1}^{N_3}$ for $(-1)^{n_2}$ times the sum over the residues at the poles where $\{z'_{\bar{i}}\}_{\bar{i}=1}^{n_2} \subset \{u_\alpha^{\pm 1}\}_{\alpha=1}^{N_1}$. The contribution from the poles where $\{z'_{\bar{i}}\}_{\bar{i}=1}^{n_2} \subset \{u_{\bar{i}}^{\pm 1}\}_{\bar{i} \in \bar{\mathfrak{S}}}$ is precisely given by the sum (70), and the desired equality (68) follows.

APPENDIX A. THE (0, 2) TRIALITY

Let N_1, N_2, N_3 be nonnegative integers such that

$$(71) \quad N = \frac{1}{2}(N_1 + N_2 + N_3)$$

is also an integer. Let

$$(72) \quad n_\bullet = N - N_\bullet, \quad \bullet = 1, 2, 3.$$

The (0, 2) triality [16] is the infrared equivalence among a triplet of gauge theories T_{N_1, N_2, N_3} , T_{N_2, N_3, N_1} and T_{N_3, N_1, N_2} .

The theory T_{N_1, N_2, N_3} is a two-dimensional $\mathcal{N} = (0, 2)$ supersymmetric gauge theory with gauge group $U(n_3)$ and has fundamental chiral multiplets Φ_α , $\alpha = 1,$

	$U(n_3)$	$U(N_1)$	$U(N_2)$	$U(N_3)$	$U(1)_\Omega$	$U(1)_R$
Φ_α	\mathbf{n}_3	$\overline{\mathbf{N}}_1$	$\mathbf{1}$	$\mathbf{1}$	0	r_α
P_β	$\overline{\mathbf{n}}_3$	$\mathbf{1}$	\mathbf{N}_2	$\mathbf{1}$	0	s_β
Ψ_γ	$\overline{\mathbf{n}}_3$	$\mathbf{1}$	$\mathbf{1}$	\mathbf{N}_3	0	t_γ
$\Gamma_{\alpha\beta}$	$\mathbf{1}$	\mathbf{N}_1	$\overline{\mathbf{N}}_2$	$\mathbf{1}$	0	$1 - r_\alpha - s_\beta$
Ω_+	det	$\mathbf{1}$	$\mathbf{1}$	det	1	q_+
Ω_-	\det^{-1}	det	det	$\mathbf{1}$	1	q_-

TABLE 1. The theory T_{N_1, N_2, N_3} .

\dots , N_1 , antifundamental chiral multiplets P_β , $\beta = 1, \dots, N_2$, antifundamental Fermi multiplets Ψ_γ , $\gamma = 1, \dots, N_3$, and $N_1 N_2$ singlet Fermi multiplets $\Gamma_{\alpha\beta}$. They are coupled by a vector multiplet for $U(n_3)$ and the superpotential $\sum_{\alpha, \beta} \Phi_\alpha \Gamma_{\alpha\beta} P_\beta$. The rank of the gauge group is fixed by the condition for the absence of nonabelian gauge anomaly. Furthermore, a pair of determinant Fermi multiplets Ω_+ , Ω_- is introduced to cancel abelian gauge anomaly.

The flavor symmetry of T_{N_1, N_2, N_3} is $U(N_1) \times U(N_2) \times U(N_3) \times U(1)_\Omega$, under which the matter multiplets transform as summarized in Table 1. We use K_1^α , K_2^β , K_3^γ , K_Ω respectively to denote the Cartan generators of $U(N_1)$, $U(N_2)$, $U(N_3)$, $U(1)_\Omega$ such that

$$(73) \quad K_1^\alpha(\Phi_{\alpha'}) = -\delta_{\alpha'}^\alpha, \quad K_2^\beta(P_{\beta'}) = \delta_{\beta'}^\beta, \quad K_3^\gamma(\Psi_{\gamma'}) = \delta_{\gamma'}^\gamma, \quad K_\Omega(\Omega_\pm) = 1.$$

The generator \overline{J}_0 of the $U(1)$ R-symmetry in the infrared $\mathcal{N} = (0, 2)$ superconformal algebra can be identified by the c -extremization procedure and is given by

$$(74) \quad \overline{J}_0(\Phi_\alpha) = \frac{n_2}{N}, \quad \overline{J}_0(P_\beta) = \frac{n_1}{N}, \quad \overline{J}_0(\Gamma_{\alpha\beta}) = \frac{n_3}{N}, \quad \overline{J}_0(\Psi_\gamma) = \overline{J}_0(\Omega_\pm) = 0.$$

For our purposes it is convenient to shift the R-charge \overline{J}_0 with Cartan generators of the flavor symmetry so that we can choose the values of the resulting R-charge R arbitrarily for Φ_α , P_β , Ψ_γ and Ω_+ :

$$(75) \quad R = \overline{J}_0 - \sum_\alpha \left(r_\alpha - \frac{n_2}{N} \right) K_1^\alpha + \sum_\beta \left(s_\beta - \frac{n_1}{N} \right) K_2^\beta + \sum_\gamma t_\gamma K_3^\gamma + \left(q_+ - \sum_\gamma t_\gamma \right) K_\Omega.$$

The R-symmetry $U(1)_R$ generated by R generically commutes only with the maximal torus of the flavor symmetry. The absence of mixed $U(1)_R$ -gauge anomaly requires

$$(76) \quad \sum_\alpha (r_\alpha - 1) - \sum_\beta (s_\beta - 1) + \sum_\gamma t_\gamma - q_+ + q_- = 0,$$

which fixes the R-charge q_- of Ω_- .

There are various pieces of evidence for the $(0, 2)$ triality. The most convincing among them is the agreement of the elliptic genera across the three theories. Here we demonstrate that the elliptic genera in the NS-NS sectors of T_{N_1, N_2, N_3} and T_{N_3, N_1, N_2} match.

The elliptic genus in the NS-NS sector of an $\mathcal{N} = (0, 2)$ superconformal field theory T with $U(1)$ flavor symmetries $U(1)_\alpha$, $\alpha = 1, \dots, N_f$, is defined by

$$(77) \quad I_T^{\text{NSNS}}(u_1, \dots, u_{N_f}; q) = \text{Tr}_{\mathcal{H}_T^{\text{NSNS}}} \left((-1)^F q^{L_0} \overline{q}^{\overline{L}_0 - \overline{J}_0/2} \prod_\alpha u_\alpha^{K_\alpha} \right),$$

where L_0, \bar{L}_0 are generators of the Virasoro algebra, K_α is a generator of $U(1)_\alpha$ and u_α is a fugacity for K_α . The trace is taken in the Hilbert space of states $\mathcal{H}_T^{\text{NSNS}}$ in the radial quantization of the NS–NS sector of the theory. Due to supersymmetry only states with $\bar{L}_0 = \bar{J}_0/2$ contribute, hence the elliptic genus is independent of \bar{q} . Using the angular momentum $L_0 - \bar{L}_0$ the above expression can be written as

$$(78) \quad I_T^{\text{NSNS}}(u_1, \dots, u_{N_f}; q) = \text{Tr}_{\mathcal{H}_T^{\text{NSNS}}} \left((-1)^F q^{L_0 - \bar{L}_0 + \bar{J}_0/2} \prod_\alpha u_\alpha^{K_\alpha} \right).$$

By appropriate rescaling of the flavor fugacities the R-charge \bar{J}_0 entering this expression can be modified to any linear combination of the form $\bar{J}_0 + \sum_\alpha c_\alpha K_\alpha$.

When a theory has a standard Lagrangian description, we can write down its NS–NS elliptic genus using the theta function [16, 17]

$$(79) \quad \theta(z) = (z; q)(q/z; q), \quad (z; q) = \prod_{k=0}^{\infty} (1 - zq^k).$$

We have

$$(80) \quad \theta(qz) = \theta(1/z) = -\frac{1}{z}\theta(z)$$

and

$$(81) \quad \text{Res}_{z=w} \frac{(q; q)^2}{z\theta((z/w)^{\pm 1})} = \mp 1.$$

Let $u_\alpha, v_\beta, w_\gamma, x$ be the fugacities for $K_1^\alpha, K_2^\beta, K_3^\gamma, K_\Omega$, respectively. The elliptic genus of T_{N_1, N_2, N_3} is given by the integral

$$(82) \quad I_{T_{N_1, N_2, N_3}}(u, v, w, x) = \frac{1}{n_3!} \oint \mathcal{I}_{T_{N_1, N_2, N_3}}(z; u, v, w, x),$$

where

$$(83) \quad \begin{aligned} & \mathcal{I}_{T_{N_1, N_2, N_3}}(z; u, v, w, x) \\ &= \prod_i -\frac{dz_i}{2\pi i z_i} (q; q)^2 \prod_{i \neq j} \theta\left(\frac{z_i}{z_j}\right) \prod_{i, \alpha} \theta\left(q^{\frac{1}{2}r_\alpha} \frac{z_i}{u_\alpha}\right)^{-1} \prod_{i, \beta} \theta\left(q^{\frac{1}{2}s_\beta} \frac{v_\beta}{z_i}\right)^{-1} \\ & \times \prod_{i, \gamma} \theta\left(q^{\frac{1}{2}(1+t_\gamma)} \frac{w_\gamma}{z_i}\right) \prod_{\alpha, \beta} \theta\left(q^{\frac{1}{2}(2-r_\alpha-s_\beta)} \frac{u_\alpha}{v_\beta}\right) \\ & \times \theta\left(q^{\frac{1}{2}(1+q_+)} x \prod_i z_i \prod_\gamma w_\gamma\right) \theta\left(q^{\frac{1}{2}(1+q_-)} x \frac{\prod_\alpha u_\alpha \prod_\beta v_\beta}{\prod_i z_i}\right). \end{aligned}$$

For each variable $z_i, i = 1, \dots, n_3$, the integrand is a meromorphic differential on the elliptic curve $\mathbb{C}^\times/q\mathbb{Z}$. The domain of integration is chosen in such a way that the integral receives contributions only from the iterated residues at the poles coming from the factor $\prod_{i, \alpha} \theta(q^{\frac{1}{2}r_\alpha} z_i/u_\alpha)^{-1}$.

The poles contributing to the integral (82) are in one-to-one correspondence with the ordered choices of n_3 elements from the set $\{1, \dots, N_1\}$. For a given choice, there are $n_3!$ permutations of elements and the corresponding poles contribute equally.

Thus, the elliptic genus is a sum over the subsets of $\{1, \dots, N_1\}$ with n_3 elements:

$$(84) \quad I_{T_{N_1, N_2, N_3}}(u, v, w, x) = \sum_{\substack{\mathfrak{S} \subset \{1, \dots, N_1\} \\ |\mathfrak{S}| = n_3}} I_{T_{N_1, N_2, N_3}}^{\mathfrak{S}}(u, v, w, x).$$

The summands are given by

$$(85) \quad \begin{aligned} & I_{T_{N_1, N_2, N_3}}^{\mathfrak{S}}(u, v, w, x) \\ &= \prod_{i \in \mathfrak{S}} \prod_{\bar{i} \in \bar{\mathfrak{S}}} \theta \left(q^{\frac{1}{2}(r_{\bar{i}} - r_i)} \frac{u_i}{u_{\bar{i}}} \right)^{-1} \prod_{i \in \mathfrak{S}} \prod_{\gamma} \theta \left(q^{\frac{1}{2}(1+t_{\gamma} + r_i)} \frac{w_{\gamma}}{u_i} \right) \prod_{\alpha, \beta} \theta \left(q^{\frac{1}{2}(r_{\bar{i}} + s_{\beta})} \frac{v_{\beta}}{u_{\bar{i}}} \right) \\ & \quad \times \theta \left(q^{\frac{1}{2}(1+q_+ - \sum_{i \in \mathfrak{S}} r_i)} x \prod_{i \in \mathfrak{S}} u_i \prod_{\gamma} w_{\gamma} \right) \theta \left(q^{\frac{1}{2}(1+q_- + \sum_{i \in \mathfrak{S}} r_i)} x \prod_{\bar{i} \in \bar{\mathfrak{S}}} u_{\bar{i}} \prod_{\beta} v_{\beta} \right), \end{aligned}$$

where the index i now runs over the elements of \mathfrak{S} , whereas \bar{i} runs over the n_2 elements of the complement $\bar{\mathfrak{S}}$ of $\mathfrak{S} \subset \{1, \dots, N_1\}$.

Trading the sum over \mathfrak{S} for a sum over $\bar{\mathfrak{S}}$, we can rewrite the elliptic genus as a similar integral with n_2 integration variables,

$$(86) \quad I_{T_{N_1, N_2, N_3}}(u, v, w, x) = \frac{(-1)^{n_2}}{n_2!} \oint \mathcal{I}_{T_{N_3, N_1, N_2}}(z'; u, v, w, x),$$

where the integrand

$$(87) \quad \begin{aligned} & \mathcal{I}_{T_{N_3, N_1, N_2}}(z'; u, v, w, x) \\ &= \prod_{\bar{i}} -\frac{dz'_{\bar{i}}}{2\pi i z'_{\bar{i}}} (q; q)^2 \prod_{\bar{i} \neq \bar{j}} \theta \left(\frac{z'_{\bar{i}}}{z'_{\bar{j}}} \right) \prod_{\bar{i}, \gamma} \theta \left(q^{\frac{1}{2}(1-t_{\gamma} - \Delta)} \frac{z'_{\bar{i}}}{w_{\gamma}} \right)^{-1} \prod_{\bar{i}, \alpha} \theta \left(q^{\frac{1}{2}(-r_{\alpha} + \Delta)} \frac{u_{\alpha}}{z'_{\bar{i}}} \right)^{-1} \\ & \quad \times \prod_{\bar{i}, \beta} \theta \left(q^{\frac{1}{2}(s_{\beta} + \Delta)} \frac{v_{\beta}}{z'_{\bar{i}}} \right) \prod_{c, \alpha} \theta \left(q^{\frac{1}{2}(1+t_{\gamma} + r_{\alpha})} \frac{w_{\gamma}}{u_{\alpha}} \right) \\ & \quad \times \theta \left(q^{\frac{1}{2}(1+q_- + \sum_{\alpha} r_{\alpha} - n_2 \Delta)} x \prod_{\bar{i}} z'_{\bar{i}} \prod_{\beta} v_{\beta} \right) \theta \left(q^{\frac{1}{2}(1+q_+ - \sum_{\alpha} r_{\alpha} + n_2 \Delta)} x \frac{\prod_{\alpha} u_{\alpha} \prod_{\gamma} w_{\gamma}}{\prod_{\bar{i}} z'_{\bar{i}}} \right), \end{aligned}$$

is a meromorphic differential on $\mathbb{C}^{\times}/q^{\mathbb{Z}}$ for each variable $z'_{\bar{i}}$, $\bar{i} = 1, \dots, n_2$, and the domain of integration encloses the poles from $\prod_{\bar{i}, \gamma} \theta(q^{\frac{1}{2}(-r_{\alpha} + \Delta)} u_{\alpha}/z'_{\bar{i}})^{-1}$. Since the sum of the residues of a meromorphic differential on a compact Riemann surface vanishes, we can change the domain to one that picks up the poles from $\prod_{\bar{i}, \gamma} \theta(q^{\frac{1}{2}(1-t_{\gamma} - \Delta)} z'_{\bar{i}}/w_{\gamma})^{-1}$ after dropping the sign factor $(-1)^{n_2}$ in front of the integral. Doing so precisely yields the integral formula for the elliptic genus of the theory T_{N_3, N_1, N_2} , with the matter contents summarized in Table 2.

The constant Δ in the R-charge assignment accounts for the freedom of shifting the R-charge by a multiple of a generator of the overall U(1) factor of the gauge group U(n_2). Such a shift acts trivially on physical states and operators. At the level of the integral formula for the elliptic genus, the shift corresponds to rescaling the integration variables by a power of q . If we set

$$(88) \quad R(\Omega_+) = R(\Omega'_-), \quad R(\Omega_-) = R(\Omega'_+),$$

then we have

$$(89) \quad \Delta = \frac{1}{n_2} \sum_{\alpha} r_{\alpha}.$$

	$U(n_2)$	$U(N_1)$	$U(N_2)$	$U(N_3)$	$U(1)_\Omega$	$U(1)_R$
P'_α	$\overline{\mathbf{n}}_2$	\mathbf{N}_1	$\mathbf{1}$	$\mathbf{1}$	0	$-r_\alpha + \Delta$
Ψ'_β	$\overline{\mathbf{n}}_2$	$\mathbf{1}$	\mathbf{N}_2	$\overline{\mathbf{1}}$	0	$s_\beta - 1 + \Delta$
Φ'_γ	\mathbf{n}_2	$\mathbf{1}$	$\mathbf{1}$	$\overline{\mathbf{N}}_3$	0	$1 - t_\gamma - \Delta$
$\Gamma'_{\gamma\alpha}$	$\mathbf{1}$	$\overline{\mathbf{N}}_1$	$\mathbf{1}$	\mathbf{N}_3	0	$t_\gamma + r_\alpha$
Ω'_+	det	$\mathbf{1}$	det	$\mathbf{1}$	1	$q_- + \sum_\alpha r_\alpha - n_2 \Delta$
Ω'_-	det ⁻¹	det	$\mathbf{1}$	det	1	$q_+ - \sum_\alpha r_\alpha + n_2 \Delta$

TABLE 2. The theory T_{N_3, N_1, N_2} related to T_{N_1, N_2, N_3} by the $(0, 2)$ triality.

ACKNOWLEDGMENT

This work is supported by NSFC Grant Number 12375064.

REFERENCES

- [1] R. J. Baxter, *On Zamolodchikov's solution of the tetrahedron equations*, Comm. Math. Phys. **88** (1983), no. 2, 185–205. MR696804
- [2] Vladimir V. Bazhanov and Sergey M. Sergeev, *Zamolodchikov's tetrahedron equation and hidden structure of quantum groups*, J. Phys. A **39** (2006), no. 13, 3295–3310, available at [hep-th/0509181](#). MR2214211
- [3] ———, *Elliptic gamma-function and multi-spin solutions of the Yang-Baxter equation*, Nucl. Phys. B **856** (2012), no. 2, 475–496, available at [1106.5874](#). MR2868616
- [4] ———, *A master solution of the quantum Yang-Baxter equation and classical discrete integrable equations*, Adv. Theor. Math. Phys. **16** (2012), no. 1, 65–95, available at [1006.0651](#). MR3019403
- [5] Francesco Benini, Richard Eager, Kentaro Hori, and Yuji Tachikawa, *Elliptic genera of two-dimensional $\mathcal{N} = 2$ gauge theories with rank-one gauge groups*, Lett. Math. Phys. **104** (2014), no. 4, 465, available at [1305.0533](#). MR3177993
- [6] ———, *Elliptic genera of 2d $\mathcal{N} = 2$ gauge theories*, Comm. Math. Phys. **333** (2015), no. 3, 1241, available at [1308.4896](#). MR3302634
- [7] Michael Blaszczyk, Stefan Groot Nibbelink, and Fabian Ruehle, *Green-Schwarz mechanism in heterotic $(2, 0)$ gauged linear sigma models: torsion and NS5 branes*, JHEP **8** (2011), 083, 22, available at [1107.0320](#). MR2876051
- [8] Kevin Costello, *Supersymmetric gauge theory and the Yangian* (2013), available at [1303.2632](#).
- [9] ———, *Integrable lattice models from four-dimensional field theories*, String-Math 2013, 2014, pp. 3–23. MR3330282
- [10] Sebastian Franco, Dongwook Ghim, Sangmin Lee, and Rak-Kyeong Seong, *Elliptic genera of 2d $(0, 2)$ gauge theories from brane brick models*, JHEP **6** (2017), 068, front matter+46, available at [1702.02948](#). MR3760717
- [11] Sebastián Franco, Dongwook Ghim, Sangmin Lee, Rak-Kyeong Seong, and Daisuke Yokoyama, *2d $(0, 2)$ quiver gauge theories and D-branes*, JHEP **9** (2015), 072, front matter+76, available at [1506.03818](#). MR3429415
- [12] Sebastian Franco, Sangmin Lee, and Rak-Kyeong Seong, *Brane brick models and 2d $(0, 2)$ triality*, JHEP **05** (2016), 020, available at [1602.01834](#).
- [13] Sebastián Franco, Sangmin Lee, and Rak-Kyeong Seong, *Brane brick models, toric Calabi-Yau 4-folds and 2d $(0, 2)$ quivers*, JHEP **2** (2016), 047, front matter+66, available at [1510.01744](#). MR3491222
- [14] Sebastián Franco, Alessandro Mininno, Ángel M. Uranga, and Xingyang Yu, *2d $\mathcal{N} = (0, 1)$ gauge theories and Spin(7) orientifolds*, JHEP **3** (2022), Paper No. 150, 52, available at [2110.03696](#). MR4427067
- [15] ———, *Spin(7) orientifolds and 2d $\mathcal{N} = (0, 1)$ triality*, JHEP **1** (2022), Paper No. 058, 62, available at [2112.03929](#). MR4403486

- [16] Abhijit Gadde, Sergei Gukov, and Pavel Putrov, $(0, 2)$ *trialities*, JHEP **03** (2014), 076, available at [1310.0818](#).
- [17] ———, *Walls, lines, and spectral dualities in 3d gauge theories*, JHEP **05** (2014), 047, available at [1302.0015](#).
- [18] Hugo García-Compeán and Angel M. Uranga, *Brane box realization of chiral gauge theories in two dimensions*, Nucl. Phys. B **539** (1999), no. 1-2, 329, available at [hep-th/9806177](#). MR1670640 (2000e:81178)
- [19] Sergei Gukov, Du Pei, and Pavel Putrov, *Trialities of minimally supersymmetric 2d gauge theories*, JHEP **04** (2020), 079, available at [1910.13455](#). MR4096970
- [20] Rei Inoue, Atsuo Kuniba, Xiaoyue Sun, Yuji Terashima, and Junya Yagi, *Solutions of tetrahedron equation from quantum cluster algebra associated with symmetric butterfly quiver* (20242), available at [2403.08814](#).
- [21] Rei Inoue, Atsuo Kuniba, and Yuji Terashima, *Quantum cluster algebras and 3D integrability: Tetrahedron and 3D reflection equations* (202310), available at [2310.14493](#).
- [22] ———, *Tetrahedron equation and quantum cluster algebras*, J. Phys. A **57** (2024), no. 8, 085202, available at [2310.14529](#).
- [23] M. M. Kapranov and V. A. Voevodsky, *2-categories and Zamolodchikov tetrahedra equations*, Algebraic groups and their generalizations: quantum and infinite-dimensional methods (University Park, PA, 1991), 1994, pp. 177–259. MR1278735
- [24] Kenji Mohri, *D-branes and quotient singularities of Calabi-Yau four-folds*, Nuclear Phys. B **521** (1998), no. 1-2, 161–182, available at [hep-th/9707012](#). MR1646360
- [25] Callum Quigley and Savdeep Sethi, *Linear sigma models with torsion*, JHEP **11** (2011), 034, 34, available at [1107.0714](#). MR2913294
- [26] Sakura Schäfer-Nameki and Timo Weigand, *F-theory and 2d $(0, 2)$ theories*, JHEP **5** (2016), 059, front matter+103, available at [1601.02015](#). MR3521778
- [27] V. P. Spiridonov, *Elliptic beta integrals and solvable models of statistical mechanics*, Algebraic aspects of Darboux transformations, quantum integrable systems and supersymmetric quantum mechanics, 2012, pp. 181. MR2905635 (2012m:82015)
- [28] Xiaoyue Sun and Junya Yagi, *Cluster transformations, the tetrahedron equation and three-dimensional gauge theories* (202211), available at [2211.10702](#).
- [29] Junya Yagi, *Quiver gauge theories and integrable lattice models*, JHEP **10** (2015), 065, available at [1504.04055](#). MR3435574
- [30] ———, *Integrable 3D lattice model in M-theory*, JHEP **1** (2023), Paper No. 22, 37, available at [2203.09706](#). MR4534097
- [31] Masahito Yamazaki, *Quivers, YBE and 3-manifolds*, JHEP **05** (2012), 147, available at [1203.5784](#). MR3042929
- [32] A. B. Zamolodchikov, *Tetrahedra equations and integrable systems in three-dimensional space*, Soviet Phys. JETP **52** (1980), no. 2, 325–336. MR611994