

# WEIGHTED EHRHART THEORY VIA EQUIVARIANT TORIC GEOMETRY

LAURENȚIU MAXIM AND JÖRG SCHÜRMAN

**ABSTRACT.** We give a  $K$ -theoretic and geometric interpretation for a generalized weighted Ehrhart theory of a full-dimensional lattice polytope  $P$ , depending on a given homogeneous polynomial function  $\varphi$  on  $P$ , and with Laurent polynomial weights  $f_Q(y) \in \mathbb{Z}[y^{\pm 1}]$  associated to the faces  $Q \preceq P$  of the polytope. For this purpose, we calculate equivariant  $K$ -theoretic Hodge-Chern classes of an equivariant mixed Hodge module  $\mathcal{M}$  on the toric variety  $X_P$  associated to  $P$  (defined via an equivariant embedding of  $X_P$  into an ambient smooth variety). For any integer  $\ell$ , we introduce a corresponding equivariant Hodge  $\chi_y$ -polynomial  $\chi_y(X_P, \ell D_P; [\cdot, \mathcal{M}])$ , with  $D_P$  the corresponding ample Cartier divisor on  $X_P$  (defined by the facet presentation of  $P$ ). Motivic properties of the Hodge-Chern classes are used to express this equivariant Hodge polynomial in terms of weighted character sums fitting with a generalized weighted Ehrhart theory. The equivariant Hodge polynomials are shown to satisfy a reciprocity and purity formula fitting with the duality for equivariant mixed Hodge modules, and implying the corresponding properties for the generalized weighted Ehrhart polynomials. In the special case of the equivariant intersection cohomology mixed Hodge module, with the weight function corresponding to Stanley's  $g$ -function of the polar polytope of  $P$ , we recover in geometric terms a recent combinatorial formula of Beck–Gunnells–Materov.

## 1. INTRODUCTION

We provide a geometric (equivariant) perspective for the generalized weighted Ehrhart theory developed in [3] for a full-dimensional lattice polytope  $P$  with a homogeneous polynomial function  $\varphi$  defined on it. This is achieved by attaching Laurent polynomial weights  $f_Q(y) \in \mathbb{Z}[y^{\pm 1}]$  of geometric origin to the faces  $\emptyset \neq Q \preceq P$  of the polytope (with the partial order given by face inclusion). The Brion–Vergne [8] combinatorial approach to reciprocity can be linearly extended to this generalized weighted Ehrhart theory. Motivation for using such weights comes from our prior work [9] on equivariant Hodge-Chern and Hirzebruch characteristic classes of toric varieties (associated to such polytopes), as well as from the use of Stanley's  $g$ -functions in the generalized Ehrhart reciprocity theorem of [3].

**1.1. Combinatorial and geometrical posets and their identification.** We review here some basic notions explaining the relation between lattice polytopes and toric geometry; for more details see [11, 17].

Let  $M \cong \mathbb{Z}^n$  be a lattice and  $P \subset M_{\mathbb{R}} := M \otimes \mathbb{R} \cong \mathbb{R}^n$  be a full-dimensional lattice polytope. To  $P$  one associates a fan  $\Sigma = \Sigma_P \subset N_{\mathbb{R}} := N \otimes \mathbb{R}$ , with  $N$  the dual lattice of  $M$ , called

---

*Date:* May 7, 2024.

*2020 Mathematics Subject Classification.* 14M25, 14C17, 14C40, 19L47, 52B20, 55N91.

*Key words and phrases.* Toric varieties, lattice polytopes, Ehrhart polynomial, reciprocity, purity, equivariant Hodge-Chern classes, equivariant mixed Hodge module, equivariant Serre duality.

the inner normal fan of  $P$ . To the fan  $\Sigma_P$  one further associates a projective toric variety  $X = X_P$  with torus  $\mathbb{T}$  (whose character lattice is  $M$ ) and an ample Cartier divisor  $D = D_P$ . There is an order-reversing one-to-one correspondence between the faces  $\emptyset \neq Q \preceq P$  of  $P$  and the cones  $\sigma_Q$  of  $\Sigma_P$ , with  $\dim_{\mathbb{R}}(Q) = n - \dim_{\mathbb{R}}(\sigma_Q)$ . There is an order-reversing one-to-one correspondence between the cones  $\sigma_Q$  of  $\Sigma_P$  and the  $\mathbb{T}$ -orbits  $O_Q$  of the toric variety  $X_P$ , with  $\dim_{\mathbb{C}}(O_Q) = n - \dim_{\mathbb{R}}(\sigma_Q)$ . Then the identification between the faces  $\emptyset \neq Q \preceq P$  of  $P$  and the torus orbits  $O_Q$  of the toric variety  $X_P$  is order-preserving, with  $\dim_{\mathbb{R}}(Q) = \dim_{\mathbb{C}}(O_Q)$ .

If, in addition,  $P$  contains the origin in its interior, then its *polar polytope*  $P^\circ \subset N_{\mathbb{R}}$  is defined as in [17, Section 1.5], as a full-dimensional lattice polytope with respect to the lattice  $N$ , and containing the origin in the interior. By taking cones at the origin of  $N_{\mathbb{R}}$  over the proper faces of  $P^\circ$ , with the empty set  $\emptyset$  corresponding to the origin, one gets the same lattice fan  $\Sigma_P$  (hence the same toric variety  $X_P$ ). This correspondence between the proper faces  $\emptyset \preceq Q^\circ \prec P^\circ$  of  $P^\circ$  and the cones of  $\Sigma_P$  is order-preserving, increasing real dimension by 1. (Here  $\dim_{\mathbb{R}} \emptyset = -1$  by convention.) This induces an order-reversing one-to-one correspondence between the faces  $Q$  of  $P$ , and the faces  $Q^\circ$  of the polar polytope  $P^\circ$ , switching the roles of polytopes and emptysets (seen as faces). Moreover, for a proper face  $\emptyset \neq Q \prec P$  of  $P$ , one has  $\dim_{\mathbb{R}}(Q) + \dim_{\mathbb{R}}(Q^\circ) = n - 1$ .

The original polytope  $P \subset M_{\mathbb{R}}$  is classically used for counting lattice points (and the corresponding Ehrhart theory), fitting with sections of the ample line bundle  $\mathcal{O}_{X_P}(D_P)$  on  $X_P$  [12, 17, 11]. The torus orbits of  $X_P$  give a natural  $\mathbb{T}$ -invariant Whitney stratification, which is particularly useful in our geometric interpretation of weighted lattice points counting via the theory of Hirzebruch homology classes  $T_{y^*}$  of [6] and the corresponding Hirzebruch-Riemann-Roch theorem; see [21, 22]. The polar polytope  $P^\circ$  appears in Stanley's work [29] for recursively defining his *g-polynomials*  $g_{Q^\circ}(t) \in \mathbb{Z}[t]$ , with  $g(\emptyset) = 1$ , for the Eulerian graded poset given by the faces  $\emptyset \preceq Q^\circ \preceq P^\circ$ . It is well-known [29, 5, 13, 16, 27] that these *g-polynomials* are related to the intersection cohomology complex of  $X_P$ . In the use of the *g-polynomials*, we will (implicitly) assume that  $P$  contains the origin in its interior (so that the polar polytope is well-defined).

**1.2. Generalized weighted Ehrhart theory.** Let  $\varphi: M_{\mathbb{R}} \cong \mathbb{R}^n \rightarrow \mathbb{C}$  be a homogeneous polynomial function. We assign to the posets mentioned in the previous subsection Laurent polynomial weights via a *weight vector*  $f = \{f_Q\}$ , with  $f_Q(y) \in \mathbb{Z}[y^{\pm 1}]$  indexed here by the non-empty faces  $\emptyset \neq Q \preceq P$  of  $P$ . Motivated by [3], we assemble these weights, the combinatorics of  $P$ , and the polynomial function  $\varphi$  into the *generalized weighted Ehrhart polynomial*  $E_{P,f}^\varphi(\ell, y)$ , resp.,  $\tilde{E}_{P,f}^\varphi(\ell, y)$ , as follows.

**Definition 1.1.** For  $\ell \in \mathbb{Z}_{>0}$ , define the generalized weighted Ehrhart polynomial of  $P$ ,  $f$  and  $\varphi$  by

$$(1.1) \quad \begin{aligned} E_{P,f}^\varphi(\ell, y) &:= \sum_{Q \preceq P} f_Q(y) \cdot (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m) \\ &=: (1+y)^{\deg(\varphi)} \cdot \tilde{E}_{P,f}^\varphi(\ell, y), \end{aligned}$$

with  $\text{Relint}(\ell Q)$  denoting the relative interior of the face  $\ell Q$  of the dilated polytope  $\ell P$ .

The renormalization  $\tilde{E}_{P,f}^\varphi(\ell, y)$  is more natural to work with from the geometric perspective considered in this paper. The Brion-Vergne [8] combinatorial approach to reciprocity can be linearly extended to this generalized weighted Ehrhart theory, so that  $E_{P,f}^\varphi(\ell, y)$ , resp.,  $\tilde{E}_{P,f}^\varphi(\ell, y)$ , have the following properties:

- (1)  $E_{P,f}^\varphi(\ell, y)$  is obtained by evaluating a polynomial  $E_{P,f}^\varphi(z, y)$  at  $z = \ell \in \mathbb{Z}_{>0}$ . Similarly for  $\tilde{E}_{P,f}^\varphi(\ell, y)$ .
- (2) (Constant term) For  $\ell = 0$ ,

$$(1.2) \quad E_{P,f}^\varphi(0, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1 - y)^{\dim(Q) + \deg(\varphi)} \cdot \varphi(0),$$

that is, evaluating  $\sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m)$  at  $\ell = 0$  as  $(-1)^{\dim(Q) + \deg(\varphi)} \cdot \varphi(0)$  (see [8, Prop.4.1]). Similarly,

$$(1.3) \quad \tilde{E}_{P,f}^\varphi(0, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1 - y)^{\dim(Q)} \cdot (-1)^{\deg(\varphi)} \cdot \varphi(0).$$

- (3) (Reciprocity formula) For  $\ell \in \mathbb{Z}_{>0}$ , using [8, Prop.4.1] for each face  $Q$  of  $P$ , one gets

$$(1.4) \quad E_{P,f}^\varphi(-\ell, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1 - y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m).$$

Similarly,

$$(1.5) \quad \tilde{E}_{P,f}^\varphi(-\ell, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1 - y)^{\dim(Q)} \cdot (-1)^{\deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m).$$

Assume now that the origin is an interior point of  $P$ , so that we can consider the weight vector given by Stanley's  $g$ -polynomials

$$(1.6) \quad f_Q(y) = g_{Q^\circ}(-y) =: \tilde{g}_Q(-y)$$

for the polar polytope of  $P$ , with  $g_\emptyset(-y) = \tilde{g}_P(-y) = 1$ . In this setup, it was shown in [3, Thm.1.3, Thm.2.6] that the following *purity* property holds:

$$(1.7) \quad E_{P,f}^\varphi(-\ell, y) = (-y)^{\dim(P) + \deg(\varphi)} \cdot E_{P,f}^\varphi(\ell, 1/y).$$

This property can be reformulated as

$$(1.8) \quad \tilde{E}_{P,f}^\varphi(-\ell, y) = (-y)^{\dim(P)} \cdot (-1)^{\deg(\varphi)} \cdot \tilde{E}_{P,f}^\varphi(\ell, 1/y).$$

Let us note here that our definition (1.1) for  $E_{P,f}^\varphi(\ell, y)$  equals  $G_\varphi(\ell, y)$  from [3, formula (14)]. In the case when  $P$  is a simple polytope, the polar polytope  $P^\circ$  is simplicial, so that  $g_{Q^\circ}(-y) = 1$ , for all faces  $Q$  of  $P$ . In this case, as explained in [3], the equality (1.7) implies the Dehn-Sommerville relations for  $P$ .

If  $\varphi = 1$  in (1.1), we recover the weighted Ehrhart polynomials studied in [22] via the theory of mixed Hodge modules on the toric variety  $X_P$ .

**1.3. Geometric viewpoint on generalized weighted Ehrhart theory.** In this paper, we give a geometric interpretation and proofs of the above properties for any homogeneous polynomial function  $\varphi$  and any weight vector  $f$  via the theory of *equivariant* mixed Hodge modules on a projective toric variety and their  $K$ -theoretic equivariant characteristic classes. Moreover, using the equivariant intersection Hodge module  $IC_{X^p}^H$  and its purity, we give a geometric proof of formula (1.7).

1.3.1. *Geometric realization of weight vectors and duality.* As already indicated in [22], we identify the weight vectors  $\{f_Q\}$ , seen as  $\mathbb{Z}[y^{\pm 1}]$ -valued functions on the set of faces  $Q$  of  $P$ , with the  $\mathbb{T}$ -invariant  $\mathbb{Z}[y^{\pm 1}]$ -valued constructible functions on  $X = X_P$ , via

$$\{f_Q\} \mapsto \sum_{Q \preceq P} 1_{O_Q} \cdot f_Q \in F^{\mathbb{T}}(X)[y^{\pm 1}].$$

Note that  $F^{\mathbb{T}}(X)[y^{\pm 1}]$  is a  $\mathbb{Z}[y^{\pm 1}]$ -module freely generated by the  $\mathbb{T}$ -invariant constructible functions  $1_{O_Q} \in F^{\mathbb{T}}(X) \subset F^{\mathbb{T}}(X)[y^{\pm 1}]$ . In what follows, we adapt the formulations from [22] in motivic and Hodge-theoretic terms to the equivariant context.

There is a tautological (injective) homomorphism

$$\tau^{\mathbb{T}} : F^{\mathbb{T}}(X)[y^{\pm 1}] \longrightarrow K_0^{\mathbb{T}}(\text{var}/X)[\mathbb{L}^{-1}]$$

to the equivariant Grothendieck group of  $\mathbb{T}$ -varieties over  $X$ , localized at the Lefschetz motive  $\mathbb{L} = [\mathbb{C} \rightarrow pt]$  (with the trivial  $\mathbb{T}$ -action on  $\mathbb{C}$ ), given by

$$1_{O_Q} \cdot f_Q(y) \mapsto [O_Q \hookrightarrow X] \cdot f_Q(-\mathbb{L}).$$

Here,  $K_0^{\mathbb{T}}(\text{var}/X)$  is a  $K_0^{\mathbb{T}}(\text{var}/pt)$ -module via the identification  $X \times pt \simeq X$ .

In order to identify the equivariant intersection cohomology complex and duality we need to go one step further, and work with  $\mathbb{T}$ -equivariant mixed Hodge modules. For technical reasons, here we need to choose a  $\mathbb{T}$ -equivariant closed embedding  $i : X \hookrightarrow Z$  into an ambient smooth  $\mathbb{T}$ -variety  $Z$ , and work with the Grothendieck group  $K_0(\text{MHM}_X^{\mathbb{T}}(Z))$  of equivariant mixed Hodge modules on  $Z$  with support on  $X$ . There is a canonical homomorphism

$$\chi_{\text{Hdg}}^{\mathbb{T}} : K_0^{\mathbb{T}}(\text{var}/X)[\mathbb{L}^{-1}] \rightarrow K_0(\text{MHM}_X^{\mathbb{T}}(Z))$$

such that

$$\chi_{\text{Hdg}}^{\mathbb{T}}([O_Q \hookrightarrow X] \cdot f_Q(-\mathbb{L})) = [(j_Q)! \mathbb{Q}_{O_Q}^H] \cdot f_Q(-u),$$

with  $j_Q : O_Q \hookrightarrow X \hookrightarrow Z$  the orbit inclusion in the ambient smooth  $\mathbb{T}$ -variety  $Z$ , and  $\mathbb{Q}_{O_Q}^H$  denoting the equivariant constant Hodge module on the orbit  $O_Q$ . Here,

$$u = [\mathbb{Q}(-1)] \in K_0(\text{MHM}^{\mathbb{T}}(pt)) \simeq K_0(\text{MHM}(pt)) \simeq K_0(\text{MHS}^p),$$

and the module structure on  $K_0(\text{MHM}_X^{\mathbb{T}}(Z))$  is induced from the exterior product for the identification  $Z \simeq pt \times Z$ . Also,  $\text{MHS}^p$  is the abelian category of graded polarizable  $\mathbb{Q}$ -mixed Hodge structures, with  $\mathbb{Q}(-1)$  the corresponding Tate Hodge structure of weight 2.

In our geometric applications, we recover the corresponding weight vector  $\{f_Q\}$  via

$$f_Q(y) := \chi_y(i_{x_Q}^*[\mathcal{M}])$$

as the Hodge  $\chi_y$ -polynomial of the stalk class  $i_{x_Q}^*[\mathcal{M}]$ , with  $[\mathcal{M}] \in K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$ , and  $i_{x_Q} : \{x_Q\} \hookrightarrow Z$  the inclusion (in the ambient smooth  $\mathbb{T}$ -variety) of a point  $x_Q$  chosen in the orbit  $O_Q$ . Then our aim is to show that many invariants of  $[\mathcal{M}]$  depend only of this choice of a weight function.

Specific to the toric context is the fact that the stabilizers of the torus action are connected, so that, using [31, Lem.1.2], any class  $[\mathcal{M}] \in K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$  can be decomposed as (see Corollary 3.2)

$$(1.9) \quad [\mathcal{M}] = \sum_{Q \preceq P} [(j_Q)_! \mathbb{Q}_{O_Q}^H] \cdot i_{x_Q}^*[\mathcal{M}],$$

and the stalk classes do not see the  $\mathbb{T}$ -action, i.e.,

$$i_{x_Q}^*[\mathcal{M}] \in K_0(\mathrm{MHM}^{\mathbb{T}}(pt)) \simeq K_0(\mathrm{MHM}(pt)) \simeq K_0(\mathrm{MHS}^p).$$

Since the composed map

$$\mathbb{Z}[u^{\pm 1}] \longrightarrow K_0(\mathrm{MHS}^p) \xrightarrow{\chi_y} \mathbb{Z}[y^{\pm 1}]$$

is the isomorphism given by  $u \mapsto -y$ , we get the injectivity of the composition

$$F^{\mathbb{T}}(X)[y^{\pm 1}] \longrightarrow K_0^{\mathbb{T}}(\mathrm{var}/X)[\mathbb{L}^{-1}] \longrightarrow K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)).$$

In this way we also see that any weight vector  $f$  is realized by the stalk class of a certain equivariant mixed Hodge module complex via  $f_Q(y) := \chi_y(i_{x_Q}^*[\mathcal{M}])$ .

Let  $IC_X^H \in \mathrm{MHM}_X^{\mathbb{T}}(Z)$  be the equivariant intersection cohomology Hodge module of  $X$ , and  $IC_X'^H := IC_X^H[-n]$  with  $n = \dim_{\mathbb{C}}(X) = \dim_{\mathbb{R}}(P)$ . Since  $IC_X^H$  is pure of weight  $n$ , we get that  $\mathbb{D}_Z^{\mathbb{T}} IC_X'^H \simeq IC_X'^H[2n](n)$ , with  $\mathbb{D}_Z^{\mathbb{T}}$  the equivariant duality functor of equivariant mixed Hodge modules on the ambient smooth  $\mathbb{T}$ -variety  $Z$ . This implies the following key purity property:

$$(1.10) \quad \mathbb{D}_Z^{\mathbb{T}}[IC_X'^H] = u^{-n} \cdot [IC_X'^H] \in K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)).$$

As already explained in the non-equivariant context of [22], the weight vector associated to  $[IC_X'^H]$  is the collection of Stanley's  $g$ -functions  $\tilde{g} := \{\tilde{g}_Q(-y)\}$ , i.e.,  $\tilde{g}_Q(-y) = \chi_y(i_{x_Q}^*[IC_X'^H])$  (cf. also [5, 16, 13, 27]).

**1.3.2. Twisted equivariant Hodge-Chern classes and generalized weighted Ehrhart theory.** We can now explain our geometric interpretation of the generalized weighted Ehrhart theory introduced in Subsection 1.2, in terms of suitable twists (by the line bundle of the ample Cartier divisor  $D_P$  of  $P$ ) of the equivariant Hodge-Chern class transformation

$$\mathrm{DR}_y^{\mathbb{T}} : K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)) \longrightarrow K_{0,X}^{\mathbb{T}}(Z)[y^{\pm 1}] \simeq K_0^{\mathbb{T}}(X)[y^{\pm 1}],$$

generalizing the corresponding non-equivariant Hodge-Chern classes of [6, 28]. This equivariant Hodge-Chern transformation is defined in terms of the equivariant filtered De Rham complex of  $\mathcal{M} \in \mathrm{MHM}_X^{\mathbb{T}}(Z)$ , which equivariantly can only be defined on the ambient smooth  $\mathbb{T}$ -variety  $Z$ . Here,  $K_{0,X}^{\mathbb{T}}(Z) \simeq K_0^{\mathbb{T}}(X)$  is the Grothendieck group of  $\mathbb{T}$ -equivariant coherent sheaves on  $(Z$  with support in)  $X$ .

The equivariant Hodge-Chern class transformation  $\mathrm{DR}_y^{\mathbb{T}}$  commutes with cross product  $\boxtimes$  for the identification  $Z \times pt \simeq Z$ . Moreover, it is functorial for  $\mathbb{T}$ -equivariant morphisms  $f: (Z, X) \rightarrow (Z', X')$  of pairs, with  $Z, Z'$  smooth complex algebraic  $\mathbb{T}$ -varieties, and  $f: Z \rightarrow Z'$  restricting to a proper morphism  $f: X \rightarrow X'$  of closed  $\mathbb{T}$ -invariant subvarieties. In this paper, the functoriality is used for the closed inclusions  $k_Q: V_Q \hookrightarrow Z$  of the closure of the  $\mathbb{T}$ -orbits  $O_Q$  into the ambient smooth  $\mathbb{T}$ -variety  $Z$ , and their compositions with toric resolutions of these orbit closures, as well as for taking degrees of these classes (i.e., pushing down to a point space). These properties are then used to prove the following basic identity (see (5.7)):

$$(1.11) \quad \mathrm{DR}_y^{\mathbb{T}}([(j_Q)! \mathbb{Q}_{O_Q}^H]) = (1+y)^{\dim(O_Q)} \cdot (k_Q)_*[\omega_{V_Q}]_{\mathbb{T}},$$

with  $\omega_{V_Q}$  the  $\mathbb{T}$ -equivariant canonical (and dualizing) sheaf of the orbit closure  $V_Q$ .

As explained in [14, Sect.3], the equivariant Hodge-Chern class transformation  $\mathrm{DR}_y^{\mathbb{T}}$  commutes with equivariant duality for Grothendieck classes of equivariant mixed Hodge modules on  $Z$  with support in  $X$ , which are in the image of  $\chi_{Hdg}^{\mathbb{T}}$ , i.e., on such elements one has

$$(1.12) \quad \mathrm{DR}_y^{\mathbb{T}} \circ \mathbb{D}_Z^{\mathbb{T}} = \mathbb{D}^{\mathbb{T}} \circ \mathrm{DR}_y^{\mathbb{T}},$$

Here,  $\mathbb{D}_Z^{\mathbb{T}}$  is induced from the exact duality functor of equivariant mixed Hodge modules (on  $Z$ , with support on  $X$ ). For  $\mathbb{D}^{\mathbb{T}}$ , one can use either the equivariant Grothendieck duality on the Grothendieck group  $K_{0,X}^{\mathbb{T}}(Z)$  induced from Grothendieck duality on  $Z$  (remembering the support in  $X$ ), or the equivariant Grothendieck duality on the Grothendieck group  $K_0^{\mathbb{T}}(X)$  induced from Grothendieck duality on  $X$  (see [20] for the foundations of equivariant Grothendieck duality). As in the nonequivariant context,  $\mathbb{D}^{\mathbb{T}}$  is extended to  $K_0^{\mathbb{T}}(X)[y^{\pm 1}]$  by  $y \mapsto 1/y$ . With the above notations, this duality property of  $\mathrm{DR}_y^{\mathbb{T}}$  implies our second basic formula (cf. (5.13)):

$$(1.13) \quad \mathrm{DR}_y^{\mathbb{T}}([(j_Q)_* \mathbb{Q}_{O_Q}^H]) = (1+y)^{\dim(O_Q)} \cdot (k_Q)_*[\mathcal{O}_{V_Q}]_{\mathbb{T}},$$

Let now  $D'$  be a  $\mathbb{T}$ -invariant Cartier divisor on  $X = X_P$ , e.g.,  $D' = \ell D = \ell D_P$  for  $\ell \in \mathbb{Z}$ . Let  $[\mathcal{M}] \in K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$  be fixed. Then

$$\mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(D') := \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]) \cdot [\mathcal{O}_X(D')] \in K_0^{\mathbb{T}}(X)[y^{\pm 1}],$$

where we extend the  $K_{\mathbb{T}}^0(X)$ -module structure of  $K_0^{\mathbb{T}}(X)$  induced from the tensor product linearly over  $\mathbb{Z}[y^{\pm 1}]$ . For  $a_X: X \rightarrow pt$  the constant map as above, we define the  $\mathbb{T}$ -equivariant Hodge polynomial of  $(X, D', [\mathcal{M}])$  as:

$$(1.14) \quad \begin{aligned} \chi_y^{\mathbb{T}}(X, D'; [\mathcal{M}]) &:= (a_X)_* \left( \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(D') \right) \in K_0^{\mathbb{T}}(pt)[y^{\pm 1}] = K_{\mathbb{T}}^0(pt)[y^{\pm 1}] \\ &=: \chi^{\mathbb{T}}(X; \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(D')) \in \mathbb{Z}[M][y^{\pm 1}], \end{aligned}$$

which only depends on  $[\mathcal{M}]$ . In the second equality of the above definition, we use the isomorphism  $K_{\mathbb{T}}^0(pt) \simeq \mathbb{Z}[M]$  induced by the eigenspace decomposition by characters  $\chi^m \in \mathbb{Z}[M]$  (for  $m \in M$  an element in the character lattice of  $\mathbb{T}$ ), see (4.2).

With these notations, we can now formulate the first main result of this paper (see Theorems 6.4, 6.8, 5.2):

**Theorem 1.2.** *Let  $X = X_P$  be the projective toric variety with ample Cartier divisor  $D = D_P$  associated to a full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ , and let  $\ell \in \mathbb{Z}_{>0}$ . Let  $Z$  be an ambient  $\mathbb{T}$ -manifold containing  $X$  as a  $\mathbb{T}$ -invariant subvariety. For any face  $Q$  of  $P$ , let  $x_Q \in O_Q$  be a point in the orbit  $O_Q$  of  $X$ , with inclusion map  $i_{x_Q} : \{i_{x_Q}\} \hookrightarrow Z$ , and set  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ . Then for any Grothendieck class  $[\mathcal{M}] \in K_0(\mathbf{MHM}_X^{\mathbb{T}}(Z))$  of a mixed Hodge module (complex) on  $Z$  with support in  $X$ , we have:*

(a)

$$(1.15) \quad \chi_y^{\mathbb{T}}(X, \ell D; [\mathcal{M}]) = \sum_{Q \leq P} f_Q(y) \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \chi^{-m},$$

with  $\text{Relint}$  denoting the relative interior of a face.

(b)

$$(1.16) \quad \chi_y^{\mathbb{T}}(X, -\ell D; [\mathcal{M}]) = \sum_{Q \leq P} f_Q(y) \cdot (-1-y)^{\dim(Q)} \cdot \sum_{m \in \ell Q \cap M} \chi^m.$$

(c)

$$(1.17) \quad \chi_y^{\mathbb{T}}(X, 0 \cdot D; [\mathcal{M}]) = \sum_{Q \leq P} \left( f_Q(y) \cdot (-1-y)^{\dim(Q)} \right) \cdot \chi^0.$$

Here, for  $\ell \in \mathbb{Z}_{>0}$ ,  $\ell D_P = D_{\ell P}$  is the ample Cartier divisor corresponding to the dilated polytope  $\ell P$ , with the same toric variety  $X_{\ell P} = X_P$ .

All the formulae of the above theorem rely on the decomposition formula (1.9), together with well-known character decompositions of  $H^*(X; \mathcal{O}(\ell D))$  and  $H^*(X; \mathcal{O}(\ell D) \otimes \omega_X)$ , for any integer  $\ell$ , as in the classical work of Danilov [12]. In addition, we also use the two basic identities (1.11) and (1.13).

Furthermore, using the duality property (2.11) for  $\text{DR}_y^{\mathbb{T}}$  as in Lemma 6.14, together with equivariant Serre duality, and combining this with (b), we obtain for  $\ell \in \mathbb{Z}_{>0}$  the following duality formula for the equivariant Hodge polynomial:

$$(1.18) \quad \chi_y^{\mathbb{T}}(X, \ell D; \mathbb{D}_Z^{\mathbb{T}}[\mathcal{M}]) = \chi_{1/y}(X, -\ell D; [\mathcal{M}])|_{m \mapsto -m}.$$

Here, the involution  $m \mapsto -m$  of  $\mathbb{Z}[M]$  corresponds to the involution  $(-)^{\vee}$  on  $K_{\mathbb{T}}^0(pt)$  induced from duality of equivariant vector bundles on a point space, i.e., representation spaces. Together with the purity property (1.10), the duality formula (1.18) applied to  $\mathcal{M} = IC_X^H$  yields

$$(1.19) \quad \chi_y^{\mathbb{T}}(X, -\ell D; [IC_X^H]) = (-y)^n \cdot \chi_{1/y}(X, \ell D; [IC_X^H])|_{m \mapsto -m}.$$

Let us finally explain the role of the chosen homogeneous polynomial function  $\varphi : M_{\mathbb{R}} \cong \mathbb{R}^n \rightarrow \mathbb{C}$  for proving the properties of the generalized weighted Ehrhart polynomials  $E_{P,f}^{\varphi}(\ell, y)$  and, resp.,  $\tilde{E}_{P,f}^{\varphi}(\ell, y)$ , stated in Subsection 1.2. The homogeneous polynomial  $\varphi$  defines group homomorphisms

$$\tilde{E}^{\varphi} : K_{\mathbb{T}}^0(pt) \simeq \mathbb{Z}[M] \longrightarrow \mathbb{C}, \quad E^{\varphi} : K_{\mathbb{T}}^0(pt) \simeq \mathbb{Z}[M] \longrightarrow \mathbb{C}[y]$$

via  $\tilde{E}^{\varphi}(\chi^m) := \varphi(-m)$ , resp.,  $E^{\varphi}(\chi^m) := \varphi(-(1+y) \cdot m)$ . Since  $\varphi$  is homogeneous, the duality involution  $m \mapsto -m$  on the left-hand side corresponds to the multiplication by

$(-1)^{\deg(\varphi)}$  on the right-hand side. Linearly extending  $\tilde{E}^\varphi, E^\varphi$  over Laurent polynomials in  $y$ , we then have that

$$\tilde{E}_{P,f}^\varphi(\ell, y) = \tilde{E}^\varphi(\chi_y^\mathbb{T}(X, \ell D; [\mathcal{M}]))$$

and

$$E_{P,f}^\varphi(\ell, y) = E^\varphi(\chi_y^\mathbb{T}(X, \ell D; [\mathcal{M}]))$$

with  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ . Then the formulae for  $\chi_y^\mathbb{T}(X, \ell D; [\mathcal{M}])$  stated in this subsection translate via  $E^\varphi$  and  $\tilde{E}^\varphi$  into the corresponding properties for the generalized weighted Ehrhart polynomials  $E_{P,f}^\varphi(\ell, y)$  and, resp.,  $\tilde{E}_{P,f}^\varphi(\ell, y)$ , as stated in Subsection 1.2 (see also Section 6).

To explain geometrically the polynomial behavior in  $\ell$  of  $\tilde{E}_{P,f}^\varphi(\ell, y)$ , we have to go from  $K$ -theory to equivariant homology via the equivariant Todd class transformation, using in addition equivariant localization at the torus fixed points (as explained in the Appendix).

**1.4. Structure of the paper.** The paper is structured as follows.

In Section 2, we give a general introduction to the equivariant  $K$ -theoretic Hodge-Chern class transformation  $\mathrm{DR}_y^\mathbb{T}$  of (classes of) equivariant mixed Hodge modules. We explain here the basic calculus (functoriality, module property, and duality), comparing it also with its motivic counterpart from [2, 9].

Preparing the ground for the toric context, in Section 3 we compute these equivariant Hodge-Chern classes in terms of a stratification, reducing them to their motivic counterparts from [9], see Proposition 3.5.

In Section 4, for a  $\mathbb{T}$ -invariant Cartier divisor  $D$  on  $\mathbb{T}$ -invariant compact algebraic subvariety  $X$  of a smooth ambient  $\mathbb{T}$ -variety  $Z$ , we define for the class  $[\mathcal{M}] \in K_0(\mathrm{MHM}_X^\mathbb{T}(Z))$  of an equivariant mixed Hodge module the corresponding equivariant Hodge polynomial  $\chi_y(X, D; [\mathcal{M}])$  of the triple  $(X, D; [\mathcal{M}])$ .

In Section 5, we specialize to the toric context, and make the characteristic class formulae from the previous sections much more explicit. We also prove here the two basic formulae (1.11) and (1.13), as well as formula (1.17).

In Section 6, we prove parts (a) and (b) of Theorem 1.2, as well as a technical result (Lemma 6.14) used to obtain the duality formula (1.18) in the toric context.

Finally, in the Appendix we explain in a geometric way the polynomial behavior in  $\ell$  of the generalized weighted Ehrhart polynomials.

*Acknowledgements.* L. Maxim acknowledges support from the project ‘‘Singularities and Applications’’ - CF 132/31.07.2023 funded by the European Union - NextGenerationEU - through Romania’s National Recovery and Resilience Plan. J. Schürsmann was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 427320536 – SFB 1442, as well as under Germany’s Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics – Geometry – Structure. L. Maxim and J. Schürsmann also thank the Isaac Newton Institute for Mathematical Sciences for the support and hospitality during the program ‘‘Equivariant methods in geometry’’ when work on this paper was undertaken.

## 2. EQUIVARIANT HODGE-CHERN CLASSES VIA MIXED HODGE MODULES

Let  $X$  be a complex algebraic variety with an algebraic  $\mathbb{T} = (\mathbb{C}^*)^n$ -action (e.g., the toric variety  $X_P$  associated to a full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ ), and assume that  $i : X \hookrightarrow Z$  is a closed  $\mathbb{T}$ -equivariant embedding of  $X$  into an ambient complex algebraic manifold  $Z$  endowed with an algebraic  $\mathbb{T}$ -action (if  $X$  is a normal quasi-projective  $\mathbb{T}$ -variety, such an embedding into a  $\mathbb{T}$ -equivariant smooth quasi-projective variety exists by [30, Thm.1], see also [10, Thm.5.1.25] for any linear algebraic group action on such a normal quasi-projective variety).

Consider the abelian category  $\mathrm{MHM}_X^{\mathbb{T}}(Z) \subset \mathrm{MHM}^{\mathbb{T}}(Z)$  of  $\mathbb{T}$ -equivariant mixed Hodge modules on  $Z$  with support in  $X$ , see [1, 31]. There is an exact forgetful functor  $\mathrm{For} : \mathrm{MHM}^{\mathbb{T}}(Z) \rightarrow \mathrm{MHM}(Z)$ , (which for the torus  $\mathbb{T}$  is even fully faithful since  $\mathbb{T}$  is connected, see [1, Thm.5.2]). Let  $K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)) \rightarrow K_0(\mathrm{MHM}^{\mathbb{T}}(Z))$  be the corresponding Grothendieck groups, and note that, via cross-product,  $K_0(\mathrm{MHM}^{\mathbb{T}}(Z))$  is a module over  $K_0(\mathrm{MHM}^{\mathbb{T}}(pt))$ . In [1], Achar introduced a triangulated category  $D_{\mathrm{MHM}}^{b,\mathbb{T}}(Z)$ , together with a bounded non-degenerate t-structure whose heart is  $\mathrm{MHM}^{\mathbb{T}}(Z)$  (note that this is only defined for a smooth  $\mathbb{T}$ -variety  $Z$ , and it is *not* the derived category of  $\mathrm{MHM}^{\mathbb{T}}(Z)$ ). So one has a canonical isomorphism of Grothendieck groups  $K_0(\mathrm{MHM}^{\mathbb{T}}(Z)) \simeq K_0(D_{\mathrm{MHM}}^{b,\mathbb{T}}(Z))$ , and similarly for objects with support contained in  $X$ . The category  $D_{\mathrm{MHM}}^{b,\mathbb{T}}(Z)$  has the same 6-functor formalism as developed in [25] for  $D^b\mathrm{MHM}(Z)$ . All of these functors commute with a corresponding forgetful functor  $\mathrm{For} : D_{\mathrm{MHM}}^{b,\mathbb{T}}(Z) \rightarrow D^b\mathrm{MHM}(Z)$ .

We next discuss a  $\mathbb{T}$ -equivariant version of the Hodge-Chern class of  $X$ , see also [14]. In the non-equivariant case, this was initially introduced in [6], but see also [28] for a definition and study via mixed Hodge modules.

Let  $X$  be as before a closed  $\mathbb{T}$ -invariant subvariety of a complex algebraic  $\mathbb{T}$ -manifold  $Z$ , and let  $\mathcal{M} \in \mathrm{MHM}_X^{\mathbb{T}}(Z)$  be a  $\mathbb{T}$ -equivariant mixed Hodge module on  $Z$  with support in  $X$ , e.g.,  $\mathcal{M} = IC_X^H$  the  $\mathbb{T}$ -equivariant intersection cohomology module (defined by the equivariant intermediate extension of the constant Hodge module from the regular part of  $X$  to the ambient  $\mathbb{T}$ -variety  $Z$ ). For  $p \in \mathbb{Z}$  consider de Rham complex  $\mathrm{DR}_Z(\mathcal{M})$  of the underlying algebraic left  $\mathcal{D}_Z$ -module  $\mathcal{M}$  with its integrable connection  $\nabla$ :

$$\mathrm{DR}_Z(\mathcal{M}) = [\mathcal{M} \xrightarrow{\nabla} \mathcal{M} \otimes_{\mathcal{O}_Z} \Omega_Z^1 \xrightarrow{\nabla} \cdots \xrightarrow{\nabla} \mathcal{M} \otimes_{\mathcal{O}_Z} \Omega_Z^{\dim(Z)}]$$

with  $\mathcal{M}$  in degree  $-\dim(Z)$ , filtered by

$$F_p\mathrm{DR}_Z(\mathcal{M}) = [F_p\mathcal{M} \xrightarrow{\nabla} F_{p+1}\mathcal{M} \xrightarrow{\nabla} \cdots \xrightarrow{\nabla} F_{p+n}\mathcal{M} \otimes_{\mathcal{O}_Z} \Omega_Z^{\dim(Z)}].$$

The complex  $Gr_p^F\mathrm{DR}_Z(\mathcal{M})$  associated to the above filtered de Rham complex  $\mathrm{DR}_Z(\mathcal{M})$  is a bounded complex of  $\mathbb{T}$ -equivariant coherent sheaves on  $Z$  with support in  $X$ . (We use the same symbol for both the mixed Hodge module and the underlying filtered (left)  $\mathcal{D}_Z$ -module.) Moreover,  $Gr_p^F\mathrm{DR}_Z(\mathcal{M}) \simeq 0$  for all but finitely many integers  $p$  and  $\mathcal{M}$  fixed (as can be checked already nonequivariantly).

*Remark 2.1.* It follows from [24, Lem.3.2.6] that the complex  $Gr_p^F\mathrm{DR}_Z(\mathcal{M})$  is a bounded complex of (equivariant) coherent  $\mathcal{O}_X$ -modules (not just supported on  $X$ ), i.e., these (equivariant) coherent  $\mathcal{O}_X$ -modules are annihilated by the ideal sheaf  $\mathcal{I}_X$  of  $X$  in  $Z$ .  $\square$

The transformation  $Gr_p^F DR_Z$  induces a corresponding map on Grothendieck groups since morphisms of (equivariant) mixed Hodge modules are strict with respect to the Hodge filtration. For an equivariant complex  $\mathcal{M} \in D_{\text{MHM},X}^{b,\mathbb{T}}(Z)$  on  $Z$  with support in  $X$ , we perform the above construction for each cohomology object of  $\mathcal{M}$ . So one gets induced Grothendieck group transformations

$$Gr_p^F DR_Z : K_0(D_{\text{MHM},X}^{b,\mathbb{T}}(Z)) \simeq K_0(\text{MHM}_X^{\mathbb{T}}(Z)) \longrightarrow K_{0,X}^{\mathbb{T}}(Z) \simeq K_0^{\mathbb{T}}(X),$$

where  $K_{0,X}^{\mathbb{T}}(Z) \simeq K_0^{\mathbb{T}}(X)$  is the Grothendieck group of  $\mathbb{T}$ -equivariant coherent sheaves on  $(Z$  with support in)  $X$ . The fact that the transformation  $Gr_p^F DR_Z$  takes values in  $K_0^{\mathbb{T}}(X)$  follows directly from Remark 2.1. Alternatively, using the ambient variety  $Z$ , this can also be deduced from the identification  $K_{0,X}^{\mathbb{T}}(Z) \simeq K_0^{\mathbb{T}}(X)$  induced by the pushforward  $i_* : K_0^{\mathbb{T}}(X) \rightarrow K_{0,X}^{\mathbb{T}}(Z)$  for the closed embedding  $i : X \hookrightarrow Z$ . Indeed, the right inverse of  $i_*$  is  $i^* : K_{0,X}^{\mathbb{T}}(Z) \rightarrow K_0^{\mathbb{T}}(X)$ , which is induced from the derived pullback (i.e., alternating sum of equivariant tor-sheaves), since  $i$  is of finite tor-dimension (as  $Z$  is smooth). So  $i_*$  is injective. Moreover, if  $\mathcal{F}$  is an equivariant coherent sheaf on  $Z$  supported on  $X$ , then  $\mathcal{F}$  is annihilated by a finite power of the ideal sheaf  $\mathcal{I}_X$  of  $X$ , and

$$[\mathcal{F}] = \sum_{i \geq 0} [\mathcal{I}_X^{i+1} \cdot \mathcal{F} / \mathcal{I}_X^i \cdot \mathcal{F}] \in i_* K_0^{\mathbb{T}}(X) \subset K_{0,X}^{\mathbb{T}}(Z),$$

proving the surjectivity of  $i_*$ .

We can now make the following definition.

**Definition 2.2** (Equivariant Hodge-Chern transformation). The  $\mathbb{T}$ -equivariant Hodge-Chern class transformation of a complex algebraic  $\mathbb{T}$ -variety  $X$  equivariantly embedded in an ambient complex algebraic  $\mathbb{T}$ -manifold  $Z$  is defined as:

$$\begin{aligned} DR_y^{\mathbb{T}} : K_0(\text{MHM}_X^{\mathbb{T}}(Z)) &\longrightarrow K_0^{\mathbb{T}}(X)[y^{\pm 1}] \\ DR_y^{\mathbb{T}}([\mathcal{M}]) &:= \sum_p [Gr_{-p}^F DR_Z([\mathcal{M}])]_{\mathbb{T}} \cdot (-y)^p. \end{aligned}$$

*Remark 2.3.* By definition,  $DR_y^{\mathbb{T}}([\mathcal{M}])$  depends only on the Grothendieck class  $[\mathcal{M}]$ , and not on the choice of a representative  $\mathcal{M} \in D_{\text{MHM},X}^{b,\mathbb{T}}(Z)$  for this class. Forgetting the  $\mathbb{T}$ -action, one recovers by definition the Hodge-Chern class transformation  $DR_y$  from [6, 28].  $\square$

**Proposition 2.4** (Functoriality of  $DR_y^{\mathbb{T}}$ ). *Let  $f : (Z, X) \rightarrow (Z', X')$  be a  $\mathbb{T}$ -equivariant morphism of smooth complex algebraic  $\mathbb{T}$ -varieties  $f : Z \rightarrow Z'$  restricting to a proper morphism  $f : X \rightarrow X'$  of closed  $\mathbb{T}$ -invariant subvarieties. Then*

$$DR_y^{\mathbb{T}} \circ f_* = f_* \circ DR_y^{\mathbb{T}} : K_0(\text{MHM}_X^{\mathbb{T}}(Z)) \rightarrow K_{0,X'}^{\mathbb{T}}(Z')[y^{\pm 1}].$$

*Proof.* This can be done exactly as explained in [14, Section 3].  $\square$

*Remark 2.5.* Under the identifications  $K_{0,X}^{\mathbb{T}}(Z) \simeq K_0^{\mathbb{T}}(X)$  and  $K_{0,X'}^{\mathbb{T}}(Z') \simeq K_0^{\mathbb{T}}(X')$ , the pushforward  $f_* : K_{0,X}^{\mathbb{T}}(Z) \rightarrow K_{0,X'}^{\mathbb{T}}(Z')$  corresponds to the usual pushforward  $f_* : K_0^{\mathbb{T}}(X) \rightarrow K_0^{\mathbb{T}}(X')$ .  $\square$

*Remark 2.6.* The transformation  $\mathrm{DR}_y^{\mathbb{T}}$  is (in the following sense) independent of the choice of an  $\mathbb{T}$ -equivariant embedding of  $X$  in a smooth ambient  $\mathbb{T}$ -manifold  $Z$ . In the context of Proposition 2.4, with  $f: X \rightarrow X'$  an isomorphism, the pushforward  $f_*: K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)) \rightarrow K_0(\mathrm{MHM}_{X'}^{\mathbb{T}}(Z'))$  is an isomorphism, e.g., using Proposition 3.1 below. In particular, the transformation  $\mathrm{DR}_y^{\mathbb{T}}$  commutes with this isomorphism and the isomorphism  $f_*: K_0^{\mathbb{T}}(X) \rightarrow K_0^{\mathbb{T}}(X')$ , as in the above remark. If we now consider two equivariant embeddings  $i: X \hookrightarrow Z$  and  $i': X \hookrightarrow Z'$ , the above observation can be applied to the diagonal embedding  $X \hookrightarrow Z \times Z'$  (with the diagonal  $\mathbb{T}$ -action) and the projections on the two factors to show that  $\mathrm{DR}_y^{\mathbb{T}}$  is (up to isomorphism) independent of the choice of an equivariant embedding of  $X$  in a smooth ambient  $\mathbb{T}$ -manifold.  $\square$

*Example 2.7* (Degree of  $\mathrm{DR}_y^{\mathbb{T}}$ ). Let us illustrate the definition of the equivariant Hodge-Chern transformation when  $X = Z = pt$  is a point space. In this case, since  $\mathbb{T}$  is connected, we get by [31, Lem.1.2] that

$$\mathrm{MHM}^{\mathbb{T}}(pt) = \mathrm{MHM}(pt) = \mathrm{MHS}^p,$$

where  $\mathrm{MHS}^p$  is the category of graded-polarizable mixed  $\mathbb{Q}$ -Hodge structures, with switching the increasing  $\mathcal{D}$ -module filtration to a decreasing Hodge filtration so that on a point space this identification gives  $\mathrm{Gr}_{-p}^F \mathrm{DR}_Z = \mathrm{Gr}_F^p$ .

By definition, for  $H^\bullet \in D_{\mathrm{MHM}}^{b,\mathbb{T}}(pt)$  we have that

$$[H^\bullet] \in K_0(D_{\mathrm{MHM}}^{b,\mathbb{T}}(pt)) = K_0(\mathrm{MHM}^{\mathbb{T}}(pt)) = K_0(\mathrm{MHS}^p),$$

so  $\mathrm{DR}_y^{\mathbb{T}}([H^\bullet])$  does not see the  $\mathbb{T}$ -action, i.e.,

$$(2.1) \quad \mathrm{DR}_y^{\mathbb{T}}([H^\bullet]) = \sum_{j,p} (-1)^j \dim_{\mathbb{C}} \mathrm{Gr}_F^p H_{\mathbb{C}}^j(-y)^p =: \chi_y([H^\bullet])$$

is the corresponding Hodge  $\chi_y$ -polynomial, defining a ring homomorphism

$$\chi_y: K_0(\mathrm{MHS}^p) \longrightarrow K_0(pt)[y^{\pm 1}] = \mathbb{Z}[y^{\pm 1}] \hookrightarrow K_0^{\mathbb{T}}(pt)[y^{\pm 1}].$$

E.g., for  $n \in \mathbb{Z}$ ,  $\chi_y(\mathbb{Q}(n)) = (-y)^{-n}$ , where  $\mathbb{Q}(n)$  denotes the Tate Hodge structure of weight  $-2n$  on the vector space  $\mathbb{Q}$ . Similarly,

$$\chi_y(X) := \chi_y([H_c^\bullet(X; \mathbb{Q}_X^H)])$$

is the (compactly supported) Hodge polynomial of  $X$ . E.g.,  $\chi_y((\mathbb{C}^*)^n) = (-y-1)^n$ .

By functoriality, if  $Z$  (and hence  $X$ ) is compact, and  $a_Z: Z \rightarrow pt$  is the constant map to a point, we have for  $\mathcal{M} \in D_{\mathrm{MHM},X}^{b,\mathbb{T}}(Z)$  that  $k_* \mathcal{M} \in D_{\mathrm{MHM}}^{b,\mathbb{T}}(pt)$ , so

$$(2.2) \quad \chi_y(X; [\mathcal{M}]) := \chi_y([(a_Z)_* \mathcal{M}]) = \chi_y([H^\bullet(Z; \mathcal{M})])$$

is the degree of the equivariant Hodge-Chern class of  $[\mathcal{M}]$  (and depends only of  $[\mathcal{M}]$ ).  $\square$

**Definition 2.8.** Let  $K_0(\mathrm{MHS}^p)^{\mathrm{Tate}}$  be the subring of  $K_0(\mathrm{MHS}^p) = K_0(\mathrm{MHM}^{\mathbb{T}}(pt))$  generated by  $u := [\mathbb{Q}(-1)]$  and  $u^{-1} := [\mathbb{Q}(1)]$ , with a surjective ring homomorphism  $\mathbb{Z}[u^{\pm 1}] \rightarrow K_0(\mathrm{MHS}^p)^{\mathrm{Tate}}$ . This is an isomorphism since its composition with  $\chi_y$  is the isomorphism  $\mathbb{Z}[u^{\pm 1}] \rightarrow \mathbb{Z}[y^{\pm 1}]$ ,  $u \mapsto -y$ .

*Remark 2.9.* Via the description of  $\mathrm{DR}_y^{\mathbb{T}}$  over a point space, it follows that the transformation  $\mathrm{DR}_y^{\mathbb{T}}$  commutes with the cross-product  $\boxtimes$  with a point space for the identification  $Z \times pt \simeq Z$ , since the equivariant Hodge filtration on the product  $\mathcal{M} \boxtimes H$  (with  $\mathcal{M} \in \mathrm{MHM}_X^{\mathbb{T}}(Z)$  and  $H \in \mathrm{MHS}$ ) is the product filtration on the underlying filtered  $\mathcal{D}$ -module, e.g., see the discussion in [31, Sections 1.4 and 1.5]. Let us also note that the effect of the Tate twist  $(n)$  on  $\mathrm{DR}_y^{\mathbb{T}}$  is (like in the case of  $\mathrm{DR}_y$ ) just multiplication by  $(-y)^{-n}$ , i.e., for  $\mathcal{M} \in D_{\mathrm{MHM},X}^{b,\mathbb{T}}(Z)$  with  $\mathcal{M}(n) := \mathcal{M} \boxtimes \mathbb{Q}(n)$ , one has

$$(2.3) \quad \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}(n)]) = (-y)^{-n} \cdot \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]).$$

□

*Remark 2.10.* Note that for  $i : X \hookrightarrow Z$  a singular closed embedding into an ambient smooth  $\mathbb{T}$ -variety, Achar's 6-functor calculus for equivariant mixed Hodge modules cannot be used to define the constant mixed Hodge complex on  $X$  as a  $\mathbb{T}$ -equivariant object, e.g., via the pullback of  $\mathbb{Q}_Z^H$ . Nevertheless, its Grothendieck class  $[\mathbb{Q}_X^H] \in K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$  can be introduced via stratifications as follows. Let  $\mathcal{S}_X$  be a  $\mathbb{T}$ -invariant algebraic stratification of  $X \subset Z$  with smooth strata, and denote by  $j_{S,Z} : S \hookrightarrow Z$  the inclusion maps of strata  $S \in \mathcal{S}_X$ . Then

$$(2.4) \quad [\mathbb{Q}_X^H] := \sum_{S \in \mathcal{S}_X} [(j_{S,Z})!(j_{S,Z})^*\mathbb{Q}_S^H] = \sum_{S \in \mathcal{S}_X} [(j_{S,Z})!\mathbb{Q}_S^H] \in K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)),$$

where  $\mathbb{Q}_S^H \in D_{\mathrm{MHM}}^{b,\mathbb{T}}(S)$  is the constant mixed Hodge complex on the stratum  $S$ . Then, by additivity,  $[\mathbb{Q}_X^H]$  is independent of the choice of the stratification. Moreover,  $\chi_y(X; [\mathbb{Q}_X^H]) = \chi_y(X)$  is the Hodge polynomial of  $X$ , as defined above. Similarly, we can introduce

$$i^* : K_0(\mathrm{MHM}^{\mathbb{T}}(Z)) \rightarrow K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)),$$

with  $i^* \circ i_* = \mathrm{id}$  on  $K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$  and  $i_* : K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z)) \rightarrow K_0(\mathrm{MHM}^{\mathbb{T}}(Z))$  defined by forgetting the support. So  $i_*$  is injective.

Alternatively, if  $j : U = Z \setminus X \hookrightarrow Z$  is the inclusion map, one may define the equivariant constant Hodge module complex of  $X$  in the ambient  $\mathbb{T}$ -manifold  $Z$  (up to non-canonical isomorphism) as the cone of the adjunction map

$$(2.5) \quad \mathbb{Q}_X^H := \mathrm{Cone}(j!j^*\mathbb{Q}_Z^H \rightarrow \mathbb{Q}_Z^H) \in D_{\mathrm{MHM},X}^{b,\mathbb{T}}(Z).$$

By construction, the equivariant Grothendieck class of  $\mathbb{Q}_X^H$  agrees with (2.4). □

Let us now relate the transformation  $\mathrm{DR}_y^{\mathbb{T}}$  to its motivic counterpart, as used e.g., in [9]. Let  $X$  be a  $\mathbb{T}$ -equivariant complex algebraic variety. The relative equivariant motivic Grothendieck group  $K_0^{\mathbb{T}}(\mathrm{var}/X)$  of varieties over  $X$  is the free abelian on isomorphism classes of  $\mathbb{T}$ -varieties over  $X$ , modulo the usual scissor relations

$$[Y \rightarrow X] = [U \rightarrow X] + [Y \setminus U \rightarrow X],$$

for  $U \subset X$  an open  $\mathbb{T}$ -invariant subvariety. If  $X = pt$  is a point space, then  $K_0^{\mathbb{T}}(\mathrm{var}/pt)$  is a ring with product given by the cross-product of morphisms for  $pt \times X \simeq X$ , and the group  $K_0^{\mathbb{T}}(\mathrm{var}/X)$  is a module over  $K_0^{\mathbb{T}}(\mathrm{var}/pt)$  with respect to the cross product.

For any equivariant morphism  $f : Y \rightarrow X$  of  $\mathbb{T}$ -varieties, there is a well-defined push-forward  $f_! : K_0^{\mathbb{T}}(\text{var}/Y) \rightarrow K_0^{\mathbb{T}}(\text{var}/X)$  defined by composition, which is compatible with the module structure over  $K_0^{\mathbb{T}}(\text{var}/pt)$ .

Assume now that  $i : X \hookrightarrow Z$  is a  $\mathbb{T}$ -equivariant closed embedding into an ambient smooth  $\mathbb{T}$ -manifold  $Z$ . Then, using the same argument as in [14], there is a natural group homomorphism

$$\chi_{Hdg}^{\mathbb{T}} : K_0^{\mathbb{T}}(\text{var}/X) \rightarrow K_0(\text{MHM}_X^{\mathbb{T}}(Z))$$

uniquely fixed by

$$[g : Y \rightarrow X] \mapsto [(i \circ g)_! \mathbb{Q}_Y^H],$$

where  $Y$  is a smooth  $\mathbb{T}$ -variety. For singular  $Y$ , one decomposes it into smooth  $\mathbb{T}$ -invariant subvarieties, and uses the usual scissor relations to define  $\chi_{Hdg}^{\mathbb{T}}([f : Y \rightarrow X])$ , e.g., in the notations of Remark 2.10, we have

$$\chi_{Hdg}^{\mathbb{T}}([i : X \hookrightarrow Z]) = [\mathbb{Q}_X^H] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z)).$$

The transformation  $\chi_{Hdg}^{\mathbb{T}}$  commutes with pushdown  $f_!$  (with proper support) for morphisms of pairs  $f : (Z, X) \rightarrow (Z', X')$ , with  $Z, Z'$  smooth. Hence, the composition

$$mC_y^{\mathbb{T}} := DR_y^{\mathbb{T}} \circ \chi_{Hdg}^{\mathbb{T}} : K_0^{\mathbb{T}}(\text{var}/X) \rightarrow K_0(\text{MHM}_X^{\mathbb{T}}(Z)) \rightarrow K_0^{\mathbb{T}}(X)[y^{\pm 1}]$$

is independent of the ambient smooth  $\mathbb{T}$ -variety  $Z$ , and it is covariant for  $\mathbb{T}$ -equivariant proper morphisms. Moreover, if  $X$  is smooth and  $\mathcal{M} = \mathbb{Q}_X^H \in D_{\text{MHM}}^{b, \mathbb{T}}(X)$  is the constant Hodge module complex on  $X$ , then the underlying  $\mathbb{T}$ -equivariant left  $\mathcal{D}$ -module is  $\mathcal{O}_X$  with the trivial Hodge filtration, so that

$$Gr_{-p}^F DR_X(\mathbb{Q}_X^H) \simeq \Omega_X^p[-p]$$

for  $0 \leq p \leq \dim(X)$ , and it is 0 otherwise. So,

$$mC_y^{\mathbb{T}}([id_X]) = \sum_{p=0}^{\dim(X)} [\Omega_X^p]_{\mathbb{T}} \cdot y^p =: \Lambda_y(\Omega_X^1),$$

the total  $\Lambda$ -class of the cotangent bundle of  $X$ . It then follows that, for  $X$  quasi-projective,  $mC_y^{\mathbb{T}}$  induces the equivariant motivic Chern class transformation defined in [2, Thm.4.2] on the  $\mathbb{T}$ -equivariant relative Grothendieck group  $K_0^{\mathbb{T}}(qproj/X)$  of  $\mathbb{T}$ -equivariant quasi-projective varieties over  $X$ , as studied in the quasi-projective toric context in [9].

In general, if  $i : X \hookrightarrow Z$  is a  $\mathbb{T}$ -equivariant closed embedding into an ambient smooth  $\mathbb{T}$ -variety, we set

$$mC_y^{\mathbb{T}}(X) := mC_y^{\mathbb{T}}([id_X]) = DR_y^{\mathbb{T}}([\mathbb{Q}_X^H])$$

for  $[\mathbb{Q}_X^H] \in K_0^{\mathbb{T}}(\text{MHM}_X(Z))$  the class defined as in (2.4). Moreover, if  $X$  is pure-dimensional we set

$$IC_y^{\mathbb{T}}(X) := DR_y^{\mathbb{T}}([IC_X^H]),$$

with  $IC_X^H := IC_X^H[-\dim(X)]$  the (shifted) equivariant intersection cohomology module of  $X$ , viewed as a (shifted)  $\mathbb{T}$ -equivariant Hodge module on  $Z$ . If  $X$  is compact, the degree of

$IC_y^{\mathbb{T}}(X)$  is the  $I\chi_y$ -polynomial

$$I\chi_y(X) := \chi_y(X; IC_X^H).$$

If  $X$  is moreover a rational homology manifold (e.g., a complex algebraic  $V$ -manifold like a simplicial toric variety), then  $[IC_X^H] \simeq [\mathbb{Q}_X^H]$  (which also motivates our shift convention).

*Remark 2.11.* As shown in [26, Thm.4.2], one has the following identification in  $D_{\text{coh}}^b(X)$ :

$$(2.6) \quad \underline{\underline{\Omega}}_X^p[-p] \simeq Gr_{-p}^F DR_Z(\mathbb{Q}_X^H),$$

where

$$\underline{\underline{\Omega}}_X^p := Gr_F^p(\underline{\underline{\Omega}}_X^\bullet)[p] \in D_{\text{coh}}^b(X)$$

are the graded parts of the filtered Du Bois complex  $(\underline{\underline{\Omega}}_X^\bullet, F)$  on  $X$ . Viewing the class  $[\mathbb{Q}_X^H] \in K_0(\text{MHM}_{\mathbb{T}}^{\mathbb{T}}(Z))$ , and considering  $Gr_{-p}^F DR_Z$  as a transformation on this Grothendieck group, one can take (2.6) as the definition of the equivariant classes  $[\underline{\underline{\Omega}}_X^p]_{\mathbb{T}}$  of the graded parts of the Du Bois complex on  $X$ . This yields the formula

$$(2.7) \quad mC_y^{\mathbb{T}}(X) = \sum_{p=0}^{\dim(X)} [\underline{\underline{\Omega}}_X^p]_{\mathbb{T}} \cdot y^p \in K_0^{\mathbb{T}}(X)[y]$$

in the language of equivariant mixed Hodge modules. Moreover, for such a  $\mathbb{T}$ -invariant closed subvariety  $X$  in an ambient smooth  $\mathbb{T}$ -variety  $Z$ , we can use the equivariant de Rham complex and Remark 2.1 together with (2.5) to define

$$\underline{\underline{\Omega}}_X^p[-p] \simeq Gr_{-p}^F DR_Z(\mathbb{Q}_X^H) \in D_{\text{coh}}^{b, \mathbb{T}}(X),$$

as graded pieces of an equivariant Du Bois complex on  $X$ . If  $X$  is, moreover, a rational homology manifold, we have  $IC_X^H \simeq \mathbb{Q}_X^H \in D_{\text{MHM}, X}^{b, \mathbb{T}}(Z)$ , defining an equivariant Du Bois complex on  $X$  in terms of the equivariant intersection cohomology module, as used e.g., in [19]. In the case of a rational homology manifold or if  $X$  is any quasi-projective toric variety,  $\underline{\underline{\Omega}}_X^p$  is just the  $\mathbb{T}$ -equivariant coherent sheaf (sitting in degree zero, since this is the case nonequivariantly) given by the  $\mathbb{T}$ -equivariant Zariski sheaf  $\widehat{\Omega}_X^p$  of  $p$ -forms on  $X$ . This holds even equivariantly, since the adjunction map  $\underline{\underline{\Omega}}_X^p \rightarrow \widehat{\Omega}_X^p = j_* j^* \underline{\underline{\Omega}}_X^p = j_* \Omega_{X_{\text{reg}}}^p$  for the open inclusion  $j$  of the regular part  $X_{\text{reg}}$  of  $X$  is nonequivariantly an isomorphism of reflexive sheaves.

In [9], another approach based on simplicial resolutions was used to derive (2.7), which in the quasi-projective toric situation yields (cf. [9, Prop.3.5])

$$(2.8) \quad mC_y^{\mathbb{T}}(X) = \sum_{p=0}^{\dim(X)} [\widehat{\Omega}_X^p]_{\mathbb{T}} \cdot y^p \in K_0(X)[y].$$

□

Denote by  $\mathbb{L} := [\mathbb{C} \rightarrow pt] \in K_0^{\mathbb{T}}(var/pt)$  the Lefschetz motive (with  $\mathbb{T}$  acting trivially on  $\mathbb{C}$ ). Then there is a natural group homomorphism on the localized Grothendieck group

$$\chi_{Hdg}^{\mathbb{T}} : K_0^{\mathbb{T}}(var/X)[\mathbb{L}^{-1}] \rightarrow K_0(\mathbf{MHM}_X^{\mathbb{T}}(Z)),$$

mapping  $\mathbb{L}$  to  $u = [\mathbb{Q}(-1)] \in K_0(\mathbf{MHS}^p) = K_0(\mathbf{MHM}^{\mathbb{T}}(pt))$ . Under the natural forgetful functors, this reduces to the corresponding localized nonequivariant transformation (as used in [22]).

*Remark 2.12* (Effect of duality). As shown in [28, Cor.5.19, Rem.5.20], the non-equivariant transformation  $\mathbf{DR}_y$  of [6, 28] commutes with duality, i.e.,

$$(2.9) \quad \mathbf{DR}_y \circ \mathbb{D}_X = \mathbb{D} \circ \mathbf{DR}_y,$$

where  $\mathbb{D}_X$  is induced from the duality of mixed Hodge modules, and  $\mathbb{D}$  on  $K_0(X)$  is induced by Grothendieck duality (e.g., see [18, Part 1, Sect.7], [20, Part 1]) extended to  $K_0(X)[y^{\pm 1}]$  by  $y \mapsto 1/y$ . A similar statement holds for the motivic transformation  $mC_y$ , with duality on the localization  $K_0(var/X)[\mathbb{L}^{-1}]$  defined as in [28, Section 4B] and [4]. All these duality transformations are functorial for proper maps, so that for  $X$  compact with  $a_X : X \rightarrow pt$  proper, one gets

$$(2.10) \quad \chi_y(X; \mathbb{D}_X[\mathcal{M}]) = \chi_{1/y}(X; [\mathcal{M}]).$$

Indeed, over a point space, Grothendieck duality and homological duality are trivial, while the mixed Hodge module duality is the usual duality of mixed Hodge structures, e.g.,  $\mathbb{D}_{pt}\mathbb{Q}(n) = \mathbb{Q}(-n)$ .  $\square$

As explained in [14, Sect.3], the duality formula (2.9) also holds equivariantly for complexes of mixed Hodge modules on  $Z$  with support in  $X$  (with  $\mathbb{D}_Z$  instead of  $\mathbb{D}_X$ ), at least on (Tate twists of) elements in the image of  $\chi_{Hdg}^{\mathbb{T}}$  (and this is the only case we'll be interested in later on), namely on such elements one has

$$(2.11) \quad \mathbf{DR}_y^{\mathbb{T}} \circ \mathbb{D}_Z^{\mathbb{T}} = \mathbb{D}^{\mathbb{T}} \circ \mathbf{DR}_y^{\mathbb{T}},$$

Here,  $\mathbb{D}_Z^{\mathbb{T}}$  is induced from the exact Verdier duality functor of equivariant mixed Hodge modules (on  $Z$ , with support on  $X$ ), see [1]. For  $\mathbb{D}^{\mathbb{T}}$ , one can use either the equivariant Grothendieck duality on the Grothendieck group  $K_{0,X}^{\mathbb{T}}(Z)$  induced from Grothendieck duality on  $Z$  (remembering the support in  $X$ ), or the equivariant Grothendieck duality on the Grothendieck group  $K_0^{\mathbb{T}}(X)$  induced from Grothendieck duality on  $X$ . These are identified by the isomorphism  $i_* : K_0^{\mathbb{T}}(X) \rightarrow K_{0,X}^{\mathbb{T}}(Z)$ , since equivariant Grothendieck duality commutes with proper pushforwards (see [20] for more details). As in the nonequivariant context,  $\mathbb{D}^{\mathbb{T}}$  is extended to  $K_0^{\mathbb{T}}(X)[y^{\pm 1}]$  by  $y \mapsto 1/y$ .

*Remark 2.13.* To the authors' knowledge, a corresponding duality transformation on the localized equivariant group  $K_0^{\mathbb{T}}(var/X)[\mathbb{L}^{-1}]$  is not currently available.  $\square$

## 3. COMPUTATIONS VIA STRATIFICATIONS

Let us now indicate how to compute the equivariant Hodge-Chern classes in terms of a  $\mathbb{T}$ -invariant stratification.

Let as before  $X$  be a closed  $\mathbb{T}$ -invariant algebraic subvariety of an algebraic  $\mathbb{T}$ -manifold  $Z$ , with inclusion map  $i : X \hookrightarrow Z$ . Let  $\mathcal{M} \in D_{\text{MHM}, X}^{b, \mathbb{T}}(Z)$  be an equivariant complex of mixed Hodge modules on  $Z$  with support in  $X$ , with underlying bounded constructible complex  $K := \text{rat}(\text{For}(\mathcal{M})) \in D_c^b(Z)$  supported on  $X$ . Let  $\mathcal{S}_X = \{S\}$  be a  $\mathbb{T}$ -invariant complex algebraic stratification of  $X$  so that for any stratum  $S \in \mathcal{S}$ ,  $S$  is smooth,  $\bar{S} \setminus S$  is a union of strata, and the sheaves  $L_{S, \ell} := \mathcal{H}^\ell K|_S$  are local systems on  $S$  for any  $\ell \in \mathbb{Z}$ . Let  $U = Z \setminus X$  and  $\mathcal{S}_Z = \mathcal{S}_X \cup \{U\}$ . Then  $\mathcal{S}_Z$  is an algebraic stratification of  $Z$  satisfying the above properties for each  $S \in \mathcal{S}_X$ , together with  $L_{U, \ell} := \mathcal{H}^\ell K|_U \simeq 0$ . For each  $S \in \mathcal{S}_X$ , consider the inclusions

$$j_{S, Z} : S \xrightarrow{i_{S, \bar{S}}} \bar{S} \xrightarrow{i_{\bar{S}, X}} X \xrightarrow{i} Z$$

together with  $i_{\bar{S}, Z} = i \circ i_{\bar{S}, X}$ , so that

$$j_{S, Z} = i_{\bar{S}, Z} \circ i_{S, \bar{S}}.$$

Under the above notations and assumptions, and using the 6-functor formalism from [1] we get as in [23, Prop.5.1.2] the following:

**Proposition 3.1.**

$$(3.1) \quad [\mathcal{M}] = \sum_{S \neq U, \ell} (-1)^\ell \cdot [(j_{S, Z})! L_{S, \ell}^H] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z)),$$

where  $L_{S, \ell}^H = H^{\ell + \dim(S)}(j_{S, Z})^* \mathcal{M}[-\dim(S)]$  is the shifted equivariant smooth mixed Hodge module on  $S$  with  $L_{S, \ell}$  the underlying local system of  $\text{For}(L_{S, \ell}^H)$ , and the summation is over strata  $S \in \mathcal{S}_X$ .

In the notations of Proposition 3.1, we get the following.

**Corollary 3.2.** *Assume, moreover, that for any  $S \in \mathcal{S}_X$ , the stabilizer  $\mathbb{T}_s \subset \mathbb{T}$  at a point  $s \in S$  is connected. Then each  $L_{S, \ell}^H \simeq \mathbb{Q}_S^H \boxtimes L_{S, s, \ell}^H$  is a constant variation of mixed Hodge structures on  $S$  with stalk the mixed Hodge structure  $L_{S, s, \ell}^H$ , and we have*

$$(3.2) \quad \begin{aligned} [\mathcal{M}] &= \sum_{S, \ell} (-1)^\ell \cdot [(j_{S, Z})! \mathbb{Q}_S^H] \cdot [L_{S, s, \ell}^H] \\ &= \sum_S [(j_{S, Z})! \mathbb{Q}_S^H] \cdot i_s^*[\mathcal{M}], \end{aligned}$$

with  $i_s : \{s\} \hookrightarrow Z$  the inclusion, for  $s \in S$  chosen, and the summation is over the strata  $S \in \mathcal{S}_X$ .

In particular,

$$\begin{aligned}
(3.3) \quad \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]) &= \sum_{S \in \mathcal{S}_X} \mathrm{DR}_y^{\mathbb{T}}[(j_{S,Z})! \mathbb{Q}_S^H] \cdot \chi_y(i_s^*[\mathcal{M}]) \\
&= \sum_{S \in \mathcal{S}_X} m\mathcal{C}_y^{\mathbb{T}}([j_{S,Z} : S \hookrightarrow Z]) \cdot \chi_y(i_s^*[\mathcal{M}]) \\
&= \sum_{S \in \mathcal{S}_X} (i_{\bar{S},Z})_* m\mathcal{C}_y^{\mathbb{T}}([i_{S,\bar{S}} : S \hookrightarrow \bar{S}]) \cdot \chi_y(i_s^*[\mathcal{M}]).
\end{aligned}$$

*Proof.* The fact that each  $L_{S,\ell}^H$  is a constant variation of mixed Hodge structures on  $S$  follows by [31, Lem.1.2] using the assumption that the stabilizer  $\mathbb{T}_s$  of  $s \in S$  is connected. This proves (3.2). To show (3.3), we apply the equivariant Hodge-Chern class transformation  $\mathrm{DR}_y^{\mathbb{T}}$  to (3.2), using the fact that  $K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$  is a  $K_0(\mathrm{MHM}^{\mathbb{T}}(pt)) \simeq K_0(\mathrm{MHS}^p)$ -module, together with the functoriality of the equivariant motivic Chern class transformation  $m\mathcal{C}_y^{\mathbb{T}}$  for the closed embedding  $i_{\bar{S},Z}$ .  $\square$

*Remark 3.3.* Formula (3.2) shows that under the connected  $\mathbb{T}$ -stabilizers assumption, the class  $[\mathcal{M}]$  belongs to the (free)  $K_0(\mathrm{MHM}^{\mathbb{T}}(pt)) = K_0(\mathrm{MHS}^p)$ -submodule (respectively,  $K_0(\mathrm{MHS}^p)^{\mathrm{Tate}}$ -submodule, if the stalks  $i_s^*[\mathcal{M}]$  are of Tate type) of  $K_0(\mathrm{MHM}_X^{\mathbb{T}}(Z))$  generated by classes  $[(j_{S,Z})! \mathbb{Q}_S^H] = \chi_{\mathrm{Hdg}}^{\mathbb{T}}([j_{S,Z} : S \hookrightarrow Z])$ ,  $S \in \mathcal{S}_X$ . In what follows, we regard  $\chi_y(i_s^*[\mathcal{M}]) \in \mathbb{Z}[y^{\pm 1}]$  as a weight (depending only on  $[\mathcal{M}]$ ) attached to the stratum  $S \in \mathcal{S}_X$  containing  $s$ . In the case of a projective toric variety  $X = X_P$  associated to a full dimensional lattice polytope  $P$  (as in the Introduction), the subgroup

$$\chi_{\mathrm{Hdg}}^{\mathbb{T}}\left(\tau^{\mathbb{T}}(F^{\mathbb{T}}(X)[y^{\pm 1}])\right) \subset K_0^{\mathbb{T}}(\mathrm{MHM}_X(Z))$$

is the  $K_0(\mathrm{MHS}^p)^{\mathrm{Tate}}$ -submodule generated by classes  $[\mathcal{M}]$  whose stalks are of Tate type.  $\square$

*Remark 3.4.* Under the assumption of connected  $\mathbb{T}$ -stabilizers, formula (3.3) reduces the calculation of equivariant Hodge-Chern classes to the motivic calculus of [9], up to assigning the above mentioned Laurent polynomial weights  $\chi_y(i_s^*[\mathcal{M}])$  to each stratum  $S$ , with  $s \in S$  chosen. For instance, the connected stabilizers assumption is automatically satisfied in the toric context, where such stabilizers are tori.  $\square$

By functoriality, the terms  $m\mathcal{C}_y^{\mathbb{T}}([S \hookrightarrow \bar{S}])$  appearing in formula (3.3) are computed via resolutions of singularities as follows (see [9, Thm.3.3], adapting ideas from [32, Thm.5.1, Rem.5.2] to the equivariant context):

**Proposition 3.5.** *Let  $i_{S,V} : S \hookrightarrow V$  be a smooth partial  $\mathbb{T}$ -equivariant compactification of a stratum  $S$  so that  $D := V \setminus S$  is a  $\mathbb{T}$ -invariant simple normal crossing divisor, and  $i_{S,\bar{S}} = \pi_V \circ i_{S,V}$  for a proper morphism  $\pi_V : V \rightarrow \bar{S}$ . Then we have*

$$(3.4) \quad m\mathcal{C}_y^{\mathbb{T}}([S \hookrightarrow \bar{S}]) = (\pi_V)_* [\mathcal{O}_V(-D) \otimes \Lambda_y \Omega_V^1(\log(D))]_{\mathbb{T}} \in K_0^{\mathbb{T}}(\bar{S})[y].$$

These formulae become more explicit and much simplified in the toric context, as it will be discussed in Section 5 below.

## 4. TWISTED EQUIVARIANT HODGE-CHERN CLASSES AND THEIR DEGREES

Assume now that a torus  $\mathbb{T}$  with character lattice  $M$  acts linearly on a finite dimensional complex vector space  $W$ , with eigenspace

$$(4.1) \quad W_{\chi^m} := \{w \in W \mid t \cdot w = \chi^m(t)w \text{ for all } t \in \mathbb{T}\},$$

for  $\chi^m$  a character of  $\mathbb{T}$ . Then, as in [11, Prop.1.1.2], one has an eigenspace decomposition

$$W = \bigoplus_{m \in M} W_{\chi^m}.$$

This induces an isomorphism

$$(4.2) \quad K_{\mathbb{T}}^0(pt) \xrightarrow{\cong} \mathbb{Z}[M], \quad [W]_{\mathbb{T}} \mapsto \sum_{m \in M} \dim_{\mathbb{C}} W_{\chi^m} \cdot \chi^m,$$

identifying the duality involution  $(-)^{\vee}$  on the left hand side with the involution  $m \mapsto -m$  on characters. If  $X$  is a compact algebraic  $\mathbb{T}$ -variety, with a  $\mathbb{T}$ -equivariant coherent sheaf  $\mathcal{F}$ , then the cohomology spaces  $H^i(X; \mathcal{F})$  are finite dimensional  $\mathbb{T}$ -representations, vanishing for  $i$  large enough. The *K-theoretic Euler characteristic* of  $\mathcal{F}$  is then defined as:

$$(4.3) \quad \begin{aligned} \chi^{\mathbb{T}}(X, \mathcal{F}) &:= (a_X)_* [\mathcal{F}]_{\mathbb{T}} = [H^{\bullet}(X; \mathcal{F})]_{\mathbb{T}} := \sum_i (-1)^i [H^i(X; \mathcal{F})]_{\mathbb{T}} \in K_{\mathbb{T}}^0(pt) \\ &= \sum_{m \in M} \sum_{i=0}^{\dim(X)} (-1)^i \dim_{\mathbb{C}} H^i(X; \mathcal{F})_{\chi^m} \cdot \chi^m \in \mathbb{Z}[M], \end{aligned}$$

with  $a_X : X \rightarrow pt$  the constant map to a point. Note that  $\chi^{\mathbb{T}}(X, \mathcal{F})$  depends only on the class  $[\mathcal{F}] \in K_0^{\mathbb{T}}(X)$ .

Let now  $X$  be a closed  $\mathbb{T}$ -invariant compact algebraic subvariety of an algebraic  $\mathbb{T}$ -manifold  $Z$ , and let  $D$  be a  $\mathbb{T}$ -invariant Cartier divisor on  $X$ . Let  $[\mathcal{M}] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z))$  be fixed. Then

$$\text{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(D) := \text{DR}_y^{\mathbb{T}}([\mathcal{M}]) \cdot [\mathcal{O}_X(D)] \in K_0^{\mathbb{T}}(X)[y^{\pm 1}],$$

where we extend the  $K_0^{\mathbb{T}}(X)$ -module structure of  $K_0^{\mathbb{T}}(X)$  induced from the tensor product linearly over  $\mathbb{Z}[y^{\pm 1}]$ . For  $a_X : X \rightarrow pt$  the constant map as above, we can then define the  *$\mathbb{T}$ -equivariant Hodge polynomial of  $(X, D, [\mathcal{M}])$*  as:

$$(4.4) \quad \begin{aligned} \chi_y^{\mathbb{T}}(X, D; [\mathcal{M}]) &:= (a_X)_* \left( \text{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(D) \right) \in K_0^{\mathbb{T}}(pt)[y^{\pm 1}] = K_{\mathbb{T}}^0(pt)[y^{\pm 1}] \\ &=: \chi^{\mathbb{T}}(X; \text{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(D)) \in \mathbb{Z}[M][y^{\pm 1}], \end{aligned}$$

which only depends on  $[\mathcal{M}]$ . In other words,  $\chi_y^{\mathbb{T}}(X, D; [\mathcal{M}])$  is an incarnation (in terms of the characters of  $\mathbb{T}$ ) of the degree of the Hodge-Chern class of  $[\mathcal{M}]$  twisted by  $\mathcal{O}_X(D)$ .

*Remark 4.1.* If  $D = 0$ ,

$$(4.5) \quad \begin{aligned} \chi_y^{\mathbb{T}}(X, 0; [\mathcal{M}]) &= (a_X)_* \text{DR}_y^{\mathbb{T}}([\mathcal{M}]) = \chi_y(X; [\mathcal{M}]) \in \mathbb{Z}[y^{\pm 1}] \\ &= \chi_y(X; [\mathcal{M}]) \cdot \chi^0 \in \mathbb{Z}[M][y^{\pm 1}], \end{aligned}$$

is the usual Hodge polynomial of  $[\mathcal{M}]$  as in (2.2).  $\square$

*Example 4.2.* If  $X$  is smooth and  $\mathcal{M} = \mathbb{Q}_X^H$  is the constant Hodge module complex (viewed as a Hodge module complex on  $Z$ ), then

$$\chi_y^{\mathbb{T}}(X, D; [\mathbb{Q}_X^H]) = \sum_{p=0}^{\dim(X)} \chi^{\mathbb{T}}(X, \Omega_X^p \otimes \mathcal{O}_X(D)) \cdot y^p,$$

which is exactly  $\chi_y^{\mathbb{T}}(X, \mathcal{O}_X(D))$ , the equivariant Hirzebruch polynomial of  $D$ , e.g., see [9, Sect.3.2].  $\square$

*Example 4.3.* Assume  $X$  is a toric variety or a complex algebraic  $V$ -manifold. Then

$$\chi_y^{\mathbb{T}}(X, D; [\mathbb{Q}_X^H]) = \sum_{p=0}^{\dim(X)} \chi^{\mathbb{T}}(X, \widehat{\Omega}_X^p \otimes \mathcal{O}_X(D)) \cdot y^p,$$

where  $\widehat{\Omega}_X^p$  denotes the corresponding sheaf of Zariski differential  $p$ -forms, is the Hodge polynomial of  $D$  considered in [9, Sect.3.2] in the toric context.  $\square$

## 5. CHARACTERISTIC CLASS FORMULAE IN TORIC GEOMETRY

In this section, we specialize the formulae of previous sections to the case of toric varieties (e.g., associated to a full-dimensional lattice polytope), which are known to be Whitney stratified by the orbits of the torus action.

Let us first introduce some notation. Let  $X = X_{\Sigma}$  be a quasi-projective toric variety associated to the fan  $\Sigma \subset N_{\mathbb{R}} \cong N \otimes \mathbb{R} = \mathbb{R}^n$ , with torus  $\mathbb{T} = N \otimes_{\mathbb{Z}} \mathbb{C}^* = (\mathbb{C}^*)^n$  having  $M$ , the dual of the lattice  $N \cong \mathbb{Z}^n$ , as the character lattice. The action of the torus  $\mathbb{T}$  on itself extends to an algebraic action of  $\mathbb{T}$  on  $X$  with finitely many orbits  $O_{\sigma}$ , one for each cone  $\sigma \in \Sigma$ . In fact, to a  $k$ -dimensional cone  $\sigma \in \Sigma$  there corresponds an  $(n-k)$ -dimensional torus-orbit  $O_{\sigma} \cong (\mathbb{C}^*)^{n-k}$ , and the orbit closure  $V_{\sigma}$  of  $O_{\sigma}$  is itself a toric variety and a  $\mathbb{T}$ -invariant subvariety of  $X$ . Let  $N(\sigma) = N/N_{\sigma}$ , with  $N_{\sigma}$  denoting the sublattice of  $N$  spanned by the points in  $\sigma \cap N$ . Let  $\mathbb{T}_{N(\sigma)} = N(\sigma) \otimes_{\mathbb{Z}} \mathbb{C}^*$  be the torus associated to  $N(\sigma)$ . For each cone  $\nu \in \Sigma$  containing  $\sigma$  (in short,  $\sigma \preceq \nu$ ), let  $\bar{\nu}$  be the image cone in  $N(\sigma)_{\mathbb{R}}$  under the quotient map  $N_{\mathbb{R}} \rightarrow N(\sigma)_{\mathbb{R}}$ . Then

$$(5.1) \quad \text{Star}(\sigma) = \{\bar{\nu} \subseteq N(\sigma)_{\mathbb{R}} \mid \sigma \preceq \nu\}$$

is a fan in  $N(\sigma)_{\mathbb{R}}$ , with associated toric variety isomorphic to  $V_{\sigma}$  (see, e.g., [11, Prop.3.2.7]), and  $\mathbb{T}$  acts on  $V_{\sigma}$  via the morphism

$$(5.2) \quad \mathbb{T} \rightarrow \mathbb{T}_{N(\sigma)}$$

induced by the quotient map  $N \rightarrow N(\sigma)$ .

Before specializing formula (3.3) to this toric context, we need the following key additional input derived in [9]:

**Proposition 5.1.** [9, Prop.3.8] *Let  $X = X_{\Sigma}$  be a quasi-projective toric variety defined by the fan  $\Sigma$ . For any cone  $\sigma \in \Sigma$ , with orbit  $O_{\sigma}$  and inclusion  $i_{\sigma} : O_{\sigma} \hookrightarrow V_{\sigma}$  in the orbit closure, we have:*

$$(5.3) \quad mC_y^{\mathbb{T}}([i_{\sigma}]) = (1+y)^{\dim(O_{\sigma})} \cdot [\omega_{V_{\sigma}}]_{\mathbb{T}},$$

where  $\omega_{V_\sigma} = \widehat{\Omega}_{V_\sigma}^{\dim(O_\sigma)}$  is the canonical sheaf on the toric variety  $V_\sigma$ , viewed as a  $\mathbb{T}$ -equivariant sheaf via the quotient map  $\mathbb{T} \rightarrow \mathbb{T}_{N(\sigma)}$  given by (5.2).

*Proof.* This follows from Proposition 3.5 by using a toric resolution  $V$  of  $V_\sigma$ , together with the fact that the locally free sheaf  $\Omega_V^1(\log(D))$  is  $\mathbb{T}_{N(\sigma)}$ -equivariantly, hence also  $\mathbb{T}$ -equivariantly, a trivial sheaf of rank  $\dim(O_\sigma)$ , and there is a  $\mathbb{T}_{N(\sigma)}$ -equivariant, hence also  $\mathbb{T}$ -equivariant isomorphism  $R(\pi_V)_* \mathcal{O}_V(-D) \simeq \omega_{V_\sigma}$ .  $\square$

In the following we assume that  $X = X_\Sigma$  is a quasi-projective toric variety. Then [30] guarantees the existence of some smooth  $\mathbb{T}$ -equivariant ambient variety  $Z$  with a closed  $\mathbb{T}$ -equivariant embedding  $i : X \hookrightarrow Z$ . Note also that the torus stabilizer at any point in a torus orbit of  $X$  is itself a torus, hence *connected*.

Together with formula (3.3), Proposition 5.1 then gives the following:

**Theorem 5.2.** *Let  $X = X_\Sigma$  be a quasi-projective toric variety defined by the fan  $\Sigma$ , with torus  $\mathbb{T}$ . For each cone  $\sigma \in \Sigma$  with orbit  $O_\sigma$ , let  $k_\sigma : V_\sigma \hookrightarrow Z$  be the inclusion of the orbit closure in the smooth ambient  $\mathbb{T}$ -manifold  $Z$  mentioned above, and fix a point  $x_\sigma \in O_\sigma$  with inclusion  $i_{x_\sigma} : \{x_\sigma\} \hookrightarrow Z$ . Let  $\mathcal{M} \in D_X^{b,\mathbb{T}}(Z)$  be a  $\mathbb{T}$ -equivariant mixed Hodge module complex on  $Z$  with support in  $X$ . Then:*

$$(5.4) \quad \mathrm{DR}_y^\mathbb{T}([\mathcal{M}]) = \sum_{\sigma \in \Sigma} \chi_y(i_{x_\sigma}^*[\mathcal{M}]) \cdot (1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\omega_{V_\sigma}]_{\mathbb{T}}.$$

In particular, by taking degrees,

$$(5.5) \quad \chi_y(X; [\mathcal{M}]) = \sum_{\sigma \in \Sigma} \chi_y(i_{x_\sigma}^*[\mathcal{M}]) \cdot (-1-y)^{\dim(O_\sigma)}.$$

Theorem 5.2 yields the following:

**Corollary 5.3.** *Let  $X = X_\Sigma$  be the quasi-projective toric variety defined by a fan  $\Sigma$ , with a chosen closed  $\mathbb{T}$ -equivariant embedding into a smooth ambient  $\mathbb{T}$ -variety  $Z$ . For each cone  $\sigma \in \Sigma$  with orbit  $O_\sigma$ , fix a point  $x_\sigma \in O_\sigma$ . Then we have the following formula:*

$$(5.6) \quad \mathrm{DR}_y^\mathbb{T}([IC_X^H]) = \sum_{\sigma \in \Sigma} \chi_y(IC_X^H|_{x_\sigma}) \cdot (1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\omega_{V_\sigma}]_{\mathbb{T}}.$$

Here,  $IC_X^H \in \mathrm{MHM}_X^\mathbb{T}(Z)$  is regarded as a  $\mathbb{T}$ -equivariant Hodge module on the ambient  $\mathbb{T}$ -manifold  $Z$ , and each class  $[\omega_{V_\sigma}]_{\mathbb{T}}$  is pushed into  $K_{0,X}^\mathbb{T}(Z) \simeq K_0^\mathbb{T}(X)$ .

*Remark 5.4.* In particular, if  $X_\Sigma$  is a simplicial toric variety then  $\chi_y(IC_X^H|_{x_\sigma}) = \chi_y(\mathbb{Q}_X^H|_{x_\sigma}) = 1$  for all  $\sigma \in \Sigma$ , and one recovers in this case a result from [9]. Also note that formula (5.6) for  $X$  an affine toric variety also fits with recent results of the paper [19].  $\square$

For future reference, we reformulate here (5.3) in the language of equivariant mixed Hodge modules. Let  $k_\sigma : V_\sigma \hookrightarrow Z$  be the orbit closure embedding into the smooth ambient  $\mathbb{T}$ -manifold  $Z$  constructed in [30]. Let  $j_\sigma := k_\sigma \circ i_\sigma : O_\sigma \hookrightarrow Z$ .

**Corollary 5.5.** *In the above notations,*

$$(5.7) \quad \mathrm{DR}_y^\mathbb{T}([(j_\sigma)! \mathbb{Q}_{O_\sigma}^H]) = (1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\omega_{V_\sigma}]_{\mathbb{T}}.$$

*Proof.* This follows immediately by applying the proper pushforward  $(k_\sigma)_*$  to both sides of (5.3), using the functoriality of  $mC_y^\mathbb{T}$  and the definition of  $\chi_{Hdg}^\mathbb{T}$ . Indeed,

$$(1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\omega_{V_\sigma}]_\mathbb{T} = (k_\sigma)_*mC_y^\mathbb{T}([i_\sigma]) = mC_y^\mathbb{T}([j_\sigma]) = \mathrm{DR}_y^\mathbb{T}([(j_\sigma)! \mathbb{Q}_{O_\sigma}^H]).$$

□

Before computing  $\mathrm{DR}_y^\mathbb{T}([(j_\sigma)_* \mathbb{Q}_{O_\sigma}^H])$ , we need the following lemma.

**Lemma 5.6.** *For  $\sigma \in \Sigma$  with orbit  $O_\sigma$  and orbit closure  $V_\sigma$ , we have for the equivariant dualizing complex that*

$$(5.8) \quad \mathbb{D}^\mathbb{T}(\mathcal{O}_{V_\sigma}) \simeq \omega_{V_\sigma}[\dim(O_\sigma)].$$

*In particular, by equivariant biduality, we get*

$$(5.9) \quad \mathbb{D}^\mathbb{T}[\omega_{V_\sigma}]_\mathbb{T} = (-1)^{\dim(O_\sigma)} \cdot [\mathcal{O}_{V_\sigma}]_\mathbb{T}.$$

*Proof.* Let  $\tilde{V}_\sigma \xrightarrow{f_\sigma} V_\sigma$  be a toric resolution of singularities of  $V_\sigma$ . Then  $\tilde{V}_\sigma$  is a toric variety obtained by refining the fan of  $V_\sigma$ . Let  $O_\sigma \hookrightarrow \tilde{V}_\sigma$  be the natural open inclusion, with complement the simple normal crossing divisor  $D_\sigma$  whose irreducible components correspond to the rays in the fan of  $\tilde{V}_\sigma$ . Note that  $D_\sigma$  is a  $\mathbb{T}_{N(\sigma)}$ -invariant (hence a  $\mathbb{T}$ -invariant) divisor.

Furthermore, note that the canonical sheaf  $\omega_{\tilde{V}_\sigma}$  on the toric variety  $\tilde{V}_\sigma$  is precisely given by  $\mathcal{O}_{\tilde{V}_\sigma}(-D_\sigma) \simeq \omega_{\tilde{V}_\sigma}$  as  $\mathbb{T}_{N(\sigma)}$ -equivariant, hence also  $\mathbb{T}$ -equivariant sheaves, e.g., see [11, Thm.8.2.3]. And since the toric morphism  $f_\sigma : \tilde{V}_\sigma \rightarrow V_\sigma$  is induced by a refinement of the fan of  $V_\sigma$ , it follows from [11, Thm.8.2.15] that there is a  $\mathbb{T}_{N(\sigma)}$ -equivariant, hence also  $\mathbb{T}$ -equivariant, isomorphism:

$$(5.10) \quad f_{\sigma*} \omega_{\tilde{V}_\sigma} \simeq \omega_{V_\sigma}.$$

Altogether, we obtain the following sequence of isomorphisms:

$$(5.11) \quad \begin{aligned} \mathbb{D}^\mathbb{T}(\mathcal{O}_{V_\sigma}) &\simeq \mathbb{D}^\mathbb{T}\left((Rf_\sigma)_*(\mathcal{O}_{\tilde{V}_\sigma})\right) \\ &\simeq (Rf_\sigma)_*(\mathbb{D}^\mathbb{T}(\mathcal{O}_{\tilde{V}_\sigma})) \\ &\simeq (Rf_\sigma)_*(\omega_{\tilde{V}_\sigma}[\dim(O_\sigma)]) \\ &\simeq \omega_{V_\sigma}[\dim(O_\sigma)], \end{aligned}$$

where the first isomorphism follows since  $V_\sigma$  has only rational singularities (as in the proof of [9, Prop.3.5]); the second isomorphism follows since equivariant Grothendieck duality commutes with proper pushforward (cf. [20, Thm.25.2]); the third isomorphism follows by

$$(5.12) \quad \mathbb{D}^\mathbb{T}(\mathcal{O}_{\tilde{V}_\sigma}) \simeq \omega_{\tilde{V}_\sigma}[\dim(O_\sigma)],$$

for the smooth variety  $\tilde{V}_\sigma$ , cf. [20, Thm.28.11]; finally, the last isomorphism is a consequence of (5.10) and the vanishing of the higher derived image sheaves of  $\omega_{\tilde{V}_\sigma}$ , i.e.,  $R^i(f_\sigma)_* \omega_{\tilde{V}_\sigma} = 0$ , for all  $i > 0$ , see [11, Thm.9.3.12].

By using equivariant biduality, cf. [20, Prop.31.9, Lem.31.11], we get from (5.8) that

$$\mathcal{O}_{V_\sigma} \simeq (\mathbb{D}^\mathbb{T} \omega_{V_\sigma})[-\dim(O_\sigma)],$$

which yields (5.9).  $\square$

In the notations of the above Corollary 5.5, we then have the following

**Proposition 5.7.** *For  $j_\sigma : O_\sigma \hookrightarrow Z$  the orbit inclusion into the ambient  $\mathbb{T}$ -manifold  $Z$ ,*

$$(5.13) \quad \mathrm{DR}_y^\mathbb{T}([(j_\sigma)_* \mathbb{Q}_{O_\sigma}^H]) = (1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\mathcal{O}_{V_\sigma}]_{\mathbb{T}}.$$

*Proof.* Since  $O_\sigma$  is smooth,  $\mathbb{Q}_{O_\sigma}^H$  is a (shifted)  $\mathbb{T}$ -equivariant pure Hodge module of weight  $\dim(O_\sigma)$ , so

$$\mathbb{D}_{O_\sigma}^\mathbb{T} \mathbb{Q}_{O_\sigma}^H \simeq \mathbb{Q}_{O_\sigma}^H[2 \dim(O_\sigma)(\dim(O_\sigma))].$$

Using (2.3) and the fact that the even shift does not affect the Grothendieck class, we get

$$(5.14) \quad \mathrm{DR}_y^\mathbb{T}([(j_\sigma)_* \mathbb{Q}_{O_\sigma}^H]) = (-y)^{\dim(O_\sigma)} \cdot \mathrm{DR}_y^\mathbb{T}(\mathbb{D}_Z^\mathbb{T}[(j_\sigma)! \mathbb{Q}_{O_\sigma}^H]).$$

Since  $(j_\sigma)! \mathbb{Q}_{O_\sigma}^H \in \mathrm{Image}(\chi_{\mathrm{Hdg}}^\mathbb{T})$ , Remark 2.12 gives

$$(5.15) \quad \begin{aligned} \mathrm{DR}_y^\mathbb{T}(\mathbb{D}_Z^\mathbb{T}[(j_\sigma)! \mathbb{Q}_{O_\sigma}^H]) &= \left( \mathbb{D}^\mathbb{T} \left( \mathrm{DR}_y^\mathbb{T}([(j_\sigma)! \mathbb{Q}_{O_\sigma}^H]) \right) \right)_{|y \rightarrow 1/y} \\ &\stackrel{(5.7)}{=} \left( \mathbb{D}^\mathbb{T} \left( (1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\omega_{V_\sigma}]_{\mathbb{T}} \right) \right)_{|y \rightarrow 1/y} \\ &= \left( 1 + \frac{1}{y} \right)^{\dim(O_\sigma)} \cdot (k_\sigma)_* \mathbb{D}^\mathbb{T}[\omega_{V_\sigma}]_{\mathbb{T}} \\ &\stackrel{(5.9)}{=} y^{-\dim(O_\sigma)} \cdot (1+y)^{\dim(O_\sigma)} \cdot (-1)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\mathcal{O}_{V_\sigma}]_{\mathbb{T}} \\ &= (-y)^{-\dim(O_\sigma)} \cdot (1+y)^{\dim(O_\sigma)} \cdot (k_\sigma)_*[\mathcal{O}_{V_\sigma}]_{\mathbb{T}}. \end{aligned}$$

Here, in the first two equalities we use a linear extension of equivariant Grothendieck duality over  $\mathbb{Z}[y^{\pm 1}]$ . Formula (5.13) is now a consequence of (5.14) and (5.15).  $\square$

## 6. APPLICATIONS TO WEIGHTED EHRHART THEORY

Let now  $M \cong \mathbb{Z}^n$  be a lattice and  $P \subset M_\mathbb{R} \cong \mathbb{R}^n$  be a full-dimensional lattice polytope with associated projective toric variety  $X = X_P$ , inner normal fan  $\Sigma = \Sigma_P$  and ample Cartier divisor  $D = D_P$ . One has (e.g., see [11, Prop.4.3.3])

$$(6.1) \quad \Gamma(X_P; \mathcal{O}_{X_P}(D_P)) = \bigoplus_{m \in P \cap M} \mathbb{C} \cdot \chi^m \subset \mathbb{C}[M],$$

where  $\chi^m$  denotes the character defined by  $m \in M$ , and  $\mathbb{C}[M]$  is the coordinate ring of the torus  $\mathbb{T}$ . Also, since  $D_P$  is ample,

$$(6.2) \quad H^i(X_P; \mathcal{O}_{X_P}(D_P)) = 0, \text{ for all } i > 0.$$

Moreover, one has (e.g., see [11, Prop.9.2.7])

$$(6.3) \quad H^i(X_P; \mathcal{O}_{X_P}(-D_P)) = 0, \text{ for all } i \neq n,$$

and

$$(6.4) \quad H^n(X_P; \mathcal{O}_{X_P}(-D_P)) = \bigoplus_{m \in \mathrm{Int}(P) \cap M} \mathbb{C} \cdot \chi^{-m},$$

with  $\text{Int}(P)$  the interior of  $P$ . Also, by equivariant toric Serre duality [11, Thm.9.2.10] (fitting with our description of the equivariant dualizing complex in (5.8)),

$$(6.5) \quad \bigoplus_{m \in \text{Int}(P) \cap M} \mathbb{C} \cdot \chi^m = (H^n(X_P; \mathcal{O}_{X_P}(-D_P)))^\vee \simeq H^0(X_P; \mathcal{O}_{X_P}(D_P) \otimes \omega_{X_P}),$$

with  $\omega_{X_P} = \widehat{\Omega}_{X_P}^{\dim(P)}$  the equivariant dualizing and canonical sheaf of  $X_P$ , together with  $H^i(X_P; \mathcal{O}_{X_P}(D_P) \otimes \omega_{X_P}) = 0$  for all  $i \neq 0$ .

*Remark 6.1.* For any  $\mathbb{T}$ -invariant Cartier divisor  $D$  on an  $n$ -dimensional complete toric variety  $X$  with torus  $\mathbb{T}$ , one has the nondegenerate equivariant pairing

$$H^i(X; \mathcal{O}_X(-D)) \times H^{n-i}(X; \mathcal{O}_X(D) \otimes \omega_X) \longrightarrow H^n(X; \omega_X) \longrightarrow \mathbb{C}$$

induced by the  $\mathbb{T}$ -equivariant sheaf isomorphism

$$\mathcal{O}_X(-D) \otimes \mathcal{O}_X(D) \otimes \omega_X \longrightarrow \omega_X$$

and the  $\mathbb{T}$ -equivariant trace map  $\omega_X[n] \rightarrow \omega_{pt} = \mathbb{C}$ . This induces the equivariant Serre duality

$$(H^i(X; \mathcal{O}_X(-D)))^\vee \simeq H^{n-i}(X; \mathcal{O}_X(D) \otimes \omega_X).$$

□

*Remark 6.2.* For a cone  $\sigma_Q \in \Sigma_P$  associated to a face  $Q$  of  $P$ , the corresponding orbit closure  $V_{\sigma_Q}$  can be identified with the toric variety  $X_Q$ , with corresponding lattice polytope defined as follows (cf. [11, Prop.3.2.9]): translate  $P$  by a vertex  $m_0$  of  $Q$  so that the origin is a vertex of  $Q_0 := Q - m_0$ ; while this translation by  $m_0$  does not change  $\Sigma_P$  or  $X_P$ ,  $Q_0$  is now a full-dimensional polytope in  $\text{Span}(Q_0)$  relative to the lattice  $\text{Span}(Q_0) \cap M$ , and  $X_Q$  is the associated toric variety. However, if one wishes to apply the above formulas to the toric variety  $X_Q$  associated to a face  $Q$  of  $P$ , one needs to work with the corresponding divisor

$$(6.6) \quad D_{Q_0} = D_{P-m_0}|_{X_Q} = D_P|_{X_Q} + \text{div}(\chi^{m_0}).$$

□

Formulae (6.1)-(6.4) correspond to the weight decomposition of the respective cohomology groups, e.g.,

$$(6.7) \quad H^i(X_P; \mathcal{O}_{X_P}(D_P)) = \bigoplus_{m \in M} H^i(X_P; \mathcal{O}_{X_P}(D_P))_m,$$

as in [11, Sect.9.1]. To relate this weight decomposition with the eigenspace decomposition of (4.1), we recall that the torus  $\mathbb{T}$  acts on  $\mathbb{C}[M]$  as follows: if  $t \in \mathbb{T}$  and  $f \in \mathbb{C}[M]$ , then  $t \cdot f \in \mathbb{C}[M]$  is given by  $p \mapsto f(t^{-1} \cdot p)$ , for  $p \in \mathbb{T}$  (see [11, pag.18]). In particular,  $t \cdot \chi^m = \chi^m(t^{-1})\chi^m$ , so that

$$H^i(X; \mathcal{O}_X(D))_m = H^i(X; \mathcal{O}_X(D))_{\chi^{-m}}.$$

Let us note that if for a positive integer  $\ell \in \mathbb{Z}_{>0}$  we consider the dilated polytope

$$\ell P := \{\ell \cdot u \mid u \in P\},$$

then the toric varieties corresponding to  $P$  and, resp.,  $\ell P$ , are the same, but  $D_{\ell P} = \ell D_P$ . Together with (6.1)-(6.5), this gives for such  $\ell \in \mathbb{Z}_{>0}$ :

$$(6.8) \quad \chi^{\mathbb{T}}(X_P, \mathcal{O}_{X_P}(\ell D_P)) = \sum_{m \in \ell P \cap M} \chi^{-m},$$

$$(6.9) \quad \chi^{\mathbb{T}}(X_P, \mathcal{O}_{X_P}(-\ell D_P)) = (-1)^n \cdot \sum_{m \in \text{Int}(\ell P) \cap M} \chi^m$$

and

$$(6.10) \quad \chi^{\mathbb{T}}(X_P, \mathcal{O}_{X_P}(\ell D_P) \otimes \omega_{X_P}) = \sum_{m \in \text{Int}(\ell P) \cap M} \chi^{-m}.$$

More generally, we have by [9, Rem.4.9], applied to the polytope  $\ell P$ , the following identity:

$$(6.11) \quad \chi^{\mathbb{T}}(X_P, mC_y^{\mathbb{T}}(X_P) \otimes \mathcal{O}_{X_P}(\ell D_P)) = \sum_{Q \preceq P} (1+y)^{\dim(Q)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \chi^{-m} \in \mathbb{Z}[M] \otimes_{\mathbb{Z}} \mathbb{Z}[y].$$

where  $\text{Relint}$  denotes the relative interior, and the first summation is over the faces  $Q$  of the lattice polytope  $P$ .

For further reference, let us also include here the following.

**Lemma 6.3.** *In the above notations, with  $\ell \in \mathbb{Z}_{>0}$ ,*

$$(6.12) \quad \chi^{\mathbb{T}}(X_P, \mathcal{O}_{X_P}(-\ell D_P) \otimes \omega_{X_P}) = (-1)^n \cdot \sum_{m \in \ell P \cap M} \chi^m$$

*Proof.* By toric equivariant Serre duality [11, Thm.9.2.10] and Remark 6.1, together with (6.1) and (6.2),

$$(6.13) \quad \bigoplus_{m \in P \cap M} \mathbb{C} \cdot \chi^{-m} = (H^0(X_P; \mathcal{O}_{X_P}(D_P)))^{\vee} \simeq H^n(X_P; \mathcal{O}_{X_P}(-D_P) \otimes \omega_{X_P}),$$

and  $H^i(X_P; \mathcal{O}_{X_P}(-D_P) \otimes \omega_{X_P}) = 0$  for all  $i \neq n$ . The assertion follows by reverting to the weight decomposition of these cohomology groups, as indicated above.  $\square$

Making use of Theorem 5.2, we can now generalize formula (6.11) as follows.

**Theorem 6.4.** *Let  $X = X_P$  be the projective toric variety with ample Cartier divisor  $D = D_P$  associated to a full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ , and let  $\ell \in \mathbb{Z}_{>0}$ . Let  $Z$  be an ambient  $\mathbb{T}$ -manifold containing  $X$  as a  $\mathbb{T}$ -invariant subvariety. Then for any Grothendieck class  $[\mathcal{M}] \in K_0(\text{MHM}_{\mathbb{T}}^{\mathbb{T}}(Z))$  of a mixed Hodge module (complex) on  $Z$  with support in  $X$ , we have*

$$(6.14) \quad \chi_y^{\mathbb{T}}(X, \ell D; [\mathcal{M}]) = \sum_{Q \preceq P} \chi_y(i_{x_Q}^*[\mathcal{M}]) \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \chi^{-m},$$

where the first summation on the right is over the faces  $Q$  of  $P$ , and  $x_Q \in O_{\sigma_Q}$  is a point in the orbit of  $X$  associated to (the cone  $\sigma_Q$  in the inner normal fan of  $P$  corresponding to) the face  $Q$ , with inclusion map  $i_{x_Q} : \{i_{x_Q}\} \hookrightarrow Z$ .

*Proof.* For a face  $Q$  of  $P$ , denote by  $k_Q : V_{\sigma_Q} := X_Q \hookrightarrow X \hookrightarrow Z$  the inclusion in the ambient  $\mathbb{T}$ -manifold  $Z$  of the orbit closure associated to the (cone  $\sigma_Q$  of the) face  $Q$ . Note that we have  $\dim_{\mathbb{R}}(Q) = \dim_{\mathbb{R}}(O_{\sigma_Q})$ . Let  $\mathbb{T}_Q$  be the quotient torus of  $\mathbb{T}$  corresponding to  $X_Q$ , as in (5.2).

In what follows, we will denote all constant maps to a point by  $a$ , regardless of their domains (which should be understood from the context). Then by (4.4) and (5.4) the following equality holds:

$$\begin{aligned} \chi_y^{\mathbb{T}}(X, \ell D; \mathcal{M}) &= a_* \left( \mathrm{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(\ell D) \right) \\ &= \sum_{Q \preceq P} \chi_y(i_{X_Q}^*[\mathcal{M}]) \cdot (1+y)^{\dim(Q)} \cdot a_* \left( (k_Q)_* [\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D) \right). \end{aligned}$$

So it remains to prove that, for any face  $Q$  of  $P$ , we have that:

$$(6.15) \quad \chi^{\mathbb{T}}(X, (k_Q)_* [\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D)) = a_* \left( (k_Q)_* [\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D) \right) = \sum_{m \in \mathrm{Relint}(\ell Q) \cap M} \chi^{-m}.$$

This follows from the projection formula and (6.10), by making use of formula (6.6). Indeed,

$$\begin{aligned} a_* \left( (k_Q)_* [\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D) \right) &= a_* (k_Q)_* \left( [\omega_{X_Q}]_{\mathbb{T}} \otimes k_Q^* \mathcal{O}_X(\ell D) \right) \\ &\stackrel{(6.6)}{=} a_* \left( [\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_{X_Q}(\ell D_{Q_0} - \mathrm{div}(\chi^{\ell m_0})) \right) \\ &= \chi^{-\ell m_0} \cdot a_* \left( [\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_{X_Q}(\ell D_{Q_0}) \right) \\ &\stackrel{(*)}{=} \chi^{-\ell m_0} \cdot a_* \left( [\omega_{X_Q}]_{\mathbb{T}_Q} \otimes \mathcal{O}_{X_Q}(\ell D_{Q_0}) \right) \\ &\stackrel{(6.10)}{=} \chi^{-\ell m_0} \cdot \left( \sum_{m \in \mathrm{Relint}(\ell(Q-m_0)) \cap M} \chi^{-m} \right) \\ &= \sum_{m \in \mathrm{Relint}(\ell Q) \cap M} \chi^{-m}, \end{aligned}$$

where  $(*)$  uses [9, Prop.3.1]. Here  $m_0$  is a vertex of  $Q$  as in (6.6), so that  $Q_0 := Q - m_0$  is a full-dimensional lattice polytope in  $\mathrm{Span}(Q_0)$  relative to the lattice  $\mathrm{Span}(Q_0) \cap M$ , with  $X_Q$  the associated toric variety.  $\square$

*Remark 6.5.* For future use, we include here the following formula, which one gets as in the proof of (6.15), but using  $\mathcal{O}_{X_Q}$  instead of  $\omega_{X_Q}$ , together with (6.8):

$$(6.16) \quad \chi^{\mathbb{T}}(X, (k_Q)_* [\mathcal{O}_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D)) = a_* \left( (k_Q)_* [\mathcal{O}_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D) \right) = \sum_{m \in \ell Q \cap M} \chi^{-m}.$$

$\square$

**Definition 6.6** (Generalized weighted Ehrhart polynomials). Let  $\varphi : M_{\mathbb{R}} \cong \mathbb{R}^n \rightarrow \mathbb{C}$  be a homogeneous polynomial function, and  $\ell \in \mathbb{Z}_{>0}$ . With the above notations, the expression

$$(6.17) \quad E_{P, \mathcal{M}}^{\varphi}(\ell, y) := \sum_{Q \preceq P} \chi_y(i_{X_Q}^*[\mathcal{M}]) \cdot (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \mathrm{Relint}(\ell Q) \cap M} \varphi(m)$$

is obtained from  $\chi_y^{\mathbb{T}}(X, \ell D; \mathcal{M})$  by substituting  $\chi^{-m}$  by  $\varphi((1+y)m) = (1+y)^{\deg(\varphi)} \cdot \varphi(m)$ , for each  $m$ . As we will show later on in the Appendix (but see also [8, Prop.4.1]),  $E_{P, \mathcal{M}}^{\varphi}(\ell, y)$  is obtained by evaluating a polynomial  $E_{P, \mathcal{M}}^{\varphi}(z, y)$  at  $z = \ell$ . We call this polynomial the *generalized weighted Ehrhart polynomial* of  $P$  and  $\varphi$  with coefficients induced from  $[\mathcal{M}] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z))$ . Similarly, one can define

$$(6.18) \quad \tilde{E}_{P, \mathcal{M}}^{\varphi}(\ell, y) := \sum_{Q \preceq P} \chi_y(i_{x_Q}^*[\mathcal{M}]) \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m)$$

which obtained from  $\chi_y^{\mathbb{T}}(X, \ell D; \mathcal{M})$  by substituting  $\chi^{-m}$  by  $\varphi(m)$ , for each  $m$ . Since  $\varphi$  is homogeneous, this substitution fits with the duality involution  $\chi^m \mapsto \chi^{-m}$  for  $m \in M$ , via  $\varphi(-m) = (-1)^{\deg(\varphi)} \cdot \varphi(m)$ .

More generally, for a Laurent polynomial *weight function*

$$f : \{\text{faces of } P\} \rightarrow \mathbb{Z}[y^{\pm 1}], \quad Q \mapsto f_Q(y)$$

defined on the set of faces of  $P$ , we define as in the Introduction an associated generalized weighted Ehrhart polynomial by

$$(6.19) \quad \begin{aligned} E_{P, f}^{\varphi}(\ell, y) &:= \sum_{Q \preceq P} f_Q(y) \cdot (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m) \\ &=: (1+y)^{\deg(\varphi)} \cdot \tilde{E}_{P, f}^{\varphi}(\ell, y). \end{aligned}$$

*Remark 6.7.* Note that any Grothendieck class  $[\mathcal{M}] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z))$  of a equivariant mixed Hodge module (complex) on  $Z$  with support in  $X = X_P$ , induces a weight function on the faces of  $P$  defined by  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ , with  $x_Q \in O_{\sigma_Q}$  a point in the orbit of  $X_P$  associated to (the cone  $\sigma_Q$  in the inner normal fan of  $P$  corresponding to) the face  $Q$  and  $i_{x_Q} : \{x_Q\} \rightarrow Z$  the inclusion map. Moreover, any weight function  $f$  can be obtained in this way from the class of such a mixed Hodge module complex  $\mathcal{M}$ , e.g., by using direct sums and shifts of classes of  $(j_Q)! \mathbb{Q}_{O_{\sigma_Q}}^H(n_Q)$ , with  $n_Q \in \mathbb{Z}$  and  $j_Q : O_{\sigma_Q} \hookrightarrow Z$  the inclusion, regarded as  $\mathbb{T}$ -equivariant objects on  $Z$  (or, using the subgroup  $\chi_{Hdg}^{\mathbb{T}}(\tau(F^{\mathbb{T}}(X)[y^{\pm 1}]))$  as in the Introduction). In particular, the expressions  $E_{P, f}^{\varphi}(\ell, y)$  and  $\tilde{E}_{P, f}^{\varphi}(\ell, y)$  are then also obtained by evaluating a *polynomial*  $E_{P, f}^{\varphi}(z, y)$ , resp.,  $\tilde{E}_{P, f}^{\varphi}(z, y)$  at  $z = \ell$ .  $\square$

Before studying the generalized weighted Ehrhart polynomial  $E_{P, f}^{\varphi}(\ell, y)$ , resp.,  $\tilde{E}_{P, f}^{\varphi}(\ell, y)$ , in more detail, let us also mention the following result, similar to Theorem 6.4.

**Theorem 6.8.** *In the notations of Theorem 6.4, we have for  $\ell \in \mathbb{Z}_{>0}$ :*

$$(6.20) \quad \chi_y^{\mathbb{T}}(X, -\ell D; [\mathcal{M}]) = \sum_{Q \preceq P} \chi_y(i_{x_Q}^*[\mathcal{M}]) \cdot (-1-y)^{\dim(Q)} \cdot \sum_{m \in \ell Q \cap M} \chi^m.$$

*Proof.* The proof is very similar to that of Theorem 6.4, so we only sketch it below. In view of (5.4), this amounts to showing that for any face  $Q$  of  $P$ , and using notations from loc.cit.,

we have the following formula:

$$(6.21) \quad \begin{aligned} \chi^{\mathbb{T}}(X, (k_Q)_*[\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(-\ell D)) &= a_*((k_Q)_*[\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(-\ell D)) \\ &= (-1)^{\dim(Q)} \cdot \sum_{m \in \ell Q \cap M} \chi^m. \end{aligned}$$

Indeed,

$$\begin{aligned} a_*((k_Q)_*[\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(-\ell D)) &= a_*(k_Q)_*([\omega_{X_Q}]_{\mathbb{T}} \otimes k_Q^* \mathcal{O}_X(-\ell D)) \\ &\stackrel{(6.6)}{=} a_*([\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_{X_Q}(-\ell D_{Q_0} + \text{div}(\chi^{\ell m_0}))) \\ &= \chi^{\ell m_0} \cdot a_*([\omega_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_{X_Q}(-\ell D_{Q_0})) \\ &\stackrel{(*)}{=} \chi^{\ell m_0} \cdot a_*([\omega_{X_Q}]_{\mathbb{T}_Q} \otimes \mathcal{O}_{X_Q}(-\ell D_{Q_0})) \\ &\stackrel{(6.12)}{=} \chi^{\ell m_0} \cdot \left( (-1)^{\dim(Q)} \cdot \sum_{m \in (\ell Q - m_0) \cap M} \chi^m \right) \\ &= (-1)^{\dim(Q)} \cdot \sum_{m \in \ell Q \cap M} \chi^m, \end{aligned}$$

where  $(*)$  uses again [9, Prop.3.1].  $\square$

We can now justify the reciprocity formula (1.4) without relying on Brion-Vergne's combinatorial work [8], as well as the constant term formula (1.2):

**Corollary 6.9.** *Let  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$  be a full-dimensional lattice polytope, with  $\varphi : M_{\mathbb{R}} \cong \mathbb{R}^n \rightarrow \mathbb{C}$  a homogeneous polynomial function. For any Laurent polynomial weight vector  $\{f_Q\}_{Q \preceq P}$  defined on the set of faces of  $P$ , we have*

$$(6.22) \quad E_{P,f}^{\varphi}(-\ell, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1-y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m)$$

and

$$(6.23) \quad \tilde{E}_{P,f}^{\varphi}(-\ell, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1-y)^{\dim(Q)} \cdot (-1)^{\deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m).$$

In particular,

$$(6.24) \quad E_{P,f}^{\varphi}(0, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1-y)^{\dim(Q) + \deg(\varphi)} \cdot \varphi(0)$$

and

$$(6.25) \quad \tilde{E}_{P,f}^{\varphi}(0, y) = \sum_{Q \preceq P} f_Q(y) \cdot (-1-y)^{\dim(Q)} \cdot (-1)^{\deg(\varphi)} \cdot \varphi(0).$$

*Proof.* As in Remark 6.7, choose a  $[\mathcal{M}] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z))$  with  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ , such that  $E_{P,f}^{\varphi}(\ell, y) = E_{P, \mathcal{M}}^{\varphi}(\ell, y)$ . From the definition of  $E_{P, \mathcal{M}}^{\varphi}(\ell, y)$  in (6.17),  $E_{P, \mathcal{M}}^{\varphi}(-\ell, y)$  is obtained from  $\chi_y^{\mathbb{T}}(X, -\ell D; \mathcal{M})$  by substituting each  $\chi^m$  by  $\varphi((1+y) \cdot (-m)) = (-1-y)^{\deg(\varphi)} \cdot \varphi(m)$ . Formula (6.22) follows then by applying this rule to the expression for

$\chi_y^{\mathbb{T}}(X, -\ell D; \mathcal{M})$  given by (6.20). The constant term formula (6.24) is then an easy consequence, by setting  $\ell = 0$  in (6.22), or by combining (4.5) and (5.5). Similarly, for  $\tilde{E}_{P,f}^{\varphi}(-\ell, y)$  and  $\tilde{E}_{P,f}^{\varphi}(0, y)$ .  $\square$

*Example 6.10.* Let us next give some examples for Theorem 6.4, formulated in terms of our generalized weighted Ehrhart polynomials  $E_{P,\mathcal{M}}^{\varphi}(\ell, y)$  via Definition 6.6. Let  $X = X_P$  be the projective toric variety with inner normal fan  $\Sigma = \Sigma_P$  and ample Cartier divisor  $D = D_P$  associated to a full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ , and let  $\ell \in \mathbb{Z}_{>0}$ . For a face  $Q$  of  $P$  denote by  $j_Q : O_{\sigma_Q} \hookrightarrow Z$  the inclusion of the orbit associated to the (cone of the) face  $Q$ .

(a) For  $\mathcal{M} = (j_Q)_! \mathbb{Q}_{O_{\sigma_Q}}^H$ , we get

$$(6.26) \quad E_{P,\mathcal{M}}^{\varphi}(\ell, y) = (1+y)^{\dim(Q)+\deg(\varphi)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m),$$

with  $E_{P,\mathcal{M}}^{\varphi}(0, y) = (-y-1)^{\dim(Q)+\deg(\varphi)} \cdot \varphi(0) = (-y-1)^{\deg(\varphi)} \cdot \chi_y(O_{\sigma_Q}) \cdot \varphi(0)$ .

In particular, by evaluating  $E_{P,\mathcal{M}}^{\varphi}(\ell, y)$  at  $y = 0$ , we get the polynomial

$$E_{P,\mathcal{M}}^{\varphi}(\ell, 0) = \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m)$$

summing up the values of  $\varphi$  at lattice points in the relative interior of the dilated face  $\ell Q$  (i.e., the Euler-Maclaurin sum for the relative interior of  $\ell Q$ ), with  $E_{P,\mathcal{M}}^{\varphi}(0, 0) = (-1)^{\dim(Q)+\deg(\varphi)} \cdot \varphi(0)$ .

(b) Let  $X' := X_{P'}$  be a torus-invariant closed algebraic subset of  $X = X_P$  corresponding to a polytopal subcomplex  $P' \subseteq P$  (i.e., a closed union of faces of  $P$ ), with  $[\mathcal{M}] = [\mathbb{Q}_{X'}^H]$  pushed forward to the ambient  $\mathbb{T}$ -manifold  $Z$  containing  $X_P$  as a  $\mathbb{T}$ -invariant closed subvariety. Then

$$(6.27) \quad E_{P,\mathcal{M}}^{\varphi}(\ell, y) = \sum_{Q \preceq P'} (1+y)^{\dim(Q)+\deg(\varphi)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m) \in \mathbb{Z}[y],$$

where the summation is over the faces  $Q$  of  $P'$ . Hence,

$$E_{P,\mathcal{M}}^{\varphi}(\ell, 0) = \sum_{m \in \ell P' \cap M} \varphi(m)$$

is the corresponding Euler-Maclaurin sum for  $\ell P'$ , and

$$E_{P,\mathcal{M}}^{\varphi}(0, y) = \sum_{Q \preceq P'} (-y-1)^{\dim(Q)+\deg(\varphi)} \cdot \varphi(0) = (-1)^{\deg(\varphi)} \cdot \chi_y(X') \cdot \varphi(0),$$

which gives that

$$E_{P,\mathcal{M}}^{\varphi}(0, 0) = (-1)^{\deg(\varphi)} \cdot \chi(P') \cdot \varphi(0),$$

with  $\chi(P')$  the topological Euler characteristic of  $P'$ .

(c) For  $\mathcal{M} = (j_Q)_* \mathbb{Q}_{O_{\sigma_Q}}^H$ , we get by (5.13) and (6.16) that

$$(6.28) \quad E_{P,\mathcal{M}}^{\varphi}(\ell, y) = (1+y)^{\dim(Q)+\deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m),$$

with  $E_{P,\mathcal{M}}^\varphi(0,y) = (1+y)^{\deg(\varphi)} \cdot \chi_y(V_{\sigma_Q}) \cdot \varphi(0)$ ,

$$E_{P,\mathcal{M}}^\varphi(\ell,0) = \sum_{m \in \ell Q \cap M} \varphi(m)$$

the Euler-Maclaurin sum for  $\ell Q$ , and  $E_{P,\mathcal{M}}^\varphi(0,0) = \chi(Q) \cdot \varphi(0)$ .  $\square$

Let  $X = X_P$  be as before the projective toric variety with inner normal fan  $\Sigma = \Sigma_P$  and ample Cartier divisor  $D = D_P$  associated to a full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ . Let  $X \hookrightarrow Z$  be the equivariant embedding of  $X$  into an ambient  $\mathbb{T}$ -manifold, as in [30]. Then another important example is provided by  $\mathcal{M} = IC_X^H$ . This is a shifted  $\mathbb{T}$ -equivariant pure Hodge module on  $Z$  with support in  $X$ . As explained in the Introduction, in the case when  $P$  contains the origin in its interior (so that its polar polytope  $P^\circ$  is defined), the stalk contributions  $\chi_y(IC_X^H|_{x_Q})$  have the following combinatorial description (see, e.g., [5, 16, 13, 27], etc):

$$(6.29) \quad \chi_y(IC_X^H|_{x_Q}) = \tilde{g}_Q(-y) := g_{Q^\circ}(-y),$$

with  $Q$  a face of  $P$  and  $Q^\circ$  the corresponding face of the polar polytope  $P^\circ$ , and  $g_{Q^\circ}(t) \in \mathbb{Z}[t]$  the Stanley  $g$ -polynomial of  $Q^\circ$ , cf. [29, Sect.3]. In the notations of Definition 6.6, we then get for  $E_{P,IC_X^H}^\varphi(\ell,y) := E_{P,IC_X^H}^\varphi(\ell,y)$  the following.

**Corollary 6.11.** *Let  $X = X_P$  is the projective toric variety with ample Cartier divisor  $D = D_P$  associated to a full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ , with  $\varphi : M_{\mathbb{R}} \cong \mathbb{R}^n \rightarrow \mathbb{C}$  a homogeneous polynomial function. Assume that  $P$  contains the origin in its interior. Then, for  $\ell \in \mathbb{Z}_{>0}$ ,*

$$(6.30) \quad E_{P,IC^H}^\varphi(\ell,y) = \sum_{Q \preceq P} \tilde{g}_Q(-y) \cdot (1+y)^{\dim(Q)+\deg(\varphi)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m),$$

and

$$(6.31) \quad E_{P,IC^H}^\varphi(-\ell,y) = \sum_{Q \preceq P} \tilde{g}_Q(-y) \cdot (-1-y)^{\dim(Q)+\deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m).$$

where the summation is over the faces  $Q$  of  $P$ . In particular,

$$(6.32) \quad E_{P,IC^H}^\varphi(0,y) = \sum_{Q \preceq P} \tilde{g}_Q(-y) \cdot (-1-y)^{\dim(Q)+\deg(\varphi)} \cdot \varphi(0).$$

Similar formulas hold for  $\tilde{E}_{P,IC^H}^\varphi(\ell,y)$  by deleting the factor  $(1+y)^{\deg(\varphi)}$ .

*Remark 6.12.* (a) Formula (6.30) equals  $G_\varphi(\ell,y)$  from [3, formula (14)].

(b) If the full-dimensional polytope  $P$  is simple, then  $X = X_P$  is a simplicial toric variety, hence  $[IC_X^H] = [\mathbb{Q}_X^H] \in K_0^{\mathbb{T}}(\text{MHM}_X(Z))$ . So, in this case,  $\chi_y(IC_X^H|_{x_Q}) = 1$  (hence  $\tilde{g}_Q(-y) = 1$ , if  $P$  contains the origin in its interior).  $\square$

We next discuss the effect of duality on generalized weighted Ehrhart polynomials, proving a generalized *reciprocity formula*. This takes the form of a *purity* statement for pure equivariant Hodge modules (like  $IC_X^H$ ).

**Theorem 6.13** (Reciprocity and Purity). *In the above notations, for any Grothendieck class  $[\mathcal{M}] \in \text{Image}(\chi_{\text{Hdg}}^{\mathbb{T}}) \subset K_0(\text{MHM}_X^{\mathbb{T}}(Z))$ , we have*

$$(6.33) \quad E_{P, \mathcal{M}}^{\varphi}(-\ell, y) = (-y)^{\deg(\varphi)} \cdot E_{P, \mathbb{D}_Z^{\mathbb{T}}, \mathcal{M}}^{\varphi}\left(\ell, \frac{1}{y}\right),$$

and

$$(6.34) \quad \tilde{E}_{P, \mathcal{M}}^{\varphi}(-\ell, y) = (-1)^{\deg(\varphi)} \cdot \tilde{E}_{P, \mathbb{D}_Z^{\mathbb{T}}, \mathcal{M}}^{\varphi}\left(\ell, \frac{1}{y}\right).$$

In particular, if  $\mathcal{M}$  is such a self-dual pure equivariant Hodge module of weight  $n$  on  $X = X_P$ , then the following purity property holds:

$$(6.35) \quad E_{P, \mathcal{M}}^{\varphi}(-\ell, y) = (-y)^{n+\deg(\varphi)} \cdot E_{P, \mathcal{M}}^{\varphi}\left(\ell, \frac{1}{y}\right),$$

and

$$(6.36) \quad \tilde{E}_{P, \mathcal{M}}^{\varphi}(-\ell, y) = (-y)^n \cdot (-1)^{\deg(\varphi)} \cdot \tilde{E}_{P, \mathcal{M}}^{\varphi}\left(\ell, \frac{1}{y}\right).$$

Before proving Theorem 6.13, we need the following.

**Lemma 6.14.** *Under the assumptions of Theorem 6.13 and for  $\ell \in \mathbb{Z}_{>0}$ , we have:*

$$(6.37) \quad \chi_y^{\mathbb{T}}(X, \ell D; \mathbb{D}_Z^{\mathbb{T}}[\mathcal{M}]) = \sum_{Q \preceq P} f_Q\left(\frac{1}{y}\right) \cdot (-y)^{-\dim(Q)} \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \ell Q \cap M} \chi^{-m},$$

with  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ .

*Proof.* First, by (5.4) and (5.7), we have

$$(6.38) \quad \begin{aligned} \text{DR}_y^{\mathbb{T}}([\mathcal{M}]) &= \sum_{Q \preceq P} \chi_y(i_{x_Q}^*[\mathcal{M}]) \cdot (1+y)^{\dim(Q)} \cdot (k_Q)_*[\omega_{V_{\sigma_Q}}] \\ &= \sum_{Q \preceq P} f_Q(y) \cdot \text{DR}_y^{\mathbb{T}}\left([\!(j_Q)! \mathbb{Q}_{O_{\sigma_Q}}^H]\right), \end{aligned}$$

with  $j_Q : O_{\sigma_Q} \hookrightarrow Z$  and  $k_Q : V_{\sigma_Q} = X_Q \hookrightarrow Z$  the inclusion maps. Moreover, as in (5.14), we have:

$$(6.39) \quad \text{DR}_y^{\mathbb{T}}\left(\mathbb{D}_Z^{\mathbb{T}}\left([\!(j_Q)! \mathbb{Q}_{O_{\sigma_Q}}^H]\right)\right) = (-y)^{-\dim(Q)} \cdot \text{DR}_y^{\mathbb{T}}\left([\!(j_Q)_* \mathbb{Q}_{O_{\sigma_Q}}^H]\right).$$

Since  $\text{DR}_y^{\mathbb{T}}$  commutes with dualities, up to exchanging  $y$  by  $1/y$ , we then get from (6.38) and (6.39) that

$$(6.40) \quad \begin{aligned} \text{DR}_y^{\mathbb{T}}(\mathbb{D}_Z^{\mathbb{T}}[\mathcal{M}]) &= \sum_{Q \preceq P} f_Q\left(\frac{1}{y}\right) \cdot (-y)^{-\dim(Q)} \cdot \text{DR}_y^{\mathbb{T}}\left([\!(j_Q)_* \mathbb{Q}_{O_{\sigma_Q}}^H]\right) \\ &\stackrel{(5.13)}{=} \sum_{Q \preceq P} f_Q\left(\frac{1}{y}\right) \cdot (-y)^{-\dim(Q)} \cdot (1+y)^{\dim(Q)} \cdot (k_Q)_*[\mathcal{O}_{X_Q}]_{\mathbb{T}}. \end{aligned}$$

Therefore, with  $a$  denoting as before the constant map to a point,

$$\begin{aligned}
(6.41) \quad \chi_y^\mathbb{T}(X, \ell D; \mathbb{D}_Z^\mathbb{T}[\mathcal{M}]) &= a_* \left( \mathrm{DR}_y^\mathbb{T}(\mathbb{D}_Z^\mathbb{T}[\mathcal{M}]) \otimes \mathcal{O}_X(\ell D) \right) \\
&= \sum_{Q \preceq P} f_Q \left( \frac{1}{y} \right) \cdot (-y)^{-\dim(Q)} \cdot (1+y)^{\dim(Q)} \cdot a_* \left( (k_Q)_* [\mathcal{O}_{X_Q}]_{\mathbb{T}} \otimes \mathcal{O}_X(\ell D) \right) \\
&\stackrel{(6.16)}{=} \sum_{Q \preceq P} f_Q \left( \frac{1}{y} \right) \cdot (-y)^{-\dim(Q)} \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \ell Q \cap M} \chi^{-m}.
\end{aligned}$$

□

*Proof of Theorem 6.13.* First, we get from (6.37) that

$$E_{P, \mathbb{D}_Z^\mathbb{T} \mathcal{M}}^\varphi(\ell, y) = \sum_{Q \preceq P} f_Q \left( \frac{1}{y} \right) \cdot (-y)^{-\dim(Q)} \cdot (1+y)^{\dim(Q)} \cdot (1+y)^{\deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m),$$

with  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$  as before. Substituting  $1/y$  in place of  $y$ , we then get

$$\begin{aligned}
(6.42) \quad E_{P, \mathbb{D}_Z^\mathbb{T} \mathcal{M}}^\varphi\left(\ell, \frac{1}{y}\right) &= \sum_{Q \preceq P} f_Q(y) \cdot (-y)^{\dim(Q)} \cdot \left(1 + \frac{1}{y}\right)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m) \\
&= \sum_{Q \preceq P} f_Q(y) \cdot y^{-\deg(\varphi)} \cdot (-1)^{\dim(Q)} \cdot (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m) \\
&= \sum_{Q \preceq P} f_Q(y) \cdot y^{-\deg(\varphi)} \cdot (-1)^{\deg(\varphi)} \cdot (-1-y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m) \\
&= (-y)^{-\deg(\varphi)} \cdot \sum_{Q \preceq P} f_Q(y) \cdot (-1-y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \ell Q \cap M} \varphi(m) \\
&\stackrel{(6.22)}{=} (-y)^{-\deg(\varphi)} \cdot E_{P, f}^\varphi(-\ell, y).
\end{aligned}$$

Formula (6.35) follows immediately by combining (6.33) with (2.3).

The formulas for the polynomials  $\tilde{E}$  are obtained similarly. □

*Example 6.15.* The reciprocity formula (6.33) can be applied to all cases of Example 6.10. In the (dual) cases (a) and (c), for  $\mathcal{M} = (j_Q)! \mathbb{Q}_{\mathcal{O}_{\sigma_Q}}^H$  with  $\mathbb{D}_Z^\mathbb{T}[\mathcal{M}] = [(j_Q)_* \mathbb{Q}_{\mathcal{O}_{\sigma_Q}}^H(\dim(Q))] \in K_0^\mathbb{T}(\mathrm{MHM}_X(Z))$ , formula (6.33) specializes to the *classical Ehrhart reciprocity* for  $Q$  from [8, Prop.4.1]. Indeed,

$$E_{P, \mathcal{M}}^\varphi(\ell, y) = (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \mathrm{Relint}(\ell Q) \cap M} \varphi(m)$$

and

$$E_{P, \mathbb{D}_X^\mathbb{T} \mathcal{M}}^\varphi\left(\ell, \frac{1}{y}\right) = (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \left( (-1)^{\dim(Q) + \deg(\varphi)} \sum_{m \in \ell Q \cap M} \varphi(m) \right).$$

□

*Example 6.16 (IC-reciprocity).* Assume  $P$  contains the origin in its interior. Then for  $\mathcal{M} = IC_X^{IH}$ , corresponding to the weight function  $f_Q(y) = \tilde{g}_Q(-y)$ , our general purity formula (6.35) recovers via Corollary 6.11 the reciprocity formula proved [3, Theorems 1.3 and 2.6] by combinatorial methods.  $\square$

## 7. APPENDIX

In this section we give a geometric proof of the following result (without relying on the combinatorial work of Brion-Vergne [8]).

**Theorem 7.1.** *Let  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$  be a full-dimensional lattice polytope. Let  $\varphi : M_{\mathbb{R}} \cong \mathbb{R}^n \rightarrow \mathbb{C}$  be a homogeneous polynomial function, and  $\ell \in \mathbb{Z}_{>0}$ . Then for any Laurent polynomial weight function*

$$f : \{\text{faces of } P\} \rightarrow \mathbb{Z}[y^{\pm 1}], \quad Q \mapsto f_Q(y)$$

defined on the set of faces of  $P$ , the expression

$$(7.1) \quad E_{P,f}^{\varphi}(\ell, y) := \sum_{Q \preceq P} f_Q(y) \cdot (1+y)^{\dim(Q) + \deg(\varphi)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} \varphi(m) = (1+y)^{\deg(\varphi)} \cdot \tilde{E}_{P,f}^{\varphi}(\ell, y)$$

is obtained by evaluating a polynomial  $E_{P,f}^{\varphi}(z, y)$  at  $z = \ell$ .

Let  $X = X_P$  be the toric variety with torus  $\mathbb{T}$  and ample Cartier divisor  $D = D_P$  associated to the lattice polytope  $P$ . Let  $Z$  be an ambient smooth  $\mathbb{T}$ -variety. In view of Remark 6.7, it suffices to prove the assertion in the above theorem for  $E_{P,\mathcal{M}}^{\varphi}(\ell, y)$ , as defined in (6.17), i.e., assuming that  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ , for a certain class  $[\mathcal{M}] \in K_0(\text{MHM}_X^{\mathbb{T}}(Z))$  of a mixed Hodge module (complex) on  $Z$  with support in  $X = X_P$ , with  $x_Q \in O_{\sigma_Q}$  a point in the orbit of  $X$  associated to (the cone  $\sigma_Q$  in the inner normal fan of  $P$  corresponding to) the face  $Q$ , and  $i_{x_Q} : \{x_Q\} \rightarrow Z$  the inclusion map. Our proof in this case is based on localization techniques developed in [9]. We first introduce some terminology.

Recall that the (Borel-type) rational equivariant cohomology of  $X$  is defined as

$$(7.2) \quad H_{\mathbb{T}}^*(X; \mathbb{Q}) := H^*(E\mathbb{T} \times_{\mathbb{T}} X; \mathbb{Q}),$$

where  $\mathbb{T} \hookrightarrow E\mathbb{T} \rightarrow B\mathbb{T}$  is the universal principal  $\mathbb{T}$ -bundle, i.e.,  $B\mathbb{T} = (B\mathbb{C}^*)^n = (\mathbb{C}P^{\infty})^n$  and  $E\mathbb{T}$  is contractible. In particular, if  $X = pt$  is a point space,

$$(7.3) \quad H_{\mathbb{T}}^*(pt; \mathbb{Q}) = H^*(B\mathbb{T}; \mathbb{Q}) \simeq \mathbb{Q}[t_1, \dots, t_n] =: (\Lambda_{\mathbb{T}})_{\mathbb{Q}}.$$

Similarly, one can define equivariant homology groups  $H_*^{\mathbb{T}}(X; \mathbb{Q})$  as in [7, 15], using suitable finite-dimensional approximations of  $B\mathbb{T}$ . Note that these equivariant homology groups can be non-zero also in negative degrees. As in the non-equivariant context, there is a cap product operation

$$\cap : H_{\mathbb{T}}^*(X; \mathbb{Q}) \times H_*^{\mathbb{T}}(X; \mathbb{Q}) \longrightarrow H_*^{\mathbb{T}}(X; \mathbb{Q}),$$

together with an equivariant Poincaré duality if  $X$  is simplicial. In particular,  $H_{\mathbb{T}}^*(X; \mathbb{Q})$ , resp.,  $H_*^{\mathbb{T}}(X; \mathbb{Q})$ , is a  $(\Lambda_{\mathbb{T}})_{\mathbb{Q}}$ -algebra, resp.,  $(\Lambda_{\mathbb{T}})_{\mathbb{Q}}$ -module.

For  $m \in M$ , an element in the character lattice of the torus  $\mathbb{T}$ , one can view the corresponding character  $\chi^m : \mathbb{T} \rightarrow \mathbb{C}^*$  as a  $\mathbb{T}$ -equivariant line bundle  $\mathbb{C}\chi^m$  over a point space  $pt$ , where the  $\mathbb{T}$ -action on  $\mathbb{C}$  is induced via  $\chi^m$ . This gives rise to an abelian group isomorphism

$$(M, +) \simeq (Pic_{\mathbb{T}}(pt), \otimes).$$

Recall that  $m \mapsto -m$  corresponds to the duality involution  $(-)^{\vee}$ . Taking the first equivariant Chern class  $c_{\mathbb{T}}^1$  (or the dual  $-c_{\mathbb{T}}^1 = c_{\mathbb{T}}^1 \circ (-)^{\vee}$ ) of  $\mathbb{C}\chi^m$  gives an isomorphism

$$(7.4) \quad c = c_{\mathbb{T}}^1, \text{ resp., } s = -c_{\mathbb{T}}^1 : M \simeq H_{\mathbb{T}}^2(pt; \mathbb{Z})$$

and

$$(7.5) \quad c, \text{ resp., } s : Sym_{\mathbb{Q}}(M) \simeq H_{\mathbb{T}}^*(pt; \mathbb{Q}) = (\Lambda_{\mathbb{T}})_{\mathbb{Q}},$$

with  $Sym_{\mathbb{Q}}(M) = \bigoplus_{k=0}^{\infty} Sym_{\mathbb{Q}}^k(M)$  the (rational) symmetric algebra of  $M$ . So if  $m_i$  ( $i = 1, \dots, n$ ) is a basis of  $M \simeq \mathbb{Z}^n$ , then  $H_{\mathbb{T}}^*(pt; \mathbb{Q}) = (\Lambda_{\mathbb{T}})_{\mathbb{Q}} \simeq \mathbb{Q}[t_1, \dots, t_n]$ , with  $t_i = \pm c_{\mathbb{T}}^1(\mathbb{C}\chi^{m_i})$  for  $i = 1, \dots, n$ .

Let

$$\mathbb{Q}\{t_1, \dots, t_n\} \simeq (H_{\mathbb{T}}^*(pt; \mathbb{Q}))^{an} =: (\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}} \subset (\widehat{\Lambda}_{\mathbb{T}})_{\mathbb{Q}} \simeq \mathbb{Q}[[t_1, \dots, t_n]]$$

be the subring of convergent power series (around zero) with rational coefficients, i.e., after pairing with  $z \in N_{\mathbb{K}} = N \otimes_{\mathbb{Z}} \mathbb{K}$  (for  $\mathbb{K} = \mathbb{R}, \mathbb{C}$ ) one gets a convergent power series *function* in  $z$  around zero, whose corresponding Taylor polynomials have rational coefficients. Here, the degree completion  $(\widehat{\Lambda}_{\mathbb{T}})_{\mathbb{Q}}$  of  $(\Lambda_{\mathbb{T}})_{\mathbb{Q}}$  corresponds to formal power series with rational coefficients.

Recall that if  $\mathcal{F}$  is a  $\mathbb{T}$ -equivariant coherent sheaf on  $X$ , then the  $K$ -theoretic Euler characteristic of  $\mathcal{F}$  is defined as in (4.3) by:

$$(7.6) \quad \chi^{\mathbb{T}}(X, \mathcal{F}) = \sum_{m \in M} \sum_{i=0}^{\dim(X)} (-1)^i \dim_{\mathbb{C}} H^i(X; \mathcal{F})_{\chi^m} \cdot \chi^m \in \mathbb{Z}[M],$$

By applying the equivariant Chern character ring homomorphism

$$\text{ch}^{\mathbb{T}} : \mathbb{Z}[M] \simeq K_{\mathbb{T}}^0(pt) \hookrightarrow (\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}} \subset (\widehat{\Lambda}_{\mathbb{T}})_{\mathbb{Q}}$$

given by  $m \mapsto e^{c(m)}$ , we further get the *cohomological Euler characteristic* of  $\mathcal{F}$ :

$$(7.7) \quad \chi^{\mathbb{T}}(X, \mathcal{F}) = \sum_{m \in M} \sum_{i=0}^{\dim(X)} (-1)^i \dim_{\mathbb{C}} H^i(X; \mathcal{F})_{\chi^m} \cdot e^{c(m)} \in (\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}} \subset (\widehat{\Lambda}_{\mathbb{T}})_{\mathbb{Q}}.$$

Consider now the fixed-point set  $X^{\mathbb{T}}$  of the torus action; this can be identified with the set of vertices of the lattice polytope  $P$ . By [7, Lem.8.4, Lem.8.5], the inclusion  $\alpha : X^{\mathbb{T}} \hookrightarrow X$  induces an injective homomorphism of  $H_{\mathbb{T}}^*(pt; \mathbb{Q})$ -modules

$$\alpha_* : \widehat{H}_*^{\mathbb{T}}(X^{\mathbb{T}}; \mathbb{Q}) \hookrightarrow \widehat{H}_*^{\mathbb{T}}(X; \mathbb{Q})$$

which becomes an isomorphism

$$\alpha_* : \widehat{H}_*^{\mathbb{T}}(X^{\mathbb{T}}; \mathbb{Q})_L \simeq \widehat{H}_*^{\mathbb{T}}(X; \mathbb{Q})_L$$

upon localization at the multiplicative subset  $L \subset (\Lambda_{\mathbb{T}})_{\mathbb{Q}} = H_{\mathbb{T}}^*(pt; \mathbb{Q})$  generated by the elements  $\pm c(m)$ , for  $0 \neq m \in M$  (cf. [7, Cor.8.9]). Here, we use the completion

$$\widehat{H}_*^{\mathbb{T}}(X; \mathbb{Q}) := \prod_{i \leq \dim(X)} H_i^{\mathbb{T}}(X; \mathbb{Q}).$$

Since

$$\widehat{H}_*^{\mathbb{T}}(X^{\mathbb{T}}; \mathbb{Q}) = \bigoplus_{x \in X^{\mathbb{T}}} \widehat{H}_*^{\mathbb{T}}(x; \mathbb{Q}),$$

one gets via the isomorphism  $\alpha_*$  a projection map of  $H_{\mathbb{T}}^*(pt; \mathbb{Q})$ -modules, called the homological localization map at  $x$ ,

$$pr_x : \widehat{H}_*^{\mathbb{T}}(X; \mathbb{Q})_L \simeq \bigoplus_{x \in X^{\mathbb{T}}} \widehat{H}_*^{\mathbb{T}}(x; \mathbb{Q})_L \longrightarrow \widehat{H}_*^{\mathbb{T}}(x; \mathbb{Q})_L = L^{-1}(\widehat{\Lambda}_{\mathbb{T}})_{\mathbb{Q}}.$$

Moreover, as in [9, Sect.4.2], if  $i_x : \{x\} \hookrightarrow X$  is the inclusion map, then for any  $a \in \widehat{H}_{\mathbb{T}}^*(X; \mathbb{Q})_L$  and  $b \in \widehat{H}_*^{\mathbb{T}}(X; \mathbb{Q})_L$ , we have:

$$(7.8) \quad pr_x(a \cap b) = i_x^*(a) \cap pr_x(b).$$

We then have the following result from [9, Sect.4.2], adapted to the notations of our setting.

**Proposition 7.2.** *Let  $X = X_P$  be the projective toric variety associated to the full-dimensional lattice polytope  $P \subset M_{\mathbb{R}} \cong \mathbb{R}^n$ , with ample Cartier divisor  $D = D_P$ . Then for  $\mathcal{F}$  a  $\mathbb{T}$ -coherent sheaf on  $X$ , the cohomological Euler characteristic  $\chi^{\mathbb{T}}(X, \mathcal{O}_X(D) \otimes \mathcal{F})$  can be calculated via localization at the  $\mathbb{T}$ -fixed points as:*

$$(7.9) \quad \chi^{\mathbb{T}}(X, \mathcal{O}_X(D) \otimes \mathcal{F}) = \sum_{v \text{ vertex of } P} pr_{x_v}(\text{td}_*^{\mathbb{T}}([\mathcal{O}_X(D) \otimes \mathcal{F}]),$$

with each summand

$$(7.10) \quad pr_{x_v}(\text{td}_*^{\mathbb{T}}([\mathcal{O}_X(D) \otimes \mathcal{F}]) \in L^{-1}(\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}},$$

where  $x_v$  is the  $\mathbb{T}$ -fixed point of  $X$  corresponding to the vertex  $v$  of  $P$ , and  $\text{td}_*^{\mathbb{T}} : K_0^{\mathbb{T}}(X) \longrightarrow \widehat{H}_{2*}^{\mathbb{T}}(X; \mathbb{Q})$  is the equivariant Riemann-Roch map of Edidin-Graham [15] (see also [7]).

*Remark 7.3.* The above homological localization formula is in fact a consequence of a similar localization formula in equivariant  $K$ -theory, using the fact that the equivariant Chern character induces a ring homomorphism of localized rings

$$\text{ch}^{\mathbb{T}} : S^{-1}\mathbb{Z}[M] \simeq S^{-1}K_{\mathbb{T}}^0(pt) \longrightarrow L^{-1}(\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}},$$

with  $S$  the multiplicative system generated by elements  $1 - \chi^m$  for  $0 \neq m \in M$ . Note that for a homogeneous polynomial function  $\varphi : M_{\mathbb{R}} \rightarrow \mathbb{C}$ , this property does not hold for the corresponding group homomorphisms  $\widetilde{E}^{\varphi}$  and  $E^{\varphi}$  as defined in the Introduction. For this reason, we use exponential sums and their Taylor expansions (as in [8]), as we shall explain below.  $\square$

Recall the identification  $s : \text{Sym}_{\mathbb{Q}}(M) \simeq H_{\mathbb{T}}^*(pt; \mathbb{Q}) =: (\Lambda_{\mathbb{T}})_{\mathbb{Q}}$ . Let us fix  $z \in N_{\mathbb{C}} := N \otimes_{\mathbb{Z}} \mathbb{C} = \text{Hom}_{\mathbb{R}}(M_{\mathbb{R}}, \mathbb{C})$ . By the universal property of  $\text{Sym}$ ,  $z$  induces an  $\mathbb{R}$ -algebra homomorphism

$$\langle -, z \rangle : \text{Sym}_{\mathbb{R}}(M) \rightarrow \mathbb{C},$$

by which we can view  $\langle p, z \rangle$  for  $z$  now variable and  $p \in \text{Sym}_{\mathbb{R}}(M)$ , resp.,  $p \in (\text{Sym}_{\mathbb{R}}(M))^{an}$  fixed, as a  $\mathbb{C}$ -valued polynomial on  $N_{\mathbb{R}}$ , resp., as a convergent power series function around zero in  $N_{\mathbb{R}}$ .

We can now complete the proof of Theorem 7.1.

*Proof of Theorem 7.1.* From the geometric viewpoint, it is more natural to prove the assertion for  $\tilde{E}_{P,f}^{\varphi}(\ell, y)$ .

By applying the Chern character  $\text{ch}^{\mathbb{T}}$  to formula (6.14), we obtain its cohomological counterpart as follows:

$$\begin{aligned}
(7.11) \quad \sum_{Q \preceq P} f_Q(y) \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} e^{s(m)} &= \chi_y^{\mathbb{T}}(X, \ell D; [\mathcal{M}]) \\
&= \chi^{\mathbb{T}}(X; \text{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(\ell D)) \\
&\stackrel{(7.9)}{=} \sum_{v \text{ vertex}} pr_{x_v}(\text{td}_*^{\mathbb{T}}(\text{DR}_y^{\mathbb{T}}([\mathcal{M}]) \otimes \mathcal{O}_X(\ell D))) \\
&\stackrel{(*)}{=} \sum_{v \text{ vertex}} pr_{x_v}(e^{\ell[D]} \cap \text{td}_*^{\mathbb{T}}(\text{DR}_y^{\mathbb{T}}([\mathcal{M}]))) \\
&\stackrel{(7.8)}{=} \sum_{v \text{ vertex}} e^{\ell \cdot i_{x_v}^*[D]} \cap pr_{x_v}(\text{td}_*^{\mathbb{T}}(\text{DR}_y^{\mathbb{T}}([\mathcal{M}]))),
\end{aligned}$$

where  $i_{x_v} : \{x_v\} \hookrightarrow X$  is the inclusion of the  $\mathbb{T}$ -fixed point of  $X$  corresponding to the vertex  $v$  of  $P$ , with  $f_Q(y) = \chi_y(i_{x_Q}^*[\mathcal{M}])$ , and  $(*)$  makes use of the module property for  $\text{td}_*^{\mathbb{T}}$ .

Applying  $\langle -, z \rangle$  to both sides of (7.11), we obtain

$$(7.12) \quad \sum_{Q \preceq P} f_Q(y) \cdot (1+y)^{\dim(Q)} \cdot \sum_{m \in \text{Relint}(\ell Q) \cap M} e^{\langle m, z \rangle} = \sum_{v \text{ vertex}} e^{\langle \ell \cdot i_{x_v}^*[D], z \rangle} \cdot \langle pr_{x_v}(\text{td}_*^{\mathbb{T}}(\text{DR}_y^{\mathbb{T}}([\mathcal{M}]))), z \rangle.$$

Let us now fix a vertex  $v$  of  $P$ . By (7.10), one also has that

$$pr_{x_v}(\text{td}_*^{\mathbb{T}}(\text{DR}_y^{\mathbb{T}}([\mathcal{M}]))) \in L^{-1}(\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}},$$

since  $e^{\ell \cdot i_{x_v}^*[D]} \in (\Lambda_{\mathbb{T}}^{an})_{\mathbb{Q}}$  is invertible. Then  $\langle pr_{x_v}(\text{td}_*^{\mathbb{T}}(\text{DR}_y^{\mathbb{T}}([\mathcal{M}]))), z \rangle$  is a finite linear combination of terms of the form  $\frac{h(z)}{\prod_{i=1}^k \langle m_i, z \rangle}$ , with  $0 \neq m_i \in M$  and  $h$  analytic in  $z$  near 0.

As in [8, Prop.4.1], we next replace  $z$  by  $tz$  in (7.12), with  $0 \neq t \in \mathbb{R}$  small and  $z$  generic (in the sense that  $\langle m_i, z \rangle \neq 0$ ,  $i = 1, \dots, k$ ), and consider the expansion into Laurent series in  $t$  on both sides of the equality. We then obtain exactly as in [8, Prop.4.1] the polynomial behavior in  $\ell$  of  $\tilde{E}_{P,\mathcal{M}}^{\varphi}(\ell, y)$  for the polynomial function  $\varphi(m) = \langle m, z \rangle^r$ , for any given non-negative integer  $r$  and  $z$  outside a finite number of hyperplanes. This then implies the statement for any polynomial  $\varphi(m)$ .  $\square$

*Remark 7.4.* If we want to estimate the degree in  $\ell$  of the generalized weighted Ehrhart polynomial in Theorem 7.1 to be at most  $\deg(\varphi) + d$  for some  $d \in \mathbb{Z}_{>0}$ , we need to control uniformly the number  $k \leq d$  of linear factors appearing in the above localization. In fact, we can choose  $d = n = \dim(P)$ , as we will now explain. By linearity and using the previous notations, it suffices to work with  $\mathcal{M} = (j_Q)_! \mathbb{Q}_{O_{\sigma_Q}}^H$ , for all faces  $Q$  of  $P$ . By using a resolution of singularities of the orbit closure and functoriality, one can further reduce to the case of a smooth projective toric variety  $X'$  of dimension at most  $n' \leq n$ . In this smooth context, a more precise formula for the homological localization  $pr_x$  to a torus fixed point  $x$  is available via equivariant Poincaré duality as (e.g., see [9, Sect.4.2]):

$$pr_x = \frac{i_x^*}{c_n^{\mathbb{T}}(N_x)},$$

with  $i_x : \{x\} \hookrightarrow X'$  the inclusion map and  $c_n^{\mathbb{T}}(N_x)$  the top equivariant Chern class of the normal bundle  $N_x$  of  $x$  in  $X'$ . Taking a character decomposition of  $N_x = \bigoplus_{i=1}^{n'} \mathbb{C}\chi^{m_i}$ , we get  $c_n^{\mathbb{T}}(N_x) = \prod_{i=1}^{n'} c(m_i)$ .  $\square$

## REFERENCES

- [1] P. Achar, *Equivariant mixed Hodge modules*, preprint, Lecture notes from the Clay Mathematics Institute workshop on Mixed Hodge Modules and Applications (2013), available at <https://www.math.lsu.edu/pramod/docs/emhm.pdf>. [9](#), [15](#), [16](#)
- [2] P. Aluffi, L. Mihalcea, J. Schürmann, C. Su, *Motivic Chern classes of Schubert cells, Hecke algebras, and applications to Casselman’s problem*, arXiv:1902.10101. To appear in *Annales Scientifiques de l’École Normale Supérieure*. [8](#), [13](#)
- [3] M. Beck, P. Gunnells, E. Materov, *Weighted lattice point sums in lattice polytopes, unifying Dehn–Sommerville and Ehrhart–Macdonald*, *Discrete Comput. Geom.* 65 (2021), no. 2, 365–384. [1](#), [2](#), [3](#), [29](#), [32](#)
- [4] F. Bittner, *The universal Euler characteristic for varieties of characteristic zero*, *Compos. Math.* 140 (2004), no. 4, 1011–1032. [15](#)
- [5] T. Braden, R. MacPherson, *Intersection homology of toric varieties and a conjecture of Kalai*, *Comment. Math. Helv.* 74 (1999), no. 3, 442–455. [2](#), [5](#), [29](#)
- [6] J.-P. Brasselet, J. Schürmann, S. Yokura, *Hirzebruch classes and motivic Chern classes for singular spaces*, *J. Topol. Anal.* 2 (2010), no. 1, 1–55. [2](#), [5](#), [9](#), [10](#), [15](#)
- [7] J.L. Brylinski, B. Zhang, *Equivariant Todd Classes for toric varieties*, arXiv:math/0311318. [32](#), [33](#), [34](#)
- [8] M. Brion, M. Vergne, *Lattice points in simple polytopes*, *J. Amer. Math. Soc.* 10 (1997), no. 2, 371–392. [1](#), [3](#), [26](#), [27](#), [31](#), [32](#), [34](#), [35](#)
- [9] S. Cappell, L. Maxim, J. Schürmann, J. Shaneson, *Equivariant toric geometry and Euler–Maclaurin formulae*, arXiv:2303.16785. [1](#), [8](#), [12](#), [13](#), [14](#), [17](#), [19](#), [20](#), [21](#), [24](#), [25](#), [27](#), [32](#), [34](#), [36](#)
- [10] N. Chriss, V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser Boston, Inc., Boston, MA, 1997. [9](#)
- [11] D. A. Cox, J. B. Little, H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, 124. American Mathematical Society, Providence, RI, 2011. [1](#), [2](#), [18](#), [19](#), [21](#), [22](#), [23](#), [24](#)
- [12] V. I. Danilov, *The geometry of toric varieties*, *Russian Math. Surveys* 33 (1978), 97–154. [2](#), [7](#)
- [13] J. Denef, F. Loeser, *Weights of exponential sums, intersection cohomology, and Newton polyhedra*, *Invent. Math.* 106 (1991), no. 2, 275–294. [2](#), [5](#), [29](#)
- [14] B. Dirks, M. Mustață, *The Hilbert series of Hodge ideals of hyperplane arrangements*, *J. Singul.* 20 (2020), 232–250. [6](#), [9](#), [10](#), [13](#), [15](#)

- [15] D. Edidin, W. Graham, *Riemann-Roch for quotients and Todd classes of simplicial toric varieties*, *Comm Algebra* 31 (2003), 3735–3752. [32](#), [34](#)
- [16] K.-H. Fieseler, *Rational intersection cohomology of projective toric varieties*, *J. Reine Angew. Math.* 413 (1991), 88–98. [2](#), [5](#), [29](#)
- [17] W. Fulton, *Introduction to toric varieties*, *Annals of Mathematics Studies*, 131. Princeton University Press, Princeton, NJ, 1993. [1](#), [2](#)
- [18] W. Fulton, R. MacPherson, *Categorical framework for the study of singular spaces*, *Memoirs of the AMS* 243 (1981). [15](#)
- [19] H. Kim, S. Venkatesh, *The intersection cohomology Hodge module of toric varieties*, arXiv:2404.04767. [14](#), [20](#)
- [20] J. Lipmann, M. Hashimoto, *Foundations of Grothendieck Duality for Diagrams of Schemes*, *Lecture Notes in Mathematics*, Vol. 1960. Springer-Verlag, 2009. [6](#), [15](#), [21](#)
- [21] L. Maxim, J. Schürmann, *Characteristic classes of singular toric varieties*, *Comm. Pure Appl. Math.* 68 (2015), no. 12, 2177–2236. [2](#)
- [22] L. Maxim, J. Schürmann, *Weighted Ehrhart theory via mixed Hodge modules on toric varieties*, arXiv:2403.17747. [2](#), [3](#), [4](#), [5](#), [15](#)
- [23] L. Maxim, M. Saito, J. Schürmann, *Hirzebruch-Milnor classes of complete intersections*, *Adv. Math.* 241 (2013), 220–245. [16](#)
- [24] M. Saito, *Modules de Hodge polarisables*, *Publ. Res. Inst. Math. Sci.* 24 (1988), no. 6, 849–995. [9](#)
- [25] M. Saito, *Mixed Hodge modules*, *Publ. Res. Inst. Math. Sci.* 26 (1990), 221–333. [9](#)
- [26] M. Saito, *Mixed Hodge complexes on algebraic varieties*, *Math. Ann.* 316 (2000), 283–331. [14](#)
- [27] M. Saito, *Intersection complexes of toric varieties and mixed Hodge modules*, arXiv:2006.04081. [2](#), [5](#), [29](#)
- [28] J. Schürmann, *Characteristic classes of mixed Hodge modules*, in *Topology of stratified spaces*, 419–470, *Math. Sci. Res. Inst. Publ.*, 58, Cambridge Univ. Press, Cambridge, 2011. [5](#), [9](#), [10](#), [15](#)
- [29] R. Stanley, *Generalized H-vectors, intersection cohomology of toric varieties, and related results*, in *Commutative algebra and combinatorics (Kyoto, 1985)*, 187–213, *Adv. Stud. Pure Math.*, 11, North-Holland, Amsterdam, 1987. [2](#), [29](#)
- [30] H. Sumihiro, *Equivariant completion*, *J. Math. Kyoto Univ.* 14 (1974), 1–28. [9](#), [20](#), [29](#)
- [31] T. Tanisaki, *Hodge modules, equivariant K-theory and Hecke algebras*, *Publ. Res. Inst. Math. Sci.* 23 (1987), no. 5, 841–879. [5](#), [9](#), [11](#), [12](#), [17](#)
- [32] A. Weber, *Equivariant Hirzebruch class for singular varieties*, *Selecta Math. (N.S.)* 22 (2016), no. 3, 1413–1454. [17](#)

L. MAXIM : DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, 480 LINCOLN DRIVE, MADISON WI 53706-1388, USA,

AND

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, 70700 BUCHAREST, ROMANIA.

*Email address:* maxim@math.wisc.edu

J. SCHÜRMAN : MATHEMATISCHE INSTITUT, UNIVERSITÄT MÜNSTER, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY.

*Email address:* jschuerm@math.uni-muenster.de