

All S_p notions of quantum expansion are equivalent

Francisco Escudero Gutiérrez*

Garazi Muguruza†

Abstract

In a recent work Li, Qiao, Wigderson, Wigderson and Zhang introduced notions of quantum expansion based on S_p norms and posed as an open question if they were all equivalent. We give an affirmative answer to this question.

1 Introduction

In this note we consider the problem of showing that certain notions of quantum expansion are equivalent. Quantum expansion is an analogue of the well-studied concept of graph expansion. The latter has many equivalent versions that let it play an important role in many fields such as geometry, group-theory, combinatorics and theoretical computer science (for a survey see [HLW06]).

Given a d -regular graph $G = ([n], E)$, its *edge expansion* is given by

$$h(G) = \min_{W \subset [n], |W| \leq n/2} \frac{|\partial W|}{d|W|},$$

where $|W|$ stands for the size of W and $|\partial W|$ is the number of edges that go from W to its complement. Let $\mathbf{B} = (B_1, \dots, B_d)$ be permutation matrices that decompose the adjacency matrix A of G as $\sum_{i \in [d]} B_i$ ¹. In particular, \mathbf{B} is a bistochastic tuple, meaning that $\sum_{i \in [d]} B_i^* B_i = \sum_{i \in [d]} B_i B_i^* = d \cdot \text{Id}$. Thus, it induces a quantum unital channel via

$$\Phi_{\mathbf{B}}(X) = \frac{1}{d} \sum_{i \in [d]} B_i X B_i^*.$$

A quantum channel is a linear operator from $M_n(\mathbb{C})$ to $M_n(\mathbb{C})$ that maps quantum states (positive semidefinite matrices with trace 1) to quantum states. It is unital if it maps the identity to itself. Note that if \mathbf{B} is a permutation tuple, W a subset of $[n]$ and P_W the projector onto the space spanned by the canonical basis vectors indicated by W , then $\langle \text{Id} - P_W, \Phi_{\mathbf{B}}(P_W) \rangle = |\partial W|$. Here $\langle \cdot, \cdot \rangle$ stands for the inner product $\langle M, N \rangle = \text{Tr}[M^* N]$. Motivated by this, Hastings introduced the *quantum edge expansion* of a bistochastic tuple $\mathbf{B} \in M_n(\mathbb{C})^d$, which is given by

$$h_Q(\mathbf{B}) = \min_{V \subset \mathbb{C}^n, \dim V \leq n/2} \frac{\langle \text{Id} - P_V, \Phi_{\mathbf{B}}(P_V) \rangle}{d \dim V} \quad (1)$$

*Qusoft and CWI feg@cwi.nl

†Qusoft and UvA g.muguruzalasa@uva.nl

¹It follows from Hall's theorem that this decomposition always exists.

where $V < \mathbb{C}^n$ means that V is a subspace of \mathbb{C}^n and P_V is the projector onto V [Has07]. Note that if \mathbf{B} is a permutation tuple determined by a graph G and we restricted the minimum in Eq. (1) to subspaces spanned by vectors of the canonical basis we would recover the edge expansion of G . This is just one of the notions of quantum expansion introduced in the last two decades, which have found multiple applications in cryptography, learning theory and information theory [Has07, BASTS08, HH08, ABW09, KLR19, FM20].

Another analogue of edge expansion in the quantum setting was recently introduced by Li, Qiao, Wigderson, Wigderson and Zhang [LQW⁺22]. Given a bistochastic tuple $\mathbf{B} \in M_n(\mathbb{C})^d$, its *dimension edge expansion* is defined via

$$h_D(\mathbf{B}) = \min_{V < \mathbb{C}^n, \dim V \leq n/2} \frac{\sum_{i \in [n]} \text{rk}(B_i|_{V^\perp, V})}{d \dim V}, \quad (2)$$

where $B_i|_{V^\perp, V} = P_V B_i (\text{Id} - P_V)$. Again, if \mathbf{B} is a permutation tuple determined by a graph G and we restricted the minimum in Eq. (2) to subspaces spanned by vectors of the canonical basis, we would recover the edge expansion of G . Li et al. showed that $d \cdot h_D \geq h_Q$ and they exhibited a sequence \mathbf{B}_m of bistochastic d -tuples such that $\inf h_D(\mathbf{B}_m) > 0$, but $\inf h_Q(\mathbf{B}_m) = 0$ [LQW⁺22, Theorems 1.8 and 1.9]. To do that they considered a different expression of $h_Q(\mathbf{B})$ in terms of the Schatten-2 norm²

$$h_Q(\mathbf{B}) = \underbrace{\min_{V < \mathbb{C}^n, \dim V \leq n/2} \frac{\sum_{i \in [n]} \|B_i|_{V^\perp, V}\|_{S_2}^2}{d \dim V}}_{:= h_{S_2}(\mathbf{B})}.$$

If one substitutes 2 in expression above by other $p \in [1, \infty)$ other natural notions of expansion, namely

$$h_{S_p}(\mathbf{B}) = \min_{V < \mathbb{C}^n, \dim V \leq n/2} \frac{\sum_{i \in [n]} \|B_i|_{V^\perp, V}\|_{S_p}^p}{d \dim V}.$$

These S_p notions of expansion behave in a similar way as the case of $p = 2$, as $d^{p/2} \cdot h_D \geq h_{S_p}$ and the aforementioned sequence \mathbf{B}_m of bistochastic d -tuples also satisfies $\inf h_D(\mathbf{B}_m) > 0$, but $\inf h_{S_p}(\mathbf{B}_m) = 0$. This led Li et. al to ask whether all the S_p notions of expansions are equivalent; h_{S_p} and h_{S_q} are equivalent if for every sequence \mathbf{B}_m of bistochastic tuples d -tuples $\inf h_{S_p}(\mathbf{B}_m) > 0$ if and only if $\inf h_{S_q}(\mathbf{B}_m) > 0$. We give an affirmative answer to this question.

Theorem 1. *Let $\mathbf{B} \in M_n(\mathbb{C})^d$ be a bistochastic tuple. Let $p \geq q \geq 1$. Then,*

1. $h_{S_p}(\mathbf{B}) \leq d^{\frac{p-q}{2}} h_{S_q}(\mathbf{B})$,
2. $h_{S_p}(\mathbf{B}) \geq [h_{S_q}(\mathbf{B})]^{\frac{p}{q}}$,

In particular, h_{S_p} and h_{S_q} are equivalent for every $p, q \in [1, \infty)$.

Li et al. pointed out that Theorem 1 could be a quantum analogue of the equivalence between certain notions of graph expansion based on ℓ_p norms, a result by Matoušek [Mat97], who used this equivalence to prove that the metric spaces defined by graph expanders are hard to embed into ℓ_p . However, we believe the analogy between our result and the one of Matoušek to be more notational than operational, so we suspect that Theorem 1 will not help to construct finite metric spaces hard to embed into S_p . Nevertheless, we consider the latter an interesting open problem and propose a route towards solving it. We discuss this in Section 3.

²For $p \in [1, \infty)$ the Schatten- p norm of a matrix is the ℓ_p norm of its singular values.

2 All S_p notions of expansion are equivalent

Proof of Theorem 1.1: As B_i belongs to a d -stochastic tuple, it satisfies $B_i^* B_i = d \cdot \text{Id} - \sum_{j \neq i} B_j^* B_j \preceq d \cdot \text{Id}$, so all the singular values (s_1^i, \dots, s_n^i) of $B_i|_{V^\perp, V}$ are at most \sqrt{d} . Hence,

$$\frac{\|B_i|_{V^\perp, V}\|_{S_p}^p}{d^{\frac{p}{2}}} = \sum_{l \in [n]} \left(\frac{s_l^i}{\sqrt{d}} \right)^p \leq \sum_{l \in [n]} \left(\frac{s_l^i}{\sqrt{d}} \right)^q = \frac{\|B_i|_{V^\perp, V}\|_{S_q}^q}{d^{\frac{q}{2}}},$$

where in the inequality we have used that all s_l^i/\sqrt{d} are at most 1 and that $p \geq q$. From here, the desired inequality follows immediately. \square

Proof of Theorem 1.2: Let $(s_1^i, \dots, s_{\dim V}^i)$ be the singular values of $B_i|_{V^\perp, V}$. Then by Hölder's inequality,

$$\|B_i|_{V^\perp, V}\|_{S_q}^q = \sum_{l \in [\dim V]} |s_l^i|^q \leq \left(\sum_{l \in [\dim V]} |s_l^i|^p \right)^{q/p} \dim V^{1-\frac{q}{p}} = \|B_i|_{V^\perp, V}\|_{S_p}^q \dim V^{1-\frac{q}{p}}. \quad (3)$$

We conclude that for every subspace $V < \mathbb{C}^n$, we have that

$$\sum_{i \in [d]} \frac{\|B_i|_{V^\perp, V}\|_{S_q}^q}{d \dim V} \leq \sum_{i \in [d]} \frac{\|B_i|_{V^\perp, V}\|_{S_p}^q}{d \dim V^{q/p}} = d^{\frac{q}{p}-1} \sum_{i \in [d]} \left(\frac{\|B_i|_{V^\perp, V}\|_{S_p}^p}{d \dim V} \right)^{q/p} \leq \left(\sum_{i \in [d]} \frac{\|B_i|_{V^\perp, V}\|_{S_p}^p}{d \dim V} \right)^{q/p},$$

where in the first inequality we used Eq. (3) and in the second Hölder's inequality. From here, the desired inequality follows immediately. \square

3 How much distortion is needed to embed metric spaces into S_p ?

3.1 Embeddings into ℓ_p spaces

A metric space (M, ρ) can be embedded into ℓ_p with distortion D if there is a map $f : M \rightarrow \ell_p$ such that

$$\frac{1}{D} \rho(x, y) \leq \|f(x) - f(y)\|_{\ell_p} \leq \rho(x, y), \quad \forall x, y \in M. \quad (4)$$

Finding the minimum distortion D_{n, ℓ_p} needed to embed any n -point metric space into ℓ_p was a long-standing question. The case $p = 1$ was fully resolved by Bourgain (who proved the upper bound) and Linial, London and Rabinovich (who showed the lower bound). Together they concluded that $D_{\ell_1} = \Theta(\log n)$ [Bou85, LLR95]. The key ingredient of the lower bound of Linial et al. is that for a d -regular graph G it holds that

$$\underbrace{\inf_{f: G \rightarrow \ell_1} \frac{\frac{1}{|E|} \sum_{(i,j) \in E} \|f(i) - f(j)\|_{\ell_1}}{\frac{1}{n^2} \sum_{i,j \in [n]} \|f(i) - f(j)\|_{\ell_1}}}}_{:= h_{\ell_1}(G)} \geq h(G). \quad (5)$$

Matoušek realized that generalizing this result to other ℓ_p norms would allow one to prove similar lower bounds. With that purpose he (implicitly) introduced notions of graph expansion based on

other ℓ_p norms, namely

$$h_{\ell_p}(G) := \min_{f:G \rightarrow \ell_p} \left(\frac{\frac{1}{|E|} \sum_{(i,j) \in E} \|f(i) - f(j)\|_{\ell_p}^p}{\frac{1}{n^2} \sum_{i,j \in [n]} \|f(i) - f(j)\|_{\ell_p}^p} \right)^{1/p}.$$

Then, he showed that

$$h_{\ell_p}(G) \geq \frac{h_{\ell_1}(G)}{4dp}, \quad (6)$$

which, together with (5) and the existence of a sequence G_n of n -vertex d -regular graphs such that $\inf h(G_n) = C > 0$, implies that

$$\inf h_{\ell_p}(G_n) \geq \frac{C}{4dp}. \quad (7)$$

From Eq. (7), Matoušek showed with an elegant argument that the metric spaces defined by these graphs require distortion $D_n = \Omega(\log(n)/p)$ to be embedded into ℓ_p . Indeed, consider the metric spaces $([n], \rho_n)$, where $\rho_n(i, j)$ is the length of the shortest path in G_n from i to j . Define

$$R_{\rho_n} = \left(\frac{\frac{1}{|E|} \sum_{(i,j) \in E} \rho_n(i, j)^p}{\frac{1}{n^2} \sum_{i,j \in [n]} \rho_n(i, j)^p} \right)^{1/p}.$$

The numerator of R_{ρ_n} is 1, while the denominator is $\Omega(\log_d n)$, because there are at most $d^1 + \dots + d^{\log_d(n/4)} \leq 2d^{\log_d(n/4)} = n/2$ pairs of vertices whose shortest-path distance is at most $\log_d(n/4)$. Hence, treating d as a constant, it follows that $R_{\rho_n} = O(1/\log(n))$. Now let f_n be embeddings of $([n], \rho_n)$ into ℓ_p with distortion D_n and define R_{f_n} in an analogue way. As f_n are embeddings with distortion D_n it follows that $R_{f_n} = O(D_n/\log(n))$ and Eq. (7) implies that $R_{f_n} = \Omega(1/p)$. Putting both bounds together yields $D_{n, \ell_p} = \Omega(\log(n)/p)$. Matoušek also proved the matching upper bound by mimicking the argument of Bourgain, giving $D_{n, \ell_p} = \Theta(\log(n)/p)$.

3.2 What about S_p ?

As pointed out by Li et al. [LQW⁺22], our Theorem 1 can be regarded as a quantum analogue of Eq. (6) proved by Matoušek, in the sense that both prove relations between notions of expansion based on similar norms. However, we believe that this resemblance is notational, rather than operational, and so our Theorem 1 does not seem to help to construct metric spaces that are hard to embed into S_p . To the best of our knowledge there is no literature about this problem, so we pose it as an open question.

Question 2. *Let $p \in [1, \infty)$. What is the minimum distortion D_{n, S_p} needed, as a function of n and p , to embed any n -point metric space into S_p ?*

To answer this question it would suffice to show that the following family of notions of graph expansion based on S_p norms for $p \in [1, \infty)$ are related in a certain way

$$\tilde{h}_{S_p}(G) := \min_{f:G \rightarrow S_p} \left(\frac{\frac{1}{|E|} \sum_{(i,j) \in E} \|f(i) - f(j)\|_{S_p}^p}{\frac{1}{n^2} \sum_{i,j \in [n]} \|f(i) - f(j)\|_{S_p}^p} \right)^{1/p}.$$

Indeed, as S_2 is isometric to ℓ_2 , we have that there exists a sequence of n -vertex d -regular graphs G_n such that $\inf \tilde{h}_{S_2}(G_n) > 0$. Hence, if one manages to show that for every d -regular graph

G it holds that

$$\tilde{h}_{S_p}(G) = \Omega_d \left(\frac{\tilde{h}_{S_2}(G)}{p} \right),$$

we would have that $\inf \tilde{h}_{S_p}(G_n) = \Omega(1/p)$, from which we could reproduce the argument of Matoušek and prove that $D_{n,S_p} = \Omega(\log(n)/p)$. The matching upper bound follows from the fact that ℓ_p can be embedded isometrically into S_p .

Acknowledgements. We thank Jop Briët for detailed comments and discussions. We thank Davi Castro-Silva, Yinan Li, Youming Qiao, Yuval Wigderson and Chuanqi Zhang for useful comments and discussions. This research was supported by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 945045, and by the NWO Gravitation project NETWORKS under grant no. 024.002.003.

References

- [ABW09] Andris Ambainis, Jan Bouda, and Andreas Winter. Nonmalleable encryption of quantum information. *Journal of Mathematical Physics*, 50(4), 2009.
- [BASTS08] Avraham Ben-Aroya, Oded Schwartz, and Amnon Ta-Shma. Quantum expanders: Motivation and constructions. In *2008 23rd Annual IEEE Conference on Computational Complexity*, pages 292–303, 2008.
- [Bou85] Jean Bourgain. On lipschitz embedding of finite metric spaces in Hilbert space. *Israel Journal of Mathematics*, 52:46–52, 1985.
- [FM20] William Cole Franks and Ankur Moitra. Rigorous guarantees for Tyler’s M-estimator via quantum expansion. In *Conference on Learning Theory*, pages 1601–1632. PMLR, 2020.
- [Has07] Matthew B Hastings. Random unitaries give quantum expanders. *Physical Review A*, 76(3):032315, 2007.
- [HH08] Matthew B Hastings and Aram W Harrow. Classical and quantum tensor product expanders. *arXiv preprint arXiv:0804.0011*, 2008.
- [HLW06] Shlomo Hoory, Nathan Linial, and Avi Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(4):439–561, 2006.
- [KLR19] Tsz Chiu Kwok, Lap Chi Lau, and Akshay Ramachandran. Spectral analysis of matrix scaling and operator scaling. In *2019 IEEE 60th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1184–1204. IEEE, 2019.
- [LLR95] Nathan Linial, Eran London, and Yuri Rabinovich. The geometry of graphs and some of its algorithmic applications. *Combinatorica*, 15:215–245, 1995.
- [LQW⁺22] Yinan Li, Youming Qiao, Avi Wigderson, Yuval Wigderson, and Chuanqi Zhang. On linear-algebraic notions of expansion. *arXiv preprint arXiv:2212.13154*, to appear in *Theory of Computing*, 2022.

- [Mat97] Jiří Matoušek. On embedding expanders into l_p spaces. *Israel Journal of Mathematics*, 102:189–197, 1997.