On a Completion of Cohomological Functors Generalising Tate Cohomology I

Max Gheorghiu

7 June 2024

Abstract

Viewing group cohomology as a so-called cohomological functor, G. Mislin has generalised Tate cohomology from finite groups to all discrete groups by defining a completion for cohomological functors in [28]. If $T^{\bullet}: \mathcal{C} \to \mathcal{D}$ is any cohomological functor, then we construct its Mislin completion $\widehat{T}^{\bullet}: \mathcal{C} \to \mathcal{D}$ under the assumption that the abelian category \mathcal{C} has enough projectives and that in the abelian category \mathcal{D} all countable direct limits exist and are exact. This takes Tate cohomology to settings where it has never been introduced such as in condensed mathematics. Through the latter, one can define Tate cohomology for any T1 topological group. More specifically, we generalise four constructions of Mislin completions from the literature, prove that they yield isomorphic cohomological functors and provide explicit formulae for their connecting homomorphisms. For any morphism $f: M \to N$ in \mathcal{C} we develop formulae for $\widehat{T}^n(f): \widehat{T}^n(M) \to \widehat{T}^n(N)$ in terms of each construction.

Contents

1	Introduction 1.1 Notation and terminology	$rac{2}{7}$
2	Outline of constructions	7
3	The satellite functor construction and Mislin completions	11
4	The resolution and the satellite functor construction	16
	4.1 Isomorphism of cohomology groups	17
	4.2 Induced morphisms and connecting homomorphisms	
	4.3 Isomorphisms of cohomological functors	
	4.4 Countably many distinct constructions of Mislin completions	
5	The naïve and the resolution construction	30
	5.1 Performing the naïve construction	30
	5.2 Isomorphism of cohomology groups	33
	5.3 Connecting homomorphisms of the naïve construction	37

6	The hypercohomology and the resolution construction		
	6.1	Defining Ext-functors through chain maps	40
	6.2	Connecting homomorphisms through chain maps	43
	6.3	Isomorphism of cohomology groups	47
	6.4	Isomorphism of cohomological functors	51
7	Con	nplete cohomology in condensed mathematics	57

61

References

1 Introduction

Because Tate cohomology and its generalisations are the main motivation of this work, we first outline Tate cohomology for a finite group G based on [1, p. 78-79]. For any module M over the group ring $\mathbb{Z}[G]$ and any integer $n \in \mathbb{Z}$ there is a Tate cohomology group $\widehat{H}^n(G, M)$. Note that $\widehat{H}^n(G, M) = H^n(G, M)$ for $n \geq 1$ and $\widehat{H}^n(G, M) = H_{-n-1}(G, M)$ for $n \leq -2$, whence Tate cohomology unites both group cohomology and group homology. According to [6, p. 136], Tate cohomology of G satisfies dimension shifting and possesses cup products. The former means that for any M there are $\mathbb{Z}[G]$ -modules M_* , M^* such that $\widehat{H}^{n-1}(G, M_*) \cong \widehat{H}^n(G, M)$ and $\widehat{H}^{n+1}(G, M^*) \cong \widehat{H}^n(G, M)$ for every $n \in \mathbb{Z}$. Thus, if one desires to establish a property for every degree $\widehat{H}^n(G, -)$ of $\widehat{H}^{\bullet}(G, -)$, then one only needs to establish it for a carefully chosen degree n. If N is a $\mathbb{Z}[G]$ -module and $m \in \mathbb{Z}$ another integer, then a cup product is a homomorphism of the form

$$\smile: \widehat{H}^n(G, M) \otimes \widehat{H}^m(G, N) \to \widehat{H}^{m+n}(G, M \otimes N)$$

where the tensor product $M \otimes N$ is a $\mathbb{Z}[G]$ -module endowed with the diagonal action of G. Importantly for us, Tate cohomology for G vanishes on projectives according to [1, p. 79], meaning that $\widehat{H}^n(G, P) = 0$ for any projective $\mathbb{Z}[G]$ -module P and $n \in \mathbb{Z}$. Then complete cohomology generalises Tate cohomology for finite groups and was introduced for all discrete groups by G. Mislin, P. Vogel, D. J. Benson and J. F. Carlson using different constructions all yielding isomorphic cohomology groups [25, p. 196–197].

On the one hand we generalise these constructions to a previously unknown extent. On the other hand, we develop for each construction explicit formulae for connecting homomorphisms and induced morphisms that have not been covered in the literature. In particular, since our generalisation takes Tate cohomology to condensed mathematics, one can construct Tate cohomology for any topological group in which every point is closed, meaning for any T1 topological group. More specifically, condensed mathematics is a novel theory developed by D. Clausen and P. Scholze in 2018 [35]. In [36], P. Scholze writes that he wants "to make the strong claim that in the foundations of mathematics, one should replace topological spaces with condensed sets". Independently of whether this claim turns out to be true, it is a very promising tool for the study of continuous cohomology of topological groups and the study of representations of topological groups to topological abelian groups [35, p. 6]. It also serves as a foundation for a very general form of analytic geometry [9, p. 6]. More fundamentally, it provides a unified approach for studying topological groups, rings and modules [35, p. 6]. In the subsequent paper [16], we show that complete cohomology detects whether a group has finite cohomological dimension, establish a version of dimension shifting, an Eckmann-Shapiro result as well as cohomology products such as external products, cup products and Yoneda products.

Let us provide more detail on the nature of our constructions. As in the case of G. Mislin's work [28], the notion of a cohomological functor is essential. More specifically, if \mathcal{C}, \mathcal{D} denote two abelian categories, then a family of additive functors $(T^n: \mathcal{C} \to \mathcal{D})_{n \in \mathbb{Z}}$ is a cohomological functor if there are so-called connecting homomorphisms $(\delta^n: T^n \to T^{n+1})_{n \in \mathbb{Z}}$ satisfying two (natural) axioms [25, p. 201–202]. In particular, if G is a discrete group, R a discrete ring and A a discrete R-module, then setting $H^n_R(G, -) = 0$ and $\operatorname{Ext}^n_R(A, -) = 0$ for n < 0 renders group cohomology and Ext-functors into cohomological functors [25, p. 201], [28, p. 295]. We refer the reader unfamiliar with abelian categories to the account in [27, p. 249–257], [38, Tag 00ZX] or [42, Section A.4]. G. Mislin generalised Tate cohomology to all discrete groups by defining complete cohomology $\widehat{H}^{\bullet}_{R}(G,-)$ as a specific completion of ordinary group cohomology $H^{\bullet}_{R}(G, -)$ in his paper [28]. Following the convention from [25, p. 197/202], we call such a completion of a cohomological functor a Mislin completion. Formally, a Mislin completion of $T^{\bullet}: \mathcal{C} \to \mathcal{D}$ is a cohomological functor $\widehat{T}^{\bullet}: \mathcal{C} \to \mathcal{D}$ together with a morphism $\Phi^{\bullet}: T^{\bullet} \to \widehat{T}^{\bullet}$ such that $\widehat{T}^n(P) = 0$ for any projective $P \in obj(\mathcal{C})$ and $n \in \mathbb{Z}$ and such that any morphism $T^{\bullet} \to V^{\bullet}$ to a cohomological functor $V^{\bullet}: \mathcal{C} \to \mathcal{D}$ also vanishing on projectives factors uniquely through Φ^{\bullet} [28, Definition 2.1]. By its universal property, any Mislin completion is unique up to isomorphism [25, p. 202].

We generalise two constructions of Mislin completions due to G. Mislin [28] under the assumption that the domain category \mathcal{C} has enough projective objects and that in the codomain category \mathcal{D} all countable direct limits exist and are exact (Lemma 3.2 and Theorem 4.5). Direct limits are a specific kind of colimits [34, p. 14] and are vital to these constructions. In doing so, we demonstrate that there are countably many distinct constructions of Mislin completions (Lemma 4.10). If Ab denotes the category of abelian groups, then we generalise P. Vogel's [17] as well as D. J. Benson and J. F. Carlson's construction [5] mentioned above so that they give rise to the Mislin completion of (unenriched) Ext-functors $\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -) : \mathcal{C} \to \operatorname{Ab}$ that we call completed (unenriched) Ext-functors (Theorem 5.10 and Corollary 6.15). Especially in the context of condensed mathematics, we also consider Hom-functors of the form $\operatorname{Hom}_{\mathcal{C}}: \mathcal{C}^{\operatorname{op}} \times \mathcal{C} \to \mathcal{D}$ called enriched Hom-functors where the category \mathcal{D} does not need to be **Ab** and the corresponding enriched Ext-functors $\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -) : \mathcal{C} \to \mathcal{D}$ together with their completed enriched Extfunctors $\widehat{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(A, -)$. For any morphism $f: M \to N$ in \mathcal{C} we develop formulae for the induced morphism $\widehat{T}^n(f): \widehat{T}^n(M) \to \widehat{T}^n(N)$ of the Mislin completion in terms of each construction (Proposition 3.4, Proposition 4.2, Definition 5.4, Definition 6.2). Using each construction, we provide explicit formulae for the connecting homomorphisms of \hat{T}^{\bullet} . (Lemma 3.3, Definition 4.3, Lemma 5.9, Definition 6.5).

Before presenting some examples, we note that that complete cohomology generalises Tate-Farrell cohomology, which we prove in [16]. More specifically, F. T. Farrell generalised Tate cohomology of finite groups to discrete groups having a finite index subgroup of finite cohomological dimension in his paper [14] by using so-called complete resolutions. A complete resolution of an object A in C is a particular acyclic chain complex $(\overline{A}_n)_{n\in\mathbb{Z}}$ of projective objects in C that agrees with a projective resolution of M in sufficiently high degree [10, Definition 1.1]. Following [14, p. 158], Tate-Farrell Ext-functors are defined as the cohomology of the cochain complex $\operatorname{Hom}_{\mathcal{C}}(\overline{A}_{\bullet}, B)$ where we the Hom-functor is unenriched. Using F. T. Farrell's work, P. Symonds deduces in [40, p. 34] that Tate-Farrell cohomology exists for every profinite group having an open subgroup of finite cohomological dimension. In [39], he computes the Tate-Farrell cohomology of the Morava stabiliser group S_{p-1} with coefficients in the moduli space E_{p-1} for odd primes p.

However, there are groups for which their complete cohomology cannot be calculated by a complete resolution. For instance, a free abelian group of countable rank $\bigoplus_{n \in \mathbb{N}} \mathbb{Z}$, $GL_n(K)$ for K a subfield of the algebraic closure of \mathbb{Q} , the Thompson group F and the free product $\bigstar_{n \in \mathbb{N}} \mathbb{Z}^n$ do not admit complete resolutions [10, Example 5.3], [11, p. 119–120]. By [28, p. 297–298], the cohomology of the first three examples is isomorphic to their complete cohomology while all cohomology groups of the last group are distinct from its complete cohomology groups [23, p. 432]. The zeroeth complete cohomology group of this last example is calculated in [11, Corollary 2.4 and Theorem A]. In [12], F. Dembegioti calculates the zeroeth complete cohomology group of a class of discrete polycyclic groups, but also explains why one cannot determine a general formula calculating the zeroeth complete cohomology groups for all polycyclic groups. Lastly, if p is a prime number, then pro-p groups are a class of profinite groups for which there is a rich theory of their group cohomology. See for instance [37, Chapter I] or [41]. One example of a pro-p group that is also a pro-p ring are the p-adic integers \mathbb{Z}_p which are studied in [43, Section 1.5]. These examples give rise to the following questions.

Question 1.1 As in [34, Example 3.3.8(c)], let $\prod_{\mathbb{N}} \mathbb{Z}_p$ denote the free abelian pro-p group over \mathbb{N} . Is its cohomology $H^{\bullet}_{\mathbb{Z}_p}(\prod_{\mathbb{N}} \mathbb{Z}_p, -)$ isomorphic to its complete cohomology $\widehat{H}^{\bullet}_{\mathbb{Z}_p}(\prod_{\mathbb{N}} \mathbb{Z}_p, -)$ as a cohomological functor?

Question 1.2 In accordance with [33, p. 137–140], denote by $G = \bigsqcup_{n \in \mathbb{N} \cup \{\infty\}} G_n$ the following free pro-p product over the one-point compactification of the integers $\mathbb{N} \cup \{\infty\}$. Set $G_n = \mathbb{Z}_p^n$ for $n \in \mathbb{N}$ and $G_\infty = \{1\}$. Can one compute the zeroeth complete cohomology $\widehat{H}^0_{\mathbb{Z}_n}(G, A)$ for an $\mathbb{Z}_p[[G]]$ -module A?

Let us elaborate on the implementation of complete cohomology into condensed mathematics. If **Pro** denotes the category of profinite spaces, then, informally, a condensed set can be thought as a contravariant functor $X : \mathbf{Pro} \to \mathbf{Set}$ satisfying certain compatibility conditions [35, p. 7]. Given any T1 topological space Y, one can turn it into a condensed set

$$\underline{Y}: \mathbf{Pro}^{\mathrm{op}} \to \mathbf{Set}, S \mapsto \{f: S \to Y \text{ continuous}\}$$

called the condensate of T [35, p. 15/16 and Proposition 1.7]. Condensed groups (resp. rings, modules etc.) are defined analogously [35, p. 7] and condensates of T1 topological groups are condensed groups (resp. rings, modules etc) [35, p. 8]. Because we define condensed sets and modules by using sheaves, they possess excellent category theoretic properties such as the existence of all small limits and colimits [35, p. 11/15] as well as enough projectives [9, p. 16]. Moreover, all products, direct sums and direct limits of condensed modules are exact [35, Theorem 2.2]. J. Anschütz and A.-C. Le Bras have defined condensed group cohomology over any condensed group ring $\mathcal{R}[\mathcal{G}]$ in [2, p. 5] since condensed abelian groups form an abelian category contrary to topological abelian groups [35, p. 6/11]. We note that for condensed \mathcal{R} -modules over a condensed ring \mathcal{R} there is not only an unenriched Hom-functor $\operatorname{Hom}_{\operatorname{Cond}(\mathcal{R})}(-,-)$ mapping to abelian groups [35, p. 13].

Theorem 1.3 (= Theorem 7.1) Let \mathcal{R} be a condensed ring and A a condensed \mathcal{R} -module. Then there are completed condensed unenriched Ext-functors $\widehat{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$ and completed condensed internal Ext-functors $\widehat{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$. If \mathcal{G} is a condensed group, then complete condensed unenriched group cohomology $\widehat{H}^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ and complete condensed internal group cohomology $\widehat{H}^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ can be defined.

Corollary 1.4 One can define group cohomology and thus complete cohomology for any T1 topological group G by taking its condensate <u>G</u>.

Specifically for profinite groups, it is more natural to consider so-called solid modules. In the same manner as a completed group ring of a profinite group over a profinite ring can be seen as a completion of an ordinary group ring [43, Proposition 7.1.2], one can introduce a notion of completion for condensed modules over a condensed ring \mathcal{R} . We call such completed condensed modules solid [35, p. 32/44] and \mathcal{R} analytic if it admits a choice of solid modules [35, Definition 7.4].

Lemma 1.5 (= Lemma 7.3) Let \mathcal{A} be an analytic ring and M a solid \mathcal{A} -module. Then there are completed solid unenriched Ext-functors $\widehat{\operatorname{Ext}}_{\operatorname{Solid}(\mathcal{A})}^{\bullet}(M, -)$.

For a profinite group G and for analytic rings \mathcal{R} of a specific form, J. Anschütz and A.-C. Le Bras define in [2, p. 2/5] solid group cohomology $H^{\bullet}_{\text{Solid}(\mathcal{R})}(\underline{G}, -)$ of the condensate of G taking coefficients in solid $\mathcal{R}[\underline{G}]$ -modules. They mention some profinite rings Rwhose condensates \underline{R} are analytic rings of this form [2, p. 2]. As any condensate \underline{M} of a profinite or discrete R-module M is a solid \underline{R} -module for such rings [2, p. 4], one has $H^{\bullet}_{\text{Solid}(\underline{R})}(\underline{G},\underline{M}) \cong H^{\bullet}_{R}(G,M)$ [2, Lemma 2.1]. We prove that there exists complete solid cohomology $\widehat{H}^{\bullet}_{\text{Solid}(\mathcal{R})}(\underline{G},-)$ mapping to abelian groups and complete solid internal group cohomology $\widehat{H}^{\bullet}_{\text{Solid}(\mathcal{R})}(\underline{G},-)$ mapping to solid $\underline{\mathbb{Z}}$ -modules whenever the analytic ring \mathcal{R} is of the required form (Lemma 7.5). In particular, the following question arises.

Question 1.6 When is the complete cohomology of a profinite group $\widehat{H}^{\bullet}_{R}(G, -)$ isomorphic to the complete solid cohomology of its condensate $\widehat{H}^{\bullet}_{\text{Solid}(\underline{R})}(\underline{G}, -)$ as a cohomological functor?

We refer the reader to M. Land's notes [26] and P. Scholze's notes [35] for more detail on condensed mathematics. For results on condensed and solid group cohomology the reader is referred to J. Anschütz and A.-C. Le Bras' preprint [2].

Let us detail how our work compares to the literature. Recently, S. Guo and L. Liang have generalised in [19] the above constructions by G. Mislin [28], by D. J. Benson and J. F. Carlson [5] and by P. Vogel [17] for unenriched relative Ext-functors over any abelian category \mathcal{C} with a special precovering subcategory \mathcal{W} and only proved that the resulting completed relative Ext-functors $\widehat{\operatorname{Ext}}_{\mathcal{WC}}^n(A, -) : \mathcal{C} \to \operatorname{Ab}$ are naturally isomorphic for every $n \in \mathbb{Z}$. Although they prove that their completed Ext-functors $\widehat{\operatorname{Ext}}_{\mathcal{WC}}^n(A, -)$ form a cohomological functor, we do not know whether they also form a Mislin completion of the Ext-functors $\operatorname{Ext}_{\mathcal{WC}}^n(A, -)$ (Question 6.19). On the other hand, A. Beligiannis and I. Reiten establish specific Mislin completions in their book on torsion theories [4]. In the elegant account from their Section IX.2, they consider an abelian category \mathcal{C} containing a particular pair of subcategories $(\mathcal{X}, \mathcal{Y})$, define a completed unenriched relative Ext-functor $\widehat{\operatorname{Ext}}_{(\mathcal{X},\mathcal{Y})}^{\bullet}(-,-)$ using a version of D. J. Benson and J. F. Carlson's construction from [5] and prove that $\operatorname{Ext}_{\mathcal{X}\cap\mathcal{Y}}^{\bullet}(A,-) \to \widehat{\operatorname{Ext}}_{(\mathcal{X},\mathcal{Y})}^{\bullet}(A,-)$ forms a Mislin completion for any $A \in \operatorname{obj}(\mathcal{C})$. Lastly, J. Hu et al. use in [22] a version of P. Vogel's construction from [17] to establish completed unenriched Ext-functors $\widehat{\operatorname{Ext}}_{\mathcal{C}}^{\bullet}(-,-)$ for an extriangulated category \mathcal{C} where extriangulated categories are a generalisation of exact categories and triangulated categories. Among their applications, they provide a criterion for the validity of the Wakamatsu Tilting Conjecture. As a common application of completed unenriched Ext-functors, all above mentioned authors consider various notions of homological dimension for which they characterise when a homological dimension of an object in \mathcal{C} is finite.

Apart from providing different applications in [16], our contribution mainly consists in establishing a Mislin completion whenever we are given any cohomological functor between two particular abelian categories. For instance, we can define completed internal condensed Ext-functors which cannot be established through the previous frameworks (Remark 7.2). Moreover, in none of the above sources, connecting homomorphisms or induced morphisms of Mislin completions are constructed explicitly.

For an overview of various other generalisations of Tate cohomology and their applications, the reader is referred to M. Paganin's survey paper [31]. To mention two developments that are not contained in it, a version of completed Ext-functors for complexes of modules over a ring is defined in [3] and a generalisation of Tate cohomology in the language of spectra can be found in [24]. B. E. A. Nucinkis observed first in [29] that all construction can be performed dually by assuming that there are enough injectives instead of enough projectives. More specifically, if \mathcal{C} is an abelian category with enough injectives and \mathcal{D} is an abelian category in which all countable direct limits exist and are exact, then one can define for a cohomological functor $T^{\bullet}: \mathcal{C} \to \mathcal{D}$ a "Nucinkis completion" $T^{\bullet} \to \widetilde{T}^{\bullet}$ such that \widetilde{T}^{\bullet} vanishes on injectives and is universal with respect to this property. For Extfunctors, this completion takes the form $\operatorname{Ext}^{\bullet}_{\mathcal{C}}(-, A) \to \widetilde{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(-, A)$. S. Guo and L. Liang perform all their constructions of completed unenriched relative Ext-functors in a dual manner in [19]. Together with X. Yang they provide in [20, Main Theorem] a criterion for when $\widetilde{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(-,-)$ is isomorphic to $\widetilde{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(-,-)$ as well as A. Beligiannis and I. Reiten in [4, Theorem IX.4.4] as well as S. Hu et al. in [21, Theorem 4.4].

After outlining four constructions of Mislin completions in Section 2, we prove in the subsequent four sections that they result in isomorphic cohomological functors and provide explicit formulae of their connecting homomorphisms and induced morphisms. More specifically, in Section 3 we demonstrate that one construction in [28] by G.Mislin gives rise to Mislin completions. In Section 4 we prove that another construction of a cohomological functor from [28] is isomorphic to the former. Thereby we deduce that there are countably many distinct constructions of Mislin completions. Then in Section 5 we show that P. Vogel's construction in [17] gives rise to completed unenriched Ext-functors that are isomorphic to the ones from the latter construction by G. Mislin. In Section 6 we prove that the completed unenriched Ext-functors resulting from D.J.Benson and J.F.Carlson's construction in [5] are also isomorphic to the ones from the latter construction by G. Mislin. Lastly, in Section 7, we outline how one can introduce completed Ext-functors and complete cohomology in condensed mathematics.

1.1 Notation and terminology


We adopt the convention that the natural numbers \mathbb{N} include 1, but do not include 0. We write \mathbb{N}_0 for $\mathbb{N} \cup \{0\}$. Moreover, we adopt B. Poonen's convention from [32] that a ring is an abelian group together with a totally associative product, meaning a binary associative relation admitting an identity element. In particular, ring homomorphisms are understood to map identity elements to identity elements. We use the symbol $\varinjlim_{\substack{i \in I}}$ only to denote direct limits. If \mathcal{D} is a category, then $\varinjlim_{\substack{i \in I}}$ denotes a direct limit in this category and if I is a directed set, then $\varinjlim_{\substack{i \in I}}$ denotes a direct limit indexed over I. We write $\varinjlim_{k \in \mathbb{N}} (M_k, \mu_k)$ if $\mu_k : M_k \to M_{k+1}$ are the morphisms giving rise to the direct limit. Lastly, we use the same numbering to label diagrams and equations.

2 Outline of constructions

To showcase the relevant constructions of Mislin completions, let us axiomatically define cohomological functors. For two abelian categories \mathcal{C} , \mathcal{D} we call a family of additive functors $(T^n : \mathcal{C} \to \mathcal{D})_{n \in \mathbb{Z}}$ a cohomological functor if it satisfies the following two axioms [25, p. 201–202].

Axiom 2.1 For every $n \in \mathbb{Z}$ and short exact sequence $0 \to A \to B \to C \to 0$ in \mathcal{C} , there is natural connecting homomorphism $\delta^n : T^n(C) \to T^{n+1}(A)$.

Being natural means in this context that for every commuting diagram in C



with exact rows there is a commuting diagram

$$T^{n}(A) \xrightarrow{\delta^{n}} T^{n+1}(A)$$

$$\downarrow^{T^{n}(g)} \qquad \qquad \downarrow^{T^{n+1}(f)}$$

$$T^{n}(A') \xrightarrow{\delta^{n}} T^{n+1}(C')$$

in \mathcal{D} .

Axiom 2.2 For every short exact sequence $0 \to A \xrightarrow{\iota} B \xrightarrow{\pi} C \to 0$ in C there is a long exact sequence

$$\dots \xrightarrow{T^{n-1}(\pi)} T^{n-1}(C) \xrightarrow{\delta^{n-1}} T^n(A) \xrightarrow{T^n(\iota)} T^n(B) \xrightarrow{T^n(\pi)} T^n(C) \xrightarrow{\delta^n} T^{n+1}(A) \xrightarrow{T^{n+1}(\iota)} \dots$$

Assume that $(T^{\bullet}, \delta^{\bullet})$ and $(U^{\bullet}, \varepsilon^{\bullet})$ are cohomological functors from \mathcal{C} to \mathcal{D} . Then a family of natural transformations $(\nu^n : T^n \to U^n)_{n \in \mathbb{Z}}$ is a morphism of cohomological functors if it satisfies the axiom [25, p. 202]

Axiom 2.3 For every $n \in \mathbb{Z}$ and short exact sequence $0 \to A \to B \to C \to 0$ in C, the following square commutes:

$$T^{n}(C) \xrightarrow{\delta^{n}} T^{n+1}(A)$$

$$\downarrow^{\nu^{n}} \qquad \qquad \downarrow^{\nu^{n+1}}$$

$$U^{n}(C) \xrightarrow{\varepsilon^{n}} U^{n+1}(A)$$

Let us generalise G. Mislin's Definition 2.1 from [28].

Definition 2.4 (Mislin completion) Let $(T^{\bullet}, \delta^{\bullet})$ be a cohomological functor from \mathcal{C} to \mathcal{D} . Then its Mislin completion is a cohomological functor $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ from \mathcal{C} to \mathcal{D} together with a morphism $\nu^{\bullet} : T^{\bullet} \to \widehat{T}^{\bullet}$ satisfying the following two conditions:

1. For every $n \in \mathbb{Z}$ and every projective $P \in obj(\mathcal{C})$ we have $\widehat{T}^n(P) = 0$.

2. If $(U^{\bullet}, \varepsilon^{\bullet})$ is any cohomological functor vanishing on projectives, then each morphism $T^{\bullet} \to U^{\bullet}$ factors uniquely as $T^{\bullet} \xrightarrow{\nu^{\bullet}} \widehat{T}^{\bullet} \to U^{\bullet}$.

Hence, if $(U^{\bullet}, \varepsilon^{\bullet})$ is another Mislin completion of $(T^{\bullet}, \delta^{\bullet})$, then there is an isomorphism $\mu^{\bullet} : \widehat{T}^{\bullet} \to U^{\bullet}$, meaning that $\mu^{n}(M) : \widehat{T}^{n}(M) \to U^{n}(M)$ is an isomorphism for any $n \in \mathbb{Z}$ and $M \in obj(\mathcal{C})$ [25, p. 202]. This allows us to state G. Mislin's definition from [28, p. 297] in greater generality.

Definition 2.5 (Axiomatic, Mislin) For any $A \in obj(\mathcal{C})$ we extend the (enriched or unenriched) Ext-functors to cohomological functor by setting $\operatorname{Ext}^n_{\mathcal{C}}(A, -) = 0$ for n < 0and define completed Ext-functors as the Mislin completion ($\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -), \widehat{\delta}^{\bullet}$). Analogously, if G is a group object in C and R a ring object, then we extend group cohomology to a cohomological functor ($H^{\bullet}_{R}(G, -), \delta^{\bullet}$) by imposing $H^n_{R}(G, -) = 0$ for n < 0 and define complete cohomology as the Mislin completion ($\widehat{H}^{\bullet}_{R}(G, -), \widehat{\delta}^{\bullet}$).

To ensure that complete cohomology exists, we introduce left satellite functors to present a construction due to G. Mislin. As we assume that \mathcal{C} has enough projective objects, there is for any $M \in \text{obj}(\mathcal{C})$ a short exact sequence $0 \to K \to P \to M \to 0$ in \mathcal{C} with P projective. For a cohomological functor $(T^{\bullet}, \delta^{\bullet})$, we define the zeroeth left satellite functor of T^n as $S^0T^n := T^n$, the first left satellite functor as

$$S^{-1}T^n(M) := \operatorname{Ker}(T^n(K) \to T^n(P))$$

and the k^{th} left satellite functor as $S^{-k}T^n := S^{-1}(S^{-k+1}T^n)$ for $k \geq 2$ [7, p. 36]. It is shown in [7, Section III.1] that left satellite functors do not depend on the choice of short exact sequence. Since they have been defined as kernels, it follows from Axiom 2.2 that $\delta^n : T^n \to T^{n+1}$ induces a morphism $\underline{\delta}^n : T^n(M) \to S^{-1}T^{n+1}(M)$ and therefore $S^{-k}\underline{\delta}^{n+k} : S^{-k}T^{n+k}(M) \to S^{-k-1}T^{n+k+1}(M)$ for any $k \in \mathbb{N}$ [25, p. 207–208]. Here, our assumption hits in that all countable direct limits exist in the codomain category \mathcal{D} of T^{\bullet} .

Definition 2.6 A partially ordered set (I, \leq) is a directed set if for every $i, j \in I$ there is $k \in I$ such that $i, j \leq k$ [34, p. 1]. According to [34, p. 14], a diagram $\{D_i\}_{i\in I}$ in \mathcal{D} indexed over a directed set is called a direct system in \mathcal{D} . More formally, I can be turned into a category whose objects are its elements and there is a unique morphism $i \to j$ whenever $i \leq j \in I$. Then a direct system is a covariant functor $I \to \mathcal{D}, i \mapsto D_i$. A direct limit $\varinjlim_{i\in I} D_i$ in \mathcal{D} is a colimit of a direct system $\{D_i\}_{i\in I}$. Direct limits in \mathcal{D} are called exact if for every direct system of short exact sequence $\{0 \to A_i \to B_i \to C_i \to 0\}_{i\in I}$ also

$$0 \to \varinjlim_{i \in I} A_i \to \varinjlim_{i \in I} B_i \to \varinjlim_{i \in I} C_i \to 0$$

is a short exact sequence [38, Tag 079A].

Hence, we can extend G. Mislin's construction from [28, p. 293].

Definition 2.7 (Via satellite functors, Mislin) The Mislin completion of a cohomological functor $(T^{\bullet}, \delta^{\bullet})$ can be defined as $\widehat{T}^{n}(M) := \varinjlim_{k \in \mathbb{N}_{0}} (S^{-k}T^{n+k}(M), S^{-k}\underline{\delta}^{n+k})$ for any $M \in \operatorname{obj}(\mathcal{C})$ and $n \in \mathbb{Z}$. Accordingly, $\widehat{H}^{n}_{R}(G, M) := \varinjlim_{k \in \mathbb{N}_{0}} (S^{-k}H^{n+k}_{R}(G, M), S^{-k}\underline{\delta}^{n+k})$ is a definition of complete cohomology.

In order that the above forms a cohomological functor, one needs to impose that all direct limits in \mathcal{D} are exact. The reader might be aware that one can define cohomology of discrete groups and cohomology of profinite groups taking discrete coefficients by using that the respective category \mathcal{C} has enough injectives [6, p. 61], [37, p. 9]. One could assume instead that \mathcal{C} has enough injectives and in the category \mathcal{D} all countable inverse limits exist and are exact. One could then perform a construction via right satellite functors dual to the above. This completion of group cohomology would have a universal property as a Mislin completion, except that it would vanish on injective objects instead of projective ones. However, cohomology of discrete groups as well as of profinite groups taking discrete coefficients already vanishes on injectives [6, p. 61], [37, p. 9]. Thus, such a completion would not yield Tate cohomology whence we assume that \mathcal{C} has enough injectives.

Notation 2.8 For the rest of the paper, C always denotes an abelian category with enough projectives and D an abelian category in which all countable direct limits exist and are exact.

Let us go over to what we term the resolution construction that occurs in [8, Lemma B.3] and can be retrieved from page 299 in G. Mislin's paper [28]. If $(M_n)_{n\in\mathbb{N}_0}$ is a projective resolution of $M \in \operatorname{obj}(\mathcal{C})$, let us define $\widetilde{M}_0 := M$ and $\widetilde{M}_k := \operatorname{Ker}(M_{k-1} \to \widetilde{M}_{k-1})$ for $k \in \mathbb{N}$. This is called the k^{th} syzygy of M_{\bullet} in the Gorenstein context [31, p. 89]. The choice of our notation is meant to reflect that our syzygies do not necessarily arise from a specific choice of projective resolution as in [25] and [28]. Since for every $k \in \mathbb{N}_0$ we have the short exact sequence $0 \to \widetilde{M}_{k+1} \to M_k \to \widetilde{M}_k \to 0$, there is a connecting homomorphism $\delta^{n+k} : T^{n+k}(\widetilde{M}_k) \to T^{n+k+1}(\widetilde{M}_{k+1})$ for every $n \in \mathbb{Z}$. Then the following definition makes it more apparent why Mislin completions vanish on projective objects.

Definition 2.9 (Resolutions, Mislin) The Mislin completion of a cohomological functor $(T^{\bullet}, \delta^{\bullet})$ can be defined as $\widehat{T}^{n}(M) := \varinjlim_{k \in \mathbb{N}_{0}} (T^{n+k}(\widetilde{M}_{k}), \delta^{n+k})$ for any $n \in \mathbb{Z}$ and $M \in \operatorname{obj}(\mathcal{C})$. Accordingly, $\widehat{H}^{n}_{R}(G, M) := \varinjlim_{k \in \mathbb{N}_{0}} (H^{n+k}_{R}(G, \widetilde{M}_{k}), \delta^{n+k})$ is a definition of complete cohomology.

The next two constructions only give rise to completed unenriched Ext-functors. Let A_{\bullet} , B_{\bullet} are projective resolutions of $A, B \in \text{obj}(\mathcal{C})$ and let $\tilde{f}_{n+k} : \tilde{A}_{n+k} \to \tilde{B}_k$ be a morphism for $n \in \mathbb{Z}$ and $k \in \mathbb{N}_0$ such that $n+k \geq 0$. Then we can write the commuting diagram

$$0 \longrightarrow \widetilde{A}_{n+k+1} \longrightarrow A_{n+k} \longrightarrow \widetilde{A}_{n+k} \longrightarrow 0$$
$$\downarrow \widetilde{f}_{k+1} \qquad \qquad \downarrow f_{k+1} \qquad \qquad \downarrow \widetilde{f}_{k}$$
$$0 \longrightarrow \widetilde{B}_{k+1} \longrightarrow B_{k+1} \longrightarrow \widetilde{B}_{k} \longrightarrow 0$$

whose terms arise as follows. Since the bottom row is exact and the term A_{n+k} projective, there is a lift f_k of \tilde{f}_k making the right-hand square commute. Because $\tilde{B}_{k+1} \to B_{k+1}$ is a kernel, there is a morphism \tilde{f}_{k+1} making the left-hand side commute. If $\operatorname{Hom}_{\mathcal{C}}(-,-)$ denotes the (unenriched) Hom-functor in \mathcal{C} , then $\operatorname{Hom}_{\mathcal{C}}(\tilde{A}_{n+k}, \tilde{B}_k)$ is an abelian group by virtue of \mathcal{C} being an abelian category. We define $\mathcal{P}_{\mathcal{C}}(\tilde{A}_{n+k}, \tilde{B}_k)$ to be the subgroup of $\operatorname{Hom}_{\mathcal{C}}(\tilde{A}_{n+k}, \tilde{B}_k)$ consisting of all morphisms factoring through a projective object and write the quotient as $[\tilde{A}_{n+k}, \tilde{B}_k]_{\mathcal{C}} := \operatorname{Hom}_{\mathcal{C}}(\tilde{A}_{n+k}, \tilde{B}_k)/\mathcal{P}_{\mathcal{C}}(\tilde{A}_{n+k}, \tilde{B}_k)$ [25, p. 203]. As in the case of modules over a ring covered by [25, p. 204], we prove that

$$t_{\widetilde{A}_{n+k},\widetilde{B}_{k}} : [\widetilde{A}_{n+k},\widetilde{B}_{k}]_{\mathcal{C}} \to [\widetilde{A}_{n+k+1},\widetilde{B}_{k+1}]_{\mathcal{C}},$$

$$\widetilde{f}_{k} + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k},\widetilde{B}_{k}) \mapsto \widetilde{f}_{k+1} + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k+1},\widetilde{B}_{k+1})$$

is a well defined homomorphism. Through this we generalise the following definition from D. J. Benson and J. F. Carlson's paper [5, p. 109].

Definition 2.10 (Naïve construction, Benson & Carlson) We can define the n^{th} completed unenriched Ext-functor as $\widehat{\operatorname{Ext}}_{\mathcal{C}}^{n}(A,B) := \varinjlim_{k \in \mathbb{N}_{0}, n+k \geq 0} ([\widetilde{A}_{n+k}, \widetilde{B}_{k}]_{\mathcal{C}}, t_{\widetilde{A}_{n+k}, \widetilde{B}_{k}})$ for $n \in \mathbb{Z}$. In particular, if R_{\bullet} is a projective R[G]-resolution of $R \in \operatorname{obj}(\mathcal{C}_{R,G})$, we can define complete unenriched cohomology as $\widehat{H}_{R}^{n}(G, B) := \varinjlim_{k \in \mathbb{N}_{0}, n+k \geq 0} ([\widetilde{R}_{n+k}, \widetilde{B}_{k}]_{\mathcal{C}}, t_{\widetilde{R}_{n+k}, \widetilde{B}_{k}}).$

Lastly, we present what we call the hypercohomology construction of complete cohomology. We define the chain complex $(A'_n)_{n\in\mathbb{Z}}$ by $A'_n = A_n$ for $n \ge 0$ and $A'_n = 0$ for n < 0 and similarly $(B'_n)_{n\in\mathbb{Z}}$ [25, p. 209]. Define the hypercohomology complex $(\operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_n, d^n)_{n\in\mathbb{Z}}$ by having *n*-cochains $\operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_n = \prod_{k\in\mathbb{Z}} \operatorname{Hom}_{\mathcal{C}}(A'_{k+n}, B'_k)$. To ease notation in the following, we view abelian groups as \mathbb{Z} -modules. If we denote by $a_n : A'_n \to A'_{n-1}$ and $b_n : B'_n \to B'_{n-1}$ the differentials induced from the respective projective resolution, we define for $n \in \mathbb{Z}$ the differential

$$d^{n}: \operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_{n} \to \operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_{n+1}$$

$$(\varphi_{n+k})_{k\in\mathbb{Z}} \mapsto (b_{k+1} \circ \varphi_{n+k+1} - (-1)^{n} \varphi_{n+k} \circ a_{n+k+1})_{k\in\mathbb{Z}}$$

$$(2.1)$$

Let us define the bounded complex $\operatorname{Bdd}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_{n \in \mathbb{Z}}$ as the subcomplex of $\operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_{n \in \mathbb{Z}}$ given by $\operatorname{Bdd}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_n = \bigoplus_{k \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{C}}(A'_{k+n}, B'_k)$. Now we define the Vogel complex $\operatorname{Vog}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_{n \in \mathbb{Z}}$ as the quotient complex $\operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_n/\operatorname{Bdd}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_n$ [25, p. 209]. By this, we generalise Definition 1.2 from F. Goichot's paper [17] where he attributes it to P. Vogel on page 39.

Definition 2.11 (Hypercohomology, Vogel) For $n \in \mathbb{Z}$ we can define the n^{th} completed unenriched Ext-functor as $\widehat{\operatorname{Ext}}_{\mathcal{C}}^{n}(A, B) := H^{n}((\operatorname{Vog}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})_{k}, d^{k})_{k \in \mathbb{Z}})$. We can thus define complete unenriched cohomology as $\widehat{H}_{R}^{n}(G, M) := H^{n}((\operatorname{Vog}_{R}(R'_{\bullet}, B'_{\bullet})_{k}, d^{k})_{k \in \mathbb{Z}})$.

Let us remark why it is unlikely that the previous two constructions could yield Mislin completions of more general enriched Ext-functors. For the naïve construction, one aims to find a morphism $\operatorname{Hom}_{\mathcal{C}}(\widetilde{A}_{n+k}, \widetilde{B}_k) \to \operatorname{Hom}_{\mathcal{C}}(\widetilde{A}_{n+k+1}, \widetilde{B}_{k+1})$. If one tries to lift along the morphisms induced by $A_{n+k} \to \widetilde{A}_{n+k}$ and $B_{n+k} \to \widetilde{B}_{n+k}$, one requires that $\operatorname{Hom}_{\mathcal{C}}(A_{n+k}, -)$ preserves epimorphisms. For unenriched Hom-functors, this role is exactly played by projective objects [42, Lemma 2.2.3]. For the hypercohomology construction we require that the coproduct $\operatorname{Bdd}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})$ maps into the product $\operatorname{Hyp}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})$. Moreover, as we shall see in Subsection 6.1, the cohomology groups of the Vogel complex $\operatorname{Vog}_{\mathcal{C}}(A'_{\bullet}, B'_{\bullet})$ correspond exactly to so-called almost chain maps modulo almost chain homotopy. However, this implies that the objects $\operatorname{Hom}_{\mathcal{C}}(A_p, B_q) \in \operatorname{obj}(\mathcal{D})$ describe morphisms of the form $A_p \to B_q$ in \mathcal{C} .

3 The satellite functor construction and Mislin completions

Satellite functors are treated in the third chapter of H. Cartan and S. Eilenberg's book on homological algebra [7]. By using morphisms of cohomological functors, G. Mislin constructs Mislin completions through satellite functors in the proof of Theorem 2.2 from [28] while P. H. Kropholler provides more explicit descriptions of the terms of such a Mislin completion on pages 206–208 of [25]. In this section we generalise G. Mislin and P. H. Kropholler's work that is performed only using modules over a ring. As we also aim to determine explicit formulae for the induced morphisms and for the connecting homomorphisms of the resulting Mislin completions, we provide full details on the satellite functor construction. In turn, these details are needed later to manufacture the induced morphisms and connecting homomorphisms of all other constructions of Mislin completions.

For the reader's convenience, we summarise relevant results on left satellite functors from the third chapter of [7]. They may be working explicitly in a category of modules over a ring, but A. Grothendieck has generalised their satellite functors in his Tôhoku paper [18, p. 140–143] and thus, their arguments pertain to our abelian categories. In the first section, they show that left satellite functors are well defined and additive functors. More specifically, let $f: C' \to C$ be a morphism in \mathcal{C} . As we assume that the abelian category \mathcal{C} has enough projectives, one can take short exact sequences

$$0 \to M' \to P' \to C' \to 0$$
 and $0 \to M \to P \to C \to 0$

with P' and P projective. Using this and that every monomorphism is normal in an abelian category, we deduce the existence of morphisms $\overline{f}^* : P' \to P$ and $\overline{f}' : M' \to M$ rendering the diagram

commutative. To ease future notation, let $T^{\bullet} : \mathcal{C} \to \mathcal{D}$ be a cohomological functor with connecting homomorphisms δ^{\bullet} . The morphism $T^n(\overline{f}^*) : T^n(M') \to T^n(M)$ induces the corresponding morphism between the kernels $S^{-1}T^n(f) : S^{-1}T^n(C') \to S^{-1}T^n(C)$. Pictorially,

$$S^{-1}T^{n}(C') \xrightarrow{\varepsilon^{-1}} T^{n}(M')$$
$$\downarrow^{S^{-1}T^{n}(f)} \qquad \downarrow^{T^{n}(\overline{f}^{*})}$$
$$S^{-1}T^{n}(C) \xrightarrow{\varepsilon^{-1}} T^{n}(M)$$

where the morphisms ε^{-1} are the canonical monomorphisms from the kernels. Being independent of the choices of short exact sequences, the morphism $S^{-1}T^n(f)$ is well defined. The morphisms $S^{-k}T^n(f): S^{-k}T^n(C') \to S^{-k}T^n(C)$ for higher satellite functors are defined in the same manner. In the second section, they construct for any exact sequence $0 \to A \to B \to C \to 0$ in \mathcal{C} and any $k \in \mathbb{N}_0$ a connecting homomorphism $\varepsilon^{-k-1}: S^{-k-1}T^n(C) \to S^{-k}T^n(A)$ in the following manner. If B is projective, then ε^{-k-1} is the monomorphism from the kernel $S^{-k-1}T^n(C)$. Otherwise let $0 \to M \to P \to C \to 0$ be a short exact sequence with P projective. As we have seen above, there are morphisms $h: P \to B$ and $h^*: M \to A$ extending the identity morphism $id_C: C \to C$ making the diagram

commute. Then $\varepsilon^{-k-1}: S^{-k-1}T^n(C) \to S^{-k}T^n(M) \xrightarrow{S^{-k}T^n(h^*)} S^{-k}T^n(A)$ is a connecting homomorphism where the first morphism is the monomorphism of the kernel $S^{-n-1}T^n(C)$. Moreover they show that the every connecting homomorphism is a natural transformation. Extending into the third section, they use these to construct the long exact sequence

$$\dots \to S^{-k-1}T^n(C) \xrightarrow{\varepsilon^{-k-1}} S^{-k}T^n(A) \to S^{-k}T^n(B) \to S^{-k}T^n(C) \xrightarrow{\varepsilon^{-k}} S^{-k+1}T^n(A) \to \dots$$

where the unlabelled morphisms are induced from the underlying short exact sequence. In the fifth section, they provide the following crucial result corresponding to Proposition 5.2 on page 46:

Lemma 3.1 Let T^{\bullet} and U^{\bullet} be cohomological functors and $\Phi^{0}: T^{0} \to U^{0}$ be a natural transformation. If for every short exact sequence $0 \to K \to P \to M \to 0$ with P projective and every $n \in \mathbb{N}$ the sequence $0 \to U^{-n-1}(M) \to U^{-n}(K)$ involving the connecting homomorphism is exact, then Φ^{0} extends uniquely to a (partial) morphism of cohomological functors $(\Phi^{n}: T^{n} \to U^{n})_{n \leq 0}$ only defined for $n \leq 0$. In particular, if also T^{\bullet} has the property that the sequence $0 \to T^{-n-1}(M) \to T^{-n}(K)$ is exact for $n \in \mathbb{N}$ and Φ^{0} is an equivalence, then the extension Φ^{\bullet} is an isomorphism.

G. Mislin uses this lemma to perform his satellite functor construction on pages 295–296 in [28] which we generalise in the following. For any $m \in \mathbb{N}_0$ we can define the cohomological functor

$$T^{k}\langle m \rangle(M) := \begin{cases} S^{k-m}T^{m}(M) & \text{if } k < m\\ T^{k}(M) & \text{if } k \ge m \end{cases}$$
(3.3)

where the connecting homomorphisms are given by $\delta^k \langle m \rangle = \delta^k : T^k \to T^{k+1}$ for $k \geq m$ and $\delta^k \langle m \rangle = \varepsilon^{k-m} : S^{k-m}T^m \to S^{k-m+1}T^m$ for k < m. Consider the partial morphism $(\Phi_m^k := \mathrm{id} : T^k \to T^k \langle m \rangle)_{m \leq k}$ of cohomological functors defined only in degrees $m \leq k$. According to the previous lemma we can extend Φ_m^{\bullet} uniquely to a morphism of cohomological functors defined in all degrees. Using the same arguments we have for $m \leq n$ a unique morphism $\Phi_{m,n}^{\bullet} : T^{\bullet} \langle m \rangle \to T^{\bullet} \langle n \rangle$ that is equal to the identity morphism in any degree $n \leq k \in \mathbb{Z}$. Using the uniqueness of all these morphisms, we conclude for $m \leq n \leq o \in \mathbb{N}_0$ that $\Phi_n^{\bullet} = \Phi_{m,n}^{\bullet} \circ \Phi_m^{\bullet}$ and $\Phi_{m,o}^{\bullet} = \Phi_{n,o}^{\bullet} \circ \Phi_{m,n}^{\bullet}$. This yields a morphism to a direct system $T^{\bullet} \to (T^{\bullet} \langle m \rangle, \Phi_{m,n}^{\bullet})_{m \leq n \in \mathbb{N}_0}$. This is where countable direct limits $\varinjlim_{\mathcal{D}}$ in the codomain category \mathcal{D} come into play. For every $n \in \mathbb{Z}$ we define the n^{th} term of the satellite functor construction as

$$\widehat{T}^n(M) := \lim_{k \in \mathbb{N}_0} T^n \langle k \rangle(M) \,.$$

Unravelling the definitions and noting that $\{k \in \mathbb{N}_0 \mid k \ge n\}$ is cofinal in \mathbb{N}_0 , we obtain

$$\widehat{T}^{n}(M) = \lim_{k \in \mathbb{N}_{0}, k \ge n} \left(S^{n-k} T^{k}(M), \Phi^{n}_{k,k+1}(M) \right) = \lim_{k \in \mathbb{N}_{0}} \left(S^{-k} T^{n+k}(M), \Phi^{n}_{n+k,n+k+1}(M) \right).$$
(3.4)

As T^{\bullet} is a cohomological functor and left satellite functors are additive, each functor $S^{-k}T^{n+k}$ is additive. Since $\varinjlim_{\mathcal{D}}$ is additive, \widehat{T}^n is an additive functor. Every short exact sequence $0 \to A \to B \to \overrightarrow{C} \to 0$ in the category \mathcal{C} yields a long exact sequence in the cohomological functors T^{\bullet} and $T^{\bullet}\langle m \rangle$ for any $m \in \mathbb{N}_0$ by Axiom 2.2. Hence we obtain a long exact sequence in $\widehat{T}^{\bullet} = \varinjlim_{\mathcal{D}, m \in \mathbb{N}_0} T^{\bullet}\langle m \rangle$ and in particular, a connecting homomorphism $\widehat{\delta}^{\bullet}$ only if the direct limit functor $\varinjlim_{\mathcal{D}, m \in \mathbb{N}_0}$ is exact.

More specifically, to demonstrate that $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ is a cohomological functor it suffices to show that $\widehat{\delta}^{\bullet}$ is a natural transformation and how a long exact sequence in \widehat{T}^{\bullet} arises from a short exact sequence $0 \to A \xrightarrow{f} B \xrightarrow{g} C \to 0$ in \mathcal{C} . As for every $m \leq n \in \mathbb{N}_0$ we have a morphism of cohomological functors $\Phi^{\bullet}_{m,n}$, the following diagram commutes for every $k \in \mathbb{Z}$:

As $\Phi_{m,n}^k(-)$ (resp. $\Phi_{m,n}^{k+1}(-)$) is a functor and $\delta^k \langle m \rangle$ (resp. $\delta^{k+1} \langle n \rangle$) is natural, the connecting homomorphism $\widehat{\delta}^k = \varinjlim_{\mathcal{D},n \in \mathbb{N}_0} \delta^k \langle n \rangle : \widehat{T}^k(C) \to \widehat{T}^{k+1}(A)$ is natural by the above square. Using again that $\Phi_{m,n}^{\bullet}$ is a morphism of cohomological functors, also the following diagram commutes:

$$\dots \longrightarrow T^{k-1}\langle m \rangle(C) \xrightarrow{\delta^{k-1}\langle m \rangle} T^k \langle m \rangle(A) \xrightarrow{T^k \langle m \rangle(f)} T^k \langle m \rangle(B) \xrightarrow{T^k \langle m \rangle(g)} T^k \langle m \rangle(C) \xrightarrow{\delta^k \langle m \rangle} \dots \\ \downarrow \Phi^{k-1}_{m,n}(C) \qquad \qquad \downarrow \Phi^k_{m,n}(A) \qquad \qquad \downarrow \Phi^k_{m,n}(B) \qquad \qquad \downarrow \Phi^k_{m,n}(C) \\ \dots \longrightarrow T^{k-1}\langle n \rangle(C) \xrightarrow{\delta^{k-1}\langle n \rangle} T^k \langle n \rangle(A) \xrightarrow{T^k \langle n \rangle(f)} T^k \langle n \rangle(B) \xrightarrow{T^k \langle n \rangle(g)} T^k \langle n \rangle(C) \xrightarrow{\delta^k \langle n \rangle} \dots$$

Since the above constitutes a direct system of long exact sequences and countable direct limits $\varinjlim_{\mathcal{D}}$ are exact, we obtain the corresponding long exact sequence in $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$

$$\dots \longrightarrow \widehat{T}^{k-1}(C) \xrightarrow{\widehat{\delta}^{k-1}} \widehat{T}^k(A) \xrightarrow{\widehat{T}^k(f)} \widehat{T}^k(B) \xrightarrow{\widehat{T}^k(g)} \widehat{T}^k(C) \xrightarrow{\widehat{\delta}^k} \dots$$

Lemma 3.2 The cohomological functor $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ constructed via satellite functors is a Mislin completion of $(T^{\bullet}, \delta^{\bullet})$.

Proof As for G. Mislin's satellite functor construction, the proof is an adaptation of material found on page 296 of his paper [28]. Let P be projective projective object in \mathcal{C} . If we choose the short exact sequence $0 \to (0 \to P \to P) \to 0$, then we see that $S^{-1}T^n(P) = 0$ and thus $S^{-k}T^{n+k}(P) = 0$ for any $k \ge 1$. In particular, $(\widehat{T}^{\bullet}, \delta^{\bullet})$ vanishes on projectives. Let us define a canonical morphism $\Phi^{\bullet}: (T^{\bullet}, \delta^{\bullet}) \to (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ such that any morphism $\Psi^{\bullet}: (T^{\bullet}, \delta^{\bullet}) \to (U^{\bullet}, \zeta^{\bullet})$ to a cohomological functor vanishing on projectives factors uniquely as

$$\Psi^{\bullet}: (T^{\bullet}, \delta^{\bullet}) \xrightarrow{\Phi^{\bullet}} (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet}) \to (U^{\bullet}, \zeta^{\bullet}) \,.$$

By Lemma 3.1 there is a morphism $\Phi_m^{\bullet} : (T^{\bullet}, \delta^{\bullet}) \to (T^{\bullet}\langle m \rangle, \delta^{\bullet}\langle m \rangle)$ for any $m \in \mathbb{N}_0$ and denote by $\widehat{\Phi}_m^{\bullet} : (T^{\bullet}\langle m \rangle, \delta^{\bullet}\langle m \rangle) \to (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ the morphism to the direct limit. Defining $\Phi^{\bullet} := \widehat{\Phi}_m^{\bullet} \circ \Phi_m^{\bullet} : (T^{\bullet}, \delta^{\bullet}) \to (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ is then independent of $m \in \mathbb{N}_0$ since

$$\widehat{\Phi}_{m}^{\bullet} \circ \Phi_{m}^{\bullet} = (\widehat{\Phi}_{n}^{\bullet} \circ \Phi_{m,n}^{\bullet}) \circ \Phi_{m}^{\bullet} = \widehat{\Phi}_{n}^{\bullet} \circ (\Phi_{m,n}^{\bullet} \circ \Phi_{m}^{\bullet}) = \widehat{\Phi}_{n}^{\bullet} \circ \Phi_{n}^{\bullet}$$

for any $m \leq n$. We can also define a morphism $\Psi_m^{\bullet} : (T^{\bullet}\langle m \rangle, \delta^{\bullet}\langle m \rangle) \to (U^{\bullet}, \zeta^{\bullet})$ by setting $\Psi_m^n = \Psi^n$ for any $m \leq n$. Using Lemma 3.1 we extend Ψ_m^{\bullet} uniquely to a morphism of cohomological functors. Given that $\Phi_m^n = \mathrm{id}_{\mathcal{D}}$ for $m \leq n \in \mathbb{Z}$, we have $\Psi^n = \Psi_m^n \circ \Phi_m^n$ and Lemma 3.1 implies that

$$\Psi^{\bullet} = \Psi^{\bullet}_m \circ \Phi^{\bullet}_m : (T^{\bullet}, \delta^{\bullet}) \to (T^{\bullet} \langle m \rangle, \delta^{\bullet} \langle m \rangle) \to (U^{\bullet}, \zeta^{\bullet}) .$$

Deduce analogously $\Psi_m^{\bullet} = \Psi_n^{\bullet} \circ \Phi_{m,n}^{\bullet}$ for any $m \leq n \in \mathbb{N}_0$. We apply the universal property of direct limits to obtain the unique morphism $\widehat{\Psi}^{\bullet} : (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet}) \to (U^{\bullet}, \zeta^{\bullet})$ such that $\Psi_m^{\bullet} = \widehat{\Psi}^{\bullet} \circ \widehat{\Phi}_m^{\bullet}$. Therefore

$$\Psi^{\bullet} = \Psi^{\bullet}_{m} \circ \Phi^{\bullet}_{m} = \widehat{\Psi}^{\bullet} \circ \widehat{\Phi}^{\bullet}_{m} \circ \Phi^{\bullet}_{m} = \widehat{\Psi}^{\bullet} \circ \widehat{\Phi}^{\bullet} : (T^{\bullet}, \delta^{\bullet}) \to (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet}) \to (U^{\bullet}, \zeta^{\bullet}).$$

We want to show that this factorisation is unique. So assume that $\Psi^{\bullet} = X^{\bullet} \circ \widehat{\Phi}^{\bullet}$ is another factorisation. Thus

$$\widehat{\Psi}^{\bullet} \circ \widehat{\Phi}^{\bullet}_{m} \circ \Phi^{\bullet}_{m} = \widehat{\Psi}^{\bullet} \circ \widehat{\Phi}^{\bullet} = X^{\bullet} \circ \widehat{\Phi}^{\bullet} = X^{\bullet} \circ \widehat{\Phi}^{\bullet}_{m} \circ \Phi^{\bullet}_{m}.$$

As $\Phi_m^n = \mathrm{id}_{\mathcal{D}}$ for $m \leq n \in \mathbb{N}_0$, $\widehat{\Psi}^n \circ \widehat{\Phi}_m^n = X^n \circ \widehat{\Phi}_m^n$ and

$$\Psi_m = \widehat{\Psi}^{\bullet} \circ \widehat{\Phi}_m^{\bullet} = X^{\bullet} \circ \widehat{\Phi}_m^{\bullet} : (T^{\bullet} \langle m \rangle, \delta^{\bullet} \langle m \rangle) \to (U^{\bullet}, \zeta^{\bullet})$$

by Lemma 3.1. Since $\widehat{\Psi}^{\bullet}$ is the unique morphism such that $\Psi_m^{\bullet} = \widehat{\Psi}^{\bullet} \circ \widehat{\Phi}_m^{\bullet}$ for any $m \in \mathbb{N}_0$, we conclude that $X^{\bullet} = \widehat{\Psi}^{\bullet}$.

Let us provide an explicit formula for the connecting homomorphisms of the satellite functor construction. For this, we require H. Cartan and S. Eilenberg's extension of the concept of left satellite functors to natural transformations from [7, Corollary 5.3] that we present in the following. As before, their considerations for modules over a ring pass to abelian categories. Let $\varphi: F \to G$ be a natural transformation between two half-exact functors $F, G: \mathcal{C} \to \mathcal{D}$, meaning that for any short exact sequence $0 \to A \to B \to C \to 0$ in \mathcal{C} the induced sequences in \mathcal{D}

$$F(A) \to F(B) \to F(C)$$
 and $G(A) \to G(B) \to G(C)$

are exact [25, p. 205]. For $C \in obj(\mathcal{C})$, let $0 \to M \to P \to C \to 0$ be a short exact sequence with P projective. This induces the commutative diagram

$$F(M) \longrightarrow F(P) \longrightarrow F(C)$$

$$\downarrow^{\varphi(M)} \qquad \qquad \downarrow^{\varphi(P)} \qquad \qquad \downarrow^{\varphi(C)}$$

$$G(M) \longrightarrow G(P) \longrightarrow G(C)$$

In the same manner as we have defined morphisms $S^{-1}T^n(f): S^{-1}T^n(C') \to S^{-1}T^n(C)$ between first left satellite functors, we denote by $S^{-1}\varphi(C): S^{-1}F(C) \to S^{-1}G(C)$ the morphism induced from $\varphi(M): F(M) \to G(M)$ between the corresponding kernels. Pictorially

Write $S^0 \varphi := \varphi$ and

$$S^{-k}\varphi := S^{-1}(S^{-k+1}\varphi) : S^{-k}F \to S^{-k}G \,,$$

for any $k \in \mathbb{N}$. We term this the k^{th} left satellite transformation of φ . Now remember that any connecting homomorphism $\delta^n : T^n \to T^{n+1}$ is a natural transformation. We have seen in Subsection 2 that it factors through $\underline{\delta}^n : T^n(M) \to S^{-1}T^{n+1}(M)$ for any $M \in \text{obj}(\mathcal{C})$. Then P. H. Kropholler's account [25, p. 206–208] suggests the following result.

Lemma 3.3 For any $n \in \mathbb{Z}$, $\underline{\delta}^n : T^n \to S^{-1}T^{n+1}$ is a natural transformation. In case that $0 \to A \to B \to C \to 0$ is a short exact sequence in \mathcal{C} , we take the connecting homomorphism $\varepsilon^{-k} : S^{-k}T^{n+k}(C) \to S^{-k+1}T^{n+k}(A)$ for $k \in \mathbb{N}_0$. Then the n^{th} connecting homomorphism of the satellite functor construction is given by

$$\widehat{\delta}^n = \varinjlim_{k \in \mathbb{N}} \varepsilon^{-k} : \widehat{T}^n(C) \to \widehat{T}^{n+1}(A)$$

where the morphisms ε^{-k} are connected over $S^{-k} \underline{\delta}^{n+k} : S^{-k} T^{n+k} \to S^{-k-1} T^{n+k+1}$.

Proof The proof is based on determining all terms of each morphism of cohomological functors $\Phi_{k,k+1}^{\bullet}: T^{\bullet}\langle k \rangle \to T^{\bullet}\langle k+1 \rangle$ that give rise to the satellite functor construction. We already know that $\Phi_{k,k+1}^{n} = \mathrm{id}_{\mathcal{D}}$ for k < n. To describe $\Phi_{k,k+1}^{k}$, consider a short exact sequence $0 \to M \to P \to C \to 0$ with P projective. We note that $\Phi_{k,k+1}^{\bullet}$ is meant to commute with the connecting homomorphisms of $T^{\bullet}\langle k \rangle$ and $T^{\bullet}\langle k+1 \rangle$, in particular with the respective k^{th} connecting homomorphisms

$$\delta^k \langle k \rangle = \delta^k : T^k(C) \to T^{k+1}(A) \text{ and } \delta^k \langle k+1 \rangle = \varepsilon^{-1} : S^{-1}T^{k+1}(C) \to T^{k+1}(A).$$

Since $(T^{\bullet}, \delta^{\bullet})$ is a cohomological functor, we obtain from Diagram 3.2 the commutative diagram

$$\dots \longrightarrow T^{k}(P) \longrightarrow T^{k}(C) \xrightarrow{\delta^{k}} T^{k+1}(M) \longrightarrow T^{k+1}(P) \longrightarrow \dots$$

$$\downarrow^{T^{k}(h)} \qquad \downarrow^{\operatorname{id}_{T^{k}(C)}} \qquad \downarrow^{T^{k+1}(h^{*})} \qquad \downarrow^{T^{k+1}(h)} \qquad (3.5)$$

$$\dots \longrightarrow T^{k}(B) \longrightarrow T^{k}(C) \xrightarrow{\delta^{k}} T^{k+1}(A) \longrightarrow T^{k+1}(B) \longrightarrow \dots$$

Recalling that $S^{-1}T^{k+1}(C)$ is a kernel and $\varepsilon^{-1}: S^{-1}T^{k+1}(C) \to T^{k+1}(M)$ its canonical monomorphism, we can write the middle square as

By definition, $\delta^k \langle k+1 \rangle = \varepsilon^{-1} : S^{-1}T^{k+1}(C) \xrightarrow{\varepsilon^{-1}} T^{k+1}(M) \xrightarrow{T^{k+1}(h^*)} T^{k+1}(A)$. Tilting the above diagram, we obtain the desired commutative diagram

We conclude that $\underline{\delta}^k = \Phi_{k,k+1}^k : T^k \to S^{-1}T^{k+1}$ is a natural transformation by the proof of Lemma 3.1 found in [7, p. 46–47]. By the construction of left satellite transformations, we infer that $\Phi_{k,k+1}^n = S^{n-k}\underline{\delta}^k : S^{n-k}T^k \to S^{n-k+1}T^{k+1}$ for any n < k. In particular, the square

$$S^{n-k}T^{k}(C) \xrightarrow{\varepsilon^{n-k}} S^{-k+1}T^{n}(A)$$

$$\downarrow^{S^{n-k}\underline{\delta}^{k}} \qquad \qquad \downarrow^{S^{n-k+1}\underline{\delta}^{k}}$$

$$S^{n-k-1}T^{k+1}(C) \xrightarrow{\varepsilon^{n-k-1}} S^{n-k}T^{k+1}(A)$$

$$(3.7)$$

commutes. In the direct limit, these squares give rise to the connecting homomorphism $\hat{\delta}^n$ and the formula in the lemma follows from reindexing as in Equation 3.4.

Lastly, we provide an explicit formula for the induced morphisms in the satellite functor construction.

Proposition 3.4 Let $f : A \to B$ be a morphism in C. Then for any $n \in \mathbb{Z}$ the commuting square

give rise to $\widehat{T}^n(f) = \varinjlim_{\mathcal{D}, k \in \mathbb{N}_0} S^{-k} T^{n+k}(f) : \widehat{T}(A) \to \widehat{T}(B).$

Proof This follows from Equation 3.4 and the description of the cohomological functors $\Phi_{n+k,n+k+1}^{\bullet}$ found in the proof of Lemma 3.3.

4 The resolution and the satellite functor construction

The main goal of this section is to prove that the resolution construction gives rise to Mislin completions of cohomological functors. First, we show in Subsection 4.1 that the cohomology groups resulting from the satellite functor construction are isomorphic to those from the resolution construction. In Subsection 4.2, we provide explicit formulae for the induced morphisms and connecting homomorphisms of the resolution construction. Then, in Subsection 4.3, we prove that the isomorphisms between the cohomology groups extend to isomorphisms of cohomological functors. This sets the ground for Subsection 4.4 where we demonstrate that there are countably many distinct constructions of Mislin completions.

4.1 Isomorphism of cohomology groups

To set up notation for the resolution construction, let $M \in obj(\mathcal{C})$, $(M_n)_{n \in \mathbb{N}_0}$ be a projective resolution of M and \widetilde{M}_k be the k^{th} syzygy of M_{\bullet} for $k \in \mathbb{N}_0$. Keeping our notation from last section, we denote for $n \in \mathbb{Z}$ the n^{th} term of the satellite functor construction by $\widehat{T}^n(M)$. If the connecting homomorphisms $\delta^{n+k} : T^{n+k}(\widetilde{M}_k) \to T^{n+k+1}(\widetilde{M}_{k+1})$ are taken as in Subsection 2, we write the n^{th} term of the resolution construction as $T^n_{Res}(M) := \varinjlim_{k \in \mathbb{N}_0} \left(T^{n+k}(\widetilde{M}_k), \delta^{n+k}\right)$. However it depends on a choice of projective resolution a priori. This issue is resolved by demonstrating

Lemma 4.1 For any $n \in \mathbb{Z}$ there is an isomorphism $\omega_n(M) : \widehat{T}^n(M) \to T^n_{Res}(M)$ in \mathcal{D} .

A proof of this lemma can be extracted from [28, p. 299] for modules over rings and can be found in [19, p. 20] for unenriched relative Ext-functors. In the following, we present an analogous proof for the cohomological functor $T^{\bullet}: \mathcal{C} \to \mathcal{D}$. The manner in which we conduct it allows us in Subsection 4.3 to extend the resulting isomorphism to an isomorphism of cohomological functors.

Proof The proof consists of a two step endeavour where we introduce the relevant notation throughout its course. In the first step, we construct an isomorphism

$$\eta_n(M) : \lim_{k \in \mathbb{N}} (S^{-1}T^{n+k}(\widetilde{M}_{k-1}), \Sigma^{-1}\delta^{n+k}) \to T^n_{Res}(M)$$

and in the second step we extend it to an isomorphism $\omega_n(M) : \widehat{T}^n(M) \to T^n_{Res}(M)$.

Step 1 Defining an isomorphism $\eta_n(M) : \varinjlim_{k \in \mathbb{N}} (S^{-1}T^{n+k}(\widetilde{M}_{k-1}), \Sigma^{-1}\delta^{n+k}) \to T^n_{Res}(M)$ By Axiom 2.2, the short exact sequence $0 \to \widetilde{M}_{k+1} \xrightarrow{\iota_{k+1}} M_k \xrightarrow{\pi_k} \widetilde{M}_k \to 0$ induces a long exact sequence in the cohomological functor $(T^{\bullet}, \delta^{\bullet})$ one of whose parts is

$$T^{n+k}(\widetilde{M}_k) \xrightarrow{\delta^{n+k}} T^{n+k+1}(\widetilde{M}_{k+1}) \xrightarrow{T^{n+k+1}(\iota_{k+1})} T^{n+k+1}(M_k) \xrightarrow{T^{n+k+1}(\pi_k)} T^{n+k+1}(\widetilde{M}_k)$$

Because M_k is projective, it follows from the exactness of the above and from the definition of left satellite functors that

$$\operatorname{Im}(\delta^{n+k}) = \operatorname{Ker}(T^{n+k+1}(\iota_{k+1})) = S^{-1}T^{n+k+1}(\widetilde{M}_k).$$

This allows us to think of $\underline{\delta}^{n+k}$: $T^{n+k}(\widetilde{M}_k) \to S^{-1}T^{n+k+1}(\widetilde{M}_k)$ as being obtained by restricting the codomain of δ^{n+k} onto the subobject $S^{-1}T^{n+k+1}(\widetilde{M}_k)$ of $T^{n+k+1}(\widetilde{M}_{k+1})$. Because $\varepsilon^{-1}: S^{-1}T^{n+k}(\widetilde{M}_{k-1}) \to T^{n+k}(\widetilde{M}_k)$ is the canonical monomorphism from a kernel, we can interpret

$$\Sigma^{-1}\delta^{n+k} := \underline{\delta}^{n+k} \circ \varepsilon^{-1} : \quad S^{-1}T^{n+k}(\widetilde{M}_{k-1}) \to T^{n+k}(\widetilde{M}_k) \to S^{-1}T^{n+k+1}(\widetilde{M}_k)$$

as arising by restricting the domain of δ^{n+k} onto $S^{-1}T^{n+k}(\widetilde{M}_{k-1})$ and its codomain onto $S^{-1}T^{n+k+1}(\widetilde{M}_k)$. Using this and Diagram 3.6 we obtain the commuting diagram

Since the above squares give rise to a direct system of morphisms, we define

$$\eta_n(M) = \varinjlim_{k \in \mathbb{N}} \varepsilon^{-1} : \varinjlim_{k \in \mathbb{N}} (S^{-1}T^{n+k}(\widetilde{M}_{k-1}), \Sigma^{-1}\delta^{n+k}) \to T^n_{Res}(M) .$$

If N_{n+k} is the cokernel of the monomorphism $\varepsilon^{-1} : S^{-1}T^{n+k}(\widetilde{M}_{k-1}) \to T^{n+k}(\widetilde{M}_k)$, then it follows from its universal property that we can extend Diagram 4.1 by a morphism $m_{n+k} : N_{n+k} \to N_{n+k+1}$ in a commutative manner as

$$0 \longrightarrow S^{-1}T^{n+k}(\widetilde{M}_{k-1}) \xrightarrow{\varepsilon^{-1}} T^{n+k}(\widetilde{M}_{k}) \xrightarrow{q_{n+k}} N_{n+k} \longrightarrow 0$$

$$\downarrow^{\Sigma^{-1}\delta^{n+k}} \xrightarrow{\delta^{n+k}} \downarrow^{\delta^{n+k}} \qquad \downarrow^{m_{n+k}} \qquad (4.2)$$

$$0 \longrightarrow S^{-1}T^{n+k+1}(\widetilde{M}_{k}) \xrightarrow{\varepsilon^{-1}} T^{n+k+1}(\widetilde{M}_{k+1}) \xrightarrow{q_{n+k+1}} N_{n+k+1} \longrightarrow 0$$

However, note that

$$m_{n+k} \circ q_{n+k} = q_{n+k+1} \circ \varepsilon^{-1} \circ \underline{\delta}^{n+k} : T^{n+k}(\widetilde{M}_k) \to N_{n+k+1}$$

is the zero morphism. Since q_{n+k} is surjective, we deduce that $m_{n+k} = 0$. Given that these diagrams form a direct system of short exact sequences and countable direct limits in \mathcal{D} are exact, we obtain a short exact sequence

$$0 \to \varinjlim_{k \in \mathbb{N}} (S^{-1}T^{n+k}(\widetilde{M}_{n+k}), \Sigma^{-1}\delta^{n+k}) \xrightarrow{\eta_n(M)} T^n_{Res}(M) \to 0 \to 0.$$

Hence, $\eta_n(M)$ is an isomorphism.

Step 2 Defining an isomorphism $\omega_n(M) : \widehat{T}^n(M) \to T^n_{Res}(M)$

The way we construct this isomorphism is by connecting the terms of the direct system $(S^{-k}T^{n+k}(M), S^{-k}\underline{\delta}^{n+k})_{k\in\mathbb{N}_0}$ to the ones of $(T^{n+k}(\widetilde{M}_k), \delta^{n+k})_{k\in\mathbb{N}_0}$. This requires a larger commutative diagram for which we introduce all the necessary morphisms. By the previous step, we can extend Diagram 3.7 to the commutative diagram

$$S^{-1}T^{n+k+2}(\widetilde{M}_{k+1}) \xrightarrow{\varepsilon^{-1}} T^{n+k+2}(\widetilde{M}_{k+2})$$

$$\downarrow_{S^{-1}\underline{\delta}^{n+k+2}} \qquad \qquad \downarrow_{\underline{\delta}^{n+k+2}} \qquad \qquad \downarrow_{\underline{\delta}^{n+k+2}}$$

$$S^{-2}T^{n+k+3}(\widetilde{M}_{k+1}) \xrightarrow{\varepsilon^{-2}} S^{-1}T^{n+k+3}(\widetilde{M}_{k+2})$$

Since ε^{-2} : $S^{-2}T^{n+k+3}(\widetilde{M}_{k+1}) \to S^{-1}T^{n+k+3}(\widetilde{M}_{k+2})$ is the monomorphism of a kernel, we can think of $S^{-1}\underline{\delta}^{n+k+2}$: $S^{-1}T^{n+k+2}(\widetilde{M}_{k+1}) \to S^{-2}T^{n+k+3}(\widetilde{M}_{k+1})$ as being obtained by restricting the codomain of $\Sigma^{-1}\delta^{n+k+2}$ onto the subobject $S^{-2}T^{n+k+3}(\widetilde{M}_{k+1})$ of $S^{-1}T^{n+k+3}(\widetilde{M}_{k+2})$. Furthermore, we can interpret the morphism

$$\Sigma^{-2}\delta^{n+k+2} := S^{-1}\underline{\delta}^{n+k+2} \circ \varepsilon^{-2} :$$

$$S^{-2}T^{n+k+2}(\widetilde{M}_k) \to S^{-1}T^{n+k+2}(\widetilde{M}_{k+1}) \to S^{-2}T^{n+k+3}(\widetilde{M}_{k+1})$$

as arising from restricting the domain of $\Sigma^{-1}\delta^{n+k+2}$ onto $S^{-2}T^{n+k+2}(\widetilde{M}_k)$ and its codomain onto $S^{-2}T^{n+k+3}(\widetilde{M}_{k+1})$. As for left satellite functors, we may define $\Sigma^0\delta^{n+k} := \delta^{n+k}$ and

$$\Sigma^{-l}\delta^{n+k+l} := S^{-l+1}\underline{\delta}^{n+k+l} \circ \varepsilon^{-l} :$$

$$S^{-l}T^{n+k+l}(\widetilde{M}_k) \to S^{-l+1}T^{n+k+l}(\widetilde{M}_{k+1}) \to S^{-l}T^{n+k+l+1}(\widetilde{M}_{k+1})$$
(4.3)

for any $l \in \mathbb{N}$. We can think of $\Sigma^{-l} \delta^{n+k+l}$ as being obtained by restricting the domain of $\Sigma^{-l+1} \delta^{n+k+l-1}$ onto $S^{-l}T^{n+k+l}(\widetilde{M}_k)$ and its codomain onto $S^{-l}T^{n+k+l+1}(\widetilde{M}_{k+1})$. We can again extend Diagram 3.7 to another commutative diagram of the form

$$S^{-l}T^{n+k+l}(\widetilde{M}_{k}) \xrightarrow{\varepsilon^{-l}} S^{-l+1}T^{n+k+l}(\widetilde{M}_{k+1})$$

$$\downarrow_{S^{-l}\underline{\delta}^{n+k+l}} \qquad \downarrow_{S^{-l+1}\underline{\delta}^{n+k+l}} \qquad (4.4)$$

$$S^{-l-1}T^{n+k+l+1}(\widetilde{M}_{k}) \xrightarrow{\varepsilon^{-l-1}} S^{-l}T^{n+k+l+1}(\widetilde{M}_{k+1})$$

By [7, Section III.2], for every $2 \le l \in \mathbb{N}$ and $k \in \{0, \ldots, l-1\}$ the short exact sequence $0 \to \widetilde{M}_{k+1} \to M_k \to \widetilde{M}_k \to 0$ gives rise to a long exact sequence one of whose parts is


$$S^{-l}T^{n+k+l}(M_k) \to S^{-l}T^{n+k+l}(\widetilde{M}_k) \xrightarrow{\varepsilon^{-l}} S^{-l+1}T^{n+k+l}(\widetilde{M}_{k+1}) \to S^{-l+1}T^{n+k+l}(M_k).$$

By construction, left satellite functors vanish on projective objects such as M_k . Hence the morphism $\varepsilon^{-l}: S^{-l}T^{n+k+l}(\widetilde{M}_k) \to S^{-l+1}T^{n+k+l}(\widetilde{M}_{k+1})$ is an isomorphism. Define the isomorphisms

$$\widetilde{o}^{k} := \varepsilon^{-3} \circ \dots \circ \varepsilon^{-k+1} \circ \varepsilon^{-k} : \quad S^{-k} T^{n+k}(M) \to S^{-2} T^{n+k}(\widetilde{M}_{k-2})$$


and
$$o^{k} := \varepsilon^{-2} \circ \widetilde{o}^{k} : \quad S^{-k} T^{n+k}(M) \to S^{-1} T^{n+k}(\widetilde{M}_{k-1}).$$
(4.5)

We can form the following diagram with all morphisms we have introduced thus far:



We need to argue why this diagram commutes before we can demonstrate how it yields the desired isomorphism. The triangles at the very right side containing the morphisms

 $\delta^{n+k}: T^{n+k}(\widetilde{M}_k) \to T^{n+k+1}(\widetilde{M}_{k+1})$ commute by Diagram 3.6. All other triangles commute due to Diagram 4.4. Finally, due to Diagram 3.7 and 4.4 the following diagram commutes for any $k \geq 3$.



We conclude from it that $\Sigma^{-2}\delta^{n+k} \circ \tilde{o}^k = \tilde{o}^{k-1} \circ S^{-k} \underline{\delta}^{n+k}$. In particular, Diagram 4.6 commutes as all its stand-alone squares also do so.

We see in Diagram 4.6 that the right most column corresponds to the direct system $(T^{n+k}(\widetilde{M}_k), \delta^{n+k})_{k \in \mathbb{N}_0}$ while the top diagonal together with the left most column forms the direct system $(S^{-k}T^{n+k}(M), S^{-k}\underline{\delta}^{n+k})_{k \in \mathbb{N}_0}$. The right most squares in the diagram recover the isomorphism

$$\eta_n(M) = \varinjlim_{k \in \mathbb{N}} \varepsilon^{-1} : \varinjlim_{k \in \mathbb{N}} (S^{-1}T^{n+k}(\widetilde{M}_{k-1}), \Sigma^{-1}\delta^{n+k}) \to T^n_{Res}(M)$$

For $k \geq 3$, all the remaining squares yield together squares of the form

$$S^{-k}T^{n+k}(M) \xrightarrow{o^{k}} S^{-1}T^{n+k}(\widetilde{M}_{k-1})$$
$$\downarrow^{S^{-k}\underline{\delta}^{n+k}} \qquad \qquad \downarrow^{\Sigma^{-1}\delta^{n+k}}$$
$$S^{-k-1}T^{n+k+1}(M) \xrightarrow{o^{k+1}} S^{-1}T^{n+k+1}(\widetilde{M}_{k})$$

As the morphisms o^k are isomorphisms and countable direct limits in \mathcal{D} are exact, this gives rise to the isomorphism

$$\varinjlim o^k : \widehat{T}^n(M) \to \varinjlim_{k \in \mathbb{N}} (S^{-1}T^{n+k}(\widetilde{M}_{k-1}), \Sigma^{-1}\delta^{n+k}) \,.$$

Due to the universal property of direct limits, we observe that $\varinjlim o^k \circ \varinjlim \varepsilon^{-1} = \varinjlim o^k \circ \varepsilon^{-1}$. Because $\eta_n(M) = \varinjlim \varepsilon^{-1}$ is already an isomorphism,

$$\omega_n(M) = \varinjlim o^k \circ \varepsilon^{-1} : \widehat{T}^n(M) \to T^n_{Res}(M)$$

is our desired isomorphism. Since the morphisms $o^k \circ \varepsilon^{-1}$ span the entire width of Diagram 4.6, every single morphism was needed to construct this isomorphism.

4.2 Induced morphisms and connecting homomorphisms

The first step to turn $\omega_n(M) : \widehat{T}^n(M) \to T_{Res}^n(M)$ into an isomorphism of cohomological functor is to show that it is natural. Let $f: M \to N$ be a morphism in \mathcal{C} and M_{\bullet} , N_{\bullet} be projective resolutions of M, N. Set $\widetilde{f}_0 := f: \widetilde{M}_0 \to \widetilde{N}_0$ and assume that $\widetilde{f}_{k+1} : \widetilde{M}_k \to \widetilde{N}_k$ has been defined for $k \geq 0$. Let us denote by $\pi_k : M_k \to \widetilde{M}_k$ the epimorphism from the projective M_k onto \widetilde{M}_k and by $\iota_{k+1} : \widetilde{M}_{k+1} \to M_k$ the monomorphism from the $(k+1)^{\text{st}}$ syzygy \widetilde{M}_{k+1} into M_k . We adopt an analogous convention for \widetilde{N}_{k+1} , N_k and \widetilde{N}_k . Then we can lift \widetilde{f}_k to fit in the commutative diagram

since M_0 is a projective object and $\iota_{k+1} : \widetilde{N}_{k+1} \to N_k$ can be taken to be the kernel of π_k .

Proposition 4.2 Let $f: M \to N$ and $(\widetilde{f}_k: \widetilde{M}_k \to \widetilde{N}_k)_{k \in \mathbb{N}_0}$ be a sequence of morphism as arising from Diagram 4.8. As $(T^{\bullet}, \delta^{\bullet})$ is a cohomological functor, we denote the morphism in the direct limit of the commuting squares

by $T_{Res}^n(f) := \varinjlim_{k \in \mathbb{N}_0} T^{n+k}(\widetilde{f}_k)$. Then the square


$$\widehat{T}^{n}(M) \xrightarrow{\omega_{n}(M)} T^{n}_{Res}(M)$$

$$\downarrow^{\widehat{T}^{n}(f)} \qquad \downarrow^{T^{n}_{Res}(f)}$$


$$\widehat{T}^{n}(N) \xrightarrow{\omega_{n}(N)} T^{n}_{Res}(N)$$

commutes. In particular, every term $T_{Res}^n(-)$ forms an additive functor and the homomorphisms $\omega_n : \widehat{T}^n \to T_{Res}^n$ a natural isomorphism.

Proof Recall that $\widehat{T}^n(f) = \varinjlim_{k \in \mathbb{N}_0} S^k T^{n+k}(f)$ by Proposition 3.4. Let $k \in \mathbb{N}_0, l \in \mathbb{N}$ and write $\Lambda = n + k + l$. Consider the diagram



The front and back side commute by Diagram 4.4. The left hand and right hand side commute since $S^{-l}\underline{\delta}^{\Lambda}$ and $S^{-l+1}\underline{\delta}^{\Lambda}$ are natural transformations by Lemma 3.3. Lastly, the top and bottom side commute by Equation 3.3. Remember that we have defined the morphisms $o^k = \varepsilon^{-2} \circ \varepsilon^{-3} \circ \ldots \circ \varepsilon^{-k}$ in the proof of Lemma 4.1. Because δ^{n+k} is also a natural transformation, we conclude from Diagram 4.6 and the above cube that the diagram



is commutative. Referring again to the proof of Lemma 4.1, we recall that we have defined $\omega_n = \varinjlim_{k \in \mathbb{N}} \varepsilon^{-1} \circ o^k : \widehat{T}^n \to T_{Res}^n$. Therefore, the bottom square of the proposition commutes and $T_{Res}^n(f)$ is well defined in the sense that it does not depend on the choice of sequence $(\widetilde{f}_k : \widetilde{M}_k \to \widetilde{N}_k)_{k \in \mathbb{N}_0}$. Because every term \widehat{T}^n is an additive functor, any T_{Res}^n is so too. Or more directly, for any $g : M \to N$ and any correspondign sequence $(\widetilde{g}_k : \widetilde{M}_k \to \widetilde{N}_k)_{k \in \mathbb{N}_0}$ we see that

$$T_{Res}^n(f+g) = \varinjlim_{k \in \mathbb{N}_0} T^{n+k}(\widetilde{f}_k + \widetilde{g}_k) = \varinjlim_{k \in \mathbb{N}_0} \left(T^{n+k}(\widetilde{f}_k) + T^{n+k}(\widetilde{g}_k) \right) = T_{Res}^n(f) + T_{Res}^n(g) .$$

In order to turn T_{Res}^{\bullet} into a cohomological functor, we need to construct connecting homomorphisms. For this let $0 \to \widetilde{A}_k \to \widetilde{B}_k \to \widetilde{C}_k \to 0$ be a short exact sequence in \mathcal{C} . By the Horseshoe Lemma [42, p. 37], there is a commutative diagram

where A_k , B_k and C_k are projective and all rows and columns are exact. Then one can generalise Proposition III.4.1 of H. Cartan and S. Eilenberg's book [7] from modules over a ring to abelian categories either by turning all diagram chases into diagrams involving the respective kernels, cokernels and morphism or by invoking the Freyd-Mitchell Embedding Theorem that can be found in [42, p. 25]. This says that the square

anticommutes for any $n \in \mathbb{Z}$ and $k \in \mathbb{N}_0$, meaning that the composite morphisms in the abelian group $\operatorname{Hom}_{\mathcal{D}}(T^{n+k}(\widetilde{C}_k), T^{n+k+2}(\widetilde{A}_{k+1}))$ have opposite signs.

Definition 4.3 For a short exact sequence $0 \to A \to B \to C \to 0$ in C we define the (proposed) connecting homomorphism of T^{\bullet}_{Res} to be

$$\widetilde{\delta}^n := \lim_{k \in \mathbb{N}_0} (-1)^k \delta^{n+k} : T^n_{Res}(C) \to T^{n+1}_{Res}(A)$$

where the morphisms $(-1)^k \delta^{n+k}$ are connected over the morphisms δ^{n+k} and δ^{n+k+1} from Diagram 4.10.

4.3 Isomorphisms of cohomological functors


The key step in establishing that $(T^{\bullet}_{Res}, \tilde{\delta}^{\bullet})$ is a cohomological functor and proving that $\omega_{\bullet}: \hat{T}^{\bullet} \to T^{\bullet}_{Res}$ extends to an isomorphism of cohomological functors is

Lemma 4.4 Denote by $\hat{\delta}^{\bullet}$ the connecting homomorphism from the satellite functor construction \hat{T}^{\bullet} . Then for any short exact sequence $0 \to A \to B \to C \to 0$ in C and any $n \in \mathbb{Z}$ the diagram

$$\widehat{T}^{n}(C) \xrightarrow{\omega_{n}} T^{n}_{Res}(C) \downarrow_{\widehat{\delta}^{n}} \qquad \qquad \downarrow_{\widetilde{\delta}^{n}} \\ \widehat{T}^{n+1}(A) \xrightarrow{\omega_{n+1}} T^{n+1}_{Res}(A)$$

commutes.

Proof The proof is based on showing that the diagram on the next page is commutative. The front side is a copy of Diagram 4.6 as well as the back side up to its commuting right most column. The right hand side of the diagram gives rise to $\delta^n : T_{Res}^n(C) \to T_{Res}^{n+1}(A)$ in the direct limit and the top and left hand side to $\delta^n : \hat{T}^n(C) \to \hat{T}^{n+1}(A)$. In particular, all squares and triangles from these sides commute. Using the cohomological functors defined in Equation 3.3 we see that the horizontal squares commute by [7, Proposition III.4.1].




To investigate the vertical squares running from back to front, we can insert commuting triangles of the form

$$S^{-l}T^{n+k}(\widetilde{M}_{k-l})$$

$$S^{-l}\underline{\delta}^{n+k} \downarrow^{\Sigma^{-l}\delta^{n+k}}$$

$$S^{-l-1}T^{n+k+1}(\widetilde{M}_{k-l}) \xrightarrow{\varepsilon^{-l-1}} S^{-l}T^{n+k+1}(\widetilde{M}_{k-l+1})$$

into the front and back side by Diagram 4.6. These give rise to prisms of the form



where their right hand square correspond to the vertical squares in Diagram 4.11. These squares commute because the bottom square of the prism does so and the top left square by Diagram 3.7. In particular, Diagram 4.11 commutes.

Denote by $\chi = \varinjlim \delta^{n+k+1} : T^{n+1}_{Res}(A) \to T^{n+1}_{Res}(A)$ the morphism resulting from the right most column of the back side of Diagram 4.11 and write $\Psi^k_{n+1} : T^{n+k+1}(\widetilde{A}_k) \to T^{n+1}_{Res}(A)$ for the morphism to the direct limit. Since every square in this column can be written as

we deduce that the diagram

$$T^{n+k+2}(\widetilde{A}_{k+1}) \xrightarrow{\Psi_{n+1}^{k+1}} T^{n+1}_{Res}(A)$$

$$\downarrow^{\text{id}} \qquad \qquad \downarrow_{\delta^{n+k+2}} \Psi_{n+1}^{k+2} \xrightarrow{\Psi_{n+1}^{k+2}} T^{n+k+3}(\widetilde{A}_{k+2}) \xrightarrow{\Phi^{n+k+2}} T^{n+k+3}(\widetilde{A}_{k+2})$$

$$(4.13)$$

commutes. Observing that $(T^{n+k+1}(\widetilde{A}_k), \delta^{n+k+1})_{k \in \mathbb{N}}$ is cofinal in $(T^{n+k+1}(\widetilde{A}_k), \delta^{n+k+1})_{k \in \mathbb{N}_0}$, we conclude from the previous two diagrams that $\mathrm{id} = \chi : T^{n+1}_{Res}(A) \to T^{n+1}_{Res}(A)$.

Let us consider the top and left most edge in the front side of Diagram 4.11 yielding $\widehat{T}^n(C)$. Going over the front side yields the morphism $\omega_n : \widehat{T}^n(C) \to T^n_{Res}(C)$. Through the right hand side we obtain $\widetilde{\delta}^n : T^n_{Res}(C) \to T^{n+1}_{Res}(A)$. Thus, going around these sides yields $\widetilde{\delta}^n \circ \omega_n$. Instead, if we go over the left hand side first, we obtain the connecting homomorphism $\widehat{\delta}^n : \widehat{T}^n(C) \to \widehat{T}^{n+1}(A)$. Going subsequently over the back side yields

first $\omega_{n+1}: \widehat{T}^{n+1}(A) \to T_{Res}^{n+1}(A)$ and then $\chi: T_{Res}^{n+1}(A) \to T_{Res}^{n+1}(A)$. Hence, going around those sides, we obtain $\chi \circ \omega_{n+1} \circ \widehat{\delta}^n$. Because Diagram 4.11 commutes, we conclude that $\widetilde{\delta}^n \circ \omega_n = \chi \circ \omega_{n+1} \circ \widehat{\delta}^n$.

Theorem 4.5 The resolution construction $(T^{\bullet}_{Res}, \widetilde{\delta}^{\bullet})$ forms a cohomological functor and $\omega_{\bullet} : (\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet}) \to (T^{\bullet}_{Res}, \widetilde{\delta}^{\bullet})$ is an isomorphism of cohomological functors.

Proof First, we show that the proposed connecting homomorphisms $\tilde{\delta}^{\bullet}$ are natural. For this let $n \in \mathbb{Z}$ and let

be a commutative diagram in \mathcal{C} with exact rows.Consider the diagram

By Proposition 4.2, the isomorphism ω_n (resp. ω_{n+1}) is natural and thus the front and back squares of the cube commute. According to Lemma 4.4, the top and bottom squares of the cube commute. As $(\hat{T}^{\bullet}, \hat{\delta}^{\bullet})$ is a cohomological functor, the left hand side commutes. Due to this and the fact that $\omega_n : \hat{T}^n(A) \to T^n_{Res}(A)$ is an isomorphism, we infer that also the right hand side commutes. Hence $(T^{\bullet}_{Res}, \tilde{\delta}^{\bullet})$ satisfies Axiom 2.1 of a cohomological functor. Using this together with the fact $\omega_n : \hat{T}^n \to T^n_{Res}$ is a natural transformation for every $n \in \mathbb{Z}$, we see that the diagram

commutes. Using again that every $\omega_n : \widehat{T}^n \to T_{Res}^n$ is an isomorphism, we infer by the Five Lemma [7, Proposition I.1.1] that every image and kernel in the bottom row is isomorphic to the corresponding image or kernel in the top row. Because $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ is a cohomological functor, the bottom row forms a long exact sequence and thus the top row does so too. In particular, $(T_{Res}^{\bullet}, \widetilde{\delta}^{\bullet})$ satisfies Axiom 2.2 and is a cohomological functor. From this we conclude that $\omega_{\bullet} : \widehat{T}^{\bullet} \to T_{Res}^{\bullet}$ is an isomorphism of cohomological functors.

4.4 Countably many distinct constructions of Mislin completions

Based on the material of this section thus far, we can demonstrate that there are countably many distinct constructions of Mislin completions.

Lemma 4.6 Let $a = (a_k)_{k \in \mathbb{N}} \in \{0,1\}^{\mathbb{N}}$ be a sequence taking values in 0 and 1. Define $(P(a)_k)_{k \in \mathbb{N}_0}$ by $P(a)_0 := 0$ and $P(a)_k := \sum_{i=1}^k a_i$ and $(D(a)_k)_{k \in \mathbb{N}_0}$ by $D(a)_0 := 0$ and $D(a)_k := \sum_{i=1}^k (1-a_i)$. For any $M \in obj(\mathcal{C})$ and $n \in \mathbb{Z}$ form the direct system $(S^{-P(a)_k}T^{n+k}(\widetilde{M}_{D(a)_k}), \delta_a^{n+k})_{k \in \mathbb{N}_0}$ where

$$\delta_{a}^{n+k} := \begin{cases} \Sigma^{-P(a)_{k}} \delta^{n+k} : S^{-P(a)_{k}} T^{n+k}(\widetilde{M}_{D(a)_{k}}) \to S^{-P(a)_{k}} T^{n+k+1}(\widetilde{M}_{D(a)_{k}+1}) & \text{if } P(a)_{k+1} = P(a)_{k} \\ S^{-P(a)_{k}} \underline{\delta}^{n+k} : S^{-P(a)_{k}} T^{n+k}(\widetilde{M}_{D(a)_{k}}) \to S^{-P(a)_{k}-1} T^{n+k+1}(\widetilde{M}_{D(a)_{k}}) & \text{if } P(a)_{k+1} = P(a)_{k} + 1 \end{cases}$$

and the morphisms $\Sigma^{-P(a)_k} \delta^{n+k}$ are taken as in Equation 4.3. Write

$$T_a^n(M) := \lim_{\mathcal{D}, k \in \mathbb{N}_0} S^{-P(a)_k} T^{n+k}(\widetilde{M}_{D(a)_k}) \,.$$

Then there exists an isomorphism $\omega_{a,n}: T_a^n(M) \to T_{Res}^n(M)$.

Proof For any $k \in \mathbb{N}_0$ we define the morphism

$$\widehat{o}_a^k := \begin{cases} \varepsilon^{-1} \circ \dots \circ \varepsilon^{-P(a)_k} : S^{-P(a)_k} T^{n+k}(\widetilde{M}_{D(a)_k}) \to T^{n+k}(\widetilde{M}_k) & \text{if } P(a)_k \ge 1\\ \text{id} : T^{n+k}(\widetilde{M}_k) \to T^{n+k}(\widetilde{M}_k) & \text{if } P(a)_k = 0 \end{cases}$$

similarly to the morphism o^k from Equation 4.5. By Diagram 4.6 we infer that the square

commutes. We conclude as in the proof of Lemma 4.1 that $\omega_{a,n} := \lim_{k \in \mathbb{N}_0} \widehat{o}_a^k$ is an isomorphism.

We can reiterate the proof of Proposition 4.2 to demonstrate

Proposition 4.7 Let $f : M \to N$ be a morphism in \mathcal{C} and $(\widetilde{f}_k : \widetilde{M}_k \to \widetilde{N}_k)_{k \in \mathbb{N}_0}$ be a sequence of morphism as in Diagram 4.8. Then the squares

$$S^{-P(a)_{k}}T^{n+k}(\widetilde{M}_{D(a)_{k}}) \xrightarrow{S^{-P(a)_{k}}T^{n+k}(f_{D(a)_{k}})} S^{-P(a)_{k}}T^{n+k}(\widetilde{N}_{D(a)_{k}}) \xrightarrow{\delta_{a}^{n+k}} \delta_{a}^{n+k}$$

$$S^{-P(a)_{k+1}}T^{n+k+1}(\widetilde{M}_{D(a)_{k+1}}) \xrightarrow{S^{-P(a)_{k+1}}T^{n+k+1}(\widetilde{f}_{D(a)_{k+1}})} S^{P(a)_{k+1}}T^{n+k+1}(\widetilde{N}_{D(a)_{k+1}})$$

commute. If we define $T_a^n(f) := \varinjlim_{k \in \mathbb{N}_0} S^{-P(a)_k} T^{n+k}(\widetilde{f}_{D(a)_k})$, then the square

$$\begin{array}{ccc} T_a^n(M) & \xrightarrow{\omega_{a,n}(M)} & T_{Res}^n(M) \\ & & \downarrow^{T_a^n(f)} & & \downarrow^{T_{Res}^n(f)} \\ & T_a^n(N) & \xrightarrow{\omega_{a,n}(N)} & T_{Res}^n(N) \end{array}$$

is also commutative. In particular, every term $T_a^n(-)$ forms an additive functor and the homomorphisms $\omega_{a,n}: \widehat{T}^n \to T_{Res}^n$ a natural isomorphism.

Lemma 4.8 If $0 \to A \to B \to C \to 0$ is a short exact sequence, define for any $k \in \mathbb{N}_0$ the morphism

$$\delta_{a,k}^{n+k} := \varepsilon^{-P(a)_k - 1} \circ S^{-P(a)_k} \underline{\delta}^{n+k} :$$

$$S^{-P(a)_k} T^{n+k} (\widetilde{C}_{D(a)_k}) \to S^{-P(a)_k - 1} T^{n+k+1} (\widetilde{C}_{D(a)_k}) \to S^{-P(a)_k} T^{(n+1)+k} (\widetilde{A}_{D(a)_k})$$

Then the squares

commute whence we define $\widetilde{\delta}_a^n := \varinjlim_{k \in \mathbb{N}_0} \delta_{a,k}^{n+k} : T_a^n(C) \to T_a^{n+1}(A)$. Therefore, the square

is also commutative, $(T_a^{\bullet}, \widetilde{\delta}_a^{\bullet})$ forms a cohomological functor and $\omega_{a,\bullet} : (T_a^{\bullet}, \widetilde{\delta}_a^{\bullet}) \to (T_{Res}^{\bullet}, \widetilde{\delta}^{\bullet})$ is an isomorphism of cohomological functors.

Proof Regarding the Diagram 4.14, the term $S^{-P(a)_k}T^{n+k}(\widetilde{C}_{D(a)_k})$ lies in the front side of Diagram 4.11. Through the morphism $\widehat{\sigma}_a^k : S^{-P(a)_k}T^{n+k}(\widetilde{C}_{D(a)_k}) \to T^{n+k}(\widetilde{C}_k)$ we reach the very right hand side of the latter diagram. The morphism $\delta^{n+k} : T^{n+k}(\widetilde{C}_k) \to T^{(n+1)+k}(\widetilde{A}_k)$ leads us to the back side and we move by id $: T^{(n+1)+k}(\widetilde{A}_k) \to T^{(n+1)+k}(\widetilde{A}_k)$ from Diagram 4.12 across a square in the very right hand column of the back side. On the other had, the morphism $\delta^{n+k}_{a,k} : S^{-P(a)_k}T^{n+k}(\widetilde{C}_{D(a)_k}) \to S^{-P(a)_k}T^{(n+1)+k}(\widetilde{A}_{D(a)_k})$ takes us first down the front side of Diagram 4.11 and then over to the back side. Lastly, we reach the left hand side of the right most column of the back side by the morphism $\widehat{\sigma}_a^{k+1} : S^{-P(a)_k}T^{(n+1)+k}(\widetilde{A}_{D(a)_k}) \to T^{(n+1)+k}(\widetilde{A}_k)$. Since Diagram 4.11 commutes, we conclude that

$$\operatorname{id} \circ \delta^{n+k} \circ \widehat{o}_a^k = \widehat{o}_a^{k+1} \circ \delta_{a,k}^{n+k} : S^{-P(a)_k} T^{n+k} (\widetilde{C}_{D(a)_k}) \to S^{-P(a)_k} T^{(n+1)+k} (\widetilde{A}_{E_k}) .$$
(4.16)

Therefore, Diagram 4.14 commutes. Using again that Diagram 4.11 is commutative, we observe that the squares of Diagram 4.14 form a direct system in whose direct limit we obtain Diagram 4.15. The remainder of the lemma is proved identically to Theorem $4.5.\square$

Corollary 4.9 Let us denote by $e \in \{0,1\}^{\mathbb{N}}$ the sequence all whose terms are set to 0 and by $f \in \{0,1\}^{\mathbb{N}}$ the sequence all whose terms are set to 1. Then $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet}) = (T_f^{\bullet}, \widetilde{\delta}_f^{\bullet})$ and $(T_{Res}^{\bullet}, \widetilde{\delta}^{\bullet}) = (T_e^{\bullet}, \widetilde{\delta}_e^{\bullet}).$

Proof By definition, $\widehat{T}^n = T_f^n$ and $T_{Res}^n = T_e^n$ for every $n \in \mathbb{Z}$. In the same manner as we have concluded by Diagram 4.12 and 4.13 that the morphism χ in the proof of Lemma 4.4 agrees with the identity, we see that

$$\widetilde{\delta}_{f}^{n} = \lim_{k \in \mathbb{N}_{0}} \delta_{f,k}^{n+k} = \lim_{k \in \mathbb{N}} \varepsilon^{-k-1} \circ S^{-k} \underline{\delta}^{n+k} = \lim_{k \in \mathbb{N}} \varepsilon^{-k} = \widehat{\delta}^{n}$$

where $\varepsilon^{-k-1} \circ S^{-k} \underline{\delta}^{n+k} : S^{-k} T^{n+k} \to S^{-k} T^{(n+1)+k}$ and $\varepsilon^{-k} : S^{-k} T^{n+k} \to S^{-k+1} T^{n+k}$. By Equation 4.16, we observe that $\widetilde{\delta}_e^n = \widetilde{\delta}^n$.

Lemma 4.10 For $a, b \in \{0, 1\}^{\mathbb{N}}$ we say that the constructions of the Mislin completions $(T_a^{\bullet}, \widetilde{\delta}_a^{\bullet})$ and $(T_b^{\bullet}, \widetilde{\delta}_b^{\bullet})$ agree if there is an infinite (cofinal) subset $\Lambda \subseteq \mathbb{N}$ such that $S^{-P(a)_{\lambda}}T^{n+\lambda}(\widetilde{M}_{D(a)_{\lambda}}) = S^{-P(b)_{\lambda}}T^{n+\lambda}(\widetilde{M}_{D(b)_{\lambda}})$ for every $\lambda \in \Lambda$, $n \in \mathbb{Z}$ and $M \in obj(\mathcal{C})$. The basic idea is that the constructions agree whenever their terms

$$T_a^n(M) = \lim_{k \in \mathbb{N}_0} S^{-P(a)_k} T^{n+k}(\widetilde{M}_{D(a)_k}) \text{ and } T_b^n(M) = \lim_{k \in \mathbb{N}_0} S^{-P(b)_k} T^{n+k}(\widetilde{M}_{D(b)_k})$$

are isomorphic in a very straightforward manner. If this is not the case, we say that the constructions of $(T_a^{\bullet}, \widetilde{\delta}_a^{\bullet})$ and $(T_b^{\bullet}, \widetilde{\delta}_b^{\bullet})$ are distinct. Then there are countably many distinct constructions of Mislin completions of the form $(T_a^{\bullet}, \widetilde{\delta}_a^{\bullet})$.

Proof Define the binary relation '~' on $\{0,1\}^{\mathbb{N}}$ by setting $a \sim b$ if for every $k \in \mathbb{N}$ there is $K \geq k$ such that $P(a)_K = P(b)_K$. If \sim' denotes the transitive closure of \sim , then the proof reduces to showing that there are countably many elements in $\{0,1\}^{\mathbb{N}}/\sim'$. Hence, define for any $m \in \mathbb{N}$ the sequences $e^m, f^m \in \{0,1\}^{\mathbb{N}}$ by

$$e_k^m := \begin{cases} 1 & \text{if } k < m \\ 0 & \text{if } k \ge m \end{cases} \quad \text{and} \quad f_k^m := \begin{cases} 0 & \text{if } k < m \\ 1 & \text{if } k \ge m \end{cases}$$

Comparing with Corollary 4.9, we see that $e = e_0$ and $f = f_0$. For any $g \in \{0, 1\}^{\mathbb{N}}$ taking finitely many times either the value 0 or the value 1 there is $m \in \mathbb{N}$ such that $g \sim e_m$ or $g \sim f_m$. For any distinct $m, p \in \mathbb{N}$ we note that $e_m \not\sim' e_p$, $e_m \not\sim' f_p$ and $f_m \not\sim' f_p$. To complete the proof, we demonstrate that for any $a, b \in \{0, 1\}^{\mathbb{N}}$ taking both the values 0 and 1 infinitely often there is another sequence c such that $a \sim c$ and $c \sim b$. If $a \not\sim b$, then assume that there is $K \in \mathbb{N}$ such that $P(a)_k < P(b)_k$ for every $k \ge K$. We construct c inductively as follows. For the base case, set $c_k := a_k$ for any $1 \le k \le K$ and let $c_l := 1$ for the subsequent $K \le l$. Since P(c) grows faster than P(b), there is $l_1 \in \mathbb{N}$ such that $P(c)_{l_1} = P(b)_{l_1}$. Assume for $i \in \mathbb{N}$ that we have found $k_i < l_i \in \mathbb{N}$ such that $P(a)_{k_i} = P(c)_{k_i}$ and $P(c)_{l_i} = P(b)_{l_i}$. Setting $c_k := 0$ for $k \ge l_i$ yields $k_{i+1} \in \mathbb{N}$ and defining $c_l := 1$ for $l \ge k_{i+1}$ yields $l_{i+1} \in \mathbb{N}$. Then $a \sim c \sim b$ as desired.

Notation 4.11 In the rest of the paper, we write $(\widehat{T}^{\bullet}, \widehat{\delta}^{\bullet})$ for the Mislin completion of a cohomological functor $(T^{\bullet}, \delta^{\bullet})$ independently from which construction it arises. When required, we specify the construction or use a special notation for it all of which we keep consistent throughout.

5 The naïve and the resolution construction

In this section we prove that the naïve construction gives rise to Mislin completions of unenriched Ext-functors by defining an isomorphism of cohomological functors to the respective terms of the resolution construction. For this, we rigorously perform the naïve construction in Subsection 5.1 in a different manner than in [4] and [19]. Thereafter, we show in Subsection 5.2 that its cohomology groups are naturally isomorphic to the ones of the resolution construction. Lastly, in Subsection 5.3, we construct a connecting homomorphism for the naïve construction and extend the isomorphism between the cohomology groups to an isomorphism of cohomological functors.

5.1 Performing the naïve construction

To start, we consider the unenriched Hom-functors

F

$$\operatorname{Hom}_{\mathcal{C}}(-,-): \mathcal{C}^{\operatorname{op}} \times \mathcal{C} \to \operatorname{Ab}$$

which are additive in both variables. As in Definition 2.5, we turn the corresponding unenriched Ext-functors into cohomological functors by setting $\operatorname{Ext}_{\mathcal{C}}^{n}(A, -) = 0$ for n < 0and any $A \in \operatorname{obj}(\mathcal{C})$. Both A. Beligiannis and I. Reiten in [4, p. 37–38/163–165] as well as S. Guo and L. Liang in [19, p. 14–15] have established relative homological versions of the naïve construction in great generality. However, in order to construct Yoneda products in [16, Theorem 6.6], we require an (absolute homological) generalisation faithful to D. J. Benson and J. F. Carlson's original construction found in [5, p. 109], which is why our account differs from the previous two. First, we define a bifunctor $[-, -]_{\mathcal{C}}$ that we have already encountered in Subsection 2. Let $\mathcal{P}_{\mathcal{C}}(A, B) \subseteq \operatorname{Hom}_{\mathcal{C}}(M, N)$ be the subset of all morphisms factoring through a projective object. Since the zero object $0 \in \mathcal{C}$ is projective, the zero morphism is in $\mathcal{P}_{\mathcal{C}}(A, B)$. Let $\varphi : M \to P \to N$ and $\psi : M \to Q \to N$ be morphisms factoring though projectives. If $\Delta_A : A \to A \oplus A$ is the diagonal map and $\nabla_B : B \oplus B \to B$ the codiagonal, then $\varphi + \psi = \nabla_B \circ (\varphi \oplus \psi) \circ \Delta_A$ by [27, p. 253]. Factoring the map $\varphi \oplus \psi : A \oplus A \to B \oplus B$ further, we obtain

$$\varphi + \psi : A \xrightarrow{\Delta_A} A \oplus A \to P \oplus Q \to B \oplus B \xrightarrow{\nabla_B} B.$$

As $P \oplus Q$ is a projective object [27, p. 250–251], $\varphi + \psi \in \mathcal{P}_{\mathcal{C}}(M, N)$ and $\mathcal{P}_{\mathcal{C}}(A, B)$ is a subgroup of $\operatorname{Hom}_{\mathcal{C}}(A, B)$. Define $[A, B]_{\mathcal{C}} := \operatorname{Hom}_{\mathcal{C}}(A, B)/\mathcal{P}_{\mathcal{C}}(A, B)$. The following proposition is not only relevant to the construction, but also to the construction of Yoneda products in [16, Theorem 6.6].

Proposition 5.1 The bifunctor \circ : Hom_{\mathcal{C}} $(B, C) \times$ Hom_{\mathcal{C}} $(A, B) \rightarrow$ Hom_{\mathcal{C}}(A, C) descends to a bifunctor \circ : $[B, C]_{\mathcal{C}} \times [A, B]_{\mathcal{C}} \rightarrow [A, C]_{\mathcal{C}}$ that is associative and additive in both variables.

Proof First we compose the composition bifunctor with the projection homomorphism $\operatorname{Hom}_{\mathcal{C}}(A, C) \to [A, C]_{\mathcal{C}}$. Then we quotient out $\operatorname{Hom}_{\mathcal{C}}(B, C) \times \mathcal{P}_{\mathcal{C}}(A, B)$ in the domain yielding $\operatorname{Hom}_{\mathcal{C}}(B, C) \times [M, N]_{\mathcal{C}} \to [A, C]_{\mathcal{C}}$. Lastly, we quotient out $\mathcal{P}_{\mathcal{C}}(B, C) \times [A, B]_{\mathcal{C}}$ in the domain. Pictorially, this can be summarised in the commutative diagram

Since the original composition functor is associative as well as additive in both variables and the new one is obtained by subsequent quotients, also the resulting composition functor is associative as well as additive in both variables. $\hfill \Box$

If $(A_n)_{n \in \mathbb{N}_0}$, $(B_n)_{n \in \mathbb{N}_0}$ are projective resolutions of A, B, then the first step of the naïve construction is given by the following generalisation of results found in [25, p. 204]

Proposition 5.2 There exists a homomorphism $t_{A,B} : [A, B]_{\mathcal{C}} \to [\widetilde{A}_1, \widetilde{B}_1]_{\mathcal{C}}$ with the property that $t_{B,C}(-) \circ t_{A,B}(-) = t_{A,C}(-\circ -)$.

Proof Consider the following version of Diagram 4.8:

If $f'_0: A_0 \to B_0$ and $\widetilde{f}'_1: \widetilde{A}_1 \to \widetilde{B}_1$ are different lifts, then the diagram

$0 \longrightarrow \widetilde{A}_1 \xrightarrow{\iota_A} A_0 \xrightarrow{\pi_A} A \longrightarrow 0$
$\left \widetilde{f_1} - \widetilde{f_1'} \right = \left f_0 - f_0' \right = 0$
$0 \longrightarrow \widetilde{B}_1 \xrightarrow{\iota_B} B_0 \xrightarrow{\pi_B} B \longrightarrow 0$

commutes. Because ι_B is the kernel of π_B , there is a morphism $e: A_0 \to \widetilde{B}_1$ making the bottom right triangle of


commute. As the above square also commutes and ι_B is a monomorphism, we observe that $\tilde{f}_1 - \tilde{f}'_1 = e \circ \iota_A$. Using that A_0 is projective, \tilde{f}_1 and \tilde{f}'_1 agree in $[\tilde{A}_1, \tilde{B}_1]_{\mathcal{C}}$. Therefore, there is a well define map

$$s_{A,B}$$
: Hom _{\mathcal{C}} $(A, B) \to [\widetilde{A}_1, \widetilde{B}_1]_{\mathcal{C}}, f \mapsto \widetilde{f}_1 + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_1, \widetilde{B}_1).$

We infer by construction that

$$s_{A,B}(f+g) = \widetilde{f}_1 + \widetilde{g}_1 + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_1, \widetilde{B}_1)$$


and $s_{A,B}$ is a homomorphism. If $h \in \operatorname{Hom}_{\mathcal{C}}(B,C)$, then one can concatenate the corresponding diagrams of the above form to obtain that $s_{B,C}(h) \circ s_{A,B}(f) = s_{A,C}(h \circ f)$. In particular, if $f \in \mathcal{P}_{\mathcal{C}}(A, B)$, then one can choose a projective resolution of the respective projective such that $s_{A,B}(f) = 0$. Thus, $s_{A,B}$ descends to the desired homomorphism $t_{A,B} : [A, B]_{\mathcal{C}} \to [\widetilde{A}_1, \widetilde{B}_1]_{\mathcal{C}}$. Using again that $s_{B,C}(-) \circ s_{A,B}(-) = s_{A,C}(-\circ -)$, we can factor the respective maps through quotient homomorphisms as in the commutative diagram



Note that Diagram 5.1 can be fitted into the left hand side of the above rectangle. Since the quotient homomorphisms from the proof of Proposition 5.1 give rise to the surjective map

 $\operatorname{Hom}_{\mathcal{C}}(B,C) \times \operatorname{Hom}_{\mathcal{C}}(A,B) \to \operatorname{Hom}_{\mathcal{C}}(B,C) \times [A,B]_{\mathcal{C}} \to [B,C]_{\mathcal{C}} \times [A,B]_{\mathcal{C}},$

we conclude that the square



commutes, meaning that $t_{B,C}(-) \circ t_{A,B}(-) = t_{A,C}(-\circ -)$.

Definition 5.3 Let $n \in \mathbb{Z}$. Then there exists for any $k \in \mathbb{N}_0$ with $n + k \ge 0$ a homomorphism $t_{\widetilde{A}_{n+k},\widetilde{B}_{n+k}}$: $[\widetilde{A}_{n+k},\widetilde{B}_k]_{\mathcal{C}} \to [\widetilde{A}_{n+k+1},\widetilde{B}_{k+1}]_{\mathcal{C}}$. We define the n^{th} term of the naïve construction to be $BC^n_{\mathcal{C}}(A,B) := \varinjlim_{\mathcal{D}} ([\widetilde{A}_{n+k},\widetilde{B}_k]_{\mathcal{C}}, t_{\widetilde{A}_{n+k},\widetilde{B}_k})_{k \in \mathbb{N}_0, n+k \ge 0}$.

In order to define induced homomorphisms for $BC_{\mathcal{C}}^n(-,-)$, let A, B, C be objects in \mathcal{C} and consider corresponding projective resolutions $A_{\bullet}, B_{\bullet}, C_{\bullet}$. For a morphism $f: B \to C$ in \mathcal{C} let us construct a homomorphism $BC_{\mathcal{C}}^n(A, f)$ using these projective resolutions. Lifting fas in the proof of Proposition 5.2, we obtain a sequence of morphisms $(\tilde{f}_k: \tilde{B}_k \to \tilde{C}_k)_{k \in \mathbb{N}_0}$ where

$$t_{\widetilde{B}_k,\widetilde{C}_k}\big(\widetilde{f}_k + \mathcal{P}_{\mathcal{C}}(\widetilde{B}_k,\widetilde{C}_k)\big) = \widetilde{f}_{k+1} + \mathcal{P}_{\mathcal{C}}(\widetilde{B}_{k+1},\widetilde{C}_{k+1}).$$

Since $f = \tilde{f}_0$, the element $(\tilde{f}_k + \mathcal{P}_{\mathcal{C}}(\tilde{B}_k, \tilde{C}_k))_{k \in \mathbb{N}_0}$ of $([\tilde{B}_k, \tilde{C}_k]_{\mathcal{C}})_{k \in \mathbb{N}_0}$ is independent of the lifts of f. Again by Proposition 5.2, $t_{\tilde{A}_{n+k},\tilde{C}_k}(-\circ -) = t_{\tilde{B}_k,\tilde{C}_k}(-) \circ t_{\tilde{A}_{n+k},\tilde{B}_k}(-)$ and thus the square of homomorphisms of abelian groups

$$[\widetilde{A}_{n+k}, \widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{\left(\widetilde{f}_{k} + \mathcal{P}_{\mathcal{C}}(\widetilde{B}_{k}, \widetilde{C}_{k})\right) \circ -}} [\widetilde{A}_{n+k}, \widetilde{C}_{k}]_{\mathcal{C}} \xrightarrow{\left(\widetilde{f}_{k+1}, \widetilde{P}_{\mathcal{C}}(\widetilde{B}_{k+1}, \widetilde{C}_{k+1})\right) \circ -}} [\widetilde{A}_{n+k+1}, \widetilde{C}_{k+1}]_{\mathcal{C}} \xrightarrow{\left(\widetilde{f}_{k+1} + \mathcal{P}_{\mathcal{C}}(\widetilde{B}_{k+1}, \widetilde{C}_{k+1})\right) \circ -}} [\widetilde{A}_{n+k+1}, \widetilde{C}_{k+1}]_{\mathcal{C}}$$

$$(5.3)$$

commutes. Analogously, if $g: C \to A$ is another morphism, then we can construct an element $(\tilde{g}_k + \mathcal{P}_{\mathcal{C}}(\tilde{C}_k, \tilde{A}_k))_{k \in \mathbb{N}_0}$ of $([\tilde{C}_k, \tilde{A}_k]_{\mathcal{C}})_{k \in \mathbb{N}_0}$ that is independent of the lifts of g.

Definition 5.4 Taking the notation from before, we define

$$BC^n_{\mathcal{C}}(A,f) := \lim_{k \in \mathbb{N}_0, n+k \ge 0} \left(\left(\widetilde{f}_k + \mathcal{P}_{\mathcal{C}}(\widetilde{B}_k, \widetilde{C}_k) \right) \circ - \right) : BC^n_{\mathcal{C}}(A,B) \to BC^n_{\mathcal{C}}(A,C)$$

and

$$BC^n_{\mathcal{C}}(g,B) := \lim_{k \in \mathbb{N}_0, n+k \ge 0} \left(-\circ \left(\widetilde{g}_{n+k} + \mathcal{P}_{\mathcal{C}}(\widetilde{C}_{n+k}, \widetilde{A}_{n+k}) \right) \right) : BC^n_{\mathcal{C}}(A,B) \to BC^n_{\mathcal{C}}(C,B) \,.$$

Proposition 5.5 For every $n \in \mathbb{Z}$, $BC^n_{\mathcal{C}}(-,-) : \mathcal{C}^{\text{op}} \times \mathcal{C} \to Ab$ forms a well defined bifunctor that is additive in both variables.

Proof We demonstrate that $BC_{\mathcal{C}}(A, B)$ is independent of choices of projective resolutions. Hence let A_{\bullet} and A'_{\bullet} be projective resolutions of A. As in Definition 5.4, take $(\tilde{\iota}_k + \mathcal{P}_{\mathcal{C}}(\tilde{A}_k, \tilde{A}'_k))_{k \in \mathbb{N}_0}$ and $(\tilde{\iota}'_k + \mathcal{P}_{\mathcal{C}}(\tilde{A}'_k, \tilde{A}_k))_{k \in \mathbb{N}_0}$ lifting $\mathrm{id}_A : A \to A$. Given that $(\tilde{\iota}'_k \circ \tilde{\iota}_k + \mathcal{P}_{\mathcal{C}}(\tilde{A}_k, \tilde{A}_k))_{k \in \mathbb{N}_0}$ and $(\tilde{\iota}_k \circ \tilde{\iota}'_k + \mathcal{P}_{\mathcal{C}}(\tilde{A}'_k, \tilde{A}'_k))_{k \in \mathbb{N}_0}$ are also lifts of id_A , we conclude by Proposition 5.2 that

$$\iota'_k \circ \iota_k + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_k, \widetilde{A}_k) = \operatorname{id}_{\widetilde{A}_k} + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_k, \widetilde{A}_k) \text{ and } \iota_k \circ \iota'_k + \mathcal{P}_{\mathcal{C}}(\widetilde{A}'_k, \widetilde{A}'_k) = \operatorname{id}_{\widetilde{A}'_k} + \mathcal{P}_{\mathcal{C}}(\widetilde{A}'_k, \widetilde{A}'_k)$$

for any $k \in \mathbb{N}_0$. As there is an analogous statement for any two projective resolutions of $B, BC_{\mathcal{C}}^n(A, B)$ is unique up to isomorphism. Knowing that composition of morphisms is associative by Proposition 5.1, we see that

$$BC_{\mathcal{C}}(A, f' \circ f) = BC_{\mathcal{C}}(A, f') \circ BC_{\mathcal{C}}(A, f)$$
 and $BC_{\mathcal{C}}(g \circ g', B) = BC_{\mathcal{C}}(g, B) \circ BC_{\mathcal{C}}(g', B)$

for any $f': C \to D$ and $g': D \to C$. Thus, $BC_{\mathcal{C}}^{m}(-,-)$ a well defined bifunctor. Because composition of morphisms is bi-additive also by Proposition 5.1 and Diagram 5.3 commutes, we infer that

$$BC^n_{\mathcal{C}}(A, f+f'') = BC^n_{\mathcal{C}}(A, f) + BC^n_{\mathcal{C}}(A, f'') \text{ and } BC^n_{\mathcal{C}}(g+g', B) = BC^n_{\mathcal{C}}(g, B) + BC^n_{\mathcal{C}}(g'', B)$$

for any $f, f'': B \to C$ and $g, g'': C \to A$. In particular, $BC_{\mathcal{C}}^n(-, -)$ is additive in both variables.

5.2 Isomorphism of cohomology groups

Let us construct a homomorphism from completed Ext-functors $\widehat{\operatorname{Ext}}_{\mathcal{C}}^n(A, B)$ arising from the resolution construction to the terms $BC_{\mathcal{C}}^n(A, B)$ of the naïve construction. The idea of the construction and of the proof that it yields an isomorphism can be retrieved from page 299 of G. Mislin's paper [28] and are generalised to our setting in the following. Again, this differs from S. Guo and L. Liang's account containing a similar construction and proof on pages 18–19 of their paper [19]. If $\partial_{n+1} : A_{n+1} \to A_n$ denotes a morphism of a projective resolution A_{\bullet} of A, then

$$\operatorname{Ker}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+1}, B)) = \{ f \in \operatorname{Hom}_{\mathcal{C}}(A_n, B) \mid f \circ \partial_{n+1} = 0 \}.$$

Note that the cokernel of ∂_{n+1} can be realised by the morphism $\pi_n : A_n \to \widetilde{A}_n$ occurring in the factorisation $\partial_n : A_n \xrightarrow{\pi_n} \widetilde{A}_n \xrightarrow{\iota_n} A_{n-1}$. Thus, for every $f \in \text{Ker}(\text{Hom}_{\mathcal{C}}(\partial_{n+1}, B))$ there is a unique morphism $f': \widetilde{A}_n \to B$ such that $f = f' \circ \pi_n$. Therefore, there is a bijection

$$\alpha_n(B) : \operatorname{Ker}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+1}, B)) \to \operatorname{Hom}_{\mathcal{C}}(A_n, B), \ f \mapsto f'.$$

As $(\alpha_n(B)(f) + \alpha_n(B)(g)) \circ \pi_n = f + g$, $\alpha_n(B)$ is an isomorphisms of abelian groups. Pictorially, one can summarise the content of this isomorphism as

$$\begin{array}{ccc} A_n & \xrightarrow{\pi_n} & \widetilde{A}_n \\ & \searrow f & \downarrow_{\alpha_n(B)(f)} \\ & & B \end{array} \tag{5.4}$$

Before we proceed we need to observe that in the case n = 0 this yields an isomorphism

$$\alpha_0(B) : \operatorname{Ext}^0_{\mathcal{C}}(A, B) \to \operatorname{Hom}_{\mathcal{C}}(A, B)$$

since Ker(Hom_{\mathcal{C}}(∂_0, B)) = Ext⁰_{\mathcal{C}}(A, B). If $n \geq 1$, assume that f is an element in Im(Hom_{\mathcal{C}}(∂_n, B)) \leq Ker(Hom_{\mathcal{C}}(∂_{n+1}, B)). Being of the form $f : A_n \xrightarrow{\partial_n} A_{n-1} \to B$, it follows from the above factorisation of ∂_n that $\alpha_n(B)(f) : \widetilde{A}_n \xrightarrow{\iota_n} A_{n-1} \to B$. As A_{n-1} is a projective object, $\alpha_n(B)(f) \in \mathcal{P}_R(\widetilde{A}_n, B)$. Thus $\alpha_n(B)$ descends to a homomorphism of abelian groups

$$\beta_n(B) \colon \operatorname{Ext}^n_{\mathcal{C}}(A, B) \to [A_n, B]_{\mathcal{C}}$$
$$f + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_n, B)) \mapsto \alpha_n(B)(f) + \mathcal{P}_R(\widetilde{A}_n, B).$$

Due to the above, we remark that $\beta^n(B)$ is only defined for $n \ge 1$ while we already have the isomorphism $\alpha_0(B)$ in the case n = 0. The last ingredient to construct our desired isomorphism between the cohomology groups is

Proposition 5.6 ([30, p. 261]) In the category Ab of abelian groups, the direct limit of a direct system $\{M_i, \psi_{i,j}\}_{i \leq j \in I}$ can be given as $\bigoplus_{i \in I} M_i / \sim$ where $m_i \in M_i \sim m_j \in M_j$ if there is $k \in I$ such that $\psi_{i,k}(m_i) = \psi_{j,k}(m_j) \in M_k$. In particular, if $\{m_i \in M_i\}_{i \in I}$ is a collection of nonzero elements such that $\psi_{i,j}(m_i) = m_j$ for any $i \leq j$, then the element $(m_i)_{i \in I} \in \varinjlim_{i \in I} M_i$ is nonzero.

Hence we can generalise G. Mislin's Theorem 4.1 and its proof from [28].

Lemma 5.7 For every $n \in \mathbb{Z}$ and $k \in \mathbb{N}_0$ with $n + k \ge 1$ the square

$$\operatorname{Ext}_{\mathcal{C}}^{n+k}(A, \widetilde{B}_{k}) \xrightarrow{\beta_{n+k}(B_{k})} [\widetilde{A}_{n+k}, \widetilde{B}_{k}]_{\mathcal{C}}$$

$$\downarrow_{\delta^{n+k}} \qquad \qquad \qquad \qquad \downarrow^{t_{\widetilde{A}_{n+k}, \widetilde{B}_{k}}} \qquad (5.5)$$

$$\operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A, \widetilde{B}_{k+1}) \xrightarrow{\beta_{n+k+1}(\widetilde{B}_{k+1})} [\widetilde{A}_{n+k+1}, \widetilde{B}_{k+1}]_{\mathcal{C}}$$

commutes. The induced homomorphism

$$\beta^n(B) := \lim_{k \in \mathbb{N}_0, n+k \ge 1} \beta_{n+k}(\widetilde{B}_k) : \widehat{\operatorname{Ext}}^n_{\mathcal{C}}(A, B) \to BC^n_{\mathcal{C}}(A, B)$$

is an isomorphism.

Proof Note that we impose $n + k \ge 1$ in Diagram 5.5 and in the definition of $\beta^n(B)$ so that the morphisms $\beta_{n+k}(B)$ are well defined. To prove that the diagram is commutative, let $f + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{A}_k))$ be an element in $\operatorname{Ext}_{\mathcal{C}}^{n+k}(A, \widetilde{B}_k)$. It suffices to show that the diagram

commutes where we explain in the following how to construct it. The right most triangle corresponds to Diagram 5.4. We can lift $\alpha_{n+k}(\widetilde{B}_k)(f)$ to $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)}$ and $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)}^*$ as we have done in Diagram 5.2. In particular, the morphism $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)}^*$ is a representative of $t_{\widetilde{A}_{n+k},\widetilde{B}_k} \circ \beta_{n+k}(\widetilde{B}_k)(f) + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k},\widetilde{B}_k)))$. Writing F for $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)}^* \circ \pi_{n+k+1}$, let us deduce that

$$\delta^{n+k} \left(f + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k)) \right) = F + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k+1}, \widetilde{B}_{k+1}))$$

in $\operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A, \widetilde{B}_{k+1})$. Following the construction of the connecting homomorphism δ^{n+k} by means of the Snake Lemma as in [42, p. 11–12], we see that $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)}$ is a lift of \underline{f} along π_k . As the boundary map ∂_{n+k+1} equals $\iota_{n+k+1} \circ \pi_{n+k+1} : A_{n+k+1} \to A_{n+k}$ and $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)} \circ \partial_{n+k+1}$ equals $\iota_{k+1} \circ F$, F is a representative of $\delta^{n+k}(f+\operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k},\widetilde{B}_k)))$. Thus, $\overline{\alpha_{n+k}(\widetilde{B}_k)(f)}^*$ is a representative of $\beta_{n+k+1}(\widetilde{B}_{k+1}) \circ \delta^{n+k}(f+\operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k},\widetilde{B}_k)))$ and Diagram 5.5 commutes.

To show that $\beta^n(B)$ is an isomorphism, let us first prove that it is surjective. For this, let $g + \mathcal{P}_R(\widetilde{A}_{n+k}, \widetilde{B}_k) \in [\widetilde{A}_{n+k}, \widetilde{B}_k]_{\mathcal{C}}$. Due to the diagram

$$\begin{array}{ccc} A_{n+k} & \xrightarrow{\pi_{n+k}} \widetilde{A}_{n+k} \\ & & & & \downarrow^g \\ & & & & \downarrow^g \\ & & & & \widetilde{B}_k \end{array}$$

and Diagram 5.4, we see that $\alpha_{n+k}(\widetilde{B}_k)(g \circ \pi_{n+k}) = g$ and thus

$$\beta_{n+k}(\widetilde{B}_k)(g \circ \pi_{n+k} + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k))) = g + \mathcal{P}_R(\widetilde{A}_{n+k}, \widetilde{B}_k).$$

Since $\beta_{n+k}(\widetilde{B}_k)$ is surjective and direct limits in the category of abelian groups are exact, $\beta^n(B) = \varinjlim_{k \in \mathbb{N}_0, n+k \ge 1} \beta_{n+k}(\widetilde{B}_k)$ is surjective.

In order to demonstrate that $\beta^n(B)$ is injective, let $\varphi \in \widehat{\operatorname{Ext}}_{\mathcal{C}}^n(A, B)$ be an element such that $\beta^n(B)(\varphi) = 0$. By the resolution construction, $\widehat{\operatorname{Ext}}_{\mathcal{C}}^n(A, B)$ can be defined as the direct limit $\varinjlim_{k \in \mathbb{N}_0} \operatorname{Ext}_{\mathcal{C}}^{n+k}(A, \widetilde{B}_k)$ in the category **Ab**. By Proposition 5.6, there exists $k \in \mathbb{N}_0$ and $h \in \operatorname{Ker}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k+1}, \widetilde{B}_k)))$ such that the element $h + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k))$ of $\operatorname{Ext}_{\mathcal{C}}^{n+k}(A, \widetilde{B}_k)$ is mapped to $\varphi \in \operatorname{Ext}_{\mathcal{C}}^n(A, B)$ by the homomorphism to the direct limit. Still using Proposition 5.6, we may assume $\beta_{n+k}(\widetilde{B}_k)(h + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k))) = 0$. This implies that the morphism $\alpha_{n+k}(\widetilde{B}_k)(h) : \widetilde{A}_{n+k} \to \widetilde{B}_k$ factors through a projective.

Because the morphism π_k at the bottom of Diagram 5.6 is an epimorphism, it follows from the lifting property of the previous projective object that there is a morphism l making the diagram

$$B_k \xrightarrow{\widetilde{A}_{n+k}} \widetilde{B}_k$$

commute. From this diagram and Diagram 5.6 we infer that

$$\pi_k \circ (l \circ \pi_{n+k}) = \alpha_{n+k}(\widetilde{B}_k)(h) \circ \pi_{n+k} = \pi_k \circ \overline{\alpha_{n+k}(\widetilde{B}_k)(h)}$$

In particular, $\pi_k \circ (\alpha_{n+k}(\widetilde{B}_k)(h) - l \circ \pi_{n+k}) = 0$. Note that the kernel of π_k is given by the morphism $\iota_{k+1} : \widetilde{B}_{k+1} \to B_k$ whence there is a L such that

$$\widetilde{B}_{k+1} \xrightarrow{\iota_{k+1}} B_k \xrightarrow{A_{n+k}} B_k$$

commutes. Using the morphism $\iota_{n+k+1}: \widetilde{A}_{n+k+1} \to A_{n+k}$ from Diagram 5.6, we see that

$$\iota_{k+1} \circ L \circ \iota_{n+k+1} = (\overline{\alpha_{n+k}(\widetilde{B}_k)(h)} - l \circ \pi_{n+k}) \circ \iota_{n+k+1}.$$

As $\pi_{n+k} \circ \iota_{n+k+1} = 0$, we deduce from Diagram 5.6 that

$$\iota_{k+1} \circ L \circ \iota_{n+k+1} = \overline{\alpha_{n+k}(\widetilde{B}_k)(h)} \circ \iota_{n+k+1} = \iota_{k+1} \circ \overline{\alpha_{n+k}(\widetilde{B}_k)(h)}^*.$$

Because ι_{n+k+1} is a monomorphism, we conclude that $L \circ \iota_{n+k+1} = \overline{\alpha_{n+k}(\widetilde{B}_k)(f)}^*$. This conclusion together with the previous diagram and Diagram 5.6 can be pictorially summarised as

Recalling that $\pi_{n+k+1} \circ \iota_{n+k+1} = \partial_{n+k+1}$, we have $H = L \circ \partial_{n+k+1} \in \operatorname{Hom}_{\mathcal{C}}(\partial_{n+k+1}, \widetilde{B}_{k+1})$. Since this is a representative in $\operatorname{Im}(\delta^{n+k})$, we infer that $\delta^{n+k}(h+\operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_{k}))) = 0$. According to the resolution construction, the latter as well as $h + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_{k}))$ are both mapped to $\varphi \in \operatorname{Ext}^{n}_{\mathcal{C}}(A, B)$. Therefore $\varphi = 0$, $\operatorname{Ker}(\beta^{n}(B)) = 0$ and $\beta^{n}(B)$ is injective.

Proposition 5.8 For every $n \in \mathbb{Z}$ the isomorphism $\beta^n(-)$ is natural. That is, for every $f \in \operatorname{Hom}_{\mathcal{C}}(B, C)$ the square

$$\begin{array}{cccc}
\widehat{\operatorname{Ext}}^{n}_{\mathcal{C}}(A,B) &\xrightarrow{\beta^{n}(B)} & BC^{n}_{\mathcal{C}}(A,B) \\
& & & \downarrow^{\widehat{\operatorname{Ext}}^{n}_{\mathcal{C}}(A,f)} & \downarrow^{BC^{n}_{\mathcal{C}}(A,f)} \\
& & & \widehat{\operatorname{Ext}}^{n}_{\mathcal{C}}(A,C) \xrightarrow{\beta^{n}(C)} & BC^{n}_{\mathcal{C}}(A,C)
\end{array}$$

commutes.

Proof By construction of $\widehat{\operatorname{Ext}}_{\mathcal{C}}^{n}(A, f)$ through the resolution construction from Proposition 4.2 and of $BC_{\mathcal{C}}^{n}(A, f)$ from Definition 5.4, it suffices to consider the cube

for any $k \in \mathbb{N}_0$ with $n + k \ge 1$. The right hand side commutes due to the choice of the sequence of morphisms $(\tilde{f}_k : \tilde{B}_k \to \tilde{C}_k)_{k \in \mathbb{N}_0}$ and the left hand side since connecting homomorphisms are natural. The front and back side correspond to Diagram 5.5. Regarding its top and bottom side, let $g \in \text{Ker}(\text{Hom}_{\mathcal{C}}(\partial_{n+k+1}, \tilde{B}_k))$. Then

$$([\widetilde{A}_{n+k}, \widetilde{f}_k]_{\mathcal{C}} \circ \beta_{n+k}(\widetilde{B}_k)) (g + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k)))$$
$$= [\widetilde{A}_{n+k}, \widetilde{f}_k]_{\mathcal{C}} (\alpha_{n+k}(g) + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k}, \widetilde{B}_k))$$
$$= \widetilde{f}_k \circ \alpha_{n+k}(g) + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k}, \widetilde{C}_k).$$

Note that

$$\widetilde{f}_k \circ \alpha_{n+k}(\widetilde{B}_k)(g) \circ \pi_{n+k} = \widetilde{f}_k \circ g = \alpha_{n+k}(\widetilde{C}_k)(\widetilde{f}_k \circ g) \circ \pi_{n+k}.$$

By definition of the morphism $\alpha_{n+k}(\widetilde{C}_k) : \operatorname{Ext}_{\mathcal{C}}^{n+k}(A, \widetilde{C}_k) \to [\widetilde{A}_{n+k}, \widetilde{C}_k]_{\mathcal{C}}$ pictorially given in Diagram 5.4, we deduce that $\widetilde{f}_k \circ \alpha_{n+k}(\widetilde{B}_k)(g) = \alpha_{n+k}(\widetilde{C}_k)(\widetilde{f}_k \circ g)$. In particular,

$$([\widetilde{A}_{n+k}, \widetilde{f}_k]_{\mathcal{C}} \circ \beta_{n+k}(\widetilde{B}_k)) (g + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k))) = \alpha_{n+k}(\widetilde{C}_k) (\widetilde{f}_k \circ g) + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k}, \widetilde{C}_k) = \beta_{n+k}(\widetilde{C}_k) (\widetilde{f}_k \circ g + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{C}_k))) = (\beta_{n+k}(\widetilde{C}_k) \circ \operatorname{Ext}_{\mathcal{C}}^{n+k}(A, \widetilde{f}_k)) (g + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(\partial_{n+k}, \widetilde{B}_k)))) .$$

Thus, the top and bottom side of the above cube also commute and form a direct system of commuting squares. The statement of the proposition follows by passing to the direct limit. $\hfill \Box$

5.3 Connecting homomorphisms of the naïve construction

To render the naïve construction into a cohomological functor, a connecting homomorphism is needed. It is formed similarly to the morphisms $\varepsilon^{-m} : S^{-m}T \to S^{-m+1}T$ between satellite functors that we have established at the start of Section 3. Namely, let $0 \to B \xrightarrow{f} C \xrightarrow{g} D \to 0$ be a short exact sequence in \mathcal{C} . By Diagram 4.9, we can lift the morphisms f and g such that the diagram

commutes and has short exact sequences as rows for every $k \in \mathbb{N}_0$. As the domain of the connecting homomorphism is supposed to be $BC_R^n(A, D) = \varinjlim_{k \in \mathbb{N}_0, n+k \ge 0} [\widetilde{A}_{n+k}, \widetilde{D}_k]_C$, let $l \in \operatorname{Hom}_{\mathcal{C}}(\widetilde{A}_{n+k}, \widetilde{D}_k)$. According to Diagram 5.2, we can lift l such that the diagram

commutes and $t_{\widetilde{A}_{n+k},\widetilde{D}_k}(l + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k},\widetilde{D}_k)) = \overline{l}^* + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k+1},\widetilde{D}_{k+1})$. Using Diagram 3.2, we can extend it in a commutative manner to

We define the map

$$\tau_{n+k,k} := [\widetilde{A}_{n+k+1}, h^*]_{\mathcal{C}} \circ t_{\widetilde{A}_{n+k}, \widetilde{D}_k} : \quad [\widetilde{A}_{n+k}, \widetilde{D}_k]_{\mathcal{C}} \to [\widetilde{A}_{n+k+1}, \widetilde{B}_k]_{\mathcal{C}}$$
$$l + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k}, \widetilde{D}_k) \mapsto h^* \circ \overline{l}^* + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{n+k+1}, \widetilde{B}_k)$$

Following the proof of Proposition 5.2, it is well defined in the sense that it does not depend on the particular lift $h^* : \widetilde{D}_{k+1} \to \widetilde{B}_k$ of $\mathrm{id}_{\widetilde{D}_k}$. By Proposition 5.1 and 5.2, $\tau_{n+k,k}$ is a homomorphism.

Lemma 5.9 Both the squares

$$\operatorname{Ext}_{\mathcal{C}}^{n+k}(A,\widetilde{D}_{k}) \xrightarrow{\beta_{n+k}(\widetilde{D}_{k})} [\widetilde{A}_{n+k},\widetilde{D}_{k}]_{\mathcal{C}} \qquad [\widetilde{A}_{n+k},\widetilde{D}_{k}]_{\mathcal{C}} \xrightarrow{(-1)^{k}\tau_{n+k,k}} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(-1)^{k}\tau_{n+k,k}} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(-1)^{k}\tau_{n+k,k}} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(\tau_{n+k},\varepsilon_{n+k})} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(\tau_{n+k},\varepsilon_{n+k})} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(-1)^{k+1}\tau_{n+k+1,k+1}} [\widetilde{A}_{n+k+2},\widetilde{B}_{k+1}]_{\mathcal{C}}$$

commute for every $k \in \mathbb{N}_0$ with $n + k \ge 1$. In particular, the induced homomorphism

$$\tau^n := \varinjlim_{k \in \mathbb{N}_0, n+k \ge 1} (-1)^k \tau_{n+k,k} : BC^n_{\mathcal{C}}(A, D) \to BC^{n+1}_{\mathcal{C}}(A, B)$$

fits into the commuting square

$$\begin{array}{ccc} & \widehat{\operatorname{Ext}}^{n}_{\mathcal{C}}(A,D) \xrightarrow{\beta^{n}(D)} BC^{n}_{\mathcal{C}}(A,D) \\ & & & \downarrow^{\widehat{\delta}^{n}} & & \downarrow^{\tau^{n}} \\ & & \widehat{\operatorname{Ext}}^{n+1}_{\mathcal{C}}(A,B) \xrightarrow{\beta^{n+1}(B)} BC^{n+1}_{\mathcal{C}}(A,B) \\ \end{array}$$

Therefore, $\tau^n : BC^n_{\mathcal{C}}(A, -) \to BC^{n+1}_{\mathcal{C}}(A, -)$ represents the (proposed) connecting homomorphism of $BC^{\bullet}_{\mathcal{C}}(A, -)$.

Proof Taking a morphism $h^* : \widetilde{D}_{k+1} \to \widetilde{B}_k$ lifting $\operatorname{id}_{\widetilde{D}_k} : \widetilde{D}_k \to \widetilde{D}_k$ as before, we apply the Ext-functor $\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -)$ to obtain the commutative square

Using this and the definition of the morphism $\tau_{n+k,k}$, we can factorise the top left square in the statement of the Lemma as

$$\operatorname{Ext}_{\mathcal{C}}^{n+k}(A,\widetilde{D}_{k}) \xrightarrow{\beta_{n+k}(\widetilde{D}_{k})} [\widetilde{A}_{n+k},\widetilde{D}_{k}]_{\mathcal{C}} \downarrow_{\delta^{n+k}} \downarrow^{t_{\widetilde{A}_{n+k},\widetilde{D}_{k}}} \operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A,\widetilde{D}_{k+1}) \xrightarrow{\beta_{n+k+1}(\widetilde{D}_{k+1})} [\widetilde{A}_{n+k+1},\widetilde{D}_{k+1}]_{\mathcal{C}} \downarrow_{\operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A,h^{*})} \downarrow^{[\widetilde{A}_{n+k+1},h^{*}]_{\mathcal{C}}} \operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A,\widetilde{B}_{k}) \xrightarrow{\beta_{n+k+1}(\widetilde{B}_{k})} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}}$$

It commutes by Lemma 5.7 and Proposition 5.8. As for the top right square in the statement of the lemma, we can fit it as the right hand side of the cube

$$\operatorname{Ext}_{\mathcal{C}}^{n+k}(A,\widetilde{D}_{k}) \xrightarrow{\beta_{n+k}(\widetilde{D}_{k})} [\widetilde{A}_{n+k},\widetilde{D}_{k}]_{\mathcal{C}}$$

$$\operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A,\widetilde{B}_{k}) \xrightarrow{(-1)^{k}\delta^{n+k}} \beta_{n+k+1}(\widetilde{B}_{k}) \xrightarrow{(-1)^{k}\tau_{n+k,k}} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(-1)^{k}\tau_{n+k,k}} t_{\widetilde{A}_{n+k},\widetilde{D}_{k}}$$

$$\downarrow^{\delta^{n+k+1}} \operatorname{Ext}_{\mathcal{C}}^{n+k+1}(A,\widetilde{D}_{k+1}) \xrightarrow{(-1)^{k+1}\delta^{n+k+1}} \underbrace{\beta_{n+k+1}(\widetilde{D}_{k+1})}_{(-1)^{k+1}\delta^{n+k+1}} [\widetilde{A}_{n+k+1},\widetilde{B}_{k}]_{\mathcal{C}} \xrightarrow{(-1)^{k+1}\tau_{n+k+1,k+1}}$$

$$\operatorname{Ext}_{\mathcal{C}}^{n+k+2}(A,\widetilde{B}_{k+1}) \xrightarrow{\beta_{n+k+2}(\widetilde{B}_{k+1})} [\widetilde{A}_{n+k+1},\widetilde{B}_{k+1}]_{\mathcal{C}}$$

We have just seen that the top and bottom side commutes. The left hand side corresponds to Diagram 4.10 while the front and back side commute by Lemma 5.7. The right hand commutes as desired since $\beta_{n+k+1}(\tilde{B}_k)$ is an epimorphism by the proof of Lemma 5.7. Lastly, the top and bottom sides of these cubes form a direct system of commuting squares in whose direct limit we retrieve the bottom square from the statement of the lemma.

Theorem 5.10 The naïve construction $(BC^{\bullet}_{\mathcal{C}}(A, -), \tau^{\bullet})$ forms a cohomological functor and $\beta^{\bullet} : (\widehat{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(A, -), \widehat{\delta}^{\bullet}) \to (BC^{\bullet}_{\mathcal{C}}(A, -), \tau^{\bullet})$ is an isomorphism of cohomological functors.

Proof We already know that $(\widehat{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(A, -), \widehat{\delta}^{\bullet})$ is a cohomological functor by Lemma 3.2, that β^{\bullet} is a natural transformation by Proposition 5.8 and that every β^n is an isomorphism by Lemma 5.7. Hence, we can reiterate the proof of Theorem 4.5 where we invoke Lemma 5.9 instead of Lemma 4.4 whenever required.

Remark 5.11 On page 299 of [28], G. Mislin constructs for every $n \in \mathbb{Z}$ an isomorphism $\lim_{k \in \mathbb{N}_0} S^{-k} \operatorname{Ext}^{n+k}(A, -) \to BC^n(A, -)$ in the case of modules over a ring. Then he asserts that one can turn $BC^{\bullet}(A, -)$ into a Mislin completion by transferring the structure of $\widehat{\operatorname{Ext}}^{\bullet}(A, -)$ and claims that homomorphisms of the form $[\widetilde{A}_{n+k}, \widetilde{D}_k] \to [\widetilde{A}_{n+k+1}, \widetilde{B}_k]$ induce the connecting homomorphism $\tau^n : BC^n(A, D) \to BC^{n+1}(A, B)$. However, he does not explain how to construct the latter homomorphisms and in particular, does not indicate that there alternating signs involved in the direct system giving rise to τ^n as can be seen in Lemma 5.9.

6 The hypercohomology and the resolution construction

We demonstrate in this section that the hypercohomology construction gives rise to the Mislin completion of unenriched Ext-functors in the same way as the naïve construction. For this, we relate in Subsection 6.1 the hypercohomology complex with chain maps and the Vogel complex with so-called almost chain maps. It is known that unenriched Extgroups can be retrieved as cohomology groups of the former complex. In Subsection 6.2 we establish an explicit isomorphism between these two forms of Ext-groups and use it to construct connecting homomorphisms through (almost) chain maps. This sets the base for Subsection 6.3 where we construct a natural isomorphism from the cohomology groups of the resolution construction to the ones of the hypercohomology construction. In Subsection 6.4, we extend it to an isomorphism of cohomological functors. Moreover, we provide an explicit isomorphism of cohomological functors from the Mislin completion resulting from the hypercohomology construction to the one resulting from the naïve construction. Lastly, although S. Guo and L. Liang have turned the cohomology groups of the Vogel complex into a cohomological functor in [19, Proposition 4.8], we do not know whether their construction forms a Mislin completion (Question 6.19). Similarly, J. Hu et al. introduce the Vogel complex in [22, p. 7] in a far more general setting than we or S. Guo and L. Liang consider, but we also do not know whether it yields a Mislin completion.

6.1 Defining Ext-functors through chain maps

There is a reformulation of the hypercohomology construction occurring in D. J. Benson and J. F. Carlson's paper [5, p. 109] in terms of almost chain maps. For this, we introduce the following notation. If $(M_k, \mu_k)_{k \in \mathbb{Z}}$ is a chain complex in \mathcal{C} with boundary maps $\mu_k : M_k \to M_{k-1}$ and $n \in \mathbb{Z}$, then we define $(M[n]_k, \mu[n]_k)_{k \in \mathbb{Z}}$ to be $M[n]_k := M_{k+n}$ and $\mu[n]_k := (-1)^n \mu_{k+n}$ [15, p. 154], [42, p. 9–10]. Recall that we have constructed the hypercohomoology complex $\operatorname{Hyp}_{\mathcal{C}}(M_{\bullet}, N_{\bullet})_{\bullet}$ for any other chain complexes $(N_k, \nu_k)_{k \in \mathbb{Z}}$ before Definition 2.11. According to [42, p. 62–63], an *n*-cocycle of $\operatorname{Hyp}_{\mathcal{C}}(M_{\bullet}, N_{\bullet})_{\bullet}$ is exactly a chain map of the form $M[n]_{\bullet} \to N_{\bullet}$ where *n*-coboundaries are nullhomotopic chain maps of this form. In the same way, we observe that an *n*-cocycle of the Vogel complex $\operatorname{Vog}_{\mathcal{C}}(M_{\bullet}, N_{\bullet})_{\bullet}$ is a collection of morphisms $(f_{k+n} : M[n]_k \to N_k)_{k \in \mathbb{Z}}$ such that all but finitely many of the squares

$$\begin{array}{c} M_{k+n} \xrightarrow{f_{k+n}} N_k \\ \downarrow^{(-1)^n \mu_{n+k}} \qquad \downarrow^{\nu_k} \\ M_{k-1+n} \xrightarrow{f_{k-1+n}} N_{k-1} \end{array}$$

commute. We follow the convention in [5, p. 109] by calling this an almost chain map of degree n. We call $f_{\bullet+n}: M[n]_{\bullet} \to N_{\bullet}$ nullhomotopic if there is a collection of morphisms $(e_{k+n}: M[n]_k \to N_{k+1})_{k\in\mathbb{Z}}$ such that $f_{k+n} = e_{k-1+n} \circ \mu[n]_k + \mu[n]_{k+1} \circ e_{k+n}$ for all but finitely many $k \in \mathbb{Z}$. Then we observe that an n-cboundary of $\operatorname{Vog}_{\mathcal{C}}(M_{\bullet}, N_{\bullet})_{\bullet}$ is exactly such a nullhomotopic almost chain map. Denote by $\operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})$ the set of chain maps $M[n]_{\bullet} \to N_{\bullet}$ and by $\operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})$ the subset of nullhomotopic maps. Analogously, write $\operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})$ for the set of almost chain maps $M[n]_{\bullet} \to N_{\bullet}$ and by $\operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})$ the subset of nullhomotopic almost chain maps $M[n]_{\bullet}$ and by $\operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})$ the subset of nullhomotopic almost chain maps. Above, we have just seen that

$$H^{n}(\mathrm{Hyp}_{\mathcal{C}}(M_{\bullet}, N_{\bullet})_{\bullet}) = \mathrm{Hom}_{\mathrm{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})/\mathrm{Null}_{\mathrm{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet}) \text{ and}$$
$$H^{n}(\mathrm{Vog}_{\mathcal{C}}(M_{\bullet}, N_{\bullet})_{\bullet}) = \widehat{\mathrm{Hom}}_{\mathrm{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet})/\widehat{\mathrm{Null}}_{\mathrm{Ch}(\mathcal{C})}(M[n]_{\bullet}, N_{\bullet}).$$

If M_{\bullet} is a projective resolution of an object $M \in obj(\mathcal{C})$ and N_{\bullet} a projective resolution of $N \in obj(\mathcal{C})$, then the former equals the unenriched Ext-group $\operatorname{Ext}^n_{\mathcal{C}}(M, N)$ according to [15, p. 166] and [42, Section 10.7]. The proper setting to illustrate this are derived categories. The idea and basic properties of the derived category $\mathcal{D}(\mathcal{C})$ of \mathcal{C} are summarised on pages 143–144 in S. I. Gelfand and Yu. I. Manin's book on homological algebra [15]. Morally, objects of $\mathcal{D}(\mathcal{C})$ are chain complexes $(M_k)_{n\in\mathbb{Z}}$ in \mathcal{C} and morphisms are chain maps where any chain map $(f_k : M_k \to N_k)_{k \in \mathbb{Z}}$ is taken to be an isomorphism if for every $n \in \mathbb{Z}$ the induced morphism in homology $H_n(f_{\bullet}) : H_n(M_{\bullet}) \to H_n(N_{\bullet})$ is an isomorphism. Every $A \in obj(\mathcal{C})$ can be turned into an object $\iota(A)_{\bullet} \in obj(\mathcal{D}(\mathcal{C}))$ where $\iota(A)_0 = A, \,\iota(A)_k = 0$ for $k \neq 0$ and all boundary maps are zero. If $(A_k)_{k \in \mathbb{N}_0}$ is a projective resolution of A, then we can extend it to a chain complex $(A_k)_{k\in\mathbb{Z}}$ by setting $A_k = 0$ for k < 0 and all other boundary maps to be zero. By means of the augmentation map $A_0 \to A$ we see that $\iota(A)_{\bullet}$ is isomorphic to A_{\bullet} in $\mathcal{D}(\mathcal{C})$. As we index our projective resolutions to have positive degrees, our notation in the following agrees with the one found in [42] and disagrees with the one in [15]. According to [15, p. 166] and [42, p. 399], one can define unenriched Ext-groups as $\operatorname{Ext}^n_{\mathcal{C}}(A,B) := \operatorname{Hom}_{\mathcal{D}(\mathcal{C})}(\iota(A)[n]_{\bullet},\iota(B)_{\bullet})$. Note that $\operatorname{Ext}^n_{\mathcal{C}}(A, B) \cong \operatorname{Hom}_{\mathcal{D}(\mathcal{C})}(A[n]_{\bullet}, \iota(B)_{\bullet})$. Since homotopic chain maps are identified in $\mathcal{D}(\mathcal{C})$ [15, p. 159], this agrees with the definition of $\operatorname{Ext}^n_{\mathcal{C}}(-,B)$ as the nth right derived functors of the unenriched Hom-functor $\operatorname{Hom}_{\mathcal{C}}(-, B)$. We refer the reader to Sections 2.4 and 2.5 of [42] for a thorough account on derived categories. In order to distinguish the two emerging notions of Ext-functors, we introduce the following notation.

Notation 6.1 From this point on, if $(A_k)_{k \in \mathbb{Z}}$ is a chain complex in \mathcal{C} , then it is assumed that $(A_k)_{k \in \mathbb{N}_0}$ is a projective resolution of an object A in \mathcal{C} and $A_k = 0$ for k < 0. We write

$$\mathcal{E}xt^n_{\mathcal{C}}(A,B) := \operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})/\operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$$

whenever we consider the n^{th} Ext-group as arising from chain maps modulo chain homotopy. Analogously, we write

$$\widehat{\mathcal{Ext}}^n_{\mathcal{C}}(A,B) := \widehat{\operatorname{Hom}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet}) / \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet}) \,.$$

These agree with the n^{th} cohomology group of the hypercohomology complex and of the Vogel complex respectively. If $a_k : A_k \to A_{k-1}$ denote the boundary maps of A_{\bullet} , then we continue to write

$$\operatorname{Ext}^{n}_{\mathcal{C}}(A, B) = \operatorname{Ker}(\operatorname{Hom}_{\mathcal{C}}(a_{n+1}, B)) / \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_{n}, B))$$

for the n^{th} Ext-group defined as the n^{th} derived functor of Hom_{\mathcal{C}}(-, B).

For morphisms $f: B \to C$ and $g: C \to A$ in \mathcal{C} , let us construct $\mathcal{E}xt^n_{\mathcal{C}}(A, f)$ and $\mathcal{E}xt^n_{\mathcal{C}}(g, B)$ as well as $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, f)$ and $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(g, B)$. Consider projective resolutions $A_{\bullet}, B_{\bullet}, C_{\bullet}$ of A, B, C. By the Comparison Theorem [42, Theorem 2.2.6] there is a chain map $f_{\bullet}: B_{\bullet} \to C_{\bullet}$ that is unique up to chain homotopy such that the diagram

$$\dots \longrightarrow B_2 \longrightarrow B_1 \longrightarrow B_0 \longrightarrow B \downarrow_{f_2} \qquad \downarrow_{f_1} \qquad \downarrow_{f_0} \qquad \downarrow_f \dots \longrightarrow C_2 \longrightarrow C_1 \longrightarrow C_0 \longrightarrow C$$

commutes. There is an analogous chain map $g_{\bullet} : C_{\bullet} \to A_{\bullet}$ that is unique up to chain homotopy. Define $g[n]_{\bullet} : C[n]_{\bullet} \to A[n]_{\bullet}$ by $g[n]_k := g_{k+n}$ for every $k \in \mathbb{Z}$ [42, p. 10]. Since the category of chain complexes $Ch(\mathcal{C})$ with chain maps as morphisms forms an abelian category [42, Theorem 1.2.3], there are homomorphisms of abelian groups

$$\operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, f_{\bullet}) : \operatorname{Hom}_{\mathcal{C}}(A[n]_{\bullet}, B_{\bullet}) \to \operatorname{Hom}_{\mathcal{C}}(A[n]_{\bullet}, C_{\bullet}), \ \varphi_{\bullet} \mapsto f_{\bullet} \circ \varphi_{\bullet} \text{ and} \\ \operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(g[n]_{\bullet}, B_{\bullet}) : \operatorname{Hom}_{\mathcal{C}}(A[n]_{\bullet}, B_{\bullet}) \to \operatorname{Hom}_{\mathcal{C}}(C[n]_{\bullet}, B_{\bullet}), \ \varphi_{\bullet} \mapsto \varphi_{\bullet} \circ g[n]_{\bullet}$$

It follows by definition and [42, p. 5] that any $\operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$ is an abelian group and that the maps

$$\widehat{\operatorname{Hom}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, f_{\bullet}) : \widehat{\operatorname{Hom}}_{\mathcal{C}}(A[n]_{\bullet}, B_{\bullet}) \to \widehat{\operatorname{Hom}}_{\mathcal{C}}(A[n]_{\bullet}, C_{\bullet}), \ \varphi_{\bullet} \mapsto f_{\bullet} \circ \varphi_{\bullet} \text{ and} \\
\widehat{\operatorname{Hom}}_{\operatorname{Ch}(\mathcal{C})}(g[n]_{\bullet}, B_{\bullet}) : \widehat{\operatorname{Hom}}_{\mathcal{C}}(A[n]_{\bullet}, B_{\bullet}) \to \widehat{\operatorname{Hom}}_{\mathcal{C}}(C[n]_{\bullet}, B_{\bullet}), \ \varphi_{\bullet} \mapsto \varphi_{\bullet} \circ g[n]_{\bullet}$$

are also homomorphisms of abelian groups.

Definition 6.2 The above descend to homomorphisms

$$\mathcal{E}xt^{n}_{\mathcal{C}}(A,f): \mathcal{E}xt^{n}_{\mathcal{C}}(A,B) \to \mathcal{E}xt^{n}_{\mathcal{C}}(A,C)$$
$$\varphi_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet},B_{\bullet}) \mapsto f_{\bullet} \circ \varphi_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet},C_{\bullet})$$

and

$$\begin{aligned} & \mathcal{E}xt^n_{\mathcal{C}}(g,B): \ \mathcal{E}xt^n_{\mathcal{C}}(A,B) \to \mathcal{E}xt^n_{\mathcal{C}}(C,B) \\ & \varphi_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet},B_{\bullet}) \mapsto \varphi_{\bullet} \circ g[n]_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(C[n]_{\bullet},B_{\bullet}) \,. \end{aligned}$$

The homomorphisms

$$\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,f):\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,B)\to\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,C) \text{ and } \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(g,B):\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,B)\to\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(C,B)$$

are defined analogously.

Although F. Goichot only works with modules over a ring, we generalise his remark from [17, p. 41] that $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, B)$ does not depend on choices of projective resolutions of A and B.

Proposition 6.3 For any $n \in \mathbb{Z}$, both

$$\mathcal{E}xt^n_{\mathcal{C}}(-,-): \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \to \mathbf{Ab} \text{ and } \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(-,-): \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \to \mathbf{Ab}$$

form well defined functors that are additive in both variables with respect to the induced morphisms from Definition 6.2.

Proof Let B_{\bullet} , B'_{\bullet} be two projective resolutions of B. According to [42, Theorem 2.2.6] there are chain maps $\iota_{\bullet} : B_{\bullet} \to B'_{\bullet}$ and $\iota'_{\bullet} : B'_{\bullet} \to B_{\bullet}$ lifting $\mathrm{id}_B : B \to B$ that are unique up to chain homotopy. In particular, we see that

$$\iota'_{\bullet} \circ \iota_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(B_{\bullet}, B_{\bullet}) = \operatorname{id} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(B_{\bullet}, B_{\bullet}) \text{ and } \iota_{\bullet} \circ \iota'_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(B'_{\bullet}, B'_{\bullet}) = \operatorname{id} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(B'_{\bullet}, B'_{\bullet}).$$

This constitutes a direct proof that the abelian groups $\mathcal{E}xt_{\mathcal{C}}(A, B)$ and $\widehat{\mathcal{E}xt}_{\mathcal{C}}(A, B)$ do not depend on a choice of projective resolution of B. We may conclude analogously that they do not depend on the choice of projective resolution of A. If $f: A \to B$ and $g: C \to A$ are morphisms in \mathcal{C} , then this also proves that $\mathcal{E}xt_{\mathcal{C}}(A, f)$ and $\widehat{\mathcal{E}xt}_{\mathcal{C}}(A, f)$ as well as $\mathcal{E}xt_{\mathcal{C}}(g, B)$ and $\widehat{\mathcal{E}xt}_{\mathcal{C}}(g, B)$ are well defined homomorphisms. We conclude from [42, p. 5] that $\mathcal{E}xt^n_{\mathcal{C}}(-,-)$ and $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(-,-)$ are additive in both variables.

6.2 Connecting homomorphisms through chain maps

We construct an explicit isomorphism from the Ext-group $\operatorname{Ext}^n_{\mathcal{C}}(A, B)$ to $\mathcal{Ext}^n_{\mathcal{C}}(A, B)$ which allows us to determine connecting homomorphisms for $\mathcal{Ext}^n_{\mathcal{C}}(A, -)$ and $\widehat{\mathcal{Ext}}^\bullet_{\mathcal{C}}(A, -)$ in terms of (almost) chain maps. The construction of this isomorphism is of similar nature to pages 63–64 of C. A. Weibel's book [42] where he compares the chain complex $\operatorname{Hom}_{\mathcal{C}}(A_{\bullet}, B)$ to $\operatorname{Hyp}_{\mathcal{C}}(A_{\bullet}, B_{\bullet})_{\bullet,\bullet}$ seen as a double complex. However, as our techniques differ and we have not found this isomorphism in the literature otherwise, we provide full details.

Given that Ext-groups vanish in negative degrees, let $n \in \mathbb{N}_0$. Let $(A_{\bullet}, a_{\bullet})$ be a projective resolution of A, $(B_{\bullet}, b_{\bullet})$ is a projective resolution of B and $\varphi \in \text{Ker}(\text{Hom}_{\mathcal{C}}(a_{k+1}, B))$. By Diagram 5.4 and the Comparison Theorem [42, Theorem 2.2.6], there exists a chain map $(\varphi[n]_k : A[n]_k \to B_k)_{k \in \mathbb{N}_0}$ that is unique up to chain homotopy making the diagram

$$\dots \longrightarrow A_{n+2} \xrightarrow{(-1)^n a_{n+2}} A_{n+1} \xrightarrow{(-1)^n a_{n+1}} A_n \xrightarrow{\pi_n} \widetilde{A}_n$$

$$\downarrow^{\varphi[n]_2} \qquad \downarrow^{\varphi[n]_1} \qquad \downarrow^{\varphi[n]_0} \xrightarrow{\varphi} \qquad \downarrow^{\alpha_n(B)(\varphi)}$$

$$\dots \longrightarrow B_2 \xrightarrow{b_2} B_1 \xrightarrow{b_1} B_0 \xrightarrow{b_1} B_0 \xrightarrow{b} B$$


$$(6.1)$$

commute. Since $B_k = 0$ for k < 0, there is an extension $(\varphi[n]_k : A[n]_k \to B_k)_{k \in \mathbb{Z}}$. Hence the map

$$\zeta'_{n} : \operatorname{Ker}(\operatorname{Hom}_{\mathcal{C}}(a_{n+1}, B)) \to \mathcal{E}xt^{n}_{\mathcal{C}}(A, B)$$
$$\varphi \mapsto \varphi[n]_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$$

is a well defined homomorphism of abelian groups.

Assume that $n \ge 1$ and $\psi \in \text{Im}(\text{Hom}_{\mathcal{C}}(a_n, B))$. Then there is a morphism $\psi : A_{n-1} \to B$ such that $\varphi = \psi \circ a_n$. Because A_{n-1} is projective and $b : B_0 \to B$ an epimorphism, there $e_{n-1} : A_{n-1} \to B_0$ such that $\psi = b \circ e_{n-1}$. Knowing that $a_n \circ a_{n+1} = 0$, we see that the diagram




is commutative. Set $\varphi[n]_{\bullet} \in \operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$ to be zero in every degree except for k = 0 where $\varphi[n]_0 = e_{n-1} \circ a_n$. Similarly, set $e[n]_{\bullet} : A[n]_{\bullet} \to B_{\bullet+1}$ to be also zero in every degree except for k = -1 where $e[n]_{-1} = (-1)^n e_{n-1} : A_{n-1} \to B_0$. Then $e[n]_{\bullet}$ is a chain homotopy between $\varphi[n]_{\bullet}$ and the zero chain map. Therefore, ζ'_n descends to a homomorphism

$$\zeta_n : \operatorname{Ext}^n_{\mathcal{C}}(A, B) \to \mathcal{E}xt^n_{\mathcal{C}}(A, B)$$
$$\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_n, B)) \mapsto \varphi[n]_{\bullet} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$$

As we have assumed $n \ge 1$ for this, we define $\zeta_0 := \zeta'_0$ where we exploit the fact that $\operatorname{Ker}(\operatorname{Hom}_{\mathcal{C}}(a_1, B)) = \operatorname{Ext}^0_{\mathcal{C}}(A, B)$. Given that unenriched Ext-functors of negative degree vanish, we set $\zeta_n := 0$ for n < 0.

Proposition 6.4 For every $n \in \mathbb{Z}$, $\zeta_n : \operatorname{Ext}^n(A, B) \to \mathcal{E}xt^n_{\mathcal{C}}(A, B)$ is a isomorphism that is natural in both variables A and B.

Proof Since ζ_n is a natural isomorphism for n < 0, assume that $n \in \mathbb{N}_0$ for the rest of the proof. To prove that ζ_n is surjective, let $\psi_{\bullet+n} \in \operatorname{Hom}_{\mathcal{C}}(A[n]_{\bullet}, B_{\bullet})$. As the diagram



commutes, $B_k = \text{for } k < 0 \text{ and } b_1 \circ b = 0$, we infer that $b \circ \psi_n \in \text{Ker}(\text{Hom}_{\mathcal{C}}(a_{n+1}, B))$ and that


$$\zeta_n(b \circ \psi_0 + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_n, B))) = \psi_{\bullet+n} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet}) + \mathcal{O}(A[n]_{\bullet}, B_{\bullet}) + \mathcal{O}(A[n]_{\bullet}) + \mathcal{O}(A[n]_{\bullet}$$

Let us show that ζ_n is injective. Assume first that $n \geq 1$. If $\varphi \in \text{Ker}(\text{Hom}_{\mathcal{C}}(a_{n+1}, B))$ is such that $\zeta_n(\varphi + \text{Im}(\text{Hom}_{\mathcal{C}}(a_n, B))) = 0$, then there are morphisms $\varphi_n : A_n \to B_0$, $e_n : A_n \to B_1$ and $e_{n-1} : A_{n-1} \to B_0$ such that

$$\varphi = b \circ \varphi_n = b \circ \left(e_{n-1} \circ (-1)^n a_n + b_1 \circ e_n \right) = (-1)^n (b \circ e_n) \circ a_n.$$

In particular, $\varphi \in \text{Im}(\text{Hom}_{\mathcal{C}}(a_n, B))$ and ζ_n is injective. Suppose that n = 0 and that $\varphi \in \text{Ker}(\text{Hom}_{\mathcal{C}}(a_1, B))$ is such that $\zeta_0(\varphi) = 0$. Because $A_{-1} = 0$ and $a_0 = 0$, we conclude that $\varphi = 0$ and that ζ_0 is also injective.

Let us demonstrate that ζ_n is natural in both variables. Let $g: C \to A$ and $f: B \to D$ be a morphism in \mathcal{C} and $\varphi \in \text{Ker}(\text{Hom}_{\mathcal{C}}(a_{n+1}, B))$. Taking lifts $f_{\bullet}: B_{\bullet} \to D_{\bullet}$ and $g[n]_{\bullet}: C[n]_{\bullet} \to A[n]_{\bullet}$ as needed in Definition 6.2, we see that the diagram



commutes. This shows that

$$\zeta_n \Big(\operatorname{Ext}^n_{\mathcal{C}}(A, f) \big(\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_{n+1}, B)) \big) \Big) = \mathcal{E}xt^n_{\mathcal{C}}(A, f) \Big(\zeta_n \big(\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_{n+1}, B)) \big) \Big)$$

and that

$$\zeta_n\Big(\operatorname{Ext}^n_{\mathcal{C}}(g,B)\big(\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_{n+1},B))\big)\Big) = \mathcal{E}xt^n_{\mathcal{C}}(g,B)\Big(\zeta_n\big(\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_{n+1},B))\big)\Big).$$

In short, $\zeta_n \circ \operatorname{Ext}^n_{\mathcal{C}}(A, -) = \mathcal{E}xt^n_{\mathcal{C}}(A, -) \circ \zeta_n$ and $\zeta_n \circ \operatorname{Ext}^n_{\mathcal{C}}(-, B) = \mathcal{E}xt^n_{\mathcal{C}}(-, B) \circ \zeta_n$. \Box

Our construction of the desired connecting homomorphisms for $\mathcal{E}xt^{\bullet}_{\mathcal{C}}(A, -)$ and $\widehat{\mathcal{E}xt}^{\bullet}_{\mathcal{C}}(A, -)$ is similar to the definitions found in [13, p. 2202]. We consider a projective resolution $(D_{\bullet}, d_{\bullet})$ of D. For any $k \in \mathbb{N}_0$ define the chain complex $(D^{+1}_{\bullet}, d^{+1}_{\bullet})$ by setting $(D^{+1}_m, d^{+1}_m)_{m \in \mathbb{N}} := (D_{m+1}, d_{m+1})_{m \in \mathbb{N}}, D^{+1}_0 := D_1$ and $D^{+1}_m := 0$ for m < 0. Observe that D^{+1}_{\bullet} is a projective resolution of \widetilde{D}_1 . We define a chain map $\pi^n_{\bullet} : D_{\bullet} \to D^{+1}[-1]_{\bullet}$ by setting

$$\pi_m^n := \begin{cases} (-1)^{m-1+n} \mathrm{id}_{D_m} : D_m \to D_m & \text{if } m \ge 1 \\ 0 & \text{if } m \le 0 \end{cases}$$

Pictorially

$$\dots \longrightarrow D_4 \xrightarrow{d_4} D_3 \xrightarrow{d_3} D_2 \xrightarrow{d_2} D_1 \xrightarrow{d_1} D_0$$

$$\downarrow^{(-1)^{n+3}\text{id}} \downarrow^{(-1)^{n+2}\text{id}} \downarrow^{(-1)^{n+1}\text{id}} \downarrow^{(-1)^n\text{id}} \downarrow \qquad (6.2)$$

$$\dots \longrightarrow D_4 \xrightarrow{-d_4} D_3 \xrightarrow{-d_3} D_2 \xrightarrow{-d_2} D_1 \xrightarrow{-d_2} 0$$

As is noted in [15, p. 166], we can equally well express the $(n + 1)^{st}$ Ext-group as

$$\mathcal{E}xt_{\mathcal{C}}^{n+1}(A,D) = \operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, D[-1]_{\bullet})/\operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, D[-1]_{\bullet})$$

with an analogous statement true for $\widehat{\mathcal{Ext}}_{\mathcal{C}}^{n+1}(A, D)$. The following definition bears similarity with the connecting homomorphisms $\varepsilon^{-m}: S^{-m}T \to S^{-m+1}T$ established at the start of Section 3 and with the morphisms used in the construction of the connecting homomorphisms of the naïve construction found in Lemma 5.9.

Definition 6.5 We set

$$\Delta^{n} : \mathcal{E}xt^{n}_{\mathcal{C}}(A, D) \to \mathcal{E}xt^{n+1}_{\mathcal{C}}(A, \widetilde{D}_{1})$$
$$\varphi_{\bullet+n} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, D_{\bullet}) \mapsto \pi^{n}_{\bullet} \circ \varphi_{\bullet+n} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, D^{+1}[-1]_{\bullet})$$

to be the (proposed) connecting homomorphism associated too the short exact sequence $0 \to \widetilde{D}_1 \to D_0 \to D \to 0$. Analogously, we define the (proposed) connecting homomorphism $\widehat{\Delta}^n : \widehat{\mathcal{Ext}}^n_{\mathcal{C}}(A, D) \to \widehat{\mathcal{Ext}}^{n+1}_{\mathcal{C}}(A, \widetilde{D}_1).$

If $0 \to B \to C \to D \to 0$ is another short exact sequence in \mathcal{C} , then consider the lift $h^*: \widetilde{D}_1 \to B$ of $\mathrm{id}_D: D \to D$ from Diagram 3.2 and a chain map $h^*_{\bullet+1}: D^{+1}_{\bullet} \to B_{\bullet}$ as in [42, Theorem 2.2.6]. We define the (proposed) connecting homomorphism for this short exact sequence as

$$\Delta^{n} : \mathcal{E}xt^{n}_{\mathcal{C}}(A, D) \to \mathcal{E}xt^{n+1}_{\mathcal{C}}(A, B)$$

$$\varphi_{\bullet+n} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, D_{\bullet}) \mapsto h^{*}[-1]_{\bullet+1} \circ \pi^{n}_{\bullet} \circ \varphi_{\bullet+n} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B[-1]_{\bullet})$$

where the definition of $\widehat{\Delta}^n : \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, D) \to \widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A, B)$ is analogous.


Because the maps in the above definition are defined by compositions of (almost) chain maps, they are homomorphisms of abelian groups. However, they are not well defined a priori because they depend on a choice of morphism $h^* : \widetilde{D}_1 \to B$.

Lemma 6.6 The family $(\mathcal{E}xt^{\bullet}_{\mathcal{C}}(A, -), \Delta^{\bullet})$ forms a cohomological functor. Furthermore, $\zeta_{\bullet} : (\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -), \delta^{\bullet}) \to (\mathcal{E}xt^{\bullet}_{\mathcal{C}}(A, -), \Delta^{\bullet})$ is an isomorphism of cohomological functors.

Proof It suffices to prove that $\zeta_n : \operatorname{Ext}^n_{\mathcal{C}}(A, -) \to \mathcal{E}xt^{\bullet}_{\mathcal{C}}(A, -)$ commutes with the (proposed) connecting homomorphisms. Let $0 \to B \to C \to D \to 0$ be a short exact sequence in \mathcal{C} and let $h^* : \widetilde{D}_1 \to B$ be a lift of id_D as before. We have seen at the start of the proof of Lemma 5.9 that the associated connecting homomorphism factors through the one associated to $0 \to \widetilde{D}_1 \to D_0 \to D \to 0$ as

$$\delta^n : \operatorname{Ext}^n_{\mathcal{C}}(A, D) \xrightarrow{\delta^n} \operatorname{Ext}^{n+1}_{\mathcal{C}}(A, \widetilde{D}_1) \xrightarrow{\operatorname{Ext}^{n+1}_{\mathcal{C}}(A, h^*)} \operatorname{Ext}^{n+1}_{\mathcal{C}}(A, B).$$

Thus, the left hand triangle in the diagram



commutes. By Definition 6.2 and 6.5, the right hand triangle triangle also commutes. According to Proposition 6.4, the bottom trapezium is commutative. The top trapzium can be seen as a generalisation of Proposition 1.1 from [13] where we use a different method to prove that it commutes. Let $\varphi \in \text{Ker}(\text{Hom}_{\mathcal{C}}(a_n, D))$. We explain how one can combine Diagram 5.6 and 6.1 as

$$\dots \longrightarrow A_{n+2} \xrightarrow{(-1)^n a_{n+2}} A_{n+1} \xrightarrow{\pi_{n+1}} \widetilde{A}_{n+1} \xrightarrow{\iota_{n+1}} A_n \xrightarrow{\pi_n} \widetilde{A}_n$$

$$\downarrow \varphi[n]_2 \qquad \qquad \downarrow \varphi[n]_1 \xrightarrow{\Phi} \downarrow \qquad \qquad \downarrow \varphi[n]_0 \xrightarrow{\varphi} \downarrow \alpha_{n+k}(B)(\varphi) \qquad (6.4)$$

$$\dots \longrightarrow D_2 \xrightarrow{d_2} D_1 \xrightarrow{\pi_1} \widetilde{D}_1 \xrightarrow{\iota_1} D_0 \xrightarrow{\pi_0} D$$

The right most morphisms up to $\Phi : A_{n+1} \to \widetilde{B}_1$ are taken from Diagram 5.6 while $\varphi : A_n \to D$ and $\varphi[n]_{\bullet} : A[n]_{\bullet} \to D_{\bullet}$ are the morphisms as they occur in Diagram 6.1. In particular,

$$\delta^n (\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_n, D))) = \Phi + (\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_n, \widetilde{D}_1)).$$

However, note that $\iota_{n+1} \circ \pi_{n+1} = a_{n+1}$ might not agree with $(-1)^n a_{n+1} : A_{n+1} \to A_n$. This together with the construction of the Diagram 5.6 and the proof of the Comparison from [42, p. 36] implies that $\Phi = (-1)^n \pi_1 \circ \varphi[n]_1$. Therefore, we can choose the representative $\Phi[n+1]_{\bullet} : A[n+1] \to \widetilde{D}_{\bullet}^{+1}$ of a lift of Φ as in Diagram 6.1 which is defined as

$$\Phi[n+1]_m := \begin{cases} (-1)^{n+m} \varphi[n]_{m+1} : A_{m+n+1} \to D_{m+1} & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$

The alternating signs are due to the fact that the boundary maps of $A[n]_{\bullet}$ and A[n+1] have opposite signs. We conclude by Diagram 6.2, Definition 6.5 and Diagram 6.4 that

$$\zeta_{n+1}\Big(\delta^n\big(\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_n, D))\big)\Big) = \Delta^n\Big(\zeta^n\big(\varphi + \operatorname{Im}(\operatorname{Hom}_{\mathcal{C}}(a_n, D))\big)\Big).$$

In particular, the top trapezium and with it all of Diagram 6.3 commutes. As for Theorem 5.10, we can now reiterate the proof of Theorem 4.5 with Diagram 6.3, the fact that $(\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -), \delta^{\bullet})$ is a cohomological functor and that $\zeta_{\bullet} : \operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -) \to \mathcal{E}xt^{\bullet}_{\mathcal{C}}(A, -)$ is a natural isomorphism by Proposition 6.4.

6.3 Isomorphism of cohomology groups

In this subsection we construct for every $n \in \mathbb{Z}$ a natural isomorphism from the completed unenriched Ext-functor $\operatorname{Ext}_{\operatorname{Res},\mathcal{C}}^n(A,-)$ defined through the resolution construction to $\widehat{\operatorname{Ext}}_{\mathcal{C}}^n(A,-)$. In order to do so, we first reformulate the former completed Ext-functor.

Definition 6.7 For any $B \in obj(\mathcal{C})$, $f : B \to C$ and $n \in \mathbb{Z}$ define

$$\mathcal{E}xt^{n}_{Res,\mathcal{C}}(A,B) := \lim_{k \in \mathbb{N}_{0}} (\mathcal{E}xt^{n+k}_{\mathcal{C}}(A,\widetilde{B}_{k}), \Delta^{n+k})$$

and

$$\mathcal{E}xt^n_{Res,\mathcal{C}}(A,f) := \lim_{k \in \mathbb{N}_0} \mathcal{E}xt^{n+k}_{\mathcal{C}}(A,\widetilde{f}_k) : \mathcal{E}xt^n_{\mathcal{C}}(A,B) \to \mathcal{E}xt^n_{\mathcal{C}}(A,C)$$

where $\tilde{f}_k : \tilde{B}_k \to \tilde{C}_k$ are taken as in Proposition 4.2. For any short exact sequence $0 \to B \to C \to D \to 0$ we set

$$\widetilde{\Delta}^{n} := \lim_{k \in \mathbb{N}_{0}} (-1)^{k} \Delta^{n+k} : \mathcal{E}xt^{n}_{Res,\mathcal{C}}(A,D) \to \mathcal{E}xt^{n+1}_{Res,\mathcal{C}}(A,B)$$

where $\Delta^{n+k} : \mathcal{E}xt^{n+k}_{\mathcal{C}}(A, \widetilde{D}_k) \to \mathcal{E}xt^{n+k+1}_{\mathcal{C}}(A, \widetilde{B}_k)$ are taken as in Diagram 4.10. Using the isomorphisms of cohomological functors $\zeta_{n+k} : \operatorname{Ext}^{n+k}_{\mathcal{C}}(A, -) \to \mathcal{E}xt^{n+k}_{\mathcal{C}}(A, -)$ from Lemma 6.6, we can extend the natural isomorphisms

$$\zeta^{n} := \varinjlim_{k \in \mathbb{N}_{0}} \zeta_{n+k} : \operatorname{Ext}^{n}_{\operatorname{Res}, \mathcal{C}}(A, -) \to \mathcal{E}xt^{n}_{\operatorname{Res}, \mathcal{C}}(A, -)$$

to ζ^{\bullet} : $(\operatorname{Ext}_{\operatorname{Res},\mathcal{C}}^{\bullet}(A,-),\widetilde{\delta}^{\bullet}) \to (\operatorname{\mathcal{E}xt}_{\operatorname{Res},\mathcal{C}}^{\bullet}(A,-),\widetilde{\Delta}^{\bullet})$ which is an isomorphism of cohomological functors.

We only consider the cofinal system $(\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{B}_{2k}), \Delta^{n+2k+1} \circ \Delta^{n+2k})_{k \in \mathbb{N}_0}$ and specifically form $\mathcal{E}xt_{Res,\mathcal{C}}^n(A, B)$ as its direct limit in order to ease notation when constructing a homomorphism from any cohomology group $\mathcal{E}xt_{Res,\mathcal{C}}^n(A, B)$ to $\widehat{\mathcal{E}xt}_{\mathcal{C}}^n(A, B)$. Nonetheless, we require

Notation 6.8 If $(B_{\bullet}, b_{\bullet})$ is a projective resolution of B and $k \in \mathbb{N}_0$, then we define the projective resolution $(B_{\bullet}^{+k}, b_{\bullet}^{+k})_{m \in \mathbb{Z}}$ of \widetilde{B}_k by declaring for $m \in \mathbb{Z}$ that

$$(B_m^{+k}, b_{m+1}^{+k}) := \begin{cases} (B_{m+k}, b_{m+1+k} : B_{m+1+k} \to B_{m+k}) & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$

Lemma 6.9 There is an isomorphism $\vartheta^n(B) : \mathcal{E}xt^n_{\operatorname{Res},\mathcal{C}}(A,B) \to \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,B)$ for any $n \in \mathbb{Z}$.

Proof For any $k \in \mathbb{N}_0$ we construct a map

$$\vartheta_n^{2k}(B) : \mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{B}_{2k}) \to \widehat{\mathcal{E}xt}_{\mathcal{C}}^n(A, B)$$
$$\varphi_{\bullet+n+2k} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k]_{\bullet}, B_{\bullet}^{+2k}) \mapsto \Phi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$$

as follows. For a representative $\varphi_{\bullet+n+2k} \in \operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k]_{\bullet}, B_{\bullet}^{+2k})$ of an element in $\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{B}_{2k})$ we define $\Phi_{\bullet+n} : A[n]_{\bullet} \to B_{\bullet}$ by


$$\Phi_{m+n} := \begin{cases} \varphi_{m+n} : A_{m+n} \to B_m & \text{if } m \ge 2k \\ 0 & \text{if } m < 2k \end{cases}$$

As the boundary maps of A[n] and A[n+2k] agree modulo a shift of 2k in their indices, all squares of the form

$$\begin{array}{c} A_{m+n} & \xrightarrow{\Phi_{m+n}} & B_m \\ \downarrow^{(-1)^n a_{m+n}} & \downarrow^{b_m} \\ A_{m-1+n} & \xrightarrow{\Phi_{n+m-1}} & B_{m-1} \end{array}$$

commute for $m \in \mathbb{Z} \setminus \{2k\}$. We conclude that $\Phi_{\bullet+n} \in \widehat{\operatorname{Hom}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$. If the chain map $\varphi_{\bullet+n+2k}$ is nullhomotopic, then $\Phi_{\bullet+n}$ is genuinely nullhomotopic in any degree $m \in \mathbb{Z} \setminus \{2k-1\}$, hence nullhomotopic as an almost chain map. Thus, ϑ_n^{2k} is a well defined map. It follows by construction that it is also a homomorphism of abelian groups.

We argue that the triangle



commutes. By Definition 6.5, Δ^{n+2k} and Δ^{n+2k+2} are constructed using the chain maps $\pi^n_{\bullet}: B^{+2k}_{\bullet} \to B^{+2k+1}[-1]_{\bullet}$ and $\pi^{n+1}[-1]_{\bullet}: B^{+2k+1}[-1]_{\bullet} \to B^{+2k+2}[-2]_{\bullet}$ which are pictorially rendered in Diagram 6.2. Then $\pi^{n+1}[-1]_{\bullet} \circ \pi^n_{\bullet}$ is given by

$$\pi^{n+1}[-1]_m \circ \pi^n_m = \begin{cases} \operatorname{id}_{B_m} : B_m \to B_m & \text{if } m \ge 2\\ 0 & \text{if } m \le 1 \end{cases}$$

since $(-1)^{(m-1)-1+(n+1)} \cdot (-1)^{m-1+n} = 1$. Hence, the m^{th} degree of an almost chain map representative of the image of an element $\varphi_{\bullet+n+2k} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k], B_{\bullet}^{+2k})$ in $\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{B}_{2k})$ under $\Delta^{n+2k+1} \circ \Delta^{n+2k}$ is

$$\begin{cases} \varphi_{m+n+2k+2} : A_{m+n+2k+2} \to B_{m+2k+2} & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$
(6.6)

We infer by definition of $\vartheta_n^{2k}(B)$ and $\vartheta_n^{2k+2}(B)$ that

$$\vartheta_n^{2k}(B) \big(\varphi_{\bullet+n+2k} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k], B_{\bullet}^{+2k}) \big) = \\ \big(\vartheta_n^{2k+2} \circ \Delta^{n+2k+1} \circ \Delta^{n+2k} \big) \big(\varphi_{\bullet+n+2k} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k], B_{\bullet}^{+2k}) \big)$$

The resulting homomorphism in the direct limit of the above commuting triangles is

$$\vartheta^n(B) := \varinjlim_{k \in \mathbb{N}_0} \vartheta_n^{2k}(B) : \ \mathcal{E}xt^n_{\operatorname{Res},\mathcal{C}}(A,B) \to \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,B) \,.$$

To show that it is surjective, let $\Phi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$ be in $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, B)$. Since it is an almost chain map, there is $k \in \mathbb{N}_0$ such that $(\Phi_{m+n} : A_{m+n} \to B_m)_{m \geq 2k}$ is a chain map. Defining $\varphi_{\bullet+n+2k} : A[n+2k]_{\bullet} \to B^{+2k}_{\bullet}$ by setting

$$\varphi_{m+n+2k} := \begin{cases} \Phi_{m+n+2k} : A_{m+n+2k} \to B_{m+2k} & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$

gives an element in $\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{B}_{2k})$ that $\vartheta_n^{2k}(B)$ maps to $\Phi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$. To prove that $\vartheta^n(B)$ is injective, let $x \in \varinjlim \mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{B}_{2k})$ with $\vartheta^n(B)(x) = 0$. According to Proposition 5.6, there exists $k \in \mathbb{N}_0$ and an element

$$\psi_{\bullet+n+2k} \in \operatorname{Hom}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k]_{\bullet}, B_{\bullet}^{+2k})$$

mapping to x in the direct limit that lies in the kernel of the homomorphism $\vartheta_n^{2k}(B)$. If we construct an almost chain map representative $\Psi_{\bullet+n} : A[n]_{\bullet} \to B_{\bullet}$ from $\psi_{\bullet+n+k}$ as above, then $\Psi_{\bullet+n} \in \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$. Therefore, there is $K \geq k$ such that $(\Psi_m)_{m \geq 2K}$ is nullhomotopic as a chain map and not just as an almost chain map. The chain map $\overline{\psi}_{\bullet+n+2K} : A[n+2K]_{\bullet} \to B_{\bullet}^{+2K}$ that we define by

$$\overline{\psi}_{m+n+2K} := \begin{cases} \Psi_{m+n+2K} = \psi_{m+n+2K} : A_{m+n+2K} \to B_{m+2K} & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$


lies in $\operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n+2K]_{\bullet}, B_{\bullet}^{+2K})$. By Equation 6.6, $\overline{\psi}_{\bullet+n+2K}$ is a representative of the image of $\psi_{\bullet+n+2k} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})}(A[n+2k]_{\bullet}, B_{\bullet}^{+2k})$ under $\Delta^{n+2K-1} \circ \ldots \circ \Delta^{n+2k}$, which also maps to the element x in the direct limit. As x = 0, this implies that $\vartheta^n(B)$ is injective and thus an isomorphism. \Box

Proposition 6.10 For every $n \in \mathbb{Z}$ the isomorphism $\vartheta^n(-)$ is natural. That is, for every morphism $f: B \to C$ in \mathcal{C} the square

$$\begin{aligned} \mathcal{E}xt^{n}_{Res,\mathcal{C}}(A,B) & \xrightarrow{\vartheta^{n}(B)} \widehat{\mathcal{E}xt}^{n}_{\mathcal{C}}(A,B) \\ & \downarrow^{\mathcal{E}xt^{n}_{Res,\mathcal{C}}(A,f)} & \downarrow^{\widehat{\mathcal{E}xt}^{n}_{\mathcal{C}}(A,f)} \\ \mathcal{E}xt^{n}_{Res,\mathcal{C}}(A,C) & \xrightarrow{\vartheta^{n}(C)} \widehat{\mathcal{E}xt}^{n}_{\mathcal{C}}(A,C) \end{aligned}$$

commutes.

Proof Let $(\widetilde{f}_k : \widetilde{M}_k \to \widetilde{N}_k)_{k \in \mathbb{N}_0}$ be a sequence of morphism as arising from Diagram 4.8. By Definition 6.7 it suffices to consider the diagram



The left hand side of the prism commutes because Δ^{n+2k} and Δ^{n+2k+1} are connecting homomorphisms. The triangles in the front and the back commute by Diagram 6.5. In Definitino 6.2, we have constructed $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, f)$ by lifting $f: B \to C$ to a chain map $f_{\bullet}: B_{\bullet} \to C_{\bullet}$ through the Comparison Theorem [42, Theorem 2.2.6] and by composing any almost chain map representative of an element in $\widehat{\mathcal{Ext}}^n_{\mathcal{C}}(A, B)$ with f_{\bullet} . Note that we can lift $\widetilde{f}_{2k}: \widetilde{B}_{2k} \to \widetilde{C}_{2k}$ to a chain map $f_{\bullet}^{+2k}: B_{\bullet}^{+2k} \to C_{\bullet}^{+2k}$ given by

$$f_m^{+2k} := \begin{cases} f_{m+2k} : B_{m+2k} \to C_{m+2k} & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$

In particular, the square

commutes and with it Diagram 6.7. The top right and bottom right squares of the latter diagram form a direct system in whose direct limit we obtain the commuting square in the statement of the proposition. $\hfill \Box$

Corollary 6.11 For every $n \in \mathbb{Z}$ we define the natural isomorphism

$$\sigma^n := \vartheta^n \circ \zeta^n : \widehat{\operatorname{Ext}}^n_{\mathcal{C}}(A, -) \to \mathcal{E}xt^n_{\operatorname{Res}, \mathcal{C}}(A, -) \to \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, -)$$

where ζ^n is taken from Definition 6.7.

6.4 Isomorphism of cohomological functors

In order to prove that the natural isomorphisms $\vartheta^n : \mathcal{E}xt^n_{Res,\mathcal{C}}(A,-) \to \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,-)$ extend to an isomorphism of cohomological functors, we need the following technical result.

Proposition 6.12 Let $0 \to B \xrightarrow{f} C \xrightarrow{g} D \to 0$ be a short exact sequence in \mathcal{C} . If h^* : $\widetilde{D}_1 \to B$ denotes a lift of $\mathrm{id}_D : D \to D$ as in Diagram 3.2 and D_{\bullet} a projective resolution of D, consider a lift of h^* to a chain map $h_{\bullet+1} : D_{\bullet}^{+1} \to B_{\bullet}$ given by [42, Theorem 2.2.6]. Denote by $\widetilde{h}_{k+1}^* : \widetilde{D}_{k+1} \to \widetilde{B}_k$ the induced morphism between the corresponding syzygies where $\widetilde{h}_1^* = h^*$. Then there is a projective resolution \underline{D}'_{\bullet} of D such that for any $k \in \mathbb{N}_0$, $\widetilde{D}_k = \underline{\widetilde{D}}_k$ and there is a commutative diagram

where the morphisms ι_{k+1} , π_k stem from \underline{D}_{\bullet} . In particular, any \tilde{h}_{k+1}^* is a lift of $\mathrm{id}_{\tilde{D}_k}$ as in Diagram 3.2.

Let us explain the context in which we need this result. In Definition 6.5, we have first constructed a connecting homomorphism $\widehat{\Delta}^n : \widehat{\mathcal{Ext}}^n_{\mathcal{C}}(A, D) \to \widehat{\mathcal{Ext}}^{n+1}_{\mathcal{C}}(A, \widetilde{D}_1)$ associated to the short exact sequence $0 \to \widetilde{D}_1 \to D_0 \to D \to 0$. Then we defined


$$\widehat{\mathcal{E}xt}^{n}_{\mathcal{C}}(A,D) \xrightarrow{\widehat{\Delta}^{n}} \widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A,\widetilde{D}_{1}) \xrightarrow{\widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A,h^{*})} \widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A,B)$$

to be the connecting homomorphism associated to $0 \to B \to C \to D \to 0$. Later, we shall compare this to


$$\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A,\widetilde{D}_{2k}) \xrightarrow{\Delta^{n}} \mathcal{E}xt_{\mathcal{C}}^{n+1+2k}(A,\widetilde{D}_{2k+1}) \xrightarrow{\mathcal{E}xt_{\mathcal{C}}^{n+1+2k}(A,\widetilde{h}_{2k+1}^{*})} \mathcal{E}xt_{\mathcal{C}}^{n+1+2k}(A,\widetilde{B}_{2k})$$

which we would like to be the connecting homomorphism associated to the short exact sequence $0 \to \widetilde{B}_{2k} \to \widetilde{C}_{2k} \to \widetilde{D}_{2k} \to 0$. This is where Diagram 6.8 comes into play.


Proof (Proposition 6.12) We proceed by induction over $k \in \mathbb{N}_0$. If k = 0, then we take $\underline{D}_0 := D_0$ where Diagram 3.2 yields the desired diagram for this case. Assume that $\underline{D}_0, \ldots, \underline{D}_k$ have been defined and that there is a diagram of the required form for a number $k \geq 0$. In order to construct \underline{D}_{k+1} and establish Diagram 6.8 with $h_{k+1} : \underline{D}_{k+1} \to \widetilde{C}_k$ and $\widetilde{h}_{k+2}^* : \widetilde{D}_{k+2} \to \widetilde{B}_{k+1}$, we consider two instance of Diagram 4.9 from the Horseshoe Lemma. For this, we follow the construction found in [42, p. 37–38] that yields the proof of this so-called lemma. We explain how to obtain the diagram



The morphisms π_k , ι_{k+1} stem from the projective resolution $(\underline{D}_m)_{m=1}^k$ and the morphisms π'_{k+1} , ι'_{k+2} from D_{\bullet} . In particular, the triangle in the bottom left corner and the one in the bottom right corner commute. The morphisms of the middle row are the ones of a direct product as they can be found in [27, p. 250]. Moreover, $\varepsilon = (\iota_{k+1} \circ \pi'_{k+1}) \circ p_1^1 + \mathrm{id} \circ p_2^1$ by [27, p. 250] and [42, p. 37]. Since ι'_{k+2} and ι_{k+1} can be taken as kernels of π'_{k+1} and π_k , the morphisms ι'_{k+2} and π''_{k+1} are induced by the morphisms i_1^1 and p_2^1 . All rows and columns are exact. Because \underline{D}_k is projective, the middle column is a split exact sequence by [38, Tag 010G]. This together with $D_{k+1} \oplus \underline{D}_k$ being projective implies that $\mathrm{Ker}(\varepsilon)$ is also projective. Similarly, we construct with the morphisms \tilde{f}_k and \tilde{g}_k from Diagram 6.8 the diagram



Note that $\eta = (\tilde{f}_k \circ \check{\pi}_k) \circ p_1^2 + h_k \circ p_2^2$ and that the triangle at the bottom right corner commutes by Diagram 6.8. Now consider the diagram



The bottom side stems from Diagram 6.8, the back side from the lower half of Diagram 6.9 and the front side from the lower half of Diagram 6.10. The left hand side arises from the chain map $h_{\bullet}^*: D_{\bullet}^{+1} \to B_{\bullet}$ while the right hand side commutes trivially. The morphism $h_{k+1}^* \oplus \text{id}: D_{k+1} \oplus \underline{D}_k \to B_k \oplus \underline{D}_k$ is constructed as in [27, p. 251] and renders the top side commutative. In particular, all squares on the outside of Diagram 6.11 commute. The slanted squares in the interior give rise to the commuting diagram

$$D_{k+1} \xrightarrow{\iota_{k+1} \circ \pi'_{k+1}} \underline{D}_k \xleftarrow{\text{id}} \underline{D}_k$$

$$\downarrow h_{k+1} \qquad \qquad \downarrow h_k \qquad \qquad \downarrow \text{id}$$

$$B_k \xrightarrow{\tilde{f}_k \circ \tilde{\pi}_k} C_k \xleftarrow{h_k} \underline{D}_k$$

We can use it together with Diagram 6.9, Diagram 6.10 and the content of [27, p. 251] to prove that the middle square in the interior of Diagram 6.11 also commutes. Namely,

$$\eta \circ (h_{k+1}^* \oplus \operatorname{id}) = \left((\widetilde{f}_k \circ \check{\pi}_k) \circ p_1^2 + h_k \circ p_2^2 \right) \circ (h_{k+1}^* \oplus \operatorname{id}) \\ = (\widetilde{f}_k \circ \check{\pi}_k) \circ p_1^2 \circ (h_{k+1}^* \oplus \operatorname{id}) + h_k \circ p_2^2 \circ (h_{k+1}^* \oplus \operatorname{id}) \\ = (\widetilde{f}_k \circ \check{\pi}_k) \circ h_{k+1}^* \circ p_1^1 + h_k \circ \operatorname{id} \circ p_2^1 \\ = h_k \circ (\iota_{k+1} \circ \pi_{k+1}') \circ p_1^1 + h_k \circ \operatorname{id} \circ p_2^1 \\ = h_k \circ \left((\iota_{k+1} \circ \pi_{k+1}') \circ p_1^1 + \operatorname{id} \circ p_2^1 \right) \\ = h_k \circ \varepsilon.$$

Therefore, all squares in Diagram 6.11 commute. Its top side connected by epimorphisms to its bottom side according to Diagram 6.9 and 6.10. Still using the latter two diagrams, we can take kernels to obtain the commuting diagram

$$0 \longrightarrow \widetilde{D}_{k+2} \xrightarrow{\iota_{k+2}''} \operatorname{Ker}(\varepsilon) \xrightarrow{\pi_{k+1}''} \widetilde{D}_{k+1} \longrightarrow 0$$
$$\downarrow \widetilde{h}_{k+2}^{*} \qquad \qquad \downarrow H \qquad \qquad \downarrow \operatorname{id} \\ 0 \longrightarrow \widetilde{B}_{k+1} \xrightarrow{\widetilde{f}_{k+1}} \widetilde{C}_{k+1} \xrightarrow{\widetilde{g}_{k+1}} \widetilde{D}_{k+1} \longrightarrow 0$$


where the rows are short exact sequences. Having seen that $\operatorname{Ker}(\varepsilon)$ is projective, we complete the inductive step by setting $\underline{D}_{k+1} := \operatorname{Ker}(\varepsilon)$, $\iota_{k+2} := \iota''_{k+2}$, $\pi_{k+1} := \pi''_{k+1}$ and $h_{k+1} := H$.

Theorem 6.13 For any short exact sequence $0 \rightarrow B \rightarrow C \rightarrow D \rightarrow 0$ in C the square


$$\begin{aligned} \mathcal{E}xt^{n}_{Res,\mathcal{C}}(A,D) & \stackrel{\vartheta^{n}}{\longrightarrow} \widehat{\mathcal{E}xt}^{n}_{\mathcal{C}}(A,D) \\ & \downarrow_{\widetilde{\Delta}^{n}} & \downarrow_{\widehat{\Delta}^{n}} \\ \mathcal{E}xt^{n+1}_{Res,\mathcal{C}}(A,B) & \stackrel{\vartheta^{n+1}}{\longrightarrow} \widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A,B) \end{aligned}$$

commutes for every $n \in \mathbb{Z}$. In particular, $(\widehat{\mathcal{Ext}}^{\bullet}_{\mathcal{C}}(A, -), \widehat{\Delta}^{\bullet})$ forms a cohomological functor and $\vartheta^{\bullet} : \mathcal{E}xt^{\bullet}_{Res,\mathcal{C}}(A, -) \to \widehat{\mathcal{Ext}}^{\bullet}_{\mathcal{C}}(A, -)$ an isomorphism of cohomological functors.

Proof Analogously to the proof of Proposition 6.10, it suffices by Definition 6.7 to show that the diagram



is commutative in order to conclude that the square in the statement of the theorem commutes. If we write N := n + 2k and K := 2k, we can reformulate the above prism in form of the diagram



The left hand side of Diagram 6.12 corresponds to the top four squares of Diagram 6.13 while the bottom right hand side of the former corresponds to the bottom two squares

of the latter. In Diagram 6.13, the top and bottom row together with the morphisms connecting them directly retrieve the top right side of Diagram 6.12.

Let us explain how to construct Diagram 6.13 by making use of Proposition 6.12 and its proof. We take the morphism $h^*: \widetilde{D}_1 \to B$ as in that proposition, meaning that any term \widetilde{h}_{2k+1}^* originates from a chain map $h_{\bullet+1}^*: D_{\bullet}^{+1} \to B_{\bullet}$. By Definition 6.5, the connecting homomorphism $\widehat{\Delta}^n: \widehat{\mathcal{Ext}}_{\mathcal{C}}^n(A, D) \to \widehat{\mathcal{Ext}}_{\mathcal{C}}^{n+1}(A, B)$ associated to the short exact sequence $0 \to B \to C \to D \to 0$ is rendered in the very bottom row. The top left square arises from Diagram 4.10 and 6.9. More specifically, its top connecting homomorphism is associated to the short exact sequence at the bottom of Diagram 6.9, its right hand connecting homomorphism to the short exact sequence at the left hand side of the latter diagram, its bottom one to the top short exact sequence and its left hand one to the right hand short exact sequence. Because the connecting homomorphism at the right hand side of the top right square is associated to the short exact sequences at the left hand side of Diagram 6.10, the top two squares commute. By Proposition 6.12, the homomorphism

$$\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A,\widetilde{D}_{2k}) \xrightarrow{\Delta^{n+2k}} \mathcal{E}xt_{\mathcal{C}}^{n+2k+1}(A,\widetilde{D}_{2k+1}) \xrightarrow{\mathcal{E}xt_{\mathcal{C}}^{n+2k+1}(A,\widetilde{h}_{2k+1}^{*})} \mathcal{E}xt_{\mathcal{C}}^{n+2k+1}(A,\widetilde{B}_{2k})$$

at the very top of the diagram is the connecting homomorphism Δ^{n+2k} associated to the short exact sequence $0 \to \widetilde{B}_k \to \widetilde{C}_k \to \widetilde{D}_k \to 0$. This does not only imply that the entire top third of the diagram commutes, but also that the middle third does so too. Due to Proposition 6.10 the entire right hand side of the diagram commutes. Regarding the left hand side, since we are in the same situation as in the proof of the latter proposition, we are left to show that the diagram

$$\begin{aligned} & \mathcal{E}xt^{n+2k}_{\mathcal{C}}(A,\widetilde{D}_{2k}) \xrightarrow{\Delta^{n+2k}} \mathcal{E}xt^{n+1+2k}_{\mathcal{C}}(A,\widetilde{D}_{2k+1}) \\ & \downarrow^{\vartheta^{2k}_n(D)} & \downarrow^{\vartheta^{2k}_{n+1}(\widetilde{D}_1)} \\ & \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A,D) \xrightarrow{\widehat{\Delta}^n} \widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A,\widetilde{D}_1) \end{aligned}$$

commutes. If $\varphi_{\bullet+n+2k}$ + Hom_{Ch(C)} $(A[n+2k], D_{\bullet}^{+2k})$ is a an element in $\mathcal{E}xt_{\mathcal{C}}^{n+2k}(A, \widetilde{D}_{2k})$. Then

$$\begin{cases} (-1)^{m-1+n}\varphi_{m+n} : A_{m+n} \to D_m & \text{if } m \ge 2k+1 \\ 0 & \text{if } m \le 2k \end{cases}$$

is an almost chain map representative of the image under the homomorphism $\widehat{\Delta}^n \circ \vartheta_n^{2k}(D)$ while

$$\begin{cases} (-1)^{m-2k-1+n}\varphi_{m+n} : A_{m+n} \to D_m & \text{if } m \ge 2k+1\\ 0 & \text{if } m \le 2k \end{cases}$$

is a representative of the image under $\vartheta_{n+1}^{2k} \circ \Delta^{n+2k}$. Thus, Diagram 6.12 and 6.13 commute as well as the square in the statement of the theorem.

By Definition 6.7, $(\mathcal{E}xt^{\bullet}_{Res,\mathcal{C}}(A, -), \widetilde{\Delta}^{\bullet})$ forms a cohomological functor and by Lemma 6.9 and Proposition 6.10, ϑ^n is a natural isomorphism for any $n \in \mathbb{Z}$. As for Theorem 5.10 and Lemma 6.6, we can reiterate the proof of Theorem 4.5 to demonstrate that $(\widehat{\mathcal{E}xt}^{\bullet}_{\mathcal{C}}(A, -), \widehat{\Delta}^{\bullet})$ is a cohomological functor and that $\vartheta^{\bullet} : \mathcal{E}xt^{\bullet}_{Res,\mathcal{C}}(A, -) \to \widehat{\mathcal{E}xt}^{\bullet}_{\mathcal{C}}(A, -)$ is an isomorphism of cohomological functors. \Box **Remark 6.14** Given any short exact sequence $0 \to B \to C \to D \to 0$ in C, we conclude by means of Theorem 6.13 that every associated connecting homomorphism $\widehat{\Delta}^n : \widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, D) \to \widehat{\mathcal{E}xt}^{n+1}_{\mathcal{C}}(A, B)$ is well defined in the sense that it does not depend on a choice of a morphism $h^* : \widetilde{D}_1 \to B$ as it occurs in Definition 6.5. We have not been able to prove this directly.

Corollary 6.15 We can extend the natural isomorphisms σ^n from Corollary 6.11 to isomorphism of cohomological functors

$$\sigma^{\bullet} = \vartheta^{\bullet} \circ \zeta^{\bullet} : \left(\widehat{\operatorname{Ext}}^{n}_{\mathcal{C}}(A, -), \widehat{\delta}^{\bullet}\right) \to \left(\mathcal{E}xt^{n}_{Res, \mathcal{C}}(A, -), \widetilde{\Delta}^{\bullet}\right) \to \left(\widehat{\mathcal{E}xt}^{n}_{\mathcal{C}}(A, -), \widehat{\Delta}^{\bullet}\right)$$

In [19, p. 21–22], S. Guo and L. Liang establish an isomorphism $\widehat{\mathcal{Ext}}_{\mathcal{C}}^n(A, B) \to BC_{\mathcal{C}}^n(A, B)$ from the cohomology groups of the hypercohomology construction to the ones of the naïve construction for every $n \in \mathbb{Z}$. As it is used in the construction of Yoneda products in [16, Theorem 6.6], we present it below and prove that it is an isomorphism of cohomological functors.

Definition 6.16 ([19, p. 21–22]) For any $A, B \in obj(\mathcal{C})$ and $n \in \mathbb{Z}$ let the map

$$\rho^n:\widehat{\mathcal{Ext}}^n_{\mathcal{C}}(A,B)\to BC^n_{\mathcal{C}}(A,B)$$

be given as follows. If $x = \varphi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$ is an element in $\widehat{\mathcal{E}xt}^n_{\mathcal{C}}(A, B)$, then there exists $K \in \mathbb{N}_0$ such that $(\varphi_{m+n} : A_{m+n} \to B_m)_{m \geq 2K}$ forms a chain map. Denote by $\widetilde{\varphi}_{2K+n} : \widetilde{A}_{2K+n} \to \widetilde{B}_{2K}$ the morphism induced between the corresponding syzygies. We set $\rho^n(x)$ to be the element in $BC^n_{\mathcal{C}}(A, B)$ to which the element $\widetilde{\varphi}_{2K} + \mathcal{P}_{\mathcal{C}}(\widetilde{A}_{2K+n}, \widetilde{B}_{2K})$ in $[\widetilde{A}_{2K+n}, B_{2K}]_{\mathcal{C}}$ is mapped by the corresponding homomorphism going to the direct limit.

Lemma 6.17 The maps ρ^n form an isomorphism of cohomological functors such that

$$\rho^{\bullet} = \beta^{\bullet} \circ (\sigma^{\bullet})^{-1} : (\widehat{\mathcal{E}xt}^{\bullet}_{\mathcal{C}}(A, -), \widehat{\Delta}^{\bullet}) \to (\operatorname{Ext}^{n}_{\operatorname{Res}, \mathcal{C}}(A, -), \widetilde{\delta}^{\bullet}) \to (BC^{\bullet}_{\mathcal{C}}(A, -), \tau^{\bullet})$$

where β^{\bullet} is taken from Theorem 5.10.

Proof Since β^{\bullet} and σ^{\bullet} are already isomorphisms of cohomological functors, it suffices to prove that $\rho^n = \beta^n \circ (\sigma^n)^{-1}$ for every $n \in \mathbb{Z}$. Let $\varphi_{\bullet+n} + \widetilde{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet})$ be an element in $\widehat{\operatorname{Ext}}^n_{\mathcal{C}}(A, B)$. If $K \in \mathbb{N}_0$ is such that $(\varphi_{m+n})_{m \geq 2K}$ is chain map, then for any $k \geq K$ we define $\varphi_{\bullet+n+2k}^{\{2k\}} : A[n+2k]_{\bullet} \to B_{\bullet}^{+2k}$ by setting

$$\varphi_{m+n+2k}^{\{2k\}} := \begin{cases} \varphi_{m+n+2k} : A_{m+n+2k} \to B_{m+2k} & \text{if } m \ge 0\\ 0 & \text{if } m < 0 \end{cases}$$

By Corollary 6.11, $(\sigma^n)^{-1} = (\zeta^n)^{-1} \circ (\vartheta^n)^{-1}$. As $\operatorname{Ext}_{\operatorname{Res},\mathcal{C}}^n(A, B)$ is formed as a direct limit of abelian groups, we introduce the following notation. Let $(A_i)_{i\in\mathbb{N}}$ be a direct system of abelian groups with homomorphisms given by $a_{i,j} : A_i \to A_j$ for $i \leq j$. We write $(x_k)_{k\geq K}$ if $x_k \in A_k$ and $a_{i,j}(x_i) = x_j$ for any $j \geq i \geq K$. This results in an element in $\varinjlim_{i\in\mathbb{N}} A_i$ that we identify with $(x_k)_{k\geq K}$. Then we see that

$$(\vartheta^n)^{-1} \big(\varphi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})} (A[n]_{\bullet}, B_{\bullet}) \big) = \big(\varphi_{\bullet+n+2k}^{\{2k\}} + \operatorname{Null}_{\operatorname{Ch}(\mathcal{C})} (A[n+2k]_{\bullet}, B_{\bullet}^{+2k}) \big)_{k \ge K}$$

Denote by $a_m : A_m \to A_{m-1}$ the boundary maps in the projective resolution A_{\bullet} and by $\pi_m : B_m \to \tilde{B}_m$ the morphisms to the m^{th} syzygy. According to the construction of ζ_m in the proof of Lemma 6.6 and of σ^n in Definition 6.7, we infer that

$$(\sigma^{n})^{-1} \big(\varphi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet}) \big) = \Big(\pi_{n+2k} \circ \varphi_{n+2k} + \operatorname{Im} \big(\operatorname{Hom}_{\mathcal{C}}(a_{n+2k}, \widetilde{B}_{2k}) \big) \Big)_{k \ge K}$$

We deduce from the construction of β^n in Lemma 5.7 that

$$\beta^n \circ (\sigma^n)^{-1} \big(\varphi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet}) \big) = \big(\alpha_{n+2k}(\widetilde{B}_{2k})(\varphi_{n+2k}) + [\widetilde{A}_{2k+n}, \widetilde{B}_{2k}]_{\mathcal{C}} \big)_{k \ge K}$$

where $\alpha_{n+2k}(\widetilde{B}_{2k})$ denotes the homomorphism from Diagram 5.4. Because

$$\beta^n \circ (\sigma^n)^{-1} \big(\varphi_{\bullet+n} + \widehat{\operatorname{Null}}_{\operatorname{Ch}(\mathcal{C})}(A[n]_{\bullet}, B_{\bullet}) \big) = \big(\widetilde{\varphi}_{n+2k} + [\widetilde{A}_{2k+n}, \widetilde{B}_{2k}]_{\mathcal{C}} \big)_{k \ge K}$$

by Diagram 6.1, we conclude that $\rho^n = \beta^n \circ (\sigma^n)^{-1}$.

We close our account on constructions of Mislin completions by posing a question. In Notation 6.1 we have remarked that any completed Ext-group $\widetilde{\mathcal{E}xt}^n_{\mathcal{C}}(A, B)$ can be defined as the n^{th} cohomology group of the Vogel complex $\operatorname{Vog}^n_{\mathcal{C}}(A_{\bullet}, B_{\bullet})_{\bullet}$ whence one obtains

Lemma 6.18 ([19, Proposition 4.8]) If $0 \to B \to C \to D \to 0$ is a short exact sequence in C, one can lift it to a short exact sequence $0 \to B_{\bullet} \to C_{\bullet} \to D_{\bullet} \to 0$ of chain complexes by the Horseshoe Lemma [42, p. 37]. Then, by the Snake Lemma [42, p. 11– 12], there are associated connecting homomorphisms $\overline{\delta}^n : \widehat{\operatorname{Ext}}^n_{\mathcal{C}}(A, D) \to \widehat{\operatorname{Ext}}^{n+1}_{\mathcal{C}}(A, B)$ and $\overline{\delta}^n : \widehat{\operatorname{Ext}}^n_{\mathcal{C}}(B, A) \to \widehat{\operatorname{Ext}}^{n+1}_{\mathcal{C}}(D, A)$. In particular, $(\widehat{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(A, -), \overline{\delta}^{\bullet})$ and $(\widehat{\operatorname{Ext}}^{\bullet}_{\mathcal{C}}(-, A), \overline{\delta}^{\bullet})$ form cohomological functors.

Question 6.19 Does the connecting homomorphism $\widehat{\Delta}^{\bullet}$ constructed in Definition 6.5 agree with $\overline{\delta}^{\bullet}$? More specifically, is $(\widehat{\mathcal{E}xt}^{\bullet}_{\mathcal{C}}(A, -), \overline{\delta}^{\bullet})$ a Mislin completion of the unenriched Ext-functors $(\operatorname{Ext}^{\bullet}_{\mathcal{C}}(A, -), \delta^{\bullet})$?

7 Complete cohomology in condensed mathematics

As we implement complete cohomology in condensed mathematics, we provide hereby more detail on the latter. First, we define condensed sets following D. Clausen and P. Scholze's account [9, Lecture I and II]. After giving a definition that is sheaf-theoretic in nature, we provide a more down-to-earth definition. A finite collection of (continuous) maps $\{f_i : S_i \to S\}_{i=1}^n$ in the category **Pro** of profinite spaces is called a covering if the induced map $\bigsqcup_{i=1}^n S_i \to S$ is surjective. Although this turns **Pro** into a Grothendieck site, it is not well suited for sheaf theoretic purposes as it is not essentially small. A cardinal number κ is called a strong limit cardinal if for every $\lambda < \kappa$ we also have $2^{\lambda} < \kappa$. Then we define the site of κ -small profinite spaces **Pro**_{\kappa} for an uncountable strong limit cardinal κ as the full subcategory of **Pro** consisting of all profinite spaces with less than κ clopen subsets and restrict the Grothendieck topology to it. Note that the Grothendieck site **Pro**_{\kappa} is essentially small. Then a κ -condensed set is a sheaf of sets on **Pro**_{\kappa}. In more down-to-earth terms, a (contravariant) functor

$$X : \operatorname{Pro}_{\kappa}^{\operatorname{op}} \to \operatorname{Set}$$

is a κ -condensed set if the following two conditions are satisfied.

- 1. For every finite collection $\{S_i\}_{i=1}^n$ of objects in the category $\operatorname{Pro}_{\kappa}$ there is a bijection $X(\bigsqcup_{i=1}^n S_i) \to \prod_{i=1}^n X(S_i).$
- 2. Let $f: T \to S$ be a (continuous) surjection in $\operatorname{Pro}_{\kappa}$ and denote the projections from the fibre product by $p_1, p_2: T \times_S T \to T$. Then the function $X(f): X(S) \to X(T)$ is injective with image

$$Im(X(f)) = \{x \in X(T) \mid X(p_1)(x) = X(p_2)(x) \in X(T \times_S T)\}.$$

One can think of X as encoding a topological space and of X(S) as the continuous maps " $S \to X$ " from a κ -profinite space. If $\kappa < \kappa'$ is another strong limit cardinal, then there is a fully faithful functor

$$\operatorname{Cond}_{\kappa}(\operatorname{\mathbf{Set}}) \to \operatorname{Cond}_{\kappa'}(\operatorname{\mathbf{Set}})$$

due to our choice of the cardinal numbers. This means that one can extend any κ condensed set to a κ' -condensed set. If K denotes the class of uncountable strong limit
cardinals, then the category of condensed sets is defined as the colimit category

$$\operatorname{Cond}(\operatorname{\mathbf{Set}}) := \varinjlim_{\kappa \in K} \operatorname{Cond}_{\kappa}(\operatorname{\mathbf{Set}}).$$

Hence, a condensed set is an equivalence class of κ -condensed sets. According to [35, p. 15/16 and Proposition 1.7] we can establish the following relation with topological spaces. If T1-Top denotes the category of T1 topological spaces, then there is a well-defined faithful functor T1-Top \rightarrow Cond(Set) mapping any Y to its so-called condensate

$$\underline{Y}: \mathbf{Pro}^{\mathrm{op}}_{\kappa} \to \mathbf{Set}, \ S \mapsto \{f: S \to Y \text{ continuous}\}$$

where κ is chosen to be a sufficiently large strong limit cardinal. This functor is fully faithful when restricted to T1 compactly generated topological spaces. It follows from the proofs of Proposition 2.9 and Proposition 2.15 in [35] that the functor does not depend on κ . Namely, for every strong limit cardinal $\kappa < \kappa'$ and κ' -profinite space $S = \lim_{i \in I} S_i$ written as a κ -inverse limit of κ -profinite spaces S_i , the function $\lim_{i \in I} \underline{Y}(S_i) \to \underline{Y}(S)$ is a bijection. Although Cond(**Set**) does not form the category of sheaves over any site [35, Remark 2.12], it nevertheless inherits good sheaf-theoretic properties from the categories Cond_{κ}(**Set**) [9, p. 13].

Condensed groups (resp. rings, modules etc.) are defined analogously [35, p. 7] and condensates of T1 topological groups are condensed groups (resp. rings, modules etc) [35, p. 8]. In the category of condensed abelian groups Cond(Ab) all limits and colimits exist where arbitrary products, arbitrary direct sums and direct limits are exact [35, Theorem 1.10]. According to [35, p. 12], Cond(Ab) has enough projectives. Following the source, these assertions also hold for the category of condensed \mathcal{R} -modules Cond($Mod(\mathcal{R})$) where \mathcal{R} is a condensed ring. If S is a projective object in **Pro**, then one can form the free condensed \mathcal{R} -module $\mathcal{R}[\underline{S}]$ as the sheafification of the presheaf of \mathcal{R} -modules

$$\operatorname{Pro}_{\kappa}^{\operatorname{op}} \to \operatorname{Ab}, T \mapsto \mathcal{R}(T)[\underline{S}(T)]$$

where $\mathcal{R}(T)[\underline{S}(T)]$ denotes the free $\mathcal{R}(T)$ -module over the set $\underline{S}(T)$ and κ is a sufficiently large strong limit cardinal [35, p. 12]. Since the Grothendieck site \mathbf{Pro}_{κ} is essentially small, sheafification exists and again by the proof of [35, Proposition 2.9], the construction of $\mathcal{R}[\underline{S}]$ is independent of κ . Moreover, the condensed \mathcal{R} -modules $\mathcal{R}[\underline{S}]$ are compact projective generators, meaning that for every condensed \mathcal{R} -module A there is a collection of projective profinite spaces S_i such that there is an epimorphism $\bigoplus_{i \in I} \mathcal{R}[\underline{S}_i] \to A$ [35, p. 12]. For condensed \mathcal{R} -modules A, B one does not only have the "usual" Hom-set $\operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, B) \in \operatorname{obj}(\operatorname{Ab})$, but also an internal Hom-set which is constructed in [35, p. 13]. Namely, for any $C \in \operatorname{obj}(\operatorname{Cond}(Mod(\mathcal{R})))$ one can form the tensor product $C \otimes_{\mathcal{R}} A \in \operatorname{Cond}(Mod(\mathcal{R}))$ as the sheafification of

$$\operatorname{Pro}_{\kappa}^{\operatorname{op}} \to \operatorname{Ab}, T \mapsto C(T) \otimes_{\mathcal{R}(T)} A(T).$$

Then define the internal Hom-set $\underline{\text{Hom}}_{\text{Cond}(Mod(\mathcal{R}))}(A, B) \in \text{obj}(\text{Cond}(\mathbf{Ab}))$ by

 $\operatorname{\mathbf{Pro}}_{\kappa}^{\operatorname{op}} \to \operatorname{\mathbf{Ab}}, T \mapsto \operatorname{Hom}_{\operatorname{Cond}(\operatorname{Mod}(\mathcal{R}))}(A \otimes_{\mathcal{R}} \mathcal{R}[\underline{T}], B).$

By definition, it satisfies the adjunction

 $\operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{R}))}(C, \operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, B)) \cong \operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{R}))}(C \otimes_{\mathcal{R}} A, B).$

Because the category $Cond(Mod(\mathcal{R}))$ has enough projectives, we can define the unenriched Ext-functors

$$\operatorname{Ext}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -) : \operatorname{Cond}(Mod(\mathcal{R})) \to \operatorname{Ab}$$

and the internal Ext-functors

$$\underline{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -) : \operatorname{Cond}(Mod(\mathcal{R})) \to \operatorname{Cond}(\operatorname{\mathbf{Ab}})$$

to be the derived functors of $\operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$ and $\operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$. This can be used to define cohomology of a condensed group \mathcal{G} . Following [2, p. 2–3] one can form the condensed group ring $\mathcal{R}[\mathcal{G}]$ such that the category $\operatorname{Cond}(Mod(\mathcal{R}[\mathcal{G}]))$ is equivalent to the subcategory of $\operatorname{Cond}(\mathcal{R})$ of all \mathcal{R} -modules M with a \mathcal{G} -action $\mathcal{G} \times M \to M$. By [2, p. 5/8], one can define condensed unenriched group cohomology $H^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ and the condensed internal group cohomology $\underline{H}^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ as the Ext-functors $\operatorname{Ext}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}[\mathcal{G}]))}(\mathcal{R}, -)$ and $\underline{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}[\mathcal{G}]))}(\mathcal{R}, -)$. By the above, we conclude

Theorem 7.1 Let \mathcal{R} be a condensed ring and A a condensed \mathcal{R} -module. Then there are completed condensed unenriched Ext-functors $\widehat{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$ and completed condensed internal Ext-functors $\widehat{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$. If \mathcal{G} is a condensed group, then complete condensed unenriched group cohomology $\widehat{H}^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ and complete condensed internal group cohomology $\widehat{H}^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ can be defined.

Remark 7.2 Because both $\widehat{\operatorname{Ext}}^{\bullet}_{\operatorname{Cond}(Mod(\mathcal{R}))}(A, -)$ and $\widehat{H}^{\bullet}_{\mathcal{R}}(\mathcal{G}, -)$ have as codomain category $\operatorname{Cond}(\operatorname{Ab})$, they are an example of completed Ext-functors that are not covered under the previous frameworks such as the one by A. Beligiannis and I. Reiten in [4], the one by S. Guo and L. Liang's in [19] and the one by J. Hu et al. in [22].

Having discussed condensed modules, let us consider solid modules. We have already mentioned in Section 1 that one can think of these as being a form of completed condensed modules where a condensed ring \mathcal{A} is called analytic if it admits a choice of solid modules. More formally, let **ProjPro** denote the full subcategory of **Pro** consisting of projective objects. Then an analytic ring \mathcal{A} comes with a particular functor

$$c_{\mathcal{A}}: \mathbf{ProjPro} \to \mathrm{Cond}(Mod(\mathcal{A})), \ S \mapsto \mathcal{A}[S]_{\blacksquare}$$

and a condensed \mathcal{A} -module homomorphism $C_{\mathcal{A}}(S) : \mathcal{A}[\underline{S}] \to \mathcal{A}[\underline{S}]_{\blacksquare}$ for every projective profinite space S [35, p. 44]. One can think of $c_{\mathcal{A}}$ as assigning to every projective profinite space S a canonical choice of free solid \mathcal{A} -module $\mathcal{A}[S]_{\blacksquare}$. According to [35, Proposition 7.5], analytic rings have the following properties. The full subcategory Solid(\mathcal{A}) of Cond($Mod(\mathcal{A})$) with objects M such that

$$\operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{A}))}(C_{\mathcal{A}}(S), M) : \operatorname{Hom}_{\operatorname{Cond}(Mod(\mathcal{A}))}(\mathcal{A}[S]_{\blacksquare}, M) \to M(S)$$

is an isomorphism for all $S \in \operatorname{\mathbf{ProjPro}}$ is an abelian category stable under limits, colimits and extensions. The objects $\mathcal{A}[S]_{\blacksquare}$ are compact projective generators in $\operatorname{Solid}(\mathcal{A})$. Moreover, there is an extension of the morphisms $C_{\mathcal{A}}(S) : \mathcal{A}[\underline{S}] \to \mathcal{A}[\underline{S}]_{\blacksquare}$ to a functor $\operatorname{Cond}(\operatorname{Mod}(\mathcal{A})) \to \operatorname{Solid}(\mathcal{A})$ left adjoint to the inclusion of categories that we call the solidification functor. Lastly, if \mathcal{A} is commutative, then there exists a unique symmetric monoidal tensor product on $\operatorname{Solid}(\mathcal{A})$ rendering the solidification functor symmetric monoidal. In particular, since the category $\operatorname{Solid}(\mathcal{A})$ has enough projectives, we can define the Ext functors

$$\operatorname{Ext}^{\bullet}_{\operatorname{Solid}(\mathcal{A})}(M, -) : \operatorname{Solid}(\mathcal{A}) \to \operatorname{Ab}$$

as the derived functors of $\operatorname{Hom}_{\operatorname{Solid}(\mathcal{A})}(M, -)$.

Lemma 7.3 Let \mathcal{A} be an analytic ring and M a solid \mathcal{A} -module. Then there are completed solid Ext-functors $\widehat{\operatorname{Ext}}_{\operatorname{Solid}(\mathcal{A})}^{\bullet}(M, -)$.

Example 7.4 ([2, p. 2]) Let $S = \varprojlim_{i \in I}$ be a profinite space and define the condensed $\underline{\mathbb{Z}}$ -module $\mathbb{Z}[S]^{\blacksquare} := \varprojlim_{i \in I} \underline{\mathbb{Z}}[\underline{S}_i]$. Then the condensed ring $\underline{\mathbb{Z}}$ together with the assignment $S \mapsto \mathbb{Z}[S]^{\blacksquare}$ for every projective profinite space S forms an analytic ring. By extension, we denote by $(-)^{\blacksquare}$: Cond(Ab) \rightarrow Solid($\underline{\mathbb{Z}}$) the solidification functor. If Λ denotes one of the profinite rings \mathbb{Z}_p , $\mathbb{Z}/p\mathbb{Z}$ or $\mathbb{Z}/p^n\mathbb{Z}$, then also $(\underline{\Lambda}, S \mapsto (\underline{\Lambda}[\underline{S}])^{\blacksquare})$ is an analytic ring.

Lemma 7.5 Let G be a profinite group and $(L, S \mapsto L[S]^{\bullet})$ an analytic ring where $(-)^{\bullet}$ denotes the solidification functor of the analytic ring $\underline{\mathbb{Z}}$ from Example 7.4. Then the group ring $(L[\underline{G}], S \mapsto L[\underline{G}][S]^{\bullet})$ is also an analytic ring and one can thus define complete solid cohomology $\widehat{H}^{\bullet}_{\mathrm{Solid}(L)}(\underline{G}, -)$: $\mathrm{Solid}(L[\underline{G}]) \to \mathrm{Ab}$ and complete solid internal cohomology $\widehat{H}^{\bullet}_{\mathrm{Solid}(L)}(\underline{G}, -)$: $\mathrm{Solid}(L[\underline{G}]) \to \mathrm{Solid}(\underline{\mathbb{Z}}).$

Proof It follows from Corollary 5.5 and Corollary 6.1 of [35] that $L[S]^{\blacksquare}$ is a projective solid *L*-module for every profinite (but not necessarily projective) space *S*. Then, by [2, Lemma 1.3], the group ring $(L[\underline{G}], S \mapsto L[\underline{G}][S]^{\blacksquare})$ is analytic. It follows from [2, p. 3–4] that the category of solid $L[\underline{G}]$ is equivalent to the category of solid *L*-modules together with an action $\underline{G} \times M \to M$ in the category of solid *L*-modules. In particular, there is an unenriched Hom-functor $\operatorname{Hom}_{\operatorname{Solid}(L[\underline{G}])}(L, -)$ from which we can form the solid group cohomology $H^{\bullet}_{\operatorname{Solid}(L)}(\underline{G}, -)$ [2, p. 5] and thus the corresponding complete solid cohomology. On the other hand, the internal Hom-functor of condensed $L[\underline{G}]$ -modules restricted onto the full subcategory Solid($L[\underline{G}]$)

$$\operatorname{Solid}(L[\underline{G}]) \to \operatorname{Cond}(\operatorname{\mathbf{Ab}}), M \to \operatorname{\underline{Hom}}_{\operatorname{Cond}(L[\underline{G}])}(L, M)$$

maps into solid $\underline{\mathbb{Z}}$ -modules according to [2, p. 8]. Therefore, we can also form the complete solid internal cohomology.

Acknowledgements

A special thanks goes to Peter H. Kropholler for sharing his expertise with me. He has discussed examples of complete cohomology groups of discrete groups with me and pointed out his joint paper with Jonathan Cornick "On Complete Resolutions". In particular, he introduced me to condensed mathematics. I would also like to express my gratitude to Emma Brink for her great help in condensed mathematics.

I would like to acknowledge Alejandro Adem, Jon F. Carlson, Joel Friedman, Kalle Karu, Zinovy Reichstein and Ben Williams who assessed my PhD thesis and whose invaluable suggestions have led to an improvement of the introduction. Moreover, I am thankful to Andrew Fisher for pointing out the paper "Complete Homology over Associative Rings" by Olgur Celikbas et al. Lastly, I am grateful to Ged Corob Cook for insightful discussions.

References

- A. Adem and R. J. Milgram. Cohomology of Finite Groups, volume 309 of Grundlehren der mathematischen Wissenschaften / A Series of Comprehensive Studies in Mathematics. Springer-Verlag Berlin Heidelberg New York, 2nd edition, 2004.
- [2] J. Anschütz and A.-C. Le Bras. Solid Group Cohomology. https://www.math. uni-bonn.de/people/ja/homology_profinite.pdf, September 2020. This website has been consulted on 15 July 2023.
- [3] J. Asadollahi and S. Salarian. Cohomology Theories for Complexes. *Journal of Pure* and Applied Algebra, 210(3):771–787, 2007.
- [4] A. Beligiannis and I. Reiten. Homological and Homotopical Aspects of Torsion Theories, volume 188 of Memoirs of the American Mathematical Society. American Mathematical Society, 2007.
- [5] D. J. Benson and J. F. Carlson. Products in Negative Cohomology. Journal of Pure and Applied Algebra, 82(2):107–129, 1992.
- [6] K. S. Brown. Cohomology of Groups, volume 87 of Graduate Texts in Mathematics. Springer-Verlag New York, Inc., 1982.
- [7] H. Cartan and S. Eilenberg. Homological Algebra, volume 19 of Princeton Mathematical Series. Princeton University Press, 1956.
- [8] O. Celikbas, L. W. Christensen, L. Liang, and G. Piepmeyer. Complete Homology over Associative Rings. Israel Journal of Mathematics, 221(1):1–24, 2017.
- [9] D. Clausen and P. Scholze. Condensed Mathematics and Complex Geometry. https: //people.mpim-bonn.mpg.de/scholze/Complex.pdf, April 2022. This website has been consulted on 8 April 2023.
- [10] J. Cornick and P. H. Kropholler. On Complete Resolutions. Topology and its Applications, 78(3):235—250, 1997.
- [11] F. Dembegioti. On the Zeroeth Complete Cohomology. Journal of Pure and Applied Algebra, 203(1–3):119–132, 2005.

- [12] F. Dembegioti. On the Zeroeth Complete Cohomology of Certain Polycyclic Groups. Communications in Algebra, 36(5):1927–1941, 2008.
- [13] I. Emmanouil and P. Manousaki. On the Stable Homology of Modules. Journal of Pure and Applied Algebra, 221(9):2198–2219, 2017.
- [14] F. T. Farrell. An Extension of Tate Cohomology to a Class of Infinite Groups. Journal of Pure and Applied Algebra, 10(2):153–161, 1977.
- [15] S. I. Gelfand and Yu. I. Manin. Methods of Homological Algebra. Springer Monographs in Mathematics. Springer-Verlag Berlin Heidelberg, 2nd edition, 2003.
- [16] M. Gheorghiu. On a Completion of Cohomological Functors Generalising Tate Cohomology II. arXiv e-prints, page arXiv:2405.03634v1, May 2024.
- [17] F. Goichot. Homologie de Tate-Vogel Équivariante. Journal of Pure and Applied Algebra, 82(1):39–64, 1992.
- [18] A. Grothendieck. Sur Quelques Points d'Algèbre Homologique. Tohoku Mathematical Journal, 9(2):119–221, 1957.
- [19] S. Guo and L. Liang. Stable Functors and Cohomology Theory in Abelian Categories. arXiv e-prints, page arXiv:2006.13562v3, October 2023.
- [20] S. Guo, L. Liang, and X. Yang. Balancedness of Complete Cohomology with respect to Subcategories. https://sites.google.com/site/lliangnju/publications, 2023. This website has been consulted on 7 June 2024.
- [21] J. Hu, D. Zhang, T. Zhao, and P. Zhou. Balance of Complete Cohomology in Extriangulated Categories. *Electronic Research Archive*, 29(5):3341–3359, 2021.
- [22] J. Hu, D. Zhang, T. Zhao, and P. Zhou. Complete Cohomology for Extriangulated Categories. Algebra Colloquium, 28(4):701–720, 2021.
- [23] J. H. Jo. Complete Homology and Related Dimensions of Groups. Journal of Group Theory, 12(3):431–448, 2009.
- [24] J. R. Klein. Axioms for Generalized Farrell–Tate Cohomology. Journal of Pure and Applied Algebra, 172(2–3):225–238, 2002.
- [25] P. H. Kropholler. Hierarchical Decompositions, Generalized Tate Cohomology, and Groups of Type (FP)∞. In Combinatorial and Geometric Group Theory, Edinburgh 1993, volume 204 of London Mathematical Society Lecture Note Series, pages 190– 216. Cambridge University Press, January 1995.
- [26] M. Land. Condensed Mathematics. https://www.markus-land.de/teaching/, December 2022. This website has been consulted on 22 September 2023.
- [27] S. Mac Lane. *Homology*. Classics in Mathematics. Springer-Verlag Berlin Heidelberg, 1995.
- [28] G. Mislin. Tate Cohomology for Arbitrary Groups via Satellites. Topology and its Applications, 56:293–300, 1994.

- [29] B. E. A. Nucinkis. Complete Cohomology for Arbitrary Rings Uusing Injectives. Journal of Pure and Applied Algebra, 131(3):297–318, 1998.
- [30] M. S. Osborne. Basic Homological Algebra, volume 196 of Graduate Texts in Mathematics. Springer-Verlag New York, Inc., 2000.
- [31] M. Paganin. On Some Generalizations of Tate Cohomology: an Overview. Pro Mathematica, 29(58):63–115, 2016.
- [32] B. Poonen. Why All Rings should have a 1. https://math.mit.edu/~poonen/ papers/ring.pdf, June 2018. This website has been consulted on 25 October 2023.
- [33] L. Ribes. Profinite Graphs and Groups, volume 66 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer Nature, 2017.
- [34] L. Ribes and P. Zalesskii. Profinite Groups, volume 40 of Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics. Springer-Verlag Berlin Heidelberg, 2nd edition, 2010.
- [35] P. Scholze. Lectures on Condensed Mathematics. https://www.math.uni-bonn. de/people/scholze/Condensed.pdf, May 2019. This website has been consulted on 21 February 2023.
- [36] P. Scholze. Liquid Tensor Experiment. https://xenaproject.wordpress.com/ 2020/12/05/liquid-tensor-experiment/, December 2020. This website has been consulted on 25 September 2023.
- [37] J.-P. Serre. *Cohomologie Galoisienne*, volume 5 of *Lecture Notes in Mathematics*. Springer-Verlag Berlin Heidelberg, 5th edition, 1994.
- [38] The Stacks project authors. The Stacks Project. https://stacks.math.columbia. edu, 2024.
- [39] P. Symonds. The Tate-Farrell Cohomology of the Morava Stabilizer Group s_{p-1} with Coefficients in e_{p-1}. In Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-Theory, volume 346 of Contemporary Mathematics, pages 485–402. American Mathematical Society, 2004.
- [40] P. Symonds. Permutation Complexes for Profinite Groups. Commentarii Mathematici Helvetici, 82(1):1–37, 2007.
- [41] P. Symonds and T. Weigel. Cohomology of p-adic Analytic Groups. In New Horizons in Pro-p Groups, volume 184 of Progress in Mathematics, pages 347–408. Birkhäuser, Boston, 2000.
- [42] C. A. Weibel. An Introduction to Homological Algebra, volume 38 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, 1994.
- [43] J. S. Wilson. Profinite Groups, volume 19 of London Mathematical Society Monographs New Series. Oxford University Press Inc., New York, 1998.