Investigation of sample paths properties of sub-Gaussian type random fields, with application to stochasic heat equations

Olha Hopkalo

Department of Economic Cybernetics, Faculty of Economics, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine, E-mail: olha.hopkalo@knu.ua

Lyudmyla Sakhno

Department of Probability Theory, Statistics and Actuarial Mathematics, Faculty of Mechanics and Mathematics, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine, E-mail: lyudmyla.sakhno@knu.ua

Abstract

The paper presents bounds for the distributions of suprema for a particular class of sub-Gaussian type random fields defined over spaces with anisotropic metrics. The results are applied to random fields related to stochastic heat equations with fractional noise: bounds for the tail distributions of suprema and estimates for the rate of growth are provided for such fields.

Key words: sub-Gaussian type random fields, distribution of supremum, rate of growth, stochastic heat equation, fractional noise

1 Introduction and preliminaries

Aim and motivation. In this paper, we study sample paths properties of a class of sub-Gaussian type random fields X(t), $t \in T$, focusing on the case where a parameter set T is endowed with an anisotropic metric and imposing some kind of Hölder continuity condition on the field X. Our aim is to establish upper bounds for the distribution of the supremum $P\{\sup_{t\in T} |X(t)| > u\}$ for bounded T and to evaluate a rate of growth of X over an unbounded domain V by considering upper bounds for $P\{\sup_{t\in V} \frac{|X(t)|}{f(t)} > u\}$ for a properly chosen continuous function f. The study is motivated by applications to random fields related to stochastic heat equations. Extensive recent investigations of such equations resulted, in particular, in establishing the Hölder continuity of solutions in various settings. It is quite natural and appealing to make further steps towards considerarion of different functionals of solutions. We evaluate the distribution of suprema of solutions and their asymptotic rate of growth.

Approach and tools. We present bounds for distributions of suprema assuming X to belong to a particular class of φ -sub-Gaussian random fields (to be defined below), which provides a generalization of Gaussian and sub-Gaussian fields. To derive the results we apply entropy methods. Recall that the entropy approach in studing sample paths of a stochastic process X(t), $t \in T$, requires to evaluate entropy characteristics of the set T with respect to a particular metrics generated by the process X. The origins of this approach are due to Dudley, who stated coditions for boundedness of Gaussian processes in the form of convergence of metric entropy integrals (which we call now Dudley entropy integrals). We address for corresponding references, e.g., to [1] and [4], where in the latter one the entropy approach was extended to different classes of processes, more general than Gaussian ones. Some facts from the general theory of φ -sub-Gaussian random variables and fields. Note that it is important for applications to go beyond the Gaussianity assumption in considered models and possible extensions are provided by sub-Gaussian and φ -sub-Gaussian random processes and fields. Recall that a random variable ξ is sub-Gaussian if its moment generating function is majorized by that of a Gaussian centered random variable $\eta \sim N(0, \sigma^2)$:

$$\mathsf{E}\exp(\lambda\xi) \le \mathsf{E}\exp(\lambda\eta) = \exp(\sigma^2\lambda^2/2).$$

The generalization of this notion to the classes of φ -sub-Gaussian random variables is introduced as follows (see, [4, Ch.2], [7], [16], [20]).

Consider a continuous even convex function φ such that $\varphi(0) = 0$, $\varphi(x) > 0$ as $x \neq 0$ and $\lim_{x \to 0} \frac{\varphi(x)}{x} = 0$, $\lim_{x \to \infty} \frac{\varphi(x)}{x} = \infty$. Note that such functions are called Orlicz N-functions. Suppose that φ satisfies additionally: $\lim_{x \to 0} \inf_{x \to 0} \frac{\varphi(x)}{x^2} = c > 0$, where the case $c = \infty$ is possible.

Let φ be the function with the above properties and $\{\Omega, \mathcal{F}, \mathsf{P}\}$ be a standard probability space. The random variable ζ is φ -sub-Gaussian, or belongs to the space $\operatorname{Sub}_{\varphi}(\Omega)$, if $\mathsf{E}\zeta = 0$, $\mathsf{E}\exp\{\lambda\zeta\}$ exists for all $\lambda \in \mathbb{R}$ and there exists a constant a > 0 such that the following inequality holds for all $\lambda \in \mathbb{R}$

$$\mathsf{E}\exp\{\lambda\zeta\} \le \exp\{\varphi(\lambda a)\}. \tag{1.1}$$

The random field $X(t), t \in T$, is called φ -sub-Gaussian if the random variables $\{X(t), t \in T\}$ are φ -sub-Gaussian.

The space $\operatorname{Sub}_{\varphi}(\Omega)$ is a Banach space with respect to the norm (see [7, 16]):

$$\tau_{\varphi}(\zeta) = \inf\{a > 0 : \mathsf{E}\exp\{\lambda\zeta\} \le \exp\{\varphi(a\lambda)\}.$$

For φ -sub-Gaussian random variable ζ the following estimate for its tail probability holds:

$$P\{|\zeta| > u\} \le 2\exp\left\{-\varphi^*\left(\frac{u}{\tau_{\varphi}(\zeta)}\right)\right\}, \ u > 0,$$
(1.2)

where the function φ^* defined by $\varphi^*(x) = \sup_{y \in \mathbb{R}} (xy - \varphi(y))$ is called the Young-Fenchel transform (or Legendre transform, or convex conjugate) of the function φ . It was stated in [4] (Corollary 4.1, p. 68) that, moreover, a random variable ζ is a φ -sub-Gaussian if and only if $\mathsf{E}\zeta = 0$ and there exist constants C > 0, D > 0 such that

$$P\{|\zeta| > u\} \le C \exp\left\{-\varphi^*\left(\frac{u}{D}\right)\right\}.$$
(1.3)

This second characterization of φ -sub-Gaussian random variable by the tail behavior of its distribution is important for practical applications.

The class of φ -sub-Gaussian random variables includes centered compactly supported distributions, reflected Weibull distributions, centered bounded distributions, Gaussian, Poisson distributions. In the case when $\varphi = \frac{x^2}{2}$, the notion of φ -sub-Gaussianity reduces to the classical sub-Gaussianity. The main theory for the spaces of φ -sub-Gaussian random variables and stochastic processes was presented in [4, 7, 16] followed by numerous further studies. Various classes of φ -sub-Gaussian processes and fields were studied, in particular, in [3, 9, 13, 14, 15, 17].

The property of φ -sub-Gaussianity allows to evaluate different functionals of the stochastic processes, in particular, the behavior of their suprema.

Estimates for distribution of supremum $P\{\sup_{t \in T} |X(t)| > u\}$ of φ -sub-Gaussian stochastic process X were derived in various forms in the monograph [4] basing on entropy methods.

We will base our study on the following theorem (see [4], Theorems 4.1–4.2, pp. 100, 105).

Theorem 1.1 ([4]). Let X(t), $t \in T$, be a φ -sub-Gaussian process and ρ_X be the pseudometrics generated by X, that is, $\rho_X(t,s) = \tau_{\varphi}(X(t) - X(s))$, $t, s \in T$. Suppose further that

(i) the pseudometric space (T, ρ_X) is separable, the process X is separable on (T, ρ_X) ; (ii) $\varepsilon_0 := \sup_{t \in T} \tau_{\varphi}(X(t)) < \infty$; (iii)

$$I_{\varphi}(\varepsilon_0) := \int_{0}^{\varepsilon_0} \Psi(\ln(N(v))) \, dv < \infty, \tag{1.4}$$

where $\Psi(v) = \frac{v}{\varphi^{(-1)}(v)}$ and $N(v) = N_{\rho_X}(v)$, v > 0, is the metric massiveness of the pseudometric space (T, ρ_X) , that is, the smallest number of elements in a v-covering of T by closed balls, w.r.t. the metric ρ_X , of a radius at most v.

Then for all $\lambda > 0$ and $0 < \theta < 1$

$$\operatorname{Eexp}\{\lambda \sup_{t \in T} |X(t)|\} \le 2Q(\lambda, \theta), \tag{1.5}$$

where

$$Q(\lambda,\theta) = \exp\left\{\varphi\left(\frac{\lambda\varepsilon_0}{1-\theta}\right) + \frac{2\lambda}{\theta(1-\theta)}I_{\varphi}(\theta\varepsilon_0)\right\},\tag{1.6}$$

and for $0 < \theta < 1$, $u > \frac{2I_{\varphi}(\theta \varepsilon_0)}{\theta(1-\theta)}$

$$P\left\{\sup_{t\in T} |X(t)| > u\right\} \le 2A(u,\theta),\tag{1.7}$$

where

$$A(u,\theta) = \exp\left\{-\varphi^*\left(\frac{1}{\varepsilon_0}\left(u(1-\theta) - \frac{2}{\theta}I_{\varphi}(\theta\varepsilon_0)\right)\right)\right\}.$$
(1.8)

In the above theorem and in what follows we denote by $f^{(-1)}$ the inverse function for a function f.

The integrals of the form (1.4) are called entropy integrals. Entropy characteristics of the parameter set T with respect to the pseudometric ρ_X generated by the process X and the rate of growth of the metric massiveness $N(v) = N_{\rho_X}(v), v > 0$, or metric entropy $H(v) := \ln(N(v))$ are important for the study of sample paths properties of the underlying process X (see [4]).

Consider now a metric space (T, d), with an arbitrary metric d and suppose that this metric space is separable. Suppose that we can evaluate the metric massivness N_d of T with respect to the metric d and also have a bound for the function $\rho_X(t, s) = \tau_{\varphi}(X(t) - X(s))$ in terms of d(t, s). Then Theorem 1.1 implies the following result.

Theorem 1.2. Let X(t), $t \in T$, be a φ -sub-Gaussian process and T be supplied with a metric d. Assume that

- (i) the metric space (T, d) is separable, the process X is separable on (T, d);
- (*ii*) $\varepsilon_0 := \sup_{t \in T} \tau_{\varphi}(X(t)) < \infty;$
- (iii) there exists a monotonically increasing continuous function $\sigma(h)$, $0 < h \leq \sup_{t,s \in T} d(s,t)$ such that $\sigma(h) \to 0$ as $h \to 0$ and

$$\sup_{\substack{d(t,s) \le h, \\ t,s \in T}} \tau_{\varphi}(X(t) - X(s)) \le \sigma(h), \tag{1.9}$$

and for $0 < \varepsilon \leq \gamma_0$

$$\widetilde{I}_{\varphi}(\varepsilon) := \int_{0}^{\varepsilon} \Psi(\ln(N_d(\sigma^{(-1)}(v))) \, dv < \infty, \tag{1.10}$$

where $\Psi(v) = \frac{v}{\varphi^{(-1)}(v)}$, $N_d(v)$, v > 0, is the metric massiveness of the metric space (T, d), $\gamma_0 = \sigma(\sup_{t,s \in T} d(t,s)).$

Then the statement of Theorem 1.1 holds for $\lambda > 0$ and $0 < \theta < 1$ such that $\theta \varepsilon_0 \leq \gamma_0$ with $Q(\lambda, \theta)$ and $A(u, \theta)$ changed for the $\widetilde{Q}(\lambda, \theta)$ and $\widetilde{A}(u, \theta)$ which correspond to the integral \widetilde{I}_{φ} :

$$\operatorname{\mathsf{Eexp}}\{\lambda \sup_{t \in T} |X(t)|\} \le 2\widetilde{Q}(\lambda, \theta) \tag{1.11}$$

and for $u > \frac{2\tilde{I}_{\varphi}(\theta\varepsilon_0)}{\theta(1-\theta)}$

$$P\left\{\sup_{t\in T} |X(t)| > u\right\} \le 2\widetilde{A}(u,\theta),\tag{1.12}$$

where

$$\widetilde{Q}(\lambda,\theta) = \exp\left\{\varphi\left(\frac{\lambda\varepsilon_0}{1-\theta}\right) + \frac{2\lambda}{\theta(1-\theta)}\widetilde{I}_{\varphi}(\theta\varepsilon_0)\right\},\tag{1.13}$$

$$\widetilde{A}(u,\theta) = \exp\left\{-\varphi^*\left(\frac{1}{\varepsilon_0}\left(u(1-\theta) - \frac{2}{\theta}\widetilde{I}_{\varphi}(\theta\varepsilon_0)\right)\right)\right\}.$$
(1.14)

Proof. Theorem 1.2 follows immediately from Theorem 1.1. We have from (2.5) that $\sup_{\substack{d(t,s) \leq h, \\ t,s \in T}} \rho_X(t,s) \leq \sigma(h)$, therefore, the smallest number of elements in an ε -covering of (T, ρ_X) can be bounded by the smallest number of elements in a $\sigma^{(-1)}(\varepsilon)$ -covering of (T,d): $N_{\rho_X}(\varepsilon) \leq N_d(\sigma^{(-1)}(\varepsilon))$. This implies the estimate $I_{\varphi}(\varepsilon) \leq \tilde{I}_{\varphi}(\varepsilon)$, as $\varepsilon \leq \gamma_0$, and the statement of the theorem follows.

Theorem 1.2 has been mainly used in the literature with a choice of the metric space (T, d) of the form: $T = \{a_i \leq t_i \leq b_i, i = 1, 2\}$ and $d(t, s) = \max_{i=1,2} |t_i - s_i|, t = (t_1, t_2), s = (s_1, s_2)$. (see, for example, [3, 14, 10] for application to the analysis of solutions to heat equation and higher order heat-type equations with random initial conditions).

In the paper we study a particular class of φ -sub-Gaussian random fields with $\varphi = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$, which as a natural generalization of Gaussian and sub-Gaussian random fileds. Gaussian and sub-Gaussian cases are involved in our consideration, with the choice $\alpha = 2$. We study the sample paths of such fields for the case of the parameter set T of the form $T = [a_1, b_1] \times [a_2, b_2]$ or $T = [0, +\infty) \times [-A, A]$ with the so-called anisotropic metric $d(t, s) = \sum_{i=1,2} |t_i - s_i|^{H_i}$, $H_i \in (0, 1]$, i = 1, 2. Theorem 1.2 will serve as the main tool in our study.

Stochastic heat equations with fractional noises. Stochastic heat equations have been studied in various settings: with a time-space white noise, with generalizations of noise in space and/or in time, and also by considering differential operators more general than the Laplacian. The case of fractional noises was considered, e.g., in the recent papers [2], [8], among many others (see references therein). In the present paper we consider the stochastic heat equation:

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 W}{\partial t \partial x}, \quad (t, x) \in (0, T] \times \mathbb{R}, \tag{1.15}$$

$$u(0,x) = u_0(x), x \in \mathbb{R},$$
 (1.16)

where W is a centered Gaussian process which is white in time and is fractional Brownian motion in space with index $H \leq \frac{1}{2}$. The problem (1.15)–(1.16) was considered, for example, in [8] and it was stated that under some continuity and boundedness conditions on $u_0(x)$ there exists a unique mild solution $u(t, x), (t, x) \in (0, T] \times \mathbb{R}$, satisfying the Hölder condition

$$\|u(t,x) - u(s,y)\|_{L^p} \le c\Delta((t,x),(s,y))^{\rho \wedge H}$$
(1.17)

with some constant c = c(p, T, H) and where $\Delta((t, x), (s, y)) = |t - s|^{\frac{1}{2}} + |x - y|$ is a parabolic metric, ρ is an index in the Hölder condition imposed on $u_0(x)$. For our consideration, the bounds for the increments u(t, x) - u(s, y) in L_2 norm will be important. In view of this, we restate the bound (1.17) in another form for the case of p = 2, with the constant c given by a closed expression, and then use this bound to derive the results on the distribution of suprema and on the rate of growth for random fields representing the solution.

Contents. The paper is organized as follows. In Section 2 we study φ -sub-Gaussian random fields with $\varphi = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$. In Section 2.1 we present the estimates for the tail distribution of suprema on the bounded domain and in Section 2.2 we state the results on the rate of growth of random fields over unbounded domains. Section 3 presents applications of the results of sections 2.1 and 2.2 to random fields related to stochastic heat equations with fractional noise.

2 $Sub_{\varphi}(\Omega)$ processes with $\varphi = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$, defined on spaces with anisotropic metrics

Consider the process X(t), $t \in T$, from the class of φ -sub-Gaussian processes with $\varphi = \frac{|x|^{\alpha}}{\alpha}$, $1 < \alpha \leq 2$. This class is a natural generalisation of Gaussian and sub-Gaussian processes, which correspond to $\alpha = 2$.

For the function $\varphi(x) = \frac{|x|^{\alpha}}{\alpha}$, $1 < \alpha \leq 2$, we have $\varphi^{(-1)}(x) = (\alpha x)^{1/\alpha}$, x > 0, and the Young-Fenchel transform $\varphi^*(x) = \frac{|x|^{\beta}}{\beta}$, where $\beta > 0$ is such that $\frac{1}{\beta} + \frac{1}{\alpha} = 1$, that is, $\beta = \frac{\alpha}{\alpha-1}$.

The entropy integrals (1.4) and (1.10) become of the form

$$I_{\varphi}(\varepsilon) = \int_{0}^{\varepsilon} \left(\ln(N_{\rho_X}(u))) \right)^{\frac{1}{\beta}} du$$

and

$$\widetilde{I}_{\varphi}(\varepsilon) = \int_{0}^{\varepsilon} \left(\ln(N_d(\sigma^{(-1)}(u))) \right)^{\frac{1}{\beta}} du, \qquad (2.1)$$

and the bounds in Theorems 1.1 and 1.2 will be based on these integrals.

As we can see, for such function φ the integrals appear in a quite simple form and can be evaluated for particular metrics d. Note that for more general φ sometimes it is more convenient to use entropy integrals of another form (see, e.g., [9, 13, 15]).

We next consider φ -sub-Gaussian fields $X(t), t \in T$, with the parameter set $T = [a_1, b_1] \times [a_2, b_2]$ endowed with the anisotropic metric

$$d(t,s) = \sum_{i=1,2} |t_i - s_i|^{H_i}, H_i \in (0,1], i = 1,2.$$
(2.2)

2.1Estimates for the distribution of suprema

Theorem 2.1. Let $X(t), t \in T$, be a φ -sub-Gaussian field with $\varphi(x) = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$, $T = [a_1, b_1] \times [a_2, b_2]$ with the metric d(t, s) defined by (2.2), $\beta = \frac{\alpha}{\alpha - 1}$. Suppose that the field X satisfies conditions (i)-(iii) of Theorem 1.2 with $\sigma(h) = ch^{\gamma}, c > 0, \gamma \in (0, 1]$ and $\gamma \beta \neq 1$.

Then for all $\lambda > 0$ and $\theta \in (0, 1)$

$$\operatorname{\mathsf{E}exp}\left\{\lambda\sup_{t\in T}|X(t)|\right\} \le 2\exp\left\{\frac{1}{\alpha}\left(\frac{\lambda\varepsilon_0}{1-\theta}\right)^{\alpha} + \frac{2\lambda}{\theta(1-\theta)}\left(\theta\varepsilon_0\right)^{1-\frac{1}{\gamma\beta}}c_1\right\}$$
(2.3)

and for all $\theta \in (0,1), \theta \varepsilon_0 < \gamma_0$ and $u > \frac{2}{\theta(1-\theta)} (\theta \varepsilon_0)^{1-\frac{1}{\gamma\beta}} c_1$, we have

$$\mathsf{P}\{\sup_{t\in T} |X(t)| > u\} \le 2\exp\Big\{-\frac{1}{\beta}\Big(\frac{u(1-\theta)}{\varepsilon_0} - \frac{2}{\theta}\big(\theta\varepsilon_0\big)^{1-\frac{1}{\gamma\beta}}c_1\Big)^{\beta}\Big\},\tag{2.4}$$

where $c_1 = \frac{2^{\frac{1}{\beta}} c^{\frac{1}{\gamma\beta}}}{1 - \frac{1}{\gamma\beta}} \sum_{i=1,2} \frac{1}{H_i} \left(\frac{T_i}{2}\right)^{\frac{H_i}{\beta}}$, with $T_i = b_i - a_i$, i = 1, 2.

Proof. We apply Theorem 1.2. We need to estimate the entropy integral $\widetilde{I}_{\varphi}(\varepsilon)$ given by (1.10) for the particular φ, σ, d under consideration. For the metric d given by (2.2) we can write the following estimate for the metric massivenes

$$N_d(\varepsilon) \le \prod_{i=1,2} \left(\frac{T_i}{2(\frac{\varepsilon}{2})^{\frac{1}{H_i}}} + 1 \right) = \prod_{i=1,2} \left(\frac{2^{\frac{1}{H_i}} T_i}{2\varepsilon^{\frac{1}{H_i}}} + 1 \right).$$

This estimate can be deduced from the observation that a rectangle $\left[-\left(\frac{\varepsilon}{2}\right)^{\frac{1}{H_1}}, \left(\frac{\varepsilon}{2}\right)^{\frac{1}{H_1}}\right] \times$ $\left[-\left(\frac{\varepsilon}{2}\right)^{\frac{1}{H_2}}, \left(\frac{\varepsilon}{2}\right)^{\frac{1}{H_2}}\right]$ is contained in the ball in metric d with center (0,0) and radius ε , which is given as $B(\varepsilon) = \{(x_1, x_2) : |x_1|^{H_1} + |x_2|^{H_2} \le \varepsilon\}.$

Now, for the given functions φ and σ (note that $\sigma^{(-1)}(u) = (\frac{u}{c})^{\frac{1}{\gamma}}$), we consider the entropy integral (2.1):

$$\widetilde{I}_{\varphi}(\varepsilon) = \int_{0}^{\varepsilon} \left(\ln(N_{d}(\sigma^{(-1)}(u))) \right)^{\frac{1}{\beta}} du \leq \int_{0}^{\varepsilon} \left(\ln\prod_{i=1,2} \left(\frac{2^{\frac{1}{H_{i}}}T_{i}}{2(\sigma^{(-1)}(u))^{\frac{1}{H_{i}}}} + 1 \right) \right)^{\frac{1}{\beta}} du$$
$$= \int_{0}^{\varepsilon} \left(\ln\prod_{i=1,2} \left(\frac{T_{i}2^{\frac{1}{H_{i}}}c^{\frac{1}{\gamma H_{i}}}}{2u^{\frac{1}{\gamma H_{i}}}} + 1 \right) \right)^{\frac{1}{\beta}} du.$$

For any $0 < \varkappa \leq 1$, $\ln(1+x) = \frac{1}{\varkappa} \ln(1+x)^{\varkappa} \leq \frac{x^{\varkappa}}{\varkappa}$, we apply this inequality for each term in the product in the above formula choosing $\varkappa = H_i$, i = 1, 2:

$$\begin{split} \widetilde{I}_{\varphi}(\varepsilon) &\leq \int_{0}^{\varepsilon} \sum_{i=1,2} \frac{1}{H_{i}} \Big(\frac{T_{i} 2^{\frac{1}{H_{i}}} c^{\frac{1}{\gamma H_{i}}}}{2u^{\frac{1}{\gamma H_{i}}}} \Big)^{\frac{H_{i}}{\beta}} du = \sum_{i=1,2} \int_{0}^{\varepsilon} \frac{1}{H_{i}} \Big(\frac{T_{i}}{2} \Big)^{\frac{H_{i}}{\beta}} 2^{\frac{1}{\beta}} c^{\frac{1}{\gamma \beta}} \frac{du}{u^{\frac{1}{\gamma \beta}}} \\ &= \frac{\varepsilon^{1-\frac{1}{\gamma \beta}}}{1-\frac{1}{\gamma \beta}} 2^{\frac{1}{\beta}} c^{\frac{1}{\gamma \beta}} \sum_{i=1,2} \frac{1}{H_{i}} \Big(\frac{T_{i}}{2} \Big)^{\frac{H_{i}}{\beta}} = \varepsilon^{1-\frac{1}{\gamma \beta}} c_{1}. \end{split}$$

2.2 Estimates for the rate of growth

Consider now the field $X(t_1, t_2), (t_1, t_2) \in V$, defined over the unbounded domain $V = [0, +\infty) \times [-A, A]$.

Let $f(t) > 0, t \ge 0$, be a continuous strictly increasing function and $f(t) \to \infty$ as $t \to \infty$. Introduce the sequence $b_0 = 0, b_{k+1} > b_k, b_k \to \infty, k \to \infty$. We will use the following notations: $l_k = b_{k+1} - b_k, V_k = [b_k, b_{k+1}] \times [-A, A], k = 0, 1, \dots, f_k = f(b_k),$ $\varepsilon_k = \sup_{(t_1, t_2) \in V_k} \tau_{\varphi}(X(t_1, t_2)), \text{ and suppose that } 0 < \varepsilon_k < \infty;$ $\gamma_k = c_k \max_{(t_1, t_2), (s_1, s_2) \in V_k} (d((t_1, t_2), (s_1, s_2)))^{\gamma}, \text{ where } c_k \text{ are from } (2.5) \text{ below, } \tilde{\theta} = \inf_k \frac{\gamma_k}{\varepsilon_k},$ $\beta = \frac{\alpha}{\alpha - 1}.$

Theorem 2.2. Let $X(t_1, t_2)$, $(t_1, t_2) \in V$, be a φ -sub-Gaussian separable field with $\varphi = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$. Suppose further that

(i)

$$\sup_{\substack{d((t_1,t_2),(s_1,s_2)) \le h, \\ (t_1,t_2),(s_1,s_2) \in V_k}} \tau_{\varphi}(X(t_1,t_2) - X(s_1,s_2)) \le c_k h^{\gamma}, c_k > 0, 0 < \gamma \le 1;$$
(2.5)

(*ii*)
$$C = \sum_{k=0}^{\infty} \frac{\varepsilon_k}{f_k} < \infty.$$

(*iii*) $S = \sum_{k=0}^{\infty} \frac{\varepsilon_k^{1-\frac{1}{\gamma\beta}} c_1(k)}{f_k} < \infty, \text{ where } c_1(k) = \left(\frac{1}{H_1} (l_k/2)^{\frac{H_1}{\beta}} + \frac{1}{H_2} A^{\frac{H_2}{\beta}}\right) \frac{2^{\frac{1}{\beta}} c_k^{\frac{1}{\gamma\beta}}}{1-\frac{1}{\gamma\beta}}.$

Then

(i) for any
$$\theta \in (0, \min(1, \tilde{\theta}))$$
 and any $\lambda > 0$

$$\operatorname{\mathsf{E}exp}\left\{\lambda\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)}\right\} \le 2\exp\left\{\frac{\lambda^{\alpha}}{\alpha}\left(\frac{C}{1-\theta}\right)^{\alpha} + \frac{2\lambda}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}}S\right\};\tag{2.6}$$

(ii) for any
$$\theta \in (0, \min(1, \widetilde{\theta}))$$
 and $u > \frac{2S}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}}$

$$\mathsf{P}\left\{\sup_{(t_1, t_2)\in V} \frac{|X(t_1, t_2)|}{f(t_1)} > u\right\} \le 2\exp\left\{-\frac{1}{\beta C^{\beta}}\left(u(1-\theta) - 2S\theta^{-\frac{1}{\gamma\beta}}\right)^{\beta}\right\}.$$
(2.7)

Proof. We use the scheme of the proof which is similar that in [6] (Theorem 2.4), [9] (Theorem 4). Using (2.3) we can write the estimate:

$$\begin{split} I(\lambda) &= \mathsf{E} \exp\left\{\lambda \sup_{(t_1, t_2) \in V} \frac{|X(t_1, t_2)|}{f(t_1)}\right\} \le \mathsf{E} \exp\left\{\lambda \sum_{k=0}^{\infty} \sup_{(t_1, t_2) \in V_k} \frac{|X(t_1, t_2)|}{f_k}\right\} \le \\ &\leq \prod_{k=0}^{\infty} \left(\mathsf{E} \exp\left\{\lambda \frac{r_k}{f_k} \sup_{(t_1, t_2) \in V_k} |X(t_1, t_2)|\right\}\right)^{\frac{1}{r_k}} \le 2 \prod_{k=0}^{\infty} Q_k(\lambda, \theta)^{\frac{1}{r_k}}, \end{split}$$

where

$$Q_k(\lambda,\theta) = \exp\left\{\frac{1}{\alpha} \left(\frac{\lambda\varepsilon_k r_k}{(1-\theta)f_k}\right)^{\alpha} + 2\lambda \frac{r_k}{f_k} \frac{1}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}} \varepsilon_k^{1-\frac{1}{\gamma\beta}} c_1(k)\right\},\,$$

$$c_1(k) = \left(\frac{1}{H_1}(l_k/2)^{\frac{H_1}{\beta}} + \frac{1}{H_2}A^{\frac{H_2}{\beta}}\right)\frac{2^{\frac{1}{\beta}}c_k^{\frac{\gamma}{\gamma\beta}}}{1 - \frac{1}{\gamma\beta}}$$

and let here $r_k, k \ge 0$ are such that $\sum_{k=0}^{\infty} \frac{1}{r_k} = 1$. This implies that

$$I(\lambda) \le 2 \exp\left\{\frac{1}{\alpha} \left(\frac{\lambda}{1-\theta}\right)^{\alpha} \sum_{k=0}^{\infty} \left(\frac{\varepsilon_k r_k}{f_k}\right)^{\alpha} r_k^{-1} + \frac{2\lambda}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}} \sum_{k=0}^{\infty} \frac{\varepsilon_k^{1-\frac{1}{\gamma\beta}} c_1(k)}{f_k}\right\}.$$

Choose $r_k = \frac{f_k}{\varepsilon_k}C$, where $C = \sum_{k=0}^{\infty} \frac{\varepsilon_k}{f_k}$, then

$$I(\lambda) \le 2 \exp\left\{\frac{1}{\alpha} \left(\frac{\lambda}{1-\theta}\right)^{\alpha} C^{\alpha} + \frac{2\lambda}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}} \sum_{k=0}^{\infty} \frac{\varepsilon_k^{1-\frac{1}{\gamma\beta}} c_1(k)}{f_k}\right\}.$$

Therefore,

$$\mathsf{E}\exp\left\{\lambda\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)}\right\} \le 2\exp\left\{\frac{\lambda^{\alpha}}{\alpha}\left(\frac{C}{1-\theta}\right)^{\alpha} + \frac{2\lambda}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}}S\right\},$$

and for all $\lambda > 0, \, u > 0, \, 0 < \theta < \widetilde{\theta}$

$$P\left\{\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)} > u\right\} \le \exp\{-\lambda u\} \mathsf{E}\exp\left\{\lambda \sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)}\right\} \le \\ \le 2\exp\left\{\frac{\lambda^{\alpha}}{\alpha} \left(\frac{C}{1-\theta}\right)^{\alpha} + \frac{2\lambda S}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}} - \lambda u\right\}.$$

We minimise the right-hand side with respect to λ and obtain for $u > \frac{2S}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}}$

$$P\left\{\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)} > u\right\} \le 2\exp\left\{-\left(u - \frac{2S}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}}\right)^{\frac{\alpha}{\alpha-1}}\left(\frac{C}{1-\theta}\right)^{-\frac{\alpha}{\alpha-1}}\frac{\alpha-1}{\alpha}\right\} = 2\exp\left\{-\frac{\alpha-1}{\alpha}C^{-\frac{\alpha}{\alpha-1}}\left(u(1-\theta) - 2S\theta^{-\frac{1}{\gamma\beta}}\right)^{\frac{\alpha}{\alpha-1}}\right\}.$$

Theorem 2.3. Let $X(t_1, t_2)$, $(t_1, t_2) \in V$, be a φ -sub-Gaussian separable field with $\varphi = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$. Suppose further that

(i)

$$\sup_{\substack{d((t_1,t_2),(s_1,s_2)) \le h, \\ (t_1,t_2),(s_1,s_2) \in V_k}} \tau_{\varphi}(X(t_1,t_2) - X(s_1,s_2)) \le c_k h^{\gamma}, c_k > 0, 0 < \gamma \le 1;$$

(ii) there exists $\delta > 0$ and a constant $c(\delta)$ such that for $(t_1, t_2) \in V$

$$\tau_{\varphi}(X(t_1, t_2)) \le c(\delta) t_1^{\delta};$$

(iii) $\widetilde{C} = c(\delta) \sum_{k=0}^{\infty} \frac{b_{k+1}^{\delta}}{f_k} < \infty$,

$$(iv) \ \widetilde{S} = (c(\delta))^{1 - \frac{1}{\gamma\beta}} \sum_{k=0}^{\infty} \frac{b_{k+1}^{1 - \frac{1}{\gamma\beta}} c_1(k)}{f_k} < \infty, \ where \ c_1(k) = \left(\frac{1}{H_1} (l_k/2)^{\frac{H_1}{\beta}} + \frac{1}{H_2} A^{\frac{H_2}{\beta}}\right) \frac{2^{\frac{1}{\beta}} c_k^{\frac{1}{\gamma\beta}}}{1 - \frac{1}{\gamma\beta}}.$$

Then

(i) for any $\theta \in (0, \min(1, \widetilde{\theta}))$ and any $\lambda > 0$

$$\operatorname{\mathsf{E}exp}\left\{\lambda\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)}\right\} \le 2\exp\left\{\frac{\lambda^{\alpha}}{\alpha}\left(\frac{\widetilde{C}}{1-\theta}\right)^{\alpha} + \frac{2\lambda}{(1-\theta)\theta}\widetilde{S}\right\}; \qquad (2.8)$$

(*ii*) for any $\theta \in (0, \min(1, \widetilde{\theta}))$ and $u > \frac{2(c(\delta))^{1-\frac{1}{\gamma\beta}}\widetilde{S}}{(1-\theta)\theta^{\frac{1}{\gamma\beta}}}$

$$\mathsf{P}\left\{\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)} > u\right\} \le 2\exp\left\{-\frac{1}{\beta\widetilde{C}^{\beta}}\left(u(1-\theta) - 2\widetilde{S}\theta^{\frac{1}{\gamma\beta}}\right)^{\beta}\right\}.$$
 (2.9)

The proof is analogous to that of Theorem 2.2.

Remark 2.1. The bound (2.7) can be rewritten in another form. We can choose $\theta = u^{-\frac{\gamma\beta}{\gamma\beta+1}}$ and then under conditions of Theorem 2.1 for $u > (1+2S)^{\frac{\gamma\beta}{\gamma\beta+1}}$ the following bound holds

$$\mathsf{P}\left\{\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)} > u\right\} \le 2\exp\left\{-\frac{1}{\beta C^{\beta}}\left(u - u^{\frac{1}{\gamma\beta+1}}(1+2S)\right)^{\beta}\right\}.$$

Correspondingly, under conditions of Theorem 2.3 for $u > (1 + 2\tilde{S})^{\frac{\gamma\beta}{\gamma\beta+1}}$ we can write the bound

$$\mathsf{P}\left\{\sup_{(t_1,t_2)\in V}\frac{|X(t_1,t_2)|}{f(t_1)} > u\right\} \le 2\exp\left\{-\frac{1}{\beta\widetilde{C}^{\beta}}\left(u - u^{\frac{1}{\gamma\beta+1}}(1+2\widetilde{S})\right)^{\beta}\right\}.$$

Theorem 2.4. Suppose that for the field $X(t_1, t_2)$, $(t_1, t_2) \in V$, and function f conditions of Theorem 2.3 are satisfied. Then there exists a random variable ξ such that with probability one

$$|X(t_1, t_2)| \le f(t_1)\xi, \tag{2.10}$$

and ξ satisfies the following assumption:

$$\mathsf{P}\left\{\xi > u\right\} \le 2\exp\left\{-\frac{1}{\beta \widetilde{C}^{\beta}}\left(u - u^{\frac{1}{\gamma\beta+1}}(1+2\widetilde{S})\right)^{\beta}\right\}$$
(2.11)

for $u > (1+2\widetilde{S})^{\frac{\gamma\beta}{\gamma\beta+1}}$.

Proof. The theorem is a corollary of Theorem 2.3. We denote $\xi = \sup_{(t_1,t_2)\in V} \frac{|X(t_1,t_2)|}{f(t_1)}$, then ξ satisfies (2.11) and for all ω we have (2.10).

3 Application to stochastic heat equation with fractional noise

Consider the stochastic heat equation:

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 W}{\partial t \partial x}, \quad (t, x) \in (0, T] \times \mathbb{R}, \tag{3.1}$$

$$u(0,x) = u_0(x), x \in \mathbb{R}, \tag{3.2}$$

with the following assumption about the noise:

A.1. W is a centered Gaussian process which is white in time and is fractional Brownian motion in space with index $H \leq \frac{1}{2}$.

Note that there exists an extensive literature on SPDEs driven by Gaussian noises and involving various differential operators.

As for the stochastic heat equation, it has been studied in various settings: as that one driven by time-space white noise, further with generalization of noise in space or in time, and also by considering differential operators more general than the Laplacian.

Since we are going to apply the results of the previous section, we are interested in the results for solution to the problem (3.1)-(3.2), which allow to write down the bounds for $\mathsf{E}|u(t,x) - u(s,y)|^2$.

We will base our study on the results from the recent papers [2, 8] on regularity properties of solution to stochastic heat equation.

We consider the problem (3.1)-(3.2) with assumption A.1. on the noise and imposing the following assumption on the initial condition u_0 .

A.2. The process $u_0(x), x \in \mathbb{R}$, is continuous, possesses uniformly bounded *p*-th moments for $p \geq 2$ and is stochastically Hölder continuous with $\rho \in (0, 1]$, i.e., for all $p \geq 1$

$$||u_0(x) - u_0(y)||_{L^p} \le L_0(p)|x - y|^{\rho}, \quad x, y \in \mathbb{R},$$

where $L_0(p)$ is a constant and $\|.\|_{L^p}$ denotes the norm in $L^p(\Omega, \mathbb{R})$: $\|u\|_{L^p} = (\mathsf{E}|u|^p)^{\frac{1}{p}}, p \ge 1.$

From [8] (see Theorem 1.1) it follows that under assumptions A.1. and A.2. equation (3.1) has a unique mild solution $u(t,x), (t,x) \in (0,T] \times \mathbb{R}$, satisfying $\sup_{(t,x)\in(0,T]\times\mathbb{R}} \mathsf{E}|u(t,x)|^p < \infty$ and the Hölder are different.

the Hölder condition

$$||u(t,x) - u(s,y)||_{L^p} \le c\Delta((t,x),(s,y))^{\rho \wedge H}$$

with some constant c = c(p, T, H) and where $\Delta((t, x), (s, y)) = |t - s|^{\frac{1}{2}} + |x - y|$ is the parabolic metric.

The mild solution is defined as the random field

$$u(t,x) = \int_{\mathbb{R}} G_t(x-y)u_0(y) \, dy + \int_0^t \int_{\mathbb{R}} G_{t-\theta}(x-\eta)W(d\theta, d\eta) = \omega(t,x) + V(t,x), \quad (3.3)$$

where $G_t(x)$ is the Green's function (fundamental solution) of the equation $\left(\frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2}\right)u = 0$, that is,

$$G_t(x) = \frac{1}{\sqrt{4\pi t}} \exp\left(-\frac{|x|^2}{4t}\right).$$

We refer for the rigorous definitions of the integrals in (3.3), for example, to [8], [2].

In the next theorem we state the result on Hölder continuity of the fields $\omega(t, x)$ and V(t, x) in the form, where explicit expressions for the constants are given.

Theorem 3.1. Let assumptions A.1. and A.2. hold. Then the following bounds hold:

$$\|\omega(t,x)\|_{L^2} \le \sup_{x \in \mathbb{R}} \|u_0(x)\|_{L^2} \le c_0;$$
(3.4)

$$\|V(t,x)\|_{L^2} \le A(H)t^{\frac{H}{2}}; \tag{3.5}$$

$$\|\omega(t,x) - \omega(s,y)\|_{L^2} \le c_\omega \left(|t-s|^{\frac{\rho}{2}} + |x-y|^{\rho}\right);$$
(3.6)

$$\|V(t,x) - V(s,y)\|_{L^2} \le c_V \left(|t-s|^{\frac{H}{2}} + |x-y|^H\right),\tag{3.7}$$

where the constants $c_{\omega}, c_V, A(H)$ are given by formulas (3.8), (3.9), (3.10) below.

Proof. For the proof we use the reasoning similar to that applied for more general case in [8] (Theorem 1.1) (see also [2]). Therefore, we present only the main steps, our interest is in keeping the values of constants involved on each step through all the chain of bounds. We have

$$\begin{split} \mathsf{E}|\omega(t,x) - \omega(s,x)|^2 &= \mathsf{E} \left| \int_{\mathbb{R}} \left(G_t(x-y) - G_s(x-y) \right) u_0(y) \, dy \right|^2 \\ &= \mathsf{E} \left| \int_{\mathbb{R}} G_{t-s}(y) \left(\int_{\mathbb{R}} G_s(x-z) (u_0(z-y) - u_0(z)) \, dz \right) \, dy \right|^2 \\ &\leq \int_{\mathbb{R}} G_{t-s}(y) \int_{\mathbb{R}} G_s(x-z) \mathsf{E} \left| u_0(z-y) - u_0(z) \right|^2 \, dz dy \\ &\leq L \int_{\mathbb{R}} G_{t-s}(y) |y|^{2\rho} \, dy = L C_1 |t-s|^{\rho}, \end{split}$$

since

$$\int_{\mathbb{R}} G_h(y) |y|^{2\rho} \, dy = \int_{\mathbb{R}} \frac{1}{\sqrt{4\pi h}} \exp\left(-\frac{y^2}{4h}\right) |y|^{2\rho} \, dy = \frac{1}{\sqrt{4\pi h}} 2 \int_0^\infty \exp\left(-\frac{y^2}{4h}\right) y^{2\rho} \, dy$$
$$= \frac{4^{\rho}}{\sqrt{\pi}} \Gamma(\rho + \frac{1}{2}) h^{\rho} = C_1 h^{\rho},$$

where $C_1 = \frac{4^{\rho}}{\sqrt{\pi}} \Gamma(\rho + \frac{1}{2})$, $L = L_0(2)$, and we have used above the property $\int_{\mathbb{R}} G_t(y) \, dy = 1$ and assumption A.2. We now consider the more general increment:

$$\begin{split} \mathsf{E}|\omega(t,x) - \omega(s,y)|^2 &\leq 2\left(\mathsf{E}|\omega(t,x) - \omega(s,x)|^2 + \mathsf{E}|\omega(s,x) - \omega(s,y)|^2\right) \\ &\leq 2LC_1|t-s|^\rho + 2\mathsf{E}\left|\int_{\mathbb{R}} \left(G_s(x-z) - G_s(y-z)\right)u_0(z)\,dz\right|^2 \\ &= 2LC_1|t-s|^\rho + 2\mathsf{E}\left|\int_{\mathbb{R}} G_s(z)\left(u_0(x-z) - u_0(y-z)\right)\,dz\right|^2 \\ &\leq 2LC_1|t-s|^\rho + 2L^2|x-y|^{2\rho} \leq c\left(|t-s|^\rho + |x-y|^{2\rho}\right), \end{split}$$

where $c = 2L \max(C_1, L)$, and using again assumption A.2. Therefore,

$$\left(\mathsf{E}|\omega(t,x) - \omega(s,y)|^2\right)^{\frac{1}{2}} \le c_{\omega} \left(|t-s|^{\frac{\rho}{2}} + |x-y|^{\rho}\right), \ c_{\omega} = \sqrt{c}.$$
(3.8)

Considering V(t, x), we follow again the similar lines as those in [8] keeping the track of constants. We can write:

$$\begin{split} \mathsf{E} \left| V(t,x) - V(s,y) \right|^2 = & \mathsf{E} \left| \int_0^t \int_{\mathbb{R}} G_{t-\theta}(x-\eta) W(d\theta, d\eta) - \int_0^s \int_{\mathbb{R}} G_{s-\theta}(y-\eta) W(d\theta, d\eta) \right|^2 \\ = & \mathsf{E} \left| \int_0^t \int_{\mathbb{R}} G_{t-\theta}(x-\eta) W(d\theta, d\eta) \right| \\ & + \int_0^s \int_{\mathbb{R}} \left(G_{t-\theta}(x-\eta) - G_{s-\theta}(x-\eta) \right) W(d\theta, d\eta) \\ & + \int_0^s \int_{\mathbb{R}} \left(G_{s-\theta}(x-\eta) - G_{s-\theta}(y-\eta) \right) W(d\theta, d\eta) \Big|^2 \\ \leq & 3C_H \left(\int_0^{t-s} \int_{\mathbb{R}} |FG_{\theta}(\xi)|^2 |\xi|^{1-2H} d\xi d\theta \\ & + \int_0^s \int_{\mathbb{R}} |FG_{t-s+\theta}(\xi) - FG_{\theta}(\xi)|^2 |\xi|^{1-2H} d\xi d\theta \\ & + \int_0^s \int_{\mathbb{R}} (1 - \cos(\xi(x-y))) |FG_{\theta}(\xi)|^2 |\xi|^{1-2H} d\xi d\theta \Big) \\ = & 3C_H (I_1 + I_2 + I_3), \end{split}$$

here above by F attached to a function we mean the Fourier transform, we use that $FG_t(\xi) = \exp(-t\xi^2)$ and also the isometry relation (see, e.g., [8, Theorem 2.1], [2, Theorem 2.7]):

$$\mathsf{E}\left|\int_{0}^{t}\int_{\mathbb{R}}\varphi(s,y)W(ds,dy)\right|^{2} = \int_{0}^{t}\int_{\mathbb{R}}\left|F\varphi(s,\cdot)(y)\right|^{2}\mu(dy)ds,$$

where μ is the spectral measure involved in the representation of the covariance of W and

$$\mu(dy) = C_H |y|^{1-2H} dy, \ C_H = \frac{\Gamma(2H+1)\sin(\pi H)}{2\pi}$$

(see for details [8], [2]). Evaluate now the integrals I_1, I_2, I_3 .

$$I_1 = \int_0^t \int_{\mathbb{R}} \exp(-2s\xi^2) |\xi|^{1-2H} d\xi ds = 2^{-(H+1)} \frac{\Gamma(H+1)}{H} t^H = c_{1,H} t^H,$$

where $c_{1,H} = 2^{-(H+1)} \frac{\Gamma(H+1)}{H};$

$$\begin{split} I_2 &= \int_0^s \int_{\mathbb{R}} |FG_{t-s+\theta}(\xi) - FG_{\theta}(\xi)|^2 |\xi|^{1-2H} d\xi d\theta \\ &= \int_0^s \int_{\mathbb{R}} \exp(-2\theta\xi^2) \left(1 - \exp(-(t-s)\xi^2)\right)^2 \times |\xi|^{1-2H} d\xi d\theta \\ &= \int_{\mathbb{R}} \frac{1 - \exp(-2s\xi^2)}{2|\xi|^2} \left(1 - \exp(-(t-s)\xi^2)\right)^2 |\xi|^{1-2H} d\xi \\ &\leq \frac{1}{2} \int_{\mathbb{R}} \frac{\left(1 - \exp(-(t-s)\xi^2)\right)^2}{|\xi|^{2-1+2H}} d\xi \\ &= (t-s)^H \frac{1}{2} \int_{\mathbb{R}} \frac{\left(1 - \exp(-u^2)\right)^2}{u^{1+2H}} du = c_{2,H}(t-s)^H, \end{split}$$

where $c_{2,H} = \frac{1}{2} \int_{\mathbb{R}} \frac{(1 - \exp(-u^2))^2}{u^{1+2H}} du;$

$$I_{3} = \int_{0}^{s} \int_{\mathbb{R}} (1 - \cos(\xi(x - y))) \exp(-2\theta\xi^{2}) |\xi|^{1-2H} d\xi d\theta$$

= $\int_{\mathbb{R}} \frac{1 - \exp(-2s\xi^{2})}{2|\xi|^{2}} (1 - \cos(\xi(x - y))) |\xi|^{1-2H} d\xi$
 $\leq \int_{\mathbb{R}} (1 - \cos(\xi(x - y))) \frac{1}{2} |\xi|^{-1-2H} d\xi = c_{3,H} (x - y)^{2H}$

where

for
$$H < \frac{1}{2}$$
: $c_{3,H} = \int_0^\infty (1 - \cos x) x^{-1-2H} dx = (2H)^{-1} \Gamma(1 - 2H) \cos(H\pi)$,
for $H = \frac{1}{2}$: $c_{3,H} = \frac{\pi}{2}$.

(We used here Lemma D.1. from $\lfloor 2 \rfloor$).

Therefore,

$$\mathsf{E} |V(t,x) - V(s,y)|^2 \le 3C_H((c_{1,H} + c_{2,H})|t - s|^H + c_{3,H}|x - y|^{2H})$$

and

$$\left(\mathsf{E} |V(t,x) - V(s,y)|^2\right)^{\frac{1}{2}} \le c_V \left(|t-s|^{\frac{H}{2}} + |x-y|^H\right),$$

where

$$c_V = \sqrt{3C_H \max((c_{1,H} + c_{2,H}), c_{3,H})}.$$
(3.9)

We also have the bound

$$\left(\mathsf{E} |V(t,x)|^2\right)^{\frac{1}{2}} \le A(H)t^{\frac{H}{2}}$$

where

$$A(H) = \sqrt{C_H c_{1,H}}.$$
(3.10)

For the next results we will use an additional assumption on the initial condition u_0 and will need the following definition([12]):

A family Δ of φ -sub-Gaussian random variables is called strictly φ -sub-Gaussian if there exists a constant C_{Δ} (called determining constant) such that for all countable sets I of random variables $\zeta_i \in \Delta$, $i \in I$, the inequality holds: $\tau_{\varphi} \left(\sum_{i \in I} \lambda_i \zeta_i \right) \leq C_{\Delta} \left(\mathsf{E} \left(\sum_{i \in I} \lambda_i \zeta_i \right)^2 \right)^{1/2}$. Random field $\zeta(t)$, $t \in T$, is called strictly φ -sub-Gaussian if the family of random variables $\{\zeta(t), t \in T\}$ is strictly φ -sub-Gaussian.

Let K be a deterministic kernel and $\eta(x) = \int_T K(x, y) d\xi(y), x \in X$, where $\xi(y), y \in Y$, is a strictly φ -sub-Gaussian field and the integral is defined in the mean-square sense. Then $\eta(x)$, $x \in X$, is strictly φ -sub-Gaussian field with the same determining constant (see [12]).

Theorem 3.2. Let assumptions A.1 and A.2 hold and assume that $u_0(x)$ is strictly φ -sub-Gaussian random field with $\varphi(x) = \frac{|x|^{\alpha}}{\alpha}$, $\alpha \in (1, 2]$, and let c_{φ} be the determining constant. Then for fields $\omega(t, x)$ and V(t, x), $(t, x) \in D = [a_1, b_1] \times [a_2, b_2]$, the following estimates hold:

(i)

$$\mathsf{P}\left\{\sup_{(t,x)\in D}|\omega(t,x)| > u\right\} \leq 2\exp\left\{-\frac{1}{\beta}\left(\frac{u(1-\theta)}{c_0c_{\varphi}} + 2(\theta c_0c_{\varphi})^{-\frac{1}{\beta}}\widetilde{c}_1\right)^{\beta}\right\}$$
(3.11)
for $u > \frac{2}{(1-\theta)\theta}(\theta c_0c_{\varphi})^{1-\frac{1}{\beta}}\widetilde{c}_1$, where $\widetilde{c}_1 = \frac{2^{\frac{1}{\beta}}(c_{\omega}c_{\varphi})^{\frac{1}{\beta}}}{1-\frac{1}{\beta}}\left(\frac{2}{\rho}\left(\frac{T_1}{2}\right)^{\frac{\rho}{2\beta}} + \frac{1}{\rho}\left(\frac{T_2}{2}\right)^{\frac{\rho}{\beta}}\right), T_i = b_i - a_i, i = 1, 2, \beta = \frac{\alpha}{\alpha-1}.$
(ii)

$$\mathsf{P}\left\{\sup_{(t,x)\in D}|V(t,x)| > u\right\} \leq 2\exp\left\{-\frac{1}{2}\left(\frac{u(1-\theta)}{\varepsilon_V} + 2(\theta\varepsilon_V)^{-\frac{1}{2}}\widetilde{c}_1\right)^2\right\}$$
for $u > \frac{2}{(1-\theta)\theta}(\theta\varepsilon_V)^{\frac{1}{2}}\widetilde{c}_1$, where $\varepsilon_V = A(H)b_1^{H/2}, \widetilde{c}_1 = 2\sqrt{2c_V}\left(\frac{2}{H}\left(\frac{T_1}{2}\right)^{\frac{H}{4}} + \frac{1}{H}\left(\frac{T_2}{2}\right)^{\frac{H}{2}}\right), T_i = b_i - a_i, i = 1, 2.$

Proof. Statement (i) follows from Theorem 2.1 and the estimates (3.4), (3.6) and we use that $\tau_{\varphi}(\omega(t,x)) \leq c_{\varphi} \|\omega(t,x)\|_{L^2}$. Statement (ii) follows from Theorem 2.1 with $\alpha = 2$ (since the field V(t, x) is Gaussian) and the estimate (3.7).

The next theorem presents the power upper bound for the asymptotic growth of the trajectories of the field V(t, x).

Theorem 3.3. Let assumptions A.1 and A.2 hold. Then for any p > 1 there exist a random variable $\xi(p)$ such that for any $(t, x) \in V$, with probability one

$$|V(t,x)| \le ((t^{\frac{H}{2}}(\log t)^p) \lor 1)\xi(p),$$

where $\xi(p)$ satisfies assumption (2.11) with $\gamma = 1, \beta = 2$ and some constants \widetilde{C} and \widetilde{S} .

Proof. We apply Theorem 2.4 and Theorem 2.3. Condition (i) holds since V(t, x) is continuous, condition (ii) holds in view of (3.5).

Consider conditions (*iii*) and (*iv*). Let $f(t) = (t^{\frac{H}{2}} |\log t|^p) \vee 1$ for t > 0 and some p > 1. Let us choose $b_k = e^k, k \ge 0$. Then $f_k = b_k^{\frac{H}{2}} (\log b_k)^p = e^{k\frac{H}{2}} (\log e^k)^p = e^{k\frac{H}{2}} k^p$ and we have

$$\widetilde{C} = A(H) \sum_{k=0}^{\infty} \frac{b_{k+1}^{\frac{H}{2}}}{f_k} = A(H) \left(e^{\frac{H}{2}} + \sum_{k=1}^{\infty} \frac{e^{\frac{H}{2}}}{k^p} \right) = A(H) e^{\frac{H}{2}} \left(1 + \sum_{k=1}^{\infty} \frac{1}{k^p} \right) < \infty.$$

Consider

$$\widetilde{S} = (A(H))^{\frac{1}{2}} \sum_{k=0}^{\infty} \frac{b^{\frac{H}{4}}_{k+1}c_1(k)}{f_k} = (A(H))^{\frac{1}{2}} \sum_{k=0}^{\infty} \frac{e^{(k+1)\frac{H}{4}}c_1(k)}{e^{k\frac{H}{2}}k^p},$$

where $c_1(k) = \frac{2}{H}e^{k\frac{H}{4}}(\frac{e-1}{2})^{\frac{H}{4}} + A^{\frac{H}{2}}$, and we can see that $\tilde{S} < \infty$. Therefore, conditions of Theorem 2.4 hold true, and applying this theorem with f(t) = $(t^{\frac{H}{2}}|\log t|^p) \vee 1$ we obtain the statement of Theorem 3.3.

We consider now an assumption on the initial condition which can be used instead of assumption A.2.

A.3. The process $u_0(x), x \in \mathbb{R}$ is a real, measurable, mean-square continuous stationary stochastic process.

Let $B(x), x \in \mathbb{R}$, is a covariance function of the process $u_0(x), x \in \mathbb{R}$, with the representation

$$B(x) = \int_{\mathbb{R}} \cos(\lambda x) dF(\lambda), \qquad (3.12)$$

where $F(\lambda)$ is a spectral measure, and for the process itself we can write the spectral representation

$$u_0(x) = \int_{\mathbb{R}} e^{i\lambda x} Z(d\lambda), \qquad (3.13)$$

The stochastic integral is considered as $L_2(\Omega)$ integral. Orthogonal random measure Z is such that $\mathsf{E}|Z(d\lambda)|^2 = F(d\lambda)$.

Then the field ω can be written in the form

$$\omega(t,x) = \int_{\mathbb{R}} \exp\left\{i\lambda x - \mu t\lambda^2\right\} Z(d\lambda)$$
(3.14)

and its covariance function has the following representation

$$Cov\Big(\omega(t,x),\omega(s,y)\Big) = \int_{\mathbb{R}} \exp\left\{i\lambda(x-y) - \mu\lambda^2(t-s)\right\} F(d\lambda)$$
(3.15)

(see, [9]).

Theorem 3.4. Let assumption A.3 holds. Then

$$\|\omega(t,x)\|_{L^2} \le \left(\int_{\mathbb{R}} F(d\lambda)\right)^{\frac{1}{2}},\tag{3.16}$$

and if for some $\varepsilon \in (0, \frac{1}{2}]$

$$c^{2}(\varepsilon) := \int_{\mathbb{R}} \lambda^{2\varepsilon} F(d\lambda) < \infty, \qquad (3.17)$$

then the following estimate holds:

$$\|\omega(t,x) - \omega(s,y)\|_{L^2} \le c(\varepsilon) \left(4^{1-\varepsilon} |x-y|^{2\varepsilon} + |t-s|^{\varepsilon}\right)^{1/2}.$$
(3.18)

Proof. We have

$$\mathsf{E}\big(\omega(t,x) - \omega(s,y)\big)^2 = \int_{\mathbb{R}} |b(\lambda)|^2 F(d\lambda),$$

where

$$b(\lambda) = \exp\{i\lambda x\} \exp\{-\mu\lambda^2 t\} - \exp\{i\lambda y\} \exp\{-\mu\lambda^2 s\},$$

and we can estimate

$$\begin{aligned} |b(\lambda)|^2 &\leq \left(1 - \exp\left\{-\mu\lambda^2|t-s|\right\}\right)^2 + 4\sin^2\left(\frac{1}{2}\lambda(x-y)\right) \\ &\leq \left(\min(\lambda^2|t-s|,1)\right)^2 + 4\min\left(\frac{1}{2}|\lambda||x-y|,1\right)^2 \\ &\leq \left(\lambda^2|t-s|\right)^{2\varepsilon_1} + 4\left(\frac{1}{2}|\lambda||x-y|\right)^{2\varepsilon_2} \end{aligned}$$

for any $\varepsilon_1, \varepsilon_2 \in (0, 1]$. Let us choose $\varepsilon := \varepsilon_1 = \varepsilon_2/2, \varepsilon \in (0, 1/2]$, and suppose $\int_{\mathbb{R}} \lambda^{2\varepsilon} F(d\lambda) < \infty$. Then we can write the bound:

$$\int_{\mathbb{R}} |b(\lambda)|^2 F(d\lambda) \le \Big(\int_{\mathbb{R}} \lambda^{2\varepsilon} F(d\lambda)\Big) \Big(4^{1-\varepsilon} |x-y|^{2\varepsilon} + |t-s|^{\varepsilon}\Big),$$

which implies (3.18). The estimate (3.16) follows from (3.15).

In view of Theorem 3.4, under assumption A.3 and assuming u_0 to be strictly φ -sub-Gaussian, we can write the estimate for the tail distribution of supremum of $\omega(t, x)$ which is analogous to (3.11), where the constants c_0 and c_{ω} will come now from (3.16), (3.17).

In the example below we present the process which can be used as initial condition, for which (3.17) is satisfied.

Example 3.1. Let the initial condition $u_0(x)$, $x \in \mathbb{R}$, be a Gaussian stationary process with the spectral density

$$f(\lambda) = \frac{\sigma^2}{(1+\lambda^2)^{2\alpha}}, \ \lambda \in \mathbb{R}.$$
(3.19)

The corresponding covariance function is of the form:

$$B_{\eta}(x) = \frac{\sigma^2}{\sqrt{\pi}\Gamma(2\alpha)} \left(\frac{|x|}{2}\right)^{2\alpha - 1/2} K_{2\alpha - 1/2}(|x|), \ x \in \mathbb{R},$$
(3.20)

where K_{ν} is the modified Bessel function of the second kind, in particular, $K_{1/2}(x) = \sqrt{\frac{\pi}{2x}}e^{-x}$. Covariances (3.20) constitute the so-called Matérn class, a parameter $\nu = 2\alpha - 1/2 > 0$ controls the level of smoothness of the stochastic process.

Note that the Gaussian stochastic process with the above covariance and spectral density can be obtained as solution to the following fractional partial differential equation:

$$\left(1 - \frac{d^2}{dx^2}\right)^{\alpha} \eta(x) = w(x), \ x \in \mathbb{R},$$

with w being a white noise: $\mathsf{E}w(x) = 0$ and $\mathsf{E}w(x)w(y) = \sigma^2\delta(x-y)$ (see, e.g. [5, Thm. 3.1]).

The Matérn model is popular in spatial statistics and modeling random fields (with corresponding adjustment of (3.20) for *n*-dimensional case). The relation between the spatial Matérn covariance model and stochastic partial differential equation $(\mu - \Delta)^{\alpha} \eta(x) = w(x), x \in \mathbb{R}^n$, was established by Whittle in 1963 and since then has been widely used in various applied and theoretical contexts.

For the stationary initial condition u_0 with spectral density (3.19) the condition (3.17) holds and we are able to calculate the constant $c(\varepsilon)$ defined in (3.17). We have

$$\int_{\mathbb{R}} \frac{\lambda^{2\varepsilon}}{(1+\lambda^2)^{2\alpha}} = \int_0^\infty \frac{t^{\varepsilon+1/2-1}}{(1+t)^{\varepsilon+1/2+2\alpha-\varepsilon-1/2}} dt = \mathcal{B}(\varepsilon+1/2, 2\alpha-\varepsilon-1/2),$$

where \mathcal{B} is Beta-function, $\varepsilon \in (0, 1/2], 2\alpha - \varepsilon - 1/2 > 0$ and we used the formula $\int_0^\infty \frac{t^{\mu-1}}{(1+t)^{\mu+\nu}} dt = \mathcal{B}(\mu, \nu).$

Therefore, in this case we obtain $c^2(\varepsilon) = \sigma^2 \mathcal{B}(\varepsilon + 1/2, 2\alpha - \varepsilon - 1/2)$. In particular, having in (3.19) $\alpha > 1$ and choosing $\varepsilon = 1/2$ we get $c(1/2) = \sigma^2 \frac{1}{\alpha - 1}$.

References

- [1] Adler R.J., Taylor J.E. Random Fields and Geometry. Springer, New York. 472 p. (2007)
- [2] Balan R., Jolis M., Quer-Sardanyons L. SPDEs with affine multiplicative fractional noise in space with index $\frac{1}{4} < H < \frac{1}{2}$. Electron. J. Probab. 20(54), 1–36 (2015)
- [3] Beghin L., Kozachenko Yu., Orsingher E., Sakhno L. On the Solutions of Linear Odd-Order Heat-Type Equations with Random Initial Conditions. J. Stat. Phys., 127(4), 721–739 (2007).
- [4] Buldygin V.V., Kozachenko Yu.V. Metric characterization of random variables and random processes. Translations of Mathematical Monographs. 188. Providence, RI: AMS, American Mathematical Society. 257 p. (2000)
- [5] D'Ovidio M., Orsingher E., Sakhno L. Spectral densities related to some fractional stochastic differential equations. Electron. Commun. Probab. 21(18), 1–15 (2016)
- [6] Dozzi M., KozachenkoYu., Mishura Yu., Ralchenko K. Asymptotic growth of trajectories of multifractional Brownian motion with statistical applications to drift parameter estimation. Statistical Inference for Stochastic processes, 21, No. 1, 21–52 (2018)
- [7] Giuliano Antonini R., Kozachenko Yu.V., Nikitina T. Spaces of φ-subgaussian random variables. Rendiconti Accademia Nazionale delle Scienze XL. Memorie di Matematica e Applicazioni 121. Vol. XXVII, 95–124 (2003)
- [8] Hong J., Liu Z., Sheng D. Optimal Hölder continuity and hitting probabilities for SPDEs with rough fractional noises. J. Math. Anal. Appl., 512(1), 126125 (2022)
- [9] Hopkalo O., Sakhno L. Investigation of sample paths properties for some classes of φ -sub-Gaussian stochastic processes. Modern Stoch. Theory Appl. 8(1), 41–62 (2021)
- [10] Hopkalo O.M., Sakhno L.M., Vasylyk O.I. Properties of solutions to linear KdV equations with φ -sub-Gaussian initial conditions. Bulletin of the Taras Shevchenko National University of Kyiv. Physics and Mathematics, 2022(2), 11–19 (2022)
- [11] Hopkalo O.M., Sakhno L.M., Vasylyk O.I. Bounds for the Tail Distributions of Suprema of Sub-Gaussian Type Random Fields. Austrian Journal of Statistics, 52(SI), 54–70 (2023)
- [12] Kozachenko Yu.V., Koval'chuk Yu.A. Boundary value problems with random initial conditions and series of functions of $Sub_{\varphi}(\Omega)$. Ukrainian Math. J. 50(4), 572–585 (1998)
- [13] Kozachenko Yu., Olenko A. Whitaker-Kotelnikov-Shanon approximation of φ -sub-Gaussian random processes. J. Math. Analysis Appl. 442(2), 924–946 (2016)
- [14] Kozachenko Yu., Orsingher E., Sakhno L., Vasylyk O. Estimates for functional of solution to higher-order heat-type equation with random initial condition. J. Stat. Phys. 72, 1641–1662 (2018)
- [15] Kozachenko Yu., Orsingher E., Sakhno L., Vasylyk O. Estimates for distribution of suprema of solutions to higher-order partial differential equations with random initial conditions. Modern Stoch. Theory Appl. 7(1), 79–96 (2020)
- [16] Kozachenko Yu.V., Ostrovskij E.I. Banach spaces of random variables of sub-Gaussian type. Theory Probab. Math. Stat. 32, 45–56 (1986)
- [17] Sakhno L. Estimates for distributions of suprema of spherical random fields. Statistics, Optimization & Information Computing, 11(2), 186–195 (2022)
- [18] Sakhno L. Investigation of Airy equations with random initial conditions. Electron. Commun. Probab. 28, article no. 16, 1-14 (2023)
- [19] Sakhno L.M., Vasylyk O.I. Investigation of solutions to higher-order dispersive equations with φ sub-Gaussian initial conditions. Bulletin of Taras Shevchenko National University of Kyiv. Series: Physics & Mathematics, Vol.2(2021), 78–84 (2021)
- [20] Vasylyk O., Kozachenko Yu., Yamnenko R. $\varphi\text{-sub-Gaussian stochastic processes. Kyiv University, Kyiv (2008) (in Ukrainian)$