Variety of mutual-visibility problems in hypercubes

Danilo Korže[†] and Aleksander Vesel^{*}

 [†]Faculty of Electrical Engineering and Computer Science,
 University of Maribor, Koroška cesta 46, SI-2000 Maribor, Slovenia
 *Faculty of Natural Sciences and Mathematics, University of Maribor, Koroška cesta 160, SI-2000 Maribor, Slovenia

May 10, 2024

Abstract

Let G be a graph and $M \subseteq V(G)$. Vertices $x, y \in M$ are M-visible if there exists a shortest x, y-path of G that does not pass through any vertex of $M \setminus \{x, y\}$. We say that M is a mutual-visibility set if each pair of vertices of M is M-visible, while the size of any largest mutual-visibility set of G is the mutual-visibility number of G. If some additional combinations for pairs of vertices x, y are required to be M-visible, we obtain the total (every $x, y \in V(G)$ are M-visible), the outer (every $x \in M$ and every $y \in V(G) \setminus M$ are M-visible), and the dual (every $x, y \in V(G) \setminus M$ are M-visible) mutual-visibility set of G. The cardinalities of the largest of the above defined sets are known as the total, the outer, and the dual mutual-visibility number of G, respectively.

We present results on the variety of mutual-visibility problems in hypercubes.

1 Introduction and preliminaries

In graph theory, a mutual-visibility set refers to a collection of vertices within a graph where for every pair of vertices there exists a shortest path between them that avoids any other vertex in the set. The mutual-visibility number of a graph represents the maximum size of such a mutual-visibility set within the graph.

Mutual-visibility sets have been examined across diverse fields, encompassing wireless sensor networks, mobile robot networks, and distributed computing. In wireless sensor networks, they aid in sensor deployment to ensure interferencefree communication among sensors. In mobile robot networks, they facilitate collision avoidance strategies for robot control. In distributed computing, they contribute to the development of efficient algorithms for tasks such as consensus and broadcasting [3,5,9,11,17]. In 2022, Di Stefano [12] established the foundation of the preceding problem within graph theory, aiming to maximize the size of the largest mutual-visibility set.

Research highlighted in the cited papers explores visibility problems in graphs, indicating specific adjustments to visibility properties of significance. In this context, [8] presents a range of novel mutual-visibility problems, including the total mutual-visibility problem, the dual mutual-visibility problem, and the outer mutual-visibility problem. The total mutual-visibility number on Cartesian products of complete graphs, also known as Hamming graphs, is studied in [18]. The (total) mutual-visibility sets in hypercubes, a subset of Hamming graphs, are investigated in [7] as well as in [4].

In this paper, we build upon previous research of mutual-visibility on hypercubes. Subsequently, we present definitions and results crucial for the remainder of the paper in the following section. Section 2 delves into mathematical optimization techniques capable of yielding novel insights into the mutual-visibility set problem in hypercubes. In Section 3, we investigate the mutual-visibility number and its variations within hypercubes. We establish new upper bounds for these numbers and provide exact values for hypercubes of smaller dimensions. Additionally, we extend the concept introduced in [4], demonstrating that the upper bound on the total mutual-visibility number in hypercubes can be determined by leveraging the size of the largest binary code. This is achieved by showing that the task of finding the largest total mutual-visibility set in an h-cube is equivalent to determining the maximum size of a binary code of length h with a minimum Hamming distance of 4.

Let G = (V(G), E(G)) is a graph, $M \subseteq V(G)$ and $u, v \in V(G)$. We say that a u, v-path P is M-free, if P does not contain a vertex of $M \setminus \{u, v\}$. Vertices $u, v \in V(G)$ are M-visible if G admits an M-free shortest u, v-path.

Let $M \subseteq V(G)$ and $\overline{M} = V(G) \setminus M$. Then we say that M is a

- mutual-visibility set, if every $u, v \in M$ are M-visible,
- total mutual-visibility set, if every $u, v \in V(G)$ are M-visible,
- outer mutual-visibility set, if every $u, v \in M$ are M-visible, and every $u \in M, v \in \overline{M}$ are M-visible,
- dual mutual-visibility set, if every $u, v \in M$ are M-visible, and every $u, v \in \overline{M}$ are M-visible.

The cardinality of a largest mutual-visibility set, a largest total mutualvisibility set, a largest outer mutual-visibility set, and a largest dual mutualvisibility set will be respectively denoted by $\mu(G)$, $\mu_t(G)$, $\mu_o(G)$, and $\mu_d(G)$. Also, these graph invariants will be respectively called the *mutual-visibility num*ber, the total mutual-visibility number, the outer mutual-visibility number, and the dual mutual-visibility number of G.

Let $B = \{0, 1\}$. If b is a word of length h over B, that is, $b = (b_1, \ldots, b_h) \in B^h$, then we will briefly write b as $b_1 \ldots b_h$. Elements of B^h are also called

binary strings of length h. If $x, y \in B^h$, then the Hamming distance H(x, y) between x and y is the number of positions in which x and y differ.

The hypercube of order h or simply h-cube, denoted by Q_h , is the graph G = (V, E), where the vertex set V(G) is the set of all binary strings of length h, while two vertices $x, y \in V(G)$ are adjacent in Q_h if and only if the Hamming distance between x and y is equal to one.

distance between x and y is equal to one. The weight of $u \in B^h$ is $w(u) = \sum_{i=1}^h u_i$, in other words, w(u) is the number of 1s in the word u. For the concatenation of bits the power notation will be used, for instance $0^h = 0 \dots 0 \in B^h$.

If G is a connected graph, then the distance $d_G(u, v)$ (or simply d(u, v)) between vertices u and v is the length of a shortest u, v-path (that is, a shortest path between u and v) in G. The set of vertices lying on all shortest u, v-paths is called the *interval* between u and v and denoted by $I_G(u, v)$ We will also write I(u, v) when G will be clear from the context.

If G is a graph and $X \subseteq V(G)$, then G[X] denotes the subgraph of G induced by X.

If u is a vertex of a graph G, let $N_G(u)$ (or simply $N_G(u)$) denote the set of neighbors of u. Moreover, let $N[u] = N(u) \cup \{u\}$.

For a positive integer n we will use the notation $[n] = \{1, 2, ..., n\}$.

The Cartesian product of graphs G and H is the graph $G \Box H$ with vertex set $V(G) \times V(H)$ and $(x_1, x_2)(y_1, y_2) \in E(G \Box H)$ whenever $x_1y_1 \in E(G)$ and $x_2 = y_2$, or $x_2y_2 \in E(H)$ and $x_1 = y_1$. It is well-known, that the Cartesian product is commutative and associative, having the trivial graph as a unit.

Since for $h \ge 2$ we have $Q_h = Q_{h-1} \Box K_2$, it was observed in [7] that the mutual-visibility number of Q_h is bounded above by twice the mutual-visibility number of Q_{h-1} . Clearly, this observation can be generalized as follows.

Observation 1.1. If $h \ge 2$, then

(i) $\mu(Q_h) \le 2\mu(Q_{h-1}),$ (ii) $\mu_t(Q_h) \le 2\mu_t(Q_{h-1}),$ (iii) $\mu_o(Q_h) \le 2\mu_o(Q_{h-1}),$ (vi) $\mu_d(Q_h) \le 2\mu_d(Q_{h-1}).$

We will need the following well known result, see for example [14].

Proposition 1.2. Let $u, v \in V(Q_h)$.

(i) If $u_i = v_i$ for some $i \in [h]$, then $x_i = u_i = v_i$ for every $x \in I(u, v)$.

(ii) If d(u, v) = d, then the subgraph of Q_h induced by I(u, v) is isomorphic to the d-cube.

The diameter diam(G) of a connected graph G is the maximum distance between two vertices of G. Let G be a graph, $d \leq diam(G)$ and $M \subseteq V(G)$. Vertices $u, v \in V(G)$ are M_d -visible if d(u, v) > d or G admits an M-free shortest u, v-path. It is evident that M is not a mutual-visibility set if M admits a pair of vertices that are not M_d -visible. Furthermore, this concept can be naturally extended to total, outer and dual mutual-visibility sets of G.

2 Mathematical optimization methods

We applied two well-known techniques for computing the mutual-visibility sets of hypercubes: Integer linear programming (ILP) and a reduction to SAT. Both methods have been previously successfully applied for distance constrained coloring problems for various finite and infinite graphs. used for distanceconstrained coloring problems in various finite and infinite graphs. For instance, [19] demonstrates the application of these concepts in the radio coloring problem.

2.1 Integer linear programming (ILP) model

Let G = (V, E) be a graph. For a vertex $v \in V(G)$ we introduce the Boolean variable x_v such that $x_v = 1$ if and only if v belongs to the mutual-visibility set M of G. The problem of finding the maximal mutual-visibility set can be formulated as an integer linear program as follows:

maximize
$$\sum_{v \in V(G)} x_v$$
 (2.1)

subject to:

$$x_u + x_v - \sum_{P \in \mathcal{P}(u,v)} z_{u,v,P} \le 1, \qquad \forall u, v \in V(G);$$
(2.2)

$$z_{u,v,P} + x_z \le 1, \qquad \forall u, v \in V(G); \qquad (2.3)$$
$$\forall P \in \mathcal{P}(u,v); \forall z \in V(P) \setminus \{u,v\}.$$

where $\mathcal{P}(u, v)$ denotes the set of all different shortest paths between vertices u and v, while for $P \in \mathcal{P}(u, v)$ the set $V(P) \setminus \{u, v\}$ comprises all intermediary vertices in the corresponding shortest path P between u and v. Clearly, all shortest paths between all pairs of vertices in G should be computed in order to establish the model for G. The additional variable $z_{u,v,P}$ equals 1 if and only if the shortest path P between u and v enables mutual visibility of these two vertices. That is to say, for each pair of vertices u and v, the number of corresponding additional variables equals the number of different shortest paths between u and v.

Integer linear programming was also employed for searching the dual mutualvisibility and outer mutual-visibility sets of hypercubes.

For the dual visibility sets, the following additional constraints have to be added to constraints (2.2):

$$-x_u - x_v - \sum_{P \in \mathcal{P}(u,v)} z_{u,v,P} \le -1, \ \forall u, v \in V(G);$$

$$(2.4)$$

In order to compute outer visibility sets, constraints (2.2) has to be replaced

with constraints (2.5):

$$x_u + x_v - 2\sum_{P \in \mathcal{P}(u,v)} z_{u,v,P} \le 0, \ \forall u, v \in V(G);$$

$$(2.5)$$

Constraints (2.3) are applied for all three varieties of mutual-visibility set.

To reduce the computation time, we can apply equations only for paths shorter than a predefined length. In this scenario, we can only confirm the non-existence of a visibility set of a given cardinality. However, it has been observed that solutions obtained with path length restrictions often also constitute mutual-visibility sets, provided that the distance length limitation is not too profound.

We applied Gurobi Optimization [13] for solving ILP models.

2.2 Reduction to SAT model

Let G = (V, E) be a graph with n vertices and ℓ a positive integer (the size of a potential mutual-visibility set). For every $v \in V(G)$ we introduce an atom x_v . Intuitively, this atom expresses whether vertex v is inside the mutual-visibility set M or not. More precisely, $x_v = 0$ if and only if v belongs to the mutual-visibility visibility set M.

First collection of propositional formulas define an encoding for cardinality constraints (known as $\geq k(x_1, ..., x_n)$ constraints) which ensure that at most $k = n - \ell$ atoms are assigned value 1. We do not give the details of the applied encoding here; the interested reader is referred to [20], where the implemented encoding based on the so called sequential counter was introduced.

In order to define mutual-visibility constraints, consider the following propositional formulas:

$$x_u \vee x_v \vee \bigvee_{P \in \mathcal{P}(u,v)} \left[\bigwedge_{x_z \in V(P) \setminus \{u,v\}} x_z \right] \; (\forall u, v \in V(G)) \tag{2.6}$$

where $\mathcal{P}(u, v)$ denotes the set of all different shortest paths between vertices u and v, while for $P \in \mathcal{P}(u, v)$ the set $V(P) \setminus \{u, v\}$ comprises all intermediary vertices in the corresponding shortest path P between u and v.

Clearly, all shortest paths between all pairs of vertices in G should be computed in order to establish the clauses for G. Before using the above formulas in a SAT solver, they have to be transformed to the conjunctive normal form. These propositional formulas transform a mutual-visibility problem into a propositional satisfiability test (SAT). We can confirm that a SAT instance is satisfiable if and only if G has a mutual-visibility set of size at least ℓ . Note that for a given *h*-cube only cardinality constrains $\geq k(x_1, ..., x_n)$ depend on ℓ .

In the case of outer and dual visibility set problem, the formulas (2.6) have to be slightly modified. For the outer visibility set, we obtain

$$x_u \wedge x_v \vee \bigvee_{p \in \mathcal{P}(u,v)} \left[\bigwedge_{x_z \in V(P) \setminus \{u,v\}} x_z \right] \ (\forall u, v \in V(G))$$
(2.7)

while for the dual visibility set problem the clauses are of the following form:

$$(x_u \wedge \neg x_v) \vee (\neg x_u \wedge x_v) \vee \bigvee_{P \in \mathcal{P}(u,v)} \left[\bigwedge_{x_z \in V(P) \setminus \{u,v\}} x_z \right] \quad (\forall u, v \in V(G)). \quad (2.8)$$

To keep the model smaller and the computation more efficient, we may include in formulas only the paths of length up to given constant s. As we noted in the above subsection, only the non-existence of a mutual-visibility set of a given cardinality can be confirmed if s < diam(G). However, the tests performed for h-cubes showed that solutions obtained with path length restrictions very often also constitute mutual-visibility sets if s is close to h.

We used the Criptominisat5 SAT-solver [10] to find the solutions of the above presented clauses for hypercubes.

3 Theoretical and computational results

3.1 Mutual-visibility

The mutual-visibility number of hypercubes has been studied in [7]. Obtained exact values and bounds are summarized in the following proposition.

Proposition 3.1.

$$\mu(Q_h) = \begin{cases} 2, & h = 1\\ 3, & h = 2\\ 5, & h = 3\\ 9, & h = 4\\ 16, & h = 5 \end{cases}$$

Moreover, if $h \ge 6$, then $\max_{i \in [h-3]} \left(\binom{h}{i} + \binom{h}{i+3} \right) \le \mu(Q_h) \le 2^{h-1}$.

For a vertex $v \in V(G)$ and $i \ge 0$ let L_i^v (also called an *i*-layer with respect to v) denote the set of vertices of V(G) at distance i from v.

It is shown in [7], that $L_i^v \cup L_{i+3}^v$ is a mutual-visibility set of Q_h for every $v \in V(Q_H)$ and $1 \le i \le h-3$. We will show in the sequel that $L_i^v \cup L_{i+k}^v$ cannot form a mutual-visibility set if $k \le 2$.

Proposition 3.2. Let $h \ge 3$ and $v \in V(Q_h)$. If $1 \le i \le h-2$, then $L_i^v \cup L_{i+1}^v$ is not a mutual-visibility set of Q_h .

Proof. Let $X := L_i^v \cup L_{i+1}^v$. We can set w.l.o.g. that $v = 0^h$. It follows that for every $u \in L_i^v$ and every $z \in L_{i+1}^v$ we have w(u) = i and w(z) = i + 1. Since $|N(z) \cap L_i^v| = i + 1 < h$, there exist a vertex $u \in L_i^v$ such that $uz \notin E(Q_h)$. Suppose that there exists an X-free z, u-path P. Let x denote the vertex adjacent to z in P. Obviously, $x \in L_{i+2}^v$ or $x \in L_i^v$. Since $L_i^v \subset X$, we have $x \in L_{i+2}^v$. Moreover, $N(u) \subset L_{i+1}^v \cup L_{i-1}^v$. But since from Proposition 1.2 it follows that P cannot posses a vertex from L_{i-1}^v , it follows that P admits a vertex from L_{i+1}^v and we obtained a contradiction.

Proposition 3.3. Let $h \ge 3$ and $v \in V(Q_h)$. If $1 \le i \le h-3$, then $L_i^v \cup L_{i+2}^v$ is not a mutual-visibility set of Q_h .

Proof. Let $X := L_i^v \cup L_{i+2}^v$. We can set w.l.o.g. that $v = 0^h$. It follows that for every $u \in L_i^v$ and every $z \in L_{i+2}^v$ we have w(u) = i and w(z) = i + 2. Suppose that $u = 0^{h-i}1^i$ and $z = 1^{i+2}0^{h-i-2}$. Since $h - i \ge 3$ and $i + 2 \ge 3$, it holds that d(u, z) > 2. Suppose that there exists an X-free z, u-path P. Let x denote the vertex adjacent to z in P. Obviously, $x \in L_{i+3}^v$ or $x \in L_{i+1}^v$. If $x \in L_{i+1}^v$, let $y \ne z$ denote the vertex adjacent to x in P. Note that $N(x) \subset L_i^v \cup L_{i+2}^v$. It follows that $y \in X$. Since by d(u, z) > 2 we have $y \ne u$, we obtain a contradiction.

If $x \in L_{i+3}^v$, let w denote the vertex adjacent to u in P. Since $N(u) \subset L_{i+1}^v \cup L_{i-1}^v$, we have $w \in L_{i+1}^v$. But $N(w) \subset L_i^v \cup L_{i+2}^v = X$ and we obtained a contradiction.

Let $u \in V(Q_h)$ and $X \subseteq N(v)$ such that $|X| = d \ge 1$. Note that the vertices of $X \cup \{u\}$ belong to a unique *d*-cube, say Q, a subgraph of Q_h , which is induced by I(u, v), where v is the vertex of Q_h at distance d - 1 from all vertices of X. We will say that the sub-cube Q is raised by $X \cup \{u\}$.

Let M be a mutual-visibility set of Q_h . If $u \in M$, then u and the vertices of M adjacent to u restrict the number of vertices of M in the corresponding sub-cube, as shown in the next proposition.

Proposition 3.4. Let u be a vertex of a mutual-visibility set M of Q_h . If $X = N[u] \cap M$ and Q the sub-cube of Q_h raised by X, then $V(Q) \cap M = X$.

Proof. As noted above, the sub-cube of Q_h raised by X is the subgraph of Q_h induced by I(u, v), where v is the vertex of Q_h at distance d - 1 from all vertices of $X \setminus \{u\}$. Since from Proposition 1.2 then it follows that for every $z \in I(u, v) \setminus M$ a shortest u, z-path contains a vertex from $X \setminus \{u\}$, the proof is complete.

The above proposition states that if M contains a subset $Y \subseteq M$ such that $Q_h[Y]$ is isomorphic to $K_{1,d}$ then the *d*-cube raised by Y cannot have other vertices in M.

The following corollary of the above proposition can be utilized for a computer search of large mutual-visibility sets of Q_h .

Proposition 3.5. Let $h \ge 6$. If Y is a subset of a mutual-visibility set M of Q_h such that $Q_h[Y]$ is isomorphic to $K_{1,4}$, then $|M| \le 2^{h-1} - 2$.

Proof. Let u denote the vertex of degree 4 in $Q_h[Y]$. We can assume without loss of generality that $u = 0^h$, while the other four vertices of Y are 10^{h-1} , 010^{h-2} , 0010^{h-3} and 00010^{h-4} .

The vertices of Y raise the 4-cube, denoted as Q^0 , where Q^0 is induced by the set of vertices of the form $b_1b_2b_3b_40^{h-4}$, $b_i \in \{0,1\}$. Consider now the sequence of cubes $Q^1 \dots Q^{h-4}$ induced by the subsets of $V(Q_h)$ as follows:

 $V(Q^1) = \{b_1 b_2 \dots b_{h-1} 1, b_i \in \{0, 1\}\},\$

 $V(Q^2) = \{b_1 b_2 \dots b_{h-2} 10, b_i \in \{0, 1\}\},\$

 $V(Q^{h-4}) = \{b_1 b_2 b_3 b_4 10^{h-5}, b_i \in \{0, 1\}\}.$

Note that for $i \in [h-4]$ the set $V(Q^i)$ possesses exactly 2^{h-i} vertices, while $V(Q^0)$ possesses exactly 2^4 vertices. Moreover, the vertices of $V(Q^0), \ldots, V(Q^{h-4})$ partition the vertices of Q_h .

By Proposition 3.4, the cube Q^0 contains exactly 5 vertices in M (since $M \cap V(Q^0) = Y$). By Proposition 3.1, for $i \in [h-3]$ we have that $|M \cap V(Q^i)| \le 2^{h-i-1}$, while $|M \cap V(Q^4)| \le 9 = 2^3 + 1$. It follows that $M \le 2^{h-2} + 2^{h-3} + \ldots + 2^3 + 6 = 2^{h-1} - 2$.

Theorem 3.6.

$$\mu(Q_h) = \begin{cases} 2, & h = 1\\ 3, & h = 2\\ 5, & h = 3\\ 9, & h = 4\\ 16, & h = 5\\ 32, & h = 6\\ 59, & h = 7 \end{cases}$$

 $Moreover, \ if \ h \geq 8, \ then \ {\binom{h}{\lfloor \frac{h}{2} \rfloor - 1}} + {\binom{h}{\lfloor \frac{h}{2} \rfloor + 2}} \leq \mu(Q_h) \leq 59 \cdot 2^{h-7}.$

Proof. The results for $h \leq 5$ and the general lower bound are from Proposition 3.1 (note that $i = \lfloor \frac{h}{2} \rfloor - 1$ maximizes $\binom{h}{i} + \binom{h}{i+3}$ for $i \geq 8$). For Q_6 and Q_7 we found a mutual-visibility set with 32 and 59 vertices,

For Q_6 and Q_7 we found a mutual-visibility set with 32 and 59 vertices, respectively, by using an ILP model. To improve the efficiency of computing a mutual-visibility set M, we imposed a constraint on the number of vertices in $M \cap N[u]$ to be 4 for every $u \in M$, as implied by Proposition 3.5. These computations confirmed that the mutual-visibility number of Q_6 is 32. However, for Q_7 , the ILP model did not refute the existence of a larger mutual-visibility set, as the computations did not finish after more than one month.

To verify that a larger mutual-visibility set of Q_7 does not exist, we employed a reduction to SAT. Since the search space in a straightforward approach was too large, we imposed additional restrictions.

First, we searched for the largest mutual-visibility set of Q_7 using an ILP model while prohibiting triples $u, v, z \in M$ such that $Q_7[\{u, v, z\}]$ is isomorphic to $K_{1,2}$. We determined that the size of a mutual-visibility set of Q_7 with this restriction cannot exceed 49.

The above result indicates that every mutual-visibility set M of Q_7 of size greater than 49 must include at least one triple $u, v, z \in M$ such that $Q_7[\{u, v, z\}]$ is isomorphic to $K_{1,2}$. This fact allows us to fix the vertices 0000000, 0000001, 0000010 to be included in M in our SAT model. Additionally, we limited the length of the paths involved in the SAT formulas to 5.

The resulting model comprised 254268 variables and 874806 clauses. After computation lasting 962,000 seconds on a powerful desktop computer with 16 threads, we confirmed that the presented reduction is unsatisfiable. Consequently, we can conclude that a mutual-visibility set of Q_7 with 60 vertices does not exist.

Since the upper bound on $\mu(Q_h)$ for $h \ge 8$ follows from the fact that $\mu_o(Q_7) = 59$ and Observation 1.1, the proof is complete.

3.2 Total mutual-visibility

Let $V_1 \cup V_2$ be the partition of $V(Q_h)$ such that V_1 and V_2 comprise the set of vertices of Q_h with even and odd weights, respectively. Then the *halved cube* Q_h^e (resp. Q_h^o) is the graph with $V(Q_h^e) = V_1$ (resp. $V(Q_h^o) = V_2$), where u and v are adjacent in Q_h^e (resp. Q_h^o) if $d(u, v)_{Q_h} = 2$ (see [15] for more details). Clearly, graphs Q_h^e and Q_h^o are isomorphic. Therefore, we can denote either of them as Q_h' when it is more convenient.

An independent vertex set of a graph G is a subset of the vertices of G such that no two vertices in the subset are adjacent in G. The size of a largest independent set of G is called the *independent number* of G and denoted $\alpha(G)$.

An *independent* vertex set of a graph G is a subset of its vertices where no two vertices are adjacent. The size of the largest independent set of G is referred to as the *independent number* of G, denoted as $\alpha(G)$.

The following result is (in more general form) given in [6].

Theorem 3.7. *M* is a total mutual-visibility set of Q_h if and only if for every $u, v \in M$, it holds that $d(u, v) \neq 2$.

We will demonstrate that determining the largest total mutual-visibility set in an h-cube is tantamount to identifying the largest independent set in the corresponding halved cube.

Proposition 3.8. If $h \ge 1$, then $\mu_t(Q_h) = 2\alpha(Q'_h)$.

Proof. Let I_e (resp. I_o) be an independent set in Q_h^e (resp. Q_h^o). Since Q'_h is connected, for every $u, v \in I_e$ (resp. $u, v \in I_o$) we have $d_{Q_h}(u, v) \ge 4$. Furthermore for every $u \in I_e$ and $v \in I_o$ it holds that $d_{Q_h}(u, v)$ is odd. Hence, by Theorem 3.7 it follows that $I_e \cup I_o$ is a total mutual-visibility set of Q_h . Additionally, since $I_e \cap I_o = \emptyset$, we have $|I_e \cup I_o| = |I_e| + |I_o| \le 2\alpha(Q'_h) \le \mu_t(Q_h)$.

Now, consider M as a total mutual-visibility set of Q_h and let M_e (resp. M_o) denote all vertices of M with (resp. odd) weight. By Theorem 3.7, for every $u, v \in M_e$ (resp. $u, v \in M_o$) we have $d_{Q_h}(u, v) \neq 2$. Moreover, since all weights of vertices in M_e (resp. M_o) are of the same parity, we have $d_{Q_h}(u, v) \geq 4$ for

every $u, v \in M_e$ (resp. $u, v \in M_o$). It is evident that M_e (resp. M_o) forms an independent set in Q_h^e (resp. Q_h^o). Therefore, $|M_e| + |M_o| \le \mu_t(Q_h) \le 2\alpha(Q'_h)$, completing the proof.

If $C \subseteq B^n$ such that for every $x, y \in C$ we have $H(x, y) \geq d$, then we say that C is a binary code of length n and minimum Hamming distance d.

Let A(n,d) denote the maximum size of a binary code of length n and minimum Hamming distance d. It is well known, e.g. [16], that A(n,4) = A(n-1,3) for every $n \ge 6$.

Proposition 3.9. If $h \ge 1$, then $\alpha(Q'_h) = A(h, 4)$.

Proof. Let C denote a binary code of length h - 1 and minimum Hamming distance 3. Let us define the sets C_e and C_o as follows:

$$C_e = \{u0 \mid u \in C \text{ and } w(u) \text{ even}\} \cup \{u1 \mid u \in C \text{ and } w(u) \text{ odd}\}.$$

$$C_o = \{u0 \mid u \in C \text{ and } w(u) \text{ odd}\} \cup \{u1 \mid u \in C \text{ and } w(u) \text{ even}\}.$$

Clearly, $C_e \subseteq V(Q_h^e)$ and $C_o \subseteq V(Q_h^o)$. Moreover, for every $u' \in C_e$, there exists exactly one $u \in C$ such that either u' = u0 or u' = u1. Since for every $u, v \in C$ we have $H(u, v) \geq 3$, for the corresponding vertices $u', v' \in C_e$ it holds $d_{Q_h}(u', v') \geq 3$. Moreover, since d(u', v') is even, we have $d_{Q_h}(u', v') \geq 4$. It follows that C_e is an independent set in Q_h^e . Since the situation for C_o and Q_h^o is analogous, we have $\alpha(Q'_h) \geq A(n, 4)$.

Let I be an independent set of Q'_h . Since for every $u, v \in I$ we have $d_{Q_h}(u, v) \geq 4$, it follows that I is a binary code of length h and minimum Hamming distance 4. Thus, $\alpha(Q'_h) \leq A(n, 4)$. This assertion completes the proof.

From Propositions 3.8 and 3.9 it immediately follows that the problem of finding a largest total mutual-visibility set in a h-cube is equivalent to finding the maximum size of a binary code of length h and minimum Hamming distance 4.

Theorem 3.10. If $h \ge 1$, then $\mu_t(Q_h) = 2 \cdot A(h, 4)$.

Note that the exact values of A(h, 4) are known for every $h \leq 16$, while for bigger dimension only upper and lower bounds have been obtained (for more information on the subject please refer to [1,2]). Total mutual-visibility numbers of Q_h , $2 \leq h \leq 16$ are presented in Table 1.

h $\mathbf{2}$ $\mu_t(Q_h)$

Table 1: Total mutual-visibility numbers of Q_h .

3.3 Outer mutual-visibility

Proposition 3.11. Let $h \ge 2$, $v \in V(Q_h)$ and $i \in [h]$. Then L_i^v is an outer mutual-visibility set of Q_h .

Proof. It is shown in the proof of [12, Theorem 1] that L_i^v is a mutual-visibility set of Q_h . Thus, we have to show that for every $z \notin L_i^v$ and every $u \in L_i^v$ there exists a L_i^v -free shortest u, z-path. Let $z \in L_j^v, j \neq i$. Obviously, $d(u, z) \geq |i-j|$.

Suppose first that j < i. If d(u, z) = i - j, then it is not difficult to find a shortest u, z-path P such that every vertex in P apart of u and z belongs to L_k^v , where j < k < i. It follows that P is L_i^v -free. If d(u, z) > i - j, then let S (resp. S') denote the set of indices from [h], where for every $\ell \in S$ we have $u_\ell \neq z_\ell$ and $z_\ell = 0$ (resp. $z_\ell = 1$). Let $S = \{i_1, \ldots, i_s\}$ and $S' = \{i_{s+1}, \ldots, i_{s+t}\}$. Clearly, d(u, z) = s + t. Let $x^0 = u$, while for $k \ge 1$ let x^k be obtained from u by replacing $u_{i_1}, u_{i_2}, \ldots, u_{i_k}$ by $z_{i_1}, z_{i_2}, \ldots, z_{i_k}$. We can see that $x^{s+t} = z$. We now construct a shortest u, z-path P, where $P = x^0, x^1, \ldots, x^s, x^{s+1}, \ldots, x^{s+t}$. Obviously, for every vertex $y \in V(P) \setminus \{u\}$ we have $y \in L_k^v$, where k < i. If follows that P is L_i^v -free.

The proof for j > i is analogous.

Theorem 3.12.

$$\mu_o(Q_h) = \begin{cases} 2, & h = 1\\ 2, & h = 2\\ 4, & h = 3\\ 6, & h = 4\\ 12, & h = 5\\ 24, & h = 6\\ 40, & h = 7 \end{cases}$$

Moreover, if $h \ge 8$, then $\binom{h}{\lfloor \frac{h}{2} \rfloor} \le \mu(Q_h) \le 40 \cdot 2^{h-7}$.

Proof. For $h \leq 6$, we found outer mutual-visibility numbers of h-cubes by using an ILP model and a reduction to SAT as presented in Section 2.

Moreover, we found an outer mutual-visibility set of Q_7 with 40 vertices. In order to confirm that an outer mutual-visibility set of cardinality 41 does not exist, some additional restrictions where needed.

First, we searched (by using an ILP model) a largest outer mutual-visibility set M of Q_7 such that pairs $u, v \in M$ with $uv \in E(Q_7)$ are forbidden and established that a largest outer mutual-visibility set of Q_7 with this restriction is of size 36.

The above result assures that every outer mutual-visibility set M of Q_7 of size bigger than 36 must contain at least one pair $u, v \in M$ with $uv \in Q_7$. This fact allows us to preset the vertices 0000000 and 0000001 to be included in M in the applied SAT model. Moreover, we restricted the length of the paths involved in the SAT formulas to 4. The obtained model comprises 86889 variables and 230460 clauses. We confirmed (the computation lasted 5562 seconds on a powerful desktop computer with 6 threads) that the above reduction is



Figure 1: An outer mutual-visibility set of Q_4

unsatisfiable. Hence, we may establish that an outer mutual-visibility set of Q_7 with 41 vertices does not exist.

In order to prove the lower bound, remind that L_i^v is an outer mutualvisibility set for every $i \in [h]$ by Proposition 3.11. If $v = 0^h$, then L_i^v contains vertices of Q_h with exactly *i* ones. It follows that $|L_i^v| = {h \choose i}$, which is maximized for $i = |\frac{h}{2}|$.

Since $\mu_o(Q_7) = 40$, the upper bound on $\mu_o(Q_h)$ for $h \ge 8$ follows from Observation 1.1. This assertion completes the proof.

An example of a largest outer mutual-visibility set of Q_4 is presented in Fig. 1.

3.4 Dual mutual-visibility

Given a graph G, it is easy to see that every dual mutual-visibility set of G is also a mutual-visibility set of G.

In this subsection, we first establish a more restricted relation between a mutual-visibility set and a dual mutual-visibility set in hypercubes. More precisely, we present necessary and sufficient conditions that a mutual-visibility set of hypercube must satisfy in order to be a dual mutual-visibility set.

Proposition 3.13. Let M be a mutual-visibility set of Q_h . Then M is a dual mutual-visibility set of Q_h if and only if for every $u, v \in V(Q_h)$ with d(u, v) = 2 it holds that $|I(u, v) \cap M| \neq 2$ or $I(u, v) \cap M = \{w, z\}$ and $wz \in E(Q_h)$.

Proof. If M is a dual mutual-visibility set of Q_h , then M cannot admit vertices u, v at distance 2 such that $I(u, v) \cap M = \{u, v\}$, as in this case, for $I(u, v) \setminus M = \{x, y\}$ we cannot find an M-free shortest x, y-path.

Let for every $u, v \in V(Q_h)$ with d(u, v) = 2 it holds that $|I(u, v) \cap M| \neq 2$ or $I(u, v) \cap M = \{w, z\}$ and $wz \in E(Q_h)$. Since M is a mutual-visibility set of Q_h , for every $u, v \in V(Q_h)$ with d(u, v) = 2 it holds that $|I(u, v) \cap M| \leq 3$. Now, we'll demonstrate that for every $x, y \in Q_h \setminus M$ there exists an M-free shortest x, y-path. If $d(x, y) \leq 2$, the claim is evident. Suppose, for contradiction, that there exist $x, y \in \overline{M}$ with d(x, y) > 2 that are not M-visible. Moreover, let x and y be vertices of \overline{M} that are not M-visible with the minimal distance d(x, y) = i.

By Proposition 1.2 we have that $Q_h[I(x,y)]$ is isomorphic to Q_i . Let L_j^x denote the set of vertices of I(x,y) that are at distance j from x.

We first show that vertices of $N_{Q_h[I(u,v)]}(y) = L_{i-1}^x$ must belong to M. Assuming otherwise, let $z \in L_{i-1}^x \setminus M$. By minimality of i, it follows that z is M-visible, leading to the conclusion that y is also M-visible, contradicting the assumption.

Therefore, we may assume that vertices of L_{i-1}^x belong to M. On other hand, by Proposition 3.2, there exists a vertex $z \in L_{i-2}^x$ that does not belong to M. Since d(u, z) = 2, vertices y and z admit two common neighbours from the set $L_{i-1}^x \subset M$, leading to a contradiction.

1 0

Theorem 3.14.

$$\mu_d(Q_h) = \begin{cases} 2, & h = 1\\ 3, & h = 2\\ 4, & h = 3\\ 8, & h = 4\\ 10, & h = 5\\ 20, & h = 6\\ 29, & h = 7 \end{cases}$$

Moreover, if $h \ge 8$, then $2 \cdot A(h, 4) \le \mu(Q_h) \le 29 \cdot 2^{h-7}$.

Proof. For $h \leq 6$, we determined dual mutual-visibility numbers of h-cubes using an ILP model and a reduction to SAT, as presented in Section 2.

Additionally, we determined a dual mutual-visibility set of Q_7 with 29 vertices. However, the applied models did not rejected the existence of a larger dual mutual-visibility set since the computations did not finish in a reasonable time.

Thus, in order to confirm that a dual mutual-visibility set of cardinality 30 does not exist, some additional restrictions where needed.

First, we search (by using an ILP model) a largest outer mutual-visibility set M of Q_7 such that triples $u, v, z \in M$ with $Q_7[\{u, v, z\}]$ isomorphic to $K_{1,2}$ are forbidden, and we established that a largest dual mutual-visibility set of Q_7 with this restriction has a size of 16.

This result indicates that every dual mutual-visibility set M of Q_7 of size greater than 16 must contain at least one triple $u, v, z \in M$ with $Q_7[\{u, v, z\}]$ isomorphic to $K_{1,2}$. This fact allowed us to preselect the vertices 0000000, 0000001 and 0000010 to be included in M in the applied SAT model. Moreover,



Figure 2: A dual mutual-visibility set of Q_4

we restricted the length of the paths involved in the SAT formulas to 4. The obtained model comprises 94110 variables and 244892 clauses. We confirmed (the computation lasted 5562 seconds on a powerful desktop computer with 6 threads) that the above presented reduction is unsatisfiable. Hence, we may establish that an dual mutual-visibility set of Q_7 with 30 vertices does not exist.

By definition, every total mutual-visibility set is also a dual mutual-visibility set. Hence, the lower bound readily follows from Theorem 3.10. Since the upper bound on $\mu_o(Q_h)$ for $h \ge 8$ follows from the fact that $\mu_d(Q_d) = 29$ and Observation 1.1, the proof is complete.

It is worth to mention that the existence of a dual mutual-visibility set of cardinality k in Q_h does not guarantee that there exists a dual mutual-visibility set of cardinality k - 1 in Q_h . In particular, there is no dual mutual-visibility set of cardinality 7 in Q_4 although the dual mutual-visibility number of Q_4 is 8. An example of a largest dual mutual-visibility set of Q_4 is presented in Fig. 2.

3.5 Summary of results and concluding remarks

This section summarize all known values and bounds on the mutual-visibility, dual mutual-visibility and outer mutual-visibility number of hypercubes. The corresponding values as well as total mutual-visibility numbers for hypercubes of dimensions up to 11 are presented in Table 2. (As already noted, the problem of finding a largest total mutual-visibility set in a h-cube is equivalent to finding the maximum size of a binary code of length h and minimum Hamming distance 4. For that reason, this invariant is presented more in detail in Subsection 3.2.) With respect to the previous subsections, the table presents some additional lower bounds on the invariants studied in this paper. These bounds were improved by using a reduction to SAT. We provide the results of our computations, including the cardinalities of the obtained sets (as well as the time needed and the number of threads used), in the sequel.

| h | $\mu(Q_h)$ | $\mu_t(Q_h)$ | $\mu_d(Q_h)$ | $\mu_o(Q_h)$ |
|----|------------|--------------|--------------|--------------|
| 3 | 5 | 2 | 4 | 4 |
| 4 | 9 | 4 | 8 | 6 |
| 5 | 16 | 4 | 10 | 12 |
| 6 | 32 | 8 | 20 | 24 |
| 7 | 59 | 16 | 29 | 40 |
| 8 | 116-118 | 32 | 52-58 | 72-80 |
| 9 | 222-236 | 40 | 86-116 | 126-160 |
| 10 | 432-472 | 80 | 148-232 | 252-320 |
| 11 | 820-944 | 144 | 210-464 | 462-640 |

 Table 2: Lower bounds, upper bounds and exact values on mutual-visibility

 number varieties

In particular, we found a mutual-visibility set of Q_8 of size 116 (2 seconds, 2 threads), a dual mutual-visibility set of Q_8 of size 52 (27140 seconds, 2 threads), an outer mutual-visibility set of Q_8 of size 72 (3100 seconds, 10 threads), a mutual-visibility set of Q_9 of size 222 (840 seconds, 2 threads), a dual mutual-visibility set of Q_9 of size 86 ($7.7 \cdot 10^5$ seconds, 10 threads), a mutual-visibility set of Q_{10} of size 432 (4500 seconds, 32 threads), a dual mutual-visibility set of Q_{10} of size 148 (610000 seconds, 8 threads), a mutual-visibility set of Q_{11} of size 820 (23400 seconds, 2 threads) and a dual mutual-visibility set of Q_{11} of size 210 (27600 seconds, 16 threads).

In order to narrow down the search space for hypercubes of dimensions 8 or higher, we employed two additional heuristics, leading to the discovery of some large mutual-visibility and dual mutual-visibility sets.

In the first heuristic, we search a mutual-visibility set M (or its variety) by presetting a "substantial" subset of vertices of Q_h to be included in M in the applied SAT model. In this context, we may utilize the fact that every total mutual-visibility set also functions as a (standard) mutual-visibility set, as well as an outer and dual mutual-visibility set. Additionally, every outer (or dual) mutual-visibility set is also a (standard) mutual-visibility set. However, the best results in searching large mutual-visibility sets of Q_h , $h \ge 8$, by this approach were obtained by preselecting vertices of the set $L_{\lfloor \frac{h}{2} \rfloor -1}^{v} \cup L_{\lfloor \frac{h}{2} \rfloor +2}^{v}$, where $v = 0^h$.

The second heuristic leveraged the observation that for $h \leq 6$, every vertex $u = u_1, \ldots, u_h$ in the largest computed dual mutual-visibility set M of Q_h has its corresponding *antipode* in M, denoted as $v = v_1, \ldots, v_h$, where $v_i = 1 - u_i$, $i \in [h]$. Inspired by this, we devised a SAT model wherein the antipode of every

vertex in the form $u = u_1, \ldots, u_{h-1}0$ included in M is always inserted into M. This approach significantly accelerated the search for large dual mutualvisibility sets. Notably, we managed to find dual mutual-visibility sets of Q_{10} and Q_{11} with 148 and 210 vertices, respectively.

Data availability

The datasets generated during and/or analysed during the current study are available on the web page https://omr.fnm.um.si/wp-content/uploads/2024/04/hypercubesMV.pdf and from the corresponding author on reasonable request.

Funding

This work was supported by the Slovenian Research Agency under the grants P1-0297, J1-2452, J1-1693 and J2-1731.

References

- E. Agrell, A. Vardy, and K. Zeger, A table of upper bounds for binary codes, IEEE Trans. Inf. Theory 47 (2001) 3004–3006.
- [2] A. Brouwer, Table of general binary codes. Available: https://www.win.tue.nl/ aeb/codes/binary-1.html
- [3] A. Aljohani, G. Sharma, Complete visibility for mobile robots with lights tolerating faults, Int. J. Netw. Comput. 8 (2018) 32–52.
- [4] M. Axenovich, D. Liu, Visibility in Hypercubes, arXiv:2402.04791
- [5] S. Bhagat, Optimum algorithm for the mutual visibility problem, Lect. Notes Comput. Sci. 12049 (2020) 31–42.
- [6] C. Bujtás, S. Klavžar, and J. Tian, Total mutual-visibility in Hamming graphs, arXiv:2307.05168 (2023)
- [7] S. Cicerone, A. Di Fonso1, G. Di Stefano, A. Navarra, F. Piselli, Mutual visibility in hypercube-like graphs, arXiv:2308.14443v1
- [8] S. Cicerone, G. Di Stefano, L. Droždek, J. Hedžet, S. Klavžar, I. G. Yero, Variety of mutual-visibility problems in graphs, Theor. Comput. Sci., DOI: 10.1016/j.tcs.2023.114096.
- [9] S. Cicerone, A. Di Fonso, G. Di Stefano, A Navarra, The geodesic mutual visibility problem for oblivious robots: The case of trees, ICDCN '23: Proceedings of the 24th International Conference on Distributed Computing and Networking, 2023, 150–159.
- [10] https://www.msoos.org/cryptominisat5/.

- [11] G.A. Di Luna, P. Flocchini, S.G. Chaudhuri, F. Poloni, N. Santoro, G. Viglietta, Mutual visibility by luminous robots without collisions, Inf. Comput. 254 (2017) 392–418.
- [12] G. Di Stefano, Mutual visibility in graphs, Appl. Math. Comput. 419 (2022) 126850.
- [13] https://www.gurobi.com/
- [14] R. Hammack, W. Imrich, S. Klavžar, Handbook of Product Graphs, Second ed., CRC Press, Boca Raton, 2011.
- [15] W. Imrich, S. Klavźar, A. Vesel, Recognizing halved cubes in a constant time per edge, Eur. J. Comb. 16 (1995) 617-621.
- [16] F. J. Macwilliams and N. J. A Sloane, The theory of error-correcting codes, North-Holland, 2000.
- [17] P. Poudel, A. Aljohani, G. Sharma, Fault-tolerant complete visibility for asynchronous robots with lights under one-axis agreement, Theor. Comput. Sci. 850 (2021) 116–134.
- [18] J. Tian, S. Klavžar, Graphs with total mutual-visibility number zero and total mutual-visibility in Cartesian products, http://arxiv.org/abs/2212.07193v1.
- [19] D. Korže, Z. Shao, A. Vesel, New results on radio k-labelings of distance graphs, Discrete Appl. Math 319 (2022) 472–479.
- [20] C. Sinz, Towards an Optimal CNF Encoding of Boolean Cardinality Constraints, In: van Beek, P. (eds) Principles and Practice of Constraint Programming - CP 2005. CP 2005. Lect. Notes Comput. Sci., 3709 (2015) 827– 831, Springer, Berlin, Heidelberg, https://doi.org/10.1007/11564751.73.