

# Multilevel Regression and Poststratification Interface: Application to Track Community-level COVID-19 Viral Transmission\*

Yajuan Si,<sup>†</sup> Toan Tran,<sup>†</sup> Jonah Gabry,<sup>‡</sup> Mitzi Morris,<sup>‡</sup> and Andrew Gelman<sup>‡</sup>

## Abstract

In the absence of comprehensive or random testing throughout the COVID-19 pandemic, we have developed a proxy method for synthetic random sampling to estimate the actual viral incidence in the community, based on viral RNA testing of asymptomatic patients who present for elective procedures within a hospital system. The approach collects routine testing data on SARS-CoV-2 exposure among outpatients and performs statistical adjustments of sample representation using multilevel regression and poststratification (MRP). MRP adjusts for selection bias and yields stable small area estimates. We have developed an open-source, user-friendly MRP interface for public implementation of the statistical workflow. We illustrate the MRP interface with an application to track community-level COVID-19 viral transmission in the state of Michigan.

**Keywords:** Selection bias, Small area estimation, Surveillance system, Bayesian workflow

## 1. Introduction

Early and accurate knowledge of incidence and trends of SARS-CoV-2 transmission in communities is crucial to monitoring the pandemic and supporting policymakers in assessing the effects of restrictive measures on individual and community behaviors. However, in the absence of universal screening or random testing, government policy and healthcare implementation responses relied on the testing of COVID symptomatic or presumed-exposed patients and were guided by officially reported positivity rates and counts of positive cases in the community. These data are biased in concept and flawed in practice. It is essential to set up a data collection and analysis procedure in an operational surveillance system that allows prompt assessments of mitigation efforts and future predictions of clinical burdens to trigger an effective healthcare response and also informs other epidemics. We need to be able to abstract from universal random testing the key characteristics that would make it useful and find ways to realize many of such advantages in a proxy measure. Such a metric would need to respond to increases in viral incidence before they become clinically relevant and identify decreases in incidence that can enable safe suspension of mitigation strategies

---

\*The interface is available here <https://github.com/mrp-interface/shinymp> for local installation and here <https://mrpinterface.shinyapps.io/shinymp/> for web-based implementation.

<sup>†</sup>University of Michigan, Ann Arbor.

<sup>‡</sup>Columbia University, New York

to enhance economic and social recovery. Ideally, the method would be practically applicable at the community level or generalizable nationwide and also able to focus on burdens within specific demographics. It needs to be reliable, statistically verifiable, inexpensive, and automatic, being ready with prompt use for future pandemics.

In this paper, we present a user-friendly surveillance interface with an automatic implementation of multilevel regression and poststratification (MRP, Gelman and Little (1997)) to track community-level COVID-19 viral transmission. Previous work has developed the foundation of such a proxy metric for data collection and statistical adjustment of demographic representation (Covello et al., 2021; Si et al., 2022). The approach collected electronic health records (EHRs) on routine viral testing of patients who present for elective procedures within a hospital system and performed MRP to estimate actual viral trends in a diverse urban-suburban-rural setting in Indiana. The findings show that this model predicted the clinical burden of SARS-CoV-2 earlier and more accurately than currently accepted metrics. In contrast, the reliance on current data was neither nimble to predict increases early enough nor to display decreases until they have passed. We extend the previous work by tracking community-level transmission—across geographic areas and demographic subgroups—to monitor the epidemic over time as an operational surveillance system.

The approach collects EHRs of prospective surgical (and other invasive procedure) patients who are COVID asymptomatic and have tested for acute SARS-CoV-2 infection before performing the procedure (Covello et al., 2021). All patients used the same test administered within a health system by similar healthcare professionals. The group has a broad age, racial/ethnic, and economic diversity, and its only overt correlation to disease status is that it is specifically selected for surgical appointments with a lack of symptoms and a negative exposure history. We assume that, for any uniform demographic, the ratio of asymptomatic-to-symptomatic viral infection is constant. The asymptomatic population in a community would then vary in a strict ratio with overall prevalence and serve as a proxy for true viral incidence. To the degree to which health utilizations or other factors might contribute to variation in this ratio across the sample, we anticipate that much of this variation would be captured by our adjustments. MRP adjusts the demographic (sex, age, and race) and geographic (five-digit zip code) distributions to the target population. The target population is defined as U.S. residents dwelling in the catchment areas of the collected zip codes. The interface links the input patient EHR data with zip codes to the American Community Survey (ACS), the largest household survey of U.S. population (U.S. Bureau of the Census, 2024), and uses the ACS aggregated summaries of sociodemographic and socioeconomic characteristics as geographic predictors at the zip level.

As a prediction approach to modeling the test outcomes with individual-level and geographic predictors, MRP has become increasingly popular for subgroup estimation. Originally applied to estimate state-level public opinions from sociodemographic subgroups using sample surveys, MRP has two key components: (1) multilevel regression for small area estimation by setting up a predictive model with a large number of covariates and regularizing with Bayesian prior specifications, and (2) poststratification to adjust for selection bias and correct for imbalances in the sample composition. Flexible modeling of survey outcomes

can capture complex data structures conditional on poststratification cells, which are determined by the cross-tabulation of categorical auxiliary variables and calibrate the sample discrepancy with population control information. Besides applications in social sciences, especially in election forecasting (for example, Wang et al. (2015); Lauderdale et al. (2020)), MRP has also shown promise in public health research (for example, Zhang et al. (2015); Downes et al. (2018); Downes and Carlin (2020); Si et al. (2020)). Si (2024) show that the key to the success of MRP in applications is the inclusion of highly predictive covariates, and Li and Si (2024) discuss estimation approaches when the population distribution of the poststratification variables is unknown.

The interface extends MRP with two novel aspects as methodological contributions: time-varying and granular geographic level estimation. Currently most application studies with MRP are based on cross-sectional data, which can be collected via a probability sample survey (e.g., Zhang et al. (2015); Si et al. (2020)) or a nonprobability sample (e.g., Wang et al. (2015); Si (2024)). We use cross-sectional data to refer to the dataset with measures collected at a specific time point that does not account for temporal variation in the modeling. We extend MRP to time-varying data and track prevalence by week in the interface. Because of sample size limitations, existing MRP estimates are often at the state level. The interface analyzes data from each zip code, includes zip-level predictors, and aggregates to yield county-level estimates. We use spatiotemporal data to refer to the data collected over time by granular geography. The interface can apply to both cross-sectional and spatiotemporal datasets. In this paper, we focus on the application of spatialtemporal data analysis and track community-level COVID-19 viral transmission.

We demonstrate the interface workflow via demographic and county-level COVID-19 viral tracking in Michigan. We describe the MRP methods in Section 2. Section 3 presents the workflow of statistical analyses, from data preprocessing, descriptive summaries, model fitting and diagnostics, to result presentation. Section 4 concludes with discussions and future extensions.

## 2. Methods

The EHR data include patients' information on the polymerase chain reaction (PCR) test results, result date, sex, age, race, and five-digit zip codes. The PCR tests are presumed to have around 70% clinical sensitivity and essentially 100% specificity. Previous work treated the sensitivity and specificity as unknown parameters, borrowed information from prior studies, and accounted for the estimation uncertainty in the final MRP estimates (Gelman and Carpenter, 2020; Covello et al., 2021; Si et al., 2022). In the interface, we fix the sensitivity at 70% and the specificity at 100% to accommodate the computational implementation with the R package `brms` (Bürkner, 2020).

We track viral infections by week. We first introduce the conventional MRP framework without time-varying factors and then describe its extensions for viral tracking over time.

## 2.1. MRP for cross-sectional data

First MRP fits a multilevel regression model to predict the outcome measure based on a set of factors, then poststratifies the categorical factors to match their distributions with those in the target population. We use a binary outcome of interest as an example. Let  $y_i (= 0/1)$  be the binary indicator of the positive response  $y_i = 1$  for individual  $i$ . We consider a logistic regression and include varying intercepts for age, race/ethnicity, and zip code, where the variation for the zip-varying intercepts is explained by the zip-level predictors.

$$\Pr(y_i = 1) = \text{logit}^{-1}(\beta_1 + \beta_2 \text{male}_i + \alpha_{a[i]}^{\text{age}} + \alpha_{r[i]}^{\text{race}} + \alpha_{s[i]}^{\text{zip}}), \quad (2.1.1)$$

where  $\text{male}_i$  is an indicator for men,  $\alpha_a^{\text{age}}$  is the effect of subject  $i$ 's age on the log-odds function of the probability of having a positive response,  $\alpha_r^{\text{eth}}$  is the coefficient for subject  $i$ 's race/ethnicity, and  $\alpha_s^{\text{zip}}$  is the coefficient for the zip code of residence. As we have zip-level predictors  $\vec{Z}_s^{\text{zip}}$ , we need to build another model in which  $\alpha_s^{\text{zip}}$  is the outcome of a linear regression with zip-level predictors. For zip  $s$ ,

$$\alpha_s^{\text{zip}} = \vec{\alpha} \vec{Z}_s^{\text{zip}} + e_s,$$

where  $e_s$  is a random error.

In the Bayesian framework, we assign hierarchical priors to varying intercepts  $\alpha^{\text{name}}$  or error terms  $e_s$ :

$$\begin{aligned} \alpha^{\text{name}} &\sim \text{normal}(0, \sigma^{\text{name}}) \\ \sigma^{\text{name}} &\sim \text{normal}_+(0, 2.5), \end{aligned} \quad (2.1.2)$$

for  $\text{name} \in \{\text{age}, \text{race}\}$ . Here,  $\text{normal}_+(0, 2.5)$  represents a half-normal distribution with the mean 0 and standard deviation 2.5 restricted to positive values. For the zip-level error,  $e_s \sim \text{normal}(0, \sigma^{\text{zip}})$ , and  $\sigma^{\text{zip}} \sim \text{normal}_+(0, 2.5)$ .

Since (2.1.1) assumes the individuals in the same poststratification cell have the same probability of obtaining a positive response, we can replace the microdata modeling with a Binomial model of cell-wise aggregates. We can model the sum of the responses in cell  $j$  as  $y_j^* \sim \text{Binomial}(n_j, \theta_j)$ , where  $n_j$  is the sample cell size and  $\theta_j = \text{logit}^{-1}(\beta_1 + \beta_2 \text{male}_j + \alpha_{a[j]}^{\text{age}} + \alpha_{r[j]}^{\text{race}} + \alpha_{s[j]}^{\text{zip}})$  using the cell-wise effects of all factors.

To generate overall population or subgroup estimates, we obtain the poststratification cells—in the contingency table of sex, age, race/ethnicity, and zip—and weight the model predictions by the population cell frequency  $N_j$ 's. Suppose the cell-wise estimate based on model (2.1.1) is  $\theta_j$  in cell  $j$ , the MRP estimate can be expressed as:

$$\theta^{\text{MRP}} = \frac{\sum_j N_j \theta_j}{\sum_j N_j}.$$

The MRP estimator for zip  $s$  is

$$\theta_s^{MRP} = \frac{\sum_{j \in s} N_j \theta_j}{\sum_{j \in s} N_j}.$$

We implement Bayesian inference for the estimates, where the variance is based on the posterior samples.

## 2.2. MRP for spatiotemporal data

We account for the PCR testing sensitivity and specificity and apply MRP to test results for population representation. For spatiotemporal data, MRP has two key steps: (1) fit a multilevel model to the testing data for the prevalence with time and the adjustment variables, and (2) poststratify using the population distribution of the adjustment variables: the biological variable of sex, age, race, and zip codes, where we assume the population distribution is the same during the study period. Hence, the poststratification cell is defined by the cross-tabulation of sex, age, race, zip code, and indicators of time in weeks based on the test result dates.

We denote the PCR test result for individual  $i$  as  $y_i$ , where  $y_i = 1$  indicates a positive result and  $y_i = 0$  indicates a negative. Similarly, with poststratification cells, we assume that individuals in the same cell have the same infection probability and can directly model cell-wise summaries. We obtain aggregated counts as the number of tests  $n_j$  and the number of positive cases  $y_j^*$  in cell  $j$ . Let  $p_j = \Pr(y_{j[i]} = 1)$  be the probability that person  $i$  in cell  $j$  tests positive. The analytic incidence  $p_j$  is a function of the test sensitivity  $\delta$ , specificity  $\gamma$ , and the true incidence  $\pi_j$  for individuals in cell  $j$ :

$$p_j = (1 - \gamma)(1 - \pi_j) + \delta\pi_j.$$

We fit a binomial model for  $y_j^*, y_j^* \sim \text{binomial}(n_j, p_j)$  with a logistic function for  $\pi_j$  with covariates—sex, age, race, zip codes, and time in weeks—to allow time-varying prevalence in the multilevel model.

$$\text{logit}(\pi_j) = \beta_1 + \beta_2 \text{male}_j + \alpha_{a[j]}^{\text{age}} + \alpha_{r[j]}^{\text{race}} + \alpha_{s[j]}^{\text{zip}} + \alpha_{t[j]}^{\text{time}}, \quad (2.2.1)$$

where  $\text{male}_j$  is an indicator for men;  $a[j]$ ,  $r[j]$ , and  $s[j]$  represent age, race, and zip levels; and  $t[j]$  denotes the time in weeks when the test result is collected for cell  $j$ . We include zip-level predictors  $\vec{Z}_s^{\text{zip}}$  for zip code  $s$ ,

$$\alpha_s^{\text{zip}} = \vec{\alpha} \vec{Z}_s^{\text{zip}} + e_s.$$

Here the zip-level error  $e_s$  can follow a normal distribution or a spatial model such as conditional autoregression to capture the geospatial dependency.

We assign the same hierarchical priors in (2.1.2) to varying intercepts  $\alpha^{\text{name}}$  or error terms  $e_s$ . Using the estimated incidence  $\hat{\pi}_j$  based on the Bayesian model in (2.2.1), we adjust for selection bias by applying the sociodemographic distributions in the community

to generate the population-level prevalence estimates, as the poststratification step in MRP. We calculate the weekly prevalence estimate in the population,

$$\hat{\pi}_t = \sum_{j \in \text{week } t} N_j \hat{\pi}_j / \sum_{j \in \text{week } t} N_j,$$

which can be restricted to subdomains of interest, as another property of MRP is to yield robust estimates for small areas.

### 3. Bayesian Workflow with MRP

The interface implements a complete Bayesian MRP workflow of statistical analyses, from data preprocessing, descriptive summaries, model fitting and diagnostics, to result presentation.

#### 3.1. Data preprocessing

The input data can be individual patient test records or aggregated summaries across post-stratification cells. As the initial step, the interface reads and displays the input data. The data cleaning and linking process is illustrated by the flowchart in Figure 3.1. The procedure automatically imputes missing values in the predictors, converts categorical variables as factors, and standardizes continuous predictors on the zip level. The MRP integrates three data sources: (1) PCR test results, (2) poststratification cell counts obtained from ACS, and (3) zip-level predictors linked from ACS.




Figure 3.1: Data preprocessing flowchart.

The interface analyzes COVID-19 PCR test records from hospitals. The program identifies the relevant measures in the data frame used for the modeling step. Specifically, the

input data frame must contain columns corresponding to the following demographic, geographic, and temporal measures: sex, race, age, zip code (five digits), PCR test result, and result date.

We link the patient data to the 2020 five-year ACS via residence zip codes. The linking has two goals: (1) identify the target population as individuals who live in the collected zip codes' catchment areas and obtain the poststratification cells' population counts, and (2) include area-level predictors of viral infection to account for geographic variation. The geography of the ACS data covers census tracts, counties, and states, while zip codes are defined by the U.S. Postal Service (USPS). We use the USPS crosswalk table to link zip codes and tracts and compute the zip code-level measures by aggregating all the tracts covered by each zip code.

To yield population representation, we construct poststratification cells based on cross-tabulating sex, race, age, and zip code. The population cell counts are obtained from the weighted distributions of ACS individual records in the catchment areas. We assume they are constant during the studied period (2020-2022).

The geographic measures of interest are defined for individuals (education, employment, and income), or at the tract level (for example, urbanicity and the Area Deprivation Index; ADI, Kind and Buckingham (2018)). We aggregate these to the zip level as the following geographic-area level measures: (1) the percentage of covered census tracts classified as urban, weighted by tract population counts; (2) the percentage of the residing population who have earned an Associate's degree or higher; (3) the percentage of the residing population whose ratio of income to poverty level in the past 12 months is below 100%; (4) the percentage of the civilian labor force who are employed; (5) the average of tract-level medians of household income values of the residing population in the past 12 months, weighted by tract population counts; and (6) the average ADI across covered census tracts, weighted by tract population counts.

### 3.2. Descriptive statistics

We examine descriptive statistics of the observed positivity across time and counties, demographics based on the individual records, and characteristics of the covered geographic areas. The observed viral infection shows variation across time, geography, and demographic groups.

Figure 3.2 presents the highest value among weekly positivities and collected sample sizes across counties, exemplifying the large geographic variation. Most individuals are from the four counties in Southeast Michigan, where the medical center is located. However, the sample catchment area covers 94 counties. The test positivity among asymptomatic patients is often lower than 1%, but some counties observe higher than 80% positivity, showing unstable results because of small numbers of tests. The geographically adjacent areas may not share similar peak values.

Figure 3.3 compares the sex, race, and age distributions between the hospital patients ( $n = 128,222$ ) and the population residing in the catchment areas ( $N = 18,794,558$ ). The




Figure 3.2: Highest values among weekly positivities and available sample sizes across 94 counties in the catchment area.

hospital patients have larger proportions of females, White, and youths than the population, and this sample discrepancy will be adjusted by the poststratification step in MRP.

Figure 3.4 presents the distributions of geographic characteristics. The catchment area of the hospital patients' residence covers 848 zip codes and has a broad and diverse representation in terms of urban/rural areas, area deprivation status, higher education attainment, employment rate, income, and poverty. We expect that socioeconomic measures at the zip level would affect individual behaviors and health and be related to viral transmission. The geographic characteristics would explain the spatial variation. The poststratification uses the population counts by zip code but does not adjust the geographic characteristics.

### 3.3. Model fitting

The interface allows users to specify and fit different models with various choices of individual/geographic covariates and fixed/varying effects. The model fitting is via Bayesian computation with Markov chain Monte Carlo algorithms in Stan (Stan Development Team, 2024). Currently the interface uses the R package *brms* (Bürkner, 2020) to facilitate model specification and estimation. Users can specify prior distributions or choose the default prior setups that are weakly informative.

The *brms* package allows users to specify customized likelihood functions. To account for the PCR test sensitivity and specificity (*sens*, *spec*), we modify the binomial model likelihood (numbers of tests and cases) by connecting the test positivity to the true prevalence, which is reflected in the formula for the outcome: `cases | vint(tests) + vreal(sens, spec)`.

The model outputs include summaries and convergence assessments of the posterior sam-






Figure 3.3: Comparisons of demographic distributions between the input data of hospital patients and the linked American Community Survey data in the catchment area.

Table 3.1: Model comparisons with the leave-one-out cross-validation.

	expected log predictive density differences (elpd_diff)	standard errors (se_diff)
Model A	0.0	0.0
Model B	-1.5	1.7
Model C	-5.1	3.5

ple of model parameters. We give an example for model (2.2.1) in Section A of the Appendix.

### 3.4. Model diagnostics

As an important feature, the interface compares different models and presents model diagnostics results. We employ the leave-one-out cross-validation (LOO-CV) implemented in the R package *loo* (Vehtari et al., 2024) and posterior predictive check (PPC; Gelman et al. (1996)). The LOO-CV assesses the posterior predictive performances of Bayesian models and compares different models on expected log predictive density (elpd) for new data.

We have compared the following three models with different mean structure and variance specifications. Model A includes the fixed effects of sex and geographic predictors and varying effects of age, race, time in weeks, and zip code. Model B adds high-order interactions, between race and college attainment status, to Model A. Model C removes the varying effects of zip code areas from Model A.

Model A: 1 + ADI + college + employment + income + poverty + urbanicity + sex + (1 | age) + (1 | race) + (1 | time) + (1 | zip)

Model B: 1 + ADI + college + employment + income + poverty + urbanicity + sex + race:college + (1 | age) + (1 | race) + (1 | time) + (1 | zip)

Model C: 1 + ADI + college + employment + income + poverty + urbanicity + sex + (1 | age) + (1 | race) + (1 | time)




Figure 3.4: Distributions of geographic characteristics based on the linked American Community Survey data in the catchment area.

Table 3.1 gives the LOO-CV outputs on the model comparison. The difference,  $elpd\_diff$ , will be positive if the expected predictive accuracy for Model B or Model C is higher than that for Model A. The negative  $elpd\_diff$  values show that Model A has the best predictive performance. The  $se\_diff$  values support whether the improvement of Model A is substantial. A rule of thumb is to check whether the interval  $(elpd\_diff - 2se\_diff, elpd\_diff + 2se\_diff)$  covers the value 0. Hence, we select Model A for inference.

If the model fits the observed data well and preserves the correlation structure, we expect the model to generate replicated data of the observations that mimic the raw values. The PPC in Figure 3.5 examines the weekly positivity and compares the raw values to the replicates based on 200 posterior predictive samples from Model A. Across time, the replicates are close to the observations, showing that the model fits the data well without red flags of




Figure 3.5: Posterior predictive check comparing the replicated positivity generated from the estimated model to the raw positivity by week, where shaded areas indicate the 95% posterior predictive intervals.

failing to capture important structures.

### 3.5. Estimation results

Based on the selected Model A, we present the estimated infection prevalence over time for the target population and demographic and geographic subpopulations.




Figure 3.6: Estimated weekly prevalence in the community. The shaded areas represent one standard deviation of uncertainty.

Figure 3.6 shows the estimated viral transmission rate by week in the catchment areas. We observe spikes in March 2020, November 2020, January 2022, and August 2022. The

MRP estimates are generally higher than the raw positivity, mainly because of the test sensitivity, where 70% of infections are tested positive. MRP matches the sample demographics with those in the population.




Figure 3.7: Estimated weekly prevalence across racial groups. The shaded areas represent one standard deviation of uncertainty.

MRP stabilizes small group estimates and adjusts for the sample discrepancy within each group. Figure 3.7 presents the estimated prevalence for White, Black, and other race categories. Whites tend to have lower infection rates than Black and other racial groups, even though most weekly differences are small and not significant. The model does not include time trends varying across racial groups, i.e., without racial moderation effects. The estimated trends are close to paralleling with similar spike and flat periods. The same observation of trends applies to the sex and age group estimates, given in Figures B.1 and B.2 of the Appendix.

We select one week that observes a spike of infection, 01/31/2022-02/06/2022, and present the county-level estimated prevalence in Figure 3.8. The collected test records across counties




Figure 3.8: Observed and estimated county-level prevalence during the week of 01/31/2022-02/06/2022 in the catchment area.

are sparse, where 33 out of 94 counties do not have any tests and 27 counties have only one test during the selected week. The county with the largest number of tests (223) is where the health system is located. The observed positivity values are unreliable, and 51 out of 61 collected values are zero. The MRP estimates are available for all 94 counties based on the predictions with the ACS data. The multilevel model smooths county-level prevalence estimates with a range of 2.8%–7.3%, with a median value of 5%. The standard error (SE) values of the 94 county-wise prevalence estimates are between 0.007 and 0.035, and the variation generally increases with the estimated prevalence.

## 4. Discussion

With generalizability as the goal, the MRP method adjusts for selection bias and stabilizes small group estimates. The interface allows applying MRP to cross-sectional data without time-varying measures and spatiotemporal data to track estimates over time by geography. The interface supports statistical computation and visualization, based on the statistical language R and the shiny application. Users can download and install the interface on local machines and conduct the computation. Alternatively, users can upload data to the web-based interface and apply cloud-based computation resources. The web-based implementation is under development for data privacy protection and security concerns. The interface analyzes aggregated data by poststratification cells, which also facilitates disclosure risk control. The open-source interface will be easily accessed and regularly maintained,

and will continue to evolve and develop over time.

The interface tracks the epidemic and delivers substantive findings of health and sociodemographic disparities in time. We present the results analyzing data from the Michigan health system, which covers a broad distribution of sociodemographics. Our previous work used data from a community hospital in Indiana (Covello et al., 2021), and the results show differences between the states. As shown in our county-level estimates in Michigan, geographic variation can be substantial. To further enhance national generalizability, we need to analyze national data from more states. The web-based interface development will allow users with general backgrounds to apply our method to the local data.

With respect to our current sampling method for COVID-19 viral tracking, in accordance with accepted American Society of Anesthesiology standards, all preoperative patients in the hospital system are subjected to surgical risk evaluation. Hence, routine hospital testing is already implemented uniformly across the U.S., increasing the operational feasibility of our proposed surveillance system. Nonetheless, our further research goals are not tied to the above sampling method. MRP normalization of sample demographics anticipates applicability to many different sampling methods.

The MRP interface can analyze data beyond COVID-19 viral tracking and be applied to broad health and social science studies. Illustrated with the cross-sectional and spatiotemporal data examples, the interface adjusts for bias on known variables using poststratification to the target population distribution and can analyze probability survey data, nonprobability samples, and a combination of multiple data sources. Further work will implement advanced modeling and prior specifications, complex spatiotemporal dependency structure, and unknown population distributions of the poststratification variables.

## Acknowledgements

This work is supported by the National Institutes of Health through grant U01MD017867.

## References

- Bürkner, P.-C. (2020). brms: Bayesian regression models using Stan. <https://cran.r-project.org/package=brms>.
- Covello, L., A. Gelman, Y. Si, and S. Wang (2021). Routine hospital-based SARS-CoV-2 testing outperforms state-based data in predicting clinical burden. *Epidemiology* 32(6), 792–799.
- Downes, M. and J. Carlin (2020). Multilevel regression and poststratification versus survey sample weighting for estimating population quantities in large population health studies. *American Journal of Epidemiology* 189(7), 717–725.

- Downes, M., L. C. Gurrin, D. R. English, J. Pirkis, D. Currier, M. J. Spittal, and J. B. Carlin (2018). Multilevel regression and poststratification: A modeling approach to estimating population quantities from highly selected survey samples. *American Journal of Epidemiology* 187(8), 1780–1790.
- Gelman, A. and B. Carpenter (2020). Bayesian analysis of tests with unknown specificity and sensitivity. *Applied Statistics* 69(5), 1269–1283.
- Gelman, A. and T. C. Little (1997). Poststratification into many categories using hierarchical logistic regression. *Survey Methodology* 23, 127–135.
- Gelman, A., X. L. Meng, and H. Stern (1996). Posterior predictive assessment of model fitness via realized discrepancies. *Statistica Sinica* 6, 733–807.
- Kind, A. J. and W. R. Buckingham (2018). Making neighborhood-disadvantage metrics accessible—the neighborhood atlas. *New England Journal of Medicine* 378(26), 2456.
- Lauderdale, B. E., D. Bailey, J. Blumenau, and D. Rivers (2020). Model-based pre-election polling for national and sub-national outcomes in the US and UK. *International Journal of Forecasting* 36, 399–413.
- Li, K. and Y. Si (2024). Embedded multilevel regression and poststratification: Model-based inference with incomplete poststratifier information. *Statistics in Medicine* 43(2), 256–278.
- Si, Y. (2024). On the use of auxiliary variables in multilevel regression and poststratification. *Statistical Science*.
- Si, Y., L. Covello, S. Wang, T. Covello, and A. Gelman (2022). Beyond vaccination rates: A synthetic random proxy metric of total SARS-CoV-2 immunity seroprevalence in the community. *Epidemiology* 33(4), 457–464.
- Si, Y., R. Trangucci, J. S. Gabry, and A. Gelman (2020). Bayesian hierarchical weighting adjustment and survey inference. *Survey Methodology* 46(2), 181–214.
- Stan Development Team (2024). Stan: A C++ library for probability and sampling. <http://mc-stan.org>.
- U.S. Bureau of the Census (2024). The American Community Survey. <https://www.census.gov/programs-surveys/acs>.
- Vehtari, A., J. Gabry, M. Magnusson, Y. Yao, P.-C. Bürkner, T. Paananen, and A. Gelman (2024). loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models. R package version 2.7.0, <https://mc-stan.org/loo/>.
- Wang, W., D. Rothschild, S. Goel, and A. Gelman (2015). Forecasting elections with non-representative polls. *International Journal of Forecasting* 31(3), 980–991.

Zhang, X., J. B. Holt, S. Yun, H. Lu, K. J. Greenlund, and J. B. Croft (2015). Validation of multilevel regression and poststratification methodology for small area estimation of health indicators from the behavioral risk factor surveillance system. *American Journal of Epidemiology* 182(2), 127–137.

## A. Model fitting results

```
Family: binomial_sens_spec
Links: mu = logit
Formula: cases | vint(tests) + vreal(sens, spec) ~ 1 + ADI + college +
employment + income + poverty + urbanicity + sex +
(1 | age) + (1 | race) + (1 | time) + (1 | zip)
Data: input_data (Number of observations: 96535)
Draws: 4 chains, each with iter = 10000; warmup = 5000; thin = 1;
total post-warmup draws = 20000
```

Multilevel Hyperparameters:

```
~age (Number of levels: 5)
      Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
sd(Intercept)    0.63     0.35    0.26    1.56 1.00    5555    7979
```

```
~race (Number of levels: 3)
      Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
sd(Intercept)    0.44     0.40    0.04    1.53 1.00    3352    2261
```

```
~time (Number of levels: 137)
      Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
sd(Intercept)    1.10     0.11    0.90    1.34 1.00    4828    7538
```

```
~zip (Number of levels: 848)
      Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
sd(Intercept)    0.37     0.11    0.12    0.58 1.00    3011    3011
```

Regression Coefficients:

```
      Estimate Est.Error l-95% CI u-95% CI Rhat Bulk_ESS Tail_ESS
Intercept   -5.17     0.45    -6.07   -4.23 1.00    4784    3877
ADI          0.18     0.16    -0.13    0.50 1.00   13351   10822
college     -0.07     0.11    -0.29    0.15 1.00   11813    8374
employment   0.04     0.10    -0.16    0.24 1.00   14137   14073
income      -0.00     0.13    -0.27    0.25 1.00   12066    9108
poverty     -0.09     0.10    -0.29    0.12 1.00   11149    8952
```



urbanicity	-0.17	0.09	-0.33	0.00	1.00	7576	3222
sexmale	0.19	0.09	0.01	0.37	1.00	14455	5400

Draws were sampled using sampling(NUTS). For each parameter, Bulk\_ESS and Tail\_ESS are effective sample size measures, and Rhat is the potential scale reduction factor on split chains (at convergence, Rhat = 1).

## B. Model estimates




Figure B.1: Estimated weekly prevalence by sex. The shaded areas represent one standard deviation of uncertainty.




Figure B.2: Estimated weekly prevalence by age group. The shaded areas represent one standard deviation of uncertainty.