# Gradient Descent for Noisy Optimization

Annie Hu and Mathieu Gerber

School of Mathematics

University of Bristol

### Abstract

We study the use of gradient descent with backtracking line search (GD-BLS) to solve the noisy optimization problem  $\theta_* := \operatorname{argmi}_{\theta \in \mathbb{R}^d} \mathbb{E}[f(\theta, Z)]$ , imposing that the function  $F(\theta) := \mathbb{E}[f(\theta, Z)]$  is strictly convex. Assuming that  $\mathbb{E}[||\nabla_{\theta}f(\theta_*, Z)||^2] < \infty$  and that objective function is locally *L*-smooth, we first prove that GD-BLS allows to estimate  $\theta_*$  with an error of size  $\mathcal{O}_{\mathbb{P}}(B^{-0.25})$ , where *B* is the available computational budget. We then show that we can improve upon this rate by stopping the optimization process earlier when the gradient of the objective function is sufficiently close to zero, and use the residual computational budget to optimize, again with GD-BLS, a finer approximation of *F*. By iteratively applying this strategy *J* times, we establish that we can estimate  $\theta_*$  with an error of size  $\mathcal{O}_{\mathbb{P}}(B^{-\frac{1}{2}(1-\delta^J)})$ , where  $\delta \in (1/2, 1)$  is a user-specified parameter. More generally, we show that if  $\mathbb{E}[||\nabla_{\theta}f(\theta_*, Z)||^{1+\alpha}] < \infty$  for some known  $\alpha \in (0, 1]$  then this approach allows to learn  $\theta_*$  with an error of size  $\mathcal{O}_{\mathbb{P}}(B^{-\frac{1}{2}(1-\delta^J)})$ , where  $\delta \in (2\alpha/(1+3\alpha), 1)$  is a tuning parameter. Beyond knowing  $\alpha$ , achieving the aforementioned convergence rates do not require to tune the algorithms parameters according to the specific functions *F* and *f* at hand, and we exhibit a simple noisy optimization problem for which stochastic gradient is not guaranteed to converge while the algorithms discussed in this work are.

# 1 Introduction

### 1.1 Set-up and problem formulation

We consider the problem of computing

$$\theta_{\star} := \operatorname*{argmin}_{\theta \in \mathbb{R}^d} F(\theta), \quad F(\theta) = \mathbb{E}[f(\theta, Z)]$$
(1)

where Z is a random variable taking its values in a Polish space Z and where  $f : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$  is a measurable function such that  $\nabla_{\theta} f(\theta, z)$  exists for all  $(\theta, z) \in \Theta \times \mathbb{Z}$ . We assume that only the function f can be evaluated point-wise and that to solve (1) we have at our disposal a sequence  $(Z_i)_{i\geq 1}$  of i.i.d. copies of Z. All the random variables are assumed to be defined on a complete probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and the function F is assumed to be strictly convex on  $\mathbb{R}^d$ . Remark that we do not assume that the function  $\theta \mapsto f(\theta, Z)$  is  $\mathbb{P}$ -almost surely convex.

Due to their ease of implementation and low computational cost, stochastic gradient (SG) algorithms arguably constitute the main set of tools used in practice for solving this type of optimization problems (Patel et al., 2022). Stochastic gradient methods are for instance widely applied for training machine learning algorithms (Bottou et al., 2018) or for performing statistical inference in large datasets (Toulis and Airoldi, 2017). In order to guarantee that a SG algorithm is stable when deployed on an unbounded search space it is commonly assumed that the variance of the norm of the stochastic gradient  $\nabla_{\theta} f(\theta, Z)$  is bounded and that the function F is L-smooth, that is that the gradient of Fis L-Lipschitz (see for instance Patel, 2022; Garrigos and Gower, 2023, for a review of convergence results for SG). Under the additional assumption that F is twice differentiable, the condition that Fis L-smooth can be replaced by the weaker assumption that F is  $(L_0, L_1)$ -smooth, in the sense that  $\|\nabla^2 F(\theta)\| \leq L_0 + L_1 \|\nabla F(\theta)\|$  for all  $\theta \in \mathbb{R}^d$  and some finite constants  $L_0$  and  $L_1$  (see Zhang et al., 2019, Assumption 3). The assumption that the variance of  $\|\nabla_{\theta} f(\theta, Z)\|$  is bounded can also be weakened. Notably, letting C be a finite constant, it can instead be assumed that this variance is bounded by  $C + C \|\nabla F(\theta)\|^2$  for all  $\theta \in \mathbb{R}^d$  (see e.g. Assumption 4.3 in Bottou et al., 2018 and Assumption 2 in Faw et al., 2022) or that  $\mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^2] \leq C + C \|\nabla F(\theta)\|^2$  for all  $\theta \in \mathbb{R}^d$  (see e.g Khaled and Richtárik, 2020, Assumption 2). Convergence results for SG that hold under the weaker condition that  $\mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^{1+\alpha}]$  for some  $\alpha \in (0, 1]$  have been obtained by some authors, see for instance Wang et al. (2021); Patel et al. (2022).

These assumptions used to establish the convergence of SG algorithms may however be violated even for simple optimization problems. For instance, as shown in Patel et al. (2022), both the assumption that F is L-smooth and the assumption that the variance of  $\|\nabla_{\theta} f(\theta, Z)\|$  is bounded, as well as their variants mentioned in the previous paragraph, may not hold for parameter inference in Poisson regression models. More precisely, Patel et al. (2022, Proposition 3) show the following result:

**Proposition 1.** Consider the d = 1 dimensional optimization problem (1) where Z = (X, Y), with X and Y two independent Poisson random variables such that  $\mathbb{E}[X] = \mathbb{E}[Y] = 1$ , and where the function  $f : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$  is defined by

$$f(\theta, z) = -z_2 z_1 \theta + \exp(\theta z_1), \quad \theta \in \mathbb{R}, \quad z \in \mathsf{Z} := \mathbb{R}^2.$$

Then, the function F is strictly convex, twice continuously differentiable and  $\operatorname{Var}(\nabla_{\theta}f(\theta, Z)) < \infty$  for all  $\theta \in \mathbb{R}$ . However, F is neither globally L-smooth nor  $(L_0, L_1)$ -smooth, and there exists no finite constant C such that  $\operatorname{Var}(\nabla_{\theta}f(\theta, Z)) \leq C + C|\nabla F(\theta)|^2$  for all  $\theta \in \mathbb{R}$  or such that  $\mathbb{E}[|\nabla_{\theta}f(\theta, Z)|^2] \leq C + C|\nabla F(\theta)|^2$  for all  $\theta \in \mathbb{R}$  (and thus  $\sup_{\theta \in \mathbb{R}} \operatorname{Var}(\nabla_{\theta}f(\theta, Z)) = \infty$ ).

To the best of our knowledge, Patel et al. (2022) provide the only existing general convergence result for SG on unbounded spaces that holds under the assumption that F is only locally L-smooth, and which allows the variance of  $\|\nabla_{\theta} f(\theta, Z)\|$  to be unbounded. Patel et al. (2022) actually only assume that, for some  $\alpha \in (0, 1]$ , the gradient of F is locally  $\alpha$ -Hölder continuous and that  $\mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^{1+\alpha}]$ is finite for all  $\theta \in \mathbb{R}^d$ . However, to guarantee the stability of SG, their result requires that the Hölder constant of  $\nabla F$  on a ball centred at  $\theta$  and of size  $\mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^{1+\alpha}]^{\frac{1}{1+\alpha}}$  does not increase too quickly as  $\|\theta\| \to \infty$  (see Assumption 5 in Patel et al., 2022). This condition appears to be quite strong since, as we show in the following proposition, it is not satisfied for the optimization problem considered in Proposition 1.

**Proposition 2.** Consider the set-up of Proposition 1. Then, there exists no  $\alpha \in (0,1]$  such that Assumption 5 used in Patel et al. (2022) to establish the convergence of SG holds.

Finally, we note that Fehrman et al. (2020) study SG algorithms under the assumption that F is locally *L*-smooth and allowing  $\mathbb{E}[||\nabla_{\theta} f(\theta, Z)||^2]$  to be unbounded but, informally speaking, this reference only shows that SG will converge to  $\theta_{\star}$  if the starting value of the algorithm is sufficiently close to that point.

Consequently, as illustrated by Propositions 1-2, even for seemingly simple noisy optimization problems on  $\mathbb{R}^d$ , where F is strictly convex and  $\mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^2]$  is finite for all  $\theta \in \mathbb{R}^d$ , SG is not guaranteed to converge.

### **1.2** Contribution of the paper

In this paper we first show that gradient descent with backtracking line search (GD-BLS) can be used to solve (1) without facing the aforementioned difficulties of SG methods. In particular, the backtracking line search procedure for choosing the step-sizes ensures the stability of the algorithm assuming neither that F is L-smooth nor that  $\sup_{\theta \in \mathbb{R}^d} \mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^{1+\alpha}] < \infty$  for some  $\alpha \in (0, 1]$ .

More precisely, we assume throughout this work that there is a computational cost  $C_{\text{grad}} \in \mathbb{N}$  for computing  $\nabla_{\theta} f(\theta, z)$  and a computational cost  $C_{\text{eval}} \in \mathbb{N}$  for computing  $f(\theta, z)$  for a pair  $(\theta, z) \in \mathbb{R}^d \times \mathbb{Z}$ . Then, we first show that, given the available computational budget  $B \in \mathbb{N}$ , by applying GD-BLS to minimize the function  $F_{n_1(B)} := \frac{1}{n_1(B)} \sum_{i=1}^{n_1(B)} f(\cdot, Z_i)$  we obtain a consistent estimator  $\hat{\theta}_{1,B}$  of  $\theta_{\star}$ . The convergence rate of this estimator depends on the choice of the function  $n_1$  and, assuming for now that  $\mathbb{E}[\|\nabla_{\theta} f(\theta_{\star}, Z)\|^2] < \infty$  to simplify the discussion, the upper bound we obtain for this rate is minimized when  $n_1(B) \approx B^{1/2}$ , in which case  $\|\hat{\theta}_{1,B} - \theta_{\star}\| = \mathcal{O}_{\mathbb{P}}(B^{-0.25})$ . Remark that the  $B^{-0.25}$  convergence rate obtained for  $\hat{\theta}_{1,B}$  is slower than the standard  $B^{-0.5}$  convergence rate.

In order to save computational resources gradient descent algorithms are often stopped if a prespecified convergence criterion is reached, typically when the gradient of the objective function is sufficiently close to zero. Following this approach, we propose to allow the optimization of  $F_{n_1(B)}$  to stop earlier (i.e. before the computational budget B is exhausted) and, if any, to use the remaining computational resources to improve the estimation of  $\theta_{\star}$ . In this work, the optimization of  $F_{n_1(B)}$  is stopped if its gradient falls below some threshold  $\tau_1(B)$  and the residual computational budget is used to apply GD-BLS on a finer approximation  $F_{n_2(B)}$  of F (i.e.  $n_2(B) > n_1(B)$ ), using as initial value the estimate of  $\theta_{\star}$  obtained by minimizing  $F_{n_1(B)}$ . With a careful choice for the functions  $n_1$ ,  $\tau_1$  and  $n_2$ , we show that the resulting estimate  $\hat{\theta}_{2,B}$  of  $\theta_{\star}$  has an error of size  $B^{-0.5(1-\delta^2)}$ , with  $\delta \in (1/2, 1)$  a user specified parameter.

The approach outlined in the previous paragraph can be generalized as follows to any  $J \geq 2$ applications of GD-BLS on increasingly finer approximations of F: For  $j = 1, \ldots, J - 1$  we stop earlier the optimization of the function  $F_{n_j(B)}$  by GD-BLS if its gradient becomes smaller than some threshold  $\tau_j(B)$  and, if any, the remaining computational resources are used to repeat this process to the function  $F_{n_{j+1}(B)}$ , using as starting value the estimate  $\tilde{\theta}_{j,B}$  of  $\theta_{\star}$  obtained by minimizing  $F_{n_j(B)}$ . For a suitable choice of functions  $(n_j)_{j\geq 1}$  and  $(\tau_j)_{j\geq 1}$ , and a user specified parameter  $\delta \in (1/2, 1)$ , we prove that the estimator  $\hat{\theta}_{J,B}$  of  $\theta_{\star}$  defined by this approach has an error of size  $\mathcal{O}_{\mathbb{P}}(B^{-\frac{1}{2}(1-\delta^J)})$  while, for all  $j \in \{1, \ldots, J-1\}$ , the estimator  $\tilde{\theta}_{j,B}$  converges to  $\theta_{\star}$  at rate  $B^{-\frac{1}{2}(1-\delta^J)}$ . The assumptions under which these results are obtained are shown to be satisfied for the optimization problem considered in Proposition 1, and to achieve the aforementioned convergence rates it is not needed to tune the parameters of the algorithms according to the specific functions F and f at hand.

In practice, to observe the  $B^{-\frac{1}{2}(1-\delta^J)}$  convergence rate of  $\hat{\theta}_{J,B}$  it is necessary that the optimization of  $F_{n_j(B)}$  stops earlier for all  $j \in \{1, \ldots, J-1\}$ , and thus that GD-BLS is applied to the function  $F_{n_J(B)}$ . Informally speaking, if we denote by  $J_B$  the smallest value of  $j \in \{1, \ldots, J\}$  such that  $\tilde{\theta}_{j,B} = \hat{\theta}_{J,B}$ , we therefore expect the observed convergence rate for  $\hat{\theta}_{J,B}$  to be  $B^{-\frac{1}{2}(1-\delta^{J_B})}$ . In this work we do not study theoretically the behaviour of  $J_B$  but the numerical experiments suggest that, as B increases, this random variable 'grows' at speed log B. In some applications the variance of the noisy gradient is not finite, and in this work the procedure described above is studied under the weaker assumption that  $\mathbb{E}[\|\nabla_{\theta}f(\theta_{\star}, Z)\|^{1+\alpha}] < \infty$  for some  $\alpha \in (0, 1]$ . In this scenario, the user needs to choose a noise parameter  $\alpha' \in (0, 1]$  and a  $\delta \in (2\alpha'/(1+3\alpha'), 1)$ , and the proposed approach for computing  $\theta_{\star}$  is proved to be consistent for any  $\alpha' \in (0, 1]$ . In addition, if  $\alpha' = \alpha$  then its convergence rate for computing  $\theta_{\star}$  is shown to be  $B^{-\frac{\alpha}{1+\alpha}(1-\delta^J)}$ . We recall the reader that  $B^{-\frac{\alpha}{1+\alpha}}$  is the best possible rate we can have for a first-order stochastic convex optimization algorithm under the assumption that  $\mathbb{E}[\|\nabla_{\theta}f(\theta_{\star}, Z)\|^{1+\alpha}] < \infty$  (see Vural et al., 2022, and references therein).

To contrast these results with the known properties of SG we note that, recently, Nguyen et al. (2024) showed that clipped SG achieves the optimal  $B^{-\frac{\alpha}{1+\alpha}}$  rate, assuming that  $\alpha$  is known. However, the results obtained in this reference rely on the strong assumption that  $\sup_{\theta \in \mathbb{R}^d} \mathbb{E}[\|\nabla_{\theta} f(\theta, Z)\|^{1+\alpha}] < \infty$  and require the function F to be L-smooth. The behaviour of standard SG algorithms under infinite noise variance is studied in Wang et al. (2021), but a very particular structure on the noisy gradient  $\nabla_{\theta} f(\theta, Z)$  is imposed in addition to an unusual smoothness condition on F.

### 1.3 Outline of the paper

The rest of the paper is organized as follows. In Section 2 we state the assumptions we impose on (1) and present some preliminary results. The use of GD-BLS for solving noisy optimization problems is studied in Section 3. In Section 4 we illustrate the convergence results derived in this work with some numerical experiments and Section 5 concludes. All the proofs are gathered in the appendix.

# 2 Notation, assumptions and some preliminary results

# 2.1 Notation

To simplify the notation, in what follows the operator  $\nabla$  is assumed to be such that  $\nabla g(\theta, z) = \frac{\partial}{\partial \theta} g(\theta, z)$  for all differentiable functions  $g : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$ , so that  $\nabla f(\theta, z) = \nabla_{\theta} f(\theta, z)$ . Similarly, the operator  $\nabla^2$  is assumed to be such that  $\nabla^2 g(\theta, z) = \frac{\partial^2}{\partial \theta \partial \theta^+} g(\theta, z)$  for all twice differentiable functions  $g : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$ , and we use the convention that  $\nabla^0$  is the identity operator, that is that  $\nabla^0 g = g$  for any function  $g : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$ .

Next, for all  $n \in \mathbb{N}$  and  $\theta \in \mathbb{R}^d$  we let  $F_n(\theta) = \frac{1}{n} \sum_{i=1}^n f(\theta, Z_i)$ ,

$$\nabla^k F_n(\theta) = \frac{1}{n} \sum_{i=1}^n \nabla^k f(\theta, Z_i), \quad \forall k \in \{0, 1, 2\}$$

and, for all non-empty compact set  $\Theta \subset \mathbb{R}^d$  and  $n \in \mathbb{N}$ , we let  $\hat{\theta}_{\Theta,n} \in \operatorname{argmin}_{\theta \in \Theta} F_n(\theta)$ . Finally, we denote by  $\lambda_{\min}(M)$  and  $\lambda_{\max}(M)$  the smallest and largest eigenvalue of a  $d \times d$  matrix M, respectively.

### 2.2 Assumptions

We consider the following two assumptions:

Assumption A1. The function F is locally strongly convex.

Remark that Assumption A1 implies that F is strictly convex on  $\mathbb{R}^d$  and thus that  $\theta_*$  is well-defined.

**Assumption A2.** The function  $\theta \mapsto f(\theta, z)$  is twice continuously differentiable on  $\mathbb{R}^d$  for all  $z \in \mathbb{R}^d$ , and

$$\mathbb{E}[\|\nabla^2 f(\theta_\star, Z)\|] < \infty, \quad \mathbb{E}[\|\nabla f(\theta_\star, Z)\|^{1+\alpha}] < \infty$$

for some  $\alpha \in (0,1]$ . Moreover, for all (non-empty) compact set  $\Theta \subset \mathbb{R}^d$  there exists a measurable function  $M_{\Theta} : \mathsf{Z} \mapsto [0,\infty)$  such that  $\mathbb{E}[M_{\Theta}(Z)] < \infty$  and such that

$$\|\nabla^2 f(\theta, Z) - \nabla^2 f(\theta', Z)\| \le M_{\Theta}(Z) \|\theta - \theta'\|, \quad \forall (\theta, \theta') \in \Theta^2, \quad \mathbb{P}-a.s.$$

These two assumptions are quite weak. In particular, as shown in the following proposition, they are satisfied for the optimization problem considered in Propositions 1-2.

**Proposition 3.** Consider the set-up of Proposition 1. Then, Assumptions A1-A2 are satisfied with  $\alpha = 2$ .

### 2.3 Some implications of Assumptions A1-A2

The following lemma notably implies that, under A2, the random function  $\theta \mapsto f(\theta, Z)$  is  $\mathbb{P}$ -a.s. locally *L*-smooth.

**Lemma 1.** Assume that A2 holds. Then, for all (non-empty) compact set  $\Theta \subset \mathbb{R}^d$  there exists a measurable function  $M'_{\Theta} : \mathsf{Z} \mapsto [0, \infty)$  such that  $\mathbb{E}[M'_{\Theta}(Z)] < \infty$  and such that, for all  $k \in \{0, 1, 2\}$ , we have

$$\|\nabla^k f(\theta, Z) - \nabla^k f(\theta', Z)\| \le M'_{\Theta}(Z) \|\theta - \theta'\|, \quad \forall (\theta, \theta') \in \Theta^2, \quad \mathbb{P}-a.s.$$

Remark that, in the set-up of Lemma 1, for any  $\epsilon > 0$  we may have  $\mathbb{E}[M'_{\Theta}(Z)^{1+\epsilon}] = \infty$ . Therefore, under A2, the quantity  $\mathbb{E}[\|\nabla f(\theta, Z)\|^{1+\alpha}]$  is assumed to be finite for some  $\alpha \in (0, 1]$  only if  $\theta = \theta_{\star}$ . In addition, it is worth noting that no conditions are imposed on the behaviour of  $M'_{\Theta}(\cdot)$  as the size of the set  $\Theta$  increases.

The next lemma shows that, under A2, we can swap the expectation and the differential operators.

**Lemma 2.** Assume that A2 holds. Then, the function  $F(\cdot)$  is twice continuously differentiable on  $\mathbb{R}^d$  and

$$\nabla^k F(\theta) = \mathbb{E}[\nabla^k f(\theta, Z)], \quad \forall \theta \in \mathbb{R}^d, \quad \forall k \in \{0, 1, 2\}.$$

By combining Lemma 1 and Lemma 2, it follows that under A2 the function F is locally L-smooth. More precisely, under A2, for any (non-empty) compact set  $\Theta \subset \mathbb{R}^d$  we have

$$\|\nabla F(\theta) - \nabla F(\theta')\| \le \mathbb{E}[M'_{\Theta}(Z)] \|\theta - \theta'\|, \quad \forall (\theta, \theta') \in \Theta^2$$

with  $M'_{\Theta}(\cdot)$  as in Lemma 1. We stress that, unlike in Patel et al. (2022), no conditions are imposed on the behaviour of the Lipschitz constant  $\mathbb{E}[M'_{\Theta}(Z)]$  as the size of the set  $\Theta$  increases.

The following lemma shows that the random function  $F_n(\cdot)$  converges uniformly to F on any compact subset of  $\mathbb{R}^d$ .

**Lemma 3.** Assume that A2 holds. Then, for all (non-empty) compact set  $\Theta \subset \mathbb{R}^d$  we have

$$\lim_{n \to \infty} \sup_{\theta \in \Theta} \left\| \nabla^k F_n(\theta) - \nabla^k F(\theta) \right\| = 0, \quad \mathbb{P} - a.s. \quad \forall k \in \{0, 1, 2\}.$$

The next lemma provides a useful result on the asymptotic behaviour of the largest and smallest eigenvalues of the random matrix  $\nabla^2 F_n(\theta)$ .

**Lemma 4.** Assume that A2 holds. Then, for all (non-empty) compact set  $\Theta \subset \mathbb{R}^d$  there exists a  $\theta_{\Theta} \in \Theta$  such that

$$\liminf_{n \to \infty} \left( \inf_{\theta \in \Theta} \lambda_{\min} \left( \nabla^2 F_n(\theta) \right) \right) \ge \lambda_{\min} \left( \mathbb{E} \left[ \nabla^2 f(\theta_{\Theta}, Z) \right] \right), \quad \mathbb{P} - a.s.$$

and a  $\theta'_{\Theta} \in \Theta$  such that

$$\limsup_{n \to \infty} \left( \sup_{\theta \in \Theta} \lambda_{\max} \left( \nabla^2 F_n(\theta) \right) \right) \le \lambda_{\max} \left( \mathbb{E} \left[ \nabla^2 f(\theta'_{\Theta}, Z) \right] \right), \quad \mathbb{P} - a.s$$

By combining Lemma 2 and Lemma 4, we readily obtain the following result which notably shows that if  $\Theta \subset \mathbb{R}^d$  is a (non-empty) compact set then the random function  $F_n$  is strictly convex on  $\Theta$  with  $\mathbb{P}$ -probability tending to one.

**Corollary 1.** Assume that A1-A2 hold. Then, for all (non-empty) compact set  $\Theta \subset \mathbb{R}^d$  there exists a constant  $c_{\Theta} \in (0,1)$  such that

$$\liminf_{n \to \infty} \left( \inf_{\theta \in \Theta} \lambda_{\min} \left( \nabla^2 F_n(\theta) \right) \right) \ge c_{\Theta}, \quad \limsup_{n \to \infty} \left( \sup_{\theta \in \Theta} \lambda_{\max} \left( \nabla^2 F_n(\theta) \right) \right) \le \frac{1}{c_{\Theta}}, \quad \mathbb{P}-a.s.$$

The last lemma establishes that, under A1 and A2, by using the first n elements of  $(Z_i)_{i\geq 1}$  we can learn  $\theta_{\star}$  at rate  $n^{-\frac{\alpha}{1+\alpha}}$ , with  $\alpha \in (0, 1]$  as in A2. In addition to be used to establish our main results, this lemma will allow us to discuss the convergence rate of the proposed estimators, obtained in the next section.

**Lemma 5.** Assume that A1-A2 hold and let  $\Theta \subset \mathbb{R}^d$  be a non-empty compact and convex set such that  $\theta_{\star} \in \mathring{\Theta}$ . Then,  $\|\theta_{\star} - \hat{\theta}_{\Theta,n}\| = \mathcal{O}_{\mathbb{P}}(n^{-\frac{\alpha}{1+\alpha}})$ , with  $\alpha \in (0,1]$  as in A2.

### 2.4 Gradient descent with backtracking line search

Algorithm 1 provides a generic description of a gradient descent algorithm that can be used to compute the global minimum  $g_{\star} := \inf_{x \in \mathbb{R}^d} g(x)$  of a convex and differentiable function  $g : \mathbb{R}^d \to \mathbb{R}$ . The convergence behaviour of Algorithm 1 depends on the value of the step-size  $v_t$  chosen on its Line 2 and, in particular, we have the following well-known result (whose proof is given in the appendix for sake of completeness).

**Proposition 4.** Assume that  $g_* > -\infty$ . Let  $x_0 \in \mathbb{R}^d$  and, for all  $T \in \mathbb{N}$ , let  $x_T$  be as computed by Algorithm 1 with  $(v_t)_{t>1}$  such that, for some constant c > 0,

$$g(x_t) \le g(x_{t-1}) - \frac{v_t}{2} \|\nabla g(x_{t-1})\|^2, \quad v_t > c, \quad \forall t \ge 1.$$
(2)

Then,

$$g(x_T) - g_\star \le \frac{\|x_0 - x_\star\|^2}{2cT}, \quad \forall T \ge 1.$$

If the function g is L-smooth (assuming without loss of generality that L > 1) then (2) holds for instance when  $v_t = v$  for all  $t \ge 1$  and some  $v \in (0, 1/L]$  (see e.g. the proof of Nesterov, 2018, Theorem 2.1.14, page 80). In practice, the value of L is however rarely known and backtracking line search, described in Algorithm 2, is a simple procedure that can be used to automatically compute, at each

Algorithm 1 (Generic gradient descent algorithm)

**Input:** Starting value  $x_0 \in \mathbb{R}^d$  and number of iterations  $T \in \mathbb{N}$ . 1: for t = 1, ..., T do 2: Choose a set-size  $v_t$ 3:  $x_t = x_{t-1} - v_t \nabla g(x_{t-1})$ 4: end for **Return:**  $x_T$ 

Algorithm 2 (Backtracking line search)

**Input:**  $x_{t-1} \in \mathbb{R}^d$  and backtracking parameter  $\beta \in (0, 1)$ . 1:  $g_{t-1} = \nabla g(x_{t-1})$ 2:  $v_t = 1$ 3: while  $g(x_{t-1} - v_t ||g_{t-1}||) > g(x_{t-1}) - \frac{v_t}{2} ||g_{t-1}||^2$  do 4:  $v_t \leftarrow \beta v_t$ 5: end while 6: return:  $v_t$ 

iteration t, a value for  $v_t$  which ensures that (2) holds with  $c = \min(1, \beta/L)$ , where  $\beta \in (0, 1)$  is a parameter of Algorithm 2 (see e.g. Boyd and Vandenberghe, 2004, page 465).

Interestingly, for the backtracking line search procedure to ensure that (2) holds, it is enough to assume that the function g is locally L-smooth, as shown in the next proposition.

**Proposition 5.** Assume that  $g_* > -\infty$  and that the function g is locally L-smooth. Let  $x_0 \in \mathbb{R}^d$ ,  $\beta \in (0,1)$  and, for all  $T \in \mathbb{N}$ , let  $x_T$  be as computed by Algorithm 1 where Algorithm 2 is used to compute  $v_t$  for all  $t \geq 1$ . In addition, let  $\tilde{K} \subset \mathbb{R}^d$  be a compact and convex set such that  $\{x \in \mathbb{R}^d : g(x) \leq g(x_0)\} \subseteq \tilde{K}$ , let

$$K = \left\{ x \in \mathbb{R}^d : \exists x' \in \tilde{K} \text{ such that } \|x - x'\| \le \sup_{\tilde{x} \in \tilde{K}} \|\nabla g(\tilde{x})\| \right\}$$

and, noting that K is a compact set, let  $L_K \in [1, \infty)$  be such that  $\|\nabla g(x) - \nabla g(x')\| \le L_K \|x - x'\|$  for all  $x, x' \in K$ . Then, (2) holds with  $\beta/L_K$ .

Given the large literature on convex optimization, the conclusion of Proposition 5 is probably known. We however failed to find a reference establishing a similar result and therefore the proposition is proved in the appendix.

# **3** Using gradient descent for noisy optimization

### 3.1 A simple application of gradient descent to noisy optimization

Recall that it is assumed in this work that there is a computational cost  $C_{\text{grad}} \in \mathbb{N}$  for computing  $\nabla f(\theta, z)$  and a computational cost  $C_{\text{eval}} \in \mathbb{N}$  for computing  $f(\theta, z)$  for a given pair  $(\theta, z) \in \mathbb{R}^d \times \mathbb{Z}$ . Then, for a given computational budget  $B \in \mathbb{N}$ , a simple way to estimate  $\theta_{\star}$  is to apply Algorithm 1 to minimize the function  $g = F_{n(B)}$  for some  $n(B) \in \mathbb{N}$ , using at each iteration the backtracking

### Algorithm 3

**Input:** Starting value  $\theta_0 \in \mathbb{R}^d$ , sample size  $n \in \mathbb{N}_0$ , computational budget  $B \in \mathbb{N}_0$ , tolerance parameter  $\tau \in [0, \infty)$  and backtracking parameter  $\beta \in (0, 1)$ . 1: Let t = 0,  $\theta_{B,t} = \theta_0$  and  $\tilde{B} = B$ 2: if  $\tilde{B} \ge nC_{\text{grad}}$  then Let  $G_t = \nabla F_n(\theta_{B,t})$ 3:  $\tilde{B} \leftarrow \tilde{B} - nC_{\text{grad}}$ 4: 5: end if while  $||G_t|| > \tau$  and  $\tilde{B} \ge 2nC_{\text{eval}}$  do 6: Let  $g_t = F_n(\theta_{B,t}), g'_t = F_n(\theta_{B,t} - G_t)$  and  $v_t = 1$ 7:  $\tilde{B} \leftarrow \tilde{B} - 2nC_{\text{eval}}$ 8: while  $g'_t > g_t - \frac{v_t}{2} \|G_t\|^2$  and  $\tilde{B} \ge nC_{\text{eval}}$  do 9:  $v_t \leftarrow \beta v_t$ 10: $g_t' \leftarrow F_n(\theta_{B,t} - v_t G_t)$ 11: $\tilde{B} \leftarrow \tilde{B} - nC_{\text{eval}}$ 12:end while 13:if  $g'_t \leq g_t - \frac{v_t}{2} \|G_t\|^2$  then 14:Let  $\theta_{B,t+1} = \theta_{B,t} - v_t G_t$ 15: $t \leftarrow t + 1$ 16:if  $\tilde{B} \ge nC_{\text{grad}}$  then 17:Let  $G_t \leftarrow \nabla F_n(\theta_{B,t})$ 18: $\tilde{B} \leftarrow \tilde{B} - nC_{\text{grad}}$ 19:else 20:break 21:end if 22:end if 23:24: end while 25: return:  $(\hat{\theta}_{B,t}, \tilde{B} \vee 0)$ 

procedure described in Algorithm 2 for choosing the step-size, and running the algorithm until the computational resources are exhausted. Algorithm 3 (with  $\tau = 0$ ) provides a precise description of this procedure.

In order for the resulting estimator  $\tilde{\theta}_B$  of  $\theta_*$  to be consistent as  $B \to \infty$ , it is clear that we must have  $n(B) \to \infty$ , so that  $F_{n(B)} \to F$ , and  $n(B)/B \to 0$ , so that the number of iterations of the algorithm converges to infinity. The following theorem formalizes this intuitive reasoning and provides an upper bound on the convergence rate of  $\|\tilde{\theta}_B - \theta_*\|$ .

**Theorem 1.** Assume that A1-A2 hold. Let  $\theta_0 \in \mathbb{R}^d$ ,  $\beta \in (0,1)$  and  $n : \mathbb{N} \to \mathbb{N}$  be such that  $\lim_{B\to\infty} n(B)/B = 0$  and such that  $\lim_{B\to\infty} n(B) = \infty$ . In addition, for all  $B \in \mathbb{N}$  let

$$(\tilde{\theta}_B, \tilde{B}) = \text{Algorithm } 3(\theta_0, n(B), B, 0, \beta), \quad r(B) = \max\left(\sqrt{n(B)/B}, n(B)^{-\frac{\alpha}{1+\alpha}}\right)$$

with  $\alpha \in (0,1]$  as in A2. Then,  $\|\tilde{\theta}_B - \theta_\star\| = \mathcal{O}_{\mathbb{P}}(r(B))$  and  $\liminf_{B\to\infty} B^{\frac{\alpha}{1+3\alpha}}r(B) > 0$ .

The expression for the rate r(B) has a very intuitive and simple explanation. Letting  $\Theta$  be a

compact set containing  $\theta_{\star}$ , we have

$$\|\tilde{\theta}_B - \theta_\star\| \le \|\tilde{\theta}_B - \hat{\theta}_{\Theta, n(B)}\| + \|\hat{\theta}_{\Theta, n(B)} - \theta_\star\|$$
(3)

where the first term on the right-hand side is the numerical error resulting from using GD-BLS to minimize  $F_{n(B)}$ . Under the assumptions of the theorem, if T iterations of GD-BLS are performed then the numerical error is of size  $T^{-1/2}$ . In the proof of the theorem we show that, under A1-A2, the cost per iteration of Algorithm 3 is approximatively constant, implying that  $\tilde{\theta}_B$  is computed by running  $T \approx B/n(B)$  iterations of GD-BLS. Consequently, the numerical error  $\|\tilde{\theta}_B - \hat{\theta}_{\Theta,n(B)}\|$  is of size  $\sqrt{B/n(B)}$ , which is the first term appearing in the definition of r(B). The second term in (3) is the statistical error which, by Lemma 5, is of size  $\mathcal{O}_{\mathbb{P}}(n(B)^{-\frac{\alpha}{1+\alpha}})$ . Therefore, the rate r(B) given in Theorem 1 is simply the maximum between the numerical error and the statistical error.

The last part of Theorem 1 follows from the fact that the fastest rate at which r(B) can decrease to zero is obtained when the numerical error and the statistical error vanish at the same speed. It is direct to see that this happens when  $n(B) \approx B^{\frac{1+\alpha}{1+3\alpha}}$ , in which case  $r(B) = \mathcal{O}(B^{-\frac{\alpha}{1+3\alpha}})$ . This result is formalized in the following corollary.

**Corollary 2.** Consider the set-up of Theorem 1 with  $n : \mathbb{N} \to \mathbb{N}$  such that

$$0 < \liminf_{B \to \infty} \frac{n(B)}{B^{\frac{1+\alpha}{1+3\alpha}}} \le \limsup_{B \to \infty} \frac{n(B)}{B^{\frac{1+\alpha}{1+3\alpha}}} < \infty.$$

Then,  $\|\tilde{\theta}_B - \theta_\star\| = \mathcal{O}_{\mathbb{P}}(n(B)^{-\frac{\alpha}{1+\alpha}}) = \mathcal{O}_{\mathbb{P}}(B^{-\frac{\alpha}{1+3\alpha}}).$ 

# 3.2 An improved application of gradient descent to noisy optimization

### 3.2.1 The algorithm

The procedure sketched in the introductory section to improve upon the convergence rate for  $\tilde{\theta}_B$  obtained in Corollary 2 is presented in Algorithm 4. Before providing, in Theorem 2 below, a convergence result for this algorithm some comments are in order.

Firstly, in Algorithm 4, for all j a specific form is imposed on the number  $n_j(B)$  of elements of  $(Z_i)_{i\geq 1}$  and on the tolerance parameter  $\tau_j(B)$  that are used in the *j*-th call of Algorithm 3. This is to simplify the presentation of the algorithm and the conclusions of Theorem 2 hold for any alternative functions  $\tilde{n}_i$  and  $\tilde{\tau}_i$  satisfying

$$0 < \liminf_{B \to \infty} \frac{n_j(B)}{\tilde{n}_j(B)} \leq \limsup_{B \to \infty} \frac{n_j(B)}{\tilde{n}_j(B)} < \infty, \quad 0 < \liminf_{B \to \infty} \frac{\tau_j(B)}{\tilde{\tau}_j(B)} \leq \limsup_{B \to \infty} \frac{\tau_j(B)}{\tilde{\tau}_j(B)} < \infty.$$

Secondly, in Algorithm 4 we have  $\tau_j(B) > 0$  even when j = J, that is even in the last call of Algorithm 3 the optimization of the function  $F_{n_j(B)}$  is stopped earlier when its gradient is small. However, there is no reasons for stopping earlier the minimization of  $F_{n_J(B)}$  and our convergence result (Theorem 2) remains valid when, for j = J, line 5 of Algorithm 4 is replaced by

5: 
$$(\hat{\theta}_{j,B}, B_j) = \text{Algorithm } 3\Big(\hat{\theta}_{j-1,B}, \lceil \kappa B^{\gamma_j} \rceil, B_{j-1}, 0, \beta\Big).$$

Nevertheless, in its current form, Algorithm 4 has the following advantage: if we let  $J = J_1$  for some  $J_1 > 1$  but, for some  $J_2 \in \{1, \ldots, J_1 - 1\}$ , the computational budget is exhausted after the  $J_2$ -th

### Algorithm 4

Input: Starting value θ<sub>0</sub> ∈ ℝ<sup>d</sup>, computational budget B ∈ N, rate parameters α' ∈ (0, 1] and δ ∈ [0, 1), maximum number of GD-BLS calls J ∈ N, sample size parameters κ ∈ (0, ∞), tolerance parameter τ ∈ (0, ∞) and backtracking parameter β ∈ (0, 1).
1: Let j = 0, B<sub>0</sub> = B and θ̂<sub>0,B</sub> = θ<sub>0</sub>
2: while B<sub>j</sub> > 0 and j < J do</li>
3: Let j = j + 1 and γ<sub>j</sub> = 1 − δ<sup>j</sup>
4: Let (θ̂<sub>j,B</sub>, B<sub>j</sub>) = Algorithm 3(θ̂<sub>j-1,B</sub>, [κB<sup>γ<sub>j</sub></sup>], B<sub>j-1</sub>, τB<sup>- α'/(1+α')γ<sub>j</sub>, β)
5: end while
</sup>

6: return:  $\hat{\theta}_{j,B}$ 

call of Algorithm 3 then the estimate of  $\theta_{\star}$  returned by Algorithm 4 is the one we would have obtained if we had chosen  $J = J_2$ . From a practical point of view, this means that we can choose a large value for J and let the algorithm determines how many calls to Algorithm 3 can be made given the available computational resources.

### 3.2.2 Convergence result

In fact, expanding on the latter point, the only reason why Algorithm 4 imposes an upper bound J on the number of calls of Algorithm 3 is to prove the following result:

**Theorem 2.** Assume that A1-A2 hold. Let  $\alpha' \in (0,1]$ ,  $\theta_0 \in \mathbb{R}^d$ ,  $J \in \mathbb{N}$ ,  $(\kappa, \tau) \in (0,\infty)^2$ ,  $\beta \in (0,1)$ ,  $\delta \in (2\alpha'/(1+3\alpha'), 1)$  and

 $(\hat{\theta}_{J,B}, \tilde{B}) = \text{Algorithm} 4(\theta_0, B, \alpha', \delta, J, \kappa, \tau, \beta), \quad \forall B \in \mathbb{N}.$ 

Then,  $\|\hat{\theta}_{J,B} - \theta_{\star}\| = o_{\mathbb{P}}(1)$  and if A2 holds for  $\alpha = \alpha'$  then

$$\|\hat{\theta}_{J,B} - \theta_{\star}\| = \mathcal{O}\left(B^{-\frac{\alpha'}{1+\alpha'}(1-\delta^J)}\right).$$

Assuming that A2 holds for  $\alpha = \alpha'$ , it follows from Theorem 2 that taking a small value for J can only make slower the speed at which  $\hat{\theta}_{J,B}$  converges to  $\theta_{\star}$ . However, as mentioned in the introductory section, to observe in practice the  $\mathcal{O}(B^{-\frac{\alpha'}{1+\alpha'}(1-\delta^J)})$  convergence rate it is needed to minimize the function  $F_{n_J(B)}$ , which may not be the case if J is too large given the available computational budget. For this reason, we expect the observed convergence rate to be  $B^{-\frac{\alpha'}{1+\alpha'}(1-\delta^{J_B})}$ , with  $J_B$  the number of calls of Algorithm 3 needed for computing  $\hat{\theta}_{J,B}$ , that is with  $J_B$  the smallest value of  $j \in \{1, \ldots, J\}$ such that  $\hat{\theta}_{J_B,B} = \hat{\theta}_{J,B}$ . A sensible approach is therefore to let the algorithm decides how many calls to Algorithm 3 are needed by choosing J large enough so that  $J_B < J$  with probability close to one. The numerical experiments presented in the next section suggest that  $J_B$  increases with B at speed log B, and verifying theoretically this observation is left for future research.

Noting that, for any  $j \ge 1$ , increasing  $\delta$  makes the *j*-th call of Algorithm 3 computationally less expensive, it follows that  $J_B$  increases with  $\delta$ . Hence, the larger  $\delta$  the slower is the convergence rate of  $\hat{\theta}_{j,B}$  for any given  $j \ge 1$ , but the larger the number of calls to Algorithm 3 can be made. According to the experiments of Section 4, the value of  $\delta$  does not influence the observed convergence rate as long

as  $J_B < J$  with probability close to one, but the convergence behaviour of  $\hat{\theta}_{J,B}$  appears to be better when  $\delta$  is close to one.

Finally, if A2 does not hold for  $\alpha = \alpha'$  then the proof of Theorem 2 shows that  $\|\hat{\theta}_{J,B} - \theta_{\star}\|$  converges to zero at least at speed  $B^{-\frac{\alpha}{1+\alpha}(1-\delta)}$ , for any  $\alpha$  such that A2 holds. Studying more precisely the convergence rate of  $\hat{\theta}_{J,B}$  in this context is left for future research, but we expect this rate to be slower than the one obtained when A2 holds for the chosen value of  $\alpha'$ . To understand why, remark that in Algorithm 4 the parameter  $\alpha'$  only impacts the value of the tolerance parameter used in each call of Algorithm 3, and that the larger  $\alpha'$  the smaller the tolerance parameter. Then, if  $\alpha'$  is too large (i.e. if A2 does not hold for  $\alpha = \alpha'$ ), for any  $j \in \{1, \ldots, J\}$  too much computational budget is allocated to the *j*-th call of Algorithm 3, resulting in a waste of computational resources and thus in a slower convergence rate as  $B \to \infty$ . In particular, it may be the case that for any values of *B* we have  $J_B < J$  with high probability.

#### 3.2.3 Understanding where the fast convergence rate comes from

To understand why Algorithm 4 allows to significantly improve upon the convergence rate obtained in Section 3.1, consider the set-up of Theorem 1 with the fixed starting value  $\theta_0$  replaced by a sequence  $(\theta_{0,B})_{B>1}$  of starting values such that  $\|\theta_{0,B} - \theta_{\star}\| = \mathcal{O}(B^{-c})$  for some constant  $c \geq 0$ , that is let

$$(\hat{\theta}_{c,B}, \hat{B}) = \text{Algorithm } \mathbf{3}(\theta_{0,B}, n_c(B), B, 0, \beta), \quad \forall B \in \mathbb{N}$$

with  $n_c(\cdot)$  satisfying the assumptions of Theorem 1. In this case, our computations show that

$$\|\tilde{\theta}_{c,B} - \theta_{\star}\| = \mathcal{O}_{\mathbb{P}}(r_c(B)), \quad r_c(B) = \max\left(B^{-c}\sqrt{n_c(B)/B}, n_c(B)^{-\frac{\alpha}{1+\alpha}}\right)$$
(4)

where, as in Section 3.1 and for some  $\Theta \subset \mathbb{R}^d$ , the first term in the definition of  $r_c(B)$  is the numerical error  $\|\hat{\theta}_{c,B} - \hat{\theta}_{\Theta,n_c(B)}\|$  and the second one is the statistical error  $\|\hat{\theta}_{\Theta,n_c(B)} - \theta_{\star}\|$ . From (4) we observe that the value of c modifies the trade-off computational error versus statistical error we are facing when choosing  $n_c(B)$ , and these two sources of errors converge to zero at the same speed when

$$0 < \liminf_{B \to \infty} \frac{n_c(B)}{B^{\frac{(1+\alpha)(1+2c)}{1+3\alpha}}} \le \limsup_{B \to \infty} \frac{n_c(B)}{B^{\frac{(1+\alpha)(1+2c)}{1+3\alpha}}} < \infty.$$
(5)

in which case we have

$$r_c(B) = B^{-\frac{\alpha(1+2c)}{1+3\alpha}}.$$

Therefore, initializing Algorithm 1 with a starting value that converges to  $\theta_{\star}$  (i.e. having c > 0) allows to improve the rate at which we learn  $\theta_{\star}$ . In particular, for c = 0 we recover the result of Corollary 2, and the larger c the faster is the convergence rate of  $\tilde{\theta}_{c,B}$ .

### 3.2.4 Understanding the convergence rate given in Theorem 2

The convergence rate given in Theorem 2 is more easily understood by studying an alternative (but less practical, see below) estimator of  $\theta_{\star}$ .

To define this estimator let  $\delta_{\alpha'} = 2\alpha'/(1+3\alpha')$ ,  $c_j = (1-\delta_{\alpha'}^j)\alpha'/(1+\alpha')$  for all  $j \ge 0$  and  $\theta_0 \in \mathbb{R}^d$ . In addition, let  $J \in \mathbb{N}$  and  $\{\kappa_{J,j}\}_{j=1}^J$  be such that  $\sum_{j=1}^J \kappa_{J,j} = 1$  and such that  $\kappa_{J,j} > 0$  for all  $j \in \{1, \ldots, J\}$ . Finally, for all  $j \in \{1, \ldots, J\}$  and using the convention that  $\tilde{\theta}_{j-1,B,J} = \theta_0$  for all  $B \in \mathbb{N}$  when j = 1, let

$$(\tilde{\theta}_{j,B,J}, \tilde{B}_j) = \text{Algorithm } 3\Big(\tilde{\theta}_{j-1,B,J}, n_{c_{j-1}}(B), \lfloor \kappa_{J,j}B \rfloor, 0, \beta\Big), \quad \forall B \in \mathbb{N}$$

with the function  $n_{c_{j-1}}$  such that (5) holds (with  $c = c_{j-1}$ ). Then, assuming that A2 holds for  $\alpha = \alpha'$ and using the computations of Section 3.2.3, it is easily checked that for all  $j \in \{1, \ldots, J\}$  we have  $\|\tilde{\theta}_{j,B,J} - \theta_{\star}\| = \mathcal{O}_{\mathbb{P}}(B^{-c_j})$ , where

$$B^{-c_{j}} = \max\left(B^{-c_{j-1}}\sqrt{n_{c_{j-1}}(B)/B}, n_{c_{j-1}}(B)^{-\frac{\alpha'}{1+\alpha'}}\right), \quad \forall B \in \mathbb{N}.$$

Informally speaking, and following the reasoning of Section 3.2.3, this implies that  $\hat{\theta}_{j,B,J}$  is computed from  $\tilde{\theta}_{j-1,B,J}$  using the number  $n_{c_{j-1}}(B)$  of elements of  $(Z_i)_{i\geq 1}$  that maximizes the speed at which  $\|\tilde{\theta}_{j,B,J} - \theta_{\star}\|$  converges to zero. Using the definition of  $(c_j)_{j\geq 1}$ , it is readily checked that, for all  $j \in \{1, \ldots, J\}$ ,

$$\|\tilde{\theta}_{j,B,J} - \theta_{\star}\| = \mathcal{O}_{\mathbb{P}}\left(B^{-\frac{\alpha'}{1+\alpha'}}(1-\delta^{j}_{\alpha'})\right)$$

while Theorem 2 establishes that  $\|\hat{\theta}_{j,B} - \theta_{\star}\| = \mathcal{O}_{\mathbb{P}}\left(B^{-\frac{\alpha'}{1+\alpha'}(1-\delta^i)}\right)$  for  $\delta \in (\delta_{\alpha'}, 1)$ .

In addition to give some insight on this latter result, the above calculations show that  $\tilde{\theta}_{j,B,J}$  converges to  $\theta_{\star}$  (slightly) faster than  $\hat{\theta}_{j,B}$  for any  $j \in \{1, \ldots, J\}$ . However, these alternative estimators have the important drawback to require that the computational budget allocated to each call of Algorithm 3 is chosen beforehand, and in a way that depends on the chosen value of J. By contrast, in Algorithm 4 this allocation is done automatically and does not depend on J, and if only j < J calls to Algorithm 3 are made then the algorithm returns the estimate of  $\theta_{\star}$  we would have obtained if we had chosen J = j. In our calculations, this desirable properties of Algorithm 4 are obtained at the price of imposing that  $\delta$  is strictly bigger than  $\delta_{\alpha'}$ .

# 4 Numerical experiments

In this section we study the behaviour of  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  and of  $\mathbb{E}[J_B]$  as B increases, as estimated from 100 independent realizations of  $(Z_i)_{i\geq 1}$ . In all the experiments presented below Algorithm 4 is implemented with  $\beta = 0.5$ ,  $C_{\text{grad}} = C_{\text{eval}} = \kappa = 1$  and  $\tau = 0.1$ . In addition, following the discussion of Section 3.2, we let  $J = 10^4$  so that  $J_B < J$  with probability close to one. Finally, in what follows we denote by  $\rho_{J_B}$  the coefficient of correlation between the variables  $(\mathbb{E}[J_B])_{B\geq 1}$  and  $(\log B)_{B\geq 1}$ , as estimated from 100 independent realizations of  $(Z_i)_{i\geq 1}$ . Remark that a value of  $\rho_{J_B}$  close to one implies that  $\mathbb{E}[J_B] \approx a + b \log B$  for some constants  $a \in \mathbb{R}$  and  $b \in (0, \infty)$ .

### 4.1 Example of Proposition 1

We first consider the univariate optimization problem defined in Proposition 1, which amounts to solving the d = 1 dimensional noisy optimization problem (1) with Z = (X, Y), where X and Y are two independent Poisson random variables such that  $\mathbb{E}[X] = \mathbb{E}[Y] = 1$ , and with the function  $f : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$  defined by

$$f(\theta, z) = -z_2 z_1 \theta + \exp(\theta z_1), \quad \theta \in \mathbb{R}, \quad z \in \mathsf{Z} := \mathbb{R}^2.$$



Figure 1: Results for the example of Section 4.1. The left plot shows  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  as a function of B, where the solid line is for  $\delta = 0.95$  and the dashed line for  $\delta = 0.51$ , and with the two dotted lines representing the  $B^{-1/2}$  convergence rate. The middle and right plots show the evolution of  $\mathbb{E}[J_B]$  as a function of B for  $\delta = 0.95$  (middle plot) and for  $\delta = 0.51$  (right plot). All the results are obtained from 100 realizations of  $(Z_i)_{i>1}$ .

We recall that for this optimization problem Assumptions A1-A2 hold with  $\alpha = 2$  (see Proposition 3) and that, to the best of our knowledge, SG is not guaranteed to converge when used to solve it (see Propositions 1-2). Simple computations show that the global minimizer of the resulting objective function F is  $\theta_{\star} = 0$ . For this example we let  $\theta_0 = 1$  and  $\alpha' = 2$ , in which case Theorem 2 imposes to have  $\delta \in (1/2, 1)$ .

Figure 1 shows, for  $\delta = 0.51$  and for  $\delta = 0.95$ , the evolution of  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  and of  $\mathbb{E}[J_B]$  as B increases. For the two considered values of  $\delta$  the observed rate at which  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  decreases to zero is indistinguishable from the optimal  $B^{-1/2}$  convergence rate, and for any B the value of  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  is smaller for  $\delta = 0.95$  than for  $\delta = 0.51$ . As expected, we see in Figure 1 that the number of calls to Algorithm 3 is much larger for  $\delta = 0.95$  than for  $\delta = 0.95$  than for  $\delta = 0.95$  than for  $\delta = 0.51$ , with a value of  $\mathbb{E}[J_B]$  ranging from 27 to 38 for the former choice of  $\delta$  and ranging from 2.4 to 3.8 for the latter. Visually, we observe (at least for  $\delta = 0.95$ ) that  $\mathbb{E}[J_B]$  grows with B as log B. The coefficient of correlation between the variables  $(\mathbb{E}[J_B])_{B\geq 1}$  and  $(\log B)_{B\geq 1}$  confirms this observation, since for this experiment we have  $\rho_{J_B} = 0.994$  when  $\delta = 0.95$  and  $\rho_{J_B} = 0.799$  when  $\delta = 0.51$ .

### 4.2 A multivariate extension of the example of Proposition 1

We now consider the d = 20 dimensional extension of the example of Proposition 1 obtained by letting, in (1), Z = (W, X, Y) be such that

$$X \sim \text{Poisson}(1), \quad W|X \sim \mathcal{U}([-1,1]^{d-1}), \quad Y|(W,X) \sim \text{Poisson}(a_{\star}^{\top}W)$$

for some  $a_{\star} \in \mathbb{R}^{d-1}$ , and the function  $f : \mathbb{R}^d \times \mathsf{Z} \to \mathbb{R}$  be defined by

$$f(\theta, z) = -z_2 \theta^\top z_1 + \exp(\theta^\top z_1), \quad \theta \in \mathbb{R}^d, \quad z \in \mathsf{Z} := \mathbb{R}^{d+1}.$$

It is easily checked that the resulting objective function F is such that A1-A2 hold with  $\alpha = 2$ and that  $\theta_{\star} = (0, a_{\star})$ . Compared to the example of Proposition 1, we expect that by increasing the dimension of  $\theta$  we make the optimization problem more challenging and thus that the conclusions of



Figure 2: Results for the example of Section 4.2. The left plot shows  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  as a function of B, where the solid line is for  $\delta = 0.95$  and the dashed line for  $\delta = 0.51$ , and with the two dotted lines representing the  $B^{-1/2}$  convergence rate. The middle and right plots show the evolution of  $\mathbb{E}[J_B]$  as a function of B for  $\delta = 0.95$  (middle plot) and for  $\delta = 0.51$  (right plot). All the results are obtained from 100 realizations of  $(Z_i)_{i>1}$ 

Proposition 1-2 hold for this multivariate optimization problem. For this experiment the value  $a_{\star}$  is a random draw from the  $\mathcal{N}_{d-1}(0, \mathbb{1}_{d-1})$  distribution and, as in the previous subsection, the results are presented for  $(\alpha', \delta) = (1, 0.51)$  and for  $(\alpha', \delta) = (1, 0.95)$ . Finally, Algorithm 4 is implemented using  $\theta_0 = (1, \ldots, 1)$  as starting value.

From Figure 2, which shows the evolution of  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|]$  and of  $\mathbb{E}[J_B]$  as *B* increases, we see that the behaviour of these two quantities is similar to what we observed in the previous subsection. The main difference is however that, because of the larger dimension of  $\theta$ , the  $B^{-1/2}$  convergence rate for  $\hat{\theta}_{J,B}$  is observed only for *B* large enough (roughly for  $B > 10^6$  in the plots). The study of the dependence of the performance of Algorithm 4 to the dimension *d* of  $\theta$  is left for future research. Finally, in Figure 2 we also observe that  $\mathbb{E}[J_B]$  appears to increase with *B* as log *B*, and indeed, for this example, the coefficient of correlation between  $(\mathbb{E}[J_B])_{B\geq 1}$  and  $(\log B)_{B\geq 1}$  is  $\rho_{J_B} = 0.986$  for  $\delta = 0.95$ and  $\rho_{J_B} = 0.663$  for  $\delta = 0.51$ .

### 4.3 Example of Proposition 1 with an extra heavy tailed noise

We finally consider the example of Proposition 1 with an extra heavy tailed noise. More precisely, we aim at solving the d = 1 noisy optimization problem (1) with Z = (W, X, Y), where  $W \sim t_{\nu}(0, 1)$  and where X and Y are two independent Poisson random variables which are independent of W and such that  $\mathbb{E}[X] = \mathbb{E}[Y] = 1$ , and with the function  $f : \mathbb{R}^d \times \mathbb{Z} \to \mathbb{R}$  defined by

$$f(\theta, z) = -z_3 z_2 \theta + \exp(\theta z_2) + z_1 \theta, \quad \theta \in \mathbb{R}, \quad z \in \mathsf{Z} := \mathbb{R}^3.$$

We let  $\nu = 1.501$  so that A2 holds for any  $\alpha < 1 - \nu = 0.501$ . Remark that since  $\mathbb{E}[W] = 0$  it follows that the resulting objective function F is the same as for the example of Proposition 1, and thus  $\theta_{\star} = 0$ . It is also immediate to see that the conclusions of Proposition 1-2 remain valid for this example, and therefore to be best of our knowledge SG is not guaranteed to converge when applied to solve it. In addition, the results of Nguyen et al. (2024) for clipped SG do not apply to this optimization problem, notably because the function F is not L-smooth (see Proposition 1). For this example we implement Algorithm 4 with  $\theta_0 = 1$  and the results are presented for  $\alpha' = 0.5$  and for  $\alpha' = 1$ . Remark that A2



Figure 3: Results for the example of Section 4.3. The top plots shows  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_*\|_{0.1}]$  as a function of B and the bottom plots show  $\mathbb{E}[J_B]$  as a function of B. The left plots are for  $(\alpha', \delta) = (0, 5, 0.41)$ , the middle plots for  $(\alpha', \delta) = (0.5, 0.95)$  and the right plots for  $(\alpha', \delta) = (1, 0.95)$ . In the top plots the dotted lines represent the  $B^{-1/3}$  convergence rate and all the results are obtained from 100 realizations of  $(Z_i)_{i>1}$ 

holds with  $\alpha = \alpha'$  for the former value of  $\alpha'$  but not for the latter. Remark also that, by Theorem 2, we must have  $\delta \in (0.4, 1)$  when  $\alpha' = 0.5$  and  $\delta \in (0.5, 1)$  when  $\alpha' = 1$ .

Due to the heavy tailed noise, for this example it is not guaranteed that we have  $\mathbb{E}[\|\hat{\theta}_{J,B}\|] < \infty$ and the numerical experiments suggest that this may indeed not be the case. For this reason, below we study, in Figure 3, the behaviour as *B* increases of  $\mathbb{E}[\|\hat{\theta}_{J,B} - \theta_{\star}\|_{0.1}]$ , the expected 10% trimmed mean of  $\|\hat{\theta}_{J,B} - \theta_{\star}\|$ .

The plots in the first two columns of Figure 3 are obtained for  $(\alpha', \delta) = (0.5, 0.41)$  and for  $(\alpha', \delta) = (0.5, 0.95)$ . From these plots we see that the convergence rate of  $\hat{\theta}_{J,B}$  is undistinguishable from the  $B^{-\frac{\alpha'}{1+\alpha'}} = B^{-1/3}$  convergence rate, confirming the result of Theorem 2. As for the examples of Sections 4.1-4.2, the estimation error tends to be the smallest when  $\delta$  is large, that is when  $\delta = 0.95$ . We also observe that for the two considered values of  $\delta$  the quantity  $\mathbb{E}[J_B]$  appears to grow with B as log B, an observation which is confirmed by the fact that  $\rho_{J_B} = 0.840$  for  $\delta = 0.41$  and  $\rho_{J_B} = 0.982$  for  $\delta = 0.95$ . The plots in last column of Figure 3 study the behaviour of  $\hat{\theta}_{J,B}$  when  $(\alpha', \delta) = (1, 0.95)$ , in which case A2 does not hold for  $\alpha = \alpha'$ . If  $\hat{\theta}_{J,B}$  appears to roughly converge to  $\theta_{\star}$  at speed  $B^{-1/3}$ , we observe that the convergence becomes slower for values of B greater than  $10^{6.7}$ . Remark that, from the discussion of Section 3.2.2, it was expected that the convergence rate of  $\hat{\theta}_{J,B}$  is slower when A2 does not hold

for the chosen value of  $\alpha'$  than when it does. Finally, for this value of  $(\alpha', \delta)$  the quantity  $\mathbb{E}[J_B]$  also seems to grow as log B, an observation which is confirmed by the large value  $\rho_{J_B} = 0.968$  obtained for the coefficient of correlation between  $(\mathbb{E}[J_B])_{B\geq 1}$  and  $(\log B)_{B\geq 1}$ .

# 5 Concluding remarks

In this paper we show, theoretically and empirically, that gradient descent with backtracking line search can be used to solve noisy optimization problems with a nearly optimal convergence rate, and without assuming that the objective function is globally *L*-smooth.

The proposed algorithm (Algorithm 4) requires that GD-BLS can be called at most J times, with J a user-specified parameter. We conjecture that this condition can be removed and, assuming that A2 holds for  $\alpha = \alpha'$ , that the resulting estimator  $\hat{\theta}_B$  of the global minimiser  $\theta_{\star}$  converges at speed  $B^{-\frac{\alpha'}{1+\alpha'}(1-j_B)}$  for some sequence  $(j_B)_{B\geq 1}$  such that  $\lim_{B\to\infty} j_B = \infty$ . Based on our numerical experiments, we conjecture that  $j_B = a + b \log B$  for some constants  $a \in [0, \infty)$  and  $b \in (0, \infty)$ depending on the choice of the  $\delta$  parameter of Algorithm 4. Confirming these results theoretically is left for future research.

Another interesting question that deserves further work is the possibility to replace, in Algorithm 4, GD-BLS by GD-BLS with Nesterov momentum (GD-BLS-NM). We recall the reader that, while the convergence of the former algorithm for computing  $\min_{x \in \mathbb{R}^d} g(x)$  for a convex and *L*-smooth function g is 1/T, with T the number of iterations, GD-BLS-NM converges to the global minimum of g at the faster (and optimal)  $1/T^2$  rate (see Nesterov, 2018, Section 2.2). Based on this result, it is reasonable to expect that replacing GD-BLS by GD-BLS-NM will improve the non-asymptotic behaviour of the proposed procedure for computing  $\theta_{\star}$ . The extension of our convergence results to the case where GD-BLS-NM is used is however non-trivial, notably because it is unclear if the aforementioned fast convergence for this algorithm remains valid when the function g is only locally *L*-smooth. Indeed, the key reason why the 1/T convergence rate of GD-BLS holds in this context is that the backtracking procedure ensures that the estimate of  $\min_{x \in \mathbb{R}^d} g(x)$  improves from one iteration to the next, and this property of backtracking does not apply when Nesterov momentum is used.

# References

Amann, H., Escher, J., and Brookfield, G. (2008). Analysis II. Springer.

- Bottou, L., Curtis, F. E., and Nocedal, J. (2018). Optimization methods for large-scale machine learning. SIAM review, 60(2):223–311.
- Boyd, S. P. and Vandenberghe, L. (2004). Convex optimization. Cambridge university press.
- Bubeck, S., Cesa-Bianchi, N., and Lugosi, G. (2013). Bandits with heavy tail. IEEE Transactions on Information Theory, 59(11):7711–7717.
- Faw, M., Tziotis, I., Caramanis, C., Mokhtari, A., Shakkottai, S., and Ward, R. (2022). The power of adaptivity in sgd: Self-tuning step sizes with unbounded gradients and affine variance. In *Conference* on Learning Theory, pages 313–355. PMLR.

- Fehrman, B., Gess, B., and Jentzen, A. (2020). Convergence rates for the stochastic gradient descent method for non-convex objective functions. *The Journal of Machine Learning Research*, 21(1):5354– 5401.
- Garrigos, G. and Gower, R. M. (2023). Handbook of convergence theorems for (stochastic) gradient methods. arXiv preprint arXiv:2301.11235.
- Khaled, A. and Richtárik, P. (2020). Better theory for sgd in the nonconvex world. arXiv preprint arXiv:2002.03329.
- Nesterov, Y. (2018). Lectures on convex optimization, volume 137. Springer.
- Nguyen, T. D., Nguyen, T. H., Ene, A., and Nguyen, H. (2024). Improved convergence in high probability of clipped gradient methods with heavy tailed noise. Advances in Neural Information Processing Systems, 36.
- Patel, V. (2022). Stopping criteria for, and strong convergence of, stochastic gradient descent on bottou-curtis-nocedal functions. *Mathematical Programming*, 195(1-2):693-734.
- Patel, V., Zhang, S., and Tian, B. (2022). Global convergence and stability of stochastic gradient descent. Advances in Neural Information Processing Systems, 35:36014–36025.
- Toulis, P. and Airoldi, E. M. (2017). Asymptotic and finite-sample properties of estimators based on stochastic gradients. *The Annals of Statistics*, 45(4):1694 1727.
- Van der Vaart, A. W. (2000). Asymptotic Statistics. Cambridge series in statistical and probabilistic mathematics.
- Vural, N. M., Yu, L., Balasubramanian, K., Volgushev, S., and Erdogdu, M. A. (2022). Mirror descent strikes again: Optimal stochastic convex optimization under infinite noise variance. In *Conference* on Learning Theory, pages 65–102. PMLR.
- Wang, H., Gurbuzbalaban, M., Zhu, L., Simsekli, U., and Erdogdu, M. A. (2021). Convergence rates of stochastic gradient descent under infinite noise variance. Advances in Neural Information Processing Systems, 34:18866–18877.
- Zhang, J., He, T., Sra, S., and Jadbabaie, A. (2019). Why gradient clipping accelerates training: A theoretical justification for adaptivity. arXiv preprint arXiv:1905.11881.

# A Proofs

# A.1 Proof of Proposition 2

*Proof of Proposition 2.* For all  $\theta \in \mathbb{R}$  we have

$$F(\theta) = -\theta + \exp\left(\exp(\theta) - 1\right)$$
  

$$\nabla F(\theta) = -1 + \exp\left(\exp(\theta) + \theta - 1\right)$$
  

$$\nabla^2 F(\theta) = \left(\exp(\theta) + 1\right) \exp\left(\exp(\theta) + \theta - 1\right).$$
  
(6)

We now show that there exists no  $\alpha \in (0, 1]$  such that Assumption 5 of Patel et al. (2022) holds. To this aim, for all  $\alpha \in (0, 1]$  we let

$$G_{\alpha}(\theta) = \mathbb{E}[|\nabla f(\theta, Z)|^{1+\alpha}], \quad \forall \theta \in \mathbb{R}$$
(7)

and, for all  $\epsilon > 0$  and all  $\theta \in \mathbb{R}$ , we let  $\mathcal{L}_{\alpha,\epsilon}(\theta) \in (0,\infty)$  be the smallest constant  $L \in (0,\infty)$  such that

$$\left|\nabla F(\theta) - \nabla F(\theta')\right| \le L \|\theta - \theta'\|^{1+\alpha}, \quad \forall \theta' \in B_{r_{\alpha,\epsilon,\theta}}(\theta)$$

where  $r_{\alpha,\epsilon,\theta} = (G_{\alpha}(\theta) \vee \epsilon)^{\frac{1}{1+\alpha}}$  and where  $B_r(\theta)$  denote the closed ball centred at  $\theta \in \mathbb{R}$  and with radius r > 0.

Then, to show that Assumption 5 of Patel et al. (2022) does not hold it is enough to show that

$$\liminf_{\theta \to \infty} \frac{\mathcal{L}_{\alpha,\epsilon}(\theta) G_{\alpha}(\theta)}{F(\theta) + |\nabla F(\theta)|^2} = \infty, \quad \forall \epsilon > 0, \quad \forall \alpha \in (0,1].$$
(8)

To show (8) let  $\alpha \in (0, 1]$  and  $\epsilon > 0$ , and we first compute a lower bound for  $G_{\alpha}(\theta)$  and for  $\mathcal{L}_{\alpha,\epsilon}(\theta)$ . Using Hölder inequality, for all  $\theta \in \mathbb{R}$  we have

$$\mathbb{E}[|\nabla f(\theta, Z)|] \le \mathbb{E}[|\nabla f(\theta, Z)|^{1+\alpha}]^{\frac{1}{1+\alpha}}$$

where

$$\mathbb{E}[|\nabla f(\theta, Z)|] = \mathbb{E}[|X \exp(\theta X) - XY|] \ge \mathbb{E}[X \exp(\theta X)] - \mathbb{E}[XY]$$
$$= \mathbb{E}[X \exp(\theta X)] - 1$$
$$= \exp(\exp(\theta) + \theta - 1)$$

and thus

$$G_{\alpha}(\theta) \ge \left(\exp\left(\exp(\theta) + \theta - 1\right) - 1\right)^{1+\alpha}, \quad \forall \theta \ge 1.$$
(9)

To proceed further let  $\theta \in \mathbb{R}$ . To obtain a lower bound for  $\mathcal{L}_{\alpha,\epsilon}(\theta)$  let  $r > \log 2$  and, using the fact that

$$\exp(\theta') \ge \exp(\theta) + (\theta' - \theta) \exp(\theta), \quad \forall \theta' \in \mathbb{R},$$

remark that

$$\exp\left(\exp(\theta) - \exp(\theta') + \theta - \theta'\right) \le \frac{1}{2}, \quad \forall \theta' \in (\theta + r, \infty).$$

Therefore, for all  $\theta' \in (\theta + r, \infty)$  we have, using (6),

$$\begin{split} \left|\nabla F(\theta) - \nabla F(\theta')\right| &= \left|\exp\left(\exp(\theta) + \theta - 1\right) - \exp\left(\exp(\theta') + \theta' - 1\right)\right| \\ &= \exp\left(\exp(\theta') + \theta' - 1\right)\left(1 - \exp\left(\exp(\theta) - \exp(\theta') + \theta - \theta'\right)\right) \\ &\geq \frac{1}{2}\exp\left(\exp(\theta') + \theta' - 1\right) \\ &= \frac{\exp\left(\exp(\theta') + \theta' - 1\right)}{2|\theta' - \theta|^{\alpha}}|\theta' - \theta|^{\alpha} \end{split}$$

and thus

$$\left|\nabla F(\theta) - \nabla F(\theta')\right| \ge \frac{\exp\left(\exp(\theta + r) + \theta + r - 1\right)}{2(2r)^{\alpha}} |\theta' - \theta|^{\alpha}, \ \forall \theta' \in \left(B_{2r}(\theta) \setminus B_r(\theta)\right) \cap (\theta, \infty).$$
(10)

This shows that if  $D_{r,\alpha}(\theta) \in (0,\infty)$  is such that

$$\left|\nabla F(\theta) - \nabla F(\theta')\right| \le D_{r,\alpha}(\theta)|\theta - \theta'|^{\alpha}, \quad \forall \theta' \in B_{2r}(\theta)$$

then

$$D_{r,\alpha}(\theta) \ge K_{r,\alpha}(\theta) := \frac{\exp\left(\exp(\theta + r) + \theta + r - 1\right)}{2(2r)^{\alpha}}$$

To proceed further let  $c_{\alpha,\epsilon} \in (1,\infty)$  be such that

$$\exp\left(\exp(\theta) + \theta - 1\right) - 1 \ge R(\theta) := \frac{1}{2}\exp\left(\exp(\theta) + \theta - 1\right) \ge \epsilon^{\frac{1}{1+\alpha}}, \quad \forall \theta \ge c_{\alpha,\epsilon}$$

and note that, by (9), we have  $r_{\alpha,\epsilon,\theta} \ge R(\theta)$  for all  $\theta \ge c_{\alpha,\epsilon}$ .

We now let  $c'_{\alpha,\epsilon} \in \mathbb{R}$  be such that  $c'_{\alpha,\epsilon} > c_{\alpha,\epsilon}$  and such that  $R(\theta) > 2$  for all  $\theta \ge c'_{\alpha,\epsilon}$ . Then,  $r_{\alpha,\epsilon,\theta} > 2 > \log 2$  for all  $\theta \ge c'_{\alpha,\epsilon}$  and thus, using the above computations,

$$\mathcal{L}_{\alpha,\epsilon}(\theta) \ge K_{\frac{1}{2}r_{\alpha,\epsilon,\theta},\alpha}(\theta), \quad \forall \theta \ge c'_{\alpha,\epsilon}$$
(11)

Therefore, using (9) and (11), and noting that for all  $\theta > 0$  we have  $K_{r',\alpha}(\theta) \ge K_{r,\alpha}(\theta)$  for all  $r' \ge r \ge 1$ , it follows that for all  $\theta \ge c'_{\alpha,\epsilon}$  we have

$$\mathcal{L}_{\alpha,\epsilon}(\theta)G_{\alpha}(\theta) \geq R(\theta)^{1+\alpha}K_{\frac{1}{2}r_{\alpha,\epsilon,\theta},\alpha}$$
  
$$\geq R(\theta)^{1+\alpha}K_{\frac{1}{2}R(\theta),\alpha}$$
  
$$= R(\theta)^{1+\alpha}\frac{\exp\left(\exp(\theta + R(\theta)/2) + \theta + R(\theta)/2 - 1\right)}{2R(\theta)^{\alpha}}$$
  
$$= \frac{R(\theta)}{2}\exp\left(\exp(\theta + R(\theta)/2) + \theta + R(\theta)/2 - 1\right)$$

and thus

$$\liminf_{\theta \to \infty} \frac{\mathcal{L}_{\alpha,\epsilon}(\theta) G_{\alpha}(\theta)}{F(\theta) + |\nabla F(\theta)|^2} = \infty$$

Since  $\alpha \in (0, 1]$  and  $\epsilon > 0$  are arbitrary, (8) follows and the proof of the proposition is complete.  $\Box$ 

# A.2 Proofs of the results of Section 1

### A.2.1 Proof of Proposition 3

Proof of Proposition 3. From (6) we observe that  $\nabla^2 F(\theta) > 0$  for all  $\theta \in \mathbb{R}$  and thus the function F satisfies A1. Remark also that, from (6), the unique minimizer of F is  $\theta_* = 0$ .

The function  $\theta \mapsto f(\theta, z)$  is twice continuously differentiable on  $\mathbb{R}$  for all  $z \in \mathbb{R}^2$  where, for all  $\theta \in \mathbb{R}$  and  $z \in \mathbb{R}^2$ ,

$$\nabla f(\theta, z) = -z_1 z_2 + z_1 \exp(\theta z_1), \quad \nabla^2 f(\theta, z) = z_1^2 \exp(\theta z_1).$$
(12)

Using (12) and the fact that  $\theta_{\star} = 0$ , we obtain

$$\mathbb{E}[(\nabla f(\theta_{\star}, Z))^{2}] = \mathbb{E}\left[\left(-XY + X\exp(\theta_{\star}X)\right)^{2}\right] = \mathbb{E}\left[X^{2}(1-Y)^{2}\right] = 2$$
(13)

and

$$\mathbb{E}[|\nabla^2 f(\theta_\star, Z)|] = \mathbb{E}[X^2 \exp(\theta_\star X)] = 2.$$
(14)

We now let  $\Theta \subset \mathbb{R}$  be a non-empty compact set,  $C_{\Theta} \in (0, \infty)$  be such that  $|\theta| \leq C_{\Theta}$  and such that  $|\theta - \theta'| \leq C_{\Theta}$  for all  $\theta, \theta' \in \Theta$ , and we let  $M_{\Theta} : \mathbb{R}^2 \to \mathbb{R}$  be defined by

$$M_{\Theta}(z) = |z_1^3| \exp(2C_{\Theta}|z_1|), \quad z \in \mathbb{R}^2$$

Then, using the fact that  $|e^x - 1| \leq |x|e^{|x|}$  for all  $x \in \mathbb{R}$ , for all  $(\theta, \theta') \in \Theta^2$  and  $z \in \mathbb{R}^2$  we have

$$\begin{aligned} \left| \nabla^2 f(\theta, z) - \nabla^2 f(\theta', z) \right| &= z_1^2 \left| \exp(\theta z_1) - \exp(\theta' z_1) \right| \\ &= z_1^2 \exp(\theta z_1) \left| 1 - \exp((\theta' - \theta) z_1) \right| \\ &\leq |z_1^3| \exp(\theta z_1) \left| \theta' - \theta \right| \exp(|\theta' - \theta| |z_1|) \\ &\leq M_{\Theta}(z) \left| \theta' - \theta \right| \end{aligned}$$

where, using Cauchy-Schwartz inequality,

$$\mathbb{E}[M_{\Theta}(Z)] = \mathbb{E}\left[X^3 \exp(2C_{\Theta}X)\right] \le \left(\mathbb{E}[X^6]\mathbb{E}\left[\exp(4C_{\Theta}X)\right]\right)^{1/2}$$
$$= \mathbb{E}[X^6]^{1/2} \exp\left(\exp(4C_{\Theta}) - 1\right)^{1/2}$$
$$< \infty.$$

Together with (13)-(14) this shows that A2 holds and the proof of Proposition 3 is complete.

### A.2.2 Proof of Lemma 1

Proof of Lemma 1. Let  $\Theta \subset \mathbb{R}^d$  be a (non-empty) closed ball containing  $\theta_{\star}$  and let  $(\theta, \theta') \in \Theta^2$ . Then, under A2, by Taylor's theorem there exists a  $\Theta$ -valued random variable  $\tilde{\theta}$  such that,  $\mathbb{P}$ -a.s.,

$$\begin{aligned} \left\|\nabla f(\theta, Z) - \nabla f(\theta', Z)\right\| &= \left\|\nabla^2 f(\bar{\theta}, Z)(\theta - \theta')\right\| \\ &\leq \left\|\theta - \theta'\right\| \left\|\nabla^2 f(\bar{\theta}, Z)\right\| \\ &\leq \left\|\theta - \theta'\right\| \left(\left\|\nabla^2 f(\bar{\theta}, Z) - \nabla^2 f(\theta_\star, Z)\right\| + \left\|\nabla^2 f(\theta_\star, Z)\right\|\right) \\ &\leq \left\|\theta - \theta'\right\| \left(\left\|\tilde{\theta} - \theta_\star\right\| M_{\Theta}(Z) + \left\|\nabla^2 f(\theta_\star, Z)\right\|\right) \\ &\leq \left\|\theta - \theta'\right\| M_{\Theta}^{(1)}(Z) \end{aligned}$$
(15)

where

$$M_{\Theta}^{(1)}(z) = \sup_{\theta_1, \theta_2 \in \Theta} \|\theta_1 - \theta_2\| M_{\Theta}(z) + \|\nabla^2 f(\theta_\star, z)\|, \quad \forall z \in \mathsf{Z}.$$

Under A2 we have both  $\mathbb{E}[M_{\Theta}(Z)] < \infty$  and  $\mathbb{E}[\|\nabla^2 f(\theta_{\star}, Z)\|] < \infty$ , and thus  $\mathbb{E}[M_{\Theta}^{(1)}(Z)] < \infty$ .

Similarly, under A2, by Taylor's theorem there exists a  $\Theta$ -valued random variable  $\tilde{\theta}$  such that,  $\mathbb{P} - a.s.$ ,

$$\begin{split} \left| f(\theta, Z) - f(\theta', Z) \right| &= \left| (\theta - \theta')^{\top} \nabla f(\tilde{\theta}, Z) \right| \\ &\leq \left\| \theta - \theta' \right\| \left\| \nabla f(\tilde{\theta}, Z) \right\| \\ &\leq \left\| \theta - \theta' \right\| \left( \left\| \nabla f(\tilde{\theta}, Z) - \nabla f(\theta_{\star}, Z) \right\| + \left\| \nabla f(\theta_{\star}, Z) \right\| \right) \\ &\leq \left\| \theta - \theta' \right\| \left( \left\| \tilde{\theta} - \theta_{\star} \right\| M_{\Theta}^{(1)}(Z) + \left\| \nabla f(\theta_{\star}, Z) \right\| \right) \\ &\leq \left\| \theta - \theta' \right\| M_{\Theta}^{(2)}(Z) \end{split}$$

where the third inequality uses (15) and where

$$M_{\Theta}^{(2)}(z) = \sup_{\theta_1, \theta_2 \in \Theta} \|\theta_1 - \theta_2\| M_{\Theta}^{(1)}(z) + \|\nabla f(\theta_\star, z)\|, \quad \forall z \in \mathsf{Z}.$$

Since, as shown above,  $\mathbb{E}[M_{\Theta}^{(1)}(Z)] < \infty$  while, by assumption,  $\mathbb{E}[\|\nabla f(\theta_{\star}, Z)\|] < \infty$ , it follow that  $\mathbb{E}[M_{\Theta}^{(2)}(Z)] < \infty$ .

The result of the lemma follows.

### A.2.3 Proof of Lemma 2

Proof of Lemma 2. Let  $\theta_{\star}$  be as in A2 and  $\Theta$  be a (non-empty) closed ball containing  $\theta_{\star}$ . Next, let  $\theta \in \Theta$ ,  $M'_{\Theta}(\cdot)$  be as in Lemma 1 and

$$G(z) = \sup_{\theta_1, \theta_2 \in \Theta} \|\theta_1 - \theta_2\| M'_{\Theta}(z) + \|\nabla f(\theta_\star, z)\| + \|\nabla^2 f(\theta_\star, z)\|, \quad \forall z \in \mathsf{Z}.$$

Then, for all  $k \in \{1, 2\}$  we  $\mathbb{P}$ -a.s. have, by Lemma 1,

$$\begin{split} \sup_{\theta \in \Theta} \|\nabla^k f(\theta, Z)\| &\leq \sup_{\theta \in \Theta} \|\nabla^k f(\theta, Z) - \nabla^k f(\theta_\star, Z)\| + \|\nabla^k f(\theta_\star, Z)\| \\ &\leq \sup_{\theta_1, \theta_2 \in \Theta} \|\theta_1 - \theta_2\| M'_{\Theta}(Z) + \|\nabla^k f(\theta_\star, Z)\| \\ &\leq G(Z). \end{split}$$

Since by Lemma 1 and under A2 we have  $\mathbb{E}[G(Z)] < \infty$ , it follows from the dominated convergence theorem that, for all  $k \in \{1, 2\}$ , we have  $\nabla^k F(\theta) = \mathbb{E}[\nabla^k f(\theta, Z)]$  for all  $\theta \in \mathring{\Theta}$  and thus, since  $\Theta$  is arbitrary,  $\nabla^k F(\theta) = \mathbb{E}[\nabla^k f(\theta, Z)]$  for all  $\theta \in \mathbb{R}^d$ .

Since F is twice differentiable on  $\mathbb{R}^d$ , the mapping  $\nabla F : \mathbb{R}^d \to \mathbb{R}^d$  is continuous and thus to conclude the proof of the lemma it remains to show that the mapping  $\nabla^2 F : \mathbb{R}^d \to \mathbb{R}^{d \times d}$  is continuous as well. To this aim, let  $\Theta$  be a (non-empty) closed ball containing  $\theta_{\star}$  and let  $\theta, \theta' \in \Theta$ . Then,

$$\begin{aligned} \left\| \nabla^2 F(\theta) - \nabla^2 F(\theta') \right\| &= \left\| \mathbb{E} \left[ \nabla^2 f(\theta, Z) \right] - \mathbb{E} \left[ \nabla^2 f(\theta', Z) \right] \right| \\ &\leq \mathbb{E} \left[ \left\| \nabla^2 f(\theta, Z) - \nabla^2 f(\theta', Z) \right\| \right] \\ &\leq \left\| \theta - \theta' \right\| \mathbb{E} [M_{\Theta}(Z)] \end{aligned}$$

where the first inequity holds by Jensen's inequality and the second one holds under A2. Since  $\mathbb{E}[M_{\Theta}(Z)] < \infty$ , this shows that the mapping  $\nabla^2 F : \mathbb{R}^d \to \mathbb{R}^{d \times d}$  is locally Lipschitz and thus continuous. The proof of the lemma is complete.

### A.2.4 Proof of Lemma 3

Proof of Lemma 3. Let  $k \in \{0, 1, 2\}$  and note that, using Lemma 1, we have  $\mathbb{E}[\|\nabla^k f(\theta, Z)\|] < \infty$  for all  $\theta \in \mathbb{R}^d$  under A2. Therefore, by the law of large numbers, for all  $\theta \in \mathbb{R}^d$  we have

$$\lim_{n \to \infty} \left\| \nabla^k F_n(\theta) - \mathbb{E} \big[ \nabla^k f(\theta, Z) \big] \right\| = 0, \quad \mathbb{P} - a.s.$$

To proceed further let  $\Theta \subset \mathbb{R}^d$  be a (non-empty) compact set. Then, by Lemma 1, there exists a measurable function  $M'_{\Theta}(\cdot)$  such that  $\mathbb{E}[M'_{\Theta}(Z)] < \infty$  and such that

$$\left\|\nabla^k f(\theta, Z) - \nabla^k f(\theta', Z)\right\| \le M'_{\Theta}(Z) \|\theta - \theta'\|, \quad \forall \theta, \theta' \in \Theta, \quad \mathbb{P}-a.s.$$

Therefore,  $\mathbb{P}$ -a.s., for all  $\theta, \theta' \in \mathbb{R}^d$  and  $n \ge 1$  we have

$$\|\nabla^{k}F_{n}(\theta) - \nabla^{k}F_{n}(\theta')\| = \frac{1}{n} \left\| \sum_{i=1}^{n} \left( \nabla^{k}f(\theta, Z_{i}) - \nabla^{k}f(\theta', Z_{i}) \right) \right\|$$
  
$$\leq \frac{1}{n} \sum_{i=1}^{n} \|\nabla^{k}f(\theta, Z_{i}) - \nabla^{k}f(\theta', Z_{i})\|$$
  
$$\leq \|\theta - \theta'\| \frac{1}{n} \sum_{i=1}^{n} M_{\Theta}'(Z_{i}).$$
 (16)

By the law of large numbers,  $\lim_{n\to\infty} \frac{1}{n} \sum_{i=1}^n M'_{\Theta}(Z_i) = \mathbb{E}[M'_{\Theta}(Z)] < \infty$ ,  $\mathbb{P}$ -.a.s, and thus

$$\frac{1}{n}\sum_{i=1}^{n}M'_{\Theta}(Z) = \mathcal{O}(1), \quad \mathbb{P}-a.s.$$
(17)

Using (16)-(17), as well as the fact that  $\Theta$  is compact and the fact that, by Lemma 2, the mapping  $\nabla^k F$  is uniformly continuous on the compact set  $\Theta$ , it is readily checked that

$$\lim_{n \to \infty} \sup_{\theta \in \Theta} \left\| \nabla^k F_n(\theta) - \mathbb{E} \left[ \nabla^k f(\theta, Z) \right] \right\| = 0.$$

The proof is complete.

### A.2.5 Proof of Lemma 4

Proof of Lemma 4. To simplify the notation in what follows for all  $\theta \in \mathbb{R}^d$  we let  $H(\theta) = \mathbb{E}[\nabla^2 f(\theta, Z)]$ and  $H_n(\theta) = \nabla^2 F_n(\theta)$  for all  $n \ge 1$ .

Let  $\Theta$  be a (non-empty) compact set,  $\theta \in \Theta$  and  $n \geq 1$ . Note that the matrices  $H_n(\theta)$  and  $H(\theta)$  are symmetric and thus, by the min-max theorem,

$$\lambda_{\min}(H_n(\theta)) = \min\left\{x^\top H_n(\theta)x : ||x|| = 1\right\},$$
  
$$\lambda_{\min}(H(\theta)) = \min\left\{x^\top H(\theta)x : ||x|| = 1\right\}.$$
 (18)

Since for all  $x \in \mathbb{R}^d$  such that ||x|| = 1 we have

$$x^{\top}H_{n}(\theta)x = x^{\top}H(\theta)x + x^{\top}(H_{n}(\theta) - H(\theta))x \ge \lambda_{\min}(H(\theta)) - \|H_{n}(\theta) - H(\theta)\|$$

it follows from Lemma 3 that

$$\liminf_{n \to \infty} \inf_{\theta \in \Theta} \lambda_{\min} (H_n(\theta)) \geq \inf_{\theta \in \Theta} \lambda_{\min} (H(\theta)) - \limsup_{n \to \infty} \sup_{\theta \in \Theta} \|H_n(\theta) - H(\theta)\| = \inf_{\theta \in \Theta} \lambda_{\min} (H(\theta)).$$

By Lemma 2, the mapping  $H : \mathbb{R}^d \to \mathbb{R}^{d \times d}$  is continuous and thus, using (18) and the maximum theorem, it follows that the mapping  $\theta \mapsto \lambda_{\min}(H(\theta))$  is continuous. Therefore, since  $\Theta$  is compact, there exists a  $\theta_{\Theta} \in \Theta$  such that

$$\inf_{\theta \in \Theta} \lambda_{\min} (H(\theta)) = \lambda_{\min} (H(\theta_{\Theta}))$$

which concludes the proof of the first part of the lemma.

Using the fact, by the min-max theorem,

$$\lambda_{\max}(H_n(\theta)) = \max\left\{x^\top H_n(\theta)x : \|x\| = 1\right\},\$$
$$\lambda_{\max}(H(\theta)) = \max\left\{x^\top H(\theta)x : \|x\| = 1\right\}$$

the second part of the lemma is proved in a similar way and its proof is therefore omitted to save space.  $\hfill \Box$ 

### A.2.6 Proof of Proposition 4

Proof of Proposition 4. To prove the proposition remark first since by assumption  $g_* > -\infty$  and g is convex then there exists an  $x_* \in \mathbb{R}^d$  such that  $g_* = g(x_*)$ .

Next, remark that since g is convex for all  $t \ge 0$  we have

$$g(x_{\star}) \ge g(x_t) + \nabla g(x_t)^{\top} (x_{\star} - x_t)$$

and thus

$$g(x_t) \le g(x_\star) - \nabla g(x_t)^\top (x_\star - x_t) = g(x_\star) + \nabla g(x_t)^\top (x_t - x_\star), \quad \forall t \ge 1.$$
(19)

In addition, remark that

$$\|x_{t+1} - x_{\star}\|^{2} = \|x_{t} - x_{\star}\|^{2} + v_{t}^{2} \|\nabla g(x_{t})\|^{2} - 2v_{t} \nabla g(x_{t})^{\top} (x_{t} - x_{\star}), \quad \forall t \ge 0.$$
(20)

Let  $t \ge 0$  and note that

$$g(x_{t+1}) - g(x_{\star}) \leq g(x_{t}) - g(x_{\star}) - \frac{v_{t}}{2} \|g(x_{t})\|^{2}$$

$$\leq \nabla g(x_{t})^{\top} (x_{t} - x_{\star}) - \frac{v_{t}}{2} \|g(x_{t})\|^{2}$$

$$= \frac{1}{2v_{t}} \left( 2v_{t} \nabla g(x_{t})^{\top} (x_{t} - x_{\star}) - v_{t}^{2} \|g(x_{t})\|^{2} \right)$$

$$= \frac{1}{2v_{t}} \left( 2v_{t} \nabla g(x_{t})^{\top} (x_{t} - x_{\star}) - v_{t}^{2} \|g(x_{t})\|^{2} - \|x_{t} - x_{\star}\|^{2} + \|x_{t} - x_{\star}\|^{2} \right)$$

$$= \frac{1}{2v_{t}} \left( \|x_{t} - x_{\star}\|^{2} - \|x_{t+1} - x_{\star}\|^{2} \right)$$

$$(21)$$

where the first inequality uses (2), the second inequality uses (19) and the first equality uses (20).

In addition, since  $g(x_{t+1}) - g(x_{\star}) \ge 0$  for all  $t \ge 0$ , it follows from (21) that

$$\|x_t - x_\star\|^2 - \|x_{t+1} - x_\star\|^2 \ge 0, \quad \forall t \ge 1$$
(22)

and thus, since for some constant c > 0 we have  $\inf_{t \ge 1} v_t \ge c > 0$  by assumption, it follows from (21)-(22) that

$$g(x_{t+1}) - g(x_{\star}) \le \frac{1}{2c} \Big( \|x_t - x_{\star}\|^2 - \|x_{t+1} - x_{\star}\|^2 \Big), \quad \forall t \ge 0.$$

To conclude the proof let  $T \in \mathbb{N}$ . Then, using this latter result, we have

$$\sum_{t=1}^{T} \left( g(x_t) - g(x_\star) \right) \le \frac{1}{2c} \sum_{t=1}^{T} \left( \|x_{t-1} - x_\star\|^2 - \|x_t - x_\star\|^2 \right)$$
$$= \frac{1}{2c} \left( \|x_0 - x_\star\|^2 - \|x_T - x_\star\|^2 \right)$$
$$\le \frac{\|x_0 - x_\star\|^2}{2c}$$

and thus, since by (2) we have  $g(x_t) - g(x_*) \leq g(x_{t-1}) - g(x_*)$  for all  $t \geq 1$ , it follows that

$$g(x_T) - g(x_\star) \le \frac{1}{T} \sum_{t=1}^T \left( g(x_t) - g(x_\star) \right) \le \frac{\|x_0 - x_\star\|^2}{2cT}$$

and the proof is complete.

### A.2.7 Proof of Proposition 5

Proof of Proposition 5. Let K and  $\tilde{K}$  be as in the statement of the proposition.

Remark first that, since g is  $L_K$ -smooth on the convex set K, we have

$$g(y) \le g(x) + \nabla g(x)^{\top} (y - x) + \frac{L_K}{2} ||y - x||^2, \quad \forall x, y \in K$$
(23)

and remark that  $x + v\nabla g(x) \in K$  for all  $x \in \tilde{K}$  and  $v \in (0, 1]$ . We now let  $x \in \tilde{K} \subset K$  and  $v \in (0, 1]$ . Then, since by (23) we have

$$g(x - v\nabla g(x)) \le g(x) - v\left(1 - \frac{L_K}{2}v\right) \|\nabla g(x)\|^2$$

where

$$v\left(1-\frac{L_K}{2}v\right) \ge \frac{v}{2} \Leftrightarrow v \le \frac{1}{L_K},$$

it follows that if  $v = \beta^{k_x}$  with

$$k_x = \min\left\{k \in \mathbb{N}_0 : g\left(x - \beta^k \|\nabla g(x)\|\right) \le g(x) - \frac{\beta^k}{2} \|\nabla g(x)\|^2\right\}$$

then  $v \ge \beta/L_K$ . Since we trivially have  $x_t \in \tilde{K}$  for all  $t \ge 0$ , this shows that  $v_t \ge \beta/L_K$  for all  $t \ge 1$  and the proof of the proposition is complete.

#### A.2.8 Proof of Lemma 5

Proof of Lemma 5. Remark first that under the assumptions of the lemma we have  $\mathbb{E}[\nabla f(\theta_{\star}, Z)] = 0$ by Lemma 2. Hence, since  $\mathbb{E}[\|\nabla f(\theta_{\star}, Z)\|^{1+\alpha}] < \infty$  by assumption, it follows from Bubeck et al. (2013, Lemma 3) that there exists a constant  $C^* < \infty$  such that

$$\mathbb{P}(\|\nabla F_n(\theta_\star)\| \ge M) \le \frac{C^*}{n^{\alpha} M^{1+\alpha}}, \quad \forall M > 0, \quad \forall n \ge 1.$$
(24)

Remark also that, since  $\sup_{\theta \in \Theta} ||F_n(\theta) - F(\theta)|| = o_{\mathbb{P}}(1)$  by Lemma 3 while, under A1, there exists a constant  $c_{\Theta} > 0$  such that  $F(\theta) - F(\theta_{\star}) \ge c_{\Theta} ||\theta - \theta_{\star}||$  for all  $\theta \in \Theta$ , it follows from Van der Vaart (2000, Theorem 5.7, page 45) that

$$\|\theta_{\star} - \hat{\theta}_{\Theta,n}\| = o_{\mathbb{P}}(1). \tag{25}$$

We now let

$$\lambda_{\Theta} = \frac{1}{2} \inf_{\theta \in \Theta} \lambda_{\min} \big( \mathbb{E} \big[ \nabla^2 f(\theta, Z) \big] \big)$$

and, to simplify the notation in what follows, for all  $\theta \in \mathbb{R}^d$  and  $n \geq 1$  we let  $H_n(\theta) = \nabla^2 F_n(\theta)$ . Finally, for all  $\epsilon > 0$  and  $n \geq 1$  we let

$$\Omega_{\epsilon,n} = \Big\{ \omega \in \Omega : \ \hat{\theta}_{\Theta,n}^{\omega} \in \mathring{\Theta}, \ \left\| \nabla F_n^{\omega}(\theta_{\star}) \right\| \le n^{-\frac{\alpha}{1+\alpha}} (C^*/\epsilon)^{\frac{1}{1+\alpha}}, \ \inf_{\theta \in \Theta} \lambda_{\min} \big( H_n^{\omega}(\theta) \big) \ge \lambda_{\Theta} \Big\}.$$

Remark that, by (24)-(25) and Lemma 4, and since  $\theta_{\star} \in \mathring{\Theta}$ , we have  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,n}) \geq 1-\epsilon$  for all  $\epsilon \in (0,1)$ . Remark also that  $\lambda_{\Theta} > 0$  under A1 and by Lemma 2, and thus for all  $n \geq 1$  and  $\epsilon > 0$  the function  $F_n^{\omega}$  is strictly convex on  $\Theta$  for all  $\omega \in \Omega_{\epsilon,n}$ .

Let  $n \geq 1$ ,  $\epsilon \in (0, 1)$  and  $\omega \in \Omega_{\epsilon,n}$ , and note that  $\nabla F_n^{\omega}(\hat{\theta}_{\Theta,n}^{\omega}) = 0$  since  $F_n^{\omega}$  is strictly convex on  $\Theta$ and  $\hat{\theta}_{\Theta,n}^{\omega} \in \mathring{\Theta}$ . Therefore, since  $\theta_{\star} \in \Theta$ , it follows from Taylor's theorem that there exists a  $\check{\theta}_{n_B}^{\omega} \in \mathring{\Theta}$  such that

$$\nabla F_n^{\omega}(\theta_{\star}) = H_n^{\omega}(\check{\theta}_n^{\omega})(\theta_{\star} - \hat{\theta}_{\Theta,n}^{\omega}) \Leftrightarrow \theta_{\star} - \hat{\theta}_{\Theta,n}^{\omega} = \left(H_n^{\omega}(\check{\theta}_n^{\omega})\right)^{-1} \nabla F_n^{\omega}(\theta_{\star}).$$
(26)

Remark that the matrix  $H_n^{\omega}(\check{\theta}_n^{\omega})$  is indeed invertible since all its eigenvalues are bounded below by  $\lambda_{\Theta}$ , where  $\lambda_{\Theta} > 0$  under A1 and by Lemma 2.

Using (26), we obtain that

$$\|\theta_{\star} - \hat{\theta}_{\Theta,n}^{\omega}\| \le \|H_n^{\omega}(\check{\theta}_n^{\omega})\|^{-1} \|\nabla F_n^{\omega}(\theta_{\star})\| \le \left(\frac{C_{\star}}{\epsilon}\right)^{\frac{1}{1+\alpha}} \lambda_{\Theta}^{-1} n^{-\frac{\alpha}{1+\alpha}}$$
(27)

and the result of the lemma follows.

### A.3 Preliminary results for proving Theorems 1-2

### A.3.1 A technical lemma

**Lemma 6.** Let  $\alpha' \in (0,1]$ ,  $\delta \in (0, 2\alpha'/(1+3\alpha'))$  and, for all  $j \in \mathbb{N}$ , let  $\gamma_j = 1 - \delta^j$ . Next, let  $w_{\alpha'} = (1+\alpha')/(1+3\alpha')$  and

$$\gamma'_j = w_{\alpha'} + (1 - w_{\alpha'})\gamma_{j-1}, \quad \forall j \in \mathbb{N} \setminus \{1\}.$$

Then, there exists a constant  $\pi \in (0,1)$  such that

$$\gamma_j = \pi \gamma'_j + (1 - \pi) \gamma_{j-1}, \quad \forall j \in \mathbb{N} \setminus \{1\}.$$

In addition, if for some constants  $(\kappa, \tau) \in (0, \infty)^2$  and all  $j \in \mathbb{N}$  and  $B \in \mathbb{N}$  we let  $n_{j,B} = \lceil \kappa B^{\gamma_j} \rceil$ ,  $\tau_{j,B} = \tau B^{-\frac{\alpha'}{1+\alpha'}\gamma_j}$  and  $r_{j,B} = B^{-\frac{\alpha'}{1+\alpha'}\gamma_j}$ , then

$$\lim_{B \to \infty} \frac{n_{j,B} r_{j-1,B}^2}{\tau_{j,B}^2 B} = \lim_{B \to \infty} \frac{n_{j,B}}{B} = 0, \quad \forall j \in \mathbb{N}$$

with the convention that  $r_{j-1,B} = 1$  for all  $B \in \mathbb{N}$  when j = 1.

Proof of Lemma 6. We show that the first part of the lemma holds for

$$\pi = (1 - \delta) \frac{1 + 3\alpha'}{1 + \alpha'}.$$

Remark that the conditions on  $\delta$  ensure that  $\pi \in (0,1)$  and that, by definition,  $\pi$  is such that

$$\frac{2\alpha' + (1-\pi)(1+\alpha')}{1+3\alpha'} = \delta.$$
 (28)

Then, the first part of the lemma holds if and only if for all  $j \ge 2$  we have

$$\gamma_{j} = \pi w_{\alpha'} + \pi (1 - w_{\alpha'})\gamma_{j-1} + (1 - \pi)\gamma_{j-1}$$

$$= \pi w_{\alpha'} + (1 - \pi w_{\alpha'})\gamma_{j-1}$$

$$= \pi w_{\alpha'} \sum_{i=0}^{j-2} (1 - \pi w_{\alpha'})^{i} + (1 - \pi w_{\alpha'})^{j-1}\gamma_{1}$$

$$= 1 - (1 - \pi w_{\alpha'})^{j-1} (1 - \gamma_{1})$$

$$= 1 - \left(\frac{2\alpha' + (1 - \pi)(1 + \alpha')}{1 + 3\alpha'}\right)^{j-1} (1 - \gamma_{1})$$

$$= 1 - \delta^{j-1} (1 - \gamma_{1})$$

$$= 1 - \delta^{j}$$
(29)

where the penultimate equality uses (28) and the last one the fact that  $\gamma_1 = 1 - \delta$ . This shows the first part of the lemma.

To show the second part of the lemma note that, by the first part of the lemma, we have

$$\gamma_j < \gamma_{j+1} < \gamma'_{j+1}, \quad \forall j \in \mathbb{N}$$

$$\tag{30}$$

and to simplify the notation in the following let  $c_{\alpha'} = \alpha'/(1+\alpha')$ .

Firstly, the conditions on  $\delta$  ensuring that  $\gamma_1(1+2c_{\alpha'})-1<0$ , it follows that

$$\lim_{B \to \infty} \frac{n_{1,B}}{\tau_{1,B}^2 B} = 0.$$
(31)

Next, using the first part of the lemma, we remark that for all  $j \in \mathbb{N} \setminus \{1\}$  we have

$$(1+2c_{\alpha'})\gamma_j - 2c_{\alpha'}\gamma_{j-1} - 1 = (1+2c_{\alpha'})\gamma'_j - 2c_{\alpha'}\gamma_{j-1} - 1 + (1+2c_{\alpha'})(1-\pi)(\gamma_{j-1} - \gamma'_j)$$
  
=  $(1+2c_{\alpha'})(1-\pi)(\gamma_{j-1} - \gamma'_j)$   
<  $0$ 

where the last inequality uses (30), and thus

$$\lim_{B \to \infty} \frac{n_{j+1,B} r_{j,B}^2}{\tau_{j+1,B}^2 B} = 0, \quad \forall j \in \mathbb{N}.$$
(32)

By combining (31) and (32) we obtain that

$$\lim_{B \to \infty} \frac{n_{j,B} r_{j-1,B}^2}{\tau_{j,B}^2 B} = 0, \quad \forall j \in \mathbb{N}.$$
(33)

Finally, since  $\gamma_j < 1$  for all  $j \in \mathbb{N}$ , we have

$$\lim_{B \to \infty} \frac{n_{j,B}}{B} = 0, \quad \forall j \in \mathbb{N}$$
(34)

and the second part of the lemma follows from (33)-(34). The proof is complete.

#### A.3.2 A first key property of Algorithm 3

**Lemma 7.** Assume that A2 holds. Let  $\beta \in (0,1)$ ,  $(n_B)_{B\geq 1}$  be a sequence in  $\mathbb{N}$ ,  $(\tau_B)_{B\geq 1}$  be a sequence in  $[0,\infty)$  and  $(\tilde{B}_B)_{B\geq 1}$  be a sequence of  $\mathbb{N}_0$ -valued random variables. In addition, for all  $B \in \mathbb{N}$  and  $\theta_0 \in \mathbb{R}^d$ , let

$$(\hat{\theta}_{\theta_0,B}, \tilde{B}_{\theta_0}) = \text{Algorithm } \mathbf{3}(\theta_0, n_B, \tilde{B}_B, \tau_B, \beta)$$

and denote by  $\{\theta_{\theta_0,B,t}\}_{t=0}^{T_{\theta_0,B}}$  the sequence in  $\mathbb{R}^d$  generated by Algorithm 3 to compute  $\hat{\theta}_{\theta_0,B}$  from  $\theta_0$ and (which is therefore such that  $\theta_{\theta_0,B,0} = \theta_0$  and such that  $\theta_{\theta_0,B,T_{\theta_0,B}} = \hat{\theta}_{\theta_0,B}$ ). Finally, let  $x \in \mathbb{R}^d$ and, for all  $j \in \mathbb{N}$ , let

$$\Theta_{j,x} = \left\{ \theta \in \mathbb{R}^d : F(\theta) \le F(x) + j \right\}.$$

Then, for all  $j \in \mathbb{N}$ , there exists a sequence  $(\Omega_{j,B}^{(1)})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\lim_{B\to\infty} \mathbb{P}(\Omega_{j,B}^{(1)}) = 1$  and such that

$$\theta^{\omega}_{\theta_0,B,t} \in \Theta_{j+1,x}, \quad \forall t \in \{0,\dots,T^{\omega}_{\theta_0,B}\}, \quad \forall \omega \in \Omega^{(1)}_{j,B}, \quad \forall \theta_0 \in \Theta_{j,x}, \quad \forall B \in \mathbb{N}$$

Proof of Lemma 7. Let  $j \in \mathbb{N}$ ,

$$\widetilde{\Theta}_{j,x} = \left\{ \theta \in \mathbb{R}^d : \exists \theta' \in \Theta_{j+1,x} \text{ such that } \|\theta - \theta'\| \le \sup_{\widetilde{\theta} \in \Theta_{j+1,x}} \|\nabla F(\widetilde{\theta})\| + 1 \right\}$$

and, for all  $B \in \mathbb{N}$ , let

$$\Omega_{j,B}^{(1)} = \left\{ \omega \in \Omega :, \max_{k \in \{0,1\}} \sup_{\theta \in \tilde{\Theta}_{j,x}} \|\nabla^k F_{n_B}^{\omega}(\theta) - \nabla^k F(\theta)\| \le 1/2 \right\}$$

By Lemma 2, the function F is continuous on  $\mathbb{R}^d$  and thus the set  $\Theta_{j+1,x}$  is compact. Therefore, using again Lemma 2, we have  $\sup_{\tilde{\theta}\in\Theta_{j+1,x}} \|\nabla F(\tilde{\theta})\| < \infty$  and thus the set  $\Theta_{j,x}$  is also compact. Consequently, by Lemma 3, we have  $\lim_{B\to\infty} \mathbb{P}(\Omega_{j,B}^{(1)}) = 1$ .

To proceed further we let  $B \in \mathbb{N}$ ,  $\omega \in \Omega_{j,B}^{(1)}$  and  $\theta_0 \in \Theta_{j,x}$ , and we show that  $\theta_{\theta_0,B,t}^{\omega} \in \Theta_{j+1,x}$  for all  $t \in \{0, \ldots, T_{\theta_0,B}^{\omega}\}$ . We assume below that  $T_{\theta_0,B}^{\omega} > 0$  since otherwise the result is trivial (since by  $\theta_0 \in \Theta_{j,x} \subset \Theta_{j+1,x}$ ).

To do so let  $\theta \in \widetilde{\Theta}_{j,x}$  be such that  $F_{n_B}^{\omega}(\theta) \leq F_{n_B}^{\omega}(\theta_0)$  and note that

$$F(\theta) - 1/2 \le F_{n_B}^{\omega}(\theta) \le F_{n_B}^{\omega}(\theta_0) \le F(\theta_0) + 1/2 \le F(x) + j + 1/2$$

showing that  $F(\theta) \leq F(x) + j + 1$  and thus that

$$\{\theta \in \widetilde{\Theta}_{j,x}: F_{n_B}^{\omega}(\theta) \le F_{n_B}^{\omega}(\theta_0)\} \subset \Theta_{j+1,x}.$$
(35)

We now let  $t \in \{0, \ldots, T^{\omega}_{\theta_0, B} - 1\}$  and remark that Algorithm 3 ensures that we have

$$\|\theta_{\theta_0,B,t+1}^{\omega} - \theta_{\theta_0,B,t}^{\omega}\| \le \|\nabla F_{n_B}^{\omega}(\theta_{\theta_0,B,t}^{\omega})\|.$$

Hence, if  $\theta_{\theta_0,B,t}^{\omega} \in \Theta_{j+1,x} \subset \tilde{\Theta}_{j,x}$ , we have

$$\|\theta_{\theta_0,B,t+1}^{\omega} - \theta_{\theta_0,B,t}^{\omega}\| \le \|\nabla F(\theta_{\theta_0,B,t}^{\omega})\| + 1/2 \le \sup_{\tilde{\theta} \in \Theta_{j+1,x}} \|\nabla F(\tilde{\theta})\| + 1$$

showing that  $\theta_{\theta_0,B,t+1}^{\omega} \in \widetilde{\Theta}_{j,x}$ . Therefore, since Algorithm 3 ensures that we have

$$F_{n_B}^{\omega}(\theta_{\theta_0,B,t+1}^{\omega}) \le F_{n_B}^{\omega}(\theta_{\theta_0,B,t}^{\omega}) \le F_{n_B}^{\omega}(\theta_0)$$

it follows from (35) that  $\theta_{\theta_0,B,t+1}^{\omega} \in \Theta_{j+1,x}$  if  $\theta_{\theta_0,B,t}^{\omega} \in \Theta_{j+1,x}$ . Consequently, since  $\theta_0 \in \Theta_{j,x} \subset \Theta_{j+1,x}$ , it follows that  $\theta_{\theta_0,B,t}^{\omega} \in \Theta_{j+1,x}$  for all  $t \in \{0, \ldots, T_{\theta_0,B}^{\omega}\}$  and the proof of the lemma is complete.  $\Box$ 

### A.3.3 A second key property of Algorithm 3

**Lemma 8.** Consider the set-up of Lemma 7. In addition, assume that A1 holds and for all  $B \in \mathbb{N}$ and  $t \in \{1, \ldots, T_{\theta_0, B}\}$  let  $C_{\theta_0, B, t}$  denotes the computational cost needed for computing  $\theta_{\theta_0, B, t}^{\omega}$  from  $\theta_{\theta_0, B, t-1}^{\omega}$ , with the convention that  $C_{\theta_0, B, t}^{\omega} = 0$  when  $T_{\theta_0, B}^{\omega} = 0$ . Then, for all  $j \in \mathbb{N}$ , there exist (i) a compact and convex set  $K_{j+1,x} \subset \mathbb{R}^d$  such that  $\Theta_{j+1,x} \subset \mathring{K}_{j+1,x}$ , (ii) a constant  $D_j \in (0, \infty)$  and (iii) a sequence  $(\Omega_{j,B}^{(2)})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\lim_{B\to\infty} \mathbb{P}(\Omega_{j,B}^{(2)}) = 1$ , such that we have, for all  $B \in \mathbb{N}$ ,  $\theta_0 \in \Theta_{j,x}$ and  $\omega \in \Omega_{j,B}^{(2)}$ ,

$$F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F_{n_B}^{\omega}(\hat{\theta}_{K_{j+1,x},n_B}^{\omega}) \le \frac{D_j \|\theta_0 - \hat{\theta}_{K_{j+1,x},n_B}\|^2}{T_{\theta_0,B}^{\omega}}, \quad C_{\theta_0,B,t}^{\omega} \le D_j n_B$$

Proof of Lemma 8. Let  $j \in \mathbb{N}$ ,  $\Theta'_{j+1,x}$  be a compact and convex set containing  $\mathring{\Theta}_{j+1,x}$  and let  $\widetilde{\Theta}_{j+1,x}$  be a compact and convex set such that

$$\left\{\theta \in \mathbb{R}^d : \exists \theta' \in \Theta'_{j+1,x} \text{ such that } \|\theta - \theta'\| \le \sup_{\tilde{\theta} \in \Theta'_{j+1,x}} \|\nabla F(\tilde{\theta})\| + 1\right\} \subset \widetilde{\Theta}_{j+1,x}.$$

Remark that such a compact set  $\Theta_{j+1,x}$  indeed exists since  $F \in \mathcal{C}^2(\mathbb{R}^2)$  by Lemma 2.

Then, for all  $B \in \mathbb{N}$  we let

$$\begin{split} \Omega_{j,B}^{(2)} &= \Omega_{j,B}^{(1)} \cap \left\{ \omega \in \Omega : F_{n_B}^{\omega} \text{ is strictly convex on } \widetilde{\Theta}_{j+1,x}, \\ \| \nabla F_{n_B}^{\omega}(\theta) - \nabla F_{n_B}^{\omega}(\theta') \| \leq 2\mathbb{E}[M'_{\widetilde{\Theta}_{j+1,x}}(Z)] \| \theta - \theta' \| \text{ for all } (\theta, \theta') \in \widetilde{\Theta}_{j+1,x}^2 \right\} \end{split}$$

with  $\Omega_{j,B}^{(1)}$  as in the statement of Lemma 7 and with the function  $M'_{\tilde{\Theta}_{j+1,x}}(\cdot)$  as in Lemma 1 (when  $\Theta = \tilde{\Theta}_{j+1,x}$ ). Remark that, by Lemma 1, Corollary 1 and Lemma 7, we have  $\lim_{B\to\infty} \mathbb{P}(\Omega_{j,B}^{(2)}) = 1$ .

To prove the result of the lemma let  $L_{j+1,x} = (2\mathbb{E}[M'_{\tilde{\Theta}_{j+1,x}}(Z)] \vee 1), \theta_0 \in \Theta_{j,x}, B \in \mathbb{N}$  and  $\omega \in \Omega_{j,B}^{(2)}$ . Remark that the function  $F_{n_B}^{\omega}$  is strictly convex and  $L_{j+1,x}$ -smooth on the set  $\tilde{\Theta}_{j+1,x}$ , and that since  $\omega \in \Omega_{j,B}^{(1)}$  we have  $\theta_{\theta_0,B,t}^{\omega} \in \Theta'_{j+1,x}$  for all  $t \in \{0, \ldots, T_{\theta_0,B}^{\omega}\}$ . Next, assume first that  $T_{\theta_0,B}^{\omega} > 0$  and for all  $t \in \{0, \ldots, T_{\theta_0,B}^{\omega} - 1\}$  let  $v_t^{\omega}$  be such that  $\theta_{\theta_0,B,t+1}^{\omega} = \theta_{\theta_0,B,t}^{\omega} + v_t^{\omega} \nabla F_{n_B}(\theta_{\theta_0,B,t}^{\omega})$ . Then, by using the computations done to proof Proposition 5 with  $g = F_{n_B}^{\omega}, K = \tilde{\Theta}_{j+1,x}, \tilde{K} = \Theta'_{j+1,x}$  and  $L_K = L_{j+1,x}$ , it follows that  $v_t^{\omega} \ge \beta/L_{j+1}$  for all  $t \in \{0, \ldots, T_{\theta_0,B}^{\omega} - 1\}$  and thus that  $C_{\theta_0,B,t}^{\omega} \le D_j n_B$ , with  $D_j = \beta^{k_j} + 1$  where  $k_j = \inf\{k \in \mathbb{N} : \beta^k < \beta/L_{j+1}\}$ . In addition, by Proposition 4 we have

$$F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F_{n_B}^{\omega}(\hat{\theta}_{K_{j+1,x},n_B}^{\omega}) \le \frac{\beta \|\theta_0 - \hat{\theta}_{K_{j+1,x},n_B}\|^2}{2L_{j+1}T_{\theta_0,B}^{\omega}}$$

Noting that this latter inequality also holds if  $T^{\omega}_{\theta_0,B}$ , the proof of the proposition is complete.

#### A.3.4 A third key property of Algorithm 3

**Lemma 9.** Consider the set-up of Lemma 8. Assume in addition that there exists a constant  $c \in (0,1)$  such that  $\mathbb{P}(\tilde{B}_B \ge cB) = 1$  for all  $B \in \mathbb{N}$  and that

$$\lim_{B \to \infty} \frac{n_B}{B} = 0, \quad \lim_{B \to \infty} \tau_B = 0.$$

Then, for all  $j \in \mathbb{N}$ , there exist a constant  $D'_j \in (0,\infty)$  and a sequence  $(\Omega_{j,B}^{(3)})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\lim_{B\to\infty} \mathbb{P}(\Omega_{j,B}^{(3)}) = 1$  and such that

$$\|\hat{\theta}_{\theta_0,B}^{\omega} - \hat{\theta}_{K_{j+1,x},n_B}^{\omega}\| \le D_j' \frac{\|\theta_0 - \hat{\theta}_{K_{j+1,x},n_B}^{\omega}\|}{\sqrt{T_{\theta_0,B}^{\omega}}}, \quad \forall \omega \in \Omega_B^{(3)}, \quad \forall \theta_0 \in \Theta_{j,x}, \quad \forall B \in \mathbb{N}.$$

Proof of Lemma 9. Let  $j \in \mathbb{N}$  and, to simplify the notation in what follows, let  $K_x = K_{j+1,x}$  and, for all  $\theta \in \mathbb{R}^d$  and  $n \ge 1$ , let  $H_n(\theta) = \nabla^2 F_n(\theta)$ .

We start by showing that

$$\sup_{\theta_0 \in \Theta_{j,x}} \|\hat{\theta}_{\theta_0,B} - \hat{\theta}_{K_x,n_B}\| = o_{\mathbb{P}}(1).$$
(36)

To this aim let  $M'_{K_x}(\cdot)$  be as defined in Lemma 1 when  $\Theta = K_x$ ,

$$\lambda_{K_x} = \frac{1}{2} \inf_{\theta \in K_x} \lambda_{\min} \left( \mathbb{E} \left[ \nabla^2 f(\theta, Z) \right] \right)$$

and, for all  $\epsilon > 0$  and  $B \in \mathbb{N}$ , let

$$\begin{split} \Omega_{\epsilon,B} &= \Omega_B^{(1)} \cap \Omega_B^{(2)} \\ &\cap \left\{ \omega \in \Omega : \ \tilde{B}_B^{\omega} \ge cB, \ \inf_{\theta \in K_x} \lambda_{\min} \left( H_{n_B}^{\omega}(\theta) \right) \ge \lambda_{K_x}, \ \|\hat{\theta}_{K_x,n_B}^{\omega} - \theta_\star\| \le \epsilon, \ \hat{\theta}_{K_x,n_B}^{\omega} \in \mathring{K}_x, \\ &\sup_{\theta \in K_x} |F_{n_B}^{\omega}(\theta) - F(\theta)| \le \epsilon, \ \frac{1}{n_B} \sum_{i=1}^{n_B} M'_{K_x}(Z^{\omega}) \le 2\mathbb{E}[M'_{K_x}(Z)] \right\} \end{split}$$

with  $\Omega_B^{(1)}$  as in Lemma 7 and  $\Omega_B^{(2)}$  as in Lemma 8. Remark that  $\theta_{\star} \in \mathring{K}_x$  and thus, by Lemmas 1, 3, 4, 5, 7 and 8, we have  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,B}) = 1$  for all  $\epsilon > 0$ .

We now let  $\theta_0 \in \Theta_{j,x}$ ,  $\epsilon > 0$ ,  $B \in \mathbb{N}$  and  $\omega \in \Omega_{\epsilon,B}$ , and assume first that  $\|\nabla F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})\| \leq \tau_B$ . By Lemma 2 and under A1 we have  $\lambda_{K_x} > 0$ , implying that the function  $F_{n_B}^{\omega}$  is strictly convex on  $K_x$  and thus, since  $\hat{\theta}_{K_x,n_B}^{\omega} \in \mathring{K}_x$ , we have  $\nabla F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) = 0$ . Therefore, using Taylor's theorem, the fact that  $\|Ax\| \geq \lambda_{\min}(A)\|x\|$  for any  $d \times d$  symmetric matrix A and  $x \in \mathbb{R}^d$ , and noting that  $\hat{\theta}_{\theta_0,B}^{\omega} \in \Theta_{j+1,x} \subset K_x$ , we have

$$\tau_B \ge \|\nabla F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})\| \ge \lambda_{K_x} \|\hat{\theta}_{\theta_0,B}^{\omega} - \hat{\theta}_{K_x,n_B}\|$$

showing that

$$\|\hat{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\| \leq \frac{\tau_{B}}{\lambda_{K_{x}}}, \quad \text{if } \|\nabla F^{\omega}_{n_{B}}(\hat{\theta}^{\omega}_{\theta_{0},B})\| \leq \tau_{B}.$$

$$(37)$$

Assume now that  $\|\nabla F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})\| > \tau_B$ , in which case

$$T^{\omega}_{\theta_0,B} \ge \frac{c\,B}{D_j n_B} \tag{38}$$

with  $D_j < \infty$  as in Lemma 8. Under A1 there exists a constant  $\mu_x > 0$  such that

$$|F(\theta) - F(\theta_{\star})| \ge \frac{\mu_x}{2} \|\theta - \theta_{\star}\|^2, \quad \forall \theta \in K_x$$

and thus

$$\frac{\mu_x}{2} \|\hat{\theta}^{\omega}_{\theta_0,B} - \theta_\star\|^2 \le \left| F(\hat{\theta}^{\omega}_{\theta_0,B}) - F(\theta_\star) \right| \le \left| F(\hat{\theta}^{\omega}_{\theta_0,B}) - F^{\omega}_{n_B}(\hat{\theta}^{\omega}_{\theta_0,B}) \right| + \left| F^{\omega}_{n_B}(\hat{\theta}^{\omega}_{\theta_0,B}) - F(\theta_\star) \right|$$

$$\le \epsilon + \left| F^{\omega}_{n_B}(\hat{\theta}^{\omega}_{\theta_0,B}) - F(\theta_\star) \right|.$$
(39)

Noting that  $\theta_{\star} \in K_x$ , we have

$$\begin{aligned} \left| F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F(\theta_{\star}) \right| &\leq \left| F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) \right| + \left| F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) - F_{n_B}^{\omega}(\theta_{\star}) \right| \\ &+ \left| F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) \right| + \left( \frac{1}{n_B} \sum_{i=1}^{n_B} M'_{K_x}(Z_i^{\omega}) \right) \right\| \hat{\theta}_{K_x,n_B}^{\omega} - \theta_{\star} \| \\ &+ \sup_{\theta \in K_x} \left| F_{n_B}^{\omega}(\theta) - F(\theta) \right| \\ &\leq \left| F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) \right| + \epsilon \left( 2\mathbb{E}[M'_{K_x}(Z)] + 1 \right) \\ &\leq \frac{D_j \sup_{(\theta,\theta') \in K_x^2} \|\theta - \theta'\|^2}{T_{\theta_0,B}^{\omega}} + \epsilon \left( 2\mathbb{E}[M'_{K_x}(Z)] + 1 \right) \\ &\leq \frac{n_B D_j^2 \sup_{(\theta,\theta') \in K_x^2} \|\theta - \theta'\|^2}{cB} + \epsilon \left( 2\mathbb{E}[M'_{K_x}(Z)] + 1 \right) \end{aligned}$$

where the last inequality holds by (38).

By combining (39) and (40) we obtain that if  $||F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})|| > \tau_B$  then

$$\|\hat{\theta}_{\theta_0,B}^{\omega} - \hat{\theta}_{K_x,n_B}^{\omega}\| \le \epsilon + \sqrt{\frac{2}{\mu_x}} \Big(\epsilon + \frac{n_B D_j^2 \sup_{(\theta,\theta') \in K_x^2} \|\theta - \theta'\|^2}{cB} + \epsilon \Big(2\mathbb{E}[M'_{K_x}(Z)] + 1\Big)\Big)^{\frac{1}{2}}.$$

Together with (37), this shows that for all  $\epsilon > 0, B \in \mathbb{N}$  and  $\omega \in \Omega_{\epsilon,B}$ , we have

$$\sup_{\theta_0 \in \Theta_{j,x}} \|\hat{\theta}^{\omega}_{\theta_0,B} - \hat{\theta}^{\omega}_{K_x,n_B}\| \le \epsilon + \sqrt{\frac{2}{\mu_x}} \Big(\epsilon + \frac{n_B D_j^2 \sup_{(\theta,\theta') \in K_x^2} \|\theta - \theta'\|^2}{cB} + \epsilon \Big(2\mathbb{E}[M'_{K_x}(Z)] + 1\Big)\Big)^{\frac{1}{2}} + \frac{\tau_B}{\lambda_{K_x}}\Big)^{\frac{1}{2}} + \frac{\tau_B}{cB}\Big(\epsilon + \frac{n_B D_j^2 \sup_{(\theta,\theta') \in K_x^2} \|\theta - \theta'\|^2}{cB} + \epsilon \Big(2\mathbb{E}[M'_{K_x}(Z)] + 1\Big)\Big)^{\frac{1}{2}} + \frac{\tau_B}{\lambda_{K_x}}\Big)^{\frac{1}{2}} + \frac{\tau_B}{cB}\Big)^{\frac{1}{2}} + \frac$$

and (36) follows since, by assumption,  $\lim_{B\to\infty} (n_B/B) = \lim_{B\to\infty} \tau_B = 0$  while  $\epsilon > 0$  is arbitrary. To proceed further, let

$$\delta = \frac{\lambda_{K_x}}{4\mathbb{E}[M'_{K_x}(Z)]}$$

and, for all  $B \in \mathbb{N}$ , let

$$\Omega_B^{(3)} = \Omega_B^{(1)} \cap \Omega_B^{(2)} \cap \left\{ \omega \in \Omega : \hat{\theta}_{K_x, n_B}^{\omega} \in \mathring{K}_x, \sup_{\theta_0 \in \Theta_{j,x}} \| \hat{\theta}_{\theta_0, B}^{\omega} - \hat{\theta}_{K_x, n_B}^{\omega} \| \le \delta, \quad \inf_{\theta \in K_x} \lambda_{\min} \left( H_{n_B}^{\omega}(\theta) \right) \ge \lambda_{K_x}, \sup_{(\theta, \theta') \in K_x^2 : \theta \neq \theta'} \frac{\| \nabla^2 F_{n_B}^{\omega}(\theta) - \nabla^2 F_{n_B}^{\omega}(\theta') \|}{\| \theta - \theta' \|} \le 2\mathbb{E}[M'_{K_x}(Z)] \right\}$$

Note that, by Lemmas 1, 4, 7 and 8, and using (36), we have  $\liminf_{B\to\infty} \mathbb{P}(\Omega_B^{(3)}) = 1$ . We now let  $B \in \mathbb{N}$ ,  $\theta_0 \in \Theta_{j,x}$  and  $\omega \in \Omega_B^{(3)}$ . Then, noting that  $\nabla F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) = 0$  since  $\hat{\theta}_{K_x,n_B}^{\omega} \in \mathring{K}_x$  and the function  $F_{n_B}^{\omega}$  is strictly convex on  $K_x$ , it follows from Taylor's theorem (see Amann et al., 2008, Theorem 5.8 and Remark 5.9) that

$$F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega}) - F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) = \frac{1}{2} \left( \hat{\theta}_{\theta_0,B}^{\omega} - \hat{\theta}_{K_x,n_B}^{\omega} \right)^\top H_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) \left( \hat{\theta}_{\theta_0,B}^{\omega} - \hat{\theta}_{K_x,n_B}^{\omega} \right)^\top + R_{\theta_0,B}^{\omega}$$

where

$$|R^{\omega}_{\theta_{0},B}| \leq \|\hat{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\|^{2} \delta \mathbb{E}[M'_{K}(Z)]$$

Therefore, with  $D_j < \infty$  as in Lemma 8, we have

$$\frac{D_{j} \|\theta_{0} - \hat{\theta}_{K_{x},n_{B}}^{\omega}\|^{2}}{T_{\theta_{0},B}^{\omega}} \ge F_{n_{B}}^{\omega}(\hat{\theta}_{\theta_{0},B}^{\omega}) - F_{n_{B}}^{\omega}(\hat{\theta}_{K_{x},n_{B}}^{\omega}) \\
\ge \|\hat{\theta}_{\theta_{0},B}^{\omega} - \hat{\theta}_{K_{x},n_{B}}^{\omega}\|^{2} \left(\frac{\lambda_{K_{x}}}{2} - \delta \mathbb{E}[M_{\Theta}^{\prime}(Z)]\right) \\
= \frac{\lambda_{K_{x}}}{4} \|\hat{\theta}_{\theta_{0},B}^{\omega} - \hat{\theta}_{K_{x},n_{B}}^{\omega}\|^{2}$$

and the result follows from the fact that  $\lambda_{K_x} > 0$  under A1 and by Lemma 2.

### A.3.5 A fourth key property of Algorithm 3

**Lemma 10.** Consider the set-up of Lemma 9. Assume in addition that for some sequence  $(r_B)_{B\geq 1}$  in  $(0,\infty)$  we have

$$\lim_{B \to \infty} \frac{n_B r_B^2}{\tau_B^2 B} = 0.$$

Then, for all  $j \in \mathbb{N}$  and  $\epsilon > 0$ , there exist a constant  $\tilde{C}_{j,\epsilon} \in [1,\infty)$  and a sequence  $(\Omega_{j,\epsilon,B}^{(4)})_{B\geq 1}$  in  $\mathcal{F}$ , such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{j,\epsilon,B}^{(4)}) \geq 1-\epsilon$ , such that for all  $B \in \mathbb{N}$ ,  $\omega \in \Omega_{j,\epsilon,B}^{(4)}$  and  $\theta_0 \in \Theta_{j,x}$  verifying  $\|\theta_0 - \hat{\theta}_{K_{j+1,x},n_B}^{\omega}\| \leq r_B$ , we have

$$\hat{\theta}^{\omega}_{\theta_0,B} \in \Theta_{j+1,x},\tag{41}$$

$$\tilde{B}^{\omega}_{\theta_0} \ge \frac{cB}{2},\tag{42}$$

$$\|\hat{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{j+1,x},n_{B}}\| \le \tilde{C}_{\epsilon} \left(\tau_{B} + r_{B} \sqrt{\frac{n_{B}}{B}}\right)$$

$$\tag{43}$$

$$\|\hat{\theta}^{\omega}_{\theta_{0},B} - \theta_{\star}\| \leq \tilde{C}_{\epsilon} \left(\tau_{B} + n_{B}^{-\frac{\alpha}{1+\alpha}} + r_{B}\sqrt{\frac{n_{B}}{B}}\right)$$

$$\tag{44}$$

where  $c \in (0,1)$  is as in Lemma 9 and  $\alpha \in (0,1]$  is as in A2.

Proof of Lemma 10. To simplify the notation in what follows we let  $K_x = K_{j+1,x}$  and, for all  $\theta \in \mathbb{R}^d$ and  $n \geq 1$ , we let  $H_n(\theta) = \nabla^2 F_n(\theta)$ .

To show the result of the lemma, for all  $B \in \mathbb{N}$  and  $\theta_0 \in \mathbb{R}^d$  we let

$$(\hat{\theta}_{\theta_0,B}, B'_{\theta_0}) = \text{Algorithm } \mathbf{3}(\theta_0, n_B, \tilde{B}_B, 0, \beta)$$

and we denote by  $\{\theta'_{\theta_0,B,t}\}_{t=0}^{\tilde{T}_{\theta_0,B}}$  the sequence in  $\mathbb{R}^d$  generated by Algorithm 3 to compute  $\tilde{\theta}_{\theta_0,B}$  from  $\theta_0$  (which is therefore such that  $\theta'_{\theta_0,B,0} = \theta_0$  and such that  $\theta'_{\theta_0,B,\tilde{T}_{\theta_0,B}} = \tilde{\theta}_{\theta_0,B}$ ). In addition, we let

$$\lambda_{K_x} = \frac{1}{2} \inf_{\theta \in K_x} \lambda_{\min} \big( \mathbb{E} \big[ \nabla^2 f(\theta, Z) \big] \big), \quad \bar{\lambda}_{K_x} = 2 \sup_{\theta \in K_x} \lambda_{\max} \big( \mathbb{E} \big[ \nabla^2 f(\theta, Z) \big] \big)$$

and, for all  $\epsilon > 0$ , we let  $D_{\epsilon} < \infty$  be such that

$$\liminf_{B \to \infty} \left( \|\hat{\theta}_{K_x, n_B} - \theta_\star\| \le D_\epsilon n_B^{-\frac{\alpha}{1+\alpha}} \right) \ge 1 - \epsilon.$$
(45)

Remark that such a finite constant  $D_{\epsilon}$  exists for all  $\epsilon > 0$  by Lemma 5.

Finally, for all  $B \in \mathbb{N}$  and  $\epsilon > 0$  we let

$$\begin{split} \Omega_{\epsilon,B}^{(4)} &= \Omega_B^{(1)} \cap \Omega_B^{(2)} \cap \Omega_B^{(3)} \cap \left\{ \omega \in \Omega : \hat{\theta}_{K_x,n_B}^{\omega} \in \mathring{K}_x, \quad \tilde{B}_B^{\omega} \ge cB, \; \inf_{\theta \in K_x} \lambda_{\min} \big( H_{n_B}^{\omega}(\theta) \big) \ge \lambda_{K_x}, \\ & \| \hat{\theta}_{K_x,n_B}^{\omega} - \theta_\star \| \le D_{\epsilon} n_B^{-\frac{\alpha}{1+\alpha}}, \\ & \sup_{\theta \in K_x} \lambda_{\max} \big( H_{n_B}^{\omega}(\theta) \big) \le \bar{\lambda}_{K_x} \right\} \end{split}$$

with  $c \in (0, 1)$  as in Lemma 9,  $\Omega_B^{(1)}$  as in Lemma 7,  $\Omega_B^{(2)}$  as in Lemma 8 and  $\Omega_B^{(3)}$  as in Lemma 9. Remark that  $\theta_{\star} \in \mathring{K}_x$  and thus, by Lemma 4, Lemmas 7-9 and by (45), we have  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,B}^{(4)}) \geq 1-\epsilon$ for all  $\epsilon > 0$ . In addition, remark that there exists a constant  $D' < \infty$  such that

$$\|\tilde{\theta}^{\omega}_{\theta_0,B} - \hat{\theta}^{\omega}_{K_x,n_B}\| \le D' \|\theta_0 - \hat{\theta}^{\omega}_{K_x,n_B}\| \sqrt{\frac{n_B}{B}}, \quad \forall \omega \in \Omega^{(4)}_{\epsilon,B}, \quad \forall \theta_0 \in \Theta_{j,x}, \quad \forall B \in \mathbb{N}, \quad \forall \epsilon > 0.$$
(46)

We now let  $\epsilon > 0, B \in \mathbb{N}, \omega \in \Omega_{\epsilon,B}^{(4)}$  and  $\theta_0 \in \Theta_{j,x}$  be such that  $\|\theta_0 - \hat{\theta}_{K_x,N_{\mu}^{\omega}}^{\omega}\| \leq r_B$ . Assume first that

$$\left\|\nabla F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})\right\| \le \tau_B$$

and remark that  $\nabla F_{n_B}^{\omega}(\hat{\theta}_{K_x,n_B}^{\omega}) = 0$  since  $F_{n_B}^{\omega}$  is strictly convex on  $K_x$  (as  $\lambda_{K_x} > 0$  under A1 and by Lemma 2) and since  $\hat{\theta}_{K_x,n_B}^{\omega} \in \mathring{K}_x$ . Then, using Taylor's theorem, the fact that  $||Ax|| \ge \lambda_{\min}(A)||x||$  for any  $d \times d$  symmetric matrix A

and  $x \in \mathbb{R}^d$ , we have

$$\tau_B \ge \|\nabla F^{\omega}_{n_B}(\hat{\theta}^{\omega}_{\theta_0,B})\| \ge \lambda_{K_x} \|\hat{\theta}^{\omega}_{\theta_0,B} - \hat{\theta}_{K_x,n_B}\| \Leftrightarrow \|\hat{\theta}^{\omega}_{\theta_0,B} - \hat{\theta}_{K_x,n_B}\| \le \frac{\tau_B}{\lambda_{K_x}}.$$
(47)

Assume now that  $\|\nabla F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})\| > \tau_B$ . In this case,  $\hat{\theta}_{\theta_0,B}^{\omega} = \tilde{\theta}_{\theta_0,B}^{\omega}$  and thus

$$\|\hat{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\| = \|\tilde{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\| \le D' \|\theta_{0} - \hat{\theta}^{\omega}_{K_{x},N^{\omega}_{B}}\| \sqrt{\frac{n_{B}}{B}} \le D' r_{B} \sqrt{\frac{n_{B}}{B}}$$
(48)

where the first inequality holds by (46). Then, (43) follows from (47) and (48), implying that (44)holds since

$$\|\hat{\theta}^{\omega}_{\theta_{0},B} - \theta_{\star}\| \leq \|\hat{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\| + \|\theta_{\star} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\| \leq \|\hat{\theta}^{\omega}_{\theta_{0},B} - \hat{\theta}^{\omega}_{K_{x},n_{B}}\| + D_{\epsilon}n_{B}^{-\frac{\alpha}{1+\alpha}}.$$

To show (42) let  $D'_j < \infty$  be as in Lemma 9 and remark that, using Taylor's theorem,

$$\|\nabla F_{n_B}^{\omega}(\hat{\theta}_{\theta_0,B}^{\omega})\| \leq \bar{\lambda}_{K_x} \|\hat{\theta}_{\theta_0,B}^{\omega} - \hat{\theta}_{K_x,n_B}^{\omega}\| \leq \frac{\bar{\lambda}_{K_x} D'_j \|\theta_0 - \hat{\theta}_{K_x,n_B}^{\omega}\|}{\sqrt{T_{\theta_0,B}^{\omega}}}$$

$$\leq \tau_B \bar{\lambda}_{K_x} D'_j \sqrt{\frac{r_B^2}{\tau_B^2 T_{\theta_0,B}^{\omega}}}$$

$$\tag{49}$$

so that

$$T^{\omega}_{\theta_0,B} \le \min\left(\tilde{T}^{\omega}_{\theta_0,B}, (r_B/\tau_B)^2 (\bar{\lambda}_{K_x} D'_j)^2 + 1\right).$$
 (50)

To proceed further let  $D_j < \infty$  be as in Lemma 8 and remark that

$$\tilde{T}^{\omega}_{\theta_0,B} \geq \frac{\tilde{B}^{\omega}_B}{n_B D_j} \geq \frac{cB}{n_B D_j}$$

Since by assumption,  $\lim_{B\to\infty} (r_B/\tau_B)^2 n_B/B = 0$ , it follows from (50) that for B large enough we have

$$T^{\omega}_{\theta_0,B} \le (r_B/\tau_B)^2 (\bar{\lambda}_{K_x} D'_j)^2$$

and thus

$$\tilde{B}_{\theta_0}^{\omega} \ge \tilde{B}_B^{\omega} - T_{\theta_0,B}^{\omega} n_B D_j \ge c B - \frac{n_B r_B^2}{\tau_B^2} D_j (\bar{\lambda}_{K_x} D'_j)^2 = B \Big( c - \frac{n_B r_B^2}{\tau_B^2 B} D_j (\bar{\lambda}_{K_x} D'_j)^2 \Big).$$

Therefore, since  $\lim_{B\to\infty} n_B r_B^2/(B\tau_B^2) = 0$  by assumption, the proof of the lemma is complete.

## A.4 Proof of Theorem 1

Proof of Theorem 1. Let  $\epsilon \in (0, 1)$ . Then, Lemma 5 and by applying Lemmas 8-9 with  $j = 1, x = \theta_0$ , c = 1 and with

$$n_B = n(B), \quad \tau_B = 0, \quad \forall B \in \mathbb{N},$$
(51)

it directly follows that there exists a sequence  $(\Omega_{\epsilon,B})_{B\geq 1}$  in  $\mathcal{F}$ , such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,B}) \geq 1-\epsilon$ , and a constant  $C_{\epsilon} \in [1,\infty)$ , for which we have

$$\|\tilde{\theta}_B^{\omega} - \theta_{\star}\| \le C_{\epsilon} \left( n(B)^{-\frac{\alpha}{1+\alpha}} + \sqrt{\frac{n_B}{B}} \right), \quad \forall \omega \in \Omega_{\epsilon,B}, \quad \forall B \in \mathbb{N}.$$
(52)

The result of the theorem follows.

### A.5 Proof of Theorem 2

### A.5.1 Proof of the first part of Theorem 2

Proof of the first part of Theorem 2. Noting that if A2 holds for some  $\alpha \in (0, 1]$  then it also holds for any  $\tilde{\alpha} \in (0, \alpha]$ , below we can without loss of generality assume that A2 holds for some  $\alpha \leq \alpha'$ .

To simplify the notation in what follows, for all  $\theta \in \mathbb{R}^d$  and  $n \ge 1$  we let  $H_n(\theta) = \nabla^2 F_n(\theta)$  and, for all  $j \in \mathbb{N}$  and  $B \in \mathbb{N}$  we let  $n_{j,B} = \lceil \kappa B^{\gamma_j} \rceil$  and  $\tau_{j,B} = \tau B^{-\frac{\alpha'}{1+\alpha'}\gamma_j}$ . In addition, for all  $x \in \mathbb{R}^d$  and  $j \in \mathbb{N}$  we let the sets  $\Theta_{j,x}$  and  $K_{j,x}$  be as defined in Lemma 7 and in Lemma 8, respectively.

Let  $\epsilon \in (0, 1)$ . Then, by applying Lemma 10 with  $j = 1, x = \theta_0, c = 1$  and with

$$n_B = n_{1,B}, \quad \tau_B = \tau_{1,B}, \quad r_B = \sup_{(\theta, \theta') \in K_{2,\theta_0}} \|\theta - \theta'\|, \quad \forall B \in \mathbb{N},$$
 (53)

it follows that there exists a sequence  $(\Omega_{\epsilon,B})_{B\geq 1}$  in  $\mathcal{F}$ , such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,B}) \geq 1-\epsilon$ , and a constant  $C_{\epsilon} \in [1,\infty)$  for which we have

$$\max\left(\|\hat{\theta}_{1,B}^{\omega} - \theta_{\star}\|, \|\hat{\theta}_{1,B}^{\omega} - \hat{\theta}_{K_{2,\theta_{0}},n_{1,B}}^{\omega}\|\right) \le C_{\epsilon} B^{-c_{\alpha,\delta}}, \quad \forall \omega \in \Omega_{\epsilon,B}, \quad \forall B \in \mathbb{N}$$
(54)

with

$$c_{\alpha,\delta} = \min\left(\frac{(1-\delta)\alpha}{1+\alpha}, \frac{\delta}{2}\right) = \frac{(1-\delta)\alpha}{1+\alpha}$$

Remark that, by Lemma 6, the sequences  $(n_B)_{B\geq 1}$ ,  $(\tau_B)_{B\geq 1}$  and  $(r_B)_{B\geq 1}$  defined in (53) satisfy the assumptions of Lemma 10. Then, the result of the first part of the theorem for J = 1 follows from (54).

We assume now that J > 1 and, without loss of generality, we assume that  $K_{j+1,\theta_0} = K$  for all  $j \in \{1, \ldots, J\}$  and for some compact and convex set  $K \subset \mathbb{R}^d$ . In addition, we let

$$\lambda_J = \frac{1}{2} \inf_{\theta \in K} \lambda_{\min} \big( \mathbb{E} \big[ \nabla^2 f(\theta, Z) \big] \big)$$

and, for all  $\epsilon \in (0, 1)$ , we let  $C'_{\epsilon} < \infty$  be such that

$$\liminf_{B \to \infty} \mathbb{P}\Big( \|\hat{\theta}_{K,n_{j,B}} - \theta_{\star}\| \le C_{\epsilon}' n_{j,B}^{-\frac{\alpha}{1+\alpha}} \Big) \ge 1 - \frac{\epsilon}{J}, \quad \forall j \in \{1, \dots, J\}$$

Remark that, for all  $\epsilon \in (0, 1)$ , such a constant  $C'_{\epsilon} < \infty$  exists by Lemma 5. Remark also that, using Fréchet's inequality,

$$\liminf_{B \to \infty} \mathbb{P}\Big(\max_{j \in \{1, \dots, J\}} \|\hat{\theta}_{K, n_{j, B}} - \theta_{\star}\| \le C'_{\epsilon} n_{j, B}^{-\frac{\alpha}{1+\alpha}}\Big) \ge 1 - \epsilon.$$
(55)

Then, for all  $\epsilon \in (0, 1)$  and  $B \in \mathbb{N}$ , we let

$$\Omega_{\epsilon,B}' = \Omega_{\epsilon,B} \cap_{j=1}^{J} \left( \Omega_{j,B}^{(1)} \cap \Omega_{j,B}^{(2)} \right)$$
$$\cap \left\{ \omega \in \Omega : \min_{j \in \{1, \dots, J\}} \inf_{\theta \in K} \lambda_{\min} \left( H_{n_{j,B}}^{\omega}(\theta) \right) \ge \lambda_{J}, \max_{j \in \{1, \dots, J\}} n_{j,B}^{\frac{\alpha}{1+\alpha}} \| \hat{\theta}_{K,n_{j,B}}^{\omega} - \theta_{\star} \| \le C_{\epsilon}',$$
$$\hat{\theta}_{K,n_{j}}^{\omega} \in \mathring{K} \text{ for all } j \in \{1, \dots, J\} \right\}$$

with  $\Omega_{\epsilon,B}$  as above,  $\Omega_{j,B}^{(1)}$  as in Lemma 7 and  $\Omega_{j,B}^{(2)}$  as in Lemma 8. Remark that, by Lemmas 4, 7 and 8, and by (55), we have  $\liminf_{B\to\infty} \mathbb{P}(\Omega'_{\epsilon,B}) \geq 1-\epsilon$ .

Let  $\epsilon \in (0, 1), B \in \mathbb{N}$  and  $\omega \in \Omega'_{\epsilon,B}$ . Remark that  $\nabla F^{\omega}_{n_{j,B}}(\hat{\theta}^{\omega}_{K,n_{j,B}}) = 0$  for all  $j \in \{1, \ldots, J\}$ , since for all  $j \in \{1, \ldots, J\}$  the function  $F^{\omega}_{n_{j,B}}$  is strictly convex on K (since  $\lambda_J > 0$  under A1 and by Lemma 2) while  $\hat{\theta}^{\omega}_{K,n_j} \in \mathring{K}$ . Therefore, using Taylor's theorem, the fact that  $||Ax|| \ge \lambda_{\min}(A)||x||$  for any  $d \times d$ symmetric matrix A and  $x \in \mathbb{R}^d$ , we have

$$F_{n_{j,B}}^{\omega}(\theta) - F_{n_{j,B}}^{\omega}(\hat{\theta}_{K,n_{j,B}}^{\omega}) \ge \lambda_J \|\theta - \hat{\theta}_{K,n_{j,B}}^{\omega}\|^2, \quad \forall \theta \in K.$$
(56)

For all  $j \in \mathbb{N}$  let  $D_j$  be as in Lemma 8 and assume without loss of generality that  $D_j \leq D_{j+1}$  for all  $j \in \mathbb{N}$  and that  $D_J/\lambda_J \geq 1$ . In addition, let  $J_B^{\omega}$  denote the smallest  $j \in \{1, \ldots, J\}$  such that  $\hat{\theta}_{j,B}^{\omega} = \hat{\theta}_{J,B}^{\omega}$ . Then, noting that  $n_{j,B} \leq n_{j+1,B}$  for all  $j \in \mathbb{N}$ , if  $J_B^{\omega} > 1$  we have

$$\begin{split} \|\hat{\theta}_{J_{B}^{\omega},B}^{\omega} - \hat{\theta}_{K,n_{J_{B}^{\omega},B}}^{\omega}\|^{2} &\leq \frac{1}{\lambda_{J}} \Big( F_{n_{J_{B}^{\omega},B}}^{\omega} (\hat{\theta}_{J_{B}^{\omega},B}^{\omega}) - F_{n_{J_{B}^{\omega},B}}^{\omega} (\hat{\theta}_{K,n_{J_{B}^{\omega},B}}^{\omega}) \Big) \\ &\leq \frac{D_{J} \|\hat{\theta}_{J_{B}^{\omega}-1,B}^{\omega} - \hat{\theta}_{K,n_{J_{B}^{\omega},B}}^{\omega}\|^{2}}{T_{J_{B}^{\omega},B}^{\omega} \lambda_{J}} \\ &\leq \frac{2D_{J}}{\lambda_{J}} \Big( \|\hat{\theta}_{J_{B}^{\omega}-1,B}^{\omega} - \hat{\theta}_{K,n_{J_{B}^{\omega}-1,B}}^{\omega}\|^{2} + \|\hat{\theta}_{K,n_{J_{B}^{\omega},B}}^{\omega} - \hat{\theta}_{K,n_{J_{B}^{\omega}-1,B}}^{\omega}\|^{2} \Big) \\ &\leq \frac{4D_{J}}{\lambda_{J}} \Big( \|\hat{\theta}_{J_{B}^{\omega}-1,B}^{\omega} - \hat{\theta}_{K,n_{J_{B}^{\omega}-1,B}}^{\omega}\|^{2} + (C_{\ell}')^{2} n_{1,B}^{-\frac{2\alpha}{1+2\alpha}} \Big) \\ &\leq \Big(\frac{4D_{J}}{\lambda_{J}}\Big)^{J} \|\hat{\theta}_{1,B}^{\omega} - \hat{\theta}_{K,n_{1,B}}^{\omega}\|^{2} + (C_{\ell}')^{2} n_{1,B}^{-\frac{2\alpha}{1+2\alpha}} \sum_{j=1}^{J} (4D_{J}/\lambda_{J})^{j} \\ &\leq \Big(\frac{4D_{J}}{\lambda_{J}}\Big)^{J} C_{\ell}^{2} B^{-2c_{\alpha,\delta}} + (C_{\ell}')^{2} n_{1,B}^{-\frac{2\alpha}{1+2\alpha}}} \sum_{j=1}^{J} (4D_{J}/\lambda_{J})^{j} \end{split}$$

where the first inequality uses (56) and the last inequality uses (54).

If  $J_B^{\omega} = 1$  then, from (54), we have

$$\|\hat{\theta}_{J_B^{\omega},B}^{\omega} - \hat{\theta}_{K,n_{J_B^{\omega},B}}^{\omega}\|^2 \le C_{\epsilon}^2 B^{-2c_{\alpha,\delta}}.$$

Therefore,

$$\|\hat{\theta}_{J,B}^{\omega} - \hat{\theta}_{K,n_{J}\omega,B}^{\omega}\| = \|\hat{\theta}_{J_{B},B}^{\omega} - \hat{\theta}_{K,n_{J}\omega,B}^{\omega}\| \le C_{\epsilon}^{\prime\prime} B^{-c_{\alpha,\delta}}$$

where

$$C_{\epsilon}^{\prime\prime} = \left( \left( \frac{4D_J}{\lambda_J} \right)^J C_{\epsilon}^2 + (C_{\epsilon}^{\prime})^2 \sum_{j=1}^J (4D_J/\lambda_J)^j \right)^{1/2}.$$

Therefore,

$$\begin{aligned} \|\hat{\theta}_{J,B}^{\omega} - \theta_{\star}\| &\leq \|\hat{\theta}_{K,n_{J}\omega_{,B}}^{\omega} - \theta_{\star}\| + \|\hat{\theta}_{J,B}^{\omega} - \hat{\theta}_{K,n_{J}\omega_{,B}}^{\omega}\| \\ &\leq C_{\epsilon}' n_{1,B}^{-\frac{\alpha}{1+\alpha}} + C_{\epsilon}'' B^{-c_{\alpha,\delta}} \\ &\leq (C_{\epsilon}' + C_{\epsilon}'') B^{-c_{\alpha,\delta}}. \end{aligned}$$

The result of the first part of the theorem follows.

### A.5.2 Proof of the second part of Theorem 2

Proof of the 2nd part of Theorem 2. Let  $c_{\alpha} = \alpha/(1+\alpha)$  and, for all  $j \in \mathbb{N}$  and  $B \in \mathbb{N}$ , let  $n_{j,B} = \lceil \kappa B^{\gamma_j} \rceil$ ,  $\tau_{j,B} = \tau B^{-c_{\alpha}\gamma_j}$  and  $r_{j,B} = B^{-c_{\alpha}\gamma_j}$ . Finally, let  $r_{0,B} = 1$  for all  $B \in \mathbb{N}$  and, for all  $x \in \mathbb{R}^d$  and  $j \in \mathbb{N}$ , let  $\Theta_{j,x}$  be as defined in Lemma 7 and  $K_{j+1,x}$  be as defined in Lemma 8.

To prove the second part of the theorem we show by induction that for all  $j \in \mathbb{N}$  there exist, for all  $\epsilon \in (0, \infty)$ , a sequence  $(\Omega_{\epsilon,j,B})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,j,B}) \geq 1-\epsilon$ , and a constant  $C_{\epsilon,j} \in [1,\infty)$  for which we have

$$B_{j}^{\omega} \geq \frac{B}{C_{\epsilon,j}}, \quad \|\hat{\theta}_{j,B}^{\omega} - \theta_{\star}\| \leq C_{\epsilon,j} r_{j,B}, \quad \|\hat{\theta}_{j,B}^{\omega} - \hat{\theta}_{K_{j+1,\theta_{0}},n_{j,B}}^{\omega}\| \leq C_{\epsilon,j} r_{j,B}, \quad \hat{\theta}_{j,B}^{\omega} \in \Theta_{j+1,\theta_{0}}$$
(57)

for all  $w \in \Omega_{\epsilon,j,B}$  and  $B \in \mathbb{N}$ .

To do so for all  $B \in \mathbb{N}$  let  $n_B = n_{1,B}$ ,  $\tau_B = \tau_{1,B}$  and  $r_B = \sup_{(\theta,\theta') \in K_{2,\theta_0}} \|\theta - \theta'\|$ , and note that, by Lemma 6, these sequences  $(n_B)_{B \ge 1}$ ,  $(\tau_B)_{B \ge 1}$  and  $(r_B)_{B \ge 1}$  satisfy the assumptions of Lemma 10. Therefore, by applying Lemma 10 with c = 1,  $x = \theta_0$  and j = 1, it follows that for all  $\epsilon \in (0, \infty)$  there exist a sequence  $(\Omega_{\epsilon,1,B})_{B \ge 1}$  in  $\mathcal{F}$  such that  $\liminf_{B \to \infty} \mathbb{P}(\Omega_{\epsilon,j,B}) \ge 1 - \epsilon$ , and a constant  $C_{\epsilon,1} \in [1, \infty)$ , such that (57) holds for all  $w \in \Omega_{\epsilon,1,B}$  and all  $B \in \mathbb{N}$  when j = 1.

Assume now that for some  $s \in \mathbb{N}$  there exist, for all  $\epsilon \in (0, \infty)$ , a sequence  $(\Omega_{\epsilon,s,B})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,s,B}) \geq 1-\epsilon$  and a constant  $C_{\epsilon,s} \in [1,\infty)$  for which (57) holds for all  $w \in \Omega_{\epsilon,s,B}$  and all  $B \in \mathbb{N}$  when j = s.

For all  $\epsilon > 0$  and  $B \in \mathbb{N}$  we now let  $\tilde{B}_{\epsilon,s}^{\omega} = \max(B_s^{\omega}, B/C_{\epsilon,s})$  for all  $\omega \in \Omega$ . In addition, for all  $B \in \mathbb{N}$ , we let  $n_B = n_{s+1,B}$ ,  $\tau_B = \tau_{s+1,B}$  and  $r_B = r_{s,B}$  and remark that, by Lemma 6, these sequences  $(n_B)_{B\geq 1}$ ,  $(\tau_B)_{B\geq 1}$  and  $(r_B)_{B\geq 1}$  satisfy the assumptions of Lemma 10. Let  $\epsilon > 0$ . Then, by applying Lemma 10 with  $c = 1/C_{\frac{\epsilon}{2},s}$ ,  $x = \theta_0$  and j = s+1, it follows that there exist a sequence  $(\Omega'_{\epsilon/2,s+1,B})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega'_{\epsilon/2,s+1,B}) \geq 1 - \epsilon/2$  and a constant  $\tilde{C}_{\epsilon/2,s+1} \in [1,\infty)$  such that, for all  $B \in \mathbb{N}$  and  $\omega \in \Omega_{\epsilon/2,s,B} \cap \Omega'_{\epsilon/2,s+1,B}$ , we have

$$\hat{\theta}_{s+1,B}^{\omega} \in \Theta_{s+2,\theta_0}, \quad B_{s+1}^{\omega} \ge \frac{B}{2C_{\frac{\epsilon}{2},s}}, \\ \|\hat{\theta}_{s+1,B}^{\omega} - \hat{\theta}_{K_{s+1,\theta_0},n_{s+1,B}}^{\omega}\| \le \tilde{C}_{\epsilon/2,s+1} \left(\tau_{s+1,B} + r_{s,B} \sqrt{\frac{n_{s+1,B}}{B}}\right)$$
(58)  
$$\|\hat{\theta}_{s+1,B}^{\omega} - \theta_{\star}\| \le \tilde{C}_{\epsilon/2,s+1} \left(\tau_{s+1,B} + r_{s,B} \sqrt{\frac{n_{s+1,B}}{B}}\right).$$

To proceed further remark that, for some constant  $D < \infty$ , f

$$\tau_{s+1,B} + r_{s,B} \sqrt{\frac{n_{s+1,B}}{B}} \le D\left(B^{-c_{\alpha}\gamma_{s+1}} + B^{\frac{\gamma_{s+1}-1}{2} - c_{\alpha}\gamma_s}\right), \quad \forall B \in \mathbb{N}.$$
(59)

To proceed further let  $\gamma'_{s+1}$  be as defined in Lemma 6 when  $\alpha' = \alpha$ , and note that  $\gamma'_{s+1}$  is such that

$$B^{-c_{\alpha}\gamma'_{s+1}} = B^{\frac{\gamma'_{s+1}-1}{2} - c_{\alpha}\gamma_s}, \quad \forall B \in \mathbb{N}.$$

By Lemma 6, we have  $\gamma_{s+1} < \gamma'_{s+1}$  and thus, since for all  $B \in \mathbb{N} \setminus \{1\}$  the mapping  $\gamma \mapsto B^{-c_{\alpha}\gamma}$  is decreasing in  $\gamma$  while the mapping  $\gamma \mapsto B^{\frac{\gamma-1}{2}-c_{\alpha}\gamma_s}$  is increasing in  $\gamma$ , it follows that

$$B^{\frac{\gamma_{s+1}-1}{2}-c_{\alpha}\gamma_{s}} \le B^{-c_{\alpha}\gamma_{s+1}}, \quad \forall B \in \mathbb{N}$$

and thus, by (59),

$$\tau_{s+1,B} + r_{s,B} \sqrt{\frac{n_{s+1,B}}{B}} \le 2DB^{-c_{\alpha}\gamma_{s+1}} = 2Dr_{s+1,B}, \quad \forall B \in \mathbb{N}.$$
(60)

By combining (58) and (60), it follows that there exists a constant  $C_{\epsilon,s+1} \in [1,\infty]$  such that, for j = s + 1, (57) holds for all  $\omega \in \Omega_{\epsilon/2,s,B} \cap \Omega'_{\epsilon/2,s+1,B}$  and  $B \in \mathbb{N}$  where, using Fréchet's inequality,

$$\liminf_{B \to \infty} \mathbb{P}\big(\Omega_{\epsilon/2,s,B} \cap \Omega'_{\epsilon/2,s+1,B}\big) \ge \liminf_{B \to \infty} \mathbb{P}\big(\Omega_{\epsilon/2,s,B}\big) + \liminf_{B \to \infty} \mathbb{P}\big(\Omega'_{\epsilon/2,s+1,B}\big) - 1 \ge 1 - \epsilon.$$

Since  $\epsilon > 0$  is arbitrary, this concludes to show that for all  $j \in \mathbb{N}$  there exit, for all  $\epsilon \in (0, \infty)$ , a sequence  $(\Omega_{\epsilon,j,B})_{B\geq 1}$  in  $\mathcal{F}$  such that  $\liminf_{B\to\infty} \mathbb{P}(\Omega_{\epsilon,j,B}) \geq 1-\epsilon$ , and a constant  $C_{\epsilon,j} \in [1,\infty)$  such that (57) holds for all  $\omega \in \Omega_{\epsilon,j,B}$  and all  $B \in \mathbb{N}$ . The second part of the theorem follows.