

# MODULARITY OF CERTAIN PRODUCTS OF THE ROGERS-RAMANUJAN CONTINUED FRACTION

RUSSELLE GUADALUPE

ABSTRACT. We study the modularity of the functions of the form  $r(\tau)^a r(2\tau)^b$ , where  $a$  and  $b$  are integers with  $(a, b) \neq (0, 0)$  and  $r(\tau)$  is the Rogers-Ramanujan continued fraction, which may be considered as companions to the Ramanujan's function  $k(\tau) = r(\tau)r(2\tau)^2$ . In particular, we show that under some condition on  $a$  and  $b$ , there are finitely many such functions generating the field of all modular functions on the congruence subgroup  $\Gamma_1(10)$ . Furthermore, we establish certain arithmetic properties of the function  $l(\tau) = r(2\tau)/r(\tau)^2$ , which can be used to evaluate these products. We employ the methods of Lee and Park, and some properties of  $\eta$ -quotients and generalized  $\eta$ -quotients to prove our results.

## 1. INTRODUCTION

For an element  $\tau$  in the complex upper half-plane  $\mathbb{H}$  and  $q := e^{2\pi i\tau}$ , the Rogers-Ramanujan continued fraction is defined by

$$r(\tau) = \frac{q^{1/5}}{1 + \frac{q}{1 + \frac{q^2}{1 + \frac{q^3}{1 + \dots}}}} = q^{1/5} \prod_{n=1}^{\infty} \frac{(1 - q^{5n-1})(1 - q^{5n-4})}{(1 - q^{5n-2})(1 - q^{5n-3})}.$$

In his first correspondence [2, pp. 21–30] to Hardy in 1913, Ramanujan asserted that

$$r(i) = \sqrt{\frac{5 + \sqrt{5}}{2}} - \frac{\sqrt{5} + 1}{2}.$$

He later claimed in his second letter [2, pp. 53–62] to Hardy that the values of  $r(\frac{\sqrt{-n}}{2})$  “can be exactly found” for any positive rational number  $n$ . Ramanujan's observations were settled by Gee and Honsbeek [10], who treated  $r(\tau)$  as a modular function, and used Watson's formulas [18] to show that  $r(\tau)$  generates the field of all modular functions on the congruence subgroup  $\Gamma(5)$ . Consequently, they determined the exact values of  $r(\tau)$  at some imaginary quadratic points  $\tau \in \mathbb{H}$  as nested radicals.

Recently, Lee and Park [12] proved that the Ramanujan's function  $k(\tau) := r(\tau)r(2\tau)^2$  generates the field of all modular functions on the congruence subgroup  $\Gamma_1(10)$ . They also proved that if  $K$  is an imaginary quadratic field with ring of integers  $\mathbb{Z}[\tau]$  and  $\tau \in K \cap \mathbb{H}$ , then  $k(\tau)$  generates the ray class field  $K_{(10)}$  modulo 10 over  $K$ . As a result, they computed the exact values of  $r(\tau)$  at some imaginary quadratic points  $\tau \in \mathbb{H}$ . We refer the reader to [1, 6, 19, 22] for other arithmetic properties of  $k(\tau)$ .

In this paper, we partially extend Lee and Park's result by exploring the functions  $r(\tau)^a r(2\tau)^b$  for integers  $a$  and  $b$  with  $(a, b) \neq (0, 0)$ , which may be considered as companions of  $k(\tau)$ . More precisely, we show that under some condition on  $a$  and  $b$ , there

---

2020 *Mathematics Subject Classification*: Primary 11F03; 11F11, 11R37.

*Key words and phrases*: Rogers-Ramanujan continued fraction,  $\eta$ -quotient, generalized  $\eta$ -quotient, modular equations, ray class fields

are finitely many such functions generating the field of all modular functions on  $\Gamma_1(10)$ . We also show that under the same condition, the above functions can be expressed in terms of  $l(\tau) := r(2\tau)/r(\tau)^2$ . We further establish certain arithmetic properties of  $l(\tau)$ , which will be useful in computing exactly the values of  $r(\tau)^a r(2\tau)^b$  at imaginary quadratic points  $\tau \in \mathbb{H}$ . We apply the methods of Lee and Park [12], as well as some properties of  $\eta$ -quotients and generalized  $\eta$ -quotients defined by Yang [20], to prove our results.

We organize the paper as follows. In Section 2, we give a brief background about modular functions, particularly  $\eta$ -quotients and generalized  $\eta$ -quotients, on some congruence subgroups of  $\mathrm{SL}_2(\mathbb{Z})$ . In Section 3, we discuss the modularity of the functions  $r(\tau)^a r(2\tau)^b$  on  $\Gamma_1(10)$ , where  $a$  and  $b$  are integers with  $(a, b) \neq (0, 0)$  and  $5 \mid a - 3b$ . We show that these functions can be expressed as a rational function of  $l(\tau)$ . In Section 4, we prove that modular equations for  $l(\tau)$  of any level always exist, and derive such equations for  $l(\tau)$  of levels two and five. We prove in Section 5 that if  $K$  is an imaginary quadratic field with ring of integers  $\mathbb{Z}[\tau]$  for some  $\tau \in K \cap \mathbb{H}$ , then  $l(\tau)$  generates  $K_{(10)}$ . We also prove that  $l(\tau)$  and the functions  $r(\tau)^a r(2\tau)^b$  are algebraic units for  $\tau \in K \cap \mathbb{H}$ . Finally, we give an example of finding the class polynomial for  $K_{(10)}$  when the discriminant of  $K$  is divisible by 4 using Shimura reciprocity law due to Cho and Koo [4] and Gee [9] to evaluate  $r(\tau)^a r(2\tau)^b$ . We perform most of our calculations via *Mathematica*.

## 2. PRELIMINARIES

In this section, we recall some facts about modular functions on some congruence subgroups, which are subgroups of  $\mathrm{SL}_2(\mathbb{Z})$  containing

$$\Gamma(N) := \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : a - 1 \equiv b \equiv c \equiv d - 1 \equiv 0 \pmod{N} \right\}$$

for some integer  $N \geq 1$ . Examples of congruence subgroups we focus on in this paper are

$$\begin{aligned} \Gamma_1(N) &:= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : a - 1 \equiv c \equiv d - 1 \equiv 0 \pmod{N} \right\}, \\ \Gamma_0(N) &:= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z}) : c \equiv 0 \pmod{N} \right\}. \end{aligned}$$

Any element of a given congruence subgroup  $\Gamma$  acts on the extended upper half-plane  $\mathbb{H}^* := \mathbb{H} \cup \mathbb{Q} \cup \{\infty\}$  by a linear fractional transformation. We define the cusps of  $\Gamma$  as the equivalence classes of  $\mathbb{Q} \cup \{\infty\}$  under this action, and it is known that there are finitely many inequivalent cusps of  $\Gamma$ . We define the modular function on  $\Gamma$  by a function  $f(\tau) : \mathbb{H} \rightarrow \mathbb{C}$  such that

- $f(\tau)$  is meromorphic on  $\mathbb{H}$ ,
- $f(\gamma\tau) = f(\tau)$  for all  $\gamma \in \Gamma$ , and
- $f(\tau)$  is meromorphic at all cusps of  $\Gamma$ .

The last condition implies that for every cusp  $r$  of  $\Gamma$  and an element  $\gamma \in \mathrm{SL}_2(\mathbb{Z})$  with  $\gamma(\infty) = r$ , the  $q$ -expansion of  $f \circ \gamma$  given by  $f(\gamma\tau) = \sum_{n \geq n_0} a_n q^{n/h}$ , for some integers  $h$  and  $n_0$  with  $a_{n_0} \neq 0$ , has finitely many negative exponents. We call the least such  $n_0$  the order of  $f(\tau)$  at  $r$ , denoted by  $\mathrm{ord}_r f(\tau)$ , and we say that  $f(\tau)$  has a zero (resp., a pole) at  $r$  if  $\mathrm{ord}_r f(\tau)$  is positive (resp., negative). We also call  $h$  the width of  $r$ , which is the smallest positive integer  $k$  such that  $\gamma \begin{bmatrix} 1 & k \\ 0 & 1 \end{bmatrix} \gamma^{-1} \in \pm\Gamma$ . We note that  $h$  is independent on the choice of  $\gamma$ .

Let  $A_0(\Gamma)$  be the field of all modular functions on  $\Gamma$  and let  $A_0(\Gamma)_{\mathbb{Q}}$  be its subfield consisting of all modular functions on  $\Gamma$  with rational  $q$ -expansion. We identify  $A_0(\Gamma)$  with the field of all meromorphic functions on the modular curve  $X(\Gamma) := \Gamma \backslash \mathbb{H}^*$ . If

$f(\tau) \in A_0(\Gamma)$  has no zeros nor poles on  $\mathbb{H}$ , then the degree  $[A_0(\Gamma) : \mathbb{C}(f(\tau))]$  is the total degree of poles of  $f(\tau)$  given by  $-\sum_r \text{ord}_r f(\tau)$ , where the sum ranges over the inequivalent cusps  $r$  of  $\Gamma$  for which  $f(\tau)$  has a pole at  $r$  (see [17, Prop. 2.11]).

We describe a systematic way of getting the set of inequivalent cusps of the congruence subgroup  $\Gamma := \Gamma_1(N) \cap \Gamma_0(mN)$  for positive integers  $m$  and  $N$ , following Cho, Koo, and Park [5]. Let  $\Gamma(1)_\infty$  be the stabilizer of  $\infty$  in  $\text{SL}_2(\mathbb{Z})$  and let  $\Gamma\gamma_i\Gamma(1)_\infty$ ,  $i = 1, \dots, g$ , be the representatives for the double coset  $\Gamma \backslash \text{SL}_2(\mathbb{Z}) / \Gamma(1)_\infty$ . Then the set  $\{\gamma_i(\infty) : i = 1, \dots, g\}$  is the set of all inequivalent cusps of  $\Gamma$ . Consider the set

$$M = \{(c, d) \in (\mathbb{Z}/mN\mathbb{Z})^2 : \gcd(c, d, mN) = 1\}$$

and the subgroup  $\Delta$  of  $(\mathbb{Z}/mN\mathbb{Z})^\times$  given by

$$\Delta = \{\pm(1 + Nk) \in (\mathbb{Z}/mN\mathbb{Z})^\times : k = 0, \dots, m-1\}.$$

We define a relation  $\sim$  on  $M$  as follows: if  $(c_1, d_1), (c_2, d_2) \in M$ , then  $(c_1, d_1) \sim (c_2, d_2)$  if  $(c_2, d_2) \equiv (sc_1, sd_1 + nc_1) \pmod{mN}$  for some  $s \in \Delta$  and  $n \in \mathbb{Z}/mN\mathbb{Z}$ . This is an equivalence relation, and the map  $\phi : \Gamma \backslash \text{SL}_2(\mathbb{Z}) / \Gamma(1)_\infty \rightarrow M / \sim$  sending  $\Gamma \begin{bmatrix} a & b \\ c & d \end{bmatrix} \Gamma(1)_\infty$  to the equivalence class of  $(c, d)$  is well-defined and bijective (see [8, Prop. 3.8.5]).

**Lemma 2.1** ([5]). *Suppose  $a, c, a', c' \in \mathbb{Z}$  such that  $\gcd(a, c) = \gcd(a', c') = 1$ . We denote  $\infty := \pm 1/0$ . Then  $a/c$  and  $a'/c'$  are equivalent under  $\Gamma_1(N) \cap \Gamma_0(mN)$  if and only if there exist  $s \in \Delta$  and  $n \in \mathbb{Z}$  such that  $\begin{bmatrix} a' \\ c' \end{bmatrix} \equiv \begin{bmatrix} s^{-1}a+nc \\ sc \end{bmatrix} \pmod{mN}$ .*

We now consider the surjective canonical projection  $\pi_x : (\mathbb{Z}/mN\mathbb{Z})^\times \rightarrow (\mathbb{Z}/x\mathbb{Z})^\times$  for a positive divisor  $x$  of  $mN$ . For each positive divisor  $c$  of  $mN$ , let  $s'_{c,i}, i = 1, \dots, n_c$ , be the distinct coset representatives of  $\pi_{mN/c}(\Delta)$  in  $(\mathbb{Z}/\frac{mN}{c}\mathbb{Z})^\times$  with  $n_c = \varphi(\frac{mN}{c}) / |\pi_{mN/c}(\Delta)|$  (here,  $\varphi$  is the Euler's totient function). We then choose  $s_{c,i} \in (\mathbb{Z}/mN\mathbb{Z})^\times$  such that  $\pi_{mN/c}(s_{c,i}) = s'_{c,i}$  for each  $i$  and define  $S_c = \{s_{c,i} \in (\mathbb{Z}/mN\mathbb{Z})^\times : i = 1, \dots, n_c\}$ . We next let  $a'_{c,j}, j = 1, \dots, m_c$ , be the distinct coset representatives of  $\pi_c(\Delta \cap \ker \pi_{mN/c})$  in  $(\mathbb{Z}/c\mathbb{Z})^\times$  with

$$m_c = \frac{\varphi(c)}{|\pi_c(\Delta \cap \ker \pi_{mN/c})|} = \frac{\varphi(c) |\pi_{mN/c}(\Delta)|}{|\pi_{mN/\gcd(c, mN/c)}(\Delta)|}.$$

We then choose a positive integer  $a_{c,j}$  less than  $mN$  and coprime to  $mN$  such that  $\pi_c(a_{c,j}) = a'_{c,j}$  for each  $j$  and define  $A_c = \{a_{c,j} : j = 1, \dots, m_c\}$ .

**Lemma 2.2** ([5]). *With the notations as above, let*

$$S = \{(cs_{c,i}, a_{c,j}) \in (\mathbb{Z}/mN\mathbb{Z})^2 : 0 < c \mid mN, s_{c,i} \in S_c, a_{c,j} \in A_c, \gcd(cs_{c,i}, a_{c,j}, mN) = 1\}.$$

*For a given  $(cs_{c,i}, a_{c,j}) \in S$ , choose coprime integers  $x$  and  $y$  such that  $x \equiv cs_{c,i} \pmod{mN}$  and  $y \equiv a_{c,j} \pmod{mN}$ . Then the set of  $y/x$  for such  $x$  and  $y$  is the set of all inequivalent cusps of  $\Gamma_1(N) \cap \Gamma_0(mN)$  and the number of such cusps is*

$$|S| = \sum_{0 < c \mid mN} n_c m_c = \sum_{0 < c \mid mN} \frac{\varphi(c) \varphi(mN/c)}{|\pi_{mN/\gcd(c, mN/c)}(\Delta)|}.$$

We now obtain the width of a cusp of  $\Gamma_1(N) \cap \Gamma_0(mN)$  as the following result shows.

**Lemma 2.3** ([5]). *The width  $h$  of a cusp  $a/c$  of  $\Gamma_1(N) \cap \Gamma_0(mN)$ , where  $a$  and  $c$  are coprime integers, is*

$$h = \begin{cases} \frac{m}{\gcd(c^2/4, m)} & \text{if } N = 4, \gcd(m, 2) = 1, \gcd(c, 4) = 2, \\ \frac{mN}{\gcd(c, N) \gcd(m, c^2/\gcd(c, N))} & \text{otherwise.} \end{cases}$$

For the congruence subgroups  $\Gamma_0(N)$  and  $\Gamma_1(N)$ , we obtain a nice description for their respective sets of inequivalent cusps.

**Corollary 2.4** ([5]). *Suppose  $a, c, a', c' \in \mathbb{Z}$  such that  $\gcd(a, c) = \gcd(a', c') = 1$ . We denote  $\infty := \pm 1/0$ . Let  $S_\Gamma$  be the set of all inequivalent cusps of a congruence subgroup  $\Gamma$ .*

- (1) *The cusps  $a/c$  and  $a'/c'$  are equivalent under  $\Gamma_0(m)$  if and only if there exist  $s \in (\mathbb{Z}/m\mathbb{Z})^\times$  and  $n \in \mathbb{Z}$  such that  $\begin{bmatrix} a' \\ c' \end{bmatrix} \equiv \begin{bmatrix} s^{-1}a+nc \\ sc \end{bmatrix} \pmod{m}$ . Moreover, we have*

$$S_{\Gamma_0(m)} = \{a_{c,j}/c \in \mathbb{Q} : 0 < c \mid m, 0 < a_{c,j} \leq m, \gcd(a_{c,j}, m) = 1 \\ a_{c,j} = a_{c,j'} \iff a_{c,j} \equiv a_{c,j'} \pmod{\gcd(c, m/c)}\}$$

*and the width of the cusp  $a/c$  in  $\Gamma_0(m)$  is  $m/\gcd(c^2, m)$ .*

- (2) *The cusps  $a/c$  and  $a'/c'$  are equivalent under  $\Gamma_1(N)$  if and only if there exists  $n \in \mathbb{Z}$  such that  $\begin{bmatrix} a' \\ c' \end{bmatrix} \equiv \pm \begin{bmatrix} a+nc \\ c \end{bmatrix} \pmod{N}$ . Moreover, we have*

$$S_{\Gamma_1(N)} = \{y_{c,j}/x_{c,i} \in \mathbb{Q} : 0 < c \mid N, 0 < s_{c,i}, a_{c,j} \leq N, \\ \gcd(s_{c,i}, N) = \gcd(a_{c,j}, N) = 1, \\ s_{c,i} = s_{c,i'} \iff s_{c,i} \equiv \pm s_{c,i'} \pmod{N/c}, \\ a_{c,j} = a_{c,j'} \iff a_{c,j} \equiv \begin{cases} \pm a_{c,j'} \pmod{c} & \text{if } c \in \{N/2, N\}, \\ a_{c,j'} \pmod{c} & \text{otherwise,} \end{cases} \\ (x_{c,i}, y_{c,j}) \in \mathbb{Z}^2, \gcd(x_{c,i}, y_{c,j}) = 1, \\ x_{c,i} \equiv cs_{c,i} \pmod{N}, y_{c,j} \equiv a_{c,j} \pmod{N}\}$$

*and the width of the cusp  $a/c$  in  $\Gamma_1(N)$  is 1 if  $N = 4$  and  $\gcd(c, 4) = 2$  and  $N/\gcd(c, N)$  otherwise.*

We now introduce an  $\eta$ -quotient, which is a function of the form

$$f(\tau) = \prod_{\delta \mid N} \eta(\delta\tau)^{r_\delta}$$

for some indexed set  $\{r_\delta \in \mathbb{Z} : \delta \mid N\}$ , where  $\eta(\tau) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n)$  is the Dedekind eta function. The next two lemmas provide necessary conditions on the modularity and the behavior at the cusps of an  $\eta$ -quotient on  $\Gamma_0(N)$ .

**Lemma 2.5** ([14, 15, 16]). *Let  $f(\tau) = \prod_{\delta \mid N} \eta(\delta\tau)^{r_\delta}$  be an  $\eta$ -quotient with  $k = \frac{1}{2} \sum_{\delta \mid N} r_\delta \in \mathbb{Z}$  such that*

$$\sum_{\delta \mid N} \delta r_\delta \equiv 0 \pmod{24} \quad \text{and} \quad \sum_{\delta \mid N} \frac{N}{\delta} r_\delta \equiv 0 \pmod{24}.$$

*Then for all  $\begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \Gamma_0(N)$ ,*

$$f\left(\frac{a\tau + b}{c\tau + d}\right) = \left(\frac{(-1)^k \prod_{\delta \mid N} \delta^{r_\delta}}{d}\right) (c\tau + d)^k f(\tau).$$

**Lemma 2.6** ([13, 16]). *Let  $c, d$  and  $N$  be positive integers with  $d \mid N$  and  $\gcd(c, d) = 1$  and let  $f(\tau) = \prod_{\delta \mid N} \eta(\delta\tau)^{r_\delta}$  be an  $\eta$ -quotient satisfying the conditions of Lemma 2.5. Then the order of vanishing of  $f(\tau)$  at the cusp  $c/d$  is*

$$\frac{N}{24d \gcd(d, \frac{N}{d})} \sum_{\delta \mid N} \gcd(d, \delta)^2 \cdot \frac{r_\delta}{\delta}.$$

We next define, following Yang [20], generalized  $\eta$ -quotients with analogous properties on  $\Gamma_1(N)$ . These are functions of the form

$$f(\tau) = \prod_{1 \leq g \leq \lfloor N/2 \rfloor} \eta_{N,g}(\tau)^{r_g}$$

for some indexed set  $\{r_g \in \mathbb{Z} : 1 \leq g \leq \lfloor N/2 \rfloor\}$  of integers, where

$$\eta_{N,g}(\tau) = q^{NB(g/N)/2} \prod_{m=1}^{\infty} (1 - q^{N(m-1)+g})(1 - q^{Nm-g})$$

is the generalized Dedekind eta function with  $B(t) := t^2 - t + \frac{1}{6}$ . The following results give the transformation formula of  $\eta_{N,g}$  on  $\Gamma_0(N)$  and conditions for the modularity of a generalized  $\eta$ -quotient on  $\Gamma_1(N)$ .

**Lemma 2.7** ([20]). *The function  $\eta_{N,g}(\tau)$  satisfies  $\eta_{N,g+N}(\tau) = \eta_{N,-g}(\tau) = -\eta_{N,g}(\tau)$ . Moreover, let  $\gamma = \begin{bmatrix} a & b \\ cN & d \end{bmatrix} \in \Gamma_0(N)$  with  $c \neq 0$ . Then we have*

$$\eta_{N,g}(\gamma\tau) = \varepsilon(a, bN, c, d) e^{\pi i(g^2 ab/N - gb)} \eta_{N,ag}(\tau),$$

where

$$\varepsilon(a, b, c, d) = \begin{cases} e^{\pi i(bd(1-c^2)+c(a+d-3))/6}, & \text{if } c \equiv 1 \pmod{2}, \\ -ie^{\pi i(ac(1-d^2)+d(b-c+3))/6}, & \text{if } c \equiv 0 \pmod{2}. \end{cases}$$

**Lemma 2.8** ([20]). *Suppose  $f(\tau) = \prod_{1 \leq g \leq \lfloor N/2 \rfloor} \eta_{N,g}(\tau)^{r_g}$  is a generalized  $\eta$ -quotient such that*

$$\sum_{1 \leq g \leq \lfloor N/2 \rfloor} r_g \equiv 0 \pmod{12} \quad \text{and} \quad \sum_{1 \leq g \leq \lfloor N/2 \rfloor} gr_g \equiv 0 \pmod{2}.$$

*Then  $f(\tau)$  is a modular function on  $\Gamma(N)$ . Moreover, if in addition,  $f(\tau)$  satisfies*

$$\sum_{1 \leq g \leq \lfloor N/2 \rfloor} g^2 r_g \equiv 0 \pmod{2N}$$

*then  $f(\tau)$  is a modular function on  $\Gamma_1(N)$ .*

**Lemma 2.9** ([20]). *Let  $N$  be a positive integer and  $\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_2(\mathbb{Z})$ . Then the first term of the  $q$ -expansion of  $\eta_{N,g}(\gamma\tau)$  is  $\varepsilon q^\delta$ , where  $|\varepsilon| = 1$  and*

$$\delta = \frac{\gcd(c, N)^2}{2N} P_2\left(\frac{ag}{\gcd(c, N)}\right)$$

*with  $P_2(t) := B(\{t\})$  and  $\{t\}$  is the fractional part of  $t$ .*

Using Lemma 2.3, Corollary 2.4, and Lemma 2.9, we can compute the order of a generalized  $\eta$ -quotient at a cusp of  $\Gamma_1(N) \cap \Gamma_0(mN)$ .

### 3. MODULARITY OF $r(\tau)^a r(2\tau)^b$

In this section, we show that under a congruence condition on integers  $a$  and  $b$  with  $(a, b) \neq (0, 0)$ , there are finitely many functions of the form  $r(\tau)^a r(2\tau)^b$  generating the field  $A_0(\Gamma_1(10))$  of all modular functions on  $\Gamma_1(10)$ , partially extending the result of Lee and Park [12] that  $k(\tau)$  is one such function. To prove this, we use the fact that  $r(\tau) = \eta_{5,1}(\tau)/\eta_{5,2}(\tau)$ .

**Theorem 3.1.** *Let  $a$  and  $b$  be integers with  $(a, b) \neq (0, 0)$  such that  $5 \mid a - 3b$ . Then the function  $r(\tau)^a r(2\tau)^b$  is modular on  $\Gamma_1(10)$ . In addition, if  $r(\tau)^a r(2\tau)^b$  generates  $A_0(\Gamma_1(10))$ , then there are two such functions up to taking reciprocals, namely  $k(\tau)$  and  $l(\tau)$ .*

*Proof.* We write  $F(\tau) := r(\tau)^a r(2\tau)^b$  as a generalized  $\eta$ -quotient given by

$$F(\tau) = \eta_{10,1}(\tau)^a \eta_{10,2}(\tau)^{b-a} \eta_{10,3}(\tau)^{-a} \eta_{10,4}(\tau)^{a-b}.$$

By Lemma 2.8 and the given divisibility condition, we see that  $F(\tau)$  is modular on  $\Gamma_1(10)$ . Using Corollary 2.4(2), we choose the set of inequivalent cusps of  $\Gamma_1(10)$  to be  $S_{\Gamma_1(10)} = \{\infty, 0, 1/2, 1/3, 1/5, 1/6, 3/5, 3/10\}$ . We then calculate using Lemma 2.9 the order of  $F(\tau)$  at each element of  $S_{\Gamma_1(10)}$ , as shown in Table 1.

cusps $r$	$\infty$	$1/5$	$3/5$	$3/10$	$0, 1/2, 1/3, 1/6$
$\text{ord}_r F(\tau)$	$\frac{1}{5}(a+2b)$	$\frac{1}{5}(2a-b)$	$\frac{1}{5}(-2a+b)$	$\frac{1}{5}(-a-2b)$	0

TABLE 1. The orders of  $F(\tau)$  at the cusps of  $\Gamma_1(10)$

Observe that the orders of  $F(\tau)$  at the cusps  $\infty, 1/5, 3/5$  and  $3/10$  are integers, and that  $\text{ord}_{3/5} F(\tau) = -\text{ord}_{1/5} F(\tau)$  and  $\text{ord}_{3/10} F(\tau) = -\text{ord}_{\infty} F(\tau)$ . Since  $F(\tau)$  generates  $A_0(\Gamma_1(10))$ , exactly one of  $\text{ord}_{1/5} F(\tau)$  and  $\text{ord}_{\infty} F(\tau)$  is zero and the other is  $\pm 1$  (otherwise, the total degree of poles of  $F(\tau)$  would have been at least 2). We now proceed by doing casework:

- If  $0 = \text{ord}_{1/5} F(\tau) = \frac{1}{5}(2a-b)$  and  $\pm 1 = \text{ord}_{\infty} F(\tau) = \frac{1}{5}(a+2b)$ , then  $(a, b) \in \{(1, 2), (-1, -2)\}$ .
- If  $0 = \text{ord}_{\infty} F(\tau) = \frac{1}{5}(a+2b)$  and  $\pm 1 = \text{ord}_{1/5} F(\tau) = \frac{1}{5}(2a-b)$ , then  $(a, b) \in \{(-2, 1), (2, -1)\}$ .

Thus, we get four possible ordered pairs  $(a, b)$ , corresponding to the functions  $k(\tau), k(\tau)^{-1}, l(\tau)$  and  $l(\tau)^{-1}$ . The desired conclusion now follows.  $\square$

As a consequence of Theorem 3.1, we prove that the field  $A_0(\Gamma_0(10))$  of all modular functions on  $\Gamma_0(10)$  can be generated by a rational function of  $l(\tau)$ . We use the fact that  $\Gamma_0(10)$  is generated by  $\Gamma_1(10)$  and the matrix  $\gamma := \begin{bmatrix} 3 & -1 \\ 10 & -3 \end{bmatrix}$ .

**Theorem 3.2.** *The function  $l(\tau) - 1/l(\tau)$  generates  $A_0(\Gamma_0(10))$ .*

*Proof.* Since  $l(\gamma\tau) = -1/l(\tau)$  by Lemma 2.7,  $l(\tau) - 1/l(\tau) \in A_0(\Gamma_0(10))$ . We take  $S_{\Gamma_0(10)} = \{\infty, 0, 1/2, 1/5\}$  as the set of inequivalent cusps of  $\Gamma_0(10)$  by Corollary 2.4(1). Using Lemma 2.9, we compute  $\text{ord}_{1/5} l(\tau) = -1$  and  $\text{ord}_r l(\tau) = 0$  for any  $r \in S_{\Gamma_0(10)}$  with  $r \neq 1/5$ . We see that

$$\text{ord}_{1/5}(l(\tau) - 1/l(\tau)) = \min\{\text{ord}_{1/5} l(\tau), \text{ord}_{1/5} 1/l(\tau)\} = -1$$

and

$$\text{ord}_r(l(\tau) - 1/l(\tau)) \geq \min\{\text{ord}_r l(\tau), \text{ord}_r 1/l(\tau)\} = 0$$

for any  $r \in S_{\Gamma_0(10)}$  with  $r \neq 1/5$ . Thus, we obtain that  $l(\tau) - 1/l(\tau)$  has a simple pole only at  $1/5$  and holomorphic elsewhere, which yields the desired conclusion.  $\square$

We now consider modular functions on  $\Gamma_0(10)$  involving  $r(\tau)^a r(2\tau)^b$  for integers  $a$  and  $b$  with  $(a, b) \neq (0, 0)$ . The following results, which extend Theorem 3.2, imply that under the condition of Theorem 3.1, the functions  $r(\tau)^a r(2\tau)^b$  can be expressed explicitly in terms of  $l(\tau)$ .

**Theorem 3.3.** *Let  $a$  and  $b$  be integers with  $(a, b) \neq (0, 0)$  such that  $5 \mid a - 3b$ . Then the function  $f_{a,b}(\tau) := r(\tau)^a r(2\tau)^b + (-1)^{b-a} / (r(\tau)^a r(2\tau)^b)$  is modular on  $\Gamma_0(10)$ . In addition, the function  $(l(\tau) - 1/l(\tau))^{|a+2b|/5} f_{a,b}(\tau)$  is a polynomial in  $l(\tau) - 1/l(\tau)$  with rational coefficients.*



*Proof.* We know from Theorem 3.1 that  $F(\tau) := r(\tau)^a r(2\tau)^b$  is a modular function on  $\Gamma_1(10)$ . Since  $F(\gamma\tau) = (-1)^{b-a}/F(\tau)$  by Lemma 2.7, we see that  $f_{a,b}(\tau) = F(\tau) + F(\gamma\tau)$  is a modular function on  $\Gamma_0(10)$ . We again choose  $S_{\Gamma_0(10)} = \{\infty, 0, 1/2, 1/5\}$  as the set of inequivalent cusps of  $\Gamma_0(10)$  by Corollary 2.4(1). By Lemma 2.9, we find that  $\text{ord}_\infty F(\tau) = (a + 2b)/5$ ,  $\text{ord}_{1/5} F(\tau) = (2a - b)/5$  and  $\text{ord}_r F(\tau) = 0$  for  $r \in \{0, 1/2\}$ . Thus, we get

$$\begin{aligned}\text{ord}_\infty f_{a,b}(\tau) &= \min\{\text{ord}_\infty F(\tau), \text{ord}_\infty 1/F(\tau)\} = -|a + 2b|/5, \\ \text{ord}_{1/5} f_{a,b}(\tau) &= \min\{\text{ord}_{1/5} F(\tau), \text{ord}_{1/5} 1/F(\tau)\} = -|2a - b|/5,\end{aligned}$$

and

$$\text{ord}_r f_{a,b}(\tau) \geq \min\{\text{ord}_r F(\tau), \text{ord}_r 1/F(\tau)\} = 0,$$

for  $r \in \{0, 1/2\}$ . We now claim that  $l(\tau) - 1/l(\tau)$  has a zero at  $\infty$ . Indeed, we have the  $q$ -expansion of  $l(\tau)$  given by

$$(1) \quad l(\tau) = 1 + 2q - 4q^3 - 2q^4 + 6q^5 + 8q^6 - 4q^7 + O(q^8)$$

Observe that when  $\tau = \infty$ ,  $q = 0$  and  $l(\tau) = 1$ , which proves our claim. Consequently, we see that  $\text{ord}_\infty(l(\tau) - 1/l(\tau)) \geq 1$ . We deduce from Theorem 3.2 that the nonconstant function  $(l(\tau) - 1/l(\tau))^{|a+2b|/5} f_{a,b}(\tau)$  has a pole of order  $d := (|a + 2b| + |2a - b|)/5$  only at  $1/5$  and holomorphic elsewhere. Thus, by [21, Lem. 2], we obtain that  $(l(\tau) - 1/l(\tau))^{|a+2b|/5} f_{a,b}(\tau)$  is a polynomial in  $l(\tau) - 1/l(\tau)$  of degree  $d$ . Moreover, this polynomial has rational coefficients since the  $q$ -expansions of  $r(\tau)$  and  $l(\tau)$  are rational.  $\square$

**Corollary 3.4.** *We have the identities*

$$r(\tau)^5 = l^{-2} \left( \frac{l-1}{l+1} \right), \quad r(2\tau)^5 = l \left( \frac{l-1}{l+1} \right)^2,$$

where  $l := l(\tau)$ . Consequently, under the conditions of Theorem 3.3, we have

$$r(\tau)^a r(2\tau)^b = l^{(b-2a)/5} \left( \frac{l-1}{l+1} \right)^{(a+2b)/5}.$$

*Proof.* Using the notations of Theorem 3.3 and following its proof, we know that  $(l - 1/l)f_{5,0}(\tau)$  is a third-degree polynomial in  $l - 1/l$ . Writing

$$(l - 1/l)f_{5,0}(\tau) = \sum_{i=0}^3 a_i (l - 1/l)^i$$

for some constants  $a_0, a_1, a_2$  and  $a_3$ , and applying the  $q$ -expansions

$$r(\tau)^5 = q - 5q^2 + 15q^3 - 30q^4 + 40q^5 - 26q^6 + O(q^7)$$

and (1), we get  $(a_0, a_1, a_2, a_3) = (-4, -4, -2, -1)$  and

$$f_{5,0}(\tau) = r(\tau)^5 - \frac{1}{r(\tau)^5} = -\frac{(1-l+l^2+l^3)(-1+l+l^2+l^3)}{l^2(-1+l)(1+l)}.$$

Solving the above equation for  $r(\tau)^5$  and applying again the  $q$ -expansions of  $r(\tau)^5$  and  $l(\tau)$  yield the first identity. The identities involving  $r(2\tau)^5$  and  $r(\tau)^a r(2\tau)^b$  now follow from the first identity and the definition of  $l(\tau)$ .  $\square$

4. MODULAR EQUATIONS FOR  $l(\tau)$ 

We infer from Theorem 3.1 that  $l(\tau)$  generates  $A_0(\Gamma_1(10))$ . The following shows that we can define an affine model for the modular curve  $X(\Gamma_1(10) \cap \Gamma_0(10n))$  over  $\mathbb{Q}$  using  $l(\tau)$ . This affine model gives rise to the modular equation for  $l(\tau)$  of suitable level. We omit the proof as it is similar to that of [12, Lem. 3.2].

**Lemma 4.1.** *We have  $A_0(\Gamma_1(10) \cap \Gamma_0(10n))_{\mathbb{Q}} = \mathbb{Q}(l(\tau), l(n\tau))$  for all positive integers  $n$ .*

According to Table 1,  $l(\tau)$  has a simple pole at  $1/5$  and a simple zero at  $3/5$ . We use this to describe the behavior of  $l(\tau)$  at the cusps.

**Lemma 4.2.** *Let  $a, c, a', c'$  and  $n$  be integers with  $n$  positive. Then:*

- (1) *The modular function  $l(\tau)$  has a pole at  $a/c \in \mathbb{Q} \cup \{\infty\}$  if and only if  $\gcd(a, c) = 1, a \equiv \pm 1 \pmod{5}$  and  $c \equiv 5 \pmod{10}$ .*
- (2) *The modular function  $l(n\tau)$  has a pole at  $a'/c' \in \mathbb{Q} \cup \{\infty\}$  if and only if there exist integers  $a$  and  $c$  such that  $a/c = na'/c', \gcd(a, c) = 1, a \equiv \pm 1 \pmod{5}$  and  $c \equiv 5 \pmod{10}$ .*
- (3) *The modular function  $l(\tau)$  has a zero at  $a/c \in \mathbb{Q} \cup \{\infty\}$  if and only if  $\gcd(a, c) = 1, a \equiv \pm 3 \pmod{5}$  and  $c \equiv 5 \pmod{10}$ .*
- (4) *The modular function  $l(n\tau)$  has a zero at  $a'/c' \in \mathbb{Q} \cup \{\infty\}$  if and only if there exist integers  $a$  and  $c$  such that  $a/c = na'/c', \gcd(a, c) = 1, a \equiv \pm 3 \pmod{5}$  and  $c \equiv 5 \pmod{10}$ .*

*Proof.* Note that  $l(\tau)$  has a pole (resp., zero) at  $a/c \in \mathbb{Q} \cup \{\infty\}$  which is equivalent to  $1/5$  (resp.,  $3/5$ ) on  $\Gamma_1(10)$ . By Corollary 2.4(2),  $a/c$  is equivalent to  $1/5$  on  $\Gamma_1(10)$  if and only if  $\begin{bmatrix} a \\ c \end{bmatrix} \equiv \pm \begin{bmatrix} 1+5^m \\ 5 \end{bmatrix} \pmod{10}$  for some integer  $m$ . Thus, we get  $\gcd(a, c) = 1, a \equiv \pm 1 \pmod{5}$  and  $c \equiv 5 \pmod{10}$ , proving (1). On the other hand,  $a/c$  is equivalent to  $3/5$  on  $\Gamma_1(10)$  if and only if  $\begin{bmatrix} a \\ c \end{bmatrix} \equiv \pm \begin{bmatrix} 3+5^m \\ 5 \end{bmatrix} \pmod{10}$  for some integer  $m$ . This yields  $\gcd(a, c) = 1, a \equiv \pm 3 \pmod{5}$  and  $c \equiv 5 \pmod{10}$ , proving (3). Statement (2) (resp. (4)) now follows from (1) (resp. (3)).  $\square$

To obtain explicitly modular equations for  $l(\tau)$  of any level, we present the following result by Ishida and Ishii [11], which gives information on the coefficients of the modular equation between two functions  $f_1(\tau)$  and  $f_2(\tau)$  that generate the field  $A_0(\Gamma')$  for some congruence subgroup  $\Gamma'$  of  $SL_2(\mathbb{Z})$ .

**Proposition 4.3** ([3, 11]). *Let  $\Gamma'$  be a congruence subgroup of  $SL_2(\mathbb{Z})$ , and  $f_1(\tau)$  and  $f_2(\tau)$  be nonconstant functions such that  $A_0(\Gamma') = \mathbb{C}(f_1(\tau), f_2(\tau))$ . For  $k \in \{1, 2\}$ , let  $d_k$  be the total degree of poles of  $f_k(\tau)$ . Let*

$$F(X, Y) = \sum_{\substack{0 \leq i < d_2 \\ 0 \leq j \leq d_1}} C_{i,j} X^i Y^j \in \mathbb{C}[X, Y]$$

*satisfy  $F(f_1(\tau), f_2(\tau)) = 0$ . Let  $S_{\Gamma'}$  be the set of all the inequivalent cusps of  $\Gamma'$ , and for  $k \in \{1, 2\}$ , define*

$$\begin{aligned} S_{k,0} &= \{r \in S_{\Gamma'} : f_k(\tau) \text{ has a zero at } r\}, \\ S_{k,\infty} &= \{r \in S_{\Gamma'} : f_k(\tau) \text{ has a pole at } r\}. \end{aligned}$$

*Further let*

$$a = - \sum_{r \in S_{1,\infty} \cap S_{2,0}} \text{ord}_r f_1(\tau) \quad \text{and} \quad b = \sum_{r \in S_{1,0} \cap S_{2,0}} \text{ord}_r f_1(\tau)$$

*with the convention that  $a = 0$  (resp.  $b = 0$ ) whenever  $S_{1,\infty} \cap S_{2,0}$  (resp.  $S_{1,0} \cap S_{2,0}$ ) is empty. Then*



- (1)  $C_{d_2,a} \neq 0$  and if, in addition,  $S_{1,\infty} \subseteq S_{2,\infty} \cup S_{2,0}$ , then  $C_{d_2,j} = 0$  for all  $j \neq a$ ;  
 (2)  $C_{0,b} \neq 0$  and if, in addition,  $S_{1,0} \subseteq S_{2,\infty} \cup S_{2,0}$ , then  $C_{0,j} = 0$  for all  $j \neq b$ .

By interchanging the roles of  $f_1(\tau)$  and  $f_2(\tau)$ , one may obtain properties analogous to (1) - (2).

We apply Lemma 4.2 and Proposition 4.3 to deduce the following result, which asserts that modular equations for  $l(\tau)$  of any level always exist, which serve as affine models for  $X(\Gamma_1(10) \cap \Gamma_0(10n))$ .

**Theorem 4.4.** *One can explicitly obtain modular equation for  $l(\tau)$  of level  $n$  for all positive integers  $n$ .*

*Proof.* The proof is similar to that of [12, Thm. 1.1](1), so we omit the details. We instead note that if  $L_n(l(\tau), l(n\tau)) = 0$  is the modular equation for  $l(\tau)$  of level  $n$  for some polynomial  $L_n(X, Y) \in \mathbb{C}[X, Y]$ , then  $L_n(X, Y)$  is irreducible as a polynomial of  $X$  (resp.  $Y$ ) over  $\mathbb{C}(Y)$  (resp.  $\mathbb{C}(X)$ ), in view of [3, Lem. 3.1].  $\square$

We now use Theorem 4.4 to derive the modular equations for  $l(\tau)$  of levels two and five, and find which coefficients of the modular equations for  $l(\tau)$  of odd prime level  $p \neq 5$  are nonzero.

**Theorem 4.5** (modular equation of level 2). *We have*

$$1 + l(\tau) - 2l(\tau)l(2\tau) - l(\tau)l(2\tau)^2 + l(\tau)^2l(2\tau)^2 = 0.$$

*Proof.* In view of Lemmata 2.2 and 4.2, we work on the congruence subgroup  $\Gamma_1(10) \cap \Gamma_0(20)$  and the following cusps:  $1/5, 3/5, 1/10$  and  $3/10$ . Recall that

$$l(\tau) = \frac{\eta_{10,2}(\tau)^3 \eta_{10,3}(\tau)^2}{\eta_{10,1}(\tau)^2 \eta_{10,4}(\tau)^3}$$

is a generalized  $\eta$ -quotient. Using Lemmata 2.3 and 2.9, we know that  $l(\tau)$  has a double pole at  $1/5$  and a double zero at  $3/5$ . We now compute the order of  $l(2\tau)$  at  $3/10$  (which is a zero) as follows. Since

$$l\left(2 \begin{bmatrix} 3 & -1 \\ 10 & -3 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 6 & -2 \\ 10 & -3 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 3 & -2 \\ 5 & -3 \end{bmatrix} (2\tau)\right),$$

by Lemma 2.9, the first term of the  $q$ -expansion of  $l(2[\frac{3}{10} \ -1]{\tau})$  is  $\varepsilon q^\delta$ , where  $|\varepsilon| = 1$  and

$$\delta = 2 \cdot \frac{25}{20} \left[ -2P_2\left(\frac{3}{5}\right) + 3P_2\left(\frac{6}{5}\right) + 2P_2\left(\frac{9}{5}\right) - 3P_2\left(\frac{12}{5}\right) \right] = 1.$$

As the width of the cusp  $3/10$  is 1 by Lemma 2.3, we see that  $\text{ord}_{3/10} l(2\tau) = 1 \cdot 1 = 1$ , so  $l(2\tau)$  has a simple zero at  $3/10$ . Similarly,  $l(2\tau)$  has another simple zero at  $1/5$ , and two simple poles at  $1/10$  and  $3/5$ . We deduce that the total degree of poles of both  $l(\tau)$  and  $l(2\tau)$  are 2, so by Proposition 4.3, there is a polynomial

$$L_2(X, Y) = \sum_{0 \leq i, j \leq 2} C_{i,j} X^i Y^j \in \mathbb{C}[X, Y]$$

such that  $L_2(l(\tau), l(2\tau)) = 0$ . In addition, we get that  $C_{2,2} \neq 0, C_{0,0} \neq 0$  and  $C_{2,0} = C_{2,1} = C_{0,1} = C_{0,2} = 0$ . Using the  $q$ -expansion (1) of  $l(\tau)$ , we may take  $C_{2,2} = 1$ . Hence, we arrive at  $L_2(X, Y) = 1 + X - 2XY - XY^2 + X^2Y^2$  and the desired identity follows.  $\square$

**Theorem 4.6** (modular equation of level 5). *We have*

$$(l(\tau)^5 - l(5\tau))(l(5\tau)^2 - l(5\tau) - 1)^2 = -5l(\tau)l(5\tau)P(l(\tau), l(5\tau)),$$

where  $P(X, Y) := (Y^2 - Y - 1)YX^3 + (3Y^2 + 2Y - 3)X^2 - (3Y^2 + 2Y - 3)YX + Y^2 - Y - 1$ .

*Proof.* Appealing to Lemmata 2.2 and 4.2, we focus on the congruence subgroup  $\Gamma_1(10) \cap \Gamma_0(50)$  and the following cusps:

$$1/5, 2/5, 3/5, 4/5, 1/15, 2/15, 4/15, 8/15, 1/25, 3/25.$$

Applying Lemmata 2.3 and 2.9, we see that  $l(\tau)$  has five simple poles at  $1/5, 4/5, 1/15, 4/15$  and  $1/25$ , and five simple zeros at  $2/5, 3/5, 2/15, 8/15$  and  $3/25$ . We now compute the order of  $l(5\tau)$  at the cusp  $1/25$  (which is a pole) as follows. Observe that

$$l\left(5 \begin{bmatrix} 1 & 0 \\ 25 & 1 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 5 & 0 \\ 25 & 1 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix} (5\tau)\right),$$

so by Lemma 2.9, the  $q$ -expansion of  $l(5[\frac{1}{25} \ 0] \tau)$  starts with  $\varepsilon q^\delta$  with  $|\varepsilon| = 1$  and

$$\delta = 5 \cdot \frac{25}{20} \left[ -2P_2\left(\frac{1}{5}\right) + 3P_2\left(\frac{2}{5}\right) + 2P_2\left(\frac{3}{5}\right) - 3P_2\left(\frac{4}{5}\right) \right] = -\frac{5}{2}.$$

Since the width of the cusp  $1/25$  is 2 by Lemma 2.3, we see that  $\text{ord}_{1/25} l(5\tau) = 2 \cdot -5/2 = -5$ . Similarly, we get that  $\text{ord}_{3/25} l(5\tau) = 5$ , so the total degree of poles of both  $l(\tau)$  and  $l(5\tau)$  are 5. Thus, by Proposition 4.3, there is a polynomial

$$L_5(X, Y) = \sum_{0 \leq i, j \leq 5} C_{i, j} X^i Y^j \in \mathbb{C}[X, Y]$$

such that  $L_5(l(\tau), l(5\tau)) = 0$ . In addition, we get that  $C_{5,0} \neq 0$  and  $C_{0,1} \neq 0$ . Switching the roles of  $l(\tau)$  and  $l(5\tau)$  yields  $C_{0,5} \neq 0, C_{j,5} = 0$  for  $j \in \{1, 2, 3, 4, 5\}$  and  $C_{k,0} = 0$  for  $k \in \{0, 1, 2, 3, 4\}$ . Using the  $q$ -expansion (1) of  $l(\tau)$  leads us to

$$L_5(X, Y) = (X^5 - Y)(Y^2 - Y - 1)^2 + 5XYP(X, Y)$$

and the desired identity follows.  $\square$

**Theorem 4.7.** *Let  $0 = L_p(X, Y) = \sum_{0 \leq i, j \leq p+1} C_{i, j} X^i Y^j \in \mathbb{Q}[X, Y]$  be the modular equation for  $l(\tau)$  of odd prime level  $p \neq 5$ .*

- (1) *If  $p \equiv \pm 1 \pmod{10}$ , then  $C_{p+1,0} \neq 0, C_{0,p+1} \neq 0$ , and  $C_{i,p+1} = C_{p+1,i} = 0$  for  $i \in \{1, \dots, p+1\}$  and  $C_{0,j} = C_{j,0} = 0$  for  $j \in \{0, \dots, p\}$ .*
- (2) *If  $p \equiv \pm 3 \pmod{10}$ , then  $C_{p+1,p} \neq 0, C_{0,1} \neq 0, C_{1,p+1} \neq 0, C_{p,0} \neq 0$ , and  $C_{p+1,i} = C_{i,0} = 0$  for  $i \in \{0, \dots, p-1, p+1\}$  and  $C_{0,j} = C_{j,p+1} = 0$  for  $j \in \{0, 2, \dots, p+1\}$ .*

*Proof.* We adapt the notations of Lemma 2.2, and in view of Lemma 4.2, we need to determine the sets  $S_c$  and  $A_c$  for  $c \in \{5, 5p\}$ . We set  $I := \{0, 1, \dots, p-1\}$  and choose a unique integer  $k_0 \in I$  such that  $p \mid 1 + 10k_0$ . Then

$$\Delta = \{\pm(1 + 10k) \in (\mathbb{Z}/10p\mathbb{Z})^\times : k \in I \setminus \{k_0\}\}.$$

For each  $a \in I \setminus \{k_0\}$ , choose  $b \in I \setminus \{k_0\}$  such that  $p \mid 1 + 5(a + b)$ . Then  $\pi_{2p}(1 + 10a) = \pi_{2p}(-1 - 10b)$ , so  $\pi_{2p}(\Delta)$  has exactly  $p - 1$  distinct elements modulo  $2p$ . We see that  $\pi_{2p}(\Delta) = (\mathbb{Z}/2p\mathbb{Z})^\times$  and thus,  $S_5 = \{1\}$ . On the other hand, we choose  $d \in I \setminus \{k_0\}$  such that  $p \mid 1 + 5d$ . Then  $\pi_{2p}(-1 - 10d) = 1$ , so  $\Delta \cap \ker \pi_{2p} = \{1, -1 - 10d\}$  and  $\pi_5(\Delta \cap \ker \pi_{2p}) = \{\pm 1\}$ . Thus, we take  $A_5 = \{1, 2\}$ . We next observe that  $\pi_2(\Delta) = \{1\} = (\mathbb{Z}/2\mathbb{Z})^\times$ , so we have  $S_{5p} = \{1\}$  and  $\pi_{5p}(\Delta \cap \ker \pi_2) = \pi_{5p}(\Delta) \subset (\mathbb{Z}/5p\mathbb{Z})^\times$ . We claim that  $\pi_{5p}(\Delta)$  has  $2(p - 1)$  distinct elements modulo  $5p$ ; otherwise, we would have  $1 + 10r \equiv \pm(1 + 10s) \pmod{5p}$  for some distinct  $r, s \in I \setminus \{k_0\}$ , and all cases lead to a contradiction. As  $(\mathbb{Z}/5p\mathbb{Z})^\times$  has  $4(p - 1)$  elements, we take  $A_{5p} = \{1, 2\}$ .

We see that  $l(\tau)$  and  $l(p\tau)$  has poles and zeros at the following cusps of  $\Gamma_1(10) \cap \Gamma_0(10p)$ :  $1/5, 2/5, 1/5p$  and  $2/5p$ . Applying Lemma 2.9, we get  $\text{ord}_{1/5} l(\tau) = -p = -\text{ord}_{2/5} l(\tau)$  and

$\text{ord}_{1/5p}l(\tau) = -1 = -\text{ord}_{2/5p}l(\tau)$ . We next compute  $\text{ord}_{1/5p}l(p\tau)$  and  $\text{ord}_{2/5p}l(p\tau)$ . We note that

$$l\left(p \begin{bmatrix} 1 & 0 \\ 5p & 1 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} p & 0 \\ 5p & 1 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix} (p\tau)\right),$$

so by Lemma 2.9, the first term of the  $q$ -expansion of  $l(p[\frac{1}{5p} \ 0] \tau)$  is  $\varepsilon q^\delta$  with  $|\varepsilon| = 1$  and

$$\delta = p \cdot \frac{25}{20} \left[ -2P_2\left(\frac{1}{5}\right) + 3P_2\left(\frac{2}{5}\right) + 2P_2\left(\frac{3}{5}\right) - 3P_2\left(\frac{4}{5}\right) \right] = -\frac{p}{2}.$$

Since the width of the cusp  $1/5p$  is 2 by Lemma 2.3, we get  $\text{ord}_{1/5p}l(p\tau) = 2 \cdot -p/2 = -p$ . On the other hand, we choose integers  $u$  and  $v$  such that  $2v - 5pu = 1$ . We have

$$l\left(p \begin{bmatrix} 2 & u \\ 5p & v \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 2p & up \\ 5p & v \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 2 & up \\ 5 & v \end{bmatrix} (p\tau)\right),$$

so Lemma 2.9 shows that the  $q$ -expansion of  $l(p[\frac{2}{5p} \ \frac{u}{v}] \tau)$  starts with  $\varepsilon q^\delta$  with  $|\varepsilon| = 1$  and

$$\delta = p \cdot \frac{25}{20} \left[ -2P_2\left(\frac{2}{5}\right) + 3P_2\left(\frac{4}{5}\right) + 2P_2\left(\frac{6}{5}\right) - 3P_2\left(\frac{8}{5}\right) \right] = \frac{p}{2}.$$

As the cusp  $2/5p$  has width 2 by Lemma 2.3, we obtain  $\text{ord}_{2/5p}l(p\tau) = 2 \cdot p/2 = p$ .

We now compute  $\text{ord}_{1/5}l(p\tau)$  and  $\text{ord}_{2/5}l(p\tau)$ . We choose an integer  $g$  with  $pg \equiv 1 \pmod{5}$ , so that

$$l\left(p \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} p & 0 \\ 5 & 1 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} p & (pg-1)/5 \\ 5 & g \end{bmatrix} \frac{\tau + (1-pg)/5}{p}\right),$$

By Lemma 2.9, we deduce that the  $q$ -expansion of  $l(p[\frac{1}{5} \ 0] \tau)$  starts with  $\varepsilon q^{\delta_p}$  with  $|\varepsilon| = 1$  and

$$\delta_p := \frac{1}{p} \cdot \frac{25}{20} \left[ -2P_2\left(\frac{p}{5}\right) + 3P_2\left(\frac{2p}{5}\right) + 2P_2\left(\frac{3p}{5}\right) - 3P_2\left(\frac{4p}{5}\right) \right].$$

Since the width of  $1/5$  is  $2p$  by Lemma 2.3, we see that

$$\text{ord}_{1/5}l(p\tau) = 2p \cdot \delta_p = \begin{cases} -1 & \text{if } p \equiv \pm 1 \pmod{10}, \\ 1 & \text{if } p \equiv \pm 3 \pmod{10}. \end{cases}$$

We next take an integer  $h$  with  $2ph \equiv 1 \pmod{5}$ , so that

$$l\left(p \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 2p & p \\ 5 & 3 \end{bmatrix} \tau\right) = l\left(\begin{bmatrix} 2p & (2ph-1)/5 \\ 5 & h \end{bmatrix} \frac{\tau + (3-ph)/5}{p}\right).$$

Using Lemma 2.9, the first term of the  $q$ -expansion of  $l(p[\frac{2}{5} \ \frac{3}{1}] \tau)$  is  $\varepsilon q^{\delta_p}$  with  $|\varepsilon| = 1$  and

$$\delta_p := \frac{1}{p} \cdot \frac{25}{20} \left[ -2P_2\left(\frac{2p}{5}\right) + 3P_2\left(\frac{4p}{5}\right) + 2P_2\left(\frac{6p}{5}\right) - 3P_2\left(\frac{8p}{5}\right) \right].$$

Since the width of  $2/5$  is  $2p$  by Lemma 2.3, we arrive at

$$\text{ord}_{2/5}l(p\tau) = 2p \cdot \delta_p = \begin{cases} 1 & \text{if } p \equiv \pm 1 \pmod{10}, \\ -1 & \text{if } p \equiv \pm 3 \pmod{10}. \end{cases}$$

Using the notations of Proposition 4.3 with  $f_1(\tau) = l(\tau)$  and  $f_2(\tau) = l(p\tau)$ , we obtain  $S_{1,\infty} = \{1/5, 1/5p\}$ ,  $S_{1,0} = \{2/5, 2/5p\}$  and

$$S_{2,\infty} = \begin{cases} \{1/5, 1/5p\} & \text{if } p \equiv \pm 1 \pmod{10}, \\ \{2/5, 1/5p\} & \text{if } p \equiv \pm 3 \pmod{10}, \end{cases}$$

$$S_{2,0} = \begin{cases} \{2/5, 2/5p\} & \text{if } p \equiv \pm 1 \pmod{10}, \\ \{1/5, 2/5p\} & \text{if } p \equiv \pm 3 \pmod{10}, \end{cases}$$

so that  $S_{1,\infty} \cup S_{1,0} = S_{2,\infty} \cup S_{2,0}$ . In addition, we deduce that the total number of poles of both  $l(\tau)$  and  $l(p\tau)$  is  $p + 1$ , so there is a polynomial

$$L_p(X, Y) = \sum_{0 \leq i, j \leq p+1} C_{i,j} X^i Y^j \in \mathbb{Q}[X, Y]$$

with  $L_p(l(\tau), l(p\tau)) = 0$ . We find that if  $p \equiv \pm 1 \pmod{10}$ , then  $C_{p+1,0} \neq 0$  and  $C_{0,p+1} \neq 0$ , so that  $C_{p+1,i} = 0$  for  $i \in \{1, \dots, p+1\}$  and  $C_{0,j} = 0$  for  $j \in \{0, \dots, p\}$ . Switching the roles of  $l(\tau)$  and  $l(p\tau)$  yields  $C_{i,p+1} = 0$  for  $i \in \{1, \dots, p+1\}$  and  $C_{j,0} = 0$  for  $j \in \{0, \dots, p\}$ . On the other hand, if  $p \equiv \pm 3 \pmod{10}$ , then  $C_{p+1,p} \neq 0$  and  $C_{0,1} \neq 0$ , implying that  $C_{p+1,i} = 0$  for  $i \in \{0, \dots, p-1, p+1\}$  and  $C_{0,j} = 0$  for  $j \in \{0, 2, \dots, p+1\}$ . Swapping the roles of  $l(\tau)$  and  $l(p\tau)$  gives  $C_{1,p+1} \neq 0$ ,  $C_{p,0} \neq 0$ ,  $C_{i,0} = 0$  for  $i \in \{0, \dots, p-1, p+1\}$  and  $C_{j,p+1} = 0$  for  $j \in \{0, 2, \dots, p+1\}$ .  $\square$

We give the modular equations  $L_n(X, Y)$  for  $l(\tau)$  of level  $n$  with  $n \in \{2, 4, 5, 6, 13\}$  as shown in Table 2. On the other hand, we verify via *Mathematica* that the modular equations for  $l(\tau)$  of prime levels  $p \in \{3, 7, 11, 19\}$  are the same as those for  $k(\tau)$ , which appeared as a supplementary material (Appendix A) to [12], so we will not present them here and instead remark that they satisfy the Kronecker congruence depending on the values of  $p$  modulo 10.

We now consider modular equations for  $l(\tau)$  of levels  $n$  coprime to 10. For any integer  $a$  coprime to 10, we choose a matrix  $\sigma_a \in \mathrm{SL}_2(\mathbb{Z})$  such that  $\sigma_a \equiv \begin{bmatrix} a^{-1} & 0 \\ 0 & a \end{bmatrix} \pmod{10}$ . We take

$$\sigma_{\pm 1} = \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \sigma_{\pm 3} = \pm \begin{bmatrix} -3 & -10 \\ 10 & 33 \end{bmatrix}.$$

We know that  $l(\sigma_{\pm 1}\tau) = l(\tau)$  and  $l(\sigma_{\pm 3}\tau) = -1/l(\tau)$  by Lemma 2.7. Moreover, by [17, Prop. 3.36], we have the disjoint union

$$\Gamma_1(10) \begin{bmatrix} 1 & 0 \\ 0 & n \end{bmatrix} \Gamma_1(10) = \bigsqcup_{0 < a | n} \bigsqcup_{\substack{0 \leq b < n/a \\ \gcd(a, b, n/a) = 1}} \Gamma_1(10) \sigma_a \begin{bmatrix} a & b \\ 0 & n/a \end{bmatrix}$$

and  $\Gamma_1(10) \setminus \Gamma_1(10) \begin{bmatrix} 1 & 0 \\ 0 & n \end{bmatrix} \Gamma_1(10)$  has  $\psi(n) := n \prod_{p|n} (1 + 1/p)$  elements. We now define the polynomial

$$\Phi_n(X, \tau) := \prod_{0 < a | n} \prod_{\substack{0 \leq b < n/a \\ \gcd(a, b, n/a) = 1}} (X - (l \circ \alpha_{a,b})(\tau))$$

where  $\alpha_{a,b} := \sigma_a \begin{bmatrix} a & b \\ 0 & n/a \end{bmatrix}$ . Then the coefficients of  $\Phi_n(X, \tau)$  are elementary symmetric functions of  $l \circ \alpha_{a,b}$ , so these are invariant under the action of  $\Gamma_1(10)$  and thus are in  $A_0(\Gamma_1(10)) = \mathbb{C}(l(\tau))$ . Thus, we may write  $\Phi_n(X, \tau)$  as  $\Phi_n(X, l(\tau)) \in \mathbb{C}(l(\tau))[X]$ .

$n$	$L_n(X, Y)$
2	$1 + X - 2XY - XY^2 + X^2Y^2$
4	$-X - X^2 + X^3 + X^4 + 4XY + 4X^2Y - 8X^3Y + 2XY^2 + 8X^2Y^2 - 2X^3Y^2 - 8XY^3 - 4X^2Y^3 + 4X^3Y^3 + Y^4 - XY^4 - X^2Y^4 + X^3Y^4$
5	$(X^5 - Y)(Y^2 - Y - 1)^2 + 5XY(1 - 3X^2 + Y + 3XY + 2X^2Y - X^3Y - Y^2 - 2XY^2 + 3X^2Y^2 - X^3Y^2 - 3XY^3 + X^3Y^3)$
6	$-X - 2X^2 + 2X^4 + X^5 + 6XY - 12X^3Y + 8X^4Y - 2X^5Y - 3XY^2 + 33X^2Y^2 + 27X^3Y^2 - 53X^4Y^2 - 13X^5Y^2 + 3X^6Y^2 + 5X^7Y^2 + X^8Y^2 - 22XY^3 + 24X^2Y^3 + 96X^3Y^3 - 40X^4Y^3 - 78X^5Y^3 + 24X^6Y^3 + 12X^7Y^3 + 9XY^4 - 30X^2Y^4 - 39X^3Y^4 + 92X^4Y^4 + 39X^5Y^4 - 30X^6Y^4 - 9X^7Y^4 + 12XY^5 - 24X^2Y^5 - 78X^3Y^5 + 40X^4Y^5 + 96X^5Y^5 - 24X^6Y^5 - 22X^7Y^5 + Y^6 - 5XY^6 + 3X^2Y^6 + 13X^3Y^6 - 53X^4Y^6 - 27X^5Y^6 + 33X^6Y^6 + 3X^7Y^6 - 2X^3Y^7 - 8X^4Y^7 - 12X^5Y^7 + 6X^7Y^7 - X^3Y^8 + 2X^4Y^8 - 2X^6Y^8 + X^7Y^8$
13	$(X^{13} - Y)(XY^{13} + 1) + 13XY(1 - 4X + X^2 + 26X^3 - 33X^4 - 57X^5 + 88X^6 + 53X^7 - 77X^8 - 18X^9 + 21X^{10} + X^{11} - X^{12} + Y + 6XY - 88X^2Y + 76X^3Y + 580X^4Y - 214X^5Y - 990X^6Y + 226X^7Y + 608X^8Y - 100X^9Y - 110X^{10}Y + 6X^{11}Y + 4X^{12}Y - 21Y^2 + 110XY^2 + 178X^2Y^2 - 1147X^3Y^2 - 629X^4Y^2 + 3003X^5Y^2 + 396X^6Y^2 - 2915X^7Y^2 + 255X^8Y^2 + 1053X^9Y^2 - 178X^{10}Y^2 - 88X^{11}Y^2 - X^{12}Y^2 - 18Y^3 - 100XY^3 + 1053X^2Y^3 + 114X^3Y^3 - 6447X^4Y^3 - 586X^5Y^3 + 11132X^6Y^3 + 382X^7Y^3 - 6808X^8Y^3 + 114X^9Y^3 + 1147X^{10}Y^3 + 76X^{11}Y^3 - 26X^{12}Y^3 + 77Y^4 - 608XY^4 - 255X^2Y^4 + 6808X^3Y^4 + 1291X^4Y^4 - 18366X^5Y^4 - 484X^6Y^4 + 18077X^7Y^4 - 1291X^8Y^4 - 6447X^9Y^4 + 629X^{10}Y^4 + 580X^{11}Y^4 + 33X^{12}Y^4 + 53Y^5 + 226XY^5 - 2915X^2Y^5 + 382X^3Y^5 + 18077X^4Y^5 + 96X^5Y^5 - 30602X^6Y^5 + 96X^7Y^5 + 18366X^8Y^5 - 586X^9Y^5 - 3003X^{10}Y^5 - 214X^{11}Y^5 + 57X^{12}Y^5 - 88Y^6 + 990XY^6 - 396X^2Y^6 - 11132X^3Y^6 + 484X^4Y^6 + 30602X^5Y^6 - 30602X^7Y^6 + 484X^8Y^6 + 11132X^9Y^6 - 396X^{10}Y^6 - 990X^{11}Y^6 - 88X^{12}Y^6 - 57Y^7 - 214XY^7 + 3003X^2Y^7 - 586X^3Y^7 - 18366X^4Y^7 + 96X^5Y^7 + 30602X^6Y^7 + 96X^7Y^7 - 18077X^8Y^7 + 382X^9Y^7 + 2915X^{10}Y^7 + 226X^{11}Y^7 - 53X^{12}Y^7 + 33Y^8 - 580XY^8 + 629X^2Y^8 + 6447X^3Y^8 - 1291X^4Y^8 - 18077X^5Y^8 - 484X^6Y^8 + 18366X^7Y^8 + 1291X^8Y^8 - 6808X^9Y^8 - 255X^{10}Y^8 + 608X^{11}Y^8 + 77X^{12}Y^8 + 26Y^9 + 76XY^9 - 1147X^2Y^9 + 114X^3Y^9 + 6808X^4Y^9 + 382X^5Y^9 - 11132X^6Y^9 - 586X^7Y^9 + 6447X^8Y^9 + 114X^9Y^9 - 1053X^{10}Y^9 - 100X^{11}Y^9 + 18X^{12}Y^9 - Y^{10} + 88XY^{10} - 178X^2Y^{10} - 1053X^3Y^{10} + 255X^4Y^{10} + 2915X^5Y^{10} + 396X^6Y^{10} - 3003X^7Y^{10} - 629X^8Y^{10} + 1147X^9Y^{10} + 178X^{10}Y^{10} - 110X^{11}Y^{10} - 21X^{12}Y^{10} - 4Y^{11} + 6XY^{11} + 110X^2Y^{11} - 100X^3Y^{11} - 608X^4Y^{11} + 226X^5Y^{11} + 990X^6Y^{11} - 214X^7Y^{11} - 580X^8Y^{11} + 76X^9Y^{11} + 88X^{10}Y^{11} + 6X^{11}Y^{11} - X^{12}Y^{11} - Y^{12} - XY^{12} + 21X^2Y^{12} + 18X^3Y^{12} - 77X^4Y^{12} - 53X^5Y^{12} + 88X^6Y^{12} + 57X^7Y^{12} - 33X^8Y^{12} - 26X^9Y^{12} + X^{10}Y^{12} + 4X^{11}Y^{12} + X^{12}Y^{12})$

TABLE 2. Modular equations  $L_n(X, Y) = 0$  for  $l(\tau)$  of level  $n \in \{2, 4, 5, 6, 13\}$

Let  $S_{n,\infty}$  (resp.  $S_{n,0}$ ) be the set of inequivalent cusps of  $\Gamma_1(10) \cap \Gamma_0(10n)$  for which  $l(n\tau)$  has a pole (resp. a zero). We define

$$r_n := - \sum_{\tau \in S_{1,\infty} \cap S_{n,0}} \text{ord}_\tau l(\tau),$$

with the convention that  $r_n = 0$  if  $S_{1,\infty} \cap S_{n,0}$  is empty, and set

$$L_n(X, l(\tau)) = l(\tau)^{r_n} \Phi_n(X, l(\tau)).$$

The following result gives the properties of the polynomial  $L_n(X, l(\tau))$  when  $\gcd(n, 10) = 1$ . We omit the proof as it is similar to that of [12, Thm. 1.1]. We note that  $L_n(l(\frac{\tau}{n}), l(\tau)) = 0$ , which can be seen as the modular equation for  $l(\tau)$  of level  $n$ .

**Theorem 4.8.** *Let  $n$  be a positive integer coprime to 10 and let  $L_n(X, Y)$  be the polynomial as above. Then*

- (1)  $L_n(X, Y) \in \mathbb{Z}[X, Y]$  and  $\deg_X L_n(X, Y) = \psi(n)$ .
- (2)  $L_n(X, Y)$  is irreducible both as a polynomial in  $X$  over  $\mathbb{C}(Y)$  and as a polynomial in  $Y$  over  $\mathbb{C}(X)$ .
- (3) We have

$$L_n(X, Y) = \begin{cases} L_n(Y, X) & \text{if } n \equiv \pm 1 \pmod{10}, \\ Y^{\psi(n)} L_n(-Y^{-1}, X) & \text{if } n \equiv \pm 3 \pmod{10}. \end{cases}$$

- (4) (Kronecker congruence) If  $p \neq 5$  is an odd prime, then

$$L_p(X, Y) \equiv \begin{cases} (X^p - Y)(X - Y^p) \pmod{p\mathbb{Z}[X, Y]} & \text{if } p \equiv \pm 1 \pmod{10}, \\ (X^p - Y)(XY^p + 1) \pmod{p\mathbb{Z}[X, Y]} & \text{if } p \equiv \pm 3 \pmod{10}. \end{cases}$$

## 5. ARITHMETIC OF $l(\tau)$ AND EVALUATION OF $r(\tau)^a r(2\tau)^b$

We study in this section some arithmetic properties of  $l(\tau)$  at imaginary quadratic points  $\tau \in \mathbb{H}$ , which will be useful in evaluating the functions of the form  $r(\tau)^a r(2\tau)^b$  satisfying the conditions of Theorem 3.1. We first state the following result of Cho and Koo [4] about generating ray class fields by a singular value of a modular function on some congruence subgroup between  $\Gamma(N)$  and  $\Gamma_1(N)$ .

**Lemma 5.1** ([4]). *Let  $K$  be an imaginary quadratic field with discriminant  $d_K$  and  $\tau \in K \cap \mathbb{H}$  be a root of a primitive equation  $aX^2 + bX + c = 0$  such that  $b^2 - 4ac = d_K$  and  $a, b, c \in \mathbb{Z}$ . Let  $\Gamma'$  be a congruence subgroup such that  $\Gamma(N) \subset \Gamma' \subset \Gamma_1(N)$ . Suppose that  $\gcd(a, N) = 1$ . Then the field generated over  $K$  by all values of  $h(\tau)$ , where  $h \in A_0(\Gamma')_{\mathbb{Q}}$  is defined and finite at  $\tau$ , is the ray class field  $K_{(N)}$  modulo  $N$  over  $K$ .*

We now apply Lemma 5.1 to prove the following.

**Theorem 5.2.** *Let  $K$  be an imaginary quadratic field with discriminant  $d_K$  and  $\tau \in K \cap \mathbb{H}$  be a root of the primitive polynomial  $aX^2 + bX + c \in \mathbb{Z}[X]$  such that  $b^2 - 4ac = d_K$  and  $\gcd(a, 10) = 1$ . Then  $K(l(\tau))$  is the ray class field modulo 10 over  $K$ .*

*Proof.* In view of Theorem 3.1, we have that  $A_0(\Gamma_1(10))_{\mathbb{Q}} = \mathbb{Q}(l(\tau))$  since  $l(\tau)$  has a rational  $q$ -expansion. Suppose  $\tau \in K \cap \mathbb{H}$  is a root of the primitive polynomial  $aX^2 + bX + c \in \mathbb{Z}[X]$  such that  $b^2 - 4ac = d_K$  and  $\gcd(a, 10) = 1$ . Then  $l(\tau)$  is well-defined and finite, so the desired conclusion follows from Lemma 5.1.  $\square$



The following result shows that the modular  $j$ -invariant defined by

$$j(\tau) = \eta^{-24}(\tau) \left( 1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n} \right)^3$$

can be written as a rational function of  $l(\tau)$ , which implies that  $l(\tau)$  is an algebraic unit for imaginary quadratic points  $\tau \in \mathbb{H}$ . In general, we infer from [10, Prop. 3] that the functions  $r(\tau)^a r(2\tau)^b$  are algebraic units for all integers  $a$  and  $b$  and for such points  $\tau \in \mathbb{H}$ .

**Theorem 5.3.** *Let  $K$  be an imaginary quadratic field and  $\tau \in K \cap \mathbb{H}$ . Then*

$$j(\tau) = \frac{P(l)^3}{l^2(l^2 - 1)(l^2 + 4l - 1)^5(l^2 - l - 1)^{10}},$$

where  $l := l(\tau)$  and

$$\begin{aligned} P(X) := & 1 - 4X + 234X^2 - 460X^3 + 495X^4 + 456X^5 - 1444X^6 \\ & - 456X^7 + 495X^8 + 460X^9 + 234X^{10} + 4X^{11} + X^{12}. \end{aligned}$$

*Proof.* We first remark that the function  $f(\tau) := \eta^6(\tau)\eta^{-6}(5\tau)$  generates the field of all modular functions on  $\Gamma_0(5)$  by Lemmata 2.5 and 2.6, and satisfies the identity [7, Thm. 5.26]

$$(2) \quad j(\tau) = \frac{(f^2(\tau) + 250f(\tau) + 3125)^3}{f^5(\tau)}.$$

As a modular function on  $\Gamma_0(10)$ , we deduce from Lemma 2.6 that  $f(\tau)$  has a double pole at  $1/5$  and a simple pole at  $\infty$ . Since  $A_0(\Gamma_0(10)) = \mathbb{C}(f(\tau), l(\tau) - 1/l(\tau))$  by Theorem 3.2, we infer from Proposition 4.3 that there is a polynomial

$$F(X, Y) = \sum_{\substack{0 \leq i \leq 1 \\ 0 \leq j \leq 3}} C_{i,j} X^i Y^j \in \mathbb{C}[X, Y]$$

such that  $F(f(\tau), l(\tau) - 1/l(\tau)) = 0$ . Using (1) and the  $q$ -expansion of  $f(\tau)$  leads to  $F(X, Y) = 4 - 7Y - XY + 2Y^2 + Y^3$  and

$$(3) \quad f(\tau) = \frac{(l^2 - l - 1)^2(l^2 + 4l - 1)}{l^2(l^2 - 1)},$$

where  $l := l(\tau)$ . Combining (2) and (3) gives the desired identity.  $\square$

The next result concerns about the singular values of  $l(r\tau)$  for positive rational numbers  $r$ , which can be written as radicals using the values of  $l(\tau)$ . We omit the proof as it is analogous to that of [12, Thm. 1.4].

**Theorem 5.4.** *For any positive  $r \in \mathbb{Q}$ , if  $l(\tau)$  can be expressed in terms of radicals, then  $l(r\tau)$  can also be expressed in terms of radicals.*

We now discuss a method of evaluating singular values of  $l(\theta)$ , where  $\theta \in K \cap \mathbb{H}$  is a generator for the ring of integers  $\mathcal{O}_K$  of an imaginary quadratic field  $K$  with discriminant  $d_K \equiv 0 \pmod{4}$ , by finding its minimal polynomial over  $K$ , which is the class polynomial of the ray class field  $K_{(10)}$  modulo 10 over  $K$ , using Shimura reciprocity law. For more details, we refer the reader to [4, 9].

We denote by  $\mathcal{F}_N$  the field modular functions of level  $N$  whose coefficients in their  $q$ -expansions lie in the cyclotomic field  $\mathbb{Q}(\zeta_N)$ . Then  $\mathcal{F}_N$  is a Galois extension of  $\mathcal{F}_1 = \mathbb{Q}(j)$  with Galois group  $\mathrm{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm 1\}$ . We recall the action of this Galois group on  $\mathcal{F}_N$  as follows. Given  $h \in \mathcal{F}_N$  and  $A_N \in \mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$ , we take a lift  $A \in \mathrm{SL}_2(\mathbb{Z})$  of  $A_N$  and define the action  $h^{A_N} = h \circ A$ , which induces the isomorphism  $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm 1\} \simeq$

$\text{Gal}(\mathcal{F}_N/\mathbb{Q}(\zeta_N)(j))$ . We next define the action of an element  $d \in (\mathbb{Z}/N\mathbb{Z})^\times$  on  $h$  by applying the  $\mathbb{Q}(\zeta_N)$ -automorphism  $\zeta_N \mapsto \zeta_N^d$  on the  $q$ -expansion of  $h$ , which yields the isomorphism  $(\mathbb{Z}/N\mathbb{Z})^\times \simeq \text{Gal}(\mathbb{Q}(\zeta_N)(j)/\mathbb{Q}(j))$ . Combining these isomorphisms induces the exact sequence

$$(4) \quad 1 \longrightarrow \{\pm 1\} \longrightarrow \text{GL}_2(\mathbb{Z}/N\mathbb{Z}) \longrightarrow \text{Gal}(\mathcal{F}_N/\mathcal{F}_1) \longrightarrow 1.$$

We know from the main theorem of complex multiplication that if  $h \in \mathcal{F}_N$  is a modular function with  $h(\theta)$  finite, then  $h(\theta)$  generates the ray class field  $K_{(N)}$  over  $K$ . Consequently, class field theory gives the exact sequence

$$(5) \quad 1 \longrightarrow \mathcal{O}_K^\times \longrightarrow (\mathcal{O}_K/N\mathcal{O}_K)^\times \xrightarrow{\bar{\theta}} \text{Gal}(K_{(N)}/K(j(\theta))) \longrightarrow 1,$$

where  $K(j(\theta))$  is the Hilbert class field of  $K$ , and  $\bar{\theta}$  is the reduction modulo  $N$  of the restriction of the Artin map to the profinite completion of  $\mathcal{O}_K$ . The Shimura reciprocity law relates (4) with (5) via the map  $g_\theta : (\mathcal{O}_K/N\mathcal{O}_K)^\times \rightarrow \text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  defined by

$$g_\theta(s\theta + t) = \begin{bmatrix} t - Bs & -Cs \\ s & t \end{bmatrix} \in \text{GL}_2(\mathbb{Z}/N\mathbb{Z}),$$

where  $s\theta + t \in (\mathcal{O}_K/N\mathcal{O}_K)^\times$  and  $X^2 + BX + C \in \mathbb{Z}[X]$  is the minimal polynomial of  $\theta$ . We denote by  $W_{N,\theta}$  the image of  $(\mathcal{O}_K/N\mathcal{O}_K)^\times$  under  $g_\theta$ . We see that there is a surjective homomorphism from  $W_{N,\theta}$  to  $\text{Gal}(K_{(N)}/K(j(\theta)))$  that sends  $\alpha$  to the map  $h(\theta) \mapsto h^{\alpha^{-1}}(\theta)$ , whose kernel  $T$  is the image  $g_\theta(\mathcal{O}_K^\times)$  given by

$$T = \begin{cases} \left\{ \left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \pm \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \right\} \right\} & \text{if } K = \mathbb{Q}(\sqrt{-1}), \\ \left\{ \left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \pm \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix}, \pm \begin{bmatrix} 0 & -1 \\ 1 & 1 \end{bmatrix} \right\} \right\} & \text{if } K = \mathbb{Q}(\sqrt{-3}), \\ \left\{ \left\{ \pm \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\} \right\} & \text{otherwise.} \end{cases}$$

To get the conjugates of  $h(\theta) \in K_{(N)}$  over  $K$ , we now exploit the action of the form class group  $C(d_K)$  on  $h(\theta)$  as follows. Suppose  $x := [a, b, c] \in C(d_K)$  is a reduced primitive binary quadratic form and  $\tau_x := (-b + \sqrt{d_K})/(2a) \in \mathbb{H}$ . We choose a matrix  $u_x \in \text{GL}_2(\mathbb{Z}/N\mathbb{Z})$  such that

$$u_x \equiv \begin{cases} \begin{bmatrix} a & b/2 \\ 0 & 1 \end{bmatrix} \pmod{p^{r_p}} & \text{if } \gcd(p, a) = 1, \\ \begin{bmatrix} -b/2 & -c \\ 1 & 0 \end{bmatrix} \pmod{p^{r_p}} & \text{if } p \mid a \text{ and } \gcd(p, c) = 1, \\ \begin{bmatrix} -b/2 - a & -b/2 - c \\ 1 & -1 \end{bmatrix} \pmod{p^{r_p}} & \text{if } p \mid a \text{ and } p \mid c \end{cases}$$

for each prime power  $p^{r_p}$  dividing  $N$ . We have the action  $h(\theta)^{x^{-1}} = h^{u_x}(\tau_x)$ , and note that if  $f$  is a modular function such that  $f(\theta) \in K(j(\theta))$ , then there is an isomorphism from  $C(d_K)$  to  $\text{Gal}(K(j(\theta))/K)$  that sends the class  $x^{-1}$  to the map  $f(\theta) \mapsto f^{u_x}(\tau_x)$ . In view of the isomorphism

$$\text{Gal}(K_{(N)}/K)/\text{Gal}(K_{(N)}/K(j(\theta))) \simeq \text{Gal}(K(j(\theta))/K),$$

we obtain the following.

**Proposition 5.5** ([4]). *Using the above notations, the set of all conjugates of  $h(\theta) \in K_{(N)}$  over  $K$  is given by*

$$\{h^{\alpha \cdot u_x}(\tau_x) : \alpha \in W_{N,\theta}/T, x \in C(d_K)\}.$$

After obtaining all conjugates of  $h(\theta)$ , we can now define the class polynomial of  $K_{(N)}$  given by

$$(6) \quad F_N(X) := \prod_{\substack{\alpha \in W_{N,\theta}/T \\ x \in C(d_K)}} (X - h^{\alpha \cdot u_x}(\tau_x)) \in K[X].$$

We now let  $\theta := \frac{\sqrt{d_K}}{2}$ ; then  $l(\theta)$  is a real number since  $e^{2\pi i \sqrt{d_K}/2} \in \mathbb{R}$ . Thus,

$$0 = F_{10}(l(\theta)) = \overline{F_{10}(l(\theta))} = \overline{F_{10}(\overline{l(\theta)})} = \overline{F_{10}(l(\theta))},$$

so that  $F_{10}(X) \in (K \cap \mathbb{R})[X] = \mathbb{Q}[X]$ . We see from Theorem 5.3 that  $l(\theta)$  is an algebraic integer, which shows that  $F_{10}(X) \in \mathbb{Z}[X]$ . In this case, we can numerically obtain  $F_{10}(X)$  using an accurate approximation of  $l(\theta)$  via (1). We illustrate an example of finding  $F_{10}(X)$ , which can be used to evaluate  $l(\theta)$  and the functions  $r(\theta)^a r(2\theta)^b$  using Corollary 3.4.

**Example 5.6.** Consider the field  $K = \mathbb{Q}(i)$  with  $d_K = -4$  and  $\mathcal{O}_K = \mathbb{Z}[i]$ , where  $i = \sqrt{-1}$ . We wish to find the class polynomial  $F_{10}(X)$  with  $F_{10}(l(i)) = 0$ . We choose  $(B, C) = (0, 1)$  since  $X^2 + 1$  is the minimal polynomial of  $i$ . Since  $x := [1, 0, 1]$  is the only reduced primitive binary quadratic form of discriminant  $-4$ , we have  $\tau_x = i$  and  $u_x = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \in \text{GL}_2(\mathbb{Z}/10\mathbb{Z})$ . We compute

$$W_{10,\theta}/T = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}, \begin{bmatrix} -1 & -4 \\ 4 & -1 \end{bmatrix}, \begin{bmatrix} 2 & -3 \\ 3 & 2 \end{bmatrix}, \right. \\ \left. \begin{bmatrix} 5 & 2 \\ -2 & 5 \end{bmatrix}, \begin{bmatrix} -4 & -5 \\ 5 & -4 \end{bmatrix}, \begin{bmatrix} -3 & 2 \\ -2 & -3 \end{bmatrix}, \begin{bmatrix} 4 & 1 \\ -1 & 4 \end{bmatrix} \right\}.$$

Using the  $q$ -expansion (1) of  $l(\tau)$ , we get

$$F_{10}(X) = (X - l(i))(X - l^{\begin{bmatrix} 3 & -10 \\ 10 & -33 \end{bmatrix}}(i))(X - l^{\begin{bmatrix} -1 & -4 \\ 2 & 7 \end{bmatrix}}(i))(X - l^{\begin{bmatrix} 2 & 7 \\ 1 & 4 \end{bmatrix}}(i)) \\ \times (X - l^{\begin{bmatrix} 5 & 12 \\ 2 & 5 \end{bmatrix}}(i))(X - l^{\begin{bmatrix} 6 & -5 \\ 5 & -4 \end{bmatrix}}(i))(X - l^{\begin{bmatrix} 7 & 2 \\ -4 & -1 \end{bmatrix}}(i))(X - l^{\begin{bmatrix} 4 & 1 \\ 7 & 2 \end{bmatrix}}(i)) \\ \approx X^8 + 26X^7 + 62X^6 + 458X^5 - 130X^4 - 458X^3 + 62X^2 - 26X + 1.$$

As  $l(i) \approx 1.00373486$ , we deduce that

$$l(i) = \frac{1}{4} \left( -13 - 5\sqrt{5} + (15 - \sqrt{5})\alpha + 2\sqrt{5}\sqrt{25 + 13\sqrt{5} - (15 + 5\sqrt{5})\alpha} \right),$$

where  $\alpha := \sqrt{2 + \sqrt{5}}$ . In view of Corollary 3.4 and the approximations  $r(2i)^3/r(i) \approx 0.00187091$  and  $r(2i)/r(i)^7 \approx 542.52907744$ , we arrive at the exact values of  $r(2i)^3/r(i)$  and  $r(2i)/r(i)^7$  respectfully given by

$$\frac{1}{4} \left( 251 + 115\sqrt{5} - (115 + 59\sqrt{5})\alpha + 2\sqrt{5}\sqrt{15725 + 7033\sqrt{5} - (7645 + 3415\sqrt{5})\alpha} \right), \\ \frac{1}{4} \left( -6259 - 2815\sqrt{5} + (3145 + 1557\sqrt{5})\alpha + 26\sqrt{5}\sqrt{50825 + 22733\sqrt{5} - (24615 + 11005\sqrt{5})\alpha} \right).$$

On the other hand, using the modular equation for  $l(\tau)$  of level two from Theorem 4.5 and  $l(i/2) \approx 1.08609902$ , we obtain the exact value of  $l(i/2)$ :

$$\frac{1}{2} \left( -134 - 60\sqrt{5} - (65 + 29\sqrt{5})\alpha + \sqrt{10}\sqrt{7225 + 3231\sqrt{5} + (3510 + 1570\sqrt{5})\alpha} \right).$$

We can then obtain the exact values of  $r(i)^3/r(i/2)$  and  $r(i)/r(i/2)^7$  using Corollary 3.4 and the value of  $l(i/2)$ .

## REFERENCES

1. G. E. Andrews and B. C. Berndt, *Ramanujan's Lost Notebook, Part I*, Springer Science+Business Media, New York, 2005.
2. Bruce C. Berndt and Robert A. Rankin, *Ramanujan: Letters and commentary*, History of Mathematics, vol. 9, American Mathematical Society, Rhode Island, 2000.
3. B. Cho, N. M. Kim, and J. K. Koo, *Affine models of the modular curves  $X(p)$  and its application*, Ramanujan J. **24** (2011), 235–257.
4. B. Cho and J. K. Koo, *Construction of class fields over imaginary quadratic fields and applications*, Q. J. Math. **61** (2010), 199–216.
5. B. Cho, J. K. Koo, and Y. K. Park, *Arithmetic of the Ramanujan-Göllnitz-Gordon continued fraction*, J. Number Theory **129** (2009), 922–947.
6. S. Cooper, *On Ramanujan's function  $k(q) = r(q)r(q^2)^2$* , Ramanujan J. **20** (2009), 311–328.
7. S. Cooper, *Ramanujan's Theta Functions*, Springer, Switzerland, 2017.
8. F. Diamond and J. Shurman, *A First Course in Modular Forms*, Graduate Texts in Mathematics, vol. 228, Springer, New York, 2005.
9. A. Gee, *Class invariants by Shimura's reciprocity law*, J. Théor. Nombres Bordeaux **11** (1999), 45–72.
10. A. Gee and M. Honsbeek, *Singular values of the Rogers-Ramanujan continued fraction*, Ramanujan J. **11** (2006), 267–284.
11. N. Ishida and N. Ishii, *The equations for modular function fields of principal congruence subgroups of prime level*, Manuscripta Math. **90** (1996), 271–285.
12. Y. Lee and Y. K. Park, *Ramanujan's function  $k(\tau) = r(\tau)r(2\tau)^2$  and its modularity*, Open Math. **18** (2020), 1727–1741.
13. G. Ligozat, *Courbes modulaires de genre 1*, Mémoires de la S. M. F. **43** (1975), 5–80.
14. M. Newman, *Construction and application of a class of modular functions*, Proc. London Math. Soc. **s3-7** (1957), 334–350.
15. ———, *Construction and application of a class of modular functions (II)*, Proc. London Math. Soc. **s3-9** (1959), 373–387.
16. K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and  $q$ -series*, CBMS Regional Conference Series in Mathematics, vol. 102, American Mathematical Society, Rhode Island, 2004.
17. G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton University Press, New Jersey, 1971.
18. G. Watson, *Theorems stated by Ramanujan (VII): Theorems on continued fraction*, J. London Math. Soc. **4** (1929), 39–48.
19. E. X. W. Xia and O. Y. M. Yao, *Some modular relations for Ramanujan's function  $k(q) = r(q)r^2(q^2)$* , Ramanujan J. **35** (2014), 243–251.
20. Y. Yang, *Transformation formulas for generalized Dedekind eta functions*, Bull. Lond. Math. Soc. **36** (2004), 671–682.
21. Y. Yang, *Defining equations of modular curves*, Adv. Math. **204** (2006), 481–508.
22. D. Ye, *Ramanujan's function  $k$ , revisited*, Ramanujan J. **56** (2021), 931–952.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF THE PHILIPPINES-DILIMAN, QUEZON CITY 1101, PHILIPPINES

*Email address:* rguadalupe@math.upd.edu.ph