


Modulus of continuity of Kerov transition measure for continual Young diagrams^{*†}

Piotr Śniady[‡] 

Abstract

The *transition measure* is a foundational concept introduced by Sergey Kerov to represent the shape of a Young diagram as a centered probability measure on the real line. Over a period of decades the transition measure turned out to be an invaluable tool for many problems of the asymptotic representation theory of the symmetric groups. Kerov also showed how to expand this notion for a wider class of *continual* diagrams so that the transition measure provides a homeomorphism between a subclass of continual diagrams (having a specific support) and a class of centered probability measures with a support contained in a specific interval. We quantify the modulus of continuity of this homeomorphism. More specifically, we study the dependence of the cumulative distribution function of Kerov transition measure on the profile of a diagram at the locations where the profile is not too steep.

Keywords: Kerov transition measure; Young diagrams; continual Young diagrams; Markov–Krein transform.

MSC2020 subject classifications: Primary 20C30, Secondary 44A60; 60E10; 05E10; 20C32.

1 Introduction

1.1 Young diagrams

1.1.1 The French convention

A Young diagram is a finite collection of boxes arranged in the positive quarter-plane, aligned to the left and bottom (see Figure 1a). This particular way of drawing Young diagrams is known as *the French convention*. For a Young diagram with ℓ rows, we associate an integer partition $\lambda = (\lambda_1, \dots, \lambda_\ell)$, where λ_j represents the number of boxes

^{*}Research was supported by Narodowe Centrum Nauki, grant number 2017/26/A/ST1/00189.

[†]We thank Maciej Dołęga, Łukasz Maślanka and Dan Romik for discussions and suggestions concerning the bibliography.

[‡]Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, 00-656 Warszawa, Poland E-mail: psniady@impan.pl




Figure 1: The Young diagram $\lambda = (4, 3, 1)$ shown in (a) the French convention, and (b) the Russian convention. The thick solid red zigzag line represents the *profile* ω_λ of the Young diagram. The coordinates system (u, v) corresponding to the Russian convention and the coordinate system (x, y) corresponding to the French convention are shown.

in the j -th row (counted from bottom to top). Essentially, we identify a Young diagram with its corresponding partition λ .

The motivation for studying Young diagrams lies in their natural occurrence in representation theory. Specifically, they play a crucial role in the representation theory of symmetric groups as well as in the representation theory of general linear groups [7].

1.1.2 The Russian convention

For asymptotic problems, it is convenient to draw the Young diagrams in the *Russian convention*, as shown in Figure 1b. This convention corresponds to the coordinate system (u, v) that is related to the usual (French) Cartesian coordinates by

$$u = x - y, \quad v = x + y. \tag{1.1}$$

The boundary of a Young diagram λ is called its *profile*, as depicted in Figure 1a. In the Russian coordinate system, the profile can be seen as the plot of the function $\omega_\lambda: \mathbb{R} \rightarrow \mathbb{R}_+$, as shown in Figure 1b.

1.2 Transition measure of a Young diagram

For a given Young diagram λ we denote by $x_0 < \dots < x_L$ the u -coordinates of its *concave corners* and by $y_1 < \dots < y_L$ the u -coordinates of its *convex corners*, see Figure 2. The *Cauchy transform* of λ is defined as the rational function

$$\mathbf{G}_\lambda(z) = \frac{(z - y_1) \cdots (z - y_L)}{(z - x_0) \cdots (z - x_L)}, \tag{1.2}$$

see [8] and [9, Chapter 4, Section 1]. Note that in the work of Kerov this function is called *the generating function* of λ . The Cauchy transform can be uniquely written as a sum of simple fractions:

$$\mathbf{G}_\lambda(z) = \sum_{0 \leq i \leq L} \frac{p_i}{z - x_i}$$

with the positive coefficients $p_0, \dots, p_L > 0$ such that $p_0 + \dots + p_L = 1$. The *transition measure* of λ is defined as the discrete measure

$$\mu_\lambda = p_0 \delta_{x_0} + \dots + p_L \delta_{x_L} \tag{1.3}$$




Figure 2: Concave corners (blue and empty) and convex corners (red and filled) of the Young diagram $(4, 2, 2, 2)$ and their u -coordinates.

which to the atom x_i associates the probability p_i . In this way the rational function defined by (1.2) fulfills

$$G_\lambda(z) = \int_{\mathbb{R}} \frac{1}{z - x} d\mu_\lambda(x),$$

so it is indeed the Cauchy transform of the measure μ_λ .

1.3 Motivations and context for transition measure

The definition of the transition measure presented in Section 1.2 has an advantage of being self-contained, but it does not provide much insight into the motivations behind this concept. We will recall some equivalent definitions of the transition measure which show its relevance for probabilistic and algebraic methods of the representation theory of the symmetric groups. We will also recall briefly some of its applications.

1.3.1 Plancherel growth process

The *Plancherel growth process* is a natural Markov chain $(\lambda^{(n)})$ on the set of Young diagrams. This random walk starts with $\lambda^{(0)} = \emptyset$ being the empty diagram. At each step, the random Young diagram $\lambda^{(n+1)}$ is obtained by adding a single box to one of the concave corners of the previous Young diagram $\lambda^{(n)}$. If μ is a Young diagram obtained from λ by adding a single box, the transition probability for this Markov chain

$$\mathbb{P} \left[\lambda^{(n+1)} = \mu \mid \lambda^{(n)} = \lambda \right] = \frac{f^\mu}{(n + 1)f^\lambda} \tag{1.4}$$

is defined in a natural way in terms of the dimensions f^λ and f^μ of the irreducible representations of the symmetric groups defined by, respectively, λ and μ .

Equivalently, the Plancherel growth process can be described as the outcome of the Robinson–Schensted–Knuth algorithm (RSK) applied to a sequence of independent, identically distributed random variables with the uniform distribution on the unit interval $[0, 1]$.

For a pedagogical introduction to this topic we refer to the book of Romik [13, Section 1.19].

1.3.2 Plancherel growth process and the transition measure

Let us fix a Young diagram λ ; we will keep the notations from Figure 2. The collection of the transition probabilities (1.4) over all valid choices of μ can be encoded by a discrete probability measure on the real line, given as follows. To a real number x_i which corresponds to some concave corner of λ we associate the transition probability (1.4) from λ to the diagram μ obtained from λ by adding a single box at the corner x_i .

Thanks to the hook-length formula for the dimensions of the irreducible representations, the atoms of this discrete measure (each atom is given by the right-hand side of (1.4)) can be explicitly calculated. After a heavy cancellation of the factors in the hook-length formula for f^λ and f^μ this transition probability turns out to coincide with the corresponding atom p_i prescribed in Section 1.2.

This was the original motivation for introducing the transition measure. Indeed, transition measure is a convenient tool for the probabilistic and ergodic approach to the problems of the (asymptotic) representation theory of the symmetric groups.

1.3.3 Transition measure and Jucys–Murphy elements

The *Jucys–Murphy element*

$$J_{n+1} = (1, n + 1) + \cdots + (n, n + 1) \in \mathbb{C}[S_{n+1}]$$

is an element of the group algebra of the symmetric group S_{n+1} , given by the formal sum of the transpositions interchanging $n + 1$ with the smaller natural numbers. Jucys–Murphy elements are fundamental for the modern approach to the representation theory of the symmetric groups, see [3] for a pedagogical introduction.

It was noticed by Biane [1, Section 3.3] that the transition measure of a Young diagram λ with n boxes can be equivalently defined as the spectral measure of the Jucys–Murphy element J_{n+1} with respect to some specific state defined in terms of the irreducible character of λ . The result of Biane implies that each moment

$$\int x^k d\mu_\lambda(x)$$

of the transition measure is directly related to the character of λ evaluated on (some specific conditional expectation applied to) the power J_{n+1}^k of the Jucys–Murphy element.

1.3.4 Transition measure and the characters of symmetric groups

This link opened the possibility to look for exact formulas which would express the irreducible characters of the symmetric group in terms of the transition measure of the corresponding Young diagram. The advantage of such formulas (over some classical tools such as *the Murnaghan–Nakayama rule*) would lie in their low computational complexity which would make them perfect for some asymptotic problems of the representation theory.

It was soon noticed that the best quantities for this endeavor are not the moments of the transition measure, but *free cumulants*, slightly more complex quantities which have their origin in the random matrix theory and Voiculescu’s non-commutative free probability [12]. The irreducible characters can be expressed as polynomials, called *Kerov polynomials*, in terms of the free cumulants of the corresponding transition measure [4].

This explains why transition measure and its free cumulants turned out to be important tools for probabilistic problems of the asymptotic representation theory of the symmetric groups [1, 2, 14].

1.4 Continual diagrams

1.4.1 Motivations: limit theorems

One of the branches of the asymptotic representation theory concerns limit theorems for large random Young diagrams. For example, for each integer $n \geq 1$ let $\lambda^{(n)}$ be a random Young diagram with n boxes, with some specified probability distribution. Informally speaking, we would like to consider a ‘*resized Young diagram*’

$$\frac{1}{\sqrt{n}}\lambda^{(n)} \tag{1.5}$$

which is obtained by drawing the boxes of $\lambda^{(n)}$ as squares with the size $\frac{1}{\sqrt{n}}$, so that the total area of (1.5) is equal to 1. Unfortunately, this object does not make sense as a Young diagram.

Even though the ‘*resized Young diagram*’ (1.5) does not exist, it makes sense to speak about its *profile* which is defined as the homothety of the profile of the original diagram $\lambda^{(n)}$:

$$\omega\left(\frac{1}{\sqrt{n}}\lambda^{(n)}\right)(z) = \frac{1}{\sqrt{n}} \omega_{\lambda^{(n)}}(\sqrt{n} z). \tag{1.6}$$

A typical problem in the asymptotic representation theory would be to ask if the sequence of such resized profiles (1.6) converges to some limit [10, 15]. This was the original motivation for the notion of *continual diagrams* which encompasses the resized profiles (1.6) as well as their limits. We present the details in the following.

1.4.2 Continual diagrams

We say that $\omega: \mathbb{R} \rightarrow \mathbb{R}_+$ is a *continual diagram* if the following two conditions are satisfied:

- $|\omega(z_1) - \omega(z_2)| \leq |z_1 - z_2|$ holds true for any $z_1, z_2 \in \mathbb{R}$,
- $\omega(z) = |z|$ for sufficiently large $|z|$.

In the literature, such continual diagrams are also referred to as *generalized diagrams*, *continuous diagrams*, or simply *diagrams*. Notably, an important class of examples arises from the profiles ω_λ associated with the usual Young diagrams.

Remark 1.1. In his original paper [8], Kerov considered a more general form for the second condition. Specifically, he required that $\omega(z) = |z - z_0|$ should hold for some fixed value of $z_0 \in \mathbb{R}$ and for sufficiently large $|z|$. The definition we use corresponds to the specific choice of $z_0 = 0$. Consequently, the *continual diagrams* discussed in the current paper align with what Kerov terms *centered diagrams*.

1.5 Transition measure for continual diagrams

1.5.1 Transition measure for zigzag diagrams

We will say that a continual diagram ω is a *zigzag* if $\omega: \mathbb{R} \rightarrow \mathbb{R}_+$ is a piece-wise affine function with the slopes being only ± 1 . Heuristically, this means that ω is a zigzag line as depicted on Figure 2. The only difference is that now the u -coordinates of its concave corners $x_0 < \dots < x_L$ as well as the u -coordinates of its convex corners $y_1 < \dots < y_L$ are no longer assumed to be integer numbers.

For a zigzag diagram ω we define its Cauchy transform G_ω in the same way as before, by the formula (1.2). The transition measure of ω , denoted by μ_ω is defined in the same way as before, by the formula (1.3).

1.5.2 Transition measure for continual diagrams

Let ω be a zigzag, we keep the notations from above. The second derivative ω'' is well-defined as a Schwartz distribution and can be identified with a signed measure on the real line. The positive part of this measure is supported in the concave corners $\mathbb{x}_0, \dots, \mathbb{x}_L$ while the negative part of this measure is supported in the convex corners $\mathbb{y}_1, \dots, \mathbb{y}_L$, with each atom having equal weight 2. An application of logarithm transforms the product on the right-hand side of (1.2) into a sum. It was observed by Kerov that it can be conveniently written in the form

$$\log [z\mathbf{G}_\omega(z)] = - \int_{-\infty}^{\infty} \log(z-w) \left(\frac{\omega(w) - |w|}{2} \right)'' dw = - \int_{-\infty}^{\infty} \frac{1}{z-w} \left(\frac{\omega(w) - |w|}{2} \right)' dw, \quad (1.7)$$

where z is a complex variable.

We can drop the assumption that ω is a zigzag and use (1.7) to define the Cauchy transform \mathbf{G}_ω for an arbitrary continual diagram ω . The transition measure μ_ω is then defined from the Cauchy transform \mathbf{G}_ω via Stieltjes inversion formula. The *cumulative function* K_ω of a continual diagram ω is defined as the cumulative distribution function of the corresponding transition measure

$$K_\omega(z) = \mu_\omega((-\infty, z]) \quad \text{for } z \in \mathbb{R}.$$

1.5.3 Transition measure for continual diagrams as a homeomorphism

Kerov (see [8] and [9, Chapter 4, Section 1]) proved that (for each constant $C > 0$) the map

$$\omega \mapsto \mu_\omega \quad (1.8)$$

is a homeomorphism between the following two topological spaces:

- the set of continual diagrams ω with the property that

$$\omega(z) = |z| \quad \text{for each } z \notin [-C, C],$$

equipped with the topology given by the supremum distance, and

- the set of probability measures with support contained in the interval $[-C, C]$, equipped with the weak topology of probability measures.

The fact that the map (1.8) is a homeomorphism allows to retrieve the transition measure of an arbitrary continual diagram by approximating it by a sequence of zigzag diagrams and then by taking the limit of their transition measures. We will use this idea later on.

1.6 The main problem: what is the modulus of continuity of the transition measure?

In the current paper we address the problem which can be stated informally as follows.

Problem 1.2 (The main problem). What is the modulus of continuity of the map (1.8)?

Given two continual diagrams ω_1 and ω_2 such that their distance (with respect to the supremum distance) is 'small', how small is the distance between the cumulative functions K_{ω_1} and K_{ω_2} (in some unspecified metric)?

Kerov’s result that (1.8) is a homeomorphism implies that the distance between K_{ω_1} and K_{ω_2} converges to zero (in some metric which corresponds to the weak convergence of probability measures) as the distance between ω_1 and ω_2 tends to zero. However, Kerov’s original proof [8] is not very constructive in the sense that it does not provide much information about the speed of the convergence, and hence it does not provide quantitative information about the ‘discrepancy’ between the cumulative functions K_{ω_1} and K_{ω_2} .

1.7 The motivations

Our motivation for considering Theorem 1.2 is related to our recent joint work with Marciniak [11] about the statistical properties of the *Schensted row insertion* applied to random tableaux. We recall that the Schensted row insertion is the basic component of the Robinson–Schensted–Knuth algorithm [6].

Up to some minor simplifications, for a given large Young diagram λ we consider a uniformly random *standard Young tableau* T with the prescribed shape λ . We ask about the position of the new box created by the row insertion $T \leftarrow z$, when a deterministic number z is inserted into T . The position of the new box is clearly random, located in one of the concave corners of λ . The statistical properties of its probability distribution can be expressed in terms of the transition measure μ_λ of the diagram λ .

The problem becomes even more interesting asymptotically, when the single Young diagram λ is replaced by a *sequence* of Young diagrams $(\lambda^{(n)})$. In this setup one can ask whether the (suitably rescaled) fluctuations of the position of the new box around the mean value converge to some limit distribution. As an extra bonus, the Young diagram $\lambda^{(n)}$ may be chosen to be deterministic or random.

The main result of the paper [11] is that the fluctuations of the position of the new box indeed converge to an explicit Gaussian distribution, provided that the cumulative function of the rescaled diagram $\lambda^{(n)}$

$$z \mapsto K\left(\frac{1}{\sqrt{n}} \omega_{\lambda^{(n)}}\right)(z) = K_{\lambda^{(n)}}(\sqrt{n} z)$$

converges in probability to a deterministic limit which is sufficiently smooth. Additionally, the rate of convergence must be fast enough, more specifically the error must be at most $o(n^{-\frac{1}{4}})$.

The goal of the current paper is to provide tools for producing explicit natural examples of large (deterministic or random) Young diagrams λ for which the cumulative function K_λ can be written as a sum of a known smooth function and a small error term with an explicit asymptotic bound. For such examples the results of [11] are applicable. We will come back to this topic in Section 1.9.

1.8 The metric d on the set of continual diagrams

For the purposes of this section we will use the following notational shorthand. For real numbers $x, y \geq 0$ and a continual diagram $\omega: \mathbb{R} \rightarrow \mathbb{R}_+$ we will write $(x, y) \in \omega$ if the point (x, y) belongs to the graph of ω drawn in the *French coordinate system* or, equivalently, if

$$\omega(x - y) = x + y,$$

cf. (1.1). Additionally, for $x \geq 0$ we denote by

$$\Pi_Y^\omega(x) = \{y : (x, y) \in \omega\} \subseteq \mathbb{R}_+$$

the projection on the y -axis of the intersection of the plot of ω with the line having a specified x -coordinate. One can verify that $\Pi_Y^\omega(x)$ is a non-empty closed set. Similarly, for $y \geq 0$ we denote by

$$\Pi_X^\omega(y) = \{x : (x, y) \in \omega\} \subseteq \mathbb{R}_+$$

the projection on the x -axis.

If ω_1, ω_2 are continual diagrams, we define their y -distance

$$d_Y(\omega_1, \omega_2) = \sup_{x \geq 0} d_H(\Pi_Y^{\omega_1}(x), \Pi_Y^{\omega_2}(x))$$

as the supremum (over all choices of the x -coordinate) of the *Hausdorff distance* d_H between their y -projections. Heuristically, the metric d_Y is a way to quantify the discrepancy between continual diagrams along the y -coordinate. In a similar manner we define the x -distance between continual diagrams as

$$d_X(\omega_1, \omega_2) = \sup_{y \geq 0} d_H(\Pi_X^{\omega_1}(y), \Pi_X^{\omega_2}(y)).$$

Finally, we define the distance between ω_1 and ω_2 as the maximum of the x - and the y -distance:

$$d(\omega_1, \omega_2) = \max(d_X(\omega_1, \omega_2), d_Y(\omega_1, \omega_2)).$$

1.9 The main result

The following result (for the exponent $\alpha = \frac{1}{4}$) provides a wide class of examples for the problem which we discussed in Section 1.7. Informally speaking, it says that if Ω is locally a ‘nice’ continual diagram and a sequence (ω_n) converges to Ω in the metric d with some prescribed speed then the cumulative functions of (ω_n) converge uniformly to the cumulative function of Ω with almost the same speed.

Theorem 1.3 (The main result). *Let $\alpha > 0$ be fixed. Let $\Omega: \mathbb{R} \rightarrow \mathbb{R}_+$ be a continual diagram, let $a < a_0 < b_0 < b$ be such that:*

- (i) *the function Ω restricted to the interval $[a, b]$ is a contraction (i.e., it is $(1 - \delta)$ -Lipschitz for some constant $\delta > 0$);*
- (ii) *the transition measure μ_Ω restricted to the interval $[a, b]$ is absolutely continuous and has a density which is bounded from above by some constant.*

Let (ω_n) be a sequence of continual diagrams such that

$$d(\Omega, \omega_n) = o\left(\frac{1}{n^\alpha \log n}\right).$$

Then

$$\sup_{z \in [a_0, b_0]} |K_{\omega_n}(z) - K_\Omega(z)| = o\left(\frac{1}{n^\alpha}\right).$$

The proof is postponed to Section 7.

1.10 Toy example

In this section we introduce a toy example to which the main result (Theorem 1.3) applies. Although this example is straightforward and elementary, allowing us to calculate the transition measures explicitly, it is still valuable to see how the general tool applies to a specific context.

1.10.1 The staircase diagrams

In the context of Theorem 1.3 we will pass to a subsequence defined as

$$n = n_N = 1 + \dots + N = \frac{N(N+1)}{2}.$$

We define

$$\lambda^{(n_N)} = (N, N-1, \dots, 3, 2, 1) \tag{1.9}$$

to be a *staircase Young diagram* with n_N boxes. We define

$$\omega_n = \omega_{\frac{1}{\sqrt{n}}\lambda^{(n)}}$$

to be the rescaled profile of the staircase diagram.

We define the *triangle diagram*

$$\Omega(z) = \begin{cases} \sqrt{2} & \text{for } |z| \leq \sqrt{2}, \\ |z| & \text{for } |z| \geq \sqrt{2}. \end{cases}$$

Then the sequence (ω_n) converges to Ω in the metric d , with the rate of convergence given by

$$d(\omega_n, \Omega) = O\left(\frac{1}{N}\right) = O\left(\frac{1}{\sqrt{n}}\right). \tag{1.10}$$

1.10.2 The transition measures

A straightforward calculation of the residues of the Cauchy transform shows that the transition measure of $\lambda^{(n)}$ is supported on the set of even, respectively odd integer numbers

$$\{-N, -N+2, \dots, N-2, N\}$$

with the probabilities

$$\mu_{\lambda^{(n)}}(2k-N) = \frac{1}{2^{2N}} \binom{2k}{k} \binom{2N-2k}{N-k} \quad \text{for } k \in \{0, \dots, N\}. \tag{1.11}$$

This probability distribution appears naturally in the context of random walks and the *arcsine theorem*, see [5, Chapter III].

On the other hand, the transition measure μ_Ω of the triangle diagram is the arcsine law supported on the interval $I = [-\sqrt{2}, \sqrt{2}]$ with the density

$$f_{AS}(z) = \frac{1}{\pi\sqrt{2-z^2}} \quad \text{for } z \in I.$$

1.10.3 Application of Theorem 1.3

By (1.10) it follows that the assumptions of Theorem 1.3 are fulfilled for the exponent $\alpha = \frac{1}{2} - \epsilon$, for any $\epsilon > 0$, and for any quadruple $a < a_0 < b_0 < b$ from the open interval $(-\sqrt{2}, \sqrt{2})$. As a consequence, the rate of convergence of the cumulative function of the rescaled staircase diagram ω_n towards the cumulative function of the triangle diagram Ω is bounded from above by

$$\sup_{z \in [a_0, b_0]} |K_{\omega_n}(z) - K_\Omega(z)| = o\left(\frac{1}{n^{\frac{1}{2}-\epsilon}}\right) = o\left(\frac{1}{N^{1-2\epsilon}}\right) \tag{1.12}$$

for each $\epsilon > 0$.

1.10.4 What is the optimal rate of convergence?

A straightforward calculation based on the Stirling approximation provides an upper bound of the form

$$\sup_{z \in [a_0, b_0]} |K_{\omega_n}(z) - K_{\Omega}(z)| = O\left(\frac{1}{\sqrt{n}}\right) = O\left(\frac{1}{N}\right). \quad (1.13)$$

On the other hand, the transition measure of ω_n is a discrete measure which in the interval (a_0, b_0) has atoms which are of order $\Theta\left(\frac{1}{N}\right)$. It follows that also the left-hand side of (1.13) is at least of order $\Theta\left(\frac{1}{N}\right) = \Theta\left(\frac{1}{\sqrt{n}}\right)$, hence the bound (1.13) is optimal.

We can see that Theorem 1.3 provided us with an upper bound (1.12) which is very close to the optimal result (1.13). We conclude that there is not much room for improvement left in Theorem 1.3.

1.11 The main result for random continual diagrams

Theorem 1.3 allows the following extension for random diagrams.

Theorem 1.4 (The main result, version for random diagrams). *We keep the assumptions from Theorem 1.3, but now (ω_n) is a sequence of random continual diagrams such that*

$$n^\alpha \log n \cdot d(\Omega, \omega_n) \xrightarrow[n \rightarrow \infty]{P} 0.$$

Then

$$n^\alpha \sup_{z \in [a_0, b_0]} |K_{\omega_n}(z) - K_{\Omega}(z)| \xrightarrow[n \rightarrow \infty]{P} 0.$$

The proof is postponed to Section 7.

1.12 Content of the paper

We start in Section 2 with a simple monotonicity result (Theorem 2.1) which, roughly speaking, says that when one ‘adds new boxes’ to a continual diagram on the right-hand side, or ‘removes some boxes’ on the left-hand side, the values of the cumulative function in the middle can only increase. Thanks to this monotonicity result we will later estimate the cumulative function of generic diagrams in terms of the cumulative function of some nice continual diagram over which we have a good control.

Section 3 is devoted to a small technical result about vanishing of the transition measure of a continual diagram ω on the intervals for which the derivative ω' is constantly equal to ± 1 .

In Sections 4 and 5 we address the following problem. In the metric space of the continual diagrams (equipped with the metric d) we consider the ball $B_{\epsilon, \Omega}$ which has radius $\epsilon > 0$ and is centered in some continual diagram Ω . Inside this ball we consider a subset

$$\{\omega \in B_{\epsilon, \Omega} : \omega(z_0) = v\} \quad (1.14)$$

of diagrams which at given $z_0 \in \mathbb{R}$ take a specified value v . *What can we say about the supremum of the map $\omega \mapsto K_{\omega}(z_0)$ over this set (1.14)?*

In Section 4 we will construct a diagram ω in (1.14) for which the cumulative function $K_{\omega}(z_0)$ takes the maximal value. As we shall see, this diagram is given by the ϵ -shift of Ω along some specific affine function f , cf. Theorem 4.1.

In Section 5 we will find an upper bound for the supremum (Theorem 5.5) in terms of Ω and its transition measure.

In Section 6 we prove a twin pair of results: Theorem 6.1 and Theorem 6.2. They give an upper bound for the difference

$$|K_\Omega(z_0) - K_\omega(z_0)| \tag{1.15}$$

of the cumulative functions of two continual diagrams Ω and ω . The roles played by these diagrams are quite different: in the applications Ω is ‘nice’ (for example, $\Omega: \mathbb{R} \rightarrow \mathbb{R}_+$ may be given by an explicit analytic expression, additionally it may have a derivative which locally is away from ± 1 , etc.), while ω is quite generic and we not have much information about it. The aforementioned upper bound for (1.15) is expressed in terms of the distance $d(\Omega, \omega)$ between the diagrams, as well as in terms of the shape of Ω , and the transition measure of Ω . Notably, this bound does not depend on some fine details of ω .

Finally, in Section 7 we apply Theorems 6.1 and 6.2 in order to prove Theorems 1.3 and 1.4.

2 Steeper diagram has a larger cumulative function

Proposition 2.1. *Let $\omega_1, \omega_2: \mathbb{R} \rightarrow \mathbb{R}_+$ be two continual diagrams and let $z_0 \in \mathbb{R}$ be fixed. Assume that the following two conditions hold true:*

$$\begin{cases} \omega_1(z) \geq \omega_2(z) & \text{for each } z \leq z_0, \\ \omega_1(z) \leq \omega_2(z) & \text{for each } z \geq z_0. \end{cases} \tag{2.1}$$

Then the left and the right limits at z_0 of the cumulative distribution functions of the continual diagrams ω_1 and ω_2 fulfill

$$\lim_{\tau \rightarrow 0^+} K_{\omega_1}(z_0 - \tau) \leq K_{\omega_2}(z_0). \tag{2.2}$$

The remaining part of this section is devoted to the proof.

2.1 Special case: Young diagrams.

We start with the special case when $\omega_1 = \omega_{\lambda^1}$ and $\omega_2 = \omega_{\lambda^2}$ are profiles of some Young diagrams λ^1 and λ^2 . We will show a stronger result

$$K_{\omega_1}(z_0) \leq K_{\omega_2}(z_0). \tag{2.3}$$

The assumption (2.1) implies that the diagram λ^1 can be transformed to λ^2 by a sequence of *elementary changes*. Each elementary change is either:

- (a) an *addition of a box* with the u -coordinate of the two central vertices equal to $\xi \in \mathbb{Z}$ with $\xi \geq z_0 + 1$, see Figure 3, or
- (b) a *removal of a box* with the u -coordinate of the two central vertices equal to $\xi \in \mathbb{Z}$ with $\xi \leq z_0 - 1$.

Each neighboring pair of Young diagrams along this sequence of elementary changes fulfills an analogue of the inequalities (2.1) and we will show below that an analogue of (2.3) holds true for each such a pair. By chaining the sequence of the inequalities of the form (2.3) along the sequence of the elementary changes, the desired inequality for the endpoints ω_1 and ω_2 would follow.

We proceed now with an elementary change of type (a). Let $\omega = \omega_\lambda$ be a profile of a Young diagram and let $\bar{\omega}$ be the profile of the diagram obtained from ω by adding a




Figure 3: A Young diagram λ (white boxes) and its profile ω_λ (the thick zigzag line). The u -coordinate of the two central vertices of the blue box is equal to ξ .

single box, as in Figure 3. The elementary change transforms the profile ω to $\bar{\omega}$. By comparing the sets of convex and the concave corners for these two profiles it follows that their Cauchy transforms are related by the following equality:

$$\mathbf{G}_{\bar{\omega}}(z) = \frac{(z - \xi)^2}{(z - \xi - 1)(z - \xi + 1)} \mathbf{G}_{\omega}(z).$$

It follows that the atoms of the transition measures of ω and $\bar{\omega}$ are related as follows:

- for any integer $z \notin \{\xi - 1, \xi, \xi + 1\}$

$$\mu_{\bar{\omega}}(\{z\}) = \frac{(z - \xi)^2}{(z - \xi - 1)(z - \xi + 1)} \mu_{\omega}(\{z\}) = \left[1 + \frac{1}{(z - \xi)^2 - 1} \right] \mu_{\omega}(\{z\}) \geq \mu_{\omega}(\{z\}); \quad (2.4)$$

- additionally, the transition measure $\mu_{\bar{\omega}}$ potentially may have atoms in $\xi \pm 1$ while the transition measure μ_{ω} does not have any atoms there, so the inequality

$$\mu_{\bar{\omega}}(\{z\}) \geq \mu_{\omega}(\{z\}) \quad (2.5)$$

holds true also for $z \in \{\xi - 1, \xi + 1\}$.

The cumulative distribution functions can be expressed as sums of the appropriate atoms, hence

$$K_{\bar{\omega}}(z_0) = \sum_{\substack{z \in \mathbb{Z} \\ z \leq z_0}} \mu_{\bar{\omega}}(\{z\}) \geq \sum_{\substack{z \in \mathbb{Z} \\ z \leq z_0}} \mu_{\omega}(\{z\}) = K_{\omega}(z_0)$$

where we used the inequalities (2.4) and (2.5), as well as the fact that if the integer z contributes to the above sums then $z \leq \xi - 1$. This concludes the proof for an elementary change of type (a).

We proceed now with an elementary change of type (b). We keep the above notations; the elementary change transforms the continual diagram $\bar{\omega}$ to ω . The cumulative distribution functions can be expressed in terms of the tails of the respective distributions therefore

$$K_{\omega}(z_0) = 1 - \sum_{\substack{z \in \mathbb{Z} \\ z > z_0}} \mu_{\omega}(\{z\}) \geq 1 - \sum_{\substack{z \in \mathbb{Z} \\ z > z_0}} \mu_{\bar{\omega}}(\{z\}) = K_{\bar{\omega}}(z_0)$$

where we used the inequalities (2.4) and (2.5), as well as the fact that if the integer z contributes to the above sums then $z \geq \xi + 1$. This concludes the proof for an elementary change of type (b).

2.2 Rescaled Young diagrams.

For any $c > 0$ and any Young diagram λ , the transition measure of the rescaled profile

$$\omega_{c\lambda}(z) = \frac{1}{c} \omega(cz)$$

is a dilation of the transition of the original Young diagram λ . As a consequence, the inequality (2.3) holds true also when $\omega_1 = \omega_{c\lambda^1}$ and $\omega_2 = \omega_{c\lambda^2}$ are rescaled profiles of two Young diagrams.

2.3 The general case.

For each $i \in \{1, 2\}$ and an integer $n \geq 1$ we define $\lambda^{i,n}$ to be the largest Young diagram λ with the property that

$$\omega_{\frac{1}{n}\lambda}(z) \leq \omega_i(z)$$

holds true for each $z \in \mathbb{R}$. The pair of rescaled profiles

$$\omega_{1,n} := \omega_{\frac{1}{n}\lambda^{1,n}} \quad \text{and} \quad \omega_{2,n} := \omega_{\frac{1}{n}\lambda^{2,n}}$$

fulfills an analogue of the system of inequalities (2.1), hence the above discussion is applicable and therefore

$$K_{\omega_{1,n}}(z_0) \leq K_{\omega_{2,n}}(z_0). \quad (2.6)$$

For each $i \in \{1, 2\}$, the sequence of generalized diagrams $(\omega_{i,n})_{n \geq 1}$ converges to the generalized diagram ω_i in the suitable topological space described by Kerov [8]. From the result of Kerov [8] it follows that the sequence of transition measures $(\mu_{\omega_{i,n}})_{n \geq 1}$ converges in the weak topology to the transition measure μ_{ω_i} .

Let $z_- < z_0$ be a continuity point of K_{ω_1} and let $z_+ > z_0$ be a continuity point of K_{ω_2} . It follows that

$$\begin{aligned} K_{\omega_1}(z_-) &= \lim_{n \rightarrow \infty} K_{\omega_{1,n}}(z_-) \leq \liminf_{n \rightarrow \infty} K_{\omega_{1,n}}(z_0) \leq \\ &\liminf_{n \rightarrow \infty} K_{\omega_{2,n}}(z_0) \leq \lim_{n \rightarrow \infty} K_{\omega_{2,n}}(z_+) = K_{\omega_2}(z_+). \end{aligned}$$

Since the set of the continuity points of K_{ω_1} and K_{ω_2} is dense, we may consider the left limit of the left-hand side at $z_- = z_0$ and the right limit of the right-hand side at $z_+ = z_0$. As a consequence,

$$\lim_{\tau \rightarrow 0} K_{\omega_1}(z_0 - \tau) \leq K_{\omega_2}(z_0),$$

as required. This completes the proof of Theorem 2.1.

3 Vanishing of the transition measure if the slope is ± 1

Lemma 3.1. *Let ω be a continual diagram such that its derivative ω' is constant, equal to ± 1 , on some open interval $I = (a, b)$.*

Then the restriction of transition measure μ_ω to this open interval I is equal to zero.

Proof. In the case when ω is a zigzag diagram, this result is an immediate consequence of the definition from Section 1.2.

For the general case we will use the method similar to the one from Section 2.3. In order to keep the discussion focused we consider the case when $\omega'(z) = 1$ for $z \in I$.

For an integer $n \geq 1$ let $\lambda^{(n)}$ be the largest Young diagram such that

$$\omega_{\frac{1}{n}\lambda^{(n)}}(z) \leq \omega(z)$$

holds true for each $z \in \mathbb{R}$. We denote $\omega_n := \omega_{\frac{1}{n}\lambda^{(n)}}$; by the same argument as in Section 2.3 the sequence (μ_{ω_n}) of the corresponding transition measures converges weakly to μ_ω .

Just as in Section 1.2, let $\mathbb{x}_0, \dots, \mathbb{x}_L$ and $\mathbb{y}_1, \dots, \mathbb{y}_L$ be the concave and the convex corners of the zigzag diagram ω_n . From the following on we assume that n is large enough so that $\frac{1}{n} < \frac{1}{3}|I|$. There is an open interval between some concave corner \mathbb{x}_i and the subsequent convex corner \mathbb{y}_{i+1} which (up to a discretization error of size $\frac{1}{n}$) contains the interval I ; more specifically

$$\mathbb{x}_i \leq a + \frac{1}{n}, \quad b - \frac{1}{n} \leq \mathbb{y}_{i+1}.$$

We denote $a_n = \mathbb{x}_i$ and $b_n = \mathbb{y}_{i+1}$. It follows that

$$\limsup a_n \leq a, \quad b \leq \liminf b_n.$$

The transition measure of the zigzag diagram ω_n is zero on the interval (a_n, b_n) . By the weak convergence of measures it follows that μ_ω is zero on (a, b) , as required.

The proof in the case when $\omega'(z) = -1$ for $z \in I$ is analogous and we skip it. \square

4 Shifted diagrams

4.1 The input data

Let $\Omega: \mathbb{R} \rightarrow \mathbb{R}_+$ be a continual diagram, let $\epsilon > 0$ be a real number and let $f: \mathbb{R} \rightarrow \mathbb{R}$ be an affine function of the form

$$f(x) = x + b$$

for some constant $b > 0$. For this input data we will construct in the current section a new continual diagram $\bar{\Omega} = \bar{\Omega}_{\Omega, \epsilon, f}$, called ϵ -shift of Ω along f .

4.2 The upper and the lower bound for the ϵ -ball

Let ω be an arbitrary continual diagram such that $d(\Omega, \omega) \leq \epsilon$. Since ϵ is an upper bound on the deviation between the curves Ω and ω along the x -coordinate, it follows that

$$\omega(z) \leq \Omega(z - \epsilon) + \epsilon \tag{4.1}$$

holds true for any $z \in \mathbb{R}$. The function which appears on the right hand side, i.e.,

$$z \mapsto \Omega(z - \epsilon) + \epsilon$$

is depicted in Figure 4 as the red dashed line. Analogously, since ϵ is an upper bound on the deviation between the curves Ω and ω along the y -coordinate, it follows that

$$\omega(z) \geq \Omega(z - \epsilon) - \epsilon \tag{4.2}$$

holds true for any $z \in \mathbb{R}$. The function which appears on the right hand side, i.e.,

$$z \mapsto \Omega(z - \epsilon) - \epsilon$$

is depicted in Figure 4 as the blue dashed line. In this way any continual diagram within the ϵ -ball around Ω (with respect to the metric d) lies within the area between the red and the blue zigzag lines.




Figure 4: The solid black zigzag line is the continual diagram Ω . The upper red dashed zigzag line was obtained by translation of Ω by the vector (in the French coordinates) $x = \epsilon, y = 0$. The bottom blue dashed zigzag line was obtained by translation of Ω by the vector $x = 0, y = -\epsilon$. The red and the blue dotted arrows indicate these translation vectors.

The densely dotted diagonal black line is the plot of the affine function f . The two black circles indicate the intersection points of the function f with, respectively, the blue and the red zigzag lines. Their u -coordinates are equal to, respectively, z_- and z_+ .

The thick green zigzag line is the continual diagram $\bar{\Omega}$ which was obtained by combining a segment of the blue dashed line (to the left of the first black circle), a segment of the function f (between the black circles), and a segment of the red dashed line (to the right of the second black circle).

4.3 The intersection points z_- and z_+

Let z_- be the minimal number such that

$$f(z_-) = \Omega(z_- - \epsilon) - \epsilon \quad (4.3)$$

and let z_+ be the maximal number such that

$$f(z_+) = \Omega(z_+ - \epsilon) + \epsilon. \quad (4.4)$$

With the notations of Figure 4 these are the u -coordinates of the intersection of the dotted line f with the blue zigzag line and with the red zigzag line. Clearly, the intersection points exist and the numbers z_- , z_+ are well-defined and fulfill

$$z_- \leq z_+,$$

see Figure 4.

4.4 The shifted diagram

We define a continual diagram $\bar{\Omega} = \bar{\Omega}_{\Omega, \epsilon, f}$ given by

$$\bar{\Omega}(z) = \begin{cases} \Omega(z - \epsilon) - \epsilon & \text{for } z \leq z_-, \\ f(z) & \text{for } z_- \leq z \leq z_+, \\ \Omega(z - \epsilon) + \epsilon & \text{for } z \geq z_+. \end{cases}$$

Equivalently,

$$\bar{\Omega}(z) = \min \left[\max \left[\Omega(z - \epsilon) - \epsilon, f(z) \right], \Omega(z - \epsilon) + \epsilon \right]. \quad (4.5)$$

We refer to $\bar{\Omega}$ as ϵ -shift of Ω along f . In Figure 4 this diagram is shown as the thick green zigzag line; it is a piece-wise combination of the red zigzag line, the dotted line, and the blue zigzag line.

This definition has the following heuristic interpretation in the special case when ϵ and $b = f(0)$ are positive integers, and Ω is a profile of some Young diagram λ . In this case, in the French coordinate system the curve f is a horizontal straight line with the equation $y = b$. Then the shifted diagram $\bar{\Omega}$ is the profile of the Young diagram obtained from Ω by the following procedure:

- we extend each of the rows of λ lying below the horizontal line f by ϵ additional boxes,
- among the rows which are lying above the horizontal line f , we remove the bottom ϵ rows, and we shift the remaining rows down, so that they are placed on top of the horizontal line f .

4.5 Shifted diagram as a reference point for the ϵ -ball

Proposition 4.1. *Let $z_0 \in \mathbb{R}$, $\epsilon > 0$, let ω, Ω be continual diagrams such that $d(\omega, \Omega) \leq \epsilon$ and such that $\omega(z_0) > z_0$. Let*

$$f(z) = z - z_0 + \omega(z_0) \quad (4.6)$$

be the unique affine function with slope 1 and such that $f(z_0) = \omega(z_0)$. Let $\bar{\Omega} = \bar{\Omega}_{\Omega, \epsilon, f}$ be the ϵ -shifted diagram Ω along f .

Then the left limit of the cumulative function of ω fulfills

$$\lim_{\tau \rightarrow 0^+} K_\omega(z_0 - \tau) \leq K_{\bar{\Omega}}(z_0).$$

Proof. From the construction of $\overline{\Omega}$ it follows that

$$\begin{aligned}\omega(z) &\geq \overline{\Omega}(z) && \text{for each } z \leq z_0, \\ \omega(z) &\leq \overline{\Omega}(z) && \text{for each } z \geq z_0.\end{aligned}$$

Theorem 2.1 completes the proof. \square

5 Transition measure for shifted diagrams

The main result of the current section is Theorem 5.5 which provides an upper bound for the tail of the transition measure of a shifted diagram. Theorem 5.1 will serve as a technical intermediate step.

5.1 Function $P_{z_+}^{\min}$

In the remaining part of this section $\epsilon > 0$ is a fixed constant. It will be convenient to treat the dependence of $P_{z_+}^{\min}$ (see below) on ϵ as implicit.

For a real number z_+ we define

$$\begin{aligned}P_{z_+}^{\min}(z) &= \begin{cases} 1 - \frac{\epsilon}{z + \epsilon - z_+} & \text{for } z \geq z_+, \\ 0 & \text{otherwise,} \end{cases} \\ &= \begin{cases} \frac{z - z_+}{z + \epsilon - z_+} & \text{for } z \geq z_+, \\ 0 & \text{otherwise.} \end{cases}\end{aligned}\tag{5.1}$$

The function $z \mapsto P_{z_+}^{\min}(z)$ is continuous and weakly increasing; it takes values from the interval $[0, 1)$.

5.2 Lower bound for the transition measure of $\overline{\Omega}$, only zigzag diagrams

We start with an additional assumption that the continual diagram Ω is a zigzag, cf. Section 1.5.1. In this way we can use the notations from Section 1.2, in particular let $\mathfrak{x}_0 < \dots < \mathfrak{x}_L$ be the u -coordinates of the concave corners of Ω and let $\mathfrak{y}_1 < \dots < \mathfrak{y}_L$ be the convex corners, see Figure 2.

Proposition 5.1. *Let Ω be a zigzag diagram, and let $\epsilon > 0$, f and z_+ be as described in Sections 4.1 and 4.3, and let $\overline{\Omega} = \overline{\Omega}_{\Omega, f, \epsilon}$ be the corresponding shifted diagram.*

Then for any non-negative function $\phi: \mathbb{R} \rightarrow \mathbb{R}_+$

$$\int_{-\infty}^{\infty} \phi(z) d\mu_{\overline{\Omega}}(z) \geq \int_{-\infty}^{\infty} \phi(z + \epsilon) P_{z_+}^{\min}(z + \epsilon) d\mu_{\Omega}(z).$$

Above, ϕ plays the role of a *test function* and this result can be interpreted informally as inequality

$$\mu_{\overline{\Omega}} \geq P_{z_+}^{\min}(\cdot + \epsilon) \mu_{\Omega}$$

between measures.

The remaining part of the current section is devoted to the proof.

5.2.1 The Cauchy transform of $\overline{\Omega}$.

The shifted convex and the concave corners

$$\mathfrak{x}_0 + \epsilon < \mathfrak{y}_1 + \epsilon < \mathfrak{x}_1 + \epsilon < \dots < \mathfrak{y}_L + \epsilon < \mathfrak{x}_L + \epsilon$$




Figure 5: A simplified version of Figure 4. The upper red zigzag line as well as the lower blue zigzag line are, as before, translations of Ω . The u -coordinates of their convex and the concave corners coincide, up to a shift by ϵ , with their counterparts for Ω , see Figure 2. The convention for convex and concave corners is the one from Figure 2. The two black circles maintain their meaning from Figure 4.

split the real line into a number of (finite or semi-infinite) intervals; the number z_- belongs to one of them. Note that the case when $x_i + \epsilon < z_- \leq y_{i+1} + \epsilon$ is not possible since this would contradict the minimality of z_- , see Figures 4 and 5. There exists therefore an index $i \in \{0, \dots, \mathbb{L}\}$ such that

$$y_i + \epsilon < z_- \leq x_i + \epsilon,$$

see Figure 5. In the case $i = 0$ we use the convention that $y_0 = -\infty$ so that the first inequality is automatically fulfilled. Similarly, there exists an index $j \in \{1, \dots, \mathbb{L}\}$ such that

$$y_j + \epsilon \leq z_+ < x_j + \epsilon,$$

see Figure 5.

Lemma 5.2. *The relationship between the Cauchy transforms of $\bar{\Omega}$ and Ω is given by*

$$\mathbf{G}_{\bar{\Omega}}(z) = \mathbf{G}_{\Omega}(z - \epsilon) P(z), \quad (5.2)$$

where

$$P(z) = \frac{z - z_+}{z - z_-} \prod_{i \leq k < j} \frac{z - (x_k + \epsilon)}{z - (y_{k+1} + \epsilon)}. \quad (5.3)$$

Proof. An inspection of Figures 4 and 5 shows that $\bar{\Omega}$ has the following set of concave corners:

$$x_0 + \epsilon, \dots, x_{i-1} + \epsilon, \quad z_-, \quad x_j + \epsilon, \dots, x_{\mathbb{L}} + \epsilon,$$

and the following set of concave corners:

$$y_1 + \epsilon, \dots, y_i + \epsilon, \quad z_+, \quad y_{j+1} + \epsilon, \dots, y_L + \epsilon.$$

The proof is completed by comparing the zeros and the poles of the rational functions which appear on both sides of (5.2). \square

Our strategy is to use this formula in order to express the cumulative function of $\overline{\Omega}$ in terms of the cumulative function for Ω .

5.2.2 Identity fulfilled by convex and concave corners.

Lemma 5.3. *With the above notations,*

$$z_+ - z_- = \epsilon + \sum_{i \leq k < j} y_{k+1} - x_k. \quad (5.4)$$

Proof. We will calculate the difference of the x -coordinates of the two black circles (the intersection points) in Figure 4 in two ways.

On one hand, this difference is equal to the difference $z_+ - z_-$ of their u -coordinates which is the left-hand side of (5.4).

On the other hand, the difference of the x -coordinates would decrease by ϵ if we transform the intersection points as follows:

- we shift the left circle by the vector with the French coordinates $x = 0, y = \epsilon$ (which is the opposite to the blue vector on Figure 4);
- we shift the right circle by the vector with the French coordinates $x = -\epsilon, y = 0$ (which is the opposite to the red vector on Figure 4).

The transformed black circles both lie on the continual diagram Ω . The difference of their x -coordinates is equal to the sum of the segments of Ω which lie between the transformed circles and are parallel to the x -axis, see Figures 4 and 5, which is the second summand on the right-hand side of (5.4). \square

5.2.3 The lower bound for the product P .

Lemma 5.4. *With the above notations, for each $z \geq z_+$*

$$P(z) \geq P_{z_+}^{\min}(z),$$

cf. (5.1).

Proof. In order to find a lower bound for $P = P(z)$, we shall view it as a function of the variables

$$x_i \leq y_{i+1} \leq x_{i+1} \leq y_{i+2} \leq \dots \leq y_{j-1} \leq x_{j-1} \leq y_j \in [z_- - \epsilon, z_+ - \epsilon] \quad (5.5)$$

which are subject to the constraint (5.4). Let the variables (5.5) be such that the minimal value of P is attained (such a minimum is attained by the compactness argument). Let us modify two of these variables by replacing x_m by $x_m + \tau$, as well as y_{m+1} by $y_{m+1} + \tau$, where $\tau < 0$ is very close to zero. Since

$$P(z) = \frac{z - z_+}{z - z_-} \prod_{i \leq k < j} \left[1 + \frac{y_{k+1} - x_k}{z - (y_{k+1} + \epsilon)} \right]$$

is an increasing function of τ , the minimality of P implies that such a modification should not be possible. As a consequence, the appropriate inequalities in (5.5) must be saturated and

$$\mathbb{x}_i = z_- - \epsilon, \quad \mathbb{y}_{i+1} = \mathbb{x}_{i+1}, \quad \mathbb{y}_{i+2} = \mathbb{x}_{i+2}, \quad \dots, \quad \mathbb{y}_{j-1} = \mathbb{x}_{j-1}.$$

Additionally, by (5.4),

$$\mathbb{y}_j = z_+ - 2\epsilon.$$

In this way we proved that for $z \geq z_+$ the product (5.3) is bounded from below by

$$P_{z_+}^{\min}(z) = \frac{z - z_+}{z + \epsilon - z_+} = 1 - \frac{\epsilon}{z + \epsilon - z_+},$$

as required. \square

5.2.4 Proof of Theorem 5.1

Proof of Theorem 5.1. The atoms of the transition measure $\mu_{\bar{\Omega}}$ (respectively, the atoms of μ_{Ω}) are given by the residues of the rational function $\mathbf{G}_{\bar{\Omega}}$ (respectively, \mathbf{G}_{Ω}). We use (5.2) in order to relate the residues of $\mathbf{G}_{\bar{\Omega}}$ to the residues to \mathbf{G}_{Ω} . Since the residues of these two functions coincide up to a shift by ϵ , it follows that

$$\int_{-\infty}^{\infty} \phi(z) d\mu_{\bar{\Omega}}(z) \geq \int_{z_+}^{\infty} \phi(z) d\mu_{\bar{\Omega}}(z) = \int_{z_+ - \epsilon}^{\infty} \phi(z + \epsilon) P(z + \epsilon) d\mu_{\Omega}(z) \geq \int_{-\infty}^{\infty} \phi(z + \epsilon) P_{z_+}^{\min}(z + \epsilon) d\mu_{\Omega}(z). \quad \square \quad (5.6)$$

5.3 Tail of the transition measure of $\bar{\Omega}$, generic diagrams

The content of the following result is very similar to Theorem 5.1; the key difference is that now we do *not* assume that Ω is a zigzag.

Proposition 5.5. *Let $\epsilon > 0$, Ω , f and z_+ be as described in Sections 4.1 and 4.3, and let $\bar{\Omega} = \bar{\Omega}_{\Omega, f, \epsilon}$ be the corresponding shifted diagram. Then*

$$\mu_{\bar{\Omega}}((z_+, \infty)) \geq \int_{-\infty}^{\infty} P_{z_+}^{\min}(z + \epsilon) d\mu_{\Omega}(z),$$

cf. (5.1).

The remaining part of this section is devoted to the proof.

5.3.1 Approximation by zigzags

Let $(\Omega^{(n)})$ be a sequence of zigzag diagrams which converges pointwise to Ω , and such that the continual diagrams $(\Omega^{(n)})$ have a common compact support, i.e., there exists a constant C such that

$$\Omega^{(n)}(z) = |z| \quad \text{if } |z| > C \quad (5.7)$$

holds for any integer $n \geq 1$. Such a sequence can be explicitly constructed similarly as in the proof of the general case of Theorem 2.1. Since the transition measure provides a homeomorphism between appropriate topological spaces, it follows that the sequence of transition measures $(\mu_{\Omega^{(n)}})$ converges weakly to the transition measure μ_{Ω} .

5.3.2 Asymptotics of $z_+^{(n)}$

The constructions from Sections 4.3 and 4.4 can be applied to $\Omega := \Omega^{(n)}$. We denote by $\overline{\Omega}^{(n)} = \overline{\Omega}_{\Omega^{(n)}, \epsilon, f}$ and $z_+^{(n)}$ and the corresponding values of $\overline{\Omega}$ and z_+ . Let

$$z_+^{(\infty)} := \limsup_{n \rightarrow \infty} z_+^{(n)}. \quad (5.8)$$

Lemma 5.6. *With the above notations,*

$$z_- + \epsilon \leq z_+^{(\infty)} \leq z_+. \quad (5.9)$$

Proof. In the context of $\Omega := \Omega^{(n)}$, the defining property of $z_+^{(n)}$, i.e., equation (4.4), takes the form

$$f(z_+^{(n)}) = \Omega^{(n)}(z_+^{(n)} - \epsilon) + \epsilon.$$

By continuity it follows that

$$f(z_+^{(\infty)}) = \Omega(z_+^{(\infty)} - \epsilon) + \epsilon.$$

In other words, $z_+^{(\infty)}$ is a solution of (4.4); by the maximality of z_+ , the second inequality in (5.9) follows.

On the other hand, any solution of the equation (4.4) is bigger by at least ϵ than any solution of (4.3). The first inequality in (5.9) follows. \square

5.3.3 Asymptotics of $\overline{\Omega}^{(n)}$

From the alternative definition (4.5) of the shifted diagram it follows that the sequence of shifted diagrams $(\overline{\Omega}^{(n)})$ has a common compact support and converges pointwise to the shifted diagram $\overline{\Omega}$. As a consequence, the sequence of their transition measures $(\mu_{\overline{\Omega}^{(n)}})$ converges to $\mu_{\overline{\Omega}}$ in the weak topology of probability measures.

The shifted diagram $\overline{\Omega}$ restricted to the interval $[z_-, z_+]$ is an affine function with the slope 1. For this reason, Theorem 3.1 is applicable. The weak convergence of the transition measures implies therefore

$$\mu_{\overline{\Omega}}([z_+, \infty)) = \mu_{\overline{\Omega}}([z, \infty)) = \lim_{n \rightarrow \infty} \mu_{\overline{\Omega}^{(n)}}([z, \infty)) \quad \text{for each } z \in I. \quad (5.10)$$

5.3.4 Applying Theorem 5.1 to zigzag diagrams

Let $\phi_k: \mathbb{R} \rightarrow \mathbb{R}_+$ be a continuous function which will be specified later. Theorem 5.1 implies that

$$\int_{-\infty}^{\infty} \phi_k(z) d\mu_{\overline{\Omega}^{(n)}}(z) \geq \int_{-\infty}^{\infty} \phi_k(z) P_{z_+^{(n)}}^{\min}(z + \epsilon) d\mu_{\Omega^{(n)}}(z) \quad (5.11)$$

holds true for each n .

Theorem 5.6 implies that for each $\delta > 0$

$$z_+^{(n)} \leq \delta + z_+$$

holds true for all $n \geq n_0$ for some value of n_0 . Since for each fixed value of z , the map $t \mapsto P_t^{\min}(z)$ is weakly decreasing, this implies that the right hand side of (5.11) can be bounded from below as follows

$$\int_{-\infty}^{\infty} \phi_k(z + \epsilon) P_{z_+^{(n)}}^{\min}(z + \epsilon) d\mu_{\Omega^{(n)}}(z) \geq \int_{-\infty}^{\infty} \phi_k(z + \epsilon) P_{\delta + z_+}^{\min}(z + \epsilon) d\mu_{\Omega^{(n)}}(z) \quad (5.12)$$

for $n \geq n_0$.

We chain the inequalities (5.11) and (5.12) and take iterated limit: first $n \rightarrow \infty$ and then $\delta \rightarrow 0$ and get:

$$\int_{-\infty}^{\infty} \phi_k(z) d\mu_{\overline{\Omega}}(z) \geq \int_{-\infty}^{\infty} \phi_k(z + \epsilon) P_{z_+}^{\min}(z + \epsilon) d\mu_{\Omega}(z). \quad (5.13)$$

5.3.5 Approximating the step function

For an integer $k \geq 1$ we will specify now the function ϕ_k as follows:

$$\phi_k(z_+ + \Delta) = \begin{cases} 0 & \text{for } \Delta \leq 0, \\ k\Delta & \text{for } 0 \leq \Delta \leq \frac{1}{k}, \\ 1 & \text{for } \Delta > \frac{1}{k}. \end{cases}$$

In this way (ϕ_k) is a sequence of continuous functions which converges pointwise to

$$\phi(z) = \begin{cases} 0 & \text{for } z \leq z_+, \\ 1 & \text{for } z > z_+. \end{cases}$$

We consider (5.13) for this specific choice of ϕ_k and take the limit $k \rightarrow \infty$:

$$\mu_{\overline{\Omega}}((z_+, \infty)) \geq \int_{-\infty}^{\infty} P_{z_+}^{\min}(z + \epsilon) d\mu_{\Omega}(z),$$

which completes the proof of Theorem 5.5.

6 The bounds for the cumulative function

6.1 The upper bound

Theorem 6.1. *Let $\Omega: \mathbb{R} \rightarrow \mathbb{R}_+$ be a continual diagram and let $\epsilon > 0$ and $z_0 \in \mathbb{R}$ be fixed. Assume that z_+^{\max} is the maximal solution of the equation*

$$\Omega(z_+^{\max} - \epsilon) - \Omega(z_0 - \epsilon) = z_+^{\max} - z_0 - 2\epsilon \quad (6.1)$$

see Figure 6 for an illustration.

Then for any continual diagram ω such that $d(\Omega, \omega) \leq \epsilon$ the following upper bound for the cumulative function of ω holds true:

$$\lim_{\tau \rightarrow 0^+} K_{\omega}(z_0 - \tau) \leq 1 - \int_{z_+^{\max} - \epsilon}^{\infty} \left[1 - \frac{\epsilon}{z + 2\epsilon - z_+^{\max}} \right] d\mu_{\Omega}(z) = \int_{-\infty}^{\infty} \left(\mathbb{1}_{(-\infty, z_+^{\max} - \epsilon)}(z) + \mathbb{1}_{[z_+^{\max} - \epsilon, \infty)}(z) \frac{\epsilon}{z + 2\epsilon - z_+^{\max}} \right) d\mu_{\Omega}(z). \quad (6.2)$$

Note that equation (6.1) may not have any solutions, or the set of solutions may be unbounded from above; the above result does not concern such cases. Since Ω is 1-Lipschitz, it follows that each solution of the equation (6.1) fulfills $z_+^{\max} \geq z_0 + \epsilon$, see Figure 6.

The integrand on the right-hand side of (6.2) can be heuristically interpreted as a continuous, smoothed-out version of the indicator function $\mathbb{1}_{(-\infty, z_0]}$.

Some readers may find it aesthetically more pleasing to replace in (6.1) and (6.2) the quantity z_+^{\max} by $\hat{z} := z_+^{\max} - \epsilon$.




Figure 6: The solid black zigzag line is the continual diagram Ω . The densely dotted diagonal line is the plot of the affine function $z \mapsto z - (z_0 - \epsilon) + \Omega(z_0 - \epsilon) - 2\epsilon$. The rightmost, red circle indicates the intersection of this line with the plot of Ω .

Proof. Our strategy will be to apply Theorem 4.1. We revisit the setup considered in Sections 4.1 to 4.3. Let the affine function f be given by (4.6) and, as before, let z_+ be the maximal solution of (4.4). In order to find an upper bound for z_+ we notice that such a maximal value corresponds to the affine function f with the smallest possible v -intercept, see Figure 4.

The inequality Equation (4.2) taken at $z = z_0$ implies that the affine function f is bounded from below by

$$f_{\min}(z) = z - z_0 + \Omega(z_0 - \epsilon) - \epsilon.$$

With these notations, (6.1) can be rewritten in the form

$$f_{\min}(z_+^{\max}) = \Omega(z_+^{\max} - \epsilon) + \epsilon. \quad (6.3)$$

By comparing it with the defining property (4.4) it follows that z_+^{\max} is a special case of z_+ evaluated for the minimal affine function f_{\min} . In this way we proved that $z_+ \leq z_+^{\max}$.

Furthermore, the degenerate case $\omega(z_0) = z_0$ is not possible because it would imply that $f_{\min}(z) \leq f(z) = z$ lies below the x -axis. As a consequence, the set of solutions of (6.3) (which is equivalent to (6.1)) would be empty or not bounded from above which contradicts the assumption on z_+^{\max} . It follows that without the loss of generality we may assume $\omega(z_0) > z_0$ and Theorem 4.1 is applicable.

We apply Theorem 4.1; we denote by $\bar{\Omega}$ the shifted diagram which is defined there. The left limit of the cumulative function of ω is bounded by

$$\lim_{\tau \rightarrow 0^+} K_{\omega}(z_0 - \tau) \leq K_{\bar{\Omega}}(z_0) = 1 - \mu_{\bar{\Omega}}((z_0, \infty)). \quad (6.4)$$

We will use the notations from Figure 4. Let P be the point on the plane which has the Russian coordinates $u = z_0$ and $v = \omega(z_0)$. From the way the affine function f was defined in Theorem 4.1 it follows that P belongs both to the plot of the function ω as

well as to the plot of the affine function f . We can rephrase it as follows: P belongs to the intersection of the area between the red and the blue curve (cf. Section 4.2) and the plot of f . It follows immediately that

$$z_0 \leq z_+,$$

hence

$$\mu_{\overline{\Omega}}((z_0, \infty)) \geq \mu_{\overline{\Omega}}((z_+, \infty)). \quad (6.5)$$

The right-hand side can be bounded from below by Theorem 5.5. Since for each fixed value of z , the map $t \mapsto P_t^{\min}(z)$ is weakly decreasing, it follows that

$$\mu_{\overline{\Omega}}((z_+, \infty)) \geq \int_{-\infty}^{\infty} P_{(z_+^{\max})}^{\min}(z + \epsilon) d\mu_{\Omega}(z). \quad (6.6)$$

We complete the proof by combining the inequalities provided by Equation (6.4), (6.5) and (6.6). \square

6.2 Transpose of a diagram

If $\omega: \mathbb{R} \rightarrow \mathbb{R}_+$ is a continual diagram we define its *transpose*

$$\omega^T(z) = \omega(-z).$$

This definition extends the usual notion of a transposed (or *conjugate*) Young diagram.

It is easy to check that the transition measure of a transposed diagram $\mu_{(\omega^T)}$ is the push-forward of the transition measure of the original diagram μ_{ω} under the involution $\mathbb{R} \ni z \mapsto -z \in \mathbb{R}$.

6.3 The lower bound

The following is a mirror image of Theorem 6.1.

Theorem 6.2. *Let $\Omega: \mathbb{R} \rightarrow \mathbb{R}_+$ be a continual diagram and let $\epsilon > 0$ and $z_0 \in \mathbb{R}$ be fixed. Assume that z_{-}^{\min} is the minimal solution of the equation*

$$\Omega(z_{-}^{\min} + \epsilon) - \Omega(z_0 + \epsilon) = z_0 - z_{-}^{\min} - 2\epsilon.$$

Then for any continual diagram ω such that $d(\Omega, \omega) \leq \epsilon$ the following lower bound for the cumulative function of ω holds true:

$$K_{\omega}(z_0) \geq \int_{-\infty}^{z_{-}^{\min} + \epsilon} \left[1 - \frac{\epsilon}{z_{-}^{\min} + 2\epsilon - z} \right] d\mu_{\Omega}(z). \quad (6.7)$$

Proof. Consider the involution of the real line $z \mapsto -z$ and apply Theorem 6.1 for transposed diagrams

$$\omega' := \omega^T, \quad \Omega' := \Omega^T,$$

and for

$$z'_0 := -z_0, \quad z_{-}^{\min'} := -z_{+}^{\max}. \quad \square$$

7 Proof of Theorems 1.3 and 1.4

Lemma 7.1. *We keep the assumptions from Theorem 1.3. We denote by $z_{+}^{\max} = z_{+}^{\max}(z, \epsilon)$ the quantity defined in Theorem 6.1.*

There exists a constant $\epsilon_0 > 0$ with the property that for each $z \in [a_0, b_0]$ and for each $\epsilon \in (0, \epsilon_0)$

$$z_{+}^{\max} - z_0 \leq \frac{2\epsilon}{\delta}.$$

Modulus of continuity of Kerov transition measure

Proof. Let us fix $z_0 \in [a_0, b_0]$ and $\epsilon > 0$. We consider a weakly decreasing function

$$F(z) = \Omega(z - \epsilon) - z - \Omega(z_0 - \epsilon) + z_0 + 2\epsilon;$$

the motivation for this definition is that the condition (6.1) is equivalent to $F(z_+^{\max}) = 0$.

Assume that $\epsilon \in (0, a_0 - a)$ is sufficiently small. The assumption (i) implies that if $z \in [z_0, b]$ then

$$F(z) \leq F(z_0) - (z - z_0) \delta = 2\epsilon - (z - z_0) \delta. \quad (7.1)$$

We denote by

$$z_{\max} := z_0 + \frac{2\epsilon}{\delta}$$

the value of the variable z for which the right-hand side of (7.1) is equal to zero. Note that for $z > z_{\max}$ this right hand side is strictly negative. We also define

$$\epsilon_0 = \min \left(\frac{(b - b_0) \delta}{2}, a_0 - a \right).$$

In this way, if $0 < \epsilon < \epsilon_0$ then

$$z_{\max} < z_0 + b - b_0 \leq b.$$

As a consequence, (7.1) is applicable for $z := z_{\max}$ and $F(z_{\max}) \leq 0$. Furthermore, F is strictly negative on the interval (z_{\max}, ∞) . As a consequence $z_+^{\max} \leq z_{\max}$ which completes the proof. \square

Proof of Theorem 1.3. We denote by ρ_{\max} any upper bound for the density of the measure μ_{Ω} on the interval $[a, b]$. We apply Theorem 6.1 for $\omega := \omega_n$ and $\epsilon_n := d(\Omega, \omega_n)$. We will keep the dependence of the corresponding z_+^{\max} on the variable n implicit. We obtain the following upper bound:

$$\begin{aligned} \lim_{\tau \rightarrow 0^+} K_{\omega_n}(z_0 - \tau) - K_{\Omega}(z_0) &\leq \\ \mu_{\Omega}((z_0, z_+^{\max} - \epsilon]) + \epsilon_n \int_{z_+^{\max} - \epsilon_n}^b \frac{1}{z + 2\epsilon_n - z_+^{\max}} d\mu_{\Omega}(z) &+ \epsilon_n \int_b^{\infty} \frac{1}{z + 2\epsilon_n - z_+^{\max}} d\mu_{\Omega}(z). \end{aligned} \quad (7.2)$$

The first summand on the right hand side is of order $O(\epsilon_n)$ by Theorem 7.1, uniformly over all choices of $z_0 \in [a_0, b_0]$. The second summand is bounded from above by

$$\epsilon_n \rho_n \log \frac{b - a_0}{\epsilon_n} = O \left(\epsilon_n \log \frac{1}{\epsilon_n} \right).$$

The third summand is bounded from above by

$$\frac{\epsilon_n}{b - z_+^{\max}} = O(\epsilon_n)$$

because of the bound provided by Theorem 7.1. This shows that the right-hand side converges to zero at the right asymptotic speed. Additionally, since this bound is uniform over z_0 it follows that on the left-hand side we can replace the left limit of the cumulative function by the cumulative function $K_{\omega_n}(z_0)$ itself.

The proof of the lower bound is analogous and we skip it. \square

Proof of Theorem 1.4. We will revisit the above proof of Theorem 1.3. Let $C > 0$ be fixed; we modify the definition of ε_n by setting

$$\varepsilon_n = \frac{C}{n^\alpha \log n}.$$

In this way the probability of the event $d(\Omega, \omega_n) < \varepsilon_n$ converges to 1 as $n \rightarrow \infty$. As a consequence, also the probability that the inequality (7.2) holds true, converges to 1. The right-hand side of (7.2) is, as before, at most of order $O\left(\varepsilon_n \log \frac{1}{\varepsilon_n}\right)$. Since we have the freedom to choose $C > 0$ arbitrarily small, this completes the proof. \square

References

- [1] P. Biane, *Representations of symmetric groups and free probability*, Adv. Math. **138** (1998), no. 1, 126–181. MR1644993
- [2] ———, *Approximate factorization and concentration for characters of symmetric groups*, Internat. Math. Res. Notices **2001** (2001), no. 4, 179–192. MR1813797
- [3] T. Ceccherini-Silberstein, F. Scarabotti, and F. Tolli, *Representation theory of the symmetric groups: The Okounkov-Vershik approach, character formulas, and partition algebras*, Cambridge Studies in Advanced Mathematics, vol. 121, Cambridge University Press, Cambridge, 2010. MR2643487
- [4] Maciej Dołęga, Valentin Féray, and Piotr Śniady, *Explicit combinatorial interpretation of Kerov character polynomials as numbers of permutation factorizations*, Adv. Math. **225** (2010), no. 1, 81–120.
- [5] William Feller, *An introduction to probability theory and its applications. Vol. I*, third edition ed., John Wiley & Sons, Inc., New York-London-Sydney, 1968. MR0228020
- [6] W. Fulton, *Young tableaux: With applications to representation theory and geometry*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997. MR1464693
- [7] William Fulton and Joe Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, A first course, Readings in Mathematics. MR1153249
- [8] S. Kerov, *Transition probabilities for continual Young diagrams and the Markov moment problem.*, Funct. Anal. Appl. **27** (1993), 104–117.
- [9] S. V. Kerov, *Asymptotic representation theory of the symmetric group and its applications in analysis*, Translations of Mathematical Monographs, vol. 219, American Mathematical Society, Providence, RI, 2003, Translated from the Russian manuscript by N. V. Tsilevich, With a foreword by A. Vershik and comments by G. Olshanski. MR1984868
- [10] B. F. Logan and L. A. Shepp, *A variational problem for random Young tableaux*, Advances in Math. **26** (1977), no. 2, 206–222. MR1417317
- [11] Mikołaj Marciniak and Piotr Śniady, *Fluctuations of Schensted row insertion*, arXiv:2302.03762, 2023.
- [12] Alexandru Nica and Roland Speicher, *Lectures on the combinatorics of free probability*, Lond. Math. Soc. Lect. Note Ser., vol. 335, Cambridge: Cambridge University Press, 2006.
- [13] Dan Romik, *The surprising mathematics of longest increasing subsequences*, Institute of Mathematical Statistics Textbooks, vol. 4, Cambridge University Press, New York, 2015. MR3468738
- [14] P. Śniady, *Gaussian fluctuations of characters of symmetric groups and of Young diagrams*, Probab. Theory and Rel. Fields **136** (2006), no. 2, 263–297. MR2240789
- [15] A. M. Vershik and S. V. Kerov, *Asymptotics of the Plancherel measure of the symmetric group and the limit form of Young tableaux*, Soviet Math. Dokl. **18** (1977), 527–531.