

ON UNIQUENESS OF KP SOLITON STRUCTURES

FRANCISCO ALEGRÍA, GONG CHEN, CLAUDIO MUÑOZ, FELIPE POBLETE,
AND BENJAMÍN TARDY

ABSTRACT. We consider the Kadomtsev-Petviashvili II (KP) model placed in $\mathbb{R}_t \times \mathbb{R}_{x,y}^2$, in the case of smooth data that are not necessarily in a Sobolev space. In this paper, the subclass of smooth solutions we study is of “soliton type”, characterized by a phase $\Theta = \Theta(t, x, y)$ and a unidimensional profile F . In particular, every classical KP soliton and multi-soliton falls into this category with suitable Θ and F . We establish concrete characterizations of KP solitons by means of a natural set of nonlinear differential equations and inclusions of functionals of Wronskian, Airy and Heat types, among others. These functional equations only depend on the new variables Θ and F . A distinct characteristic of this set of functionals is its special and rigid structure tailored to the considered soliton. By analyzing Θ and F , we establish the uniqueness of line-solitons, multi-solitons, and other degenerate solutions among a large class of KP solutions. Our results are also valid for other 2D dispersive models such as the quadratic and cubic Zakharov-Kuznetsov equations.

1. INTRODUCTION AND MAIN RESULTS

1.1. **Setting of the problem.** Let $t \in \mathbb{R}$, and $(x, y) \in \mathbb{R}^2$. In this work we consider the KP-II model

$$(-4u_t + u_{xxx} + 6uu_x)_x + 3u_{yy} = 0, \quad (1.1)$$

where $u = u(t, x, y) \in \mathbb{R}$ is the unknown. The KP equations are canonical integrable models in 2D and were first introduced by Kadomtsev and Petviashvili in 1970 [14] for modeling “long and weakly nonlinear waves” propagating essentially along the x direction, but with a small dependence on the y variable. A rigorous derivation of both models from the Boussinesq system was obtained by Lannes and Lannes-Saut [24, 25].

The KP-II model (1.1) (KP from now on, if there is no confusion) has an important set of symmetries. If $u = u(t, x, y)$ is a solution to (1.1), then $u(t + t_0, x + x_0, y + y_0)$, with $t_0, x_0, y_0 \in \mathbb{R}$, $cu(c^{3/2}t, c^{1/2}x, cy)$, if $c > 0$, and if $\beta \in \mathbb{R}$ is a given speed,

$$u\left(t, x - \frac{4\beta}{3}y + \frac{4\beta^2}{3}t, y - 2\beta t\right)$$

Date: May 14, 2024.

F.A.’s work is partially supported by ANID Exploration project 13220060.

G. C. was partially funded by Simons fundation MP-TSM-00002258.

C.M. was partially funded by Chilean research grants FONDECYT 1231250, ANID Exploración 13220060, MathAmSud WAFFLE 23-MATH-18, and Basal CMM FB210005.

F.P.’s work is partially supported by ANID Exploration project 13220060, ANID project FONDECYT 1221076 and MathAmSud WAFFLE 23-MATH-18.

B.T. was partially funded by Chilean research grants FONDECYT 1231250 and 1221076 and Basal CMM FB210005.

(Galilean invariance) define new solutions to KP.

Our purpose here is to establish uniqueness results for KP solitons in the class of smooth solutions that is of soliton type, complementing the results obtained in [6]. More precisely, assume that u in (1.1) is sufficiently smooth and has the form

$$u(t, x, y) = 2\partial_x^2 F(\Theta(t, x, y)), \quad (1.2)$$

where, for some $s_0 > 0$, $F : [s_0, \infty) \rightarrow \mathbb{R}$ and $\Theta = \Theta(t, x, y) \in [s_0, \infty)$ are smooth functions. Later, we will justify that without loss of generality, one can choose $s_0 = 1$. Consequently, we shall assume this particular choice throughout this paper. Since F can be changed by any linear affine function, we can also assume that $F(1) = 0$ and $F'(1) = 1$. We shall call Θ the *phase* of the solution u , and F will be the *profile*. Every classical KP soliton is in this class for suitable F and Θ . Indeed, one of the most important examples of solutions in the form of (1.2) is provided by classical solitons and multi-solitons (see, e.g., Kodama [22]), namely solutions of the KP equation in the form of

$$\begin{aligned} u(t, x, y) &= 2\partial_x^2 \log(\Theta(t, x, y)), \\ \Theta(t, x, y) &:= \text{Wr}(\Theta_1, \dots, \Theta_n)(t, x, y), \\ \Theta_i(t, x, y) &:= \sum_{j=1}^{M_i} a_{ij} \exp(k_{ij}x + k_{ij}^2 y + k_{ij}^3 t), \end{aligned} \quad (1.3)$$

where Wr represents the classical Wronskian of n functions, and k_{ij}, a_{ij} are in principle just real-valued, although specific values determine precise solutions. In this case, $F = \log$ and Θ has the additional scaling symmetry: if Θ is a valid phase, then $\lambda\Theta$ in (1.3) also does, for any $\lambda > 0$. This fact motivates the reason why requiring $\Theta > 1$ is in principle not extremely restrictive. Note that, in the particular case where $F = \log$, given $u = u(t, x, y)$ solution of (1.1), the formula $\Theta = \exp\left(\int_0^x \int_0^t \frac{1}{2}u(s, r, y) dr ds\right)$ returns a valid phase Θ . Consequently, despite some loss of regularity, solving an equation for the profile F and the phase Θ may be considered as general as dealing with the original KP model (1.1) for u .

Taking into account the great diversity of KP solutions, our study will focus on a simple but useful characterization of soliton solutions. It turns out that this question is interesting and quite challenging, since Θ can assume complicated values, from simple linear functions to periodic ones. Also, the question of whether or not profile $F = \log$ is the only possibility for F is also extremely interesting.

The first step is to rewrite KP (1.1) in terms of F and Θ as in (1.2). After rearrangements, and assuming convergence to zero at infinity, we obtain the following fourth order equation for (F, Θ) :

$$\begin{aligned} 0 &= (F'''' + 6F''^2)(\Theta)\Theta_x^4 + 6(F'' + F'^2)'(\Theta)\Theta_x^2\Theta_{xx} + 3(F'' + F'^2)(\Theta)(\Theta_{xx}^2 + \Theta_y^2) \\ &\quad - 4F''(\Theta)\Theta_x(\Theta_t - \Theta_{xxx}) + F'(\Theta)(-4(\Theta_t - \Theta_{xxx})_x) \\ &\quad + 3F'(\Theta)(\Theta_{yy} - \Theta_{xxxx}) + 3F'^2(\Theta)(\Theta_{xx}^2 - \Theta_y^2). \end{aligned} \quad (1.4)$$

This is the equation that will be worked in this paper. It represents a highly nonlinear equation for the two unknowns F and Θ , but its nature is certainly better than (1.1), since it possesses hidden structures. The most important is a splitting phenomenon between some parts concerning only F -motivated terms, and others only dealing with Θ -related

terms. Of course, this is far from being an exact “separation of variables” as in classical linear PDEs, but we shall mention important similarities. Indeed, it is possible to divide (1.4) into three somehow well-defined sub-equations:

- (a) The first line, that can be written in terms of

$$\rho(s) := (F'' + F'^2)(s), \quad (1.5)$$

corresponding to those terms that are equal to 0 when $\rho \equiv 0$. And $\rho \equiv 0$ when $F = \log$ and some particular initial conditions are met.

- (b) The second line involves a modified Airy function $Ai(\Theta) := \Theta_t - \Theta_{xxx}$ and its derivative with respect to the variable x . This is a reminiscent of the 1D variable x that has a natural Airy structure for Θ (see Appendix B for further details).
(c) The third line has a complex structure represented by a Heat type term defined by $H(\Theta) := \Theta_y - \Theta_{xx}$. Notice that H is an operator in the y and x variables only.
(d) Additionally, there is a hidden structure in (1.4) represented by Wronskian type functions. Later, in Definition 1.1 we will explain better this structure.

The Heat and Airy functions are classical in the KP literature, see e.g. Kodama [22], but Wronskians and ρ functions are, as far as we understand, not so well-understood. Translated to the equation (1.4), the purpose of this work is to find suitable conditions on the Airy, Heat, Wronskian and ρ functions that characterize different soliton solutions. We consider the following definitions:

Definition 1.1 (Classification of phases Θ). We shall say that Θ as in (1.2)

- (i) is of Airy type if for all $(t, x, y) \in \mathbb{R}^3$,

$$Ai(\Theta) := \Theta_t - \Theta_{xxx} = 0;$$

- (ii) is of Heat type if for all $(t, x, y) \in \mathbb{R}^3$,

$$H(\Theta) := \Theta_y - \Theta_{xx} = 0;$$

- (iii) is of x -Wronskian type and y -Wronskian type if $\Theta > 0$, and

$$W_x(\Theta) := \Theta_{xxxx} - \frac{\Theta_{xx}^2}{\Theta} = 0, \quad W_y(\Theta) := \Theta_{yy} - \frac{\Theta_y^2}{\Theta} = 0, \quad (1.6)$$

respectively, for all $(t, x, y) \in \mathbb{R}^3$;

- (iv) is of \mathcal{T} -type if for F fixed,

$$\begin{aligned} \mathcal{T}(\Theta) := & -4F''(\Theta) Ai(\Theta) \Theta_x \\ & + F'(\Theta) \left(-4Ai(\Theta)_x + 3 \left(H(\Theta)_y + H(\Theta)_{xx} \right) \right) \\ & - 3F'^2(\Theta) H(\Theta) (\Theta_y + \Theta_{xx}) = 0, \end{aligned} \quad (1.7)$$

for all $(t, x, y) \in \mathbb{R}^3$.

Notice that we ask for equality for all $(t, x, y) \in \mathbb{R}^3$, while relaxing these conditions are interesting options not considered in this work. Before stating our main results, some important comments are necessary:

Remark 1.2 (On the classical operators). Heat and Airy operators are naturally involved in (1.4). Indeed, it can be proved (see Section 2 for further details) that (1.4) can be written as

$$\begin{aligned} & (\rho'' - 2F'(\Theta)\rho' + 4F''(\Theta)\rho)\Theta_x^4 + 6\rho'\Theta_x^2\Theta_{xx} \\ & - 4F''(\Theta)\Theta_x Ai(\Theta) + F'(\Theta)\left(-4Ai(\Theta)_x + 3\left(H(\Theta)_y + H(\Theta)_{xx}\right)\right) \\ & + 3\left(\rho(\Theta_{xx}^2 + \Theta_y^2) - F'^2(\Theta)H(\Theta)(\Theta_y + \Theta_{xx})\right) = 0, \end{aligned} \quad (1.8)$$

where from (1.5) $\rho(s) = F''(s) + F'^2(s)$. On the other hand, the operator \mathcal{T} is the natural counterpart of the ODE type satisfied by ρ , in the sense that (1.4)-(1.8) reads

$$\Theta_x^4 \rho'' + 2(6\Theta_x^2 \Theta_{xx} - 2F'(\Theta)\Theta_x^4)\rho' + (3(\Theta_{xx}^2 + \Theta_y^2) + 4F''(\Theta)\Theta_x^4)\rho + \mathcal{T}(\Theta) = 0. \quad (1.9)$$

Finally, notice that Θ being of \mathcal{T} -type is a condition depending on the profile F , and consequently is a more complex condition than being of Airy or Heat type, which are independent of the profile F .

Remark 1.3 (On the Wronskian operators). The emergence of the Wronskians (1.6) in (1.4) seems obscure and nonstandard. However, it is possible to rewrite (1.4) as

$$\begin{aligned} & (F'''' + 6F'^2)(\Theta)\Theta_x^4 + 6(F'' + F'^2)'(\Theta)\Theta_x^2\Theta_{xx} + 3(F'' + F'^2)(\Theta)(\Theta_{xx}^2 + \Theta_y^2) \\ & - 4F''(\Theta)\Theta_x Ai(\Theta) - 4F'(\Theta)Ai(\Theta)_x \\ & + 3F'(\Theta)(W_y^F(\Theta) - W_x^F(\Theta)) = 0, \end{aligned} \quad (1.10)$$

with W_x^F and W_y^F generalized Wronskian functionals,

$$\begin{aligned} W_y^F(\Theta) & := \Theta_{yy} - F'(\Theta)\Theta_y^2, \\ W_x^F(\Theta) & := \Theta_{xxx} - F'(\Theta)\Theta_{xx}^2. \end{aligned} \quad (1.11)$$

Later we will prove that under $H(\Theta) = 0$, one has $W_x^F(\Theta) - W_y^F(\Theta) = W_x(\Theta) - W_y(\Theta) = 0$, namely one can assume that $F = \log$ in (1.11), leading to the natural definitions in (1.6). In that sense, null Wronskians are naturally related to the Heat condition $H(\Theta) = 0$, however, the equivalence will not be as exact as one would prefer.

1.2. KP solitons. The soliton family stands out as one of the most distinctive features within the KP model. Distinguished by their complexity and rich character, numerous works have been dedicated to understanding them, employing integrability, algebraic, and analytic techniques. Among these we find the works by Kodama and Williams [20, 21], which provide a precise description of KP-solitons within the positive Grassmannian. For a comprehensive and detailed overview of this line of research, see also Kodama's monograph [22].

The line soliton family (see [22]) is given by

$$\Theta(t, x, y) = a_1 \exp(\theta_1) + a_2 \exp(\theta_2), \quad (1.12)$$

where $a_1, a_2 > 0$, and $\theta_j := k_j x + k_j^2 y + k_j^3 t$, $k_1, k_2 \in \mathbb{R}$. Assuming $F = \log$, the corresponding KP solution via (1.2) is given by

$$u(t, x, y) = \frac{1}{2}(k_1 - k_2)^2 \operatorname{sech}^2\left(\frac{1}{2}(\theta_1 - \theta_2)\right). \quad (1.13)$$

See Fig. 1 (left) for details. The classical KdV soliton is recovered by setting $k_1 = -k_2 = k$, and in this case u becomes

$$Q_k(t, x) := 2k^2 \operatorname{sech}^2(kx + k^3t). \quad (1.14)$$

The next case of KP solution is the *resonant multi-soliton*. This corresponds to the case

$$\Theta(t, x, y) = \sum_{i=1}^M a_i \exp(\theta_i) = \sum_{i=1}^M a_i \exp(k_i x + k_i^2 y + k_i^3 t), \quad (1.15)$$

where to ensure the positivity and nondegeneracy of Θ we impose that each $a_i > 0$ and $k_1 < k_2 < \dots < k_M$. A special case of resonant soliton is given in Fig. 1 (right).




FIGURE 1. Left: One line-soliton solution (1.13) with $k_1 = -0.5$, $k_2 = 1$ and $t = 0$. This solution divides the plane into two regions according to the sum (1.12), and on each region a different exponential dominates. Center: A 2-soliton of KP with $k_1 = -1$, $k_2 = -0.5$, $k_3 = 0.5$ and $k_4 = 1$, at time $t = 0$. Right: A Y-soliton characterized by $\Theta = \exp(\theta_1) + \exp(\theta_2) + \exp(\theta_3)$ in (1.15). Notice that the coefficients a_j are set equal to 1, so that the three solitons meet at the origin at $t = 0$. Here, $(k_1, k_2, k_3) = (-0.3, 0, 0.5)$.

Now we recall the KP 2-soliton. In this case $\Theta = \operatorname{Wr}[\Theta_1, \Theta_2]$, where Wr is the Wronskian of two functions $\Theta_1 = \exp(\theta_1) + \exp(\theta_2)$ and $\Theta_2 = \exp(\theta_3) + \exp(\theta_4)$ being 1-soliton phases. Calculating the phase Θ , one obtains the classical formula

$$\begin{aligned} \Theta = & (k_3 - k_1) \exp(\theta_1 + \theta_3) + (k_4 - k_1) \exp(\theta_1 + \theta_4) \\ & + (k_3 - k_2) \exp(\theta_2 + \theta_3) + (k_4 - k_2) \exp(\theta_2 + \theta_4). \end{aligned} \quad (1.16)$$

In order to ensure the positivity and nondegeneracy of Θ , we require $k_1 < k_2 < k_3 < k_4$. See Fig. 1 (center) for further details on the family of KP 2-solitons.

1.3. Main Results. This paper represents a departure from previous approaches, in a sense to be explained below. We adopt the perspective that each KP soliton should adhere to a specific “variational” characterization, manifested in their critical points of a suitable nonlinear functional. Our primary aim is to offer clear and straightforward characterizations of the most distinct KP solitons using simple ‘trapping’ functionals that exhibit rigidity properties. This represents an initial step in the direction previously outlined. We believe that this concept holds promising potential for applications not only in elucidating more complex KP solutions but also in other related dispersive models.

Our first result is a characterization of the KdV line soliton as a KP solution.

Theorem 1.4. *Let u be a smooth solution to KP (1.1) of the form (1.2), with a smooth profile F such that $F(1) = 0$, $F'(1) = 1$, $F''(1) = -1$, and $F'''(1) = 2$. Then u is a KdV soliton and $F = \log$ if and only if $H(\Theta) = Ai(\Theta) = W_y^F(\Theta) = 0$.*

Remark 1.5. The four initial conditions on F may seem extremely demanding, however they are naturally explained by the fourth order equation representing (1.1). Consequently, in order to determine F , the four derivatives on x at (1.1) induce corresponding initial conditions for F and its three first derivatives. It is also easy to see that different initial conditions may lead to other solutions (F periodic, for instance), as it happens in the simpler KdV case. See [16, 3] for instance for examples of space-periodic profiles F .

Remark 1.6. It will be proved below (see Lemma 3.5) that the Heat condition $H(\Theta) = 0$ implies $W_x^F(\Theta) = W_y^F(\Theta)$ and $W_x(\Theta) = W_y(\Theta)$. So, additionally, one has $W_x(\Theta) = 0$.

As a natural consequence of Theorem 1.4, we obtain the following uniqueness result of KdV solitons as extended KP solutions. As usual, we require

$$F(1) = 0, \quad F'(1) = 1, \quad F''(1) = -1, \quad \text{and} \quad F'''(1) = 2. \quad (1.17)$$

Corollary 1.7. *Assume (1.17). Let u be a nontrivial KP solution of the form (1.2) such that $\Theta > 0$ is a solution to $H(\Theta) = Ai(\Theta) = W_y^F(\Theta) = 0$. Then $F = \log$ and $u = Q_k$ in (1.14) for some $k \in \mathbb{R} - \{0\}$.*

Theorem 1.4 can be extended to general KP line-solitons (1.13) as in Fig. 1 left, and not necessarily of KdV type. In this case, we denote them oblique line-solitons. From now on, the structure $\Theta W_y(\Theta)$ from (1.6) will be essential.

Theorem 1.8. *Let u be a smooth solution to (1.1) of the form (1.2), with a smooth profile F such that (1.17) is satisfied. Then u is an oblique line-soliton of the form (1.12)-(1.13) and $F = \log$ if and only if $H(\Theta) = Ai(\Theta) = 0$, and*

$$\Theta W_x(\Theta) = \Theta W_y(\Theta) = A(t, x) \exp(k(t, x)y), \quad (1.18)$$

for some particular functions $A, k > 0$ everywhere.

It is worth to mention that the conditions $H(\Theta) = Ai(\Theta) = 0$ do not ensure the validity of Theorem 1.8. Indeed, it will be proved that the class of phases satisfying these two conditions is large enough to contain many multi-soliton solutions, such as the Y structure defined below, which is not a line soliton. Consequently, (1.18) is a necessary condition.

Remark 1.9. Notice that $\Theta W_x(\Theta)$ remains unchanged after a Galilean transformation (see (A.1) in Appendix A). This is not the case for $\Theta W_y(\Theta)$. However, one can prove that if Θ is of the form (1.12), then coincidentally for $\beta = k_1 + k_2$ one has that the Galilean transformation of Θ , denoted Θ_β (see (2.1)), satisfies $\Theta_\beta W_y(\Theta_\beta) = 0$. It is an interesting problem to fully elucidate the role of Galilean transforms in the classification of solitons as proposed in this paper.

Theorem 1.8 puts in evidence an intriguing new mathematical structure, a natural finite-dimensional cone in the y variable.

Definition 1.10 (Invariant \mathcal{W}_n cones). Given $n \in \{1, 2, 3, \dots\}$, consider the linear, positively generated cone

$$\mathcal{W}_n := \left\{ \sum_{j=1}^n a_j(t, x) \exp(k_j(t, x)y) : \begin{array}{l} \exists 0 \leq k_1(t, x) < k_2(t, x) \cdots < k_n(t, x), \\ \exists a_1(t, x), a_2(t, x), \dots, a_n(t, x) \geq 0 \end{array} \right\}. \quad (1.19)$$

Using the terminology above, (1.18) can be recast as $\Theta W_x(\Theta) = \Theta W_y(\Theta) \in \mathcal{W}_1$.

The space \mathcal{W}_n has interesting properties, in particular its behavior under the nonlinear mapping $\Theta W_y(\Theta)$ is key to understand complex multi-soliton structures. First of all, $\Theta \in \mathcal{W}_M$ implies that $\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ (Lemma 4.7). Second, there is a natural “kernel” given by the function $\exp(ky)$, k arbitrary: one has $\exp(ky) W_y(\exp(ky)) = 0$ for any $k = k(t, x)$. Additionally, under the gauge $\mathcal{W}_M \ni \Theta \mapsto \Theta_k := \exp(ky)\Theta$, one has $\Theta_k W_y(\Theta_k) = \exp(2ky)\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$, revealing that *there is no unique nontrivial solution* to the set inclusion $\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ for $\Theta \in \mathcal{W}_M$. One way to repair this gauge freedom is to ask for $\Theta(y=0)$ and $\Theta_y(y=0)$ uniquely defined, as it is done below.¹ For natural reasons, we will work in a slightly larger class of resonant phases than (1.15), given by

$$\Theta(t, x, y) = \sum_{i=1}^M (a_{i,1} \exp(-k_i x + k_i^2 y + k_i^3 t) + a_{i,2} \exp(k_i x + k_i^2 y + k_i^3 t)), \quad (1.20)$$

with coefficients $k_1 < k_2 < \dots < k_M$ and $a_{i,j} > 0$.

Theorem 1.11 (Resonant multisolitons). *Let u be a solution of (1.1) of the form (1.2) with a smooth real-valued phase $\Theta > 0$ satisfying for $k = 0, 1, 2, 3$,*

$$\partial_x^k \Theta(t, 0, 0), \quad \partial_x^k \partial_y \Theta(t, 0, 0) \quad \text{uniquely prescribed.} \quad (1.21)$$

Assume that the smooth profile F such that (1.17) holds. Then Θ corresponds to an M resonant multi-soliton (1.20) and $F = \log u$ if and only if Θ satisfies $H(\Theta) = Ai(\Theta) = 0$ and $\Theta W_y(\Theta) = \Theta W_x(\Theta)$ has a unique value in $\mathcal{W}_{\frac{1}{2}M(M-1)}$.

Remark 1.12. Notice that the condition (1.21) only requires information of Θ at $x = y = 0$. This is necessary to ensure the uniqueness of the solution Θ to $\Theta W_y(\Theta) = \Theta W_x(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ within the class \mathcal{W}_M . Due to the exponential growth in x of the functions solving the equations for Θ , a less demanding sufficient condition will require to establish a Cauchy theory for the linear Airy equation $\Theta_t - \Theta_{xxx} = 0$ with initial conditions in the distributional class $D'(\mathbb{R}_x)$, a problem that is far from trivial due to the oscillatory character of the Airy kernel.

The key in the proof of Theorem 1.11 is the property that $\Theta \in \mathcal{W}_M$ implies $\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ (Lemma 4.7). This property allows us to estimate the size of the cone representing the image of \mathcal{W}_M under the nonlinear mapping $\Theta W_y(\Theta)$. Then one has to establish a sort of uniqueness in the representation of Θ , which in the case $M = 1$ is easy to obtain (see Theorem 1.8), but in the general case it is not known to hold in generality. Under the additional prescribed data at $x = y = 0$, then uniqueness is recovered and Theorem 1.11 establishes the equivalence between resonant multi-solitons and Airy-Heat type phases with finite dimensional Wronskians.

¹On the other hand, it is easy to check that

$$\mathcal{W} := \bigcup_{n \geq 0} \mathcal{W}_n$$

is a multiplicative algebra under nonnegative coefficients. With respect to the y variable, standard topological arguments ensure that the closure of this space restricted to any compact set K of \mathbb{R} , under the uniform norm, is nothing but $C_+(K, \mathbb{R})$, the space of real-valued, nonnegative continuous functions defined on K .

Resonant solitons of Y -type, or Miles type (see Fig. 1 right), are essential KP solutions included in the previous result. These are usually given by [22] ($k_1 < k_2 < k_3$)

$$\Theta = a_1 \exp(\theta_1) + a_2 \exp(\theta_2) + a_3 \exp(\theta_3), \quad a_i > 0, \quad \theta_i := k_i x + k_i^2 y + k_i^3 t.$$

Sometimes referred as resonant interacting 3-solitons, Theorem 1.11 states that they are characterized as having zero Heat and Airy operators, but having $\Theta W_y(\Theta)$ with one more dimension than the one obtained in Theorem 1.8, measured in terms of the subspace \mathcal{W}_2 . Additionally, in this case $\Theta \in \mathcal{W}_3$ and $\Theta W_y(\Theta) \in \mathcal{W}_3$, being the only phases (as far as we understand) that have this invariance under the nonlinear mapping $\Theta \mapsto \Theta W_y(\Theta)$.

Our last result concerns the characterization of crossed 2-solitons, with Θ given in (1.16) (see also Fig. 1 right panel). Recall the subspace \mathcal{W}_n defined in (1.19).

Theorem 1.13 (2-solitons). *Let u be a solution of (1.1) of the form (1.2) with $F = \log$ and with a smooth real-valued phase $\Theta > 0$ satisfying (1.21) and being at most exponentially growing in x : there are $C_1, c_2 > 0$ such that*

$$|\Theta(t, x, y)| \leq C_1 e^{c_2|x|}.$$

Then Θ corresponds to a 2-soliton (1.16) with $k_1 < k_2 < k_3 < k_4$ if and only if

- (1) $H(\Theta), Ai(\Theta)$ are contained in \mathcal{W}_4 ,
- (2) $\Theta W_y(\Theta), \Theta W_x(\Theta)$ describe unique elements in \mathcal{W}_5 , and
- (3) $Ai(\Theta) = \frac{3}{2} \partial_x H(\Theta)$.

Remark 1.14. Notice that Theorem 1.13 does not assume that $\Theta = \text{Wr}[\Theta_1, \Theta_2]$ (standard Wronskian of Θ_1 and Θ_2), with $H(\Theta_i) = Ai(\Theta_i) = 0$ and $\Theta_1, \Theta_2 \in \mathcal{W}_2$. This is the standard and well-known definition of the 2-soliton that assumes the key Wronskian substructure. Here we lift that condition and only ask conditions on Θ itself. The Wronskian structure is recovered from the proof.

Remark 1.15. Theorem 1.13 can be recast as follows. It is easy to check that 2-solitons solve $\mathcal{T}(\Theta) = 0$ (Corollary 4.10). However, this equation has plenty of additional, more complicated solutions [22], and a suitable characterization of the 2-soliton subclass is desirable. In that sense, Theorem 1.13 and the \mathcal{W}_n structure give a precise equivalence that separates 2-solitons of other more complex KP solutions.

1.4. Further results. Our last comments are related to possible extensions of the results presented in this paper. We believe that with some work it is possible to give a suitable characterization of KP-I line-soliton and lumps in terms of particular phases. The challenge is to get a good understanding of the fact that KP-I lumps are degenerate soliton solutions, in a sense already described in [1]. However, in the Zakharov-Kuznetsov (ZK) case,

$$-4\partial_t u + \partial_{x_1}(\Delta u + 3u^2) = 0, \tag{1.22}$$

where $u = u(t, x)$, $x = (x_1, x') \in \mathbb{R}^d$, $x_1 \in \mathbb{R}$, $x' \in \mathbb{R}^{d-1}$, one can say the following:

Theorem 1.16 (ZK case). *Let $u = 2\partial_{x_1}^2 F(\Theta)$ be a smooth solution of (1.22) with $\Theta > 0$ and (1.17) satisfied. Then $u = Q_k$, $k > 0$ (the KdV soliton) as in (1.14) and $F = \log$ if and only if*

$$Ai(\Theta) = W_1(\Theta) = W_{x_j}^F(\Theta) = 0, \quad j = 2, \dots, d, \tag{1.23}$$

and where

$$W_1(\Theta) := (\partial_{x_1}^2 \Theta)^2 - \partial_{x_1} \Theta \partial_{x_1}^4 \Theta, \quad W_{x_j}^F(\Theta) := \partial_{x_j}^2 \Theta - F'(\Theta)(\partial_{x_j} \Theta)^2.$$

Similarly, in the modified Zakharov-Kuznetsov (mZK) case,

$$(-4u_t + \partial_{x_1}^3 u + 6u^2 \partial_{x_1} u) + \partial_{x_1}(\Delta_c u) = 0, \quad (1.24)$$

we have the following result:

Theorem 1.17 (mZK case). *Let $u = 2\partial_{x_1} F(\Theta)$ be a smooth solution of (1.24) with smooth profile $F : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $F(0) = 0$ and F strictly increasing in \mathbb{R} . Then a nontrivial Θ is a mKdV soliton (B.11) and $F = 2 \arctan$, and only if*

$$A_i(\Theta) = W(\Theta) = \Lambda_{x_j}^F(\Theta) = 0, \quad j = 2, \dots, d, \quad (1.25)$$

and where

$$W(\Theta) := \Theta_{xx}^2 - \Theta_x \Theta_{xx}, \quad \Lambda_{x_j}^F(\Theta) := \Theta_{x_j x_j} F'(\Theta) + \Theta_{x_j}^2 F''(\Theta).$$

Remark 1.18 (The KdV and ZK cases). An important outcome of the proofs will be its robust character. Indeed, Theorem 1.4 has natural counterparts in the case of the 1D KdV model and ZK model (Theorem 1.16), where similar notions of Airy and Heat operators are introduced. See Appendix B and Section 6 respectively, for the complete details.

Remark 1.19 (The mKdV and mZK cases). In concordance with the previous remark, it should be natural that the same ideas can be applied as well for the so-called mKdV and mZK models. It turns out that this models has a particular rich structure involving even more demanding special solutions that complicates matters. We provide for completeness a suitable short treatment of the problem in Appendix B and Subsection 6.1. See also [6] for a detailed account of the difficulties found when dealing with mKdV models.

1.5. Previous results. We mention some key results obtained for KP-II models during the past years. Bourgain [4] showed that KP is globally well-posed (GWP) in $L^2(\mathbb{R}^2)$ (see also Ukai [38] and Iório-Nunes [11] for early results). Bourgain's result was later improved by Takaoka-Tzvetkov [37], Isaza-Mejia [12], Hadac [7] and Hadac-Herr-Koch [8]. Molinet, Saut and Tzvetkov [32] proved global well-posedness of KP along the KdV line-soliton in $L^2(\mathbb{R} \times \mathbb{T})$ and $L^2(\mathbb{R}^2)$.

The long time behavior of small KP solutions has been studied by Hayashi-Naumkin-Saut and Hayashi-Naumkin [10, 9], see also recent improvements by Niizato [33]. de Bouard-Martel [5] showed that KP has no ‘‘lump’’ structures, namely compact-in-space solutions. Any KdV soliton becomes an (infinite energy) line-soliton solution of KP. This structure is stable, as proved by Mizumachi and Tzvetkov [31], and asymptotically stable, see Mizumachi [28, 29]. The linear stability of the 2-soliton was recently proved by Mizumachi [30]. Finally, Izasa-Linares-Ponce [13] showed propagation of regularity for this model.

Numerical studies of KP solutions have been performed in [19], see also [17, 18] for a detailed account of the KP literature via PDE methods. Multi-line-soliton structures are known to exist via Inverse Scattering Transforms (IST) methods [1, 23]. Their stability in rigorous terms has been recently considered by Wu [40, 41]. See also [34, 35] for a detailed theory of transversal stability and instability of PDE models of water waves, that applies to one line-solitons as the ones studied in Theorem 1.4-1.8. The description of small data can be found in [39, 36]. Recently, and following [26], in [27] it was shown that every solution u of KP obtained from arbitrary initial data u_0 in $L^2(\mathbb{R}^2)$ satisfies $\liminf_{t \rightarrow \infty} \int_K u^2(t, x, y) dx dy = 0$, with $K \subseteq \mathbb{R}^2$ compact. Finally, Kenig and Martel [15]

showed that for any $\beta > 0$ and initial data small in $L^1 \cap L^2$, $\lim_{t \rightarrow \infty} \int_{x > \beta t} u^2(t, x, y) \, dx dy = 0$.

Organization of this work. This work is organized as follows. In Section 2 we introduce the basic elements needed for the proof of the main results. Section 2.3 is devoted to recall standard results on KP solitons. In Section 3 we present general results and properties about phases Θ satisfying (1.4). On the other hand, Section 4 presents properties satisfied by classical soliton structures. Finally, in Section 5 we prove the main results, Theorems 1.4, 1.8, 1.11, and 1.13. Section 6 considers the ZK and mZK cases. Appendix A contains some useful computations needed in the paper, and finally Appendix B is devoted to the proof of similar results in the case of KdV and mKdV models.

Acknowledgments. Part of this work was done while the third and fourth authors were visiting UACH (Valdivia, Chile), Georgia Tech and Texas A&M Mathematics departments. We thank these institutions for their warm hospitality and support.

2. PRELIMINARIES

In this section we first mention some simple but important facts related to solutions of the KP model.

2.1. Invariances. First of all, it is clear that (1.4) is invariant under space and time shifts in the phase Θ . Additionally, (1.4) has the natural scaling invariance associated to KP:

$$\Theta(t, x, y) \longrightarrow \Theta(\lambda^3 t, \lambda x, \pm \lambda^2 y), \quad \lambda > 0.$$

KP-II obeys the Galilean Transform [27]

$$u(t, x, y) \longrightarrow u\left(t, x - \frac{4\beta}{3}y + \frac{4\beta^2}{3}t, y - 2\beta t\right) =: u_\beta(t, \tilde{x}, \tilde{y}), \quad \beta \in \mathbb{R}.$$

This invariance naturally translates into the phase Θ

$$\Theta_\beta(t, x, y) = \Theta\left(t, x - \frac{4\beta}{3}y + \frac{4\beta^2}{3}t, y - 2\beta t\right) = \Theta(t, \tilde{x}, \tilde{y}), \quad (2.1)$$

that also satisfies (1.4) provided Θ does.

2.2. Kernel in the solitonic representation. Notice that the formulation (1.2) involves a nontrivial kernel.

Lemma 2.1. *One has $\partial_x^2 \log(\Theta(t, x, y)) = 0$ for all $(t, x, y) \in \mathbb{R}^3$ if and only if the phase Θ satisfies $\Theta(t, x, y) = \exp(a(t, y)x + b(t, y))$, for any well-defined functions a and b .*

Proof. It is a consequence of direct integrations. \square

As a consequence of the previous result, if $F = \log$, any phase that can be expressed in the form $\Theta = \exp(f(t, x, y))$ with $f(t, x, y)$ any smooth linear affine function in the x variable gives a trivial solution. This is the kernel of the operator $\partial_x^2 \log$ which permits to construct the KP multi-soliton solutions, based on this first seed.

2.3. Quick review of the simplest KP solitons. Recall the line soliton introduced in (1.12) and (1.13). The line that separates these regions correspond to $\theta_1 = \theta_2$. In this case, this line is called $[1, 2]$ -soliton [22]. In the general case of resonant structures such as (1.15), they are called $[i, j]$ -solitons and are formed by the intersection of the corresponding exponentials i and j . Each $[i, j]$ -soliton has the same local structure as a line-soliton, which is described by the form [22, p. 5]

$$u = A_{[i,j]} \operatorname{sech}^2 \left(\frac{1}{2} \left(K_{[i,j]} \cdot (x, y) - \Omega_{[i,j]} t + \Theta_{[i,j]}^0 \right) \right),$$

with $\Theta_{[i,j]}^0$ a constant. The parameters $A_{[i,j]}$, $K_{[i,j]}$ and $\Omega_{[i,j]}$ are known as the amplitude, wave-vector and frequency, respectively, and are defined by

$$\begin{aligned} A_{[i,j]} &= \frac{1}{2} (k_j - k_i)^2, \\ K_{[i,j]} &= (k_j - k_i, k_j^2 - k_i^2) = (k_j - k_i) (1, k_j + k_i), \\ \Omega_{[i,j]} &= - (k_j^3 - k_i^3) = - (k_j - k_i) (k_i^2 + k_i k_j + k_j^2). \end{aligned}$$

If one denotes $\psi_{[i,j]}$ the angle measured counterclockwise between the $[i, j]$ -soliton and the y -axis, then $\tan(\psi_{[i,j]}) = k_i + k_j$.

Now consider the case of resonant solitons (1.15) with $M = 3$. As in the previous case, it is possible to determine the dominant exponentials and analyze the structure of the solution in the xy -plane. Indeed, the line-soliton at y positive, corresponding to the $[1, 3]$ -soliton, is located on the phase transition $x + cy = \text{constant}$ with direction parameter $c = k_1 + k_3$. In the same way, the line-soliton located at y negative, corresponding to the $[1, 2]$ -soliton and $[2, 3]$ -soliton are located over their respective phase transitions with direction parameter $c = k_1 + k_2$ and $c = k_2 + k_3$, respectively. The *resonance condition* of these three line-solitons is given by

$$K_{[1,3]} = K_{[1,2]} + K_{[2,3]}, \quad \Omega_{[1,3]} = \Omega_{[1,2]} + \Omega_{[2,3]},$$

and both are satisfied when $K_{[i,j]} = (k_j - k_i, k_j^2 - k_i^2)$ and $\Omega_{[i,j]} = - (k_j^3 - k_i^3)$.

In a simple way it is possible to extend the previous result for a general solution constructed from a Θ composed with an arbitrary number M of exponentials, as in (1.15).

Theorem 2.2 ([22], Proposition 1.2). *Let Θ be an M resonant phase as in (1.15). Then the solution u has the following asymptotic characteristics:*

- (i) *For values of $y \gg 1$, there is only one soliton of the form $[1, M]$ -soliton.*
- (ii) *For values of $y \ll -1$, there are $M - 1$ line-solitons of the form $[k, k + 1]$ -soliton, with $k = 1, 2, \dots, M - 1$, located counter-clockwise from the negative part to the positive part of the x -axis.*

Definition 2.3. Let $N < M$. The Grassmannian $\operatorname{Gr}(N, M)$ are all the matrices which represents an N -dimensional sub-vectorial space contained in a M -dimensional vectorial space.

It can be verified the following isomorphism: $\operatorname{Gr}(N, M) \cong \operatorname{GL}_N(\mathbb{R}) \setminus M_{N \times M}(\mathbb{R})$, where $\operatorname{GL}_N(\mathbb{R})$ consists of those matrices with dimension $n \times n$, real coefficients and whose determinant is non-zero.

Let $\{\Theta_i : i = 1, \dots, N\}$ be linearly independent solutions of $Ai(\Theta) = H(\Theta) = 0$. Let $\Theta := \text{Wr}(\Theta_1, \dots, \Theta_N)$ be the Wronskian of the functions Θ_i with respect to the x -variable (usually called a τ -function). It is well-known and not difficult to see (see [22]) that $u(t, x, y) = 2\partial_x^2 \ln(\Theta(t, x, y))$ satisfies the KP equation. A particular choice for Θ_i is given by

$$\Theta_i(t, x, y) = \sum_{j=1}^N a_{ij} \exp(\theta_j(t, x, y)), \quad \text{with} \quad \theta_j = k_j x + k_j^2 y + k_j^3 t,$$

where $A := (a_{ij})$ is an $N \times M$ matrix. Thus each KP soliton expressed in the previous form is parametrized by M parameters (k_1, \dots, k_M) and an $N \times M$ matrix A . The matrix A will be identified as a point of the real Grassmannian $\text{Gr}(N, M)$.

2.4. Linear ODEs related to F . In this section we describe some ODE theory related to the equations that F must satisfy. Indeed, from (1.5) consider the auxiliary variable

$$\rho(s) := F''(s) + F'^2(s).$$

Lemma 2.4. *If $F = \log$, then $\rho = 0$. If $\rho = 0$ and $F(1) = 0$, $F'(1) = 1$, then $F = \log$.*

Proof. It follows from directly solving the ODE $\rho(s) = 0$. \square

Lemma 2.5. *Let F be a smooth profile such that $F(1) = 0$ and $F'(1) = 1$. Then the following are satisfied, for any $s \geq 1$:*

- (i) *If $(F'' + F'^2)(s) = 0$, then $(F'''' + 6F''^2)(s) = 0$.*
- (ii) *If now $F''(1) = -1$ and $F'''(1) = 2$, and $(F'''' + 6F''^2)(s) = 0$, then one has $(F'' + F'^2)(s) = 0$.*
- (iii) *Assume now that $F(1) = 0$, $F'(1) = 1$, $F''(1) = -1$, $F'''(1) = 2$, $s \in [1, \infty)$ and*

$$h_1(s)(F'''' + 6F''^2)(s) + h_2(s)(F'' + F'^2)'(s) + h_3(s)(F'' + F'^2)(s) = 0, \quad (2.2)$$
for some continuous $h_1, h_2, h_3 : [1, \infty) \rightarrow \mathbb{R}$, $h_1 > 0$. Then $F = \log$.

Proof. The proof of (i) is a consequence of the following identity. One has

$$(F'' + F'^2)'' - 2F'(F'' + F'^2)' + 4F''(F'' + F'^2) = F'''' + 6F''^2. \quad (2.3)$$

Therefore, $F'' + F'^2 = 0$ implies $F'''' + 6F''^2 = 0$, proving (i).

Proof of (ii). In the case where $F'''' + 6F''^2 = 0$ one has that (2.3) can be written in terms of ρ as

$$\rho'' - 2F'\rho' + 4F''\rho = 0. \quad (2.4)$$

This is a second order linear ODE with continuous coefficients. This solution has a basis of solutions of dimension 2, say $\{\rho_1, \rho_2\}$. Consequently,

$$\rho(s) = C_1\rho_1(s) + C_2\rho_2(s), \quad C_1, C_2 \in \mathbb{R}.$$

Recall that $F(1) = 0$ and $F'(1) = 1$, and $\rho = F'' + F'^2$, $\rho' = F''' + 2F'F''$. Since $F''(1) = -1$ and $F'''(1) = 2$, one has $\rho(1) = \rho'(1) = 0$, leading to $\rho(s) = 0$ for all $s \geq 1$, proving (ii).

Proof of (iii). Thanks to (2.3), equation (2.2) is equivalent to

$$h_1(s)(\rho''(s) - 2F'(s)\rho'(s) + 4F''(s)\rho(s)) + h_2(s)\rho'(s) + h_3(s)\rho(s) = 0,$$

with $\rho = F'' + F'^2$. This system is analogous to (2.4), and proceeding as in step (ii) we get from Lemma 2.4 that $\rho(s) = 0$ for all $s \geq 1$, and $F = \log$. \square

Recall from Definition 1.1 that (Θ, F) are of \mathcal{T} -type if (1.7) is satisfied. The following corollary is a direct result of (iii) in Lemma 2.4.

Corollary 2.6. *Let $u = 2\partial_x^2 F(\Theta)$ solution of (1.1), with $F(1) = 0$, $F'(1) = 1$, $F''(1) = -1$, and $F'''(1) = 2$. Then Θ is \mathcal{T} -type if and only if $F = \log$.*

Proof. Direct from (1.9) and (iii) in Lemma 2.5. \square

3. WRONSKIAN STRUCTURES

3.1. Simple Phases. The purpose of this section is to establish simple properties for the smooth phases satisfying (1.4). We start with the following

Lemma 3.1. *Assume that $\Theta > 0$. Then the following are satisfied:*

- (i) *If $F''(\Theta)\Theta_x^2 + F'(\Theta)\Theta_{xx} = 0$ for all $(t, x, y) \in \mathbb{R}^3$, then u is the trivial solution.*
- (ii) *If $F = \log$, then a positive phase Θ that satisfies $\Theta\Theta_{xx} = \Theta_x^2$, for all $(t, x, y) \in \mathbb{R}^3$, gives the trivial solution.*
- (iii) *If $F = \log$ then every Θ of the form $\Theta(t, x, y) = A(t, y) \exp(kx)$ with $A > 0$, gives the trivial solution.*

Proof. Proof of (i). Direct from the fact that $u = 2\partial_x^2 F(\Theta) = 2(F''(\Theta)\Theta_x^2 + F'(\Theta)\Theta_{xx})$. Proof of (ii): also direct from the fact that $F''(\Theta)\Theta_x^2 + F'(\Theta)\Theta_{xx} = \frac{\Theta\Theta_{xx} - \Theta_x^2}{\Theta^2}$, and the previous result. Finally, (iii) is consequence of the fact that in this case $\Theta F''(\Theta) = -F'(\Theta)$, for all $(t, x, y) \in \mathbb{R}^3$. \square

Remark 3.2. Note that in Lemma 3.1 we are assuming that the conditions are satisfied for all $(t, x, y) \in \mathbb{R}^3$. In the case where the conditions are satisfied just for certain points $(t, x, y) \in \mathbb{R}^3$, this points will correspond to zeros of the associated solution u .

Corollary 3.3. *If $\Theta > 0$ is a smooth phase such that $\Theta\Theta_{xx} - \Theta_x^2 = 0$, then $W_x(\Theta) = 0$.*

Proof. Taking derivative in x , $0 = \Theta_x\Theta_{xx} + \Theta\Theta_{xxx} - 2\Theta_x\Theta_{xx} = \Theta\Theta_{xxx} - \Theta_x\Theta_{xx}$. Once again, taking derivative in x , $0 = \Theta_x\Theta_{xxx} + \Theta\Theta_{xxxx} - \Theta_{xx}^2 - \Theta_x\Theta_{xxx} = \Theta W_x(\Theta)$. \square

3.2. General vs. simple Wronskians. It is noted that in Definition 1.1, Wronskians (1.6) do not coincide with the expected value if taken from (1.4). Indeed, the correct definition should be

$$\begin{aligned} W_y^F(\Theta) &:= \Theta_{yy} - F'(\Theta)\Theta_y^2, \\ W_x^F(\Theta) &:= \Theta_{xxx} - F'(\Theta)\Theta_{xx}^2. \end{aligned} \tag{3.1}$$

Here F is taken general. One recovers the values stated in Definition 1.1 if $F = \log$. Notice that from (1.10) in terms of $A_i(\Theta)$, one gets

$$\begin{aligned} (F'''' + 6F''^2)(\Theta)\Theta_x^4 + 6(F'' + F'^2)'(\Theta)\Theta_x^2\Theta_{xx} + 3(F'' + F'^2)(\Theta)(\Theta_{xx}^2 + \Theta_y^2) \\ - 4(F'(\Theta)A_i(\Theta))_x + 3F'(\Theta)(W_y^F(\Theta) - W_x^F(\Theta)) = 0. \end{aligned} \tag{3.2}$$

Having this structure in mind, let us study the phases related to the Wronskian conditions (1.6).

Lemma 3.4. *Assume that $u = 2\partial_x^2 \log(\Theta)$ is solution to KP with $\Theta > 0$ and $Ai(\Theta) = 0$. Then $W_y(\Theta) = W_x(\Theta)$.*

Proof. Since $F = \log$ one has from Lemmas 2.4 and 2.5 that $\rho = F'' + F'^2 = 0$ and $F'''' + 6F''^2 = 0$. Using $Ai(\Theta) = 0$, from (3.2) and the fact that $F' \neq 0$, we get

$$W_y^F(\Theta) - W_x^F(\Theta) = 0.$$

The conclusion is obtained by recalling that $F = \log$. \square

Now we put our attention to the following rigidity property.

Lemma 3.5. *Given any everywhere smooth functions F and $\Theta > 0$, if $H(\Theta) = 0$, then $W_y^F(\Theta) - W_x^F(\Theta) = W_y(\Theta) - W_x(\Theta) = 0$.*

Proof. Fix smooth functions F and $\Theta > 0$. By hypothesis $H(\Theta) = 0$. Then $H(\Theta)_y = H(\Theta)_{xx} = 0$. Consequently, $0 = H(\Theta)_y + H(\Theta)_{xx} = \Theta_{yxx} - \Theta_{xxxx} + \Theta_{yy} - \Theta_{xxy}$. Since $\Theta_{xxy} = \Theta_{yxx}$, one obtains $\Theta_{yy} = \Theta_{xxxx}$. Finally, from (3.1),

$$W_x^F(\Theta) = \Theta_{xxxx} - F'(\Theta)\Theta_{xx}^2 = \Theta_{yy} - F'(\Theta)\Theta_y^2 = W_y^F(\Theta).$$

This proves that $W_x^F(\Theta) = W_y^F(\Theta)$. Now we prove that the previous result is independent of F . First of all, one has again from (3.1),

$$W_y^F(\Theta) - W_x^F(\Theta) = (\Theta_{yy} - \Theta_{xxxx}) + F'(\Theta)(\Theta_{xx}^2 - \Theta_y^2).$$

We compute,

$$\begin{aligned} & W_y^F(\Theta) - W_x^F(\Theta) \\ &= W_y(\Theta) - W_x(\Theta) + \frac{1}{\Theta}\Theta_y^2 - \frac{1}{\Theta}\Theta_{xx}^2 + F'(\Theta)(\Theta_{xx}^2 - \Theta_y^2) \\ &= W_y(\Theta) - W_x(\Theta) + \left(\frac{1}{\Theta} - F'(\Theta)\right)H(\Theta)(\Theta_y + \Theta_{xx}). \end{aligned}$$

Since $H(\Theta) = 0$, by the previous result $W_y^F(\Theta) - W_x^F(\Theta) = 0$ and $W_y(\Theta) - W_x(\Theta) = 0$. \square

Remark 3.6. The converse of Lemma 3.5 does not hold in general. Indeed, the condition $H(\Theta) = 0$ is sufficient to cancel the third part of the equation (1.10). However, the conditions $W_x^F(\Theta) = W_y^F(\Theta)$ are not sufficient to cancel $H(\Theta)$. See Corollary 3.10 for additional details.

Lemma 3.7. *Let $\Theta > 0$ be a everywhere smooth real-valued phase. Then Θ is of y -Wronskian type if and only if $\Theta = A(t, x) \exp(c(t, x)y)$, for arbitrary $A > 0$, c real-valued.*

Proof. The sufficient condition is clear. We prove the necessary condition. By hypothesis, $\Theta W_y(\Theta) = 0$, i.e. $\Theta\Theta_{yy} - \Theta_y^2 = 0$. Since $\Theta > 0$, this equation can be written as $\Theta^2\partial_y\left(\frac{\Theta_y}{\Theta}\right) = 0$. Hence, $\Theta_y = c(t, x)\Theta$, with $c(t, x)$ a well-defined function in \mathbb{R}^2 . Then $\Theta(t, x, y) = A(t, x)\exp(c(t, x)y)$. Note that $A(t, x) > 0$, since $\Theta(t, x, y) > 0$. \square

Remark 3.8. It is noticed that phases satisfying the y -Wronskian condition are extremely rigid. On the contrary, the x -Wronskian condition seems less demanding.

Corollary 3.9. *Let $\Theta > 0$ be a smooth phase. Then Θ is of x -Wronskian and y -Wronskian type if and only if*

$$\Theta(t, x, y) = A(t, x) \exp((c_1(t) + c_2(t)x)y),$$

with c_1, c_2 are time-dependent, smooth arbitrary functions, and $A > 0$ is an x -Wronskian type function. In the case where $c_2(t) \neq 0$, one has $A(t, x) = c_3(t) \exp(c_4(t)x)$, with $c_3(t) > 0$ and $c_4(t) \in \mathbb{R}$ is smooth and arbitrary.

Proof. As in the proof of Lemma 3.7, we only prove the necessary condition. Thanks to this last result, $\Theta(t, x, y) = A(t, x) \exp(c(t, x)y)$. Assuming now $W_x(\Theta) = 0$, i.e. $\Theta\Theta_{xxxx} - \Theta_{xx}^2 = 0$, and replacing Θ in this equation, one gets the next degree-3 polynomial in the y variable:

$$\begin{aligned} p(y) &= y^3 (4A^2 c_x^2 c_{xx}) \\ &+ y^2 (-4A_x^2 c_x^2 + 4AA_{xx} c_x^2 + 8AA_x c_x c_{xx} + 2A^2 c_{xx}^2 + 4A^2 c_x c_{xxx}) \\ &+ y (-4A_x A_{xx} c_x + 4AA_{xx} c_{xx} + 4AA_{xxx} c_x + 4AA_x c_{xxx} + A^2 c_{xxxx}) \\ &+ AA_{xxx} - A_{xx}^2 = 0. \end{aligned}$$

By the linear independence, each of the coefficients that multiplies y^i with $i \in \{0, 1, 2, 3\}$, must be equal to zero. Making the coefficient that multiplies y^3 equal to 0 one gets $4A^2 c_x^2 c_{xx} = 0$. Since $A > 0$, $(c_x^3)_x = 0$. Consequently, $c(t, x) = c_1(t) + c_2(t)x$. Now the remaining equations in powers of y return the equations

$$\begin{aligned} AA_{xxx} - A_{xx}^2 &= 0, \\ 4c_2(t) (AA_{xx} - A_x A_{xx}) &= 0, \\ 4c_2^2(t) (AA_{xx} - A_x^2) &= 0. \end{aligned}$$

Assume $c_2(t) \neq 0$. The third equation implies the first and second ones. We are left to solve $AA_{xx} = A_x^2$, or $A^2 \left(\frac{A_x}{A}\right)_x = 0$. The solution is

$$A(t, x) = c_3(t) \exp(c_4(t)x).$$

Notice that A is of x -Wronskian type, that is, $AA_{xxx} = A_{xx}^2$. Replacing A in the phase Θ ,

$$\Theta(t, x, y) = c_3(t) \exp(c_4(t)x + (c_1(t) + c_2(t)x)y),$$

as desired.

If now $c_2(t) = 0$, then A is of x -Wronskian type. This ends the proof. \square

Corollary 3.10. *Assume $\Theta > 0$ everywhere. The conditions $W_y(\Theta) = W_x(\Theta) = 0$ do not necessarily imply $H(\Theta) = 0$.*

Proof. From Corollary 3.9, and assuming $c_2(t)$ different from zero, we obtain that necessarily $\Theta(t, x, y) = c_3(t) \exp(c_4(t)x + (c_1(t) + c_2(t)x)y)$. Then

$$H(\Theta) = \Theta_y - \Theta_{xx} = \Theta \left((c_1 + c_2x) - (c_2y + c_4)^2 \right).$$

Then H is identically zero only if $c_2 = 0$ and $c_1 = c_4^2$. Therefore, in general $W_x(\Theta) = W_y(\Theta) = 0$ do not imply $H(\Theta) = 0$, except if $c_2 = 0$ and $c_1 = c_4^2$. \square

Lemma 3.11. *If $W_y(\Theta) = W_x(\Theta) = Ai(\Theta) = 0$, then $\Theta = A(t, x) \exp(cy)$, with $A(t, x)$ being of Airy type and with $c \in \mathbb{R}$.*

Proof. From Corollary 3.9, if $W_y(\Theta) = W_x(\Theta) = 0$ one has

$$\Theta(t, x, y) = A(t, x) \exp((c_1(t) + c_2(t)x)y).$$

If now $Ai(\Theta) = 0$,

$$\begin{aligned} 0 = & -y \exp(y(c_1(t) + xc_2(t))) (3c_2(t) A_{xx}(t, x) + c_1'(t) (-A(t, x)) - xc_2'(t) A(t, x)) \\ & - 3y^2 c_2^2(t) A_x(t, x) \exp(y(c_1(t) + xc_2(t))) \\ & - (A_{xxx}(t, x) - A_t(t, x)) \exp(y(c_1(t) + xc_2(t))) \\ & - y^3 c_2^3(t) A(t, x) (\exp(y(c_1(t) + xc_2(t))))). \end{aligned}$$

If $A = 0$, we are done. Assuming $c_2(t) = 0$, one gets $c_1'(t) = 0$ and then c_1 is constant, and A satisfies Airy. Then $\Theta = A(t, x) \exp(cy)$, with $A(t, x)$ being of Airy type. \square

Lemma 3.12. *Let $\Theta > 0$ be a smooth phase, and F smooth such that $u = 2\partial_x^2 F(\Theta)$ solves KP (1.1). Then $Ai(\Theta) = 0$ and $H(\Theta) = 0$ imply $\mathcal{T}(\Theta) = 0$.*

Proof. Direct from the definition of $\mathcal{T}(\Theta)$ in (1.7). \square

Classical resonant solitons are in the class $Ai(\Theta) = 0$ and $H(\Theta) = 0$. From the previous result, their phases do satisfy $\mathcal{T}(\Theta) = 0$ as well. The following is a sort of reciprocal saving Corollary 3.10.

Lemma 3.13. *Let $u = 2\partial_x^2 F(\Theta)$ solution of (1.1). Let F be a smooth profile satisfying $F(1) = 0$, $F'(1) = 1$, $F''(1) = -1$ and $F'''(1) = 2$. Let $\Theta > 0$ be a smooth phase such that $\Theta_x \neq 0$, $W_y^F(\Theta) = W_x^F(\Theta)$ and $Ai(\Theta) = 0$. Then $F = \log$ and consequently $W_y^F = W_y$, $W_x^F = W_x$.*

Proof. From (1.10) we have

$$\begin{aligned} & (\rho''(s) - 2F'(\Theta)\rho'(s) + 4F''(\Theta)\rho(s))\Theta_x^4 + 6\rho'(s)\Theta_x^2\Theta_{xx} + 3\rho(s)(\Theta_{xx}^2 + \Theta_y^2) \\ & - 4F''(\Theta)\Theta_x Ai(\Theta) - 4F'(\Theta)Ai(\Theta)_x \\ & + 3F'(\Theta)(W_y^F(\Theta) - W_x^F(\Theta)) = 0. \end{aligned}$$

Using the hypotheses, only the first part remains:

$$(\rho''(s) - 2F'(\Theta)\rho'(s) + 4F''(\Theta)\rho(s))\Theta_x^4 + 6\rho'(s)\Theta_x^2\Theta_{xx} + 3\rho(s)(\Theta_{xx}^2 + \Theta_y^2) = 0.$$

This is an ODE for ρ with variable coefficients, exactly with the form (2.2). Lemma 2.5 ensures that $\rho = 0$. Lemma 2.4 implies that $F = \log$ and by definition $W_y^F = W_y$, $W_x^F = W_x$. \square

4. SOLITON STRUCTURES

4.1. Airy and Heat structures. Recall that the Airy type condition defined in Definition 1.1 is described by the identity $Ai(\Theta) = \Theta_t - \Theta_{xxx} = 0$. Unfortunately, this condition does not coincide with linear part of the KP equation. Later (see Appendix B) we will check that in the KdV this is not the case: being of Airy type implies that Θ satisfies the linear part of KdV.

Some simple solutions of $Ai(\Theta) = 0$ are

$$\Theta_1(t, x, y) = A_0 \exp(k_1 x + k_1^2 y + k_1^3 t) + B_0 \exp(k_2 x + k_2^2 y + k_2^3 t) + C_0,$$

$$\Theta_2(t, x, y) = A(y) \exp(kx + k^3 t),$$

where $A_0, B_0, C_0, k_1, k_2, k \in \mathbb{R}$ and $A > 0$ is any well-defined function. Associated to the profile $F = \log$, the phase Θ_1 corresponds to a line-soliton phase (1.12) while Θ_2 gives the trivial solution (Lemma 3.1 (iii)). Finally, the phase Θ_1 serves as an example exhibiting both Airy and Heat type.

Lemma 4.1. *Assume that Θ is of \mathcal{T} -type and $F'(\Theta)$ is different from zero for $(t, x, y) \in \mathbb{R}^3$. If Θ is of Heat type, then there exists $c_0(t, y) \in \mathbb{R}$ such that*

$$Ai(\Theta) = \frac{c_0(t, y)}{F'(\Theta)}. \quad (4.1)$$

Proof. Notice that Θ being of \mathcal{T} -type as in (1.7) is equivalent to have

$$\begin{aligned} 0 = & -4F''(\Theta) \Theta_x Ai(\Theta) - 4F'(\Theta) Ai(\Theta)_x + 3F'(\Theta) \left(H(\Theta)_y + H(\Theta)_{xx} \right) \\ & - 3F'(\Theta)^2 H(\Theta) (\Theta_y + \Theta_{xx}). \end{aligned}$$

Equivalently,

$$(F'(\Theta) Ai(\Theta))_x = \frac{3F'(\Theta)}{4} \left((H(\Theta)_y + H(\Theta)_{xx}) - F'(\Theta) H(\Theta) (\Theta_y + \Theta_{xx}) \right).$$

Then (4.1) follows directly from $H(\Theta) = 0$ for all (t, x, y) and solving the corresponding ODE for $Ai(\Theta)$. \square

Lemma 4.2. *Let $\Theta > 0$ be a smooth phase satisfying $W_x(\Theta) = W_y(\Theta) = 0$. Then the following conditions are satisfied:*

(i) *If $H(\Theta) = 0$, then*

$$\Theta(t, x, y) = a_0(t) \exp(k_1(t)x + k_1^2(t)y) + a_1(t) \exp(-k_1(t)x + k_1^2(t)y),$$

with $a_0, a_1, k_1 \geq 0$.

(ii) *If $Ai(\Theta) = 0$, then $\Theta = A(t, x) \exp(c_1 y)$ where $c_1 \in \mathbb{R}$ and A is of Airy and W_x type.*

(iii) *If both $H(\Theta) = Ai(\Theta) = 0$, then*

$$\Theta = a_1 \exp(k_1 x + k_1^2 y + k_1^3 t) + a_2 \exp(-k_1 x + k_1^2 y - k_1^3 t),$$

for some $a_1, a_2 > 0$ and k_1 constants.

Proof. Proof of (i). Assume $H(\Theta) = 0$. From $W_x(\Theta) = W_y(\Theta) = 0$, Corollary 3.9 yields

$$\Theta(t, x, y) = A(t, x) \exp((c_1(t) + c_2(t)x)y)$$

with A of W_x type. Now if $c_2(t)$ is different from zero, from the proof of Corollary 3.10 we first have

$$\Theta(t, x, y) = c_3(t) \exp(c_4(t)x + (c_1(t) + c_2(t)x)y),$$

and from $H(\Theta) = 0$ we must have $c_2(t) = 0$ and $c_1(t) = c_4^2(t)$. We conclude

$$\Theta(t, x, y) = c_3(t) \exp(c_4(t)x + c_4^2(t)y).$$

Denoting $a_0 := c_3$, $k_1 := c_4$, we conclude this case.

In the case $c_2(t) = 0$ we first have $\Theta(t, x, y) = A(t, x) \exp(c_1(t)y)$. Since $H(\Theta) = 0$ necessarily $A(t, x)c_1(t) - A_{xx}(t, x) = 0$. Depending on the sign of $c_1(t)$, we have

$$A(t, x) = a_0(t)e^{\sqrt{c_1(t)}x} + a_1(t)e^{-\sqrt{c_1(t)}x}, \quad c_1(t) > 0,$$

$$A(t, x) = a_0(t) + a_1(t)x, \quad c_1(t) = 0,$$

or

$$A(t, x) = a_0(t) \cos(\sqrt{-c_1(t)}x) + a_1(t) \sin(\sqrt{-c_1(t)}x), \quad -c_1(t) > 0.$$

Since $\Theta > 0$ we discard the third case, and in the second case we get $c_2(t) = 0$, leading to a solution included in the first case. Note that naturally the condition A is of W_x type is satisfied. Denoting $k_1 := \sqrt{c_1}$ we conclude. This proves (i).

Proof of (ii). As in the previous item, from Corollary 3.9 we have

$$\Theta = A(t, x) \exp((c_1(t) + c_2(t)x)y).$$

This phase will assume different values depending on the value of $c_2(t)$.

Case $c_2(t) = 0$. First, we will examine the case in which $c_2(t) = 0$. In this case from Corollary 3.9 $\Theta = A(t, x) \exp(c_1(t)y)$, with $W_x(A) = 0$. First we compute

$$\Theta_t = A_t \exp(c_1 y) + A c_1' y \exp(c_1 y), \quad \Theta_{xxx} = A_{xxx} \exp(c_1 y).$$

Consequently, imposing the Airy condition,

$$\begin{aligned} Ai(\Theta) &= A_t \exp(c_1 y) + A c_1' \exp(c_1 y) - A_{xxx} \exp(c_1 y) \\ &= \exp(c_1 y) (A_t - A_{xxx}) + y \exp(c_1 y) A c_1' = 0. \end{aligned}$$

The equality leads to a first-degree polynomial in the variable y equals zero. Setting each coefficient equal to 0 yields the following system

$$A_t - A_{xxx} = 0, \quad A c_1' = 0.$$

From the first equation, $A(t, x)$ is Airy type and from the second one, $c_1(t) = c_1$ with $c_1 \in \mathbb{R}$ an arbitrary constant. This means that the phase is

$$\Theta = A(t, x) \exp(c_1 y). \tag{4.2}$$

Case $c_2(t) \neq 0$. Now, from Corollary 3.9 it is known that when $c_2(t)$ is different from zero the phase is

$$\Theta = c_3(t) \exp(c_1(t)y + c_2(t)xy + c_4(t)x).$$

One has

$$\begin{aligned} \Theta_t &= c_3' \exp(c_1(t)y + c_2(t)xy + c_4(t)x) \\ &\quad + c_3 (c_1' y + c_2' xy + c_4' x) \exp(c_1(t)y + c_2(t)xy + c_4(t)x), \\ \Theta_{xxx} &= c_3 (c_2 y + c_4)^3 \exp(c_1(t)y + c_2(t)xy + c_4(t)x). \end{aligned}$$

Imposing the Airy condition,

$$\begin{aligned} 0 &= Ai(\Theta) \\ &= \exp(c_1(t)y + c_2(t)xy + c_4(t)x) \\ &\quad \times ((c_3' - c_4^3) + y(c_3 c_1' - 3c_2 c_4^2) + x(c_3 c_4') + xy(c_3 c_2') + y^2(-3c_2^2 c_4) + y^3(-c_3 c_2^2)). \end{aligned}$$

This equality leads to a third-degree polynomial with variables x and y . Setting each coefficient equal to 0 yields the following system of six equations,

$$\begin{aligned} c_3' - c_4^3 &= 0, & c_3c_1' - 3c_2c_4^2 &= 0, & c_3c_4' &= 0, \\ c_3c_2' &= 0, & c_2^2c_4 &= 0, & c_3c_2^2 &= 0. \end{aligned}$$

From the last equation $c_3 = 0$ implies $\Theta = 0$, then we discard this trivial case. We assume $c_3 \neq 0$ and $c_2(t) = 0$. Then the system is reduced to

$$c_3' - c_4^3 = 0, \quad c_3c_1' = 0, \quad c_3c_4' = 0.$$

From the third and the second equation $c_4(t) = c_4$ and $c_1(t) = c_1$, respectively, with $c_1, c_4 \in \mathbb{R}$ arbitrary constants. Then, from the first equation $c_3(t) = c_4^3t + c_3$, with $c_3 \in \mathbb{R}$ an arbitrary constant. The condition $\Theta > 0$ implies $c_4 = 0$. Thus, the phase is

$$\Theta = c_3 \exp(c_1y + c_4x).$$

Using again the Airy condition, one has $c_4 = 0$ and

$$\Theta = c_5 \exp(c_1y),$$

which is a particular case of (4.2), and a trivial solution to (1.1). This proves (ii).

Proof of (iii). From (i) and (ii), we get

$$\Theta(t, x, y) = a_1(t) \exp(k_1x + k_1^2y) + a_2(t) \exp(-k_1x + k_1^2y).$$

Additionally, to satisfy the Airy condition,

$$(a_1' - a_1k_1^3) \exp(k_1x + k_1^2y) + (a_2' + a_2k_1^3) \exp(-k_1x + k_1^2y) = 0.$$

We obtain $a_1(t) = a_1e^{k_1^3t}$, $a_2(t) = a_2e^{-k_1^3t}$, where a_1 and a_2 are constants. Therefore,

$$A(t, x) = a_1 \exp(k_1x + k_1^2y + k_1^3t) + a_2 \exp(-k_1x + k_1^2y - k_1^3t).$$

This proves (iii). □

Finally, we provide a quick method to construct an Airy-Heat phase Θ . The relevance here is that the Airy-Heat type is a property still satisfied by infinitely many functions, specially in the KP case.

Lemma 4.3. *Let $\Theta_0 = \Theta_0(t, x)$ be any solution of Airy $Ai(\Theta_0) = 0$ such that there are $C_1, C_2 > 0$ under which $|\Theta_0(t)| \leq C_1 \exp(C_2|x|)$. Then $\Theta(t, x, y) := \exp(y\partial_x^2)\Theta_0$ solves $H(\Theta) = 0$ for $y \geq 0$.*

Proof. This result is clear from the formula

$$\Theta(t, x, y) = \exp(y\partial_x^2)\Theta_0(t, x) = \frac{1}{(4\pi y)^{1/2}} \int_{\mathbb{R}} \exp\left(-\frac{(x-s)^2}{4\pi y}\right) \Theta_0(t, s) ds$$

and the growth of Θ_0 . □

Remark 4.4. As already mentioned in the introduction (see Remark 1.12), a similar result for the case of the Airy equation $Ai(\Theta) = 0$ with initial condition exponentially growing in x is far from being obvious.

4.2. Soliton structures. Recall from (1.12) that a simple KP line soliton is obtained by the structure

$$\Theta(t, x, y) = a_1 \exp(\theta_1) + a_2 \exp(\theta_2), \quad (4.3)$$

where $a_1, a_2 > 0$, and $\theta_j := k_j x + k_j^2 y + k_j^3 t$, $k_1, k_2 \in \mathbb{R}$. We shall say that Θ represents a line-soliton if Θ has the previous form. Notice that unless $F = \log$, one does not have that $u = 2\partial_x^2 F(\Theta)$ is the standard KP line-soliton (1.13).

Lemma 4.5. *If $\Theta > 0$ smooth represents a line-soliton, then $A_i(\Theta) = H(\Theta) = 0$. Moreover, the case $k_1 = \pm k_2$ is the unique instance in which $W_x(\Theta) = W_y(\Theta) = 0$.*

Remark 4.6. Notice that the condition $k_1 = -k_2$ corresponds to the case of the KdV soliton (vertical line-soliton) and $k_1 = k_2$ the trivial solution.

Proof of Lemma 4.5. The proof comes from (4.3). Indeed

$$\begin{aligned} A_i(\Theta) &= \Theta_t - \Theta_{xxx} \\ &= k_1^3 a_1 \exp(\theta_1) + k_2^3 a_2 \exp(\theta_2) - (k_1^3 a_1 \exp(\theta_1) + k_2^3 a_2 \exp(\theta_2)) = 0. \end{aligned}$$

Additionally,

$$\begin{aligned} H(\Theta) &= \Theta_y - \Theta_{xx} \\ &= k_1^2 a_1 \exp(\theta_1) + k_2^2 a_2 \exp(\theta_2) - (k_1^2 a_1 \exp(\theta_1) + k_2^2 a_2 \exp(\theta_2)) = 0. \end{aligned}$$

Finally,

$$\begin{aligned} \Theta W_y(\Theta) &= \Theta \Theta_{yy} - \Theta_y^2 \\ &= (a_1 \exp(\theta_1) + a_2 \exp(\theta_2)) (k_1^4 a_1 \exp(\theta_1) + k_2^4 a_2 \exp(\theta_2)) \\ &\quad - (k_1^2 a_1 \exp(\theta_1) + k_2^2 a_2 \exp(\theta_2))^2 \\ &= k_1^4 a_1^2 \exp(2\theta_1) + k_2^4 a_1 a_2 \exp(\theta_1 + \theta_2) \\ &\quad + k_1^4 a_1 a_2 \exp(\theta_1 + \theta_2) + k_2^4 a_2^2 \exp(2\theta_2) \\ &\quad - k_1^4 a_1^2 \exp(2\theta_1) - 2k_1^2 k_2^2 a_1 a_2 \exp(\theta_1 + \theta_2) - k_2^4 a_2^2 \exp(2\theta_2) \\ &= (k_1^4 + k_2^4 - 2k_1^2 k_2^2) a_1 a_2 \exp(\theta_1 + \theta_2). \end{aligned}$$

We conclude that

$$\Theta W_y(\Theta) = (k_1^2 - k_2^2)^2 a_1 a_2 \exp(\theta_1 + \theta_2). \quad (4.4)$$

Notice that $W_y(\Theta) = 0$ if and only if $k_1 = \pm k_2$. Since Θ is a Heat type phase, by Lemma 3.5,

$$W_x(\Theta) = W_y(\Theta) = 0.$$

The proof is complete. \square

The previous result can be extended to smooth phases Θ of the form,

$$\Theta = \sum_{j=1}^M a_j \exp(\theta_j), \quad a_j > 0, \quad \theta_j := k_j x + k_j^2 y + k_j^3 t, \quad k_j \in \mathbb{R}. \quad (4.5)$$

Here, $k_1 < k_2 < \dots < k_M$. In Kodama [22] this phase represents a multi-soliton structure graphically represented by $M - 1$ legs on the region y negative and 1 leg in the positive part of the y -axis. Recall \mathcal{W}_n introduced in (1.19).

Lemma 4.7. *If $\Theta \in \mathcal{W}_M$, then*

$$\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}.$$

In particular, $\Theta \in \mathcal{W}_3$ implies $\Theta W_y(\Theta) \in \mathcal{W}_3$.

Proof. Assume that $\Theta \in \mathcal{W}_M$, that is for $0 \leq k_1(t, x) < k_2(t, x) < \dots < k_N(t, x)$,

$$\Theta = \sum_{j=1}^M a_j \exp(\theta_j), \quad a_j(t, x) > 0, \quad \theta_j(t, x, y) = k_j(t, x) y.$$

Note now that $\left(\sum_{n=1}^M a_n \sum_{n=1}^M b_n = \sum_{n=1}^M c_n, \text{ where } c_n = \sum_{i=0}^n a_i b_{n-i}\right)$,

$$\begin{aligned} \Theta W_y(\Theta) &= \Theta \Theta_{yy} - \Theta_y^2 \\ &= \left(\sum_{i=1}^M a_i \exp(\theta_i)\right) \left(\sum_{i=1}^M k_i^4 a_i \exp(\theta_i)\right) - \left(\sum_{i=1}^M k_i^2 a_i \exp(\theta_i)\right)^2 \\ &= \sum_{n=1}^M \sum_{i=1}^n k_{n-i+1}^4 a_i a_{n-i+1} \exp(\theta_i + \theta_{n-i+1}) \\ &\quad - \sum_{n=1}^M \sum_{i=1}^n k_i^2 k_{n-i+1}^2 a_i a_{n-i+1} \exp(\theta_i + \theta_{n-i}) \\ &= \sum_{n=1}^M \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1}^4 - k_i^2 k_{n-i+1}^2) \exp(\theta_i + \theta_{n-i+1}). \end{aligned}$$

Since $\theta_i + \theta_{n-i+1} = \theta_{n-i+1} + \theta_i$,

$$\begin{aligned} \Theta W_y(\Theta) &= \sum_{n=1}^M \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1}^4 - k_i^2 k_{n-i+1}^2) \exp(\theta_i + \theta_{n-i+1}) \\ &= \frac{1}{2} \sum_{n=1}^M \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1}^4 - k_i^2 k_{n-i+1}^2 + k_i^4 - k_{n-i+1}^2 k_i^2) \exp(\theta_i + \theta_{n-i+1}) \\ &= \frac{1}{2} \sum_{n=1}^M \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1}^4 - 2k_i^2 k_{n-i+1}^2 + k_i^4) \exp(\theta_i + \theta_{n-i+1}) \\ &= \frac{1}{2} \sum_{n=1}^M \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1}^2 - k_i^2)^2 \exp(\theta_i + \theta_{n-i+1}). \end{aligned}$$

Under the hypotheses of the lemma, in general there are $\frac{1}{2}(M^2 - M)$ linearly independent terms in the last expression, proving the required inclusion in $\mathcal{W}_{\frac{1}{2}M(M-1)}$. The M terms correspond to the diagonal, and M^2 represents all the possible elements in an $M \times M$ matrix. \square

Corollary 4.8. *If $\Theta > 0$ as in (4.5) generates an M resonant soliton, then $A_i(\Theta) = H(\Theta) = 0$ but $W_x(\Theta) = W_y(\Theta)$ are different from zero unless $M = 1$ or $M = 2$.*

Proof. The proof of $Ai(\Theta) = H(\Theta) = 0$ is direct. Indeed, computing the corresponding derivatives and replacing its values in $H(\Theta)$,

$$\begin{aligned} H(\Theta) &= \Theta_y - \Theta_{xx} \\ &= \sum_{i=1}^M k_i^2 a_i \exp(k_i x + k_i^2 y + k_i^3 t) - \sum_{i=1}^M k_i^2 a_i \exp(k_i x + k_i^2 y + k_i^3 t) = 0. \end{aligned}$$

Similarly,

$$\begin{aligned} Ai(\Theta) &= \Theta_t - \Theta_{xxx} \\ &= \sum_{i=1}^M k_i^3 a_i \exp(k_i x + k_i^2 y + k_i^3 t) - \sum_{i=1}^M k_i^3 a_i \exp(k_i x + k_i^2 y + k_i^3 t) = 0. \end{aligned}$$

Therefore, if $\Theta(t, x, y) = \sum_{i=1}^M a_i \exp(k_i x + k_i^2 y + k_i^3 t)$ then $H(\Theta) = Ai(\Theta) = 0$.

Now, from the proof of Lemma 4.7, one has $\Theta W_y(\Theta) = 0$ if and only if,

$$k_i^2 = k_j^2,$$

for all $i, j \in \{1, \dots, M\}$. If $M \geq 3$ at least one term of the sum is different to zero, since $k_1 < k_2 < \dots < k_M$ imply W_y different from zero. In the cases $M = 2$ or $M = 1$, from Lemma 4.2 (iii) one has that $W_y = 0$ is equivalent to a phase associated KdV soliton or a trivial solution, respectively. \square

4.3. 2-solitons. For the following result, recall the 2-soliton phase introduced in (1.16).

Lemma 4.9. *Let $\Theta > 0$ be a 2-soliton of scaling parameters $k_1 < k_2 < k_3 < k_4$, and where each soliton correspond to a line-soliton. Then the following are satisfied:*

- (i) *In general $Ai(\Theta)$, $H(\Theta)$, $W_y(\Theta)$ and $W_x(\Theta)$ are different from zero.*
- (ii) *In general, $Ai(\Theta)$, $H(\Theta) \in \mathcal{W}_4$ and $\Theta W_y(\Theta)$, $\Theta W_x(\Theta) \in \mathcal{W}_5$.*
- (iii) *If now $k_1 = -k_2$ and $k_3 = -k_4$ (that is, the case of 2 vertical line-solitons), then $W_y(\Theta) = 0$.*

Proof. After computing (see Appendix A.2),

$$\begin{aligned} Ai(\Theta) &= \Theta_t - \Theta_{xxx} \\ &= (k_3 - k_1)(k_3^3 + k_1^3) \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4^3 + k_1^3) \exp(\theta_1 + \theta_4) \\ &\quad + (k_3 - k_2)(k_3^3 + k_2^3) \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4^3 + k_2^3) \exp(\theta_2 + \theta_4) \quad (4.6) \\ &\quad - (k_3 - k_1)(k_3 + k_1)^3 \exp(\theta_1 + \theta_3) - (k_4 - k_1)(k_4 + k_1)^3 \exp(\theta_1 + \theta_4) \\ &\quad - (k_3 - k_2)(k_3 + k_2)^3 \exp(\theta_2 + \theta_3) - (k_4 - k_2)(k_4 + k_2)^3 \exp(\theta_2 + \theta_4). \end{aligned}$$

notice that $Ai(\Theta) = 0$ if and only if the associated exponential coefficients are zero. Thus, for each $(j, i) \in \{(1, 3), (1, 4), (2, 3), (2, 4)\}$

$$\begin{aligned} &(k_i - k_j)(k_i^3 + k_j^3) - (k_i - k_j)(k_i + k_j)^3 \\ &= k_i^4 + k_i k_j^3 - k_i^3 k_j - k_j^4 - (k_i - k_j)(k_i^3 + 3k_i^2 k_j + 3k_i k_j^2 + k_j^3) \\ &= k_i^4 + k_i k_j^3 - k_i^3 k_j - k_j^4 - 3k_i^3 k_j - 3k_i^2 k_j^2 - k_i k_j^3 + k_i^3 k_j + 3k_i^2 k_j^2 + 3k_i k_j^3 + k_j^4 \\ &= -3k_i^3 k_j + 3k_i k_j^3 = 3(k_i k_j^3 - k_i^3 k_j) = 3k_i k_j (k_j^2 - k_i^2). \end{aligned}$$

Denoting $E_{ij} = (k_j - k_i) \exp(\theta_i + \theta_j)$, then $A_i(\Theta)$ can be rewritten as

$$A_i(\Theta) = -3 \left(k_1 k_3 (k_1 + k_3) E_{13} + k_1 k_4 (k_1 + k_4) E_{14} \right. \\ \left. + k_2 k_3 (k_2 + k_3) E_{23} + k_2 k_4 (k_2 + k_4) E_{24} \right).$$

This proves that $A_i(\Theta) \in \mathcal{W}_4$.² Replacing the values of the derivatives in $H(\Theta)$,

$$H(\Theta) = \Theta_y - \Theta_{xx} \\ = (k_3 - k_1) (k_3^2 + k_1^2) \exp(\theta_1 + \theta_3) + (k_4 - k_1) (k_4^2 + k_1^2) \exp(\theta_1 + \theta_4) \\ + (k_3 - k_2) (k_3^2 + k_2^2) \exp(\theta_2 + \theta_3) + (k_4 - k_2) (k_4^2 + k_2^2) \exp(\theta_2 + \theta_4) \\ - (k_3 - k_1) (k_3 + k_1)^2 \exp(\theta_1 + \theta_3) - (k_4 - k_1) (k_4 + k_1)^2 \exp(\theta_1 + \theta_4) \\ - (k_3 - k_2) (k_3 + k_2)^2 \exp(\theta_2 + \theta_3) - (k_4 - k_2) (k_4 + k_2)^2 \exp(\theta_2 + \theta_4).$$

Repeating the procedure of the Airy condition, $H(\Theta)$ can be rewritten as

$$H(\Theta) = -2 (k_1 k_3 E_{13} + k_1 k_4 E_{14} + k_2 k_3 E_{23} + k_2 k_4 E_{24}).$$

The condition $k_1 < k_2 < k_3 < k_4$ naturally forbids $H(\Theta) = 0$.

Now, replacing the values of the derivatives in $W_y(\Theta)$,

$$\Theta W_y(\Theta) \\ = \Theta \Theta_{yy} - \Theta_y^2 \\ = \left((k_3 - k_1) \exp(\theta_1 + \theta_3) + (k_4 - k_1) \exp(\theta_1 + \theta_4) \right. \\ \left. + (k_3 - k_2) \exp(\theta_2 + \theta_3) + (k_4 - k_2) \exp(\theta_2 + \theta_4) \right) \\ \cdot \left((k_3 - k_1) (k_3^2 + k_1^2)^2 \exp(\theta_1 + \theta_3) \right. \\ \left. + (k_4 - k_1) (k_4^2 + k_1^2)^2 \exp(\theta_1 + \theta_4) + (k_3 - k_2) (k_3^2 + k_2^2)^2 \exp(\theta_2 + \theta_3) \right. \\ \left. + (k_4 - k_2) (k_4^2 + k_2^2)^2 \exp(\theta_2 + \theta_4) \right) \\ - \left((k_3 - k_1) (k_3^2 + k_1^2) \exp(\theta_1 + \theta_3) + (k_4 - k_1) (k_4^2 + k_1^2) \exp(\theta_1 + \theta_4) \right. \\ \left. + (k_3 - k_2) (k_3^2 + k_2^2) \exp(\theta_2 + \theta_3) + (k_4 - k_2) (k_4^2 + k_2^2) \exp(\theta_2 + \theta_4) \right)^2.$$

Developing and rearranging,

$$\Theta W_y(\Theta) \\ = (k_1^2 - k_2^2)^2 \left(\exp(\theta_1 + \theta_2 + 2\theta_4) + \exp(\theta_1 + \theta_2 + 2\theta_3) \right) \\ + (k_3^2 - k_4^2)^2 \left(\exp(2\theta_1 + \theta_3 + \theta_4) + \exp(2\theta_2 + \theta_3 + \theta_4) \right) \\ + k_{1234} \exp(\theta_1 + \theta_2 + \theta_3 + \theta_4), \tag{4.7}$$

²Assume $k_1 < k_2 < k_3 < k_4$. Let us study the condition $A_i(\Theta) = 0$. If $k_1 = 0$, then $k_2 = -k_3 = -k_4$, which is impossible. A similar argument holds if now $k_2 = 0$, or $k_3 = 0$, or $k_4 = 0$. Therefore, each k_i must be nonzero. However, in this case $k_1 = -k_3 = -k_4$, also impossible. Thus, $A_i(\Theta) \neq 0$ for a nondegenerate 2-soliton.

where

$$k_{1234} := (k_3 - k_1)(k_4 - k_2) \left((k_1^2 + k_3^2) - (k_2^2 + k_4^2) \right)^2 \\ + (k_3 - k_2)(k_4 - k_1) \left((k_1^2 + k_4^2) - (k_2^2 + k_3^2) \right)^2.$$

A simple observation reveals that $\Theta W_y(\Theta) = 0$ if $k_{1234} = 0$ and $k_1 = \pm k_2$, $k_3 = \pm k_4$. Since $k_1 < k_2 < k_3 < k_4$, one necessarily has $k_1 = -k_2$ and $k_3 = -k_4$. In this case, we also have $k_{1234} = 0$, making this assumption unnecessary. Except by this particular case, one naturally concludes from (4.7) that $\Theta W_y(\Theta) \in \mathcal{W}_5$. Therefore, W_y can be zero in the case of a 2-soliton when $k_1 = -k_2$ and $k_3 = -k_4$. This can be true when all constants have different values.

As mentioned earlier, $H(\Theta)$ is not equal to zero and therefore it is necessary to verify both conditions, $\Theta W_y(\Theta) = 0$ and $\Theta W_x(\Theta) = 0$, separately. Then,

$$\begin{aligned} \Theta W_x(\Theta) &= \Theta \Theta_{xxxx} - \Theta_{xx}^2 \\ &= \left((k_3 - k_1) \exp(\theta_1 + \theta_3) + (k_4 - k_1) \exp(\theta_1 + \theta_4) + (k_3 - k_2) \exp(\theta_2 + \theta_3) \right. \\ &\quad \left. + (k_4 - k_2) \exp(\theta_2 + \theta_4) \right) \cdot \left((k_3 - k_1)(k_3 + k_1)^4 \exp(\theta_1 + \theta_3) \right. \\ &\quad \left. + (k_4 - k_1)(k_4 + k_1)^4 \exp(\theta_1 + \theta_4) + (k_3 - k_2)(k_3 + k_2)^4 \exp(\theta_2 + \theta_3) \right. \\ &\quad \left. + (k_4 - k_2)(k_4 + k_2)^4 \exp(\theta_2 + \theta_4) \right) - \left((k_3 - k_1)(k_3 + k_1)^2 \exp(\theta_1 + \theta_3) \right. \\ &\quad \left. + (k_4 - k_1)(k_4 + k_1)^2 \exp(\theta_1 + \theta_4) + (k_3 - k_2)(k_3 + k_2)^2 \exp(\theta_2 + \theta_3) \right. \\ &\quad \left. + (k_4 - k_2)(k_4 + k_2)^2 \exp(\theta_2 + \theta_4) \right)^2. \end{aligned}$$

Developing and grouping terms, we arrive at

$$\begin{aligned} \Theta W_x(\Theta) &= \left((k_1 + k_4)^2 - (k_1 + k_3)^2 \right)^2 E_{13} E_{14} + \left((k_2 + k_3)^2 - (k_1 + k_3)^2 \right)^2 E_{13} E_{23} \\ &\quad + \left(\left((k_2 + k_4)^2 - (k_1 + k_3)^2 \right)^2 + \left((k_2 + k_3)^2 - (k_1 + k_4)^2 \right)^2 \right) E_{14} E_{23} \\ &\quad + \left((k_1 + k_4)^2 - (k_2 + k_4)^2 \right)^2 E_{14} E_{24} + \left((k_2 + k_3)^2 - (k_2 + k_4)^2 \right)^2 E_{23} E_{24}. \end{aligned} \tag{4.8}$$

Clearly from (4.8) one concludes that in general $\Theta W_x(\Theta) \in \mathcal{W}_5$. In order to get zero value, one should have

$$|k_1 + k_4| = |k_1 + k_3| = |k_2 + k_4| = |k_2 + k_3|.$$

Under the hypothesis $k_1 < k_2 < k_3 < k_4$, this is never satisfied. Indeed,

$$k_1 + k_3 < k_2 + k_3 < k_2 + k_4, \quad k_1 + k_3 < k_1 + k_4;$$

contradicting the equality of absolute values. Then $\Theta W_x(\Theta)$ is always different from zero to a phase of a 2-soliton. \square

Corollary 4.10. *In general, given the profile $F = \log$, any 2-soliton as in (1.16) only satisfies $\mathcal{T} = 0$.*

Proof. Direct from (1.9), Lemma 2.4 and Lemma 4.9 (i). \square

5. PROOF OF MAIN RESULTS

5.1. Proof of Theorem 1.4. Assume $\Theta > 0$ is a smooth KdV line-soliton phase with a profile $F = \log$. Its phase is given by

$$\begin{aligned}\Theta(t, x, y) &= \exp(kx + k^2y + k^3t) + \exp(-kx + k^2y - k^3t) \\ &= \exp(\theta_1) + \exp(\theta_2),\end{aligned}\tag{5.1}$$

with $\theta_1 = kx + k^2y + k^3t$ and $\theta_2 = -kx + k^2y - k^3t$. It will be shown that the KdV phase satisfies the following conditions

$$H(\Theta) = Ai(\Theta) = W_x(\Theta) = W_y(\Theta) = 0.$$

Computing derivatives, we obtain

$$\begin{aligned}\Theta_x &= k \exp(\theta_1) - k \exp(\theta_2); & \Theta_{xx} &= k^2 \exp(\theta_1) + k^2 \exp(\theta_2); \\ \Theta_{xxx} &= k^3 \exp(\theta_1) - k^3 \exp(\theta_2); & \Theta_{xxxx} &= k^4 \exp(\theta_1) + k^4 \exp(\theta_2); \\ \Theta_y &= k^2 \exp(\theta_1) + k^2 \exp(\theta_2); & \Theta_{yy} &= k^4 \exp(\theta_1) + k^4 \exp(\theta_2); \\ \Theta_t &= k^3 \exp(\theta_1) - k^3 \exp(\theta_2).\end{aligned}$$

Replacing the values of Θ_y and Θ_{xx} in $H(\Theta)$ by corresponding expressions above, one has

$$\begin{aligned}H(\Theta) &= \Theta_y - \Theta_{xx} \\ &= (k^2 \exp(\theta_1) + k^2 \exp(\theta_2)) - (k^2 \exp(\theta_1) + k^2 \exp(\theta_2)) \\ &= (k^2 \exp(\theta_1) - k^2 \exp(\theta_1)) + (k^2 \exp(\theta_2) - k^2 \exp(\theta_2)) = 0.\end{aligned}$$

Hence $H(\Theta) = 0$, then Θ is of Heat type. Since $H(\Theta) = 0$ from Lemma 3.5, $W_x(\Theta) = W_y(\Theta)$. Now, replacing the values of the derivatives of Θ in $W_x(\Theta)$,

$$\begin{aligned}W_x(\Theta) &= \Theta_{xxxx} - \frac{\Theta_{xx}^2}{\Theta} \\ &= (k^4 \exp(\theta_1) + k^4 \exp(\theta_2)) - \frac{(k^2 \exp(\theta_1) + k^2 \exp(\theta_2))^2}{(\exp(\theta_1) + \exp(\theta_2))} \\ &= k^4 \frac{(\exp(\theta_1) + \exp(\theta_2))^2}{(\exp(\theta_1) + \exp(\theta_2))} - k^4 \frac{(\exp(\theta_1) + \exp(\theta_2))^2}{(\exp(\theta_1) + \exp(\theta_2))} = 0.\end{aligned}$$

Since $H(\Theta) = 0$, this also means that $W_y(\Theta) = 0$.

Finally, replacing the derivatives of Θ in $Ai(\Theta)$

$$\Theta_t - \Theta_{xxx} = (k^3 \exp(\theta_1) - k^3 \exp(\theta_2)) - (k^3 \exp(\theta_1) - k^3 \exp(\theta_2)) = 0.$$

Then, if $u = 2\partial_x^2 \log \Theta$ is a KdV line-soliton solution of KP (i.e. $\Theta = \exp(kx + k^2y + k^3t) + \exp(-kx + k^2y - k^3t)$) then $H(\Theta) = Ai(\Theta) = W_x(\Theta) = W_y(\Theta) = 0$.

Conversely, it will now be demonstrated that if a phase Θ satisfies $H(\Theta) = W_x^F(\Theta) = W_y^F(\Theta) = Ai(\Theta) = 0$, then the corresponding solution $u = 2\partial_x^2 F(\Theta)$ is a KdV vertical line-soliton, that is, $F = \log$ and Θ as in (5.1).

Since $H(\Theta) = Ai(\Theta) = W_x^F(\Theta) = W_y^F(\Theta) = 0$, it is sufficient to look at equation (1.8), to conclude that

$$(\rho(s)'' - 2F'(\Theta)\rho(s)' + 4F''(\Theta)\rho(s))\Theta_x^4 + 6\rho(s)'\Theta_x^2\Theta_{xx} + 3\rho(s)(\Theta_{xx}^2 + \Theta_y^2) = 0.$$

By the hypothesis on F and the values of its derivatives at $s = 1$, if there exists a solution to the KP equation of the form $u = 2\partial_x^2 F(\Theta)$, (1.8) is satisfied, and by Lemma 2.5 (iii), $F = \log$.

Now, Lemma 3.5 implies that $W_y(\Theta) - W_x(\Theta) = 0$. Since $F = \log$, $W_y(\Theta) = W_x(\Theta) = 0$. Since $W_y(\Theta) = W_x(\Theta) = Ai(\Theta) = 0$, Lemma 3.11 ensures that $\Theta = A(t, x) \exp(cy)$, with $A(t, x)$ being of Airy type and with $c \in \mathbb{R}$.

Finally, the condition $H(\Theta) = 0$ implies

$$\Theta_y - \Theta_{xx} = (cA - A_{xx}) \exp(cy) = 0.$$

We first treat the case $c = 0$. In this case

$$A(t, x) = c_{A,1}(t) + c_{A,2}(t)x.$$

Since A must satisfy the Airy equation for all $(t, x, y) \in \mathbb{R}^3$, one gets $c'_{A,1}(t) + c'_{A,2}(t)x = 0$, implying that $c_{A,1}$ and $c_{A,2}$ are constants. Θ is given in this case by

$$\Theta = c_{A,1} + c_{A,2}x,$$

corresponding to a singular soliton solution, which is discarded by smoothness assumptions.

Now we assume c different from zero. Here,

$$A(t, x) = c_{A,1}(t) \exp(kx) + c_{A,2}(t) \exp(-kx), \quad k = \sqrt{c} \in \mathbb{C} - \{0\}. \quad (5.2)$$

Solving again the Airy equation for A , it follows

$$\begin{aligned} A_t - A_{xxx} &= c'_{A,1}(t) \exp(kx) + c'_{A,2}(t) \exp(-kx) \\ &\quad - (k^3 c_{A,1} \exp(kx) - k^3 c_{A,2} \exp(-kx)) = 0. \end{aligned}$$

By the linear independence,

$$\begin{aligned} c'_{A,1}(t) - k^3 c_{A,1}(t) &= 0; \\ c'_{A,2}(t) + k^3 c_{A,2}(t) &= 0. \end{aligned}$$

This are independents ODE's for $c_{A,1}(t)$ and $c_{A,2}(t)$. Solving them, it is obtained

$$c_{A,1}(t) = c_{A,1,0} \exp(k^3 t), \quad c_{A,2}(t) = c_{A,2,0} \exp(-k^3 t), \quad (5.3)$$

with $c_{A,1,0}, c_{A,2,0} \in \mathbb{R}$ arbitrary constants. We obtain from (5.2) and (5.3)

$$\Theta(t, x, y) = c_{A,1,0} \exp(kx + k^2 y + k^3 t) + c_{A,2,0} \exp(-kx + k^2 y - k^3 t).$$

The condition $\Theta \in \mathbb{R}$ implies that k is real-valued. Also, $\Theta > 0$ implies $c_{A,1,0}, c_{A,2,0} > 0$. This finally shows that Θ corresponds to the phase of a KdV vertical line-soliton. The proof is complete.

5.2. Proof of Theorem 1.8. The proof of this result is based in two lemmas. Since Theorem 1.4 considers the case of KdV line solitons, we focus on the most demanding case of oblique solitons ($A > 0$).

Lemma 5.1. *Let u be a smooth solution to (1.1) of the form (1.2), with a smooth profile $F(\Theta)$ such that $F(1) = 0$, $F'(1) = 1$, $F''(1) = -1$, and $F'''(1) = 2$. Then if u is a line-soliton of the form (1.13)-(1.12) and $F = \log$, one has that $H(\Theta) = Ai(\Theta) = 0$, and*

$$\begin{aligned} \Theta W_x(\Theta) &= \Theta W_y(\Theta) \\ &= a_1 a_2 (k_1^2 - k_2^2)^2 \exp((k_1 + k_2)x + (k_1^2 + k_2^2)y + (k_1^3 + k_2^3)t), \end{aligned} \quad (5.4)$$

for some particular $a_1, a_2 > 0$, $k_1, k_2 \in \mathbb{R}$.

Proof. By Lemma 4.5, we know that $H(\Theta) = Ai(\Theta) = 0$. Thanks to (4.4) and Lemma 3.5, we conclude (5.4). This proves Lemma 5.1. \square

Lemma 5.2. *Let u be a smooth solution to (1.1) of the form (1.2), with $\Theta > 0$ smooth and real-valued and F a smooth profile such that $F(1) = 0$, $F'(1) = 1$, $F''(1) = -1$, and $F'''(1) = 2$. If $H(\Theta) = Ai(\Theta) = 0$, and*

$$\Theta W_y(\Theta) = A(t, x) \exp(k(t, x)y), \quad (5.5)$$

for some particular $A > 0$, $k > 0$, then u is a line-soliton of the form (1.13)-(1.12) and $F = \log$.

Remark 5.3. Lemmas 5.1 and 5.2 conclude the proof of Theorem 1.8.

Proof of Lemma 5.2. Step 1. Since $H(\Theta) = Ai(\Theta) = 0$, Lemmas 3.5 and 3.13 and the hypotheses on F ensure that $F = \log$. Let us assume (5.5). Then

$$\Theta^2 \partial_y^2 (\ln(\Theta)) = A(t, x) \exp(k(t, x)y).$$

This equation can be studied like a nonlinear second order ODE on Θ and the variable y . Considering the change of variable $f = \ln(\Theta)$ one gets

$$\exp(2f) f'' = A \exp(ky), \quad \text{then} \quad \tilde{f}'' = 2A \exp(-\tilde{f}),$$

with $\tilde{f} = -ky + 2f$, and consequently $\tilde{f}'' = 2f''$. This is a classical Toda equation. The general solution \tilde{f} is given by

$$\tilde{f} = 2 \log \left(\frac{\sqrt{A}}{\sqrt{c_1}} \left(\exp\left(\frac{1}{2}\eta\right) + \exp\left(-\frac{1}{2}\eta\right) \right) \right),$$

with $\eta := \sqrt{c_1}(y + c_2)$, $c_1 > 0$, $c_2 \in \mathbb{R}$. Therefore,

$$\begin{aligned} \Theta &= \exp(f) = \exp\left(\frac{ky}{2}\right) \exp\left(\frac{\tilde{f}}{2}\right) \\ &= \frac{\sqrt{A}}{\sqrt{c_1}} \left(\exp\left(\left(\frac{k + \sqrt{c_1}}{2}\right)(y + c_2)\right) + \exp\left(\left(\frac{k - \sqrt{c_1}}{2}\right)(y + c_2)\right) \right). \end{aligned}$$

Defining

$$A_1(t, x) = \frac{\sqrt{A}}{\sqrt{c_1}} \exp\left(c_2 \left(\frac{k + \sqrt{c_1}}{2}\right)\right), \quad A_2(t, x) = \frac{\sqrt{A}}{\sqrt{c_1}} \exp\left(c_2 \left(\frac{k - \sqrt{c_1}}{2}\right)\right),$$

and

$$B_1(t, x) = \left(\frac{k + \sqrt{c_1}}{2} \right) > 0, \quad B_2(t, x) = \left(\frac{k - \sqrt{c_1}}{2} \right),$$

Θ has the form $\Theta = A_1(t, x) \exp(B_1(t, x)y) + A_2(t, x) \exp(B_2(t, x)y)$.

Step 2. Since $c_1 > 0$, one gets B_1 different from B_2 . Replacing Θ in the condition $H(\Theta) = 0$, and using Appendix A.3 in the case of two linearly independent exponentials,

$$\begin{aligned} H(\Theta) &= (A_1 B_1 \exp(B_1 y) + A_2 B_2 \exp(B_2 y)) \\ &\quad - \left(A_{1,xx} \exp(B_1 y) + A_{2,xx} \exp(B_2 y) \right. \\ &\quad \quad \left. + y \exp(B_1 y) (2A_{1,x} B_{1,x} + A_1 B_{1,xx}) + y \exp(B_2 y) (2A_{2,x} B_{2,x} + A_2 B_{2,xx}) \right. \\ &\quad \quad \left. + y^2 \exp(B_1 y) A_1 B_{1,x}^2 + y^2 \exp(B_2 y) A_2 B_{2,x}^2 \right) \\ &= (A_1 B_1 - A_{1,xx}) \exp(B_1 y) + (A_2 B_2 - A_{2,xx}) \exp(B_2 y) \\ &\quad - (2A_{1,x} B_{1,x} + A_1 B_{1,xx}) y \exp(B_1 y) - (2A_{2,x} B_{2,x} + A_2 B_{2,xx}) y \exp(B_2 y) \\ &\quad - A_1 B_{1,x}^2 y^2 \exp(B_1 y) - A_2 B_{2,x}^2 y^2 \exp(B_2 y). \end{aligned} \tag{5.6}$$

In order for $H(\Theta)$ to be null, it is necessary that each of the coefficients multiplying a term $y^i \exp(B_j y)$ with $i \in \{0, 1, 2\}$ and $j \in \{1, 2\}$ be equal to zero. Taking the expression above into consideration, the following system of equations is obtained

$$\begin{aligned} A_1 B_1 &= A_{1,xx}, & A_2 B_2 &= A_{2,xx}, \\ 2A_{1,x} B_{1,x} + A_1 B_{1,xx} &= 0, & 2A_{2,x} B_{2,x} + A_2 B_{2,xx} &= 0, \\ A_1 B_{1,x}^2 &= 0, & A_2 B_{2,x}^2 &= 0. \end{aligned}$$

Since $A_1, A_2 > 0$, from the two bottom equations, $B_i(t, x) = B_i(t)$ for $i \in \{1, 2\}$. This reduces the system to the top two equations, from which it can be concluded that $A_i(t, x) = c_i(t) \exp(\sqrt{B_i(t)}x)$. Since $\Theta > 0$ is real-valued, it is required $B_i > 0$. Therefore, the phase takes the form

$$\Theta = c_1(t) \exp(\sqrt{B_1(t)}x + B_1(t)y) + c_2(t) \exp(\sqrt{B_2(t)}x + B_2(t)y).$$

Then, using again Appendix A.3 and inserting those terms into $A_i(\Theta) = 0$, we get

$$\begin{aligned} A_i(\Theta) &= \left(c_{1,t} \exp(\sqrt{B_1}x + B_1 y) + \frac{c_1 B_{1,t}}{2\sqrt{B_1}} x \exp(\sqrt{B_1}x + B_1 y) \right. \\ &\quad \left. + c_1 B_{1,t} y \exp(\sqrt{B_1}x + B_1 y) + c_{2,t} \exp(\sqrt{B_2}x + B_2 y) \right. \\ &\quad \left. + \frac{c_2 B_{2,t}}{2\sqrt{B_2}} x \exp(\sqrt{B_2}x + B_2 y) + c_2 B_{2,t} y \exp(\sqrt{B_2}x + B_2 y) \right) \\ &\quad - \left(c_1 \sqrt{B_1^3} \exp(\sqrt{B_1}x + B_1 y) + c_2 \sqrt{B_2^3} \exp(\sqrt{B_2}x + B_2 y) \right). \end{aligned}$$

Simplifying,

$$\begin{aligned} Ai(\Theta) &= \left(c_{1,t} - c_1 \sqrt{B_1^3} \right) \exp\left(\sqrt{B_1}x + B_1y\right) + \left(c_{2,t} - c_2 \sqrt{B_2^3} \right) \exp\left(\sqrt{B_2}x + B_2y\right) \\ &\quad + \frac{c_1 B_{1,t}}{2\sqrt{B_1}} x \exp\left(\sqrt{B_1}x + B_1y\right) + \frac{c_2 B_{2,t}}{2\sqrt{B_2}} x \exp\left(\sqrt{B_2}x + B_2y\right) \\ &\quad + c_1 B_{1,t} y \exp\left(\sqrt{B_1}x + B_1y\right) + c_2 B_{2,t} y \exp\left(\sqrt{B_2}x + B_2y\right). \end{aligned}$$

To ensuring that the expression is equal to zero, it is necessary that each coefficient multiplying an exponential term be null for all values of $(x, y) \in \mathbb{R}^2$. Taking the above into consideration, the following system of equations is obtained

$$\begin{aligned} c_{1,t} - c_1 \sqrt{B_1^3} &= 0, & c_{2,t} - c_2 \sqrt{B_2^3} &= 0, \\ c_1 B_{1,t} &= 0, & c_2 B_{2,t} &= 0. \end{aligned}$$

Note that the last two equations are derived from the coefficients multiplying an exponential term, multiplied either by x or by y . From the last two equations, it is concluded that $B_i(t) = B_i$ for $i \in \{1, 2\}$ and with $B_i \in \mathbb{R}$. Taking this into account in the first two equations is obtained $c_i = a_i \exp\left(\sqrt{B_i^3}t\right)$ for $i \in \{1, 2\}$ and with $a_i > 0$. In conclusion, the phase is

$$\Theta = a_1 \exp\left(\sqrt{B_1}x + B_1y + \sqrt{B_1^3}t\right) + a_2 \exp\left(\sqrt{B_2}x + B_2y + \sqrt{B_2^3}t\right).$$

where $a_1, a_2 > 0$ are arbitrary constants. Denoting $k_i := \sqrt{B_i}$, we obtain the desired conclusion. \square

5.3. Proof of Theorem 1.11. From Corollary 4.8 we know that u solution of (1.1) of the form (1.2) with smooth real-valued phase $\Theta > 0$ corresponding to an M resonant multi-soliton (1.20) and $F = \log$ imply $H(\Theta) = Ai(\Theta) = 0$. Also, Lemma 4.7 in this particular case $(k_j^2 \geq 0)$ states that $\Theta W_y(\Theta) = \Theta W_x(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$.

Now we prove the opposite. Assume that $\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ has the form

$$\Theta W_y(\Theta) = \sum_{j=1}^{\frac{1}{2}M(M-1)} b_j \exp(m_j y). \quad (5.7)$$

Recall that each $b_j > 0$. After arranging terms as members of an upper triangular matrix, we get

$$\Theta W_y(\Theta) = \sum_{n=1}^M \sum_{i=1}^n b_{n,i} \exp(m_{n,i} y).$$

For the moment, assume

$$\Theta = \sum_{j=1}^M a_j(t, x) \exp(\theta_j), \quad (5.8)$$

with $\theta_j(t, x, y) = k_j(t, x)y$, be a phase in \mathcal{W}_M . From Corollary 4.7,

$$\Theta W_y(\Theta) = \sum_{n=1}^M \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1} - k_i)^2 \exp(\theta_i + \theta_{n-i+1}).$$

The system

$$k_i + k_{n-i+1} = m_{n,i}, \quad i \leq n - i + 1,$$

reads

$$\begin{aligned} k_1 + k_1 &= m_{1,1} \\ k_1 + k_2 &= m_{2,1} \\ k_1 + k_3 &= m_{3,1} \\ k_2 + k_2 &= m_{3,2} \\ &\dots \\ k_1 + k_n &= m_{n,1} \\ k_2 + k_{n-1} &= m_{n,2} \\ k_3 + k_{n-2} &= m_{n,3} \\ &\dots \end{aligned}$$

and has a unique solution on $k = (k_1, \dots, k_N)$, thanks to a nonsingular determinant matrix. The second system is given by

$$a_i a_{n-i+1} (k_{n-i+1} - k_i)^2 = b_{n,i},$$

which implies

$$\log a_i + \log a_{n-i+1} = \log b_{n,i} - 2 \log |k_{n-i+1} - k_i|.$$

This system also has a unique solution for $a = (a_1, a_2, \dots, a_N)$ exactly following the previous argument. Consequently, $\Theta W_y(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ has always a solution in \mathcal{W}_M . The fact that $\Theta W_x(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ is direct.

Now we prove uniqueness. The key now is to use that $\Theta W_x(\Theta) \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ and has a unique value. Let $F \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ and let Θ_1, Θ_2 be such that $\Theta_j W_x(\Theta_j) = F$. Therefore,

$$\Theta_1 \Theta_{1,xxxx} - \Theta_{1,xx}^2 = \Theta_2 \Theta_{2,xxxx} - \Theta_{2,xx}^2.$$

Let $\Pi_0(x; t) := (\Theta_1 - \Theta_2)(t, x, y = 0)$. Then Π_0 satisfies the fourth order linear ODE on x :

$$\Theta_2 \Pi_0'''' - (\Theta_{1,xx} + \Theta_{2,xx}) \Pi_0'' + \Theta_{1,xxxx} \Pi_0 = 0,$$

and thanks to the hypothesis (1.21), $\Pi_0(x = 0; t) = \Pi_0'(x = 0; t) = \Pi_0''(x = 0; t) = \Pi_0'''(x = 0; t) = 0$, leading to $\Pi_0(x; t) \equiv 0$ and consequently $\Theta_1(t, x, 0) = \Theta_2(t, x, 0)$. Additionally, $\Pi_1(x; t) := (\Theta_{1,y} - \Theta_{2,y})(t, x, y = 0)$ satisfies

$$\Theta_2 \Pi_1'''' - 2\Theta_{1,xx} \Pi_1'' + \Theta_{1,xxxx} \Pi_1 = 0.$$

(Notice that we have used that $\Pi_0(x; t) = \Pi_{0,xx}(x; t) = \Pi_{0,xxxx}(x; t) = 0$.) Again, thanks to (1.21) we conclude that $\Pi_1(x; t) \equiv 0$, leading to $\Theta_{1,y}(t, x, 0) = \Theta_{2,y}(t, x, 0)$.

Now we extend the previous uniqueness. The argument is similar to the previous case. Let $G \in \mathcal{W}_{\frac{1}{2}M(M-1)}$ unique and let Θ_1, Θ_2 be such that $\Theta_j W_y(\Theta_j) = G$. Therefore,

$$\Theta_1 \Theta_{1,yy} - \Theta_{1,y}^2 = \Theta_2 \Theta_{2,yy} - \Theta_{2,y}^2.$$

Let $\Pi_2(y; t, x) := (\Theta_1 - \Theta_2)(t, x, y)$. Then Π_2 satisfies the second order linear ODE on y :

$$\Theta_2 \Pi_2'' - (\Theta_{1,y} + \Theta_{2,y}) \Pi_2' + \Theta_{1,yy} \Pi_2 = 0,$$

and thanks to the previous step, one has $\Pi_2(0; t, x) = \Pi_{2,y}(0; t, x) = 0$. Therefore, $\Pi(y; t, x) = 0$ and the required uniqueness holds. This shows that (5.8) is the unique solution of (5.7).

Now we improve the coefficients using that Θ satisfies zero Heat and Airy, showing that Θ is a line-multi-soliton. Since Heat and Airy are linear equations, the proof is similar to the proof of Lemma 5.2, Step 2. More precisely,

$$H(\Theta) = \sum_{j=1}^M \left\{ (a_j k_j - a_{j,xx}) - (2a_{j,x} k_{j,x} + a_j k_{j,xx}) y - a_j k_{j,x}^2 y^2 \right\} \exp(k_j y) = 0. \quad (5.9)$$

By the linear independence, real-valued character and positivity of Θ , we conclude $k_j = k_j(t) \geq 0$ and

$$a_j(t, x) = a_{0,j}(t) \exp\left(\sqrt{k_j(t)}x\right) + a_{1,j}(t) \exp\left(-\sqrt{k_j(t)}x\right), \quad a_{0,j}, a_{1,j} \geq 0.$$

Computing Airy, one gets

$$\begin{aligned} Ai(\Theta) &= \sum_{j=1}^M \left\{ \left(a_{0,j,t} - a_{0,j} \sqrt{k_j^3} \right) + \frac{a_{0,j} k_{j,t}}{2\sqrt{k_j}} x + a_{0,j} k_{j,t} y \right\} \exp\left(\sqrt{k_j}x + k_j y\right) \\ &\quad + \sum_{j=1}^M \left\{ \left(a_{1,j,t} - a_{1,j} \sqrt{k_j^3} \right) - \frac{a_{1,j} k_{j,t}}{2\sqrt{k_j}} x + a_{1,j} k_{j,t} y \right\} \exp\left(-\sqrt{k_j}x + k_j y\right) \\ &= 0. \end{aligned} \quad (5.10)$$

From (5.9) and (5.10), and proceeding exactly as in the previous proof of Lemma 5.2, Step 2, one gets

$$a_{0,j} = a_{0,0,j} \exp\left(\sqrt{k_j^3}t\right), \quad a_{1,j} = a_{1,0,j} \exp\left(\sqrt{k_j^3}t\right),$$

with k_j constants. Renaming $k_j \mapsto k_j^2$, from (5.8) we obtain

$$\Theta = \sum_{j=1}^M \left(a_{0,0,j} \exp(-k_j x) + a_{1,0,j} \exp(k_j x) \right) \exp(k_j^2 y + k_j^3 t).$$

This ends the proof.

5.4. Proof of Theorem 1.13. We are now ready to prove Theorem 1.13, the case of KP 2-solitons. Let u be a solution of (1.1) of the form (1.2) with a smooth real-valued phase $\Theta > 0$ satisfying $\Theta(y=0)$ and $\Theta_y(y=0)$ uniquely prescribed and $F = \log$.

Assume that Θ corresponds to a 2-soliton (1.16) with $k_1 < k_2 < k_3 < k_4$. Let us prove that $H(\Theta), Ai(\Theta) \in \mathcal{W}_4$ and $\Theta W_y(\Theta), \Theta W_x(\Theta) \in \mathcal{W}_5$. Thanks to Lemma 4.9 (i) and (ii), this part is already proved. Finally, $Ai(\Theta) = \frac{3}{2} \partial_x H(\Theta)$ is direct.

Assume now that $H(\Theta), Ai(\Theta) \in \mathcal{W}_4$ and $\Theta W_y(\Theta), \Theta W_x(\Theta) \in \mathcal{W}_5$. We have $\Theta W_y(\Theta) \in \mathcal{W}_6$, and the proof of Theorem 1.11 and hypotheses (1.21) on Θ ensure that $\Theta \in \mathcal{W}_4$. As

in (5.8), one has

$$\Theta = \sum_{j=1}^4 a_j(t, x) \exp(\theta_j), \quad a_j > 0, \quad (5.11)$$

with $\theta_j(t, x, y) = k_j(t, x)y$. The image of (5.11) under $\Theta W_y(\Theta)$ is given by

$$\begin{aligned} \Theta W_y(\Theta) &= \sum_{n=1}^4 \sum_{i=1}^n a_i a_{n-i+1} (k_{n-i+1} - k_i)^2 \exp(\theta_i + \theta_{n-i+1}) \\ &= a_1 a_2 (k_1 - k_2)^2 E_{12} + a_1 a_3 (k_1 - k_3)^2 E_{13} + a_1 a_4 (k_1 - k_4)^2 E_{14} \\ &\quad + a_2 a_3 (k_2 - k_3)^2 E_{23} + a_2 a_4 (k_2 - k_4)^2 E_{24} + a_3 a_4 (k_3 - k_4)^2 E_{34}, \end{aligned}$$

where $E_{ij} := \exp(\theta_i + \theta_j)$. Since $\Theta W_y(\Theta) \in \mathcal{W}_5$, at least one exponential is linearly dependent with the rest of exponentials. This implies that $k_i + k_j = k_{i'} + k_{j'}$ for some i different from i' , j different from j' and therefore one term above is redundant. With no loss of generality, we assume $k_1 < k_2 < k_3 < k_4$ and $k_1 + k_4 = k_2 + k_3$. In this case we obtain $k_4 = k_2 + k_3 - k_1 > 0$, and from (5.11),

$$\Theta(t, x, y) = a_1 \exp(k_1 y) + a_2 \exp(k_2 y) + a_3 \exp(k_3 y) + a_4 \exp((k_2 + k_3 - k_1)y), \quad (5.12)$$

and

$$\begin{aligned} \Theta W_y(\Theta) &= a_1 a_2 (k_1 - k_2)^2 E_{12} + a_1 a_3 (k_1 - k_3)^2 E_{13} \\ &\quad + \left(a_1 a_4 (k_1 - k_4)^2 + a_2 a_3 (k_2 - k_3)^2 \right) E_{14} \\ &\quad + a_2 a_4 (k_2 - k_4)^2 E_{24} + a_3 a_4 (k_3 - k_4)^2 E_{34}. \end{aligned}$$

Θ in (5.12) can be written as follows: for $\tilde{k}_i \geq 0$,

$$\tilde{k}_1^2 + \tilde{k}_3^2 := k_1, \quad \tilde{k}_1^2 + \tilde{k}_4^2 := k_2, \quad \tilde{k}_2^2 + \tilde{k}_3^2 := k_3.$$

Then $k_4 = k_2 + k_3 - k_1 = \tilde{k}_2^2 + \tilde{k}_4^2$, exactly as in (1.16). Therefore, our new variables will be \tilde{k}_i , but in order to avoid too much notation, we drop the tildes. Replacing in (5.12), we obtain a new representation of Θ :

$$\begin{aligned} \Theta(t, x, y) &= a_1 \exp((k_1^2 + k_3^2)y) + a_2 \exp((k_1^2 + k_4^2)y) \\ &\quad + a_3 \exp((k_2^2 + k_3^2)y) + a_4 \exp((k_2^2 + k_4^2)y). \end{aligned} \quad (5.13)$$

(Compare with (1.16).) Repeating again (5.9) with (5.13), we obtain e.g.

$$\begin{aligned} &H(a_1 \exp((k_1^2 + k_3^2)y)) \\ &= (a_1(k_1^2 + k_3^2) - a_{1,xx}) \exp((k_1^2 + k_3^2)y) \\ &\quad + (-4a_{1,x}(k_1 k_{1,x} + k_3 k_{3,x}) - 2a_1(k_1 k_{1,x} + k_3 k_{3,x})_x) y \exp((k_1^2 + k_3^2)y) \\ &\quad - 4a_1(k_1 k_{1,x} + k_3 k_{3,x})^2 y^2 \exp((k_1^2 + k_3^2)y). \end{aligned}$$

By the linear independence among the exponentials, the nontrivial character of Θ , and the hypothesis $H(\Theta) \in \mathcal{W}_4$, it is clear that one will obtain $k_1 k_{1,x} + k_3 k_{3,x} = 0$, so that if $k_j(t, 0) =: k_j(t)$,

$$(k_1^2 + k_3^2)(t, x) = k_1^2(t) + k_3^2(t).$$

A similar argument reveals that

$$\begin{aligned}(k_1^2 + k_4^2)(t, x) &= k_1^2(t) + k_4^2(t), \\ (k_2^2 + k_3^2)(t, x) &= k_2^2(t) + k_3^2(t), \\ (k_2^2 + k_4^2)(t, x) &= k_2^2(t) + k_4^2(t).\end{aligned}$$

Consequently, we get in (5.13)

$$\begin{aligned}\Theta(t, x, y) &= a_1(t, x) \exp((k_1^2(t) + k_3^2(t))y) + a_2(t, x) \exp((k_1^2(t) + k_4^2(t))y) \\ &\quad + a_3(t, x) \exp((k_2^2(t) + k_3^2(t))y) + a_4(t, x) \exp((k_2^2(t) + k_4^2(t))y).\end{aligned}\tag{5.14}$$

Now, repeating (5.10) with (5.14) one has

$$\begin{aligned}Ai(a_1(t, x) \exp((k_1^2(t) + k_3^2(t))y)) \\ = (a_{1,t}(t, x) + 2(k_1 k_{1,t} + k_2 k_{2,t})y a_1(t, x) - a_{1,xxx}(t, x)) \exp((k_1^2(t) + k_3^2(t))y),\end{aligned}$$

revealing that $k_1^2(t) + k_2^2(t) = k_1^2(0) + k_2^2(0) =: k_1^2 + k_2^2$ are constants independent of time. Similarly,

$$\begin{aligned}(k_1^2 + k_4^2)(t) &= k_1^2(0) + k_4^2(0) =: k_1^2 + k_4^2, \\ (k_2^2 + k_3^2)(t) &= k_2^2(0) + k_3^2(0) =: k_2^2 + k_3^2, \\ (k_2^2 + k_4^2)(t) &= k_2^2(0) + k_4^2(0) =: k_2^2 + k_4^2.\end{aligned}$$

Consequently, we get in (5.14),

$$\begin{aligned}\Theta(t, x, y) &= a_1(t, x) \exp((k_1^2 + k_3^2)y) + a_2(t, x) \exp((k_1^2 + k_4^2)y) \\ &\quad + a_3(t, x) \exp((k_2^2 + k_3^2)y) + a_4(t, x) \exp((k_2^2 + k_4^2)y).\end{aligned}\tag{5.15}$$

From the hypothesis (1.21), and following $\Theta(0, x, 0)$, $\Theta_y(0, x, 0)$, $\Theta_{yy}(0, x, 0)$ and $\Theta_{yyy}(0, x, 0)$ are uniquely determined by the values from (1.16), leading to the equations

$$\begin{aligned}a_1(0, x) + a_2(0, x) + a_3(0, x) + a_4(0, x) &= \Theta(0, x, 0) \\ (k_1^2 + k_3^2)a_1(0, x) + (k_1^2 + k_4^2)a_2(0, x) \\ &\quad + (k_2^2 + k_3^2)a_3(0, x) + (k_2^2 + k_4^2)a_4(0, x) = \Theta_y(0, x, 0) \\ (k_1^2 + k_3^2)^2 a_1(0, x) + (k_1^2 + k_4^2)^2 a_2(0, x) \\ &\quad + (k_2^2 + k_3^2)^2 a_3(0, x) + (k_2^2 + k_4^2)^2 a_4(0, x) = \Theta_{yy}(0, x, 0) \\ (k_1^2 + k_3^2)^3 a_1(0, x) + (k_1^2 + k_4^2)^3 a_2(0, x) \\ &\quad + (k_2^2 + k_3^2)^3 a_3(0, x) + (k_2^2 + k_4^2)^3 a_4(0, x) = \Theta_{yyy}(0, x, 0).\end{aligned}$$

This is a classical invertible system thanks to the Vandermonde determinant and the condition $k_1 < k_2 < k_3 < k_4$. Therefore, $a_j(0, x)$, $j = 1, 2, 3, 4$ are uniquely determined:

$$a_1(0, x) = (k_3 - k_1) \exp((k_1 + k_3)x), \quad a_2(0, x) = (k_4 - k_1) \exp((k_1 + k_4)x),$$

and

$$a_3(0, x) = (k_3 - k_2) \exp((k_2 + k_3)x), \quad a_4(0, x) = (k_4 - k_2) \exp((k_2 + k_4)x).$$

Using (5.15), we compute now $Ai(\Theta) - \frac{3}{2}\partial_x H(\Theta)$:

$$\begin{aligned} Ai(\Theta) - \frac{3}{2}\partial_x H(\Theta) &= \left(a_{1,t} + \frac{1}{2}a_{1,xxx} - \frac{3}{2}(k_1^2 + k_3^2)a_{1,x} \right) \exp((k_1^2 + k_3^2)y) \\ &\quad + \left(a_{2,t} + \frac{1}{2}a_{2,xxx} - \frac{3}{2}(k_1^2 + k_4^2)a_{2,x} \right) \exp((k_1^2 + k_4^2)y) \\ &\quad + \left(a_{3,t} + \frac{1}{2}a_{3,xxx} - \frac{3}{2}(k_2^2 + k_3^2)a_{3,x} \right) \exp((k_2^2 + k_3^2)y) \\ &\quad + \left(a_{4,t} + \frac{1}{2}a_{4,xxx} - \frac{3}{2}(k_2^2 + k_4^2)a_{4,x} \right) \exp((k_2^2 + k_4^2)y) = 0. \end{aligned}$$

Therefore,

$$\begin{aligned} a_{1,t} + \frac{1}{2}\partial_x(a_{1,xx} - 3(k_1^2 + k_3^2)a_1) &= 0, \\ a_1(0, x) &= (k_3 - k_1)\exp((k_1 + k_3)x). \end{aligned} \tag{5.16}$$

We shall prove that the unique solution to this problem is

$$a_1(t, x) = (k_3 - k_1)\exp((k_1 + k_3)x + (k_1^3 + k_3^3)t). \tag{5.17}$$

If we assume this equality, it is not hard to see that a similar argument reveals that

$$\begin{aligned} a_2(t, x) &= (k_4 - k_1)\exp((k_1 + k_4)x + (k_1^3 + k_4^3)t), \\ a_3(t, x) &= (k_3 - k_2)\exp((k_2 + k_3)x + (k_2^3 + k_3^3)t), \end{aligned}$$

and

$$a_4(t, x) = (k_4 - k_2)\exp((k_2 + k_4)x + (k_2^3 + k_4^3)t).$$

This finally proves Theorem 1.13. Let us show (5.17). Clearly the RHS of (5.17) is a valid solution to (5.16) satisfying the initial condition $a_1(0, x) = (k_3 - k_1)\exp((k_1 + k_3)x)$. Let us show that it is the unique one.

First, notice that if $\tilde{a}_1(t, y)$ is a function such that $a_1(t, x) = \tilde{a}_1(t, 2^{1/3}(x + \frac{3}{2}(k_1^2 + k_3^2)t))$, then

$$\tilde{a}_{1,t} + \tilde{a}_{1,xxx} = 0.$$

Consequently, by the following uniqueness we get the desired result.

Lemma 5.4 (Uniqueness of exponentially growing Airy solutions). *Let $m_1, m_2 > 0$. There is a unique solution u of*

$$\partial_t u + \partial_x^3 u = 0, \quad u(t = 0, x) = m_1 \exp(m_2 x),$$

and it is given by

$$u(t, x) = m_1 \exp(m_2 x - m_2^3 t).$$

Proof. The existence is exactly given by the explicit formula. Let us see the uniqueness, which is equivalent to prove that

$$\partial_t u + \partial_x^3 u = 0, \quad u(t = 0, x) = 0,$$

has solution $u = 0$. Assume $x > 0$. Thanks to the exponential bound on x for Θ , this is obtained by simply taking Laplace transform, solving the corresponding obtained ODE, and using the uniqueness of the inverse Laplace transform in the exponentially growing class of solutions. The remaining case $x < 0$ is proved similarly by changing $u(t, x)$ by $u(t, -x)$. \square

6. PROOF IN THE ZK CASE

In this section we provide a proof for Theorem 1.16 for the quadratic ZK model (1.22). Notice that this equation can be recast in terms of KdV (B.1) as follows:

$$0 = (-4u_t + \partial_{x_1}^3 u + 6u\partial_{x_1} u) + \partial_{x_1}(\Delta_c u), \quad (6.1)$$

where $\Delta_c u = \sum_{j=2}^d \partial_{x_j}^2 u$. For simplicity in the notation, let $x = x_1$. Inserting $u = 2\partial_x^2 F(\Theta)$ into (6.1), and integrating once in x , we obtain

$$\begin{aligned} 0 = & (F'''' + 6F''^2)(\Theta) \Theta_x^4 + 6(F'' + F'^2)'(\Theta) \Theta_x^2 \Theta_{xx} \\ & + (F'' + F'^2)(\Theta) (3\Theta_{xx}^2 - 4\Theta_x(\Theta_t - \Theta_{xxx})) + F'(\Theta) (-4\Theta_t + \Theta_{xxx})_x \\ & + F'^2(\Theta) (3(\Theta_{xx}^2 - \Theta_x \Theta_{xxx}) - \Theta_x(-4\Theta_t + \Theta_{xxx})) \\ & + \sum_{j=2}^d \left(\Theta_{x_j x_j} F'(\Theta) + \Theta_{x_j}^2 F''(\Theta) \right)_{xx}. \end{aligned}$$

The term $\Theta_{x_j x_j} F'(\Theta) + \Theta_{x_j}^2 F''(\Theta)$ can be written as

$$F'(\Theta) W_{x_j}^F(\Theta) + \Theta_{x_j}^2 \rho(\Theta), \quad \rho = F''(\Theta) + F'^2(\Theta).$$

Therefore, following the same ideas as in the KP-II case, we arrive to the equation

$$0 = \text{EDO}_4(\rho) - F'(\Theta)(Ai(\Theta))_x + F'^2(\Theta)(3W_1(\Theta) + \Theta_x Ai(\Theta)) + \sum_{j=2}^d \left(F'(\Theta) W_{x_j}^F(\Theta) \right)_{xx},$$

where $\text{EDO}_4(\rho)$ is a fourth order linear ODE for ρ . Now we finish the proof. If u is a KdV soliton and $F = \log$, it is easy to check that (1.23) is satisfied. Now, if (1.23) is satisfied, $\text{EDO}_4(\rho) = 0$ and $F = \log$. Therefore, $W_{x_j}^F(\Theta) = W_{x_j}^{\log}(\Theta) = 0$. From $W_1(\Theta) = 0$ we get

$$\Theta = a(x_2, \dots, x_d, t) \exp(xb(x_2, \dots, x_d, t)) + c(x_2, \dots, x_d, t).$$

From $W_{x_j}^{\log}(\Theta) = 0$ we get that a, b, c are only time dependent. Finally, from Airy we obtain that $u = Q_k$, for some constant $k > 0$.

6.1. Proof in the mZK case. In this section we provide proofs for Theorem 1.17 for the cubic version ZK model (1.22). Notice that, as above, this equation can be recast in terms of mKdV (B.9) as follows:

$$0 = (-4u_t + \partial_{x_1}^3 u + 6u^2 \partial_{x_1} u) + \partial_{x_1}(\Delta_c u),$$

where $\Delta_c u = \sum_{j=2}^d \partial_{x_j}^2 u$. For simplicity in the notation, let $x = x_1$. Replacing $u = 2\partial_x F(\Theta)$ in (6.1), and integrating once in x , we obtain

$$\begin{aligned} 0 = & (\Theta_{xxx} - 4\Theta_t) F'(\Theta) + 3\Theta_{xx} \Theta_x F''(\Theta) + \Theta_x^3 (F''' + 2F'^3)(\Theta) \\ & + \sum_{j=2}^d \left(\Theta_{x_j x_j} F'(\Theta) + \Theta_{x_j}^2 F''(\Theta) \right)_x \end{aligned} \quad (6.2)$$

Notice in contrast of ZK case, the setting $Ai(\Theta) = 0$, $W(\Theta) = 0$ with $W_{x_i}^F = 0$ produces a contradiction in (6.2), when considering $F = 2 \arctan$. Thus the condition $\Lambda_{x_j}^F(\Theta) =$

$\Theta_{x_j x_j} F'(\Theta) + \Theta_{x_j}^2 F''(\Theta) = 0$ enters in action. Similarly as in the ZK case, we arrive to the equation

$$\begin{aligned} 0 = & (\Theta_{xxx} - 4\Theta_t) F'(\Theta) + 3 \left(\Theta_{xx} - \frac{\Theta_x^2}{\Theta} \right) h'(\Theta) \\ & + \Theta_x^2 \left(h'' + \frac{3}{s} h' + 8h^3 \right) (\Theta) + \sum_{j=2}^d \left(\Lambda_{x_j}^F(\Theta) \right)_x, \end{aligned} \quad (6.3)$$

where $h(s) = \frac{1}{2} F'(s)$. Now we finish the proof. If u is a mKdV soliton (see (B.11)), it is easy to check that (6.3) is satisfied. Now, if

$$Ai(\Theta) = W(\Theta) = \Lambda_{x_j}^F(\Theta) = 0, \quad j = 2, \dots, d, \quad (6.4)$$

the second term in (6.3) (being zero now) can be written as the radial solution $h = h(s)$, $s = |x| > 0$ of $\Delta h + 8h^3 = 0$ in dimension 4. Requiring positive solutions (by hypothesis, $h(s) := \frac{1}{2} F'(s) > 0$), these are in $H^1(\mathbb{R}^4)$ and by classical Talenti-Aubin arguments, one has $h(s) = \frac{1}{1+s^2}$, and finally $F = 2 \arctan$ if we assume $F(0) = 0$. Therefore,

$$\Lambda_{x_j}^F(\Theta) = \Lambda_{x_j}^{2 \arctan}(\Theta) = 0, \quad i = 2, \dots, d.$$

Now, from $W(\Theta) = 0$ we get

$$\Theta = a(x_2, \dots, x_d, t) \exp(xb(x_2, \dots, x_d, t)) + c(x_2, \dots, x_d, t).$$

Also since $\Lambda_{x_j}^{2 \arctan}(\Theta) = 0$ we will get that a, b, c are only time dependent. Indeed if $a := a(x_2, \dots, x_d, t)$, $b := b(x_2, \dots, x_d, t)$, $c := c(x_2, \dots, x_d, t)$, replacing Θ in $\Lambda_{x_j}^F(\Theta) = 0$ for each $j = 2, \dots, d$, one has

$$\begin{aligned} 0 = & A_0 + A_1 \exp(bx) + A_2 x \exp(bx) + A_3 x^2 \exp(bx) \\ & + B_1 \exp(2bx) + B_2 x \exp(2bx) + B_3 x^2 \exp(2bx) \\ & + C_1 \exp(3bx) + C_2 x \exp(3bx) + C_3 x^2 \exp(3bx), \end{aligned}$$

where the A_0, A_1, A_2, A_3 are described by

$$\begin{aligned} A_0 &= -2cc'^2 + c^2 c'' + c'' \\ A_1 &= -4ca'c' + c^2 a'' + a'' - 2ac'^2 + 2acc'' \\ A_2 &= 2c^2 a'b' + 2a'b' - 4acb'c' + ac^2 b'' + ab'' \\ A_3 &= ac^2 b'^2 + ab'^2; \end{aligned}$$

the values of B_1, B_2 and B_3 are

$$\begin{aligned} B_1 &= -4aa'c' - 2ca'^2 + 2aca'' + a^2 c'' \\ B_2 &= 2a^2 cb'' - 4a^2 b'c' \\ B_3 &= 0, \end{aligned}$$

and C_1, C_2, C_3 are given by

$$\begin{aligned} C_1 &= -2aa'^2 + a^2 a'' - x^2 a^3 b'^2 \\ C_2 &= -2aa'^2 + a^2 a'' \\ C_3 &= a^3 b'^2. \end{aligned}$$

Here the prime derivative denotes partial derivative in x_j . Notice that $A_i = B_i = C_i = 0$ for $i = 1, 2, 3$.

Since $C_3 = a^3 b_{x_j}^2$ need to be zero, we have b is constant in x_j . From $C_2 = 0$, taking in mind $w = a^{-1}$ as a change of variable of this ODE, we conclude that

$$a(x_j) = \frac{1}{Ax_j^2 + Bx_j + C}. \quad (6.5)$$

Replacing a in B_1 , after some simplifications, one has

$$\begin{aligned} 0 = & (4A^2c + 12ABc' + 2ACc'' + B^2c'')x_j^2 + (8A^2c' + 2ABc'')x_j^3 \\ & + A^2c''x_j^4 + (4ABc + 8ACc' + 4B^2c' + 2BCc'')x_j \\ & - 4ACc + 2B^2c + 4BCc' + C^2c''. \end{aligned} \quad (6.6)$$

Also recall from that A_1

$$0 = -4ca'c' + c^2a'' + a'' - 2ac'^2 + 2acc''. \quad (6.7)$$

Now, the term x_j^4 implies $A = 0$ or $c'' = 0$. If $A \neq 0$ from the term x_j^3 one has $c' = 0$ then the term x_j^2 implies $c = 0$. Thus (6.7) yields $a'' = 0$ which contradicts (6.5). Hence $A = 0$. Similarly from the term x_j^2 of (6.6) one has $B = 0$. Hence a is constant in x_j . Now, $A = B = 0$ in (6.6) implies $c'' = 0$. Notice that (6.7) now is described by

$$0 = \frac{2cc''}{C} - \frac{2c'^2}{C} = -\frac{2c'^2}{C}.$$

Thus c is constant in x_j . Finally, from Airy we obtain that $u = Q_k$ as in (B.11), for some constant $k > 0$.

APPENDIX A. PROOF OF TECHNICAL RESULTS

A.1. Galilean actions. Evaluation Θ_β , obtained after apply the Galilean Transformation to a phase Θ , in the terms that appears in (2.1). Then the terms in Definition 1.1 satisfy,

$$\begin{aligned} H(\Theta_\beta) &= -\frac{4\beta}{3}\partial_{\tilde{x}}\Theta + \partial_{\tilde{y}}\Theta - \partial_{\tilde{x}}^2\Theta = -\frac{4\beta}{3}\partial_{\tilde{x}}\Theta + H(\Theta), \\ Ai(\Theta_\beta) &= \frac{4\beta^2}{3}\partial_{\tilde{x}}\Theta - 2\beta\partial_{\tilde{y}}\Theta + \partial_{\tilde{t}}\Theta - \partial_{\tilde{x}}^3\Theta = \frac{4\beta^2}{3}\partial_{\tilde{x}}\Theta - 2\beta\partial_{\tilde{y}}\Theta + Ai(\Theta), \\ \Theta_\beta W_y(\Theta_\beta) &= \Theta W_{\tilde{y}}(\Theta) + \frac{16}{9}\beta^2(\Theta\Theta_{\tilde{x}\tilde{x}} - \Theta_{\tilde{x}}^2) - \frac{8}{3}\beta(\Theta\Theta_{\tilde{x}\tilde{y}} - \Theta_{\tilde{x}}\Theta_{\tilde{y}}), \\ \Theta_\beta W_x(\Theta_\beta) &= \Theta W_{\tilde{x}}(\Theta). \end{aligned} \quad (A.1)$$

In particular H , Ai and W_y do not cancel for a nontrivial Galilean version of the vertical soliton (1.14).

A.2. Proof of phase computations. In this section we prove (4.6). Given Θ as in (1.16), its derivatives are

$$\begin{aligned}
\Theta_x &= (k_3 - k_1)(k_3 + k_1) \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4 + k_1) \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3 + k_2) \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4 + k_2) \exp(\theta_2 + \theta_4), \\
\Theta_{xx} &= (k_3 - k_1)(k_3 + k_1)^2 \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4 + k_1)^2 \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3 + k_2)^2 \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4 + k_2)^2 \exp(\theta_2 + \theta_4), \\
\Theta_{xxx} &= (k_3 - k_1)(k_3 + k_1)^3 \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4 + k_1)^3 \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3 + k_2)^3 \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4 + k_2)^3 \exp(\theta_2 + \theta_4), \\
\Theta_{xxxx} &= (k_3 - k_1)(k_3 + k_1)^4 \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4 + k_1)^4 \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3 + k_2)^4 \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4 + k_2)^4 \exp(\theta_2 + \theta_4).
\end{aligned}$$

Also,

$$\begin{aligned}
\Theta_y &= (k_3 - k_1)(k_3^2 + k_1^2) \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4^2 + k_1^2) \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3^2 + k_2^2) \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4^2 + k_2^2) \exp(\theta_2 + \theta_4), \\
\Theta_{yy} &= (k_3 - k_1)(k_3^2 + k_1^2)^2 \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4^2 + k_1^2)^2 \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3^2 + k_2^2)^2 \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4^2 + k_2^2)^2 \exp(\theta_2 + \theta_4), \\
\Theta_t &= (k_3 - k_1)(k_3^3 + k_1^3) \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4^3 + k_1^3) \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3^3 + k_2^3) \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4^3 + k_2^3) \exp(\theta_2 + \theta_4).
\end{aligned}$$

Consequently,

$$\begin{aligned}
Ai(\Theta) &= \Theta_t - \Theta_{xxx} \\
&= (k_3 - k_1)(k_3^3 + k_1^3) \exp(\theta_1 + \theta_3) + (k_4 - k_1)(k_4^3 + k_1^3) \exp(\theta_1 + \theta_4) \\
&\quad + (k_3 - k_2)(k_3^3 + k_2^3) \exp(\theta_2 + \theta_3) + (k_4 - k_2)(k_4^3 + k_2^3) \exp(\theta_2 + \theta_4) \\
&\quad - (k_3 - k_1)(k_3 + k_1)^3 \exp(\theta_1 + \theta_3) - (k_4 - k_1)(k_4 + k_1)^3 \exp(\theta_1 + \theta_4) \\
&\quad - (k_3 - k_2)(k_3 + k_2)^3 \exp(\theta_2 + \theta_3) - (k_4 - k_2)(k_4 + k_2)^3 \exp(\theta_2 + \theta_4).
\end{aligned}$$

This equation is equal to 0 when all the coefficient which multiplies the exponentials are null. Note that all the terms are exponentials, $\exp(\theta_j + \theta_i)$ multiplies by a term

$$\begin{aligned}
&(k_i - k_j)(k_i^3 + k_j^3) - (k_i - k_j)(k_i + k_j)^3 \\
&= k_i^4 + k_i k_j^3 - k_i^3 k_j - k_j^4 - (k_i - k_j)(k_i^3 + 3k_i^2 k_j + 3k_i k_j^2 + k_j^3) \\
&= k_i^4 + k_i k_j^3 - k_i^3 k_j - k_j^4 - 3k_i^3 k_j - 3k_i^2 k_j^2 - k_i k_j^3 + k_i^3 k_j + 3k_i^2 k_j^2 + 3k_i k_j^3 + k_j^4 \\
&= -3k_i^3 k_j + 3k_i k_j^3 = 3(k_i k_j^3 - k_i^3 k_j).
\end{aligned}$$

A.3. Computation of derivatives of Θ . We perform here some of the computations required in (5.6) and subsequent lines. If $\Theta = \sum_{j=1}^M A_j(t, x) \exp(B_j(t, x)y)$, then one has

$$\begin{aligned}\Theta_x &= \sum_{j=1}^M (A_{j,x} + A_j B_{j,x}) \exp(B_j y), \\ \Theta_{xx} &= \sum_{j=1}^M (A_{j,xx} + 2A_{j,x} B_{j,x} y + A_j B_{j,xx} y + A_j B_{j,x}^2 y^2) \exp(B_j y), \\ \Theta_{xxx} &= \sum_{j=1}^M \left(A_{j,xxx} + 3A_{j,x} B_{j,x} y + 3A_{j,x} B_{j,xx} y + 3A_{j,x} B_{j,x}^2 y^2 \right. \\ &\quad \left. + A_j B_{j,xxx} y + 3A_j B_{j,x} B_{j,xx} y^2 + A_j B_{j,x}^3 y^3 \right) \exp(B_j y).\end{aligned}$$

Additionally,

$$\begin{aligned}\Theta_{xxxx} &= \sum_{j=1}^M \left(A_{j,xxxx} + 4A_{j,xxx} B_{j,x} y + 6A_{j,xx} B_{j,xx} y \right. \\ &\quad \left. + 4A_{j,x} B_{j,xxx} y + A_j B_{j,xxxx} y + 6A_{j,xx} B_{j,x}^2 y^2 \right. \\ &\quad \left. + 2A_{j,x} B_{j,x} B_{j,xx} y^2 + 3A_j B_{j,xx}^2 y^2 + A_j B_{j,x} B_{j,xxx} y^2 \right. \\ &\quad \left. + 4A_{j,x} B_{j,x}^3 y^3 + 6A_j B_{j,x}^2 B_{j,xx} y^3 + A_j B_{j,x}^4 y^4 \right) \exp(B_j y), \\ \Theta_y &= \sum_{j=1}^M A_j B_j \exp(B_j y), \quad \Theta_{yy} = \sum_{j=1}^M A_j B_j^2 \exp(B_j y), \\ \Theta_t &= \sum_{j=1}^M (A_{j,t} + A_j B_{j,t} y) \exp(B_j y).\end{aligned}$$

APPENDIX B. THE KdV AND MKdV CASES

For the sake of completeness, we provide a sketch of proofs for Remarks 1.18 and 1.19.

B.1. The KdV case. Consider the KdV model

$$-4u_t + u_{xxx} + 6uu_x = 0, \quad (\text{B.1})$$

where $u = u(t, x) \in \mathbb{R}$ and $t, x \in \mathbb{R}$. We study the solution u of the form

$$u(t, x) = 2\partial_x^2 F(\Theta(t, x)), \quad (\text{B.2})$$

where, with no loss of generality, we consider $F : [1, \infty) \rightarrow \mathbb{R}$ is smooth and $\Theta = \Theta(t, x) \in [1, \infty)$ is also smooth. Since F can be changed by any linear affine function, we can assume that $F(1) = 0$, $F'(1) = 1$. The simplest case, the KdV soliton, is found as

$$F(s) := \log s, \quad \Theta(t, x) = 1 + \exp(ax - ta^3/4), \quad a \in \mathbb{R}. \quad (\text{B.3})$$

Exactly as in the KP case, we set the following definitions:

Definition B.1 (Classification of phases Θ). We shall say that Θ as in (B.2)

(i) is of Airy type if for all $(t, x) \in \mathbb{R}^2$,

$$Ai(\Theta) := -4\Theta_t + \Theta_{xxx} = 0;$$

(ii) is of Wronskian type if $\{\Theta_x, \Theta_{xx}\}(t, x)$ are linearly dependent, for any (t, x) ;

(iii) is of \mathcal{T} -type if for F fixed,

$$\mathcal{T}(\Theta) := (Ai(\Theta)_x + F'(\Theta)(3W(\Theta) - \Theta_x Ai(\Theta))) = 0, \quad (\text{B.4})$$

$$\text{where } W(\Theta) := \Theta_{xx}^2 - \Theta_x \Theta_{xxx}.$$

Some comments are necessary.

Remark B.2. The Airy type condition naturally describes that the phase of the nonlinear KdV solution u solves the classical Airy linear equation

$$4\Theta_t - \Theta_{xxx} = 0.$$

This is an interesting coincidence that confirms the complex integrable structure of the KdV model. Standard solutions to the Airy equation have a complex oscillatory behavior for $x < 0$, but some simple solutions are

$$\Theta(t, x) = \exp\left(ax - \frac{1}{4}a^3t + C\right), \quad \Theta(t, x) = t + \frac{2}{3}x^3 + C, \quad (\text{B.5})$$

where $C \in \mathbb{R}$ is any constant. The first phase corresponds to the 1-soliton phase, and the second will represent an interesting counterexample to our main results.

Remark B.3. Unlike KP, here in KdV the Wronskian type phase is extremely restrictive. Multi-soliton solutions will not be of this type.

Remark B.4. Exactly as in KP, notice that Θ of \mathcal{T} -type is a condition depending on the profile F , and consequently is a more complex condition than being of Wronskian or Airy type, which are independent of the profile F . Additionally, the 1-soliton phase (B.3) is of Airy type as well.

Inserting (B.2) to (B.1), and arranging similar terms, one easily arrives at

$$\begin{aligned} & (F'''' + 6F''^2)(\Theta)\Theta_x^4 + 6(F'' + F'^2)'(\Theta)\Theta_x^2\Theta_{xx} \\ & + (F'' + F'^2)(\Theta)(3\Theta_{xx}^2 - 4\Theta_x(\Theta_t - \Theta_{xxx})) + F'(\Theta)(-4\Theta_t + \Theta_{xxx})_x \\ & + F'^2(\Theta)(3(\Theta_{xx}^2 - \Theta_x\Theta_{xxx}) - \Theta_x(-4\Theta_t + \Theta_{xxx})) = 0. \end{aligned} \quad (\text{B.6})$$

(Compare with (1.4).) As in KP, we set

$$\rho(s) := F''(s) + F'^2(s).$$

If $\rho = 0$ and $F(1) = 0$, $F'(1) = -1$, then $F = \log$. This follows directly from solving the ODE $\rho(s) = 0$. Notice that from (2.3) and Definition B.1, one has that (B.6) can be written as

$$\begin{aligned} & (\rho'' - 2F'\rho' + 4F''\rho)\Theta_x^4 + 6\rho'\Theta_x^2\Theta_{xx} \\ & + \rho(3\Theta_{xx}^2 - 4\Theta_x(\Theta_t - \Theta_{xxx})) + F'(\Theta)(Ai(\Theta))_x \\ & + F'^2(\Theta)(3W(\Theta) - \Theta_x Ai(\Theta)) = 0, \end{aligned}$$

and arranging terms,

$$\begin{aligned} & \Theta_x^4 \rho'' + 2(3\Theta_x^2 \Theta_{xx} - F'(\Theta)\Theta_x^4)\rho' \\ & + (3\Theta_{xx}^2 - 4\Theta_x(\Theta_t - \Theta_{xxx}) + 4F''(\Theta)\Theta_x^4)\rho + F'(\Theta)\mathcal{T}(\Theta) = 0. \end{aligned} \quad (\text{B.7})$$

Our main result is

Theorem B.5. *The following are satisfied:*

- (i) *Assume that Θ is of \mathcal{T} -type, Θ_x is different from 0, and F satisfies $F''(1) = 1$ and $F'''(1) = -2$. Then $F = \log$.*
- (ii) *Assume that $\Theta > 0$ is of Airy and Wronskian type, $\Theta_x \neq 0$, and F satisfies $F''(1) = 1$ and $F'''(1) = -2$. Then $F = \log$ and $\Theta = 1 + \exp(ax - a^3/4t)$, for any $a \in \mathbb{R}$.*
- (iii) *Assume that u as in (B.2) solves KdV with $F(s) = \log s$, and Θ is of Wronskian type. Then u is a soliton.*
- (iv) *Assume that u is a nontrivial multisoliton with $F = \log$. Then Θ is of \mathcal{T} -type, but it cannot be of Wronskian or Airy type.*

The simplest phase to characterize is that of Wronskian type.

Lemma B.6. *Assume that Θ is of Wronskian type. Then*

- (i) *One has $W(\Theta) = \Theta_{xx}^2 - \Theta_x \Theta_{xxx} = 0$.*
- (ii) *Additionally, there are $a = a(t)$ and $b = b(t)$ such that either*

$$\Theta(t, x) = 1 + \exp(a(t)x + b(t)),$$

or

$$\Theta(t, x) = a(t)x + b(t).$$

The previous result establishes that phases Θ of Wronskian type are directly related to soliton solutions. This fact is independent of the value of F , that will be determined independently.

Proof. The proof follows directly from Definition B.1 (ii). Indeed, the proof of (i) is direct from the definition of the linear dependence and the fact that

$$W(\Theta) = \Theta_{xx}^2 - \Theta_x \Theta_{xxx} = \text{Wr}(\Theta_x, \Theta_{xx}),$$

where Wr denotes the Wronskian. The proof of (ii) follows from the fact that $\eta := \Theta_x$ satisfies the ODE $\eta_x^2 = \eta \eta_{xx}$, which has solutions $\eta(x) = \exp(a(t)x + b(t))$ and $\eta = a(t)$. The final result is obtained by integrating in space. \square

Remark B.7. 1. Notice that multisolitons cannot have a phase of Wronskian type.

2. An important example of phase of Airy type but not being of Wronskian type is the one given in (B.5): $\Theta(t, x) = t + \frac{2}{3}x^3 + 1$. In this case, $W(\Theta) = 8x^2$.

3. If Θ is of Wronskian and Airy type, then is of \mathcal{T} -type. The reciprocal is clearly false. This follows directly from (B.4).

Lemma B.8. *Assume that Θ is of \mathcal{T} -type. Then*

- (i) *There exists $c_0 \in \mathbb{R}$ such that*

$$Ai(\Theta) = c_0 \exp(F(\Theta)) - 3W(\Theta) + 3F(\Theta) \int_0^x \exp(-F(\Theta)) (W(\Theta))_x(t, s) ds. \quad (\text{B.8})$$

- (ii) *If Θ is of Airy and \mathcal{T} -type, and $F'(\Theta)$ is different from 0, then it is of Wronskian type.*

(iii) If Θ is of Wronskian and \mathcal{T} -type, then there exists $c \in \mathbb{R}$ such that $Ai(\Theta) = c_0 \exp(\Theta)$.

Proof. Let us prove (i): notice that Θ being of \mathcal{T} -type as in (B.4) is equivalent to have

$$(Ai(\Theta))_x - F'(\Theta) \Theta_x Ai(\Theta) = 3F'(\Theta) W(\Theta).$$

Then (B.8) follows directly from solving the corresponding ODE for $Ai(\Theta)$. In order to prove (ii), notice that from (B.4) one has $0 = F'(\Theta) W(\Theta)$, proving (ii). Finally, to prove (iii), from (B.8) one has $Ai(\Theta) = c_0 \exp(F(\Theta))$, as required. \square

Now we are ready to prove the main result in the KdV case.

Proof of Theorem B.5. Now we are ready to prove Theorem B.5.

Proof of (i). Under $\mathcal{T}(\Theta) = 0$, we have from (B.7) an ODE of the form

$$\Theta_x^4 \rho'' + 2(3\Theta_x^2 \Theta_{xx} - F'(\Theta) \Theta_x^4) \rho' + (3\Theta_{xx}^2 - 4\Theta_x(\Theta_t - \Theta_{xxx}) + 4F''(\Theta) \Theta_x^4) \rho = 0.$$

This homogeneous ODE for ρ has zero as the unique solution provided $\rho(1) = \rho'(1) = 0$, which is indeed the case. Consequently, from Lemma 2.4, we get $F = \log$.

Proof of (ii). From Θ being of Airy and Wronskian type we have Θ of \mathcal{T} -type. The previous result ensures $F = \log$. Finally, Lemma B.6 proves the final result, after checking that Θ is of Airy type and $\Theta > 0$.

Proof of (iii). If $F = \log$ then $\rho = 0$, and from (B.7), $\Theta^{-1} \mathcal{T}(\Theta) = 0$. Consequently, Θ is of \mathcal{T} -type. Since it is additionally of Wronskian type, it is of Airy type and (ii) applies.

Proof of (iv). The multisoliton Θ is of \mathcal{T} -type, but it does not satisfy being of Airy or Wronskian type. This ends the proof of Theorem B.5. \square

B.2. The mKdV case. Let us consider the mKdV model

$$-4u_t + u_{xxx} + 6u^2 u_x = 0, \quad (\text{B.9})$$

where $u = u(t, x) \in \mathbb{R}$ and $(t, x) \in \mathbb{R}^2$. Let us extend our previous results to the mKdV case. Consider

$$u(t, x) = \partial_x F(\Theta(t, x)), \quad (\text{B.10})$$

(notice that we only consider one derivative in space). We consider $F : \mathbb{R} \rightarrow \mathbb{R}$ smooth and the phase $\Theta = \Theta(t, x) \in \mathbb{R}$ also smooth. Since F can be changed by any constant, we can assume that $F(0) = 0$. Classical mKdV solitons (see e.g. [2]) are given by

$$\begin{aligned} Q_k(t, x) &= 2\partial_x \arctan(\Theta_k(t, x)) \\ &= k \operatorname{sech}\left(kx + \frac{1}{4}k^3t + b_0\right), \quad \Theta_k := \exp\left(kx + \frac{1}{4}k^3t + b_0\right), \quad b_0 \in \mathbb{R}. \end{aligned} \quad (\text{B.11})$$

Replacing u in (B.9) by (B.10), we obtain

$$(\Theta_{xxx} - 4\Theta_t) F'(\Theta) + 3\Theta_{xx} \Theta_x F''(\Theta) + \Theta_x^3 (F''' + 2F'^3)(\Theta) = 0. \quad (\text{B.12})$$

Trivial phases $\Theta = \text{const.}$ will be discarded now, so that we assume $\Theta_x \neq 0$. Note that Θ_k in (B.11) satisfies $Ai(\Theta) = \Theta_{xxx} - 4\Theta_t = 0$. Consequently, from (B.12) one must have

$$3\Theta_{xx} F''(\Theta) + \Theta_x^2 (F''' + 2F'^3)(\Theta) = 0.$$

Let $h(s) := \frac{1}{2}F'(s)$. The equation above can be written as

$$3 \left(\Theta_{xx} - \frac{\Theta_x^2}{\Theta} \right) h'(\Theta) + \Theta_x^2 \left(h'' + \frac{3}{s}h' + 8h^3 \right) (\Theta) = 0. \quad (\text{B.13})$$

The first term contains nothing but $W_x(\Theta) := \Theta_{xx} - \frac{\Theta_x^2}{\Theta}$, which will be required to be zero. Notice that Θ_k in (B.11) does satisfy this condition, i.e., $W_x(\Theta_k) = 0$. In the opposite direction, $W_x(\Theta) = 0$ implies $\Theta \left(\frac{\Theta_x}{\Theta} \right)_x = 0$, leading to $\Theta(t, x) = \exp(k(t)x + b(t))$. From the Airy condition $\Theta_{xxx} - 4\Theta_t = 0$ one gets

$$k^3 - 4k'x - 4b' = 0,$$

leading to $k(t) = k$ for some constant k and $b = \frac{1}{4}k^3t + b_0$ with some constant b_0 . Therefore, $\Theta = \exp(kx + \frac{1}{4}k^3t + b_0)$. Notice that the case $k = 0$ returns the trivial solution, therefore it will be left out of the subsequent analysis. In particular, the image of Θ is $(0, \infty)$.

Finally, the second term in (B.13) (being zero now) can be written as the radial solution $h = h(s)$, $s = |x| > 0$ of $\Delta h + 8h^3 = 0$ in dimension 4. Requiring positive solutions (by hypothesis, $h(s) := \frac{1}{2}F'(s) > 0$), these are in $H^1(\mathbb{R}^4)$ and by classical Talenti-Aubin arguments, one has $h(s) = \frac{1}{1+s^2}$, giving, finally, $F = 2 \arctan$ if we assume $F(0) = 0$. We conclude the following result:

Theorem B.9. *Let u be a smooth solution to mKdV (B.9) of the form (B.10), with smooth profile $F : \mathbb{R} \rightarrow \mathbb{R}$ satisfying $F(0) = 0$ and F strictly increasing in \mathbb{R} . Then a nontrivial Θ is a soliton (B.11) and $F = 2 \arctan$ if and only if $W_x(\Theta) = Ai(\Theta) = 0$.*

Notice that the condition F strictly increasing in \mathbb{R} can be replaced by F strictly monotone in \mathbb{R} , since by symmetries of the equation (B.9), if u is a solution, then $-u$ also does. However, it is noticed that ‘‘cn’’ periodic solutions may appear if this condition is lifted.

Now we assume that Θ_x may take the value zero. Additionally, we will assume that $Ai(\Theta)$ need not be zero as well. Coming back to (B.12), and following the ideas that led to (B.13), we shall obtain

$$(\Theta_{xxx} - 4\Theta_t)h + 3 \left(\Theta_{xx} - \frac{\Theta_x^2}{\Theta} \right) \Theta_x h'(\Theta) + \Theta_x^3 \left(h'' + \frac{3}{s}h' + 8h^3 \right) (\Theta) = 0.$$

This equation is valid provided $\Theta \neq 0$. If F is assumed to be increasing as in the previous argument, one has $h(s) = \frac{1}{1+s^2}$, leading to the equation

$$(1 + \Theta^2)(4\Theta_t - \Theta_{xxx}) + 6\Theta_x(\Theta\Theta_{xx} - \Theta_x^2) = 0, \quad (\text{B.14})$$

valid now in the full region. One can easily see that for any $\alpha, \beta > 0$, $\delta = \frac{1}{4}(3\beta^2 - \alpha^2)$, and $\gamma = \frac{1}{4}(\beta^2 - 3\alpha^2)$, one has that

$$\Theta = \pm \frac{\beta \sin(\alpha(x + \delta t))}{\alpha \cosh(\beta(x + \gamma t))},$$

solves the previous equation, representing the classical breather solution, see [2] and references therein. Also,

$$\Theta = \frac{e^{\sqrt{c_1}(x+c_1t)} + e^{\sqrt{c_2}(x+c_2t)}}{1 - \rho^2 e^{\sqrt{c_1}(x+c_1t) + \sqrt{c_2}(x+c_2t)}}, \quad \rho := \frac{\sqrt{c_1} - \sqrt{c_2}}{\sqrt{c_1} + \sqrt{c_2}},$$

with $c_1, c_2 > 0$, is another solution, representing the 2-soliton solution. In general, characterizing the solutions to (B.14) is not a simple task. In [6] it is proposed a first result in this direction, where one assumes that the phase Θ has a particular *monochromatic* structure.

REFERENCES

- [1] M.J. Ablowitz, and H. Segur, *Solitons and the inverse scattering transform*. SIAM Studies in Applied Mathematics, 4. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, Pa., 1981. x+425 pp. ISBN: 0-89871-174-6.
- [2] M. A. Alejo, and C. Muñoz, *Nonlinear stability of mKdV breathers*, Comm. Math. Phys. (2013), Vol. 324, Issue 1, pp. 233–262.
- [3] M.A. Alejo, C. Muñoz, and J. M. Palacios, *On the variational structure of breather solutions II: Periodic mKdV equation*. Electron. J. Differential Equations 2017, Paper No. 56, 26 pp.
- [4] J. Bourgain, *On the Cauchy problem for the Kadomtsev-Petviashvili equation*, Geom. Funct. Anal., 3 (1993), pp. 315–341.
- [5] A. de Bouard, and Y. Martel, *Non existence of L^2 -compact solutions of the Kadomtsev-Petviashvili II equation*, Math. Ann. 328, 525–544 (2004).
- [6] J. Faya, P. Figueroa, C. Muñoz and F. Poblete, *On nonexistence and uniqueness of breathers for the generalized Korteweg-de Vries and Zakharov-Kuznetsov models*, preprint arXiv:2404.09100, <https://arxiv.org/abs/2404.09100> (2024).
- [7] M. Hadac, *Well-posedness for the Kadomtsev-Petviashvili II equation and generalisations*, Trans. Amer. Math. Soc., 360 (2008), pp. 6555–6572.
- [8] M. Hadac, S. Herr, and H. Koch, *Well-posedness and scattering for the KP-II equation in a critical space*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 26 (2009), pp. 917–941.
- [9] N. Hayashi, and P.I. Naumkin, *Large Time Asymptotics for the Kadomtsev-Petviashvili Equation*, Commun. Math. Phys. 332, 505–533 (2014).
- [10] N. Hayashi, P.I. Naumkin, and J.-C. Saut, *Asymptotics for large time of global solutions to the generalized Kadomtsev-Petviashvili equation*, Comm. Math. Phys. 201 (1999), 577–590.
- [11] R. Iório, and W. Nunes, *On equations of KP type*. Proc. Roy. Soc. Edinburgh, 128 A, 725–743 (1998).
- [12] P. Isaza and J. Mejía, *Local and global Cauchy problems for the Kadomtsev-Petviashvili (KP-II) equation in Sobolev spaces of negative indices*, Comm. Partial Diff. Equations, 26 (2001), pp. 1027–1054.
- [13] P. Isaza, F. Linares, and G. Ponce, *On the propagation of regularity of solutions of the Kadomtsev-Petviashvili equation*, SIAM J. Math. Anal. 48 (2016), no. 2, 1006–1024.
- [14] B. B. Kadomtsev, and V. I. Petviashvili, *On the stability of solitary waves in weakly dispersing media*, Dokl. Akad. Nauk SSSR., 192 (1970), p. 753–756.
- [15] C. E. Kenig, and Y. Martel, *Global well-posedness in the energy space for a modified KP II equation via the Miura transform*, Trans. AMS 358 (2006), 2447–2488.
- [16] P.G. Kevrekidis, A. Khare and A. Saxena, *Breather lattice and its stabilization for the modified Korteweg-de Vries equation*, Phys. Rev. E, **68**, 0477011-0477014 (2003).
- [17] C. Klein, and J.-C. Saut, *IST versus PDE: a comparative study*. Hamiltonian partial differential equations and applications, 383–449, Fields Inst. Commun., 75, Fields Inst. Res. Math. Sci., Toronto, ON, 2015.

- [18] C. Klein, and J.-C. Saut, *Nonlinear dispersive equations. Inverse scattering and PDE methods*, under preparation (2021).
- [19] C. Klein, and J.-C. Saut, *Numerical study of blow-up and stability of solutions to generalized Kadomtsev-Petviashvili equations*, J. Nonlinear Science, 22, 5, (2012) 763–811.
- [20] Y. Kodama, and L. Williams, *The Deodhar decomposition of the Grassmannian and the regularity of KP solitons*, Advances in Mathematics, Vol. 244, 10 Sept. 2013, 979–1032.
- [21] Y. Kodama, and L. Williams, *KP solitons and total positivity for the Grassmannian*, Inventiones Mathematicae Vol. 198, 637–699 (2014).
- [22] Y. Kodama. *KP Solitons and the Grassmannians*. Combinatorics and Geometry of Two-Dimensional Wave Patterns. Springer Briefs in Mathematical Physics Vol. 22, Springer 2017.
- [23] B. G. Konopelchenko, *Introduction to Multidimensional Integrable Equations: The Inverse Spectral Transform in 2+1 dimensions*, 1992 Plenum Press, NY, 292 pp., ISBN 0-306-44220-5.
- [24] D. Lannes, *Consistency of the KP Approximation*, Proceedings of the 4th. international conference on dynamical systems and differential equations, May 24-27 (2002), Wilmington, NC, USA, 517–525.
- [25] D. Lannes, and J.-C. Saut, *Weakly transverse Boussinesq systems and the KP approximation*, Nonlinearity 19 (2006), 2853–2875.
- [26] A. J. Mendez, C. Muñoz, F. Poblete and J. C. Pozo, *On local energy decay for large solutions of the Zakharov-Kuznetsov equation*, Comm. Partial Differential Equations 46 (2021), no. 8, 1440–1487.
- [27] A. J. Mendez, C. Muñoz, F. Poblete and J. C. Pozo, *Long time asymptotics of large data in the Kadomtsev-Petviashvili models*, Nonlinearity 37 005017.
- [28] T. Mizumachi, *Stability of line solitons for the KP-II equation in \mathbb{R}^2* , Memoires of the AMS., vol 238, no. 1125, (2015).
- [29] T. Mizumachi, *Stability of line solitons for the KP-II equation in \mathbb{R}^2 . II.*, Proc. Roy. Soc. Edin. Sect. A, 143, (2018), 149–198.
- [30] T. Mizumachi, *Linear stability of elastic 2-line solitons for the KP-II equation*, Quart. Appl. Math. 82 (2024), no. 1, 115–226.
- [31] T. Mizumachi, and N. Tzvetkov, *Stability of the line soliton of the KP-II equation under periodic transverse perturbations*, Math. Annalen 352 (2012), 659–690.
- [32] L. Molinet, J.-C. Saut, and N. Tzvetkov, *Global well-posedness for the KP-II equation on the background of a non-localized solution*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 28 (2011), pp. 653–676.
- [33] T. Niizato, *Large time behavior for the generalized Kadomtsev-Petviashvili equations*, Diff. Eqns. and Appl. 3 (2) (2011), 299–308.
- [34] F. Rousset, and N. Tzvetkov, *Transverse instability of the line solitary water waves*, Invent. Math. 184 (2) (2011), 257–388.
- [35] F. Rousset, and N. Tzvetkov, *Transverse nonlinear instability for two-dimensional dispersive models*, Ann. IHP. Analyse Nonlinéaire, 26 (2009), 477–496.
- [36] L. Y. Sung, *Square integrability and uniqueness of the solutions of the Kadomtsev-Petviashvili-I equation*, Mathematical Physics, Analysis and Geometry 2 (1999), 1–24.

- [37] H. Takaoka and N. Tzvetkov, *On the local regularity of the Kadomtsev-Petviashvili-II equation*, Internat. Math. Res. Notices, (2001), pp. 77–114.
- [38] S. Ukai, *Local solutions of the Kadomtsev–Petviashvili equation*. J. Fac. Sci. Univ. Tokyo, Sect. A Math. 36, 193–209 (1989).
- [39] M. V. Wickerhauser, *Inverse scattering for the heat equation and evolutions in $(2+1)$ variables*, Comm. Math. Phys. 108 (1987), 67–89.
- [40] Derchyi Wu, *Stability of Kadomtsev Petviashvili multi line solitons*, <https://arxiv.org/abs/2205.07432v2>, pp. 1–68 (2022).
- [41] Derchyi Wu, *Stability of KdV solitons*, preprint arXiv:2401.15819v1 (2024).

INSTITUTO DE CIENCIAS FÍSICAS Y MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUSTRAL DE CHILE, VALDIVIA, CHILE.

Email address: `francisco.alegria@uach.cl`

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332-0160, USA

Email address: `gc@math.gatech.edu`

DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CENTRO DE MODELAMIENTO MATEMÁTICO (UMI 2807 CNRS), UNIVERSIDAD DE CHILE, CASILLA 170 CORREO 3, SANTIAGO, CHILE.

Email address: `cmunoz@dim.uchile.cl`

INSTITUTO DE CIENCIAS FÍSICAS Y MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD AUSTRAL DE CHILE, VALDIVIA, CHILE.

Email address: `felipe.poblete@uach.cl`

DEPARTAMENTO DE INGENIERÍA MATEMÁTICA, UNIVERSIDAD DE CHILE, CASILLA 170 CORREO 3, SANTIAGO, CHILE.

Email address: `btardy@dim.uchile.cl`