

Solutions of time fractional anomalous diffusion equations with coefficients depending on both time and space variables

Ganbileg Bat-Ochir, Khongorzul Dorjgotov, Uuganbayar Zunderiya

May 14, 2024

Abstract

We derive explicit solutions for time-fractional anomalous diffusion equations with diffusivity coefficients that depend on both space and time variables. These solutions are expressed in Fox-H and generalized Wright functions, which are commonly used in anomalous diffusion equations. Our study represents a significant advancement in our understanding of anomalous diffusion with potential applications in a wide range of fields.

1 Introduction

Anomalous diffusion has been the subject of extensive research in recent years, with numerous publications that address different aspects of this phenomenon. The growing number of applications that can be described by anomalous diffusivity has led to increased interest in this area of study. Until now, previous research has explored the dependence of diffusion coefficients on space, but no published results have provided explicit solutions when the diffusion coefficients depend on both space and time variables. This work builds on previous research [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14], which provided exact solutions for anomalous diffusion equations with a diffusion coefficient function that depended only on the space variable. We extend our previous work to include the time variable in the diffusion coefficient function. This development allows for the modeling of processes that were previously inaccessible.

In [15], the authors proposed that the following diffusion equation

$$\frac{\partial}{\partial t}P(x, t) = \frac{\partial}{\partial x} \left(K(x, t) \frac{\partial}{\partial x} P(x, t) \right),$$

where the diffusivity coefficient is taken as $K(x, t) = cx^{at^b}$, $a, b, c \in \mathbb{R}$, to describe diffusion processes in turbulent media.

In [4], authors proposed time fractional version of the equation with the coefficient K depends only on x , i.e. $K(x) = cx^a$, to describe anomalous diffusion with an asymptotic behaviour which is regarded as a general property of diffusion in fractal structure. Since it has been shown that the diffusivity coefficient should not depend only on spatial variable in [15], we consider the following generalized equation

$$\frac{\partial^\alpha}{\partial t^\alpha} u(x, t) = t^m \left(Ax^d \frac{\partial^2}{\partial x^2} u(x, t) + Bx^{d-1} \frac{\partial}{\partial x} u(x, t) + Cx^{d-2} u(x, t) \right), \quad x > 0, t > 0. \quad (1)$$

Here m is a nonnegative integer, d is a real number, A is a positive real number, B and C are real numbers.

Equation (1) can be considered as a time fractional generalization of equation given in [15] as well as a generalization of the equation studied in [4] considering the diffusivity coefficient depends both on time and space variables.

The solutions that will be presented in this work will be expressed in special functions, such as generalized Wright function and Fox-H functions. So we refer the readers who are not familiar with these special functions should consult with the literature [16, 17].

2 Main result

We recall the following special functions and their properties in preparation for introducing exact invariant solutions of Eq. (1). The Fox H-function (e.g. in [10, 16, 17]) is defined by means of the Mellin-Barnes type contour integral

$$H_{p,q}^{m,l} \left[z \left| \begin{matrix} (a_i, \alpha_i)_{1,p} \\ (b_j, \beta_j)_{1,q} \end{matrix} \right. \right] = \frac{1}{2\pi i} \int_L \frac{\prod_{j=1}^m \Gamma(b_j - \beta_j s) \prod_{i=1}^l \Gamma(1 - a_i + \alpha_i s)}{\prod_{i=l+1}^p \Gamma(a_i - \alpha_i s) \prod_{j=m+1}^q \Gamma(1 - b_j + \beta_j s)} z^s ds, \quad (2)$$

for $z \in \mathbb{C} \setminus \{0\}$, where $m, l, p, q \in \mathbb{N}_0 = \{0, 1, 2, \dots\}$, $0 \leq m \leq q$, $0 \leq l \leq p$, $(m, l) \neq (0, 0)$, $\alpha_i, \beta_j \in \mathbb{R}_+$, $a_i, b_j \in \mathbb{R}$ ($i = 1, \dots, p; j = 1, \dots, q$). Here L is a suitable contour from $\gamma - i\infty$ to $\gamma + i\infty$, where γ is a real number. The integral in (2) converges if the following conditions are met

$$\omega = \sum_{i=1}^l \alpha_i - \sum_{i=l+1}^p \alpha_i + \sum_{j=1}^m \beta_j - \sum_{j=m+1}^q \beta_j > 0 \text{ and } |\arg z| < \frac{\pi\omega}{2}.$$

The Fox H-function vanishes for large z because

$$H_{p,q}^{m,0}[z] \approx O \left(\exp \left(-\nu z^{\frac{1}{\nu}} \mu^{\frac{1}{\nu}} \right) z^{\frac{2\delta+1}{2\nu}} \right),$$

where $\mu = \prod_{i=1}^p \alpha_i \prod_{j=1}^q \beta_j^{-\beta_j}$, $\delta = \sum_{j=1}^q b_j - \sum_{i=1}^p a_i + \frac{p-q}{2}$ and $\nu = \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i > 0$. The generalized Wright function is defined (e.g. in [10]) as

$${}_p\Psi_q \left[z \left| \begin{matrix} (a_i, \alpha_i)_{1,p} \\ (b_j, \beta_j)_{1,q} \end{matrix} \right. \right] = \sum_{k=0}^{\infty} \frac{\prod_{i=1}^p \Gamma(a_i + \alpha_i k)}{\prod_{j=1}^q \Gamma(b_j + \beta_j k)} \frac{z^k}{k!}, \quad (3)$$

for $z \in \mathbb{C}$, $p, q \in \mathbb{N}_0$, $a_i, b_j \in \mathbb{C}$ and $\alpha_i, \beta_j \in \mathbb{R} \setminus \{0\}$ ($i = 1, \dots, p; j = 1, \dots, q$). If $\Delta = \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i > -1$ or $\Delta = -1$, then the series in (3) is absolutely convergent for $z \in \mathbb{C}$ or

$|z| < \prod_{i=1}^p |\alpha_i|^{-\alpha_i} \prod_{j=1}^q |\beta_j|^{\beta_j}$, respectively.

We know the following identities of H-function for $z > 0$

$$H_{p,q}^{m,l} \left[z \left| \begin{matrix} (A_i, \alpha_i)_{1,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] = H_{q,p}^{l,m} \left[\frac{1}{z} \left| \begin{matrix} (1 - B_j, \beta_j)_{1,q} \\ (1 - A_i, \alpha_i)_{1,p} \end{matrix} \right. \right], \quad (4)$$

$$H_{p,q}^{m,l} \left[z \left| \begin{matrix} (A_i, \alpha_i)_{1,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] = k H_{p,q}^{m,l} \left[z^k \left| \begin{matrix} (A_i, k\alpha_i)_{1,p} \\ (B_j, k\beta_j)_{1,q} \end{matrix} \right. \right] \text{ for } k > 0, \quad (5)$$

$$z^\sigma H_{p,q}^{m,l} \left[z \left| \begin{matrix} (A_i, \alpha_i)_{1,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] = H_{p,q}^{m,l} \left[z \left| \begin{matrix} (A_i + \sigma\alpha_i, \alpha_i)_{1,p} \\ (B_j + \sigma\beta_j, \beta_j)_{1,q} \end{matrix} \right. \right] \text{ for any } \sigma \in \mathbb{C}, \quad (6)$$

$$H_{p+1, q+r}^{m+r, l} \left[z \left| \begin{matrix} (A_i, \alpha_i)_{1,p}, (1, r) \\ \left(\frac{1}{r}, 1\right)_{1,r}, (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] = \frac{(2\pi)^{\frac{r-1}{2}}}{\sqrt{r}} H_{p,q}^{m,l} \left[r^r z \left| \begin{matrix} (A_i, \alpha_i)_{1,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] \text{ for } r \in \mathbb{Z}_+, l \leq p. \quad (7)$$

Moreover the following relations hold between the special functions. The Mittag-Leffler and Wright functions can be expressed by generalized Wright functions

$$E_{\alpha,\beta}(z) = {}_1\Psi_1 \left[z \left| \begin{matrix} (1, 1) \\ (\beta, \alpha) \end{matrix} \right. \right] \quad \text{and} \quad \Psi(z; \alpha, \beta) = \sum_{k=1}^{\infty} \frac{1}{\Gamma(\alpha k + \beta)} \frac{z^k}{k!} = {}_0\Psi_1 \left[z \left| \begin{matrix} - \\ (\beta, \alpha) \end{matrix} \right. \right]. \quad (8)$$

Let us state the main result of the study as the following theorem.

Theorem 1. 1. For $0 < \alpha < 2$ and $d \neq 2$, the equation given in (1) has the following solution

$$u(x, t) = c_1 x^a H_{1,m+2}^{m+2,0} \left[\frac{x^{2-d}}{A(d-2)^2(\alpha+m)^m t^{\alpha+m}} \left| \begin{matrix} (1, \alpha+m) \\ (-\frac{s_1}{\alpha+m}, 1), (-\frac{s_2}{\alpha+m}, 1), (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right]; \quad (9)$$

2. For $\alpha > 2$ and $d \neq 2$, the equation given in (1) has the following solution

$$u(x, t) = x^a \sum_{k=1}^{[\alpha]+1} c_k x^{\frac{(d-2)(\alpha-k)}{\alpha+m}} t^{\alpha-k} \times_{m+3}\Psi_1 \left[A(d-2)^2(\alpha+m)^m x^{d-2} t^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_1}{\alpha+m}, 1), (\frac{\alpha-k-s_2}{\alpha+m}, 1), (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1+\alpha-k, \alpha+m) \end{matrix} \right. \right],$$

3. For $d = 2$, the equation given in (1) has the following solution

$$u(x, t) = x^a \sum_{k=1}^{[\alpha]+1} c_k t^{\alpha-k} {}_{m+1}\Psi_1 \left[K t^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (\alpha-1, \alpha+m) \end{matrix} \right. \right],$$

where a is any real number, $K = Aa^2 - Aa + Ba + C$ and

$$s_{1,2} = \frac{\alpha+m}{2(2-d)} \left(\frac{B}{A} + 2a - 1 \pm \sqrt{\left(1 - \frac{B}{A}\right)^2 - \frac{4C}{A}} \right). \quad (10)$$

By setting $\alpha = 1$ in (9), we get

$$u(x, t) = c_1 x^a H_{1,m+2}^{m+2,0} \left[\frac{x^{2-d}}{A(d-2)^2(1+m)^m t^{1+m}} \left| \begin{matrix} (1, 1+m) \\ (-\frac{s_1}{1+m}, 1), (-\frac{s_2}{1+m}, 1), (\frac{j}{1+m}, 1)_{1,m} \end{matrix} \right. \right]. \quad (11)$$

If we take $a = \frac{1}{2} \left(2d - \frac{B}{A} - 3 \pm \sqrt{\left(1 - \frac{B}{A}\right)^2 - \frac{4C}{A}} \right)$, (11) becomes

$$u(x, t) = c_1 x^{\frac{1}{2} \left(2d - \frac{B}{A} - 3 \pm \sqrt{\left(1 - \frac{B}{A}\right)^2 - \frac{4C}{A}} \right)} \times H_{1,m+2}^{m+2,0} \left[\frac{x^{2-d}}{A(d-2)^2(1+m)^m t^{1+m}} \left| \begin{matrix} (1, 1+m) \\ (\frac{j}{1+m}, 1)_{1,1+m}, \left(\frac{d-2 \pm \sqrt{\left(1 - \frac{B}{A}\right)^2 - \frac{4C}{A}}}{d-2}, 1 \right) \end{matrix} \right. \right].$$

Then, we obtain the following corollary using (7) and (1.125) of [17] in above the solution.

Corollary 1. For $\alpha = 1$ and $d \neq 2$, the equation given in (1) has the following solution

$$u(x, t) = cx^{-\frac{1}{2}} \left(\frac{B}{A} - 1 \pm \sqrt{\left(1 - \frac{B}{A}\right)^2 - \frac{4C}{A}} \right) t^{-\frac{(1+m)}{d-2}} \left(d - 2 \pm \sqrt{\left(1 - \frac{B}{A}\right)^2 - \frac{4C}{A}} \right) EXP \left(-\frac{(1+m)x^{2-d}}{A(d-2)^2 t^{1+m}} \right),$$

where c is a constant.

The corollary shows that for a special case of (1), we have solutions expressed in terms of exponential functions.

Let us formulate the following known results as lemmas for the generalized Wright functions [10, 18, 20] and Fox H-functions [17, 19, 20] for the convenience to prove our main result.

Lemma 1. Let $\Delta = \sum_{j=1}^q B_j - \sum_{i=1}^p A_i > -1$. Then the following equalities hold for $\alpha \in \mathbb{R}_+$ and $a \in \mathbb{R}$.

(1) If $\beta_1 > 0$, $B_1 > 0$ and $A_1 = \alpha_1 = 1$, then we have

$$\begin{aligned} & \frac{d^\alpha}{dz^\alpha} \left(z^{B_1-1} {}_p\Psi_q \left[az^{\beta_1} \left| \begin{matrix} (1, 1), (A_i, \alpha_i)_{2,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] \right) \\ &= a^m z^{B_1+m\beta_1-1-\alpha} {}_p\Psi_q \left[az^{\beta_1} \left| \begin{matrix} (1, 1), (A_i + m\alpha_i, \alpha_i)_{2,p} \\ (B_1 + m\beta_1 - \alpha, \beta_1), (B_j + m\beta_j, \beta_j)_{2,q} \end{matrix} \right. \right], \end{aligned}$$

where m is the smallest non-negative integer such that $B_1 + m\beta_1 - \alpha - 1$ is not a negative integer.

(2) For $\sigma \in \mathbb{R} \setminus \{0\}$ and $R \in \mathbb{R}$, the following equality holds

$$\begin{aligned} & \left(\frac{1}{\alpha} z \frac{d}{dz} + R \right) \left(z^{\frac{A_1\sigma}{\alpha_1} - \alpha R} {}_p\Psi_q \left[az^\sigma \left| \begin{matrix} (A_i, \alpha_i)_{1,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] \right) \\ &= \frac{\sigma}{\alpha_1 \alpha} z^{\frac{A_1\sigma}{\alpha_1} - \alpha R} {}_p\Psi_q \left[az^\sigma \left| \begin{matrix} (A_1 + 1, \alpha_1), (A_i, \alpha_i)_{2,p} \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right]. \end{aligned}$$

Lemma 2. Let $\nu = \sum_{j=1}^q \beta_j - \sum_{i=1}^p \alpha_i > 0$, $\mu = \sum_{j=1}^m \beta_j - \sum_{j=m+1}^q \beta_j - \sum_{i=1}^p \alpha_i > 0$, $\alpha \in \mathbb{R}_+$ and $a \in \mathbb{R} \setminus \{0\}$. Then the following equalities hold.

(1) If $a > 0$, then

$$\begin{aligned} & \frac{d^\alpha}{dz^\alpha} H_{p,q}^{m,0} \left[az^{-\alpha_p} \left| \begin{matrix} (A_i, \alpha_i)_{1,p-1}, (1, \alpha_p) \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] \\ &= z^{-\alpha} H_{p,q}^{m,0} \left[az^{-\alpha_p} \left| \begin{matrix} (A_i, \alpha_i)_{1,p-1}, (1 - \alpha, \alpha_p) \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right], \quad z > 0. \end{aligned}$$

(2) If $m \geq 1$, then

$$\begin{aligned} & \left(\frac{\beta_1}{\alpha_p} z \frac{d}{dz} + B_1 \right) H_{p,q}^{m,0} \left[az^{-\alpha_p} \left| \begin{matrix} (A_i, \alpha_i)_{1,p-1}, (1, \alpha_p) \\ (B_j, \beta_j)_{1,q} \end{matrix} \right. \right] \\ &= H_{p,q}^{m,0} \left[az^{-\alpha_p} \left| \begin{matrix} (A_i, \alpha_i)_{1,p-1}, (1, \alpha_p) \\ (B_1 + 1, \beta_1), (B_j, \beta_j)_{2,q} \end{matrix} \right. \right]. \end{aligned}$$

3 Proof of Main Theorem

First, we will derive solutions to the following equations

$$\frac{d^\alpha}{dz^\alpha}y(z) = z^m \left(a_n z^n \frac{d^n}{dz^n}y + a_{n-1} z^{n-1} \frac{d^{n-1}}{dz^{n-1}}y + \cdots + a_1 z \frac{d}{dz}y + a_0 y \right), \quad z > 0, \quad (12)$$

where α and a_n ($i = 0, \dots, n$) are positive real numbers, a_1, a_2, \dots, a_{n-1} are real numbers, $m \in \mathbb{N}$. Here the fractional differentiation is defined in Riemann-Liouville manner. In [20], we derived exact solutions to the class of linear fractional differential equations of (12) when $m = 0$.

In order to obtain the solutions of the equation given in (12), we consider the following characteristic polynomial

$$\tilde{P}(s) = a_n \prod_{j=0}^{n-1} (s - j) + a_{n-1} \prod_{j=0}^{n-2} (s - j) + \cdots + a_1 s + a_0 \quad (13)$$

associated with the differential operator

$$P(y) = a_n z^n \frac{d^n}{dz^n}y + a_{n-1} z^{n-1} \frac{d^{n-1}}{dz^{n-1}}y + \cdots + a_1 z \frac{d}{dz}y + a_0 y.$$

Let s_1, \dots, s_n be the roots of the characteristic polynomial (13). Then we can rewrite the right hand side of (12) in factorized form as following

$$\begin{aligned} a_n z^n \frac{d^n}{dz^n}y + a_{n-1} z^{n-1} \frac{d^{n-1}}{dz^{n-1}}y + \cdots + a_1 z \frac{d}{dz}y + a_0 y \\ = a_n \left(z \frac{d}{dz} - s_n \right) \cdots \left(z \frac{d}{dz} - s_2 \right) \left(z \frac{d}{dz} - s_1 \right) y. \end{aligned}$$

To present the solutions of the equation (12), as a preparation we show that the following identity holds.

Lemma 3. Let $a \in \mathbb{R}_+$, $m \in \mathbb{N}$ and $b \in \mathbb{C}$. Then

$$\frac{1}{\Gamma(1 + ab + m)} \prod_{i=1}^m \Gamma\left(\frac{i}{a} + b + 1\right) = \frac{1}{a^m \Gamma(1 + ab)} \prod_{i=1}^m \Gamma\left(\frac{i}{a} + b\right). \quad (14)$$

Proof. By noticing the following identities

$$\prod_{i=1}^m \left(\frac{i}{a} + b\right) \Gamma\left(\frac{i}{a} + b\right) = \frac{1}{a^m} \prod_{i=1}^m (i + ab) \prod_{i=1}^m \Gamma\left(\frac{i}{a} + b\right) = \frac{\Gamma(1 + ab + m)}{a^m \Gamma(1 + ab)} \prod_{i=1}^m \Gamma\left(\frac{i}{a} + b\right).$$

Hence, the proof is completed. \square

Now we are ready to give the solutions of (12).

Proposition 1. The equation given in (12) has the following solutions.

1. For $\alpha < n$, the solution is expressed as

$$y(z) = c_1 H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n (\alpha + m)^{m+n}} \left| \begin{matrix} (1, \alpha + m) \\ \left(-\frac{s_j}{\alpha+m}, 1\right)_{1,n}, \left(\frac{j}{\alpha+m}, 1\right)_{1,m} \end{matrix} \right. \right]; \quad (15)$$

2. For $\alpha > n$, the solution is expressed as

$$y(z) = \sum_{k=1}^{[\alpha]+1} c_k z^{\alpha-k} {}_{m+n+1}\Psi_1 \left[a_n(\alpha+m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m}, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1+\alpha-k, \alpha+m) \end{matrix} \right. \right]. \quad (16)$$

Here s_1, \dots, s_n are the roots of the characteristic polynomial (13).

Proof. Because the convergence conditions of H-functions and generalized wright functions holds the functions in (15) and (16), it is sufficient to show that the functions satisfy the equation in (12). Firstly, we rewrite the right hand side of the equation (12) in the factorized form

$$\begin{aligned} & z^m \left(a_n z^n \frac{d^n}{dz^n} + a_{n-1} z^{n-1} \frac{d^{n-1}}{dz^{n-1}} + \dots + a_1 z \frac{d}{dz} + a_0 \right) \\ & \left(c_1 H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1, \alpha+m) \\ (-\frac{s_j}{\alpha+m}, 1)_{1,n}, (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right] \right) \\ & = c_1 a_n (\alpha+m)^n z^m \left(\frac{1}{\alpha+m} z \frac{d}{dz} - \frac{s_n}{\alpha+m} \right) \dots \left(\frac{1}{\alpha+m} z \frac{d}{dz} - \frac{s_1}{\alpha+m} \right) \\ & \left(H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1, \alpha+m) \\ (-\frac{s_j}{\alpha+m}, 1)_{1,n}, (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right] \right). \end{aligned} \quad (17)$$

From Lemma 2, we get the following identity for $k = 1, \dots, n$

$$\begin{aligned} & \left(\frac{1}{\alpha+m} z \frac{d}{dz} - \frac{s_k}{\alpha+m} \right) \left(H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1, \alpha+m) \\ (-\frac{s_j}{\alpha+m}, 1)_{1,n}, (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right] \right) \\ & = H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1, \alpha+m) \\ (-\frac{s_j}{\alpha+m}, 1)_{1,k-1}, (1 - \frac{s_k}{\alpha+m}, 1), (-\frac{s_j}{\alpha+m}, 1)_{k+1,n}, (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right]. \end{aligned}$$

Then applying the above identity in (17), the right hand side of the equation becomes

$$c_1 a_n z^m (\alpha+m)^n H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1, \alpha+m) \\ (1 - \frac{s_j}{\alpha+m}, 1)_{1,n}, (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right].$$

Now, let us begin the calculation of left hand side of the equation. By applying the Lemma 2, we get

$$\frac{d^\alpha}{dz^\alpha} y = c_1 z^{-\alpha} H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1-\alpha, \alpha+m) \\ (-\frac{s_j}{\alpha+m}, 1)_{1,n}, (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right].$$

Then applying (6), the left hand side is

$$c_1 a_n z^m (\alpha+n)^{m+n} H_{1,m+n}^{m+n,0} \left[\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \left| \begin{matrix} (1+m, \alpha+m) \\ (1 - \frac{s_j}{\alpha+m}, 1)_{1,n}, (1 + \frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right].$$

Hence, the left hand side becomes

$$\frac{d^\alpha}{dz^\alpha} y = c_1 a_n z^m (\alpha+n)^{m+n} \frac{1}{2\pi i} \int_L \frac{\prod_{j=1}^n \Gamma(1 - \frac{s_j}{\alpha+m} - s) \prod_{j=1}^m \Gamma(1 + \frac{j}{\alpha+m} - s)}{\Gamma(1+m - (\alpha+m)s)} \cdot \left(\frac{z^{-(\alpha+m)}}{a_n(\alpha+m)^{m+n}} \right)^s ds$$

By substituting $a = \alpha + m$, $b = -s$ in Lemma 3, we see that

$$\frac{\prod_{j=1}^m \Gamma(1 + \frac{j}{\alpha+m} - s)}{\Gamma(1+m - (\alpha+m)s)} = \frac{1}{(\alpha+m)^m} \cdot \frac{\prod_{j=1}^m \Gamma(\frac{j}{\alpha+m} - s)}{\Gamma(1 - (\alpha+m)s)},$$

which concludes the right and left hand sides of the equations are equal.

The second assertion of the theorem consists of $[\alpha] + 1$ summands. Since the equation is linear, it is sufficient to show that it holds for single k th summand. For the fixed summand, we will take the similar steps as in the previous case. From Lemma 2, we get the following identity for $l = 1, \dots, n$

$$\left(\frac{1}{\alpha} z \frac{d}{dz} - \frac{s_l}{\alpha} \right) \left(z^{\alpha-k} {}_{m+n+1}\Psi_1 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m}, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1 + \alpha - k, \alpha + m) \end{matrix} \right. \right] \right) = \frac{\alpha + m}{\alpha} z^{\alpha-k} \\ \times {}_{m+n+1}\Psi_1 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m}, 1)_{1,l-1}, (\frac{\alpha-k-s_l}{\alpha+m} + 1, 1), (\frac{\alpha-k-s_i}{\alpha+m}, 1)_{l+1,n}, (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1 + \alpha - k, \alpha + m) \end{matrix} \right. \right],$$

the right hand side of the equation becomes

$$a_n \alpha^n z^m \left(\frac{1}{\alpha} z \frac{d}{dz} - \frac{s_n}{\alpha} \right) \cdots \left(\frac{1}{\alpha} z \frac{d}{dz} - \frac{s_2}{\alpha} \right) \left(\frac{1}{\alpha} z \frac{d}{dz} - \frac{s_1}{\alpha} \right) \\ \left(z^{\alpha-k} {}_{m+n+1}\Psi_1 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m}, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1 + \alpha - k, \alpha + m) \end{matrix} \right. \right] \right) \\ = a_n (\alpha + m)^n z^{\alpha+m-k} {}_{m+n+1}\Psi_1 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m} + 1, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1 + \alpha - k, \alpha + m) \end{matrix} \right. \right].$$

The left hand side of the equation is calculated as following

$$\frac{d^\alpha}{dz^\alpha} z^{\alpha-k} {}_{m+n+1}\Psi_1 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m}, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1 + \alpha - k, \alpha + m) \end{matrix} \right. \right] \\ = b_0 a_n (\alpha + m)^{m+n} z^{\alpha+m-k} \\ \times {}_{m+n+2}\Psi_2 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (1, 1), (\frac{\alpha-k-s_i}{\alpha+m} + 1, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m} + 1, 1)_{1,m}, (2, 1) \\ (2, 1), (1 + \alpha - k + m, \alpha + m) \end{matrix} \right. \right] \\ = b_0 a_n (\alpha + m)^{m+n} z^{\alpha+m-k} \\ \times {}_{m+n+1}\Psi_1 \left[a_n (\alpha + m)^{m+n} z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_i}{\alpha+m} + 1, 1)_{1,n}, (\frac{\alpha-k+i}{\alpha+m} + 1, 1)_{1,m}, (1, 1) \\ (1 + \alpha - k + m, \alpha + m) \end{matrix} \right. \right] \\ = b_0 a_n (\alpha + m)^{m+n} z^{\alpha+m-k} \\ \times \sum_{j=0}^{\infty} \frac{\prod_{i=1}^n \Gamma \left(\frac{\alpha-k-s_i}{\alpha+m} + 1 + j \right) \prod_{i=1}^m \Gamma \left(\frac{\alpha-k+i}{\alpha+m} + 1 + j \right) \Gamma(1 + j)}{\Gamma(1 + \alpha - k + m + (\alpha + m)j)} \cdot \frac{(a_n (\alpha + m)^{m+n} z^{\alpha+m})^j}{j!}$$

which equals to the right hand side of the equation via Lemma 3 with $a = \alpha + m$, $b = \frac{\alpha-k}{\alpha+m} + j$. \square

So, we have formulated exact solutions to the fractional differential equations given in (12). In the following, we will demonstrate how to apply the results to (1).

By applying the substitution

$$u = x^a \varphi(x^{\frac{d-2}{\alpha+m}} t), \quad x^{\frac{d-2}{\alpha+m}} t = z, \quad (a \in \mathbb{R}) \quad (18)$$

in (1), we have

$$\frac{d^\alpha}{dz^\alpha} \varphi = z^m \left[\frac{A(d-2)^2}{(\alpha+m)^2} z^2 \varphi_{zz} + \frac{d-2}{\alpha+m} \left(\frac{A(d-2)}{\alpha+m} + B \right. \right. \\ \left. \left. + 2Aa - A \right) z \varphi_z + (Aa^2 - Aa + Ba + C) \varphi \right]. \quad (19)$$

Let's take a look at the following characteristic equation of (19)

$$\frac{A(d-2)^2}{(\alpha+m)^2}s(s-1) + \frac{d-2}{\alpha+m} \left(\frac{A(d-2)}{\alpha+m} + B + A(2a-1) \right) s + Aa(a-1) + Ba + AC = 0$$

where $d \neq 2$. Finally, according to Proposition 1 we obtain the following solutions to equation (19)

1. For $0 < \alpha < 2$,

$$\varphi(z) = c_1 H_{1,m+2}^{m+2,0} \left[\frac{z^{-(\alpha+m)}}{A(d-2)^2(\alpha+m)^m} \left| \begin{matrix} (1, \alpha+m) \\ (-\frac{s_1}{\alpha+m}, 1), (-\frac{s_2}{\alpha+m}, 1), (\frac{j}{\alpha+m}, 1)_{1,m} \end{matrix} \right. \right], \quad (20)$$

2. For $\alpha > 2$,

$$\varphi(z) = \sum_{k=1}^{[\alpha]+1} c_k z^{\alpha-k} {}_{m+3}\Psi_1 \left[A(d-2)^2(\alpha+m)^m z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k-s_1}{\alpha+m}, 1), (\frac{\alpha-k-s_2}{\alpha+m}, 1), (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1+\alpha-k, \alpha+m) \end{matrix} \right. \right]. \quad (21)$$

The case $d = 2$, we get the following solutions to equation (19)

$$\varphi(z) = \sum_{k=1}^{[\alpha]+1} c_k z^{\alpha-k} {}_{m+1}\Psi_1 \left[(Aa^2 - Aa + Ba + C)(\alpha+m)^m z^{\alpha+m} \left| \begin{matrix} (\frac{\alpha-k+i}{\alpha+m}, 1)_{1,m}, (1, 1) \\ (1+\alpha-k, \alpha+m) \end{matrix} \right. \right]. \quad (22)$$

Under the substitution (18) we get our desired results in Theorem 1.

Acknowledgments

This work was partially supported by the National University of Mongolia (Grant No.P2018-3584) and by the Mongolian Foundation for Science and Technology (Grant No.SHUTBIKHKHZG-2022/164).

References

- [1] K.B. Oldham, J. Spanier, *The Fractional Calculus*, Academic Press, New York-London (1974).
- [2] W. Wyss, *The fractional diffusion equation*, J. Math. Phys. 27 (1986), no. 11, pp. 2782-2785.
- [3] W. R. Schneider, W. Wyss, *Fractional diffusion and wave equations*, J. Math. Phys. 30 (1989), no. 1, pp. 134-144.
- [4] R. Metzler, W. G. Glockle, T. F. Nonnenmacher *Fractional model equation for anomalous diffusion*, Physica A 211 (1994) 13-24.
- [5] F. Mainardi, *The fundamental solutions for the fractional diffusion-wave equation*, Appl. Math. Lett. 9 (1996), no. 6, 23-28.
- [6] E. Buckwar, Yu. Luchko *Invariance of a partial differential equation of fractional order under Lie group of scaling transformations*, J. Math. Anal. Appl. 227 (1998), 81-97.

- [7] I. Podlubny, *Fractional Differential Equations*, Academic Press (1999).
- [8] R. Metzler, J. Klafter, *The random walk's guide to anomalous diffusion: A fractional dynamics approach*, Physics Report 339, No 1 (2000), 1–77.
- [9] F. Mainardi, G. Pagnini, The Wright functions as solutions of the time-fractional diffusion equation, Appl. Math. Comput. 141 (2003), no. 1, 51–62.
- [10] A. A. Kilbas, *Fractional calculus of the generalized Wright function*, Fract. Calc. Appl. Anal. 8 (2) (2005), 113—126.
- [11] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier Science B.V., Amsterdam (2006).
- [12] F. Mainardi, G. Pagnini, The role of Fox-Wright functions in fractional sub-diffusion of distributed order, J. Comput. Appl. Math. 207 (2007), no. 2, 245–257.
- [13] F. Mainardi, A. Mura, G. Pagnini, The M-Wright function in time-fractional diffusion processes: a tutorial survey, Int. J. Differ. Equ. (2010) 104505, 29pp.
- [14] Ed. V. Tarasov, *Handook of Fractional Calculus with Applications Volumes 4 and 5: Applications in Physics*, De Gruyter, Germany (2019).
- [15] H.G.E. Hentschel, I. Procaccia *Relative diffusion in turbulent media: The fractal dimension of clouds*, Physical Review A 29 (3), 1984, 1461–1470.
- [16] A. M. Mathai, R. K. Saxena, *The H-Function with Applications in Statistics and Other Disciplines*, John Wiley, New York-London-Sidney (1978).
- [17] A. M. Mathai, R. K. Saxena, H. J. Haubold, *The H-Function, Theory and Applications*, Springer, New York (2010).
- [18] V. Kiryakova, *Fractional calculus operators of special functions? The result is well predictable!*, Chaos, Solitons and Fractals 102, 2017, 2–15.
- [19] W. G. Glockle, T. F. Nonnenmacher, *Fox function representation of Non-Debye relaxation processes*, J. Statist. Phys. 71 (3/4), 1993, 741–757.
- [20] Kh. Dorjgotov, H. Ochiai, U. Zunderiya *On solutions of linear fractional differential equations and systems thereof*, Fract. Calc. Appl. Anal. No22,(2019),479–494.
- [21] E. Awad, R. Metzler *Closed form multi-dimensional solution and asymptotic behaviours for subdiffusive processes with crossovers: II. Accelerating case*, J. Phys. A: Math. Theor. **55** (2022) 205003.
- [22] J. Klafter, A. Blumen, M. F. Shlesinger *Stochastic pathway to anomalous diffusion*, Phys. Rev. A 35 (7), 1987, 3081–3085.