# An Infra-Red Fixed Point in QCD with a Dilaton in Nuclear Dynamics

Long-Qi Shao<sup>1, 2, 3, \*</sup> and Mannque Rho<sup>4, †</sup>

<sup>1</sup>School of Fundamental Physics and Mathematical Sciences,

Hangzhou Institute for Advanced Study, UCAS, Hangzhou, 310024, China

<sup>2</sup>Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China

<sup>3</sup>University of Chinese Academy of Sciences, Beijing 100049, China

<sup>4</sup> Université Paris-Saclay, CNRS, CEA, Institut de Physique Théorique, 91191, Gif-sur-Yvette, France

(Dated: May 31, 2024)

The possible consequence of an IR fixed point in QCD for  $N_f = 2, 3$  in nuclear matter is discussed. It is shown in terms of d(ilaton)- $\chi$  effective field theory (d $\chi$ EFT) incorporated in a generalized effective field theory (G*n*EFT) implemented with hidden local symmetry and hidden scale symmetry that the recently measured superallowed Gamow-Teller transition in the doubly-magic-shell nucleus <sup>100</sup>Sn indicates a surprisingly large anomaly-induced quenching (AIQ) of  $g_A$  from the free-space value of 1.276 to  $\approx 0.8$ . Combined with the quenching expected from strong nuclear correlations, the effective coupling in nuclei,  $g_A^{\text{eff}}$  would come to  $\sim 1/2$ . If this result were reconfirmed, it would impact not only nuclear structure and dense compact-star matter – where  $g_A$  figures in  $\pi$ -N coupling via the Goldberger-Treiman relation – but also in search for physics Beyond the Standard Model (BSM), e.g.  $0\nu\beta\beta$  decay, where 4th power of  $g_A$  figures.

# I. INTRODUCTION

It was argued [1, 2] that the "quenched  $g_A^{\text{eff}} \approx 1$ " with the quenching factor (usually denoted in the literature as)  $q \equiv g_A^{\text{eff}}/g_A \approx 0.78$  (where  $g_A = 1.276$  is the free-space value) observed in Gamow-Teller transitions in light nuclei [3] fails to account for the large reduction of transition strength in the doubly magic nucleus <sup>100</sup>Sn making superallowed Gamow-Teller transition recently measured in RIKEN [4]. It would require a big quenching factor  $q \sim$ 1/2 compared to what's observed in light nuclei. This result would raise havoc not only for nuclear physics but also particle physics.

Shell-model analyses have rather persuasively suggested that for light nuclei, say, with  $A \leq 21$ , if the nuclear correlation is fully considered with proper wave function and effective operator, no *further* quenching is needed [5]. However in heavier nuclei, higher-order correlations involving large configuration space were difficult to put under control, making it difficult to reliably address the RIKEN result. There are two major issues raised in this conundrum, one in nuclear manybody problem touching on the structure of dense neutron stars and the other a hidden symmetry in QCD that gets un-hidden in nuclear strong correlations. It can also have serious impact on the searches for physics beyond the SM. Now the question that arises is: Does the quenching involve a "fundamental" phenomenon in QCD or just mundane nuclear effect? It turns out fortunately that for superallowed Gamow-Teller transitions in doubly-closed magic shell-nuclei, there is a possibility of closely "mapping" what happens in heavy nuclei in shellmodel to what can be reliably treated in Landau-Fermiliquid (LFL) theory [6] in the Fermi-liquid fixed-point (FLFP) approximation [7] to address this issue.

In addressing the problem concerned, one cannot avoid an inherent fuzziness in what is "fundamental" and what is not in the observable quantities. We will try to give as precise and concise a definition as feasible of what is meant by "fundamental." This issue arises because we will be working in the framework of an effective field theory (EFT).

The nuclear EFT we will be exploiting is defined with the energy-momentum scale set by the "chiral scale"  $\Lambda_{\chi} \sim 4\pi f_{\pi}$  at which the QCD degrees of freedom are to be integrated out. For the processes concerned in nuclear physics, the scale can in practice be brought down for nuclear physics to the mass scale of the lowest-lying vector mesons  $(\rho, \omega)$ . We further need to implement the scalar dilaton as the pseudo-Goldstone boson for broken scale symmetry. Although the existence of an IR fixed in QCD with  $N_f = 2$  or 3 involved in nuclear processes remains still in controversy, we will be arguing it should equally figure in dense nuclear interactions [8–10] as well as particle physics involving scalar degrees of freedom in QCD [11, 12]. One of the early role of the dilaton was recognized in the BR scaling [10]. It has figured importantly in nuclear many-body approaches in the form of generalized nuclear effective field theory (referred to as GnEFT [1, 2, 13]) involving both hidden local (HLS) [14] and hidden scale (HSS) symmetries.

We give a brief summary of the principal results of this paper: From a recent, startling, new argument put forward by Zwicky [15] that there can exist an IR fixed point for 2 or 3 flavor QCD with the derivative of  $\beta$  function with respect to the QCD coupling constant  $\alpha_s$  (at the leading scale symmetry order),  $\beta'_*$ , is zero, we find from the most recent RIKEN experiment that the anomalyinduced quenching (AIQ) of  $g_A$  can be surprisingly big, making the value of  $g_A$  drop from the free-space value of 1.276 to ~ 0.8. If the RIKEN result were reconfirmed

<sup>\*</sup> shaolongqi22@mails.ucas.ac.cn

<sup>&</sup>lt;sup>†</sup> mannque.rho@ipht.fr

by new measurements with accurately controlled theoretical inputs, then it would bring a totally new development not only in nuclear theory and nuclear astrophysics but also in searches for BSM. Since the axial coupling  $g_A$  is connected to the pion-nucleon coupling in nuclear medium via the Goldberger-Treiman relation, the strong AIQ would basically revamp nuclear dynamics that is controlled by pion-nuclear interactions, that is, the nuclear physics anchored on chiral dynamics, the currently widely accepted paradigm in nuclear physics community.

### II. THE GnEFT

We begin with the EFT framework we are working in. The concept of dilaton we are interested in has appeared in various different contexts in the literature with different definitions and different goals. It is out of range of our expertise to give appropriate reviews here. We eschew giving them. Among them, we adopt the "genuine dilaton (GD)" scheme proposed by Crewther [11], the characteristic of which is shared to certain extent with the "conformal dilaton" phase in QCD (CD-QCD) developed by Zwicky [15]. There are some differences between them but they do not affect in our scheme exploiting the notion of emergence of the scale symmetry [16]. The scale symmetry that figures in GD (and also in CD-QCD) is hidden, emerging only in deep IR region. The process we are concerned with, as our work indicates, takes place not far from – though not on – the IR fixed point, so the strong controversy in the field does not seriously affect our reasoning that figures in nuclear dynamics. Unless otherwise noted, we will follow the notations of the GD framework. The part with which we are dealing, we think, overlaps with the CD-QCD scheme.

In the Lagrangian constructed by Crewther and Tunstall [11, 12], the scale and chiral symmetry are treated on the same footing. The chiral-scale Lagrangian is of the form

$$\mathcal{L}_{d\chi EFT} =: \mathcal{L}_{inv}^{d=4} + \mathcal{L}_{anom}^{d>4} + \mathcal{L}_{mass}^{d<4},$$
(1)

where d is the scaling dimension. For  $\mathcal{L}_{anom}^{d>4}$ ,  $d = 4 + \gamma_{G^2}(\alpha_s)$  with  $\alpha_s$  the QCD coupling constant and  $\gamma_{G^2}(\alpha_s)$  the anomalous dimension of the gluonic operator  $G_{\mu\nu}G^{\mu\nu}$ . For  $\mathcal{L}_{mass}^{d<4}$ ,  $d = 3 - \gamma_m(\alpha_s)$  where  $\gamma_m(\alpha_s)$  is the anomalous dimension of the bilinear quark operator  $\bar{q}q$ . It is assumed that we are near the IR fixed point in the low-energy and density regime of QCD,  $\alpha_s \leq \alpha_{\rm IR}$  with  $\alpha_{\rm IR}$  the QCD coupling constant at the IR fixed point, so the anomalous dimension has the expansion with respect to  $\delta_{\alpha s} \equiv O(\alpha_s - \alpha_{\rm IR})$ .

$$\gamma_{G^2}(\alpha_s) \equiv \beta'(\alpha_s) - \beta(\alpha_s)/\alpha_s = \beta'_* + O(\delta_{\alpha s}).$$
 (2)

with  $\beta'_*$  the derivative of  $\beta$  function at the IR fixed point.

Ignoring higher order terms in  $\delta_{\alpha s}$ , we have

$$\mathcal{L}_{\text{inv}}^{d=4} = \{c_1 \mathcal{K} + c_2 \mathcal{K}_{\sigma} + c_3 (\chi/f_{\sigma})^2\} (\chi/f_{\sigma})^2,$$
  

$$\mathcal{L}_{\text{anom}}^{d>4} = \{(1-c_1)\mathcal{K} + (1-c_2)\mathcal{K}_{\sigma} + c_4 (\chi/f_{\sigma})^2\} (\chi/f_{\sigma})^{2+\beta'_*}$$
  

$$\mathcal{L}_{\text{mass}}^{d<4} = \text{Tr}(MU^{\dagger} + UM^{\dagger}) (\chi/f_{\sigma})^{3-\gamma_m}.$$
(3)

Here

$$\mathcal{K} = \frac{1}{4} f_{\pi}^2 \text{Tr}(\partial_{\mu} U \partial^{\mu} U^{\dagger}) \quad \text{and} \quad \mathcal{K}_{\sigma} = \frac{1}{2} \partial_{\mu} \sigma \partial^{\mu} \sigma \qquad (4)$$

and  $U = exp(i\pi/f_{\pi})$  is the nonlinear realization of pion field,  $\chi/f_{\sigma} = exp(\sigma/f_{\sigma})$  is the dilaton field,  $c_i$  for i =1,2,3,4 are parameters and M is quark mass matrix. Two counting schemes are involved: the scale and chiral power counting in the Lagrangian. Near the IR fixed point,  $c_{1,2} = 1 + O(M), c_{3,4} = O(M)$  where  $O(M) \sim$  $O(m_{\pi}^2) \sim O(p^2)$  in the chiral power counting. When approaching the IR fixed point,  $M \to 0, c_{1,2} \to 1, c_{3,4} \to$ 0 such that both the anomalous dimension term and the mass term in Eq. (3) vanish.

In GnEFT [13], the many-body problem in nuclear matter is handled by a renormalization-group (RG) approach to interacting fermions on the Fermi sphere [7]. Implemented with the hidden local symmetric [14] and hidden scale-invariant fields, the GnEFT Lagrangian renders the mean field theory with the suitable BR scaling parameters equal to what corresponds to the Landau-Fermi-liquid fixed point theory of many-nucleon systems [6, 21]. With the nucleons put on a Fermi sphere, the LFL fixed point approximation corresponds to taking  $1/\bar{N}$  to zero where  $\bar{N} = k_F/(\Lambda_{fs} - k_F)$  with  $\Lambda_{fs}$ the cut-off on top of the Fermi-surface measured with respect to the origin and  $k_F$  the Fermi momentum. This approach becomes more reliable as density increases. Being a bona-fide EFT, one can do higher-order corrections in  $1/\bar{N}$  in what's known as " $V_{\text{lowK}}$  RG-expansion." For the  $g_A$  problem explicit  $1/\bar{N}$  corrections are found to be unnecessary. This makes higher-order calculations in the Fermi-liquid approach much simpler than doing the higher-order calculation in nuclear matter with  $d\chi EFT$ with many undetermined parameters.

Let us briefly describe the leading scale-chiral order axial current involved in the baryonic sector  $J_{5\mu}$  coupled to the weak external field  $\mathcal{W}^{\mu}$ . The weak Lagrangian is

$$\mathcal{L}_{weak} = J^a_{5\mu} \mathcal{W}^{a_\mu} \tag{5}$$

with

$$J_{5\mu}^{\pm} = Q_{ssb}(\chi) g_A \bar{\psi} \tau^{\pm} \gamma_{\mu} \gamma_5 \psi \tag{6}$$

where

$$Q_{ssb}(\chi) = c_A + (1 - c_A) \left(\frac{\chi}{f_\sigma}\right)^{\beta'_*} \tag{7}$$

stands for the anomaly-induced effect inherited from QCD. Here  $c_A$  is an undetermined parameter. Apart from  $Q_{ssb}$  in (6),  $\mathcal{L}_{weak}$  is scale-invariant. Deviation from

possible scale-invariance resides in (7). In GnEFT with the "vacuum" given by the medium with the baryon density n, we expand  $\chi = \langle \chi \rangle + \chi'$  with  $\chi'$  the fluctuation dilaton field and  $f_{\sigma} = \langle \chi \rangle_{n=0}$  in the medium-free space n = 0. As it stands, the exponent  $\beta'_*$  is O(1) in the scale-chiral counting but as discussed below, it can vanish, so one would have to consider the O(M) corrections reflecting the location of  $\alpha_s$  relative to  $\alpha_{\rm IR}$  which could be non-negligible. If we ignore the fluctuating dilatonfield contributions that enter at higher loop orders, we have  $Q_{ssb}(\langle \chi \rangle_{n=0}) = 1$  at zero density, so there is no scale symmetry breaking effect at the leading order on  $g_A$  in the matter-free vacuum. One can therefore say that the anomaly-induced effect, hidden in the matterfree vacuum, can be revealed primarily by the presence of baryonic matter,

Going into finite density,

$$\langle \chi \rangle^* / f_\sigma = f_\sigma^* / f_\sigma = \Phi(n)$$
 (8)

can be considered to be an "order parameter" that characterizes the vacuum structure modified by medium, the uppercase \* standing for in-medium quantity. The quantity  $\Phi(n)$  is referred to in the literature as "BR scaling factor." It governs how parameters in the  $d\chi EFT$  scale with density.

In the order we are considering, the axial current we are dealing with is given by

$$J_{5\mu}^{\pm} = q_{ssb} g_A \bar{\psi} \tau^{\pm} \gamma_{\mu} \gamma_5 \psi \tag{9}$$

with

$$q_{ssb} = c_A + (1 - c_A)\Phi^{\beta'_*}.$$
 (10)

For simplicity, we replaced the  $\Phi(n)$  with  $\Phi$ .  $q_{ssb}$  is a density-dependent factor that multiplies the coupling constant  $g_A$  in nuclear axial processes, hence represents a "fundamental renormalization" of the coupling constant  $g_A \to g_A^{inherit} \equiv q_{ssb}g_A$  inherited from QCD which gets manifested importantly in medium, to be distinguished from what's given in standard many-body nuclear correlations. It is worth noting that the effect of  $\beta'_*$  which is hidden in the vacuum is " exposed" in finite density in the axial current for  $\Phi \neq 1$ . The physical transition matrix element of the current (9) in nuclei will then be the *full* nuclear matrix element  $M_{\text{nucl}}$  of the operator  $j_5^{\text{nucl}} = \bar{\psi} \tau^{\pm} \gamma_{\mu} \gamma_5 \psi$  multiplied by the axial constant  $g_A^{inherit}$ . These two quantities are not entirely separate from each other because the BR scaling  $\Phi$  must figure in both  $M_{\text{nucl}}$  and the effective axial constant. Now the question raised is how it figures in  $g_A^{\rm eff}$  and  $M_{\rm nucl},$  such as does it depend on density interplaying in both etc.? It turns out fortunately, as we will show, that there is very little interplay between the two.

There are two quantities to be considered. The first is the full nuclear correlations that we refer to as "nuclear effect" (NE in short) and the second, the main issue in this paper, is the "fundamental quenching factor"  $q_{ssb} - 1 \neq 0$  referred to as "AIQ" (anomaly-induced quenching). Now in order to minimize the possible overlap between NE and AIQ we focus on the superallowed Gamow-Teller transition involving zero momentum transfer q = 0 and zero energy transfer  $\omega = 0$ . The strategy adopted [2] is to map the "Extreme Single Particle (shell) Model" (ESPM for short) in doubly magic closed shell nuclei to "Fermi-liquid fixed-Point" (FLFP) approximation [7] at  $(q, \omega) \rightarrow (0, 0)$  with  $q/\omega \rightarrow 0$  on the Fermi surface. The FLFP approximation is best applicable for a quasi-particle making the superallowed transition on the Fermi surface with loop contributions suppressed in nuclear matter. The resulting full GT matrix element is given by

$$M^{\rm L} = \frac{1}{2} q^L_{snc} g_A \langle \sigma \tau \rangle \tag{11}$$

where  $\langle \sigma \tau \rangle$  is the single quasi-nucleon matrix element of the GT operator. The factor  $q_{snc}^L$  given in (13) below, with the subscript *snc* representing strong nuclear correlation, is the factor that captures the *total* matrix element in the EFT adopted. The product

$$g_A^L = q_{snc}^L g_A \tag{12}$$

is then LFL fixed-point prediction for the singlequasiparticle coupling constant that captures the complete nuclear correlation. It is related to the  $g_A^{\text{eff}}$  mentioned in Introduction.

We now turn to the ESPM. For this we take the superallowed GT transition in <sup>100</sup>Sn nucleus which has the proton and neutron shells completely filled at 50/50 magic shells. The transition involved in the ESPM is the pure superallowed transition of a proton (denoted  $\pi$ )  $\pi g_{9/2}$ in the completely filled orbital to a neutron ( $\nu$ ) in the empty spin-orbit partner  $\nu g_{7/2}$  orbital of <sup>100</sup>In. This offers the simplest possible structure of the daughter state that is of a pure  $\nu g_{7/2}$  particle- $\pi g_{9/2}$  hole state to which the ESPM can be applied. Now if one assumes that the final (daughter) state reached in the GT process has ignorable mixing with other particle-hole states, then the mapping from the FLFP approximation in the Fermiliquid for  $1/\bar{N} \rightarrow 0$  to the ESPM would be as exact as feasible.

# III. AIQ: OBSERVATION AND PREDICTION

### A. Experiments

Here we look at what Nature says in the <sup>100</sup>Sn transition. Equation (10) for  $q_{ssb}$  involves two unknowns,  $c_A$ and  $\beta'_*$ . Neither is available by lattice QCD or experimental. Therefore an unambiguous prediction with the formula (10) is not feasible at present. To proceed, we need to resort to experimental decay rates to arrive at the nuclear matrix element  $M_{nucl}$  so as to get  $q_{snc}$ .

For this, an *ab initio* no-core shell-model calculation that takes into as full an account of nuclear correlations as feasible would be the best one can hope for. Up to date, one crucially important theoretical has not been properly accounted for. For accurately calculating superallowed GT transitions in heavy nuclei, the nuclear tensor force with its strength decreasing with density following the BR scaling will play an essential role [22]<sup>1</sup>. However up to date there are no such calculations available in the literature [23]. Here the ESPM-Landau FLFP mapping enters giving

$$q_{snc}^L = (1 - \frac{1}{3}\Phi \tilde{F}_1^{\pi})^{-2} \simeq 1/g_A \simeq 0.78.$$
 (13)

It turns out that this quantity is highly insensitive to density, so it can be applied to heavy nuclei as well as to nuclear matter. It was given numerically in [1, 6]. The robustness of the FLFP approximation for (13) reflected in the EFTs used in [1, 6] has been further confirmed by the nonlinear bosonization approach with the coadjoint orbit method [25]. The correction to (13) is found to come only at  $O(10^{-4})$ . Note that this is more or less what's found in light nuclei for  $g_{\rm eff}^{\rm eff}$  which indicates that with no AIQ effect there is little dependence on density [3].

Let's now look at the experiments available in <sup>100</sup>Sn where the mapping between shell-model and Landau Fermi-liquid model could be made. There are two experiments, one from GSI [26] and another from RIKEN [4].

• **GSI**: In this experiment, the transition zeroing-in on ~ 95% of the final daughter state of the pure  $(\nu g_{7/2})$ particle- $(\pi g_{9/2})$ hole configuration has been reported. The resulting  $q_{snc}^{GSI}$  comes out close to  $q_{snc}^{L}$  [2] leading to

$$q_{ssb}^{GSI} \approx (0.9 - 1.0).$$
 (14)

A detailed account of the mixing in the final state involving theoretical inputs seems to give a somewhat smaller  $q_{ssb}$  but it is not clear how reliable the mixing can be estimated. We choose not to rely on this analysis. Modulo the  $\sim 5\%$  uncertainty, however, it seems safe to conclude  $q_{ssb}^{GSI}$ given by (14) indicates no AIQ. This gives then what one might call "pure quasi-nucleon constant"  $g_A^{pqn} = q_{snc} \times g_A = 0.78 \times 1.276 = 1$ . This is equal to  $g_A^L$  (12) predicted by FLFP. It is worth noting that one obtains the same result as what is called "dilaton-limit fixed-point" (DLFP) [27] but at a much higher density in GnEFT where an intricate interplay between the attraction due to the dilaton exchange and the  $\omega$  repulsion plays a crucial role [9, 28].

• **RIKEN**: The more recent measurement at RIKEN for the same transition, proclaimed as improved

over the GSI result, comes out to be drastically different and points to a major AIQ effect. It is difficult to assess the accuracy with which the nuclear correlations between the neighboring states near the pure  $(\nu g_{7/2})$ particle- $(\pi g_{9/2})$ hole configuration are taken into account in arriving at the experimental result of  $q_{snc}$ . It would require highly accurate theoretical inputs to  $q_{snc}$  which we, not involved in the analyses, are unable to assess. Assuming the  $q_{snc}^L$  given by the mapping for  $q_{snc}$ , the RIKEN result implies an extremely significant AIQ effect,

$$q_{ssb}^{RIKEN} \approx (0.6 - 0.7).$$
 (15)

This RIKEN result is saying that the "fundamental"  $g_A$  inherited from QCD is ~ 0.8 – 0.9, not 1.276. Given its fundamentality, it should apply to ALL weak processes at all kinematics in nuclear medium, not just superallowed Gamow-Teller transitions at  $(q, \omega) \rightarrow (0, 0)$ . Even more significantly, in the process  $0\nu\beta\beta$  transitions in nuclei where the momentum transfer can be of order ~ 100 MeV, the axial coupling constant appears at fourth power in the cross-section.

So the big question is: Which one is correct?

# B. Prediction by GD/CD-QCD

As noted, neither the coefficient  $c_A$  nor the anomalous dimension  $\beta'_*$  in (10) has been worked out for  $N_f = 2$ or 3. However the recent development on the IR fixed point in the CD-QCD scheme anchored on soft dilaton theorems in  $d\chi \text{EFT}$  has led Zwicky [15] to predict the anomalous dimension

$$\beta'_* = 0. \tag{16}$$

This predicts

$$q_{ssb} \approx 1.$$
 (17)

This would apply not just to the axial current but also to other channels in the GnEFT Lagrangian to be mentioned below. We put an approximate equality because of the caveat that follows from (10). Here the higher scale-order term  $O(\alpha_s - \alpha_{\rm IR})$  is taken to be negligible. But it is not obvious that the system lies very near the IR fixed point, so that  $O(\alpha_s - \alpha_{\rm IR})$  may not be ignorable. In fact the would-be free-space dilaton mass, if identified with  $f_0(500)$  in the CD-QCD scheme, must be higher than the pion mass, so the correction to  $\beta'_*$  in the exponent to  $\Phi$  may not be ignorable unless the system is driven close to the IR fixed point by external disturbance, say, baryon density. We cannot give a convincing simple argument at the moment but the fact that the phenomenology in GnEFT with the topology change as hadron-quark crossover gives the precocious pseudoconformal sound velocity  $v_s^2/c^2 \approx 1/3$  in compact stars at a density as low as  $n \gtrsim 3n_0$  is consistent with  $\beta'_* \approx 0$ .

 $<sup>^1</sup>$  In the scheme, three-body forces are integrated out together with the  $\Delta$  resonance, as justified in the Gamow-Teller channel.

# C. Evidences for and against $q_{ssb}^{RIKEN} \approx (0.6 - 0.7)$

The dilaton-limit fixed point (DLFP) in GnEFT is expected to appear at a much higher density than normal  $n_0$ , encompassing what's expected in massive compact stars. How the effect relevant to normal nuclear matter persists from  $n_0$  where the  $q_A$  problem lies to  $\sim (5-7)n_0$ probed in compact stars cannot be given a simple description. But as defined, the AIQ should apply to the axial coupling constant  $q_A$  figuring in ALL processes involving the nucleon axial current independently of the kinematics. Furthermore this applies to ALL processes where the pion-nucleon coupling figures as expected from the Goldberger-Treiman relation between  $g_A$  and  $g_{\pi NN}$ . One may then ask how the currently successful  $\chi EFT$  in nuclear physics which is anchored on soft pions - which includes nuclear astrophysical observations - escapes "torpedos" of the fundamentally quenched  $q_A$ ?

Among numerous processes in nuclear dynamics, there is one case that seems to be strongly against  $q_{ssb}^{RIKEN}$ . It is the first-forbidden beta decay in nuclei dominated by the axial-charge operator  $J_{05}^{\pm}$ . For this process, the soft-pion prediction [29] with  $q_{ssb} = 1$  is well confirmed by an experiment in Pb nuclei [30]. This result has been further supported with A = 12 nuclei [31], indicating relatively weak dependence on density. Note that the AIQ factor for the axial current controlled by soft-dilaton effects [15] should apply exactly as in the superallowed GT transition and the axial charge operator that enters in the first-forbidden process is primarily controlled by soft-pion effects. It is intriguing that both cases involve soft theorems in providing nuclear model independence and could be given a "first-principle" test.

There are however certain axial processes being discussed in the literature that may be indicative for an appreciable quenching of  $q_{ssb}$ . As stressed by Suhonen<sup>2</sup>, there is a suggestion that measuring  $\beta$ -decay spectral shape instead of superallowed GT transitions would allow the AIQ to be better extracted. Indeed there seems to be a signal [32] for the  $\delta = 1 - q_{ssb} \sim (0.3 - 0.4)$  comparable to what's indicated by the RIKEN data. Here the nucleon wavefunction involved probes different kinematics from the superallowed GT transition, so the LFL fixed-point-to-ESPM mapping where the nuclear tensor forces play a key role for determining  $q_{snc}^L$  cannot be applied. In fact the spectral shape involves nuclear matrix elements with the structure of the current operator (i.e., many-body meson-exchange currents) and the BR-scaled tensor forces (mentioned in [23]) appreciably different from what enters in the superallowed GT transitions, so sorting out the nuclear effect to handle the intricacy between "fundamental" and "mundane" nuclear effects would seem to require a drastically different approach.

# IV. CONCLUSION

In this paper, we provided what we consider to be a solution to the long-standing mystery of the quenched  $g_A$  in nuclear GT transitions. What turns out to play a key role here is the presence of an IR fixed-point in QCD with 2 or 3 flavors, so far unobserved, at which hidden scale symmetry with GD [11] or CD-QCD [15] intervenes. Combined with hidden local symmetry [14], implemented with a hadron-quark continuity in terms of a topology change, this GD/CD-QCD scheme has met with no tension with modern developments with compact-star observables [8, 9].

A most striking observation in the scheme with the IR fixed point is that the anomalous dimension at the fixed point,  $\beta'_*$ , comes out to be zero [15]. As shown in this paper, if higher-order scale-chiral contributions could not be ignored as assumed so far - and are being looked into [33], this would imply that there can be an important renormalization of the axial-current coupling constant from the free-space value 1.276 to  $\sim 1/2$ in nuclear matter leading to a dramatic quenching of  $g_A$ in nuclear medium – which has not so far been detected in nuclear processes where pion-nuclear coupling is involved – and to a ~  $(1/2)^4$  quenching in  $2\nu\beta\beta$  and  $0\nu\beta\beta$ processes - highly crucially relevant for on-going BSM search. The culprit for this big "anomaly-induced" effect is the recent RIKEN experiment in the superallowed Gamow-Teller transition in the doubly-closed magic-shell nucleus <sup>100</sup>Sn. An unambiguous reconfirmation or invalidation of this result is called for. If it turned out to be either definitively reconfirmed or invalidated, then it would either support or rule out the possible IR fixed point with GD/QCD-CD, this coming from nuclear physics.

#### Acknowledgments

We are grateful for correspondences with helpful comments from F. Sannino, J.T. Suhonen, K. Yamawaki and R. Zwicky.

- Y. L. Ma and M. Rho, Phys. Rev. Lett. **125**, 142501 (2020).
- [2] M. Rho, Symmetry 15, no.9, 1648 (2023).

- [3] J. T. Suhonen, Front. in Phys. 5, 55 (2017).
- [4] D. Lubos *et al* Phys. Rev. Lett. **122**, no.22, 222502 (2019).
- [5] D. H. Wilkinson, "Nuclear physics with heavy ions and mesons," R. Balian, M. Rho, and G. Ripka, Eds, North-Holland: Amsterdam, The Netherlands, (1978).
- [6] B. Friman and M. Rho, Nucl. Phys. A 606, 303 (1996).

 $<sup>^2</sup>$  Private communication from J. Suhonen.

- [7] R. Shankar, Rev. Mod. Phys. 66, 129 (1994).
- [8] Y. L. Ma and M. Rho, Prog. Part. Nucl. Phys. 113, 103791 (2020).
- [9] M. Rho and Y. L. Ma, Mod. Phys. Lett. A 36, 2130012 (2021).
- [10] G. E. Brown and M. Rho, Phys. Rev. Lett. 66, 2720 (1991).
- [11] R. J. Crewther, Universe 6, no.7, 96 (2020).
- [12] R. J. Crewther and L. C. Tunstall, Phys. Rev. D 91, 034016 (2015).
- [13] W. G. Paeng, T. T. S. Kuo, H. K. Lee, Y. L. Ma and M. Rho, Phys. Rev. D 96, 014031 (2017).
- [14] M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. 54, 1215 (1985); K. Yamawaki, Symmetry 15, no.12, 2209 (2023).
- [15] R. Zwicky, [arXiv:2312.13761 [hep-ph]]; L. Del Debbio and R. Zwicky, JHEP 08, 007 (2022).
- [16] Briefly stated, the phase CD-QCD is different from the conformal window [17–20] in the  $N_c$ — $N_f$  phase diagrams in that the quark condensate  $\langle \bar{q}q \rangle$  exists in the CD-QCD phase, thus it can cause spontaneous breaking of scale symmetry in the chiral limit and give a large decay constant to dilaton. This gives mass to other hadrons through the Goldberger-Treiman type relation. The presence of other massive particles does not break scale invariance in the CD-QCD phase.
- [17] W. E. Caswell, Phys. Rev. Lett. 33, 244 (1974).
- [18] T. Banks and A. Zaks, Nucl. Phys. B 196, 189 (1982).
- [19] K. Yamawaki, M. Bando and K. I. Matumoto, Phys. Rev. Lett. 56, 1335 (1986).

- [20] D. D. Dietrich, F. Sannino and K. Tuominen, Phys. Rev. D 72, 055001 (2005).
- [21] M. Rho, Phys. Rev. Lett. 66, 1275 (1991).
- [22] J. W. Holt, G. E. Brown, T. T. S. Kuo, J. D. Holt and R. Machleidt, Phys. Rev. Lett. **100**, 062501 (2008).
- [23] There is what's heralded as "first-principles" calculation [24] but it fails for the superallowed <sup>100</sup>Sn decay as pointed out in [1]. The problem is two-fold: First the nuclear tensor forces in GnEFT controlled by the BR scaling as a function of density are not properly taken into account and second the many-body currents in the Gamow-Teller channel are not protected by chiral symmetry and hence cannot be controlled at low order of chiral expansion, this in stark contrast to the axial-charge transition mentioned in Sect. III C.
- [24] P. Gysbers et al, Nature Phys. 15, 428 (2019).
- [25] L. Q. Shao and M. Rho, [arXiv:2311.15085 [nucl-th]].
- [26] C. B. Hinke et al., Nature (London) 486, 341 (2012).
- [27] S. R. Beane and U. van Kolck, Phys. Lett. B 328, 137 (1994)
- [28] W. G. Paeng, H. K. Lee, M. Rho and C. Sasaki, Phys. Rev. D 88, 105019 (2013).
- [29] K. Kubodera and M. Rho, Phys. Rev. Lett. 67, 3479 (1991).
- [30] E. K. Warburton, Phys. Rev. C 44, 233 (1991).
- [31] K. Minamisono et al, Phys. Rev. C 65, 015209 (2002).
- [32] A. F. Leder et al, Phys. Rev. Lett. 129, 232502 (2022).
- [33] L.Q. Shao, "IR fixed point emergent in dense nuclear matter via distorted dilaton potential," to appear.