Toeplitz Operators and Berezin-type Operators on Different Bergman Spaces

Lvchang Li * Haichou Li †

Abstract

In the present paper, we study the boundedness and compactness of Toeplitz operators and Berezin-type operators between different weighted Bergman spaces over tubular domains in \mathbb{C}^n . We establish their connection with Carleson measures and provide some characterizations.

Key words: Bergman space; Toeplitz operator; Berezin-type operator; Carleson measure

1 Introduction

Carleson measures are a powerful tool in modern complex analysis, initially introduced by Carleson [6] while studying the Corona problem on the Hardy space structure of the unit disk. Subsequently, it became the most essential tool in the study of operator theory on function spaces. Regarding the further applications of Carleson measures in operator theory on function spaces, refer to [3, 5, 8, 10, 11].

It is an interesting topic to study the analytic properties of operators from one function space to another function space. In particular, it is also meaningful to study the boundedness and compactness of parametric operators from one weighted Bergman space to another weighted Bergman space. In the present paper, We study the analytic properties of two kinds of operators on the Bergman spaces over tubular domains. Specifically, we study the relationship between the boundedness and compactness of the operators and the Carleson measures. Finally, the characterization between them is given.

^{*}College of Mathematics and Informatics, South China Agricultural University, Guangzhou, 510640, China Email: 20222115006@stu.scau.edu.cn.

[†]Corresponding author, College of Mathematics and Informatics, South China Agricultural University, Guangzhou, 510640, China Email: hcl2016@scau.edu.cn. Li is supported by NSF of China (Grant No. 12326407 and 12071155).

The main motivation of this work comes from the work of many scholars. In some simple domains, such as the unit ball, Zhao et al.[15]studied the boundedness and compactness of Toeplitz operators on this domain, and connected it with Carleson measure, and gave an equivalent characterization. Subsequently, zhou et al.[14] improved the results of zhao et al.[15] on this basis, and studied a class of operators similar to Toeplitz operators, later called Berezin-type operators, which evolved from Berezin transform. At the end of the article, Zhou et al.[14] further discussed the case when the index is infinite. For the Toeplitz operator theory on the unit disk and the unit ball, see [18, 19]. On bounded strongly pseudoconvex domains, Abate et al.[1, 2, 3] also studied similar problems.

The previous studies are all on bounded domains, and some scholars have also carried out related research on unbounded domains of interest to the author of this paper. Si et al. [17] studied the relationship between the Toeplitz operators and the Carleson measures on the Bergman spaces on the Siegel upper half spaces. In this paper, we will study the tubular domains. Deng et al. [7] provided the reproducing kernels of Bergman spaces on tubular domains by a new method, and Liu et al. [13] presented fundamental properties of the Bergman metric on these domains, laying the groundwork for subsequent theoretical studies. The first author Li [12] studied the Toeplitz operators on the weighted Bergman spaces over the tubular domains, and established the relationship between the boundedness and compactness of the operators and the Carleson measures. This paper is the subsequent work of [12]. The operators studied in this paper are Berezin-type operators and Toeplitz operators with parameters. The Berezin-type operator is derived from the Berezin transform. The Berezin transform is an important tool for studying operator theory on function spaces. For details, please refer to [19, 20].

The organization of the paper is as follows: The second part introduces some basic terms; The third part gives some instrumental lemmas; The fourth part gives the characterization of Carleson measures and vanishing Carleson measures on the tubular domains; In the last part, we discuss Toeplitz operators and Berezin type operators. We link their boundedness and compactness with Carleson measure, and obtain the main results, including Theorem 5.1, 5.2 and 5.3.

2 Preliminaries

Let \mathbb{C}^n be the *n* dimensional complex Euclidean space. For any two points $z = (z_1, \dots, z_n)$ and $w = (w_1, \dots, w_n)$ in \mathbb{C}^n , we write

$$z \cdot \bar{w} := z_1 \bar{w}_1 + \dots + z_n \bar{w}_n,$$

 $z^2 = z \cdot z := z_1^2 + z_2^2 + \dots + z_n^2$

and

$$|z| := \sqrt{z \cdot \bar{z}} = \sqrt{|z_1|^2 + \dots + |z_n|^2}.$$

The set $\mathbb{B}_n = \{z \in \mathbb{C}^n : |z| < 1\}$ will be called the unit ball of \mathbb{C}^n . The tubular domain T_B of \mathbb{C}^n with base B, is defined as follows:

$$T_B = \{ z = x + iy \in \mathbb{C}^n | x \in \mathbb{R}^n, y \in B \subseteq \mathbb{R}^n \},$$

where

$$B = \{ (y', y_n) = (y_1, \dots, y_{n-1}, y_n) \in \mathbb{R}^n \mid y'^2 := y_1^2 + \dots + y_{n-1}^2 < y_n \}.$$

We define the spaces $L_{\alpha}^{p}(T_{B})$, which is composed of all Lebesgue measurable functions f on T_{B} , and its norm

$$||f||_{p,\alpha} = \left\{ \int_{T_B} |f(z)|^p dV_\alpha(z) \right\}^{\frac{1}{p}}$$

is finite, where $dV_{\alpha}(z) = (y_n - |y'|^2)^{\alpha} dV(z)$, $\alpha > -1$, dV(z) denotes the Lebesgue measure on \mathbb{C}^n .

The space $L^{\infty}(T_B)$ consists of all bounded Lebesgue measurable functions f on T_B such that

$$||f||_{\infty} = \operatorname{ess sup}_{z \in T_{R}} |f(z)|$$

is finite.

The Bergman spaces $A_{\alpha}^{p}(T_{B})$ on tube T_{B} is a set composed of all holomorphic functions in $L_{\alpha}^{p}(T_{B})$. We know that when $1 \leq p < \infty$ the space $A_{\alpha}^{p}(T_{B})$ is a Banach space with the norm $\|\cdot\|_{p,\alpha}$. In particular, when p=2, $A_{\alpha}^{p}(T_{B})$ is a Hilbert space.

An very important orthogonal projection from $L_{\alpha}^{2}\left(T_{B}\right)$ to $A_{\alpha}^{2}\left(T_{B}\right)$ is the following integral operator:

$$P_{\alpha}f\left(z\right) = \int_{T_{B}} K_{\alpha}\left(z, w\right) f\left(w\right) dV_{\alpha}\left(w\right),$$

with the Bergman kernel

$$K_{\alpha}(z,w) = \frac{2^{n+1+2\alpha}\Gamma(n+1+\alpha)}{\Gamma(\alpha+1)\pi^{n}} \left(\left(z'-\overline{w'}\right)^{2} - 2i\left(z_{n}-\overline{w_{n}}\right) \right)^{-n-\alpha-1}.$$

For convenience, we introduce the following notations:

$$\rho(z, w) = \frac{1}{4} \left(\left(z' - \overline{w'} \right)^2 - 2i \left(z_n - \overline{w_n} \right) \right)$$

and let $\rho(z) := \rho(z, z) = y_n - y'^2$.

With the above notion $\rho(z, w)$, the weighted Bergman kernel of T_B becomes

$$K_{\alpha}(z,w) = \frac{\Gamma(n+\alpha+1)}{2^{n+1}\pi^{n}\Gamma(\alpha+1)\rho(z,w)^{n+\alpha+1}}.$$

We state some explanations regarding the boundary of tubes T_B .

Recall $\rho(z) = y_n - y'^2$ and let $\partial T_B := \{z \in \mathbb{C}^n : \rho(z) = 0\}$ denote the boundary of T_B . Then

$$\widehat{T_B} := T_B \cup \partial T_B \cup \{\infty\}$$

is the one-point compactification of T_B .

Also, let $\partial \widehat{T}_B := \partial T_B \cup \{\infty\}$. Thus, $z \to \partial \widehat{T}_B$ means $\rho(z) \to 0$ or $|z| \to \infty$.

We review the Bergman metric ball on domains in \mathbb{C}^n .

We will use the important transform $\Phi: \mathbb{B}_n \to T_B$ given by

$$\Phi(z) = \left(\frac{\sqrt{2}z'}{1+z_n}, \ i\frac{1-z_n}{1+z_n} - i\frac{z' \cdot z'}{(1+z_n)^2}\right), \ z \in \mathbb{B}_n$$

and it is not hard to calculate that

$$\Phi^{-1}(w) = \left(\frac{2iw'}{i + w_n + \frac{i}{2}w' \cdot w'}, \frac{i - w_n - \frac{i}{2}w' \cdot w'}{i + w_n + \frac{i}{2}w' \cdot w'}\right), \ w \in T_B.$$

The mapping Φ is a biholomorphic map from \mathbb{B}_n to T_B and a key tool for this paper.

In Krantz's book, there is the following proposition [9, proposition 1.4.12]:

Let Ω_1 , $\Omega_2 \subseteq \mathbb{C}^n$ be domains and $f: \Omega_1 \to \Omega_2$ a biholomorphic mapping. Then f induces an isometry of Bergman metrics:

$$\beta_{\Omega_{1}}(z, w) = \beta_{\Omega_{2}}(f(z), f(w))$$

for all $z, w \in \Omega_1$.

Hence, taking $\Omega_1 = T_B$ and $\Omega_2 = \mathbb{B}_n$, we have:

$$\beta_{T_{B}}\left(z,w\right)=\beta_{\mathbb{B}_{n}}\left(\Phi^{-1}\left(z\right),\Phi^{-1}\left(w\right)\right)=\tanh^{-1}\left(\left|\varphi_{\Phi^{-1}\left(z\right)}\left(\Phi^{-1}\left(w\right)\right)\right|\right).$$

A computation shows that

$$\beta_{T_B}(z, w) = \tanh^{-1} \sqrt{1 - \frac{\rho(z)\rho(w)}{|\rho(z, w)|^2}}.$$

Let D(z,r) denote the Bergman metric ball at z with radius r, that is

$$D(z,r) = \{ w \in T_B : \beta(z,w) < r \}.$$

Let \mathcal{M}_+ be the set of all positive Borel measure μ such that

$$\int_{T_B} \frac{d\mu\left(z\right)}{\left|\rho\left(z,\mathbf{i}\right)\right|^t} < \infty$$

for some t > 0, where $\mathbf{i} = (0', i)$.

Next, we introduce the two operators studied in the present paper. For s,t>0, we denote

$$B_{s,t}(\mu)(z) := \rho(z)^t \int_{T_B} \frac{1}{|\rho(z,w)|^{(n+1+\alpha)s+t}} d\mu(w).$$

This is called a Berezin-type transform for the measure μ , where $\alpha > -1$. Notice that $B_{1,n+\alpha+1}$ is the canonical Berezin transform of μ except a constant.

Let $\xi > -1$ and $\mu \in \mathcal{M}_+$ a positive Borel measure on T_B . For $f \in H(T_B)$, where $H(T_B)$ is the space consisting of all holomorphic functions on T_B , we consider the following sublinear operator:

$$B_{\mu}^{\xi} f(z) = \int_{T_B} \frac{|f(w)|}{|\rho(z, w)|^{n+1+\xi}} d\mu(w), \ z \in T_B.$$

We refer to B_{μ}^{ξ} as a Berezin-type operator.

The Toeplitz operator T^{ξ}_{μ} is defined by:

$$T_{\mu}^{\xi} f\left(z\right) = \int_{T_{B}} \frac{f\left(w\right)}{\rho\left(z,w\right)^{n+1+\xi}} d\mu\left(w\right), \ z \in T_{B}.$$

It is clear that $|T_{\mu}^{\xi}f| \leq B_{\mu}^{\xi}f$ for $f \in H(T_B)$. Therefore, The boundedness of Berezin-type operators implies the boundedness of Toeplitz operators.

For any positive numbers $\lambda > 0$ and $\alpha > -1$, a positive Borel measure μ is considered a (λ, α) -Carleson measure, if for all positive integers p and q where $q/p = \lambda$, there exists a positive constant C > 0 such that

$$\int_{T_R} |f(z)|^q \ d\mu(z) \le C ||f||_{p,\alpha}^q$$

for every $f \in A^p_{\alpha}(T_B)$.

A positive Borel measure μ as a vanishing (λ, α) -Carleson measure, if for any positive numbers p and q such that $q/p = \lambda$, and for any bounded sequence f_k in $A^p_{\alpha}(T_B)$ that converges uniformly to 0 on every compact subset of T_B , we have

$$\lim_{k \to \infty} \int_{T_R} |f_k|^q \, d\mu = 0.$$

We also denote by

$$\|\mu\|_{\lambda,\alpha} = \sup_{f \in A^p_{\alpha}(T_B), \|f\|_{p,\alpha} \le 1} \int_{T_B} |f(z)|^q d\mu(z).$$

Finally, we recall the definition of a compact operator: Given two Banach spaces X and Y, a bounded linear operator is called compact if it maps any bounded set in X to a set that is relatively compact in the topology of Y. When the Banach space X is reflexive, a compact operator on X is equivalent to being completely continuous. Complete continuity refers to an operator that maps weakly convergent sequences to sequences that converge in norm. For sublinear operators, the boundedness and compactness of operators are also defined according to the boundedness and compactness of linear operators

Throughout the paper we use C to denote positive constants whose value may change from line to line but does not depend on the functions being considered. The notation $A \lesssim B$ means that there is a positive constant C such that $A \leq CB$, and the notation $A \simeq B$ means that $A \lesssim B$ and $B \lesssim A$.

3 Main lemmas

To prove our main results, we need the following key lemmas, where Lemmas 3.1-3.4 are from [13]. They play a crucial role as instrumental lemmas in the present paper.

The following lemma 3.1 is the so-called Bergman decomposition on tubular domains. It plays an important role in the topic of operator theory on function spaces.

Lemma 3.1 There exists a positive integer N such that for any $0 < r \le 1$ we can find a sequence $\{a_k\}$ in T_B with the following properites:

- $(1) T_B = \bigcup_{k=1}^{\infty} D(a_k, r);$
- (2) The sets $D(a_k, r/4)$ are mutually disjoint;
- (3) Each point $z \in T_B$ belongs to at most N of the sets $D(a_k, 2r)$.

Any sequence $\{a_k\}$ satisfying the conditions of the above lemma is called a lattice.

The following lemmas 3.2 and 3.3 give the control of the size of $|\rho(z, w)|$ in the Bergman ball D(z, r) and the size of the Bergman ball D(z, r) under the Euclidean metric, respectively.

Lemma 3.2 For any r > 0, the inequalities

$$|\rho(z,u)| \simeq |\rho(z,v)|$$

hold for all $z, u, v \in T_B$ with $\beta(u, v) < r$.

Lemma 3.3 For any $z \in T_B$ and r > 0 we have

$$V_{\alpha}\left(D\left(z,r\right)\right) \simeq \rho\left(z\right)^{n+\alpha+1}$$
.

The following lemma 3.4 plays a key role in the integral operation in this paper.

Lemma 3.4 Let r, s > 0, t > -1 and r + s - t > n + 1, then

$$\int_{T_B} \frac{\rho(w)^t}{\rho(z,w)^r \rho(w,u)^s} dV(w) = \frac{C_1(n,r,s,t)}{\rho(z,u)^{r+s-t-n-1}}$$

for all $z, u \in T_B$, where

$$C_1(n, r, s, t) = \frac{2^{n+1}\pi^n\Gamma(1+t)\Gamma(r+s-t-n-1)}{\Gamma(r)\Gamma(s)}.$$

In particular, let $s, t \in \mathbb{R}$, if t > -1, s - t > n + 1, then

$$\int_{T_{B}} \frac{\rho(w)^{t}}{|\rho(z,w)|^{s}} dV(w) = \frac{C_{1}(n,s,t)}{\rho(z)^{s-t-n-1}}.$$

Otherwise, the above equation is infinity.

The following lemma is commonly encountered in the operator theory on function spaces topic, illustrating that the growth of functions in Bergman spaces is controlled.

Lemma 3.5 On the Bergman space $A_{\alpha}^{p}(T_{B})$, every valuation functional is a bounded linear functional. More specifically, each function $f \in T_{B}$ has the property

$$|f(z)|^p \le \frac{C}{\rho(z)^{n+\alpha+1}} \int_{D(z,r)} |f(w)|^p dV_{\alpha}(w).$$

The following lemma 3.6 is a utility lemma, and its proof can be found in [12].

Lemma 3.6 As |z| approaches infinity, $\rho(z, \mathbf{i})$ also tends to infinity.

We provide several lemmas related to the properties of reproducing kernels, as seen in Lemma 3.7 and Lemma 3.8.

Lemma 3.7 Let 1 -1 for each z > 0, the Bergman kernel function $K_{\alpha}(z, w)$ is in $A^{p}_{\alpha}(T_{B})$, and its norm is $C_{\alpha}(z)^{-(n+\alpha+1)/p'}$, where p' = p/(p-1) and C is a positive constant only depending on n, α , and p.

Lemma 3.8 For $1 and <math>\alpha > -1$, we have $K_{\alpha,z} \| K_{\alpha,z} \|_{p,\alpha}^{-1} \to 0$ weakly in $A_{\alpha}^{p}(T_{B})$ as $z \to \partial \widehat{T_{B}}$.

4 Carleson measures on T_B

Since the Carleson measure is a very important tool in operator theory, we first need to characterize the Carleson measures and the vanishing Carleson measures. Carleson measures and vanishing Carleson measures on the unit disk and unit ball have numerous characterizations, as detailed in [18, 19]. We now provide characterizations of Carleson measures and vanishing Carleson measures on tubular domains, which will serve as basis results for our subsequent theorems.

In order to get the characterization of Carleson measures, we need the help of the following lemma 4.1.

Lemma 4.1 For $1 \le \lambda < \infty$, $-1 < \alpha < \infty$, and μ being a (λ, α) -Carleson measure on T_B , satisfying $\mu(D(z,r)) \lesssim \rho(z)^{(n+1+\alpha)\lambda}$ for any Bergman ball D(z,r), the following inequality holds for any $f \in H(T_B)$ and 0 :

$$\int_{T_R} |f(z)|^p d\mu(z) \lesssim \int_{T_R} |f(z)|^p \rho(z)^{(n+1+\alpha)\lambda - (n+1)} dV(z).$$

Proof:

By Lemma 3.5, for 0 < r < 1, we have

$$|f(z)|^p \le \frac{C}{\rho(z)^{n+1}} \int_{D(z,r)} |f(w)|^p dV(w).$$

Hence, using Fubini's theorem along with Lemma 3.2, we obtain

$$\int_{T_B} |f(z)|^p d\mu(z) \lesssim \int_{T_B} \frac{1}{\rho(z)^{n+1}} \int_{D(z,r)} |f(w)|^p dV(w) d\mu(z)
= \int_{T_B} |f(w)|^p \int_{D(w,r)} \frac{d\mu(z)}{\rho(z)^{n+1}} dV(w)
\lesssim \int_{T_B} |f(w)|^p \int_{D(w,r)} \frac{d\mu(z)}{\rho(w)^{n+1}} dV(w)
= \int_{T_B} |f(w)|^p \frac{\mu(D(w,r))}{\rho(w)^{n+1}} dV(w)
\lesssim \int_{T_B} |f(w)|^p \rho(w)^{(n+1+\alpha)\lambda - (n+1)} dV(w).$$

This completes the proof of Lemma 4.1.

Theorem 4.1 Suppose $1 \le \lambda < \infty$ and $-1 < \alpha < \infty$, the following conditions are equivalent:

(1) μ is a (λ, α) -Carleson measure.

(2) For any real number r with 0 < r < 1 and any $a \in T_B$

$$\mu\left(D\left(a,r\right)\right) \lesssim \rho\left(a\right)^{(n+1+\alpha)\lambda}$$
.

(3) For some (every) t > 0, the Berezin-type transform of μ

$$B_{\lambda,t}(\mu) \in L^{\infty}(T_B)$$
.

Proof:

(1) \Rightarrow (2) First assume that μ is a (λ, α) -Carleson measure for $A^p_{\alpha}(T_B)$. There is a constant C > 0 such that

$$\int_{T_B} |f(z)|^q d\mu(z) \le C ||f||_{p,\alpha}^q$$

for all $f \in A^p_\alpha(T_B)$.

In particular, if $a \in T_B$ and

$$g(w) = \left(\frac{\rho(a)^{n+1+\alpha}}{|\rho(w,a)|^{2(n+1+\alpha)}}\right)^{1/p}, \ z \in T_B,$$

then $||g(w)||_{p,\alpha}$ is a constant, so

$$\int_{D(a,r)} |g(w)|^q d\mu(w) \le \int_{T_B} |g(w)|^q d\mu(w) \le C.$$

According to Lemma 3.2, $|\rho(w, a)|$ is comparable to $|\rho(a)|$ when $w \in D(a, r)$, we have $\mu(D(a, r)) \lesssim \rho(a)^{(n+1+\alpha)\lambda}$.

 $(2) \Rightarrow (3)$ It follows from Lemma 4.1 that

$$B_{\lambda,t}(\mu)(z) \lesssim \rho(z)^t \int_{T_B} \frac{1}{|\rho(z,w)|^{(n+1+\alpha)\lambda+t}} \rho(w)^{(n+1+\alpha)\lambda-(n+1)} dV(w).$$

By calculation, we find that the parameters in the above expression satisfy the conditions of Lemma 3.4, so the above expression is less than a constant.

 $(3) \Rightarrow (1)$ Assume (3) is ture. For any $f \in A^p_\alpha(T_B)$, we have

$$\int_{T_B} |f(z)|^q d\mu(z) \le \sum_{k=1}^{+\infty} \int_{D(a_k,r)} |f|^q d\mu \le \sum_{k=1}^{+\infty} \mu(D(a_k,r)) \sup_{z \in D(a_k,r)} |f(z)|^q.$$

for all $k \geq 1$.

It follows from Lemma 3.5 that

$$\sup_{z \in D(a_k, r)} |f(z)|^p \lesssim \frac{1}{\rho(a_k)^{n+\alpha+1}} \int_{D(a_k, 2r)} |f(w)|^p dV_{\alpha}(w).$$

Then we obtain

$$\int_{T_B} |f\left(z\right)|^q d\mu\left(z\right) \lesssim \sum_{k=1}^{\infty} \frac{\mu\left(D\left(a_k,r\right)\right)}{\rho\left(a_k\right)^{(n+\alpha+1)\lambda}} \left(\int_{D\left(a_k,2r\right)} |f\left(z\right)|^p dV_{\alpha}\left(z\right)\right)^{\lambda},$$

where $\lambda = q/p$.

From (3), when $z \in D\left(a_k, r\right)$ holds, $\mu\left(D\left(a_k, r\right)\right) / \rho\left(a_k\right)^{(n+\alpha+1)\lambda} \lesssim C$ is true.

Since $\lambda > 1$, it follows that

$$\int_{T_B} |f(z)|^q d\mu(z) \lesssim \sum_{k=1}^{+\infty} \left(\int_{D(a_k, 2r)} |f(z)|^p dV_\alpha(z) \right)^{\lambda}
\lesssim \left(\sum_{k=1}^{+\infty} \int_{D(a_k, 2r)} |f(z)|^p dV_\alpha(z) \right)^{\lambda}
\leq N^{\lambda} ||f||_{p,\alpha}^q,$$

where N is as in Lemma 3.1. The proof is complete.

Theorem 4.2 Suppose $1 \le \lambda < \infty$ and $-1 < \alpha < \infty$, the following statements are equivalent:

- (1) μ is a vanishing (λ, α) -Carleson measure.
- (2) For some(any) t > 0

$$\lim_{a \to \partial \widehat{T_B}} \int_{T_B} \frac{\rho\left(a\right)^t}{\left|\rho\left(z,a\right)\right|^{(n+1+\alpha)\lambda+t}} d\mu\left(z\right) = 0.$$

(3) For any real number r with 0 < r < 1 and any $a \in T_B$

$$\lim_{a \to \partial \widehat{T}_{B}} \frac{\mu\left(D\left(a, r\right)\right)}{\rho\left(a\right)^{(n+1+\alpha)\lambda}} = 0.$$

Proof:

 $(1) \Rightarrow (2)$ If μ is a vanishing (λ, α) -Carleson measure. Let t > 0 and

$$f_a(z) = \frac{\rho(a)^{t/q}}{\rho(z, a)^{(n+1+\alpha)/p+t/q}},$$

We calculate to obtain that

$$\sup_{a \in T_B} \|f_a(z)\|_{p,\alpha} < \infty.$$

So, we have

$$\int_{T_B} \frac{\rho(a)^t}{\left|\rho(z,a)\right|^{(n+1+\alpha)\lambda+t}} d\mu(z) \to 0$$

as $a \to \partial \widehat{T}_B$.

We denote $Q_k = \overline{D(\mathbf{i}, r)}$, if we can prove that f_a converges uniformly to 0 on every Q_k as $a \to \partial \widehat{T}_B$, then the conclusion holds. For a given Q_k , we have

$$\sup_{z \in Q_k} |f_a(z)| \lesssim \frac{\rho(a)^{t/q}}{|\rho(\mathbf{i}, a)|^{(n+1+\alpha)/p+t/q}}.$$

Since $|\rho(\mathbf{i}, a)| \geq 1/2$ for all $a \in T_B$, it follows that $\sup_{z \in Q_k} |f_a(z)| \lesssim \rho(z)^{t/q}$. Hence $f_a \to 0$ uniformly on Q_k as $a \to bT_B$. Due to Lemma 3.6, therefore $f_a \to 0$ uniformly on Q_k as $|a| \to \infty$. Thus, the condition (2) holds.

 $(2) \Rightarrow (3)$ The inequality

$$\int_{D(a,r)} \frac{\rho(a)^t}{\rho(z,a)^{(n+1+\alpha)\lambda+t}} d\mu(z) \le \int_{T_R} \frac{\rho(a)^t}{\rho(z,a)^{(n+1+\alpha)\lambda+t}} d\mu(z) \to 0$$

combined with Lemma 3.2 yields the desired result.

 $(3) \Rightarrow (1)$ Assuming $\{f_j\}$ is a bounded sequence in $A^p_{\alpha}(T_B)$ and uniformly converges to 0 on every compact subset of T_B , according to the definition of vanishing Carleson measure, it suffices to prove the following statement:

$$\lim_{j \to \infty} \int_{T_B} |f_j(z)|^q d\mu(z) = 0.$$

For any r > 0 we fix an r-lattice $\{a_j\}$ in the Bergman metric. Since $a_j \to \partial \widehat{T}_B$ as $j \to \infty$,

$$\lim_{j \to \infty} \frac{\mu\left(D\left(a_{j}, r\right)\right)}{\rho\left(a_{j}\right)^{(n+1+\alpha)\lambda}} = 0.$$

Given $\varepsilon > 0$, there is a positive integer N_0 such that

$$\frac{\mu\left(D\left(a_{j},r\right)\right)}{\rho\left(a_{j}\right)^{(n+1+\alpha)\lambda}} < \varepsilon, \quad j \ge N_{0}.$$

Therefore, the inequality to be proved can be divided into the following two parts:

$$\int_{T_{B}} |f_{j}(z)|^{q} d\mu(z) \lesssim \sum_{k=1}^{N_{0}-1} \frac{\mu(D(a_{k}, r))}{\rho(a_{k})^{(n+1+\alpha)\lambda}} \left(\int_{D(a_{k}, 2r)} |f_{j}(z)|^{p} dV_{\alpha}(z) \right)^{\lambda} + \sum_{k=N_{0}}^{\infty} \frac{\mu(D(a_{k}, r))}{\rho(a_{k})^{(n+1+\alpha)\lambda}} \left(\int_{D(a_{k}, 2r)} |f_{j}(z)|^{p} dV_{\alpha}(z) \right)^{\lambda}.$$

The first inequality must converge to 0 as $j \to \infty$, since $\{f_j\}$ converges to 0 on compact subsets. The second inequality, due to $\lambda > 1$, implies that

$$\sum_{k=N_0}^{\infty} \frac{\mu\left(D\left(a_k,r\right)\right)}{\rho\left(a_k\right)^{(n+1+\alpha)\lambda}} \left(\int_{D\left(a_k,2r\right)} \left|f_j\left(z\right)\right|^p dV_{\alpha}\left(z\right)\right)^{\lambda} \leq \varepsilon N^{\lambda} \|f_j\|_p^q$$

holds.

Combining the arbitrariness of ε , we conclude that μ is a vanishing (λ, α) -Carleson measure.

Remark 4.1 The theorem in this section is a generalization of theorem 4.1 and theorem 4.2 in [12]. Theorem 4.1 and Theorem 4.2 in this section characterize the Carleson measure and the vanishing Carleson measure, respectively. They will characterize the boundedness and compactness of the operator in the subsequent theorems 5.1 and 5.2, respectively.

5 Boundedness and compactness of operators

Now, we present the characterizations of the boundedness and compactness of Toeplitz operators and Berezin-type operators on Bergman spaces over tubular domains.

First, we present two basic lemma 5.1 and 5.2.

Let's recall the definition of a separated sequence. A sequence of points $\{z_k\}$ in T_B is called a separated sequence if there exists $\delta > 0$, such that $\beta(z, w) > \delta$ for any $i \neq j$.

Lemma 5.1 Given a separated sequence $\{z_k\}$ in T_B , with n < t < s, for any $z \in T_B$, the following holds.

$$\sum_{k=1}^{\infty} \frac{\rho(z_k)^t}{|\rho(z, z_k)|^s} \lesssim \rho(z)^{t-s}, \ z \in T_B.$$

Based on Bergman's decomposition (lemma 3.1) and combined with Lemma 3.4, we can derive the result. We omit it's proof here.

The following Lemma 5.2 provides a good estimate for the norm of functions in the spaces $L_{\alpha}^{p}(T_{B})$.

Lemma 5.2 Consider a positive sequence $\{c_j\}$ and a separated sequence $\{a_j\}$ in T_B . Let $b \in T_B$ satisfy

$$b>n\max\left(1,\frac{1}{p}\right)+\frac{1+\alpha}{p}.$$

Suppose f is a measurable function on T_B such that

$$|f(z)| \le \sum_{j=1}^{\infty} \frac{c_j}{|\rho(z, a_j)|^b}.$$

Then, $f \in L^p_{\alpha}(T_B)$, and we have the estimate

$$||f||_{p,\alpha}^p \lesssim \sum_{j=1}^{\infty} \frac{c_j^p}{\rho(a_j)^{bp-(n+1+\alpha)}}.$$

Proof:

When 0 , we have the inequality

$$|f(z)|^p \le \sum_{j=1}^{\infty} \frac{c_j}{|\rho(z, a_j)|^{bp}}.$$

By Lemma 3.4, it follows that

$$\int_{T_B} |f(z)|^p dV_{\alpha}(z) \leq \sum_{j=1}^{\infty} c_j^p \int_{T_B} \frac{1}{|\rho(z, a_j)|^{bp}} dV_{\alpha}(z)$$

$$\lesssim \sum_{j=1}^{\infty} \frac{c_j^p}{\rho(a_j)^{bp-(n+1+\alpha)}}.$$

For p > 1, we choose $b > n + \frac{1+\alpha}{p}$. Let p' be the conjugate exponent of p, such that $\frac{1}{p} + \frac{1}{p'} = 1$. By Hölder's inequality and Lemma 5.1, we obtain

$$|f(z)|^{p} = \left(\sum_{j=1}^{\infty} \frac{c_{j}}{|\rho(z, a_{j})|^{b}}\right)^{p}$$

$$\leq \left(\sum_{j=1}^{\infty} \frac{\rho(a_{j})^{b-(1+\alpha)/p}}{|\rho(z, a_{j})|^{b}}\right)^{p-1} \left(\sum_{j=1}^{\infty} \frac{c_{j}^{p} \rho(a_{j})^{b(1-p)+(1+\alpha)/p'}}{|\rho(z, a_{j})|^{b}}\right)$$

$$\lesssim \rho(z)^{-(1+\alpha)/p'} \left(\sum_{j=1}^{\infty} \frac{c_{j}^{p} \rho(a_{j})^{b(1-p)+(1+\alpha)/p'}}{|\rho(z, a_{j})|^{b}}\right).$$

This implies

$$||f||_{p,\alpha}^{p} \lesssim \sum_{j=1}^{\infty} c_{j}^{p} \rho\left(a_{j}\right)^{b(1-p)+(1+\alpha)/p'} \int_{T_{B}} \frac{\rho\left(z\right)^{-(1+\alpha)/p'}}{\left|\rho\left(z,a_{j}\right)\right|^{b}} dV_{\alpha}\left(z\right).$$

Since
$$\alpha - (1 + \alpha)/p' = (1 + \alpha)/p - 1 > -1$$
, and

$$b - (n+1+\alpha) + (1+\alpha)/p' = s - n - (1+\alpha)/p > 0,$$

we have completed the proof of the lemma according to Lemma 3.4.

Due to the above lemmas 5.1 and 5.2, combine with the characterization of Carleson measure. Now, we can prove the first main theorem 5.1 of this paper. The theorem mainly states that under the condition $p_1 \leq p_2$, the boundedness of Toeplitz operators T_{μ}^{ξ} , Berezin-type operators B_{μ}^{ξ} , and μ being a (λ, γ) -Carleson measure are equivalent, and it further states that their norms are mutually controlled.

Theorem 5.1 Let $0 < p_1 \le p_2 < \infty$, $-1 < \alpha_1, \alpha_2, \xi < \infty$ and $\mu \in \mathcal{M}_+$. Suppose that

$$n + 1 + \xi > n \max\left(1, \frac{1}{p_i}\right) + \frac{1 + \alpha_i}{p_i}, \quad i = 1, 2.$$

Let

$$\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}, \ \gamma = \frac{1}{\lambda} \left(\xi + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right).$$

Then the following statements are equivalent:

- (1) B^{ξ}_{μ} is bounded from $A^{p_1}_{\alpha_1}(T_B)$ to $L^{p_2}_{\alpha_2}(T_B)$. (2) T^{ξ}_{μ} is bounded from $A^{p_1}_{\alpha_1}(T_B)$ to $A^{p_2}_{\alpha_2}(T_B)$. (3) The measure μ is a (λ, γ) -Carleson measure.
- Moreover, we have

$$||B_{\mu}^{\xi}||_{A_{\alpha_1}^{p_1} \to L_{\alpha_2}^{p_2}} \simeq ||T_{\mu}^{\xi}||_{A_{\alpha_1}^{p_1} \to A_{\alpha_2}^{p_2}} \simeq ||\mu||_{\lambda,\gamma}.$$

Proof:

- $(1) \Rightarrow (2)$ The inequality $|T_{\mu}^{\xi}f| \leq B_{\mu}^{\xi}f$ implies that (2) holds trivially.
- $(2) \Rightarrow (3)$ Fix $a \in T_B$ and define $f_a(z) = \rho(z,a)^{-(n+1+\xi)}$. Given the condition $(n+1+\xi)p_1 > n+1+\alpha_1$, according to Lemma 3.4 we know that $f_a \in A_{\alpha_1}^{p_1}(T_B)$ with

$$||f_a||_{p_1,\alpha_1}^{p_1} \lesssim \rho(a)^{(n+1+\alpha_1)-(n+1+\xi)p_1}$$

Since

$$T_{\mu}^{\xi} f_{a}(z) = \int_{T_{B}} \frac{f_{a}(w)}{\rho(z, w)^{n+1+\xi}} d\mu(w)$$
$$= \int_{T_{B}} \frac{d\mu(w)}{\rho(z, w)^{n+1+\xi} \rho(w, a)^{n+1+\xi}},$$

we have

$$T_{\mu}^{\xi} f_{z}(z) = \int_{T_{B}} \frac{d\mu(w)}{|\rho(z, w)|^{2(n+1+\xi)}} \gtrsim \frac{\mu(D(z, r))}{\rho(z)^{2(n+1+\xi)}}.$$

Moreover, using the pointwise estimate for functions in Bergman spaces and the boundedness of the Toeplitz operator T^{ξ}_{μ} , we obtain

$$T_{\mu}^{\xi} f_{z}(z) = \left| T_{\mu}^{\xi} f_{z}(z) \right| \leq \frac{\| T_{\mu}^{\xi} f_{z} \|_{p_{2}, \alpha_{2}}}{\rho(z)^{(n+1+\alpha_{2})/p_{2}}}$$

$$\leq \| T_{\mu}^{\xi} \| \| f_{z} \|_{p_{1}, a_{1}} \rho(z)^{-(n+1+\alpha_{2})/p_{2}}$$

$$\lesssim \| T_{\mu}^{\xi} \| \rho(z)^{(n+1+\alpha_{1})/p_{1}-(n+1+\alpha_{2})/p_{2}-(n+1+\xi)}.$$

Hence, we have

$$\mu (D(z,r)) \lesssim \|T_{\mu}^{\xi}\| \rho(z)^{(n+1+\xi)+(n+1+\alpha_1)/p_1-(n+1+\alpha_2)/p_2}$$

= $\rho(z)^{(n+1+\gamma)\lambda}$.

This implies, by Theorem 4.1, that μ is a (λ, γ) -Carleson measure with

$$\|\mu\|_{\lambda,\gamma} \lesssim \|T^{\xi}_{\mu}\|.$$

(3) \Rightarrow (1) Suppose that μ is a (λ, γ) -Carleson measure. Since the condition $0 < p_1 \le p_2 < \infty$ implies that $\lambda > 1$ and $p_2/p_1 \ge 1$. According to Lemma 3.2, we know that when $w \in D(a_j, r)$, we have $|\rho(z, w)| \simeq |\rho(z, a_j)|$.

Therefore, we obtain

$$(B_{\mu}^{\xi}f)(z) = \int_{T_{B}} \frac{|f(w)|}{|\rho(z,w)|^{n+1+\xi}} d\mu(w)$$

$$\lesssim \sum_{j=1}^{\infty} \int_{D(a_{j},r)} \frac{|f(w)|}{|\rho(z,w)|^{n+1+\xi}} d\mu(w)$$

$$\lesssim \sum_{j=1}^{\infty} \left(\sup_{w \in D(a_{j},r)} |f(w)| \right) \int_{D(a_{j},r)} \frac{1}{|\rho(z,w)|^{n+1+\xi}} d\mu(w)$$

$$\lesssim \sum_{j=1}^{\infty} \left(\sup_{w \in D(a_{j},r)} |f(w)| \right) \frac{\mu(D_{j})}{|\rho(z,a_{j})|^{n+1+\xi}}.$$

According to Lemma 3.5, we have

$$|f(w)|^{p_1} \le \frac{C}{\rho(a_j)^{n+1+\alpha_1}} \int_{D(a_j,2r)} |f(w)|^{p_1} dV_{\alpha_1}(w)$$

holds for any $w \in D(a_i, r)$.

We get that

$$\left| \left(B_{\mu}^{\xi} f \right)(z) \right| \lesssim \sum_{j=1}^{\infty} \left(\frac{1}{\rho(a_{j})^{n+1+\alpha_{1}}} \int_{D(a_{j},2r)} |f(w)|^{p_{1}} dV_{\alpha_{1}}(w) \right)^{1/p_{1}} \frac{\mu(D(a_{j},r))}{\left| \rho(z,a_{j}) \right|^{n+1+\xi}}.$$

According to Lemma 5.2, we can obtain

$$||B_{\mu}^{\xi}f||_{p_{2},\alpha_{2}}^{p_{2}} \lesssim \sum_{j=1}^{\infty} \left(\int_{D(a_{j},2r)} |f(w)|^{p_{1}} dV_{\alpha_{1}}(w) \right)^{p_{2}/p_{1}} \left(\frac{\mu(D(a_{j},r))}{|\rho(a_{j})|^{(n+1+\gamma)\lambda}} \right)^{p_{2}}.$$

it follows Theorem 4.1, we have

$$||B_{\mu}^{\xi}f||_{p_{2},\alpha_{2}}^{p_{2}} \lesssim ||\mu||_{\lambda,\gamma}^{p_{2}} \sum_{j=1}^{\infty} \left(\int_{D(a_{j},2r)} |f(z)|^{p_{1}} dV_{\alpha_{1}}(z) \right)^{p_{2}/p_{1}}$$

$$\lesssim ||\mu||_{\lambda,\gamma}^{p_{2}} \left(\sum_{j=1}^{\infty} \int_{D(a_{j},2r)} |f(z)|^{p_{1}} dV_{\alpha_{1}}(z) \right)^{p_{2}/p_{1}}$$

$$\lesssim ||\mu||_{\lambda,\gamma}^{p_{2}} ||f||_{p_{1},\alpha_{1}}^{p_{2}}.$$

Hence, B_{μ}^{ξ} is bounded from $A_{\alpha_1}^{p_1}(T_B)$ to $L_{\alpha_2}^{p_2}(T_B)$.

Next, by combining the characterization of the vanishing Carleson measures in Theorem 4.2, we prove the second main theorem 5.2 of this paper. Before that, we present a lemma 5.3, which provides a sufficient condition for an operator to be compact.

Lemma 5.3 Let $0 < p_1, p_2 < \infty, -1 < \alpha_1, \alpha_2, \xi < \infty$ and $\mu \in \mathcal{M}_+$. Assume that $T^{\xi}_{\mu}: A^{p_1}_{\alpha_1}(T_B) \to A^{p_2}_{\alpha_2}(T_B)$ is a bounded operator. Suppose that for every bounded sequence $\{f_k\}$ in $A^{p_1}_{\alpha_1}(T_B)$ such that $f_k \to 0$ uniformly on every compact subset of T_B as $k \to \infty$, we have

$$\lim_{k \to \infty} \left\| T_{\mu}^{\xi} f_k \right\|_{p_2, \alpha_2} = 0.$$

Then, T^{ξ}_{μ} is compact from $A^{p_1}_{\alpha_1}(T_B)$ to $L^{p_2}_{\alpha_2}(T_B)$.

Proof: Let $\{f_k\}$ be a bounded sequence in $A_{\alpha_1}^{p_1}(T_B)$. Thus, there exists a constant M > 0 such that

$$||f_k||_{p_1,\alpha_1} \leq M$$

for all $k \geq 1$.

By Lemma 3.5, $\{f_k\}$ is uniformly bounded on every compact subset of T_B .

By Montel's Theorem, there exists a subsequence of $\{f_{k_j}\}$, $j=1,2,3\ldots$, such that $f_{k_j} \to f$ uniformly on every compact subset of T_B for some holomorphic function f on T_B , as $j \to \infty$.

Applying Fatou's Lemma we have

$$\int_{T_B} |f(z)|^{p_1} dV_{\alpha_1}(z) = \int_{T_B} \lim_{j \to \infty} |f_{k_j}(z)|^{p_1} dV_{\alpha_1}(z)
\leq \lim_{j \to \infty} \int_{T_B} |f_{k_j}(z)|^{p_1} dV_{\alpha_1}(z)
\leq \lim_{j \to \infty} ||f_{k_j}||_{p_1,\alpha_1}^{p_1} \leq M.$$

Thus, we have

$$f \in A^{p_1}_{\alpha_1}(T_B).$$

Therefore, $f_{k_j} - f \to 0$ uniformly on every compact subset of T_B as $j \to \infty$.

Given our assumption, we have

$$\lim_{j \to \infty} \left\| T_{\mu}^{\xi} \left(f_{k_j} - f \right) \right\|_{p_2, \alpha_2} = 0.$$

It is straightforward to verify that

$$||T_{\mu}^{\xi} f_{k_j} - T_{\mu}^{\xi} f||_{p_2,\alpha_2} \le ||T_{\mu}^{\xi} (f_{k_j} - f)||_{p_2,\alpha_2}.$$

From this inequality, we obtain that

$$\lim_{j \to \infty} \| T_{\mu}^{\xi} f_{k_j} - T_{\mu}^{\xi} f \|_{p_2, \alpha_2} = 0,$$

implying that $T_{\mu}^{\xi} f \in L_{\alpha_2}^{p_2}(T_B)$.

This implies that a subsequence of $\{T_{\mu}^{\xi}f_{k_j}\}$ converges in $L_{\alpha_2}^{p_2}(T_B)$, stating the compactness of T_{μ}^{ξ} from $A_{\alpha_1}^{p_1}(T_B)$ to $L_{\alpha_2}^{p_2}(T_B)$.

The following Lemma 5.4 is a well-known result, and its proof can also be found in [12].

Lemma 5.4 Suppose $\{f_j\}$ is a sequence in $A^p_{\alpha}(T_B)$ with $1 . Then <math>f_j \to 0$ weakly in $A^p_{\alpha}(T_B)$ as $j \to \infty$ if and only if $\{f_j\}$ is bounded in $A^p_{\alpha}(T_B)$ and converges to 0 uniformly on each compact subset of T_B .

With the above lemmas 5.3 and 5.4, we can now prove the main theorem 5.2.

Theorem 5.2 Let $1 < p_1 \le p_2 < \infty$, $-1 < \alpha_1, \alpha_2, \xi < \infty$ and let $\mu \in \mathcal{M}_+$. Suppose that

$$n+1+\xi > n \max\left(1, \frac{1}{p_i}\right) + \frac{1+\alpha_i}{p_i}, \quad i = 1, 2.$$

Let

$$\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}, \ \gamma = \frac{1}{\lambda} \left(\xi + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right).$$

Then the following statements are equivalent:

- B^ξ_μ is compact from A^{p₁}_{α₁}(T_B) to L^{p₂}_{α₂}(T_B).
 T^ξ_μ is compact from A^{p₁}_{α₁}(T_B) to A^{p₂}_{α₂}(T_B).
 The measure μ is a vanishing (λ, γ)-Carleson measure.

Proof:

- $(1) \Rightarrow (2)$ The inequality $|T_{\mu}^{\gamma}f| \leq B_{\mu}^{\gamma}f$ implies that (2) holds trivially.
- $(2) \Rightarrow (3)$ Let T^{ξ}_{μ} be a compact linear map, and consider $\{a_k\} \in T_B$ such that $|a_k| \to \partial \widehat{T}_B$ as $k \to \infty$.

We consider the function

$$f_k(z) = \frac{\rho(z)^{(n+1+\xi)-(n+1+\alpha_1)/p_1}}{\rho(z, a_k)^{n+1+\xi}}.$$

By Lemma 3.4, we have $\sup_{k} ||f_{k}||_{p_{1},\alpha_{1}} < \infty$.

Let $Q_k = \overline{D(i,r)}$ and by Lemma 3.6, we know that f_k uniformly converges to 0 on the compact subset Q_k of T_B .

Since

$$\frac{\mu(D(a_k, r))}{\rho(a_k)^{(n+1+\gamma)\lambda}} \lesssim \rho(a_k)^{(n+1+\alpha_2)/p_2} \int_{T_B} \frac{\rho(a_k)^{(n+1+\xi)-(n+1+\alpha_1)/p_1}}{|\rho(w, a_k)|^{2(n+1+\xi)}} d\mu(w)
\simeq \rho(a_k)^{(n+1+\alpha_2)/p_2} T_{\mu}^{\xi} f_k(a_k),$$

by lemma 3.5 we obtain

$$T_{\mu}^{\xi} f_k(a_k) \lesssim \frac{1}{\rho(a_k)^{(n+1+\alpha_2)/p_2}} \left(\int_{D(a_k,r)} \left| T_{\mu}^{\xi} f_k(z) \right|^{p_2} dV_{\alpha_2}(z) \right)^{1/p_2}.$$

Since T^{ξ}_{μ} is compact, Lemma 5.4 implies that $f_k(z)$ weakly converges to 0.

Therefore, we obtain $||T_{\mu}^{\xi}f_{k}||_{p_{2},\alpha_{2}} \to 0$. Hence, we have

$$\mu\left(D\left(a_{k},r\right)\right)/\rho\left(a_{k}\right)^{(n+1+\gamma)\lambda}\to 0.$$

By Theorem 4.2, μ is a vanishing (λ, γ) -Carleson measure.

 $(3) \Rightarrow (1)$ Given the conditions, μ is a vanishing (λ, γ) -Carleson measure. According to Lemma 5.3, we only need to prove that for any bounded sequence $\{f_k\}$ in $A^{p_1}_{\alpha_1}(T_B)$ that uniformly converges to 0 on compact subsets of T_B , the fact $\|B^{\varepsilon}_{\mu}f_k\|_{p_2,\alpha_2} \to 0$ holds.

According to the proof of $(3) \Rightarrow (1)$ in Theorem 5.1, we establish the following equation:

$$||B_{\mu}^{\xi} f_{k}||_{p_{2},\alpha_{2}}^{p_{2}} \lesssim \sum_{j=1}^{\infty} \left(\int_{D(a_{j},2r)} |f_{k}(w)|^{p_{1}} dV_{\alpha_{1}}(w) \right)^{p_{2}/p_{1}} \left(\frac{\mu(D(a_{j},r))}{|\rho(a_{j})|^{(n+1+\gamma)\lambda}} \right)^{p_{2}}.$$

Since $p_2/p_1 \ge 1$ and $\{f_k\}$ is a bounded sequence in $A_{\alpha_1}^{p_1}(T_B)$, it follows that:

$$\sup_{k} \sum_{j=1}^{\infty} \left(\int_{D(a_{j},2r)} |f_{k}(w)|^{p_{1}} dV_{\alpha_{1}}(w) \right)^{p_{2}/p_{1}} \lesssim \sup_{k} ||f_{k}||_{p_{1},\alpha_{1}}^{p_{2}} < \infty.$$

Let $E_n = T_B - Q_n$ and $Q_n = \overline{D(\mathbf{i}, n)}$ be a compact subset in T_B . Since $\{f_k\}$ converges to 0 uniformly on compact subsets of T_B as $k \to \infty$, it follows that:

$$\lim_{k \to \infty} \sum_{j: a_j \in Q_n} \left(\int_{D(a_j, 2r)} |f_k(w)|^{p_1} dV_{\alpha_1}(w) \right)^{p_2/p_1} = 0.$$

According to Theorem 4.2, we have

$$\lim_{|a_j| \to \partial \widehat{T}_B} \frac{\mu\left(D\left(a_j, r\right)\right)}{\rho\left(a_j\right)^{(n+1+\alpha)\lambda}} = 0.$$

So, for any $\varepsilon > 0$, there is an $n_0 \in \mathbb{N}$, such that

$$\left| \frac{\mu\left(D\left(a_{j}, r\right)\right)}{\rho\left(a_{j}\right)^{(n+1+\alpha)\lambda}} \right| < \varepsilon$$

for all $a_j \in E_{n_0}$.

Therefore

$$\begin{split} \|B_{\mu}^{\xi} f_{k}\|_{p_{2},\alpha_{2}}^{p_{2}} &\leq \sum_{j:j \in Q_{n_{0}}} \left| \left(\int_{D(a_{j},2r)} |f_{k}\left(w\right)|^{p_{1}} dV_{\alpha_{1}}\left(w\right) \right)^{p_{2}/p_{1}} \left(\frac{\mu\left(D\left(a_{j},r\right)\right)}{|\rho\left(a_{j}\right)|^{(n+1+\gamma)\lambda}} \right)^{p_{2}} \right| \\ &+ \sum_{j:j \in E_{n_{0}}} \left| \left(\int_{D(a_{j},2r)} |f_{k}\left(w\right)|^{p_{1}} dV_{\alpha_{1}}\left(w\right) \right)^{p_{2}/p_{1}} \left(\frac{\mu\left(D\left(a_{j},r\right)\right)}{|\rho\left(a_{j}\right)|^{(n+1+\gamma)\lambda}} \right)^{p_{2}} \right| \\ &\leq \sup_{j:j \in Q_{n_{0}}} \left| \left(\frac{\mu\left(D\left(a_{j},r\right)\right)}{|\rho\left(a_{j}\right)|^{(n+1+\gamma)\lambda}} \right)^{p_{2}} \left| \sum_{j:j \in Q_{n_{0}}} \left| \left(\int_{D(a_{j},2r)} |f_{k}\left(w\right)|^{p_{1}} dV_{\alpha_{1}}\left(w\right) \right)^{p_{2}/p_{1}} \right| \\ &+ \varepsilon \sum_{j:j \in E_{n_{0}}} \left| \left(\int_{D(a_{j},2r)} |f_{k}\left(w\right)|^{p_{1}} dV_{\alpha_{1}}\left(w\right) \right)^{p_{2}/p_{1}} \right|. \end{split}$$

Letting $\varepsilon \to 0$ and then letting $k \to \infty$, we have the above the result. \Box

Finally, we used a specific method to prove the last main theorem 5.3 in the present paper. The method used in the following theorem was pioneered by Luecking and utilizes the Khinchine's inequality. The detailed description is as follows.

Define the Rademacher functions r_k by

$$r_0(t) = \begin{cases} 1 & 0 \le t - [t] < \frac{1}{2}, \\ -1 & \frac{1}{2} \le t - [t] < 1; \end{cases}$$
$$r_n(t) = r_0(2^n t), \quad n > 0.$$

Here [t] denotes the largest integer not greater than t.

Then Khinchine's inequality is the following.

For $0 there exist constants <math>0 < A_p \le B_p < \infty$ such that, for all natural numbers m and all complex numbers c_1, c_2, \ldots, c_m , we have

$$A_p \left(\sum_{j=1}^m |c_j|^2 \right)^{p/2} \le \int_0^1 \left| \sum_{j=1}^m c_j r_j(t) \right|^p dt \le B_p \left(\sum_{j=1}^m |c_j|^2 \right)^{p/2}.$$

Theorem 5.3 states that under the assumption of $p_2 < p_1$, the boundedness and compactness of Toeplitz operators are unified, and a connection with an integrable function is established.

Theorem 5.3 Let $0 < p_2 < p_1 < \infty$, $-1 < \alpha_1, \alpha_2, \xi < \infty$ and let $\mu \in \mathcal{M}_+$. Suppose that

$$n+1+\xi > n \max\left(1, \frac{1}{p_i}\right) + \frac{1+\alpha_i}{p_i}, \quad i = 1, 2.$$

Let

$$\lambda = 1 + \frac{1}{p_1} - \frac{1}{p_2}, \ \gamma = \frac{1}{\lambda} \left(\xi + \frac{\alpha_1}{p_1} - \frac{\alpha_2}{p_2} \right).$$

Then the following statements are equivalent:

$$\begin{array}{l} (1)\ T^{\xi}_{\mu}\ is\ bounded\ from\ A^{p_1}_{\alpha_1}(T_B)\ to\ A^{p_2}_{\alpha_2}(T_B).\\ (2)\ \left\{\frac{\mu(D(a_k,r))}{\rho(a_k)^{(n+1+\gamma)\lambda}}\right\}\in l^{1/(1-\lambda)}\ for\ any\ r>0\ and\ for\ any\ r\text{-lattice}\ \{a_k\}. \end{array}$$

(3) T^{ξ}_{μ} is compact from $A^{p_1}_{\alpha_1}(T_B)$ to $A^{p_2}_{\alpha_2}(T_B)$.

Proof: (1) \Rightarrow (2) Let $r_k(t)$ be a sequence of Rademacher functions and $\{a_k\}$ be an r-lattice on T_B .

Since

$$n+1+\xi > n \max\left(1, \frac{1}{p_1}\right) + \frac{1+\alpha_1}{p_1},$$

it follows from [4, Theorem 2] that for any sequence $\{\lambda_k\} \in l^{p_1}$, the function

$$f_t(z) = \sum_{k=1}^{\infty} \lambda_k r_k(t) \frac{\rho(a_k)^{n+1+\xi-(n+1+\alpha_1)/p_1}}{\rho(z, a_k)^{n+1+\xi}}$$

is in $A_{\alpha_1}^{p_1}(T_B)$ with $||f_t||_{p,\alpha_1} \lesssim ||\{\lambda_k\}||_{l^p}$ for almost every $t \in (0,1)$. Denote

$$f_k(z) = \frac{\rho (a_k)^{n+1+\xi - (n+1+\alpha_1)/p_1}}{\rho (z, a_k)^{n+1+\xi}}.$$

Since T^{ξ}_{μ} is bounded from $A^{p_1}_{\alpha_1}(T_B)$ to $A^{p_2}_{\alpha_2}(T_B)$.

it follows that

$$||T_{\mu}^{\xi} f_{t}||_{p_{2},\alpha_{2}}^{p_{2}} = \int_{T_{B}} \left| \sum_{k=1}^{\infty} \lambda_{k} r_{k}(t) T_{\mu}^{\xi} f_{k}(z) \right|^{p_{2}} dV_{\alpha_{2}}(z)$$

$$\lesssim ||T_{\mu}^{\xi}||^{p_{2}} ||f_{t}||_{p_{2},\alpha_{2}}^{p_{2}} \lesssim ||T_{\mu}^{\xi}||^{p_{2}} \left(\sum_{k=1}^{\infty} |\lambda_{k}|^{p_{1}} \right)^{p_{2}/p_{1}}.$$

Integrating both sides with respect to t from 0 to 1, then using Fubini's theorem and Khinchine's inequality, we obtain

$$\int_{T_{B}} \left(\sum_{k=1}^{\infty} \left| \lambda_{k} \right|^{2} \left| T_{\mu}^{\xi} f_{k}\left(z\right) \right|^{2} \right)^{p_{2}/2} dV_{\alpha_{2}}\left(z\right) \lesssim \|T_{\mu}^{\xi}\|^{p_{2}} \left(\sum_{k=1}^{\infty} \left| \lambda_{k} \right|^{p_{1}} \right)^{p_{2}/p_{1}}.$$

Through the Hölder inequality and Lemma 3.1, we have

$$\sum_{k=1}^{\infty} |\lambda_{k}|^{p_{2}} \int_{D(a_{k},2r)} |T_{\mu}^{\xi} f_{k}(z)|^{p_{2}} dV_{\alpha_{2}}(z)
\leq \max \left\{ 1, N^{1-p_{2}/2} \right\} \int_{T_{B}} \left(\sum_{k=1}^{\infty} |\lambda_{k}|^{2} |T_{\mu}^{\xi} f_{k}(z)|^{2} \right)^{p_{2}/2} dV_{\alpha_{2}}(z)
\lesssim ||T_{\mu}^{\xi}||^{p_{2}} \left(\sum_{k=1}^{\infty} |\lambda_{k}|^{p_{1}} \right)^{p_{2}/p_{1}} .$$

In the proof of that $(2) \Rightarrow (3)$ in Theorem 5.2, we establish the following equation:

$$\frac{\mu\left(D\left(a_{k},r\right)\right)}{\rho\left(a_{k}\right)^{(n+1+\gamma)\lambda}} \simeq \rho\left(a_{k}\right)^{(n+1+\alpha_{2})/p_{2}} T_{\mu}^{\xi} f_{k}\left(a_{k}\right).$$

Based on Lemma 3.5, we obtain

$$\sum_{k=1}^{\infty} |\lambda_k|^{p_2} \frac{\mu(D(a_k, r))}{\rho(a_k)^{(n+1+\gamma)\lambda}} \lesssim ||T_{\mu}^{\xi}||^{p_2} \left(\sum_{k=1}^{\infty} |\lambda_k|^{p_1}\right)^{p_2/p_1}.$$

Then the duality yields that

$$\left\{\frac{\mu\left(D\left(a_{k},r\right)\right)}{\rho\left(a_{k}\right)^{(n+1+\gamma)\lambda}}\right\} \in l^{1/(1-\lambda)}.$$

(2) \Rightarrow (3) In order to prove that T_{μ}^{ξ} is a compact mapping, according to Lemma 5.3, it suffices to show that for any bounded sequence $\{f_k\}$ in $A_{\alpha_1}^{p_1}(T_B)$ that uniformly converges to 0 on any compact subset of T_B , the property $\|T_{\mu}^{\xi}f_k\|_{p_2,\alpha_2} \to 0$ holds.

Similar to the proof of $(3) \Rightarrow (1)$ in Theorem 5.1, we establish the following equation

$$||T_{\mu}^{\xi} f_{k}||_{p_{2},\alpha_{2}}^{p_{2}} \lesssim \sum_{j=1}^{\infty} \left(\int_{D(a_{j},2r)} |f_{k}(w)|^{p_{1}} dV_{\alpha_{1}}(w) \right)^{p_{2}/p_{1}} \left(\frac{\mu(D(a_{j},r))}{|\rho(a_{j})|^{(n+1+\gamma)\lambda}} \right)^{p_{2}}.$$

Given the known conditions, we deduce that

$$\left\{ \frac{\mu\left(D\left(a_{k},r\right)\right)}{\rho\left(a_{k}\right)^{(n+1+\gamma)\lambda}} \right\} \in l^{1/(1-\lambda)}.$$

Denote by

$$A = \frac{\mu(D(a_k, r))}{\rho(a_k)^{(n+1+\gamma)\lambda}}, \quad B_k = \int_{D(a_j, 2r)} |f_k(w)|^{p_1} dV_{\alpha_1}(w), \quad E_k = T_B - Q_k,$$

Where $Q_k = \overline{D(\mathbf{i}, k)}$.

Since $A \in l^{1/(1-\lambda)}$, it follows that, for any $\varepsilon > 0$, there is an $k_0 \in \mathbb{N}$, such that

$$\sum_{j: a_j \in E_{k_0}} |A|^{1/(1-\lambda)} < \varepsilon.$$

Since Q_{k_0} is a compact subsets of T_B , so we have

$$\lim_{k \to \infty} \sum_{j: j \in Q_{k_0}} |B_k| = 0.$$

On the other hand, since $p_1 > p_2$, by Hölder inequality, we have that

$$\sum_{j=1}^{\infty} A^{p_2} B_k^{p_2/p_1} \leq \sum_{j:j \in Q_{k_0}} |A|^{p_2} |B_k|^{p_2/p_1} + \sum_{j:a_j \in E_{k_0}} |A|^{p_2} |B_k|^{p_2/p_1}$$

$$\leq \left(\sum_{j:j \in Q_{k_0}} |A|^{1/(1-\lambda)}\right)^{1-p_2/p_1} \left(\sum_{j:j \in Q_{k_0}} |B_k|\right)^{p_2/p_1}$$

$$+ \left(\sum_{j:j \in E_{k_0}} |A|^{1/(1-\lambda)}\right)^{1-p_2/p_1} \left(\sum_{j:j \in E_{k_0}} |B_k|\right)^{p_2/p_1}$$

$$\leq \left(\sum_{j:j \in Q_{k_0}} |A|^{1/(1-\lambda)}\right)^{1-p_2/p_1} \left(\sum_{j:j \in Q_{k_0}} |B_k|\right)^{p_2/p_1}$$

$$+ \varepsilon^{1-p_2/p_1} \left(\sum_{j:j \in E_{k_0}} |B_k|\right)^{p_2/p_1}.$$

Letting $\varepsilon \to 0$ and then letting $k \to \infty$, we obtain $||T_{\mu}^{\xi} f_k||_{p_2,\alpha_2} \to 0$.

 $(3) \Rightarrow (1)$ Since compact operators are bounded operators, so the result holds trivially.

References

- [1] M. Abate, S. Mongodi and J. Raissy, Toeplitz operators and skew carleson measures for weighted bergman spaces on strongly pseudoconvex domains, *J. Operat. Theor.* **84** (2020), 339-364.
- [2] M. Abate, J. Raissy and A. Saracco, Toeplitz operators and carleson measures in strongly pseudoconvex domains, J. Funct. Anal. 263 (2012), 3449–3491.
- [3] M. Abate and A. Saracco, Carleson measures and uniformly discrete sequences in strongly pseudoconvex domains, *J. Lond. Math. Soc.-Second Ser.* **83** (2011), 587–605.
- [4] R.R. Coifman and R. Rochberg, Representation theorems for holomorphic and harmonic functions in L^p , Astérisque. **77** (1980), 11-66.
- [5] J. Cima and W. Wogen. A carleson measure theorem for the bergman space on the ball. J. Operat. Theor. 7 (1982), 157–165.
- [6] L. Carleson, An interpolation problem for bounded analytic functions, Am. J. Math. 80 (1958), 921–930.
- [7] G.T. Deng, Y. Huang and T. Qian. Reproducing kernels of some weighted bergman spaces, J. Geom. Anal. 31 (2021), 9527–9550.
- [8] W.W. Hastings, A carleson measure theorem for bergman spaces, *Proc. Amer. Math. Soc.* **52** (1975), 237-241.
- [9] S.G. Krantz, Function theory of several complex variables, volume 340, American Mathematical Soc, 2001.
- [10] D. Luecking. A technique for characterizing carleson measures on bergman spaces. *Proc. Amer. Math. Soc.* **87** (1983), 656–660.
- [11] H.C. Li, J.S. Liu and H.Y. Wang, Carleson measures on convex domains with smooth boundary of finite type. *Math. Nachr.* **297** (2024), 694-706.
- [12] L.C. Li, J.Q. Ding and H.C. Li, Toeplitz Operators on Weighted Bergman Spaces over Tubular Domains, preprint (2024), https://arxiv.org/abs/2404.16439.
- [13] J.X. Liu, G.T. Deng and H.H. Yin, Bergman metric on a class of tubular domains, *Beijing Shifan Daxue Xuebao* **59** (2023), 353–357.
- [14] G.T. Prăjitură, R.H. Zhao and L.F. Zhou. On berezin type operators and toeplitz operators on bergman spaces, *Banach J. Math. Anal.* (2023), DOI:10.1007/s43037-023-00280-3.

- [15] J. Pau and R.H. Zhao, Carleson measures and toeplitz operators for weighted bergman spaces on the unit ball, *Michigan Math. J.* **64** (2015), 759–796.
- [16] R. Peng and Y.Q. Fan, Toeplitz operators from hardy spaces to weighted bergman spaces in the unit ball of \mathbb{C}^n . Acta Math. Sci. 42 (2022), 349–363.
- [17] J.J. Si, Y. Zhang and L.F. Zhou, Carleson measures and toeplitz operators between bergman spaces on the siegel upper half-space. *Complex Anal. Oper. Theory* (2022), DOI:10.1007/s11785-022-01214-x.
- [18] K.H. Zhu, Spaces of holomorphic functions in the unit ball, volume 226, Springer, 2005.
- [19] K.H. Zhu, Operator theory in function spaces, Number 138, American Mathematical Soc, 2007.
- [20] K.H. Zhu, The Berezin Transform and Its Applications, *Acta Math Sci.* **41** (2021), 1839–1858.