A note on the topological synchronization of unimodal maps

Michele Gianfelice Dipartimento di Matematica e Informatica Università della Calabria Campus di Arcavacata Ponte P. Bucci - cubo 30B I-87036 Arcavacata di Rende gianfelice@mat.unical.it

Abstract

In this note we complete the analysis carried on in [CGSV] about the topological synchronisation of unimodal maps of the interval coupled in a master-slave configuration, by answering to the questions raised in that paper. Namely, we compute the weak limits of the invariant measure of the coupled system as the coupling strength $k \in (0, 1)$ tends to 0 and to 1 and discuss the uniqueness of the invariant measure of its random dynamical system counterpart.

Contents

1 Introduction

2 Results

2.1 On the uniqueness of the stationary measure of the MC analogue to the slave system 4

2.2 The weak limits of the invariant measures of the deterministic system in the very weak and in t

 $\mathbf{2}$

4

AMS subject classification: 37A10, 60J10.

Keywords and phrases: coupled dynamical systems, unimodal maps, master-slave system, Markov chains, Random Dynamical Systems, Topological synchronisation.

1 Introduction

In [CGSV] it has been studied the asymptotic behaviour of the trajectories of the components of a dynamical system realized by coupling two unimodal map of the interval in a master/slave configuration. Namely, denoting by I the interval [-1, 1],

$$I^{2} \ni (x_{n}, y_{n}^{(k)}) \longmapsto \begin{cases} x_{n+1} = T_{1}(x_{n}) \\ y_{n+1}^{(k)} = (1-k)T_{2}(y_{n}) + kT_{1}(x_{n}) \end{cases} \in I^{2},$$
(1)

where $k \in [0, 1]$.

A system of this type were introduced in the physics literature [Lahav et al.] as a simple example of a dynamical system undergoing to the so called *topological synchronisation*, which refers to the numerically observed phenomenon, in asymmetrically coupled dynamical systems, of the slave component attractor to become very similar to the master's one for sufficiently large values of the coupling. In particular, to gauge the emergence this phenomenon the authors of [Lahav et al.] suggested to compare the spectrum of the generalized dimensions of the empirical measures of the slave component $\left(D_q\left(\nu_n^{(k)}\right), \ q \in \mathbb{R}\right)$, where $\nu_n^{(k)} := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{y_i^{(k)}}$ and

$$D_{q}\left(\nu_{n}^{(k)}\right) := \begin{cases} \frac{1}{q-1} \lim_{r \downarrow 0} \frac{\log \int_{\Sigma} \nu_{n}^{(k)}(dx) \left[\nu_{n}^{(k)}(B_{r}(x))\right]^{q-1}}{\log r} & q \neq 1\\ \lim_{r \downarrow 0} \frac{\int_{\Sigma} \nu_{n}^{(k)}(dx) \log \nu_{n}^{(k)}(B_{r}(x))}{\log r} & q = 1 \end{cases},$$
(2)

provided the limits exist, with that of the master one $\mu_n := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{x_i}$, for very large values of n, as the coupling increases. In particular, in the case of a system given in (1) where the coupled maps belong to the logistic family, they observed numerically that, for very large values of n, the difference $\Delta D_q^{(k)} := \left| D_q \left(\nu_n^{(k)} \right) - D_q \left(\mu_n \right) \right|$ converges to zero as the coupling parameter k approaches to 1 first for negative values of q. This lead them to deduce that, assuming n sufficiently large such that $\rho_n^{(k)} := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{\left(x_n, y_i^{(k)}\right)}$ is close to one of its possible weak limits with larger basin of attraction $\rho^{(k)}$ (the physical measure if it exists), as the coupling increases, the synchronisation of the trajectories of slave component with those of the master one would start first in the regions of the configurations space of the coupled system where the invariant measure $\rho^{(k)}$ has low density.

In [CGSV] it has been shown that the topological synchronization phenomenon can easily be interpreted as the emergence of an invariant set in a neighbourhood of the diagonal of I^2 to which the trajectories of the system components converge in the strong coupling limit. Moreover, it has been shown that there is a large set of maps, in the family of those of quadratic type used in [Lahav et al.], such that the physical measure μ of T_1 has density bounded away from zero and consequently $D_q(\mu)$ equal to 1 for all negative values of q. Therefore, the numerical evidence that $\Delta D_q^{(k)}$ can be equal to zero for such values of the parameter q even for values of the coupling large but bounded away from 1 can be explained by the fact that for these values of k also the weak limit $\nu^{(k)}$ of $\nu_n^{(k)}$ has a density bounded away from zero and so $D_q(\nu^{(k)}) = 1$ for q < 0.

Furthermore, in the case, T_1 is exponentially mixing w.r.t. its invariant measure μ , the asymptotic properties of the evolution of the slave component of (1) has been compared with those of the Markov chain (MC) $\left\{Y_n^{(k)}\right\}_{n\geq 0}$ such that $Y_{n+1}^{(k)} = (1-k)T_2\left(Y_n^{(k)}\right) + k\omega_n$, where $\{\omega_n\}_{n\geq 1}$ is a sequence of i.i.d. random variables sampled according to μ . In particular, if μ is a.c. w.r.t. the Lebesgue measure, which in the following we will denote by λ , it has been shown that, for any $k \in [0, 1]$, the stationary measures of $\left\{Y_n^{(k)}\right\}_{n\geq 0}$ are a.c. w.r.t. λ and there are at most finitely many ergodic probability measures. It has also been shown that any stationary measure weakly converges to μ as k tends to 1 and, as k tends to 0, to a probability measure invariant under the evolution defined by T_2 .

In this note we complete the analysis just presented on the coupled dynamical system (1) answering to the following questions that were left open [CGSV].

First, in the case T_1 and T_2 are continuous maps, for any $k \in (0, 1)$, we can construct an invariant measure $\rho^{(k)}$ for the skew-product system defined by (1). Considering the random vector (η_1, η_2) on I^2 distributed according to $\rho^{(k)}$, where μ is the marginal distribution of the first component representing the master system, and denoting by $\rho^{(k)}(\cdot|\eta_1)$ the conditional distribution of the r.v. η_2 representing the slave component, w.r.t. η_1 a natural question arose; denoting by $\rho^{(k)}(\cdot|x)$ the regular version of $\rho^{(k)}(\cdot|\eta_1)$,

Question I: Does $\{\rho^{(k)}(\cdot|x), k \in [0,1]\}$ converges weakly, when $k \to 1$, to μ for $\mu - a.e. x$?

Second, assuming that the invariant measure μ of T_1 has density h and considering the MC $\left\{Y_n^{(k)}\right\}_{n\geq 0}$ described above one may wonder about the uniqueness of the stationary measure. More precisely, for any $k \in [0, 1]$, denoting by $p_k(y, z)$ the transition probability kernel of the MC and by L_k the associated Kolmogorov forward operator

$$L^{1}(\lambda) \ni g \longmapsto (L_{k}g)(z) := \int_{I} dyg(y)p_{k}(y,z) \in L^{1}(\lambda) , \qquad (3)$$

one can ask:

Question II: Is there a unique solution to $L_k g^{(k)} = g^{(k)}$, for any $k \in (0, 1)$?

Since numerical simulations ([CGSV] fig. 3A) show that if μ is a.c. w.r.t. the Lebesgue measure then, for sufficiently large values of k, the weak limit as $n \uparrow \infty$ of the empirical measure $\nu_n^{(k)}$ is also a.c. w.r.t. the Lebesgue measure, it is reasonable to ask if, in the case Question II has a positive answer at least for k sufficiently large, a stronger notion of convergence to μ than the weak convergence as k tends to 1 of $\limsup_{n\to\infty} \nu_n^{(k)}$ proven in [CGSV] Proposition 3.1 is in force. More precisely,

Question III: Assume that for any $k \in [0, 1]$ the solution of (3) is unique. Denoting by $\Delta_n^{(k)}$ the total variation distance between $\nu_n^{(k)}$ and $\bar{\nu}_n^{(k)} := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{Y_i^{(k)}}$, do we have

$$\limsup_{k \to 1} \limsup_{n \to \infty} \Delta_n^{(k)} = 0 , \qquad (4)$$

for Lebesgue almost all choices of the initial data x_0, y_0 and \mathbb{P} almost all ω ?

In the next section we will first answer to the second question in the case the slave map T_2 is of the form considered in [Lahav et al.] and [CGSV], then turn to the answer of Question I and deduce from this the answer in the negative to the third question.

2 Results

2.1 On the uniqueness of the stationary measure of the MC analogue to the slave system

Let us assume that μ , the invariant measure of the master map T_1 , has density h whose support is strictly contained in I, namely T_1 can be chosen to satisfy the hypothesis of Proposition 2.7 in [BS] (see Proposition 5.1 in [CGSV]), and consider the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that $\Omega := I^{\mathbb{N}}, \mathcal{F}$ is the σ -algebra generated by the cylinder sets

$$\{\omega \in \Omega : (\omega_1, ..., \omega_n) \in B\}, \ n \in \mathbb{N}, \ B \in \mathcal{B}(I^n) \ , \tag{5}$$

where $\mathcal{B}(I^n)$ is the Borel σ -field on I^n , and $\mathbb{P} := \mu^{\otimes \mathbb{N}}$.

In [CGSV], for any $k \in [0, 1]$, we considered the Random Dynamical System (RDS) defined by the skew-product

$$I \times \Omega \ni (y, \omega) \longmapsto \Psi_k(y, \omega) := (F_{\pi\omega}^{(k)}(y), S\omega) \in I \times \Omega , \qquad (6)$$

where $\pi: \Omega \to I$ is such that $\pi \omega = \omega_1, S: \Omega \to \Omega$ is the left shift, so that $\forall n \ge 1, \omega_n = \pi S^n \omega$, and

$$I \ni y \longmapsto F_{\pi\omega}^{(k)}(y) := (1-k)T_2(y) + k\omega_1 \in I , \qquad (7)$$

with

$$I \ni y \longmapsto T_2(y) := c_2(1 - 2x^2) \in I , \ c_2 \in (0, 1)$$
 (8)

Denoting by $F_j^{(k)} := F_{\pi S^j \omega}^{(k)}, j \ge 1$, the sequence $\left\{Y_n^{(k)}\right\}_{n\ge 0}$ such that $Y_0^{(k)}$ is a r.v. and, for any $n \ge 0, Y_{n+1}^{(k)} = F_{n+1}^{(k)}(Y_n^{(k)})$, is a homogeneous MC (see e.g. [Ki] or [Ar]) whose transition probability kernel, computed in [CGSV] section 6.2, is

$$p_k(y,z)dz = \frac{1}{k}h\left(\frac{z - (1-k)T_2(y)}{k}\right)\mathbf{1}_{[-1,1]}\left(\frac{z - (1-k)T_2(y)}{k}\right)dz .$$
(9)

As already remarked in [CGSV], if T_1 mixes exponentially w.r.t. μ , the asymptotic properties of the evolution of the slave component of (1) can be compared with those of the MC $\left\{Y_n^{(k)}\right\}_{n\geq 0}$. From Proposition 6.1 there are at most finitely many stationary measures of the MC

From Proposition 6.1 there are at most finitely many stationary measures of the MC $\left\{Y_n^{(k)}\right\}_{n\geq 0}$ and their densities are fixed points of (3). Here we prove that in fact there is just one, which provide a positive answer to Question II. The proof rely on the existence of a Lyapunov function for the MC $\left\{Y_n^{(k)}\right\}_{n\geq 0}$, for $k \in (0,1)$, that is a measurable function $V_k : \mathbb{R} \longrightarrow \mathbb{R}_+$ such that $\lim_{|x|\uparrow+\infty} V_k(x) = +\infty$, satisfying

$$\int_{\mathbb{R}} dz p_k(y, z) \, V_k(z) \le \gamma_k V_k(y) + K_k \, , \, \gamma_k \in [0, 1), \, K_k \ge 0 \, . \tag{10}$$

Lemma 1 For any $k \in (0, 1)$ the function

$$\mathbb{R} \ni x \longmapsto V_k(x) = \mathbf{1}_I(x) \cosh\left(c_2^{\frac{1}{k}} \left(1 - c_2^2\right) \frac{x}{1 - x^2}\right) \in [1, +\infty)$$
(11)

is a Lyapunov function for the MC $\left\{Y_n^{(k)}\right\}_{n\geq 0}$.

Proof. Setting $x = \frac{z - (1-k)T_2(y)}{k}$, by the convexity of V_k ,

$$\int_{I} dz p_{k}(y, z) \cosh\left(c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{z}{1-z^{2}}\right) = \int_{I} dz \frac{1}{k}h\left(\frac{z-(1-k)T_{2}(y)}{k}\right) \times$$
(12)

$$\times \mathbf{1}_{I}\left(\frac{z-(1-k)T_{2}(y)}{k}\right) \cosh\left(c_{2}^{\frac{1}{k}}\frac{z}{1-z^{2}}\right)$$

$$= \int_{I} dxh(x) \cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{(kx+(1-k)T_{2}(y))}{1-(kx+(1-k)T_{2}(y))^{2}}\right]$$

$$\leq \int_{I} dxh(x) \left\{k\cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\left(\frac{x}{1-x^{2}}\right)\right] + (1-k)\cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{T_{2}(y)}{1-(T_{2}(y))^{2}}\right]\right\}$$

$$\leq (1-k)\cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{T_{2}(y)}{1-(T_{2}(y))^{2}}\right] + k\int_{I} dxh(x)\cosh\left(c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{x}{1-x^{2}}\right)$$

$$\leq (1-k)\cosh\left(c_{2}^{\frac{1}{k}+1}\right)\cosh\left(c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{y}{1-y^{2}}\right) + k\int_{I} dxh(x)\cosh\left(c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{x}{1-x^{2}}\right)$$

where we used that, since $T_2(I) \in [-c_2, c_2]$,

$$\sup_{y \in I} \frac{\cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{T_{2}(y)}{1-(T_{2}(y))^{2}}\right]}{\cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{y}{1-y^{2}}\right]} \leq \frac{\sup_{y \in I} \cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{T_{2}(y)}{1-(T_{2}(y))^{2}}\right]}{\inf_{y \in I} \cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\frac{y}{1-y^{2}}\right]} \qquad (13)$$

$$\leq \cosh\left[c_{2}^{\frac{1}{k}}\left(1-c_{2}^{2}\right)\sup_{y \in I}\frac{T_{2}\left(y\right)}{1-(T_{2}\left(y\right))^{2}}\right]$$

$$\leq \cosh\left(c_{2}^{\frac{1}{k}+1}\right).$$

Hence, (10) holds with $\gamma_k := (1-k) \cosh\left(c_2^{\frac{1}{k}+1}\right) < 1$ and $K_k := k \int_I dx h(x) \cosh\left(c_2^{\frac{1}{k}} \frac{x}{1-x^2}\right)$ which is finite because the support of h is strictly contained in I.

Proposition 2 For any $k \in (0, 1)$ there exists a unique solution to (3).

Proof. The thesis will follow from Theorem 5.6.1 in [LM] once we have shown that L_k is constrictive and that exists a set $A \subseteq I$ with positive Lebesgue measure such that for any density f, there exists $n_0(f)$ such that $\forall n > n_0(f)$, $(L_k^n f)(x) > 0$ for Lebesgue almost all $x \in A$.

By Proposition 6.1 in [CGSV] L_k is quasi-compact hence, by the theorem in [Ba], uniformly constrictive and therefore, by definition, constrictive (see also [BNNT] fig. 1). Moreover, denoting by $\left\{ \mathcal{F}_m^{(k)} \right\}_{m \ge 0}$ the natural filtration generated by the Markov chain $\left\{ Y_m^{(k)} \right\}_{m \ge 0}, \forall n \ge 1, k \in (0, 1), \text{ by (10)},$ $\mathbb{E} \left[V_k \left(Y_n^{(k)} \right) \middle| Y_0^{(k)} = y \right] = \mathbb{E} \left[\mathbb{E} \left[V_k \left(Y_n^{(k)} \right) \middle| \mathcal{F}_{n-1}^{(k)} \right] \middle| Y_0^{(k)} = y \right]$ (14) $= \mathbb{E} \left[\mathbb{E} \left[V_k \left(Y_{n+1}^{(k)} \right) \middle| Y_0^{(k)} = y \right]$ $\leq \gamma_k \mathbb{E} \left[V_k \left(Y_{n-1}^{(k)} \right) \middle| Y_0^{(k)} = y \right] + K_k.$

Iterating the previous inequality we get

$$\mathbb{E}\left[V_k\left(Y_n^{(k)}\right)\middle|Y_0^{(k)} = y\right] \le K_k \frac{1 - \gamma_k^{n+1}}{1 - \gamma_k} + \gamma_k^n V_k\left(y\right) .$$
(15)

By the Markov inequality, $\forall R > 0$,

$$\mathbb{E}\left[V_k\left(Y_n^{(k)}\right)\middle|Y_0^{(k)} = y\right] \ge R\mathbb{E}\left[\mathbf{1}_{\{V_k > R\}}\left(Y_n^{(k)}\right)\middle|Y_0^{(k)} = y\right]$$
(16)

On the other hand, defining, for any $n \ge 1$ and any bounded Lebesgue measurable function ϕ ,

$$((L_k^*)^n \phi)(y) =: \int_I dz p_k^n(y, z) \phi(z) , \qquad (17)$$

(16) reads

$$((L_k^*)^n V_k)(y) \ge R\left((L_k^*)^n \mathbf{1}_{\{V_k > R\}}\right)(y) .$$
(18)

Consequently, for any probability density f supported on I,

$$\int_{I} dx \left(L_{k}^{n} f \right) \mathbf{1}_{\{V_{k} > R\}} (x) = \int_{I} dy f (y) \left((L_{k}^{*})^{n} \mathbf{1}_{\{V_{k} > R\}} \right) (y)$$

$$\leq \frac{1}{R} \int_{I} dy f (y) \left((L_{k}^{*})^{n} V_{k} \right) (y) = \frac{1}{R} \int_{I} dx \left(L_{k}^{n} f \right) V_{k} (x)$$
(19)

and, by (15),

$$\int_{I} dx \left(L_{k}^{n} f \right) \mathbf{1}_{\{V_{k} > R\}} (x) \leq \frac{1}{R} \int_{I} dx \left(L_{k}^{n} f \right) V_{k} (x)$$

$$\leq \frac{1}{R} \left(K_{k} \frac{1 - \gamma_{k}^{n+1}}{1 - \gamma_{k}} + \gamma_{k}^{n} \int_{I} dy f(y) V_{k}(y) \right) .$$
(20)

Denoting by D the set of probability density, let $D_0 := \{f \in D : \int_I dx f(x) V_k(x) < \infty\}$. Given $g \in D_0$, let $n_0 \ge 1$ be such that $\gamma_k^{n_0} \int_I dyg(y) V_k(y) < 1$. Then, by (20), for any $n \ge n_0$,

$$\int_{I} dx \left(L_{k}^{n} g \right) \mathbf{1}_{\{V_{k} > R\}} \left(x \right) \leq \frac{1}{R} \left(\frac{K_{k}}{1 - \gamma_{k}} + 1 \right)$$

$$(21)$$

and so, for $R > \left(\frac{K_k}{1-\gamma_k} + 1\right)$,

$$\int_{I} dx \left(L_{k}^{n} g \right) \mathbf{1}_{\{V_{k} \le R\}} \left(x \right) \ge 1 - \frac{1}{R} \left(\frac{K_{k}}{1 - \gamma_{k}} + 1 \right) > 0 .$$
(22)

Since $V_k \in C(I)$, D_0 is dense in D; therefore, setting $\epsilon := 1 - \frac{1}{R} \left(\frac{K_k}{1 - \gamma_k} + 1 \right)$, for any $f \in D$ there exists $g \in D_0$ such that $\|f - g\|_{L^1(dx)} \leq \frac{\epsilon}{2}$, which implies that, $\forall n \geq n_0$,

$$\int_{I} dx \left(L_{k}^{n} f \right) \mathbf{1}_{\{V_{k} \leq R\}} \left(x \right) = \int_{I} dx \left(L_{k}^{n} g \right) \mathbf{1}_{\{V_{k} \leq R\}} \left(x \right) + \int_{I} dx \left(L_{k}^{n} \left(f - g \right) \right) \mathbf{1}_{\{V_{k} \leq R\}} \left(x \right) \quad (23)$$

$$\geq \epsilon - \| f - g \|_{L^{1}(dx)} \geq \frac{\epsilon}{2} ,$$

but, because $(L_k^n f) \ge 0$, there exists $A \subseteq \{x \in I : V_k(x) \le R\}$ of positive Lebesgue measure such that $(L_k^n f)(x) > 0$ for Lebesgue almost all $x \in A$.

We stress that Proposition 2 do not give back any estimate on the rate of convergence the Markov chain $\{Y_n^{(k)}\}_{n\geq 0}$ to its unique stationary measure, what instead the next result will provide to the price of proving uniqueness only for values of k larger that a given one depending on T_1 and T_2 under the assumption that there exists $f \in D$ such that h > f. Although the proof of the following result it is easily seen to hold in this more general case, we will present it for the case where T_1 satisfies the hypothesis of Proposition 2.7 in [BS], which, as shown in [CGSV], can have its own interest.

Proposition 3 Let $c_2 := \max_{x \in I} T_2(x)$. There exists $k_* := k_*(T_1, T_2)$ such that, for any $k > k_* \in [0, 1)$, the Markov chain $\{Y_n^{(k)}\}_{n\geq 0}$ admits a unique stationary probability density $g^{(k)}$ such that, for any probability density f, denoting by L_k^n the Kolmogorov forward operator defined by the n-fold composition of the transition probability kernel p_k given in (9), the sequence $\{L_k^n f\}_{n\geq 1}$ converges geometrically to $g^{(k)}$ in $L^1(\lambda)$ at rate $(1-k)\cosh c_2^{\frac{1}{k}}$.

Proof. The proof follows from Proposition 6.1 in [CGSV] Theorem 1.2 in [HM] once we have shown that the transition probability kernel p_k of the Markov chain $\left\{Y_n^{(k)}\right\}_{n\geq 0}$ satisfies the assumptions 1 and 2 given in that paper. Namely:

- 1. there exists a Lyapunov function V_k for the MC $\left\{Y_n^{(k)}\right\}_{n\geq 0}$ such that (10) holds;
- 2. the transition probability kernel of the MC $\{Y_n^{(k)}\}_{n\geq 0}$ satisfies a Doeblin's type condition on a subset of I of the form $\{x \in I : V_k(x) \leq R\} =: \mathcal{V}_k(R)$ for sufficiently large R. More specifically, there exist a probability measure $\tilde{\nu}_k$ on $(I, \mathcal{B}(I))$ and $\alpha_k \in (0, 1)$ such that

$$\inf_{y \in \mathcal{V}_k(R)} p_k\left(y, A\right) \ge \alpha_k \tilde{\nu}\left(A\right) \ , \ R > \frac{2K_k}{1 - \gamma_k} \ , \ A \in \mathcal{B}\left(I\right) \ , \tag{24}$$

where the constants γ_k and K_k are those appearing in (10).

Assumption 1 is satisfied by Lemma 1, therefore we are left with the proof of Assumption 2.

Let ψ_0 be the non negative $C^1(I)$ function appearing in the representation of h given in Proposition 2.7 formula (11) of [BS]. Then, $h \ge \psi_0$ and, for any $[a, b) \subseteq I$,

$$\int_{I} dz p_{k}(y, z) \mathbf{1}_{[a,b)}(z) = \int_{I} dz \frac{1}{k} h\left(\frac{z - (1 - k)T_{2}(y)}{k}\right) \mathbf{1}_{I}\left(\frac{z - (1 - k)T_{2}(y)}{k}\right) \mathbf{1}_{[a,b)}(z)$$
(25)
$$= \int_{I} dx h(x) \mathbf{1}_{[a,b)}(kx + (1 - k)T_{2}(y))$$
$$\geq \int_{I} dx \psi_{0}(x) \mathbf{1}_{[a,b)}(kx + (1 - k)T_{2}(y)) .$$

Denoting by $[a_0, b_0]$ the support of ψ_0 it follows that $[a_0, b_0] \subseteq \text{supp} h \subset I^1$. Moreover, setting $c_R := T_2(y_R(k)) \wedge T_2(-y_R(k))$, where, since V_k is symmetric, $y_R(k) \in I$ is such that $V_k(y_R(k)) = V_k(-y_R(k)) = R$, choosing R sufficiently large such that $c_R \leq a_0$, we have

$$\inf_{y \in \mathcal{V}_k(R)} \int_I dx \psi_0(x) \, \mathbf{1}_{[a,b)} \left(kx + (1-k) \, T_2(y) \right) \ge \inf_{w \in [c_R,c_2]} \int_I dx \psi_0(x) \, \mathbf{1}_{[a,b)} \left(kx + (1-k) \, w \right) \\
= \inf_{w \in [c_R,c_2]} \int_{(ka_0 + (1-k)w) \lor a}^{(kb_0 + (1-k)w) \land b} dx \psi_0(x) \, . \quad (26)$$

Clearly there exist Borel sets [a, b) such that $\inf_{w \in [c_R, c_2]} \int_{(ka_0 + (1-k)w) \lor a}^{(kb_0 + (1-k)w) \land b} dx \psi_0(x) > 0$ if either

$$ka_{0} + (1-k)c_{2} < b_{0} = kb_{0} + (1-k)b_{0} , \qquad (27)$$

$$k(b_{0} - a_{0}) > (1-k)(c_{2} - b_{0}) ,$$

or

$$kb_{0} + (1-k)c_{R} > a_{0} = ka_{0} + (1-k)a_{0} , \qquad (28)$$
$$k(b_{0} - a_{0}) > (1-k)(a_{0} - c_{R}) ,$$

that is

$$\frac{k}{1-k} (b_0 - a_0) > (c_2 - b_0) \wedge (a_0 - c_R)$$

$$k > \frac{\frac{(c_2 - b_0) \wedge (a_0 - c_R)}{(b_0 - a_0)}}{1 + \frac{(c_2 - b_0) \wedge (a_0 - c_R)}{(b_0 - a_0)}} \vee 0 =: k_* .$$
(29)

Then, for any $[a, b) \subseteq I$,

$$\inf_{w \in [c_R, c_2]} \int_{\left(kT_1^2(c) + (1-k)w\right) \lor a}^{\left(kT_1(c) + (1-k)w\right) \lor a} dxh\left(x\right) \ge \left\|\psi_0\right\|_{L^1} \inf_{w \in [c_R, c_2]} \frac{\int_{\left(kT_1^2(c) + (1-k)w\right) \lor a}^{\left(kT_1(c) + (1-k)w\right) \lor a} dx\psi_0\left(x\right)}{\left\|\psi_0\right\|_{L^1}} , \quad (30)$$

which implies $\alpha_k = \|\psi_0\|_{L^1}$ and $\mathcal{B}(I) \ni A \mapsto \tilde{\nu}(A) = \inf_{w \in [c_R, c_2]} \frac{\int_I dx \psi_0(x) \mathbf{1}_A(kx + (1-k)w)}{\|\psi_0\|_{L^1}} \in [0, 1]$.

To sum up what it has been proved so far: Propositions 3 and 2 together, joint with Proposition 6.1 in [CGSV], imply that $\{Y_n^{(k)}\}_{n\geq 0}$ has always a unique invariant measure $\bar{\nu}^{(k)} := g^{(k)}\lambda$ for all k and that for a suitable choice of T_1 and T_2 such that $k_*(T_1, T_2) = 0$, the convergence to $\bar{\nu}^{(k)}$ is geometric, i.e. L_k has a spectral gap in L^1 , for all k, which might not be the case for any choice of the coupled unimodal maps.

¹As an example of T_1 one can consider a map of the same form of T_2 , namely $I \ni x \mapsto T_1(x) := c_1(1-2x^2)$, for a suitable choice of $c_1 \in (0,1)$, which can be proven to satisfy the assumption of Proposition 2.7 in [BS] (see [CGSV] section 5.1). In this case supp $h \subseteq [T_1^2(0), T_1(0)] = [T_1(c_1), c_1] \subset I$.

2.2 The weak limits of the invariant measures of the deterministic system in the very weak and in the very strong coupling regimes

For any $k \in [0, 1]$, setting

$$I^2 \ni (x, y) \longmapsto \psi_x^{(k)}(y) = (1 - k)T_2(y) + kT_1(x) \in I^2$$
, (31)

the coupled system (1) can be written as a skew-product system on the product space I^2 , that is

$$I^2 \ni (x,y) \longmapsto \Theta^{(k)}(x,y) = \left(T_1(x), \psi_x^{(k)}(y)\right) \in I^2 .$$

$$(32)$$

Assuming that T_1 and $T_2 : I \oslash$ are continuous and that T_1 has an invariant measure μ with density h, since $\psi^{(k)}$ is also continuous we can construct an invariant measure $\rho^{(k)}$ for $\Theta^{(k)}$ (see e.g. [Ro]), such that μ is the marginal of the first component of the random vector $\left(\eta_1^{(k)}, \eta_2^{(k)}\right)$ with joint probability measure $\rho^{(k)}$, and a probability kernel $\mathcal{B}(I) \times I \ni (A, x) \longmapsto \rho^{(k)}(A|x) \in [0, 1]$ such that

$$\int_{I^2} \rho^{(k)}(dx, dy) f(x, y) = \int_{I} \mu(dx) \int_{I} \rho^{(k)}(dy|x) f(x, y) , \qquad (33)$$

for any $\rho^{(k)}$ -summable function f.

The answer to Question I is the object of the following

Proposition 4 The random vector $(\eta_1^{(k)}, \eta_2^{(k)})$ with joint probability measure $\rho^{(k)}$ converges in distribution, as $k \uparrow 1$, to the random vector (η_1, η_2) such that, for any Borel set $A \subseteq I^2, \mathbb{P}\{(\eta_1, \eta_2) \in A\} = \int_A dxdyh(x) \,\delta(y-x)$. In other words, the family of conditional probability distribution $\{\rho^{(k)}(\cdot|\eta_1), k \in [0,1]\}$ converges weakly, when $k \to 1$, to the law of a degenerate random variable constantly equal to η_1 , where η_1 has law μ .

Moreover, if T_2 admits a unique invariant measure $\bar{\nu}$, then, in the limit as $k \downarrow 0, \{\rho^{(k)}, k \in [0,1]\}$ converges weakly to $\mu \otimes \bar{\nu}$, that is $\{\rho^{(k)}(\cdot|\eta_1), k \in [0,1]\}$ converges weakly to $\bar{\nu}$.

Proof. We follow the same strategy of Proposition 6.2 in [CGSV].

Let us consider the characteristic function

$$\mathbb{R}^2 \ni (t_1, t_2) \longmapsto \varphi_{\left(\eta_1^{(k)}, \eta_2^{(k)}\right)}(t_1, t_2) := \mathbb{E}\left[e^{i\left\langle (t_1, t_2), \left(\eta_1^{(k)}, \eta_2^{(k)}\right)\right\rangle}\right] \in \mathbb{C}$$
(34)

of the random vector $\left(\eta_1^{(k)}, \eta_2^{(k)}\right)$. Since $\rho^{(k)}$ is invariant for $\Theta^{(k)} := \left(T_1, \psi_{\cdot}^{(k)}\right)$, i.e.

$$\Theta_{*}^{(k)}\rho^{(k)} = \rho^{(k)},$$

$$\varphi_{\left(\eta_{1}^{(k)},\eta_{2}^{(k)}\right)}\left(t_{1},t_{2}\right) = \int_{I^{2}}\rho^{(k)}\left(dx,dy\right)\exp\left\{i\left(t_{1}x+t_{2}y\right)\right\}$$

$$= \int_{I^{2}}\left(\Theta_{*}^{(k)}\rho^{(k)}\right)\left(dx,dy\right)\exp\left\{i\left(t_{1}x+t_{2}y\right)\right\}$$

$$= \int_{I^{2}}\rho^{(k)}\left(dx,dy\right)\exp\left\{i\left(t_{1}T_{1}\left(x\right)+t_{2}\psi_{x}^{(k)}\left(y\right)\right)\right\}$$

$$= \int_{I^{2}}\rho^{(k)}\left(dx,dy\right)e^{it_{1}T_{1}\left(x\right)}e^{it_{2}\left[kT_{1}\left(x\right)+\left(1-k\right)T_{2}\left(y\right)\right]}$$

$$= \int_{I}\mu\left(dx\right)e^{i\left(t_{1}+t_{2}k\right)T_{1}\left(x\right)}\int_{I}\rho^{(k)}\left(dy|x\right)e^{it_{2}\left(1-k\right)T_{2}\left(y\right)}.$$
(35)

Since $\Theta : I^2 \circlearrowleft, \mathbb{E}\left[T_2\left(\eta_2^{(k)}\right) \middle| \eta_1^{(k)}\right]$, the conditional expectation of the random variable $T_2\left(\eta_2^{(k)}\right)$ w.r.t. $\eta_1^{(k)}$, is bounded, then $(1-k)T_2\left(\eta_2^{(k)}\right) \xrightarrow[k\uparrow 1]{} 0$ μ -almost surely. Thus, for any $t_2 \in \mathbb{R}$,

$$\lim_{k \to 1} \int_{I} \left((T_{2})_{*} \rho^{(k)} \right) (dy|x) e^{it_{2}(1-k)y} = \lim_{k \to 1} \int_{I} \rho^{(k)} (dy|x) e^{it_{2}(1-k)T_{2}(y)}$$
(36)
$$= \lim_{k \to 1} \mathbb{E} \left[e^{i(1-k)t_{2}T_{2}\left(\eta^{(k)}_{2}\right)} \middle| \eta^{(k)}_{1} \right] = 1 .$$

Moreover, the family of measures $\{\rho^{(k)}, k \in [0,1]\}$ on $(I^2, \mathcal{B}(I^2))$ is tight, hence sequentially compact; therefore, given a sequence $\{\rho^{(k_n)}\}_{n\geq 1} \subset \{\rho^{(k)}, k \in [0,1]\}$, where $\{k_n\}_{n\geq 1} \uparrow 1$, let us consider $\{\rho^{(k_{n_l})}\}_{l\geq 1}$, with $\{k_{n_l}\}_{l\geq 1} \subset \{k_n\}_{n\geq 1}$, a weakly convergent subsequence of $\{\rho^{(k_n)}\}_{n\geq 1}$. Denoting by $\bar{\rho}$ the weak limit of $\{\rho^{(k_{n_l})}\}_{l\geq 1}$ and by $\bar{\rho}(dy|\cdot)$ the conditional probability distribution such that, for any bounded $\bar{\rho}$ -measurable function f on I^2 ,

$$\int_{I^2} \bar{\rho}\left(dx, dy\right) = \int_{I} \mu\left(dx\right) \int_{I} \bar{\rho}\left(dy|x\right) f\left(x, y\right) , \qquad (37)$$

by the invariance of μ w.r.t. the evolution defined be $T_1, \forall (t_1, t_2) \in \mathbb{R}^2$,

$$\int_{I^2} \bar{\rho} (dx, dy) \exp \{i (t_1 x + t_2 y)\} = \int_{I} \mu (dx) \int_{I} \bar{\rho} (dy|x) e^{i(t_1 x + t_2 y)}$$

$$= \lim_{l \to \infty} \varphi_{\left(\eta_1^{(k_{n_l})}, \eta_2^{(k_{n_l})}\right)} (t_1, t_2)$$

$$= \int_{I} \mu (dx) \lim_{l \to \infty} e^{i(t_1 + k_{n_l} t_2) T_1(x)} \int_{I} \rho^{(k_{n_l})} (dy|x) e^{it_2(1 - k_{n_l}) T_2(y)}$$

$$= \int_{I} \mu (dx) e^{i(t_1 + t_2) T_1(x)} \lim_{l \to \infty} \int_{I} \rho^{(k_{n_l})} (dy|x) e^{it_2(1 - k_{n_l}) T_2(y)}$$

$$= \int_{I} \mu (dx) e^{i(t_1 + t_2) T_1(x)} = \int_{I} \mu (dx) e^{i(t_1 + t_2) x} .$$
(38)

Hence, $\bar{\rho}(dy|x) = dy\delta(y-x)$, μ -a.s.. Since this result holds for any weakly convergent subsequence $\{\rho^{(k_{n_l})}\}_{l\geq 1}$ of any sequence $\{\rho^{(k_n)}\}_{n\geq 1}$ in $\{\rho^{(k)}, k\in[0,1]\}$, we get the thesis.

Proceeding as in the strong coupling limit $(k \to 1)$, by the tightness of $\{\rho^{(k)}, k \in [0,1]\}$, given a sequence $\{\rho^{(k_n)}\}_{n\geq 1} \subset \{\rho^{(k)}, k \in [0,1]\}$, where $\{k_n\}_{n\geq 1} \downarrow 0$, let $\{\rho^{(k_n)}\}_{l\geq 1}$, with $\{k_{n_l}\}_{l\geq 1} \subset \{k_n\}_{n\geq 1}$, be a weakly convergent subsequence of $\{\rho^{(k_n)}\}_{n\geq 1}$ converging to $\bar{\rho}$. Then, setting

$$I^{2} \ni (x, y) \longmapsto \Theta (x, y) := (T_{1}(x), T_{2}(y)) \in I , \qquad (39)$$

 $\forall (t_1, t_2) \in \mathbb{R}$, we have

$$\int_{I^2} \bar{\rho} (dx, dy) \exp \{i (t_1 x + t_2 y)\} = \lim_{l \to \infty} \int_{I^2} \rho^{(k_{n_l})} (dx, dy) \exp \{i (t_1 x + t_2 y)\}$$
(40)
$$= \lim_{l \to \infty} \int_{I^2} \left(\Theta_*^{(k_{n_l})} \rho^{(k_{n_l})}\right) (dx, dy) \exp \{i (t_1 x + t_2 y)\}$$
$$= \lim_{l \to \infty} \int_{I^2} \rho^{(k_{n_l})} (dx, dy) e^{i(t_1 + k_{n_l} t_2)T_1(x) + it_2(1 - k_{n_l})T_2(y)}$$
$$= \int_{I^2} \bar{\rho} (dx, dy) \exp \{i (t_1 T_1(x) + t_2 T_2(y))\}$$
$$= \int_{I^2} (\Theta_* \bar{\rho}) (dx, dy) \exp \{i (t_1 x + t_2 y)\} ,$$

that is the invariance of $\bar{\rho}$ for the evolution defined by Θ . In particular, if (η_1, η_2) is a random vector with law $\bar{\rho}$, by setting $t_2 = 0$ in the r.h.s. of the last expression we get that the marginal of η_1 is invariant under the evolution defined by T_1 and therefore must be equal to μ , while, setting $t_1 = 0$, we get that $\bar{\nu}$, the marginal of η_2 , is invariant under the evolution defined by T_2 . Then, if $\bar{\nu}$ is the unique invariant measure for the evolution defined by $T_2, \mu \otimes \bar{\nu}$ is the unique measure left invariant by Θ .

This result also prove that the answer to Question III is negative.

Indeed, if Question III had a positive answer then a fortiori, for any $f \in C(I)$,

$$\lim \sup_{k \to 1} \lim \sup_{n \to \infty} \left| \nu_n^{(k)}(f) - \bar{\nu}_n^{(k)}(f) \right| = 0 , \qquad (41)$$

for Lebesgue almost all choices of the initial data x_0, y_0 and \mathbb{P} almost all ω , which would imply, denoting by $\bar{\rho}_n^{(k)} := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{(Y_i,\omega_i)}$,

$$\limsup_{k \to 1} \sup_{n \to \infty} \sup_{n \to \infty} \left| \rho_n^{(k)}(\varphi) - \bar{\rho}_n^{(k)}(\varphi) \right| = 0 , \qquad (42)$$

for any $\varphi \in C(I^2)$. But from Proposition 6.2 in [CGSV] and Theorem 2.1.7 in [Ar] any weak limit $\bar{\rho}^{(k)}$ of $\left\{\bar{\rho}_n^{(k)}\right\}_{n\geq 1}$ weakly converges, as $k\uparrow 1$, to $\mu\otimes\mu$ while, denoting by $\rho^{(k)}$ any weak limit of $\left\{\rho_n^{(k)}\right\}_{n\geq 1}$, by the previous result, for any $\varphi \in C(I^2)$, $\lim_{k\to 1} \rho^{(k)}(\varphi) = \int_I \mu(dx) \varphi(x, x)$.

References

[Ar]	L. Arnold Random Dynamical Systems, Springer, (1998).
[Ba]	W. Bartoszek On uniformly smoothing stochastic operators Commenta- tiones Mathematicae Universitatis Carolinae, 36 , 203–206 (1995).
[BS]	V. Baladi, D. Smania <i>Linear response for smooth deformations of generic nonuniformly hyperbolic unimodal maps</i> , Annales scientifiques de l'École Normale Supérieure, 45 , 4, 861–926 (2012).
[BNNT]	P. G. Barrientos, F. Nakamura, Y. Nakano, H. Toyokawa <i>Finitude of physical measures for random maps</i> , preprint, https://arxiv.org/pdf/2209.08714.pdf (2022).
[CGSV]	Caby Th., Gianfelice M., Saussol B., Vaienti S. <i>Topological synchronisation or a simple attractor?</i> Nonlinearity 36 , 7, 3603–3621 (2023).
[HM]	Hairer M., Mattingly J. Yet another look at Harris' ergodic theorem for Markov chains Seminar on Stochastic Analysis, Random Fields and Applications VI, Progr. Probab. 63 , 109–117 (2011).
[Ki]	Yu. Kifer Ergodic Theory of Random Trasformations, Springer (1986).
[Lahav et al.]	Lahav N., Sendina-Nadal I., Hens C., Ksherim B., Barzel B., Cohen R., Boccaletti S. <i>Topological synchronization of chaotic systems</i> Sci. Rep. 12 , 2508 (2022).
[LM]	Lasota A., Mackey M. C. <i>Chaos, Fractals and Noise - Stochastic Aspects of Dynamics</i> Second Edition, Applied Mathematica Sciences Vol. 97, Springer (1994).
[Ro]	V. A. Rohlin On the fundamental ideas of measure theory. Amer. Math. Soc. Translation, 71:27 (1952).