

NEW CRITERIA FOR STARLIKENESS IN THE UNIT DISC

MILUTIN OBRADOVIĆ AND NIKOLA TUNESKI

ABSTRACT. It is well-known that the condition $\operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] > 0$, $z \in \mathbb{D}$, implies that f is starlike function (i.e. convexity implies starlikeness). If the previous condition is not satisfied for every $z \in \mathbb{D}$, then it is possible to get new criteria for starlikeness by using $\left| \arg \left[\alpha + \frac{zf''(z)}{f'(z)} \right] \right|$, $z \in \mathbb{D}$, where $\alpha > 1$.

1. INTRODUCTION AND DEFINITIONS

Let \mathcal{A} be the class of functions f which are analytic in the open unit disc $\mathbb{D} = \{z : |z| < 1\}$ of the form $f(z) = z + a_2z^2 + a_3z^3 + \dots$, and let \mathcal{S} be the subclass of \mathcal{A} consisting of functions that are univalent in \mathbb{D} .

Also, let

$$\mathcal{C} = \left[f \in \mathcal{A} : \operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] > 0, (z \in \mathbb{D}) \right],$$
$$\mathcal{S}^* = \left[f \in \mathcal{A} : \operatorname{Re} \left[\frac{zf'(z)}{f(z)} \right] > 0, (z \in \mathbb{D}) \right],$$

denote the classes of convex and starlike functions, respectively. It is well-known that

$$f \in \mathcal{C} \quad \Rightarrow \quad f \in \mathcal{S}^*,$$

(see Duren [1]). If the condition for convexity, $\operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] > 0$, is not satisfied for every $z \in \mathbb{D}$, then we cannot conclude that $f \in \mathcal{C}$, and consequently, $f \in \mathcal{S}^*$.

In this paper $f \in \mathcal{S}^*$ in the terms of the expression

$$\left| \arg \left[\alpha + \frac{zf''(z)}{f'(z)} \right] \right|, \quad (z \in \mathbb{D})$$

where $\alpha > 1$.

2010 *Mathematics Subject Classification.* 30C45, 30C50.

Key words and phrases. starlike functions, convex functions, criteria.

2. MAINE RESULTS

For our consideration we need the next lemma given by Nunokawa in [3].

Lemma A. *Let $p(z) = 1 + p_1z + p_2z^2 + \dots$ be analytic in the unit disc \mathbb{D} and $p(z) \neq 0$ for $z \in \mathbb{D}$. Also, let suppose that there exists a point $z_0 \in \mathbb{D}$ such that $\operatorname{Re} p(z) > 0$ for $|z| < |z_0|$ and $\operatorname{Re} p(z_0) = 0$, where $p(z_0) \neq 0$, i.e. $p(z_0) = ia$, a is real and $a \neq 0$. Then we have*

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik,$$

where $k \geq \frac{1}{2} \left(a + \frac{1}{a} \right)$, when $a > 0$, and $k \leq -\frac{1}{2} \left(|a| + \frac{1}{|a|} \right)$, when $a < 0$.

Theorem 1. *Let $f \in \mathcal{A}$, $\alpha > 1$, and let*

$$(1) \quad \left| \arg \left[\alpha + \frac{z f''(z)}{f'(z)} \right] \right| < \operatorname{arctg} \frac{\sqrt{3}}{\alpha - 1} \quad (z \in \mathbb{D}).$$

Then $f \in \mathcal{S}^*$.

Proof. If we put that $p(z) = \frac{z f'(z)}{f(z)}$, then $p(0) = 1$ and $\frac{z p'(z)}{p(z)} + p(z) - 1 = \frac{z f''(z)}{f'(z)}$. So, we would need to prove that the following implication holds:

$$(2) \quad \left| \arg \left[\frac{z p'(z)}{p(z)} + p(z) + \alpha - 1 \right] \right| < \operatorname{arctg} \frac{\sqrt{3}}{\alpha - 1} \quad (z \in \mathbb{D})$$

$$\Rightarrow \operatorname{Re} p(z) > 0 \quad (z \in \mathbb{D}).$$

First we will prove that $p(z) \neq 0, z \in \mathbb{D}$. On a contrary, if there exists $z_1 \in \mathbb{D}$ such that z_1 is the zero of order m of the function p , then $p(z) = (z - z_1)^m p_1(z)$, where m is positive integer, p_1 is analytic in \mathbb{D} with $p_1(z_1) \neq 0$, and further

$$\frac{z p'(z)}{p(z)} = \frac{mz}{z - z_1} + \frac{z p_1'(z)}{p_1(z)}.$$

This means that the real part of the right hand side can tend to $-\infty$ when $z \rightarrow z_1$, which is a contradiction to the assumption of the theorem regarding the argument. Thus $p(z) \neq 0, z \in \mathbb{D}$.

Now, let suppose that the implication (2) does not hold in the unit disc. It means that there exists a point $z_0 \in \mathbb{D}$ such that $\operatorname{Re} p(z) > 0$ for $|z| < |z_0|$ and $\operatorname{Re} p(z_0) = 0$, where $p(z_0) \neq 0$. If we put $p(z_0) = ia$, a is real and $a \neq 0$, then by Lemma A we have

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik,$$

where $k \geq \frac{1}{2} \left(a + \frac{1}{a} \right)$, when $a > 0$, and $k \leq -\frac{1}{2} \left(|a| + \frac{1}{|a|} \right)$, when $a < 0$.

Next, if we put $\Phi(z) = \frac{zp'(z)}{p(z)} + p(z) + \alpha - 1$, then by the previous facts:

$$\operatorname{Re} \Phi(z_0) = \alpha - 1, \quad \operatorname{Im} \Phi(z_0) = k + a,$$

which for $a > 0$ implies:

$$\begin{aligned} \arg \Phi(z_0) &= \operatorname{arctg} \frac{k+a}{\alpha-1} \geq \operatorname{arctg} \frac{\frac{1}{2}(a + \frac{1}{a}) + a}{\alpha-1} \\ &= \operatorname{arctg} \frac{3a + \frac{1}{a}}{2(\alpha-1)} \geq \operatorname{arctg} \frac{\sqrt{3}}{\alpha-1}, \end{aligned}$$

because the function $\varphi(a) = 3a + \frac{1}{a}$ has its minimum value $\varphi(\frac{1}{\sqrt{3}}) = 2\sqrt{3}$. Similarly, for $a < 0$:

$$\begin{aligned} \arg \Phi(z_0) &= \operatorname{arctg} \frac{k+a}{\alpha-1} \leq \operatorname{arctg} \frac{-\frac{1}{2}(|a| + \frac{1}{|a|}) - |a|}{\alpha-1} \\ &= -\operatorname{arctg} \frac{3|a| + \frac{1}{|a|}}{2(\alpha-1)} \leq -\operatorname{arctg} \frac{\sqrt{3}}{\alpha-1}. \end{aligned}$$

Combining the cases $a > 0$ and $a < 0$, we receive

$$|\arg \Phi(z_0)| \geq \operatorname{arctg} \frac{\sqrt{3}}{\alpha-1},$$

which is a contradiction to the relation (1).

This show that $\operatorname{Re} p(z) = \operatorname{Re} \frac{zf'(z)}{f(z)} > 0$, for all $z \in \mathbb{D}$, i.e. that $f \in \mathcal{S}^*$. \square

Example 1. Let $f \in \mathcal{A}$ is defined by the condition

$$(3) \quad 1 + \frac{zf''(z)}{f'(z)} = (\sqrt{3} + 1) \left(\frac{1+z}{1-z} \right)^{\frac{1}{2}} - \sqrt{3},$$

where we use the principal value of the square root. For real z close to -1 we have $\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) < 0$, which means that f is not convex. On the other hand, from (3) we have

$$\frac{zf''(z)}{f'(z)} + (\sqrt{3} + 1) = (\sqrt{3} + 1) \left(\frac{1+z}{1-z} \right)^{\frac{1}{2}}$$

and from here

$$\begin{aligned} \left| \arg \left[\frac{zf''(z)}{f'(z)} + (\sqrt{3} + 1) \right] \right| &\leq \frac{1}{2} \left| \arg \frac{1+z}{1-z} \right| \\ &< \frac{\pi}{4} = \operatorname{arctg} \frac{\sqrt{3}}{(\sqrt{3} + 1) - 1}, \end{aligned}$$

which by Theorem 1 (with $\alpha = \sqrt{3} + 1$) implies that $f \in \mathcal{S}^*$.

Letting α tend to 1 in Theorem 1, we have the following well-known result.

Corollary 1. *Let $f \in \mathcal{A}$ and*

$$\left| \arg \left[\frac{zf''(z)}{f'(z)} + 1 \right] \right| < \frac{\pi}{2} \quad (z \in \mathbb{D})$$

Then $f \in \mathcal{S}^$.*

Further, it is easy to verify that the disc with center α and radius $\frac{\alpha\sqrt{3}}{\sqrt{3+(\alpha-1)^2}}$ is contained in the angle $|\arg z| < \arctg \frac{\sqrt{3}}{\alpha-1}$, and so by using the result of Theorem 1, we have that

$$\left| \frac{zf''(z)}{f'(z)} \right| = \left| \left(\frac{zf''(z)}{f'(z)} + \alpha \right) - \alpha \right| < \frac{\alpha\sqrt{3}}{\sqrt{3+(\alpha-1)^2}} \quad (z \in \mathbb{D})$$

implies $f \in \mathcal{S}^*$. Since the function $\varphi(\alpha) =: \frac{\alpha\sqrt{3}}{\sqrt{3+(\alpha-1)^2}}$, $\alpha \geq 1$, attains its maximal value 2 for $\alpha = 4$, we get

Corollary 2. *Let $f \in \mathcal{A}$ and*

$$\left| \frac{zf''(z)}{f'(z)} \right| < 2 \quad (z \in \mathbb{D})$$

Then $f \in \mathcal{S}^$.*

In a similar way as in Theorem 1, we can consider starlikeness problem in connection with the class defined by

$$\mathcal{G} = \left[f \in \mathcal{A} : \operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] < \frac{3}{2}, z \in \mathbb{D} \right].$$

Ozaki in [4] proved that if $f \in \mathcal{G}$, then f is univalent in \mathbb{D} . Later, Umezawa in [6] showed the functions from \mathcal{G} are convex in one direction. Also, it is shown in the papers [2] and [5] that \mathcal{G} is subclass of \mathcal{S}^* .

If $\operatorname{Re} \left[1 + \frac{zf''(z)}{f'(z)} \right] < \frac{3}{2}$ is not satisfied for every $z \in \mathbb{D}$, then we can pose a question if for some $\beta < 1$, such that

$$(4) \quad \operatorname{Re} \left[\beta + \frac{zf''(z)}{f'(z)} \right] < \frac{3}{2} \quad (z \in \mathbb{D}),$$

is it possible obtain sufficient condition for starlikeness, in a similar way as in Theorem 1. The condition (4) is equivalent to

$$\operatorname{Re} \left[\frac{3-2\beta}{2} - \frac{zf''(z)}{f'(z)} \right] > 0 \quad (z \in \mathbb{D}),$$

and we can use similar technique. This lead so

Theorem 2. *Let $f \in \mathcal{A}$, $\beta < 1$, and let*

$$\left| \arg \left[\frac{3 - 2\beta}{2} - \frac{zf''(z)}{f'(z)} \right] \right| < \operatorname{arctg} \frac{2\sqrt{3}}{5 - 2\beta} \quad (z \in \mathbb{D}).$$

Then $f \in \mathcal{S}^$.*

Proof. As in the proof of Theorem 1, we put that $p(z) = \frac{zf'(z)}{f(z)}$, such that $p(0) = 1$ and $\frac{zp'(z)}{p(z)} + p(z) - 1 = \frac{zf''(z)}{f'(z)}$. Now, our aim is to prove that for some $\beta < 1$ the following implication holds:

$$(5) \quad \left| \arg \left[\frac{5 - 2\beta}{2} - \frac{zp'(z)}{p(z)} - p(z) \right] \right| < \operatorname{arctg} \frac{2\sqrt{3}}{5 - 2\beta} \quad (z \in \mathbb{D})$$

$$\Rightarrow \operatorname{Re} p(z) > 0 \quad (z \in \mathbb{D}).$$

First, as in Theorem 1 we conclude that $p(z) \neq 0, z \in \mathbb{D}$.

Further, if the implication (5) is not true, there exists a point $z_0 \in \mathbb{D}$ such that $\operatorname{Re} p(z) > 0$ for $|z| < |z_0|$ and $\operatorname{Re} p(z_0) = 0$, where $p(z_0) \neq 0$. If we put $p(z_0) = ia$, a is real and $a \neq 0$, then by Lemma A we have

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik,$$

where $k \geq \frac{1}{2} \left(a + \frac{1}{a} \right)$, when $a > 0$, and $k \leq -\frac{1}{2} \left(|a| + \frac{1}{|a|} \right)$, when $a < 0$. Also, for

$$\Psi(z) = \frac{5 - 2\beta}{2} - \frac{zp'(z)}{p(z)} - p(z),$$

using the previous conclusions, we have

$$\operatorname{Re} \Psi(z_0) = \frac{5 - 2\beta}{2}, \quad \operatorname{Im} \Psi(z_0) = -(k + a).$$

Using the same method as in the proof of Theorem 1, we easily conclude that

$$\arg \Psi(z_0) = -\operatorname{arctg} \frac{2(k + a)}{5 - 2\beta} \leq -\operatorname{arctg} \frac{2\sqrt{3}}{5 - 2\beta},$$

when $a > 0$, and

$$\arg \Psi(z_0) = -\operatorname{arctg} \frac{2(k + a)}{5 - 2\beta} \geq \operatorname{arctg} \frac{2\sqrt{3}}{5 - 2\beta},$$

when $a < 0$. These facts imply contradiction of the assumption in (5).

So, we have the statement of this theorem. \square

Letting β tend to 1 in Theorem 2, we receive

Corollary 3. *Let $f \in \mathcal{A}$ and*

$$\left| \arg \left[\frac{1}{2} - \frac{zf''(z)}{f'(z)} \right] \right| < \operatorname{arctg} \frac{2}{\sqrt{3}} \approx 49.1^\circ \quad (z \in \mathbb{D}).$$

Then $f \in \mathcal{S}^$.*

As in the case of Theorem 1, we can show that the disc with center $\frac{3-2\beta}{2}$ and radius $\frac{(3-2\beta)\sqrt{3}}{\sqrt{12+(5-2\beta)^2}}$ is containing in the angle $|\arg z| \leq \operatorname{arctg} \frac{2\sqrt{3}}{5-2\beta}$. So, if

$$\left| \frac{zf''(z)}{f'(z)} \right| = \left| \left(\frac{3-2\beta}{2} - \frac{zf''(z)}{f'(z)} \right) - \frac{3-2\beta}{2} \right| < \frac{(3-2\beta)\sqrt{3}}{\sqrt{12+(5-2\beta)^2}},$$

for all $z \in \mathbb{D}$, then Theorem 2 brings that $f \in \mathcal{S}^*$. Since the function $\psi(\beta) =: \frac{(3-2\beta)\sqrt{3}}{\sqrt{12+(5-2\beta)^2}}$ is decreasing on the interval $(-\infty, 1]$, and $\psi(\beta) \rightarrow \sqrt{3}$, when $\beta \rightarrow -\infty$, we get

Corollary 4. *Let $f \in \mathcal{A}$ and*

$$\left| \frac{zf''(z)}{f'(z)} \right| < \sqrt{3} \quad (z \in \mathbb{D}).$$

Then $f \in \mathcal{S}^$.*

For a starlike function f it is not necessary that $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$. For example, for Koebe function $k(z) = \frac{z}{(1-z)^2}$ we have

$$\operatorname{Re} \frac{k(z)}{z} \Big|_{z=(1+i)/\sqrt{2}} = -(\sqrt{2} + 1) < 0,$$

and this also holds for points in \mathbb{D} close enough to $\frac{1+i}{\sqrt{2}}$.

In the next theorem we give a condition which provides $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$.

Theorem 3. *Let $f \in \mathcal{A}$, $\gamma \geq 0$, and let*

$$\left| \arg \left[\frac{zf'(z)}{f(z)} + \gamma \right] \right| < \operatorname{arctg} \frac{1}{1+\gamma} \quad (z \in \mathbb{D}).$$

Then $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$.

Proof. We apply the same method as in the previous theorems. In this case, we use $p(z) = \frac{f(z)}{z}$, such that $p(0) = 1$ and $\frac{zp'(z)}{p(z)} + 1 = \frac{zf'(z)}{f(z)}$. So, the result we need to prove can be rewritten in the following equivalent form:

$$(6) \quad \left| \arg \left[\frac{zp'(z)}{p(z)} + 1 + \gamma \right] \right| < \operatorname{arctg} \frac{1}{1+\gamma} \quad (z \in \mathbb{D})$$

$$\Rightarrow \operatorname{Re} p(z) > 0 \quad (z \in \mathbb{D}).$$

First, $p(z) \neq 0$ for all $z \in \mathbb{D}$.

Next, if the implication (6) does not hold in the unit disc, then there exists a point $z_0 \in \mathbb{D}$ such that $\operatorname{Re} p(z) > 0$ for $|z| < |z_0|$ and

Re $p(z_0) = 0$, where $p(z_0) \neq 0$. If we put $p(z_0) = ia$, a is real and $a \neq 0$, then by Lemma A we have

$$\frac{z_0 p'(z_0)}{p(z_0)} = ik,$$

where $k \geq \frac{1}{2} \left(a + \frac{1}{a} \right)$, when $a > 0$, and $k \leq -\frac{1}{2} \left(|a| + \frac{1}{|a|} \right)$, when $a < 0$. Now, for $a > 0$ we get:

$$\begin{aligned} \arg \left[\frac{z_0 p'(z_0)}{p(z_0)} + 1 + \gamma \right] &= \arg(ik + 1 + \gamma) = \operatorname{arctg} \frac{k}{1 + \gamma} \\ &\geq \frac{\frac{1}{2} \left(a + \frac{1}{a} \right)}{1 + \gamma} \geq \operatorname{arctg} \frac{1}{1 + \gamma}, \end{aligned}$$

and for $a < 0$,

$$\arg \left[\frac{z_0 p'(z_0)}{p(z_0)} + 1 + \gamma \right] \leq -\operatorname{arctg} \frac{1}{1 + \gamma}.$$

By combining the above conclusions, we receive that

$$\left| \arg \left(\frac{z_0 p'(z_0)}{p(z_0)} + 1 + \gamma \right) \right| \geq \operatorname{arctg} \frac{1}{1 + \gamma},$$

which is a contradiction to assumption in (6). \square

For $\gamma = 0$ in previous theorem we have next result.

Corollary 5. *Let $f \in \mathcal{A}$ and*

$$\left| \arg \frac{z f'(z)}{f(z)} \right| < \frac{\pi}{4} \quad (z \in \mathbb{D}).$$

Then $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$.

In Theorem 3 the function f from \mathcal{A} need not be starlike as the next example shows.

Example 2. *Let $f \in \mathcal{A}$ is defined by*

$$(7) \quad \frac{z f'(z)}{f(z)} = \sqrt{3} \left(\frac{1+z}{1-z} \right)^{\frac{1}{3}} + 1 - \sqrt{3},$$

where we use the principal value for square root. For real z close to -1 we have that $\operatorname{Re} \left[\frac{z f'(z)}{f(z)} \right] < 0$, which means that f is not starlike. But from (7) we have

$$\frac{z f'(z)}{f(z)} + (\sqrt{3} - 1) = \sqrt{3} \left(\frac{1+z}{1-z} \right)^{\frac{1}{3}},$$

and from here

$$\begin{aligned} \left| \arg \left[\frac{zf'(z)}{f(z)} + (\sqrt{3} - 1) \right] \right| &\leq \frac{1}{3} \left| \arg \left(\frac{1+z}{1-z} \right) \right| \\ &< \frac{\pi}{6} = \operatorname{arctg} \frac{1}{(\sqrt{3} - 1) + 1}, \end{aligned}$$

which by Theorem 3 (with $\gamma = \sqrt{3} - 1$) implies that $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$.

It is possible to show that the disc with center $1 + \gamma$ and radius $\frac{1+\gamma}{\sqrt{1+(1+\gamma)^2}}$ lies in the angle $|\arg z| \leq \operatorname{arctg} \frac{1}{1+\gamma}$, and so, by using the result of Theorem 3 we have that

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| = \left| \left(\frac{zf'(z)}{f(z)} + \gamma \right) - (1 + \gamma) \right| < \frac{1 + \gamma}{\sqrt{1 + (1 + \gamma)^2}} \quad (z \in \mathbb{D})$$

implies $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$. When $\gamma \rightarrow +\infty$ we receive

Corollary 6. *Let $f \in \mathcal{A}$ and*

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| < 1 \quad (z \in \mathbb{D}).$$

Then $\operatorname{Re} \frac{f(z)}{z} > 0$, $z \in \mathbb{D}$.

REFERENCES

1. P.L. Duren, *Univalent function*, Springer-Verlag, New York, 1983.
2. I. Jovanović and M. Obradović, A note on certain classes of univalent functions, *Filomat*, **9**(1) (1995), 69–72.
3. M. Nunokawa, On properties of non-Caratheodory functions, *Proc. Japan Acad.*, **68**, Ser.A (1992), 152-153.
4. S. Ozaki, On the theory of multivalent functions, II, *Sci. Rep. Tokyo, Bunrika Daigaku* **4** (1941), 45-87.
5. R. Singh, S. Singh, Some sufficient conditions for univalence and starlikeness, *Colloq. Math.*, **47** (1982), 309-314.
6. T. Umezawa, Analytic functions convex in one direction, *J. Math. Soc. Japan*, **4** (1952), 194-202.

DEPARTMENT OF MATHEMATICS, FACULTY OF CIVIL ENGINEERING, UNIVERSITY OF BELGRADE, BULEVAR KRALJA ALEKSANDRA 73, 11000, BELGRADE, SERBIA

Email address: obrad@grf.bg.ac.rs

DEPARTMENT OF MATHEMATICS AND INFORMATICS, FACULTY OF MECHANICAL ENGINEERING, Ss. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE, KARPOŠ II B.B., 1000 SKOPJE, REPUBLIC OF NORTH MACEDONIA.

Email address: nikola.tuneski@mf.edu.mk