Bound state properties, positron annihilation and hyperfine structure of the four-body positronium hydrides

Alexei M. Frolov^{*}

Department of Applied Mathematics University of Western Ontario, London, Ontario N6H 5B7, Canada

(Dated: May 16, 2024)

Abstract

Bound state properties of the ground (bound) ${}^{1}S(L = 0)$ -state(s) in the four-body positronium hydrides 1 HPs, 2 HPs (DPs), 3 HPs (TPs) and MuPs are determined and investigated. By using numerical data from our computations of these four-body systems we have determined a number of different annihilation rates for each of these positronium hydrides and evaluated the hyperfine structure splitting. The properties of the ground (bound) states in the four-body exitonic ${}^{M}h^{+}e_{2}^{-}e^{+}$ complexes, where $M \geq 1$ and $M \leq 1$, have also been evaluated numerically. The neutral four-body systems ${}^{M}h^{+}e_{2}^{-}e^{+}$ with $M \gg 1$ are similar to the HPs hydrides. In particular, each of these states has only one bound state. We also discuss applications of the exponential and semi-exponential variational expansions for accurate, bound state computations of the four-body positronium hydrides.

This version approximately corresponds to the final (journal) version published in: Journal of Molecular Physics, 120(1), e2019337 (2022) (doi.org/10.1080/00268976.2021.2019337). In any case I recommend you to read the final/journal version of this paper.

^{*}E-mail address: alex1975frol@gmail.com

I. INTRODUCTION

The positronium hydrides HPs are the Coulomb four-body systems with unit electrical charges. Each molecule of these positronium hydrides contains one heavy nucleus of hydrogen (either protium p, or deuterium d, or tritium t), two electrons e^- and one positron e^+ . Analogous four-leptonic system with the positively charged μ^+ muon is MuPs $(\mu^+ e_2^- e^+)$. This system is also considered here as a light positronium hydride. All positronium hydrides HPs and similar exitonic ${}^{M}h^{+}e_{2}^{-}e^{+}$ complexes are of increasing interest in solid state physics, stellar astrophysics and other areas of physics (see, e.g., [1], [2] and references therein). In solid state physics the positronium hydrides are important to develop and improve the general theory of four-body exitonic complexes with unit electrical charge which are needed to describe optical properties of many important semiconductors, e.g., CdS, InS, and others (see, e.g., [3], [4]). The positronium hydrides also play an important role in stellar astrophysics [1], [2], since these four-body systems are formed in very large numbers in photospheres of all hot stars, e.g., in many O_{-}, B_{-} and even in some early A_{-} stars. By studying positron annihilation in photospheres of these stars we can investigate some crucial processes and physical conditions inside of such hot stars and accurately predict their timeevolution. It is interesting to note that annihilation of positrons in positronium hydrides in photospheres of less hot stars, e.g., in our Sun (spectral class G2), is extensively studied in modern papers to understand the nature and evaluate the rates of many fundamental processes and reactions, which proceed in outer layers of these stars (see, e.g., contributions and references in [1]). Also this allows one to evaluate numerical values of some important physical parameters, e.g., the effective magnetic and electrical fields in those areas.

Stability of the model positronium hydride $^{\infty}$ HPs was shown in [5]. Then this system has been investigated a number of times (see, e.g., [6] - [9]). In 1992 the molecules of positronium ¹HPs hydride have been created (in the collisions between positrons and methane) and observed in the laboratory [10]. By replacing the methane molecule ('target' molecule) by the molecules of deuterium and/or tritium substituted methane, e.g., CD₄ and CD₂T₂, it is possible to create (in the same experiment) the positronium DPs and TPs hydrides. The most interesting and exciting property of all positronium hydrides is annihilation of the electron-positron pair(s) which is often called the positron annihilation. In our earlier paper [11] we have considered the bound state properties and positron annihilation in these

positronium hydrides. Later, different properties of the positronim hydrides have been reevaluated in a large number of papers (see, e.g., [1], [12] - [15] and references therein). Since then the overall accuracy of our variational wave functions has been improved substatially. In addition to this, our current understanding of many processes and reactions in positronium hydrides is substantially better, than it was twenty five years ago. Furthermore, for the same time the masses and other physical parameters of particles, which are included in positronium hydrides, and numerical values of other physical constants have been changed noticeably. Now, by using the updated values of these constants and particle masses we want to evaluate the bound state properties of the HPs, DPs, TPs and MuPs hydrides to much better accuracy, than it was achieved in [11]. For each of these positronium hydrides we also determine a few different annihilation rates and predict the actual hyperfine structure. We also discuss the four-body exitonic ${}^{M}h^{+}e_{2}^{-}e^{+}$ complexes which are similar to the regular positronium hydrides HPs (${}^{M}H^{+}e_{2}^{-}e^{+}$, or ${}^{M}h^{+}e_{2}^{-}e^{+}$). Here the notation H⁺ and/or h^{+} designates the positively charged hole, which has the unit electric charge, and its mass exceeds the rest mass of electron m_e . We also consider possible applications of our four-body exponential and semi-exponential variational expansions [16], [17] and [18] in the relative coordinates for accurate, bound state computations of the four-body positronium hydrides. Concluding remarks can be found in the last Section.

II. BOUND STATE WAVE FUNCTIONS AND PROPERTIES OF THE POSITRO-NIUM HYDRIDES

The Hamiltonian \hat{H} of the non-relativistic four-body $H^+e_2^-e^+$ hydride is written in the form:

$$\hat{H} = -\frac{\hbar^2}{2m_e}\Delta_1 - \frac{\hbar^2}{2m_e}\Delta_2 - \frac{\hbar^2}{2m_e}\Delta_3 - \frac{\hbar^2}{2M}\Delta_4 + \frac{e^2}{r_{12}} - \frac{e^2}{r_{13}} - \frac{e^2}{r_{14}} - \frac{e^2}{r_{23}} - \frac{e^2}{r_{24}} + \frac{e^2}{r_{34}} , \quad (1)$$

where e is the electric charge of the positron (-e is the electric charge of the electron), m_e is the mass of the electron/positron, while $\hbar = \frac{h}{2\pi}$ is the reduced Planck constant, which is also called the Dirac constant. In atomic units, where $\hbar = 1, m_e = 1$ and e = 1, the same Hamiltonian \hat{H} takes the form

$$\hat{H} = -\frac{1}{2}\Delta_1 - \frac{1}{2}\Delta_2 - \frac{1}{2}\Delta_3 - \frac{1}{2M}\Delta_4 + \frac{1}{r_{12}} - \frac{1}{r_{13}} - \frac{1}{r_{14}} - \frac{1}{r_{23}} - \frac{1}{r_{24}} + \frac{1}{r_{34}}, \qquad (2)$$

where the notations (or indexes) 1 and 2 (or -) stand for the two electrons, the index 3 (or +) means the positron, while the index p (or 4) designates the positively charged heavy nucleus of the hydrogen isotope (p, d, t), or positively charged muon μ^+ . The mass of this nucleus is denoted as Mm_e (or M in atomic units). The same system of notations is used everywhere below in this paper. As mentioned above in this study we consider the bound state properties of the following positronium hydrides ¹HPs, DPs, TPs and MuPs. The model four-body $^{\infty}$ HPs system with the infinitely heavy hydrogen nucleus is also investigated. In general, each of these hydrides has only one ground (bound) ${}^{1}S(L=0)$ -state (or $1{}^{1}S(L=0)$ -state). The first step of our procedure is to obtain solutions of the corresponding Schrödinger equation $H\Psi = E\Psi$ for bound states (E < 0). In fact, by solving this equation we determine the total energy E and wave function Ψ of the ground (bound)¹S(L = 0)-state for each of the postronium hydrides. At the second stage we determine various bound state properties of these positronium hydrides by calculating the corresponding expectation values with the wave functions Ψ which has been derived at the first stage of our procedure. In general, the expectation value $A = \langle \hat{A} \rangle$ of some physical operator \hat{A} , which does not depend explicitly upon the time t, is defined as follows

$$A = \langle \hat{A} \rangle = \frac{\langle \Psi \mid \hat{A} \mid \Psi \rangle}{\langle \Psi \mid \Psi \rangle} = \langle \tilde{\Psi} \mid \hat{A} \mid \tilde{\Psi} \rangle , \qquad (3)$$

where Ψ is the wave function of the given (bound) state and the bound state wave function $\tilde{\Psi}$ has unit norm. The expectation value $A = \langle \hat{A} \rangle$ from Eq.(3) is often called the bound state property of this system in a given (bound) state. The bound state properties of the actual positronium hydrides HPs have been determined as described above. By choosing different quantum operators \hat{X} in Eq.(3) we can evaluate numerically a large number of bound state properties, e.g., all inter-particle distances, interparticle delta-functions and other similar values. The bound state properties of the actual geometry of these four-body systems and evaluate a number of actual physical properties, including different annihilation rates, hyperfine structure splitting, etc. In addition to this, it is interesting to compare the bound state properties of different positronium hydrides and investigate their mass dependence upon variations of the proton mass.

In calculations performed for this study we have applied the four-body variational expansion written in six-dimensional gaussoids of the relative interparticle coordinates $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j| = r_{ji}$, where (ij) = (12), (13), (23), (14), (24) and (34) and \mathbf{r}_i are the Cartesian coordinates of the particle *i*, where i = 1, 2, 3, 4. Note that each of the six relative coordinates r_{ij} is translationally and rotationally invariant. This fact allows one to use the Hamiltonian written in the Cartesian coordinates in the form of Eqs.(1) and (2). The variational expansion in multi-dimensional guassoids was proposed, developed and applied to various nuclear and atomic systems by N.N. Kolesnikov and his co-workers in the middle of 1970's (see, e.g., [19], [20] and earlier references therein). The explicit form of this variational expansion in the case of four-body systems is

$$\Psi = \mathcal{A}_S \sum_{k=1}^{N} C_k \exp\left(-\alpha_{12}^{(k)} r_{12}^2 - \alpha_{13}^{(k)} r_{13}^2 - \alpha_{14}^{(k)} r_{14}^2 - \alpha_{23}^{(k)} r_{23}^2 - \alpha_{24}^{(k)} r_{24}^2 - \alpha_{34}^{(k)} r_{34}^2\right) \quad , \qquad (4)$$

where C_k are the linear variational parameters of this expansion (k = 1, ..., N), while α_{ij}^k are the non-linear parameters of the variational expansion, Eq.(4). For four-body systems we have six different interparticle distances r_{ij} which are explicitly shown in the variational expansion, Eq.(4). This explains why the basis functions in Eq.(4) are often called the six-dimensional gaussoids written in the four-body relative coordinates r_{ij} [21]. The four-body systems are relatively simple for this variational expansion, Eq.(4), since in this case we do not have any unnecessary relative coordinate(s). In contrast with this, for the five-, six- and many body systems the number of relative inter-particle coordinates r_{ij} (in our three-dimensional space) exceeds the total number of independent 'radial' coordinates. In particular, for an arbitrary five-body system one finds ten interparticle coordinates in our three-dimensional space, while only nine of them are truly independent. For six-body systems we have three additional (or unnecessary) relative coordinates. In general, the appearance of unnecessary relative coordinates leads to additional troubles and numerical instabilities in variational calculations during optimization of the non-linear parameters (for more details, see, [21]).

In applications to actual four-body systems the trial wave function, Eq.(4), must be symmetrized (or antisymmetrized) in respect to possible presence of identical particles in such systems. In each of the positronium hydrides we always have two identical electrons (particles 1 and 2). Therefore, the \mathcal{A}_S operator in Eq.(4) should include the electron-electron symmetrization operator, i.e., $\mathcal{A}_S \simeq \frac{1}{\sqrt{2}} (1 + \hat{P}_{12})$ in our current notation. All our calculations in this study have been performed in atomic unis, where $\hbar = 1$, |e| = 1 and $m_e = 1$. The masses of heavy particles used in our computations are [22]

$$M_p = 1836.15267343 \ m_e \ , \ M_d = 3670.48296788 \ m_e \ ,$$

$$M_t = 5496.92153573 \ m_e \ , \ M_\mu = 206.7682830 \ m_e \ .$$
(5)

These masses of hydrogen isotopes are currently recommended for scientific use by CO-DATA/NIST in 2021/2022 [22].

In general, the overall efficiency of the variational expansion, Eq.(4), substantially depends upon algorithms which are used to optimize the non-linear parameters in Eq.(4). Recently, for four-body systems we have developed a number of algorithms which were found to be quite effective and fast in applications to various systems. The accuracy of the constructed wave functions is usually high (and even very high) for the total energies. The expectation values of various geometrical and dynamical properties are also determined (with the same wave functions) to relatively high numerical accuracy. However, some troubles can still be found in computations of the expectation values of some delta-functions, interparticle cusp values [23] and for other similar properties (see discussion in Section VI).

Results of our numerical computations performed for the ground ${}^{1}S(L=0)$ -states in the positronium hydrides are shown in Table I. These results include the total energies and other bound state properties of all positronium hydrides discussed in this study. As mentioned above in these hydrides the index 4 designates the heavy, positively charged nucleus of the hydrogen isotope, i.e., protium p, deuterium d and tritium t. We also determine the bound state properties of the model ${}^{\infty}H^+e_2^-e^+$ hydride and four-leptonic MuPs hydride which is the $\mu^+ e_2^- e^+$ ion. In the last two cases in Table I the index p (or 4) stands for the infinitely heavy ${}^{\infty}H^+$ nucleus (model nucleus) and for the positively charged muon μ^+ , respectively. The properties shown in Table I include the $\langle r_{ij}^k \rangle$ expectation values, where k is the power of the interparticle (ij)-distances. In calculations performed for this study we used k = -2, -1, 1, 2, 3 and 4. In Table I one also finds the expectation values of single particle kinetic energies $\langle -\frac{1}{2}\nabla_i^2 \rangle$ determined for the both electron and positron. Table II contains the expectation values of all interparticle delta-functions $\langle \delta_{ij} \rangle$ and triple electronpositron delta-functions $\langle \delta_{+--} \rangle$. These delta-functions are needed in the following Sections to determine the corresponding annihilation rates. Physical meaning of all bound state properties shown in Table I is clear from the notation used. Analytical formulas for all matrix elements used in computations of these bound state properties can be found in [11]. Note that the two-particle cusp values [23] cannot be computed directly with the use of variational expansion, Eq.(4), since the corresponding expectation values equal zero identically. This is an obvious disadvantage of the variational expansion, Eq.(4), in applications to the few-and many-body Coulomb systems.

However, this disadvantage is not critical and all similar 'problematic' expectation values can be evaluated indirectly by using one of the two following procedures: (a) re-expansion of the wave functions which is already represented in terms of variational expansion, Eq.(4), and (b) calculations of the expectation values of some other operators and their commutators. Then, at the second step of the procedure (b) the expectation values, which are needed for our evaluations, are constructed as linear combinations of the expectation values of these operators. In particular, for the interparticle delta-functions in Coulomb few-body systems such an approach has been developed and tested by R.J. Drachman and his colleagues (see, e.g., [24]).

III. POSITRON ANNIHILATION

The most interesting property of each positronium hydride is annihilation of the electronpositron pair, which is often called the positron annihilation, for short. In general, annihilation of the electron-positron pair(s) in positronium hydrides proceeds mainly with the emission of two high energy photons, or two annihilation γ -quanta. In this study we restrict ourselves to the analysis of positron annihilation from the bound states of positronium hydrides only. An alternative process which is called annihilation-in-flight is not discussed here (see, e.g., [25] and [26]). Similar process proceeds, e.g., during collisions between negatively charged hydrogen ion and fast positrons. The probability of the two-photon annihilation per second is called the two-photon annihilation rate $\Gamma_{2\gamma}$. It is clear that the $\Gamma_{2\gamma}$ value is a unique property of any given bound state. For the ground (bound) ${}^{1}S(L = 0)$ state in the positronium hydrides the explicit formula for the $\Gamma_{2\gamma}$ rate can be written in the following form (see, e.g., [27] - [29] and [30])

$$\Gamma_{2\gamma} = n \ m \ \pi \ \alpha^4 c a_0^{-1} \langle \delta(\mathbf{r}_{+-}) \rangle = 2\pi \alpha^4 c a_0^{-1} \langle \delta_{+-} \rangle \quad , \tag{6}$$

which includes the expectation value of the electron-positron delta-function $\langle \delta(\mathbf{r}_{+-}) \rangle = \langle \delta_{+-} \rangle$ expressed in atomic units. This expectation value must be evaluated numerically for the ground ${}^{1}S(L = 0)$ state in the HPs hydride. In this formula the notation $\alpha = 7.2973525693 \cdot 10^{-3}$ stands for the dimensionless fine-structure constant, $c = 2.99792458 \cdot 10^{10} \ cm \cdot sec^{-1}$ is the speed of light in vacuum and $a_0 = 5.29177210903 \cdot 10^{-9} \ cm$ is the Bohr radius [22]. Here and everywhere below in this study, all numerical values for physical constants have been taken from recent set of physical constants values published by NIST [22]. Note that these values are currently recommended for scientific use by CODATA/NIST in 2021/2022 [22].

Also, in Eq.(6) and in all formulas below the notation n stands for the total number of bound electrons in atomic system, while m denotes the total number of positrons bound in the same atomic system. The product mn is the total number of electron-positron pairs in this atomic system. In fact, in all positronium hydrides HPs we have n = 2 and m = 1, i.e., the product mn equals two. The notation $\Gamma_{2\gamma}$ is very convenient and everywhere below in this study we shall apply the notation $\Gamma_{k\gamma}$ to designate the k-photon ($k \ge 0$) annihilation rate averaged over orientations of the spins of all electrons and positrons participating in the process. Analogous formula for the three-photon annihilation rate $\Gamma_{3\gamma}$ in the posironium hydrides is

$$\Gamma_{3\gamma} = nm \, \frac{4(\pi^2 - 9)}{3\pi} \pi \alpha^5 c a_0^{-1} \langle \delta(\mathbf{r}_{+-}) \rangle = \frac{8(\pi^2 - 9)}{3} \alpha^5 c a_0^{-1} \langle \delta_{+-} \rangle \quad . \tag{7}$$

These two largest annihilation rates $\Gamma_{2\gamma}$ and $\Gamma_{3\gamma}$ are crucially important to describe positron annihilation in any atomic and/or molecular system. Furthermore, if we know these two annihilation rates, then we can also evaluate the four- and five-photon annihilation rates, i.e., the $\Gamma_{4\gamma}$ and $\Gamma_{5\gamma}$ values, respectively. The corresponding approximate formulas for these values have been derived in [31]

$$\Gamma_{4\gamma} = 0.274 \left(\frac{\alpha}{\pi}\right)^2 \Gamma_{2\gamma} \quad \text{and} \quad \Gamma_{5\gamma} = 0.177 \left(\frac{\alpha}{\pi}\right)^2 \Gamma_{3\gamma} \quad .$$
 (8)

The formulas Eqs.(7) and (8) are used in this study to determine the $\Gamma_{3\gamma}$, $\Gamma_{4\gamma}$ and $\Gamma_{5\gamma}$ annihilation rates for all positronium hydrides mentioned in this study.

Currently, the formula, Eq.(6), for the two-photon annihilation rate is used only occasionally. Instead, the following formula for the $\Gamma_{2\gamma}$ annihilation rate (in the Ps⁻, Ps₂ and HPs systems) is applied in modern calculations and experimental measurements

$$\Gamma_{2\gamma} = n \ m \ \pi \alpha^4 c a_0^{-1} \left[1 - \frac{\alpha}{\pi} \left(5 - \frac{\pi^2}{4} \right) \right] \left\langle \delta_{+-} \right\rangle \ . \tag{9}$$

This formula contains the two-photon annihilation rate, Eq.(6), and also includes the lowest order QED correction to that value [32]. In our calculations of $\Gamma_{2\gamma}$ performed for this study we also apply the formula, Eq.(9). For many actual systems it is also interesting to evaluate the total annihilation rate Γ , which is the sum of all partial annihilation rates. In reality, the four-, five- and other multi-photon annihilation rates are significantly smaller than the sum of the two- and three-photon annihilation rates. This allows one to write the following approximate expression for the total annihilation rate of the electron-positron pair in arbitrary atom and/or molecule

$$\Gamma \approx \Gamma_{2\gamma} + \Gamma_{3\gamma} = n \ m \ \pi \alpha^4 c a_0^{-1} \Big[1 - \alpha \Big(\frac{17}{\pi} - \frac{19\pi}{12} \Big) \Big] \langle \delta_{+-} \rangle \quad . \tag{10}$$

This formula has a very good numerical accuracy for arbitrary n and m, and it is often used in various applications to all positron containing few-body systems [33]. The total annihilation rate Γ was measured in early 1980's [34], [35] for the three-body Ps⁻ ion. In regular experiments and observations with poly-electrons, atoms and molecules, which contain positron(s), one usually deals with the total annihilation rate Γ . This includes the three-body Ps⁻ ion (mn = 2), four-body bi-positronium molecule Ps₂ (mn = 4), positronium hydrides HPs (mn = 2) and other positron-containing atoms and molecules which are known in astrophysics and solid state physics. All annihilation rates mentioned here and below (the both $\Gamma_{1\gamma}^a$ and $\gamma_{1\gamma}^F$ rates) can be found in Table III.

A. One- and zero-photon annihilation rates

As mentioned above all positronium hydrides are the bound Coulomb four-body systems which contain three light particles and one heavy (or central), positively charged particle. In each of these hydrides all four particles interact with each other and, in general, such interactions are not small. This means that annihilation of the electron-positron pair in each of these positronium hydrides may also proceed with the emission of either a single photon (one-photon annihilation), or even with no emission of photon(s) et al. Briely, zero-photon annihilation in postronium hydrides HPs means that the both photons emitted during two-photon annihilation are absorbed by the remaining particles: one by the electron and another by the heavy nucleus. Let us briefly discuss annihilation of the electron-positron pair in the HPs hydrides which proceeds with no emission of one and zero photon(s). The corresponding annihilation rates are designated below as $\Gamma_{1\gamma}$ (with some additional indexes) and $\Gamma_{0\gamma}$, respectively. It is easy to predict that the numerical values $\Gamma_{1\gamma}$ and $\Gamma_{0\gamma}$ are very small. Nevertheless, in a number of problems and processes known in astrophysics, solid state physics, etc, these one- and zero-photon annihilations play some roles. For the first time, the one-photon annihilation rate $\Gamma_{1\gamma}$ in the three-body Ps⁻ ion was determined in [36], [37] (see, also [38])¹. In earlier paper [39] the authors took into account twice less Feynman diagrams than in [36] and obtained a different result.

The same one-photon annihilation is possible in the four-body positronium hydrides and the corresponding annihilation rate $\Gamma^a_{1\gamma}$ was evaluated in our earlier paper [11]. In this case the electron-positron pair annihilates in the nearest vicinity of another (second) electron in the HPs hydride. The value of $\Gamma^a_{1\gamma}$ is given by the formula [36] which is written in the following form (in atomic units) [37]

$$\Gamma_{1\gamma}^{a} = \frac{64\pi^{2}}{27} \,\alpha^{8} c a_{0}^{-1} \langle \delta_{+--} \rangle \approx 1065.756921 \cdot \langle \delta_{+--} \rangle \quad , \tag{11}$$

where the notation $\langle \delta_{+--} \rangle$ is the expectation value of the three-particle $e^-e^+e^-$ deltafunction which must be expressed in atomic units. The formula, Eq.(11), is exactly the same as it was derived for the three-body Ps⁻ ion. This triple delta-function is determined for the ground (bound) ${}^{1}S(L=0)$ -state of the positronium hydride(s). In general, the knowledge of the $\langle \delta_{+--} \rangle$ expectation value allows one to evaluate the one-photon annihilation rate $\Gamma_{1\gamma}^{(I)}$ in the three-body Ps⁻ ion, four-body bi-positronium molecule Ps₂, all four-body positronium hydrides HPs and other few-body compounds which contain positron(s) and more than one bound electron.

In all positronium hydrides mentioned above another one-photon annihilation of the electron-positron (e^-, e^+) -pair is also possible. Indeed, let us assume that positron approaches the electron which is bound by the Coulomb field of the heavy, positively charged atomic nucleus. Therefore, the probability (or rate) of such an annihilation can be determined exactly in the same way as we calculate the cross section(s) of atomic photoionization. During such computations the positron must be considered as an electron which moves backwards in time. This one-photon annihilation was studied and described in detail in [40], [41] (see, also [27] and references therein). For few-electron quasi-atomic systems this one-photon annihilation rate, which is often called the Fermi annihilation rate, is written in the form (in atomic units, see, e.g., [27])

$$\gamma_{1\gamma}^{F} = \frac{4\pi}{3} \ n \ m \ \alpha^{8} Q^{5} \delta_{+-} v_{+-} \left(\frac{c}{a_{0}}\right) \,, \tag{12}$$

where the notation n stands for the total number of bound electrons, while m denotes the total number of bound positrons in this atom/molecule, while Q is the electric charge of the heavy, central nucleus. Also, in this equation δ_{+-} is the electron-positron delta-function and v_{+-} is the relative velocity of the two particles (electron and positron) at the point of electron-positron coalesence. The exact formula for the $\Gamma_{1\gamma}^F$ rate takes the form

$$\Gamma_{1\gamma}^{F} = \frac{4\pi}{3} n m \alpha^{8} Q^{5} \left\langle \delta_{+-} \frac{p_{+-}}{m_{e}} \right\rangle \left(\frac{c}{a_{0}} \right) = \frac{4\pi}{3} n m \alpha^{8} Q^{5} \mid \nu_{+-} \mid \left\langle \delta_{+-} \right\rangle \left(\frac{c}{a_{0}} \right) , \tag{13}$$

where $\nu_{+-} = \frac{\langle \delta_{+-} \nabla_{+-} \rangle}{\langle \delta_{+-} \rangle}$ is the electron-positron cusp (its definition can be found, e.g., in [23] and [38]). For highly accurate, bound state wave functions the electron-postiron cusp always equals $-\frac{1}{2} = -0.5$ (to very high accuracy). In other words, it almost coincides with the analytically predicted cusp value between two point, electrically charged particles *i* and *j* which equals $\nu_{ij} = q_i q_j \frac{m_i m_j}{m_i + m_j}$ [23]. For the positronium hydrides, where Q = 1, n = 2 and m = 1, the last formula takes the form

$$\Gamma_{1\gamma}^{F} = \frac{2\pi}{3} n m \alpha^{8} \langle \delta_{+-} \rangle \left(\frac{c}{a_{0}}\right) = \frac{4\pi}{3} \alpha^{8} \langle \delta_{+-} \rangle \left(\frac{c}{a_{0}}\right), \qquad (14)$$

where all expectation values are determined for the ground (bound) state in each of the positronium hydrides. Note that this one-photon annihilation rate is proportional to the expectation value of the electron-positron delta-function $\langle \delta_{+-} \rangle$, while the $\Gamma^a_{1\gamma}$ annihilation rate, Eq.(11), is proportional to the expectation value of the triple electron-positron-electron delta-function $\langle \delta_{+--} \rangle$. Such a difference in formulas follows from the fundamental difference between these two annihilation processes. The one-photon (Fermi) annihilation rate $\Gamma^F_{1\gamma}$ has never been evaluated for the positronium hydrides HPs. In the light positronium systems, e.g., in the Ps⁻ ion and/or in the Ps₂ molecule, the one-photon Fermi annihilation is impossible.

In our previous paper [11] we also evaluated another one-photon annihilation rate $\Gamma_{1\gamma}^{(b)}$ and zero-photon annihilation rate $\Gamma_{0\gamma}^{(b)}$. These evaluations were essentially based on our earlier results obtained for the four-body, bi-positronium Ps₂ molecule [37]. Later, we have found that 'analogies' between the light Ps₂ system and heavy (one-center) positronium hydrides HPs do not work well for one-center HPs systems, and this may lead to very large errors in similar evaluations. In particular, in the HPs hydrides the Coulomb field of the central (very heavy) hydrogen nucleus always play a substantial role in these annihilation processes (in contrast with the Ps₂ system). Currently, we derive more accurate formulas which can be used to describe these annihilation rates in HPs hydrides. The new analysis of these annihilation processes in the HPs hydrides and calculations of their annihilation rates will be discussed elsewhere.

IV. HYPERFINE STRUCTURES OF THE FOUR-BODY POSITRONIUM HY-DRIDES

As mentioned above all positromium hydrides are the Coulomb four-body systems of actual, interacting particles and each of these particles has non-zero spin. In fact, all particles which form these positronium hydrides ¹HPs, ²HPs (DPs), ³HPs (TPs) and MuPs, i.e., electrons, positron, μ^+ -muonium, protium and tritium nuclei (but deuterium nucleus!) have spin which equals $\frac{1}{2}$. The spin of the deuterium nucleus equals unity. Therefore, even for the bound $1^{1}S(L=0)$ -state in each of these positronium hydrides one can observe the direct spin-spin interactions between four particles in the HPs hydrides. In general, for the bound S(L=0)-states such a spin-spin (positron-nucleus) interaction (see below) proceeds at very short distances between two particles with non-zero spin. Also, it is crucial to note that in all positronium hydrides ¹HPs, DPs, TPs and MuPs the two bound electrons can only be in the singlet ${}^{1}S(L=0)$ -state. Otherwise, i.e., for the triplet ${}^{3}S(L=0)$ electron pair, the whole four-body HPs hydride (or $H^+e_2^-e^+$ system) is not bound [11]. Therefore, in each of the positronium hydrides there is a spin-spin interaction between positively charged positron and heavy nucleus of hydrogen isotope (or μ^+ -muon). In general, such an interaction is relatively small $\simeq \alpha^2$ and it is proportional to the expectation value of the positron-nucleus delta-function. This follows from the explicit form of the Hamiltonian H_{HF} , which describes the hyperfine structure of the positronium hydrides. In atomic units this Hamiltonian H_{HF} is

$$H_{HF} = -\sum_{(ij)} a_{ij} \left(\mathbf{s}_i \cdot \mathbf{s}_j \right) = -a \left(\mathbf{s}_+ \cdot \mathbf{I}_H \right) - b \left(\mathbf{s}_+ \cdot \mathbf{S}_- \right) - c \left(\mathbf{S}_- \cdot \mathbf{I}_H \right) = -a \left(\mathbf{s}_+ \cdot \mathbf{I}_H \right)$$
$$= -\frac{8\pi\alpha^2}{3} \mu_B^2 \left(\frac{g_+}{m_e} \right) \left(\frac{g_H}{M_p} \right) \left\langle \delta_{+H} \right\rangle \left(\mathbf{s}_+ \cdot \mathbf{I}_H \right) , \qquad (15)$$

where α is the fine-structure constant, μ_B is the Bohr magneton, M_p is the proton mass (at rest), while \mathbf{S}_{-} is the total electron spin, which equals zero for the singlet electron-electron pair (for the triplet electronic states this formula, Eq.(15), has a different form). In atomic

units the absolute value of the Bohr magneton $\mu_B = \frac{e\hbar}{2m_e}$ equals $\frac{1}{2}$ (exactly). The notation $\langle \delta_{+H} \rangle$ stands for the expectation value of the positron-nucleus delta-function expressed in atomic units. This δ -function must be determined for the ground $1^1S(L=0)$ -state of the HPs hydride. The notations g_+ and g_H stand for the gyromagnetic ratios of the positron and heavy nucleus of the hydrogen isotope, M_p is the proton mass (in atomic units), m_e is the positron mass and I_H is the nuclear spin of the hydrogen isotope which is integer (or semi-integer) positive value. The nuclear g-factor (or g_H factor in Eq.(15)) is the ratio of magnetic moment of this nucleus (in nuclear magnetons) to its maximal spin value $I_N(=I_H)$ [42]. In positronium hydrides we have $I_H = \frac{1}{2}$, $I_T = \frac{1}{2}$ and $I_D = 1$, while in the positronium MuPs hydride the factor $\left(\frac{g_H}{M_p}\right)$ must be replaced by the analogous muonic factor $\left(\frac{g_\mu}{m_\mu}\right)$.

The hyperfine structure splitting ΔE_{HF} in the HPs hydrides is the difference between two bound states with the total spin $I_H + \frac{1}{2}$ and $I_H - \frac{1}{2}$, respectively. This difference is

$$\Delta E_{HF} = \frac{2\pi\alpha^2}{3} \left(\frac{g_+}{m_e}\right) \left(\frac{g_H}{M_p}\right) F_{Ry} \left\langle \delta_{+H} \right\rangle \left(\mathbf{s}_+ \cdot \mathbf{I}_H\right) \quad MHz = \mathcal{A} \left\langle \delta_{+H} \right\rangle \left(\mathbf{s}_+ \cdot \mathbf{I}_H\right) \quad MHz , \quad (16)$$

where the constant \mathcal{A} is

$$\mathcal{A} = \frac{2\pi\alpha^2}{3} \left(\frac{g_+}{m_e}\right) \left(\frac{g_H}{M_p}\right) F_{Ry}$$

Numerical values of the g-factors in this formula are [22]

$$g_e = -2.00231930436256$$
, $g_\mu = -2.0023318418$, $g_p = 5.5856946893$
 $g_d = 0.8574382338$, $g_t = 5.957924931$, (17)

while the numerical values of the fine-structure constant α and proton mass M_p have been taken from the previous Sections. The conversion factor (or Rydberg factor) is $F_{Ry} = 2Ry \cdot c = 2 \times 3.2898419602508 \cdot 10^{15} Hz = 6.5796839205016 \cdot 10^9 MHz$ [22] in Eq.(16) allows one to re-calculate the hyperfine structure splitting from atomic units to MHz (or kHz) which are the traditional units for these values. The factor ($\mathbf{s}_+ \cdot \mathbf{I}_H$) in Eq.(16) is the scalar product of the vector of positron spin \mathbf{s}_+ and nuclear spin \mathbf{I}_H . This factor equals ($\mathbf{s}_+ \cdot \mathbf{I}_H$) = $s_+(2I_H + 1) = 1$, if $I_H = \frac{1}{2}$, and $\frac{3}{2}$, if $I_H = 1$.

The formula, Eq.(16), is used in this study to determine the hyperfine structure splittings in all positronium hydrides mentioned above. Note that for all positronium hydrides we have splitting between the two groups of energy levels, e.g., the upper and lower energy levels. Indeed, if I_H is the spin of the nucleus, then one finds $I_H = \frac{1}{2}$ for the p^+ and t^+ nuclei and μ^+ muon. For the deuterium nucleus we have $I_D = 1$. The positron's spin equals $\frac{1}{2}$. Therefore, in the ¹HPs, TPs and MuPs hydrides all states from first group have the total spin equals unity (triplet states), while the state(s) from the second group have the total spin equals zero (singlet states). For the deuterium DPs hydride all states from the first group have spin equals $\frac{3}{2}$ (quartet states), while analogous states from the second group have spin equals $\frac{1}{2}$ (doublet states). Numerical values of the hyperfine structure splitting determined for each of the positronium hydrides, considered in this study, can be found in Table VI which also contains the factors \mathcal{A} used in the formula, Eq.(16), for each of these hydrides.

V. BOUND STATES IN THE FOUR-BODY EXITONIC COMPLEXES

The neutral four-body complexes (or clusters) with unit electrical charges $A^+e_2^-e^+$ are similar to the HPs hydrides, and they are of interest in various applications closely related to the physics and spectroscopy of those semiconductors, which have large and very large electric permittivity. It is shown in numerous papers and books on solid states physics (see, [3], [4]) that many unique optical properties of semiconductors can only be explained, if we allow formation of some few-body exotic systems, which include 'effective' electrons e^- , positively charged hole(s) h^+ and atom(s) A (or ion(s) A^+) which represent the impurities and/or lattice defects. A number of experimental results obtained in solid state physics of semiconductors were explained in the late 1930's with the help of two-body exitons of Wannier-Mott [43], [44]. These two-body exitons are similar to the electron-positron pairs in QED (see, e.g., [4]). In [45] Lampert suggested that exitonic complexes may also include more than two light particles, e.g., they can include a few light particles (holes and electrons) and one heavy particle. Furthermore, such few-body exitonic complexes can be either mobile (light), or heavy (immovable). The mobile exitonic complexes usually consist of a few electrons e^- and some light, positively charged hole(s) h^+ . Immovable exitonic complexes also include at least one heavy, positively charged central atom/ion which is usually associated with either impurity, or lattice defect.

Let us consider the four-body exitonic complex $A^+e_2^-e^+$ (or $h^+e_2^-e^+$) which are similar to the positronium hydrides HPs. First, we discuss the four-body exitonic systems which contains two electrons e^- , one positron e^+ and one heavy, positively charged 'quasi-nucleus' A^+ (or h^+). All these particles have unit electrical charges. The Hamiltonians of similar four-body HPs-like exitonic complexes are written in the following general form

$$H = -\frac{\hbar^2}{2m_e} \left[\nabla_1^2 + \nabla_2^2 + \nabla_3^2 + \left(\frac{m_e}{M}\right) \nabla_4^2 \right] - \frac{e^2}{r_{41}} - \frac{e^2}{r_{42}} - \frac{e^2}{r_{31}} - \frac{e^2}{r_{32}} + \frac{e^2}{r_{21}} + \frac{e^2}{r_{43}} , \qquad (18)$$

where $\hbar = \frac{\hbar}{2\pi}$ is the reduced Planck constant (or Dirac constant), m_e is the electron mass at rest and e is the electric charge of the electron. In this equation we apply the same notations which were used in Section II, i.e., the subscripts 1 and 2 designate the two bound electrons e^- , the subscript 3 denotes the positron e^+ , while the subscript 4 means the positively charged 'heavy' hole which has the mass $M(\geq 1)$. The electric charge of this hole equals to the electric charge of positron e^+ . All other notations are exactly the same as they were used in Section II above. In atomic units (where $\hbar = 1$, |e| = 1, $m_e = 1$) this Hamiltonian takes the form

$$H = -\frac{1}{2} \left[\nabla_1^2 + \nabla_2^2 + \nabla_3^2 + \frac{1}{M} \nabla_4^2 \right] - \frac{1}{r_{41}} - \frac{1}{r_{42}} - \frac{1}{r_{31}} - \frac{1}{r_{32}} + \frac{1}{r_{21}} + \frac{1}{r_{43}} \quad . \tag{19}$$

As follows from this expression the Hamiltonian operator is a regular function of the dimensionless parameter $\frac{m_e}{M} = \frac{1}{M}$, where the mass M is expressed in the electron mass m_e .

First, we investigate transformation of the light (i.e., mobile) four-body exitonic complexes into heavy exitonic complexes which are less mobile. For now, we shall assume that the mass of heavy particle (or the effective mass of the impurity) in the ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems varies (increases) from the unity $M_{h} = m_{e}$ (bi-positronium Ps₂) to relatively large mass values, e.g., up to $M = 12 m_{e}$ and larger masses. It is interesting to investigate quantitative changes in the total energies and other bound state properties of the ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems, when such a mass increases. It is clear that when the mass of heavy particle (M) increases to very large values, then the bound states properties, including the total energies, become similar to the analogous properties of the heavy HPs hydrides and MuPs system. However, the mass region $m_{e} \leq M \leq 12 m_{e}$ is critical to understand the effect of rapid changes which occur in many bound state properties of the ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems, when the heavy mass M increases. Results of our computations performed for a number of ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems can be found in Table V. For each system shown in this Table the corresponding results include the total energy $E, \langle r_{+-} \rangle, \langle r_{--} \rangle$ and $\langle r_{h-} \rangle$ distances and the expectation value of the electron-positron delta-function $\langle \delta(\mathbf{r}_{+-}) \rangle$.

Our results obtained for the exitonic ${}^{M}h^{+}e_{2}^{-}e^{+}$ complexes can be represented in the form of mass-interpolation formula which provides a very good accuracy in actual applications. This formula has been derived to approximate the mass-dependencies of actual properties of the four-body ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems is an analytical function of the following dimensionless parameter $Z = 1 - \frac{2m_{e}}{M}$, which is the inverse mass of the heaviest particle. The actual mass-dependence of the total energies of these exitonic complexes is accurately represented by the following mass-interpolation formula

$$E(Z) = \left(\frac{1+Z}{2}\right)E(1) + \left(\frac{1-Z}{2}\right)E(-1) + \left(\frac{1-Z^2}{4}\right)\left[C_0 + C_1P_1(Z) + C_2P_2(Z) + \dots + C_nP_n(Z)\right],$$
(20)

where $P_n(Z)$ are the Legendre polynomials $(P_0(Z) = 1)$, while $E(1) \approx -0.7891967666545$ a.u. and $E(-1) \approx -0.51600379066033$ a.u. are the total energies of the ground $1^{1}S(L =$ 0)-state(s) in the $^{\infty}$ HPs and Ps₂ systems, respectively. For this systems we have Z = 1and Z = -1, respectively. The unknown coefficients C_n^k $(k = 1, \ldots, K)$ in this formula are determined by using the data from direct and accurate numerical computations of the ground states in a number of different ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems with different 'nuclear' masses $M(\geq 1)$. Similar formulas can be written for other bound state properties of these fourbody systems. Numerical values of the first ten coefficients in the formula, Eq.(20), are presented in Table VI. To determine these ten coefficients we have used data obtained in numerical calculations of twenty different positronium hydrides ${}^{M}h^{+}e_{2}^{-}e^{+}$. The formula, Eq.(20), with the coefficients from Table VI provides ≈ 8 - 9 correct decimal digits for the ground (bound) $1^{1}S(L=0)$ -state energy in an arbitrary exitonic complex $^{M}h^{+}e_{2}^{-}e^{+}$ (or positronium hydride), where $M \geq 1$. Ten similar coefficients C_n^{\star} shown in Table VI have been determined by using computational data for fourteen different ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems. In reality, the formula, Eq.(20), with the C_n^{\star} coefficients provides a better accuracy. Our massinterpolation formula, Eq.(20), can be used to predict the total energies of a number of 'new' ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems. Table VI contains the predicted total energies for the five model ${}^{M}h^{+}e_{2}^{-}e^{+}$ systems in which $M = m_{\mu}$, 100 m_{e} , 75 m_{e} , 25 m_{e} and 15 m_{e} . The overall accuracy of our current predictions can be evaluated as $\approx 1 \cdot 10^{-9} a.u.$

Here we have to note that in modern solid state physics there is a strong and permanent interest to the bound states in model systems in which masses of the both 'holes' and 'electrons' can be less than unity. Formally, the original idea of Wannier-Mott was based on the assumption that in semiconductors with very large electric permittivity ε (or dielectric tensor) we can introduce the two new 'effective' particles: (1) negatively charged 'electron' which corresponds to the bottom of conduction band, and (2) positively charged 'hole' which corresponds to the top of valence band. The numerical values of masses of these effective particles are determined by assuming the exact parabolic symmetry for the minimum of the conduction band and maximum of the valence band. There is also a direct Coulomb interaction between these two effective particles. Plus, in many actual semi-conductors (e.g., in CdS) we always have a number of different conduction and valence bands (usually, two/three bands). This means that in Wannier-Mott theory the masses of 'holes' and 'electrons' are, in fact, some experimental parameters, rather than actual particle masses. In actual computations such effective masses of the holes and 'electrons' were varied between 0.00985 m_e and 2.2678 m_e (see, references in [1], [4], [46]. Note also that almost all numerical computations of few-body exitonic complexes and/or complexes with ionized impurities in solid state physics are performed in the quasi-atomic units, which are defined by choosing the energy and linear size units as follows

$$E_0 = \frac{me^4}{\hbar^2} \left(\frac{m_e^*}{m}\right) \frac{1}{\varepsilon^2} \quad \text{and} \quad a_0 = \frac{\hbar^2}{me^2} \left(\frac{m}{m_e^*}\right) \varepsilon$$

where ε is the electric permittivity of semiconductor which is, in fact, a macroscopic parameter. For the first time this system of units was used in calculations of the three- and four-body exitonic complexes performed in [46].

In this study we also consider the four-body 'positronium hydrides' $h^+e_2^-e^+$ which also contain four point particles with unit electrical charges, but the mass of positively charged hole h^+ is less than unity, i.e., less than electron mass in atomic units. Calculations of the bound states in such systems is tricky, since all our codes work only for few-body systems where the minimal particle mass is always larger (or equal) unity (or $m_e = 1$). Such an agreement has been made at early stages of development of few-body codes [50]. However, in few-body systems with unit electrical charges there is the well known CMP-invariance, which was discovered in [51] for the four-body systems with unit electrical charges. In our current four-body case this fact simply means that the energies of the two systems ${}^{m}h^+e_2^-e^+$ and $e^+Y_2^-Z^+$ (expressed in atomic units) are related by the following, exact equation: $E({}^{m}h^+e_2^-e^+) = mE(e^+Y_2^-Z^+)$. Here $m_h = m \leq 1$ and $M_{Y^-} = M_{Z^+} = \frac{1}{m} \geq 1$, where all masses are also expressed in atomic units. In other words, the total energy of the ${}^{0.1}h^+e_2^-e^+$ system equals 0.1 of the total energy of the $e^+Y_2^-Z^+$ system, where $M_{Y^-} = M_{Z^+}$ $= 10 m_e$. For the last $e^+Y_2^-Z^+$ system our accurate computational method work perfectly. The same (or very similar) mass-relations can also be found for other bound state properties in these 'mass-conjugate' four-body systems. This CMP-invariance in the four-body systems (in general, in few-body systems) opens a wide avenue for applications of our and other computational methods to accurate and highly-accurate calculations of bound states in the systems where the minimal mass is smaller than unity.

The total energies of these four-body systems are shown in Table VI. As follows from the results presented in that Table all these model four-body $h^+e_2^-e^+$ systems are stable against dissociation into two neutral (two-body) clusters, i.e., against reaction $h^+e_2^-e^+ =$ $h^+e^- + e^+e^-$ which corresponds to the actual (or physical) threshold of stability of the $h^+e_2^-e^+$ system. It is also interesting to note that the actual limit (at $m_h \to 0$) of the total energies of four-body $h^+e_2^-e^+$ systems coincides with the total energy of the three-body Ps⁻ ion (E = -0.2620050702329801077704065(55) a.u. [47]). Another fact can be formulated in the form: our mass-interpolation formula, Eq.(20), is directly applicable to the systems $h^+e_2^-e^+$ with $m_h \leq 1$, if we redefine parameters in this formula as follows: z = 1 - 2m, where $m = m_h$ is the hole mass expressed in m_e . In addition to this, we have to assume that now $E(-1) = E(Ps_2)$ and $E(1) = E(Ps^-)$ are the total energies of the ground state(s) in the bi-positronium Ps₂ and three-body Ps⁻ ion, respectively. After such a re-definition of parameters, the formula, Eq.(20), becomes a good mass-interpolation formula for the total energies of the $h^+e_2^-e^+$ systems, where $m_h \leq 1$. The explicit form of this mass-interpolation formula in this case is

$$E(z) = \left(\frac{1+z}{2}\right)E(1) + \left(\frac{1-z}{2}\right)E(-1) + \left(\frac{1-z^2}{4}\right)\left[B_0 + B_1P_1(z) + B_2P_2(z) + \dots + B_nP_n(z)\right],$$
(21)

where E(1) is the total energy of the three-body Ps⁻ ion, while E(-1) is the total energy of the four-body Ps₂ system. The unknown coefficients B_n^k (k = 1, ..., K) in this formula are determined by using the data from direct and accurate numerical computations of the ground states in a number of different ${}^m h^+ e_2^- e^+$ systems with different 'nuclear' masses $m(\leq 1)$. Analogous mass-interpolation formulas can be derived for other bound state properties of these systems, e.g., for the $\langle r_{ij} \rangle$ and $\langle \delta_{ij} \rangle$ expectation values. Note also that all Legendre polynomials of different orders, which are needed in our calculations, can be determined for one numerical value of argument x by using the known recurrence formula (see, e.g., [48]):

$$P_{n+1}(x) = \left(2 - \frac{1}{n+1}\right) x P_n(x) - \left(1 - \frac{1}{n+1}\right) P_{n-1}(x)$$
(22)

and two 'initial conditions' $P_0(x) = 1$ and $P_1(x) = x$.

Numerical values of the first seven coefficients in the formula, Eq.(21), can also be found in Table VI. To determine these ten coefficients we have used data obtained in numerical calculations of the ten 'model' positronium hydrides ${}^{M}h^{+}e_{2}^{-}e^{+}$. The formula, Eq.(21), with the coefficients from Table VI provides $\approx 8 - 9$ correct decimal digits for the ground (bound) $1^{1}S(L = 0)$ -state energy in an arbitrary exitonic complex ${}^{m}h^{+}e_{2}^{-}e^{+}$ (or positronium hydride), where $m \leq 1$. Finally, based on the results of calculations for a large number of systems we have found that that our mass-interpolation formulas, Eqs.(20) and (21) for the $h^{+}e_{2}^{-}e^{+}$ systems with $m_{h} = M \geq 1$ and $m_{h} = m \leq 1$ are reliable and numerically stable. Based on our mass-interpolation formula, Eq.(21), for the ${}^{m}h^{+}e_{2}^{-}e^{+}$ systems we have predicted the total energies for five new similar systems where $m = 0.85 m_{e}, 0.75 m_{e}, 0.55 m_{e}, 0.3 m_{e}$ and $0.15 m_{e}$. The overall accuracy of our current predictions for these systems can be evaluated as $\approx 3 \cdot 10^{-9} a.u$.

Construction of accurate mass-interpolation formulas for the total energies and other bound state properties is an absolutely new and advanced area of research in few-body physics. Currently, we can see here quite a few directions which must be investigated in the future. For instance, one can construct the united mass-interpolation formula for all four-body ions positronium hydrides ${}^{M}h^{+}e_{2}^{-}e^{+}$. Very likely, such a formula cannot be based, in principle, on the orthogonal polynomials.

VI. NEW METHODS FOR HIGHLY ACCURATE, BOUND STATE COMPUTA-TIONS OF THE POSITRONIUM HYDRIDES

In our numerical, bound state computations of the four-body positronium hydrides HPs we have used the variational expansion written in six-dimensional gaussoids, Eq.(4), each of which explicitly depends upon the six relative coordinates $r_{12}, r_{13}, r_{23}, r_{14}, r_{24}$ and r_{34} , where $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j| = r_{ji}$ and \mathbf{r}_i (i = 1, 2, 3, 4) are the Cartesian coordinates of all four (point) particles. This variational expansion is well known in few-body physics since the middle of 1970's (see, e.g., [19], [20], [49] and earlier references therein). After many years of successful calculations of different bound states in various nuclear and atomic few-body systems this variational expansion in multi-dimensional gaussoids can be recognized as very effective, fast and sufficiently accurate in 99% cases, including all positronium hydrides HPs discussed in this study.

However, some important properties of many four-body systems, e.g., all interparticle cusp values, cannot be computed directly and accurately with the use of this variational expansion, Eq.(4). This means we do not have any criterion to check the actual accuracy of our two-particle delta-functions. Furthermore, for Coulomb systems the variational expansion in multi-dimensional gaussoids, Eq.(4), always provides incorrect asymptotics at large interparticle distances r_{ij} . This statement is true for each basis function in Eq.(4) and for any trial wave function constructed with the use of Eq.(4). As a result the actual convergence rate of many bound state properties upon the total number of basis functions N used in Eq.(4) is often non-monotonic. Furthermore, such a rate substantially depends upon algorithms and subroutines which have been used for optimization of the non-linear parameters in Eq.(4). In addition to this, description of bound few-body systems in multidimensional gaussoids is always more diffuse, than similar results obtained with the use of other variational expansions. Briefly, this means that all expectation values of interparticle distances $\langle r_{ij}^n \rangle$, where $n \geq 1$, obtained by using the KT-expansion, Eq.(4), are systematically and noticeably larger than the same expectation values computed by other methods.

In order to avoid these and other similar problems in accurate, bound state computations of the four-body systems in [16] and [17] we have proposed and developed another variational expansion which is written in the form of 'linear' exponential variational expansion of the four-body relative coordinates r_{ij} :

$$\Psi_N(r_{14}, r_{24}, r_{34}, r_{12}, r_{13}, r_{23}) = \mathcal{A}_S \sum_{k=1}^N C_k \exp(-\alpha_{12}^{(k)} r_{12} - \alpha_{13}^{(k)} r_{13} - \alpha_{14}^{(k)} r_{14} - \alpha_{23}^{(k)} r_{23} - \alpha_{24}^{(k)} r_{24} - \alpha_{34}^{(k)} r_{34}) \quad , \qquad (23)$$

where C_k are the linear variational parameters of this expansion (k = 1, ..., N), while α_{ij}^k are the non-linear parameters. The operator \mathcal{A}_S is an unitary projector which produces the trial wave function of the correct permutation symmetry between all identical particles in the system. The variational expansion, Eq.(23), includes the same six interparticle distances which are included in the expansion, Eq.(4). Numerous advantages of this exponential variational expansion, Eq.(23), are quite obvious. First of all, the variational expansion, Eq.(23), and each of its basis functions has the correct asymptotics at long-range interparticle distances r_{ij} . Therefore, this variational expansion, Eq.(23), is the truly highly accurate expansion for bound state computations in all four-body systems, where particles interact with each other either by the Coulomb, or Yukawa, or exponential potential, or by a potential which is represented as a finite linear combination of these potentials. In positronium hydrides the interaction potential between each pair of particles is a pure Coulomb potential.

Second, the actual convergence of the four-body exponential variational expansion, Eq.(23), upon the total number of basis functions N used, is significantly faster than for multi-dimensional gaussoids, Eq.(4). In fact, for an arbitrary Coulomb four-body system the exponential expansion, Eq.(23), converges faster (and even significantly faster) than other variational expansions currently known and used for four-body systems. For many bound state properties the actual convergence of our exponential variational four-body expansion, Eq.(23), upon the total number of basis functions N used, is usually monotonic and onesided. Third, accurate numerical calculations of the bound state properties, or expectation values, including interparticle cusp values, various delta-functions, etc, are relatively easy to perform by applying the exponential expansion, Eq. (23). In fact, for the pure exponential basis set, Eq.(23), one finds no difference between three- and four-body cases. Other advantages of our exponential four-body variational expansion, Eq. (23), are discussed in [16] and [17]. After these papers were published, Frank Harris wrote an effective computer code for this method. By using his code and 40 basis functions (four-dimensional exponents) we have found the following total energy of the positronium $^{\infty}$ HPs hydride: E = -0.7890967075a.u. Analogous value of the total energy obtained with the use of 50 four-body exponents is $E = -0.78913367330 \ a.u.$ is very close to the 'exact' total energy of this system (see above). These results indicate clearly that our exponential variational expansion developed in [16] and [17] is highly accurate for bound state computations of arbitrary four-body systems.

Note that the exponential variational expansion, Eq.(23), has nothing to do with the traditional atomic Hyllearaas expansion (or James-Coolidge expansion [52]) used for threeelectron atoms and ions (see, e.g., [53]). Indeed, our exponential variational expansion, Eq.(23), is truly correlated four-body expansion which can be applied for highly accurate numerical computations of bound states in arbitrary four-body systems (masses of particles can be arbitrary!), and it is not restricted to the one-center, three-electron atomic systems only.

Another interesting four-body variational expansion is written in the form [18]

 $\Psi_N(r_{14},r_{24},r_{34},r_{12},r_{13},r_{23}) =$

$$\mathcal{A}_{S} \sum_{k=1}^{N} C_{k} r_{23}^{n_{1}(k)} r_{13}^{n_{2}(k)} r_{12}^{n_{3}(k)} r_{14}^{m_{1}(k)} r_{24}^{m_{2}(k)} r_{34}^{m_{3}(k)} \exp\left(-\alpha_{14}^{(k)} r_{14} - \beta_{24}^{(k)} r_{24} - \gamma_{34}^{(k)} r_{34}\right) , \quad (24)$$

where $r_{ij} = r_{ji}$ are the four-body relative coordinates, the operator \mathcal{A}_S is the symmetrization operator, while $\alpha_i, \beta_i, \gamma_i$ are the 3N non-linear parameters of this expansion which are varied in actual computations. In Eq.(24) all powers $n_i(k)$ and $m_j(k)$ of the relative coordinates are assumed to be non-negative and symbol 4 always designates the heaviest particle. The variational expansion, Eq.(24), represents the bound states with L = 0 (or S-states, for short). Possible generalizations of this variational expansion to the rotationally excited states with $L \geq 1$ are relatively simple and straightforward, but here we cannot discuss that interesting problem.

In our earlier paper [18] the variational expansion, Eq.(24), was called the semiexponential variational expansion in the relative four-body coordinates. In applications to actual four-body systems the powers of relative coordinates in Eq.(24) are chosen by the 'families' (or sub-spaces) of polynomial parts of the basis functions. As is well known (see, e.g., [54]) polynomials of degree $\leq \Omega$ form the $(\Omega + 1)$ -dimensional vector space. To construct the trial functions for the positronium hydride $^{\infty}$ HPs we have chosen two families of polynomial wave functions with $\Omega = 28$ and $\Omega = 84$. After a few optimizations our current results (total energies) for the ground S-state in the $^{\infty}$ HPs hydride are E(28) = $-0.781757154783 \ a.u.$ and $E(84) = -0.787961386805 \ a.u.$, respectively. These energies are obviously less accurate than values obtained above with the use of exponential four-body expansion, Eq. (23), but they are much better than than analogous energies obtained with the original James-Coolidge expansion for this system (see, e.g., [15]): E(28) = -0.778527035a.u. and E(84) = -0.786802750 a.u. In other words, our semi-exponential variational expansion, Eq.(24), is significantly more flexible and accurate than the standard Hylleraas (or James-Coolidge) expansion developed long ago, which is still applied for accurate bound state calculations of many atomic-like (or one-center) Coulomb four-body systems.

To conclude this Section we have to note that two our variational expansions, Eqs.(23) and (24), have a great potential for highly accurate, bound state computations in various four-body systems. However, at that time these expansions cannot compete with the KT-variational expansion in terms of overall accuracy, numerical stability and general convenience in applications. It is clear that each of these 'new' variational expansions must be substantially improved in order to reproduce 7 - 9 stable decimal digits for the ground state

energy of all positronium hydrides. Then (and only then) these variational expansions can be considered as effective and fast methods developed for highly accurate solutions of the Coulomb for-body problem (bound states). The same criterion must be applied to other 'alternative' methods developed for similar problems, e.g., for different molecular orbital methods [55] and numerous versions of exponentially correlated Hylleraas configuration Interactions (E-Hy-CI) [56]. In any case we did not want to transform this short Section into a review of all methods which can be applied, in principle, for highly accurate, variational (and non-variational) calculations of the ground state(s) in positronium hydrides.

VII. CONCLUSION

We have investigated the bound state properties of the four-body positronium hydrides ${}^{\infty}$ HPs, ¹HPs, DPs, TPs and four-lepton, muonic hydride MuPs. Annihilation of the electronpositron pair(s) in all positronum hydrides is considered in detail. The rates of the total, two- and three-photon annihilations of the (e^-, e^+) -pair in the HPs hydrides have been determined to very good accuracy. These $\Gamma, \Gamma_{2\gamma}$ and $\Gamma_{3\gamma}$ rates are the largest values in comparison to analogous rates of other annihilation processes in HPs. We have also evaluated a number of other annihilation rates, including the $\Gamma_{4\gamma}, \Gamma_{5\gamma}$ rates of the four- and five-photon annihilations and two different rates of one-photon annihilation $\Gamma_{1\gamma}^a, \Gamma_{1\gamma}^F$ of the two different kinds. The hyperfine structure of these posironium hydrides is considered. In particular, we have determined the hyperfine structure splitting in each of the positronium hydrides ¹HPs, DPs, TPs and MuPs. The bound state properties of some four-body exitonic ${}^{M}h^+e_2^-e^+$ complexes with unit electrical charges are also briefly discussed.

We have also investigated the mass-dependence of the total energies and some other bound state properties of the four-body exitonic ${}^{M}h^{+}e_{2}^{-}e^{+}$ complexes (with $M \geq 1$ and $M \leq 1$). Some of these systems are similar to the regular positronium hydrides HPs. The both cases, when $M \geq 1$ and $M \leq 1$ of these exitonic ${}^{M}h^{+}e_{2}^{-}e^{+}$ complexes have been considered. Finally, we discuss a possibility to apply our new four-body exponential and semi-exponential variational expansions in the relative coordinates, Eqs.(23) and (24), for accurate computations of the bound states in positronium hydrides. It is shown that the both these variational expansions have a great potential in applications to bound state calculations of positronium hydrides and many other four-body systems. The both these variational expansions allow one to determine interparticle cusp values and a few other properties which cannot be obtained in the direct computations with the use of KT-expansion, Eq.(4).

Acknowledgments

This paper arose out of a large number of ideas and approaches which I/we have developed over many years. During this work I have greatly benefited from numerous discussions with Anand K. Bhatia, Richard J. Drachman, Frank E. Harris, Sergrei I. Kryuchkov, Sergiy Bubin, Frederick W. King and others. I would like to thank all of them for their valuable help and inspiration. Unfortunately, when this paper was in preparation, Richard J. Drachman was tragically killed in an automobile accident on 6th of April 2021. The present contribution is dedicated to the memory of this dear friend.

- Interactions of Positrons with Matter and Radiation (A.K. Bhatia, editor, MDPI, Basel, Switzerland (ISSN 2218-2004) (available at: https://www.mdpi.com/journal/atoms/special issues/ Positron-phys) (2020)).
- [2] R.J. Drachman, Can. J. Phys. **60**, 494 (1982).
- [3] J.M. Ziman, Principles of the Theory of Solids, (2nd ed., Cambridge University Press, Cambridge, UK (1972)).
- [4] A.S. Davydov, *Theory of Solids*, (Nauka, Moscow (1980), (in Russian)).
- [5] A. Ore, Phys. Rev. 83, 665 (1951).
- [6] S.K. Houston and R.J. Drachman, Phys. Rev. A 7, 819 (1974).
- [7] Y.K. Ho, Phys. Rev. A **34**, 609 (1986).
- [8] B.A. Page and P.A. Frazer, J. Phys. B 7, L389 (1974).
- [9] D.M. Schrader, Chemical Stability and Approximate Quantum Mechanics, in: Positron Annihilation, edited by P.G. Coleman, S.C. Sharma and L.M. Diana (North-Holland, Amsterdam, 1982), p. 71 and references therein.
- [10] D.M. Schrader, F.M. Jacobsen, N.-P. Fradsen and U. Mikkelsen, Phys. Rev. Lett. 69, 57 (1992).
- [11] A.M. Frolov and V.H. Smith, Jr., Phys. Rev. A 55, 2662 (1997).
- [12] S. Bubin and L. Adamowicz, Phys. Rev. A 74, 052502 (2006).

- [13] S. Bubin, M. Stanke, D. Kedziera and L. Adamowicz, Phys. Rev. A 75, 062504 (2007).
- [14] Z.-C. Yan and Y.K. Ho, Phys. Rev. A **60**, 5098 (1999).
- [15] A.M. Frolov and V.H. Smith, Jr., Phys. Rev. A 56, 2417 (1997).
- [16] A.M. Frolov and V.H. Smith, Jr., J. Chem. Phys. 115, 1187 (2001).
- [17] F.E. Harris, A.M. Frolov and V.H. Smith, Jr., J. Chem. Phys. 119, 8833 (2003).
- [18] A.M. Frolov, Phys. Lett. A **374**, 2361 (2010).
- [19] N.N. Kolesnikov and V.I. Tarasov, Izv. Vuzov, ser. Phys. 5, 98 (1977) [Proceedings of Colleges, Physics (1977), (in Russian)].
- [20] N.N. Kolesnikov and V.I. Tarasov, Yad. Phys. 35, 609 (1982) [Sov. Phys. Nucl. Phys. 35, 354 (1982)].
- [21] A.M. Frolov, J. Phys. A **39**, 15421 (2006).
- [22] see, e.g., https://physics.nist.gov/cgi-bin/cuu/Value?
- [23] T. Kato, Theor. Commun. Pure & Appl. Math. 10, 151 (1957).
- [24] J. Sucher and R.J. Drachman, Phys. Rev. A 20, 424 (1979).
- [25] R.A. Ferrell, Rev. Mod. Phys. 28, 308 (1956).
- [26] A.K. Bhatia, Atoms 9, 27 (2016).
- [27] W. Heitler, The Quantum Theory of Radiation (3rd ed., Oxford University Press, London, UK (1954)).
- [28] A. Akhiezer and V.B. Berestetskii, *Quantum Electrodynamics* (4th ed., Science, Moscow, (1981), (in Russian)).
- [29] W. Greiner and J. Reinhardt, Quantum Electrodynamics (4th ed., Springer Verlag, Berlin, 2009).
- [30] V.B. Berestetskii, E.M. Lifshitz and L.P. Pitaevskii, *Relativistic Quantum Theory* (Pergamon Press, Oxford, (1971)).
- [31] G.P. Lepage, P.B. Mckenzie, K.H. Streng and P.M. Zerwas, Phys. Rev. A 28, 3090 (1983).
- [32] I. Harris and L.M. Brown, Phys. Rev. **105**, 1656 (1957).
- [33] A.M. Frolov, Phys. Rev. A 80, 014502 (2009).
- [34] A.P. Mills, Jr., Phys. Rev. Lett. 46, 717 (1981).
- [35] A.P. Mills, Jr., Phys. Rev. Lett. **50**, 671 (1983).
- [36] S.I. Kryuchkov, J. Phys. B **27**, L61 (1994).
- [37] A.M. Frolov, S.I. Kryuchkov and V.H. Smith, Jr., Phys. Rev. A 51, 4514 (1995).

- [38] A.M. Frolov, Phys. Rev. A 60, 2834 (1999).
- [39] Y.-C. Chu and V. Ponish, Jr., Phys. Rev. C 33, 2222 (1986).
- [40] E. Fermi and G.E. Uhlenbeck, Phys. Rev. 44, 510 (1933).
- [41] H. Bethe, Proc. Roy. Soc. A **150**, 129 (1935).
- [42] A.M. Frolov, Phys. Lett. A, **376**, 2548 (2012).
- [43] G.H. Wannier, Phys. Rev. **52**, 191 (1937).
- [44] N.F. Mott, Trans. Farad. Soc. **34**, 500 (1938).
- [45] M. Lampert, Phys. Rev. Lett. 1, 450 (1958).
- [46] R.R. Sharma and S. Rodrigues, Phys. Rev. **153**, 823 (1967).
- [47] A.M. Frolov, Chem. Phys. Lett. **626**, 49 (2015).
- [48] I.S. Gradstein and I.M. Ryzhik, Tables of Integrals, Series and Products (6th revised ed., Academic Press, New York, (2000)).
- [49] R.D. Poshusta, J. Phys. B 18, 1887 (1985).
- [50] A.M. Frolov and D.M. Bishop, Phys. Rev. A 45, 6236 (1992).
- [51] A.M. Frolov and V.H. Smith, Jr., J. Phys. B **29**, L433 (1996).
- [52] H.M. James and A.S. Coolidge, Phys. Rev. 49, 688 (1936).
- [53] F.W. King and V. Shoup, Phys. Rev. A **33**, 2940 (1986).
- [54] I. M. Gelfand, Lectures on Linear Algebra, (Dover Publ., Inc., Mineola, New York, (1989)), Chpt. I.
- [55] J. Charry, J. Romero, M. T. do N. Varella, and A. Reyes, Phys. Rev. A 89, 052709 (2014).
- [56] M.B. Ruiz, J.S. Sims and B. Padly, Advan. in Quant. Chem. 83, 171 (2021).

TABLE I: The expectation values $\langle \hat{A} \rangle$ in atomic units *a.u.* of some bound state properties for the ground (bound) ${}^{1}S(L=0)$ -states of the positronium hydrides ${}^{M}H^{+}e_{2}^{-}e^{+}$. In this Table the notations '+' and '-' stand for the positron and electron, respectively, while the symbol 'p' means the heavy, positively charged nucleus of the hydrogen isotope.

system	$^{\infty}\mathrm{H}^{+}e_{2}^{-}e^{+}$	${}^{3}H^{+}e_{2}^{-}e^{+}$	${}^{2}H^{+}e_{2}^{-}e^{+}$	${}^{1}H^{+}e_{2}^{-}e^{+}$	$\mu^+ e_2^- e^+$
$\langle r_{}^{-2} \rangle$	0.213911	0.213823	0.213779	0.213648	0.211598
$\langle r_{+-}^{-2}\rangle$	0.349144	0.349120	0.349120	0.349073	0.348519
$\langle r_{p-}^{-2} \rangle$	1.207065	1.206593	1.206358	1.205651	1.194598
$\langle r_{p+}^{-2}\rangle$	0.172164	0.172114	0.172089	0.172015	0.170851
$\langle r_{}^{-1} \rangle$	0.3705554	0.3704805	0.3704432	0.3703313	0.3685790
$\langle r_{+-}^{-1} \rangle$	0.4184965	0.4184740	0.4184628	0.4184292	0.4179026
$\langle r_{p-}^{-1} \rangle$	0.7297093	0.7295588	0.7294839	0.7292589	0.7257325
$\langle r_{p+}^{-1}\rangle$	0.3474623	0.3474090	0.3473824	0.3473027	0.3460529
$\langle r_{} \rangle$	3.574770	3.575516	3.575888	3.577003	3.594554
$\langle r_{+-} \rangle$	3.480262	3.480563	3.480713	3.481163	3.488243
$\langle r_{p-} \rangle$	2.311517	2.312063	2.312335	2.313151	2.325995
$\langle r_{p+} \rangle$	3.661608	3.662232	3.662543	3.663477	3.678178
$\langle r_{}^2 \rangle$	15.87515	15.88199	15.88539	15.89563	16.05703
$\langle r_{+-}^2 \rangle$	15.58407	15.58716	15.58870	15.59332	15.666248
$\langle r_{p-}^2 \rangle$	7.812899	7.816816	7.818765	7.824627	7.917145
$\langle r_{p+}^2 \rangle$	16.25421	16.26007	16.26299	16.27176	16.41019
$\langle r_{}^3 \rangle$	84.5457	84.6024	84.6307	84.7156	86.0593
$\langle r_{+-}^3 \rangle$	84.3699	84.3979	84.4118	84.4537	85.1158
$\langle r_{p-}^3\rangle$	35.2118	35.2397	35.2535	35.2952	35.9552
$\langle r_{p+}^3\rangle$	85.0937	85.1425	85.1669	85.2400	86.3974
$\langle r_{}^4 \rangle$	527.879	528.372	528.617	529.354	541.059
$\langle r_{+-}^4 \rangle$	532.936	533.195	533.323	533.711	539.848
$\langle r_{p-}^4\rangle$	198.871	199.089	199.198	199.525	204.726
$\langle r_{p+}^4 \rangle$	516.036	516.456	516.665	517.294	527.286
$\langle -\tfrac{1}{2}\nabla^2\rangle$	0.32617335	0.32606754	0.32601491	0.32585679	0.32338494
$\langle -\frac{1}{2} \nabla^2_+ \rangle$	0.13685044	0.13684430	0.13684125	0.13683208	0.13668953

TABLE II: The total energies E and expectation values of some interparticle delta-functions $\langle \delta_{ij} \rangle$ in atomic units *a.u.* for the ground (bound) ${}^{1}S(L=0)$ -states of the positronium hydrides ${}^{M}H^{+}e_{2}^{-}e^{+}$. In this Table the notations '+' and '-' stand for the positron and electron, respectively, while the symbol 'p' means the heavy, positively charged nucleus of the hydrogen isotope.

system	$^{\infty}H^{+}e_{2}^{-}e^{+}$	${}^{3}H^{+}e_{2}^{-}e^{+}$	${}^{2}H^{+}e_{2}^{-}e^{+}$	${}^{1}H^{+}e_{2}^{-}e^{+}$	$\mu^+ e_2^- e^+$
E	-0.7891967651	-0.789087847	-0.789033651	-0.788870705	-0.786317295
$\langle \delta_{+-} \rangle$	$2.448351 \cdot 10^{-2}$	$2.448163 \cdot 10^{-2}$	$2.448070 \cdot 10^{-2}$	$2.447789 \cdot 10^{-2}$	$2.443306 \cdot 10^{-2}$
$\langle \delta_{p+} \rangle$	$1.625353 \cdot 10^{-3}$	$1.624877 \cdot 10^{-3}$	$1.624642 \cdot 10^{-3}$	$1.623930 \cdot 10^{-3}$	$1.612725 \cdot 10^{-3}$
$\langle \delta_{+} \rangle$	$3.69845 \cdot 10^{-4}$	$3.69619 \cdot 10^{-4}$	$3.69506 \cdot 10^{-4}$	$3.69168 \cdot 10^{-4}$	$3.63833 \cdot 10^{-4}$

TABLE III: A number of the few-photon annihilation rates $\Gamma_{n\gamma}$ (in sec^{-1}) predicted for the ground (bound) $1^{1}S_{e}(L=0)$ -states in the positronium hydrides MuPs, ¹HPs, DPs, TPs and [∞]HPs.

rate	$^{\infty}\mathrm{HPs}$	TPs	DPs	$^{1}\mathrm{HPs}$	MuPs
Г	$2.5237439 \cdot 10^9$	$2.5235504 \cdot 10^9$	$2.5234541 \cdot 10^9$	$2.5231646 \cdot 10^9$	$2.5185995 \cdot 10^9$
$\Gamma_{2\gamma}$	$2.4568126{\cdot}10^9$	$2.4566243 \cdot 10^9$	$2.4565305 \cdot 10^9$	$2.4562487 \cdot 10^9$	$2.4518047 \cdot 10^9$
$\Gamma_{3\gamma}$	$6.6931229 \cdot 10^7$	$6.6926097 \cdot 10^7$	$6.6923543 \cdot 10^7$	$6.6915865 \cdot 10^7$	$6.6794796 \cdot 10^7$
$\Gamma_{4\gamma}$	$3.6320642 \cdot 10^3$	$3.6317857 \cdot 10^3$	$3.6316471 \cdot 10^3$	$3.6312305 \cdot 10^3$	$3.6246606 \cdot 10^3$
$\Gamma_{5\gamma}$	63.91944	63.91454	63.91210	63.90477	63.78915
$\Gamma^a_{1\gamma}$	0.39417	0.39392	0.39380	0.39344	0.38776
$\Gamma^F_{1\gamma}$	1.48715	1.48704	1.48698	1.48681	1.48412

TABLE IV: The hyperfine structure splittings ΔE_{hyps} (in MHz) for the ground (bound) $1^{1}S_{e}(L = 0)$ -states in the positronium hydrides MuPs, ¹HPs, DPs and TPs

system	factor \mathcal{A}	I_H	$\langle \delta_{p+} \rangle^{(a)}$	ΔE_{hyps}
MuPs	14229.176639	$\frac{1}{2}$	$1.621275 \cdot 10^{-3}$	23.0693728
$^{1}\mathrm{HPs}$	-4469.881841	$\frac{1}{2}$	$1.623930 \cdot 10^{-3}$	-7.2519050
DPs	-686.154150	1	$1.624642 \cdot 10^{-3}$	-1.6721323
TPs	-4767.754403	$\frac{1}{2}$	$1.624877 \cdot 10^{-3}$	-7.2630081

^(a)The expectation value of the positron-nucleus delta-function in atomic units.

TABLE V: The expectation values $\langle \hat{A} \rangle$ in atomic units *a.u.*) of some bound state properties for the ground (bound) ${}^{1}S(L=0)$ -states of the model positronium hydrides ${}^{M}h^{+}e_{2}^{-}e^{+}$. In this Table the notations '+' and '-' stand for the positron and electron, respectively, while the symbol 'h' means the heavy, positively charged quasi-nucleus with mass $M(=Mm_{e})^{(a)}$.

system	E	$\langle r_{+-} \rangle$	$\langle r_{} \rangle$	$\langle r_{h-} \rangle$	$\langle \delta({f r}_{+-}) angle$
$^{12}h^+e_2^-e^+$	-0.744058235415	3.612337700	3.940177538	2.553742718	0.02394728
${}^{11}h^+e_2^-e^+$	-0.740373921176	3.623763252	3.964626240	2.574977893	0.02390840
${}^{10}h^+e_2^-e^+$	-0.736033440341	3.637343176	3.993755510	2.600284075	0.02385773
${}^{9}h^{+}e_{2}^{-}e^{+}$	-0.730843941385	3.653774543	4.029094985	2.630980145	0.02380552
${}^{8}h^{+}e_{2}^{-}e^{+}$	-0.724528577928	3.674032011	4.072811334	2.668967075	0.02373458
$^{7}h^{+}e_{2}^{-}e^{+}$	-0.716675039870	3.699651073	4.128317923	2.717213004	0.02366023
${}^{6}h^{+}e_{2}^{-}e^{+}$	-0.706640804107	3.733036217	4.201037748	2.780491629	0.02355747
${}^{5}h^{+}e_{2}^{-}e^{+}$	-0.693365530158	3.778339650	4.300413160	2.867138301	0.02343085
${}^{4}h^{+}e_{2}^{-}e^{+}$	-0.674963081481	3.843250936	4.444240116	2.993047371	0.02326605
${}^{3}h^{+}e_{2}^{-}e^{+}$	-0.647714874405	3.943731008	4.670529754	3.192898318	0.02302489
${}^{2}h^{+}e_{2}^{-}e^{+}$	-0.603091391970	4.118241595	5.076718755	3.559953733	0.02266962

^(a)The total energy of the four-body bi-positronium molecule Ps₂, or ${}^{1}h^{+}e_{2}^{-}e^{+}$ positronium hydride in our current notations, is -0.51600379066033 *a.u.*

TABLE VI: The coefficients C_k in the ten-term mass-interpolation formula, Eq.(20) for the total energies of the ground (bound) $1^1S_e(L=0)$ -states in the four-body ${}^{M}h^+e_2^-e^+$ systems, where $M \ge 1$. The coefficients B_k in the seven-term mass-interpolation formula, Eq.(21) for the total energies of the ground (bound) $1^1S_e(L=0)$ -states in the four-body ${}^{m}h^+e_2^-e^+$ systems, where $m \le 1$.

k	C_k	C_k^{\star}	B_k	system	$E^{(a)}$ (predicted)
1	0.1760927482	0.2084671422	-0.1734561261	$\mu^+ e_2^- e^+$	-0.7863172536
2	0.1699737489	0.0861748927	-0.0601985845	$^{100}h^+e_2^-e^+$	-0.7832819721
3	-0.0831406229	0.0204254266	-0.0132568021	$^{75}h^+e_2^-e^+$	-0.7813436733
4	0.1018702975	0.0102296923	-0.0029164214	$^{25}h^+e_2^-e^+$	-0.7664017141
5	-0.0642471972	-0.0016461962	0.0006601439	$^{15}h^+e_2^-e^+$	-0.7523881511
6	0.0365809780	0.0031380338	0.0004238099	$^{0.85}h^+e_2^-e^+$	-0.4951481654
7	-0.0152567422	-0.0015202684	0.0010964638	$^{0.75}h^+e_2^-e^+$	-0.4792885124
8	0.0048478700	0.0007979493		$^{0.55}h^+e_2^-e^+$	-0.4416910665
9	-0.0011064536	-0.0002802924		$^{0.3}h^+e_2^-e^+$	-0.3786180518
10	0.0002520781	0.0001083263		$^{0.15}h^+e_2^-e^+$	$-0.3279908164^{(a)}$

^(a)Also, the total energy of the four-body ${}^{0.005}h^+e_2^-e^+$ system is -0.26450300278 *a.u.*, which is close to the total energy of the three-body Ps⁻ ion ($E \approx -0.2620050702329801...$ *a.u.*, see, the main text).