

Is every triangle a trajectory of an elliptical billiard?

Vladimir Dragović^{1,3} and Milena Radnović^{2,3}

¹THE UNIVERSITY OF TEXAS AT DALLAS, DEPARTMENT OF MATHEMATICAL SCIENCES

²THE UNIVERSITY OF SYDNEY, SCHOOL OF MATHEMATICS AND STATISTICS

³MATHEMATICAL INSTITUTE SANU, BELGRADE

vladimir.dragovic@utdallas.edu, milena.radnovic@sydney.edu.au

Abstract

Using Marden’s Theorem from geometric theory of polynomials, we show that for every triangle there is a unique ellipse such that the triangle is a billiard trajectory within that ellipse. Since 3-periodic trajectories of billiards within ellipses are examples of the Poncelet polygons, our considerations provide a new insight into the relationship between Marden’s Theorem and the Poncelet Porism, two gems of exceptional classical beauty. We also show that every parallelogram is a billiard trajectory within a unique ellipse. We prove a similar result for the self-intersecting polygonal lines consisting of two pairs of congruent sides, named “Darboux butterflies”. In each of three considered cases, we effectively calculate the foci of the boundary ellipses.

Keywords: Elliptical billiards; Marden’s Theorem; 3-periodic billiard trajectories; 4-periodic billiard trajectories; Darboux butterflies; conics

AMS subclass: 51N20, 30C15, 37J35, 70H06

Contents

1	Introduction	2
2	Triangles	3
2.1	Triangular billiards	3
2.2	Theorems of Ceva and Menelaus	5
3	Conics	7
3.1	Elliptical billiards	7
3.2	Pascal’s theorem	10
4	The Siebeck–Marden theorem	11
5	3-periodic trajectories of billiards within ellipses	14
6	Convex 4-periodic trajectories of billiards within ellipses	14
6.1	Is every parallelogram a trajectory of an elliptical billiard?	14
6.2	An analytic proof	16
7	Nonconvex 4-periodic elliptical billiard trajectories	18
7.1	Characterization of nonconvex 4-periodic trajectories	18
7.2	Analytic approach	21

arXiv:2405.08922v2 [math.DS] 11 Jul 2024

1 Introduction

Recall that *mathematical billiard* in a planar domain is a dynamical system where a particle moves without constraints within the domain, and obeys *the billiard reflection law* when it hits the boundary [KT1991, Tab2005]. Thus, billiard trajectories are polygonal lines with vertices at the boundary, such that two consecutive sides form congruent angles with the tangent line to the boundary at their joint vertex, see Figure 1. Mathematical billiards are idealized models in many aspects: a usual billiard




Figure 1: Billiard reflection law: the angle of incidence with the tangent line at the bouncing point on the boundary equals the angle of reflection.

ball is replaced by a material point, the friction and spin are neglected. Such models have natural applications, for example in geometric optics. Here, we assume that the billiard particle is of the unit mass and it moves under the inertia between the impacts, i.e. uniformly along straight lines.

Billiards within ellipses have been intensively studied, see for example [Bol1990, Koz2003, KT1991, DR2011, DR2014, ADSK2016, KS2018, BM2017, BM2022, Glu2021, CZ2023, GRK2021, GKR2023, DGR2022, FV2023] and references therein.

In this note, we will give the affirmative answer to the title questions. This answer may appear surprising at the first glance, if we recall that a conic is defined with various sets of five conditions, while a triangle as an inscribed billiard trajectory imposes six conditions, three points and three tangent lines. Regardless of these seemingly overdetermined conditions, the solution conic exists and it is unique always, and it is always an ellipse. The main ingredients in our considerations of triangular trajectories are Marden's theorem from the geometry of polynomials [Mar1966] and the classical Ceva's theorem from elementary geometry. Periodic trajectories of billiards within ellipses can be seen as an instance of so-called Poncelet polygons, which are closed polygonal lines inscribed in one conic and circumscribed about the other conic, see e.g. [DR2011]. The Poncelet Theorem from projective geometry of conics states that if for a given pair of conics there is one Poncelet polygon, then there are infinitely many such polygons and all have the same number of sides. Thus, we are in this work dealing with an interaction between Marden's theorem and the Poncelet theorem, which are both recognized by their exceptional classical beauty. Previously, a strong relationship between these two theorems was observed in [Dra2011] in a different context.

This paper is organised as follows. We review basic facts about billiards within triangles and the theorems of Ceva, Menelaus, and Simson in Section 2; conics and elliptical billiards in Section 3; and Marden's theorem in Section 4. In Section 5, we prove that each triangle is a billiard trajectory within a unique ellipse. In Section 6, we give a complete characterization of convex 4-periodic elliptical billiard trajectories as parallelograms and show the converse statement: that each parallelogram is a billiard trajectory within unique ellipse, see Theorem 6.3. Nonconvex 4-periodic elliptical billiard trajectories are considered in Section 7, where they are characterized as so-called *Darboux butterflies*. We also show that every Darboux butterfly is a billiard trajectory within a unique ellipse, see Theorem 7.8. In each of the three cases, of a triangle, a parallelogram, and a Darboux butterfly, we effectively calculate the foci of the boundary ellipse. We provide multiple proofs for our statements, allowing interplay of various classical and modern geometry results.

2 Triangles

In this section consists of two parts: in Subsection 2.1, we review main properties of the billiards within a triangle in the Euclidean plane, and in Subsection 2.2 we revisit the Ceva theorem.

2.1 Triangular billiards

A trajectory of such a billiard is a polygonal line, finite or infinite, with vertices on the sides of the triangle, such that consecutive edges of the trajectory satisfy the billiard reflection law. i.e. they form the same angle with the side of the triangle which their common vertex lie on, see Figure 2.




Figure 2: Billiard motion in a triangle.

The reflection is not well defined in the vertices of the triangle, thus we omit from our consideration trajectories falling in a vertex.

Remark 2.1 *Note the following minimization property of the billiard reflection: if X, Y are two points on the same side of the a given line ℓ , then the length $XL + LY$, for $L \in \ell$, will be the smallest when the segments XL and LY satisfy the billiard reflection law off ℓ , see Figure 3.*




Figure 3: The minimization property of billiard reflection: the shortest path connecting points X and Y which visits line ℓ is the billiard path XY . Its length equals the segment XY' , where Y' is symmetric to Y with respect to ℓ . For any other point $L' \in \ell$, we have $XL' + L'Y = XL' + L'Y' > XY'$.

One of the first natural questions for any dynamical system is to establish if its periodic trajectories exist and find them if they do. Thus, next we recall a proof of the existence a periodic trajectory within any acute triangle.

Theorem 2.2 (see e.g. [DR2011, Theorem 2.2]) *Let $\triangle ABC$ be an acute triangle, and K, L, M the feet of its altitudes. Then $\triangle KLM$ is the triangle with minimal perimeter inscribed in $\triangle ABC$ and it represents a unique 3-periodic trajectory of the billiard with $\triangle ABC$.*

Proof. Let M' be an arbitrary given point on side AB . In order to find points $K' \in BC$, $L' \in AC$ such that the triangle $K'L'M'$ has the minimal perimeter, denote by M_1, M_2 the points symmetric to M' with respect to sides BC and AC respectively. Then K' and L' are intersection points of M_1M_2 with sides BC and AC respectively, see Figure 4.




Figure 4: For a fixed point M' on AB , we construct points K', L' on the remaining two sides of the triangle, such that $\triangle K'L'M'$ has smallest possible perimeter.

The perimeter of $\triangle K'L'M'$ is equal to the length of segment M_1M_2 . We observe that M_1M_2 is a side of the isosceles triangle CM_1M_2 , whose angle $\angle M_1CM_2 = 2\angle BCA$ does not depend on the choice of point M' . Since $CM_1 \cong CM_2 \cong CM'$, the segment M_1M_2 is shortest when the point M' is such that CM' is the altitude of the triangle ABC from the vertex A , i.e. the minimal perimeter is attained when $M' = M$. It immediately follows that points K' and L' , as constructed above, are then also the feet of the corresponding altitudes, see Figure 5.




Figure 5: The vertices of the triangle with smallest perimeter inscribed in $\triangle ABC$ are the feet K, L, M of the altitudes.

The minimizing property from Remark 2.1 implies that $\triangle KLM$, as the triangle with smallest perimeter inscribed in $\triangle ABC$ is a billiard trajectory within $\triangle ABC$.

Now, let us show uniqueness. Suppose that $K_1L_1M_1$ is another periodic trajectory within the triangle ABC , such that $K_1 \in BC$, $L_1 \in AC$, $M_1 \in AB$. Then, according to the billiard reflection law, we can denote: $k := \angle CK_1L_1 = \angle BK_1M_1$, $l := \angle CL_1K_1 = \angle AL_1M_1$, $m := \angle AM_1L_1 = \angle BM_1K_1$. Denoting by $\angle A, \angle B, \angle C$ the angles of the triangle ABC , we get that the sums of the angles in the triangles $AL_1M_1, BK_1M_1, CK_1L_1$ are:

$$\angle A + l + m = \angle B + k + m = \angle C + k + l = 180^\circ,$$

which gives: $k = \angle A, l = \angle B, m = \angle C$.

From there, the sides of the triangle $K_1L_1M_1$ are parallel to the corresponding sides of the triangle KLM . The assumption that $K_1L_1M_1$ is a periodic billiard trajectory then implies $K_1 = K, L_1 = L, M_1 = M$, as shown in Figure 6. Namely, if K_1 is between K and C , then M_1 lies in the interior

of $\triangle ABC$. This follows from the similarity of triangles KLM and $K_1L_1M_1$, by using the Thales theorem. The same argument shows that M_1 lies outside $\triangle ABC$ if K_1 is between K and B . \square




Figure 6: The trajectories with segments parallel to the segments of KLM are 6-periodic.

The previous proof essentially relies on the assumption that $\triangle ABC$ is acute, since otherwise some of the feet of the altitudes are not inner points of the sides. Thus, additional discussion is needed for the right and obtuse triangles.

It is easy to see that there are periodic billiard trajectories within a right triangle: one of them, the polygonal line $KLMNMLK$, is shown in Figure 7.




Figure 7: A periodic trajectory of the billiard within a right triangle.

After such elementary considerations for acute and right triangles, one can stay amazed by the fact that it is not known if billiards within general obtuse triangles have any periodic trajectories! There are examples for some special cases and also an intriguing computer-based proof for the existence of periodic billiard trajectories when the obtuse angle does not exceed 100° , see [Sch2006, Sch2009].

2.2 Theorems of Ceva and Menelaus

In this section, we will formulate the two classical theorems related to the geometry of triangles.

Theorem 2.3 (Ceva's Theorem) *Let ABC be a given triangle and K, L, M points on the lines BC, AC, AB respectively, such that none of them coincides with a vertex of the triangle. Then the lines AK, BL, CM are either concurrent or parallel if and only if:*

$$\frac{AM}{MB} \cdot \frac{BK}{KC} \cdot \frac{CL}{LA} = 1. \quad (2.1)$$

Ceva's theorem is illustrated in Figure 8.

Remark 2.4 *In Ceva's Theorem, the signed lengths of segments are used in (2.1). For example, in Figure 8, the quantities AM/MB and BK/KC are negative, while CL/LA is positive.*

Remark 2.5 *Notice that Ceva's theorem can be applied to an acute triangle ABC and its 3-periodic billiard trajectory KLM , since the altitudes intersect at one point, see Figure 5 and Theorem 2.2.*




Figure 8: Ceva's theorem.

We will use Ceva's theorem to prove the following statement.

Lemma 2.6 *Let ABC be a given triangle and K, L, M its feet of the altitudes from A, B, C respectively. Then there exist real numbers m_1, m_2, m_3 such that point K divides BC in the ratio $m_2 : m_3$, point L divides CA in the ratio $m_3 : m_1$, and M divides AB in the ratio $m_1 : m_2$. Moreover*

$$m_1 : m_2 : m_3 = (b^2 + c^2 - a^2) : (a^2 + c^2 - b^2) : (a^2 + b^2 - c^2), \quad (2.2)$$

where a, b, c respectively denote the lengths of the sides BC, AB, AC .

Proof. Since the altitudes of a triangle intersect at the orthocenter, Ceva's theorem implies the existence of numbers m_1, m_2, m_3 . The similarities of right triangles $\triangle KBA \sim \triangle MBC$, $\triangle LCB \sim \triangle KCA$, $\triangle MAC \sim \triangle LAB$ imply the following relations: $KB/AB = MB/BC$, $CL/CB = CK/AC$, $AM/AC = LA/AB$, yielding

$$m_1 : m_2 : m_3 = (AL \cdot AC) : (BM \cdot BA) : (CK \cdot CB).$$

The Pythagorean theorem gives

$$CK^2 = b^2 - AK^2, \quad AL^2 = c^2 - BL^2, \quad BM^2 = a^2 - CM^2,$$

thus

$$m_1 : m_2 : m_3 = b\sqrt{c^2 - BL^2} : c\sqrt{a^2 - CM^2} : a\sqrt{b^2 - AK^2}.$$

Observing that $a \cdot AK = b \cdot BL = c \cdot CM = 2\mathcal{A}$, where \mathcal{A} is the area of the triangle, we get:

$$m_1 : m_2 : m_3 = \sqrt{c^2b^2 - 4\mathcal{A}^2} : \sqrt{a^2c^2 - 4\mathcal{A}^2} : \sqrt{a^2b^2 - 4\mathcal{A}^2}.$$

Finally, applying the Heron formula $\mathcal{A} = \sqrt{s(s-a)(s-b)(s-c)}$, with $s = (a+b+c)/2$, and simplifying, we get (2.2). \square

Theorem 2.7 (Theorem of Menelaus) *Let $\triangle ABC$ be a given triangle and p a line intersecting the lines BC, AC, AB in points P, Q, R respectively, which do not coincide with any of the vertices of the triangle. Then*

$$\frac{AR}{RB} \cdot \frac{BP}{PC} \cdot \frac{CQ}{QA} = -1. \quad (2.3)$$

Moreover, the converse is also true: if points P, Q, R are chosen on the lines AB, BC, CA , such that (2.3) is satisfied, then P, Q, R are collinear.

The theorem of Menelaus is illustrated in Figure 9.




Figure 9: Theorem of Menelaus.




Figure 10: The bisectors of exterior angles of a triangle meet the opposite sides at collinear points.

Corollary 2.8 *For a given triangle, consider the points where the bisectors of its exterior angles meet the opposite sides. Then those three points are collinear.*

Proof. Denote by a, b, c respectively the bisectors of the exterior angles at the vertices A, B, C of the triangle, and by P, Q, R the intersections of those bisectors with the lines BC, AC, AB , as shown in Figure 10.

Since the intersection of the exterior bisector externally divides the opposite side in the ratio of two remaining sides of the triangle, the Theorem of Menelaus (Theorem 2.7) gives that P, Q, R are collinear. \square

Theorem 2.9 (Simson's line) *Let a triangle ABC and a point S be given in the plane. Then the orthogonal projections of that point to the sides of the triangle are collinear if and only if point S belongs to the circumcircle of the triangle. (See Figure 11).*

3 Conics

3.1 Elliptical billiards

In this section, we review most important properties of billiards within an ellipse. The most famous one is the focal property of elliptic billiards, which is illustrated in Figure 12.




Figure 11: Simson's line.

Proposition 3.1 (First focal property of ellipses) *Let \mathcal{E} be an ellipse with foci F_1, F_2 and $A \in \mathcal{E}$ an arbitrary point. Then segments AF_1, AF_2 satisfy the billiard reflection law off \mathcal{E} .*




Figure 12: First focal property of ellipses.

If the billiard particle is launched from one focus of the ellipse, the focal property implies that the segments of the trajectory alternately contain the foci F_1 and F_2 . Next proposition characterizes the trajectories which do not contain the foci.

Proposition 3.2 *Suppose that the billiard particle is traveling along segment s which does not contain any focus F_1, F_2 of the boundary ellipse \mathcal{E} . Then we have:*

- *If both foci of \mathcal{E} are on the same side of s , then there is unique ellipse \mathcal{E}' sharing the same foci with \mathcal{E} and touching the segment. Moreover, after reflection off the boundary, the next segment of the billiard trajectory will also be tangent to \mathcal{E}' , see Figure 13;*
- *If s crosses the segment F_1F_2 , then there is a hyperbola \mathcal{H} sharing the same foci with \mathcal{E} and touching the line containing s . Moreover, after reflection off the boundary, the next segment of the billiard trajectory will also be tangent to \mathcal{H} .*

Proposition 3.2 implies the following key property of elliptical billiards: each trajectory of the billiard within ellipse has a *caustic* – a curve such that each segment of the trajectory lies on a tangent line of that curve, see Figure 14. For elliptical billiards, caustics are ellipses hyperbolas confocal with the boundary, including degenerate ones, which can be identified as horizontal and vertical axes of the ellipse \mathcal{E} .




Figure 13: Two segments satisfying the billiard reflection law of ellipse \mathcal{E} are tangent to the same conic, which is confocal with \mathcal{E} .



Figure 14: The caustics of billiard trajectories.

A direct consequence of Propositions 3.1 and 3.2 is the well-known second focal property of ellipses, illustrated in Figure 15.

Proposition 3.3 (Second focal property of ellipses) *Let \mathcal{E} be an ellipse with foci F_1, F_2 and A an arbitrary point outside ellipse \mathcal{E} . Denote by t_1 and t_2 the two tangents from A to ellipse \mathcal{E} . Then the angle between t_1 and the segment AF_1 is equal to the angle between AF_2 and t_2 .*




Figure 15: Second focal property of ellipses

In the next lemma, we note that the type of the caustic may be determined by the period of a trajectory.

Lemma 3.4 *A periodic trajectory of the elliptic billiard with odd period has an ellipse as the caustic.*

Proof. Suppose that the caustic of a given trajectory is a hyperbola. Denote by F_1, F_2 the focal points of the boundary ellipse. Then every segment of the billiard trajectory intersects the segment F_1F_2 . Thus, for a periodic trajectory there will be an even number of intersections of the trajectory with F_1F_2 . Consequently, the period of a closed trajectory with hyperbola as caustic must be even. \square

3.2 Pascal's theorem

We review here the classical Pascal's theorem. For more details, see, for example [Ber1987] or [DR2011].

Theorem 3.5 (Pascal's theorem) *Let a non-degenerate conic be given, and M, N, O, P, Q, R six points on that conic. Then the intersection points of lines MN and PQ , NO and QR , OP and RM are collinear. (See Figure 16).*




Figure 16: Pascal's theorem.

Pascal's theorem can be also applied in the cases when some of the six given points on the conic coincide. In such cases, the line containing two coinciding points is the tangent line to the conic at that point. Two instances of such limit cases of Pascal's theorem are provided below.

Corollary 3.6 *Let a non-degenerate conic be given together with its four points M, N, P, Q . The intersection points of the tangents to the conic at M and P is collinear with the points of intersection of the pairs of lines MN and PQ , NP and MQ . (See Figure 17).*




Figure 17: Pascal's theorem: a variant.

Corollary 3.7 *Let M, N, P be three points on a non-degenerate conic. Then the intersections of the tangent lines to the conic at M, N, P with the opposite sides of the triangle MNP are collinear. (See Figure 18).*




Figure 18: Simpson's line.

The classical Pappus theorem is a version of Pascal's theorem for degenerate conics.

4 The Siebeck–Marden theorem

Marden's theorem is one of the fundamental results in geometric theory of polynomials and rational functions. That theorem has a long history which is well described in the famous Marden book [Mar1966]. The earliest version of this theorem, up to our best knowledge, goes back to 1864 when Siebeck (see [Sie1864]) formulated and proved it for the case of polynomials with simple roots. Here we consider only the case $n = 3$.

Theorem 4.1 (Siebeck, [Sie1864], $n = 3$) *Let $P(z)$ be a polynomial of degree 3 with complex coefficients, such that its zeros $\alpha_1, \alpha_2, \alpha_3$ are simple and noncollinear. Then there exists an ellipse \mathcal{E} tangent to every line segment $[\alpha_i, \alpha_j]$ at the midpoint, see Figure 19. Moreover, the foci of the curve \mathcal{E} are zeros of the derivative polynomial $P'(z)$.*




Figure 19: Siebeck theorem. If $\alpha_1, \alpha_2, \alpha_3$ are zeros of a cubic polynomial $P(z)$ and β_1, β_2 zeros of its derivative $P'(z)$, then there is an ellipse with foci β_1, β_2 inscribed in triangle with vertices $\alpha_1, \alpha_2, \alpha_3$. Moreover, the ellipse is touching the sides at midpoints.

We note that the ellipse which touches the midpoints of the triangle sides is *the Steiner ellipse*, i.e. the ellipse of the maximal area inscribed in that triangle. It is interesting to note that the ratio of areas of any triangle and its Steiner ellipse equals $3\sqrt{3}/(4\pi)$, see e.g. [GSO2020].

The nontrivial and interesting result of Siebeck still attracts a lot of attention, see for example [Kal2008, Dra2011]. In [Pra2004], this theorem is attributed to van den Berg, and two proofs are presented: the original proof of van der Berg from 1888 [vdB1888] and another one from [Sch1982]. We note that the main points of the proof of Siebeck theorem for $n = 3$ from [Kal2008] are based on Propositions 3.1 and 3.2, which indicates a deep relationship with elliptic billiards. We are going to exploit that connection in next Section 5.

Siebeck's theorem can be extended to the cases with non-simple roots, namely to the functions of the form:

$$P(z) = (z - \alpha_1)^{m_1}(z - \alpha_2)^{m_2}(z - \alpha_3)^{m_3}. \quad (4.1)$$

Each zero of the derivative $P'(z)$ is then either equal to some α_i such that $m_i > 1$ or it is a zero of the logarithmic derivative of P :

$$F(z) = (\log P(z))' = \frac{m_1}{z - \alpha_1} + \frac{m_2}{z - \alpha_2} + \frac{m_3}{z - \alpha_3}. \quad (4.2)$$

Notice that $F(z)$ has two zeros. The following statement then holds, see also Figure 20.

Theorem 4.2 ([Mar1966, Theorem 4.2 for $n = 3$]) *Consider a function $P(z)$ of the form (4.1), such that $\alpha_1, \alpha_2, \alpha_3$ are noncollinear complex numbers, and m_1, m_2, m_3 non-zero real constants. Let β_1, β_2 are the zeros of the logarithmic derivative (4.2) of $P(z)$. Then there is a conic with foci β_1, β_2 touching each line of the segment $[\alpha_i, \alpha_j]$ in a point dividing that segment in the ratio $m_i : m_j$.*




Figure 20: Marden theorem. An example for $m_1 = 1, m_2 = 2, m_3 = \sqrt{3}$.

Now we are going to expand on those results from the geometric theory of polynomials, in order to give complete characterization of ellipses inscribed in triangles.

Lemma 4.3 *Suppose that F_1 is a point inside a triangle. Then there exists a unique ellipse inscribed in the triangle, such that one of its foci is point F_1 .*

Proof. Denote the triangle by ABC . Denote by F_1', F_1'', F_1''' the points symmetric to F_1 with respect to sides BC, AC, AB respectively. Then the second focal point of the ellipse inscribed in $\triangle ABC$ is obtained as the intersection of bisectors of angles $F_1''AF_1'''$ and $F_1'CF_1''$. Namely, if F_2 is a second focal point, then triangle $CF_1''F_2$ is congruent to triangle $CF_1'F_2$. Thus, angle F_2CF_1'' is equal to angle F_2CF_1' . \square

Theorem 4.4 *Let ABC be a given triangle, and \mathcal{E} an ellipse inscribed in it, touching the sides BC , AC , AB in points K , L , M respectively. Then the lines AK , BL , CM are concurrent. (See Figure 21.)*




Figure 21: The lines connecting triangle vertices with the points where the inscribed ellipse touches the opposite sides are concurrent.

Proof. Let φ be an affine transformation that maps the ellipse \mathcal{E} to a circle \mathcal{E}' . That circle is inscribed in triangle $\varphi(ABC)$.

Since segments of each pair $\varphi(A)\varphi(L)$ and $\varphi(A)\varphi(M)$, $\varphi(B)\varphi(K)$ and $\varphi(B)\varphi(M)$, $\varphi(C)\varphi(K)$ and $\varphi(C)\varphi(L)$ are of equal lengths as tangent segments to the circle from the same vertex, the Ceva's theorem (Theorem 2.3) gives that lines $\varphi(AK)$, $\varphi(BL)$, $\varphi(CL)$ are concurrent.

The statement follows immediately. \square

In order to align further this matter with the Marden theorem, we provide an alternative proof of the last theorem.

Second proof of Theorem 4.4. Let F_1 be one of the foci of \mathcal{E} . Denote the complex numbers corresponding to A , B , C , and F_1 by z_1 , z_2 , z_3 , and q respectively. We search for nonzero real numbers m_1 , m_2 , m_3 such that q is one of the zeros of the derivative of $(z - z_1)^{m_1}(z - z_2)^{m_2}(z - z_3)^{m_3}$. This imposes one complex relation on real numbers m_1 , m_2 , m_3 . That complex relation gives two real relations which uniquely determine three real numbers m_1 , m_2 , m_3 up to a nonzero real factor. According to Marden's theorem (Theorem 4.2), there is an ellipse \mathcal{E}_1 inscribed in triangle ABC which has F_1 as one of its foci, with the other focus corresponding to the second zero of the derivative. Since ellipses \mathcal{E} and \mathcal{E}_1 are both inscribed in $\triangle ABC$ and both have F_1 as one of their foci, Lemma 4.3 implies that they coincide. \square

Remark 4.5 *One can compare the last theorem with the Bradley theorem, see e.g. [Bar2015].*

Theorem 4.6 *Every ellipse inscribed in a triangle is a Marden ellipse, i.e. there exist positive real numbers m_1 , m_2 , m_3 such that the foci of the ellipse are the zeros of the logarithmic derivative of the function $(z - \alpha_1)^{m_1}(z - \alpha_2)^{m_2}(z - \alpha_3)^{m_3}$, with α_1 , α_2 , α_3 being the complex numbers corresponding to the vertices of the triangle. The numbers m_1 , m_2 , m_3 are unique up to a non-zero factor.*

Proof. Denote the triangle by ABC , the inscribed ellipse by \mathcal{E} , and by K , L , M the common points of \mathcal{E} with sides BC , AC , AB respectively. According to Theorem 4.4, the lines AK , BL , CM are concurrent, thus Ceva's theorem implies that (2.1) is satisfied. This relation determines positive numbers m_1 , m_2 , m_3 uniquely up to a non-zero factor. \square

5 3-periodic trajectories of billiards within ellipses

Now, after reviewing billiards and Marden's theory in previous sections, we are equipped and ready to address the the title question.

Theorem 5.1 *Every triangle is a 3-periodic trajectory of the billiard within an ellipse. That ellipse is uniquely determined.*

Proof. Let an arbitrary triangle KLM be given and construct a new triangle ABC , formed by the bisectors of the exterior angles of KLM , which implies that KLM is a 3-periodic billiard trajectory within triangle ABC .

According to Theorem 2.2, points K, L, M are the feet of the altitudes of triangle ABC , so the three lines AK, BL, CM intersect at one point – the orthocenter of triangle ABC . Thus, according to Ceva's Theorem (see Theorem 2.3), there exist nonzero real numbers m_1, m_2, m_3 such that point K divides BC in the ratio $m_2 : m_3$, point L divides CA in the ratio $m_3 : m_1$, and M divides AB in the ratio $m_1 : m_2$. The numbers m_1, m_2, m_3 can be calculated in terms of the lengths of the sides of triangle ABC , see Lemma 2.6 and relation (2.2).

Now, according to Marden's theorem (Theorem 4.2), there exists a conic \mathcal{E} which touches the sides AB, BC, CA respectively at M, K, L . Since K, L, M are inner points of those sides, \mathcal{E} must be an ellipse inscribed in the triangle ABC , see Figure 20. By construction, KLM is a billiard trajectory within ellipse \mathcal{E} . \square

We provide also another proof, based on theorems of Pascal and Menelaus.

Second proof of Theorem 5.1. Let KLM be a triangle, lines k, l, m the bisectors of its exterior angles at the vertices K, L, M respectively, and P, Q, R the intersections of those lines with LM, KM, KL respectively. According to Corollary 2.8, points P, Q, R are collinear.

Now, there is a unique conic touching the lines k, l at K, L and containing M . Let m' be the tangent line to that conic at M and denote $R' = m' \cap KL$. According to Corollary 3.7, P, Q, R' are collinear, which gives $R = R'$, i.e. $m = m'$. Since the conic is touching the sides of the triangle determined by k, l, m in inner points K, L, M , it must be an ellipse. \square

6 Convex 4-periodic trajectories of billiards within ellipses

6.1 Is every parallelogram a trajectory of an elliptical billiard?

In this section, in Theorem 6.3, we will show that being a convex 4-periodic elliptical billiard trajectory represents a complete characterization of parallelograms.

We start by formulating the following useful statement.

Lemma 6.1 *Let a 4-periodic convex billiard trajectory within an ellipse be given, see Figure 22. Then we have:*

- *that trajectory is a parallelogram;*
- *the tangent lines to the ellipse at the reflection points form a rectangle;*
- *the parallelogram is also a closed billiard trajectory within the rectangle;*
- *the diagonals of the rectangle are parallel to the sides of the parallelogram.*

Proof. The first statement follows, for example, from [DR2019, Theorem 4], and the rest is obtained by straightforward calculation of the angles and application of the billiard reflection law. \square




Figure 22: A convex 4-periodic trajectory within ellipse and the tangent lines at the points of reflection.

Lemma 6.2 *Any parallelogram is a closed billiard trajectory within a unique rectangle.*

Moreover, for any rectangle and any point on one of its sides, there is a unique 4-periodic billiard trajectory within the rectangle such that the given point is one of its vertices.

Proof. The sides of the rectangle are bisectors of the exterior angles of the parallelogram.

The second part follows from the fact that the segments of any 4-periodic billiard trajectory within a rectangle are parallel to its diagonals, see Figure 22. \square

Theorem 6.3 *Any parallelogram is a closed billiard trajectory within an ellipse. Moreover, such an ellipse is uniquely determined for a given parallelogram.*

We immediately provide a compact, synthetic proof of this theorem, based on the Pascal theorem (see Theorem 3.5). An alternative, analytic proof is given in Section 6.2.

Proof. Let $EFGH$ be a given parallelogram. According to Lemma 6.2, there is a unique rectangle within which $EFGH$ is a closed billiard trajectory, see Figure 23.




Figure 23: A 4-periodic billiard trajectory within a rectangle.

Denote by a the side of the rectangle containing point E and by \mathcal{C} the unique conic circumscribed about $EFGH$ such that a is its tangent line. We want to prove that the conic is touching the remaining sides of the rectangle exactly at the points F, G, H .

Applying Pascal's theorem to \mathcal{C} and its points E, E, F, F, G, H , (see Corollary 3.6) we get that the intersection points of the following pairs of lines are collinear: a and FG ; EF and GH ; b' and

EH , where b' is the tangent line to \mathcal{C} at F . Denote $X = a \cap FG$, $Y = b' \cap EH$. Since EF and GH are parallel, this implies that the line XY also must be parallel to them, thus Y is the intersection point of EH with the line which is parallel to EF and contains X .

Parallelogram $EFXY$ is rhombus, since $EF = FX$, so its diagonals are orthogonal to each other, which implies that $Y \in b$, where b is the side of the rectangle containing point F . Thus $b = b'$.

Similarly, we can prove that the remaining two sides of the rectangle are also tangent to \mathcal{C} .

It remains to show that \mathcal{C} is an ellipse. Each degenerate conic which contains points E, F, G, H is intersecting transversely the sides of the rectangle, thus \mathcal{C} must be non-degenerate. No two tangent lines of a parabola are parallel, thus \mathcal{C} is either hyperbola or ellipse. For a hyperbola, we can notice that a pair of parallel lines cannot touch the same branch, and that the touching points of the lines containing sides of the rectangle would be placed on the extensions of those sides, which does not correspond to the geometric arrangement that we got here.

We conclude that \mathcal{C} is the unique ellipse within which parallelogram $EFGH$ is a closed billiard trajectory. \square

6.2 An analytic proof

To prepare for the proof, we need the following three lemmas.

Lemma 6.4 *Consider an ellipse with foci F_1, F_2 and center O . Let M be a point on that ellipse, t the line touching ellipse at M , F'_1 the point symmetric to F_1 with respect to t , and X the projection of F_1 to t , as shown in Figure 24. Then we have:*

- Lines F'_1M and OX are parallel and $OX = \frac{1}{2}F'_1F_2$.
- Therefore, the length of the segment OX does not depend on the choice of point M on the ellipse, since $d = F_1M + F_2M = F'_1F_2 = 2OX$, where d is the defining “rope length” for the ellipse.
- Moreover, the line containing point M which is orthogonal to t and the line containing point F_1 which is parallel to t intersect each other at a point on OX .




Figure 24: Lemma 6.4: the length of segment OX does not depend on the choice of point M on the ellipse.

Proof. The first two statements are a direct application of the minimizing property of the billiard reflection (see Remark 2.1), together with focal property (Proposition 3.1).

Denote by N the intersection of two lines, as described in the last statement of the lemma. Then MNF_1X is a rectangle, thus $MN = XF_1$. Since X is the midpoint of $F_1F'_1$, we have $MN = XF'_1$. Since segments MN and XF'_1 are, in addition to that, parallel, quadrangle $MNXF'_1$ is a parallelogram. Thus, XN is parallel to the line F'_1F_2 , so $N \in OX$. \square

Lemma 6.5 Suppose that $EFGH$ is a parallelogram which represents a periodic billiard trajectory within rectangle $ABCD$ and ellipse with centre O and foci F_1, F_2 , as shown in Figure 25. Denote the sides of the rectangle by $a = AB = CD$ and $b = BC = AD$, the distances of F_1 from AD, AB by x, y respectively, and set $e = AE$. Then:

$$(a - 2x)^2 - (b - 2y)^2 = a^2 - b^2 \quad \text{and} \quad (a - 2x)(b - 2y) = b(a - 2e). \quad (6.1)$$




Figure 25: A convex 4-periodic trajectory within ellipse and rectangle.

Proof. Denote by X, Y and O_x, O_y the projections of points F_1 and O to AB, AD respectively, see the lefthandside of Figure 26. We notice that $AX = x, AY = y, AO_x = \frac{a}{2}, AO_y = \frac{b}{2}$. Since Lemma 6.4 gives $OX = OY$, the first equality from (6.1) is obtained by applying Pythagora's theorem to triangles OXO_x and OYO_y .

Now, denote by Y_1 the point on OX such that the line Y_1E is orthogonal to AB , see the righthand-side of Figure 26. According to the third statement of Lemma 6.4, we have $F_1Y_1 \parallel AB$. Denote $Y_2 = F_1Y_1 \cap OO_x$. Now, from the similarity of triangles OY_1Y_2 and OXO_x , we get the second relation of (6.1). \square




Figure 26: The proof of Lemma 6.5.

Lemma 6.6 Let $ABCD$ be a rectangle and E a point on the side AB . There is a point F_1 in the rectangle, such that it satisfies the relationships (6.1), where a, b, e, x, y are defined as in Lemma 6.5.

Moreover, point F_1 is unique up to the symmetry with respect to the center of the rectangle.

Proof. From (6.1), we get the following biquadratic relationship for $\xi = a - 2x$:

$$\xi^4 - (a^2 - b^2)\xi^2 - b^2(a - 2e)^2 = 0.$$

The discriminant of the corresponding quadratic equation:

$$\hat{\xi}^2 - (a^2 - b^2)\hat{\xi} - b^2(a - 2e)^2 = 0 \quad (6.2)$$

is:

$$d = (a^2 - b^2)^2 + 4b^2(a - 2e)^2.$$

Notice that always $d \geq 0$.

For $d = 0$ we have $a = b = 2e$, so the rectangle $ABCD$ is a square and we get from (6.1) that F_1 is its center.

For $d > 0$, the equation (6.2) has two real solutions $\hat{\xi}_1, \hat{\xi}_2$, whose product equals $\hat{\xi}_1\hat{\xi}_2 = -b^2(a - 2e)^2 \leq 0$.

If $\hat{\xi}_1\hat{\xi}_2 = 0$, we have $a - 2e = 0$, i.e. E is the midpoint of AB . The second relationship of (6.1) then gives $a - 2x = 0$ or $b - 2y = 0$. From the first relationship of (6.1), we can deduce that either F_1 is the center of the rectangle if $a = b$, or $x = \frac{1}{2}(a \pm \sqrt{a^2 - b^2})$, $y = \frac{b}{2}$ if $a > b$, or $x = \frac{a}{2}$, $y = \frac{1}{2}(b \pm \sqrt{b^2 - a^2})$ if $a < b$.

If $\hat{\xi}_1\hat{\xi}_2 < 0$, then exactly one of the solutions $\hat{\xi}_1, \hat{\xi}_2$ is positive, so suppose $\hat{\xi}_1 > 0$. Then we have that point F_1 satisfies $x = \frac{1}{2}\left(a \pm \sqrt{\hat{\xi}_1}\right)$, $y = \frac{1}{2}\left(b \mp \frac{b(a - 2e)}{\sqrt{\hat{\xi}_1}}\right)$. The choice of signs here determines two points within the rectangle, which are symmetric with respect to its center. \square

Analytic proof of Theorem 6.3. Let $EFGH$ be a given parallelogram, and $ABCD$ the rectangle determined by external bisectors of the parallelogram. We assume that $E \in AB$. According to Lemma 6.6, there is a point F_1 in the rectangle which satisfies the relations (6.1). Denote by F_2 the point symmetric to F_1 with respect to the center O of the rectangle. Applying the reasoning from the proofs of Lemmas 6.4-6.6, the ellipse with foci F_1, F_2 which contains E will be the unique ellipse within which $EFGH$ represents a closed billiard trajectory. \square

Thus, we have concluded the analytic proof of Theorem 6.3. As an outcome, we get the exact formulas for the foci of the ellipse.

7 Nonconvex 4-periodic elliptical billiard trajectories

7.1 Characterization of nonconvex 4-periodic trajectories

Following [BLPT2020, Definition 3.4], we introduce the notion of a Darboux butterfly.

Definition 7.1 *Any closed polygonal self-intersecting line consisting of four segments, such that the opposite pairs are congruent to each other, is called a Darboux butterfly (or a bow tie). See Figure 27.*




Figure 27: A Darboux butterfly.

Lemma 7.2 *Every Darboux butterfly consists of legs and diagonals of an isosceles trapezoid.*

Proof. Let $GHKL$ be a Darboux butterfly. Then $\triangle GHK$ is congruent to $\triangle KLG$ and they are placed on the same side of the “diagonal” GK . Thus, HL is parallel to GK , and GH is congruent to KL . \square

Lemma 7.3 *Every 4-periodic elliptical billiard trajectory, having a hyperbola as a caustic, is a Darboux butterfly. (See Figure 28).*




Figure 28: A 4-periodic elliptic billiard trajectory with a hyperbola as the caustic.

Proof. We know that every 4-periodic elliptical billiard trajectory with hyperbola as a caustic is symmetric with respect to the smaller axis of the boundary ellipse, see, for example, [DR2019, Theorem 4]. Thus its vertices will be the vertices of an isosceles trapezoid, while its segments are the diagonals and legs of that trapezoid. \square

Remark 7.4 *From [Sta2021, Theorem 1], it follows that for each 4-periodic trajectory of an elliptical billiard, having a hyperbola as the caustic, there exists a 4-periodic elliptical billiard trajectory having an ellipse as the caustic such that the corresponding sides of the two trajectories are congruent.*

Next, we show that every Darboux butterfly is a billiard trajectory within a kite.

Lemma 7.5 *Every Darboux butterfly is a billiard trajectory within a unique kite, consisting of two congruent acute triangles. Moreover, the Darboux butterfly is the union of two 3-periodic billiard trajectories within each of those congruent acute triangles.*

Proof. The edges of the kite are the bisectors of the exterior angles of the Darboux butterfly, see Figure 29. One of the diagonals of the kite, represented by a dashed line in the figure, is the axis of




Figure 29: The Darboux butterfly is a billiard trajectory within a kite.

symmetry, and it divides the kite into two congruent triangles.

Consider a shaded quadrangle in Figure 29, which is determined by two sides of the kite and the bisectors of the corresponding interior angles of the butterfly. That quadrangle has right angles at the joint vertices with the butterfly, while the angle at the vertex within the butterfly is obtuse. Thus, the angle of the kite at the vertex which is outside the butterfly must be acute.

Finally, since the kite and the Darboux butterfly are both symmetric with respect to the diagonal of the kite, one can immediately see that the diagonal cuts the Darboux butterfly into two 3-periodic billiard trajectories within the triangles. \square

Lemma 7.6 *For every kite consisting of two congruent acute triangles there is a unique Darboux butterfly which is a 4-periodic billiard trajectory within the kite. This trajectory is obtained as the concatenation 3-periodic billiard trajectories within each of the two acute triangles.*

Proof. Follows from the uniqueness of such triangular billiard trajectories, see Theorem 2.2. \square

Remark 7.7 *In addition to the unique Darboux butterfly from Lemma 7.6, each kite consisting of two congruent acute triangles has an infinite family of 4-periodic billiard trajectories. The segments of such trajectories are parallel to the segments of the Darboux butterfly, see the left-hand side of Figure 30.*

There is also an infinite family of 4-periodic billiard trajectories within a kite consisting of two congruent right triangles. Such trajectories are all isosceles trapezoids, see the right-hand side of Figure 30.




Figure 30: 4-periodic trajectories within kites. The kite on the left consists of two acute triangles, and the one on the right consists of two right triangles.

Now, we are ready to prove that each Darboux butterfly is an elliptic billiard trajectory.

Theorem 7.8 *For each Darboux butterfly, there is a unique ellipse, such that the butterfly is a 4-periodic billiard trajectory within that ellipse.*

Proof. Let $GKHL$ be a given Darboux butterfly. According to Lemma 7.6, there is a kite $ABCD$, within which the butterfly is a periodic billiard trajectory, see Figure 31. Let \mathcal{C} be the unique conic




Figure 31: The Darboux butterfly is a billiard trajectory within a kite.

circumscribed about $GKHL$ and touching the line BC at point G .

Applying Pascal's theorem to \mathcal{C} and its points G, G, H, K, K, L , (see Corollary 3.6) we get that the intersection points of the following pairs of lines are collinear: BC and k ; GH and KL ; HK and GL , where k is the tangent line to \mathcal{C} at K . Since $GH \cap KL$ and $HK \cap GL$ belong to the line AC , we will have that $k \cap BC \in AC$, i.e. $C \in k$, which gives that CD is tangent to \mathcal{C} at K .

This implies that \mathcal{C} is symmetric with respect to the line AC and that the pairs of segments KL, KH and GL, GH satisfy the billiard reflection law off \mathcal{C} at points K, G respectively. From there, there are conics $\mathcal{C}', \mathcal{C}''$ which are confocal with \mathcal{C} , each touching one of those two pairs of segments.

Since \mathcal{C} is symmetric with respect to AC , the same must be true for \mathcal{C}' and \mathcal{C}'' . Moreover, since the pairs of segments KL, KH and GL, GH are also symmetric to each other with respect to AC , that will imply $\mathcal{C}' = \mathcal{C}''$. Therefore, the Darboux butterfly $GHKL$ is inscribed in \mathcal{C} and circumscribed about a conic confocal to \mathcal{C} , so it will represent a billiard trajectory within \mathcal{C} .

Finally, \mathcal{C} must be an ellipse, since it is touching the sides of the kite $ABCD$ in inner points. \square

7.2 Analytic approach

Lemma 7.9 Consider an acute triangle ABC . Let G and H be the footings of its altitudes from A and C respectively. Suppose that \mathcal{E} is an ellipse whose foci are symmetric with respect to AC such that it is touching AB and BC at H and G respectively. Denote by F_1 the focus of \mathcal{E} which is placed within $\triangle ABC$, and by O, X, Y, Z, W the projections of F_1 to AC, AB, BC, AG, CH respectively, as shown in Figure 32.

Then, the points O, Z, Y are collinear as well as the points O, W, X . Moreover, $OX = OY$.




Figure 32: Lemma 7.9.

Proof. A direct application of Lemma 6.4. \square

Applying Theorem 2.9 to triangles ACG and ACH , we get the following

Lemma 7.10 Let ABC be an arbitrary triangle such that its angles at A, C are not right, and G, H the footings of its altitudes from A, C . Then the locus of all points F in the plane such that its projections to AC, AG, BC are collinear is the circle with radius AC . (See Figure 33). That circle is also the locus of all points whose projections to AC, CH, AB are collinear.

Remark 7.11 It is interesting to note that the locus from Lemma 7.10 does not depend on the vertex B .




Figure 33: Lemma 7.10.

Proposition 7.12 Consider an acute triangle ABC . Let G and H be the footings of its altitudes from A and C respectively. Then there is a unique point F_1 inside the triangle such that its projections O, X, Y, Z, W to AC, AB, BC, AG, CH respectively, satisfy the following:

- (i) O, Z, Y are collinear;
- (ii) O, W, X are collinear;
- (iii) $OX = OY$.

Proof. Without loss of generality, we may set a coordinate system in the plane such that $A = (0, 0)$ and $C = (1, 0)$. In that system, denote the coordinates of B and F by (b_1, b_2) and (f_1, f_2) respectively. Applying Lemma 7.10, we see that each of the conditions (i) and (ii) is equivalent to F belonging to the circle with radius AC , which can be written in the coordinates as:

$$-f_1 + f_1^2 + f_2^2 = 0. \quad (7.1)$$

A direct calculation for condition (iii) gives

$$b_2^2 ((1 - 2b_1)(f_1^2 + f_2^2) + 2(b_1^2 + b_2^2)f_1 - (b_1^2 + b_2^2)) = 0. \quad (7.2)$$

Observing that $b_2 \neq 0$ and substituting $f_2^2 = f_1 - f_1^2$ from (7.1) into (7.2), we get

$$f_1 = \frac{b_1^2 + b_2^2}{1 - 2b_1 + 2b_1^2 + 2b_2^2}. \quad (7.3)$$

Thus, $F_1 = (f_1, f_2)$ which satisfies the conditions (i), (ii), and (iii) is uniquely determined through (7.3) and (7.1). This completes the proof. \square

This finishes the analytic proof of Theorem 7.8.

Acknowledgements

The authors would like to thank Djordje Baralić and Miloš Djorić for useful discussions. The research was supported by the Australian Research Council, Discovery Project 190101838 *Billiards within quadrics and beyond*, the Science Fund of Serbia grant *Integrability and Extremal Problems in Mechanics, Geometry and Combinatorics*, MEGIC, Grant No. 7744592 and the Simons Foundation grant no. 854861.

References

- [ADSK2016] A. Avila, J. De Simoi, and V. Kaloshin, *An integrable deformation of an ellipse of small eccentricity is an ellipse*, *Ann. of Math. (2)* **184** (2016), no. 2, 527–558, DOI 10.4007/annals.2016.184.2.5. MR3548532
- [Bar2015] Dj. Baralić, *A short proof of the Bradley theorem*, *American Mathematical Monthly* **122** (2015), 381–385.
- [vdB1888] F. J. van der Berg, *Nogmaals over afgeleide Wortelpunten*, *Nieuw Archief voor Wiskunde* **15** (1888), 100–164.
- [Ber1987] M. Berger, *Geometry. II*, Universitext, Springer-Verlag, Berlin, 1987.
- [BM2017] M. Bialy and A. E. Mironov, *Angular billiard and algebraic Birkhoff conjecture*, *Adv. Math.* **313** (2017), 102–126, DOI 10.1016/j.aim.2017.04.001. MR3649222
- [BM2022] ———, *The Birkhoff-Poritsky conjecture for centrally-symmetric billiard tables*, *Ann. of Math. (2)* **196** (2022), no. 1, 389–413, DOI 10.4007/annals.2022.196.1.2. MR4429262
- [Bol1990] S. V. Bolotin, *Integrable Birkhoff billiards*, *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* **2** (1990), 33–36.
- [BLPT2020] G. Bor, M. Levi, R. Perline, and S. Tabachnikov, *Tire Tracks and Integrable Curve Evolution*, *Int. Math. Res. Not. IMRN* **2020** (2020), no. 9, 2698–2768.
- [CZ2023] P. Corvaja and U. Zannier, *Finiteness theorems on elliptical billiards and a variant of the Dynamical Mordell-Lang Conjecture*, *Proceedings of the London Mathematical Society* **127** (2023), no. 5, 1268–1337.
- [Dra2011] V. Dragović, *Poncelet-Darboux curves, their complete decomposition and Marden theorem*, *Int. Math. Res. Not. IMRN* **15** (2011), 3502–3523.
- [DR2011] V. Dragović and M. Radnović, *Poncelet Porisms and Beyond*, Springer Birkhauser, Basel, 2011.
- [DR2014] ———, *Pseudo-integrable billiards and arithmetic dynamics*, *Journal of Modern Dynamics* **8** (2014), no. 1, 109–132.
- [DR2019] V. Dragović and M. Radnović, *Caustics of Poncelet polygons and classical extremal polynomials*, *Regul. Chaotic Dyn.* **24** (2019), no. 1, 1–35.
- [DGR2022] V. Dragović, S. Gasiorek, and M. Radnović, *Billiard ordered games and books*, *Regul. Chaotic Dyn.* **27** (2022), no. 2, 132–150.
- [FV2023] A. Fomenko and V. Vedyushkina, *Billiards and Intregrable Systems*, *Russian Mathematical Surveys* **78** (2023), no. 5, 881–954.
- [GRK2021] R. Garcia, D. Reznik, and J. Koiller, *New properties of triangular orbits in elliptic billiards*, *Amer. Math. Monthly* **128** (2021), no. 10, 898–910, DOI 10.1080/00029890.2021.1982360. MR4343349
- [GKR2023] R. Garcia, J. Koiller, and D. Reznik, *Loci of 3-periodics in an elliptic billiard: Why so many ellipses?*, *J. Symbolic Comput.* **114** (2023), 336–358, DOI 10.1016/j.jsc.2022.06.001. MR4446194
- [GSO2020] G. Glaeser, H. Stachel, and B. Odehnal, *The universe of quadrics*, Springer, Berlin, 2020.
- [Glu2021] A. Glutsyuk, *On polynomially integrable Birkhoff billiards on surfaces of constant curvature*, *J. Eur. Math. Soc. (JEMS)* **23** (2021), no. 3, 995–1049, DOI 10.4171/jems/1027. MR4210728
- [Kal2008] D. Kalman, *An Elementary Proof of Marden’s Theorem*, *The American Mathematical Monthly* **115** (2008), 330–337.
- [KS2018] V. Kaloshin and A. Sorrentino, *On the local Birkhoff conjecture for convex billiards*, *Ann. of Math. (2)* **188** (2018), no. 1, 315–380, DOI 10.4007/annals.2018.188.1.6. MR3815464
- [KT1991] V. Kozlov and D. Treshchëv, *Billiards*, Amer. Math. Soc., Providence RI, 1991.
- [Koz2003] V. V. Kozlov, *Rationality conditions for the ratio of elliptic integrals and the great Poncelet theorem*, *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* **71** (2003), no. 4, 6–13 (Russian); English transl., *Moscow Univ. Math. Bull.* **58** (2004), no. 4, 1–7.
- [Mar1966] M. Marden, *Geometry of Polynomials*, 2nd ed., *Math. Surveys*, vol. 2, AMS, 1966.
- [Pra2004] V. V. Prasolov, *Polynomials, Algorithms and Computation in Mathematics*, vol. 11, Springer-Verlag, Berlin, 2004. Translated from the 2001 Russian second edition by Dimitry Leites.
- [Sch1982] I. J. Schoenberg, *Mathematical time exposures*, Mathematical Association of America, Washington, DC, 1982.
- [Sch2006] R. E. Schwartz, *Obtuse triangular billiards. I. Near the (2,3,6) triangle*, *Experiment. Math.* **15** (2006), no. 2, 161–182.
- [Sch2009] ———, *Obtuse triangular billiards. II. One hundred degrees worth of periodic trajectories*, *Experiment. Math.* **18** (2009), no. 2, 137–171.
- [Sie1864] J. Siebeck, *Ueber eine neue analytische Behandlungsweise der Brennpunkte*, *J. Reine Angew. Math.* **64** (1864), 175–182.
- [Sta2021] H. Stachel, *Isometric Billiards in Ellipses and Focal Billiards in Ellipsoids*, *Journal for Geometry and Graphics* **25** (2021), no. 1, 97–118.
- [Tab2005] S. Tabachnikov, *Geometry and billiards*, *Student Mathematical Library*, vol. 30, American Mathematical Society, Providence, RI, 2005.