

$\Gamma(X)$ as Metric Space

E. Ramírez

Escuela de Matemáticas

Universidad Industrial de Santander

Cl. 14 # 6 - 07, Socorro, Santander

e-mail: edwar5119@gmail.com

2024 AMS subject to classification:

Keywords: Pseudométric, topological inverse semigroup, compactly generated space or k -space, Fell Topology and polish space.

Introduction

Functions with open domains appear in a variety of contexts in mathematics. Motivated by this phenomenon, M. Allah and R. Brown set out to endow the space of continuous functions with open domains with an appropriate topology. In chronological order, in [1], they define the topology τ_{co} , which turns the inverse semigroup of partial homeomorphisms between open sets into a topological inverse semigroup. More recently, in [6], they added open sets after realizing that this topology is not T^1 , showing that this addition makes $\Gamma(X)$ a Hausdorff topological inverse semigroup. Then, in [4], they show that if X is a compact, Hausdorff, and totally disconnected space, then $(\Gamma(X), \tau_{hco})$ is completely metrizable. In what follows, we will see how to define such a topology, we will see that the space of continuous functions with open domains $(C_{od}(X, Y), \beta)$ is a Polish space when X is locally compact, second countable space and Y is a complete space. In particular, we will find a metric for $(\Gamma(X), \tau_{hco})$.

1 The inverse semigroup $(\Gamma(X), \tau_{hco})$

Our object of study in this article is the inverse semigroup under composition $\Gamma(X)$ whose elements are all partial homeomorphisms, i.e., homeomorphisms between open sets of a topological space X , which is initially assumed to be locally compact and Hausdorff.

In [1], they define the topology τ_{co} whose subbasic open sets are of the form

$$\langle K, V \rangle = \{f \in \Gamma(X) : K \subseteq D_f \text{ y } f(K) \subseteq V\}$$

where K is compact and V is open in X .¹

Note that if $f \in \langle K, V \rangle$, any restriction of f belongs to $\langle K, V \rangle$. This implies that in general, this topology is not T_1 . Additionally, in [1], they define the weak topology on $\Gamma(X)$, denoted τ_{ico} , given by the functions:

$$\iota : \Gamma(X) \longrightarrow (\Gamma(X), \tau_{co}) \text{ the inclusion map.}$$

$$\eta : \Gamma(X) \longrightarrow (\Gamma(X), \tau_{co}) \text{ the inversion map.}$$

In [1], they show that $(\Gamma(X), \tau_{ico})$ is a topological inverse semigroup. Recently, in [6], they proved that this topology turns out to be the smallest one that makes $\Gamma(X)$ a topological inverse semigroup. Additionally, by utilizing the Fell topology [Definition 1] on the hyperspace $CL(X)$ of closed sets including the empty set, they define the topology τ_{hco} by essentially adding to the weak topology of τ_{ico} the functions

$$D : \Gamma(X) \longrightarrow (CL(X), \tau_{Fell}); f \longmapsto D(f) = X \setminus D_f$$

$$I : \Gamma(X) \longrightarrow (CL(X), \tau_{Fell}); f \longmapsto I(f) = X \setminus I_f$$

This topology turns out to be the smallest Hausdorff topology that makes $\Gamma(X)$ a topological inverse semigroup [6].

1.1 $\Gamma(X)$ as metrizable space

Definition 1. $CL(X)$ is the family of closed sets of X including the empty set. This space has as subbasic open sets those of the form $X \setminus K^+ = A \in CL(X) : A \subseteq X/K$ and $V^- = A \in CL(X) : A \cap V \neq \emptyset$ where K is compact and V is open in X . This topology is called the Fell topology.

Observation 2. If A and B are closed sets in X with $A \subseteq B$, then A is closed in B . Indeed, $A \subseteq cl_B(A)$, and if $x \in cl_B(A)$, there exists a net (x_λ) in A such that $x_\lambda \longrightarrow x$ in B . Let U be an open neighborhood of x , $x \in U \cap B$, then there exists λ_0 such that $x_\lambda \in U \cap B$ for all $\lambda > \lambda_0$, i.e., $x_\lambda \longrightarrow x$ in X . As A is closed, $x \in A$, i.e., $cl_B(A) = A$.

Proposition 3. If X is locally compact, Hausdorff and second countable, then there exists a countable base in X of *relatively compact* open sets, i.e., with compact closure.

¹This topology can be defined on the larger set of continuous functions whose domains are open subset of X , called $C_{od}(X)$ as in [6], or $P_0(X)$ as in [1].

Proof. Let \mathcal{C} be a countable base of X , V be open, and $x \in V$. By local compactness, there exists an open set U with \overline{U} compact such that $x \in U \subseteq \overline{U} \subseteq V$. So there exists $B \in \mathcal{C}$ such that $x \in B \subseteq U$. Thus, $\overline{B} \subseteq \overline{U}$, and using the previous observation, \overline{B} is closed in the compact set \overline{U} , hence \overline{B} is compact in \overline{U} and therefore in X . Consider the set

$$\mathcal{B} = \{B \in \mathcal{C} : \overline{B} \text{ is compact}\}$$

It suffices to see that \mathcal{B} is a basis.

- (i) If $x \in X$, by the previous argument (taking $V = X$), there exists $B \in \mathcal{C}$ such that \overline{B} is compact and $x \in B \subseteq \overline{B} \subseteq X$. Therefore, $x \in \bigcup_{B \in \mathcal{B}} B$.
- (ii) Let $B_1, B_2 \in \mathcal{B}$ and $x \in B_1 \cap B_2$. Taking $V = B_1 \cap B_2$, by the previous argument, there exists $B_3 \in \mathcal{B}$ such that $x \in B_3 \subseteq \overline{B_3} \subseteq B_1 \cap B_2$.

□


Lemma 4. If X is locally compact, Hausdorff, and second countable, then $(\Gamma(X), \tau_{co})$ is second countable.

Proof. Fix a relatively compact base \mathcal{B} , closed under finite unions, and given $f \in \langle K, W \rangle$. For each $x \in K$, there exists $V_x \in \mathcal{B}$ such that $f(x) \in V_x \subseteq \overline{V_x} \subseteq W \cap \text{im}(f)$. Due to the compactness of $f(K)$, there exist $x_1, \dots, x_n \in K$ such that

$$f(K) \subseteq \bigcup_{i=1}^n V_{x_i} = V_m \in \mathcal{B}$$

since \mathcal{B} is closed under finite unions.

$$V_m \subseteq \overline{V_m} = \bigcup_{i=1}^m \overline{V_{x_i}} \subseteq W \cap \text{im}(f)$$



In an analogous manner, for $K \subseteq U_m$, there exists $U_n \in \mathcal{B}$ such that $K \subseteq U_n \subseteq \overline{U_n} \subseteq U_m$. So f belongs to $\langle \overline{U_n}, V_m \rangle$.

If $g \in \langle \overline{U_n}, V_m \rangle$, $K \subseteq \overline{U_n} \subseteq D_g$ and $g(K) \subseteq g(\overline{U_n}) \subseteq V_m \subseteq W$. So $\langle \overline{U_n}, V_m \rangle \subseteq \langle K, W \rangle$ and therefore

$$\mathfrak{B} = \{\langle \overline{U}, V \rangle : U, V \in \mathcal{B}\}$$

is a countable subbase of $(\Gamma(X), \tau_{co})$.

□

Observation 5. In the previous proof only the continuity of f was used, so $C_{od}(X, Y)$ is second countable when X and Y are locally compact and second countable.

Lemma 6. If X is locally compact Hausdorff, then $(\Gamma(X), \tau_{hco})$ is regular.

Proof. Fix a relatively compact base \mathcal{B} closed under finite unions. Let $f \in \langle K, W \rangle$ with $K \neq \emptyset$ (since if $K = \emptyset$, then $\langle \emptyset, W \rangle = \Gamma(X)$). If $x \in K$, there exists $V_x \in \mathcal{B}$ such that $f(x) \in V_x \subseteq \overline{V_x} \subseteq W \cap \text{im}(f)$. Due to the compactness of $f(K)$, there exist $x_1, \dots, x_n \in K$ such that

$$f(K) \subseteq \bigcup_{i=1}^n V_{x_i} = V \subseteq \overline{V} = \bigcup_{i=1}^n \overline{V_{x_i}} \subseteq W \cap \text{im}(f), \quad V \in \mathcal{B}$$

Similarly, there exists $U \in \mathcal{B}$ such that $K \subseteq U \subseteq \overline{U} \subseteq f^{-1}(\overline{V}) \subseteq \text{dom}(f)$. This is


$$f \in \langle K, V \rangle \cap D^{-1}((X \setminus \overline{U})^+) = M$$

Let $g \in \overline{M}$ ($\overline{M} = cl_{\tau_{hco}}(M)$), there exists a red (g_λ) in M such that $g_\lambda \xrightarrow{\tau_{hco}} g$. Let's see that $K \subseteq D_g$. If we assume that $K \not\subseteq D_g$, then $X \setminus D_g \cap K \neq \emptyset$, that is, $X \setminus D_g \cap U \neq \emptyset$, ($K \subseteq U$) and therefore $D(g) = X \setminus D_g \in U^-$. Since $D(g_\lambda) \xrightarrow{\tau_{Fell}} D(g)$, there exists λ_0 such that $D(g_{\lambda_0}) = X \setminus D_{g_{\lambda_0}} \in U^-$, ie.

$$X \setminus D_{g_{\lambda_0}} \cap U \neq \emptyset$$

Which is a contradiction since $g_{\lambda_0} \in D^{-1}((X \setminus \overline{U})^+)$. So $K \subseteq D_g$.

If we now assume that $g \notin \langle K, \overline{V} \rangle$, how $K \subseteq D_g$, then $g(K) \not\subseteq \overline{V}$.



There exists $a \in K$ such that $g(a) \in I_g \setminus \overline{V}$, that is, $g \in \langle \{a\}, I_g \setminus \overline{V} \rangle$, then there exists λ_0 such that $g_{\lambda_0} \in \langle \{a\}, I_g \setminus \overline{V} \rangle$, thus $g_{\lambda_0}(a) \notin \overline{V}$ But this is a contradiction because $g_{\lambda_0} \in \langle K, V \rangle$. So $g \in \langle K, \overline{V} \rangle$ and it follows that $f \in M \subseteq \overline{M} \subseteq \langle K, \overline{V} \rangle \subseteq \langle K, W \rangle$.²

If $f \in \langle K, W \rangle^{-1}$, $f^{-1} \in \langle K, W \rangle$ And from the above, there exists M τ_{hco} -open in $\Gamma(X)$ such that $f^{-1} \in M \subseteq \overline{M} \subseteq \langle K, W \rangle$. Then $f = (f^{-1})^{-1} \in M^{-1}$ that is τ_{hco} -open. Since inversion is $\tau_{hco} - \tau_{hco}$ homeomorphism, it follows that

$$(\overline{M})^{-1} = \overline{M^{-1}}$$

²So far, only the continuity of f has been used, that is, $C_{od}(X, Y)$ is regular endowed with the weak topology given by the functions ι and D , called $\tau_{\iota, D}$.

and therefore $f \in M^{-1} \subseteq \overline{M^{-1}} \subseteq \langle K, W \rangle^{-1}$. Let $f \in D^{-1}(M)$, with M τ_{Fell} -open of $CL(X)$. As X is locally compact Hausdorff space, by [2, 5.1.4 Corollary] $(CL(X), \tau_{Fell})$ is compact Hausdorff, so it is regular, and therefore there exists an open set N in $CL(X)$ such that $D(f) \in N \subseteq \overline{N}^{Fell} \subseteq M$, then

$$f \in D^{-1}(N) \subseteq \overline{D^{-1}(N)}^{hco} \subseteq D^{-1}(\overline{N}^{Fell}) \subseteq D^{-1}(M)$$

□

Theorem 7. If X is locally compact Hausdorff and second countable, then $(\Gamma(X), \tau_{hco})$ is metrizable.

Proof. For [2, 5.1.5 Theorem] $(CL(X), \tau_{Fell})$ is compact and metrizable, hence it is second countable. Let's see that $(\Gamma(X), \tau_{hco})$ is second countable.

It suffices to consider τ_{hco} as the weak topology generated by ι , η , and D (since $I = D\eta$). Let U be a τ_{hco} -open set in $\Gamma(X)$ with $f \in U$.

If $f = \emptyset$, There exist V_1, \dots, V_n non-empty open sets of X such that (the basic neighborhoods of the empty function are of this form)

$$\emptyset \in D^{-1}(V_1^-) \cap \dots \cap D^{-1}(V_n^-) \subseteq U$$

Consider a countable base \mathcal{B} , relatively compact, and closed under finite unions, and we can assume that such V_i 's are elements of \mathcal{B} .

If $f \neq \emptyset$, using \mathcal{B} and [2, 5.1.5 Theorem] taking the base of $(CL(X), \tau_{Fell})$ of your proof

$$\begin{aligned} f \in & \langle \overline{R_1}, S_1 \rangle \quad \cap \dots \cap \langle \overline{R_m}, S_m \rangle \quad \cap \\ & \langle \overline{T_1}, U_1 \rangle^{-1} \quad \cap \dots \cap \langle \overline{T_m}, U_m \rangle^{-1} \quad \cap \\ & D^{-1}(W_1^-) \quad \cap \dots \cap D^{-1}(W_r^-) \quad \cap \\ & D^{-1}(X \setminus \overline{V_1}^+) \quad \cap \dots \cap D^{-1}(X \setminus \overline{V_t}^+) \subseteq U. \end{aligned}$$

So we can consider the **countable subbase**

$$\mathfrak{B} = \{ \langle \overline{U}, V \rangle, \langle \overline{U}, V \rangle^{-1}, D^{-1}(V^-), D^{-1}(X \setminus \overline{V}^+) : U, V \in \mathcal{B} \}$$

For [6] $(\Gamma(X), \tau_{hco})$ is Hausdorff (and therefore T^1), for the Lemma 6 and using Urysohn's metrization theorem, we conclude that $(\Gamma(X), \tau_{hco})$ is metrizable. □

Theorem 8. If X and Y are locally compact Hausdorff and second countable, then $(C_{od}(X, Y), \tau_{l,D})$ is metrizable.

Proof. Lets $f, g \in C_{od}(X, Y)$ are diferents.

Case 1. $f = \emptyset$, then $D_g \neq \emptyset$, let $x \in D_g$.

$$\begin{aligned} g \in D^{-1}(X \setminus \{x\}^+) \quad & \text{and} \quad \emptyset \notin (X \setminus \{x\}^+) \\ \emptyset \in D^{-1}(D_g^-) \quad & \text{and} \quad f \notin D^{-1}(D_g^-) \end{aligned}$$

$g = \emptyset$ is analogous.

Case 2. $f, g \neq \emptyset$

2.1. $D_f \neq D_g$:

2.1.1. $D_f \setminus D_g \neq \emptyset$, let $x \in D_f \setminus D_g$, then

$$\begin{aligned} f \in D^{-1}(X \setminus \{x\}^+) \quad \text{and} \quad g \notin D^{-1}(X \setminus \{x\}^+) \\ g \in D^{-1}(D_f^-) \quad \text{and} \quad f \notin D^{-1}(D_f^-) \end{aligned}$$

2.1.2. $D_g \setminus D_f \neq \emptyset$ is analogous.

2.2. $D_f = D_g$, then there exists $x \in D_f$ such that $f(x) \neq g(x)$. In $I_f \cap I_g$, there exist U and V disjoint open sets of Y contained in $I_f \cap I_g$ such that $f(x) \in U$ and $g(x) \in V$.

$$\begin{aligned} f \in \langle \{x\}, U \rangle \quad \text{and} \quad g \notin \langle \{x\}, U \rangle \\ g \in \langle \{x\}, V \rangle \quad \text{and} \quad f \notin \langle \{x\}, V \rangle \end{aligned}$$

So $(C_{od}(X, Y), \tau_{\iota, D})$ is \mathbf{T}^1 and by Urysohn's metrization theorem, the result follows. \square

1.2 Convergence in $\Gamma(X)$

In what follows, $\mathbb{K}(X)$ will denote the hyperspace of non-empty compact subsets of a topological space X . Let X be a metric space, f and g are partial non-zero functions, and let $K \in \mathbb{K}(D_f \cap D_g)$. Define

$$d_K(f, g) := \sup_{x \in K} d(f(x), g(x))$$

Definition 9. Let $f \in \Gamma(X) \setminus \{\emptyset\} := \Gamma^*(X)$, K non-empty compact subset of D_f and $\epsilon > 0$, we define

$$B_K(f, \epsilon) := \{g \in \Gamma(X) : K \subseteq D_g \text{ y } d_K(f, g) < \epsilon\}$$

Lemma 10. $\beta = \{B_K(f, \epsilon) : f \in \Gamma(X), K \in \mathbb{K}(D_f) \text{ y } \epsilon > 0\} \cup \{\Gamma(X)\}$ is a basis for a topology.

Proof. (i) $\bigcup_{B \in \beta} B = \Gamma(X)$.

(ii) Let $f \in B_K(g, \epsilon_1) \cap B_L(h, \epsilon_2)$ ($g, h \neq \emptyset$, $K \subseteq D_f$ and $L \subseteq D_g$). Consider $\delta_1 = \epsilon_1 - d_K(f, g)$ and $\delta_2 = \epsilon_2 - d_L(f, h)$

Statement 1. $B_K(f, \delta_1) \subseteq B_K(g, \epsilon_1)$. Indeed, if $p \in B_K(f, \delta_1)$, $K \subseteq D_p$ and $d_K(f, p) < \delta_1 = \epsilon_1 - d_K(f, g)$ ie. $d_K(p, g) \leq d_K(f, p) + d_K(f, g) < \epsilon_1$, then $p \in B_K(g, \epsilon_1)$.

Statement 2. $B_L(f, \delta_2) \subseteq B_L(h, \epsilon_2)$ Analogous to the previous proof.

Take $\delta = \delta_1 \wedge \delta_2$. $f \in B_{K \cup L}(f, \delta)$ and if $p \in B_{K \cup L}(f, \delta)$, then $d_K(f, p) < \delta_1$ and $d_L(f, p) < \delta_2$ ie. $p \in B_K(f, \delta_1) \cap B_L(f, \delta_2) \subseteq B_K(g, \epsilon_1) \cap B_L(h, \epsilon_2)$. \square

Let τ_{cc} be the topology generated by β , and we'll call it ***the topology of compact convergence.***

Proposition 11. If X is locally compact metric space, then τ_{co} and τ_{cc} coincide on $\Gamma(X)$.

Proof. $\tau_{co} \subseteq \tau_{cc}$. Indeed, let $f \in \langle K, V \rangle$, ie. $K \subseteq D_f$ y $f(K) \subseteq V$. Then there exists $\epsilon > 0$ such that the cloud $N(\epsilon, f(K)) \subseteq V$. If $g \in B_K(f, \epsilon)$, then $K \subseteq D_g$ and $d_K(f, g) < \epsilon$, that is, $d(f(x), g(x)) < \epsilon, \forall x \in K$. If $x \in K$, $d(g(x), f(K)) \leq d(f(x), g(x)) < \epsilon$. So $g(x) \in N(\epsilon, f(K))$ ie. $g(K) \subseteq N(\epsilon, f(K)) \subseteq V$ and therefore

$$B_K(f, \epsilon) \subseteq \langle K, V \rangle$$

$\tau_{cc} \subseteq \tau_{co}$. Indeed, let $f \in \Gamma^*(X)$, K is non-empty compact subset of D_f , and $\epsilon > 0$. If $x \in K$, there exists V_x open neighborhood of $f(x)$ with diameter $< \epsilon$. By continuity of f and local compactness of X , there exists an open neighborhood of x and relatively compact U_x with $\overline{U_x} \subseteq D_f$ such that $f(\overline{U_x}) \subseteq V_x$, ie. $f \in \langle \overline{U_x}, V_x \rangle, \forall x \in K$.

By compactness of K , there exist $x_1, \dots, x_n \in K$ such that $K \subseteq \bigcup_{i=1}^n U_{x_i}$ ($K \cap U_{x_i} \neq \emptyset$).

Let $K_i = \overline{U_{x_i}} \cap K$, since X is compactly generated, K_i is closed in K and therefore compact. Finally, let's see that

$$W = \bigcap_{i=1}^n \langle K_i, V_{x_i} \rangle \subseteq B_K(f, \epsilon)$$

Let $g \in W$ y $x \in K$. There exists $i \in \{1, \dots, n\}$ such that $x \in U_{x_i}$, then $x \in K_i$ and as $g \in \langle K_i, V_{x_i} \rangle$, $g(x) \in V_{x_i}$. Since $f \in \langle K_i, V_{x_i} \rangle$ and $x \in K_i$, $f(x) \in V_{x_i}$ and as $diam(V_{x_i}) < \epsilon$, $d(f(x), g(x)) < \epsilon$. Therefore $g \in B_K(f, \epsilon)$. \square

Note. The previous argument holds in $(C_{od}(X, Y), \tau_{co})$ with X locally compact Hausdorff space and Y metric space, *mutatis mutandis* in the definition of the B_K 's. This is *analogous* to what happens when we define the compact-open topology on Y^X and restrict to $C(X, Y)$ where Y is a metric space [3, Theorem 46.8].

Corollary 12. Let X be a locally compact Hausdorff space and Y a metric space. If (f_n) is a sequence in $C_{od}(X, Y)$ τ_{cc} -converging to a function f , then f is continuous.

Proof. If $f \neq \emptyset$, let $x \in D_f$. By local compactness, there exists an open set U such that $x \in U \subseteq \overline{U} \subseteq D_f$ and \overline{U} is compact. Since $f_n \xrightarrow{\tau_{cc}} f$, we have that $\overline{U} \subseteq D_{f_n}$ eventually, and $f_n|_{\overline{U}} \longrightarrow f|_{\overline{U}}$ uniformly. Since \overline{U} is compact, then f is continuous at \overline{U} and therefore is continuous at U . Let V open neighborhood of $f(x)$, we can assume that V is contained in the open I_f . Since $f|_U$ is continuous, there exists W an open neighborhood of x contained in U such that $f(W) \subseteq V$. Since U is open in X , W is open in X , so we have that f is continuous at x . \square

Exercise 13. If (f_n) it is a sequence of holomorphic functions defined on an open domain τ_{cc} -converging to $f \in C_{od}(\mathbb{C})$, then f is holomorphic. This implies that the set of holomorphic functions defined on open sets of \mathbb{C} is τ_{cc} -closed in $C_{od}(\mathbb{C})$.

Note that if $K \subseteq D_f$ is a non-empty compact set, we have that

$$\bigcup_{\epsilon > 0} B_K(f, \epsilon) = \{g \in \Gamma(X) : K \subseteq D_g\} = D^{-1}(X \setminus K^+)$$

This means that the open sets added to τ_{ico} in [6] to generate the topology τ_{hco} are only subsets of the form $D^{-1}(V^-)$, that is, neighborhoods of the function \emptyset . In particular, we have that

$$\bigcap_{U \in \tau} D^{-1}(U^-) = \{\emptyset\}$$

Intuitively, this indicates that if U is an open set in X , two functions are close to \emptyset (for U) if $f, g \in D^{-1}(U^-)$.

1.3 $\Gamma(X)$ as metric space

In what follows, X will be a second countable and locally compact Hausdorff space, and (Y, d) a complete metric space with $d \leq 1$. Let's see that it is possible to define an explicit metric on $C_{od}(X, Y)$. Let $\mathcal{B} = \{U_n\}_{n \in \mathbb{N}}$ a basis of X ($U_0 = \emptyset$), and for each $n \in \mathbb{N}$ lets K_{mn} non-empty compact sets such that

$$K_{mn} \subseteq \text{int}(K_{(m+1)n}) \text{ and } \bigcup_{m \in \mathbb{N}} K_{mn} = U_n, \forall m, n \in \mathbb{N}$$

Define

$$\beta_{mn}(f, g) = \begin{cases} 0, & f, g \in D^{-1}(\text{int}(K_{(m+1)n})^-); \\ d_{K_{mn}}(f, g), & f, g \notin D^{-1}(\text{int}(K_{(m+1)n})^-); \\ 1, & \text{other cases.} \end{cases}$$

β_{mn} is a pseudometric on $C_{od}(X, Y)$. Indeed, given $f, g, h \in C_{od}(X, Y)$, Let's call $L = D^{-1}(\text{int}(K_{(m+1)n})^-)$, we have that

$$\beta_{mn}(f, g) \begin{cases} =_{f, g \in L} 0 \leq \beta_{mn}(f, h) + \beta_{mn}(h, g) \\ =_{f, g \notin L} d_{K_{mn}}(f, g) \begin{cases} \leq_{h \in L} 1 + 1 = \beta_{mn}(f, h) + \beta_{mn}(h, g) \\ \leq_{h \notin L} d_{K_{mn}}(f, h) + d_{K_{mn}}(h, g) = \beta_{mn}(f, h) + \beta_{mn}(h, g) \end{cases} \\ =_{f \in L \text{ and } g \notin L} 1 \begin{cases} =_{h \in L} 0 + 1 = \beta_{mn}(f, h) + \beta_{mn}(h, g) \\ \leq_{h \notin L} 1 + \beta_{mn}(h, g) = \beta_{mn}(f, h) + \beta_{mn}(h, g) \end{cases} \\ =_{f \notin L \text{ and } g \in L} 1 \leq \beta_{mn}(f, h) + \beta_{mn}(h, g) \text{ Análogy.} \end{cases}$$

Let $\beta_n(f, g) = \sum_{m=1}^{\infty} 2^{-m} \beta_{mn}(f, g)$, and finally

$$\beta(f, g) = \sum_{n=1}^{\infty} 2^{-n} \beta_n(f, g)$$

β is a metric on $C_{od}(X, Y)$. Indeed, if $f, g \in C_{od}(X, Y)$ are such that $\beta(f, g) = 0 \Leftrightarrow \beta_{mn}(f, g) = 0, \forall m, n \in \mathbb{N}$. If $f = \emptyset$, since $\emptyset \in D^{-1}(int(K_{mn})^-), \forall m, n \in \mathbb{N}$, by definition of β_{mn} necessarily

$$g \in \bigcap_{n \in \mathbb{N}} \bigcap_{m \in \mathbb{N}} D^{-1}(int(K_{mn})^-) \subseteq \bigcap_{n \in \mathbb{N}} D^{-1}(U_n^-) = \{\emptyset\}$$

therefore, $g = \emptyset$. If $g = \emptyset$, analogously $f = \emptyset$.

If $f, g \neq \emptyset$, let $x \in D_f$. There exist $m, n \in \mathbb{N}$ such that $x \in K_{mn} \subseteq U_n \subseteq D_f$, since $f \notin D_{-1}(int(K_{mn})^-)$ and $\beta_{mn}(f, g) = 0$, by definition of β_{mn} , necessarily $g \notin D_{-1}(int(K_{mn})^-)$ and in particular,


$$d(f(x), g(x)) \leq d_{K_{mn}}(f, g) = \beta_{mn}(f, g) = 0 \text{ ie } f(x) = g(x)$$

Analogously, if $x \in D_g, x \in D_f$ y $f(x) = g(x)$. Therefore $f = g$.

Theorem 14. β is a metric for $(C_{od}(X, Y), \tau_{l,D})$.

Proof. If $f_\lambda \xrightarrow{\tau_{l,D}} f$, given $\epsilon > 0$ and let $N \in \mathbb{N}$ such that $2^{-N} < \frac{\epsilon}{2}$. If $n \leq N$, We have two cases:

Case 1. $f \in D^{-1}(U_n^-)$



There exists $M \in \mathbb{N}$ such that $f \in D^{-1}(int(K_{(M+1)n})^-)$ and $f \notin D^{-1}(int(K_{Mn})^-)$.

On one hand, there exists σ such that $f_\lambda \in D^{-1}(int(K_{(M+1)n})^-), \forall \lambda > \sigma$. Given that $D^{-1}(int(K_{(M+1)n})^-) \subseteq D^{-1}(int(K_{mn})^-), \forall m \geq M + 1$, it follows that $\beta_{mn}(f_\lambda, f) = 0, \forall \lambda > \sigma, \forall m \geq M$. So

$$\beta_n(f_\lambda, f) = \sum_{m=1}^{M-1} 2^{-m} \beta_{mn}(f_\lambda, f), \forall \lambda > \sigma.$$

On the other hand, $K_{1n} \subseteq K_{2n} \subseteq \dots \subseteq K_{(M-1)n} \subseteq int(K_{Mn}) \subseteq D_f$, and since $f_\lambda \xrightarrow{\tau_{cc}} f$, for each $m \in \{1, \dots, M-1\}$, there exists λ_m such that

$$f \in B_{K_{mn}} \left(f, \frac{\epsilon 2^{m+n-1}}{MN} \right), \forall \lambda > \lambda_m, \forall m \in \{1, \dots, M-1\}$$

Let $\gamma_n > \lambda_1, \dots, \lambda_{M-1}, \sigma$. Thus, we have that for each $\gamma > \gamma_n$

$$\begin{aligned}\beta_n(f_\lambda, f) &= \sum_{m=1}^{M-1} 2^m \beta_{mn}(f_\lambda, f) \\ &= \sum_{m=1}^{M-1} 2^m d_{K_{mn}}(f_\lambda, f) \\ &\leq \sum_{m=1}^{M-1} 2^m \frac{\epsilon 2^{m+n-1}}{MN} \\ &= \epsilon \frac{M-1}{MN} 2^{n-1} < \frac{\epsilon 2^{n-1}}{N}\end{aligned}$$

Case 2. $f \notin D^{-1}(U_n^-)$, i.e. $U_n \subseteq D_f$. Fix $M \in \mathbb{N}$ such that $2^{-M} < \frac{\epsilon 2^{n-2}}{N}$. For each $m \in \{1, \dots, M\}$, there exists λ_m such that $f_\lambda \in B_{K_{mn}}\left(f, \frac{\epsilon 2^{m+n-2}}{MN}\right)$, $\forall \lambda > \lambda_m$. Let $\gamma_n > \lambda_1, \dots, \lambda_M$. Thus, we have that for each $\lambda > \lambda_1, \dots, \lambda_M$

$$\begin{aligned}\beta_n(f_\lambda, f) &= \sum_{m=1}^M 2^{-m} \beta_{mn}(f_\lambda, f) + \sum_{m=M+1}^{\infty} 2^{-m} \beta_{mn}(f_\lambda, f) \\ &\leq \sum_{m=1}^M 2^{-m} \beta_{mn}(f_\lambda, f) + 2^{-M} \\ &\leq \sum_{m=1}^M 2^{-m} \frac{\epsilon 2^{m+n-2}}{MN} + \frac{\epsilon 2^{n-2}}{N} \\ &= \frac{\epsilon 2^{n-2}}{N} + \frac{\epsilon 2^{n-2}}{N} = \frac{\epsilon 2^{n-1}}{N}\end{aligned}$$

Let $\gamma > \gamma_1, \dots, \gamma_n$. Thus, we have that for each $\lambda > \gamma$

$$\begin{aligned}\beta(f_\lambda, f) &= \sum_{n=1}^{\infty} 2^{-n} \beta_n(f_\lambda, f) \\ &= \sum_{n=1}^N 2^{-n} \beta_n(f_\lambda, f) + \sum_{n=N+1}^{\infty} 2^{-n} \beta_n(f_\lambda, f) \\ &\leq \sum_{n=1}^N 2^{-n} \frac{\epsilon 2^{n-1}}{N} + 2^{-N} \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon\end{aligned}$$

that is, $f_\lambda \xrightarrow{\beta} f$. Conversely, if $f_\lambda \xrightarrow{\beta} f$ and we add that \mathcal{B} is closed under finite unions, then $f_\lambda \xrightarrow{\tau_{\epsilon, D}} f$. indeed, lets $K \in \mathbb{K}(D_f)$ and $\epsilon > 0$. For each $x \in K$, there $U_x \in \mathcal{B}$ such that $x \in U_x \subseteq D_f$. By compactness of K , there exist $x_1, \dots, x_l \in K$ such that

$$K \subseteq \bigcup_{i=1}^l U_{x_i} = U_N \in \mathcal{B}.$$


Como $K_{mN} \subseteq \text{int}(K_{(m+1)N})$, $\forall m \in \mathbb{N}$, there exists $M \in \mathbb{N}$ such that $K \subseteq K_{MN} \subseteq U_N \subseteq D_f$. Due to the convergence in the hypothesis, there exists λ_0 such that $\beta_{MN}(f_\lambda, f) < \frac{\epsilon}{2^{M+N}}$, for

each $\lambda > \lambda_0$. So

$$\begin{aligned}
d_K(f_\lambda, f) &\leq d_{K_{MN}}(f_\lambda, f) \\
&= 2^{M+N} \cdot 2^{-(M+N)} \beta_{MN}(f_\lambda, f) \\
&\leq 2^{M+N} \beta(f_\lambda, f) \\
&< 2^{M+N} \frac{\epsilon}{2^{M+N}} = \epsilon
\end{aligned}$$

Therefore $f_\lambda \xrightarrow{\tau_{cc}} f$.

Let U nonempty open set in X , $f \in D^{-1}(U^-)$ and $0 < \epsilon < 1$.



Due to convergence, there exists λ_0 such that

$$\beta(f_\lambda, f) < \frac{\epsilon}{2^{M+N}}, \quad \forall \lambda > \lambda_0.$$

Thus, for each $\lambda > \lambda_0$ We have that

$$\begin{aligned}
\beta_{MN}(f_\lambda, f) &= 2^{M+N} \cdot 2^{-(M+N)} \beta_{MN}(f_\lambda, f) \\
&\leq 2^{M+N} \beta(f_\lambda, f) \\
&< 2^{M+N} \frac{\epsilon}{2^{M+N}} = \epsilon < 1
\end{aligned}$$

since $f \in D^{-1}(int(K_{MN})^-)$, By definition of β_{MN} , necessarily $f_\lambda \in D^{-1}(int(K_{MN})^-)$, $\forall \lambda > \lambda_0$. Thus

$$\emptyset \neq int(K_{MN}) \cap D(f_\lambda) \subseteq U_N \cap D(f_\lambda) \subseteq U \cap D(f_\lambda), \quad \forall \lambda > \lambda_0.$$

Therefore $f_\lambda \xrightarrow{\tau_{\iota, D}} f$. □

In particular, if (X, d) is a locally compact and second countable metric space ($d \leq 1$), for each $f, g \in \Gamma(X)$

$$d_\Gamma(f, g) = \beta(f, g) + \beta(f^{-1}, g^{-1})$$

defines a metric for $(\Gamma(X), \tau_{hco})$.

1.4 $(C_{od}(X, Y), \beta)$ as Polish space

In this section, we will see that if X is a locally compact, Hausdorff and second countable space, and (Y, d) is a complete metric space ($d \leq 1$), then $(C_{od}(X, Y), \beta)$ is a Polish space. In particular, the set of holomorphic functions from open domain in \mathbb{C} is a closed subspace and thus Polish with the metric β .

Definition 15. Let X be locally compact, Hausdorff and second countable space, and Y be a complete metric space. A sequence $(f_\lambda) \subseteq C_{od}(X, Y)$ is called Γ -Cauchy if $(D(f_\lambda))$ converges to A , and in case $A \neq X$, for each $K \in \mathbb{K}(X \setminus A)$, there exists λ_K such that $K \subseteq D_{f_\lambda}$, for each $\lambda > \lambda_K$ and $(f|_K)_{\lambda > \lambda_K}$ is uniformly Cauchy. In this case, we will say that the sequence $(f_\lambda|_K)$ is *eventually uniformly Cauchy*.

Lemma 16. Let X be locally compact, Hausdorff and second countable space, and Y is metrically complete. A sequence (f_λ) in $C_{od}(X, Y)$ is Γ -Cauchy if, and only if, (f_λ) is $\tau_{\iota, D}$ -converging to $f \in C_{od}(X, Y)$.

Proof. If (f_λ) is Γ -Cauchy, $D(f_\lambda) \xrightarrow{\tau_{Fell}} A$. If $A = X$, $f_\lambda \xrightarrow{\tau_{\iota, D}} \emptyset$. If $A \neq X$, since X is second countable and $X \setminus A$ is a non-empty open set, for each $n \in \mathbb{N}$, there exist $K_n \in \mathbb{K}(X \setminus A)$ such that

$$K_n \subseteq \text{int}(K_{n+1}) \quad \text{and} \quad \bigcup_{n \in \mathbb{N}} K_n = X \setminus A$$

Since (f_λ) is Γ -Cauchy and Y is complete, $(f_\lambda|_{K_n})$ eventually converges uniformly to $f_n : K_n \rightarrow Y$, $\forall n \in \mathbb{N}$. Let

$$f = \bigvee_{n \in \mathbb{N}} f_n$$

Since $f_1 \subseteq f_2 \subseteq \dots$, f is function. $D_f = \bigcup_{n \in \mathbb{N}} K_n = X \setminus A$ (i.e. $D(f_\lambda) \xrightarrow{\tau_{Fell}} D(f)$) and if $K \in \mathbb{K}(D_f)$ and $\epsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that $K \subseteq K_{n_0} \subseteq D_f = X \setminus A$. Since (f_λ) is Γ -Cauchy, there exists λ_0 such that $K \subseteq K_{n_0} \subseteq D_{f_\lambda}$, for each $\lambda > \lambda_0$ and $(f_\lambda|_{K_{n_0}})_{\lambda > \lambda_0}$ is uniformly Cauchy. This sequence converges to $f_{n_0} = f|_{K_{n_0}}$, i.e. there exists $\lambda_1 > \lambda_0$ such that

$$d_K(f_\lambda, f) \leq d_{K_{n_0}}(f_\lambda, f) < \epsilon, \quad \forall \lambda > \lambda_1$$

Hence $f_\lambda \in B_K(f, \epsilon), \forall n > n_2$, that is, $f_\lambda \xrightarrow{\tau_{cc}} f$ and therefore $f_\lambda \xrightarrow{\tau_{\iota, D}} f$. f is continuous by the Corollary 12. \square

Theorem 17. Let X be a locally compact, Hausdorff and second countable space, and Y be a complete metric space with $d \leq 1$. Then $(C_{od}(X, Y), \beta)$ is a Polish space.

Proof. Let (f_λ) β -Cauchy in $C_{od}(X, Y)$, since $\tau_\beta = \tau_{\iota, D}$, $(D(f_\lambda))$ is τ_{Fell} -Cauchy and since $CL(X)$ is compact, $D(f_\lambda) \xrightarrow{\tau_{Fell}} A$. Si $A = X$, $f_\lambda \xrightarrow{\tau_{\iota, D}} \emptyset$. If $A \neq X$, let $K \in \mathbb{K}(X \setminus A)$ and

$\epsilon > 0$. There exist $M, N \in \mathbb{N}$ such that $K \subseteq K_{MN} \subseteq U_N \subseteq X \setminus A$. Since (f_λ) is β -Cauchy there exists λ_0 such that $\beta(f_\lambda, f_\sigma) < \epsilon 2^{-(M+N)}$, $\forall \lambda, \sigma > \lambda_0$. Hence

$$\begin{aligned} d_K(f_\lambda, f_\sigma) &= 2^{M+N} 2^{-(M+N)} d_K(f_\lambda, f_\sigma) \\ &\leq 2^{M+N} 2^{-(M+N)} \beta_{MN}(f_\lambda, f_\sigma) \\ &\leq 2^{M+N} \beta(f_\lambda, f_\sigma) \\ &< 2^{M+N} 2^{-(M+N)} \epsilon = \epsilon. \end{aligned}$$

Therefore (f_λ) is Γ -Cauchy and for the lemma 16 $\tau_{L,D}$ -converges. \square

Congeture. *If X is a localment compact polish space, with metric $d \leq 1$, then $(\Gamma(X), d_\Gamma)$ is a polish space.*

References

- [1] A. Allah and R. Brown. A compact-open topology on partial maps with open domain. *J. London Math. Soc.* (2), 21(3):480-486, 1980.
- [2] G. Beer. Topologies on closed and closed combex sets. *Kluwer Academic Publisher, Dordercht*, 1993.
- [3] J. R. Munkres. Topología, 2^a edición. *Pearson Educación, S.A.*, Madrid, 2002.
- [4] J. Perez, C. Uzcátegui. On the Polishness of the inverse semigroup $\Gamma(X)$ on a compact metric space X . *European Journal of Mathematics* (2023) 9:113.
- [5] J. van Mill. Infinite-Dimensional Topology. Prerequisites and Introduction, vol 43 of North-Holland Mathematical Library. Elsevier Science Publishers, Amsterdam, 1989.
- [6] L. Martínez, H. Pinedo and C. Uzcátegui. *A topological correspondence between partial actions of groups and inverse semigroup actions.* *Forum Math.*, 34(2), 431-446, 2022.