

NOTE ON EXISTENCE OF PHYSICAL MEASURES FOR SYSTEMS WITH MIXED CENTRAL BEHAVIOR

VITOR ARAUJO AND LUCIANA SALGADO

ABSTRACT. We show the existence of physical measures for C^∞ smooth instances of certain partially hyperbolic dynamics, both continuous and discrete, exhibiting mixed behavior (positive and negative Lyapunov exponents) along the central non-uniformly hyperbolic multidimensional invariant direction, as a consequence of assuming the existence of certain types of “regular points” on positive volume subsets. This includes the C^3 robust class of multidimensional non-hyperbolic attractors obtained by Viana, and the C^1 robust classes of 3-sectionally hyperbolic wild strange attractors presented by Shilnikov and Turaev.

CONTENTS

1. Introduction and statement of results	2
1.1. The diffeomorphism case	3
1.2. The flow case	6
1.3. An abstract setting	8
1.4. Comments and conjectures	9
Acknowledgments	10
2. Physical measures for smooth systems	10
2.1. Proof of the diffeomorphism case	11
2.2. Proof of the flow case	12
2.3. Proof of the abstract result	13
References	13

Date: May 21, 2024.

2010 Mathematics Subject Classification. Primary: 37D45. Secondary: 37D30, 37D25, 37D35.

Key words and phrases. physical/SRB measure, partial hyperbolicity, non-uniform hyperbolicity, central mixed behavior, regular points.

V.A. was partially supported by CNPq-Brazil (grant 304047/2023-6). L.S. was partially supported by FAPERJ-Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro Projects APQ1-E-26/211.690/2021 SEI-260003/015270/2021 and JCNE-E-26/200.271/2023 SEI-260003/000640/2023, by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior CAPES - Finance Code 001 and CNPq-Brazil (grant Projeto Universal 404943/2023-3).

1. INTRODUCTION AND STATEMENT OF RESULTS

The statistical viewpoint on Dynamical Systems is one of the cornerstones of most recent developments in dynamics. Given a map f from a manifold M into itself, a central concept is that of *physical measure*, a f -invariant probability measure μ whose *ergodic basin*

$$B(\mu) = \left\{ x \in M : \lim_{n \rightarrow +\infty} \frac{1}{n} S_n \varphi(x) = \int \varphi d\mu, \quad \forall \varphi \in C(M, \mathbb{R}) \right\}$$

has positive *volume* or *Lebesgue measure*, where we write $S_n \varphi(x) = S_n^f \varphi(x) = \sum_{j=0}^{n-1} \varphi(f^j x)$ for the standard ergodic sum.

In the continuous time case of a flow X_t generated by the vector field X on the manifold M , this notion naturally translates to

$$B(\mu) = \left\{ x \in M : \lim_{T \rightarrow +\infty} \frac{1}{T} \int_0^T \varphi(X_t(x)) dt = \int \varphi d\mu \quad \forall \varphi \in C(M, \mathbb{R}) \right\}$$

has positive volume. This kind of measures provides asymptotic information on a set of trajectories that one hopes is large enough to be observable in real-world models.

In particular, the existence of these measures implies that the subset

$$\mathbb{B} := \left\{ x \in M : \left(\frac{1}{n} S_n \delta_x \right)_{n \geq 1} \text{ weak}^* \text{ converges} \right\}$$

has positive volume for both the discrete case, and the continuous case with $f = X_1$ the time-1 map of a flow, where $S_n \delta_x := \sum_{j=0}^{n-1} \delta_{f^j x}$. The existence of forward Birkhoff averages for all continuous observables $\varphi : M \rightarrow \mathbb{R}$ on a full Lebesgue measure subset of points is known for all C^2 globally hyperbolic (Axiom A with no cycles) systems; see e.g. Bowen and Ruelle [9, 10]. It is also well-known that this is not true in general; see e.g. Ruelle and Takens [36, 38]. Moreover, as shown by Jiang, Liu and Qian [18], in this setting the Lyapunov exponents also exist Lebesgue almost everywhere and are Lyapunov regular – this also holds for smooth Axiom A endomorphisms and hyperbolic SRB measures [33].

Recently, it has been shown that, on the contrary, forward Birkhoff averages might not exist on open subsets of points for locally dense subsets of smooth diffeomorphisms; see e.g. Kiriki, Soma, Nakano, Li and Vargas [22, 20, 21]. Moreover, existence of Birkhoff averages on positive volume subsets of points does not imply the existence of Lyapunov exponents on the same subset, as shown by Kiriki, Li, Nakano, Soma [19], and it may happen that the set of Lyapunov regular points has not full Lebesgue measure; see e.g. Ott [30]. In [15], the second author together with Carvalho, Coelho and Varandas, proved more general results on irregular sets with respect to Cèsaro averages, including information on the Baire genericity of irregular sets for averages of semigroup actions that take into account the group structure.

Here we explore some consequences of the *assumption of existence of forward Birkhoff averages for all continuous observables on positive volume subsets of points* together with *existence and regularity of Lyapunov exponents* on the same subset of points, motivated by the above mentioned features of Axiom A diffeomorphisms together with the results of Tsujii [39]; or the assumption of *$p\omega$ -regular points on positive volume subsets* from the results of Burguet [11], applied to some robust examples of systems, both discrete and continuous.

We obtain new conditional results on the existence of physical measures for 3-sectionally expanding attracting sets for smooth flows with center-unstable *mixed behavior*; and for partially hyperbolic attractors whose center direction exhibits non-uniformly hyperbolic *mixed behavior* along the central direction. This means that, along certain multidimensional invariant directions, we have both positive and negative Lyapunov exponents Lebesgue almost everywhere.

This admits direct applications to the existence of physical measure for the “wild strange attractors” for singular flows from Shilnikov-Turaev [40]; and also to the C^∞ smooth instances of multidimensional non-uniformly expanding attractors from the class of diffeomorphisms obtained by Viana [41].

We note that mostly expanding/contracting behavior along multidimensional invariant directions is enough to ensure the existence of physical measures, as shown by Alves, Bonatti and Viana in [8, 1]; but no such general results are known for mixed behavior, among non-conservative diffeomorphisms, along multidimensional invariant directions, apart from the case of C^∞ surface diffeomorphisms by Burguet [12] (and Buzzi-Crovisier-Sarig [13]), certain Hénon maps [6], and also certain dissipative skew-products perturbations of the conservative Berger-Carrasco maps [7] studied by Carrasco and Obata [27, 14, 28].

1.1. The diffeomorphism case. We now present preliminary definitions and results.

Let $f : M \circlearrowleft$ be a diffeomorphism of a compact Riemannian manifold having an attractor Λ with a *trapping region* U (i.e. U is an open neighborhood of $\Lambda = \Lambda_f(U) := \bigcap_{n \geq 1} f^n U$ with $fU \subset U$) satisfying

- (1) **partial hyperbolicity:** the tangent bundle $T_\Lambda M$ over Λ splits into continuous Df -invariant subbundles $T_\Lambda M = E^s \oplus E^c \oplus E^u$ such that there are constants $C > 0, 0 < \lambda < 1$ satisfying for all $x \in \Lambda$ and $n \geq 1$
 - (a) E^s is uniformly contracted: $\|Df^n|_{E_x^s}\| \leq C\lambda^n$;
 - (b) E^u is uniformly expanded: $\|(Df^n|_{E_x^u})^{-1}\| \leq C\lambda^n$;
 - (c) the splitting is dominated:

$$\max \left\{ \frac{\|Df^n|_{E_x^s}\|}{\|Df^n|_{E_x^c}\|}, \frac{\|Df^n|_{E_x^c}\|}{\|Df^n|_{E_x^u}\|} \right\} \leq C\lambda^n;$$

- (2) **dissipativeness** (volume contraction) **along the central direction:** $|\det(Df^n|_{E_x^c})| \leq C\lambda^n$ for all $x \in \Lambda, n \geq 1$.

We consider a continuous extension of each bundle of the splitting to U and assume, without loss of generality, that the extended stable bundle is forward invariant [2], and also that dissipativeness and uniform expansion are preserved. We additionally assume the following.

- (3) **Non-uniform expansion along the central direction:** there exists $c_0 > 0$ such that

$$\limsup_{n \rightarrow \infty} \log \|Df^n|_{E_x^c}\|^{1/n} > c_0, \text{ for Lebesgue a.e. } x \in U. \quad (1.1)$$

1.1.1. *An open class of examples.* A C^r open set \mathcal{U} of partially hyperbolic diffeomorphisms with *two-dimensional* center bundle E^c , in the above setting, is provided by Viana in [41] for $r \geq 3$. Using the notation from [41], we consider the following skew-product $\varphi_{\alpha,b} : T_3 \times \mathbb{R}^2 \circlearrowleft, (\Theta, X) \mapsto (\hat{g}(\Theta), \hat{f}_{\alpha,b}(\Theta, X))$ with $\Theta = (\theta, T) \in T_3 = \mathbb{S}^1 \times B^2$ the solid 3-torus, where B^2 is the unit 2-disk, $X = (x, y) \in I_0^2 \subset \mathbb{R}^2$ and

$$\hat{g} : T_3 \circlearrowleft, \quad \hat{g}(\theta, T) = (d\theta \bmod 1, G(\theta, T))$$

is a solenoid embedding; and additionally

$$f_{\alpha,b} : T_3 \times \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad f_{\alpha,b}(\Theta, X) = (a(\theta) - x^2 + by, -by)$$

where

- $a(\theta) = a_0 + \alpha\omega(\theta)$ with
 - $a_0 \in (1, 2)$ so that 0 is preperiodic for the map $x \mapsto a_0 - x^2$;
 - $\omega : \mathbb{S}^1 \rightarrow \mathbb{R}, \theta \mapsto \sin 2\pi\theta$ is a Morse function; and
- $I_0 \subset (-2, 2)$ is a compact interval so that $\varphi_{\alpha,b}(T_3 \times I_0^2) \subset \text{int}(T_3 \times I_0^2)$ for every small $\alpha, b > 0$.

Hence, setting $U := T_3 \times I_0^2$ we get a diffeomorphism (into its image) $\varphi_{\alpha,b} : U \rightarrow U$ for every $b > 0$. We note that $\det Df_{\alpha,b} = b$ and $\varphi_{\alpha,b}$ is C^∞ . Moreover, clearly we have a $\varphi_{\alpha,b}$ -invariant partially hyperbolic splitting $TU = TB^2 \oplus T\mathbb{R}^2 \oplus E^u$, where $TB^2 \oplus E^u = TT_3$ is the hyperbolic \hat{g} -invariant splitting of the standard Smale solenoid attractor, with E^u a graph $T\mathbb{S}^1 \rightarrow TB^2$ over $T\mathbb{S}^1$, $E^c = T\mathbb{R}^2$ is the dissipative central direction of the skew-product, and $E^s = TB^2$ the uniformly contracting direction.

Theorem 1.1. [41, Theorem B] *Let us assume d to be large enough. Then, there exists an open set of (small positive) values of (α, b) for which $f := \varphi_{\alpha,b}$ admits a partially hyperbolic attractor $\Lambda = \Lambda_f(U)$ satisfying the conditions (1), (2) and (3) (with a two-dimensional center subbundle and \liminf in the place of \limsup in (1.1)). Moreover the same holds for all φ in a neighborhood \mathcal{U} of $\varphi_{\alpha,b}$ in $C^3(U)$.*

1.1.2. *Some subsets of regular points.* To state the main results, we recall some notions of differential ergodic theory. As usual, we write $\log^+ a = \max\{0, \log a\}$ for all $a > 0$ in what follows. For a diffeomorphism $f : M \rightarrow M$ (or for the time-1 map $f = X_1$ of a flow generated by a vector field X), we set the forward Lyapunov exponent of $v \in T_x M \setminus \{\vec{0}\}$ as

$$\chi^+(x, v) := \limsup_{n \rightarrow \infty} \log \|Df^n(x)v\|^{1/n}.$$

The function $\chi(x, \cdot)$ admits finitely many values only $\chi_1^+(x) > \dots > \chi_{p(x)}^+(x)$ on $TM \setminus \{0\}$ and generates a filtration $0 \subsetneq V_{p(x)}(x) \subsetneq \dots \subsetneq V_1 = T_x M$ with $V_i(x) = \{v \in T_x M, \chi^+(x, v) \leq \chi_i^+(x)\}$. The functions $p, \chi_i^+(x)$ are f -invariant; the vector spaces $V_i(x), i = 1, \dots, p(x)$ are Df -invariant; and each of them depend Borel measurably on x . We write

$$\chi_1^+(x) := \limsup_{n \rightarrow \infty} \log^+ \|Df^n(x)\|^{1/n} = \max\{\chi(x, v) : v \in T_x M \setminus \{0\}\}$$

for the top Lyapunov exponent of the action of $Df : TM \rightarrow TM$ on the tangent bundle at $x \in M$. We also set

$$\chi_\wedge^+(x) := \limsup_{n \rightarrow +\infty} \frac{1}{n} \log^+ \max_k \|\wedge^k Df^n(x)\|$$

for the top k -sectional Lyapunov exponents of the bundle morphism $\wedge Df$ induced by Df on the exterior algebra bundle $\wedge TM = \oplus_k \wedge^k TM$, for $k = 1, \dots, d := \dim M$; see e.g. [37, Definition 1.11].

We finally set $p\omega(x)$ the family of all weak* accumulation points of the empirical measures $\left(\frac{1}{n} S_n \delta_x\right)_{n \geq 1}$, which is always a nonempty set of f -invariant probability measures on compact ambient spaces like M . We say that x is $p\omega$ -regular if

$$\sup_{\mu \in p\omega(x)} \int \chi_\wedge^+ d\mu = \chi_\wedge^+(x).$$

We denote by \mathcal{P} the subset of $p\omega$ -regular points.

Theorem A. *Let $f : M \rightarrow M$ be a C^∞ diffeomorphism for which the subset \mathcal{P} has positive volume in a trapping region U (i.e. $\text{Leb}(\mathcal{P} \cap U) > 0$), where $\Lambda = \Lambda_f(U)$ is a partially hyperbolic attracting set satisfying conditions (1), (2) and (3) with a two-dimensional center subbundle. Then there exists an hyperbolic ergodic physical f -invariant probability measure μ with central mixed behavior, that is, for μ -a.e. $x \in \Lambda$ and for x in a positive volume subset of U we have a positive and a negative central Lyapunov exponent: there exists $c_0 > 0$ so that*

$$\lim_{n \rightarrow \infty} \log \|Df^n|_{E_x^c}\|^{1/n} > c_0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \log \|(Df^n|_{E_x^c})^{-1}\|^{1/n} > c_0. \quad (1.2)$$

This ensures the existence of a physical probability measure μ with *mixed central behavior*, that is, for μ -a.e. $x \in \Lambda$ and for x in a positive volume subset of U , we have both a positive and a negative central Lyapunov exponent as in (1.2), where it is enough that the limits be positive (i.e. $c_0 \geq 0$).

1.2. The flow case. Recall that a *trapping region* U for a flow X_t is an open subset of the manifold M which satisfies: $X_t(U)$ is contained in U for all $t > 0$, and there exists $T > 0$ such that $\overline{X_t(U)}$ is contained in the interior of U for all $t > T$. We define $\Gamma(U) = \Gamma_X(U) := \bigcap_{t>0} \overline{X_t(U)}$ to be the *maximal positive invariant subset in the trapping region* U .

A *singularity* for the vector field X is a point $\sigma \in M$ such that $X(\sigma) = \vec{0}$ or, equivalently, $X_t(\sigma) = \sigma$ for all $t \in \mathbb{R}$. The set formed by singularities is the *singular set of X* denoted $\text{Sing}(X)$. We say that a singularity is *hyperbolic* if the eigenvalues of the derivative $DX(\sigma)$ of the vector field at the singularity σ have nonzero real part.

A *dominated splitting* over a compact invariant set Λ of X is a continuous DX_t -invariant splitting $T_\Lambda M = E \oplus F$ with $E_x \neq \{0\}$, $F_x \neq \{0\}$ for every $x \in \Lambda$ and such that there are positive constants K, λ satisfying

$$\|DX_t|_{E_x}\| \cdot \|DX_{-t}|_{F_{X_t(x)}}\| < Ke^{-\lambda t}, \text{ for all } x \in \Lambda, \text{ and all } t > 0.$$

A compact invariant set Λ is said to be *partially hyperbolic* if it exhibits a dominated splitting $T_\Lambda M = E \oplus F$ such that

- the subbundle E is *uniformly contracted*: there are $C, \lambda > 0$ such that $\|DX_t|_{E_x}\| \leq Ce^{-\lambda t}$ for $t \geq 0$ and all $x \in \Lambda$.

In this case F is the *central subbundle* of Λ . Or else, we may replace uniform contraction along E by

- the subbundle F is *uniformly expanded*: there are $C, \lambda > 0$ such that $\|(DX_t|_{F_x})^{-1}\| \leq Ce^{-\lambda t}$ for $t \geq 0$ and all $x \in \Lambda$.

We can assume, without loss of generality, that a continuous extension of the stable direction E of the splitting is still $D\phi_t$ -invariant [2] and consider also a continuous extension of the center direction F to the trapping region U . The following is an extension of the notion of 2-sectional expansion proposed in [25, Definition 2.7].

We say that a DX_t -invariant subbundle $F \subset T_\Lambda M$ is *mostly p -sectionally expanding* if $\dim F_x \geq p$ is constant for $x \in \Lambda$ and for Lebesgue almost every $x \in U$

$$\limsup_{t \rightarrow +\infty} \log |(\wedge^p DX_t|_{\wedge^p F_x})|^{1/t} > 0. \quad (1.3)$$

This means that for every p -dimensional linear subspace $L_x \subset F_x$ one has

$$\limsup_{t \rightarrow +\infty} \log |\det(DX_t|_{L_x})|^{1/t} > 0. \quad (1.4)$$

A *mostly p -sectional-hyperbolic set* is a partially hyperbolic set whose central subbundle is mostly p -sectionally expanding – this extends the notion of sectional hyperbolicity given by the second author in [37, Definition 3].

This is a particular case of the so called *singular-hyperbolicity* whose definition we recall now. A DX_t -invariant subbundle $F \subset T_\Lambda M$ is said to be *volume expanding* if we replace (1.4) by

$$|\det(DX_t|_{F_x})| > Ce^{\lambda t}, \text{ for all } t > 0.$$

A *singular-hyperbolic set* is a partially hyperbolic set whose central subbundle is volume expanding [26, Definition 1].

Clearly, in the three-dimensional case, these notions are equivalent.

Remark 1.2. An invariant subbundle F which is 2-sectionally expanding and $\dim F \geq 3$, is also p -sectionally expanding for each $p = 3, \dots, \dim F$. However, the reciprocal is not true, as shown by the open class of examples to be discussed in what follows, since sectional contraction may occur in k -subspaces with $k < p$.

To include *contracting behavior* along F , we say that an invariant subbundle F of the tangent bundle $T_\Lambda M$ is *partially k -sectionally contracting* if there are $C, \lambda > 0$ so that for every $x \in \Lambda$ we can find a k -dimensional subspace $L_x \subset F_x$ for which

$$|\det(DX_t|_{L_x})| \leq Ce^{-\lambda t}, \text{ for all } t > 0. \quad (1.5)$$

We say that the (extended) central-bundle of a partially hyperbolic splitting is *mostly k -sectionally contracting* if for Lebesgue almost every $x \in U$, there exists a k -dimensional subspace $L_x \subset F_x$ such that

$$\limsup_{t \rightarrow +\infty} \log |\det(DX_t|_{L_x})|^{1/t} < 0. \quad (1.6)$$

1.2.1. *An open class of examples.* In [40], Turaev and Shilnikov construct a *wild attractor* Γ (see Figure 1) of a vector field having a *singular hyperbolic splitting* $T_\Gamma M = E \oplus F$, where $\dim F = 3$ and $\dim E = n - 3$ for any $n \geq 4$; and the attractor contains a (hyperbolic) saddle-focus singularity at the origin. Moreover, the bundle F is *2-sectionally contracting*; see [40, p 296, formula (13)] and more recently [16] for other concrete examples of ODE's displaying similar features.




FIGURE 1. Sketch of the trapping region of Turaev-Shil'nikov's wild strange attractor around the unfolding of a saddle-focus equilibrium at the origin.

These attractors are also robust: these properties persist for a C^1 neighborhood \mathcal{U} of the original vector field. This means that the subbundle F satisfies

- 3-sectional expansion, i.e. (1.4) (or just the volume along F , since $\dim F = 3$) holds for all $x \in U$ for $p = 3$ and a fixed positive value of t ; and also
- partial 2-sectional contraction, i.e. it satisfies (1.5): there exist $T > 0$ so that for every $x \in U$ we can find a 2-dimensional subspace $L_x \subset F_x$ so that $|\det DX_T|_{L_x}| < 1$.

Moreover, the C^1 robust property ensures that in \mathcal{U} there exists a C^s dense subset of C^r vector fields exhibiting the same properties, for $2 \leq s \leq r \leq \infty$.

Theorem B. *Let $X : M \rightarrow TM$ be a C^∞ vector field for which the subset \mathcal{P} has positive volume in a trapping region U (i.e. $\text{Leb}(\mathcal{P} \cap U) > 0$), where the attracting set $\Gamma_X(U)$ is partially hyperbolic, $\dim F = 3$, the system is mostly 3-sectionally expanding (1.4) and 2-sectionally contracting (1.6); and all singularities in U are hyperbolic. Then there exists an hyperbolic ergodic physical X_t -invariant probability measure μ with central mixed behavior, that is, for μ -a.e. $x \in \Lambda$ and for x in a positive volume subset of U we have a positive and a negative central Lyapunov exponent:*

$$\lim_{t \rightarrow \infty} \log \|DX_t|_{F_x}\|^{1/t} > 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \log \|(DX_t|_{F_x})^{-1}\|^{1/t} > 0.$$

1.3. An abstract setting. Now we go back to the diffeomorphism $f : M \rightarrow M$ and recall that a point $x \in M$ is *Lyapunov regular* if, for all $1 \leq k \leq d$, the *sectional Lyapunov exponents* satisfy

$$\chi^k(x) := \limsup_{n \rightarrow +\infty} \log \|\wedge^k Df^n(x)\|^{1/n} = \sum_{i=1}^k \chi_i^+(x). \quad (1.7)$$

By Oseledets' Theorem [29], the set of Lyapunov regular points has full measure for any f -invariant probability measure.

Motivated by Tsujii [39], we say that $x \in M$ is *T -regular* for if $x \in \mathbb{B}$ (so that $p\omega(x) = \{\mu_x\}$) and x is $p\omega$ -regular, and also Lyapunov regular (1.7) for μ_x , that is, we have $\chi_i^+(x) = \chi_i^+(y)$ for μ_x -a.e. y .

We denote by \mathcal{R} the subset of T -regular points in what follows.

1.3.1. Unimodal sectional Lyapunov spectrum. We say that the *sectional Lyapunov spectrum* of a point $x \in M$, given by $\chi^i(x), i = 1, \dots, p(x)$, is *unimodal* if there exists $1 \leq k(x) < p(x)$ so that

$$\chi^{p(x)}(x) < \dots < \chi^{k(x)}(x) \quad \text{and} \quad \chi^{k(x)}(x) > \dots > \chi^1(x). \quad (1.8)$$

If in addition $x \in \mathcal{R}$, then the Lyapunov regularity ensures that the μ_x is hyperbolic, since

$$\chi_1^+(x) = \chi^1(x) \quad \text{and} \quad \chi_i^+(x) = \chi^i(x) - \chi^{i-1}(x), \quad i = 2, \dots, d = \dim M;$$

and thus $\chi_d^+(x) \leq \dots \leq \chi_{k(x)+1}^+(x) < 0 < \chi_{k(x)}^+(x) \leq \dots \leq \chi_1^+(x)$.

When $f = X_1$ is the time-1 map of a flow, then we allow for the zero Lyapunov exponent along the flow direction: we say that x has *unimodal sectional Lyapunov spectrum* for a flow if there exists $1 < k(x) < p(x)$ so that

$$\chi^{p(x)}(x) < \dots < \chi^{k(x)}(x) \ \& \ \chi^{k(x)}(x) = \chi^{k(x)-1}(x) > \dots > \chi^1(x). \quad (1.9)$$

This condition (1.8) or (1.9) is enough to obtain a physical/SRB measure for C^∞ maps or flows *assuming simultaneously that the collection of such points has positive volume inside the family \mathcal{R} of regular points.*

Theorem C. *Let $f : M \rightarrow M$ be a C^∞ diffeomorphism (time-1 map of a flow X_t induced by a C^∞ vector field) for which the subset S of points with unimodal spectrum has positive volume among the regular points \mathcal{R} , i.e. $\text{Leb}(S \cap \mathcal{R}) > 0$. Then there exists an hyperbolic ergodic physical f -invariant (X_t -invariant) probability measure.*

1.4. Comments and conjectures. The statements Theorems A, B and C are corollaries of the results of Burguet [11], and the definition of regular point is inspired by the main results from Tsujii [39]. The proofs be detailed in Section 2.

We note that the central bundle dynamics of the skew-products $\varphi_{\alpha,b}$ in the statement of Theorem 1.1 is a perturbation of the Hénon family [17]. For this polynomial family of dissipative maps, a positive Lebesgue measure of parameters was obtained by Benedicks and Carleson [5] for which the system is non-uniform expanding, that is, condition (3) is satisfied. The recent results for C^∞ smooth surface dynamics of Burguet [12] would automatically ensure the existence and uniqueness of a physical measure for these system, which was obtained with a completely different strategy, specifically adapted to these systems, by Benedicks and Young [6].

The conditional existence results of physical measures stated in Theorems A and B, together with the positive results on “Takens last problem” [22], naturally raise the following questions.

Conjecture 1. For a dense/generic subset of dynamical systems, there exists a positive volume subset in the ambient space where at least one of the following properties hold:

- (1) the points are $p\omega$ -regular;
- (2) forward Birkhoff averages exist on all points;
- (3) forward Lyapunov exponents exist and are Lyapunov regular.

This would automatically provide for the existence of physical measures among the smooth systems in the open classes of non-uniformly hyperbolic flows and diffeomorphisms described in the previous subsection, and provide for a criterion to obtain physical/SRB measures using the sectional Lyapunov spectrum.

Since Hertz, Hertz, Tahzibi and Ures [34] have shown the uniqueness of SRB measures for transitive diffeomorphisms on surfaces, and Obata [28] obtained uniqueness for non-conservative perturbations of certain skew-products

with two-dimensional center bundle with mixed behavior, the fact that the non-hyperbolic center direction of the family $\varphi_{\alpha,b}$ is two-dimensional suggests the following.

Conjecture 2. There exists at most one SRB/physical measure for each of the maps $\varphi_{\alpha,b}$ in the setting of Theorem A.

Assuming a positive answer to the previous conjectures, it is natural to consider the dependence of the physical measure on the dynamics of the open families \mathcal{U} of maps considered in Subsections 1.1.1 and 1.2.1. Ideally we would like to obtain *statistical stability*, that is, if μ_n is the physical measure for the dynamics of f_n , and f_n converges to $f \in \mathcal{U}$ in some C^r topology ($2 \leq r \leq \infty$), then μ_n weakly converges to a f -invariant probability measure μ , which is also a physical measure for f .

Conjecture 3. The physical measures of the C^∞ dynamics in \mathcal{U} are statistically stable.

This would naturally lead to obtain physical measures for most dynamics among the open classes of systems considered here. A natural extension of this work is to study statistical properties (e.g. mixing and speed of mixing) of these physical measures.

Acknowledgments. V.A. thanks the Mathematics and Statistics Institute of the Federal University of Bahia (Brazil) for its support of basic research and CNPq (Brazil) for partial financial support. L.S. thanks the Mathematics Institute of Universidade Federal do Rio de Janeiro (Brazil) for its encouraging of mathematical research, and the Mathematics and Statistics Institute of the Federal University of Bahia (Brazil) for its hospitality, as well as CAPES-Finance code 001, CNPq, FAPERJ - Auxílio Básico à Pesquisa (APQ1) Project E-26/211.690/2021 and FAPERJ - Jovem Cientista do Nosso Estado (JCNE) grant E-26/200.271/2023, for partial financial support.

2. PHYSICAL MEASURES FOR SMOOTH SYSTEMS

Here we present the main result from Burguet [11], involving the subset \mathcal{P} , needed to obtain the main statements.

Theorem 2.1. [11, Main Theorem] *Let there be given a C^∞ map $f : M \rightarrow M$. Then for Lebesgue almost every point x there exists $\mu_x \in p\omega(x)$ such that $h_{\mu_x}(f) \geq \chi_\lambda^+(x)$.*

We note that, in the setting of Theorem 2.1, we obtain for Leb-a.e. $x \in \mathcal{P}$

$$\chi_\lambda^+(x) \leq h_{\mu_x}(f) \leq \int \chi_\lambda^+ d\mu_x = \chi^+(\mu) \leq \sup_{\mu \in p\omega(x)} \int \chi_\lambda^+ d\mu = \chi_\lambda^+(x),$$

where we used the *Ruelle Inequality* [35] for the entropy

$$h_\nu(f) \leq \int \chi_\lambda^+ d\nu$$

for any f -invariant probability measure ν ; together with the definition of $p\omega$ -regularity.

Therefore, μ_x satisfies the *Pesin Entropy Formula* $h_{\mu_x}(f) = \int \chi_{\wedge}^+ d\mu_x$ with positive entropy.

We recall that, from the results of Ledrappier and Young [23], the measures μ_x satisfying the Pesin Entropy Formula are Sinai-Ruelle-Bowen (SRB) measures, which are dynamically characterized as *μ_x -almost every point has a positive Lyapunov exponent and the disintegration of μ along the unstable manifolds is absolutely continuous with respect to the volume on the unstable manifolds inherited from the Riemannian structure on the ambient manifold.*

It is well-known, from Pesin's Nonuniformly Hyperbolic Theory, that any ergodic hyperbolic SRB measure is physical [32, 4].

Consequently, if in addition μ_x has no zero Lyapunov exponents (or only one zero Lyapunov exponent along the direction of the flow), then μ_x becomes a physical measure.

2.1. Proof of the diffeomorphism case. A consequence of the assumptions of Theorem A and Theorem 2.1, is that for f a C^∞ map and $x \in U \cap \mathcal{P}$, we have that μ_x is an SRB measure. To conclude the statement of Theorem A it is enough to show that μ_x is hyperbolic with mixed behavior along the central direction.

Let then μ_x be an SRB measure with $x \in U$ satisfying (1.1). Since $T_\Lambda M = E^s \oplus E^c \oplus E^u$ and E^s is uniformly contracting, E^u is uniformly expanding, and E^c is two-dimensional and area contracting, then for $1 \leq i \leq d_u := \dim E^u$ and $n \geq 1$ we get

$$\|\wedge^i Df^n\| \geq \|\wedge^i (Df^n|_{E_x^u})\| \geq \|\wedge^i (Df^n|_{E_x^u})^{-1}\|^{-1} \quad (2.1)$$

$$\geq \|(Df^n|_{E_x^u})^{-1}\|^{-i} \geq C^{-i} \lambda^{-ni} \quad (2.2)$$

and so $\chi^{d_u}(x) \geq \chi^{d_u-1}(x) \geq \dots \geq \chi^1(x) > \log \lambda^{-1} > 0$. From domination we get, for each $u \in \wedge^{d_u} E_x^u$ and $v \in E_x^c$

$$\|\wedge^{d_u+1} Df_x^n\| \geq \|(\wedge^{d_u+1} Df_x^n)(v \wedge u)\| \approx \|Df_x^n v\| \cdot \|\wedge^{d_u} Df_x^n u\|,$$

where \approx means equality up to a bounded multiplicative constant. From (1.1) we can find $v \in E_x^c$ so that $\chi^+(x, v) > c_0 > 0$ and from (2.1) we can find $u \in \wedge^{d_u} E_x^u$ which realizes the rate of volume growth along E^u , so we obtain $\chi^{d_u+1}(x) = \chi_{d_u+1}^+(x) + \chi^{d_u}(x) > \chi^{d_u}(x) + c_0 > \chi^{d_u}(x)$. Moreover,

$$\begin{aligned} \|\wedge^{d_u+2} Df_x^n\| &\approx \|\wedge^2 (Df^n|_{E_x^c})\| \cdot \|\wedge^{d_u} (Df^n|_{E_x^u})\| \\ &= |\det(Df^n|_{E_x^c})| \cdot \|\wedge^{d_u} (Df^n|_{E_x^u})\| \leq C \lambda^n \|\wedge^{d_u} (Df^n|_{E_x^u})\|, \end{aligned}$$

and therefore we conclude

$$\chi^{d_u+2}(x) < \chi^{d_u}(x) + \log \lambda < \chi^{d_u} + c_0 < \chi^{d_u+1}(x).$$

Since E^s is uniformly contracting, then we also get $\chi^{d_u+2+i+1}(x) < \chi^{d_u+2+i}(x)$ for all $i = 0, \dots, \dim E^s$. This ensures that $\chi^{d_u+1}(x) = \chi_{\wedge}^+(x) = h_{\mu_x}(f)$.

Thus, μ_x is hyperbolic with a Lyapunov spectrum satisfying

$$\chi_{d_u+2}^+(x) = \chi^{d_u+2}(x) - \chi^{d_u+1}(x) < \log \lambda - c_0 < 0 < c_0 < \chi_{d_u+1}^+(x)$$

Hence, the minimal exponent along the subbundle E_x^c is smaller than $\log \lambda - c_0$. This concludes the proof of Theorem A.

2.2. Proof of the flow case. Similarly, to conclude the statement of Theorem B it is enough to show that μ_x is hyperbolic with mixed behavior along the central direction for Leb-a.e. $x \in U \cap \mathcal{P}$ with a C^∞ vector field X . Then, from 3-sectional expansion, we obtain

$$\chi_3^+(x) + \chi_2^+(x) + \chi_1^+(x) \geq \chi^3(x) > 0,$$

which ensures that $\chi_1^+(x) > 0$. Since the hyperbolic singularities within U are hyperbolic and are fixed points of the dynamics, then they belong to Γ and the sectional expansion assumption ensures that each singularity is of saddle-type. Hence, the union of the stable manifold of the singularities has zero volume¹, and so Leb-a.e. $x \in U$ has a zero Lyapunov exponent along the flow direction. Moreover, the flow direction is contained in F due to the following.

Lemma 2.2. [3, Lemma 3.2] *Let Λ be a compact invariant set for the vector field X .*

- (1) *Given a continuous splitting $T_\Lambda M = E \oplus F$ such that E is uniformly contracted, then $X(x) \in F_x$ for all $x \in \Lambda$.*
- (2) *Assuming that Λ is non-trivial and has a continuous and dominated splitting $T_\Lambda M = E \oplus F$ such that $X(x) \in F_x$ for all $x \in \Lambda$, then E is a uniformly contracted subbundle.*

In addition, from 2-sectional contraction, we get that there exists $v \in F_x \setminus \{0\}$ so that $\chi^+(x, v) < 0$, and thus we conclude that $\chi_3^+(x) < 0 = \chi_2^+(x)$.

This shows that $\chi_\Lambda^+(x) = \chi^1(x) = \chi^2(x)$, that μ_x is hyperbolic (since E is uniformly contracted) and, consequently, a *physical measure* with mixed central behavior. This completes the proof of Theorem B since, clearly

$$\lim_{t \rightarrow \infty} \log \|DX_t|_{F_y}\|^{1/t} = \limsup_{n \rightarrow +\infty} \log \|DX_1^n|_{F_y}\|^{1/n} = \chi_1^+(y) > 0$$

and also

$$\lim_{t \rightarrow \infty} \log \|(DX_t|_{F_y})^{-1}\|^{1/t} = \limsup_{t \rightarrow \infty} \log \|(DX_1^n|_{F_y})^{-1}\|^{1/n} = -\chi_3^+(y) > 0$$

for μ_x -a.e. $y \in B(\mu_x) \cap U$.

Remark 2.3. Up to this point in this section we have not assumed that x is Lyapunov regular.

¹Since the stable manifold of a hyperbolic critical element in an immersed submanifold [31], then it has zero volume as a subset of the ambient manifold

2.3. Proof of the abstract result. To prove Theorem C, we note that by assumption we can find $x \in \mathcal{R}$ which is simultaneously $p\omega$ -regular, Lyapunov regular and has unimodal sectional Lyapunov spectrum, so that the measure $\mu_x \in p\omega(x)$, given by Theorem 2.1, satisfies Pesin's Entropy Formula $h_{\mu_x}(f) = \int \chi_\wedge^+ d\mu_x$.

From Ergodic Decomposition [24, Chapter II, Section 6] and Jacobs' Theorem [42, Theorem 9.6.2] we can assume without loss of generality that μ_x is ergodic (otherwise, there exists an ergodic component which also satisfies Pesin's Entropy Formula) and is hyperbolic (since the Lyapunov exponents $\chi_i^+(x)$ coincide with the exponents $\chi_i^+(y)$ for μ_x -a.e. y).

Hence, μ_x is a hyperbolic ergodic physical measure as claimed.

This completes the proof of Theorem C.

REFERENCES

- [1] J. F. Alves, C. Bonatti, and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly expanding. *Invent. Math.*, 140(2):351–398, 2000.
- [2] V. Araujo and I. Melbourne. Existence and smoothness of the stable foliation for sectional hyperbolic attractors. *Bulletin of the London Mathematical Society*, 49(2):351–367, 2017.
- [3] V. Araujo and L. Salgado. Infinitesimal Lyapunov functions for singular flows. *Mathematische Zeitschrift*, 275(3-4):863–897, 2013.
- [4] L. Barreira and Y. Pesin. *Nonuniform hyperbolicity*, volume 115 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2007. Dynamics of systems with nonzero Lyapunov exponents.
- [5] M. Benedicks and L. Carleson. The dynamics of the Hénon map. *Annals of Math.*, 133:73–169, 1991.
- [6] M. Benedicks and L. S. Young. SBR-measures for certain Hénon maps. *Invent. Math.*, 112:541–576, 1993.
- [7] P. Berger and P. D. Carrasco. Non-uniformly hyperbolic diffeomorphisms derived from the standard map. *Communications in Mathematical Physics*, 329(1):239–262, Jul 2014.
- [8] C. Bonatti and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly contracting. *Israel J. Math.*, 115:157–193, 2000.
- [9] R. Bowen. *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, volume 470 of *Lect. Notes in Math*. Springer Verlag, 1975.
- [10] R. Bowen and D. Ruelle. The ergodic theory of Axiom A flows. *Invent. Math.*, 29:181–202, 1975.
- [11] D. Burguet. Entropy of physical measures for C^∞ dynamical systems. *Communications in Mathematical Physics*, 375(2):1201–1222, July 2019.
- [12] D. Burguet. SRB measures for C^∞ surface diffeomorphisms. *Preprint arxiv.org*, 2111.06651, 2022.
- [13] J. Buzzi, S. Crovisier, and O. Sarig. Another proof of Burguet's existence theorem for SRB measures of C^∞ surface diffeomorphisms. *Preprint arxiv.org*, 2201.03165, 2022.
- [14] P. D. Carrasco and D. Obata. A new example of robustly transitive diffeomorphism. *Mathematical Research Letters*, 28(3):665–679, 2021.
- [15] M. Carvalho, V. Coelho, L. Salgado, and P. Varandas. Sensitivity and historic behavior for continuous maps on baire metric spaces. *Ergodic Theory and Dynamical Systems*, page 1–30, 2023.
- [16] S. Gonchenko, A. Kazakov, and D. Turaev. Wild pseudohyperbolic attractor in a four-dimensional Lorenz system. *Nonlinearity*, 34(4):2018, feb 2021.

- [17] M. Hénon. A two dimensional mapping with a strange attractor. *Comm. Math. Phys.*, 50:69–77, 1976.
- [18] D.-q. Jiang, P.-d. Liu, and M. Qian. Lyapunov exponents of hyperbolic attractors. *manuscripta mathematica*, 108(1):43–67, May 2002.
- [19] S. Kiriki, X. Li, Y. Nakano, and T. Soma. Abundance of observable lyapunov irregular sets. *Communications in Mathematical Physics*, 391(3):1241–1269, Feb. 2022.
- [20] S. Kiriki, X. Li, Y. Nakano, T. Soma, and E. Vargas. Takens’ last problem and strong pluripotency. *Preprint arxiv.org*, 2404.17932, 2024.
- [21] S. Kiriki, Y. Nakano, and T. Soma. Pluripotency of wandering dynamics. *Preprint arxiv.org*, 2404.00337, 2024.
- [22] S. Kiriki and T. Soma. Takens’ last problem and existence of non-trivial wandering domains. *Advances in Mathematics*, 306:524 – 588, 2017.
- [23] F. Ledrappier and L. S. Young. The metric entropy of diffeomorphisms I. Characterization of measures satisfying Pesin’s entropy formula. *Annals of Mathematics*, 122:509–539, 1985.
- [24] R. Mañé. *Ergodic theory and differentiable dynamics*. Springer Verlag, New York, 1987.
- [25] R. Metzger and C. Morales. Sectional-hyperbolic systems. *Ergodic Theory and Dynamical System*, 28:1587–1597, 2008.
- [26] C. A. Morales, M. J. Pacifico, and E. R. Pujals. Singular hyperbolic systems. *Proc. Amer. Math. Soc.*, 127(11):3393–3401, 1999.
- [27] D. Obata. On the stable ergodicity of Berger–Carrasco’s example. *Ergodic Theory and Dynamical Systems*, 40(4):1008–1056, Sept. 2018.
- [28] D. Obata. Open sets of partially hyperbolic skew products having a unique srb measure. *Advances in Mathematics*, 427:109136, 2023.
- [29] V. I. Oseledets. A multiplicative ergodic theorem: Lyapunov characteristic numbers for dynamical systems. *Trans. Moscow Math. Soc.*, 19:197–231, 1968.
- [30] W. Ott and J. A. Yorke. When lyapunov exponents fail to exist. *Physical Review E*, 78(5), Nov. 2008.
- [31] J. Palis and W. de Melo. *Geometric Theory of Dynamical Systems*. Springer Verlag, 1982.
- [32] Y. B. Pesin. Characteristic Lyapunov exponents and smooth ergodic theory. *Russian Math. Surveys*, 324:55–114, 1977.
- [33] M. Qian, J.-S. Xie, and S. Zhu. *Smooth ergodic theory for endomorphisms*. Lecture Notes in Mathematics. Springer, Berlin, Germany, 2009 edition, July 2009.
- [34] F. Rodriguez Hertz, M. A. Rodriguez Hertz, A. Tahzibi, and R. Ures. Uniqueness of srb measures for transitive diffeomorphisms on surfaces. *Communications in Mathematical Physics*, 306(1):35–49, Aug 2011.
- [35] D. Ruelle. An inequality for the entropy of differentiable maps. *Bol. Soc. Bras. Mat.*, 9:83–87, 1978.
- [36] D. Ruelle. Historical behaviour in smooth dynamical systems david ruelle. In B. Krauskopf, H. Broer, and G. Vegter, editors, *Global Analysis of Dynamical Systems*, Festschrift dedicated to Floris Takens for his 60th birthday, pages 63–65. Taylor & Franci, 2001.
- [37] L. Salgado. Singular hyperbolicity and sectional Lyapunov exponents of various orders. *Proceedings of the American Mathematical Society*, 147(2):735–749, 2019.
- [38] F. Takens. Orbits with historic behaviour, or non-existence of averages. *Nonlinearity*, 21(3):T33, 2008.
- [39] M. Tsujii. Regular points for ergodic Sinai measures. *Transactions of the American Mathematical Society*, 328(2):747–766, 1991.
- [40] D. V. Turaev and L. P. Shil’nikov. An example of a wild strange attractor. *Mat. Sb.*, 189(2):137–160, 1998.

- [41] M. Viana. Multidimensional nonhyperbolic attractors. *Inst. Hautes Études Sci. Publ. Math.*, 85:63–96, 1997.
- [42] M. Viana and K. Oliveira. *Foundations of ergodic theory*, volume 151 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016.

(V.A.) UNIVERSIDADE FEDERAL DA BAHIA, INSTITUTO DE MATEMÁTICA, AV. ADHEMAR DE BARROS, S/N , ONDINA, 40170-110 - SALVADOR-BA-BRAZIL
Email address: vitor.d.araujo@ufba.br or vitor.araujo.im.ufba@gmail.com
URL: <https://sites.google.com/view/vitor-araujo-ime-ufba>

(L.S.) UNIVERSIDADE FEDERAL DO RIO DE JANEIRO, INSTITUTO DE MATEMÁTICA, AVENIDA ATHOS DA SILVEIRA RAMOS 149 CIDADE UNIVERSITÁRIA, P.O. BOX 68530, 21941-909 RIO DE JANEIRO-RJ-BRAZIL
Email address: lsalgado@im.ufrj.br, lucianasalgado@ufrj.br
URL: <http://www.im.ufrj.br/~lsalgado>