

REDUCTION THEORY FOR STABLY GRADED LIE ALGEBRAS

JACK A. THORNE

ABSTRACT. We define a reduction covariant for the representations à la Vinberg associated to stably graded Lie algebras. We then give an analogue of the LLL algorithm for the odd split special orthogonal group and show how this can be combined with our theory to effectively reduce the coefficients of vectors in a representation connected to 2-descent for odd hyperelliptic curves.

CONTENTS

1. Introduction	1
2. The reduction covariant	3
3. Example: The stable $\mathbf{Z}/2\mathbf{Z}$ -grading of \mathfrak{sl}_{2g+1}	6
4. A reduction algorithm for $\mathrm{SO}(J)$	8
5. Example: Reducing a self-adjoint linear operator	11
References	12

1. INTRODUCTION

Let \mathfrak{h} be a semisimple Lie algebra over a field k of characteristic 0, and let $m \geq 1$. By a $\mathbf{Z}/m\mathbf{Z}$ -grading, we mean a direct sum decomposition

$$\mathfrak{h} = \bigoplus_{i \in \mathbf{Z}/m\mathbf{Z}} \mathfrak{h}_i,$$

where for all i, j we have $[\mathfrak{h}_i, \mathfrak{h}_j] \subset \mathfrak{h}_{i+j}$. To such a grading, we can associate a pair (G, V) , where G is the connected component of the identity in the group of automorphisms of \mathfrak{h} which preserve the grading, and $V = \mathfrak{h}_1$ is a k -vector space which is naturally a representation of the reductive group G . This representation is coregular, in the sense that the ring $k[V]^G$ of invariant polynomials is freely generated and the geometric quotient $B = \mathrm{Spec} k[V]^G$ is isomorphic to affine space [Vin76].

We say that the grading is stable if V contains G -orbits which are stable in the sense of geometric invariant theory (i.e. closed, with finite stabilizers in G). In this case the G -invariant locus $V^s \subset V$ of stable vectors is Zariski open and non-empty, and it is of interest to study the set of $G(k)$ -orbits in a given geometric stable orbit. The stable gradings have been classified [RLYG12] (at least when k is algebraically closed) and experience shows that it is frequently the case that there is a family of projective curves $C \rightarrow B^s$ and for each $b \in B^s(k)$ an injection

$$\mathrm{Pic}^0 C_b(k)/m \mathrm{Pic}^0 C_b(k) \hookrightarrow G(k) \backslash V_b(k).$$

When $k = \mathbf{Q}$, this can often be extended to an injection (with source the m -Selmer group of the Jacobian variety $\text{Pic}^0 C_b$)

$$\text{Sel}_m \text{Pic}^0 C_b \hookrightarrow G(\mathbf{Q}) \backslash V_b(\mathbf{Q}),$$

with image is contained in the image of the map

$$G(\mathbf{Z}) \backslash V_b(\mathbf{Z}) \rightarrow G(\mathbf{Q}) \backslash V_b(\mathbf{Q})$$

(for a suitable choice of integral structures). For example, this is the case for stable $\mathbf{Z}/m\mathbf{Z}$ -gradings of the semisimple Lie algebras of types A_2 , D_4 , E_6 and E_8 for $m = 2, 3, 4$ and 5 , respectively, in which case C can be taken to be the Weierstrass family of elliptic curves (see [CFS10, Fis13]), and for the stable $\mathbf{Z}/2\mathbf{Z}$ -gradings of the semisimple Lie algebras of types A_n , D_n , and E_n (see e.g. [BG13, Sha19, Lag22]).

It is therefore of particular interest to be able to understand the sets $G(\mathbf{Z}) \backslash V_b(\mathbf{Z})$ of integral orbits. This is the goal of reduction theory, which can be understood in this context to have two steps.

In the first step, one defines a *reduction covariant*, i.e. a $G(\mathbf{R})$ -equivariant map

$$\mathcal{R} : V^s(\mathbf{R}) \rightarrow X_G,$$

where X_G is the symmetric space of G (i.e. the homogeneous space $G(\mathbf{R})/K$, where K is a maximal compact mod centre subgroup of $G(\mathbf{R})$). In the second step, one introduces a notion of reducedness for elements of the symmetric space X_G , and say that an element $T \in V^s(\mathbf{R})$ is reduced if its reduction covariant $\mathcal{R}(T)$ satisfies this condition. For example, suppose given a fundamental set $\mathcal{F} \subset X_G$ for the action of $G(\mathbf{Z})$, i.e. a subset of \mathcal{F} such that $G(\mathbf{Z}) \cdot \mathcal{F} = X_G$ and \mathcal{F} intersects each orbit of $G(\mathbf{Z})$ finitely many times. Then we could say that T is \mathcal{F} -reduced if $\mathcal{R}(T) \in \mathcal{F}$. An immediate consequence is that any $T \in V^s(\mathbf{R})$ is $G(\mathbf{Z})$ -conjugate to an \mathcal{F} -reduced element, and that if $b \in B(\mathbf{Z}) \cap B^s(\mathbf{R})$ then $V_b(\mathbf{Z})$ contains only finitely many \mathcal{F} -reduced elements.

In this paper we consider both the first and second steps described in the previous paragraph. Reduction covariants have been defined in the literature in some isolated cases for representations arising from stably graded Lie algebras (including, in the papers [CFS10, Fis13], for the representations V associated to m -descent on elliptic curves for $m = 2, 3, 4, 5$). The first main result (Theorem 2.7) of this paper is the definition of a reduction covariant $\mathcal{R} : V^s(\mathbf{R}) \rightarrow X_G$ for any stably $\mathbf{Z}/m\mathbf{Z}$ -graded Lie algebra. To define it, we think of X_G as the space of Cartan involutions of the real semisimple group $G_{\mathbf{R}}$. It turns out that for any Cartan subspace $\mathfrak{c} \subset V_{\mathbf{R}}$, there is a unique Cartan involution of $\mathfrak{h}_{\mathbf{R}}$ which respects the grading and leaves \mathfrak{c} invariant. A stable vector T is contained a unique Cartan subspace, and we define $\mathcal{R}(T)$ to be the restriction of the Cartan involution associated to this subspace to $G_{\mathbf{R}}$.

Regarding the second step, a good starting point is the reduction theory of Borel and Harish-Chandra [BHC62, Bor19]. They defined Siegel sets $\mathfrak{S}_t \subset G(\mathbf{R})$, which are rather explicit fundamental sets in the sense defined above, and which can serve to define a notion of reducedness. However, what is lacking is general is an explicit algorithm to transport elements into the Siegel set. One case where the existing theory works well is when G is isogenous to SL_n (or more generally, a product of such groups). In this case X_G may be identified with the set of lattices in \mathbf{R}^n of volume 1 and lattice reduction algorithms (such as the LLL algorithm described in [LLL82]) apply without change. The second main result of this paper, which

appears in §4, is an algorithm to transport elements of X_G into a Siegel set in the case that G is a split special orthogonal group SO_{2g+1} . This is the group that appears for the stable $\mathbf{Z}/2\mathbf{Z}$ -grading of \mathfrak{sl}_{2g+1} , used in [BG13] to study the 2-Selmer groups of Jacobians of odd hyperelliptic curves. This algorithm is based on the LLL algorithm. Roughly speaking, we observe that the two main steps of the LLL algorithm correspond either to acting by “integral unipotent transformations” or “simple reflections in S_n ”. We simply replace S_n by the Weyl group of SO_{2g+1} . We consider the analogue of this, where SO_{2g+1} is replaced by any split semisimple group, in another paper [RT24].

1.1. Outline. In §2, we define the reduction covariant \mathcal{R} using the theory of the Cartan involution. In §3, we compute the covariant explicitly in the case of the stable $\mathbf{Z}/2\mathbf{Z}$ -grading of \mathfrak{sl}_{2g+1} . In §4, we give our algorithm to move a given element of $X_{\mathrm{SO}_{2g+1}}$ into a Siegel set. Finally, in §5, we show how these ideas work together in an explicit example.

1.2. Notation. In this paper, a reductive group over a field k means a smooth connected linear algebraic group over a field k of trivial unipotent radical. In particular, by a reductive group H over \mathbf{R} we mean a connected algebraic group (although the associated set $H(\mathbf{R})$ of real points might not be connected). Similarly, a semisimple group is a reductive group of trivial radical.

We will use gothic letters to denote Lie algebras (so $\mathrm{Lie} H = \mathfrak{h}$) and subscripts to denote base extension (so if H is an algebraic group over \mathbf{R} , then $(\mathrm{Lie} H) \otimes_{\mathbf{R}} \mathbf{C} = \mathfrak{h}_{\mathbf{C}}$). We use the notations $Z_H(\cdot)$ and $Z_{\mathfrak{h}}(\cdot)$ to denote group and Lie algebra centralizer, respectively.

We use superscript 0 to denote the connected component of the identity. Thus if H is a linear algebraic group over \mathbf{R} then there is an inclusion $H(\mathbf{R})^0 \subset H^0(\mathbf{R})$ which is not in general an equality.

2. THE REDUCTION COVARIANT

We will summarise the necessary properties of Cartan involutions of real reductive groups before proving the existence of the reduction covariant described in the introduction.

2.1. Background on Cartan involutions.

Definition 2.2. *Let \mathfrak{h} be a semisimple Lie algebra over \mathbf{R} . A Cartan involution of \mathfrak{h} is a Lie algebra involution $\theta : \mathfrak{h} \rightarrow \mathfrak{h}$ satisfying the following equivalent conditions:*

- (1) *The symmetric bilinear form $B_{\theta}(X, Y) = -B(X, \theta Y) : \mathfrak{h} \times \mathfrak{h} \rightarrow \mathbf{R}$ (where B is the Killing form) is positive definite.*
- (2) *The \mathbf{R} -vector subspace $\{X \in \mathfrak{h}_{\mathbf{C}} \mid \theta(X) = \overline{X}\}$ is a real form of $\mathfrak{h}_{\mathbf{C}}$ which is compact, in the sense that its Killing form is negative definite.*

It is convenient to define Cartan involutions for disconnected groups. We do this following [AT18] (see also [Mos55]).

Definition 2.3. *Let H be a linear algebraic group over \mathbf{R} such that H^0 is reductive. A Cartan involution of H is an involution $\theta : H \rightarrow H$ such that*

$$K_{\theta} = \{h \in H(\mathbf{C}) \mid \theta(h) = \overline{h}\}$$

is a compact subgroup of $H(\mathbf{C})$ that meets every connected component of $H(\mathbf{C})$.

If θ is a Cartan involution, then $H(\mathbf{R})^\theta = K_\theta \cap H(\mathbf{R})$ is a maximal compact subgroup of $H(\mathbf{R})$ ([AT18, §4]).

Given an automorphism θ of an algebraic group H , we will also write θ for the induced automorphism of its Lie algebra \mathfrak{h} . The following proposition is well-known.

Proposition 2.4. *Let H be a semisimple group over \mathbf{R} , and let θ be an involution of H . Then $\theta : H \rightarrow H$ is a Cartan involution if and only if $\theta : \mathfrak{h} \rightarrow \mathfrak{h}$ is a Cartan involution.*

Proposition 2.5. *Let H be a reductive group over \mathbf{R} .*

- (1) *Cartan involutions of H exist. If θ, θ' are Cartan involutions of H , then there exists $h \in H(\mathbf{R})^0$ such that $\theta' = \text{Ad}(h) \circ \theta \circ \text{Ad}(h^{-1})$.*
- (2) *Suppose $M \subset H$ is a closed subgroup and θ is a Cartan involution of H such that $\theta(M) = M$. Then M^0 is reductive and $\theta|_M$ is a Cartan involution of M .*
- (3) *Suppose $M \subset H$ is a closed subgroup with M^0 reductive and that θ_M is a Cartan involution of M . Then there exists a Cartan involution θ_H of H extending θ_M .*

Proof. See [AT18, Theorem 3.12]. □

If H is a reductive group over \mathbf{R} , then we write X_H for the set of Cartan involutions of H . Proposition 2.5(1) shows that this is a homogeneous space for $H(\mathbf{R})$.

2.6. Application to graded Lie algebras.

Theorem 2.7. *Let \mathfrak{h} be a semisimple Lie algebra over \mathbf{R} . Suppose given a stable $\mathbf{Z}/m\mathbf{Z}$ -grading $\mathfrak{h} = \bigoplus_{i \in \mathbf{Z}/m\mathbf{Z}} \mathfrak{h}_i$, and let (G, V) be the associated pair. Then:*

- (1) *For any stable vector $T \in V^s(\mathbf{R})$, there exists a unique Cartan involution θ_T of H such that $[T, \theta_T(T)] = 0$ and $\theta_T(\mathfrak{h}_i) \subset \mathfrak{h}_{-i}$ for each $i \in \mathbf{Z}/m\mathbf{Z}$.*
- (2) *The map $\mathcal{R} = \mathcal{R}_\mathfrak{h} : V^s(\mathbf{R}) \rightarrow X_G$ defined by $T \mapsto \theta_T|_G$ is $G(\mathbf{R})$ -equivariant.*

Proof. Giving an $\mathbf{Z}/m\mathbf{Z}$ -grading is equivalent to giving a homomorphism $\mu_m \rightarrow \text{Aut}(\mathfrak{h})$. Let $\sigma : \mathfrak{h}_\mathbf{C} \rightarrow \mathfrak{h}_\mathbf{C}$ be the image of $e^{2\pi i/m} \in \mu_m(\mathbf{C})$. The identity $\theta_T(\mathfrak{h}_i) \subset \mathfrak{h}_{-i}$ may be equivalently expressed as $\sigma\theta_T = \theta_T\bar{\sigma}$. Let H denote the adjoint group over \mathbf{R} with Lie algebra \mathfrak{h} ; then $\text{Aut}(H) = \text{Aut}(\mathfrak{h})$, and the embedding $\text{Ad} : H \rightarrow \text{Aut}(H)$ allows us to identify H with the connected component of the identity of the linear algebraic group $\text{Aut}(H)$. Let $\mathfrak{c} = \mathfrak{z}_\mathfrak{h}(T)$, a Cartan subalgebra of \mathfrak{h} since T is regular semisimple, and let $C = Z_H(T)$ denote the maximal torus of H with Lie algebra \mathfrak{c} .

Let us first establish uniqueness. If a Cartan involution θ_T with the given properties exists, then the identity $[T, \theta_T(T)] = 0$ shows that θ_T normalises \mathfrak{c} and C . The restriction of θ_T to C must be a Cartan involution of C . The torus C has a unique Cartan involution θ_C , so if θ'_T is another Cartan involution with the given properties, then we can write $\theta'_T = t\theta_T$ for some $t \in C(\mathbf{R})$.

The identities $\sigma\theta_T = \theta_T\bar{\sigma}$, $\sigma t\theta_T = t\theta_T\bar{\sigma}$ together imply that $t \in C^\sigma(\mathbf{R})$. This group is finite (because the grading is stable), hence compact, which implies that t, θ_T commute, and so $t^2 = 1$. We now consider the positive definite symmetric bilinear forms $B_{\theta_T}, B_{t\theta_T}$ on \mathfrak{h} . Computing

$$B_{t\theta_T}(X, Y) = -B(X, t\theta_T Y) = -B(t^{-1}X, \theta_T Y) = B_{\theta_T}(t^{-1}X, Y) = B_{\theta_T}(X, tY)$$

shows that (with respect to the inner product determined by B_{θ_T}) t is an orthogonal endomorphism of \mathfrak{h} with the property that $B_{\theta_T}(tX, X) > 0$ for every non-zero $X \in \mathfrak{h}$. Since $t^2 = 1$, this is only possible if $t = 1$, hence $\theta_T = \theta'_T$.

We now establish existence of θ_T . Let $M = C \rtimes \mu_m \subset \text{Aut}(H)$, and let θ_M be the Cartan involution of M given by the formula $\theta_M(c \rtimes \sigma) = \theta_C(c)\bar{\sigma}$. To show that this is a group homomorphism, we must show that the identity $\sigma\theta_C = \theta_C\bar{\sigma}$ holds in $\text{Aut}(C_{\mathbf{C}})$. Let $K_C \subset C(\mathbf{C})$ be the unique maximal compact subgroup. If $x \in K_C$, then $\theta_C(x) = \bar{x}$ and $\sigma(x) \in K_C$. For such x , we have $\sigma\theta_C(x) = \sigma(\bar{x})$ and $\theta_C\bar{\sigma}(x) = \bar{\sigma}(x) = \sigma(\bar{x}) = \sigma\theta_C(x)$. Since K_C is Zariski dense in $C_{\mathbf{C}}$, this shows that in fact $\sigma\theta_C = \theta_C\bar{\sigma}$.

By Proposition 2.5, we can extend θ_M to a Cartan involution θ of $\text{Aut}(H)$. We set $\theta_T = \theta|_H$. We claim that θ_T has the desired properties. It normalises C by construction, so satisfies $[T, \theta_T(T)] = 0$. We must show that $\sigma\theta_T = \theta_T\bar{\sigma}$ in $\text{Aut}(H)(\mathbf{C})$. To this end, we note that if $h \in H(\mathbf{C}) \subset \text{Aut}(H)(\mathbf{C})$, and $\alpha \in \text{Aut}(H)(\mathbf{C})$, then $\alpha h \alpha^{-1} = \alpha(h)$. (More transparently, we have $\alpha \circ \text{Ad}(h) \circ \alpha^{-1} = \text{Ad}(\alpha(h))$.) Since $\theta : \text{Aut}(H) \rightarrow \text{Aut}(H)$ is a group homomorphism, we have

$$\theta(\alpha h \alpha^{-1}) = \theta(\alpha)\theta(h)\theta(\alpha)^{-1}.$$

The right-hand side equals $\theta(\alpha)(\theta_T(h))$, while the left-hand side equals

$$\theta(\alpha(h)) = \theta_T(\alpha(h)).$$

Since h is arbitrary, this gives the identity $\theta(\alpha) \circ \theta_T = \theta_T \circ \alpha$ (composition in $\text{Aut}(H)(\mathbf{C})$). Taking $\alpha = \bar{\sigma}$ and using the identity $\theta(\bar{\sigma}) = \theta_M(\bar{\sigma}) = \sigma$, we find $\sigma\theta_T = \theta_T\bar{\sigma}$, as required. This completes the proof of the first part of the Theorem. The second part follows from the properties which uniquely characterize θ_T . \square

We now record some basic properties of the reduction covariant \mathcal{R} .

Proposition 2.8. (1) *Let $\mathfrak{h}, \mathfrak{h}'$ be semisimple Lie algebras over \mathbf{R} . Suppose given stable $\mathbf{Z}/m\mathbf{Z}$ -gradings*

$$\mathfrak{h} = \bigoplus_{i \in \mathbf{Z}/m\mathbf{Z}} \mathfrak{h}_i, \quad \mathfrak{h}' = \bigoplus_{i \in \mathbf{Z}/m\mathbf{Z}} \mathfrak{h}'_i.$$

Let $\mathfrak{h}'' = \mathfrak{h} \oplus \mathfrak{h}'$ with its induced $\mathbf{Z}/m\mathbf{Z}$ -grading. Then, with the obvious notation, we have $\mathcal{R}_{\mathfrak{h}''} = \mathcal{R}_{\mathfrak{h}} \times \mathcal{R}_{\mathfrak{h}'}$.

(2) *Let \mathfrak{h} be a semisimple Lie algebra over \mathbf{R} . Suppose given a stable $\mathbf{Z}/m\mathbf{Z}$ -grading $\mathfrak{h} = \bigoplus_{i \in \mathbf{Z}/m\mathbf{Z}} \mathfrak{h}_i$, and let (G, V) be the associated pair. Let $\mathfrak{h}' = \text{Res}_{\mathbf{C}/\mathbf{R}}(\mathfrak{h}_{\mathbf{C}})$, with its induced grading $\mathfrak{h}' = \bigoplus_{i \in \mathbf{Z}/m\mathbf{Z}} \text{Res}_{\mathbf{C}/\mathbf{R}}(\mathfrak{h}_{i, \mathbf{C}})$. Then this grading is stable and its associated pair may be identified as*

$$(G', V') = (\text{Res}_{\mathbf{C}/\mathbf{R}} G_{\mathbf{C}}, \text{Res}_{\mathbf{C}/\mathbf{R}} V_{\mathbf{C}}).$$

Moreover, there is a commutative diagram

$$\begin{array}{ccc} V^s(\mathbf{R}) & \longrightarrow & V^s(\mathbf{C}) = (V')^s(\mathbf{R}) \\ \mathcal{R}_{\mathfrak{h}} \downarrow & & \downarrow \mathcal{R}_{\mathfrak{h}'} \\ X_G & \longrightarrow & X_{G'} \end{array}$$

Proof. The first part follows quickly from the definition of \mathcal{R} . For the second, we note that the stability of the grading of \mathfrak{h}' can be checked over \mathbf{C} . There is an isomorphism $\mathfrak{h}'_{\mathbf{C}} \cong \mathfrak{h}_{\mathbf{C}} \oplus \mathfrak{h}_{\mathbf{C}}$, so the grading of \mathfrak{h}' is stable. It is easy to check that (G', V') has the claimed form.

and the associated representation is

$$V = \{T \in \mathfrak{sl}_{2g+1} \mid J^t T J = T\}.$$

The open subscheme $V^s \subset V$ consists of those operators whose characteristic polynomial has no repeated roots.

The following proposition describes the map $\mathcal{R} : V^s(\mathbf{R}) \rightarrow X_G$ in terms of linear algebra.

- Proposition 3.1.** (1) *We can identify X_G with the set of inner products H on \mathbf{R}^{2g+1} which are compatible with J , in the sense that the associated Gram matrix satisfies $JH^{-1}J = H$. The associated Cartan involution θ_H acts on \mathfrak{g} by $\theta_H(T) = -H^{-1}T H$.*
- (2) *If $T \in V^s(\mathbf{R})$, then there is a unique inner product $H = H_T$ compatible with J such that T is normal with respect to H (i.e. such that T commutes with its H -adjoint $H^t T H^{-1}$). We have $\mathcal{R}(T) = \theta_{H_T}$.*

Proof. Let Y_J denote the set of positive definite symmetric matrices H satisfying $JH^{-1}J = H$. (Observe that any such matrix satisfies $\det(H) = 1$.) If $H \in Y_J$, then $\theta(X) = -H^{-1}X H$ is a Cartan involution of \mathfrak{g} . (It preserves \mathfrak{g} because H, J are compatible, and B_θ is positive definite because H is an inner product.) Thus there is a map $Y_J \rightarrow X_H$ which is easily checked to be $G(\mathbf{R})$ -equivariant, where $x \in G(\mathbf{R})$ acts on H by $H \mapsto {}^t x^{-1} H x^{-1}$. We need to check that this map is bijective. To see that it is surjective, note that Y_J is non-empty, as it contains the identity matrix H_0 (otherwise said, the standard inner product on \mathbf{R}^{2g+1}), and $G(\mathbf{R})$ acts transitively on X_G . To see that is injective, it suffices (by $G(\mathbf{R})$ -equivariance) to show that if $\theta_H = \theta_{H_0}$, then $H = H_0$. However, if $\theta_H = \theta_{H_0}$ then $-H^{-1}X H = -{}^t X$ for all $X \in \mathfrak{g}$, so H is scalar (by Schur's lemma). Since H is positive definite and $\det(H) = 1$, this forces $H = H_0$.

Since the Lie bracket on \mathfrak{sl}_{2g+1} is given by $[X, Y] = XY - YX$, the second part of the proposition is asking for a unique H such that $[T, \theta_H(T)] = 0$. The existence and uniqueness therefore follows from Theorem 2.7. \square

To complete Proposition 3.1, we explain how to compute H explicitly in terms of T . Supposing that H exists, we extend it to a Hermitian inner product on \mathbf{C}^{2g+1} . Since the characteristic polynomial of T has no repeated roots, T is diagonalisable, and we can find $P \in \mathrm{GL}_{2g+1}(\mathbf{C})$ such that $P^{-1}TP$ is diagonal. Since T is normal with respect to H , its eigenvectors are orthogonal and so $D = {}^t P H \bar{P}$ is also diagonal, with positive real diagonal entries. Since T is also self-adjoint with respect to J , ${}^t P J P$ is also diagonal, and thus ${}^t P H \bar{P}, {}^t P J P$ commute. Starting with the identity $H = JH^{-1}J$, we get

$${}^t P H \bar{P} = {}^t P J H^{-1} J \bar{P} = {}^t P J P P^{-1} H^{-1} {}^t \bar{P}^{-1} {}^t \bar{P} J \bar{P} = {}^t P J P ({}^t \bar{P} H P)^{-1} {}^t \bar{P} J \bar{P},$$

hence

$$D^2 = {}^t P H \bar{P} {}^t \bar{P} H P = {}^t P J P {}^t \bar{P} J \bar{P}.$$

This characterizes H uniquely as

$$(3.1) \quad H = {}^t P^{-1} ({}^t P J P {}^t \bar{P} J \bar{P})^{1/2} \bar{P}^{-1}.$$

where $P \in \mathrm{GL}_{2g+1}(\mathbf{C})$ is such that $P^{-1}TP$ is diagonal.

4. A REDUCTION ALGORITHM FOR $\mathrm{SO}(J)$

In this section, we will describe a reduction algorithm for the symmetric space X_G associated to the group $G = \mathrm{SO}(J)$ introduced in §3. All of the action takes place on the group G : the ambient grading plays no role. To orient the reader, we note that we will have to consider three symmetric bilinear forms on \mathbf{R}^{2g+1} simultaneously. First, the form J defining the group G ; second, the standard inner product H_0 on \mathbf{R}^{2g+1} , which defines a base point in X_G ; and third, a second inner product H , compatible with J , which we hope to bring closer to H_0 using the action of the group $G(\mathbf{Z})$.

We continue to define

$$G = \mathrm{SO}(J) = \{x \in \mathrm{SL}_{2g+1} \mid {}^t x J x = J\},$$

which we think of as a group scheme over \mathbf{Z} . We write $A, N \subset G$ for the subgroups consisting of the diagonal and unipotent upper-triangular matrices, respectively. Then $G_{\mathbf{Q}}$ is reductive, and $A_{\mathbf{Q}}, A_{\mathbf{Q}}N_{\mathbf{Q}}$ are a maximal torus and Borel subgroup.

Let H_0 denote the inner product on \mathbf{R}^{2g+1} with respect to which e_{-g}, \dots, e_g is an orthonormal basis. We define $K \subset G(\mathbf{R})$ to be subgroup of matrices which preserve H_0 .

Proposition 4.1. (1) H_0 is compatible with J , and K is a maximal compact subgroup of $G(\mathbf{R})$.

(2) The product map $K \times A(\mathbf{R})^0 \times N(\mathbf{R}) \rightarrow G(\mathbf{R})$, $(k, a, n) \mapsto kan$, is a diffeomorphism.

(3) If H is an inner product which is compatible with J , then we can find a unique pair $(a, n) \in A(\mathbf{R})^0 \times N(\mathbf{R})$ such that $H = {}^t(an)H_0an$.

Proof. Looking at Proposition 3.1 we see that H_0 is compatible with J and $\theta_0(x) = H_0^{-1} {}^t x^{-1} H_0 = {}^t x^{-1}$ is a Cartan involution of $G_{\mathbf{R}}$. Therefore $K = G(\mathbf{R})^{\theta_0}$ is a maximal compact subgroup of $G(\mathbf{R})$. The second part of the proposition is a statement of the Iwasawa decomposition of $G(\mathbf{R})$, determined by the data of θ_0 , A , and N [Kna02, Theorem 7.31]. The third part of the proposition follows from the uniqueness of the Iwasawa decomposition and the fact that $G(\mathbf{R})$ acts transitively on X_G with $\mathrm{Stab}_{G(\mathbf{R})}(\theta_0) = K$. \square

The components of the Iwasawa decomposition can be computed using the Gram–Schmidt process. Let e_{-g}^*, \dots, e_g^* denote the result of carrying out the Gram–Schmidt orthogonalization process on the basis e_{-g}, \dots, e_g with respect to the inner product defined by H . Thus we have formulae

$$e_{-g} = e_{-g}^*, \dots, e_j = e_j^* + \sum_{i=-g}^{j-1} \mu_{i,j} e_i^*, \dots, e_g = e_g^* + \sum_{i=-g}^{g-1} \mu_{i,g} e_i^*.$$

with

$$\mu_{i,j} = (e_j, e_i^*)_H / (e_i^*, e_i^*)_H.$$

In particular, we take $\mu_{j,j} = 1$ and $\mu_{i,j} = 0$ if $i > j$.

Lemma 4.2. Let $n = n_H = (\mu_{i,j})_{-g \leq i, j \leq g}$ and $a = a_H = \mathrm{diag}(\|e_i^*\|_H)_{-g \leq i \leq g}$. Then $a \in A(\mathbf{R})^0$, $n \in N(\mathbf{R})$, and $H = {}^t(an)an$.

Proof. Let θ_1 denote the Cartan involution $\theta_1 : \mathrm{SL}_{2g+1} \rightarrow \mathrm{SL}_{2g+1}$, $g \mapsto {}^t g^{-1}$. Let $K_1 = \mathrm{SL}_{2g+1}(\mathbf{R})^{\theta_1}$ and let A_1, N_1 be the subgroups of diagonal and unipotent

upper-triangular matrices in SL_{2g+1} , respectively. The Iwasawa decomposition for SL_{2g+1} is the statement that the product map

$$K_1 \times A_1(\mathbf{R})^0 \times N_1(\mathbf{R}) \rightarrow \mathrm{SL}_{2g+1}(\mathbf{R})$$

is a diffeomorphism. It is a standard fact that the Gram–Schmidt process gives the Iwasawa decomposition for SL_{2g+1} , in the sense that $H = {}^t(an)an$ and that these are the unique elements $a \in A_1(\mathbf{R})^0$, $n \in N_1(\mathbf{R})$ with this property. We need to explain why the assumption that H, J are compatible implies that in fact a, n lie in $G(\mathbf{R})$. However, we have $K \leq K_1$, $A \leq A_1$, and $N \leq N_1$, by construction, so the existence of the Iwasawa decomposition for G and the uniqueness for SL_{2g+1} implies that the Gram–Schmidt process for SL_{2g+1} must be compatible with the Iwasawa decomposition for G . \square

Lemma 4.3. *For any $n = (n_{ij}) \in N(\mathbf{R})$, there exists $m \in N(\mathbf{Z})$ satisfying the following conditions:*

- (1) $|(nm)_{ij}| \leq 1/2$ for all $-g \leq i < j$, $j = -g, \dots, -1$.
- (2) $|(nm)_{i,0}| \leq 1$ for all $-g \leq i \leq -1$.
- (3) $|(nm)_{i,j}| \leq 1/2$ for all $-g \leq i < -j$, $j = 1, \dots, g$.

Proof. Right multiplication by m corresponds to performing column operations on n . We can check that the following column operations on n are induced by elements of $N(\mathbf{Z})$:

- $A(i, j, q)$: For $q \in \mathbf{Z}$, $-g \leq j \leq -1$, and $-g \leq i < j$, add q times column i to column j and $-q$ times column $-j$ to column $-i$.
- $B(i, q)$: $q \in 2\mathbf{Z}$, $-g \leq i \leq -1$, add q times column i to column 0, and add $-q$ times column 0 and $-q^2/2$ times column i to column $-i$.
- $C(i, j, q)$: $q \in \mathbf{Z}$, $1 \leq j \leq g$, and $-g \leq i \leq -j$, add q times column i to column j and $-q$ times column $-j$ to column $-i$.

We therefore carry out column operations as follows in order to satisfy the conditions of the lemma (noting that the order of operations is chosen so that once we have forced a given n_{ij} to satisfy the conditions of the lemma, its value will not be changed by later operations):

- (1) For each $j = -g, \dots, -1$, then for each $i = j-1, j-2, \dots, -g$, let q denote the closest integer to n_{ij} and do $A(i, j, q)$.
- (2) For each $i = -1, \dots, -g$, let q denote the closest even integer to n_{i0} and do $B(i, q)$.
- (3) For each $j = 1, \dots, g$, then for each $i = -j-1, \dots, -g$, let q denote the closest integer to n_{ij} and do $C(i, j, q)$.

\square

The following definition is the analogue in our context of [LLL82, (1.4), (1.5)].

Definition 4.4. *Let $\delta \in (1/2, 1)$, and let H be an inner product on \mathbf{R}^{2g+1} compatible with J . We say that H is δ -reduced if the following conditions are satisfied:*

- (1) n_H satisfies the conditions of Lemma 4.3.
- (2) For each $i = -g+1, \dots, -1$, we have $\|e_i^* + \mu_{i,i-1}e_{i-1}^*\|_H^2 \geq \delta \|e_{i-1}^*\|_H^2$.
- (3) $\|e_1^* + \mu_{1,0}e_0^* + \mu_{1,-1}e_{-1}^*\|_H^2 \geq \delta^2 \|e_{-1}^*\|_H^2$.

Remark 4.5. The above conditions may be reformulated in terms of the matrices n_H, a_H as follows:

- (1) n_H satisfies the conditions of Lemma 4.3.
- (2) For each $i = -g + 1, \dots, -1$, we have $a_{i,i}^2 + \mu_{i,i-1}^2 a_{i-1,i-1}^2 \geq \delta a_{i-1,i-1}^2$.
- (3) $a_{-1,-1}^{-2} + \frac{1}{2} \mu_{0,-1}^2 \geq \delta$.

The following lemma is evidence that this is a useful notion.

Lemma 4.6. *If H is δ -reduced, then $a_H n_H$ lies in the Siegel set $\mathfrak{S}_\delta = KA_\delta N_c$ defined as follows:*

$$A_\delta = \{a = (a_{-g}, \dots, a_g) \in A(\mathbf{R})^0 \mid \\ \forall i = -g, \dots, -2, a_i/a_{i+1} \leq (\delta - 1/4)^{-1/2}; \\ a_{-1} \leq (\delta - 1/2)^{-1/2}\},$$

$$N_c = \{n \in N(\mathbf{R}) \mid \\ \forall j = -g, \dots, -1, -g \leq i < j, \max(|n_{ij}|, |n_{i,-j}|) \leq 1/2; \\ \forall i = -g, \dots, -1, |n_{i0}| \leq 1\}.$$

Proof. Since n_H lies in N_c by definition of being δ -reduced, we just need to check that a_H satisfies the required inequalities. Re-arranging gives

$$a_{i+1}^2/a_i^2 \geq \delta - \mu_{i+1,i}^2 \geq \delta - 1/4$$

for each $i = -g, \dots, -2$, and

$$a_1^2 \geq \delta - \mu_{0,-1}^2/2 \geq \delta - 1/2,$$

as required. \square

The Weyl group $W(G, A)$ is generated by its simple reflections (with respect to the choice of positive roots determined by N). These simple reflections can be represented by the following elements s_{-g}, \dots, s_{-1} of $G(\mathbf{Z})$, which we define by their action on basis vectors:

- For each $i = -g, \dots, -2$, s_i sends e_i to e_{i+1} , e_{i+1} to e_i , e_{-i} to $e_{-(i+1)}$, $e_{-(i+1)}$ to e_{-i} , and fixes the remaining basis vectors.
- s_{-1} sends e_{-1} to e_1 , e_1 to e_{-1} , e_0 to $-e_0$, and fixes the remaining basis vectors.

Here finally is an algorithm which, starting with any H as above, finds $\gamma \in G(\mathbf{Z})$ such that ${}^t\gamma H \gamma$ is δ -reduced:

- (1) Let $\gamma_1 = 1$, $H_1 = H$.
- (2) Set $H_1 := {}^t\gamma_1 H \gamma_1$. Compute the matrices n_{H_1}, a_{H_1} using Gram-Schmidt. If H_1 is δ -reduced, return γ_1 . Else, proceed to Step 3.
- (3) Find γ_n such that $n_{H_1} \gamma_n$ satisfies the conclusion of Lemma 4.3.
- (4) For each $i = -g, \dots, -2$: if $a_{i+1,i+1}^2 + \mu_{i+1,i}^2 a_{i,i}^2 < \delta a_{i,i}^2$, set $\gamma_1 := \gamma_1 \gamma_n s_i$ and go to Step 2.
- (5) If $a_{-1,-1}^{-2} + \frac{1}{2} \mu_{0,-1}^2 < \delta$, set $\gamma_1 := \gamma_1 \gamma_n s_{-1}$ and go to Step 2.
- (6) Set $\gamma_1 := \gamma_1 \gamma_n$ and return γ_1 .

Proposition 4.7. *The above algorithm always terminates.*

Proof. It suffices to show that the simple reflections s_i are applied only finitely many times. At each instance of Step 2 we are given a basis $f_i = \gamma_1 e_i$ of \mathbf{Z}^{2g+1} , to which we associate the flag $F_i = \bigoplus_{k=-g}^i \mathbf{Z} f_k$, $i = -g, \dots, g$. If $i = -g, \dots, -2$, then

acting by s_i changes F_i and F_{-i} but leaves the remaining F_j unchanged. Acting by s_{-1} changes F_{-1} and F_0 and leaves the remaining F_j unchanged. In particular, among F_{-g}, \dots, F_{-1} , s_i changes only F_i .

The inner product H on \mathbf{R}^{2g+1} determines one on $\wedge^i \mathbf{R}^{2g+1}$ for each $i = 1, \dots, 2g+1$ by the usual formula $(x_1 \wedge \dots \wedge x_i, y_1 \wedge \dots \wedge y_i)_H = \det((x_j, y_k)_H)$. Since $\wedge^i \mathbf{Z}^{2g+1}$ contains finitely many vectors of bounded norm, it suffices to show that applying the simple reflection s_i decreases the norm of the vector (which, up to sign, depends only on F_i) $f_{-g} \wedge \dots \wedge f_i \in \wedge^{i+g+1} \mathbf{Z}^{2g+1}$. A calculation shows that this norm is multiplied by a scalar which is strictly less than δ . As $\delta < 1$, this completes the proof. \square

5. EXAMPLE: REDUCING A SELF-ADJOINT LINEAR OPERATOR

We now show how to combine the theory of the previous two sections in an example. Take $g = 3$, and consider the group $G = \mathrm{SO}(J) \subset \mathrm{SL}_7$ and representation $V \subset \mathfrak{sl}_7$ associated to the $\mathbf{Z}/2\mathbf{Z}$ -grading of \mathfrak{sl}_7 defined in §3. We consider the genus 3 hyperelliptic curve

$$C_f : y^2 = f(x) = x^7 - x^5 - 2x^4 - x^3 + 5x - 5$$

This curve has the integral point $P = (14, 10237)$. Associated to P is the matrix

$$T = \begin{pmatrix} -14 & 1 & 0 & 0 & 0 & 0 & 0 \\ -195 & 0 & 1 & 0 & 0 & 0 & 0 \\ -2728 & 0 & 7 & 0 & -1 & 0 & 0 \\ -10237 & 0 & 0 & 14 & 0 & 0 & 0 \\ 19095 & -6 & -48 & 0 & 7 & 1 & 0 \\ 1546 & -26 & -6 & 0 & 0 & 0 & 1 \\ 390 & 1546 & 19095 & -10237 & -2728 & -195 & -14 \end{pmatrix} \in V(\mathbf{Z})$$

of characteristic polynomial $f(x)$, which we obtain e.g. from the subregular Slodowy slice studied in [Tho13]. We compute the reduction covariant $H = \mathcal{R}(T) \in X_{\mathrm{SO}(J)}$ using the formula (3.1). We obtain numerically

$$H = \begin{pmatrix} 3.74708 & 53.7691 & 750.242 & 2813.43 & -5244.78 & -421.526 & -47.2448 \\ 53.7691 & 776.143 & 10830.1 & 40612.6 & -75708.6 & -6080.03 & -681.676 \\ 750.242 & 10830.1 & 151130. & 566729. & -1.05648 \times 10^6 & -84842.6 & -9520.71 \\ 2813.43 & 40612.6 & 566729. & 2.12521 \times 10^6 & -3.96175 \times 10^6 & -318157. & -35704.6 \\ -5244.78 & -75708.6 & -1.05648 \times 10^6 & -3.96175 \times 10^6 & 7.38537 \times 10^6 & 593097. & 66564.2 \\ -421.526 & -6080.03 & -84842.6 & -318157. & 593097. & 47660.8 & 5338.34 \\ -47.2448 & -681.676 & -9520.71 & -35704.6 & 66564.2 & 5338.34 & 660.273 \end{pmatrix}.$$

We then apply the algorithm of §4 to H with $\delta = 0.9$ to obtain an element $\gamma_1 \in G(\mathbf{Z})$ such that ${}^t\gamma_1 H \gamma_1$ is δ -reduced. The resulting matrix is

$$\gamma_1 = \begin{pmatrix} -2 & -1 & -15 & 2 & -3 & -5 & 26 \\ 0 & -8 & -117 & 12 & 0 & 9 & 203 \\ -8 & -104 & -1462 & 180 & -16 & 184 & 2557 \\ 4 & 56 & 784 & -97 & 8 & -98 & -1372 \\ 1 & 15 & 209 & -26 & 2 & -26 & -366 \\ 0 & 1 & 15 & -2 & 0 & -2 & -26 \\ 0 & 0 & 1 & 0 & 0 & 0 & -2 \end{pmatrix},$$

yielding

$$\gamma_1^{-1}T\gamma_1 = \begin{pmatrix} 0 & 0 & -1 & 2 & 2 & -2 & 3 \\ 1 & 0 & 1 & 0 & 0 & 0 & -2 \\ 0 & 1 & 1 & -2 & 0 & 0 & 2 \\ 0 & 0 & 1 & -2 & -2 & 0 & 2 \\ 0 & 0 & 0 & 1 & 1 & 1 & -1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}.$$

Since the lower left-hand 3×3 submatrix is 0, we see that the orbit of T is distinguished, in the sense of [BG13]. It follows that the divisor class of $P - P_\infty$ is divisible by 2 in $J_f(\mathbf{Q})$ (where P_∞ is the point at infinity of C_f and J_f is the Jacobian of C_f).

REFERENCES

- [AT18] Jeffrey Adams and Olivier Taïbi. Galois and Cartan cohomology of real groups. *Duke Math. J.*, 167(6):1057–1097, 2018.
- [BG13] Manjul Bhargava and Benedict H. Gross. The average size of the 2-Selmer group of Jacobians of hyperelliptic curves having a rational Weierstrass point. In *Automorphic representations and L-functions*, volume 22 of *Tata Inst. Fundam. Res. Stud. Math.*, pages 23–91. Tata Inst. Fund. Res., Mumbai, 2013.
- [BHC62] Armand Borel and Harish-Chandra. Arithmetic subgroups of algebraic groups. *Ann. of Math. (2)*, 75:485–535, 1962.
- [Bor19] Armand Borel. *Introduction to arithmetic groups*, volume 73 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2019. Translated from the 1969 French original by Lam Laurent Pham, Edited and with a preface by Dave Witte Morris.
- [CFS10] John E. Cremona, Tom A. Fisher, and Michael Stoll. Minimisation and reduction of 2-, 3- and 4-coverings of elliptic curves. *Algebra Number Theory*, 4(6):763–820, 2010.
- [Fis13] Tom Fisher. Minimisation and reduction of 5-coverings of elliptic curves. *Algebra Number Theory*, 7(5):1179–1205, 2013.
- [Kna02] Anthony W. Knaapp. *Lie groups beyond an introduction*, volume 140 of *Progress in Mathematics*. Birkhäuser Boston, Inc., Boston, MA, second edition, 2002.
- [Lag22] Jef Laga. The average size of the 2-Selmer group of a family of non-hyperelliptic curves of genus 3. *Algebra Number Theory*, 16(5):1161–1212, 2022.
- [LLL82] A. K. Lenstra, H. W. Lenstra, Jr., and L. Lovász. Factoring polynomials with rational coefficients. *Math. Ann.*, 261(4):515–534, 1982.
- [Mos55] G. D. Mostow. Self-adjoint groups. *Ann. of Math. (2)*, 62:44–55, 1955.
- [RLYG12] Mark Reeder, Paul Levy, Jiu-Kang Yu, and Benedict H. Gross. Gradings of positive rank on simple Lie algebras. *Transform. Groups*, 17(4):1123–1190, 2012.
- [RT24] Beth Romano and Jack A. Thorne. An LLL algorithm with symmetries. preprint, 2024.
- [SC03] Michael Stoll and John E. Cremona. On the reduction theory of binary forms. *J. Reine Angew. Math.*, 565:79–99, 2003.
- [Sha19] Ananth N. Shankar. 2-Selmer groups of hyperelliptic curves with marked points. *Trans. Amer. Math. Soc.*, 372(1):267–304, 2019.
- [Tho13] Jack A. Thorne. Vinberg’s representations and arithmetic invariant theory. *Algebra Number Theory*, 7(9):2331–2368, 2013.
- [Tho15] Jack A. Thorne. E_6 and the arithmetic of a family of non-hyperelliptic curves of genus 3. *Forum Math. Pi*, 3:e1, 41, 2015.
- [Vin76] È. B. Vinberg. The Weyl group of a graded Lie algebra. *Izv. Akad. Nauk SSSR Ser. Mat.*, 40(3):488–526, 709, 1976.