DISTANCE INEQUALITIES OF GENERALIZED *k*-INTERSECTION BODIES

JULIÁN HADDAD

ABSTRACT. It was shown in [11] that for every origin-symmetric star body $K \subseteq \mathbb{R}^n$ of volume 1, every even continuous probability density f on K and $1 \leq k \leq n-1$, there exists a subspace $F \subseteq \mathbb{R}^n$ of codimension k such that

k

$$\int_{K\cap F} f \ge c^k (d_{\text{ovr}}(K, \mathcal{BP}^n_k))^{-1}$$

where $d_{\text{ovr}}(K, \mathcal{BP}_k^n)$ is the outer volume ratio distance from K to the class of generalized k-intersection bodies, and c > 0 is a universal constant. The upper bound $d_{\text{ovr}}(K, \mathcal{BP}_k^n) \leq c' \sqrt{n/k} \left(\log \left(\frac{en}{k}\right) \right)^{3/2}$ was established in [13] for every convex body K. In this note we show that there exist an origin-symmetric convex body K of volume 1 and an even continuous probability density f supported on K such that for every subspace F of codimension k,

$$\int_{K \cap F} f \le \left(c \sqrt{\frac{n}{k \log(n)}} \right)^{-}$$

As a consequence we obtain a lower bound for $d_{ovr}(K, \mathcal{BP}_k^n)$ with K a convex body, complementing the upper bound in [13]. This is

$$c\sqrt{n/k}(\log(n))^{-1/2} \le \sup_{K} d_{\text{ovr}}(K, \mathcal{BP}_{k}^{n}) \le c'\sqrt{n/k}\left(\log\left(\frac{en}{k}\right)\right)^{3/2}$$

The case k = 1 was obtained previously in [5, 6].

1. INTRODUCTION

The problem known as the *slicing inequality for general measures* asks which is the greatest possible value S_n such that for every origin-symmetric convex body K(convex compact with non-empty interior) and every even continuous probability density f on K, there exists a hyperplane $F \subseteq \mathbb{R}^n$ with

$$\int_{K\cap F} f \ge S_n$$

Koldobsky investigated this problem in [9, 10, 11] and showed that $S_n \geq \frac{1}{2\sqrt{n}}$, although the constant can improve if K is restricted to certain classes of convex bodies.

A star body is a set $K \subseteq \mathbb{R}^n$ of the form

$$K = \{0\} \cup \{x \in \mathbb{R}^n \setminus \{0\} : |x| \le \rho_K(x/|x|)\}$$

where the radial function $\rho_K : S^{n-1} \to (0, \infty)$ is continuous and strictly positive, and $|\cdot|$ is the euclidean norm. Every convex body containing the origin in the

Date: June 3, 2024.

²⁰¹⁰ Mathematics Subject Classification. 52A20, 44A12.

 $Key\ words\ and\ phrases.$ Radon transform, slicing inequality, k-intersection bodies, outer volume ratio distance.

J. HADDAD

interior is also a star body. The radial distance between two star bodies K, L is $\max_{v \in S^{n-1}} |\rho_K(v) - \rho_L(v)|$.

Given a star body $K \subseteq \mathbb{R}^n$, the *intersection body* of K is the star body IK defined by $\rho_{IK}(v) = |K \cap v^{\perp}|_{n-1}$, where $|\cdot|_k$ denotes the k-dimensional volume. Taking the closure in the radial metric of the sets IK with $K \subseteq \mathbb{R}^n$ a star body, we get the class of *intersection bodies*, which we denote by \mathcal{I}_n .

For any class of star bodies C the outer volume ratio distance from a star body K to C is defined by

$$d_{\rm ovr}(K,\mathcal{C}) = \inf\left\{ \left(\frac{|D|_n}{|K|_n}\right)^{1/n} : K \subseteq D, D \in \mathcal{C} \right\}.$$

In [11] the following inequality was proven:

Theorem 1.1. Let K be an origin-symmetric star body in \mathbb{R}^n . Then for any even continuous non-negative function f we have

$$\int_{K} f \leq 2d_{\operatorname{ovr}}(K, \mathcal{I}_{n}) |K|_{n}^{1/n} \max_{v \in S^{n-1}} \int_{K \cap v^{\perp}} f.$$

In particular, if K is an origin-symmetric star body in \mathbb{R}^n of volume 1 and f is an even continuous probability density in K then

$$\max_{\nu \in S^{n-1}} \int_{K \cap \nu^{\perp}} f \ge (2d_{\text{ovr}}(K, \mathcal{I}_n))^{-1}$$

Here $d_{\text{ovr}}(K, \mathcal{I}_n)$ is the outer volume ratio distance between K and the class of intersection bodies. Since origin-symmetric ellipsoids are intersection bodies, if K is an origin-symmetric convex body, John's theorem implies that $d_{\text{ovr}}(K, \mathcal{I}_n) \leq \sqrt{n}$, hence the bound $S_n \geq \frac{1}{2\sqrt{n}}$.

The upper bound is $S_n \leq c/\sqrt{n}$, in other words, there exists an origin-symmetric convex body K of volume 1 and an even continuous probability density f on K, such that

$$\max_{v \in S^{n-1}} \int_{K \cap v^{\perp}} f \le c \frac{1}{\sqrt{n}}.$$

This estimate was obtained in [5] up to a factor $\sqrt{\log(\log(n))}$, and this factor was later removed in [6]. It follows from Theorem 1.1 that the largest possible distance from a convex body to \mathcal{I}_n has order $c\sqrt{n}$, this is,

$$c\sqrt{n} \le \sup_{K} d_{\text{ovr}}(K, \mathcal{I}_n) \le \sqrt{n}$$

where the supremum runs over all origin-symmetric convex bodies $K \subseteq \mathbb{R}^n$, and c > 0 is a universal constant. A similar argument was used to show in [1] (lower bound) and [12] (upper bound) that

$$c\sqrt{n/p} \le \sup_{K} d_{\text{ovr}}(K, L_p^n) \le c'\sqrt{n/p}$$

for $p \geq 1$, where L_p is the class of convex bodies L such that $(\mathbb{R}^n, \|\cdot\|_L)$ embeds in L_p as a Banach space (see [8] for details). Also, in [4] the convex body and measure constructed in [5] were adapted to a different slicing inequality from [3] to obtain the lower bound

(1)
$$\sup_{K} d_{\text{ovr}}(K, L^{n}_{-p}) \ge cn^{\frac{1}{2p}}$$

for $p \in [1, n)$, where the supremum runs over all origin-symmetric convex bodies $K \subseteq \mathbb{R}^n, c > 0$ is a universal constant and L_{-p}^n is the class of star bodies L for which $(\mathbb{R}^n, \|\cdot\|_L)$ embeds in L_{-p} . For integer $k \in [1, n), L_{-k}$ is the class of k-intersection bodies (see [8] for details).

A version of Theorem 1.1 with hyperplanes replaced by lower dimensional subspaces (Theorem 1.2 below) was established also in [11]. Denote by $G_{n,n-k}$ the Grassmanian of subspaces of \mathbb{R}^n of codimension k. The class of convex bodies responsible for the k-codimensional problem is the class of generalized k-intersection bodies, denoted by \mathcal{BP}_k^n , which can be defined as follows: Given two star bodies K, L and p > 0, their radial p-sum is the star body $K +_p L$ whose radial function is

$$\rho^p_{K\tilde{+}_pL} = \rho^p_K + \rho^p_L$$

For $k \in [1, n]$, the class \mathcal{BP}_k^n is defined as the closure in the radial metric of finite radial k-sums of origin-symmetric ellipsoids.

Theorem 1.2 ([11, Corollary 1]). Let K be an origin-symmetric star body in \mathbb{R}^n . Then for any even continuous non-negative function f,

$$\int_{K} f \le c^{k} d_{\text{ovr}}(K, \mathcal{BP}_{k}^{n})^{k} |K|_{n}^{k/n} \max_{F \in G_{n,n-k}} \int_{K \cap F} f_{F}$$

where c > 0 is a universal constant. In particular, if K is an origin-symmetric star body in \mathbb{R}^n of volume 1 and f is an even continuous probability density in K then

$$\max_{F \in G_{n,n-k}} \int_{K \cap F} f \ge c^{-k} (d_{\text{ovr}}(K, \mathcal{BP}_k^n))^{-k}.$$

An upper bound for the distance to \mathcal{BP}_k^n was obtained in [13] (the power of the logarithmic term was later corrected in [11, Proposition 1]).

Theorem 1.3. Let K be an origin-symmetric convex body in \mathbb{R}^n , and let $1 \le k \le n-1$. Then

(2)
$$d_{\rm ovr}(K, \mathcal{BP}_k^n) \le c\sqrt{n/k} \left(\log\left(\frac{en}{k}\right)\right)^{3/2}$$

where c > 0 is a universal constant.

Moreover, since $\mathcal{BP}_k^n \subseteq L_{-k}^n$ (this was proven in [7] and [15], see also [8, Theorem 4.23]), inequalities (1) and (2) imply

$$cn^{\frac{1}{2k}} \leq \sup_{K} d_{\text{ovr}}(K, L^{n}_{-k}) \leq \sup_{K} d_{\text{ovr}}(K, \mathcal{BP}^{n}_{k}) \leq c\sqrt{n/k} \left(\log\left(\frac{en}{k}\right)\right)^{3/2}$$

The two estimates are tight for large values of k (at least proportional to n). For low values of k > 1 the lower and upper bounds are far apart as $n \to \infty$. In this paper we use the same technique as in [5] to prove that Theorem 1.2 is sharp up to a logarithmic term, thus obtaining the correct (up to logarithms) order of $\sup_{k} d_{ovr}(K, \mathcal{BP}_{k}^{n})$. Our main theorem is the following:

Theorem 1.4. There exists a universal constant c > 0 such that for every $n \ge 2$ and $1 \le k \le n - 1$, there exists an origin-symmetric convex body $K \subseteq \mathbb{R}^n$ and an even continuous probability density f in K such that for every subspace $F \subseteq \mathbb{R}^n$ of codimension k,

$$\int_F f \le \left(c \sqrt{\frac{\log(n)k}{n}} \right)^k$$

J. HADDAD

As a consequence of Theorems 1.4 and 1.2 we deduce that the estimate (2) is also asymptotically sharp, up to logarithmic terms.

Corollary 1.5. There exists a universal constant c > 0 such that for every $n \ge 2$ and $1 \le k \le n-1$, there exists an origin-symmetric convex body $K \subseteq \mathbb{R}^n$ such that

$$d_{\text{ovr}}(K, \mathcal{BP}_k^n) \ge c\sqrt{n/k}(\log(n))^{-1/2}.$$

Combining Corollary 1.5 and Theorem 1.3 we obtain

$$c\sqrt{n/k}(\log(n))^{-1/2} \le \sup_{K} d_{\text{ovr}}(K, \mathcal{BP}_{k}^{n}) \le c'\sqrt{n/k}\left(\log\left(\frac{en}{k}\right)\right)^{3/2}$$

where the supremum runs over all origin-symmetric convex bodies $K \subseteq \mathbb{R}^n$, and c, c' > 0 are universal constants.

It should be noted that our proof is a simplification of that used in [5] and [6] since we do not intend to suppress any logarithmic term. We thank prof. A. Koldobsky for suggesting this approach, and for many other useful discussions related to this article.

The author was supported by Grant RYC2021-032768-I, funded by the Ministry of Science and Innovation/State Research Agency/10.13039 /501100011033 and by the E.U. Next Generation EU/Recovery, Transformation and Resilience Plan.

2. Proof of Theorem 1.4

Theorem 1.4 shall be proven at the end of the section after some auxiliary results. Throughout the section c, c', c'' will denote universal constants that may change from one line to the next one. We denote $\varphi(t) = e^{-t^2}$, the euclidean unit ball of \mathbb{R}^n is denoted by \mathbb{B}_n , the canonical vectors of \mathbb{R}^n are e_1, \ldots, e_n and Γ is the gamma function. For $F \in G_{n,n-k}$ and $x \in \mathbb{R}^n$, the euclidean distance from x to F is denoted by d(F, x).

First we recall two elementary inequalities from [4] and [5]:

Lemma 2.1 ([4, Lemma 2.4]). For $0 \le \mu \le \lambda$ we have

$$\lambda^{\mu} \Gamma(\lambda - \mu) \ge \Gamma(\lambda).$$

Lemma 2.2 ([5, Lemma 3.1]). Let Y_1, \ldots, Y_N be independent, identically-distributed random variables attaining values in the interval [0, 1]. Let $p \in [0, 1]$ satisfy $p \ge \mathbb{E}Y_i$. Then,

$$\mathbb{P}\left(\frac{1}{N}\sum_{i=1}^{N}Y_i \ge 3p\right) \le e^{-pN}.$$

The next elementary lemma concerns the parametrization of the sphere with respect to the so-called *bispherical coordinates*. It can be found in [14, formula (3.9)].

Lemma 2.3. Consider the decomposition $\mathbb{R}^n = \mathbb{R}^m \times \mathbb{R}^k$ with m + k = n and $m, k \in [1, n]$. If $f: S^{n-1} \to \mathbb{R}$ is a measurable function, then

$$\int_{S^{n-1}} f(v)dv = \int_{S^{m-1}} \int_{S^{k-1}} \int_0^{\pi/2} f(\cos(\alpha)x, \sin(\alpha)y)\cos(\alpha)^{m-1}\sin(\alpha)^{k-1}d\alpha dy dx$$

where integration on the sphere S^l is with respect to the *l*-dimensional Hausdorff measure. The cases k = 1 and m = 1 are also included, the measure in $S^0 = \{\pm 1\}$ being the counting measure.

Lemma 2.4. Let $n \ge 1$ and θ be a random point uniformly distributed in S^{n-1} . Let F be any subspace of codimension k with $1 \le k \le n$, then

$$\mathbb{E}(\varphi(d(F, n\theta))) \le n^{-k/2}.$$

(Notice that there is no constant in the right-hand side.)

Proof. By rotational invariance, we may assume without loss of generality that F is generated by e_{k+1}, \ldots, e_n . We decompose $\mathbb{R}^n = \mathbb{R}^k \times \mathbb{R}^{n-k}$ and notice that d(F, (x, y)) = |x|.

To find the expectation we integrate, use Lemma 2.3 and then the change of variables $t = \cos(\alpha)$, $dt = -\sqrt{1-t^2}d\alpha$.

$$\mathbb{E}(\varphi(d(F, n\theta)))$$

$$= \frac{1}{\kappa_n} \int_{S^{n-1}} \varphi(nd(F, \theta)) d\theta$$

$$= \frac{1}{\kappa_n} \int_{S^{k-1}} \int_{S^{n-k-1}} \int_0^{\pi/2} \varphi(n\cos(\alpha)) \cos(\alpha)^{k-1} \sin(\alpha)^{n-k-1} d\alpha dx dy$$

$$= \frac{\kappa_k \kappa_{n-k}}{\kappa_n} \int_0^{\pi/2} \varphi(n\cos(\alpha)) \cos(\alpha)^{k-1} \sin(\alpha)^{n-k-1} d\alpha$$

$$= \frac{\kappa_k \kappa_{n-k}}{\kappa_n} \int_0^1 \varphi(nt) t^{k-1} (1-t^2)^{\frac{n-k-2}{2}} dt,$$

where $\kappa_m = \frac{2}{\Gamma(m/2)} \pi^{m/2}$ is the m-1 dimensional volume of S^{m-1} . To estimate the constant, we use Lemma 2.1.

(3)
$$\frac{\kappa_k \kappa_{n-k}}{\kappa_n} = \frac{2\Gamma(n/2)}{\Gamma(k/2)\Gamma\left(\frac{n-k}{2}\right)} \le 2\Gamma(k/2)^{-1}\left(\frac{n}{2}\right)^{k/2}$$

For the integral we use a change of variables s = nt.

(4)
$$\int_{0}^{1} \varphi(nt) t^{k-1} (1-t^{2})^{\frac{n-k-2}{2}} dt = n^{-k} \int_{0}^{n} \varphi(s) s^{k-1} \left(1 - \left(\frac{s}{n}\right)^{2}\right)^{\frac{n-k-2}{2}} ds$$
$$\leq n^{-k} \int_{0}^{n} \varphi(s) s^{k-1} ds$$
$$= n^{-k} 2^{k/2-1} \Gamma(k/2)$$

Combining (3) and (4) we get the result.

The Grassmanian $G_{n,n-k}$ of k-codimensional subspaces of \mathbb{R}^n can be constructed as the quotient space of the group of rotations O_n , by the subgroup of rotations that fix a given k-codimensional subspace. As such, $G_{n,n-k}$ inherits naturally a metric d, which is the quotient of the operator-norm metric. This is defined as

$$d_G(F_1, F_2) = \min\{\|I - T\|_{\text{op}} : T \in O_n, T(F_1) = F_2\}$$

for $F_1, F_2 \in G_{n,n-k}$.

First we need the following simple lemma.

Lemma 2.5. Let $\theta \in S^{n-1}$ be fixed, then the function $F \in G_{n,n-k} \mapsto d(F,\theta) \in \mathbb{R}$ is Lipschitz with constant 1.

Proof. Let $F_1, F_2 \in G_{n,n-k}$. By the definition of d_G , there exists $T \in O_n, T(F_1) = F_2, ||T - I||_{\text{op}} = d_G(F_1, F_2)$. We have

$$|d_G(F_1, \theta) - d_G(F_2, \theta)| = |d_G(F_2, T\theta) - d_G(F_2, \theta)|$$

$$\leq |T\theta - \theta|$$

$$\leq ||T - I||_{op}$$

$$\leq d_G(F_1, F_2),$$

and the lemma follows.

We shall use the following result by Szarek [16].

Theorem 2.6 ([16, Proposition 8]). For every $0 \le \delta \le \sqrt{2} = \text{diam}(G_{n,n-k})$ there exists a δ -net in the Grassmanian $G_{n,n-k}$ with respect to the distance d_G , of cardinality less than $(c/\delta)^{k(n-k)}$.

Proposition 2.7. There exist a universal constant c > 0 and $N = cn^{\frac{k}{2}+4}$ points $\theta_1, \ldots, \theta_N$ such that

$$\frac{1}{N}\sum_{i=1}^{N}\varphi(d(F,n\theta)) \le 4n^{-k/2}$$

for every $F \in G_{n,n-k}$.

Proof. Let $\delta = n^{-k/2-1}$ and consider the corresponding δ -net \mathcal{F} of cardinality $(c'n^{k/2+1})^{k(n-k)} \leq e^{c''n^4}$. Take $\theta_1, \ldots, \theta_N$ independent random points uniformly distributed in S^{n-1} . By Lemma 2.4 we have $\mathbb{E}(\varphi(d(F, n\theta_i))) \leq n^{-k/2}$. Applying Lemma 2.2, for every $F \in \mathcal{F}$ we have

$$\mathbb{P}\left(\frac{1}{N}\sum_{i=1}^{N}\varphi(d(F,n\theta_i)) \ge 3n^{-k/2}\right) \le e^{-n^{-k/2}N} = e^{-cn^4}$$
$$\mathbb{P}\left(\exists F \in \mathcal{F}, \frac{1}{N}\sum_{i=1}^{N}\varphi(d(F,n\theta_i)) \ge 3n^{-k/2}\right) \le e^{c''n^4}e^{-cn^4}$$

which is less than 1 for all $1 \le k \le n$, $n \ge 2$, for some universal constant c > 0 depending on c''. We get the existence of $\theta_1, \ldots, \theta_N$ such that

$$\frac{1}{N}\sum_{i=1}^{N}\varphi(d(F,n\theta_i)) \le 3n^{-k/2}$$

for every $\forall F \in \mathcal{F}$.

By Lemma 2.5, the function $F \mapsto \frac{1}{N} \sum_{i=1}^{N} \varphi(d(F, n\theta_i))$ is Lipschitz with constant n. Then for any $F \in G_{n,n-k}$,

$$\frac{1}{N}\sum_{i=1}^{N}\varphi(d(F,n\theta_i)) \le n\delta + 3n^{-k/2} = 4n^{-k/2}$$

and the proposition follows.

Let $\gamma_n(x) = (2\pi)^{-n/2} e^{-|x|^2/2}$ denote the standard Gaussian probability measure.

Definition 2.8. Let $\theta_1, \ldots, \theta_N$ be the points obtained in Proposition 2.7. Define the function $f_0(x) = \frac{1}{M} \sum_{i=1}^M \gamma_n(x-p_i)$, where

$$\{p_1,\ldots,p_M\}=\{n\theta_1,-n\theta_1,n\theta_2,-n\theta_2,\ldots,n\theta_N,-n\theta_N\},\$$

and M = 2N.

Define the convex set K_0 as the convex hull of the points p_1, \ldots, p_M , and $\pm ne_1, \ldots, \pm ne_n$.

The set K_0 and the function f_0 will satisfy the necessary properties of Theorem 1.4, after normalization and rescaling. These properties will be stated separately in the following three propositions.

Proposition 2.9. Let f_0 be as in Definition 2.8. Let F be any subspace of codimension k with $1 \le k \le n-1$, then

$$\int_{F} f_0 = (2\pi)^{-k/2} \frac{1}{M} \sum_{i=1}^{M} \varphi(d(F, p_i)) \le 4(2\pi n)^{-k/2}.$$

Proof. Let q_i be the orthogonal projection of p_i onto F. Using the change of variables $y = x - q_i$,

$$\int_{F} f_{0} = \frac{1}{M} \sum_{i=1}^{M} (2\pi)^{-n/2} \int_{F} e^{-\frac{1}{2}|x-p_{i}|^{2}} dx$$
$$= \frac{1}{M} \sum_{i=1}^{M} (2\pi)^{-n/2} \int_{F} e^{-\frac{1}{2}|x-q_{i}|^{2} - \frac{1}{2}d(F,p_{i})^{2}} dx$$
$$= (2\pi)^{-k/2} \frac{1}{M} \sum_{i=1}^{M} e^{-\frac{1}{2}d(F,p_{i})^{2}} (2\pi)^{-(n-k)/2} \int_{F} e^{-\frac{1}{2}|y|^{2}} dy$$
$$= (2\pi)^{-k/2} \frac{1}{M} \sum_{i=1}^{M} \varphi(d(F,p_{i})).$$

The inequality follows from Proposition 2.7.

Proposition 2.10. The convex body K_0 in Definition 2.8 satisfies

$$|K_0|_n^{1/n} \le c\sqrt{\log(n)k},$$

where c > 0 is a universal constant.

Proof. By a result of Gluskin [2, Theorem 2], a polytope inside \mathbb{B}_n which has less than m vertices, has volume at most

$$\left(c\frac{\sqrt{\log(m/n)}}{n}\right)^n$$

where c > 0 is a universal constant. Since $K \subseteq n\mathbb{B}_n$,

$$|K|_{n}^{1/n} \leq c\sqrt{\log((M+2n)/n)} \\ \leq c\sqrt{\log((cn^{\frac{k}{2}+4}+2n)/n)} \\ \leq c\sqrt{\log(n)(k/2+3)}$$

and the theorem follows with c > 0 independent of k and n, provided $n \ge 2$. \Box

J. HADDAD

Proposition 2.11. Let K_0 , f_0 be as in Definition 2.8, then

$$\int_{3K_0} f_0 \ge 3/4.$$

Proof. Since K_0 contains the vectors $\pm ne_i$, it contains $\sqrt{n}\mathbb{B}_n$. Then $3K_0$ contains $K_0 + 2\sqrt{n}\mathbb{B}_n$ and

$$\int_{3K_0} f_0 \ge \int_{K_0 + 2\sqrt{n}\mathbb{B}_n} f_0$$

$$\ge \frac{1}{M} \sum_{i=1}^M \int_{p_i + 2\sqrt{n}\mathbb{B}_n} \gamma_n(x - \delta_{p_i}) dx$$

$$= \int_{2\sqrt{n}\mathbb{B}_n} \gamma_n(x) dx \ge 3/4,$$

where the last inequality follows from the Markov-Chebyshev inequality and the fact that $\int_{\mathbb{R}^n} |x|^2 d\gamma_n(x) = n$.

Proof of Theorem 1.4. Consider the set $K = K_0/|K_0|_n^{1/n}$ and the function $f : \mathbb{R}^n \to \mathbb{R}$ defined by

$$f(y) = |3K_0|_n \left(\int_{3K_0} f_0\right)^{-1} f_0(|3K_0|_n^{1/n} y)$$

for $y \in K$, and f(0) = 0 otherwise.

Then clearly $|K|_n = 1$ and $\int_K f = 1$, so f is an even probability density in K, which is also continuous in K.

By Propositions 2.9, 2.10 and 2.11, for any $F \in G_{n,n-k}$,

$$\begin{split} \int_{F} f &\leq |3K_{0}|_{n} \left(\int_{3K_{0}} f_{0} \right)^{-1} \int_{F} f_{0}(3 |K_{0}|_{n}^{1/n} y) \\ &= |3K_{0}|_{n}^{k/n} \left(\int_{3K_{0}} f_{0} \right)^{-1} \int_{F} f_{0} \\ &\leq 3^{k} \left(\sqrt{\log(n)k} \right)^{k} \frac{4}{3} \times 4(2\pi n)^{-k/2} \\ &\leq \left(c \sqrt{\frac{\log(n)k}{n}} \right)^{k}. \end{split}$$

References

- S. Bobkov, B. Klartag, and A. Koldobsky. Estimates for moments of general measures on convex bodies. *Proceedings of the American Mathematical Society*, 146(11):4879–4888, 2018.
- [2] E. D. Gluskin. Extremal properties of orthogonal parallelepipeds and their applications to the geometry of banach spaces. *Mathematics of the USSR-Sbornik*, 64(1):85, 1989.
- [3] W. Gregory and A. Koldobsky. Inequalities for the derivatives of the Radon transform on convex bodies. *Israel Journal of Mathematics*, 246(1):261–280, 2021.
- [4] J. Haddad and A. Koldobsky. Radon transforms with small derivatives and distance inequalities for convex bodies. arXiv preprint arXiv:2312.16923, 2023.
- [5] B. Klartag and A. Koldobsky. An example related to the slicing inequality for general measures. Journal of Functional Analysis, 274(7):2089–2112, 2018.

DISTANCE INEQUALITIES

- [6] B. Klartag and G. V. Livshyts. The Lower Bound for Koldobsky's Slicing Inequality via Random Rounding, pages 43–63. Springer International Publishing, Cham, 2020.
- [7] A. Koldobsky. A functional analytic approach to intersection bodies. Geometric & Functional Analysis GAFA, 10:1507–1526, 2000.
- [8] A. Koldobsky. Fourier analysis in convex geometry. Number 116. American Mathematical Soc., 2005.
- [9] A. Koldobsky. A hyperplane inequality for measures of convex bodies in \mathbb{R}^n , $n \leq 4$. Discrete & Computational Geometry, 47(3):538-547, 2012.
- [10] A. Koldobsky. A \sqrt{n} estimate for measures of hyperplane sections of convex bodies. Advances in Mathematics, 254:33–40, 2014.
- [11] A. Koldobsky. Slicing inequalities for measures of convex bodies. Advances in Mathematics, 283:473–488, 2015.
- [12] A. Koldobsky, G. Paouris, and A. Zvavitch. Measure comparison and distance inequalities for convex bodies. *Indiana University Mathematics Journal*, 2019.
- [13] A. Koldobsky, G. Paouris, and M. Zymonopoulou. Isomorphic properties of intersection bodies. Journal of Functional Analysis, 261(9):2697–2716, 2011.
- [14] A. Markoe. Analytic tomography, volume 13. Cambridge University Press, 2006.
- [15] E. Milman. Generalized intersection bodies. Journal of Functional Analysis, 240(2):530–567, 2006.
- [16] S. J. Szarek. Nets of Grassmann manifold and orthogonal group. In Proceedings of research workshop on Banach space theory (Iowa City, Iowa, 1981), volume 169, page 185. University of Iowa Iowa City, IA, 1982.

(J. Haddad) Departamento de Análisis Matemático, Universidad de Sevilla, Sevilla, España

Email address: jhaddad@us.es