DETERMINISTIC COMPUTATION OF QUANTILES IN A LIPSCHITZ FRAMEWORK

Yurun Gu^{[1](#page-0-0)}, Clément Rey^{[2](#page-0-1)}

ABSTRACT. In this article, our focus lies on computing the quantiles of a random variable $f(X)$, where X is a $[0,1]^d$ -valued random variable, $d \in \mathbb{N}^*$, and $f : [0,1]^d \to \mathbb{R}$ is a deterministic Lipschitz function. We are particularly interested in scenarios where the cost of a single function call is high, while assuming the law of *X* is known. In this context, we propose a deterministic algorithm to obtain exact deterministic lower and upper bounds for the quantile of $f(X)$ at a given level $\alpha \in (0,1)$. With a fixed budget of *N* function calls, we demonstrate that our algorithm achieves exponential convergence rate for $d = 1$ $({\sim \rho^N \text{ with } \rho \in (0,1)})$ and polynomial convergence rate for $d > 1$ $({\sim N^{-\frac{1}{d-1}}})$ and show the optimality of those rates within the class of deterministic algorithms. Furthermore, we design two algorithms based on whether the Lipschitz constant of *f* is known or unknown.

Keywords : Quantile approximation, adaptive algorithm, Lipschitz functions, convergence, optimality. **AMS MSC 2020:** 65C20, 62E17, 65D15, 68W25, 68Q25.

CONTENTS

1. INTRODUCTION

In this article, we consider a $\Omega := [0, 1]^d$ -valued $(d \in \mathbb{N}^*)$ random variable *X* having a density function and consider also a Lipschitz function $f : [0,1]^d \to \mathbb{R}$. We put ourselves in the situation such that the law of *X* is known and that *f* can be evaluated at any point but with high computational cost. For instance, *f* may be a heavy black box implementation while *X* is a random variable with well-known distribution *e.g.* uniform. We propose an algorithm based on $N \in \mathbb{N}$ calls of f in order to compute the *α*-order quantile of $f(X)$, $\alpha \in (0,1)$, defined by

(1.1)
$$
q_{\alpha}(f, X) := \inf\{l \in \mathbb{R}, \mathbb{P}(f(X) \leq l) \geq \alpha\}
$$

in this work, our contribution is to provide a deterministic approximation $q_\alpha^N(f,X)$ for $q_\alpha(f,X)$ and we show that (see Theorem [2.1](#page-4-0) and Theorem [3.1\)](#page-7-0) the error made converges to zero as *N* tends to infinity. More specifically, the rate of convergence is exponential for $d = 1$ $(|q_\alpha^N(f,X) - q_\alpha(f,X)| \leq C \rho^N$ with $\rho \in (0,1)$) and is polynomial when $d > 1$ ($|q_\alpha^N(f,X) - q_\alpha(f,X)| \leqslant CN^{-\frac{1}{d-1}}$).

Quantile computation has diverse applications spanning various domains, with notable instances found in banking [\[8\]](#page-16-1), database optimization [\[13\]](#page-16-2), sociology[\[22\]](#page-17-0) , binary classification [\[21\]](#page-17-1) or sensor networks [\[19\]](#page-17-2). Additionally, related challenges like multivariate quantiles [\[15\]](#page-16-3) or estimating the difference of quantiles at two given levels [\[23\]](#page-17-3) also warrant attention. It is worth mentioning that, many existing quantile estimation methods stem from the equivalent problems of minimizing loss functions [\[18\]](#page-17-4),

$$
q_{\alpha}(X) = \underset{x \in \mathbb{R}}{\operatorname{argmin}} \{ \alpha E[(X - x)^{+}] + (1 - \alpha)E[(X - x)^{-}] \}
$$

¹address: SAMOVAR, Telecom SudParis, Institut Polytechnique de Paris, 19 rue Marguerite Perey, 91120 Palaiseau, France e-mail: yurun.gu@telecom-sudparis.eu

²address: CMAP, École Polytechnique, Institut Polytechnique de Paris, Route de Saclay, 91120 Palaiseau, France e-mail: clement.rey@polytechnique.edu

in which $x^+ := max\{x, 0\}$ $x^- := max\{-x, 0\}$. For some applications, we refer *e.g.* to quantile estimation methods designed in [\[12\]](#page-16-4) or [\[1\]](#page-16-5).

An extensive literature is dedicated to enhancing quantile computation through Monte Carlo methods. Under mild conditions, the Monte Carlo estimator of $q_\alpha(f, X)$ typically achieves a central limit theorem with a rate of $C/N^{\frac{1}{2}}$. Many endeavors aim to reduce the variance of these estimators, represented by the constant *C*. Notably, variance reduction methods can be designed for quantile computation. The importance sampling approach, initially showcased in [\[10\]](#page-16-6), has since been developed further in subsequent works such as [\[11\]](#page-16-7), [\[9\]](#page-16-8), and [\[20\]](#page-17-5). Similarly, the design of control variates can be adapted, as demonstrated in [\[17\]](#page-17-6), [\[16\]](#page-17-7), and [\[3\]](#page-16-9). Additionally, multi-level splitting methods, as described in [\[7\]](#page-16-10), [\[5\]](#page-16-11), and [\[6\]](#page-16-12), have been applied to quantile computation [\[14\]](#page-16-13).

Considering not only the rate of convergence, our algorithm differs from a standard Monte Carlo approach in the following way: Our algorithm is deterministic, meaning it does not rely on simulations of random variables with form $f(X)$. It simply calls the function f at some specific points which are selected sequentially according to the law of *X* and to previous calls of *f*. Also, it provides upper and lower bounds for $q_\alpha(f, X)$ while a Monte Carlo approach provides lower and upper bounds for the quantile by the way of a confidence interval. It is important to note that with Monte Carlo methods, there's a small, albeit nonzero, probability that $q_\alpha(f, X)$ lies outside the computed confidence interval.

We design a first algorithm that requires knowing the Lipschitz constant of *f* for its implementation while a Monte Carlo approach merely necessitates the ability to simulate $f(X)$. Nevertheless we also propose a second algorithm requiring that *f* is Lipschitz but that does not need to have access to the Lipschitz constant. These algorithms draw inspiration from methodologies outlined in [\[4\]](#page-16-14) originally devised to find the minimum of a Lipschitz function. In this case, the rate of convergence is contingent, as for our case, to the dimension with exponential convergence when $d = 1$ and polynomial convergence when $d > 1$. Notice that this algorithm was further adapted in [\[2\]](#page-16-15) in the case of the computation of a failure probability that is the computation of $\mathbb{P}(f(X) > c)$ for a given *c* when the Lipschitz constant is known. In this case, similar regimes of convergences are observed according to the value of *d*.

Our approach revolves around a dichotomous strategy, leveraging the Lipschitz property of *f* to systematically exclude certain regions $\mathcal R$ within $[0,1]^d$ where we know $q_\alpha(f,X) \notin f(\mathcal R)$. By avoiding the evaluation of *f* within these regions, we achieve convergence with the anticipated rate. In cases where the Lipschitz constant is unknown, we employ parallel computations with exponentially increasing candidate Lipschitz constants within a feasible range for region exclusion. Remarkably, this yields convergence rates of comparable order to scenarios where the Lipschitz constants are known.

Furthermore, we establish the optimality of the rates achieved by the algorithms we develop within the class of deterministic algorithms. In essence, under our framework, any deterministic algorithm providing an approximation $\tilde{q}^N_\alpha(f, X)$ (*i.e.* $\tilde{q}^N_\alpha(f, X)$ is built by calling the function *f* at *N* deterministic points) converges at best with rate at exponential, $|\tilde{q}_{\alpha}^{N}(f, X) - q_{\alpha}(f, X)| \geq \tilde{C} \tilde{\rho}^{N}, \tilde{\rho} \in (0, 1)$, when $d = 1$, and polynomial, $|\tilde{q}_{\alpha}^{N}(f, X) - q_{\alpha}(f, X)| \geq \tilde{C} N^{\frac{1}{d-1}}, \tilde{C} > 0$, when $d > 1$.

The article is presented in the following way. The algorithm for the computation of the approximation $q^N_\alpha(f, X)$ for the quantile of order *α* of $f(X)$ when the Lipschitz constant of *f* is known is exposed in Algorithm [2.1.](#page-3-0) Its convergence, along with expected rates, is addressed in Theorem [2.1.](#page-4-0) In the case of unknown Lipschitz constant, the algorithm is presented in Algorithm [3.1](#page-6-1) and the corresponding convergence result discussed in Theorem [3.1.](#page-7-0)The optimality of the rates of convergence achieved by Algorithms [2.1](#page-3-0) and [3.1](#page-6-1) is proved in Section [4.](#page-9-0) A numerical illustration of the convergence of both algorithms is given in Section [5.](#page-15-0)

2. Computation of quantile with known Lipschitz constant

In this section, we design an algorithm for the computation of $q_\alpha(f, X)$ when the Lipschitz constant of *f* is known. We also prove convergence of this algorithm towards $q_\alpha(f, X)$ with explicit rate (polynomial) or exponential regarding the value of *d*) in Theorem [2.1.](#page-4-0) This convergence is established under the following assumptions, which will also be used in the following Section when the Lipschitz constant is unknown,

(1) *f* is a Lipschitz function taking values in \mathbb{R} : There exists $L_f > 0$ such that

$$
\forall x, y \in [0,1]^d, \qquad |f(x) - f(y)| \leqslant L_f |x - y|_{\mathbb{R}^d},
$$

(2) Level set assumption : Let $\alpha \in (0,1)$. There exists $M > 0$ such that

$$
\forall \delta > 0, \quad \lambda_{\text{Leb}}(x \in \mathbb{R}^d, f(x) \in [q_\alpha(f, X) - \delta, q_\alpha(f, X) + \delta]) \leq M\delta
$$

(3) The density function of *X* is bounded by $K > 0$.

2.1. **Preliminaries.** Our first step is to subdivise the set $[0,1]^d$. We begin by introducing the center points of our subdivisions. For $k \in \mathbb{N}$, the level of the subdivision, and $i \in \{0, \ldots, 3^k - 1\}$, we define

$$
d_i^k = \frac{2i+1}{2 \times 3^k},
$$

and for $\beta \in \{0, \ldots, 3^k - 1\}^d$, we focus on the following subdivision,

$$
D_{\beta}^{k} = \{x \in [0,1]^{d}, x_{j} \in [d_{\beta_{j}}^{k} - \frac{1}{2 \times 3^{k}}, d_{\beta_{j}}^{k} + \frac{1}{2 \times 3^{k}}]\},\
$$

Notice that $\bigcup_{\beta \in \{0,\ldots,3^k-1\}^d} D^k_{\beta} = [0,1]^d$. We also define $\delta^k_{\beta} := \sup_{x \in D^k_{\beta}} |x - d^k_{\beta}| \ (d^k_{\beta} = (d^k_{\beta_1}, \ldots, d^k_{\beta_d}))$ and remark that δ^k_β does not depend on β so we simply denote δ^k in the sequel. In particular, we have

$$
\delta^{k} = \left(\sum_{i=1}^{d} (d_{\beta_{i}}^{k} - (d_{\beta_{i}}^{k} + \frac{1}{2 \times 3^{k}}))^{2}\right)^{\frac{1}{2}} = \frac{d^{\frac{1}{2}}}{2} \frac{1}{3^{k}},
$$

For $k \in \mathbb{N}$, we introduce

$$
(2.1) \qquad \mathfrak{q}^k_\alpha(f,X)=\sup\{f(d^k_\beta),\beta\in\{0,\ldots,3^k-1\}^d, \sum_{\gamma\in\{0,\ldots,3^k-1\}^d,f(d^k_\gamma)\geqslant f(d^k_\beta)}\mathbb{P}(X\in D^k_\gamma)\geqslant 1-\alpha\}
$$

with $\sup \emptyset := \inf_{\beta \in \{0, ..., 3^k-1\}^d} f(d_{\beta}^k)$. It happens that $\mathfrak{q}_\alpha^k(f, X)$ is the value returned by our algorithm if it has enough budget to reach level of discretization *k* (and not higher). When we want to emphasize that the computation depends on the Lipschitz constant considered, we will denote $\mathfrak{q}^k_{L_f,\alpha}(f,X)$ instead of $\mathfrak{q}_{\alpha}^{k}(f, X)$. Now we propose an alternative way of defining $\mathfrak{q}_{\alpha}^{k}(f, X)$.

Let us define $\Pi^0 = \{(0, \ldots, 0)\} = \{\{0\}^d\}$, and $\tilde{\mathfrak{q}}_{\alpha}^0(f, X) = f(d_{\{0\}^d}^0)$. Let us assume that for $k \in \mathbb{N}$, we compute $f_{\beta}^k := f(d_{\beta}^k), \ \beta \in \Pi^k \subset \{0, \ldots, 3^k - 1\}^d$ and if $\beta \notin \Pi^k$, no new computation is done and we choose $f_{\beta}^k \in \{f(d_{\beta}^k)\} \cup \{f(d_{\beta-\lbrack l]}^{k-l}), l \in \{1, ..., k\}, \beta^{-\lbrack l-1 \rbrack} \notin \Pi^{k-l+1}\}\$ where for $\beta \in \{0, ..., 3^k-1\}^d$, $\beta^{-[0]} = \beta$ and for $k \in \mathbb{N}^*, \ \beta^{-[1]} = \gamma \in \{0, \ldots, 3^{k-1} - 1\}^d$ such that $\gamma_i = \lfloor \frac{\beta_i}{3} \rfloor, i \in \{1, \ldots, d\},\$ and $\beta^{-[l+1]} = (\beta^{-[l]})^{-[1]}, l \in \mathbb{N}, l \leq k.$

Let define recursively

(2.2)
$$
\tilde{\mathfrak{q}}_{\alpha}^{k}(f, X) = \sup\{f(d_{\beta}^{k}), \beta \in \Pi^{k}, \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{k} \geqslant f(d_{\beta}^{k})} \mathbb{P}(X \in D_{\gamma}^{k}) \geqslant 1 - \alpha\}
$$

and

(2.3)
$$
\Pi^{k+1} = \{3\beta + \{0, 1, 2\}^d, \beta \in \Pi^k, f(d^k_\beta) \in [\tilde{\mathfrak{q}}^k_\alpha - 2L_f\delta^k, \tilde{\mathfrak{q}}^k_\alpha + 2L_f\delta^k] \}
$$

We will sometimes denote $\Pi_{L_f}^k := \Pi^k$ and $\tilde{\mathfrak{q}}_{L_f,\alpha}^k(f,X) := \tilde{\mathfrak{q}}_{\alpha}^k(f,X)$ to emphasize the fact that Π^k is computed in (2.3) using L_f as a Lipschitz constant. In particular such a notation may be relevant when we use an upper bound for L_f instead of its true value. Notice that is if we choose $L \geqslant L' \geqslant L_f$, we have $\Pi_{L_f}^k \subset \Pi_{L'}^k \subset \Pi_L^k$.

Due to the liberty granted by the choice of f_{γ}^k , one may argue that $\tilde{\mathfrak{q}}_{\alpha}^k(f, X)$ may differ *w.r.t.* the value chosen for f^k_γ . The following result shows that all the definitions are actually equivalent. It also shows that $\tilde{\mathfrak{q}}_{L,\alpha}^k(f,X) = \tilde{\mathfrak{q}}_{L_f,\alpha}^k(f,X)$ for $L \geqslant L_f$.

Lemma 2.1. For every $k \in \mathbb{N}$, we have

A. The quantity $\mathfrak{q}_{\alpha}^{k}(f, X)$ defined in [\(2.1\)](#page-2-1) satisfies

$$
(2.4) \qquad \mathfrak{q}_{\alpha}^k(f, X) = \inf\{f(d_{\beta}^k), \beta \in \{0, \dots, 3^k - 1\}^d, \sum_{\gamma \in \{0, \dots, 3^k - 1\}^d, f(d_{\gamma}^k) \leq f(d_{\beta}^k)} \mathbb{P}(X \in D_{\gamma}^k) \geq \alpha\}
$$

B. The quantity $\tilde{\mathfrak{q}}_{\alpha}^k(f, X)$ defined in [\(2.4\)](#page-3-1) satisfies

$$
\tilde{\mathfrak{q}}^k_\alpha(f,X)=\inf\{f(d^k_\beta),\beta\in\Pi^k, \sum_{\gamma\in\{0,...,3^k-1\}^d,f^k_\gamma\leqslant f(d^k_\beta)}\mathbb{P}(X\in D^k_\gamma)\geqslant \alpha\}
$$

C. Moreover,

$$
\mathfrak{q}^k_\alpha(f,X)=\tilde{\mathfrak{q}}^k_\alpha(f,X)
$$

Proof. The result is immediate for $k = 0$. We consider $k \in \mathbb{N}^*$.

Let us denote

$$
\mathfrak{p}^k_\alpha:=\inf\{f(d^k_\beta), \beta\in\{0,\ldots,3^k-1\}^d, \sum_{\gamma\in\{0,\ldots,3^k-1\}^d, f(d^k_\gamma)\leqslant f(d^k_\beta)}\mathbb{P}(X\in D^k_\gamma)\geqslant \alpha\}
$$

We remark that, (denoting shortly $\mathfrak{q}_{\alpha}^k := \mathfrak{q}_{\alpha}^k(f, X)$)

$$
\mathbb{P}(X \in \bigcup_{\gamma \in \{0,\ldots,3^k-1\}^d, f(d^k_\gamma) < q^k_\alpha} D^k_\gamma) < \alpha,
$$

so that $\mathfrak{p}_{\alpha}^{k} \geq \mathfrak{q}_{\alpha}^{k}$. Moreover, by definition of $\mathfrak{q}_{\alpha}^{k}$

$$
\mathbb{P}(X \in \bigcup_{\gamma \in \{0, \ldots, 3^k-1\}^d, f(d^k_\gamma) > \mathfrak{q}^k_\alpha} D^k_\gamma) < 1 - \alpha.
$$

It follows that

$$
\mathbb{P}(X \in \bigcup_{\gamma \in \{0,\ldots,3^k-1\}^d, f(d_{\gamma}^k) \leqslant \mathfrak{q}_{\alpha}^k} D_{\gamma}^k) \geqslant \alpha,
$$

and since there exists $\beta^* \in \{0, \ldots, 3^k - 1\}^d$, such that $f(d_{\beta^*}^k) = \mathfrak{q}_\alpha^k$, we conclude that $\mathfrak{q}_\alpha^k \geq \mathfrak{p}_\alpha^k$ and the proof of **[A.](#page-3-2)** is completed.

Now let us prove **C**. This is true for $k = 0$ and let us prove it for $k \in \mathbb{N}^*$. Using definition [2.1,](#page-2-1) we have $q_{\alpha}(f, X) \geqslant \mathfrak{q}_{\alpha}^{k} - L_{f} \delta^{k}$ and using definition [2.4](#page-3-1) we have $q_{\alpha}(f, X) \leqslant \mathfrak{q}_{\alpha}^{k} + L_{f} \delta^{k}$. Assume that there exists $\beta_0 \notin \Pi^k$ such that $\mathfrak{q}_\alpha^k = f(d_{\beta_0}^k)$. Since $\beta_0 \notin \Pi^k$, then

$$
|\mathfrak{q}_{\alpha}^{k-1}-f(d_{\beta_0^{-}[1]}^{k-1})|>2L_f\delta^{k-1}.
$$

so that

$$
|q_{\alpha}(f, X) - f(d_{\beta_0}^k)| \geqslant |q_{\alpha}^{k-1} - f(d_{\beta_0}^{k-1})| - |f(d_{\beta_0}^{k-1}) - f(d_{\beta_0}^k)| - |q_{\alpha}^{k-1} - q_{\alpha}(f, X)|
$$

$$
> 2L_f \delta^{k-1} - 2L_f \delta^k - L_f \delta^{k-1}
$$

$$
= L_f \delta^{k-1} - 2L_f \delta^k = L_f \delta^k
$$

which is impossible due to the previous observation. In particular we can replace the condition $\beta \in$ $\{0,\ldots,3^k-1\}^d$ by $\beta \in \Pi^k$ in the definition [2.1.](#page-2-1) It remains to observe that $f(d^k_\gamma)$ may be replaced by f^k_γ as soon as $\gamma \notin \Pi^k$. Indeed, for such γ , using the same calculus as above $f(d_{\gamma-1}^{k-1}) - \mathfrak{q}_{\alpha}^{k-1}$ and $f(d_{\gamma}^k) - \mathfrak{q}_{\alpha}^k$ necessarily have the same sign and the proof of **[C.](#page-3-3)** is completed. Using similar argument we obtain the proof of **B**.

Furthermore, we can obtain an upper bound for the error between $q_\alpha(f, X)$ and our estimator, regarding that our algorithm reaches level *k* (and not higher), *i.e.* \mathfrak{q}_{α}^k .

Lemma 2.2. For any $k \in \mathbb{N}$, we have

(2.5)
$$
|q_{\alpha}(f, X) - \mathfrak{q}_{\alpha}^{k}| \leqslant L_{f} \delta^{k}
$$

Proof. This result is a direct consequence of Lemma [2.1](#page-2-2) **[A.](#page-3-2)** demonstrated in the proof of Lemma [2.1](#page-2-2) Point \mathbf{C} . 2.2. **Algorithm and Main result.** We present our algorithm when the Lipschitz constant L_f of f is known. Notice that the knowledge of the constant *M* appearing in the level-set assumption [\(2\)](#page-2-3) is not necessary to implement the algorithm.

Algorithm 2.1.

Fix $N \in \mathbb{N}^*$ the maximum number of calls to f $\Pi^0 = \{(0, \ldots, 0)\}.$ $k = 0$. $N_{\text{call}} = 1.$ **While** $N_{\text{call}} \leq N$ Compute $f(d_{\beta}^k)$, $\beta \in \Pi^k$ (When $d_{\beta}^k = d_{\beta^{-1}}^{k-1}$, the computation was already done before step *k*).

Set $f_{\beta}^k = f(d_{\beta}^k)$, for every $\beta \in \Pi^k$ and $f_{\beta}^k = f_{\beta}^{k-1}$, for every $\beta \notin \Pi^k$. **Set**

$$
q_{\alpha}^{N}(f, X) = \sup\{f(d_{\beta}^{k}), \beta \in \Pi^{k}, \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{k} \geq f(d_{\beta}^{k})} \mathbb{P}(X \in D_{\gamma}^{k}) \geq 1 - \alpha\}
$$

$$
= \inf\{f(d_{\beta}^{k}), \beta \in \Pi^{k}, \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{k} \leq f(d_{\beta}^{k})} \mathbb{P}(X \in D_{\gamma}^{k}) \geq \alpha\}.
$$
and
$$
q_{\alpha}^{N}(f, X) = q_{\alpha}^{N}(f, X) - L_{f}\delta^{k}, \overline{q}_{\alpha}^{N}(f, X) = q_{\alpha}^{N}(f, X) + L_{f}\delta^{k}.
$$
Set
$$
\Pi^{k+1} = \{3\beta + \{0, 1, 2\}^{d}, \beta \in \Pi^{k}, f(d_{\beta}^{k}) \in [q_{\alpha}^{N} - L_{f}\delta^{k}, \overline{q}_{\alpha}^{N} + L_{f}\delta^{k}]\},
$$

Set

$$
\Pi^{k+1} = \{3\beta + \{0, 1, 2\}^d, \beta \in \Pi^k, f(d_\beta^k) \in [q_\alpha^N - L_f \delta^k, \overline{q}_\alpha^N + L_f \delta^k]
$$

\n
$$
N_{\text{call}} = N_{\text{call}} + \frac{3^d - 1}{3^d} \text{Card}(\Pi^{k+1}),
$$

\nSet $k = k + 1$.

End While

 $\textbf{Return } q^N_{\alpha}(f, X), \, \underline{q}^N_{\alpha}$ $\frac{N}{\alpha}(f, X), \overline{q}_{\alpha}^N(f, X).$

We observe that to apply this algorithm, we need to have access to the law of *X*, or more specifically to $\mathbb{P}(X \in D_{\gamma}^{k})$, which is why this law is supposed to be known. When the probabilities $\mathbb{P}(X \in D_{\gamma}^{k})$ are unknown but *X* may be simulated easily, one can use a Monte Carlo approach for their computation which can be done once for all independently from the function *f*. We do not discuss further this approach in this paper.

The following result establish the convergence of Algorithm [2.1](#page-3-0) with explicit upper bound for the error observing two regimes, exponential or polynomial, regarding that $d = 1$ or $d > 1$.

Theorem 2.1. Let $N \in \mathbb{N}^*$ and let $q^N_{\alpha}(f, X), q^N_{\alpha}$ $\frac{N}{\alpha}$ and \overline{q}^N_α given by the Algorithm [2.1.](#page-3-0) Assume that [1,](#page-1-1) [2](#page-2-3) and [3](#page-2-4) hold. Then

(2.6)
$$
q_{\alpha}(f,X) \in [q_{\alpha}^{N}(f,X),\overline{q}_{\alpha}^{N}(f,X)]
$$

Moreover, if $d = 1$, then

(2.7)
$$
|q_{\alpha}(f, X) - q_{\alpha}^{N}(f, X)| \leqslant C\rho^{N}
$$

with
$$
C = \frac{1}{2} L_f 3^{1 + \frac{1}{4ML_f}}
$$
, $\rho = 3^{-\frac{1}{4ML_f}}$.

If $d > 1$ and $N > 1$, then

(2.8)
$$
|q_{\alpha}(f, X) - q_{\alpha}^{N}(f, X)| \leq C(N - 1)^{-\frac{1}{d - 1}}
$$

with
$$
C = \frac{3}{2} L_f d^{\frac{1}{2}} (3^d M L_f d^{\frac{1}{2}})^{\frac{1}{d-1}}
$$

Remark 2.2. If $X \in \prod_{i=1}^d [a_i, b_i]$, then $f(x) = g(h(x))$, with $h(x)_i = \frac{x_i - a_i}{b_i - a_i}$, with $g(y) = f(v(y))$ with $v(y)_i = a_i + (b_i - a_i)y_i$ and g is defined on $[0,1]^d$ and is Lipschitz with Lipschitz constant $L_g \leqslant cL_f$. with $c = \sup\{b_i - a_i, i \in \{1, ..., d\}\}\.$ In particular we can apply Algorithm [2.1](#page-3-0) with the random variable $h(X)$ instead of *X* and the function *g* instead of *f* to produce approximations for q_α , the quantile of order α of $f(X)$. In particular

$$
q_{\alpha}(f, X) \in [q_{\alpha}^N(f, X), \overline{q}_{\alpha}^N(f, X)] := [q_{\alpha}^N(g, h(X)), \overline{q}_{\alpha}^N(g, h(X))]
$$

and [\(2.7\)](#page-4-1) and [\(2](#page-4-2).8) hold for $\overline{q}_{\alpha}^{N}(f, X) - \underline{q}_{\alpha}^{N}$ $\binom{N}{\alpha}(f, X)$ with L_f replaced by cL_f in the upper bound.

2.3. **Proof of convergence of the algorithm.** At this point, we introduce N_k the number of calls to the functions *f* satisfying [1,](#page-1-1) in order to compute \mathfrak{q}_{α}^k . Since each new subdivision of a set with form D_{β}^l yields $3^d - 1$ new calls to the function *f*, we have

$$
N_k = 1 + (3^d - 1) \sum_{l=0}^{k-1} \sum_{\gamma \in \{0, \dots, 3^l-1\}^d} \mathbf{1}_{f(d^l_\gamma) \in [\mathfrak{q}^l_\alpha - 2L_f \delta^l, \mathfrak{q}^l_\alpha + 2L_f \delta^l]}.
$$

Lemma [2](#page-2-3).3. Let $k \in \mathbb{N}$. Assume that [1](#page-1-1) and 2 hold.

If $d = 1$, then

$$
N_k \leqslant 1 + 4ML_f k.
$$

If $d > 1$, then

$$
N_k \leq 1 + 3^d 2ML_f d^{\frac{1}{2}} \frac{3^{k(d-1)} - 1}{3^{d-1} - 1}.
$$

Proof. First we remark that

$$
N_k = 1 + (3^d - 1) \sum_{l=0}^{k-1} \sum_{\gamma \in \{0, ..., 3^l-1\}^d} \mathbf{1}_{f(d_\gamma^l) \in [q_\alpha^l - 2L_f \delta^l, q_\alpha^l + 2L_f \delta^l]}
$$

= 1 + (3^d - 1) $\sum_{l=0}^{k-1} \sum_{\gamma \in \{0, ..., 3^l-1\}^d} 3^{ld} \lambda_{\text{Leb}}(D_\gamma^l) \mathbf{1}_{f(d_\gamma^l) \in [q_\alpha^l - 2L_f \delta^l, q_\alpha^l + 2L_f \delta^l]}$

$$
\leq 1 + (3^d - 1) \sum_{l=0}^{k-1} 3^{ld} \lambda_{\text{Leb}}(x \in [0, 1]^d, f(x) \in [q_\alpha^l - 3L_f \delta^l, q_\alpha^l + 3L_f \delta^l]).
$$

Now, we introduce $r_\alpha^l = 4L_f\delta^l$. Then, using the level set assumption [2](#page-2-3) and that $[\mathfrak{q}_\alpha^l - 3L_f\delta^l, \mathfrak{q}_\alpha^l + 3L_f\delta^l] \subset$ $[q_{\alpha}(f, X) - r_{\alpha}^l, q_{\alpha}(f, X) + r_{\alpha}^l]$, we obtain

$$
\lambda_{\text{Leb}}(x \in [0,1]^d, f(x) \in [\mathfrak{q}_{\alpha}^l - 3L_f \delta^k, \mathfrak{q}_{\alpha}^l + 3L_f \delta^l]) \leq M r_{\alpha}^l,
$$

and then

$$
N_k \le 1 + (3^d - 1) \sum_{l=0}^{k-1} M 3^{ld} r_\alpha^l
$$

$$
\le 1 + (3^d - 1) 2ML_f d^{\frac{1}{2}} \sum_{l=0}^{k-1} 3^{l(d-1)},
$$

and we conclude easily regarding the value of *d*.

Lemma 2.4. Let $N \in \mathbb{N}^*$. Introduce $k(N) := \sup\{k \in \mathbb{N}, N_k \leq N\}$.

If $d = 1$, then

$$
k(N)\geqslant \lfloor \frac{N-1}{4ML_f}\rfloor
$$

and

(2.9)
$$
|q_{\alpha}(f, X) - \mathfrak{q}_{\alpha}^{k(N)}| \leq \frac{3}{2} L_f 3^{-\frac{N-1}{4ML_f}}
$$

If $d > 1$, then

$$
k(N) \geqslant \lfloor \frac{\ln(N-1) - \ln(3^d ML_f d^{\frac{1}{2}})}{(d-1)\ln(3)} \rfloor
$$

and

□

(2.10)
$$
|q_{\alpha}(f, X) - \mathfrak{q}_{\alpha}^{k(N)}| \leq \frac{3}{2} L_f d^{\frac{1}{2}} \left(\frac{N-1}{3^d M L_f d^{\frac{1}{2}}}\right)^{-\frac{1}{d-1}}
$$

Notice that $k(N)$ depends on L_f . When necessary, we will emphazise this dependence denoting $k(N, L_f)$ instead of $k(N)$.

Proof. The proof is a straightforward application of Lemma [2.3](#page-5-0) together with $\delta^k = \frac{d^{\frac{1}{2}}}{2} \frac{1}{3^k}$.

Proof of Theorem [2.1.](#page-4-0) We remark that q_{∞}^N $a_{\alpha}^{N}(f, X) = \mathfrak{q}_{\alpha}^{k(N)} - L_{f} \delta^{k(N)}$ and that $\bar{q}_{\alpha}^{N}(f, X) = \mathfrak{q}_{\alpha}^{k(N)} + \bar{q}_{\alpha}^{k(N)}$ $L_f \delta^{k(N)}$. Then the proof of [\(2.6\)](#page-4-3) is a consequence of Lemma [2.2.](#page-3-5) The proof of [2.7](#page-4-1) and [2.8](#page-4-2) is a direct consequence of Lemma [2.4](#page-5-1) (see [\(2.9\)](#page-5-2) and [\(2.10\)](#page-6-2)).

3. Computation of quantile with unknown Lipschitz constant

In this Section, we design an algorithm for the computation of $q_\alpha(f, X)$ when the Lipschitz constant of *f* is unknown and prove its convergence in Theorem [3.1.](#page-7-0) Similarly as when the Lipischitz constant is known, we observe two regimes (exponential or polynomial) for the error rate regarding the value of *d*.

3.1. **Preliminaries.** We suppose in this section that L_f is unknown and for $j \in \mathbb{N}$ we introduce

$$
L_f(j) = 3^j
$$

,

the candidates for the unkown Lipischitz constant *L^f* of *f*. Let us consider a given global budget $N \in \mathbb{N}, N \geqslant \frac{\pi^2}{6}$ $\frac{\pi^2}{6}$, and introduce $N_{max}(j, N) := \lfloor \frac{6N}{\pi^2 (j+1)^2} \rfloor$ the respective budgets allocated for com-putations that mimick Algorithm [2.1](#page-3-0) in the case the true Lipschitz constant is $L_f(j)$. We also denote *j*_{*max}*(*N*) = sup{*j* ∈ N*, N*_{*max*}(*j, N*) ≥ 1 } = sup{*j* ∈ N*, j* ≤ $\frac{\sqrt{6N}}{\pi} - 1$ } = $\frac{\sqrt{6N}}{\pi} - 1$.</sub>

Let us define $\Pi^{\diamond,0}(j) = \{(0,\ldots,0)\} = \{0^d\}$, and $\tilde{\mathfrak{q}}^0_\alpha(f,X) = f(d^0_{\{0\}^d})$. Let us assume that for $k \in \mathbb{N}$, we compute (if necessary) $f_{\beta}^k := f(d_{\beta}^k)$, $\beta \in \bigcup_{j=1}^{j_{max}} \Pi^{\diamond,k}(j)$ and if $\beta \notin \bigcup_{j=1}^{j_{max}} \Pi^{\diamond,k}(j)$, no new computation is done and we choose $f_{\beta}^k = f_{\beta^{-1}}^{k-1}$.

Let define recursively

$$
(3.1) \qquad \qquad \mathfrak{q}_\alpha^{\diamond,k}(f,X)=\sup\{f(d^k_\beta),\beta\in \cup_{j=1}^{j_{max}}\Pi^{\diamond,k}(j),\sum_{\gamma\in\{0,\ldots,3^k-1\}^d,f^k_\gamma\geqslant f(d^k_\beta)}\mathbb{P}(X\in D^k_\gamma)\geqslant 1-\alpha\}
$$

and

$$
\Pi^{\diamond,k+1}(j) = \{3\beta + \{0,1,2\}^d, \beta \in \Pi^{\diamond,k}(j), f(d_{\beta}^k) \in [\mathfrak{q}_{\alpha}^{\diamond,k} - 2L_f(j)\delta^k, \mathfrak{q}_{\alpha}^{\diamond,k} + 2L_f(j)\delta^k]\}
$$
 if $k+1 \leq \ell(j,N)$ and

$$
\Pi^{\diamond,k+1}(j) = 3\Pi^{\diamond,k}(j) + (1,\ldots,1)
$$

otherwise.

Notice that similarly that in the known Lipischitz constant case, we have

$$
(3.2) \qquad \qquad \mathfrak{q}_\alpha^{\diamond,k}(f,X)=\inf\{f(d_\beta^k),\beta\in \cup_{j=1}^{j_{max}}\Pi^{\diamond,k}(j)\sum_{\gamma\in\{0,\ldots,3^k-1\}^d,f_\gamma^k\leqslant f(d_\beta^k)}\mathbb{P}(X\in D_\gamma^k)\geqslant \alpha\}
$$

and $\mathfrak{q}_\alpha^{\diamond,k}(f,X)$ is the value returned by our algorithm if it has enough budget to reach level of discretization *k* (and not higher).

3.2. **Algorithm and Main result.** We begin by introducing our algorithm when the Lipschitz constant *L^f* of *f* is unknown.

Algorithm 3.1. Fix $N \in \mathbb{N}^*$ the maximum number of calls to f $\Pi^0 = \{(0, \ldots, 0)\}.$

 $k = 0$ $J = \{1, \ldots, j_{max}(N)\}.$ **Set** $N_{\text{call}}(j) = 1$ for all $j \in J$. **While** $J \neq \emptyset$ Compute $f(d_{\beta}^k)$, $\beta \in \bigcup_{j=1}^{j_{max}(N)} \Pi^{\diamond,k}(j)$ (When $d_{\beta}^k = d_{\beta-\lfloor 1 \rfloor}^{k-1}$, the computation was already done before step *k*).

Set
$$
f_{\beta}^k = f(d_{\beta}^k)
$$
, for every $\beta \in \bigcup_{j=1}^{j_{max}} \Pi^{\diamond,k}(j)$ and $f_{\beta}^k = f_{\beta - [1]}^{k-1}$, for every $\beta \notin \bigcup_{j=1}^{j_{max}} \Pi^{\diamond,k}(j)$.
Set

Set

$$
q_{\alpha}^{\diamond, N}(f, X) = \sup\{f(d_{\beta}^k), \beta \in \bigcup_{j=1}^{j_{max}(N)} \Pi^{\diamond, k}(j), \sum_{\gamma \in \{0, \ldots, 3^k-1\}^d, f_{\gamma}^k \ge f(d_{\beta}^k)} \mathbb{P}(X \in D_{\gamma}^k) \ge 1 - \alpha\}
$$

$$
= \inf\{f(d_{\beta}^k), \beta \in \bigcup_{j=1}^{j_{max}(N)} \Pi^{\diamond, k}(j), \sum_{\gamma \in \{0, \ldots, 3^k-1\}^d, f_{\gamma}^k \le f(d_{\beta}^k)} \mathbb{P}(X \in D_{\gamma}^k) \ge \alpha\}.
$$

For $j \in J$ Set $\Pi^{\diamond,k+1}(j) = \{3\beta + \{0,1,2\}^d, \beta \in \Pi^{\diamond,k}(j), f(d_{\beta}^k) \in [q_{\alpha}^{\diamond,N} - 2L_f(j)\delta^k, q_{\alpha}^{\diamond,N} + 2L_f(j)\delta^k] \}.$ $N_{\text{call}}(j) = N_{\text{call}}(j) + \frac{3^d - 1}{3^d} \text{Card}(\Pi^{\diamond, k+1}(j)).$ **Set If** $N_{\text{call}}(j) > N_{\text{max}}(j, N)$. $J = J \setminus \{j\}.$ **For** $j \in \{1, \ldots, j_{max}(N)\} \setminus J$ $\Pi^{\diamond,k+1}(i) = 3\Pi^{\diamond,k}(i) + (1,\ldots,1).$

Set $k = k + 1$.

End While Return $q_{\alpha}^{\diamond, N}(f, X)$.

The following result establishes the convergence of Algorithm [3.1](#page-6-1) with explicit upper bound for the error. We still observe exponential or polynomial regimes regarding that $d = 1$ or $d > 1$. Nevertheless, as expected, the constant obtained (denoted *ρ* and *C* in Theorem [2.1](#page-4-0) and Theorem [3.1\)](#page-7-0) provides a faster convergence to zero for the error bound when the Lipschitz constant is known.

Theorem [3.1.](#page-6-1) Let $N \in \mathbb{N}^*$ and let $q_\alpha^{\diamond, N}(f, X)$ given by the Algorithm 3.1. Assume that [1,](#page-1-1) [2](#page-2-3) and [3](#page-2-4) hold with $L_f \geq 1$.

If $d=1$, then

$$
|q_{\alpha}(f, X) - q_{\alpha}^{\diamond, N}(f, X)| \leq C\rho^{N}
$$

with $C = 18L_f 3^{\frac{1}{2ML_f}}$ and $\rho = 3^{-\frac{1}{(\frac{\ln(L_f)}{\ln(3)} + 2)^2 2\pi^2 ML_f}}$.

If $d > 1$, and $N > \frac{\pi^2}{3} (\frac{\ln(L_f)}{\ln(3)} + 2)^2$,

$$
|q_{\alpha}(f, X) - q_{\alpha}^{\diamond, N}(f, X)| \leq C(N - \frac{\pi^2}{3} (\frac{\ln(L_f)}{\ln(3)} + 2)^2)^{-\frac{1}{d-1}}
$$

with $C = 18L_f d^{\frac{1}{2}} (3^d ML_f d^{\frac{1}{2}} \frac{\pi^2}{2} (\frac{\ln(L_f)}{\ln(3)} + 2)^2)^{\frac{1}{d-1}}$.

3.3. **Proof of convergence of the algorithm.** Recalling that $L_f(j) = 3^j$, $j \in \mathbb{N}$, we define the (unknow) quantity $j^* \in \mathbb{N}$, such that $L_f(j^* - 1) < L_f \leq L_f(j^*)$ with convention $L_f(-1) = 0$. In particular $\frac{\ln(L_f)}{\ln(3)} \leq j^* < \frac{\ln(3L_f)}{\ln(3)}$ since $L_f \geq 1$.

The proof of Theorem [3.1](#page-7-0) is a consequence of the following result combined with the fact that $L_f(j^*) \in [L_f, 3L_f)$. Indeed, let us introduce $\ell(j, N) := k(N_{max}(j, N), L_f(j))$ when $j \in \{0, \ldots, j_{max}(N)\}\$ $(\ell(j, N) := 0$ for $j > j_{max}(N)$). Then, given a budget of $N_{\text{call}}(j)$ for the Lipischtz constant candidate $L_f(j)$, it follows from Lemma [2.4](#page-5-1) that the highest level *k* attained for discretization of the space satisfies $k \geq \ell(j, N)$. Assuming this highest level is exactly $\ell(j, N)$, the proof of Theorem [3.1](#page-7-0) is a direct consequence of Lemma [3.1](#page-7-1) thereafter. When this level is higher the proof is similar and left to the reader.

Lemma 3.1. For any $k \in \mathbb{N}$, we have

(3.3)
$$
|q_{\alpha}(f, X) - \mathfrak{q}_{\alpha}^{\diamond,k}| \leq 4L_f(j^*)\delta^{\min(k, \ell(j^*, N))}
$$

Proof. First let us consider the case j^* > $j_{max}(N)$. In this case $\mathfrak{q}_{\alpha}^{\diamond,k} \in f([0,1]^d)$ and the result follows from the Lipschitz property of *f* since $\ell(j^*, N) = 0$.

Now let us assume that $j^* \in \{0, \ldots, j_{max}(N)\}$. Let us first consider the case $k \leq \ell(j^*, N)$. We remark that (using notation from [\(2.3\)](#page-2-0)), $\Pi_{L_f}^k \subset \Pi_{L_f(j^*)}^k = \Pi^k(j^*) \subset \bigcup_{j=0}^{j_{max}} \Pi^k(j)$. In particular it follows $\mathfrak{q}_{\alpha}^{\diamond,k}(f,X) = \mathfrak{q}_{L_f(j^*),\alpha}^k(f,X) = \mathfrak{q}_{L_f,\alpha}^k(f,X) = \mathfrak{q}_{\alpha}^k(f,X)$ and the result is a direct consequence of Lemma [2.2.](#page-3-5)

We now consider $k > \ell(j^*, N)$. We first notice that if $j > j^*, \ell(j^*, N) \geq \ell(j, N)$ and it is impossible that $\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,N)}(f,X) = f(d_{\gamma}^{\ell(j^*,l)})$ for $\gamma \in \Pi^{\diamond,\ell(j^*,N)}(j) \setminus \Pi^{\diamond,\ell(j^*,N)}(j^*)$. Let us show that it holds for $k > \ell(j^*,N)$ that it is impossible that $\mathfrak{q}_{\alpha}^{\diamond,k}(f,X) = f(d_{\gamma_*}^{\ell(j^*,l)})$ for $\gamma_* \in \Pi^{\diamond,\ell(j^*,N)}(j) \setminus \Pi^{\diamond,\ell(j^*,N)}(j^*)$. Indeed, since $\gamma_* \notin \Pi^{\diamond, \ell(j^*, N)}(j^*)$ we have

$$
|\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,l)-1}(f,X)-f(d_{\gamma_*^{-}[1]}^{\ell(j^*,l)-1})|>2L_f(j^*)\delta^{\ell(j^*,l)-1}.
$$

Moreover, let us notice that $(\mathfrak{q}_{L_f(j^*),\alpha}^k - L_f(j^*)\delta^k)_{k \in \mathbb{N}^*}$ is increasing while $(\mathfrak{q}_{L_f(j^*),\alpha}^k + L_f(j^*)\delta^k)_{k \in \mathbb{N}^*}$ is decreasing. In particular, we have

$$
|\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,l)}(f,X) - \mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,N)-1}(f,X)| \leqslant 2 L_f(j^*) \delta^{\ell(j^*,N)}
$$

Using the Lipischitz property of *f* yields also

$$
|f(d_{\gamma_*^{-}[1]}^{\ell(j^*,l)-1})-f(d_{\gamma_*}^{\ell(j^*,l)})|\leqslant 2L_f(j^*)\delta^{\ell(j^*,N)}
$$

and we conclude by triangle inequality that

$$
|\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,l)}(f,X)-f(d_{\gamma_*}^{\ell(j^*,l)})|>2L_f\delta^{\ell(j^*,N)}.
$$

Assume now that $\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,l)}(f,X) \leqslant f(d_{\gamma_*}^{\ell(j^*,l)}),$ so that

$$
\sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{k} \geq f(d_{\gamma_{*}}^{\ell(j^{*},l)})} \mathbb{P}(X \in D_{\gamma}^{k}) \leq \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{\ell(j^{*},N)} = [k-\ell(j^{*},l)]} \mathbb{P}(X \in D_{\gamma}^{k})
$$
\n
$$
\leq \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{\ell(j^{*},N)} = [k-\ell(j^{*},l)]} \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{\ell(j^{*},N)} = [k-\ell(j^{*},l)]} \mathbb{P}(X \in D_{\gamma}^{k})
$$
\n
$$
\leq \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{\ell(j^{*},N)} = [k-\ell(j^{*},l)]} \mathbb{P}(X \in D_{\gamma}^{k})
$$
\n
$$
\leq \sum_{\gamma \in \{0, ..., 3^{k}-1\}^{d}, f_{\gamma}^{\ell(j^{*},l)} = [k-\ell(j^{*},l)]} \mathbb{P}(X \in D_{\gamma}^{k})
$$
\n
$$
\approx \sum_{\gamma \in \{0, ..., 3^{\ell(j^{*},N)} - 1\}^{d}, f_{\gamma}^{\ell(j^{*},l)} = q_{\alpha}^{\diamond, \ell(j^{*},l)}(f,X)}
$$
\n
$$
\leq 1 - \alpha
$$
\n
$$
\mathbb{P}(X \in D_{\gamma}^{k})
$$

where the last inequality is a direct consequence of definition [\(3.1\)](#page-6-3) and we conclude that $\mathfrak{q}_{\alpha}^{\diamond,k}(f,X) \neq$ $f(d_{\gamma_*}^{\ell(j^*,l)})$. When $\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,l)}(f,X) \geq f(d_{\gamma_*}^{\ell(j^*,l)})$, we use the same approach but using definition [\(3.2\)](#page-6-4) instead of [\(3.1\)](#page-6-3) Now, we remark that for every $j \in \{0, \ldots, j^*\}$, we have $\Pi^{\diamond, \ell(j^*)}(j) \subset \Pi^{\diamond, \ell(j^*)}(j^*)$ and it follows from the previous observation that for every $k \geq \ell(j^*, N)$,

$$
\mathfrak{q}_{\alpha}^{\diamond,k}(f,X)\in f(\cup_{\beta\in\Pi^{\ell(j^*,N)}}D_{\beta}^{\ell(j^*,N)})
$$

and

$$
\begin{aligned} |\mathfrak{q}_\alpha^{\diamond,k}(f,X)-\mathfrak{q}_\alpha^{\diamond,\ell(j^*,N)}(f,X)|&\leqslant \sup\limits_{\beta\in\Pi^{\ell(j^*,N)}}|f(d_\beta^{\ell(j^*,N)})-\mathfrak{q}_\alpha^{\diamond,\ell(j^*,N)}(f,X)|\\ &+\sup\limits_{\beta\in\Pi^{\ell(j^*,N)}}\sup\limits_{x\in D_\beta^{\ell(j^*,N)}}|f(d_\beta^{\ell(j^*,N)})-f(x)|\\ &\leqslant 3L_f(j^*)\delta^{\ell(j^*,N)} \end{aligned}
$$

and we deduce the result from the triangle inequality and Lemma [2.2](#page-3-5) with $k = \ell(j^*, N)$ using that $\mathfrak{q}_{\alpha}^{\diamond,\ell(j^*,N)}(f,X) = \mathfrak{q}_{L_{\ell}(j^*,N)}^{\ell(j^*,N)}$ *L^f* (*j* [∗])*,α* (f, X) .

4. Optimality

In this section, we aim to show the optimality of our algorithm. Specifically, we demonstrate that in our framework where [\(1\)](#page-1-1), [\(2\)](#page-2-3) and [\(3\)](#page-2-4) are assumed, it is impossible to build lower and upper deterministic bounds for $q_\alpha(f, X)$ such that the error made, *i.e.* essentially the difference between upper and lower bound, converges faster than exponential or polynomial rate (given the value of *d*) as *N* tends to infinity.

Let us consider an algorithm which evaluates $N \in \mathbb{N}^*$ times the function $f : \mathbb{R}^d \to \mathbb{R}$ at some points $(x^1, \ldots, x^N) \in ([0,1]^d)^N$ and which returns a measurable function of those evaluations to compute $q_{\alpha}(f, X), \alpha \in (0, 1)$ with *f*, *X* and α satisfying [\(1\)](#page-1-1), [\(2\)](#page-2-3) and [\(3\)](#page-2-4). We are going to show that the error made in this case is at least of order ρ^N ($\rho \in (0,1)$) when $d=1$ and $N^{-\frac{1}{d-1}}$ when $d>1$.

To do so, we propose a function *f*, a random variable *X* and $\alpha \in (0,1)$ satisfying [\(1\)](#page-1-1), [\(2\)](#page-2-3) and [\(3\)](#page-2-4) such that for any choice of (x_1, \ldots, x_N) , we build \tilde{f} satisfying [\(1\)](#page-1-1), [\(2\)](#page-2-3) and [\(3\)](#page-2-4), $f(x_i) = \tilde{f}(x_i)$ and $|q_{\alpha}(f, X) - q_{\alpha}(\tilde{f}, X)| \geqslant C\rho^{N}$ ($\rho \in (0, 1)$) or $CN^{-\frac{1}{d-1}}$. This property is combined with the following observation: for any measurable $g : \mathbb{R}^N \to \mathbb{R}$,

$$
|g(f(x^{1}),...,f(x^{N})) - q_{\alpha}(f,X)| = |g(\tilde{f}(x^{1}),..., \tilde{f}(x^{N})) - q_{\alpha}(f,X)|
$$

\n
$$
\geq |q_{\alpha}(\tilde{f},X) - q_{\alpha}(f,X)| - |g(\tilde{f}(x^{1}),..., \tilde{f}(x^{N})) - q_{\alpha}(\tilde{f},X)|,
$$

it then follows that

$$
\max(|g(f(x^1),\ldots,f(x^N))-q_\alpha(f,X)|,|g(\tilde{f}(x^1),\ldots,\tilde{f}(x^N))-q_\alpha(\tilde{f},X)|)\geqslant\frac{1}{2}|q_\alpha(\tilde{f},X)-q_\alpha(f,X)|.
$$

It means that for any choice of function (we could call algorithm in this case) *g* applied to observations of the functions f at N points, considering every f satisfying (1) , (2) and (3) , the error studied in Theorems [2.1](#page-4-0) and [3.1](#page-7-0) is lower bounded by $C \rho^N$ ($\rho \in (0,1)$) or $CN^{-\frac{1}{d-1}}$

4.1. **Case** $d > 1$. In this Section, we are interested in the optimality of the polynomial convergence obtained in Theorems [2.1](#page-4-0) and [3.1](#page-7-0) in the case $d > 1$.

Proposition 4.1. Let $g : \mathbb{R}^N \to \mathbb{R}$ be a measurable function. Then, there exists $C > 0$ such that for any $(x^1, \ldots, x^N) \in ([0,1]^d)^N$, there exists f, X, α, M, L_f such that $(1), (2)$ $(1), (2)$ $(1), (2)$ and (3) hold and that

$$
|g(f(x^1),...,f(x^N))-q_{\alpha}(f,X)| \geq CN^{\frac{1}{d-1}}.
$$

Remark 4.1. This result shows that the rate of convergence (when N tends to infinity) obtained in Theorems [2.1](#page-4-0) and [3.1](#page-7-0) in the case *d >* 1 is optimal. It means that using any other algorithm than ours may only improve (lower in this case) the constant *C* obtained in our deterministic upper bound of the error, but not the polynomial rate $N^{\frac{1}{d-1}}$.

Proof. Let us consider $X \sim \mathcal{U}_{[0,1]^d}$ a uniform random variable on $[0,1]^d$ and introduce $f : [0,1]^d \to \mathbb{R}$ such that $f(x_1, \ldots, x_d) = x_1$, and choose $\alpha = \frac{1}{2}$. In this case $q_\alpha(f, X) = \frac{1}{2}$. Assumptions [\(1\)](#page-1-1) with $L_f = 1$ and [\(3\)](#page-2-4) with $K = 1$ are straightforward to verify. Let us verify assumption [\(2\)](#page-2-3) for $f :$ Let $\delta > 0$. Since $q_{\alpha}(f, X) = \alpha = \frac{1}{2}$, we have

$$
\lambda_{\text{Leb}}(f(x) \in [q_{\alpha}(f, X) - \delta, q_{\alpha}(f, X) + \delta]) = \lambda_{\text{Leb}}(x_1 \in [\frac{1}{2} - \delta, \frac{1}{2} + \delta]) = 2\delta
$$

and [\(2\)](#page-2-3) holds with $M = 2$. We now propose a construction for \tilde{f} . We introduce

$$
\mathcal{C} = \{x \in [0,1]^d, f(x) = \frac{1}{2}\} = \{x \in [0,1]^d, x_1 = \frac{1}{2}\}.
$$

We now consider the case $N = \frac{3^{j(d-1)}}{3}$ $\frac{a-1}{3}$, and define

$$
\hat{D}^j = \{D^j_\beta, D^j_\beta \cap \mathcal{C} \neq \emptyset\}.
$$

and

$$
\tilde{D}^j = \{D^j_\beta, D^j_\beta \cap \mathcal{C} \neq \emptyset, \bigcap_{i=1}^N x^i \notin D^j_\beta\}.
$$

In particular *j* is defined in a way such that $\text{Card}(\hat{D}^j) = 3^{j(d-1)} = 3N$ and then $\text{Card}(\tilde{D}^j) \geq 2N$. We are now in a position to introduce \tilde{f} . For every $x \in [0,1]^d$,

$$
\tilde{f}(x) = f(x) + 2 \sum_{D_{\beta}^j \in \tilde{D}^j} \inf_{y \in \Omega \setminus D_{\beta}^j} |x - y|_{\infty}
$$

with $|x-y|_{\infty} := \sup_{i \in \{1,\ldots,d\}} |x_i - y_i|$. Remark that functions f and $h: x \mapsto \inf_{y \in \Omega \setminus D_{\beta}^j} |x - y|_{\infty}$, are 1-Lipschitz so that [\(1\)](#page-1-1) holds with $L_{\tilde{f}} = 3$. Let us now show that [\(2\)](#page-2-3) holds for \tilde{f} with $\alpha = \frac{1}{2}$ and $M = 2^d + 1$. First we introduce the following decomposition

$$
\begin{split} \lambda_{\mathrm{Leb}}(\tilde{f}(x)\in[\frac{1}{2}-\delta,\frac{1}{2}+\delta])&=\sum_{D_{\beta}^j\notin\tilde{D}^j}\lambda_{\mathrm{Leb}}(\tilde{f}(x)\in[\frac{1}{2}-\delta,\frac{1}{2}+\delta],x\in D_{\beta}^j)\\ &+\sum_{D_{\beta}^j\in\tilde{D}^j}\lambda_{\mathrm{Leb}}(\tilde{f}(x)\in[\frac{1}{2}-\delta,\frac{1}{2}+\delta],x\in D_{\beta}^j)\\ =&\lambda_{\mathrm{Leb}}(x_1\in[\frac{1}{2}-\delta,\frac{1}{2}+\delta],x\in\cup_{D_{\beta}^j\notin\tilde{D}^j}D_{\beta}^j)\\ &+\sum_{D_{\beta}^j\in\tilde{D}^j}\lambda_{\mathrm{Leb}}(\tilde{f}(x)\in[\frac{1}{2}-\delta,\frac{1}{2}+\delta],x\in D_{\beta}^j)\\ \leqslant&2\delta+\lambda_{\mathrm{Leb}}(2h(x)\in[-\delta,\delta],x\in\cup_{D_{\beta}^j\in\tilde{D}^j}D_{\beta}^j). \end{split}
$$

Now, we notice that, since for $x \in \bigcup_{D^j_\beta \in \tilde{D}^j} D^j_\beta$, we have $2h(x) \leqslant 3^{-j}$, and then for $3^{-j} \leqslant \delta$,

$$
\begin{aligned} \lambda_{\operatorname{Leb}}(2h(x) \in [-\delta, \delta], &\, x \in \cup_{D^j_{\beta} \in \tilde{D}^j} D^j_{\beta}) \leqslant & \lambda_{\operatorname{Leb}}(\cup_{D^j_{\beta} \in \tilde{D}^j} D^j_{\beta}) \\ \leqslant & 3N 3^{-jd} = 3^{-j} \leqslant \delta. \end{aligned}
$$

Now, let us consider $3^{-j} \geq \delta$. We have, for $D^j_\beta \in \tilde{D}^j$ and $x \in D^j_\beta$,

$$
h(x) = \min_{i=1,\dots,d} \min(x_i - \frac{1}{2}(1 - 3^{-j}), \frac{1}{2}(1 + 3^{-j}) - x_i)
$$

It follows that

$$
\lambda_{\text{Leb}}(2h(x) \in [-\delta, \delta], x \in D_{\beta}^{j}) = \lambda_{\text{Leb}}(x \in D_{\beta}^{j}) - \lambda_{\text{Leb}}(x \in D_{\beta}^{j}, x_{i} \in (\frac{1}{2}(1 - 3^{-j} + \delta), \frac{1}{2}(1 + 3^{-j} - \delta)))
$$

\n
$$
= 3^{-jd} - (3^{-j} - \delta)^{d}
$$

\n
$$
= d\delta 3^{-j(d-1)} + \sum_{k=1}^{d-2} 3^{-jk} (-\delta)^{d-k} {d \choose k}
$$

\n
$$
= d\delta 3^{-j(d-1)} - 3^{-j(d-1)} \delta \sum_{k=1}^{d-2} 3^{-j(k+1-d)} (-\delta)^{d-k-1} {d \choose k}
$$

\n
$$
\leq d\delta 3^{-j(d-1)} + 3^{-j(d-1)} \delta (2^{d} - 1 - d).
$$

Therefore, using $(a+b)^d \leq 2^d(a^d + b^d)$

$$
\begin{aligned} \lambda_{\mathrm{Leb}}(2h(x)\in[-\delta,\delta],&x\in\cup_{D^j_\beta\in\tilde{D}^j}D^j_\beta)\leqslant&3N(3^{-jd}-(3^{-j}-\delta)^d)\\ =&3^{j(d-1)}(3^{-jd}-(3^{-j}-\delta)^d)\\ \leqslant&(2^d-1)\delta\end{aligned}
$$

which completes the proof of [\(2\)](#page-2-3) for \tilde{f} , $\alpha = \frac{1}{2}$ and $M = 2^d + 1$.

Moreover since for every $i \in \{1,\ldots,N\}$, $x^i \notin D^j_\beta$, for $D^j_\beta \in \tilde{D}^j$, we have $f(x^i) = \tilde{f}(x^i)$. In order to conclude the proof, Let us show that $|q_\alpha(f, X) - q_\alpha(\tilde{f}, X)| \geq CN^{-\frac{1}{d-1}}$. To do so, we prove that, there exists *C* which does not depend on *N* such that

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + CN^{-\frac{1}{d-1}}) < \frac{1}{2}
$$

This implies that $q_{\frac{1}{2}}(\tilde{f},X) > \frac{1}{2} + CN^{-\frac{1}{d-1}}$ which is the expected conclusion. From now, let us consider the fixed value $C = \frac{1}{\sqrt{2}}$ $\frac{1}{2\times 3^{\frac{1}{d-1}}}$. Using that \tilde{f} is 3-Lipschitz for the $|.|_{\mathbb{R}^d}$ -norm, we have

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + CN^{-\frac{1}{d-1}}) = \sum_{D_{\beta}^{j} \in D^{j}} \mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D_{\beta}^{j}) \mathbb{P}(X \in D_{\beta}^{j})
$$

\n
$$
= \sum_{D_{\beta}^{j} \in D^{j} \setminus \tilde{D}^{j}} \mathbb{P}(X_{1} \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D_{\beta}^{j}) \mathbb{P}(X \in D_{\beta}^{j})
$$

\n
$$
+ \sum_{D_{\beta}^{j} \in D^{j}} \mathbb{P}(X_{1} + 2 \inf_{y \in \Omega \setminus D_{\beta}^{j}} | X - y |_{\infty} \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D_{\beta}^{j}) \mathbb{P}(X \in D_{\beta}^{j})
$$

\n
$$
\leq \sum_{D_{\beta}^{j} \in D^{j} \setminus \tilde{D}^{j}} \mathbb{P}(X_{1} \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D_{\beta}^{j}) \mathbb{P}(X \in D_{\beta}^{j})
$$

\n
$$
+ \sum_{D_{\beta}^{j} \in \tilde{D}^{j}} \mathbb{P}(\tilde{f}(d_{\beta}^{j}) - 3 | X - d_{\beta}^{j} |_{\mathbb{R}^{d}} \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D_{\beta}^{j}) \mathbb{P}(X \in D_{\beta}^{j})
$$

For $D^j_\beta \notin \hat{D}^j$, we have $|d^j_{\beta_1} - \frac{1}{2}| \geqslant 3^{-j}$ and then, since $CN^{-\frac{1}{d-1}} \leqslant \frac{3^{-j}}{2}$ $\frac{1}{2}$,

$$
\mathbb{P}(X_1 \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D^j_\beta) = \mathbf{1}_{d^j_{\beta_1} < \frac{1}{2}}
$$

and

$$
\sum_{D_{\beta}^j\in D^j\backslash \hat{D}^j}\mathbb{P}(X_1\leqslant \frac{1}{2}+CN^{-\frac{1}{d-1}}|X\in D_{\beta}^j)\mathbb{P}(X\in D_{\beta}^j)=\mathbb{P}(X_1\leqslant \frac{1}{2}-\frac{3^{-j}}{2})=\frac{1-3^{-j}}{2}
$$

It follows that

$$
\sum_{\substack{D_{\beta}^j \in D^j \setminus \tilde{D}^j}} \mathbb{P}(X_1 \leq \frac{1}{2} + CN^{-\frac{1}{d-1}} | X \in D_{\beta}^j) \mathbb{P}(X \in D_{\beta}^j) \leq \frac{1 - 3^{-j}}{2} + N 3^{-jd}
$$

$$
= \frac{1 - 3^{-j}}{2} + \frac{3^{-j}}{3} = \frac{1 - 3^{-j-1}}{2}.
$$

Moreover, for $D^j_\beta \in \tilde{D}^j$,

$$
\tilde{f}(d_{\beta}^{j}) = d_{\beta_1}^{j} + 2 \inf_{y \in \Omega \setminus D_{\beta}^{j}} |d_{\beta}^{j} - y|_{\infty} = \frac{1}{2} + \frac{1}{3^{j}}.
$$

It follows that

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + CN^{-\frac{1}{d-1}}) + \sum_{D_{\beta}^{j} \in \tilde{D}^{j}} \mathbb{P}(3|X - d_{\beta}^{j}|_{\mathbb{R}^{d}} \geq 3^{-j} - CN^{-\frac{1}{d-1}}|X \in D_{\beta}^{j}) \mathbb{P}(X \in D_{\beta}^{j})
$$

$$
\leq \frac{1 - 3^{-j-1}}{2} + \sum_{D_{\beta}^{j} \in \tilde{D}^{j}} \min(1, 3^{jd}3^{-d}(3^{-j} - CN^{-\frac{1}{d-1}})^{d})3^{-jd}
$$

$$
\leq \frac{1 - 3^{-j-1}}{2} + 3N3^{-d}3^{-jd}
$$

$$
= \frac{1}{2} - \frac{1}{18}3^{-j} < \frac{1}{2}.
$$

and

$$
\mathbb{P}(\tilde{f}(X) \leqslant \frac{1}{2} + C N^{-\frac{1}{d-1}}) < \frac{1}{2}
$$

This implies that $q_{\frac{1}{2}}(\tilde{f},X) > \frac{1}{2} + CN^{-\frac{1}{d-1}}$ (with $C = \frac{1}{2 \times 3!}$ $\frac{1}{2\times 3^{\frac{1}{d-1}}}$ which is the expected conclusion. 4.2. **Case** $d = 1$. In this Section, we are interested in the optimality of the exponential convergence obtained in Theorems [2.1](#page-4-0) and [3.1](#page-7-0) in the case $d = 1$.

Proposition 4.2. Let $g : \mathbb{R}^N \to \mathbb{R}$ be a measurable function. Then, there exists $C > 0$ such that for any $(x^1, ..., x^N) \in ([0, 1]^d)^N$, there exists f, X, α, M, L_f such that [\(1\)](#page-1-1), [\(2\)](#page-2-3) and [\(3\)](#page-2-4) hold and $\rho \in (0, \frac{3}{4})$ satisfying

$$
|g(f(x^1),...,f(x^N))-q_{\alpha}(f,X)|\geqslant C\rho^N.
$$

Remark 4.2*.* This result shows that the exponential convergence obtained for the upper bound of the error studied in Theorems [2.1](#page-4-0) and [3.1](#page-7-0) in the case $d = 1$ is optimal in the sense that Proposition [4.2](#page-11-0) provides a lower bound for this error which also converges with exponential rate.

Proof. Similarly as in the case $d > 1$, we consider $X \sim \mathcal{U}_{[0,1]}$ a uniform random variable on [0, 1] and introduce $f : [0,1] \to \mathbb{R}$ such that $f(x) = x$, and choose $\alpha = \frac{1}{2}$ so that $q_{\alpha}(f, X) = \frac{1}{2}$. Recall that assumptions [\(1\)](#page-1-1) with $L_f = 1$ and [\(3\)](#page-2-4) with $K = 1$ and [\(2\)](#page-2-3) with $M = 2$ are satisfied. We now propose a construction for \tilde{f} .

For $q \in (0,1)$, let us define $I^{N+1} = \left[\frac{1}{2}(1-q^N), \frac{1}{2}(1+q^N)\right]$, $I^j = I^j_- \cup I^j_+ = \left[\frac{1}{2}(1-q^{j-1}), \frac{1}{2}(1-q^j)\right] \cup$ $\left[\frac{1}{2}(1+q^j), \frac{1}{2}(1+q^{j-1})\right], j \in \{1, \ldots, N\},\$ let us define

$$
\hat{D}^N = \{I^j, j \in \{1, \ldots, N+1\}\}.
$$

and

$$
\tilde{D}^N = \{I \in \hat{D}^N, \bigcap_{i=1}^N x^i \notin I\}.
$$

We observe that $\text{Card}(\hat{D}^N) = N + 1$ and then $\text{Card}(\tilde{D}^j) \geq 1$. We are now in a position to introduce \tilde{f} . Let us consider some $I_0 \in \tilde{D}^N$ and $\mathbf{L} > 0$ and define, for every $x \in [0,1]^d$ and

$$
\tilde{f}(x) = f(x) + \mathbf{L} \inf_{y \in \Omega \setminus I_0} |x - y|
$$

Remark that functions f and $x \mapsto \inf_{y \in \Omega \setminus I_0} |x - y|$, are 1-Lipischitz so that [\(1\)](#page-1-1) holds with $L_{\tilde{f}} = L + 1$. Let us now show that [\(2\)](#page-2-3) holds for \tilde{f} with $\alpha = \frac{1}{2}$ and $M = 2\frac{L+1}{L}$.

$$
\lambda_{\text{Leb}}(\tilde{f}(x) \in [\frac{1}{2} - \delta, \frac{1}{2} + \delta]) = \sum_{j=1; I_j \neq I_0} \lambda_{\text{Leb}}(\tilde{f}(x) \in [\frac{1}{2} - \delta, \frac{1}{2} + \delta], x \in I_j)
$$

$$
+ \lambda_{\text{Leb}}(\tilde{f}(x) \in [\frac{1}{2} - \delta, \frac{1}{2} + \delta], x \in I_0)
$$

$$
= \lambda_{\text{Leb}}(x \in [\frac{1}{2} - \delta, \frac{1}{2} + \delta], x \in \Omega \setminus I_0)
$$

$$
+ \lambda_{\text{Leb}}(\tilde{f}(x) \in [\frac{1}{2} - \delta, \frac{1}{2} + \delta], x \in I_0)
$$

$$
\leq 2\delta + \lambda_{\text{Leb}}(\text{L}h(x) \in [-\delta, \delta], x \in I_0).
$$

First, we remak that for $I_0 = I_j$, if $\frac{\delta}{\mathbf{L}} \geqslant q^j \frac{1-q}{4q}$, then

$$
\lambda_{\text{Leb}}(\mathbf{L}h(x) \in [-\delta, \delta], x \in I_0) = \lambda_{\text{Leb}}(x \in I_j) = q^j \frac{1-q}{2q} \leq 2 \frac{\delta}{\mathbf{L}}
$$

and if $I_0 = I_{N+1}$, if $\frac{\delta}{\mathbf{L}} \geqslant \frac{q^N}{2}$ $\frac{1}{2}$, then

$$
\lambda_{\operatorname{Leb}}(\mathbf{L} h(x) \in [-\delta, \delta], x \in I_0) = \lambda_{\operatorname{Leb}}(x \in I_{N+1}) = q^N \leqslant 2\frac{\delta}{\mathbf{L}}
$$

Now, we notice that for $x \in I_0 = I_j$, then $h(x) = \min(h(x) \mathbf{1}_{x \in I_j} , h(x) \mathbf{1}_{x \in I_{j,+}})$ with

$$
h(x)\mathbf{1}_{x\in I_{j,-}} = \min(x - (c_{I_{j,-}} - q^j \frac{1-q}{4q}), c_{I_{j,-}} + q^j \frac{1-q}{4q} - x)\mathbf{1}_{x\in I_{j,-}}
$$

with $c_{I_{j,-}} = \frac{1}{2} - q^j \frac{1+q}{4q}$, and similarly

$$
h(x)\mathbf{1}_{x\in I_{j,+}} = \min(x - (c_{I_{j,+}} - q^j \frac{1-q}{4q}), c_{I_{j,+}} + q^j \frac{1-q}{4q} - x)\mathbf{1}_{x\in I_{j,+}}
$$

 $\frac{1}{L}$

with $c_{I_{j,+}} = \frac{1}{2} + q^j \frac{1+q}{4q}$. It follows that, if $\frac{\delta}{\mathbf{L}} \leqslant q^j \frac{1-q}{4q}$, λ _{Leb}(**L***h*(*x*) ∈ [−*δ*, *δ*]*,x* ∈ *I*_j) = λ _{Leb}(*x* ∈ *I*_j) – λ _{Leb}(*x* ∈ (*c*_{*I*_j−</sup> − $q^j \frac{1-q}{4q}$} $\frac{\textstyle -q}{\textstyle 4q}+\frac{\textstyle \delta}{\textstyle \textbf{I}}$ $\frac{\delta}{\mathbf{L}}, c_{I_{j,-}} + q^j \frac{1-q}{4q}$ $\frac{\textstyle -q}{\textstyle 4q}-\frac{\textstyle \delta}{\textstyle \textstyle 1}$ $- \lambda$ _{Leb}(*x* ∈ (*c*_{*I*_{*j*,+} − *q*^j $\frac{1-q}{4q}$} $\frac{\textstyle -q}{\textstyle 4q}+\frac{\textstyle \delta}{\textstyle \textbf{L}}$ $\frac{\delta}{\mathbf{L}}, c_{I_{j,+}} + q^j \frac{1-q}{4q}$ $\frac{\textstyle -q}{\textstyle 4q}-\frac{\textstyle \delta}{\textstyle \textstyle 1}$ $\frac{1}{L}$ $=q^{j} \frac{1-q}{1-q}$ $\frac{q}{q} - 4q^{j} \frac{1-q}{4q}$ $\frac{-q}{4q} + 4\frac{\delta}{\mathbf{L}} = 4\frac{\delta}{\mathbf{L}}.$

Similarly, if $I_0 = I_{N+1}$, and $x \in I_{N+1}$,

$$
h(x) = \min(x - \frac{1}{2}(1 - q^N), \frac{1}{2}(1 + q^N) - x).
$$

It follows that, if $\frac{\delta}{L} \leq \frac{q^N}{2}$ $\frac{1}{2}$,

$$
\lambda_{\text{Leb}}(\text{L}h(x) \in [-\delta, \delta], x \in I_{N+1}) = \lambda_{\text{Leb}}(x \in I_{N+1}) - \lambda_{\text{Leb}}(x \in (\frac{1}{2}(1 - q^N) + \frac{\delta}{\text{L}}, \frac{1}{2}(1 + q^N) - \frac{\delta}{\text{L}}))
$$

$$
= q^N - q^N + 2\frac{\delta}{\text{L}} = 2\frac{\delta}{\text{L}}.
$$

and the proof of [\(2\)](#page-2-3) is completed.

We also observe that we have $f(x^i) = \tilde{f}(x^i)$. In order to conclude the proof, we now show that for any $\rho \in (0,1)$, we may find $\mathbf{L} \in (\frac{1+\rho}{1-\rho}, +\infty)$ and $C > 0$ (which do not depend on *N*) such that $|q_{\alpha}(f, X) - q_{\alpha}(\tilde{f}, X)| \geq C \rho^{N}$. To do so, we prove that,

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + C\rho^N) < \frac{1}{2}
$$

.

Using that \tilde{f} is \mathbf{L} + 1-Lipschitz for the $|.|_{\mathbb{R}^d}$ -norm, we have, for $I_0 \in \{I_1, \ldots I_N\}$,

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + C\rho^N) = \mathbb{P}(X \leq \frac{1}{2} + C\rho^N, X \in \Omega \setminus I_0)
$$

+
$$
\mathbb{P}(X + \mathbf{L} \inf_{y \in \Omega \setminus I_0} |X - y| \leq \frac{1}{2} + C\rho^N, X \in I_0)
$$

$$
\leq \mathbb{P}(X \leq \frac{1}{2} + C\rho^N, X \in \Omega \setminus I_0)
$$

+
$$
\mathbb{P}(\tilde{f}(c_{I_{0,-}}) - (\mathbf{L} + 1)|X - c_{I_{0,-}}|_{\mathbb{R}^d} \leq \frac{1}{2} + C\rho^N, X \in I_{0,-})
$$

+
$$
\mathbb{P}(\tilde{f}(c_{I_{0,+}}) - (\mathbf{L} + 1)|X - c_{I_{0,+}}|_{\mathbb{R}^d} \leq \frac{1}{2} + C\rho^N, X \in I_{0,+})
$$

where $c_{I_{0,-}} = \frac{\inf\{x \in I_{0,-}\} + \sup\{x \in I_{0,-}\}}{2}$ (and similarly for $c_{I_{0,+}}$) is the center point of $I_{0,-}$. When $I_0 = I_{N+1}$, we have similarly

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + C\rho^N) \leq \mathbb{P}(X \leq \frac{1}{2} + C\rho^N, X \in \Omega \setminus I_{N+1}) + \mathbb{P}(\tilde{f}(c_{I_{N+1}}) - (\mathbf{L} + 1)|X - c_{I_{N+1}}|_{\mathbb{R}^d} \leq \frac{1}{2} + C\rho^N, X \in I_{N+1})
$$

For $I_j \in \tilde{D}^N$, we have,

$$
\mathbb{P}(X \leq \frac{1}{2} + C\rho^N, X \in \Omega \setminus I_j) = \mathbb{P}(X \leq \frac{1}{2} + C\rho^N) - \mathbb{P}(X \leq \frac{1}{2} + C\rho^N, X \in I_j)
$$

\n
$$
\leq \frac{1}{2} + C\rho^N - \mathbb{P}(X \in [\frac{1}{2}(1 - q^N), \frac{1}{2}(1 + \min(q^N, 2C\rho^N))])\mathbf{1}_{j=N+1}
$$

\n
$$
- \mathbb{P}(X \in I_{j,-})\mathbf{1}_{j \neq N+1}
$$

\n
$$
= \frac{1}{2} + C\rho^N - \frac{1}{2}(\min(q^N, 2C\rho^N) + q^N)\mathbf{1}_{j=N+1} - q^{j-1}\frac{1-q}{2}\mathbf{1}_{j \neq N+1}
$$

Moreover, for $I_0 = I_j \in \{I_1, \ldots I_N\}$ $\tilde{f}(c_{I_{j,-}}) = c_{I_{j,-}} + \mathbf{L} \inf_{y \in \Omega \setminus I_{j,-}} |c_{I_{j,-}} - y| = \frac{1}{2}$ $\frac{1}{2} - q^{j-1} \frac{1+q}{4}$ $\frac{+ q}{4} + \mathbf{L}q^{j-1}\frac{1-q}{4}$ $\frac{-q}{4} = \frac{1}{2}$ $\frac{1}{2} + \frac{L(1-q) - 1 - q}{4}$ $\frac{y-1-q}{4}q^{j-1}.$ and similarly $\tilde{f}(c_{I_{j,+}}) = \frac{1}{2} + \frac{L(1-q)+1+q}{4}$ $\frac{q^{j+1+q}}{4}q^{j-1}$. Also, we have (recall that $c_{I_{N+1}} = \frac{1}{2}$) $\tilde{f}(c_{I_{N+1}}) = \frac{1}{2} + \mathbf{L} \inf_{y \in \Omega \setminus I_{N+1}} |\frac{1}{2}|$ $\frac{1}{2} - y = \frac{1}{2}$ $\frac{1}{2} + {\bf L}_{\frac{1}{2}}^{\frac{1}{2}}$ $\frac{1}{2}q^{N}$.

It follows that, for $I_0 = I_j \in \{I_1, ..., I_N\}$,

$$
\mathbb{P}(\tilde{f}(c_{I_{j,-}}) - (\mathbf{L} + 1)|X - c_{I_{j,-}}|_{\mathbb{R}^d} \leq \frac{1}{2} + C\rho^N |X \in I_{j,-})
$$

$$
\leq \mathbb{P}(|X - c_{I_{j,-}}|_{\mathbb{R}^d} \geq \frac{1}{\mathbf{L} + 1} (\frac{\mathbf{L}(1 - q) - 1 - q}{4} q^{j-1} - C\rho^N) |X \in I_{j,-})
$$

$$
= (1 - \frac{2}{q^{j-1}(1-q)} \frac{2}{\mathbf{L} + 1} (\frac{\mathbf{L}(1 - q) - 1 - q}{4} q^{j-1} - C\rho^N) +).
$$

with notation $(a)_+ = \max(0, a)$, and then, using the Bayes formula,

$$
\begin{split} \mathbb{P}(\tilde{f}(c_{I_{j,-}}) - (\mathbf{L} + 1)|X - c_{I_{j,-}}|_{\mathbb{R}^d} &\leq \frac{1}{2} + C\rho^N, X \in I_{j,-}) \\ &\leq \frac{1}{2} q^{j-1} (1-q)(1 - \frac{2}{q^{j-1}(1-q)} \frac{2}{\mathbf{L} + 1} (\frac{\mathbf{L}(1-q) - 1 - q}{4} q^{j-1} - C\rho^N)_{+})_{+} \\ &\leq \frac{1}{2} q^{j-1} (1-q)(1 - \frac{\mathbf{L}(1-q) - 1 - q}{(1-q)(\mathbf{L} + 1)} + \frac{4C\rho^N}{(1-q)(\mathbf{L} + 1)} q^{-j+1}) \\ &= q^{j-1} (\frac{1 + 2C\rho^N q^{-j+1}}{\mathbf{L} + 1}) \end{split}
$$

and similarly

$$
\begin{split} \mathbb{P}(\tilde{f}(c_{I_{j,+}}) - (\mathbf{L}+1)|X - c_{I_{j,+}}|_{\mathbb{R}^d} &\leqslant \frac{1}{2} + C\rho^N, X \in I_{j,+}) \\ &\leqslant \frac{1}{2}q^{j-1}(1-q)(1-\frac{2}{q^{j-1}(1-q)}\frac{2}{\mathbf{L}+1}(\frac{\mathbf{L}(1-q)+1+q}{4}q^{j-1}-C\rho^N)_+)_{+} \\ &\leqslant \frac{1}{2}q^{j-1}(\frac{-2q+4C\rho^Nq^{-j+1}}{\mathbf{L}+1})_{+}. \end{split}
$$

In the same way, for $I_0 = I_{N+1}$

$$
\mathbb{P}(\tilde{f}(c_{I_{N+1}}) - (\mathbf{L} + 1)|X - c_{I_{N+1}}|_{\mathbb{R}^d} \leq \frac{1}{2} + C\rho^N, X \in I_{N+1}) \leq \mathbb{P}(|X - c_{I_{N+1}}|_{\mathbb{R}^d} \geq \frac{1}{\mathbf{L} + 1}(\frac{\mathbf{L}}{2}q^N - C\rho^N), X \in I_{N+1})
$$

$$
= (q^N - \frac{2}{\mathbf{L} + 1}(\frac{\mathbf{L}}{2}q^N - C\rho^N))_{+})_{+}
$$

$$
\leq \frac{1}{\mathbf{L} + 1}q^N + \frac{2}{\mathbf{L} + 1}C\rho^N.
$$

so we conclude that, for $I_0 = I_j \in \{I_1, \ldots, I_N\}$,

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + C\rho^N) \leq \frac{1}{2} + C\rho^N - q^{j-1}\frac{1-q}{2} + q^{j-1}\frac{1}{L+1} + \frac{4}{L+1}C\rho^N
$$

$$
\leq \frac{1}{2} + \frac{L+5}{L+1}C\rho^N - q^{j-1}\frac{(L+1)(1-q) - 2}{2(L+1)}.
$$

and for $I_0 = I_{N+1}$,

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + C\rho^N) \leq \frac{1}{2} + C\rho^N - \frac{1}{2}q^N + \frac{1}{L+1}q^N + \frac{2}{L+1}C\rho^N.
$$

= $\frac{1}{2} + \frac{L+3}{L+1}C\rho^N - \frac{L-1}{2(L+1)}q^N.$

We fix *q* ∈ (0, 1) and **L** ∈ ($\frac{1+q}{1-q}$, +∞) so that $\frac{L-1}{2(L+1)} > 0$ and $(L+1)(1-q) - 2 > 0$. Now, if *C* satisfies

$$
C < \min\left(\frac{(L+1)(1-q)-2}{2q(L+5)}, \frac{L-1}{2(L+3)}\right)
$$

and $\rho \leqslant q$, then

$$
\mathbb{P}(\tilde{f}(X) \leq \frac{1}{2} + C\rho^N) < \frac{1}{2},
$$

which is the expected conclusion.

5. Numerical illustration

To conclude this article, we propose a numerical illustration of Theorem [2.1](#page-4-0) and Theorem [3.1.](#page-7-0) In both cases, we propose an application for $d = 1$ and $d = 2$. When $d = 1$ we expect to observe an exponential convergence and when $d = 2$, we expect a polynomial convergence of order $\frac{1}{N}$. Let us present our examples.

5.1. **Exponential convergence.** First for $d = 1$, we consider $X \sim \mathcal{N}(\frac{1}{5}, \frac{1}{25})\mathbf{1}_{[0,1]}$ a Gaussian distribution with mean $\frac{1}{5}$, variance $\frac{1}{25}$ and restricted to the interval [0, 1]. The function *f* is defined on $x \in [0, 1]$ by

$$
f(x) = 0.8x - 0.3 + \exp(-11.534x^{1.95}) + \exp(-2(x - 0.9)^{2})
$$

This function is studied in [\[2\]](#page-16-15) for the approximation of failure probabilities. In particular, we have $L_f \approx 1.61$. We fix $\alpha = 0.999$. In this case $q(f, X) \approx 1.3503$.

In Figures [1](#page-15-1) (A) and (B) , we represent respectively

(5.1)
$$
\ln(|q_{\alpha}(f, X) - q_{\alpha}^{N}(f, X)|)
$$

and

(5.2)
$$
\ln(|q_{\alpha}(f, X) - q_{\alpha}^{\diamond, N}(f, X)|)
$$

resulting from Algorithm [2.1](#page-3-0) and [3.1](#page-6-1) *w.r.t. N*. Those quantities appear in blue while the red line represents the linear approximation with slope respectively given by ln(0*.*8453) for Figure [1](#page-15-1) (A) and by ln(0*.*9988) for Figure [1](#page-15-1) (B). In particular we can infer the numerical approximation for *ρ* given by $\rho \approx 0.8453$ for Algorithm [2.1](#page-3-0) and $\rho \approx 0.9988$ for Algorithm [3.1](#page-6-1)

FIGURE 1. Logarithm of the error of estimation of Algorithm [2.1](#page-3-0) (see Figure (A)) and [3.1](#page-6-1) (see Figure (B)) *w.r.t.* the number of observations.

The drawings align with the results from Theorem [2.1](#page-4-0) and Theorem [3.1](#page-7-0) as a linearly decreasing behavior is observed in both cases. We thus conclude that the exponential nature of convergence is confirmed for both algorithms. Nevertheless, in this practical case, the numerical convergence of Algorithm [2.1](#page-3-0) is faster. This observation is expected as in the unknown Lipschitz case of Algorithm [3.1](#page-6-1) we consider multiple Lipschitz constant candidates and apply for each of them a similar implementation of Algorithm [2.1](#page-3-0) but with a budget which is simply a proportion of the global budget *N* allocated to all the candidates. Finally, we point out that the piecewise constant shape is also expected for both algorithms. This is due to the fact that our algorithms need a certain budget to discretize more deeply the space [0*,* 1] *i.e.* to pass from depth of discretization k to depth $k + 1$. While this budget is not attained the algorithms always return the same result.

□

5.2. **Polynomial convergence.** In a second step, we propose an example when $d = 2$. In this case, we consider $X = (X_1, X_2)$ where X_1 and X_2 are independent and identically distributed under $\mathcal{U}_{[0,1]}$, the uniform distribution on [0, 1]. The function *f* is defined on $x \in [0,1]^2$ by $f(x) = x_1 + x_2$. In this toy example, *f* is Lipschitz with Lipschitz constant $L_f = \sqrt{2}$, $f(X)$ follows the Irwin Hall distribution of degree 2 and we know that $q_\alpha(f, X) = 2 - \sqrt{2(1 - \alpha)}$. As before, we fix $\alpha = 0.999$ and we have $q_\alpha(f, X) \approx 1.9553$.

In Figures [2](#page-16-16) (A) and (B), we represent respectively the quantities (5.1) and (5.2) resulting from Algo-rithm [2.1](#page-3-0) and [3.1](#page-6-1) $w.r.t.$ ln(*N*). Those quantities appear in blue while the red line represents the linear approximation with slope respectively given by -1.4781 for Figure [2](#page-16-16) (A) and by -0.7440 for Figure [2](#page-16-16) (B).

FIGURE 2. Logarithm of the error of estimation of Algorithm [2.1](#page-3-0) (see Figure (A)) and [3.1](#page-6-1) (see Figure (B)) *w.r.t.* the Logarithm of the number of observations.

The expected linear decreasing behavior with slope $1 = \frac{1}{d-1}$ is thus observed (to some extent) aligning with the result from Theorem [2.1](#page-4-0) and Theorem [3.1.](#page-7-0) Similarly as when $d = 1$, we observe that even if the nature of convergence is similar for both algorithm, the numerical approximations converge faster when the Lipschitz constant of *f* is known, that is when we can implement Algorithm [2.1.](#page-3-0)

REFERENCES

- [1] F. Bellini, B. Klar, A. Müller, and E. Rosazza Gianin. Generalized quantiles as risk measures. *Insurance: Mathematics and Economics*, 54:41–48, 2014.
- [2] L. Bernard, A. Cohen, A. Guyader, and F. Malrieu. Recursive Estimation of a Failure Probability for a Lipschitz Function. *The SMAI Journal of computational mathematics*, 8:75–97, 2022.
- [3] C. Cannamela, J. Garnier, and B. Iooss. Controlled stratification for quantile estimation. *The Annals of Applied Statistics*, 2(4):1554–1580, 2008.
- [4] A. Cohen, R. Devore, G. Petrova, and P. Wojtaszczyk. Finding the minimum of a function. *Methods and Applications of Analysis*, 20(4):365–382, 2014.
- [5] F. Cérou and A. Guyader. Fluctuation analysis of adaptive multilevel splitting. *The Annals of Applied Probability*, 26(6):3319–3380, 2016.
- [6] F. Cérou, A. Guyader, and M. Rousset. Adaptive multilevel splitting: Historical perspective and recent results. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 29(4):043108, 04 2019.
- [7] P. Del Moral, T. Furon, and A. Guyader. Sequential monte carlo for rare event estimation. *Statistics and Computing*, 22(3):795–808, 2012.
- [8] F. Y. Edgeworth. The mathematical theory of banking. *Journal of the Royal Statistical Society*, 51(1):113–127, 1888.
- [9] D. Egloff and M. Leippold. Quantile estimation with adaptive importance sampling. *Annals of Statistics*, 38:1244–1278, 2010.
- [10] P. W. Glynn. Importance sampling for monte carlo estimation of quantiles. 2011.
- [11] B. Goffinet and D. Wallach. Optimised importance sampling quantile estimation. *Biometrika*, 83(4):791–800, 1996.
- [12] M.I. Gomes and D. Pestana. A sturdy reduced-bias extreme quantile (var) estimator. *Journal of the American Statistical Association*, 102(477):280–292, 2007.
- [13] M. B. Greenwald and S. Khanna. *Quantiles and Equi-depth Histograms over Streams*, pages 45–86. Springer Berlin Heidelberg, Berlin, Heidelberg, 2016.
- [14] A. Guyader, N Hengartner, and E. Matzner-Lober. Simulation and estimation of extreme quantiles and extreme probabilities. *Applied Mathematics & Optimization*, 64(2):171–196, 2011.
- [15] M. Hallin, D. Paindaveine, and M Siman. Multivariate quantiles and multiple-output regression quantiles: from l1 optimization to halfspace depth. *Annals of Statistics*, 38:635–669, 2010.
- [16] T. C. Hesterberg and B. L. Nelson. Control variates for probability and quantile estimation. *Management Science*, 44(9):1295–1312, 1998.
- [17] J. C. Hsu and B. L. Nelson. Control variates for quantile estimation. *Management Science*, 36(7):835–851, 1990.
- [18] R. Koenker. Quantile regression [m]. *Econometric Society Monographs, Cambridge University Press, Cambridge*, 2005.
- [19] J. Lee, C. Tepedelenlioğlu, A. Spania, and G. Muniraju. Distributed quantiles estimation of sensor network measurements. 7:38–61, 2020.
- [20] Q. Pan, E. Byon, Young M. Ko, and H. Lam. Adaptive importance sampling for extreme quantile estimation with stochastic black box computer models. *Naval Research Logistics (NRL)*, 67(7):524–547, 2020.
- [21] A. Tambwekar, A. Maiya, S. Dhavala, and S. Saha. Estimation and applications of quantiles in deep binary classification. *IEEE Transactions on Artificial Intelligence*, 3:275–286, 2021.
- [22] S. Ullah, R. Luo, T. Sunday Adebayo, and M. T. Kartal. Dynamics between environmental taxes and ecological sustainability: evidence from top-seven green economies by novel quantile approaches. *Sustainable Development*, 31(2):825–839, 2023.
- [23] H. Yang and Y. Zhao. Smoothed jackknife empirical likelihood for the one-sample difference of quantiles. *Comput. Stat. Data Anal.*, 120:58–69, 2018.