

ON THE ITERATES OF THE LAGUERRE OPERATOR

SMILJANA JAKŠIĆ, STEVAN PILIPOVIĆ, NENAD TEOFANOV, AND ĐORĐE VUČKOVIĆ

ABSTRACT. We use the iterates of the Laguerre operator to introduce Pilipović spaces on positive orthants. It is shown that such spaces coincide with G -type spaces $g_\alpha^\alpha(\mathbb{R}_+^d)$ and $G_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha > 1$, and $\alpha \geq 1$, respectively. However, in contrast to G -type spaces, Pilipović spaces are nontrivial below the critical index $\alpha = 1$. We also remark that there is a natural isomorphism between subspaces of Pilipović spaces on \mathbb{R}^d consisting of even functions, and Pilipović spaces on positive orthants.

1. INTRODUCTION

Counterparts of Schwartz functions and of Gelfand-Shilov type functions on $(0, \infty)$ were studied in [1, 2]. These results were thereafter generalized in [6], where the test spaces of ultradifferentiable functions of Roumieu and Beurling type, denoted by $G_\alpha^\alpha(\mathbb{R}_+^d)$, and $g_\alpha^\alpha(\mathbb{R}_+^d)$, $\alpha \geq 1$, respectively, are introduced and characterized via decay properties of their coefficients in Laguerre expansions.

Toft in [10] studied Pilipović spaces $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$, $\alpha > 0$, by considering the iterates of the linear Harmonic oscillator. These spaces are non trivial even if $0 < \alpha < \frac{1}{2}$, whereas when $\alpha \geq \frac{1}{2}$ they coincide with the Gelfand-Shilov spaces usually denoted by $\mathcal{S}_\alpha^\alpha(\mathbb{R}^d)$, see e.g. [4]. Recently, Toft, Bhimani and Manna demonstrated that Pilipović spaces on \mathbb{R}^d , and their distribution spaces naturally appear in the study of well-posedness of certain Schrödinger equations, which are ill-posed in the class of Gelfand-Shilov type distributions, see [11].

In this report we use the iterates of the Laguerre operator to introduce Pilipović spaces on positive orthants, and relate them with G -type spaces $g_\alpha^\alpha(\mathbb{R}_+^d)$ and $G_\alpha^\alpha(\mathbb{R}_+^d)$ of ultradifferentiable functions when $\alpha > 1$, and $\alpha \geq 1$, respectively.

The proofs will be given in our forthcoming contribution [8], where we also study natural isomorphisms between Pilipović spaces on \mathbb{R}^d consisting of even functions and Pilipović spaces on \mathbb{R}_+^d . Related questions in the context of G -type spaces were considered in [7].

1.1. Notation. By \mathbb{N} , \mathbb{Z} , \mathbb{R} and \mathbb{C} the sets of positive integers, integers, real and complex numbers, respectively; $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\mathbb{N}_0^d = (\mathbb{N} \cup \{0\})^d$, $\mathbb{R}_+ = (0, \infty)$, $\mathbb{R}_+^d = (0, \infty)^d$ and $\overline{\mathbb{R}_+^d} = [0, \infty)^d$. The standard multi-index notation will be used: for $\alpha \in \mathbb{N}_0^d$ and $x \in \mathbb{R}^d$ (or $x \in \overline{\mathbb{R}_+^d}$), $|\alpha| := \sum_{i=1}^d \alpha_i$, $x^\alpha = x_1^{\alpha_1} \dots x_d^{\alpha_d}$, and $\partial^\alpha = \partial^{x_1^{\alpha_1}} \dots \partial^{x_d^{\alpha_d}}$. If $x, \gamma \in \overline{\mathbb{R}_+^d}$, then $x^\gamma = x_1^{\gamma_1} \dots x_d^{\gamma_d}$, and if $x_j = 0$ and $\gamma_j = 0$, $j = 1, \dots, d$, we use the convention $0^0 = 1$.

2010 *Mathematics Subject Classification.* 46F05; 33C45.

Key words and phrases. Laguerre expansions; Laguerre operator; ultradifferentiable functions; Pilipović spaces.

We use the standard notation for common spaces of functions, distributions, and sequences. For example, ℓ_p , $p \geq 1$, denotes the Banach space of Lebesgue sequences, endowed with the usual norm:

$$\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_p \quad \iff \quad \|a_n\|_{\ell^p} = \left(\sum_n |a_n|^p \right)^{1/p} < \infty,$$

and with the usual modification when $p = \infty$.

A measurable function belongs to the Hilbert space $L^2(\mathbb{R}_+^d)$ if

$$\|\cdot\|_{L^2(\mathbb{R}_+^d)} := \left(\int_{\mathbb{R}_+^d} |f(x)|^2 dx \right)^{1/2} < \infty,$$

and the scalar product is given in the usual way. By $\mathcal{S}(\mathbb{R}_+^d)$ we denote the space of all smooth functions on positive orthant, $f \in \mathcal{C}^\infty(\mathbb{R}_+^d)$ such that all of its derivatives $D^p f$, $p \in \mathbb{N}_0^d$, extend to continuous functions on $\overline{\mathbb{R}_+^d}$, and

$$\sup_{x \in \mathbb{R}_+^d} x^k |D^p f(x)| < \infty, \quad k, p \in \mathbb{N}_0^d.$$

With this system of seminorms $\mathcal{S}(\mathbb{R}_+^d)$ becomes an (F) -space, see e.g. [5].

2. PRELIMINARIES

2.1. Sequence spaces. We use the notation from [10].

Let $\alpha > 0$, $h > 1$, and put $\vartheta_{h,\alpha}(n) = e^{h|n|^{1/(2\alpha)}}$, $n \in \mathbb{N}_0^d$. By $\ell_{[\vartheta_{h,\alpha}]}^p(\mathbb{N}_0^d)$, $p \geq 1$, we denote the space of all complex sequences $\{a_n\}_{n \in \mathbb{N}_0^d}$ for which

$$\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h,\alpha}]}^p(\mathbb{N}_0^d)} := \|\{|a_n| \cdot \vartheta_{h,\alpha}(n)\}_{n \in \mathbb{N}_0^d}\|_{\ell^p} < \infty. \quad (1)$$

Equipped with this norm $\ell_{[\vartheta_{h,\alpha}(n)]}^p(\mathbb{N}_0^d)$ becomes a Banach space, with the usual modification when $p = \infty$.

We also introduce

$$\ell_\alpha(\mathbb{N}_0^d) = \bigcup_{h>0} \ell_{[\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d), \quad \ell_{0,\alpha}(\mathbb{N}_0^d) = \bigcap_{h>0} \ell_{[\vartheta_{h,\alpha}]}^\infty(\mathbb{N}_0^d),$$

and endow these locally convex spaces with inductive and projective limit topology, respectively.

For consistency, if $\alpha = 0$, then $\ell_0(\mathbb{N}_0^d)$ denotes the space of all sequences $\{a_n\}_{n \in \mathbb{N}_0^d}$ such that $a_n \neq 0$ for only finitely many $n \in \mathbb{N}_0^d$, and $\ell_{0,0}(\mathbb{N}_0^d) = \{0\}$.

Observe that different choices of $p \in [1, \infty]$ give rise to equivalent norms in (1), cf [10]. In particular, if $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h_1,\alpha}]}^\infty(\mathbb{N}_0^d)$ then there exists $h_1 > 0$ and $C_1 > 0$ such that

$$\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_1,\alpha}]}^2(\mathbb{N}_0^d)} = \left(\sum_{n \in \mathbb{N}_0^d} |a_n|^2 \vartheta_{h_1,\alpha}(n) \right)^{1/2} \leq C_1 \|\{a_n \cdot \vartheta_{h,\alpha}(n)\}_{n \in \mathbb{N}_0^d}\|_{\ell^\infty},$$

and conversely, if

$$\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{[\vartheta_{h,\alpha}]}^2(\mathbb{N}_0^d)$$

then there exist $C_2, h_2 > 0$ such that

$$\|\{a_n\}_{n \in \mathbb{N}_0^d}\|_{\ell_{[\vartheta_{h_2, \alpha}]^\infty}(\mathbb{N}_0^d)} \leq C_2 \|\{a_n \cdot \vartheta_{h, \alpha}(n)\}_{n \in \mathbb{N}_0^d}\|_{l^2}.$$

2.2. Laguerre functions and Laguerre operator. For $j \in \mathbb{N}_0$ and $\gamma \geq 0$, the j -th Laguerre polynomial of order γ is defined by

$$L_j^\gamma(x) = \frac{x^{-\gamma} e^x}{j!} \frac{d^j}{dt^j} (e^{-x} x^{\gamma+j}), \quad x \geq 0.$$

The j -th Laguerre function is then given by $l_j(x) = L_j(x) e^{-x/2}$, $x \geq 0$. The d -dimensional Laguerre polynomials and Laguerre functions, are given by tensor products $L_n(x) = \prod_{l=1}^d L_{n_l}(x_l)$, and $l_n(x) = \prod_{l=1}^d l_{n_l}(x_l)$, $n \in \mathbb{N}_0^d$, respectively. We are interested in the case $\gamma = 0$, and we write L_n and l_n instead of L_n^0 and l_n^0 , respectively.

Laguerre functions $\{l_n\}_{n \in \mathbb{N}_0^d}$ form an orthonormal basis in $L^2(\mathbb{R}_+^d)$, cf. e.g. [9, 12]. Thus every $f \in L^2(\mathbb{R}_+^d)$ can be expressed via its Laguerre series expansion, and by $a_n(f)$ we denote the corresponding Laguerre coefficients:

$$a_n(f) = \int_{\mathbb{R}_+^d} f(x) l_n(x) dx, \quad n \in \mathbb{N}_0^d.$$

Similarly as it is done in [10], we may consider the spaces of formal Laguerre series expansions $f = \sum_{n \in \mathbb{N}_0^d} a_n l_n$ that correspond to the sequence $\{a_n\}_{n \in \mathbb{N}_0^d}$ from $\ell_\alpha(\mathbb{N}_0^d)$ or $\ell_{0, \alpha}(\mathbb{N}_0^d)$, $\alpha \geq 0$. These spaces will be denoted by $\mathcal{G}_\alpha(\mathbb{R}_+^d)$ and $\mathcal{G}_{0, \alpha}(\mathbb{R}_+^d)$, respectively, and the natural mapping

$$T : \ell_\alpha(\mathbb{N}_0^d) \rightarrow \mathcal{G}_\alpha(\mathbb{R}_+^d) \quad (T : \ell_{0, \alpha}(\mathbb{N}_0^d) \rightarrow \mathcal{G}_{0, \alpha}(\mathbb{R}_+^d)),$$

$$T(\{a_n\}_{n \in \mathbb{N}_0^d}) = \sum_{n \in \mathbb{N}_0^d} a_n l_n \quad (2)$$

induces topology on the target space.

Next we introduce the Laguerre operator:

$$E = - \sum_{j=1}^d \left(x_j \frac{\partial^2}{\partial x_j^2} + \frac{\partial}{\partial x_j} - \frac{x}{4} + \frac{1}{2} \right).$$

Then the Laguerre functions are the eigenfunctions of E :

$$E(l_n(x)) = \left(\sum_{i=1}^d n_i \right) \cdot l_n(x) = |n| \cdot l_n(x), \quad x > 0,$$

see [3, (11), p.188], and for $N \in \mathbb{N}$, and $n \in \mathbb{N}_0^d$ we have

$$E^N(l_n(x)) = |n|^N l_n(x). \quad (3)$$

3. PILIPOVIĆ SPACES ON POSITIVE ORTHANTS

Before we define spaces of ultradifferentiable functions in terms of the iterates of the Laguerre operator we give a simple but useful lemma.

Lemma 3.1. *Let u be a smooth and square integrable function on \mathbb{R}_+^d , i.e. $u \in L^2(\mathbb{R}_+^d) \cap C^\infty(\mathbb{R}_+^d)$. If $E^N u \in L^2(\mathbb{R}_+^d)$ for every $N \in \mathbb{N}$, then $u \in \mathcal{S}(\mathbb{R}_+^d)$.*

The proof follows from (3) and the characterization of $\mathcal{S}(\mathbb{R}_+^d)$ via coefficients in Laguerre expansions of its elements.

Definition 3.2. *Let $h > 0$, and $\alpha > 0$. Then a smooth functions f belongs to the space $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$, if*

$$\eta_h^\alpha(f) := \sup_{N \in \mathbb{N}} \frac{\|E^N f\|_{L^2(\mathbb{R}_+^d)}}{h^{|N|} N!^\alpha} < \infty. \quad (4)$$

From Lemma 3.1 it immediately follows that $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d) \subset \mathcal{S}(\mathbb{R}_+^d)$. $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$ is a Banach space with the norm η_h^α , and clearly, if $0 < \alpha < \beta$, then $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$ is continuously injected into $\mathbf{G}_{\beta,h}^{\beta,h}(\mathbb{R}_+^d)$.

The Laguerre functions l_p , $p \in \mathbb{N}_0^d$, satisfy (4) for every $h > 0$ and $\alpha > 0$, since

$$\sup_{N \in \mathbb{N}} \frac{\|E^N l_p\|_{L^2(\mathbb{R}_+^d)}}{h^{|N|} N!^\alpha} = \sup_{N \in \mathbb{N}} \frac{\binom{|p|}{h}^N}{N!^\alpha} < \infty.$$

It follows that $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$ are non-trivial spaces since they contain at least finite linear combinations of the Laguerre functions.

We notice that even if $\alpha = 0$, and if $h > 1$, then $l_p \in \mathbf{G}_{0,h}^{0,h}(\mathbb{R}_+^d)$ for all $p \in \mathbb{N}^d$ such that $|p| \leq h$.

Now we are ready to introduce Pilipović spaces on positive orthants.

Definition 3.3. *Let $\alpha > 0$. Then the Pilipović spaces on positive orthant \mathbb{R}_+^d , $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ are defined to be the union, respectively the intersection of $\mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d)$, with respect to $h > 0$:*

$$\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) = \bigcup_{h>0} \mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d), \quad \text{and} \quad \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) = \bigcap_{h>0} \mathbf{G}_{\alpha,h}^{\alpha,h}(\mathbb{R}_+^d).$$

The spaces $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ are endowed with projective and inductive limit topologies, respectively. Thus, $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ are (F) -spaces continuously injected into $\mathcal{S}(\mathbb{R}_+^d)$, and if $0 < \alpha < \beta$ we have

$$\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) \hookrightarrow \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) \hookrightarrow \mathbf{g}_\beta^\beta \hookrightarrow \mathbf{G}_\beta^\beta(\mathbb{R}_+^d), \quad (5)$$

where \hookrightarrow denotes dense and continuous inclusion.

Next we state our main result.

Theorem 3.4. *Let $\alpha > 0$, $f \in L^2(\mathbb{R}_+^d) \cap C^\infty(\mathbb{R}_+^d)$, and let $\{a_n(f)\}_{n \in \mathbb{N}_0^d}$ be the sequence of Laguerre coefficients of f . Then the following is true:*

- (i) $f \in \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ if and only if $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{\alpha/2}(\mathbb{N}_0^d)$
- (ii) $f \in \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$ if and only if $\{a_n(f)\}_{n \in \mathbb{N}_0^d} \in \ell_{0,\alpha/2}(\mathbb{N}_0^d)$.

The proof will be given in a separate contribution, [8]

By Theorem 3.4 it follows that $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) = \mathcal{G}_\alpha(\mathbb{R}_+^d)$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) = \mathcal{G}_{0,\alpha}(\mathbb{R}_+^d)$ where $\mathcal{G}_\alpha(\mathbb{R}_+^d)$ and $\mathcal{G}_{0,\alpha}(\mathbb{R}_+^d)$ are introduced in subsection 2.2. Moreover, the mapping $T : \ell_\alpha(\mathbb{N}_0^d) \rightarrow \mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d)$ (resp. $T : \ell_{0,\alpha}(\mathbb{N}_0^d) \rightarrow \mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d)$) is a well defined topological isomorphism, see (2) for the definition of T .

3.1. G -type spaces as Pilipović spaces. Following [6], we define G -type spaces of ultradifferentiable functions $G_\alpha^\alpha(\mathbb{R}_+^d)$ and $g_\alpha^\alpha(\mathbb{R}_+^d)$ as follows:

Let $A > 0$. By $g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d)$, we denote the space of all $f \in \mathcal{S}(\mathbb{R}_+^d)$ such that

$$\sup_{p,k \in \mathbb{N}_0^d} \frac{\|x^{(p+k)/2} \partial^p f(x)\|_{L^2(\mathbb{R}_+^d)}}{A^{|p+k|} k^{(\alpha/2)k} p^{(\alpha/2)p}} < \infty.$$

With the following seminorms

$$\sigma_{A,j}^{\alpha,\alpha}(f) = \sup_{p,k \in \mathbb{N}_0^d} \frac{\|x^{(p+k)/2} \partial^p f(x)\|_{L^2(\mathbb{R}_+^d)}}{A^{|p+k|} k^{(\alpha/2)k} p^{(\beta/2)p}} + \sup_{\substack{|p| \leq j \\ |k| \leq j}} \sup_{x \in \mathbb{R}_+^d} |x^k D^p f(x)|, \quad j \in \mathbb{N}_0,$$

it becomes an (F) -space.

The spaces $G_\alpha^\alpha(\mathbb{R}_+^d)$ and $g_\alpha^\alpha(\mathbb{R}_+^d)$ are then defined as union, respectively intersection of $g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d)$ with respect to A :

$$G_\alpha^\alpha(\mathbb{R}_+^d) = \bigcup_{A>0} g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d), \quad g_\alpha^\alpha(\mathbb{R}_+^d) = \bigcap_{A>0} g_{\alpha,A}^{\alpha,A}(\mathbb{R}_+^d). \quad (6)$$

Then $G_\alpha^\alpha(\mathbb{R}_+^d)$ and $g_\alpha^\alpha(\mathbb{R}_+^d)$ are endowed with inductive, respectively projective limit topology.

We note that $G_\alpha^\alpha(\mathbb{R}_+^d)$ is non-trivial when $\alpha \geq 1$, and $g_\alpha^\alpha(\mathbb{R}_+^d)$ is non-trivial when $\alpha > 1$.

Next, we rephrase and summarize results from [2, Theorem 3.6] and [6, Theorem 5.7] as follows:

Theorem 3.5. *Let $f \in \mathcal{S}(\mathbb{R}_+^d)$. Then the following is true.*

- (i) *Let $\alpha \geq 1$. If $f \in G_\alpha^\alpha(\mathbb{R}_+^d)$, and $\{a_n\}_{n \in \mathbb{N}_0^d}$ is the sequence of its Laguerre coefficients, then $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_\alpha(\mathbb{N}_0^d)$.*

Conversely if $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_\alpha(\mathbb{N}_0^d)$, then there exists $f \in G_\alpha^\alpha(\mathbb{R}_+^d)$ such that $\{a_n\}_{n \in \mathbb{N}_0^d}$ is the sequence of its Laguerre coefficients.

- (ii) *Let $\alpha > 1$. If $f \in g_\alpha^\alpha(\mathbb{R}_+^d)$, and $\{a_n\}_{n \in \mathbb{N}_0^d}$ is the sequence of its Laguerre coefficients, then $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{0,\alpha}(\mathbb{N}_0^d)$.*

Conversely if $\{a_n\}_{n \in \mathbb{N}_0^d} \in \ell_{0,\alpha}(\mathbb{N}_0^d)$, then there exists $f \in g_\alpha^\alpha(\mathbb{R}_+^d)$ such that $\{a_n\}_{n \in \mathbb{N}_0^d}$ is the sequence of its Laguerre coefficients.

By Theorems 3.5 and 3.4 we have the following:

Corollary 3.6. *G-type spaces coincide with Pilipović spaces whenever the former are non-trivial. More precisely, $\mathbf{G}_\alpha^\alpha(\mathbb{R}_+^d) = G_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha \geq 1$ and $\mathbf{g}_\alpha^\alpha(\mathbb{R}_+^d) = g_\alpha^\alpha(\mathbb{R}_+^d)$, when $\alpha > 1$.*

Therefore, G -type spaces can be also defined by the iterates of the Laguerre operator, which gives a new insight into their structure.

It should be noticed that $\mathbf{g}_1^1(\mathbb{R}_+^d)$ is of a particular interest, since it is the largest Pilipović space on orthant which is not a G -type space.

ACKNOWLEDGEMENT

This research was supported by the Science Fund of the Republic of Serbia, #GRANT No. 2727, *Global and local analysis of operators and distributions - GOALS*. The first author is supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia Grant No. 451-03-65/2024-03/ 200169. The second author is supported by the project F10 financed by the Serbian Academy of Sciences and Arts. The third author is supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Grants No. 451-03-66/2024-03/200125 & 451-03-65/2024-03/200125).

REFERENCES

- [1] Duran A. J.: Laguerre expansions of Tempered distributions and Generalized functions, *J. Math. Anal. Appl.* **150** 166–180 (1990)
- [2] Duran, A. J.: Laguerre expansions of Gel'fand-Shilov spaces, *J. Approx. Theory* **74**, 280–300 (1993)
- [3] Erdelyi, A.: *Higher Transcendentals Function*, Vol. 2, McGraw-Hill, New York (1953)
- [4] I. M. Gel'fand, G. E. Shilov, *Generalized Functions Volume 2 - Spaces of Fundamental and Generalized functions - Academic Press*, (1968)
- [5] Jakšić, S., Prangoski, B.: Extension theorem of Whitney type for $S(\mathbb{R}_+^d)$ by the use of the Kernel Theorem, *Publ. Inst. Math. Beograd*, **99**(113), 59–65 (2016)
- [6] Jakšić, S., Pilipović, S., Prangoski, B.: G -type spaces of ultradistributions over \mathbb{R}_+^d and the Weyl pseudo-differential operators with radial symbols, *RACSAM* **111**, 613–640 (2017)
- [7] Jakšić, S., Maksimović, S., Pilipović, S., Prangoski, B.: Relations between Hermite and Laguerre expansions of ultradistributions over \mathbb{R}^d and \mathbb{R}_+^d , *J. Pseudo-Differ. Oper. Appl.*, **8**, 275–296 (2017)
- [8] Jakšić, S., Pilipović, S., Teofanov, N., Vučković, Dj: Pilipović type spaces on \mathbb{R}_+^d , in preparation (2024)
- [9] Thangavelu, S.: *Lectures on Hermite and Laguerre Expansions*, Princeton University Press, Princeton New Jersey (1993)
- [10] Toft, J.: Images of function and distribution spaces under the Bargmann transform, *J. Pseudo-Differ. Oper. Appl.* **8** 83–139 (2017)
- [11] Toft, J., Bhimani, D. G. , Manna, R.: Fractional Fourier transforms, harmonic oscillator propagators and Strichartz estimates on Pilipović and modulation spaces, *Appl. Comput. Harmon. Anal.* **67**, 101580 (2023)
- [12] Wong, M. W.: *Weyl Transform*, Springer-Verlag, New York (1998)

FACULTY OF FORESTRY, UNIVERSITY OF BELGRADE, KNEZA VIŠESLAVA 1, BELGRADE, SERBIA
Email address: `smiljana.jaksic@sfb.bg.ac.rs`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF NOVI SAD, TRG DOSITEJA
OBRADOVIĆA 4, NOVI SAD, SERBIA
Email address: `stevan.pilipovic@dmi.uns.ac.rs`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, UNIVERSITY OF NOVI SAD, TRG DOSITEJA
OBRADOVIĆA 4, NOVI SAD, SERBIA
Email address: `tnenad@dmi.uns.ac.rs@dmi.uns.ac.rs`

ĐORĐE VUČKOVIĆ, TECHNICAL FACULTY “MIHAJLO PUPIN”, ĐURE ĐAKOVIĆA BB, 23000 ZREN-
JANIN, SERBIA
Email address: `djordjeplusja@gmail.com`