

Variance-reduction for Variational Inequality Problems with Bregman Distance Function

Zeinab Alizadeh^{*}, Erfan Yazdandoost Hamedani^{*}, and Afrooz Jalilzadeh^{*}

Abstract

In this paper, we address variational inequalities (VI) with a finite-sum structure. We introduce a novel single-loop stochastic variance-reduced algorithm, incorporating the Bregman distance function, and establish an optimal convergence guarantee under a monotone setting. Additionally, we explore a structured class of non-monotone problems that exhibit weak Minty solutions, and analyze the complexity of our proposed method, highlighting a significant improvement over existing approaches. Numerical experiments are presented to demonstrate the performance of our algorithm compared to state-of-the-art methods.

keywords: Variational Inequality, Variance Reduction, Bergman Distance

1 Introduction

Let $(\mathcal{X}, \|\cdot\|_{\mathcal{X}})$ be a finite-dimensional normed vector space, with dual space $(\mathcal{X}^*, \|\cdot\|_{\mathcal{X}^*})$. In this paper, consider a Variational Inequality (VI) problem to find $x^* \in X$ such that

$$\langle F(x^*), x - x^* \rangle + h(x) - h(x^*) \geq 0, \quad \forall x \in X, \quad (1)$$

where $X \subseteq \mathcal{X}$ is a nonempty, closed, and convex set; $F : \mathcal{X} \rightarrow \mathcal{X}^*$ is a Lipschitz continuous operator; and $h : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a proper convex lower-semicontinuous function. VI problems have received a lot of attention recently due to their general formulation that subsumes many well-known problems including (un)constrained optimization, saddle point (SP) problems, and Nash Equilibrium games [25, 46]. These formulations arise in various fields such as machine learning [41, 12], robotics [49], game theory [18, 35, 42, 52], etc. Developing efficient algorithms for solving large-scale VIs can immensely contribute to the computational advancement of these fields.

One of the main structures commonly arise in a large-scale setting is when the operator F consists of a large (finite) number of components, i.e., $F = \frac{1}{n} \sum_{i=1}^n F_i$. In this setting, when n is large, evaluation of the operator F (which may correspond to first-order information) can be prohibitively expensive. While stochastic methods have been extensively studied for a more general setting when the operator F is represented by the expectation of a random variable, there is a significant gap in the convergence rate guarantee compared to deterministic methods. Consequently, variance reduction techniques have been developed to fill this gap to improve the complexity of deterministic methods. While variance reduction techniques have been successfully applied to minimization problems [5, 24],

^{*}Department of Systems and Industrial Engineering, The University of Arizona, Tucson, AZ, USA.
{zalizadeh, erfany, afrooz}@arizona.edu

their naive extension to VI problems does not improve the complexity [3]. On the other hand, proximal-gradient-based methods with Bregman divergence have been shown to enhance algorithmic flexibility across various contexts, resulting in notable computational efficiencies in specific scenarios [47, 1]. The integration of Bregman distance with variance reduction for saddle point and VI problems poses an additional layer of complexity to algorithmic analysis. These challenges have been addressed by introducing retraction steps, a novel approach first implemented in the context of convex-concave saddle point problems [50] and later in [4] for monotone VIs. However, these methods capitalize on the double-loop structure inherent in the proposed algorithms to achieve optimal convergence guarantees. Consequently, this prompts the following research questions:

1. *Is it possible to develop a single-loop algorithm for solving monotone VIs with Bregman distance?*
2. *What about addressing the same in a non-monotone setting?*

In this paper, we answer the above questions. Specifically, we provide an affirmative response to the first questions by introducing a novel variance-reduced method capable of using Bregman distance and demonstrating its complexity result. Moreover, we show that under some mild assumptions, our method can also handle the case where F is non-monotone.

1.1 Related work

In the past few decades, there have been various methods developed for solving VIs with convergence rate guarantee, such as Extragradient (EG)/Mirror-Prox (MP) [23, 30], forward-backward-forward (FBF) [44], dual extrapolation [32], or reflected gradient/forward-reflected-backward (FoRB) [28]. These methods typically demonstrate a complexity of $\mathcal{O}(1/\epsilon)$ when dealing with monotone VIs in deterministic settings. The significance of stochastic VIs has grown, particularly in scenarios involving uncertainty, where methods such as Sample Average Approximation (SAA) and Stochastic Approximation (SA) come into play. SAA involves approximating the expected value of stochastic mapping by averaging over a large number of samples, while SA employs a (mini-batch) sample at each iteration. Notably, the stochastic variants of these methods are characterized by a complexity of $\mathcal{O}(1/\epsilon^2)$ when addressing monotone VIs [31, 23].

In optimization with finite-sum objectives, variance reduction techniques, pioneered by classical works such as [40, 22, 16], aim to enhance deterministic methods. They achieve this by offering unbiased estimators of gradients while minimizing the variance of the error in gradient estimation. Recently, there has been a growing interest in developing variance-reduced techniques for saddle point and VI problems which we discuss next.

SP problems: There have been various attempts to incorporate variance reduction techniques into primal-dual algorithms for solving SP problems, mostly when the objective function is strongly-convex strongly-concave such as [34, 26, 51, 14], and a few others consider a more general setting—see [27] and the reference therein. In particular, under a convex-concave setting and Euclidean normed vector space, the complexity of $\mathcal{O}(\frac{\sqrt{n}}{\epsilon} \log(1/\epsilon))$ has been shown in [27]. Later, Yazdandoost Hamedani and Jalilzadeh [50] developed stochastic variance-reduced accelerated primal-dual (SVR-APD) algorithm equipped with Bregman distance and demonstrated the complexity of $\mathcal{O}(\sqrt{n}/\epsilon)$ matching the lower-bound obtained in [20].

VI problems: Several notable papers explore the variance reduction techniques to address monotone VIs with finite-sum structure [39, 43, 4]. In particular, authors in [43] proposed a cyclic coordinate

Table 1: Compression of algorithms for monotone VI problems

Setting	Ref	Bregman	#Loop	Complexity
Deterministic	FBF [44]	✗	1	$\mathcal{O}(n/\epsilon)$
	FoRB [28]	✗		
	Prox-method [30]	✓		
	APD [19]	✓		
Stochastic	SA,SAA [31]	✓	1	$\mathcal{O}(1/\epsilon^2)$
	Mirror-Prox [23]			
Finite-sum	SVRG [50]	✓	2	$\mathcal{O}(\sqrt{n}/\epsilon)$
	Extra gradient [4]	✗	1	
	VR-MP [4]	✓	2	
	VR-CODER [43]	✗	3	
	This paper	✓	1	

dual averaging with extrapolation (CODER) method for solving VIs of the form (1). Assuming a coordinate-friendly structure and monotone operator, their method attains a complexity of $\mathcal{O}(\sqrt{m}/\epsilon)$ where m is the number of coordinate blocks. Furthermore, they proposed a variance-reduced variant of their method (VR-CODER) by combining an SVRG-type update leading to a three-loop algorithm. This method can achieve a complexity of $\mathcal{O}(\frac{\max\{\sqrt{n}, \sqrt{m}\}}{\epsilon})$ to find an ϵ -gap in monotone setting. Considering a non-Euclidean setting, a variance-reduced extra-gradient method with the Bregman distance function is proposed in [4] for the class of monotone VIs. The proposed algorithm has a double-loop structure and using a retraction step complexity of $\mathcal{O}(\sqrt{n}/\epsilon)$ to find an ϵ -gap is achieved. Furthermore, using the same technique variance-reduced version of different algorithms including forward-backward-forward and forward-reflected-backward methods in Euclidean setup is studied – see Table 1 for further comparisons of existing methods.

Exploring beyond the realm of monotone VIs, different classes of non-monotone VIs, including two-sided PL condition, pseudo monotonicity, negatively comonotonicity, cohypomonotonicity, and weak minty VI (MVI), are studied in the literature [7, 21, 48, 37, 36, 38]. Among these settings, the weak MVI assumption appears to be the most general existing assumption. With few exceptions [9, 37, 2, 36, 4], much of the existing research primarily investigates specific instances of (1), such as when $h = 0$ and $X = \mathbb{R}^n$ –as evidenced in [15, 8, 11]. Among these exceptions, [9, 37, 2] have attained a complexity of $\mathcal{O}(1/\epsilon^2)$ in the deterministic scenario, whereas solutions for (1) in a stochastic setting remain relatively scarce. Notably, [36] introduces a bias-corrected stochastic extragradient (BC-SEG+) algorithm, guaranteeing convergence with a complexity of $\mathcal{O}(1/\epsilon^4)$. In a concurrent work to ours, Alacaoglu et al. in [2] introduce multi-loop inexact variants of Halpern and Krasnoselski-Mann (KM) iterations, achieving complexities of $\mathcal{O}(\log(1/\epsilon)/\epsilon^2)$ and $\mathcal{O}(\log(1/\epsilon)/\epsilon^4)$ for deterministic and stochastic settings, respectively – see Table 2 for comparison of existing methods.

Moreover, to the best of our knowledge, no single-loop method for the monotone setting can effectively handle Bregman functions, and none of the current methods can adapt a Bregman distance function for non-monotone Variational inequalities.

Table 2: Compression of algorithms for non-monotone VI problems under weak MVI assumption

Setting	Ref	Bregman	#Loop	Complexity
Deterministic	OG[9]	✗	1	$\mathcal{O}(n^2/\epsilon^2)$
	AdaptiveEG+ [37]	✗	1	$\mathcal{O}(n^2/\epsilon^2)$
	inexact-KM[2]	✗	2	$\mathcal{O}(n^2 \log(1/\epsilon)/\epsilon^2)$
Stochastic	BC-SEG+ [36]	✗	1	$\mathcal{O}(1/\epsilon^4)$
	RAPP[38]	✗	2	$\mathcal{O}(1/\epsilon^4)$
	Inexact-KM[2]	✗	3	$\mathcal{O}(\log(1/\epsilon)/\epsilon^4)$
Finite-sum	This Paper	\checkmark (Lip.)	1	$\mathcal{O}(n/\epsilon^2)$

1.2 Contribution

Motivated by the absence of a single loop method capable of adapting a Bregman distance function to solve VIs with a finite-sum structure, we develop a unified stochastic Variance-Reduced Forward Reflected Moving Average Backward method (VR-FoRMAB) algorithm. To contend with the challenge of using a Bregman distance function with a variance reduction, we proposed a novel stochastic operator estimation as well as a new momentum term based on a nested averaging technique. In the monotone setting, we demonstrate that our proposed method can achieve the optimal complexity of $\mathcal{O}(\sqrt{n}/\epsilon)$. To the best of our knowledge, our proposed method is the first single-loop method capable of using the Bregman distance function to achieve this complexity. Furthermore, we analyze our proposed algorithm in a non-monotone setting under a weak MVI assumption. While existing deterministic algorithms exhibit a complexity that scales at n^2 with the number of operator components (i.e., achieving a complexity of $\mathcal{O}(n^2/\epsilon^2)$) their stochastic counterparts typically entail a complexity of $\mathcal{O}(1/\epsilon^4)$ and are often characterized by an intricate multi-loop structure. Bridging this gap, we show that under weak MVI assumption and Lipschitz continuity of the Bregman distance generating function, our proposed method can achieve a complexity of $\mathcal{O}(n/\epsilon^2)$ for the first time, marking a significant improvement over existing methods.

1.3 Application

From the application perspective, VIs manifest across diverse domains, spanning machine learning, signal processing, image processing, and finance. In particular, problems arising in machine learning include generative adversarial networks [17], reinforcement learning [13], and distributionally robust learning [29]. Below, we briefly outline some intriguing examples.

Distributionally robust optimization (DRO): Define $\ell_i(u) = \ell(u, \xi_i)$, where $\ell : \mathcal{X} \times \Omega \rightarrow \mathbb{R}$ is a loss function possibly nonconvex with \mathbb{P} be a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ where $\Omega = \{\xi_1, \dots, \xi_n\}$. DRO examines the worst-case performance under uncertainty to determine solutions with a particular confidence level [29]. This problem can be stated in the following way:

$$\min_{u \in \mathcal{X}} \max_{y \in \mathcal{P}} \mathbb{E}_{\xi \sim \mathbb{P}}[\ell(u, \xi_i)] = \sum_{i=1}^n y_i \ell_i(u), \quad (2)$$

where \mathcal{P} , represents the uncertainty set, for instance $\mathcal{P} = \{y \in \Delta_n : V(y, \frac{1}{n} \mathbf{1}_n) \leq \rho\}$ is an uncertainty set considered in different papers such as [29], and $V(Q, P)$ denotes the divergence measure between two sets of probability measures Q and P and $\Delta_n \triangleq \{y = [y_i]_{i=1}^n \in \mathbb{R}_+^n \mid \sum_{i=1}^n y_i = 1\}$ represents the

simplex-set. By assuming $V(y, \frac{1}{n}\mathbf{1}_n) = \sum_{i=1}^n V_i(y_i, \frac{1}{n}\mathbf{1}_n)$. The following formulation can be obtained by relaxing the constraint on divergence in equation (2)

$$\min_{\substack{u \in \mathcal{X} \\ \lambda \geq 0}} \max_{\substack{y = [y_i]_{i=1}^n \\ y \in \Delta_n}} \sum_{i=1}^n (y_i \ell_i(u) - \frac{\lambda}{n} (V_i(y_i, \frac{1}{n}\mathbf{1}_n) - \frac{\rho}{n})). \quad (3)$$

Let $x = [u^T \lambda]^T$, $L_i(x, y) = ny_i \ell_i(x) - \lambda(\frac{1}{2}V_i(y_i, \frac{1}{n}\mathbf{1}_n) - \frac{\rho}{n})$, $h_1(x) = \mathbb{I}_{X \times \mathbb{R}_+}(u, \lambda)$ and $h_2(y) = \mathbb{I}_{\Delta_n}(y)$, then the previous equation will be a special case of SP problem which can be formulated as a VI by setting $F(z) = F(x, y) = \begin{bmatrix} \nabla_x L(x, y) \\ -\nabla_y L(x, y) \end{bmatrix}$ and $h(z) = h_1(x) + h_2(y)$.

Von Neumann's Ratio Game: Consider a two-player game where the payoff function is of the form $\langle x, Ry \rangle / \langle x, Sy \rangle$, where x, y are mixed-strategy vectors and $R \in \mathbb{R}^{n \times m}$, $S \in \mathbb{R}_+^{n \times n}$ are matrices. Such a payoff function arises in stochastic games, economic models of an expanding economy, and some nonzero-sum game formulations [33]. The goal is then to solve the following min-max problem

$$\min_{x \in X} \max_{y \in \Delta_m} \frac{\langle x, Ry \rangle}{\langle x, Sy \rangle}, \quad (4)$$

where $X \subseteq \mathbb{R}^n$ is a convex and compact set, and we assume that $\langle x, Sy \rangle > 0$, for all $(x, y) \in X \times \Delta_m$. Note that problem (4) can be formulated as (1) which has a solution [33]. Moreover, it has been shown that this problem satisfies the weak MVI condition [13].

1.4 Organization of the paper

In the next section, we begin by establishing the essential notations, lemmas, and key definitions. Then, in section 3 we introduce our proposed algorithm VR-FoRMAB and provide the convergence analysis proving the main results in section 4. In particular, we discuss how VR-FoRMAB can be used to solve monotone and non-monotone VI problems. Later, in section 5 we implement our proposed method to solve instances of DRO and a two-player zero-sum game and compare them with state-of-the-art methods.

2 Preliminaries

In this section, we provide some fundamental definitions and state the lemmas that we need for convergence analysis. Throughout the paper, $\|\cdot\|_2$ denotes the Euclidean norm and $\mathbb{I}_X(\cdot)$ denotes the indicator function of set X .

Definition 1. Let $\psi_{\mathcal{X}} : \mathcal{X} \rightarrow \mathbb{R}$ be a continuously differentiable function on $\text{int}(\text{dom } h)$. In addition, $\psi_{\mathcal{X}}$ is strongly convex with respect to $\|\cdot\|_{\mathcal{X}}$. The Bregman distance function corresponding to the distance-generating function $\psi_{\mathcal{X}}$ is defined as $\mathbf{D}_X(x, \bar{x}) \triangleq \psi_{\mathcal{X}}(x) - \psi_{\mathcal{X}}(\bar{x}) - \langle \nabla \psi_{\mathcal{X}}(\bar{x}), x - \bar{x} \rangle$ for all $x \in X$ and $\bar{x} \in X^\circ \triangleq X \cap \text{int}(\text{dom } h)$. Moreover we define the Bregman diamateres $B_X \triangleq \sup_{x \in \bar{X}_R} \mathbf{D}_X(x, x_0)$.

Definition 2. For a given sequence $\{x_j\}_{j \geq 0} \subset \mathcal{X}$, an index $k \in \{1, \dots, n\}$, and $q \in \mathbb{Z}_+$, let $D_X^k(x) \triangleq \frac{1}{q} \sum_{j=(i_k-1)q+1}^{i_k q} \mathbf{D}_X(x, x_j)$ for any $x \in \mathcal{X}$ where $i_k \triangleq \lfloor k/q \rfloor$.

Lemma 1. Let $(\mathcal{U}, \|\cdot\|_{\mathcal{U}})$ be a finite-dimensional normed vector space with the dual space $(\mathcal{U}^*, \|\cdot\|_{\mathcal{U}^*})$, $f : \mathcal{U} \rightarrow \mathbb{R} \cup \{+\infty\}$ be a closed convex function, $U \subset \mathcal{U}$ is a closed convex set, and $\phi : \mathcal{U} \rightarrow \mathbb{R}$ be a distance-generating function which is continuously differentiable on an open set containing $\mathbf{dom} f$ and is l -strongly convex with respect to $\|\cdot\|_{\mathcal{U}}$, and $\mathbf{D}_U : U \times (U \cap \mathbf{dom} f) \rightarrow \mathbb{R}$ be a Bregman distance function associated with ϕ . Then, the following result holds:

a) Given $\bar{x} \in U \cap \mathbf{dom} f$, $s \in \mathcal{U}$, and $t > 0$, let $x^+ = \arg \min_{x \in U} f(x) - \langle s, x \rangle + t\mathbf{D}_U(x, \bar{x})$. Then, for all $x \in U$, the following inequality holds:

$$f(x) \geq f(x^+) + \langle s, x - x^+ \rangle + t \langle \nabla \phi(\bar{x}) - \nabla \phi(x^+), x - x^+ \rangle. \quad (5)$$

b) For all $x \in U$ and $y, z \in U \cap \mathbf{dom} f$, $\langle \nabla \phi(z) - \nabla \phi(y), x - z \rangle = \mathbf{D}_U(x, y) - \mathbf{D}_U(x, z) - \mathbf{D}_U(z, y)$.

c) Given the update of x^+ in (a), for all $x \in U$ the following inequality holds:

$$f(x^+) - f(x) + \langle s, x - \bar{x} \rangle \leq t (\mathbf{D}_U(x, \bar{x}) - \mathbf{D}_U(x, x^+)) + \frac{1}{2t} \|s\|_{\mathcal{U}^*}^2. \quad (6)$$

d) Assuming $\phi(\cdot)$ is a closed function, then $\nabla \phi^*(\nabla \phi(x)) = x$, for all $x \in \mathbf{dom} \nabla \phi \subset \mathcal{U}$; and $\nabla \phi(\nabla \phi^*(y)) = y$, for all $y \in \mathbf{dom} \nabla \phi^* \subset \mathcal{U}^*$.

Proof. For proofs see [45] and [6]. □

Lemma 2. Let $\{x_k\}_{k \geq 0} \subseteq U$ be a sequence and define $\tilde{x}_k \triangleq \frac{1}{q} \sum_{j=(i_k-1)q+1}^{i_k q} x_j$ for any $k \geq 0$. Then, for any $k \geq 0$ we have that

$$\|x_{k+1} - \tilde{x}_k\|_{\mathcal{U}}^2 \leq D_U^k(x_{k+1}), \quad (7a)$$

$$\sum_{k=0}^{K-1} D_U^k(x) \leq \sum_{k=0}^{K-1} \mathbf{D}_U(x, x_k), \quad \forall x \in U. \quad (7b)$$

Proof. Using the definition of \tilde{x}_k and the fact that for any nonnegative sequence $\{a_i\}_{i=1}^m$, $(\sum_{i=1}^m a_i)^2 \leq m \sum_{i=1}^m a_i^2$, the following can be obtained:

$$\begin{aligned} \|x_{k+1} - \tilde{x}_k\|_{\mathcal{U}}^2 &= \left\| x_{k+1} - \frac{1}{q} \sum_{j=(i_k-1)q+1}^{i_k q} x_j \right\|_{\mathcal{U}}^2 \\ &= \left\| \frac{1}{q} \sum_{j=(i_k-1)q+1}^{i_k q} (x_{k+1} - x_j) \right\|_{\mathcal{U}}^2 \\ &\leq \frac{1}{q} \sum_{j=(i_k-1)q+1}^{i_k q} \|x_{k+1} - x_j\|_{\mathcal{U}}^2. \end{aligned}$$

The result in (7a) follows from the definition of $D_U^k(x)$.

For the second inequality, by using the definition of $D_U^k(x)$, for any $x \in U$ then we have,

$$\sum_{k=0}^{K-1} D_U^k(x) = \sum_{k=0}^{K-1} \frac{1}{q} \sum_{j=[(i_k-1)q+1]_+}^{i_k q} D_U(x, x_j)$$

$$\begin{aligned}
&= \mathbf{D}_U(x, x_0) + \sum_{j=1}^q \mathbf{D}_U(x, x_j) + \cdots + \sum_{j=(i_{K-1}-1)q+1}^{i_{K-1}q} \mathbf{D}_U(x, x_j) \\
&= \sum_{j=0}^{i_{K-1}q} \mathbf{D}_U(x, x_j) \leq \sum_{k=0}^{K-1} \mathbf{D}_U(x, x_k).
\end{aligned}$$

□

3 Proposed method

In this section, we propose our novel unified variance-reduced algorithm adaptable to both monotone and non-monotone scenarios. After introducing the algorithm, we divide the convergence analysis into two main sections by laying out the underlying assumptions necessary to achieve a convergence rate guarantee. Our goal is to develop an efficient algorithm for solving (1) in which we assume that F_i 's are Lipschitz continuous. In particular, we consider the following standard assumption.

Assumption 1. For any $i \in \{1, \dots, n\}$, $F_i : \mathcal{X} \rightarrow \mathcal{X}^*$ is Lipschitz continuous, i.e., there exists $L > 0$ such that $\|F_i(x) - F_i(\bar{x})\|_{\mathcal{X}^*} \leq L \|x - \bar{x}\|_{\mathcal{X}}$ for any $x, \bar{x} \in \mathcal{X}$.

Moreover, h is possibly nonsmooth whose Bregman proximal operator, i.e., $\operatorname{argmin}_{x \in X} \{h(x) + \langle v, x \rangle + \frac{1}{\sigma} \mathbf{D}_X(x, \bar{x})\}$ for some $\bar{x}, v \in \mathcal{X}$ and $\sigma > 0$, can be computed efficiently.

Variance reduction techniques stand out as a cornerstone in revolutionizing stochastic methods and were first introduced to address large-scale finite-sum minimization problems. Several methods including SAG, SVRG, and SPIDER [40, 22, 16] among others were introduced to reduce the per-iteration cost. Unfortunately, the extension of these methods to a more general setting of using the Bregman distance function or VIs is not trivial. Among these techniques, SPIDER method generates a progressively improving sequence of gradient estimates by sampling the gradients and periodically computing the full gradient when $\operatorname{mod}(k, q) = 0$ for some integer $q > 0$. More precisely, if we let v_{k-1} be the estimator of the gradient map ∇^{k-1} at iteration k , then the next estimator can be constructed using sample gradients ∇_S^k and ∇_S^{k-1} as follows: $v_k = v_{k-1} + \nabla_S^k - \nabla_S^{k-1}$. Note that if $\mathbb{E}[v_{k-1} | x_{k-1}] = \delta^{k-1}$, then $\mathbb{E}[v_k | x_k] = \delta^k$. Inspired by this approach, we introduce a new recursive update. In particular, to accommodate the Bregman distance function we introduce a new retraction step with parameter $\gamma_k \in [0, 1]$ to find \hat{x}_k such that $\nabla \psi_{\mathcal{X}}(\hat{x}_k) = (1 - \gamma_k) \nabla \psi_{\mathcal{X}}(x_k) + \gamma_k s_k$ where s_k is a moving average of iterates in the mirror space (see line 6 of Algorithm 1). Moreover, the next iterate point is found through a Bregman proximal step by moving along a direction that is a convex combination of F at x_k and the moving average of iterates \tilde{x}_k , i.e., $(1 - \beta)F(x_k) + \beta F(\tilde{x}_k)$ plus a novel momentum based on the current and past operators denoted by $r_k \triangleq F_S(x_k) - (1 - \beta)F_S(x_{k-1}) - \beta F_S(\tilde{x}_{k-1})$. Consequently, we update the estimate of this direction denoted by v_k as follows:

$$v_k = v_{k-1} + (1 - \beta)(F_S(x_k) - F_S(x_{k-1})), \quad (8)$$

for some $\beta \in [0, 1]$. Indeed, we can show that if v_{k-1} is an unbiased estimator of $(1 - \beta)F(x_{k-1}) + \beta F(\tilde{x}_{k-1})$, then v_k is an unbiased estimator of $(1 - \beta)F(x_k) + \beta F(\tilde{x}_k)$ and the variance of error of this estimation depends on the cumulative distance of consecutive iterates – see Lemma 4. Furthermore, the direction v_k and momentum term r_k are updated using the full batch every q iteration. The steps of the proposed method are shown in Algorithm 1.

Algorithm 1 VR-ForMAB

```

1: Input:  $x_0 \in \mathcal{X}$ ,  $S \in \{1, \dots, n\}$ ,  $q \in \{1, \dots, n\}$ , and  $\beta \in [0, 1]$ 
2: for  $k = 0, \dots, K - 1$  do
3:    $i_k \leftarrow \lfloor k/q \rfloor$ 
4:   if  $\text{mod}(k, q) = 0$  then
5:      $\tilde{x}_k \leftarrow \frac{1}{q} \sum_{j=k-q+1}^k x_j$ 
6:      $s_k \leftarrow \frac{1}{q} \sum_{j=k-q+1}^k \nabla \psi_{\mathcal{X}}(x_j)$ 
7:      $v_k \leftarrow (1 - \beta)F(x_k) + \beta F(\tilde{x}_k)$ 
8:      $r_k \leftarrow F(x_k) - (1 - \beta)F(x_{k-1}) - \beta F(\tilde{x}_{k-1})$ 
9:   else
10:     $\tilde{x}_k \leftarrow \tilde{x}_{i_k q}$ 
11:    Draw  $S$  samples  $\mathcal{S} \subset \{1, \dots, n\}$  and let  $F_S \triangleq \frac{1}{S} \sum_{i \in \mathcal{S}} F_i$ 
12:     $v_k \leftarrow v_{k-1} + (1 - \beta)(F_S(x_k) - F_S(x_{k-1}))$ 
13:     $r_k \leftarrow F_S(x_k) - (1 - \beta)F_S(x_{k-1}) - \beta F_S(\tilde{x}_{k-1})$ 
14:   end if
15:    $\hat{x}_k \leftarrow \nabla \psi_{\mathcal{X}}^*((1 - \gamma_k)\nabla \psi_{\mathcal{X}}(x_k) + \gamma_k s_k)$ 
16:    $x_{k+1} \leftarrow \text{argmin}_{x \in X} \{h(x) + \langle v_k + \theta_k r_k, x \rangle + \frac{1}{\sigma_k} \mathbf{D}_X(x, \hat{x}_k)\}$ 
17: end for

```

Next, we introduce key definitions to streamline notation and establish essential properties of the variance reduction technique outlined in (8) in the consequent lemmas. These results serve as foundational elements crucial for demonstrating the convergence rate of the proposed algorithm in both monotone and non-monotone settings.

Definition 3. For any $k \geq 0$, let $r_k \triangleq F_S(x_k) - (1 - \beta)F_S(x_{k-1}) - \beta F_S(\tilde{x}_{k-1})$, $\bar{r}_k \triangleq F(x_k) - (1 - \beta)F(x_{k-1}) - \beta F(\tilde{x}_{k-1})$, and $\delta_k \triangleq v_k - ((1 - \beta)F(x_k) + \beta F(\tilde{x}_k))$.

Lemma 3. Let $\{x_k\}_{k \geq 0}$ be the sequence generated by Algorithm 1. We some define auxiliary sequence $\{w_k^x, w_k^r\} \subset \mathcal{X}$, for $k \geq 0$ as follow:

$$w_{k+1}^x \leftarrow \text{argmin}_{x \in X} \{-\langle \delta_k, x \rangle + \frac{1}{\eta_k} \mathbf{D}_X(x, w_k^x)\} \quad (9a)$$

$$w_{k+1}^r \leftarrow \text{argmin}_{x \in X} \{\langle r_k - \bar{r}_k, x \rangle + \frac{1}{\eta_k} \mathbf{D}_X(x, w_k^r)\}. \quad (9b)$$

Then, for any $k \geq 0$, $\eta_k > 0$, and $x \in X$,

$$\langle \delta_k, x - x_{k+1} \rangle \leq \frac{1}{\eta_k} (\mathbf{D}_X(x, w_k^x) - \mathbf{D}_X(x, w_{k+1}^x)) + \eta_k \|\delta_k\|_{\mathcal{X}^*}^2 + \langle \delta_k, w_k^x - x_k \rangle + \frac{1}{\eta_k} \mathbf{D}_X(x_{k+1}, x_k), \quad (10a)$$

$$\langle r_k - \bar{r}_k, x_k - x \rangle \leq \frac{1}{\eta_k} (\mathbf{D}_X(x, w_k^r) - \mathbf{D}_X(x, w_{k+1}^r)) + \frac{\eta_k}{2} \|r_k - \bar{r}_k\|_{\mathcal{Y}^*}^2 + \langle r_k - \bar{r}_k, x_k - w_k^r \rangle. \quad (10b)$$

Proof. We begin the proof of inequality in (10a) by splitting the inner product into $\langle \delta_k, x - x_k \rangle$ and $\langle \delta_k, x_k - x_{k+1} \rangle$ and providing an upper bound for each term. In particular, using lemma 1 for (9a) with $s = \delta_k$, $t = 1/\eta_k$, and $f \equiv 0$, for any $x \in X$ we conclude that

$$\langle \delta_k, x - x_k \rangle = \langle \delta_k, x - w_k^x \rangle + \langle \delta_k, w_k^x - x_k \rangle$$

$$\leq \frac{1}{\eta_k}(\mathbf{D}_X(x, w_k^x) - \mathbf{D}_X(x, w_{k+1}^x)) + \frac{\eta_k}{2} \|\delta_k\|_{\mathcal{X}^*}^2 + \langle \delta_k, w_k^x - x_k \rangle, \quad (11)$$

in addition, by using Young's inequality and the strong convexity of the Bregman distance function, we have:

$$\langle \delta_k, x_k - x_{k+1} \rangle \leq \frac{\eta_k}{2} \|\delta_k\|_{\mathcal{X}^*}^2 + \frac{1}{\eta_k} \mathbf{D}_X(x_{k+1}, x_k). \quad (12)$$

Combining the two last inequality gives (10a). The inequality in (10b) can be shown using the same line of proof. \square

Remark 1. Let $(\mathcal{U}, \|\cdot\|)$ be a finite-dimensional normed real vector space with dual space \mathcal{U}^* . Then all the norms defined in \mathcal{U} (or \mathcal{U}^*) are equivalent, i.e., for an arbitrary norm $\|\cdot\|_{\mathcal{U}}$ there exist $\underline{c}, \bar{c} > 0$ such that $\underline{c} \|\cdot\|_2 \leq \|\cdot\|_{\mathcal{U}} \leq \bar{c} \|\cdot\|_2$. This is an important property that we use in our analysis when providing a bound on the norm of the error of the gradient estimate (δ_k) .

Lemma 4. Let $\{x_k\}_{k \geq 0}$ be the sequence generated by Algorithm 1. Then, for any $k \geq 0$, $\mathbb{E}[\delta_k] = 0$ and $\mathbb{E}[r_k - \bar{r}_k] = 0$. Moreover, for $k \geq 0$

$$\mathbb{E}[\|\delta_k\|_{\mathcal{X}^*}^2] \leq 2(1 - \beta)^2 \bar{c}^2 \underline{c}^2 L^2 \sum_{t=i_k q + 1}^k \mathbb{E}[\mathbf{D}_X(x_t, x_{t-1})], \quad (13a)$$

$$\mathbb{E}[\|r_k - \bar{r}_k\|_{\mathcal{X}^*}^2] \leq 4(1 - \beta)^2 L^2 \mathbb{E}[\mathbf{D}_X(x_k, x_{k-1})] + 4\beta^2 L^2 \mathbb{E}[D_X^{k-1}(x_k)]. \quad (13b)$$

Proof. Using the definitions of δ_k and v_k in Algorithm 1, for any $k \geq 0$ and $t \in \{i_k q + 1, \dots, (i_k + 1)q - 1\}$ we have that

$$v_t = v_{t-1} + (1 - \beta)(F_S(x_t) - F_S(x_{t-1})).$$

Adding and subtracting $(1 - \beta)F(x_t) + \beta F(\tilde{x}_t)$ and defining $e_t \triangleq F_S(x_t) - F_S(x_{t-1}) - F(x_t) + F(x_{t-1})$ leads to:

$$\delta_t = \delta_{t-1} + (1 - \beta)e_t + \beta(F(\tilde{x}_{t-1}) - F(\tilde{x}_t)).$$

Therefore, summing the above relation from $t = i_k q + 1$ to k and noting that $\delta_{i_k q} = 0$ we obtain

$$\delta_k = (1 - \beta) \sum_{t=i_k q + 1}^k e_t + \beta(F(\tilde{x}_{i_k q}) - F(\tilde{x}_k)) = (1 - \beta) \sum_{t=i_k q + 1}^k e_t, \quad (14)$$

where the last equality follows from the fact that $\tilde{x}_t = \tilde{x}_{i_k q}$ for any $t \in \{i_k q + 1, \dots, (i_k + 1)q - 1\}$. Therefore, we immediately conclude that $\mathbb{E}[\delta_k] = 0$ for any $k \geq 0$.

Next, taking the norm-square from both sides of (14) and based on Remark 1 to use equivalency between the norms we can see that

$$\|\delta_k\|_{\mathcal{X}^*}^2 \leq \bar{c}^2 \|\delta_k\|_2^2 = \bar{c}^2 (1 - \beta)^2 \left\| \sum_{t=i_k q + 1}^k e_t \right\|_2^2.$$

Next, expanding the norm-square, taking expectation from both sides and using the fact that $\mathbb{E}[\langle e_j, e_\ell \rangle \mid \mathcal{F}_\ell] = 0$ for any $j < \ell \leq k$ we obtain

$$\mathbb{E}[\|\delta_k\|_{\mathcal{X}^*}^2] \leq \bar{c}^2 (1 - \beta)^2 \sum_{t=i_k q + 1}^k \mathbb{E}[\|e_t\|_2^2]. \quad (15)$$

Moreover from the definition of e_t and Assumption 1 we have that

$$\begin{aligned}\mathbb{E}[\|e_t\|_2^2] &= \mathbb{E}[\|F_S(x_t) - F(x_t) - F_S(x_{t-1}) + F(x_{t-1})\|_2^2] \\ &\leq \mathbb{E}[\|F_S(x_t) - F_S(x_{t-1})\|_2^2] \\ &\leq \underline{c}^2 L^2 \mathbb{E}[\|x_t - x_{t-1}\|_{\mathcal{X}}^2].\end{aligned}\tag{16}$$

Therefore, combining (16) with (15) and using strong convexity of the Bregman distance leads to the desired result in (13a).

Next, recall that $r_k = F_S(x_k) - (1 - \beta)F_S(x_{k-1}) - \beta F_S(\tilde{x}_{k-1})$ and $\bar{r}_k = F(x_k) - (1 - \beta)F(x_{k-1}) - \beta F(\tilde{x}_{k-1})$. It is easy to verify that $\mathbb{E}[r_k - \bar{r}_k] = 0$. Define $\tilde{e}_k \triangleq F_S(x_k) - F_S(\tilde{x}_{k-1}) + F(\tilde{x}_{k-1}) - F(x_k)$, then using the triangle inequality and Assumption 1, one can easily demonstrate that

$$\begin{aligned}\|r_k - \bar{r}_k\|_{\mathcal{X}^*} &\leq (1 - \beta) \|e_k\|_{\mathcal{X}^*} + \beta \|\tilde{e}_k\|_{\mathcal{X}^*} \\ &\leq 2(1 - \beta)L \|x_k - x_{k-1}\|_{\mathcal{X}} + 2\beta L \|x_k - \tilde{x}_{k-1}\|_{\mathcal{X}}.\end{aligned}\tag{17}$$

Then, taking a square from both sides of the above inequality and using (7a) in Lemma 2 we obtain the desired result. \square

In the following lemma, we present a one-step analysis for VR-FoRMAB, which serves as the main component for demonstrating the rate result in section 4.

Lemma 5. *Let $\{x_k\}_{k \geq 0}$ be the sequence generated by VR-FoRMAB displayed in Algorithm 1 initialized from arbitrary vectors $x_0 \in \mathcal{X}$. Let $\{w_k^x, w_k^r\}$ be the auxiliary sequence defined in (9a)-(9b). Suppose Assumption 1 holds and r_k , δ_k , and \bar{r}_k are defined in Definition 3. For any $x \in X$ and $k \geq 0$ the following result holds:*

$$\begin{aligned}h(x_{k+1}) - h(x) + \langle F(x), x_{k+1} - x \rangle \\ \leq A_k + \theta_k \langle \bar{r}_k, x - x_k \rangle + \langle \bar{r}_{k+1}, x_{k+1} - x \rangle + \frac{1 - \gamma_k}{\sigma_k} (\mathbf{D}_X(x, x_k) - \mathbf{D}_X(x, x_{k+1})) \\ + \left(\frac{1}{\eta_k} + \theta_k(\alpha_1 + \alpha_2) - \frac{1 - \gamma_k}{\sigma_k} \right) \mathbf{D}_X(x_{k+1}, x_k) \\ + \frac{\gamma_k}{\sigma_k} (D_X^k(x) - \mathbf{D}_X(x, x_{k+1}) - D_X^k(x_{k+1})) \\ + \theta_k \left(\frac{(1 - \beta)^2 L^2}{\alpha_1} + 2\eta_k(1 - \beta)^2 L^2 \right) \mathbf{D}_X(x_k, x_{k-1}) + \theta_k \left(\frac{\beta^2 L^2}{\alpha_2} + 2\eta_k \beta^2 L^2 \right) D_X^{k-1}(x_k) \\ + 2\eta_k(1 - \beta)^2 \underline{c}^2 \underline{c}^2 L^2 \sum_{t=i_k q + 1}^k \mathbf{D}_X(x_t, x_{t-1}),\end{aligned}\tag{18}$$

where $A_k \triangleq \langle \delta_k, x - x_{k+1} \rangle + \theta_k \langle r_k - \bar{r}_k, x - x_k \rangle - 2\eta_k \theta_k (1 - \beta)^2 L^2 \mathbf{D}_X(x_k, x_{k-1}) - \frac{1}{\eta_k} \mathbf{D}_X(x_{k+1}, x_k) - 2\eta_k (1 - \beta)^2 \underline{c}^2 \underline{c}^2 L^2 \sum_{t=i_k q + 1}^k \mathbf{D}_X(x_t, x_{t-1}) - 2\eta_k \theta_k \beta^2 L^2 D_X^{k-1}(x_k)$.

Proof. Applying Lemma 1 for updating the rule of x_{k+1} , implies that for any $x \in X$,

$$\begin{aligned}h(x_{k+1}) - h(x) &\leq \langle v_k + \theta_k r_k, x - x_{k+1} \rangle + \frac{1}{\sigma_k} \langle \nabla \psi_{\mathcal{X}}(x_{k+1}) - \nabla \psi_{\mathcal{X}}(\hat{x}_k), x - x_{k+1} \rangle \\ &= \langle v_k + \theta_k r_k, x - x_{k+1} \rangle \\ &\quad + \frac{1}{\sigma_k} \langle \nabla \psi_{\mathcal{X}}(x_{k+1}) - (1 - \gamma_k) \nabla \psi_{\mathcal{X}}(x_k) - \gamma_k s_k, x - x_{k+1} \rangle \\ &= \langle v_k + \theta_k r_k, x - x_{k+1} \rangle + \frac{1 - \gamma_k}{\sigma_k} \langle \nabla \psi_{\mathcal{X}}(x_{k+1}) - \nabla \psi_{\mathcal{X}}(x_k), x - x_{k+1} \rangle\end{aligned}$$

$$\begin{aligned}
& + \frac{\gamma_k^y}{\sigma_k} \langle \nabla \psi_{\mathcal{X}}(x_{k+1}) - s_k, x - x_{k+1} \rangle \\
& = \langle v_k + \theta_k r_k, x - x_{k+1} \rangle + \frac{1-\gamma_k}{\sigma_k} \langle \nabla \psi_{\mathcal{X}}(x_{k+1}) - \nabla \psi_{\mathcal{Y}}(x_k), x - x_{k+1} \rangle \\
& + \frac{\gamma_k}{\sigma_k q} \sum_{j=(i_k-1)q+1}^{i_k q} \langle \nabla \psi_{\mathcal{X}}(x_{k+1}) - \nabla \psi_{\mathcal{X}}(x_j), x - x_{k+1} \rangle.
\end{aligned}$$

Where in the first equality we used Lemma 1 part(d) and updated the rule of \hat{x}_k in Algorithm 1. Then by using the generalized three-point property of Bregman distance in Lemma 1 part (b) twice; we will have:

$$\begin{aligned}
h(x_{k+1}) - h(x) & \leq \langle v_k + \theta_k r_k, x - x_{k+1} \rangle \\
& + \frac{1-\gamma_k}{\sigma_k} (\mathbf{D}_X(x, x_k) - \mathbf{D}_X(x, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_k)) \\
& + \frac{\gamma_k}{\sigma_k q} \sum_{j=(i_k-1)q+1}^{i_k q} (\mathbf{D}_X(x, x_i) - \mathbf{D}_X(x, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_i)), \tag{19}
\end{aligned}$$

by using the definition of δ_k , and rearranging the terms, we have:

$$\begin{aligned}
h(x_{k+1}) - h(x) & \leq \langle \delta_k, x - x_{k+1} \rangle + \langle (1-\beta)F(x_k) + \beta F(\tilde{x}_k), x - x_{k+1} \rangle \\
& + \langle \theta_k r_k, x - x_k \rangle + \langle \theta_k r_k, x_k - x_{k+1} \rangle \\
& + \frac{1-\gamma_k}{\sigma_k} (\mathbf{D}_X(x, x_k) - \mathbf{D}_X(x, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_k)) \\
& + \frac{\gamma_k}{\sigma_k} (D_X^k(x) - \mathbf{D}_X(x, x_{k+1}) - D_X^k(x_{k+1})). \tag{20}
\end{aligned}$$

Now, by adding $\langle F(x_{k+1}), x_{k+1} - x \rangle$ to the both sides, and adding and subtracting $\langle \theta_k \bar{r}_k, x - x_k \rangle$ to the right-hand side of (20) one can obtain:

$$\begin{aligned}
& h(x_{k+1}) - h(x) + \langle F(x_{k+1}), x_{k+1} - x \rangle \\
& \leq \langle \delta_k, x - x_{k+1} \rangle + \theta_k \langle r_k - \bar{r}_k, x - x_k \rangle + \langle \bar{r}_{k+1}, x_{k+1} - x \rangle \\
& + \langle \theta_k \bar{r}_k, x - x_k \rangle + \langle \theta_k r_k, x_k - x_{k+1} \rangle \\
& + \frac{1-\gamma_k}{\sigma_k} (\mathbf{D}_X(x, x_k) - \mathbf{D}_X(x, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_k)) \\
& + \frac{\gamma_k}{\sigma_k} (D_X^k(x) - \mathbf{D}_X(x, x_{k+1}) - D_X^k(x_{k+1})). \tag{21}
\end{aligned}$$

Next, by using Cauchy-Schwartz inequality, followed by Assumption 1 and Young's inequality we obtain

$$\begin{aligned}
& |\langle r_k, x_k - x_{k+1} \rangle| \\
& \leq \frac{(1-\beta)^2 L^2}{2\alpha_1} \|x_k - x_{k-1}\|_{\mathcal{X}}^2 + \frac{\beta^2 L^2}{2\alpha_2} \|x_k - \tilde{x}_{k-1}\|_{\mathcal{X}}^2 + \frac{\alpha_1 + \alpha_2}{2} \|x_{k+1} - x_k\|_{\mathcal{X}}^2 \\
& \leq \frac{(1-\beta)^2 L^2}{\alpha_1} \mathbf{D}_X(x_k, x_{k-1}) + \frac{\beta^2 L^2}{\alpha_2} D_X^{k-1}(x_k) + (\alpha_1 + \alpha_2) \mathbf{D}_X(x_{k+1}, x_k), \tag{22}
\end{aligned}$$

for any $\alpha_1, \alpha_2 > 0$. Note that the left-hand side of (21) can be lower-bounded using monotonicity of F as follows $h(x_{k+1}) - h(x) + \langle F(x_{k+1}), x_{k+1} - x \rangle \geq h(x_{k+1}) - h(x) + \langle F(x), x_{k+1} - x \rangle$. Moreover, by using the previous inequality within (21) we obtain

$$h(x_{k+1}) - h(x) + \langle F(x), x_{k+1} - x \rangle$$

$$\begin{aligned}
&\leq A_k + \theta_k \langle \bar{r}_k, x - x_k \rangle + \langle \bar{r}_{k+1}, x_{k+1} - x \rangle \\
&\quad + \frac{1-\gamma_k}{\sigma_k} (\mathbf{D}_X(x, x_k) - \mathbf{D}_X(x, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_k)) \\
&\quad + \frac{\gamma_k}{\sigma_k} (D_X^k(x) - \mathbf{D}_X(x, x_{k+1}) - D_X^k(x_{k+1})) \\
&\quad + \frac{\theta_k(1-\beta)^2 L^2}{\alpha_1} \mathbf{D}_X(x_k, x_{k-1}) + \frac{\theta_k \beta^2 L^2}{\alpha_2} D_X^{k-1}(x_k) + \theta_k(\alpha_1 + \alpha_2) \mathbf{D}_X(x_{k+1}, x_k).
\end{aligned}$$

Now adding and subtracting $\frac{1}{\eta_k} \mathbf{D}_X(x_{k+1}, x_k) + 2\eta_k \theta_k (1-\beta)^2 L^2 \mathbf{D}_X(x_k, x_{k-1}) + 2\eta_k (1-\beta)^2 \bar{c}^2 \bar{c}^2 L^2 \sum_{t=i_k, q+1}^k \mathbf{D}_X(x_t, x_{t-1}) + 2\eta_k \theta_k \beta^2 L^2 D_X^{k-1}(x_k)$ to the right-hand side of the above inequality and rearranging the terms the result in (18) can obtain. \square

Remark 2. Note that the retraction parameter γ_k and damping parameter β are introduced to control the upper bounds on the operator estimates (Lemma 4). This can be seen in the result of Lemma 5 to create telescopic terms on the right-hand side of (18).

4 Convergence Analysis

4.1 Monotone VI

In this section, we demonstrate the complexity of Algorithm 1 under monotonicity of F . We first formally state the assumptions.

Assumption 2. A solution $x^* \in X$ to the VI problem in (1) exists.

Assumption 3. Assume that $F : \mathcal{X} \rightarrow \mathcal{X}^*$ is a monotone operator, i.e., for any $x, \bar{x} \in \mathcal{X}$ $\langle F(x) - F(\bar{x}), x - \bar{x} \rangle \geq 0$.

In this scenario, we consider the standard gap function defined as follows:

$$\mathcal{G}(\bar{x}) \triangleq \sup_{x \in \tilde{X}_R} \{ \langle F(x), \bar{x} - x \rangle + h(\bar{x}) - h(x) \}, \quad (23)$$

where $\tilde{X}_R \subseteq X$ is a convex and compact set containing a solution of (1) to handle the possibility of unbounded domain [32].

In the next theorem, we provide the main result of this paper by finding a bound for the expected gap function.

Theorem 1. Let $\{x_k\}_{k \geq 0}$ be the sequence generated by VR-FoRMAB displayed in Algorithm 1 initialized from arbitrary vector $x_0 \in \mathcal{X}$. Suppose Assumptions 1, 2, and 3 hold. For $k \geq 0$ let $\theta_k = 1$, $\gamma_k = \gamma \in (0, 1)$, the step-size sequence $\sigma \leq \min\{\frac{1-\gamma}{(2\sqrt{q}+4)L}, \frac{\gamma}{L\beta(1+2\beta/\sqrt{q})}\}$, and $\beta = 1 - \frac{1}{\max\{\bar{c}_1, 1\}}$. Then, for any $K \geq 1$,

$$\mathbb{E}[\mathcal{G}(\bar{x}_K)] \leq \frac{1}{K} \left(\frac{1}{\sigma} + \sqrt{q}L \right) B_X, \quad (24)$$

where $B_X \triangleq \sup_{x \in \tilde{X}_R} \mathbf{D}_X(x, x_0)$.

Proof. See Section 4.2 for the proof of the theorem.

Corollary 1. *Under the premises of Theorem 1, if the step-sizes and parameters of Algorithm 1 are selected as $q = n$, $S = 1$, $\beta = 1 - \frac{1}{\max\{\bar{c}_c, 1\}}$, $\theta_k = 1$, $\gamma_k = \beta$, $\sigma_k = \min\{\frac{1-\beta}{(2\sqrt{q}+4)L}, \frac{1}{L(1+2\beta/\sqrt{q})}\}$, then $\mathbb{E}[\mathcal{G}(\bar{x}_K)] \leq \mathcal{O}(\frac{L\sqrt{n}}{K})$. Moreover, the oracle complexity to achieve an ϵ -gap is $\mathcal{O}(\sqrt{n}L/\epsilon)$.*

Proof. Let $q = n$, then it is easy to verify that for any $\gamma \in [0, 1]$, $\sigma = \mathcal{O}(\frac{1}{\sqrt{n}L})$. From Theorem 1 we have that $\mathbb{E}[\mathcal{G}(\bar{x}_K)] \leq D/K$, where $D = (\frac{1}{\sigma} + \sqrt{q}L)B_X$, hence, we conclude that $\mathbb{E}[\mathcal{G}(\bar{x}_K)] \leq \mathcal{O}(\frac{L\sqrt{n}}{K})$. Moreover, to achieve $\mathbb{E}[\mathcal{G}(\bar{x}_K)] \leq \epsilon$ the oracle (sample) complexity can be calculated as $n\lfloor \frac{K}{q} \rfloor + S(K - \lfloor \frac{K}{q} \rfloor) = \mathcal{O}(\frac{nD}{q\epsilon} + \frac{SD}{\epsilon})$. Now by choosing $S = 1$ one can obtain $\mathcal{O}(\frac{nD}{q\epsilon} + \frac{SD}{\epsilon}) = \mathcal{O}(\sqrt{n}L/\epsilon)$ \square

\square

Remark 3. *While SVR-APD [50] and VR-MP [4] obtain the same oracle complexity $\mathcal{O}(\sqrt{n}/\epsilon)$, they are burdened with a double-loop structure. In contrast to SVR-APD, which addressed a specific min-max problem, our approach tackles a more general case from a VI perspective. Furthermore, while VR-MP algorithm requires computation of the Bregman proximal step twice within each iteration of the inner loop, ours necessitates only one computation.*

4.2 Proof of Theorem 1

Consider the result in Lemma 5 and let $\sigma_k = \sigma$, $\gamma_k = \gamma$, $\theta_k = 1$, $\alpha_1 = (1 - \beta)L$, $\alpha_2 = \beta L$, and $\eta_k = \frac{1}{(1-\beta)\bar{c}_c\sqrt{q}L}$ for $k \geq 0$, then summing the inequality over $k = 0, \dots, K-1$, dividing both sides by K , using (7b), and the fact that $(1 - \beta)\bar{c}_c \leq 1$ imply that

$$\begin{aligned}
& \frac{1}{K} \sum_{k=0}^{K-1} \left(h(x_{k+1}) - h(x) + \langle F(x), x_{k+1} - x \rangle \right) \tag{25} \\
& \leq \frac{1}{K} \left[\sum_{k=0}^{K-1} A_k - \langle \bar{r}_0, x_0 - x \rangle + \langle \bar{r}_K, x_K - x \rangle + \frac{1-\gamma}{\sigma} (\mathbf{D}_X(x, x_0) - \mathbf{D}_X(x, x_K)) \right. \\
& \quad \left. + \frac{\gamma}{\sigma} (\mathbf{D}_X(x, x_0) - \mathbf{D}_X(x, x_K)) \right] + \frac{1}{K} \sum_{k=0}^{K-1} \left(\left(1 + \frac{2\beta}{\sqrt{q}}\right) \beta L D_X^{k-1}(x_k) - \frac{\gamma}{\sigma} D_X^k(x_{k+1}) \right) \\
& \quad + \frac{1}{K} \sum_{k=0}^{K-1} \left((2\sqrt{q} + 3(1 - \beta)) L \mathbf{D}_X(x_k, x_{k-1}) + \left(L - \frac{1-\gamma}{\sigma}\right) \mathbf{D}_X(x_{k+1}, x_k) \right) \\
& \leq \frac{1}{K} \left[\sum_{k=0}^{K-1} A_k - \langle \bar{r}_0, x_0 - x \rangle + \langle \bar{r}_K, x_K - x \rangle + \frac{1-\gamma}{\sigma} (\mathbf{D}_X(x, x_0) - \mathbf{D}_X(x, x_K)) \right. \\
& \quad \left. + \frac{\gamma}{\sigma} (\mathbf{D}_X(x, x_0) - \mathbf{D}_X(x, x_K)) \right] + \frac{1}{K} \frac{\gamma}{\sigma} \left(D_X^{-1}(x_0) - D_X^{K-1}(x_K) \right) \\
& \quad + \frac{1}{K} \left(\frac{1-\gamma}{\sigma} - L \right) \left(\mathbf{D}_X(x_0, x_{-1}) - \mathbf{D}_X(x_K, x_{K-1}) \right).
\end{aligned}$$

where the last inequality follows from the fact that $\beta \in (0, 1)$ and the step-size condition $\sigma \leq \min\{\frac{1-\gamma}{(2\sqrt{q}+4)L}, \frac{\gamma}{L\beta(1+2\beta/\sqrt{q})}\}$.

Next, we provide an upper bound on $\langle \bar{r}_K, x_K - x \rangle - \langle \bar{r}_0, x_0 - x \rangle$ by using Young's inequality and noting that $\bar{r}_0 = 0$ due to the initialization of the algorithm, i.e., $x_{-1} = x_0$. Therefore,

$$\langle \bar{r}_K, x_K - x \rangle \leq (1 - \beta) L \mathbf{D}_X(x_K, x_{K-1}) + \beta L D_X^{K-1}(x_K) + L \mathbf{D}_X(x, x_K), \tag{26}$$

moreover, the left-hand side of (25) can be lower-bounded using Jensen's inequality due to the convexity of h and defining $\bar{x}_K \triangleq \frac{1}{K} \sum_{k=0}^{K-1} x_k$. Therefore, using (26) within (25) and the fact that $\mathbf{D}_X(x_0, x_{-1}) = 0$ and $D_X^{-1}(x_0) = 0$ we conclude that

$$\begin{aligned} & h(\bar{x}_K) - h(x) + \langle F(x), \bar{x}_K - x \rangle \\ & \leq \frac{1}{K} \left[\sum_{k=0}^{K-1} A_k + \frac{1}{\sigma} \mathbf{D}_X(x, x_0) + (L - \frac{1}{\sigma}) \mathbf{D}_X(x, x_K) \right. \\ & \quad \left. + ((2 - \beta)L - \frac{1 - \gamma}{\sigma}) \mathbf{D}_X(x_K, x_{K-1}) + (\beta L - \frac{\gamma}{\sigma}) D_X^{K-1}(x_K) \right]. \end{aligned} \quad (27)$$

Note that since $\beta \in [0, 1)$ and $q \geq 1$, from the stepsize condition one clearly verify that $\sigma \leq \frac{1 - \gamma}{(2 - \beta)L}$ and $\sigma \leq \frac{\gamma}{\beta L}$, hence, the last three terms in the right-hand-side of above inequality are nonpositive and can be dropped. Moreover, from the definition of A_K and Lemmas 3 and 4 we can show that

$$\begin{aligned} & \sum_{k=0}^{K-1} \mathbb{E}[A_k] \\ & = \sum_{k=0}^{K-1} \mathbb{E} \left[\langle \delta_k, x - x_{k+1} \rangle + \langle r_k - \bar{r}_k, x - x_k \rangle - 2\eta(1 - \beta)^2 \bar{c}^2 \underline{c}^2 L^2 \sum_{t=i_k q + 1}^k \mathbf{D}_X(x_t, x_{t-1}) \right. \\ & \quad \left. - \frac{1}{\eta} \mathbf{D}_X(x_{k+1}, x_k) - 2\eta(1 - \beta)^2 L^2 \mathbf{D}_X(x_k, x_{k-1}) - 2\eta\beta^2 L^2 D_X^{k-1}(x_k) \right] \\ & \leq \sum_{k=0}^{K-1} \mathbb{E} \left[\langle \delta_k, w_k^x - x_k \rangle + \langle r_k - \bar{r}_k, x_k - w_k^r \rangle \right] \\ & \quad + \frac{1}{\eta} \mathbb{E} \left[\sum_{k=0}^{K-1} (\mathbf{D}_X(x, w_k^x) - \mathbf{D}_X(x, w_{k+1}^x) + \mathbf{D}_X(x, w_k^r) - \mathbf{D}_X(x, w_{k+1}^r)) \right] \\ & \leq \frac{1}{\eta} (\mathbf{D}_X(x, w_0^x) + \mathbf{D}_X(x, w_0^r)), \end{aligned} \quad (28)$$

where the last inequality follows from unbiasedness of δ_k and $r_k - \bar{r}_k$ and dropping the non-positive terms. Therefore, combining (28) with (27) and taking supremum over $x \in \tilde{X}_R$ followed by taking the expectation of both sides of the aforementioned inequality imply that

$$\mathbb{E}[\mathcal{G}(\bar{x}_K)] \leq \frac{1}{K} \left(\frac{1}{\sigma} + \frac{1}{\eta} \right) B_X, \quad (29)$$

where $B_X = \sup_{x \in \tilde{X}_R} \mathbf{D}_X(x, x_0)$ and note that $w_0^x = w_0^r = x_0$. \square

Remark 4. [Simplified updates for structured norm] Before moving to the non-monotone setting, we would like to discuss an important special case of a monotone setting when $\bar{c} \leq \underline{c} \leq 1$. For instance, the vector space \mathcal{X} equipped with a p -norm denoted by $\|\cdot\|_p$ satisfies this condition for any $p \in [1, 2]$ – see Remark 1. In this case, one can readily verify from Corollary 1 that $\beta = \gamma = 0$ satisfies the parameter conditions. Consequently, this simplifies the updates within our proposed algorithm while maintaining the complexity of $\mathcal{O}(\sqrt{n}/\epsilon)$. This insight underscores the challenge inherent in employing the Bregman distance function with an arbitrary norm in the algorithm.

4.3 Non-monotone VIs

In this section, we analyze our proposed method in a non-monotone setting. To enable this extension, we introduce essential assumptions and lemmas for our analysis. In this scenario, we are interested in finding $\bar{x} \in X$ such that $\mathbf{0} \in F(\bar{x}) + \partial(h + \mathbb{I}_X)(\bar{x})$. Consequently, we define the following gap function to find an ϵ -approximated solution of (1).

Definition 4. $\bar{x} \in X$ is an ϵ -approximated solution when $\mathbf{dist}(\mathbf{0}, F(\bar{x}) + \partial(h + \mathbb{I}_X)(\bar{x})) \leq \epsilon$ where $\mathbf{dist}(\cdot, \cdot)$ denotes the distance function, i.e., $\mathbf{dist}(x, S) \triangleq \min_{s \in S} \|x - s\|_{\mathcal{U}}$ for a given normed vector space \mathcal{U} , a convex set $S \subset \mathcal{U}$, and a point $x \in \mathcal{U}$.

Assumption 4 (Weak Minty Solution). *There exists $x^* \in X$ and some $\rho > 0$ such that*

$$-\rho \|v\|_{\mathcal{X}^*}^2 \leq \langle v, x - x^* \rangle, \quad \forall x \in X, \forall v \in \partial(h + \mathbb{I}_X)(x) + F(x). \quad (30)$$

Assumption 5. *The Bregman distance generating function $\psi_{\mathcal{X}}$ defined in Definition 1 has a Lipschitz continuous gradient, i.e., for any $x, \bar{x} \in \mathcal{X}$ we have $\|\nabla \psi_{\mathcal{X}}(x) - \nabla \psi_{\mathcal{X}}(\bar{x})\|_{\mathcal{X}^*} \leq L_{\psi} \|x - \bar{x}\|_{\mathcal{X}}$ for some $L_{\psi} \geq 1$.*

An example of distance generating function satisfying (5) is the quadratic kernel $\psi_{\mathcal{X}}(x) = \frac{1}{2}x^{\top}Ax$ for some positive definite matrix A . One can also ensure this assumption by restricting the domain. For example, consider Itakura–Saito divergence using the logarithmic barrier distance generating function $\psi_{\mathcal{X}}(x) = -\sum_{i=1}^n \log(x_i)$ within the domain \mathbb{R}_{++}^n . The above assumption is satisfied when the domain is restricted to $\{x \in \mathbb{R}_+^n \mid x \geq c\}$ for some $c > \mathbf{0}$.

In the following lemma, we provide a one-step analysis of our proposed algorithm based on the above assumptions.

Lemma 6. *Let $\{x_k\}_{k \geq 0}$ be the sequence generated by VR-FoRMAB displayed in Algorithm 1 initialized from arbitrary vectors $x_0 \in \mathcal{X}$. Suppose Assumptions 1, 4, and 5 hold, and r_k , δ_k , and \bar{r}_k are defined in Definition 3. For any $k \geq 0$, the following result holds:*

$$\begin{aligned} 0 &\leq \mathbb{E}[\langle \bar{r}_{k+1}, x_{k+1} - x^* \rangle] - \theta_k \mathbb{E}[\langle \bar{r}_k, x_k - x^* \rangle] \\ &\quad + \frac{1-\gamma_k}{\sigma_k} \mathbb{E}[\mathbf{D}_X(x^*, x_k) - \mathbf{D}_X(x^*, x_{k+1})] + \frac{\gamma_k}{\sigma_k} \mathbb{E}[D_X^k(x^*) - \mathbf{D}_X(x^*, x_{k+1})] \\ &\quad + \mathbb{E}[\mathcal{C}_1^k \mathbf{D}_X(x_{k+1}, x_k) + \mathcal{C}_2^k D_X^k(x_{k+1}) + \mathcal{C}_3^k \mathbf{D}_X(x_k, x_{k-1}) + \mathcal{C}_4^k D_X^{k-1}(x_k) \\ &\quad + \mathcal{C}_5^k \sum_{t=i_k q+1}^k \mathbf{D}_X(x_t, x_{t-1})], \end{aligned} \quad (31)$$

where $\mathcal{C}_1^k \triangleq 8\rho((1-\beta)^2 L^2 + \frac{L_{\psi}^2(1-\gamma_k)^2}{\sigma_k^2}) + (\theta_k + (1-\beta)\bar{c}\underline{c}\sqrt{q})L - \frac{1-\gamma_k}{\sigma_k}$ and $\mathcal{C}_2^k \triangleq 8\rho(\beta^2 L^2 + \frac{L_{\psi}^2 \gamma_k^2}{\sigma_k^2}) - \frac{\gamma_k}{\sigma_k}$, $\mathcal{C}_3^k \triangleq 8\rho(1-\beta)^2 L^2 \theta_k^2 + \theta_k(1-\beta)L$, $\mathcal{C}_4^k \triangleq 8\rho\beta^2 L^2 \theta_k^2 + \theta_k \beta L$, and $\mathcal{C}_5^k \triangleq 8\rho(1-\beta)^2 \bar{c}^2 \underline{c}^2 L^2 + \frac{(1-\beta)\bar{c}\underline{c}L}{\sqrt{q}}$.

Proof. Considering the update of x_{k+1} , the optimality condition of the subproblem implies that

$$0 \in \partial(h + \mathbb{I}_X)(x_{k+1}) + v_k + \theta_k r_k + \frac{1}{\sigma_k} (\nabla \psi_{\mathcal{X}}(x_{k+1}) - \nabla \psi_{\mathcal{X}}(\hat{x}_k)),$$

where $\mathbb{I}_X(\cdot)$ denotes the indicator function of set X . Adding $F(x_{k+1})$ to both sides of above inclusion, defining $\mathfrak{U}_k \triangleq F(x_{k+1}) - (v_k + \theta_k r_k + \frac{1}{\sigma_k}(\nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(\hat{x}_k)))$ and rearranging the terms implies that

$$\mathfrak{U}_k \in F(x_{k+1}) + \partial(h + \mathbb{I}_X)(x_{k+1}). \quad (32)$$

Therefore, from Assumption 4 followed by applying Lemma 1 parts (b) and (d) we conclude that

$$\begin{aligned} & -\rho \|\mathfrak{U}_k\|_{\mathcal{X}^*}^2 \quad (33) \\ & \leq \langle \mathfrak{U}_k, x_{k+1} - x^* \rangle \\ & = \langle v_k + \theta_k r_k - F(x_{k+1}), x^* - x_{k+1} \rangle + \frac{1}{\sigma_k} \langle \nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(\hat{x}_k), x^* - x_{k+1} \rangle \\ & = \langle \delta_k, x^* - x_{k+1} \rangle + \theta_k \langle r_k - \bar{r}_k, x^* - x_k \rangle + \langle \bar{r}_{k+1}, x_{k+1} - x^* \rangle + \langle \theta_k \bar{r}_k, x^* - x_k \rangle \\ & \quad + \langle \theta_k r_k, x_k - x_{k+1} \rangle + \frac{1-\gamma_k}{\sigma_k} (\mathbf{D}_X(x, x_k) - \mathbf{D}_X(x, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_k)) \\ & \quad + \frac{\gamma_k}{\sigma_k} (D_X^k(x) - \mathbf{D}_X(x, x_{k+1}) - D_X^k(x_{k+1})) \\ & = \langle \delta_k, x^* - x_k \rangle + \theta_k \langle r_k - \bar{r}_k, x^* - x_k \rangle + \langle \bar{r}_{k+1}, x_{k+1} - x^* \rangle + \langle \theta_k \bar{r}_k, x^* - x_k \rangle \\ & \quad + \langle \delta_k + \theta_k r_k, x_k - x_{k+1} \rangle + \frac{1-\gamma_k}{\sigma_k} (\mathbf{D}_X(x^*, x_k) - \mathbf{D}_X(x^*, x_{k+1}) - \mathbf{D}_X(x_{k+1}, x_k)) \\ & \quad + \frac{\gamma_k}{\sigma_k} (D_X^k(x^*) - \mathbf{D}_X(x^*, x_{k+1}) - D_X^k(x_{k+1})). \end{aligned}$$

On the other hand, from the definitions of δ_k and \bar{r}_k in Definition 3 one can easily verify that $\mathfrak{U}_k = \bar{r}_{k+1} - \delta_k - \theta_k r_k + \frac{1}{\sigma_k}((\nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(\hat{x}_k)))$. Moreover, from Lipschitz continuity of $\nabla\psi_{\mathcal{X}}$ and definitions of \hat{x}_k and s_k we obtain

$$\begin{aligned} & \|\nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(\hat{x}_k)\|_{\mathcal{X}^*}^2 \quad (34) \\ & = \|\nabla\psi_{\mathcal{X}}(x_{k+1}) - (1 - \gamma_k)\nabla\psi_{\mathcal{X}}(x_k) - \gamma_k s_k\|_{\mathcal{X}^*}^2 \\ & \leq 2(1 - \gamma_k)^2 \|\nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(x_k)\|_{\mathcal{X}^*}^2 + 2\gamma_k^2 \|\nabla\psi_{\mathcal{X}}(x_{k+1}) - s_k\|_{\mathcal{X}^*}^2 \\ & \leq 2(1 - \gamma_k)^2 L_\psi^2 \mathbf{D}_X(x_{k+1}, x_k) + 2\gamma_k^2 L_\psi^2 \mathbf{D}_X^k(x_{k+1}). \end{aligned}$$

Therefore,

$$\begin{aligned} & \mathbb{E}[\|\mathfrak{U}_k\|_{\mathcal{X}^*}^2] \quad (35) \\ & \leq 4\mathbb{E}[\|\bar{r}_{k+1}\|_{\mathcal{X}^*}^2] + 4\mathbb{E}[\|\delta_k\|_{\mathcal{X}^*}^2] + 4\theta_k^2 \mathbb{E}[\|r_k\|_{\mathcal{X}^*}^2] + \frac{4}{\sigma_k^2} \mathbb{E}[\|\nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(\hat{x}_k)\|_{\mathcal{X}^*}^2] \\ & \leq 8((1 - \beta)^2 L^2 + \frac{L_\psi^2 (1-\gamma_k)^2}{\sigma_k^2}) \mathbb{E}[\mathbf{D}_X(x_{k+1}, x_k)] + 8(\beta^2 L^2 + \frac{L_\psi^2 \gamma_k^2}{\sigma_k^2}) \mathbb{E}[D_X^k(x_{k+1})] \\ & \quad + 8(1 - \beta)^2 \bar{c}^2 L^2 \sum_{t=i_k q+1}^k \mathbb{E}[\mathbf{D}_X(x_t, x_{t-1})] + 8(1 - \beta)^2 L^2 \theta_k^2 \mathbb{E}[\mathbf{D}_X(x_k, x_{k-1})] \\ & \quad + 8\beta^2 L^2 \theta_k^2 \mathbb{E}[D_X^{k-1}(x_k)], \end{aligned}$$

where in the second inequality we used Lipschitz continuity of operator F for the first and third terms along with strong convexity of the Bregman distance, Lemma 4 for the second term, and (34) for the last term.

Moreover, similar to (22) by using Cauchy Schwartz inequality followed by applying Assumption 1 and Young's inequality for some $\alpha_1, \alpha_2 > 0$ we obtain

$$\langle \delta_k + \theta_k r_k, x_k - x_{k+1} \rangle \leq \frac{\eta_k}{2} \|\delta_k\|_{\mathcal{X}^*}^2 + \frac{\theta_k (1-\beta)^2 L^2}{2\alpha_1} \|x_k - x_{k-1}\|_{\mathcal{X}}^2$$

$$+ \frac{\theta_k \beta^2 L^2}{2\alpha_2} \|x_k - \tilde{x}_{k-1}\|_{\mathcal{X}}^2 + \left(\frac{\alpha_1 + \alpha_2}{2} \theta_k + \frac{1}{2\eta_k}\right) \|x_{k+1} - x_k\|_{\mathcal{X}}^2.$$

Consequently, using Lemma 2 followed by taking expectation from both sides of the inequality, applying Lemma 4, and selecting $\eta_k = \frac{1}{(1-\beta)\bar{c}_c\sqrt{q}L}$, $\alpha_1 = (1-\beta)L$, and $\alpha_2 = \beta L$ we obtain

$$\begin{aligned} & \mathbb{E}[\langle \delta_k + \theta_k r_k, x_k - x_{k+1} \rangle] \\ & \leq \frac{(1-\beta)\bar{c}_c L}{\sqrt{q}} \sum_{t=i_k q+1}^k \mathbb{E}[\mathbf{D}_X(x_t, x_{t-1})] + \theta_k(1-\beta)L\mathbb{E}[\mathbf{D}_X(x_k, x_{k-1})] \\ & \quad + \theta_k \beta L\mathbb{E}[D_X^{k-1}(x_k)] + (\theta_k + (1-\beta)\bar{c}_c\sqrt{q})L\mathbb{E}[\mathbf{D}_X(x_{k+1}, x_k)]. \end{aligned} \quad (36)$$

Now, taking expectation from the inequality in (33), adding (36) as well as (35) multiplied by ρ , and rearranging the terms lead to the desired result. \square

Based on the one-step analysis derived above, we need to impose some conditions on the parameters to ensure a telescopic term in (31). In the following lemma, we specify these parameters to ensure a telescopic structure in (31) based on the constants C_i^k for $i \in \{1, \dots, 5\}$ and any $k \geq 0$. Later, in Theorem 2 we used these conditions to derive a convergence rate guarantee based on (31).

Lemma 7. *Let $\theta_k = 1$, $q = n$, $\beta = 1 - \frac{1}{2\bar{c}_c\sqrt{q+1}}$, $\gamma_k = \gamma \in (0, 1)$, and $\sigma_k = \sigma \leq \min\left\{\frac{\vartheta(1-\gamma)}{(6+\vartheta\zeta/L_\psi)L}, \frac{\vartheta\gamma}{(2+2\zeta'\vartheta/(\sqrt{2}L_\psi))\beta L}\right\}$ for any $\vartheta \in (0, 1)$ where $\zeta \triangleq 1/(1+6L_\psi/\vartheta)$ and $\zeta' \triangleq 1/(1+2L_\psi/(\sqrt{2}\vartheta))$. Assuming that $\rho \leq \frac{\vartheta\zeta'}{16\sqrt{2}\beta LL_\psi}$, then for any $k \geq 0$,*

$$C_1^k + C_3^k + qC_5^k + (1-\vartheta)\frac{1-\gamma_k}{\sigma_k} \leq 0, \quad (37a)$$

$$C_2^k + C_4^k + (1-\vartheta)\frac{\gamma_k}{\sigma_k} \leq 0, \quad (37b)$$

where C_i^k 's are defined in Lemma 6.

Proof. Consider the desired inequality (37a) and let $\sigma_k = \sigma$ and $\theta_k = 1$. Multiplying both sides by σ^2 and defining $\chi \triangleq (1-\beta)^2(\bar{c}^2\bar{c}^2q+2)$ imply that (37a) holds if $a_1\sigma^2 - a_2\sigma + a_3 \leq 0$, where $a_1 \triangleq 8\chi\rho L^2 + (1+\sqrt{\chi})L$, $a_2 \triangleq \vartheta(1-\gamma)$, and $a_3 \triangleq 8\rho L_\psi^2(1-\gamma)^2$. Therefore, assuming that $a_2^2 - 4a_1a_3 \geq 0$, σ must satisfy $\sigma \leq \frac{a_2 + \sqrt{a_2^2 - 4a_1a_3}}{2a_1}$. Note that the former condition can be rewritten as a quadratic inequality in terms of ρ as follows

$$b_1\rho^2 + b_2\rho - \vartheta^2 \leq 0, \quad (38)$$

where $b_1 \triangleq 256\chi L^2 L_\psi^2$, and $b_2 \triangleq 32\sqrt{\chi}LL_\psi^2$, moreover, the latter condition holds if

$$\sigma \leq \frac{a_2}{2a_1} = \frac{\vartheta(1-\gamma)}{16\chi\rho L^2 + 2(1+\sqrt{\chi})L} = \frac{16\vartheta(1-\gamma)L_\psi^2}{b_1\rho + b_2}. \quad (39)$$

From (38) we have that $\rho \leq \frac{-b_2 + \sqrt{b_2^2 + 4\vartheta^2 b_1}}{2b_1}$ and a simple algebra reveals $\sqrt{b_2^2 + 4b_1\vartheta^2} \geq b_2 + 2\zeta\vartheta b_1$, for any $\zeta \leq 1/(1+6L_\psi/\vartheta)$. Therefore, one can immediately conclude that $\rho \leq \frac{\vartheta\zeta}{\sqrt{b_1}}$ which implies that (39) holds if $\sigma \leq \frac{\vartheta(1-\gamma)}{(6+\vartheta\zeta/L_\psi)L}$.

Next, considering inequality (37b) one can follow the same line of proof to show the result. In particular, (37b) leads to the following quadratic inequality $c_1\sigma^2 - c_2\sigma + c_3 \leq 0$ where $c_1 \triangleq 16\rho\beta^2L^2 + \beta L$, $c_2 \triangleq \gamma\vartheta$, and $c_3 \triangleq 8\rho L_\psi^2\gamma^2$. Assuming that $d_1\rho^2 + d_2\rho - \vartheta^2 \leq 0$ where $d_1 \triangleq 512\beta^2L^2L_\psi^2$ and $d_2 \triangleq 32\beta LL_\psi^2$, we conclude that σ satisfies $\sigma \leq \frac{c_2}{2c_1} = \frac{\gamma\vartheta}{32\rho\beta^2L^2+2\beta L}$. Moreover, $\rho \leq \frac{-d_2 + \sqrt{d_2^2 + 4\vartheta^2 d_1}}{2d_1}$ holds if $\rho \leq \frac{\zeta'\vartheta}{\sqrt{d_1}}$ for any $\zeta' \leq 1/(1 + 2L_\psi/(\sqrt{2}\vartheta))$, hence, (37b) holds if $\sigma \leq \frac{\vartheta\gamma}{(2+2\zeta'\vartheta/(\sqrt{2}L_\psi))\beta L}$. Finally, comparing the two upper-bounds obtained for ρ by choosing β as in the statement of the theorem we have that $\zeta/\sqrt{\chi} \leq \zeta'/(\sqrt{2}\beta)$, hence, $\rho \leq \frac{\vartheta\zeta'}{\sqrt{d_1}} \leq \frac{\vartheta\zeta}{\sqrt{b_1}}$. \square

Theorem 2. Let $\{x_k\}_{k \geq 0}$ be the sequence generated by VR-FoRMAB displayed in Algorithm 1 initialized from arbitrary vector $x_0 \in \mathcal{X}$. For any $k \geq 0$ let $\mathfrak{U}_k \triangleq F(x_{k+1}) - (v_k + \theta_k r_k + \frac{1}{\sigma_k}(\nabla\psi_{\mathcal{X}}(x_{k+1}) - \nabla\psi_{\mathcal{X}}(\hat{x}_k)))$. Suppose Assumptions 1, 4, and 5 hold and the step-size sequence σ_k and parameters γ_k, β , and q are selected as in the statement of Lemma 7 for any $k \geq 0$. Then, for any $K > 1$,

$$\sum_{k=0}^{K-2} \mathbb{E}[\|\mathfrak{U}_k\|_{\mathcal{X}^*}^2] \leq \vartheta \max\{\mathcal{E}_1, \mathcal{E}_2\} \mathbf{D}_X(x^*, x_0) = \mathcal{O}(nL^2), \quad (40)$$

where $\mathcal{E}_1 \triangleq 8(1 + 2(1 - \beta)^2)L^2/(1 - \gamma) + 8L_\psi^2(1 - \gamma)/\sigma^2$ and $\mathcal{E}_2 \triangleq 16\beta^2L^2/\gamma + 8L_\psi^2\gamma/\sigma^2$.

Proof. Since the step size and algorithm's parameters satisfy the conditions of Lemma 7 we have that (37a) and (37b) hold. Therefore, using Lemma 2 and thanks to the telescopic summation we have

$$\begin{aligned} & \sum_{k=0}^{K-1} \mathbb{E}[\mathcal{C}_1^k \mathbf{D}_X(x_{k+1}, x_k) + \mathcal{C}_2^k D_X^k(x_{k+1}) + \mathcal{C}_3^k \mathbf{D}_X(x_k, x_{k-1}) + \mathcal{C}_4^k D_X^{k-1}(x_k) \\ & \quad + \mathcal{C}_5^k \sum_{t=i_k q+1}^k \mathbf{D}_X(x_t, x_{t-1})] \\ & \leq -(1 - \vartheta) \sum_{k=0}^{K-1} \mathbb{E}[\frac{1-\gamma}{\sigma} \mathbf{D}_X(x_{k+1}, x_k) + \frac{\gamma}{\sigma} D_X^k(x_{k+1})]. \end{aligned} \quad (41)$$

Next, summing the result of Lemma 6 over k from 0 to $K - 1$ and using (41) we conclude that

$$\begin{aligned} & \vartheta \sum_{k=0}^{K-1} \mathbb{E}[\frac{1-\gamma}{\sigma} \mathbf{D}_X(x_{k+1}, x_k) + \frac{\gamma}{\sigma} D_X^k(x_{k+1})] \\ & \leq \mathbb{E}[\langle \bar{r}_K, x_K - x^* \rangle] - \mathbb{E}[\langle \bar{r}_0, x_0 - x^* \rangle] + \frac{1-\gamma}{\sigma} \mathbb{E}[\mathbf{D}_X(x^*, x_0) - \mathbf{D}_X(x^*, x_K)] \\ & \quad + \frac{\gamma}{\sigma} \mathbb{E}[D_X^{K-1}(x^*) - \mathbf{D}_X(x^*, x_K)], \\ & \leq \mathbb{E}[(1 - \beta)L\mathbf{D}_X(x_K, x_{K-1}) + \beta L D_X^{K-1}(x_K) + L\mathbf{D}_X(x^*, x_K)] \\ & \quad + \frac{1-\gamma}{\sigma} \mathbb{E}[\mathbf{D}_X(x^*, x_0) - \mathbf{D}_X(x^*, x_K)] + \frac{\gamma}{\sigma} \mathbb{E}[D_X^{K-1}(x^*) - \mathbf{D}_X(x^*, x_K)], \end{aligned} \quad (42)$$

where in the last inequality we followed a similar step as in (26) for $x = x^*$ to provide an upper bound on $\langle \bar{r}_K, x_K - x^* \rangle - \langle \bar{r}_0, x_0 - x^* \rangle$ by using Young's inequality and noting that $\bar{r}_0 = 0$. Since σ

satisfies $\sigma \leq \min\{\frac{1-\gamma}{2L}, \frac{\gamma}{2L}\}$ we can simplify the terms in (42) to obtain

$$\vartheta \sum_{k=1}^{K-1} \mathbb{E}[\frac{1-\gamma}{\sigma} \mathbf{D}_X(x_k, x_{k-1}) + \frac{\gamma}{\sigma} D_X^{k-1}(x_k)] \leq \frac{1}{\sigma} \mathbf{D}_X(x^*, x_0).$$

Multiplying both sides by $\vartheta\sigma$ leads to

$$\sum_{k=1}^{K-1} \mathbb{E}[(1-\gamma)\mathbf{D}_X(x_k, x_{k-1}) + \gamma D_X^{k-1}(x_k)] \leq \vartheta \mathbf{D}_X(x^*, x_0). \quad (43)$$

Next, recalling the upper-bound on $\mathbb{E}[\|\mathfrak{U}_k\|_{\mathcal{X}^*}^2]$ in (35) in terms of the consecutive iterates and summing over $k = 0$ to $K - 2$ implies that

$$\begin{aligned} & \sum_{k=0}^{K-2} \mathbb{E}[\|\mathfrak{U}_k\|_{\mathcal{X}^*}^2] \leq \\ & 8((1-\beta)^2 L^2 + \frac{L_\psi^2(1-\gamma)^2}{\sigma^2}) \sum_{k=0}^{K-2} \mathbb{E}[\mathbf{D}_X(x_{k+1}, x_k)] + 8(\beta^2 L^2 + \frac{L_\psi^2 \gamma^2}{\sigma^2}) \sum_{k=0}^{K-2} \mathbb{E}[D_X^k(x_{k+1})] \\ & + 8((1-\beta)^2 \bar{c}^2 \underline{c}^2 q L^2 + (1-\beta)^2 L^2) \sum_{k=0}^{K-2} \mathbb{E}[\mathbf{D}_X(x_k, x_{k-1})] + 8\beta^2 L^2 \sum_{k=0}^{K-2} \mathbb{E}[D_X^{k-1}(x_k)]. \end{aligned}$$

Now, using $(1-\beta)\bar{c}\underline{c}\sqrt{q} \leq 1$ and definition of \mathcal{E}_1 and \mathcal{E}_2 from the statement of the theorem in the last inequality, we have that

$$\sum_{k=0}^{K-2} \mathbb{E}[\|\mathfrak{U}_k\|_{\mathcal{X}^*}^2] \leq \sum_{k=1}^{K-1} (\mathcal{E}_1(1-\gamma)\mathbb{E}[\mathbf{D}_X(x_k, x_{k-1})] + \mathcal{E}_2\gamma\mathbb{E}[D_X^{k-1}(x_k)]).$$

Finally, combining the above inequality with (43) leads to the desired result. \square

Corollary 2. *Under the premises of Theorem 2 by choosing $\gamma = \mathcal{O}(1/\sqrt{n})$ there exists $t \in \{1, \dots, K-1\}$ such that*

$$\mathbb{E}[\mathbf{dist}(\mathbf{0}, F(x_t) + \partial(h + \mathbb{I}_X)(x_t))] \leq \epsilon, \quad (44)$$

within $K = \mathcal{O}(nL^2/\epsilon^2)$ iterations. Moreover, the oracle complexity to achieve an ϵ -approximated solution is $\mathcal{O}(nL^2/\epsilon^2)$.

Proof. From Theorem 2 and defining $t = \operatorname{argmin}_{k=0}^{K-2} \{\|\mathfrak{U}_k\|_{\mathcal{X}^*}\}$ we conclude that for any $K > 1$, $\mathbb{E}[\|\mathfrak{U}_t\|_{\mathcal{X}^*}^2] \leq \frac{\vartheta \max\{\mathcal{E}_1, \mathcal{E}_2\}}{K-1} \mathbf{D}_X(x^*, x_0)$. Now, let $A(x_{k+1}) \triangleq F(x_{k+1}) + \partial(h + \mathbb{I}_X)(x_{k+1})$ and by noting that $\mathfrak{U}_k \in A(x_{k+1})$ for any $k \geq 0$ we have

$$\mathbb{E}[\|\mathfrak{U}_t\|_{\mathcal{X}^*}^2] \geq (\mathbb{E}[\|\mathfrak{U}_t\|_{\mathcal{X}^*}])^2 \geq \left(\mathbb{E}\left[\min_{u \in A(x_{t+1})} \|u\|_{\mathcal{X}^*}\right]\right)^2 = (\mathbb{E}[\mathbf{dist}(\mathbf{0}, A(x_{t+1}))])^2. \quad (45)$$

Hence, one can readily conclude that $\mathbb{E}[\mathbf{dist}(\mathbf{0}, A(x_{t+1}))] \leq \sqrt{\frac{\vartheta \max\{\mathcal{E}_1, \mathcal{E}_2\}}{K-1} \mathbf{D}_X(x^*, x_0)}$ which in light of $\gamma = \mathcal{O}(1/\sqrt{n})$ we have that $\max\{\mathcal{E}_1, \mathcal{E}_2\} = \mathcal{O}(nL^2)$ leading to the desired result. \square

Remark 5. We can simplify the parameter selection of our proposed method by choosing $\beta = 1 - \frac{1}{2\underline{c}\underline{c}\sqrt{n+1}}$, $\gamma = \beta$, and $\sigma < \frac{(1-\gamma)}{6L}$. It is worth noting that, unlike the monotone setting when $\underline{c} \leq 1$ (e.g., Euclidean norm) the parameters β and γ do not vanish. This indeed shows the impact of novel momentum terms and operator estimation in our method to ensure the obtained complexity result.

Remark 6. The condition on the weak-MVI parameter ρ in our analysis is $\rho < \frac{1}{16LL_\psi(1+2L_\psi/\sqrt{2})}$ when $n \gg 1$ which is slightly worse than some of the existing results [36]. However, it is important to note that such a loose bound stems from our consideration of an arbitrary norm in our framework and using some crude bounds such as those in (35). It is reasonable to expect that specializing our analysis for Euclidean space could lead to an improvement of the constant in the upper bound of ρ . Nevertheless, our result encompasses a broader range of problems, and notably, we have achieved an improved convergence rate compared to existing methods [37, 36] which has appeared for the first time in the literature.

5 Experimental results

In this section, we implement VR-FoRMAB for solving DRO example in section 1 and a non-monotone two-player matrix game. We compare our method with other state-of-the-art methods. All experiments are performed on a machine running 64-bit Windows 11 with Intel i5-1135G7 @2.40GHz and 8GB RAM.

5.1 Distributionally Robust Optimization

We implement our method to solve the DRO problem (3) described in Section 1 and compare it with Stochastic Variance Reduced Accelerated Primal-Dual Method (SVR-APD) [50] and Mirror-Prox with Variance Reduction (VR-MP) [4]. Following the setup in [50], we consider a set of labeled data points consisting of feature vectors $\{a_i\}_{i=1}^n \subset \mathbb{R}^d$ and labels $\{b_i\}_{i=1}^n \subseteq \{+1, -1\}^n$. Let $\ell_i(x) = \log(1 + \exp(-b_i a_i^\top x))$ and $V(y, \frac{1}{n}\mathbf{1}_n) = \frac{1}{2}\|ny - \mathbf{1}_n\|^2$ serves as the Chi-square divergence measure. Furthermore, we specify $X = [-10, 10]^d$ and $\rho = 50$. The comparison of methods is conducted across different datasets, and a summary of the datasets is presented in Table 3. For all methods, projection onto the simplex-set constraint in the maximization is computed by utilizing the Bregman distance with generating function $\psi_Y(y) = \sum_{i=1}^n y_i \log(y_i)$. We selected the VR-FoRMAB's parameters according to Remark 4, i.e., $\sigma = \frac{1-\beta}{(2\sqrt{n+4})L}$, $\gamma = \beta = 0$, $S = 1$, $q = n$. Similarly, the parameters of SVR-APD and VR-MP are selected based on their theoretical suggestion. Let us define the Lagrangian function $\mathcal{L}(x, y) \triangleq f(x) + \sum_{i=1}^n L_i(x, y) - h(y)$ and recall that $L_i(x, y) = ny_i \ell_i(x) - \lambda(\frac{1}{2}V_i(y_i, \frac{1}{n}\mathbf{1}_n) - \frac{\rho}{n})$, $f(x) = \mathbb{I}_{X \times \mathbb{R}_+}(u, \lambda)$ and $h(y) = \mathbb{I}_{\Delta_n}(y)$. The results are depicted in Figure 1, where we plot the results in terms of the difference of Lagrangian functions evaluated at a solution pair (x^*, y^*) , i.e. $\mathcal{L}(x_k, y^*) - \mathcal{L}(x^*, y_k)$ ¹, versus time. Moreover, in Table 4 we compare the gap function defined in (23) at the last iterate point for all algorithms. Our method demonstrates better performance compared to the other two approaches across all experiments.

¹Note that $0 \leq \mathcal{L}(x_k, y^*) - \mathcal{L}(x^*, y_k) \leq \sup_{z \in X} \langle F(z^*), z - z^* \rangle = \mathcal{G}(z_k)$.

	w2a	Mushroom	Phishing	a7a
samples	3470	8124	11055	16100
features	300	112	64	122

Table 3: Different datasets used in the experiment from LIBSVM [10].




Figure 1: Comparison of the methods in terms of time for different datasets: Top row from left to right w2a and Mushrooms, Bottom row left to right Phishing and a7a.

	w2a	Mushroom	Phishing	a7a
VR-FoRMAB	5.1e-2	4.7e-3	1.2e-2	1.1e-3
SVRAPD	1.3e-1	2.6e-2	9.6e-2	2.3e-2
VR-MP	1.2e-1	1.9e-2	8.6e-2	2.3e-2

Table 4: Comparing gap function $\sup_{(x,y) \in \mathcal{X} \times \mathcal{Y}} \{\mathcal{L}(x_K, y) - \mathcal{L}(x, y_K)\}$ at the last iterate for different datasets.

5.2 Matrix Game

In this section, we compare the performance of our proposed method in a non-monotone setting with the state-of-the-art algorithms for solving a two-player matrix game formulated as follows:

$$\min_{\|u\|_2 \leq 1} \max_{\|w\|_2 \leq 1} -\frac{\nu}{2} \|u\|_2^2 + \langle Au, w \rangle + \frac{\nu}{2} \|w\|_2^2. \quad (46)$$

A first-order stationary solution of the above problem can be found by solving the corresponding VI formulated as (1) by letting $x = \begin{bmatrix} u \\ w \end{bmatrix}$, $F(x) = \begin{bmatrix} A^\top w - \nu u \\ -Au - \nu w \end{bmatrix}$, $h \equiv 0$, and $X = \{u \in \mathbb{R}^n \mid \|u\|_2 \leq 1\} \times \{w \in \mathbb{R}^m \mid \|w\|_2 \leq 1\}$. It is easy to verify that F is a non-monotone and Lipschitz continuous




Figure 2: Comparison of methods for solving (46) for various dimensions. From left to right, $n = 100$, 500, and 1000.

map. In particular, similar to [37] we can show that the Lipschitz constant of F is $L = \sqrt{v^2 + \|A\|_2^2}$ and F satisfies Assumption 4 with constant $\rho = \frac{v}{v^2 + \|A\|_2^2}$. In our experiment, we let $v = 1$, generate A randomly from a standard Gaussian distribution then normalized such that $\|A\|_2 = 40$, $n = m$, and $F_i(x) = \begin{bmatrix} nA_i w_i - vu \\ -mA_{:,i} u_i - vw \end{bmatrix}$. We compare our proposed method with VR-MP and BC-SEG+ [36].

The parameters of our method are selected based on Remark 5, i.e., $\gamma = \beta = 1 - \frac{1}{2\sqrt{n}+1}$ and $\sigma = \frac{\nu}{6L}$ for some $\nu > 0$. For VR-MP we select the step size as $\sigma = \frac{\nu}{\sqrt{n}L}$ and for BC-SEG+ we used $\alpha_k = \nu/k$. In our experiment, we select $\nu \in \{0.1, 1, 10\}$ as a tuning parameter to observe the performance of the considered methods with different step-sizes. We tested the performance of the considered methods for various dimensions $n = m \in \{100, 500, 1000\}$ and the plots are depicted in Figure 2. Since $x = \mathbf{0}$ is the equilibrium of this problem, the plots are in terms of the normalized solution norm, i.e., $\|x_k\|/\sqrt{n}$, versus the number of sample operators evaluated by the algorithms. Conversely, BC-SEG+ demonstrates convergence in lower dimensions but exhibits sluggish convergence when $n = 1000$, possibly due to increased variance in sampling and a diminishing step-size. Our proposed method, VR-FoRMAB, consistently converges to the solution across all experiments, surpassing other methods.

6 Conclusions

In this paper, we studied generalized VI problems exhibiting finite-sum structure. Motivated by the absence of a single-loop method capable of adapting a Bregman distance function to tackle this category of problems, we introduce a unified stochastic method called Variance-Reduced Forward Reflected Moving Average Backward (VR-FoRMAB). We characterize the convergence rate and oracle complexity of this method for monotone settings, marking it as the first single-loop method capable of using the Bregman distance function to achieve the optimal complexity of $\mathcal{O}(\sqrt{n}/\epsilon)$, to the best of our knowledge. Additionally, we analyzed the convergence of our proposed algorithm in a non-monotone setting under a weak MVI assumption and Lipschitz continuity of the Bregman distance generating function. Our proposed method can achieve a complexity of $\mathcal{O}(n/\epsilon^2)$ which improves the oracle complexity over existing methods.

References

- [1] M. Ahoosh, A. Themelis, and P. Patrinos. A bregman forward-backward linesearch algorithm for nonconvex composite optimization: superlinear convergence to nonisolated local minima. *SIAM Journal on Optimization*, 31(1):653–685, 2021.
- [2] A. Alacaoglu, D. Kim, and S. J. Wright. Extending the reach of first-order algorithms for nonconvex min-max problems with cohyponotonicity. *arXiv preprint arXiv:2402.05071*, 2024.
- [3] A. Alacaoglu and Y. Malitsky. Stochastic variance reduction for variational inequality methods. *ArXiv*, abs/2102.08352, 2021.
- [4] A. Alacaoglu and Y. Malitsky. Stochastic variance reduction for variational inequality methods. In *Conference on Learning Theory*, pages 778–816. PMLR, 2022.
- [5] Z. Allen-Zhu and E. Hazan. Variance reduction for faster non-convex optimization. In *International conference on machine learning*, pages 699–707. PMLR, 2016.
- [6] A. Banerjee, S. Merugu, I. S. Dhillon, J. Ghosh, and J. Lafferty. Clustering with bregman divergences. *Journal of machine learning research*, 6(10), 2005.
- [7] H. H. Bauschke, W. M. Moursi, and X. Wang. Generalized monotone operators and their averaged resolvents. *Mathematical Programming*, 189:55–74, 2021.
- [8] A. Böhm. Solving nonconvex-nonconcave min-max problems exhibiting weak minty solutions. *arXiv preprint arXiv:2201.12247*, 2022.
- [9] Y. Cai and W. Zheng. Accelerated single-call methods for constrained min-max optimization. *arXiv preprint arXiv:2210.03096*, 2022.
- [10] C.-C. Chang and C.-J. Lin. Libsvm: A library for support vector machines. *ACM transactions on intelligent systems and technology (TIST)*, 2(3):1–27, 2011.
- [11] S. Choudhury, E. Gorbunov, and N. Loizou. Single-call stochastic extragradient methods for structured non-monotone variational inequalities: Improved analysis under weaker conditions. *Advances in Neural Information Processing Systems*, 36, 2024.
- [12] L. P. Cinelli, M. A. Marins, E. A. B. Da Silva, and S. L. Netto. *Variational methods for machine learning with applications to deep networks*. Springer, 2021.
- [13] C. Daskalakis, D. J. Foster, and N. Golowich. Independent policy gradient methods for competitive reinforcement learning. *Advances in neural information processing systems*, 33:5527–5540, 2020.
- [14] A. M. Devraj and J. Chen. Stochastic variance reduced primal dual algorithms for empirical composition optimization. *Advances in Neural Information Processing Systems*, 32, 2019.
- [15] J. Diakonikolas, C. Daskalakis, and M. I. Jordan. Efficient methods for structured nonconvex-nonconcave min-max optimization. In *International Conference on Artificial Intelligence and Statistics*, pages 2746–2754. PMLR, 2021.

- [16] C. Fang, C. J. Li, Z. Lin, and T. Zhang. Spider: Near-optimal non-convex optimization via stochastic path-integrated differential estimator. *Advances in neural information processing systems*, 31, 2018.
- [17] I. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio. Generative adversarial nets. *Advances in neural information processing systems*, 27, 2014.
- [18] W. Guo, M. I. Jordan, and T. Lin. A variational inequality approach to bayesian regression games. In *2021 60th IEEE Conference on Decision and Control (CDC)*, pages 795–802. IEEE, 2021.
- [19] E. Y. Hamedani and N. S. Aybat. A primal-dual algorithm with line search for general convex-concave saddle point problems. *SIAM Journal on Optimization*, 31(2):1299–1329, 2021.
- [20] Y. Han, G. Xie, and Z. Zhang. Lower complexity bounds of finite-sum optimization problems: The results and construction. *Journal of Machine Learning Research*, 25(2):1–86, 2024.
- [21] A. N. Iusem, A. Jofré, R. I. Oliveira, and P. Thompson. Extragradient method with variance reduction for stochastic variational inequalities. *SIAM Journal on Optimization*, 27(2):686–724, 2017.
- [22] R. Johnson and T. Zhang. Accelerating stochastic gradient descent using predictive variance reduction. *Advances in neural information processing systems*, 26, 2013.
- [23] A. Juditsky, A. Nemirovski, and C. Tauvel. Solving variational inequalities with stochastic mirror-prox algorithm. *Stochastic Systems*, 1(1):17–58, 2011.
- [24] A. Kavis, S. Skoulakis, K. Antonakopoulos, L. T. Dadi, and V. Cevher. Adaptive stochastic variance reduction for non-convex finite-sum minimization. *Advances in Neural Information Processing Systems*, 35:23524–23538, 2022.
- [25] A. A. Kulkarni and U. V. Shanbhag. Revisiting generalized nash games and variational inequalities. *Journal of Optimization Theory and Applications*, 154:175–186, 2012.
- [26] X. Lian, M. Wang, and J. Liu. Finite-sum composition optimization via variance reduced gradient descent. In *Artificial Intelligence and Statistics*, pages 1159–1167. PMLR, 2017.
- [27] L. Luo, G. Xie, T. Zhang, and Z. Zhang. Near optimal stochastic algorithms for finite-sum unbalanced convex-concave minimax optimization. *arXiv preprint arXiv:2106.01761*, 2021.
- [28] Y. Malitsky and M. K. Tam. A forward-backward splitting method for monotone inclusions without cocoercivity. *SIAM Journal on Optimization*, 30(2):1451–1472, 2020.
- [29] H. Namkoong and J. C. Duchi. Stochastic gradient methods for distributionally robust optimization with f-divergences. *Advances in neural information processing systems*, 29, 2016.
- [30] A. Nemirovski. Prox-method with rate of convergence $o(1/t)$ for variational inequalities with lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15(1):229–251, 2004.

- [31] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on optimization*, 19(4):1574–1609, 2009.
- [32] Y. Nesterov. Dual extrapolation and its applications to solving variational inequalities and related problems. *Mathematical Programming*, 109(2):319–344, 2007.
- [33] J. v. Neumann. A model of general economic equilibrium. *The Review of Economic Studies*, 13(1):1–9, 1945.
- [34] B. Palaniappan and F. Bach. Stochastic variance reduction methods for saddle-point problems. *Advances in Neural Information Processing Systems*, 29, 2016.
- [35] F. Parise and A. Ozdaglar. A variational inequality framework for network games: Existence, uniqueness, convergence and sensitivity analysis. *Games and Economic Behavior*, 114:47–82, 2019.
- [36] T. Pethick, O. Fercoq, P. Latafat, P. Patrinos, and V. Cevher. Solving stochastic weak minty variational inequalities without increasing batch size. *arXiv preprint arXiv:2302.09029*, 2023.
- [37] T. Pethick, P. Latafat, P. Patrinos, O. Fercoq, and V. Cevher. Escaping limit cycles: Global convergence for constrained nonconvex-nonconcave minimax problems. *arXiv preprint arXiv:2302.09831*, 2023.
- [38] T. Pethick, W. Xie, and V. Cevher. Stable nonconvex-nonconcave training via linear interpolation. *Advances in Neural Information Processing Systems*, 36, 2024.
- [39] A. Pichugin, M. Pechin, A. Beznosikov, and A. Gasnikov. Optimal analysis of method with batching for monotone stochastic finite-sum variational inequalities. *arXiv preprint arXiv:2401.07858*, 2024.
- [40] M. Schmidt, N. Le Roux, and F. Bach. Minimizing finite sums with the stochastic average gradient. *Mathematical Programming*, 162:83–112, 2017.
- [41] M. Simonetti, D. Perri, and O. Gervasi. An example of use of variational methods in quantum machine learning. In *International Conference on Computational Science and Its Applications*, pages 597–609. Springer, 2022.
- [42] V. V. Singh and A. Lisser. Variational inequality formulation for the games with random payoffs. *Journal of Global Optimization*, 72:743–760, 2018.
- [43] C. Song and J. Diakonikolas. Cyclic coordinate dual averaging with extrapolation. *SIAM Journal on Optimization*, 33(4):2935–2961, 2023.
- [44] P. Tseng. A modified forward-backward splitting method for maximal monotone mappings. *SIAM Journal on Control and Optimization*, 38(2):431–446, 2000.
- [45] P. Tseng. On accelerated proximal gradient methods for convex-concave optimization. *submitted to SIAM Journal on Optimization*, 2(3), 2008.
- [46] T. Ui. Bayesian nash equilibrium and variational inequalities. *Journal of Mathematical Economics*, 63:139–146, 2016.

- [47] X. Wang and Z. Wang. A bregman inertial forward-reflected-backward method for nonconvex minimization. *Journal of Global Optimization*, pages 1–28, 2023.
- [48] J. Yang, N. Kiyavash, and N. He. Global convergence and variance-reduced optimization for a class of nonconvex-nonconcave minimax problems. *arXiv preprint arXiv:2002.09621*, 2020.
- [49] Y. Yang, W. Li, B. Song, Y. Zou, and Y. Pan. Enhanced fault tolerant kinematic control of redundant robots with linear-variational-inequality based zeroing neural network. *Engineering Applications of Artificial Intelligence*, 133:108068, 2024.
- [50] E. Yazdandoost Hamedani and A. Jalilzadeh. A stochastic variance-reduced accelerated primal-dual method for finite-sum saddle-point problems. *Computational Optimization and Applications*, 85(2):653–679, 2023.
- [51] J. Zhang and L. Xiao. A composite randomized incremental gradient method. In *International Conference on Machine Learning*, pages 7454–7462. PMLR, 2019.
- [52] S. Zhang, M. Hadji, A. Lisser, and Y. Mezali. Variational inequality for n-player strategic chance-constrained games. *SN Computer Science*, 4(1):82, 2022.