

Some notes on an identity of Frisch

Kunle Adegoke

Department of Physics and Engineering Physics
Obafemi Awolowo University, 220005 Ile-Ife, Nigeria

adegoke00@gmail.com

Robert Frontczak

Independent Researcher

Reutlingen, Germany

robert.frontczak@web.de

Abstract

In this note, we show how a combinatorial identity of Frisch can be applied to prove and generalize some well-known identities involving harmonic numbers. We also present some combinatorial identities involving odd harmonic numbers which can be inferred straightforwardly from our results.

2010 *Mathematics Subject Classification*: 05A10; 11B65.

Keywords: Harmonic number; odd harmonic number; binomial coefficient; binomial transform.

1 Preliminaries

The binomial coefficients are defined, for non-negative integers n and m , by

$$\binom{n}{m} = \begin{cases} \frac{n!}{m!(n-m)!}, & n \geq m; \\ 0, & n < m. \end{cases}$$

More generally, for complex numbers r and s , they are defined by

$$\binom{r}{s} = \frac{\Gamma(r+1)}{\Gamma(s+1)\Gamma(r-s+1)},$$

where the Gamma function, $\Gamma(z)$, is defined for $\Re(z) > 0$ by the integral [12]

$$\Gamma(z) = \int_0^{\infty} e^{-t} t^{z-1} dt.$$

The function $\Gamma(z)$ can be extended to the whole complex plane by analytic continuation. It has a simple pole at each of the points $z = \dots, -3, -2, -1, 0$. The Gamma function extends the classical factorial function to the complex plane: $(z-1)! = \Gamma(z)$; thereby facilitating the computation of the binomial coefficients at non-integer and non-real values. Closely related to the Gamma function is the psi or digamma function defined by $\psi(z) = \Gamma'(z)/\Gamma(z)$. It possesses the infinite series expression [12, p. 14]

$$\psi(z) = -\gamma + \sum_{k=0}^{\infty} \left(\frac{1}{k+1} - \frac{1}{k+z} \right), \quad (1)$$

with γ being the Euler-Mascheroni constant.

Harmonic numbers, H_x , $x \in \mathbb{C} \setminus \mathbb{Z}^-$, defined by the relation

$$H_x = H_{x-1} + \frac{1}{x}, \quad H_0 = 0, \quad (2)$$

are connected to the digamma function through the fundamental relation

$$H_x = \psi(x+1) + \gamma. \quad (3)$$

When each x is a positive integer, say n , we have the sequence of harmonic numbers, $(H_n)_{n \in \mathbb{Z}^+}$ and the recurrence relation (2) then gives

$$H_n = \sum_{k=1}^n \frac{1}{k}, \quad H_0 = 0. \quad (4)$$

The following combinatorial identity is attributed to Frisch [7]:

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{\binom{b+k}{c}} = \frac{c}{n+c} \frac{1}{\binom{n+b}{b-c}}, \quad n \in \mathbb{Z}^+ \text{ and } b, c, b-c \in \mathbb{C} \setminus \mathbb{Z}^-. \quad (5)$$

It is listed in Gould's compendium [8] as Identity 4.2 and was used recently by Gould and Quaintance [9] to prove a new binomial transform identity. Abel [1] gave a short proof for Frisch's identity and studied its infinite variant. In this note, we report how Frisch's identity (5) can be applied to prove and generalize some well-known identities involving harmonic numbers H_n . We also present some combinatorial identities involving odd harmonic numbers O_n , which can be inferred straightforwardly from our results. Our line of approach consists essentially of utilizing the fact that derivatives of the generalized binomial coefficients yield harmonic numbers. This method is familiar and has appeared in some important earlier research of other mathematicians [5, 6, 13, 14].

2 Results

Theorem 1. For $n \in \mathbb{Z}^+$ and $b, c, b - c \in \mathbb{C} \setminus \mathbb{Z}^-$, we have

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{\binom{b+k}{c}} (H_{k+b} - H_{k+b-c}) = \frac{c}{n+c} \frac{H_{n+b} - H_{b-c}}{\binom{n+b}{n+c}} \quad (6)$$

and

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{c}{k+c} \frac{H_{k+b} - H_{b-c}}{\binom{k+b}{k+c}} = \frac{H_{n+b} - H_{n+b-c}}{\binom{b+n}{c}}. \quad (7)$$

In particular, for $n \in \mathbb{Z}^+$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{\binom{b+k}{k}} (H_{k+b} - H_k) = \frac{b}{n+b} H_{n+b} \quad (8)$$

and

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{H_{k+b}}{k+b} = \frac{H_{n+b} - H_n}{b \binom{b+n}{b}}. \quad (9)$$

Proof. In Frisch's identity (5) treat b and c as complex numbers and differentiate w.r.t. b using

$$\frac{d}{db} \binom{b+k}{c}^{-1} = \binom{b+k}{c}^{-1} (\psi(b+k+1-c) - \psi(b+k+1)) \quad (10)$$

and

$$\frac{d}{db} \binom{n+b}{n+c}^{-1} = \binom{n+b}{n+c}^{-1} (\psi(b+1-c) - \psi(b+n+1)), \quad (11)$$

where $\psi(z)$ is the psi function. Simplify, making use of the fundamental relation (3). Identity (7) is the binomial transform of (6). \square

Corollary 2. For $n \in \mathbb{Z}^+$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$, $b \neq 0$ we have

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{(b+k)^2} = \frac{1}{n+1} \frac{H_{n+b} - H_{b-1}}{\binom{n+b}{n+1}}. \quad (12)$$

Identity (12) generalizes the well-known identity

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{(k+1)^2} = \frac{H_{n+1}}{n+1}, \quad (13)$$

which can be proved directly via

$$\begin{aligned}
(n+1) \sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)^2} &= \sum_{k=0}^n \frac{n+1}{k+1} \binom{n}{k} \frac{(-1)^k}{k+1} \\
&= \sum_{k=0}^n \binom{n+1}{k+1} \frac{(-1)^k}{k+1} \\
&= \sum_{k=1}^{n+1} \binom{n+1}{k} \frac{(-1)^{k-1}}{k} \\
&= H_{n+1}.
\end{aligned}$$

Theorem 3. For $n \in \mathbb{Z}^+$ and $b, c, b-c \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=0}^n (-1)^{k+1} \frac{\binom{n}{k}}{\binom{b+k}{c}} (H_{k+b-c} - H_c) = \frac{1}{(n+c) \binom{n+b}{n+c}} \left(\frac{n}{n+c} + c(H_{n+c} - H_{b-c}) \right). \quad (14)$$

In particular, for $n \in \mathbb{Z}^+$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} H_{k+b} = \frac{1}{n \binom{n+b}{n}}. \quad (15)$$

Proof. Differentiate Frisch's identity (5) w.r.t. c using

$$\frac{d}{dc} \binom{b+k}{c}^{-1} = \binom{b+k}{c}^{-1} (\psi(c+1) - \psi(b+k+1-c)).$$

□

Identity (15) generalizes the well-known identity

$$\sum_{k=0}^n (-1)^k \binom{n}{k} H_k = -\frac{1}{n}, \quad (16)$$

which is the binomial transform of the sequence H_n (see [4, p. 34]).

Corollary 4. For $n \in \mathbb{N}_0$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=1}^n (-1)^{k+1} \frac{\binom{n}{k}}{k \binom{k+b}{k}} = H_{n+b} - H_b. \quad (17)$$

In particular,

$$\sum_{k=1}^n (-1)^{k+1} \frac{\binom{n}{k}}{k \binom{k+n}{k}} = H_{2n} - H_n. \quad (18)$$

Corollary 5. For $n \in \mathbb{Z}^+$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{\binom{k+b}{k}} H_{k+b} = \frac{b}{n+b} H_b - \frac{n}{(n+b)^2}. \quad (19)$$

Proof. Set $c = b$ in (14) to get

$$\sum_{k=0}^n (-1)^{k+1} \frac{\binom{n}{k}}{\binom{b+k}{b}} (H_k - H_b) = \frac{1}{n+b} \left(\frac{n}{n+b} + bH_{n+b} \right).$$

Use the fact that

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{\binom{b+k}{k}} = \frac{b}{n+b} \quad (20)$$

and combine with (8). □

Corollary 6. For $n \in \mathbb{N}_0$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$ we have

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k+1} \frac{k}{(k+b)^2} = \frac{H_{n+b} - H_b}{\binom{n+b}{n}}. \quad (21)$$

In particular,

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k+1} \frac{k}{(k+n)^2} = \frac{H_{2n} - H_n}{\binom{2n}{n}}. \quad (22)$$

Proof. Using the fact that the right hand-side of (19) is the binomial transform of $H_{n+b}/\binom{n+b}{n}$ we get

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \left(\frac{b}{k+b} H_b - \frac{k}{(k+b)^2} \right) = \frac{H_{n+b}}{\binom{n+b}{n}}.$$

Combine this with

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{1}{k+b} = \frac{1}{b \binom{n+b}{b}} \quad (23)$$

and the proof is completed. □

Remark. Comparing (21) with (12) we see that

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{b}{(b+k)^2} = \frac{H_{n+b} - H_{b-1}}{\binom{n+b}{n}}$$

and

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{k}{(b+k)^2} = \frac{-H_{n+b} + H_b}{\binom{n+b}{n}},$$

which upon addition again yield (23).

Theorem 7. If $n \in \mathbb{Z}^+$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$, then

$$\sum_{k=0}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{b+k}{k}} H_k = \frac{b}{n+b} (H_{n+b} - H_b) + \frac{n}{(n+b)^2}. \quad (24)$$

Proof. Set $c = b$ in (14) and use (20). \square

Theorem 8. If $n \in \mathbb{N}_0$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$, $b \neq 0$, then

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+b)^3} = \frac{1}{2n+2} \binom{n+b}{n+1}^{-1} \left((H_{n+b} - H_{b-1})^2 - H_{b-1}^{(2)} + H_{n+b}^{(2)} \right). \quad (25)$$

In particular,

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(k+1)^3} = \frac{1}{2n+2} \left(H_{n+1}^2 + H_{n+1}^{(2)} \right). \quad (26)$$

Proof. Write (12) as

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k}}{(b+k)^2} = \frac{1}{n+1} \frac{\psi(n+b+1) - \psi(b)}{\binom{n+b}{n+1}}$$

and differentiate with respect to b to obtain

$$\begin{aligned} & \sum_{k=0}^n \frac{(-1)^k}{(k+b)^3} \binom{n}{k} \\ &= \frac{1}{2n+2} \binom{n+b}{n+1}^{-1} (\psi(n+b+1) - \psi(b))^2 \\ & \quad - \frac{1}{2n+2} \binom{n+b}{n+1}^{-1} (\psi_1(n+b+1) - \psi_1(b)), \end{aligned}$$

where $\psi_1(x)$ is the trigamma function defined by

$$\psi_1(x) = \frac{d}{dx} \psi(x) = \sum_{k=0}^{\infty} \frac{1}{(k+x)^2}$$

and $H_r^{(2)}$ is the r^{th} second order harmonic number,

$$H_r^{(2)} = \sum_{j=1}^r \frac{1}{j^2}.$$

The result follows upon using the fact that

$$\psi(x+1) - \psi(y+1) = H_x - H_y \quad (27)$$

and

$$\psi_1(x+1) - \psi_1(y+1) = H_y^{(2)} - H_x^{(2)}, \quad (28)$$

for $x, y \in \mathbb{C} \setminus \mathbb{Z}^-$. \square

Theorem 9. If $n \in \mathbb{N}_0$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$, then

$$\sum_{k=0}^n (-1)^{k-1} \binom{n}{k} H_{k+b}^{(2)} = \frac{H_{n+b} - H_b}{n \binom{n+b}{n}}, \quad n \neq 0, \quad (29)$$

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \frac{H_{k+b} - H_b}{k \binom{k+b}{k}} = H_{n+b}^{(2)} - H_b^{(2)}, \quad (30)$$

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \frac{H_{k+b}}{k \binom{k+b}{k}} = (H_{n+b} - H_b) H_b + H_{n+b}^{(2)} - H_b^{(2)}. \quad (31)$$

Proof. Differentiate (15) with respect to b to get (29). Identity (30) is the inverse of (29). Identity (31) is obtained by using (17) in (30). \square

Theorem 10. If $n \in \mathbb{Z}^+$ and $b \in \mathbb{C} \setminus \mathbb{Z}^-$, then

$$\begin{aligned} \sum_{k=0}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{b+k}{b}} H_k H_{k+b} &= \frac{1}{n+b} \left(b H_b - \frac{n}{n+b} \right) (H_{n+b} - H_b) \\ &+ \frac{b}{n+b} \left(H_{n+b}^{(2)} - H_b^{(2)} \right) + \frac{n}{(n+b)^2} \left(\frac{2}{n+b} + H_b \right). \end{aligned} \quad (32)$$

Proof. Write (24) as

$$\sum_{k=0}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{b+k}{k}} H_k = \frac{b}{n+b} (\psi(n+b+1) - \psi(b+1)) + \frac{n}{(n+b)^2}$$

and differentiate with respect to b , using (10). Use (24) again in simplifying the left hand side of the resulting expression. \square

In particular, for all positive integers n , we have

$$\sum_{k=0}^n (-1)^k \binom{n}{k} H_k^2 = \frac{H_n}{n} - \frac{2}{n^2} \quad (33)$$

and

$$\begin{aligned} \sum_{k=0}^n (-1)^{k-1} \frac{\binom{n}{k}}{\binom{n+k}{k}} H_k H_{k+n} \\ = \frac{1}{2} \left(H_{2n}^{(2)} - H_n^{(2)} \right) + \frac{1}{2} H_{2n} \left(H_n - \frac{1}{2n} \right) - \frac{1}{2} H_n H_{n-1} + \frac{1}{4n^2}. \end{aligned} \quad (34)$$

Corollary 11. If n is a positive integer, then

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \frac{H_k}{k} = H_n^{(2)}. \quad (35)$$

Proof. Treat identity (33) as the binomial transform of the sequence H_n^2 . The inverse relation yields

$$\sum_{k=1}^n (-1)^k \binom{n}{k} \frac{H_k}{k} + 2 \sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k-1}}{k^2} = H_n^2.$$

But (see Boyadzhiev's book [4, p. 64])

$$\sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k-1}}{k^2} = \frac{1}{2} (H_n^2 + H_n^{(2)})$$

and the proof is completed. \square

Remark. Identity (35) can also be obtained by taking the limit of (9) as b approaches zero as well as simply setting $b = 0$ in (30) or (31).

3 Some identities involving odd harmonic numbers

This section contains some combinatorial identities involving odd harmonic numbers O_n , which are defined by

$$O_n = \sum_{k=1}^n \frac{1}{2k-1}, \quad O_0 = 0.$$

Obvious relations between harmonic numbers H_n and odd harmonic numbers O_n are given by

$$H_{2n} = \frac{1}{2}H_n + O_n \quad \text{and} \quad H_{2n-1} = \frac{1}{2}H_{n-1} + O_n. \quad (36)$$

Additional relations are contained in the next lemma.

Lemma 1. *If n is an integer, then*

$$H_{n-1/2} = 2O_n - 2 \ln 2, \quad (37)$$

$$H_{n-1/2} - H_{-1/2} = 2O_n, \quad (38)$$

$$H_{n-1/2} - H_{1/2} = 2(O_n - 1), \quad (39)$$

$$H_{n+1/2} - H_{-1/2} = 2O_{n+1}, \quad (40)$$

$$H_{n+1/2} - H_{1/2} = 2(O_{n+1} - 1), \quad (41)$$

$$H_{n+1/2} - H_{n-1/2} = \frac{2}{2n+1}, \quad (42)$$

$$H_{n-1/2} - H_{-3/2} = 2(O_n - 1), \quad (43)$$

$$H_{n+1/2} - H_{-3/2} = 2(O_{n+1} - 1). \quad (44)$$

Proof. Use (3) as the definition of the harmonic numbers for all complex n (excluding zero and the negative integers) and use the known result for the digamma function at half-integer arguments [12, Eq. (51)], namely,

$$\psi(n + 1/2) = -\gamma - 2 \ln 2 + 2 \sum_{k=1}^n \frac{1}{2k-1}.$$

□

Lemma 2 (Gould [8, Identities Z.45 and Z.51]). *If r and s are integers such that $0 \leq s \leq r$, then*

$$\binom{r+1/2}{s} = \binom{2r+1}{2s} \binom{r}{s}^{-1} 2^{-2s} \binom{2s}{s} \quad (45)$$

and

$$\binom{r-1/2}{s} = \binom{r}{s} \binom{2r-2s}{r-s}^{-1} 2^{-2s} \binom{2r}{r}. \quad (46)$$

We also have

$$\binom{r}{1/2} = \frac{2^{2r+1}}{\pi \binom{2r}{r}}, \quad [8, \text{Identity Z.48}], \quad (47)$$

$$\binom{r}{-1/2} = \frac{2^{2r+1}}{\pi(2r+1) \binom{2r}{r}}, \quad (48)$$

and

$$\binom{r-1/2}{r+1} = -\frac{1}{r+1} \binom{2r}{r} \frac{1}{2^{2r+1}}. \quad (49)$$

Proof. Identity (48) follows from the fact that

$$\binom{r}{-1/2} = \frac{r!}{(-1/2)!(r+1/2)!},$$

while (49) is a consequence of

$$\binom{r-1/2}{r+1} = -\frac{1}{2} \frac{(r-1/2)!}{(r+1)! \sqrt{\pi}}. \quad (50)$$

□

Theorem 12. *If n is a non-negative integer, then*

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(2k+1)^2} = \frac{2^{2n+1}}{n+1} \frac{O_{n+1}}{\binom{2(n+1)}{n+1}}, \quad (51)$$

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{2^{2k+1}}{k+1} \frac{O_{k+1}}{\binom{2(k+1)}{k+1}} = \frac{1}{(2n+1)^2}, \quad (52)$$

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^{k-1}}{(2k-1)^2} = 2^{2n} \frac{O_n - 1}{\binom{2n}{n}}, \quad (53)$$

and

$$\sum_{k=0}^n \binom{n}{k} (-1)^{k-1} 2^{2k} \frac{O_k}{\binom{2k}{k}} = \frac{2n}{(2n-1)^2}. \quad (54)$$

Proof. The first identity is obtained by setting $b = 1/2$ in (12) and using (40) and (45); while the second is its inverse transform. Identity (53) is obtained by setting $b = -1/2$ in (12) and using (43) and (49). To prove identity (54) use the inverse binomial transform of (53) to get

$$\sum_{k=0}^n \binom{n}{k} (-1)^{k-1} 2^{2k} \frac{O_k}{\binom{2k}{k}} - \sum_{k=0}^n \binom{n}{k} (-1)^{k-1} \frac{2^{2k}}{\binom{2k}{k}} = \frac{1}{(2n-1)^2}.$$

But

$$\sum_{k=0}^n \binom{n}{k} (-1)^{k-1} \frac{2^{2k}}{\binom{2k}{k}} = \frac{1}{2n-1} \quad (55)$$

and the proof of (54) is completed. \square

Theorem 13. *If n is a non-negative integer, then*

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} \frac{2k-1}{2^{2(k-1)}} \binom{2(k-1)}{k-1} O_k = \frac{1}{2^{2(n-1)}} \binom{2(n-1)}{n-1} \left(\frac{2n}{2n-1} - O_n \right), \quad (56)$$

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} \frac{2k+1}{2^{2k}} \binom{2k}{k} O_{k+1} = \frac{1}{2n-1} \frac{\binom{2n}{n}}{2^{2n}} \left(\frac{4n-1}{2n-1} - O_n \right), \quad (57)$$

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} 2^{-2k-2} \binom{2(k+1)}{k+1} (O_{k+1} - 1) = \frac{2^{-2n-2}}{2n+1} \binom{2(n+1)}{n+1} \left(O_{n+1} - \frac{1}{2n+1} \right), \quad (58)$$

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} 2^{-2k} \binom{2k}{k} (O_k - 1) = 2^{-2n} \binom{2n}{n} \left(\frac{2n}{2n+1} + O_{n+1} \right). \quad (59)$$

Proof. These results follow from (14). To obtain (56), set $b = -1$, $c = -1/2$ in (14) and use (38) and (48). Identity (57) comes from setting $b = 0$, $c = -1/2$ in (14) and using (40), (47) and (48). \square

Theorem 14. *If n is a positive integer, then*

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k+1} O_k = \frac{2^{2n-1}}{n \binom{2n}{n}}, \quad (60)$$

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k+1} \frac{2^{2k-1}}{k \binom{2k}{k}} = O_n, \quad (61)$$

$$\sum_{k=0}^n \binom{n}{k} (-1)^{k+1} O_{k+1} = \frac{2^{2n-1}}{n(2n+1) \binom{2n}{n}}, \quad (62)$$

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k+1} \frac{2^{2k-1}}{k(2k+1) \binom{2k}{k}} = O_{n+1} - 1. \quad (63)$$

Proof. Write (15) as

$$\sum_{k=0}^n (-1)^{k+1} \binom{n}{k} (H_{k+b} - H_c) = \frac{1}{n \binom{n+b}{n}}.$$

Evaluate at $(b, c) = (-1/2, -1/2)$ and at $(b, c) = (1/2, -1/2)$ to obtain (60) and (62). Identity (61) is the binomial transform of (60) while (63) is the transform of (62). \square

Theorem 15. *If n is a non-negative integer, then*

$$\sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k+1} k}{(2k+1)^2} = \frac{2^{2n-1}(O_{n+1} - 1)}{(2n+1) \binom{2n}{n}}, \quad (64)$$

$$\sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k+1} k}{(2k-1)^2} = \frac{2^{2n-1} O_n}{\binom{2n}{n}}, \quad (65)$$

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k-1} \frac{2^{2k-1}(O_{k+1} - 1)}{(2k+1) \binom{2k}{k}} = \frac{n}{(2n+1)^2}, \quad (66)$$

$$\sum_{k=1}^n \binom{n}{k} (-1)^{k+1} \frac{2^{2k-1} O_k}{\binom{2k}{k}} = \frac{n}{(2n-1)^2}. \quad (67)$$

Proof. The first two results follow from (21). The other two identities are the inverse binomial transforms of the former. \square

Remark. *Identity (67) is a rediscovery of identity (54). Also, combining the proof of (54) with (67) yields (55).*

Theorem 16. *If n is a non-negative integer, then*

$$\sum_{k=0}^n \binom{n}{k} (-1)^{k-1} \frac{2^{2k} H_k}{(2k+1) \binom{2k}{k}} = \frac{2O_{n+1}}{2n+1} - \frac{2}{(2n+1)^2}, \quad (68)$$

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{2^{2k} H_k}{\binom{2k}{k}} = \frac{2O_n}{2n-1} - \frac{4n}{(2n-1)^2}, \quad (69)$$

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{O_{k+1}}{2k+1} = \frac{2^{2n+1} O_{n+1}}{n+1 \binom{2n+2}{n+1}} - \frac{2^{2n-1} H_n}{2n+1 \binom{2n}{n}}, \quad (70)$$

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{O_k}{2k-1} = \frac{2^{2n-1}}{\binom{2n}{n}} (H_n - 2O_n), \quad (71)$$

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \frac{2^{2k}}{\binom{2k}{k}} H_{2k} = \frac{4n}{(2n-1)^2} - \frac{O_n}{2n-1}. \quad (72)$$

Proof. The first two results follow from (24). Identity (70) is a consequence of the inverse binomial relation of (68) in conjunction with identity (51). Identity (71) is a consequence of the inverse binomial relation of (69) in conjunction with identity (65). Identity (72) is a consequence of (67) and (69) on account of the first identity in (36). \square

Theorem 17. *If n is a non-negative integer, then*

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^{k-1}}{(2k-1)^3} = \frac{2^{2n-1}}{\binom{2n}{n}} ((O_n - 1)^2 + O_n^{(2)} + 1). \quad (73)$$

Proof. Set $b = -1/2$ in (25) and use Lemma 1 and identity 46. Use also

$$H_{n-1/2}^{(2)} = -2\zeta(2) + 4O_n^{(2)} \quad (74)$$

which follows from the known value $H_{-1/2}^{(2)} = -2\zeta(2)$ since

$$H_{n-1/2}^{(2)} - H_{-1/2}^{(2)} = 4O_n^{(2)},$$

where $O_n^{(2)}$ is the n^{th} second order odd harmonic number defined by

$$O_n^{(2)} = \sum_{k=1}^n \frac{1}{(2k-1)^2}.$$

\square

Theorem 18. *If n is a positive integer, then*

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} O_k^{(2)} = \frac{2^{2n-1} O_n}{\binom{2n}{n} n}, \quad (75)$$

$$\sum_{k=1}^n (-1)^{k-1} \binom{n}{k} \frac{2^{2k-1} O_k}{\binom{2k}{k} k} = O_n^{(2)}. \quad (76)$$

Proof. Set $b = -1/2$ in (29) and use Lemma 1 and also (46) and (74) to obtain (75). Identity (76) is the inverse of (75). \square

Theorem 19. *If n is a non-negative integer, then*

$$\sum_{k=0}^n (-1)^{k-1} \binom{n}{k} \frac{2^{2k}}{\binom{2k}{k}} H_k O_k = \frac{8n}{(2n-1)^3} - \frac{4n O_n}{(2n-1)^2} - \frac{2O_n^{(2)}}{(2n-1)}, \quad (77)$$

$$\sum_{k=0}^n (-1)^k \binom{n}{k} \frac{2^{2k}}{\binom{2k}{k}} H_k = \frac{2O_n}{(2n-1)} - \frac{4n}{(2n-1)^2}. \quad (78)$$

Proof. Set $b = -1/2$ in (32). Use Lemma 1, (46) and (74). Equate rational coefficients from both sides. Identity (78) is a rediscovery of (69). \square

4 Final comments

In these notes, we have demonstrated how a combinatorial identity attributed to Frisch can be used to prove a series of harmonic number and odd harmonic number identities. Some of the results that we derived are known and here we provide final comments and give further references.

We begin by pointing out that identity (33) can also be found in a recent article by Batir [3, Identity 18]. Next, identity (23) can be restated as

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \frac{b}{k+b} = \prod_{k=1}^n \frac{k}{b+k}.$$

This identity is known and two probabilistic proofs were given recently by Peterson [11] and Nakata [10]. Peterson [11] writes (12) in the form

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \left(\frac{b}{b+k} \right)^2 = \prod_{k=1}^n \frac{k}{b+k} \left(1 + \sum_{k=1}^n \frac{b}{b+k} \right),$$

and also states an expression for the generalization involving an additional parameter m

$$\sum_{k=0}^n \binom{n}{k} (-1)^k \left(\frac{b}{b+k} \right)^m, \quad m \geq 1. \quad (79)$$

In 2019, Bai and Luo [2] derived a new expression for Peterson's identity (79) using a partial fraction decomposition and involving generalized harmonic numbers. As applications of their main result, they stated some harmonic number identities. One of their special cases is our identity (26).

It is remarkable that a simple expression for (79) is also hidden in Frisch's identity (5). Indeed, setting $c = 1$ in (5) yields (23), i.e.,

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{b+k} = \frac{1}{b \binom{n+b}{b}}.$$

Differentiating both sides m times w.r.t. b gives

$$\sum_{k=0}^n \binom{n}{k} \frac{(-1)^k}{(b+k)^m} = \frac{(-1)^{m-1}}{(m-1)!} \frac{d^{m-1}}{db^{m-1}} \frac{1}{b \binom{n+b}{b}}, \quad (80)$$

which shows such a simple expression.

References

- [1] U. Abel, A short proof of the binomial identities of Frisch and Klamkin, *J. Integer Sequences* 23 (2020), Article 20.7.1.
- [2] T.-T. Bai and Q.-M. Luo, A simple proof of a binomial identity with applications, *Montes Taurus J. Pure Appl. Math.* 1 (2) (2019), 13–20.
- [3] N. Batir, Finite binomial sum identities with harmonic numbers, *J. Integer Sequences* 24 (2021), Article 21.4.3.
- [4] K. N. Boyadzhiev, *Notes on the Binomial Transform*, World Scientific, 2018.
- [5] W. Chu, Summation formulae involving harmonic numbers, *Filomat* 26 (1) (2012), 143–152.
- [6] W. Chu and L. De Donno, Hypergeometric series and harmonic number identities, *Adv. Appl. Math.* 34 (2005), 123–137.
- [7] R. Frisch, Sur les semi-invariants et moments employés dans l'étude des distributions statistiques, *Skifter utgitt av Det Norske Videnskaps-Akademi i Oslo, II. Historisk-Filosofisk Klasse*, 1926, No. 3, 87 pp.
- [8] H. W. Gould, *Combinatorial Identities, A Standardized Set of Tables Listing 500 Binomial Coefficient Summations, revised edition*. Published by the author, Morgantown, WV, 1972.
- [9] H. W. Gould and J. Quaintance, Bernoulli numbers and a new binomial transform identity, *J. Integer Sequences* 17 (2014), Article 14.2.2.
- [10] T. Nakata, Another probabilistic proof of a binomial identity, *Fibonacci Quart.* 52 (2) (2014), 139–140.

- [11] J. Peterson, A probabilistic proof of a binomial identity, *American Math. Monthly*, 120 (6) (2013), 558–562.
- [12] H. M. Srivastava and J. Choi, *Series Associated with the Zeta and Related Functions*, Springer Science+Media, B.V., 2001.
- [13] J. Wang and Ch. Wei, Derivative operator and summation formulae involving generalized harmonic numbers, *J. Math. Anal. Appl.* 434 (1) (2016), 315–341.
- [14] Ch. Wei, D. Gong and Q. Yan, Telescoping method, derivative operators and harmonic number identities, *Integral Transforms Spec. Funct.* 25 (3) (2014), 203–214.