

On the treewidth of generalized q -Kneser graphs

Klaus Metsch*

Abstract

The generalized q -Kneser graph $K_q(n, k, t)$ for integers $k > t > 0$ and $n > 2k - t$ is the graph whose vertices are the k -dimensional subspaces of an n -dimensional F_q -vectorspace with two vertices U_1 and U_2 adjacent if and only if $\dim(U_1 \cap U_2) < t$. We determine the treewidth of the generalized q -Kneser graphs $K_q(n, k, t)$ when $t \geq 2$ and n is sufficiently large compared to k . The imposed bound on n is a significant improvement of the previously known bound. One consequence of our results is that the treewidth of each q -Kneser graph $K_q(n, k, t)$ with $k > t > 0$ and $n \geq 3k - t + 9$ is equal to $\binom{n}{k}_q - \binom{n-t}{k-t}_q - 1$.

Keywords: generalized Kneser graph, treewidth, tree decomposition

MSC (2020): 51D05, 51E20

1 Introduction

In this paper we only consider simple *graphs* Γ without loops, that is Γ is a pair (X, E) where X is a non-empty set and E is a set of subsets of cardinality two of X . By graph we always refer to a simple graph without loops. The elements of X are called *vertices* and the elements of E are called *edges* of Γ . We write $X = V(\Gamma)$. A graph is *empty* if it has no edges, and *finite* if it has only finitely many vertices.

A *tree decomposition* of a graph Γ is a pair (T, B) where T is a tree and $B = (B_t : t \in V(T))$ is a collection of subsets B_t of $V(\Gamma)$, indexed by the vertices of T , such that

(TD1) every edge $\{u, v\}$ of Γ is contained in B_t for some $t \in V(T)$, and

(TD2) for each $v \in V(\Gamma)$, the set $\{t \in V(T) \mid v \in B_t\}$ is not empty and the graph induced by T on this set is connected.

Each graph Γ has the trivial tree decomposition where the tree has just one vertex t and $B_t = V(\Gamma)$. The *width* of a tree decomposition of a finite graph is the number $\max\{|B_t| - 1 \mid t \in V(T)\}$, and the *treewidth* $\text{tw}(\Gamma)$ of a finite graph Γ is the smallest width of its tree decompositions. The treewidth of a graph measures how tree-like a graph is. For example, the treewidth of a non-empty tree is one and the treewidth of a

*Justus-Liebig-Universität, Mathematisches Institut, Arndtstraße 2, D-35392 Gießen

graph on n vertices is at most $n - 1$ with equality if and only if the graph is complete. There is a vast literature on the treewidth of graphs, see [2, 5, 6, 9, 10, 11, 14] for some recent ones, and there are applications. A famous one is by Robertson and Seymour [13] when they proved that the treewidth of a minor of a graph cannot exceed the treewidth of the graph.

Let q be a prime power and let F_q be the finite field of order q . For integers n, k, t with $k > t \geq 1$ and $n > 2k - t$, the q -Kneser graph $K_q(n, k, t)$ is the graph whose vertices are the subspaces of dimension k of the vector space F_q^n with two vertices K and K' adjacent if and only if $\dim(K \cap K') < t$. The condition $n > 2k - t$ ensures that the graph is non-empty. Considering the vertices of $K_q(n, k, t)$ as subspaces of the vector space dual to F_q^n , one shows that the graphs $K_q(n, k, t)$ and $K_q(n, n - k, n - 2k + t)$ are isomorphic. It is therefore sufficient to consider Kneser graphs $K_q(n, k, t)$ with $n \geq 2k$. Notice that we do not allow $t = k$, which would yields only complete graphs. In 2021 Cao, Liu, Lu and Lv proved the following theorem.

Theorem 1.1 ([1]). *Let q be the order of a finite field. For integers n, k, t with $k > t \geq 1$ and*

$$n \geq 2t(k - t + 1) + k + 1 \tag{1}$$

we have $\text{tw}(K_q(n, k, t)) = \begin{bmatrix} n \\ k \end{bmatrix}_q - \begin{bmatrix} n-t \\ k-t \end{bmatrix}_q - 1$.

In this result the Gaussian coefficient $\begin{bmatrix} n \\ k \end{bmatrix}_q$ is defined for integers $n \geq k \geq 0$ and $q \geq 2$ as follows.

$$\begin{bmatrix} n \\ k \end{bmatrix}_q := \prod_{i=0}^{k-1} \frac{q^{n-i} - 1}{q^i - 1}. \tag{2}$$

The hard part of the theorem is to prove the lower bound for the treewidth. For the upper bound, the authors of [1] proved in fact that $\text{tw}(K_q(n, k, t)) \leq \begin{bmatrix} n \\ k \end{bmatrix}_q - \begin{bmatrix} n-t \\ k-t \end{bmatrix}_q - 1$ holds whenever $k > t \geq 1$ and $n \geq 2k$. In this paper we give a shorter proof for this upper bound and we will weaken the required condition (1) for the lower bound substantially. Our main result is the following. To formulate it, we define the map ϵ from the set of all primepowers to the set $\{0, 1, 2, 3, 4\}$ by $\epsilon(2) = 9$, $\epsilon(3) = 3$, $\epsilon(4) = 2$, $\epsilon(q) = 1$ for $5 \leq q \leq 8$, and $\epsilon(q) = 0$ for $q \geq 9$.

Theorem 1.2. *Consider a q -Kneser graph $K_q(n, k, t)$ with $k > t \geq 1$ and $n \geq 2k$. Suppose that one of the following condition is satisfied.*

- (1) $t \leq \epsilon(q)$ and $n > 3k - 2t + \epsilon(q)$.
- (2) $t > \epsilon(q)$ and $n > 3k - t + 1 - 2\sqrt{t - \epsilon(q)}$.

Then $\text{tw}(K_q(n, k, t)) = \begin{bmatrix} n \\ k \end{bmatrix}_q - \begin{bmatrix} n-t \\ k-t \end{bmatrix}_q - 1$.

The strongest result is obtained for $q \geq 9$, where the required bound is only $n > 3k - t + 1 - 2\sqrt{t}$. As a simpler formulation not involving $\epsilon(q)$ we can state the following.

Corollary 1.3. For all integers $k > t > 0$ and $n \geq 3k - t + 9$ and any prime power q , the q -Kneser graph $K_q(n, k, t)$ has treewidth $\binom{n}{k}_q - \binom{n-t}{k-t}_q - 1$.

For some pairs (k, t) and fields F_q our result determines the treewidth for all $n \geq 2k$. We formulate this in the following corollary. For simplicity we only consider the case when q is at least nine.

Corollary 1.4. Let $q \geq 9$ be the order of a finite field and let k and t be integers with $k > t \geq 1$, and $t > k + 3 - 2\sqrt{k + 2}$. Then

$$\text{tw}(K_q(n, k, t)) = \binom{n}{k}_q - \binom{n-t}{k-t}_q - 1$$

for all $n \geq 2k$.

Remarks 1.5. 1. For the non-modular Kneser graphs $K(n, k, t)$ similar results have been proved in [4], [10] and [12].

2. As was shown in [1], see also Corollary 2.4, we have $\text{tw}(\Gamma) \leq |V(\Gamma)| - \alpha(\Gamma) - 1$ for every q -Kneser graph Γ . Notice however that the independence number of $K_q(n, k, t)$ is $\binom{n-t}{k-t}_q$ when $n \geq 2k$ and $\binom{2k-t}{k-t}_q$ otherwise.

The Kneser graph $K_q(n, k, k-1)$ is the dual of the Grassmann graph. Part (2) of Theorem 1.2 determines its treewidth for all $k \geq 3 + \epsilon(q)$ and $n \geq 2k$. This was proved in [1] more generally for all $k \geq 3$ and also in the case $k = 2$ and $n \geq 5$. The case $K_q(4, 2, 1)$ remained open in [1]. Our last results settles this remaining open case. This requires a more detailed look at tree decompositions.

Theorem 1.6. The treewidth of $K_q(4, 2, 1)$ is $|V(\Gamma)| - (q^2 + q + 2)$ for all prime powers $q \geq 2$.

2 An upper bound for the treewidth

For every finite graph Γ its maximum vertex degree is denoted by $\Delta(\Gamma)$ and is called its *maximum degree*. The cardinality of a largest independent set of Γ is denoted by $\alpha(\Gamma)$ and is called the *independence number* of the graph. The following connection between treewidth, maximum degree and independence number is known.

Result 2.1 ([4]). For every finite graph Γ we have

$$\text{tw}(\Gamma) \leq \max\{\Delta(\Gamma), |V(\Gamma)| - \alpha(\Gamma) - 1\}. \quad (3)$$

For generalized q -Kneser graphs we have $\Delta(\Gamma) \leq |V(\Gamma)| - \alpha(\Gamma) - 1$ as was shown for $n \geq 2k$ in [1]. We give a simpler proof for this inequality. In order to do so, we need the following result of Wilson on the independence number of generalized Kneser graphs.

Result 2.2 ([3]). *For integers $k > t \geq 1$ and $n > 2k - t$ we have*

$$\alpha(K_q(n, k, t)) = \max\left\{\binom{n-t}{k-t}_q, \binom{2k-t}{k-t}_q\right\}.$$

Lemma 2.3. *For Kneser graphs $K_q(n, k, t)$ with $n > 2k - t$ and $k > t \geq 1$ we have*

$$\Delta(K_q(n, k, t)) \leq \binom{n}{k}_q - \alpha(K_q(n, k, t)) - 1.$$

Proof. Since $K_q(n, k, t)$ and $K_q(n, n-k, n-2k+t)$ are isomorphic graphs, we may assume that $n \geq 2k$. The graph $\Gamma := K_q(n, k, t)$ has $\binom{n}{k}_q$ vertices and is regular. Given a vertex A , then $r := \binom{n}{k}_q - \Delta(\Gamma)$ is the number of vertices of the graph that are not adjacent to it. Considering A as a k -subspace of F_q^n , then r is the number of k -subspaces meeting A in a subspace of dimension at least t . We have to show that this number is strictly larger than $\alpha(\Gamma)$.

From 2.2 we have $\alpha(\Gamma) = \binom{n-t}{k-t}_q$. Let T_0 be a t -subspace of A . Then T is contained in $\binom{n-t}{k-t}_q$ k -subspaces. Let T be a second t -subspace of A . Notice that T exists as $t < k$. Then there exist k -subspaces meeting A exactly in T . This shows that $r > \binom{n-t}{k-t}_q = \alpha(\Gamma)$ as desired. \square

Corollary 2.4. *For Kneser graphs $K_q(n, k, t)$ with $n > 2k - t$ and $1 \leq t < k$ we have*

$$\text{tw}(K_q(n, k, t)) \leq \binom{n}{k}_q - \alpha(K_q(n, k, t)) - 1.$$

As mentioned before, it was proved in [1] that this upper bound is sharp when n is sufficiently large compared to k and t . We will improve this result by weakening the required bound on n significantly. The approach in [1] and [4] uses a result of Robertson and Seymour on separators. We use the slightly different result (2.5) from the same paper and modify it in the following lemma. Recall that a component of a graph is a maximal connected induced subgraph of Γ . Hence, every vertex of the graph is a vertex of exactly one component.

Lemma 2.5 (based on [13]). *Let Γ be a finite graph. Then there exists a tree representation (T, B) with $B = (B_t \mid t \in V(T))$ with width $\text{tw}(\Gamma)$ and the following properties.*

- (a) *If $t, t' \in T$ are adjacent in T , then B_t is not a subset of $B_{t'}$.*
- (b) *For some $t \in T$, there exists a subset P of B_t such that $|P| \leq \text{tw}(\Gamma) + 1$ and such that every component of $\Gamma \setminus P$ has at most $\frac{1}{2}|V(\Gamma) \setminus P|$ vertices.*

Proof. Let (T, B) with $B = (B_t \mid t \in V(T))$ be a tree decomposition of Γ with width $\text{tw}(\Gamma)$. If there exist adjacent vertices x, y of T with $B_x \subseteq B_y$, let T' be the

graph obtained from T by contracting the edge $\{x, y\}$ to a new vertex t_0 , and define $B_{t_0} := B_x$. Then $(T', (B_t \mid t \in V(T')))$ is a tree decomposition of Γ with the same width as (T, B) and one vertex less than T . Repeating this construction several times if necessary we finally obtain a tree decomposition satisfying the first condition of the lemma and having still tree width $\text{tw}(\Gamma)$. We may thus assume that (T, B) satisfies the first condition.

Claim (2.5) in [13] states that there exists a subset P of $V(\Gamma)$ such that every component of $\Gamma \setminus P$ has at most $\frac{1}{2}|V(\Gamma) \setminus P|$ vertices. Using the above tree decomposition, the proof of (2.5) in [13] in fact constructs such a set P as a subset of some set B_t . Thus the second condition is satisfied. \square

3 A lower bound for the treewidth

Lemma 3.1. *Consider integers n, k, q with $n \geq k \geq 0$ and $q \geq 2$. Then*

$$\begin{bmatrix} n \\ k \end{bmatrix}_q \leq (q + \beta)q^{k(n-k)-1}.$$

where $\beta = 5$ for $q = 2$, and $\beta = 3$ for $q = 3$ and $\beta = 2$ for $q \geq 4$. Also, if $0 < k < n$, then $(q + 1)q^{k(n-k)-1} \leq \begin{bmatrix} n \\ k \end{bmatrix}_q$.

Proof. We first prove the lower bound for the Gaussian coefficient for $0 < k < n$ by induction on k . For $k = 1$ we have

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \sum_{i=0}^{n-1} q^i \geq (q + 1)q^{n-2}$$

as required. For $k \geq 2$, we have

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{q^n - 1}{q^k - 1} \begin{bmatrix} n - 1 \\ k - 1 \end{bmatrix}_q \geq q^{n-k} \begin{bmatrix} n - 1 \\ k - 1 \end{bmatrix}_q$$

and the induction hypothesis shows that this is at least $(q + 1)q^{k(n-k)-1}$.

For $q \geq 3$, the upper bound was proved in [8, Lemma 34]. Now consider $q = 2$. We have to prove the upper bound, which we do using the technique of [8, Lemma 34]. From the definition of the Gaussian coefficient, we have

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \prod_{i=1}^k \frac{q^{n-k+i} - 1}{q^i - 1} \leq \prod_{i=1}^k \frac{q^{n-k+i}}{q^i - 1} = q^{k(n-k)} \prod_{i=1}^k \frac{q^i}{q^i - 1}. \quad (4)$$

It suffices therefore to show that $\prod_{i=1}^k \frac{q^i}{q^i - 1} \leq 1 + \frac{\beta}{q} = \frac{7}{2}$. For $1 \leq k \leq 5$, one verifies this by hand. For $k \geq 6$, one proves easily by induction on k the stronger statement $\prod_{i=1}^k \frac{q^i}{q^i - 1} \leq 1 + 5 \cdot \frac{q^{k-1} - 2}{q^k - 2}$. \square

Lemma 3.2. *Let K_1 and K_2 be two subspaces of dimension k of an F_q -vector space and let s be the dimension of their intersection. Then for every integer i with $0 \leq i \leq s$ the number of pairs (T_1, T_2) consisting of a t -subspace T_1 of K_1 and a t -subspace T_2 of K_2 such that $\dim(T_1 \cap T_2) = i$ is at most*

$$\begin{bmatrix} s \\ i \end{bmatrix}_q \begin{bmatrix} k-i \\ t-i \end{bmatrix}_q^2.$$

Proof. For any such pair, the intersection of T_1 and T_2 is a subspace of the s -subspace $K_1 \cap K_2$. Now $K_1 \cap K_2$ has $\begin{bmatrix} s \\ i \end{bmatrix}_q$ i -subspaces and each such i -subspace lies in $\begin{bmatrix} k-i \\ t-i \end{bmatrix}_q$ t -subspaces of K_1 and in as many of K_2 . The statement follows. \square

Lemma 3.3. *Let f be a real quadratic polynomial with leading coefficient -1 , and let f obtain its maximum for $x = x_0$. Let q and a be integers and suppose that $q \geq 2$.*

- (a) *If $x_0 \leq a$, then $\sum_{i=a}^{\infty} q^{f(i)} < q^{f(a)}(1 + \frac{1}{q} + \frac{1}{q^3})$.*
- (b) *If $x_0 \geq a$, then $\sum_{i=-\infty}^a q^{f(i)} < q^{f(a)}(1 + \frac{1}{q} + \frac{1}{q^3})$.*
- (c) *If $2x_0$ is an integer, then $\sum_{i \in \mathbb{Z}} q^{f(i)} < q^{f(x_0)}(1 + \frac{2}{q} + \frac{2}{q^3})$.*

Proof. We use several times that $\sum_{i=0}^{\infty} q^{-i} = q/(q-1) \leq q$.

(a) Since $x_0 \leq a$, then $f(a+i) \leq f(a) - i^2$ for all $i \geq 0$ and hence

$$\sum_{i=a}^{\infty} q^{f(i)} \leq q^{f(a)} \sum_{i=0}^{\infty} q^{-i^2} < q^{f(a)} \left(1 + \frac{1}{q} + q^{-4} \sum_{i=0}^{\infty} q^{-i} \right) \leq q^{f(a)} \left(1 + \frac{1}{q} + \frac{1}{q^3} \right).$$

(b) This follows from (a) and the symmetry of f .

(c) If x_0 is an integer, this follows from (a) and (b) applied with $a = x_0$. Now consider the case when x_0 is not an integer but $2x_0$ is. If $z \in \mathbb{Z}$, then $i := z - x_0 - \frac{1}{2} \in \mathbb{Z}$ and $f(z) = f(x_0) - (\frac{1}{2} + i)^2$. Hence

$$\begin{aligned} \sum_{z \in \mathbb{Z}} q^{f(z)} &= q^{f(x_0)} \sum_{i \in \mathbb{Z}} q^{-(1/2+i)^2} = 2q^{f(x_0) - \frac{1}{4}} \sum_{i=0}^{\infty} q^{-i(i+1)} \\ &\leq 2q^{f(x_0) - \frac{1}{4}} (1 + q^{-2} + q^{-6} \sum_{i=0}^{\infty} q^{-i}) \\ &\leq 2q^{f(x_0) - \frac{1}{4}} (1 + q^{-2} + q^{-5}) \\ &< q^{f(x_0)} (1 + \frac{2}{q} + \frac{2}{q^3}) \end{aligned}$$

where the last step is obvious for $q \geq 16$ and easily checked for $2 \leq q \leq 15$. \square

The next theorem is a reformulation of Theorem 1.2.

Theorem 3.4. *Let $\Gamma = K_q(n, k, t)$ with $n \geq 2k$ and $k > t \geq 1$. Assume that $\text{tw}(\Gamma) \neq \binom{n}{k}_q - \alpha(\Gamma) - 1$. Then the following hold where ϵ is the function defined in the introduction.*

(a) *If $t \leq \epsilon(q)$, then $n \leq 3k - 2t + \epsilon(q)$.*

(b) *If $t > \epsilon(q)$, then $n \leq 3k - t + 1 - 2\sqrt{t - \epsilon(q)}$.*

Proof. Lemma 2.5 shows that there exists a subset P of $V(\Gamma)$ such that $|P| \leq \text{tw}(\Gamma) + 1$ and such that every component of $\Gamma \setminus P$ has at most $\frac{1}{2}|V(\Gamma) \setminus P|$ vertices. Consider $Y := V(\Gamma) \setminus P$.

Since $\text{tw}(\Gamma) \neq \binom{n}{k}_q - \alpha(\Gamma) - 1$, Corollary 2.4 shows that $\text{tw}(\Gamma) \leq \binom{n}{k}_q - \alpha(\Gamma) - 2$. Since $|P| \leq \text{tw}(\Gamma) + 1$, it follows that $|Y| \geq \alpha(\Gamma) + 1$. Hence Y is not an independent set of Γ and thus Γ has an edge whose vertices S_1 and S_2 are in Y . Let X be the vertex set of the component of $\Gamma \setminus P$ that contains S_1 and S_2 . Then $|X| \leq \frac{1}{2}|Y|$. Hence $|Y \setminus X| \geq \frac{1}{2}|Y| > \frac{1}{2}\alpha(\Gamma)$.

As vertices of the graph Γ , S_1 and S_2 are connected, so as k -subspaces we have $s := \dim(S_1 \cap S_2) \leq t - 1$. As vertices of the graph, the elements of $Y \setminus X$ are neither adjacent to S_1 nor to S_2 , so as k -subspaces, every element of $Y \setminus X$ meets each of the k -subspaces S_1 and S_2 in a subspace of dimension at least t . We define

$$\begin{aligned} M &:= \{(T_1, T_2) \mid T_i \text{ is a subspace of } S_i \text{ with } \dim(T_i) = t, i = 1, 2\} \\ C &:= \{((T_1, T_2), K) \in M \times (Y \setminus X) \mid T_1, T_2 \subseteq K\}. \end{aligned}$$

As every element of $Y \setminus X$ meets S_1 and S_2 in subspaces of dimension at least t , then every element of $Y \setminus X$ occurs in a pair of C and hence $|C| \geq |Y \setminus X| > \frac{1}{2}\alpha(\Gamma)$.

Consider an element (T_1, T_2) of M that occurs in at least one pair of C . Then $\dim(T_1 + T_2) \leq k$ and hence $\dim(T_1 \cap T_2) \geq 2t - k$. As $T_1 \cap T_2 \subseteq S_1 \cap S_2$ then $\dim(T_1 \cap T_2) \leq s$. Hence, if $i = \dim(T_1 \cap T_2)$, then $2t - k \leq i \leq s$ and the number of k -subspaces of F_q^n that contain T_1 and T_2 is $\binom{n-2t+i}{k-2t+i}_q$; consequently (T_1, T_2) lies in at most this many pairs of C . Hence with

$$i_{\max} := \max\{0, 2t - k\}$$

we deduce from 2.2 and 3.2 that

$$\frac{1}{2} \binom{n-t}{k-t}_q \stackrel{2.2}{=} \frac{1}{2} \alpha(\Gamma) < |C| \stackrel{3.2}{\leq} \sum_{i=i_{\max}}^s \binom{s}{i}_q \binom{k-i}{t-i}_q^2 \binom{n-2t+i}{k-2t+i}_q. \quad (5)$$

We define

$$\beta := \begin{cases} 5 & \text{if } q = 2, \\ 3 & \text{if } q = 3, \\ 2 & \text{if } q \geq 4, \end{cases}$$

and apply Lemma 3.1 to both sides of inequality (5) to find

$$\frac{1}{2}(q+1)q^{(n-k)(k-t)-1} \leq \sum_{i=i_{\max}}^s (q+\beta)^4 q^{(s-i)i-1} (q^{(k-t)(t-i)-1})^2 q^{(n-k)(k-2t+i)-1}.$$

Using $s \leq t-1$ it follows that

$$\begin{aligned} \frac{(q+1)q^3}{2(q+\beta)^4} &\leq \sum_{i=i_{\max}}^s q^{(s-i)i} q^{2(k-t)(t-i)} q^{(n-k)(-t+i)} \\ &\leq \sum_{i=i_{\max}}^{t-1} q^{(t-1-i)i} q^{2(k-t)(t-i)} q^{(n-k)(-t+i)} \\ &= \sum_{i=i_{\max}}^{t-1} q^{(t-i)(i+3k-2t-n)-i}. \end{aligned}$$

Using the definition of β and $\epsilon(q)$ it is straightforward to check that

$$q^{-\epsilon(q)-\frac{3}{4}} \left(1 + \frac{2}{q} + \frac{2}{q^3}\right) < \frac{(q+1)q^3}{2(q+\beta)^4}. \quad (6)$$

Defining the quadratic function $f(i) = (t-i)(i+3k-2t-n) - i$ we thus find

$$q^{-\epsilon(q)-\frac{3}{4}} \left(1 + \frac{2}{q} + \frac{2}{q^3}\right) < \sum_{i=i_{\max}}^{t-1} q^{f(i)}. \quad (7)$$

The parabola f reaches its maximum for $i = i_0$ where

$$i_0 := \frac{1}{2}(n-3k+3t-1) \text{ and } f(i_0) = \frac{1}{4}(3k+1-t-n)^2 - t. \quad (8)$$

Lemma 3.3 (c) implies that the right hand side of (7) is at most $(1 + \frac{2}{q} + \frac{2}{q^3})q^{f(i_0)}$. Therefore (7) implies that $f(i_0) > -\frac{3}{4} - \epsilon(q)$. Since $f(i_0)$ or $f(i_0) - \frac{1}{4}$ is an integer, it follows that $f(i_0) \geq -\epsilon(q)$. If $i_0 \leq i_{\max}$ or $i_0 \geq t-1$, we can improve this by applying (a) or (b) of Lemma 3.3. Since $f(i_{\max})$ and $f(t-1)$ are integers, we find

$$-\epsilon(q) \leq \begin{cases} f(i_0) & \text{in any case,} \\ f(t-1) & \text{if } t-1 \leq i_0, \\ f(i_{\max}) & i_0 \leq i_{\max}. \end{cases} \quad (9)$$

Case 1. Here we consider the situation when $n \geq 3k-t-1$.

Then (8) gives $i_0 \geq t-1$ so (9) implies that $-\epsilon(q) \leq f(t-1)$. Since $f(t-1) = 3k-2t-n$, it follows that $n \leq 3k-2t+\epsilon(q)$. Hence $3k-t-1 \leq n \leq 3k-2t+\epsilon(q)$, which implies that $t \leq \epsilon(q)+1$. If $t \leq \epsilon(q)$, then we are in situation (a) of the statement, and if $t = \epsilon(q)+1$, we are situation (b) of the statement. Hence, in this case the theorem is proved.

Case 2. Here we consider the case that $n \leq 3k - t - 2$.

If $t \leq \epsilon(q)$, then we have $n \leq 3k - t - 2 \leq 3k - 2t + \epsilon(q)$ and we are in situation (a) of the statement. Suppose now that $t > \epsilon(q)$. From (9) we find $-\epsilon(q) \leq f(i_0)$, that is

$$t - \epsilon(q) \leq \frac{1}{4}(3k + 1 - t - n)^2.$$

Since $n \leq 3k - t - 2$ and $t > \epsilon(q)$, it follows that $n \leq 3k - t + 1 - 2\sqrt{t - \epsilon(q)}$. Now we are situation (b) of the statement. Hence, also in this case the theorem is proved. \square

Theorem 1.2 follows immediately from Theorem 3.4. Corollary 1.3 follows from Theorem 1.2.

Proof of Corollary 1.4. Consider a Kneser graph $K_q(n, k, t)$ as in 1.4, that is with $q \geq 9$ and $t > k + 3 - 2\sqrt{k + 2}$. From $t > k + 3 - 2\sqrt{k + 2}$ it follows that $k - t + 1 < 2\sqrt{t}$. Hence in part (b) of Theorem 1.2 we have $3k - t + 1 - 2\sqrt{t} \leq 2k$. Therefore 1.4 follows from Theorem 1.2.

4 The Kneser graph $K_q(4, 2, 1)$

The graph $K_q(n, k, t)$ with $t = k - 1$ is the complement of a Grassmann graph. This graph attracted special attention in [1] where it was shown for all $n \geq k \geq 2$ with the exception $(n, k) = (4, 2)$ that the graph $K_q(n, k, k - 1)$ has treewidth $|V(\Gamma)| - \alpha(\Gamma) - 1$. The case $K_q(4, 2, 1)$ remained unsolved in [1] and also our counting argument from the last section is not strong enough to cover this case. We will choose a special model for this graph and use geometric properties to determine its treewidth. The case $q = 2$ will lead to extra difficulties, whereas the case $q \geq 3$ only requires some basic knowledge of tree decompositions.

Using projective geometry, the graph $K_q(4, 2, 1)$ can be understood as the graph whose vertices are the lines of the projective space $\text{PG}(3, q)$ with two vertices adjacent if and only if the lines are skew. Using the Klein-correspondence from $\text{PG}(3, q)$ to the hyperbolic quadric $Q^+(5, q)$, we can define the same graph as follows. Its vertices are the points of $Q^+(5, q)$ and two vertices are non-adjacent if and only if they lie on a line of $Q^+(5, q)$. We consider $Q^+(5, q)$ naturally embedded in $\text{PG}(5, q)$ and denote the related polarity of $\text{PG}(5, q)$ by \perp . Then two points v and v' of $Q^+(5, q)$ are adjacent as vertices of the graph if and only if the points are not perpendicular, that is v' is not a point of the tangent hyperplane v^\perp at v . See [7] for properties of the hyperbolic quadric and the fact that this graph is indeed isomorphic to $K_q(4, 2, 1)$. Using this model we will show that $\text{tw}(\Gamma) = |V(\Gamma)| - \alpha(\Gamma) - 1$. We need two lemmata of preparation, the first one is an easy observation of the hyperbolic quadric $Q^+(3, q)$, which occurs in $Q^+(5, q)$ as an intersection of $Q^+(5, q)$ with ℓ^\perp , where ℓ is any secant line to the quadric.

Lemma 4.1. *The largest sets of points of the hyperbolic quadric $Q^+(3, q)$ without three pairwise non-collinear points are the unions of two skew lines with $2q + 2$ points.*

Proof. Notice that the hyperbolic quadric $Q^+(3, q)$ is a $(q+1) \times (q+1)$ -grid, so it can be described as follows: Its points are (i, j) for $0 \leq i, j \leq q$ and its lines are $\{(i, j) \mid 0 \leq i \leq q\}$ and $\{(j, i) \mid 0 \leq j \leq q\}$ for $0 \leq i \leq q$. The assertion is easily proved from this. \square

The next lemma collects some properties of tree decompositions.

Lemma 4.2. *Let (T, B) with $B = (B_t \mid t \in V(T))$ be a tree decomposition of a finite graph Γ . Let t be a vertex of T . Then we have the following.*

- (a) *If $t \in V(T)$ and $v \in V(\Gamma) \setminus B_t$, then $T \setminus \{t\}$ has exactly one component such that $v \in B_s$ for some vertex s of this component. We denote the vertex set of this component by $T_t(v)$ and we denote by $\Gamma_t(v)$ the union of the sets B_s with $s \in T_t(v)$.*
- (b) *For adjacent vertices v, w of Γ with $v \notin B_t$, we have $w \in \Gamma_t(v)$.*
- (c) *For adjacent vertices v, w of Γ with $v, w \notin B_t$ we have $\Gamma_t(v) = \Gamma_t(w)$.*
- (d) *If v is a vertex of a component C of $\Gamma \setminus B_t$, then $\Gamma_t(v)$ contains every vertex of C and every vertex of Γ that has a neighbor in C .*

Proof. Part (a) follows from the second property of the definition of a tree decomposition. Part (b) then follows from the fact that the edge $\{v, w\}$ of Γ is contained in B_s for some vertex s of T , again by the definition of a tree decomposition. Part (c) follows from (a) and (b). For part (d), we first apply (c) to see that $C \subseteq \Gamma_t(v)$ and then (b) to see that every vertex with a neighbor in C is contained in $\Gamma_t(v)$. \square

Theorem 4.3. *The treewidth of $K_q(4, 2, 1)$ is $|V(\Gamma)| - (q^2 + q + 2)$ for all prime powers q .*

Proof. We represent $\Gamma := K_q(n, 2, 1)$ by $Q^+(5, q)$ as explained above. We also consider the ambient projective space $\text{PG}(5, q)$ of $Q^+(5, q)$ and the related polarity \perp . For each set X of points of $Q^+(5, q)$, we denote by $\langle X \rangle$ the subspace of $\text{PG}(5, q)$ that is spanned by the points of X . The set $\langle X \rangle^\perp \cap Q^+(5, q)$ consists of the points of $Q^+(5, q)$ that are perpendicular to all points of X . In Γ the set $\langle X \rangle^\perp \cap Q^+(5, q)$ consists of the vertices that are not adjacent to any vertex in X (notice that vertices of X may lie in this set).

From Result 2.2 we have $\alpha(\Gamma) = q^2 + q + 1$ and Corollary 2.4 shows that $\text{tw}(\Gamma) \leq |V(\Gamma)| - \alpha(\Gamma) - 1$. Assume that

$$\text{tw}(\Gamma) \leq |V(\Gamma) - \alpha(\Gamma) - 2 \leq |V(\Gamma)| - q^2 - q - 3.$$

We shall derive a contradiction.

Lemma 2.5 shows that there exists a tree decomposition (T, B) , $B = (B_t \mid t \in V(T))$, of Γ , a vertex $t_0 \in V(T)$ and a subset P of B_{t_0} with the following properties

- (T1) The tree decomposition (T, B) has width $\text{tw}(\Gamma)$.
- (T2) For adjacent vertices s and t of T we have $B_s \not\subseteq B_t$.

(T3) With $Y := V(\Gamma) \setminus P$, every component of the graph Γ_Y induced by Γ on Y has at most $\frac{1}{2}|Y|$ vertices.

As $|P| \leq |B_{t_0}| \leq \text{tw}(\Gamma) + 1$, then

$$|Y| = |V(\Gamma)| - |P| \geq |V(\Gamma)| - |B_{t_0}| \geq |V(\Gamma)| - \text{tw}(\Gamma) - 1 \geq q^2 + q + 2. \quad (10)$$

Hence $|Y| > \alpha(\Gamma)$ and thus Y contains adjacent vertices v_1 and v_2 . Let X be the vertex set of the component of Γ_Y that contains v_1 and v_2 . Then no vertex of $Y \setminus X$ is adjacent to any vertex of X . Since X is a component of Γ_Y , we have $|X| \leq \frac{1}{2}|Y|$ and hence $|Y \setminus X| \geq \frac{1}{2}|Y| \geq \frac{1}{2}(q^2 + q + 2) > q + 1$.

On $Q^+(5, q)$ the set Y is a set of points, X is a subset of Y , v_1 and v_2 are non-perpendicular points of X , and every point of $Y \setminus X$ is distinct and perpendicular to every point of X . Therefore $Y \setminus X$ is a subset of $\langle X \rangle^\perp \cap Q^+(5, q)$. Hence the subspaces $\langle X \rangle$ and $\langle Y \setminus X \rangle$ are perpendicular.

Case 1. There exists three pairwise non-perpendicular points in Y .

We may assume that these are v_1 and v_2 and a third point v_3 . Then $v_3 \in X$ and v_1, v_2, v_3 are pairwise non-collinear points of $Q^+(5, q)$. Therefore $\pi := \langle v_1, v_2, v_3 \rangle$ is a conic plane, that is a plane of $\text{PG}(5, q)$ that meets $Q^+(5, q)$ in the $q+1$ points of a conic. We have $Y \setminus X \subseteq \langle X \rangle^\perp \subseteq \pi^\perp$. As π is a conic plane, then π^\perp is also a conic plane and hence has $q+1$ points on $Q^+(5, q)$. Therefore $|Y \setminus X| \leq |\pi^\perp \cap Q^+(5, q)| = q+1$. But we have seen above that $|Y \setminus X| > q+1$. This is a contradiction.

Case 2. Y does not contain three pairwise non-perpendicular points.

The line ℓ of $\text{PG}(5, q)$ on v_1 and v_2 is a secant line of $Q^+(5, q)$ and hence ℓ^\perp is a 3-space of $\text{PG}(5, q)$ that meets $Q^+(5, q)$ in a hyperbolic quadric $Q^+(3, q)$. We have $Y \setminus X \subseteq \langle X \rangle^\perp \subseteq \ell^\perp$, so $Y \setminus X$ is contained in this hyperbolic quadric and does not contain three pairwise non-collinear points. As $|Y \setminus X| > q+1$, then $\langle Y \setminus X \rangle$ is either the solid ℓ^\perp or a plane of this solid.

Case 2.1. $\langle Y \setminus X \rangle$ is the solid ℓ^\perp .

Then $Y \setminus X$ is a subset of $\ell^\perp \cap Q^+(5, q)$, which is the hyperbolic quadric $Q^+(3, q)$ already mentioned above. Lemma 4.1 shows therefore that $Y \setminus X$ contains at most $2(q+1)$ points. As $\langle Y \setminus X \rangle = \ell^\perp$, then $\ell = \langle Y \setminus X \rangle^\perp$ and since X and $Y \setminus X$ are perpendicular, it follows that $X \subseteq \ell$, that is $X = \{v_1, v_2\}$. It follows that $|Y| = |X| + |Y \setminus X| \leq 2 + 2(q+1)$. Therefore (10) implies that $q = 2$ and $|Y| = 8$ and $|X| = 2$. Hence $Y \setminus X$ consists of six points of $Q^+(3, q)$ and Lemma 4.1 shows that these points are the points of two skew lines of $Q^+(3, q)$. The graph induced by Γ on $Y \setminus X$ is therefore a 6-cycle. Since Y has eight points, this contradicts property (T3) of the tree decomposition (T, B) .

Case 2.2 $\pi := \langle Y \setminus X \rangle$ is a plane of ℓ^\perp .

As $|Y| > q+1$, then π meets $Q^+(5, q)$ in the union of two lines. Let z be the point of intersection of these two lines. The subspace π^\perp is also a plane that meets $Q^+(5, q)$ in the union of two lines and z is also their intersection point. Since two intersecting lines contain $2q+1$ points, then X and $Y \setminus X$ have at most $2q+1$ points, but since

z belongs to at most one of these sets, we have $|Y| = |X| + |Y \setminus X| \leq 4q + 1$. Since $|Y| \geq q^2 + q + 2$, it follows that $q = 2$ and $|Y| \in \{8, 9\}$ and (10) can be written as follows

$$|V(\Gamma)| - 9 \leq |V(\Gamma)| - |Y| = |P| \leq |B_{t_0}| \leq \text{tw}(\Gamma) + 1 \leq |V(\Gamma)| - 8. \quad (11)$$

Clearly in (11) all inequalities are sharp except one that is missing sharpness by one. Recall that $V(\Gamma) \setminus Y = P \subseteq B_{t_0}$, so either $P = B_{t_0}$ and $B_{t_0} \cap Y = \emptyset$ or otherwise $|B_{t_0}| = |P| + 1$ and $|Y \cap B_{t_0}| = 1$.

Denote by \bar{Y} the set consisting of the $4q + 1 = 9$ points of $Q^+(5, q)$ in $\pi \cup \pi^\perp$, which are z and eight points of the two 4-cycles. Then $Y \subseteq \bar{Y}$. From (11) we have $|V(\Gamma) \setminus B_{t_0}| \in \{8, 9\}$. Since $V(\Gamma) \setminus B_{t_0} \subseteq V(\Gamma) \setminus P = Y \subseteq \bar{Y}$, it follows that $|B_{t_0} \cap \bar{Y}| \leq 1$. The graph induced by Γ on \bar{Y} is the disjoint union of the singleton z and two 4-cycles, each 4-cycle consisting of the four points of $Q^+(5, q)$ other than z of one of the planes π and π^\perp . We refer to these two 4-cycles as *the 4-cycles* for the rest of the proof. We denote a 4-cycle by $uvwx$ and mean hereby that $u \sim v \sim w \sim x \sim u$ but there are no other adjacencies between the vertices u, v, w and x ; in $Q^+(5, q)$, the points z, v, x as well as the points z, w, u are the points of the two lines of π or π^\perp on z . Notice that any three vertices of the 4-cycle in π span π , so that every point that is perpendicular to these three vertices lies in π^\perp . The same holds for the 4-cycle in π^\perp . Hence we have:

(E) Given three vertices of one of the 4-cycles, the only vertices that have no neighbor among these three vertices are z and the vertices of the second 4-cycle.

Case 2.2.1: $z \notin B_{t_0}$.

We have $|B_{t_0} \cap \bar{Y}| \leq 1$. Hence, if B_{t_0} contains a vertex of \bar{Y} , then $z \notin B_{t_0}$ implies that this vertex belongs to one of the two 4-cycles. If B_{t_0} contains no vertex of \bar{Y} , then we let $uvwx$ be any of the two 4-cycles, and otherwise we let $uvwx$ be the 4-cycle with a vertex in B_{t_0} and number the vertices of the 4-cycle in such a way that $x \in B_{t_0}$. Then $u, v, w \notin B_{t_0}$.

Consider the components $T_{t_0}(u)$, $T_{t_0}(v)$ and $T_{t_0}(w)$ defined in Lemma 4.2. Since $u \sim v \sim w$, part (c) of Lemma 4.2 shows that these three components are the same, and hence the corresponding sets $\Gamma_{t_0}(u)$, $\Gamma_{t_0}(v)$ and $\Gamma_{t_0}(w)$ defined in Lemma 4.2 are the same. Since the only vertices that have no neighbor in $\{u, v, w\}$ are z and the four vertices of the second 4-cycle and since these five vertices do not belong to B_{t_0} , part (b) of Lemma 4.2 shows that $B_{t_0} \subseteq \Gamma_{t_0}(u)$. Let t be the unique vertex of the component $B_{t_0}(u)$ that is adjacent to t_0 in T . Since $B_0 \subseteq \Gamma_{t_0}(u)$ and since (T, B) is a tree decomposition of Γ , property (TD2) of the definition of a tree decomposition implies that $B_{t_0} \subseteq B_t$. This contradicts property (T2) of the tree decomposition (T, B) .

Case 2.2.2. $z \in B_{t_0}$.

We have seen that $|Y| \in \{8, 9\}$. If $|Y| = 8$, then (10) shows that $P = B_{t_0}$ and $|B_{t_0}| = \text{tw}(\Gamma) + 1 = |V(\Gamma)| - 8$. Since $P \cap Y = \emptyset$, we have $z \notin Y$ in this case. If $|Y| = 9$, then $z \in \bar{Y} = Y$ and hence $z \notin P$, so (10) shows that $B_{t_0} = P \cup \{z\}$ and $|B_{t_0}| = \text{tw}(\Gamma) + 1 = |V(\Gamma)| - 8$.

In any case we have $|B_{t_0}| = \text{tw}(\Gamma) + 1 = |V(\Gamma)| - 8$ and the eight vertices that are not in B_{t_0} are the eight vertices of the two 4-cycles and these eight vertices lie in Y . Put $E := B_{t_0} \setminus \{z\}$. Then $|E| = |V(\Gamma)| - 9 = \text{tw}(\Gamma)$ and $B_{t_0} = E \cup \{z\}$.

Let uvw be one of the two 4-cycles. As before, Lemma 4.2 shows that the sets $T_{t_0}(a)$ with a in this 4-cycle are all the same, and hence that the corresponding sets $\Gamma_{t_0}(a)$ are all the same. Lemma 4.2 shows that every vertex which has a neighbor in the 4-cycle uvw lies in $\Gamma_{t_0}(a)$. Since the only vertices with no neighbor in this 4-cycle are z and the four vertices of the second 4-cycle, it follows that $E \subseteq \Gamma_{t_0}(a)$. Let t be the vertex of the component of $T_{t_0}(u)$ that is adjacent to t_0 .

We next show that $E \subseteq B_t$. To see this, consider $e \in E$. Then $e \in \Gamma_{t_0}(w)$ and hence $e \in B_i$ for some $i \in T_i(w)$. Since also $e \in B_{t_0}$ and since t lies on the unique path of the tree T from t to i , it follows that $e \in B_t$. As this holds for any $e \in E$, we find $E \subseteq B_t$.

As $B_{t_0} = E \cup \{z\}$, then (T2) implies that $B_t \neq E$ and $z \notin B_t$. Since the tree representation under consideration has treewidth $\text{tw}(\Gamma) + 1$, then $|B_t| \leq \text{tw}(\Gamma) + 1 = |E| + 1$. It follows that there exists a unique vertex r with $B_t = E \cup \{r\}$. Also $r \neq z$ and $r \notin E$.

Since $r \notin B_{t_0}$, then r is a vertex of one of the two 4-cycles. If it is not a vertex of the 4-cycle uvw but of the other 4-cycle, then the other 4-cycle is also contained in $\Gamma_{t_0}(u)$. We may thus assume that $r = u$.

Then $v, w, x \notin B_t$. Consider $\Gamma_t(w)$ and let s be the unique vertex of the component $T_t(w)$ that is adjacent to t . Since vw is a path of length two, then Lemma 4.2 shows that $v, x \in \Gamma_t(w)$ and $\Gamma_t(w) = \Gamma_t(v) = \Gamma_t(x)$. By property (E), every vertex of E is adjacent to at least one vertex of the path vw . Therefore Lemma 4.2 shows that $E \subset \Gamma_t(w)$. As $E \subseteq B_t$, the argument used above to show that $E \subseteq B_t$ shows now that $E \subseteq B_s$. As $B_t = E \cup \{u\}$, then (T2) shows that $u \notin B_s$.

As $|E| = |V(\Gamma)| - 9$ and $|B_s| \leq \text{tw}(\Gamma) + 1 = |V(\Gamma)| - 8$, it follows that B_s contains at most one of the vertices v and x , we may assume w.l.o.g. that $v \notin B_s$. Recall that v lies in $\Gamma_t(w)$, which is the union of the sets B_i with vertices i of $T_t(w)$. As $v \notin B_s$, then the fact that (B, T) is a tree decomposition implies that every vertex i of T with $v \in B_i$ must be a vertex of $T_t(w)$. Since $\{u, v\}$ is an edge, there exists a vertex i of T with $u, v \in B_i$. Then i is a vertex of $T_t(w)$. We have $u \in T_t$ and $u \in T_i$. But the path of T from t to i contains the vertex s and $u \notin T_s$. Hence the graph induced by T on the vertices j with $u \in B_j$ is disconnected. This contradicts the fact that (B, T) is a tree decomposition of Γ . \square

References

- [1] Mengyu Cao, Ke Liu, Mei Lu, and Zequn Lv. Treewidth of the q -Kneser graphs. *Discrete Appl. Math.*, 342:174–180, 2024.
- [2] David Eppstein, Daniel Frishberg, and William Maxwell. On the treewidth of Hanoi graphs. *Theoret. Comput. Sci.*, 906:1–17, 2022.

- [3] P. Frankl and R. M. Wilson. The Erdős-Ko-Rado theorem for vector spaces. *J. Combin. Theory Ser. A*, 43(2):228–236, 1986.
- [4] Daniel J. Harvey and David R. Wood. Treewidth of the Kneser graph and the Erdős-Ko-Rado theorem. *Electron. J. Combin.*, 21(1):Paper 1.48, 11, 2014.
- [5] Daniel J. Harvey and David R. Wood. Treewidth of the line graph of a complete graph. *J. Graph Theory*, 79(1):48–54, 2015.
- [6] Daniel J. Harvey and David R. Wood. The treewidth of line graphs. *J. Combin. Theory Ser. B*, 132:157–179, 2018.
- [7] J. W. P. Hirschfeld and J. A. Thas. *General Galois geometries*. Springer Monographs in Mathematics. Springer, London, 2016.
- [8] Ferdinand Ihringer and Klaus Metsch. Large $\{0, 1, \dots, t\}$ -cliques in dual polar graphs. *J. Combin. Theory Ser. A*, 154:285–322, 2018.
- [9] Nina Kamcev, Anita Liebenau, David R. Wood, and Liana Yepremyan. The size Ramsey number of graphs with bounded treewidth. *SIAM J. Discrete Math.*, 35(1):281–293, 2021.
- [10] Ke Liu, Mengyu Cao, and Mei Lu. Treewidth of the generalized kneser graphs. *Electron. J. Combin.*, 29(1):Paper 1.57, 19, 2022.
- [11] Ke Liu and Mei Lu. The treewidth of 2-section of hypergraphs. *Discrete Math. Theor. Comput. Sci.*, 23(3):Paper No. 1, 20, 2021.
- [12] Klaus Metsch. On the treewidth of generalized Kneser graphs. *Australas. J. Combin.*, 86:477–486, 2023.
- [13] Neil Robertson and P. D. Seymour. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986.
- [14] Josse van Dobben de Bruyn and Dion Gijswijt. Treewidth is a lower bound on graph gonality. *Algebr. Comb.*, 3(4):941–953, 2020.