# Tighter Bounds on the Expected Absorbing Time of Ungarian Markov Chains

Eric Shen

May 2024

#### Abstract

In 2023, Defant and Li defined the Ungarian Markov chain associated to a lattice L. This Markov chain has state space L, and from any state  $x \in L$  transitions to the meet of  $\{x\} \cup T$ , where T is a randomly selected subset of the elements of L covered by x. They defined  $\mathcal{E}(L)$  to be the expected number of steps until the maximal element  $\hat{1}$  of L transitions into the minimal element  $\hat{0}$  in the Ungarian Markov chain, and investigated its asymptotics when L is the weak order on  $S_n$ , or the nth Tamari lattice Tam<sub>n</sub>. This paper resolves a conjecture of Defant and Li and makes partial progress towards another, by proving that  $\mathcal{E}(S_n)$  is linear in n, and proving an  $n^{1-o(1)}$  lower bound on  $\mathcal{E}(\text{Tam}_n)$ .

# 1 Introduction

Let n be a positive integer. Consider any permutation  $\sigma \in S_n$ . An index  $i \in [n-1]$  is a descent of  $\sigma$  if  $\sigma(i) > \sigma(i+1)$ . Let  $\mathrm{Des}(\sigma)$  denote the set of descents of  $\sigma$ . Now, given a permutation  $\sigma \in S_n$  and a subset  $T \subset \mathrm{Des}(\sigma)$ , the Ungar move on  $\sigma$  associated with T consists of reversing every descent in T, where consecutive descents in T are treated as blocks to be reversed. For example, given  $\sigma = 416523$ , if only index 1 is selected then the Ungar move sends  $416523 \to 146523$ , while if both indices 3 and 4 are selected, then the Ungar move sends  $416523 \to 412563$ . In the case that  $T \neq \emptyset$ , we call the Ungar move nontrivial. Note that for every  $\sigma \in S_n$ , the Ungar move associated to  $\mathrm{Des}(\Sigma)$  is called the maximal Ungar move. This corresponds to the pop-stack sorting map  $\mathrm{Pop}\colon S_n \to S_n$ , which maps a permutation through the so-called pop-stack data structure in a right-greedy manner.

Ungar moves were first studied by Ungar [11] to answer the question posed by Scott [8], of the minimum number of distinct slopes formed by the lines connecting each pair of points within a set of (not all collinear) n distinct points in a plane. To this end, Ungar proved the following bounds on the number of Ungar moves necessary to transform the permutation  $n(n-1)\cdots 1$  into the identity.

**Theorem 1.1.** ([11]) Consider any integer  $n \geq 4$ .

- (a) At most n-1 maximal Ungar moves suffice to send any permutation to the identity.
- (b) Consider a sequence of Ungar moves which send  $n(n-1)\cdots 1$  to  $12\cdots n$ . Given that the first move in this sequence is nontrivial and not maximal, the total number of moves in the sequence must be at least n if n is even, and at least n-1 if n is odd.

In [3], Defant and Li studied the Ungar moves in a probabilistic setting by introducing random Ungar moves, defined as follows. Fix  $p \in (0, 1]$ . Then, given any  $\sigma \in S_n$ , select  $T \subset \text{Des}(\sigma)$  by choosing any element of  $\text{Des}(\sigma)$  with independent probability p. The random Ungar move is now the Ungar move associated with T. In particular, note that any nontrivial Ungar move decreases the number of inversions, and hence repeatedly applying any sequence of nontrivial Ungar moves to any permutation will eventually reduce it to the identity. Motivated by this, Defant and Li raised the question of the expected number of random Ungar moves that

must be applied to the decreasing permutation  $n(n-1)\cdots 1$  until we reach the identity permutation  $12\cdots n$ . To this end, they defined the *Ungarian Markov chain* on  $S_n$  (with parameter p) as the Markov chain whose states are the elements of  $S_n$ , and transitions are given by the random Ungar moves [3]. They also defined  $\mathcal{E}_p(S_n)$  to be the expected number of steps for the permutation  $n(n-1)\cdots 1$  to transition to the identity  $12\cdots n$ . In particular, they derived the following bound on  $\mathcal{E}_p(S_n)$ :

**Theorem 1.2** ([3], Theorem 1.9). For fixed  $p \in (0,1)$ , we have that as  $n \to \infty$ ,

$$n - 1 + o_p(1) \le \mathcal{E}_p(S_n) \le \frac{8}{p} n \log n + O_p(n).$$

Defant and Li conjectured that for fixed  $p \in (0,1)$ ,  $\mathcal{E}_p(S_n)$  is linear in n. The first main result of this paper proves this conjecture.

**Theorem 1.3.** For fixed  $p \in (0,1)$ , we have that as  $n \to \infty$ ,

$$\mathcal{E}_p(S_n) \le \left(\frac{1+\sqrt{1-p}}{p}\right)n + o_p(n).$$

# 1.1 Ungarian Markov Chains on Lattices

Throughout this article, we will assume all partially ordered sets (posets) to be finite. A lattice L is a poset such that any two elements  $x, y \in L$  have a greatest lower bound, denoted with  $x \wedge y$  and called their "meet," and a lowest upper bound, denoted with  $x \vee y$  and called their "join." In [3] Defant and Li noted that by considering  $S_n$  under the right weak order, the definition of the Ungar move generalizes readily to any lattice. In particular, for any  $y \in L$ , define  $\operatorname{cov}_L(y) = \{x \in L : x \lessdot y\}$  to be the set of elements that y covers in L. Then picking a subset  $T \subset \operatorname{Des}(\sigma)$  and swapping all the selected descents is equivalent to picking a subset  $T \subset \operatorname{cov}_{S_n}(\sigma)$  and sending  $\sigma \mapsto \bigwedge(\{\sigma\} \cup T)$ .

In this manner Defant and Li generalized the definition of random Ungar moves and the Ungarian Markov chain as follows.

**Definition 1.4.** Fix some  $p \in (0,1]$  and a lattice L. For any element  $x \in L$ , a random Ungar move on x is an operation that picks a subset  $T \subset \text{cov}_L(x)$ , where each element is included with independent probability p, and then maps  $x \mapsto \bigwedge(\{x\} \cup T)$ . Moreover, the Ungarian Markov chain on L is defined as the Markov chain with state space L, satisfying that for all  $x, y \in L$ , the transition probability of  $x \to y$  is given by

$$\mathbb{P}(x \to y) = \sum_{\substack{T \subset \text{cov}_L(x) \\ \bigwedge(\{x\} \cup T) = y}} p^{|T|} (1-p)^{|\text{cov}_L(x)| - |T|}.$$

It is well-known that every finite lattice L has a minimal element  $\hat{0}$  and a maximal element  $\hat{1}$ . Evidently from any state in the Ungarian Markov chain on a lattice L, we may reach  $\hat{0}$  in a finite number of steps (i.e., the Markov chain is absorbing). Hence we may generally define  $\mathcal{E}_p(L)$  to be the expected number of steps until  $\hat{1}$  transitions to  $\hat{0}$  in the Ungarian Markov chain.

In [3], outside of investigating the Ungarian Markov chain on  $S_n$ , Defant and Li mainly considered one other class of lattices; the c-Cambrian lattices associated to a Coxeter element c of a finite Coxeter group W, that Reading first introduced in [5], [6]. For the Coxeter groups  $A_n$ ,  $B_n$ , and  $D_n$ , they proved using lattice-theoretic techniques and results from the well-studied last-passage percolation with geometric weights process that for fixed  $p \in (0,1)$ , if an element  $c_n$  was picked from every  $A_n$  then  $\mathcal{E}_p(\operatorname{Camb}_{c_n})$  would be bounded above by a linear function in n (with identical results holding for  $B_n$  and  $D_n$ ). They also investigated the special case of the nth Tamari lattice  $\operatorname{Tam}_n$  (to be formally defined in Section 4), first introduced by Dov Tamari in [10], which can be understood as a Cambrian lattice of type  $A_{n-1}$ . To introduce their results on  $\mathcal{E}_p(\operatorname{Tam}_n)$ , let  $\zeta_p(n)$  be the probability that the maximum of n independent geometric random variables,

each with expected value 1/p, is attained uniquely. In [2], Bruss and O'Cinneide proved that if we define

$$\Upsilon_p(x) = \begin{cases} px \sum_{k \in \mathbb{Z}} (1-p)^k e^{-(1-p)^k x} & p < 1\\ 0 & \text{else,} \end{cases}$$

then  $\lim_{n\to\infty} \zeta_p(n) - \Upsilon_p(n) = 0$ . Now, let  $\zeta_p^- = \liminf_{n\to\infty} \zeta_p(n)$  and  $\zeta_p^+ = \limsup_{n\to\infty} \zeta_p(n)$ . Note that  $\Upsilon_p((1-p)x) = \Upsilon_p(x)$  for all x>0, so thus

$$\begin{split} \zeta_p^+ &= \max_{0 < x < 1} \Upsilon_p(x) \\ \zeta_p^- &= \min_{0 < x < 1} \Upsilon_p(x) \ge p(1-p)e^{p-1}. \end{split}$$

Defant and Li established the following bound on  $\mathcal{E}_p(\mathrm{Tam}_n)$ :

**Theorem 1.5** ([3], Theorem 1.21). For fixed  $p \in (0,1)$ , we have that

$$\mathcal{E}_p(\operatorname{Tam}_n) \le \frac{2}{p} \left( \sqrt{\zeta_p^-(1+\zeta_p^-)} - \zeta_p^- \right) n + o_p(n).$$

Defant and Li also presented the question of establishing any nontrivial lower bound on  $\mathcal{E}_p(\operatorname{Tam}_n)$ . The second main result of this document establishes the following bound on  $\mathcal{E}_p(\operatorname{Tam}_n)$ , which notably for fixed p is  $n^{1-o(1)}$ .

**Theorem 1.6.** There exists absolute constants  $c_1, c_2 > 0$  such that for all  $n \ge e^{e^{c_2/p^2}}$ ,

$$\mathcal{E}_p(\operatorname{Tam}_n) \ge \frac{n}{e^{c_1(\log \log n)^3}}.$$

The paper is now organized as follows. Section 2 presents introductory material on statistics and well-known Markov processes (notably, the multicorner growth Totally Asymmetric Exclusion Process) that will be used to prove Theorems 1.3 and 1.6. Section 3 proves Theorem 1.3. Section 4 presents an interpretation of  $Tam_n$  that will be used to prove Theorem 1.6. Section 5 proves Theorem 1.6.

# 1.2 Acknowledgements

This work was done at the University of Minnesota Duluth with support from Jane Street Capital and the National Security Agency. This work was also supported by NSF Grant 2140043. The author would like to thank Joe Gallian and Colin Defant for organizing the Duluth REU and proofreading the paper. The author would also like to thank Evan Chen, Daniel Zhu, Mitchell Lee, and Rupert Li for their many helpful comments on the paper, as well as Noah Kravitz for many helpful discussions.

# 2 Preliminaries

Throughout this paper, the sequence  $c_1, c_2, \ldots$  denotes a sequence of positive, absolute, effectively computable constants. Moreover, any implied constants are assumed to be positive and absolute, unless otherwise indicated. Implied constants that are not absolute are indicated by subscripts denoting the dependence.

## 2.1 Background on Posets

For further reference, see [9].

Let P be a poset. A subposet of P is a poset  $P' \subset P$ , such that for any  $x, y \in P'$ , we have that  $x \leq y$  in P' if and only if  $x \leq y$  in P. For any  $x, y \in P$ , we say that x covers y (which we denote using  $y \lessdot x$ ) if

x > y and there does not exist a  $z \in P$  such that x > z > y. The Hasse diagram of a poset P is a drawing, where the vertices are the elements of P, and for any two  $x, y \in P$ , if y < x, then the corresponding vertex of x is connected to the corresponding vertex of y via an edge and x is drawn higher (vertically) than y. Now, a chain of P is a subset C such that all its elements are comparable; such a chain is maximal if there does not exist another chain C' for which  $C \subsetneq C'$ . The length of such a chain C is |C| - 1. Finally, a graded poset is a poset where all maximal chains are of the same length.

# 2.2 Probabilistic Background

In this section we will prove two useful probability lemmas that will be used in the proofs of Theorem 1.3 and 1.6. These two lemmas will all follow from standard applications of well-known results in statistics (e.g., the Markov bound, Chernoff bounds, random walk theory).

#### 2.2.1 Sum of Geometric Variables

Throughout the proof of Theorem 1.6, we will often use the following tail bounds on a sum of independent geometric random variables. Note that throughout this paper we will assume all geometric random variables to take on positive integer values.

**Lemma 2.1.** Fix a  $p \in (0,1]$ . Let  $G_1, G_2, \ldots, G_k$  be independent geometric random variables with parameter p. Then for all t > 0, we have that

$$\mathbb{P}\bigg(\sum_{i=1}^k G_i > \frac{k}{p} + t\sqrt{\frac{k}{p^3}}\bigg) < e^{-\frac{t^2}{2p+2t\sqrt{p/k}}} \quad \text{and} \quad \mathbb{P}\bigg(\sum_{i=1}^k G_i < \frac{k}{p} - t\sqrt{\frac{k}{p^3}}\bigg) < e^{-\frac{t^2}{2p-2t\sqrt{p/k}}}.$$

*Proof.* We only prove the first bound, as the proof for the other bound is analogous. Extend  $G_1, G_2, \ldots$  to an infinite sequence. Let

$$B_i = \begin{cases} 1 & i = \sum_{j=1}^m G_j \text{ for some } m \\ 0 & \text{else.} \end{cases}$$

Then evidently each  $B_i$  is given by an independent Bernoulli variable with parameter p. Let  $\mu = k + t\sqrt{k/p}$  and  $\delta = \frac{t/\sqrt{kp}}{1+t/\sqrt{kp}}$ ; note that  $(1-\delta)\mu = k$ . We thus have that by a Chernoff bound,

$$\mathbb{P}\left(\sum_{i=1}^{k} G_i > \frac{k}{p} + t\sqrt{\frac{k}{p^3}}\right) = \mathbb{P}\left(\sum_{i=1}^{\left\lfloor \frac{k}{p} + t\sqrt{\frac{k}{p^3}} \right\rfloor} B_i < k\right) \le e^{-\delta^2 \mu/2} = e^{-\frac{t^2}{2p + 2t\sqrt{p/k}}},$$

as desired.  $\Box$ 

#### 2.2.2 Random Walks on $\mathbb{Z}$

Consider an infinite sequence  $X_1, X_2, \ldots$  of i.i.d. random variables in  $\mathbb{Z}$ , where

$$X_i = \begin{cases} 1 & \text{probability } \frac{1}{2} \\ -1 & \text{probability } \frac{1}{2}. \end{cases}$$

The simple random walk on  $\mathbb{Z}$  is defined as the infinite sequence of random variables  $S_1, S_2, \ldots$  where  $S_n = X_1 + \cdots + X_n$ . The hitting time of any nonzero integer m is the random variable  $\tau_m$ , denoting the smallest  $t \in \mathbb{Z}^+$  for which  $S_t = m$ . Note that  $\tau_m$  is a sum of m i.i.d. copies of  $\tau_1$  for all positive integers m (and by symmetry,  $\tau_m = \tau_{-m}$ ); just consider the random variable denoting the number of steps between

when  $S_i$  is first k and  $S_i$  is first k+1 for all  $0 \le k \le m-1$ . Now, it is well known that the generating function  $\mathbb{E}[z^{\tau_1}]$  of  $\tau_1$  satisfies

$$\mathbb{E}[z^{\tau_1}] = \sum_{i=0}^{\infty} \mathbb{P}(\tau_1 = n) z^n = \frac{1 - \sqrt{1 - z^2}}{z}$$
$$= \sum_{n=0}^{\infty} (-1)^n \binom{1/2}{n+1} z^{2n+1}.$$

In particular, by elementary calculus we may find that  $(-1)^n \binom{1/2}{n+1} = \Theta(n^{-3/2})$ . Using the information above, we may derive the following bound.

**Lemma 2.2.** There exists an absolute constant  $c_3$  such that for all positive integers n and reals c > 0,

$$\mathbb{P}(\tau_m \ge n^2) \ge 1 - e^{-c_3 m/n}.$$

*Proof.* By the above, we have that there exists an absolute constant  $c_4$  such that

$$\mathbb{P}(\tau_1 = 2n - 1) \ge c_4 n^{-3/2}$$

Thus, there exists an absolute constant  $c_3$  such that

$$\mathbb{P}(\tau_1 \ge n^2) \ge \frac{c_3}{n}$$
.

Now, for every absolute constant  $c \in (0,1]$ , note that there exists an absolute constant c' such that for all  $x \in [0,c]$ ,  $e^{-c'x} \le 1 - x \le e^{-x}$ . Adjusting  $c_3$ , we find that for all n > 1,

$$\mathbb{P}(\tau_1 < n^2) \le e^{-c_3/n}.$$

Now, recall that  $\tau_m$  may be viewed as a sum of m i.i.d. copies of  $\tau_1$ . Thus we have that

$$\mathbb{P}(\tau_m < n^2) < \mathbb{P}(\tau_1 < n^2)^m < e^{-c_3 m/n}$$

hence

$$\mathbb{P}(\tau_m \ge n^2) \ge 1 - e^{-c_3 m/n},$$

as desired.  $\Box$ 

Now, fix some parameter  $q \in (0, 1/2]$ . Consider an infinite sequence  $Y_1, Y_2, \ldots$  of i.i.d. random variables in  $\mathbb{Z}$ , where

$$Y_i = \begin{cases} 1 & \text{probability } q \\ 0 & \text{probability } 1 - 2q \\ -1 & \text{probability } q. \end{cases}$$

The lazy simple random walk on  $\mathbb{Z}$  with parameter q is defined as the infinite sequence of random variables  $T_1, T_2, \ldots$  where  $T_n = Y_1 + \cdots + Y_n$ . We may again define the hitting time of any nonzero integer m as the random variable  $\tau'_m$  denoting the smallest  $t \in \mathbb{Z}^+$ , for which  $T_t = m$ . The following corollary will be useful in the proof of Theorem 1.6.

Corollary 2.3. There exists an absolute constant  $c_3$  such that for all positive integers m and integers n > 1,

$$\mathbb{P}(\tau'_m \ge n^2) \ge 1 - e^{-c_3 m/n}.$$

*Proof.* Let  $\tau_m^*$  be the number of indices i between 1 and  $\tau_m'$  inclusive for which  $T_i \neq 0$ . Then evidently  $\tau_m^* \leq \tau_m'$ , and moreover  $\tau_m^*$  shares the same distribution as  $\tau_m$  (essentially  $\tau_m^*$  "ignores" the  $T_i$  equal to 0 in calculating the hitting time.) Hence, we have that

$$\mathbb{P}(\tau'_m \ge n^2) \ge \mathbb{P}(\tau^*_m \ge n^2) \ge 1 - e^{-c_3 m/n},$$

as desired.  $\Box$ 

# 2.3 Ungarian Markov Chains on Posets of Order Ideals

For any lattice L and element  $x \in L$ , we let  $T_L(x)$  be the random variable denoting the number of steps elapsed in the Ungarian Markov chain of L until x has transitioned into  $\hat{0}$ . We also write  $T_L(\hat{1}) = T(L)$ . We also define  $\mathcal{E}_L(x) = \mathbb{E}(T_L(x))$ . Note that by definition,  $\mathcal{E}_L(\hat{1}) = \mathcal{E}_p(L)$ .

Now, consider any poset P. An order ideal of P is a subset  $I \subset P$  such that for any  $x, y \in P$  for which  $x \geq y$ , we have that  $x \in I$  implies  $y \in I$ . We establish a partial order on the order ideals by stating that  $I \leq J$  if  $I \subset J$ ; under this partial order, the order ideals of a poset P form a (distributive) lattice, which we will denote as J(P).

Now, consider any order ideal I of a poset P. Evidently  $\operatorname{cov}_{J(P)}(I)$  is precisely the set of order ideals J that are obtained by removing a maximal element of I. Hence, we may reformulate  $\mathcal{E}_p(J(P))$  as the Markov chain with state space J(P), that given any  $I \in J(P)$  deletes a randomly selected subset of the maximal elements of I. In particular, for any  $x \in P$ , let  $G_x$  be the random variable denoting the number of steps in which x remains the maximal element of P. Note that  $G_x$  is evidently given by a geometric random variable with parameter P. Now, define  $\operatorname{MC}(P)$  to be the set of maximal chains of P. Note that when we run the Ungarian Markov chain on J(P), any maximal chain  $C \in \operatorname{MC}(P)$  containing the last maximal element to be removed from the order ideal satisfies  $T(J(P)) = \sum_{x \in C} G_x$ . Hence,

$$T(J(P)) = \max_{C \in MC(P)} \sum_{x \in C} G_x.$$

This interpretation reformulates the Ungarian Markov chain on J(P) as the well-studied last-passage percolation with geometric weights random process. In particular, under this formulation we readily retrieve the following corollary.

**Corollary 2.4.** For any order ideal  $J' \in J(P)$  and any  $t \ge 0$ ,  $\mathbb{P}(T_{J(P)}(J') \ge t) \le \mathbb{P}(T(J(P)) \ge t)$ .

Proof. Interpret  $J' \subset J(P)$  as a subposet P' of P. Then evidently the subposet  $\{x \in J(P) : x \leq J'\}$  of J(P) is isomorphic to J(P') as posets. Now, note that every maximal chain of P' is contained within a maximal chain of P, so thus simulating J(P) using the geometric random variables  $G_x$  and simulating J(P') as a subposet of J(P), we obtain that

$$T_{J(P)}(J') = T(J(P'))$$

$$= \max_{C \in MC(P')} \sum_{x \in C} G_x$$

$$\leq \max_{C \in MC(P)} \sum_{x \in C} G_x$$

$$= T(J(P)).$$

From this, we readily obtain that for all t > 0,

$$\mathbb{P}(T_{J(P)}(J') \ge t) \le \mathbb{P}(T(J(P)) \ge t),$$

as desired.  $\Box$ 

Now, for any  $n, m \in \mathbb{Z}^+$ , consider the poset  $R_{n,m}$  defined as the product of a chain of length n and a chain of length m. In this case, the last-passage percolation with geometric weights random process corresponds to the multicorner growth Totally Asymmetric Simple Exclusion Process (TASEP) with parameter p. Loosely defined, this is a random sequence of Young diagrams  $\{\lambda_k\}$ , where  $\lambda_0 = \emptyset$  is the empty set and  $\lambda_{k+1}$  is constructed by adding in each external corner of  $\lambda_k$  with independent probability p. Now, consider the Hasse diagram of  $R_{n,m}$ ; then note that the subset of elements that are not contained in the order ideal, may be viewed as the subset of a Young diagram which evolves according to the multicorner growth TASEP (specifically, the subset of the diagram within the first n rows and m columns). This correspondence is illustrated more clearly in Figure 1; in particular, note that the subsets of the corresponding Young diagrams outside of the first n rows and m columns do not influence the number of steps required for the Young diagram to contain at least m elements in each of its first n rows (which is precisely  $T_{J(R_{m,n})}$ .) For more information on the multicorner growth TASEP, see [7, Chapters 4, 5]. Now, an important ingredient towards the proof of Theorem 1.3 will be the following convergence of the rescaled distributions of  $T_{J(R_{m,n})}$ , taken from [7, Theorem 5.31] but originally appearing as [4, Theorem 1.2].

**Theorem 2.5.** Consider any 0 . For any <math>x, y > 0, define

$$\Phi_p(x,y) = \frac{1}{p}(x+y+2\sqrt{(1-p)xy}),$$

$$\eta_p(x,y) = \frac{(1-p)^{1/6}}{p}(xy)^{-1/6}(\sqrt{x}+\sqrt{(1-p)y})^{2/3}(\sqrt{y}+\sqrt{(1-p)x})^{2/3}.$$

Now, let  $\{m_k\}_{k=1}^{\infty}$  and  $\{n_k\}_{k=1}^{\infty}$  be sequences of positive integers, satisfying that

$$\lim_{k \to \infty} m_k = \lim_{k \to \infty} n_k = \infty,$$

$$0 < \liminf_{k \to \infty} \frac{m_k}{n_k} < \limsup_{k \to \infty} \frac{m_k}{n_k} < \infty.$$

Then we have that

$$\lim_{k\to\infty}\mathbb{P}\bigg(\frac{T(J(R_{m_k,n_k}))-\Phi_p(m_k,n_k)}{\eta_p(m_k,n_k)}\leq t\bigg)=1-\frac{1}{32\pi t^{3/2}}e^{-4t^{3/2}/3}(1+O(t^{-3/2})).$$

Note that in the original statement, the right hand side is given by the Tracy-Widom Distribution  $F_2(x)$ . The above slightly modified statement may be obtained by applying [1, Equation 25].

# 3 Proof of Theorem 1.3

Next we will prove a more general version of Theorem 1.3. Note that the ideas in the following proof are modeled after Ungar's argument in [11].

**Theorem 3.1.** For any  $\sigma \in S_n$ ,

$$\mathcal{E}_{S_n}(\sigma) \le \left(\frac{1+\sqrt{1-p}}{p}\right)n + o_p(n).$$

Consider any integer  $1 \le k \le n-1$ . Let  $t_{k,\sigma} = t_k$  be the random variable denoting the number of steps




Figure 1: An example of an order ideal of  $R_{3,4}$  and a corresponding Young diagram. The indigo dots form the order ideal, while the red dots display the Young diagram. Note that the Young diagram may extend beyond the  $3 \times 4$  box of dots forming  $R_{3,4}$ , but the growth of the Young diagram in this region does not influence the evolution of the corresponding order ideal.

until all the elements  $\leq k$  in the current permutation are left of all the elements > k. Our key claim is as follows.

Claim 3.2. For all  $t \geq 0$ , we have that

$$\mathbb{P}(t_k \ge t) \le \mathbb{P}(T(J(R_{k,n-k})) \ge t).$$

*Proof.* For any permutation  $\sigma \in S_n$ , let  $\pi_k(\sigma)$  be the NE (North-East) path of length n, considered as the string made when we replace any element  $x \leq k$  in  $\sigma$  with "N" and any element y > k in  $\sigma$  with "E." Note that we may associate each such path with an element of  $J(R_{k,n-k})$  by taking the set of unit squares lying to the left of the path.

Now, we in fact prove the strengthened claim that for all  $\sigma \in S_n$ ,

$$\mathbb{P}(t_k \ge t) \le \mathbb{P}(T_{J(R_k, \pi_{k-1})}(\pi_k(\sigma)) \ge t).$$

Note that this is indeed stronger than the original claim by Corollary 2.4. We prove this by induction on the number of inversions of the permutation, and on t. The claim is trivial for any permutation already satisfying that the elements  $x \leq k$  are all left of the elements y > k. Now consider any permutation  $\sigma$  for which the claim has been proven for all permutations of smaller Coxeter length. Note that any consecutive pair of elements (x, y) with x > k and  $y \leq k$  forms a descent, whilst any consecutive pair of elements (x, y) with  $x \leq k$  and y > k does not form a descent. Call all such former pairs *critical*. Now, for any subset S' of the set of critical pairs S, let  $\sigma_{S'}$  be the permutation achieved when only the pairs in S' are swapped in  $\sigma$ . Also let  $x = \pi_k(\sigma)$ ,  $x_{S'} = \pi_k(\sigma_{S'})$ , and  $x_{\sigma'} = \pi_k(\sigma')$  for any permutation  $\sigma'$ . Then, by the above properties it is clear that

- 1. The elements that x can transition to are precisely the  $x_{S'}$  for the subsets  $S' \subset S$ .
- 2. The probability that  $\sigma$  transitions in the Ungarian Markov chain of  $S_n$  to a permutation  $\sigma'$ , for which the subset of critical pairs swapped is precisely S', is precisely the same as the probability that x transitions to  $x_{S'}$  in  $J(R_{k,n-k})$ .
- 3. For any  $\sigma'$  satisfying that the subset of critical pairs swapped is precisely S', we have that  $x_{\sigma'} \leq x_{\sigma_{S'}}$  in  $J(R_{k,n-k})$ . Hence, by Corollary 2.4 we have that for all  $t \geq 0$ ,  $\mathbb{P}(T_{J(R_{k,n-k})}(x_{\sigma'}) \geq t) \leq \mathbb{P}(T_{J(R_{k,n-k})}(x_{\sigma_{S'}}) \geq t)$ .

Also, clearly the probability that  $\sigma$  transitions to itself is at most the probability that x transitions to itself.

Now, note that the claim is trivial for t = 0. For any permutation  $\sigma'$ , let  $Q_{\sigma'}$  be the set of permutations that  $\sigma'$  can transition to. For t > 0, we now have:

$$\begin{split} \mathbb{P}(t_k \geq t) &= \mathbb{P}(\sigma \to \sigma) \mathbb{P}(t_k \geq t - 1) + \sum_{\varnothing \neq S' \subset S} \sum_{\sigma_1 \in Q_{\sigma_{S'}}} \mathbb{P}(\sigma \to \sigma_1) \mathbb{P}(t_{k,\sigma_1} \geq t - 1) \\ &\leq \mathbb{P}(x \to x) \mathbb{P}(T_{J(R_{k,n-k})}(\pi_k(\sigma)) \geq t - 1) + \sum_{\varnothing \neq S' \subset S} \sum_{\sigma_1 \in Q_{\sigma_{S'}}} \mathbb{P}(\sigma \to \sigma_1) \mathbb{P}(T_{J(R_{k,n-k})}(x_{\sigma_1}) \geq t - 1) \\ &\leq \mathbb{P}(x \to x) \mathbb{P}(T_{J(R_{k,n-k})}(\pi_k(\sigma)) \geq t - 1) + \sum_{\varnothing \neq S' \subset S} \sum_{\sigma_1 \in Q_{\sigma_{S'}}} \mathbb{P}(\sigma \to \sigma_1) \mathbb{P}(T_{J(R_{k,n-k})}(x_{\sigma_{S'}}) \geq t - 1) \\ &= \mathbb{P}(x \to x) \mathbb{P}(T_{J(R_{k,n-k})}(\pi_k(\sigma)) \geq t - 1) + \sum_{\varnothing \neq S' \subset S} \mathbb{P}(\sigma \to \sigma_{S'}) \mathbb{P}(T_{J(R_{k,n-k})}(x_{\sigma_{S'}}) \geq t - 1) \\ &= \mathbb{P}(x \to x) \mathbb{P}(T_{J(R_{k,n-k})}(\pi_k(\sigma)) \geq t - 1) + \sum_{\varnothing \neq S' \subset S} \mathbb{P}(x \to x_{S'}) \mathbb{P}(T_{J(R_{k,n-k})}(x_{\sigma_{S'}}) \geq t - 1) \\ &= \mathbb{P}(T_{J(R_{k,n-k})}(x) \geq t). \end{split}$$

This thus proves the inductive step and hence the claim.

We now return to the original problem. Consider any odd  $m \in \mathbb{Z}^+$ , and for all integers c let  $a_{n,m,c} = \lceil \frac{c(n-1)}{m} \rceil$ . Assume n > 2m; then evidently for each  $1 \le k \le n-1$  there exists a unique integer  $1 \le c \le m$  such that

$$a_{n,m,c-1} + 1 \le k \le a_{n,m,c}$$
.

For all k within the range above, we thus have

$$\mathbb{P}(t_k \ge t) \le \mathbb{P}(T(J(R_{k,n-k}))) \ge t)$$
  
$$\le \mathbb{P}(T(J(R_{a_{n,m,c},a_{n,m,m-c+1}}))) \ge t),$$

as evidently  $J(R_{k,n-k})$  is a subposet of  $J(R_{a_{n,m,c},a_{n,m,m-c+1}})$ .

Now, fix m and p, and recall the definitions of  $\Phi_p$  and  $\eta_p$  from Theorem 2.5. For m and p fixed, we have that  $\eta_p(a_{n,m,c},a_{n,m,m-c+1}) = \Theta_{m,p}(n^{1/3})$ . Also, by elementary calculus we may find that  $\Phi_p(a_{n,m,c},a_{n,m,m-c+1})$  is maximized when  $c = \frac{m+1}{2}$ . Hence, let  $A_k$  be the event that

$$t_k > \Phi_p(a_{n,m,(m+1)/2}, a_{n,m,(m+1)/2}) + \sqrt{n}.$$

Then, by Theorem 2.5, we find that for all sufficiently large n, there exists constants  $c_5$ ,  $c_6$ , possibly dependent on p, m such that:

$$\mathbb{P}(A_k) \le \frac{c_5}{n^{1/4}} e^{-c_6 n^{1/4}}.$$

Hence, by the union bound,

$$\mathbb{P}\left(\bigcup_{i=1}^{n-1} A_i\right) \le c_5 n^{3/4} e^{-c_6 n^{1/4}}.$$
 (1)

Now, let  $A = \bigcup_{i=1}^{n-1} A_i$ , and let B be the event that  $T_{S_n}(\sigma) > n^2/p$ . Evidently, if it is true that for all  $1 \le k \le n-1$ , the elements less than or equal to k are all left of the elements greater than k, then the current permutation must be the identity. Hence,  $\neg A$  implies that  $T_{S_n}(\sigma) \le \Phi_p(a_{n,m,(m+1)/2}, a_{n,m,(m+1)/2}) + \sqrt{n}$ , and thus for all sufficiently large  $n, B \subset A$ . To estimate the contribution of B to  $\mathcal{E}_{\sigma}(S_n)$ , we now use the

following lemma, first proved in [3].

Lemma 3.3 ([3]). We have that

$$\mathbb{P}(B)\mathbb{E}(T_{S_n}(\sigma)|B) = O(n^2 p e^{-n^2 p/2}).$$

Proof. Recall that  $S_n$  is a graded poset, where every maximal chain is of length  $\binom{n}{2} + 1 < \frac{n^2}{2}$ . Hence, any chain from  $\sigma$  to the identity permutation has less than  $\frac{n^2}{2}$  elements. Each step, the probability that we transition to a strictly lower element in the lattice is at least p, so we evidently have that  $T_{S_n}(\sigma) \leq \sum_{i=1}^{\binom{n}{2}} G_i$ , where each  $G_i$  is an independent geometric random variable with parameter p. Thus, applying Lemma 2.1, we find that

$$\mathbb{P}(B)\mathbb{E}(T_{S_n}(\sigma)|B) = \sum_{m=\lfloor n^2/p\rfloor+1}^{\infty} m\mathbb{P}(T_{S_n}(\sigma) = m)$$

$$\leq \sum_{m=\lfloor n^2/p\rfloor}^{\infty} (m+1)\mathbb{P}(T_{S_n}(\sigma) > m)$$

$$\leq \sum_{m=\lfloor n^2/p\rfloor}^{\infty} (m+1)e^{-(2p/m)(m-n^2/2p)^2}$$

$$\leq \sum_{m=\lfloor n^2/p\rfloor}^{\infty} (m+1)e^{-mp/2}$$

$$= O_p(n^2e^{-n^2/2}),$$

as desired.  $\Box$ 

Applying Lemma 3.3 and Equation 1, we now find that for all sufficiently large n,

$$\begin{split} \mathcal{E}_{S_{n}}(\sigma) &= \mathbb{P}(\neg A)\mathbb{E}(T_{S_{n}}(\sigma)|\neg A) + \mathbb{P}(A-B)\mathbb{E}(T_{S_{n}}(\sigma)|A-B) + \mathbb{P}(B)\mathbb{E}(T_{S_{n}}(\sigma)|B) \\ &\leq \Phi_{p}(a_{n,m,(m+1)/2}, a_{n,m,(m+1)/2}) + \sqrt{n} + c_{5}n^{3/4}e^{-c_{6}n^{1/4}}\left(\frac{n^{2}}{p}\right) + O_{p}(n^{2}e^{-n^{2}/2}) \\ &= \Phi_{p}(a_{n,m,(m+1)/2}, a_{n,m,(m+1)/2}) + o_{p,m}(n) \\ &= \left(\frac{1+1\sqrt{1-p}}{p}\right)\left(\frac{n(m+1)}{m}\right) + o_{p,m}(n). \end{split}$$

The above statement is true for all odd  $m \in \mathbb{Z}^+$ , so

$$\mathcal{E}_{S_n}(\sigma) \le \left(\frac{1+\sqrt{1-p}}{p}\right)n + o_p(n),$$

as desired.

# 4 Background on Tamari Lattices

A rooted tree is a (graph-theoretic) tree where we specify one vertex to be the "root." Within a rooted tree, we may naturally assign an orientation to each of the edges by having them point "away" from the root; if two vertices v, w are connected by an edge pointing towards w, then we say v to be the parent of w, and w to be the child of v. If a vertex has no children, then we call it a leaf. An ordered tree is a rooted tree where we also specify an ordering of the children. We may view ordered trees as trees that we may draw on

a plane, such that each parent is "higher" than their child, and the ordering of the children of each vertex from left to right is precisely the ordering we have defined. Hence ordered trees are also called plane trees.

A forest is a disjoint union of trees. An ordered forest is a forest equipped with an ordering on each tree, along with an ordering on the trees of the forest. The *preorder traversal* of an ordered forest on n vertices is a labeling of the vertices from 1 to n, iteratively assigned as follows.

- 1. We assign the label 1 to the root of the leftmost tree.
- 2. If the label i was just assigned to a non-leaf vertex, we next assign i+1 to its leftmost child.
- 3. If the label j was just assigned to a leaf vertex, and i is the largest index less than j which has at least one still-unlabeled child, then we label the leftmost unlabeled child of i with the label j + 1.
- 4. If after assigning the label i, all vertices in the tree containing vertex i are labelled, then we next assign i+1 to the root of the tree immediately right of the tree containing vertex i.

Note that throughout the rest of the paper, we will label the vertices of a forest using the preorder traversal labeling, and we will identify each vertex with its label.




Figure 2: An example of a forest in  $Tam_{12}$ . The labels in the diagram are given by the preorder traversal.

We now define a poset on the ordered forests on n vertices, that we later prove to be isomorphic to  $Tam_n$ . To do so we first define an *operation* on vertices of an ordered forest as follows.

**Definition 4.1.** Consider any ordered forest F, and a vertex v of F. If v has a parent, denote it with w. An *operation* on v maps F to a new ordered forest, defined as follows.

- 1. If v is a leaf, the operation maps F to itself.
- 2. Assume v is not a leaf. Let the rightmost child of v be v'.
  - (a) If w exists, delete the edge from  $v \to v'$ , and draw a new edge from  $w \to v'$ , such that v' is immediately right of v as a child of w.
  - (b) If w does not exist, delete the edge from  $v \to v'$ . v' and its descendants hence form a new tree within the forest immediately right of the tree containing v.

Note that the preorder traversal labeling is preserved under any operation, and hence any Ungar move. Notably, when the Ungarian Markov chain is run on  $Tam_n$  starting at  $\hat{1}$ , the labels of the vertices of the forest  $F_1$  produced after any number of steps is not dependent on the random Ungar moves beforehand, but are always given by the preorder traversal labeling of  $F_1$ .

We now define the poset structure on the ordered forests on n vertices as follows.

**Definition 4.2.** Given any two ordered forests F, F' on n vertices, say that  $F \lessdot F'$  if we may obtain F by starting with F' and operating on any one of its vertices.




Figure 3: The tree on the left is labelled via the preorder traversal. The middle tree is obtained by operating on vertex 2 of the left tree, and the right tree is obtained by operating on vertex 1 of the left tree. Note that the preorder traversal labeling is preserved in both operations.

Note that every operation on a non-leaf vertex decreases the number of descendants of that vertex, and leaves the number of descendants every other vertex has unchanged. Hence this covering relation indeed defines a partial order. We now prove that this poset is isomorphic to  $Tam_n$ , and characterize its random Ungar moves.

**Theorem 4.3.** The poset defined on the set of ordered forests on n vertices via the partial order defined in Definition 4.2 is isomorphic to  $Tam_n$ . Moreover, consider any forest F within this poset, and fix a probability  $p \in (0,1]$ . For every vertex of F, select it with independent probability p. Then a random Ungar move on F operates successively on each selected vertex of F, going in order of smallest to largest index under the preorder traversal.

Note that throughout the rest of the paper, we will write that during a given Ungar move, a vertex v receives an operation or is operated on if an Ungar move is applied to the given forest and v is one of the vertices selected in said move.

To prove Theorem 4.3, we introduce an alternative formulation of the nth Tamari lattice, first introduced in [5]. Call a permutation  $\sigma \in S_n$  312-avoiding if there do not exist indices  $1 \leq i_1 < i_2 < i_3 \leq n$  such that  $\sigma(i_1) > \sigma(i_3) > \sigma(i_2)$ . Consider  $S_n$  as a lattice under the right weak order, and let  $\operatorname{Av}_n(312)$  denote the sublattice of  $S_n$  consisting of the 312-avoiding permutations. Then  $\operatorname{Av}_n(312)$  with the given poset structure is isomorphic to the nth Tamari lattice as a poset. Moreover, to describe the random Ungar moves on  $\operatorname{Av}_n(312)$ , Defant and Li defined in [3] the following projection operator  $\pi_{\downarrow}: S_n \to \operatorname{Av}_n(312)$ . Given any permutation  $\sigma \in S_n$ , if there exist indices i, j such that i+1 < j and  $\sigma(i+1) < \sigma(j) < \sigma(i)$  we can perform an allowable swap by swapping the entries of  $\sigma(i)$  and  $\sigma(i+1)$ . Then  $\pi_{\downarrow}$  sends  $\sigma$  to the permutation obtained by starting at  $\sigma$  and repeatedly applying allowable swaps until no more can be performed. Now, as shown in [3, Section 8], applying a random Ungar move to  $\sigma$  within  $\operatorname{Av}_n(312)$  is equivalent to applying a random Ungar move to  $\sigma$  within  $S_n$ , then applying the  $\pi_{\downarrow}$  operator.

Proof of Theorem 4.3. We first construct a bijection between the 312-avoiding permutations and the ordered forests on n vertices. Given a 312-avoiding permutation  $\sigma$ , consider its plot; that is, the set of points  $p_i = (i, \sigma(i))$  in  $\mathbb{R}^2$ . Given any two integers  $i, j \in [n]$  with i < j, connect the points  $p_i$  and  $p_j$  by an edge going from  $p_j$  to  $p_i$  if

- $\sigma(i) > \sigma(j)$ , and
- no points in the plot (other than  $p_i, p_j$  themselves) are contained within the rectangle  $[i, j] \times [\sigma(i), \sigma(j)]$ .

Note that  $\sigma$  being 312-avoiding implies that the graph with vertices  $p_i$  and edges defined above is indeed a forest. Now, correspond to  $\sigma$  a plane forest with vertices denoted  $q_1, \ldots, q_n$ , satisfying the following properties:

- vertices  $q_i, q_j$  are connected by an edge if and only if  $p_i, p_j$  are connected by a (directed) edge;
- if for any integers  $i, j \in [n]$  satisfying i < j, the vertices  $q_i, q_j$  are connected by an edge, then  $q_i$  is vertically underneath  $q_j$  in the plane;
- if for any three integers  $i, j, k \in [n]$  satisfying i < j < k, the pairs  $q_i, q_k$  and  $q_j, q_k$  are connected by edges, then the vertex  $q_i$  lies to the left of the vertex  $q_j$ .

Evidently such a plane forest is unique (up to isomorphism on ordered forests), and one can inductively construct a 312-avoiding permutation in  $S_n$  that corresponds to any given ordered forest on n vertices. Hence, this correspondence bijects the 312-avoiding permutations in  $S_n$  and the plane forests on n vertices. Figure 4 demonstrates one example of such a 312-avoiding permutation and the corresponding ordered forest.




Figure 4: On the left is the plot of the permutation  $\sigma = 342651 \in S_6$ , with edges drawn between the vertices of the plot as described above. On the right is the corresponding ordered forest. Note that the vertex  $q_i$  does **not** generally have the label i under the preorder traversal.

Now, consider any integer  $i \in [n]$  such that  $q_i$  is not a leaf, and suppose  $j \in [n]$  satisfies that  $q_j$  is the rightmost child of  $q_i$ . Then by the bijection above, we have that j must be equal to i-1 for the corresponding permutation to be 312-avoiding, and applying an operation at  $q_j$  corresponds to swapping the descent i-1 and then applying the  $\pi_{\downarrow}$  operator. Hence the given bijection respects the covering relations on the two posets, and is hence a poset isomorphism. Moreover, this implies that for any  $\sigma \in \operatorname{Av}_n(312)$  corresponding to a forest F on n vertices, applying a random Ungar move to  $\sigma$  in  $\operatorname{Av}_n(312)$  corresponds to selecting each vertex of F with probability p, and then operating on the vertices in the order given by the preorder traversal of F. Hence the theorem is proved.

Throughout the rest of the paper, we will use the following properties of the aforementioned operations and preorder traversal labelings, all of which may be readily checked.

**Proposition 4.4.** The preorder traversal labeling satisfies the following properties.

- The labels of a vertex and its descendants form a contiguous subset of [n].
- Consider any two vertices v < w, and consider a sequence of Ungar moves applied to  $\hat{1} \in \operatorname{Tam}_n$ . For every vertex i, let  $h_i$  be the index of the first move in which i receives an operation. If  $h_v > h_w$  and  $h_w \ge h_i$  for all  $i \in [v+1, w-1]$ , then w will become a child of v after  $h_w$  moves.
- For any  $m \in [n]$ , let G be the induced subgraph of the current state (interpreted as a graph-theoretic forest) formed by the vertices in [m, n]. Then, given any other  $G' \in \operatorname{Tam}_{n-m+1}$ , the probability that

the current state transitions to a new state with G' as the subgraph formed by the vertices [m, n], is precisely the probability that G transitions to G' in the Ungarian Markov chain on  $\operatorname{Tam}_{n-m+1}$ . In particular, this probability is independent of the current state.

# 5 Proof of Theorem 1.6.

The proof of Theorem 1.6 is organized as follows. We will first present several definitions that will be used repeatedly throughout the proof. The bulk of the proof of Theorem 1.6 will be dedicated to proving Theorem 5.7 and Lemma 5.8. We will spend Section 5.1 proving Lemma 5.8, and in Section 5.2 we will return to prove Theorem 5.7 and Theorem 1.6.

Now, as discussed in Section 4, identify the vertices of any forest in  $\mathrm{Tam}_n$  with its labels under the reverse preorder traversal labeling. The proof of Theorem 1.6 will require us to simulate the Ungarian Markov chain using a specific algorithm that will be fully defined in Lemma 5.8. However, this algorithm will simulate the *first* operation on every vertex as follows. For all  $i \in [n]$ , define  $g_i$  to be a geometric random variable of parameter p that takes on positive integer values. Then do not operate on vertex i for the first  $g_i - 1$  steps, and operate on vertex i on the  $g_i^{\mathrm{th}}$  step. Now, given any subinterval  $I = [i, j] \subset [n]$  and positive integer m, let  $E_{I,m}$  be the event that  $g_i = m$  and  $g_l < m$  for all  $l \in [i+1,j]$ .

**Definition 5.1.** A two-rowed array  $A = \begin{bmatrix} a_1 & a_2 & \dots & a_l \\ b_1 & b_2 & \dots & b_l \end{bmatrix}$  is n-childlike if:

- for all  $i \in [l]$ ,  $a_i, b_i \in [n]$ ,
- $a_1 = 1$ ,
- $n \le a_2 > \ldots > a_l = 2$ , and
- $b_1 > b_2 \ge b_3 \ge b_4 \ge \cdots \ge b_l$ .

The following random array will keep track of (a large subset of) the children of vertex 1.

**Definition 5.2.** Simulate the first operation on every vertex using the algorithm described above. Then  $J = \begin{bmatrix} a_1 & a_2 & \cdots & a_l \\ b_1 & b_2 & \cdots & b_l \end{bmatrix}$  is the random array defined as follows.

- $a_1 = 1$ .
- $a_2$  is the largest j satisfying  $g_j = \max(g_i : i \in [2, n])$ .
- For i > 2,  $a_i$  is inductively defined as the largest j satisfying  $g_j = \max(g_l : l \in [2, a_{i-1} 1])$ . In particular, we terminate the sequence at  $a_l = 2$ .
- For all  $i \in [l]$ ,  $b_i = g_i$ .

Observe that J is completely determined by the values of the terms  $g_i$ . Moreover, note that if  $g_1 > g_2$ , then J is n-childlike.

**Definition 5.3.** Given an *n*-childlike array A, define  $E_A$  to be the event that J = A.

Suppose we simulate the first operation on every vertex using the algorithm described above. Then, after  $g_1 - 1$  steps, Proposition 4.4 implies that conditioned on  $E_A$ , the current state is a tree with vertex 1 as its root and with vertices  $a_2, \ldots, a_l$  all as children of vertex 1.

**Definition 5.4.** Given any decreasing subsequence  $I = \{a_1, \ldots, a_l\}$  of integers in [n], its *n*-summary is the subsequence  $I' = \{a_{i_1}, \ldots, a_{i_{l'}}\}$  chosen inductively as follows:

- we set  $a_{i_1} = a_1$ ;
- for each j > 1, pick  $a_{i_j}$  to be the smallest integer which corresponds to a vertex and satisfies  $a_{i_j} \ge \frac{a_{i_j-1}}{\log n}$ . If no such vertex exists, the subsequence ends.

**Definition 5.5.** Given any *n*-childlike array  $A = \begin{bmatrix} a_1 & \cdots & a_l \\ b_1 & \cdots & b_l \end{bmatrix}$ , let  $\{a_{i_1}, \dots, a_{i_{l'}}\}$  be the *n*-summary of  $\{a_2, \dots a_l\}$ . Then the *n*-summary of A is the subarray  $A' = \begin{bmatrix} a_1 & a_{i_1} & a_{i_2} & \cdots & a_{i_{l'}} \\ b_1 & b_{i_1} & b_{i_2} & \cdots & b_{i_{l'}} \end{bmatrix}$ .

**Definition 5.6.** A two-rowed array  $A = \begin{bmatrix} a_1 & \cdots & a_l \\ b_1 & \cdots & b_l \end{bmatrix}$  is n-good if it is n-childlike, and its n-summary  $A' = \begin{bmatrix} a_1 & a_{i_1} & a_{i_2} & \cdots & a_{i_{l'}} \\ b_1 & b_{i_1} & b_{i_2} & \cdots & b_{i_{l'}} \end{bmatrix}$  satisfies that  $a_{i_1} \geq \frac{n}{\log n}$  and  $a_{i_{l'}} \leq (\log n)^3$ .

Intuitively, J being n-good is a length condition that ensures the Ungarian Markov chain on  $Tam_n$  does not terminate too quickly.

Note that, by definition,  $a_{i_{k+2}} \leq \frac{a_{i_k}}{\log n}$  for all positive integers k. Since  $a_{i_1} \leq n$ , this implies that  $l' \leq \frac{2 \log n}{\log \log n}$ . Moreover by the bound  $a_{i_{k+1}} \geq \frac{a_{i_k}}{\log n}$ , we clearly have that  $l' \geq \frac{\log n}{\log \log n} - 4$ .

Now, let  $T_n$  be the random variable denoting the number of steps when running the Ungarian Markov chain on  $\operatorname{Tam}_n$  starting at  $\hat{1}$  until vertex 1 is disconnected from every other vertex. For an absolute constant  $c_7 > 1$  to be chosen later, define

$$f(x) = \begin{cases} \max\left(1, \frac{x}{e^{(c_7 \log \log x)^3}}\right) & \text{if } x \ge 16, \\ 1 & \text{else.} \end{cases}$$

The main ingredient in our proof of Theorem 1.6 is the following theorem.

**Theorem 5.7.** For all positive integers n, m,

$$\mathbb{P}(T_n \ge f(n) + m - 1 | E_{[1,n],m}) \ge \frac{1}{2}.$$

The proof of Theorem 5.7 mainly relies on the following lemma, which we dedicate the next subsection to proving.

**Lemma 5.8.** Consider any n-good array A whose bottom leftmost entry is m. Then

$$\mathbb{P}(T_n \ge f(n) + m - 1|E_A) \ge 0.9 - 402(\log\log n)^2 e^{-p\sqrt{n}}.$$

### 5.1 Proof of Lemma 5.8

For any sequence  $\alpha = \{\alpha_1, \dots, \alpha_n\}$  of positive integers, let  $E_{\alpha}$  be the event that  $g_i = \alpha_i$  for all i. Call such a sequence A-creating if  $E_{\alpha}$  implies  $E_A$ . We will instead prove that for all A-creating  $\alpha$ ,

$$\mathbb{P}(T_n \ge f(n) + \alpha_1 - 1 | E_\alpha) \ge 0.9 - 402(\log \log n)^2 e^{-p\sqrt{n}}.$$

This evidently implies the lemma above. Throughout the rest of the proof of Lemma 5.8, we will treat the  $q_i$  as fixed (rather than random) variables, as the result of conditioning on  $E_{\alpha}$ .

We proceed by induction on n. Clearly, there exists an absolute constant  $c_8 > 1$  such that for all  $n \le e^{e^{c_8/p^2}}$ , f(n) = 1 and hence the theorem is trivial (in the proof that follows we will often adjust the effective range of the theorem by adjusting  $c_8$  to a new absolute constant; this will simultaneously be an adjustment of  $c_7$ ). Henceforth assume  $n > e^{e^{c_8/p^2}}$ .

We now describe an algorithm for simulating the Ungarian Markov chain on  $\operatorname{Tam}_n$ . Let  $A = \begin{bmatrix} a_1 & a_2 & \cdots & a_l \\ b_1 & b_2 & \cdots & b_l \end{bmatrix}$ ,

and let  $A' = \begin{bmatrix} a_1 & a_{i_1} & a_{i_2} & \cdots & a_{i_{l'}} \\ b_1 & b_{i_1} & b_{i_2} & \cdots & b_{i_{l'}} \end{bmatrix}$  be the *n*-summary of *A*. Recall that  $l' \ge \frac{\log n}{\log \log n} - 4$ ; adjust  $c_8$  such that  $\frac{\log n}{\log \log n} - 4 > 402(\log \log n)^2$ . Now, we will simulate the Ungarian Markov chain on  $\operatorname{Tam}_n$  as follows.

- As discussed before, for every  $i \in [n]$ , let  $g_i$  be a geometric random variable of parameter p, taking on positive integer values. Then, for every  $i \in [n]$ , let vertex i receive its first operation on the  $g_i$ th step.
- Now, for all  $t \in \mathbb{Z}^+$ ,  $i \in [2, n]$ , let  $B_{i,t}$  and  $B'_{i,t}$  be independent Bernoulli random variables of parameter p. Also, for all  $t \in \mathbb{Z}^+$ , and even  $j \in (402 \log \log n, 402(\log \log n)^2]$ , let  $C_{j,t}$  be an independent Bernoulli random variable of parameter p.
  - If  $2 \le i < a_{i_{402(\log \log n)^2}}$ , then if  $t \le g_i$ , operate on vertex i on the tth step if and only if  $t = g_i$ . Else, operate on vertex i if and only if  $B_{i,t} = 1$ .
  - If  $i \ge a_{i_{402 \log \log n}}$ , say  $i \in [a_j, a_{j-1} 1]$  for some  $j \in [2, i_{402 \log \log n}]$  (here when j = 2, replace  $a_1$  with n + 1). Then on the tth step:
    - \* if  $t \leq g_i$ , then operate on i if and only if  $t = g_i$ ;
    - \* else, if  $i \neq a_j$ , then operate on i if and only if  $B_{i,t} = 1$ ;
    - \* else, if vertex 1 is connected to vertex  $a_{j-1}$  after t-1 steps, then operate on i if and only if  $B_{i,t} = 1$ ;
    - \* else, operate on vertex i if and only if  $B'_{i,t} = 1$ .
  - Otherwise, suppose that  $i \in [a_{i_{j+2}}, a_{i_j})$  for some even  $j \in (402 \log \log n, 402(\log \log n)^2]$ . Let  $F'_j$  denote the induced subgraph of the current state (interpreted as a graph-theoretic forest) formed by the vertices in  $[a_{i_{j+2}}, a_{i_j})$ . On the tth step,
    - \* if  $t \leq g_i$ , then operate on i if and only if  $t = g_i$ ;
    - \* else, if after t-1 steps i is the largest root of  $F'_j$  that has at least one child, then operate on vertex i if and only if  $C_{j,t} = 1$ ;
    - \* else, operate on vertex i if and only if  $B_{i,t} = 1$ .
- For every  $t \in \mathbb{Z}^+$ , every  $i \leq i_{402 \log \log n}$ , and every  $402 \log \log n \leq j \leq 402 (\log \log n)^2$ , let  $D_t$ ,  $D_{i,t}$ , and  $D'_{j,t}$  be independent Bernoulli random variables of parameter p. On the tth step, say i is the largest index satisfying that vertex 1 is still connected to vertex  $a_i$ . Then
  - if  $t \leq g_1$ , then operate on vertex 1 if and only if  $t = g_1$ ;
  - else, if  $i \leq i_{402 \log \log n}$ , then operate on vertex 1 if and only if  $D_{i,t} = 1$ ;
  - else, if  $i \in [a_{i_{j+2}}, a_{i_j})$  for some even  $j \in (402 \log \log n, 402(\log \log n)^2]$ , then operate on vertex 1 if and only if  $D'_{j,t} = 1$ ;
  - else, operate on vertex 1 if and only if  $D_t = 1$ .

One may readily verify that the algorithm above indeed simulates the Ungarian Markov chain.

Now, condition on  $E_{\alpha}$ . For all  $j \in [l]$ , let  $t_j$  be the random variable denoting the number of steps after the  $g_1-1^{\text{th}}$  step until vertex 1 is no longer connected to vertex  $a_j$ . Observe that  $t_1 \geq 1$ , and since vertex 1 may only be operated on at most once every step, the number of children it has can only decrease by at most 1 every step. Hence  $t_{i+1}-t_i \geq 1$  for every  $i \in [2,l]$ . In particular,  $t_{i\log\log n} \geq i_{\log\log n} \geq \log\log n$ . We now establish the following bound on  $t_{i_{402\log\log n}}$ .

**Lemma 5.9.** There exists an absolute constant  $c_9$  and an event  $A_1$  satisfying that

• A<sub>1</sub> only depends on the random variables

- 
$$D_{j,t}$$
, for  $j \leq i_{402 \log \log n}$  and  $t \in \mathbb{Z}^+$ , and  
-  $B_{j,t}$  and  $B'_{j,t}$ , for  $j \geq a_{i_{402 \log \log n}}$  and  $t \in \mathbb{Z}^+$ ;

• A<sub>1</sub> is contained in the event that

$$t_{i_{402 \log \log n}} \ge \frac{f(n/2e^{(\log \log n)^2})}{1700c_9};$$

•  $\mathbb{P}(A_1) \geq 0.9$ .

Before proving Lemma 5.9, we will first prove the following claim.

**Lemma 5.10.** There exists an absolute constant  $c_{10}$  such that for every  $i_{\log \log n} < i \le i_{402 \log \log n}$ , there exists an event  $E_i$  satisfying that:

- 1.  $E_i$  only depends on the random variables
  - $D_{j,t}$ , for  $j \leq i$  and  $t \in \mathbb{Z}^+$ , and
  - $B_{j,t}$  and  $B'_{j,t}$ , for  $j \geq a_i$  and  $t \in \mathbb{Z}^+$ ;
- 2.  $E_i$  is contained in the event that

$$t_i - t_{i-1} \ge \min\left(c_{10}(pt_{i-1} - \sqrt{t_{i-1}})^2, \frac{f(a_{i-1} - a_i)}{17}\right);$$

- 3. we have that  $\mathbb{P}(E_i)$ , conditioned on the values of
  - $D_{j,t}$ , for  $j \leq i-1$  and  $t \in \mathbb{Z}^+$ , and
  - $B_{i,t}$  and  $B'_{i,t}$ , for  $j \geq a_{i-1}$  and  $t \in \mathbb{Z}^+$ ,

is always at least 0.15.

*Proof.* Observe that  $t_{i-1}$  depends on the random variables

- $D_{i,t}$ , for  $j \leq i-1$  and  $t \in \mathbb{Z}^+$ , and
- $B_{i,t}$  and  $B'_{i,t}$ , for  $j \geq a_{i-1}$  and  $t \in \mathbb{Z}^+$ .

Now, condition on the values of the random variables above, and define the following events.

• Let  $E_{i,1}$  be the event that

$$\sum_{u=g_{a_i}+1}^{g_1+t_{i-1}} B_{a_i,u} \ge p(t_{i-1}+g_1-g_{a_i}) - \sqrt{t_{i-1}+g_1-g_{a_i}}.$$

• Define  $\tau_0$  to be the smallest t satisfying that

$$\sum_{u=g_1+t_{i-1}+1}^{g_1+t_{i-1}+t} D_{i,u} - B'_{a_i,u} = \left\lceil p(t_{i-1} + g_1 - g_{a_i}) - \sqrt{t_{i-1} + g_1 - g_{a_i}} \right\rceil.$$
 (2)

If no such t exists, define  $\tau_0 = r(p(t_{i-1} + g_1 - g_{a_i}) - \sqrt{t_{i-1} + g_1 - g_{a_i}})^2$ . Observe that since for all u,  $D_{i,u} - B'_{a_i,u} \le 1$ , such a t must exist if it is ever true that the LHS of (2) exceeds or is equal to the RHS. Now, for every positive real r, let  $E_{i,2,r}$  be the event that

$$\tau_0 \ge r(p(t_{i-1} + g_1 - g_{a_i}) - \sqrt{t_{i-1} + g_1 - g_{a_i}})^2.$$

- Let  $t'_i$  be the number of steps after the  $g_{a_i} 1^{\text{st}}$  step until no vertex in  $[a_i + 1, a_{i-1} 1]$  is a child of  $a_i$ . Let  $E_{i,3}$  be the event that  $t'_i \geq f(a_{i-1} a_i)$ .
- Given any state of  $Tam_n$ , let  $F_i$  denote the induced subgraph of the current state (interpreted as a graph-theoretic forest) formed by the vertices in  $[a_i, a_{i-1} 1]$ . Given any  $t > g_1$ , let  $j_t$  be the random variable denoting the label of the largest j satisfying that vertex j is a root of  $F_i$  with at least one child (if no such vertex exists set  $j_t = a_i$ ). Moreover, let  $u_0 = 0$ , and let  $u_l$  be the random variable denoting the l<sup>th</sup> smallest u satisfying  $D_{a_i,u} = 1$ . Finally, let  $Y_l = u_l u_{l-1}$ .
  - Given any real  $0 < \gamma_2 < 1$ , let  $E_{i,4,\gamma_2}$  be the event that

$$\sum_{t=g_{a_i}+1}^{g_1+t_{i-1}} B_{j_t,t} > \gamma_2 p(t_{i-1}+g_1-g_{a_i}).$$

- Let  $R = \min(\gamma_2 p(t_{i-1} + g_1 - g_{a_i}), a_{i-1} - a_i)$ . Then given any real  $0 < \gamma_3 < 1$ , let  $E_{i,5,(\gamma_2,\gamma_3)}$  be the event that

$$\sum_{i=1}^{R} Y_i \ge \gamma_3 R/p.$$

Note that all of the events above depend only on the random variables  $D'_{i,t}$ ,  $B_{j,t}$ , and  $B'_{j,t}$ , where  $j \in [a_i, a_{i-1} - 1]$  and  $t \in \mathbb{Z}^+$ , along with the variables we have already conditioned the values of. Now, consider the intersection  $E_{i,r,\gamma_2,\gamma_3}$  of the events  $E_{i,1}$ ,  $E_{i,2,r}$ ,  $E_{i,3}$ ,  $E_{i,4,\gamma_2}$ , and  $E_{i,5,(\gamma_2,\gamma_3)}$ . Suppose  $E_{i,r,\gamma_2,\gamma_3}$  is true. First, note that if  $t_i + g_1 \leq t'_i + g_{\alpha_i}$ , then we claim that after t steps the number of vertices in  $[a_i, a_{i-1} - 1]$  which are children of vertex 1 satisfies

$$\#\{j \in [a_i, a_{i-1} - 1] : j \text{ child of } 1\} \ge \begin{cases} \sum_{u = g_{a_i} + 1}^t B_{a_i, u} & t \le g_1 + t_{i-1}, \\ \sum_{g_1 + t_{i-1}}^t B_{a_i, u} + \sum_{u = g_1 + t_{i-1} + 1}^t D_{i, u} - B'_{a_i, u} & t > g_1 + t_{i-1}. \end{cases}$$

Indeed, this is because the number of such children increases by 1 whenever vertex  $a_i$  is operated on and has at least one child (the latter of which is guaranteed if  $t_i + g_1 \le t'_i + g_{\alpha_i}$ ), and the number of such children decreases by 1 whenever vertex 1 is operated on whilst its rightmost child is a vertex in  $[a_i, a_{i-1} - 1]$  (in other words,  $t > g_1 + t_{i-1}$ ). Hence if  $E_{i,1}$  and  $E_{i,2,r}$  are both true, then we have that either  $t_i + g_1 > t'_i + g_{a_i}$  or

$$t_i - t_{i-1} \ge \tau_0 \ge r(p(t_{i-1} + g_1 - g_{a_i}) - \sqrt{t_{i-1} + g_1 - g_{a_i}})^2.$$

Now, if  $E_{i,3}$  is true, then

$$t_i + g_1 > t'_i + g_{a_i}$$
  

$$t_i - t_{i-1} > t'_i + g_{a_i} - g_1 - t_{i-1}$$
  

$$\geq f(a_{i-1} - a_i) - (g_1 - g_{a_i}) - t_{i-1}.$$

Moreover, suppose that after  $t_{i-1}+g_1$  steps, there exists a vertex in  $[a_i,a_{i-1}-1]$  that is not a child of vertex 1. If  $E_{i,4,\gamma_2}$  is true, then for all  $t \leq t_{i-1}+g_1$  such that  $B_{j_t,t}=1$ , we have that the number of vertices in  $[a_i,a_{i-1}-1]$  increases after the tth step. Hence, the number of vertices in  $[a_i,a_{i-1}-1]$  that are children of vertex 1 after  $t_{i-1}+g_1$  steps is at least R, and thus if  $E_{i,5,(\gamma_2,\gamma_3)}$  is true, then  $t_i-t_{i-1} \geq \gamma_3 R/p$ . Hence, we

have that

$$t_{i} - t_{i-1} \ge \max\left(\min\left(\gamma_{3}\gamma_{2}(t_{i-1} + g_{1} - g_{a_{i}}), \frac{\gamma_{3}(a_{i-1} - a_{i})}{p}\right), \\ \min(r(p(t_{i-1} + g_{1} - g_{a_{i}}) - \sqrt{t_{i-1} + g_{1} - g_{a_{i}}})^{2}, f(a_{i} - a_{i-1}) - (t_{i-1} + g_{1} - g_{a_{i}}))\right).$$

Thus, if  $E_{i,r,1/4,1/4}$  is true, then noting that  $f(a_i - a_{i-1}) \le a_i - a_{i-1}$ , we have that

$$t_{i} - t_{i-1} \ge \min\left(r(p(t_{i-1} + g_{1} - g_{a_{i}}) - \sqrt{t_{i-1} + g_{1} - g_{a_{i}}})^{2}, \frac{f(a_{i} - a_{i-1})}{17}\right)$$

$$\ge \min\left(r(pt_{i-1} - \sqrt{t_{i-1}})^{2}, \frac{f(a_{i} - a_{i-1})}{17}\right).$$

Hence it now suffices to prove that for some absolute constant  $c_{11}$  we have that  $\mathbb{P}(E_{i,c_{11},1/4,1/4}) \geq 0.15$ . To do this, we bound the probabilities of the aforementioned events as follows.

- 1. By a Chernoff bound, we have that  $\mathbb{P}(E_{i,1}) \geq 1 e^{-2}$ .
- 2. Let  $F_{a_i,u} = D_{a_i,u-g_1-t_{i-1}} B'_{a_i,u-g_1-t_{i-1}}$ . Then note that each  $F_{a_i,u}$  is independent with probability distribution

$$F_{a_i,u} = \begin{cases} 1 & \text{probability } p(1-p) \\ 0 & \text{probability } p^2 + (1-p)^2 \\ -1 & \text{probability } p(1-p). \end{cases}$$

Thus the sequence  $\{F_{a_i,1}, F_{a_i,1} + F_{a_i,2}, \ldots\}$  defines a lazy simple random walk on  $\mathbb{Z}$  with parameter p(1-p), and hence we may apply Corollary 2.3 to obtain that

$$\mathbb{P}(E_{i,2,c_{11}}) \ge 1 - e^{-c_3/\sqrt{c_{11}}}.$$

- 3. By induction, we have that  $\mathbb{P}(E_{i,3}) \geq 0.9 402(\log \log n)^2 e^{-p\sqrt{n}}$ .
- 4. Recall that for any  $i, t_{i-1} \ge \log \log n \ge c_8/p^2$ . Hence, by a Chernoff bound (applied to the  $B_{j_t,t}$ , given the fixed value of  $t_{i-1}$ ), we have that  $\mathbb{P}(E_i, 4, \gamma_2) \ge 1 e^{-(1-\gamma_2)^2 c_8/2}$ .
- 5. Observe that each  $Y_i$  is an independent geometric variable of parameter p taking values in  $\mathbb{Z}^+$ , and note that  $t_{i-1}$  and R are both independent from each  $Y_i$ . Also note that since  $t_{i-1} \geq c_8/p^2$ ,  $R \geq \min(\gamma_2 c_8/p, 1)$ . Thus, conditioning  $t_{i-1}$  to be any possible fixed value (note it definitionally must be  $\geq c_8/p^2$ ), we find that by applying Lemma 2.1 (and setting k = R,  $t = \sqrt{Rp}(1 \gamma_3)$ , we always have

$$\begin{split} \mathbb{P}(E_{i,5,\gamma_2,\gamma_3}|t_{i-1}) &\geq 1 - e^{-Rp(1-\gamma_3)^2/(2p-2p(1-\gamma_3))} \\ &\geq 1 - e^{-R(1-\gamma_3)^2/2\gamma_3} \\ &\geq 1 - e^{-\min(\gamma_2 c_8/p,1)(1-\gamma_3)^2/2\gamma_3}. \end{split}$$

Hence, we have that  $\mathbb{P}(E_{i,5,\gamma_2,\gamma_3}) \geq 1 - e^{-\min(\gamma_2 c_8/p,1)(1-\gamma_3)^2/2\gamma_3}$  in general.

Finally, by a union bound we now find that

$$\mathbb{P}(E_{i,c_{11},1/4,1/4}) \ge 1 - e^{-2} - e^{-c_3/\sqrt{c_{11}}} - (0.1 + 402(\log\log n)^2 e^{-p\sqrt{n}}) - e^{-9c_8/32} - e^{-\min(9c_8/32,9/8)}.$$

Adjusting  $c_8$  to be sufficiently large and  $c_{11}$  to be sufficiently small now gives us that  $\mathbb{P}(E_{i,c_{11},1/4,1/4}) \geq 0.15$ , as desired.

We also now prove a variant of Lemma 5.10 for specific sequences of consecutive i.

**Lemma 5.11.** For every  $\log \log n < j \le 402 \log \log n$ , there exists an event  $F_j$  satisfying that

- $F_j$  only depends on the random variables
  - $-D_{u,t}$ , for  $u \leq i_{j+2}$  and  $t \in \mathbb{Z}^+$ , and
  - $B_{u,t}$  and  $B'_{u,t}$ , for  $u \geq a_{i_{i+2}}$  and  $t \in \mathbb{Z}^+$ ;
- $F_j$  is contained in the event that

$$t_{i_{j+2}} - t_{i_j} \ge 0.01 \min\left(\frac{f(n/k^j - n/k^{j+1})}{17}, c_{10}(pt_{i_j} - \sqrt{t_{i_j}})^2\right);$$

- we have that  $\mathbb{P}(F_i)$ , conditioned on the values of
  - $-D_{u,t}$ , for  $u \leq i_i$  and  $t \in \mathbb{Z}^+$ , and
  - $B_{u,t}$  and  $B'_{u,t}$ , for  $u \geq a_{i_j}$  and  $t \in \mathbb{Z}^+$ ;

is always at least 0.1.

*Proof.* For each  $a \in \mathbb{R}_{>0}$ , define

$$g_j(a) = \min\left(c_{10}(pt_{i_j} - \sqrt{t_{i_j}})^2, \frac{f(a)}{17}\right).$$

Note that for every  $i \in [i_j + 1, i_{j+2}]$ , the event  $E_i$  is contained in the event that

$$t_i - t_{i-1} \ge g_i(a_{i-1} - a_i).$$

Also note that for all nonnegative reals a, b, we have that  $g_j(a) + g_j(b) \ge g_j(a+b)$ . Now, for each  $i \in [1, i_{j+2} - i_j]$ , let  $X_i$  be the indicator function for  $E_{i+i_j}$ . Let  $F_j$  be the event that

$$\sum_{i=1}^{i_{j+2}-i_j} g_j(a_{i+i_j-1}-a_{i+i_j})X_i \ge 0.01g_j(a_{i_j}-a_{i_{j+2}}).$$

We claim that  $F_j$  satisfies the desired properties.  $F_j$  clearly only depends on the random variables described. Now, condition on the values of the variables

- $D_{u,t}$ , for  $u \leq i_j$  and  $t \in \mathbb{Z}^+$ , and
- $B_{u,t}$  and  $B'_{u,t}$ , for  $u \ge a_{i_j}$  and  $t \in \mathbb{Z}^+$ .

By Lemma 5.10 we know that conditioning on the variables above and on the variables

- $D_{u,t}$ , for  $i_i < u \le i_i + i$  and  $t \in \mathbb{Z}^+$ , and
- $B_{u,t}$  and  $B'_{u,t}$ , for  $a_{i,j} > u \ge a_{i,j+i}$  and  $t \in \mathbb{Z}^+$ ,

always gives us that  $\mathbb{P}(X_i) \geq 0.15$ . In particular,  $\mathbb{P}(X_i|X_1,\ldots X_{i-1}) \geq 0.15$  for all  $i \in [1,i_{j+2}-i_j]$ . Hence,

for every  $0 \le r \le q = 0.15$ , we have by the Markov inequality that

$$\mathbb{P}\left(\sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})(1 - X_{i}) \geq (1 - r) \sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})\right) \\
\leq \frac{\mathbb{E}\left[\sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})(1 - X_{i})\right]}{(1 - r) \sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})} \\
\mathbb{P}\left(\sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})X_{i} \leq r \sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})\right) \\
\leq \frac{(1 - q)\left(\sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})\right)}{(1 - r) \sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})} = \frac{1 - q}{1 - r} \\
\mathbb{P}\left(\sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})X_{i} > r \sum_{i=1}^{i_{j+2}-i_{j}} g_{j}(a_{i+i_{j}-1} - a_{i+i_{j}})\right) \geq \frac{q - r}{1 - r}.$$

Now, note that

$$\sum_{i=1}^{i_{j+2}-i_j} g_j(a_{i+i_j-1}-a_{i+i_j}) \ge g_j(a_{i_{j+2}}-a_{i_j}).$$

Thus, setting r=0.01 now evidently gives us that  $\mathbb{P}(F_j) \geq 0.14/0.99 > 0.1$ , as desired. Now, note that we always have  $t_{i+i_j} - t_{i+i_j-1} \geq g(a_{i+i_j-1} - a_{i+i_j})X_i$  (as this is trivial if  $X_i = 0$ , and  $X_i = 1$  implies  $E_i$ ). Also, since  $a_{i_j} \geq n/k^j$ ,  $a_{i_{j+2}} \leq a_{i_j}/k$ , and g is nondecreasing, we have that  $g(a_{i_j} - a_{i_{j+2}}) \geq g(n/k^j - n/k^{j+1})$ . Thus,  $F_j$  is contained in the event that

$$\sum_{i=i_j+1}^{i_{j+2}} g(a_{i-1} - a_i) \ge 0.01g(a_{i_j} - a_{i_{j+2}}) \ge g(n/k^j - n/k^{j+1}),$$

as desired.  $\Box$ 

We now prove Lemma 5.9.

Proof of Lemma 5.9. First, adjust  $c_8$  to be at least 4. Then there exists an absolute constant  $c_9$  such that for all  $t \ge c_8/p^2$ ,

$$0.01c_{10}(pt - \sqrt{t})^2 \ge c_9 p^2 t^2$$

Also, fixing  $c_9$ , adjust  $c_8$  again such that  $c_8 \ge e/c_9$ . Now, let N be the number of even  $j \in (\log \log n, \le 402 \log \log n]$  for which  $F_j$  is true. Let  $A_1$  be the event that  $N \ge 2 \log \log n$ . We claim that  $A_1$  satisfies the desired properties. Evidently,  $A_1$  only depends on the variables listed. We now bound  $\mathbb{P}(A_1)$ . For each even  $j \in (\log \log n, 402 \log \log n]$ , let  $X_{j+1-2\lceil \log \log n/2 \rceil}$  be the indicator function of  $F_j$ . Also, suppose there are J

such even j in total. Then  $N = \sum_{j=1}^{J} X_j$ , and by an analogous argument to that of Lemma 5.11 we find that  $\mathbb{P}(X_j|X_{j-1},\ldots X_1) \geq 0.1$ . Thus, for every 0 < r < q = 0.1, we have by the Markov inequality that

$$\mathbb{P}\left(\sum_{j=1}^{J} (1 - X_j) \ge (1 - r)J\right) \le \frac{\mathbb{E}(\sum_{j=1}^{J} (1 - X_j))}{(1 - r)J}$$
$$\le \frac{1 - q}{1 - r}$$
$$\mathbb{P}(N \le rJ) \le \frac{1 - q}{1 - r}$$
$$\mathbb{P}(N > rJ) \ge \frac{q - r}{1 - r}.$$

Observe that since  $J \ge 200 \log \log n$ , plugging in r = 0.01 gives us that  $\mathbb{P}(A_1) \ge 0.9$ , as desired. Now, suppose  $A_1$  is true. Let  $s_i$  be the *i*th *j* for which  $F_j$  is true. First, note that for all  $j \le 402 \log \log n$ ,

$$f(n/k^{j} - n/k^{j+1}) \ge f(n/k^{402\log\log n} - n/k^{402\log\log n + 1})$$
  
 
$$\ge f(n/2k^{402\log\log n}) = f(n/2e^{(\log\log n)^{2}}).$$

Hence, each  $F_j$  is contained in the event that  $t_{i_{j+2}} - t_{i_j} \ge \min(f(n/2e^{(\log\log n)^2})/1700, c_9p^2t^2)$ . Now, recalling that  $\log\log n \ge c_8/p^2 \ge e/c_9p^2$ , we may find by induction that for all  $i, t_{i_{s_i}} \ge \min(f(n/2e^{(\log\log n)^2})/1700, e^{2^i}/c_9p^2)$ . Thus,  $A_1$  is contained in the event that

$$\begin{split} t_{i_{402\log\log n}} & \geq t_{i_{s_{2\log\log n}}} \\ & \geq \min\left(\frac{f(n/2e^{(\log\log n)^2})}{1700}, \frac{e^{2^{2\log\log n}}}{c_9p^2}\right) \\ & \geq \min\left(\frac{f(n/2e^{(\log\log n)^2})}{1700}, \frac{n}{c_9p^2}\right) \\ & \geq \frac{f(n/2e^{(\log\log n)^2})}{1700c_9}, \end{split}$$

as desired.  $\Box$ 

We now prove a bound on  $t_{i_{\ell+2}} - t_{i_{\ell}}$ , for any even  $\ell \in (402 \log \log n, 402 (\log \log n)^2]$ . Combined with the initial bound on  $t_{i_{402 \log \log n}}$  given by Lemma 5.9, this will give us the key bound on  $T_n$  that will allow us to prove Lemma 5.8.

**Lemma 5.12.** For every real  $t \ge 0$  and even  $\ell$  with  $402 \log \log n < \ell < 402 (\log \log n)^2$ , there exists an event  $V_{\ell}$  such that

- 1.  $V_{\ell}$  only depends on the random variables
  - $D_{j,t}$ , for all j,t,
  - $D'_{i,t}$ , for  $j \leq \ell + 2$  and  $t \in \mathbb{Z}^+$ ,
  - $B_{j,t}$  and  $B'_{j,t}$ , for  $j \geq a_{i_{\ell+2}}$  and  $t \in \mathbb{Z}^+$ ,
  - $C_{j,t}$ , for for  $j \leq \ell + 2$  and  $t \in \mathbb{Z}^+$ .
- 2.  $A_1 \cap V_\ell$  is contained in the event that

$$t_{i_{\ell+2}} - t_{i_{\ell}} \ge \frac{\min(t_{i_{\ell}}, n/k^{\ell})}{4}.$$

- 3. Conditioned on the random variables
  - $D_{j,t}$ , for all j,t,
  - $D'_{j,t}$ , for  $j \leq \ell$  and  $t \in \mathbb{Z}^+$ ,
  - $B_{j,t}$  and  $B'_{j,t}$ , for  $j \geq a_{i_{\ell}}$  and  $t \in \mathbb{Z}^+$ ,
  - $C_{j,t}$ , for for  $j \leq \ell$  and  $t \in \mathbb{Z}^+$ ,

(which determine  $t_{i_{\ell}}$ ), we have that  $\mathbb{P}(V_{\ell}) \geq (1 - e^{-pt_{i_{\ell}}/8})(1 - e^{-\min(pt_{i_{\ell}}, n/k^{\ell})/12})$ .

*Proof.* Suppose that after m-1 steps, vertex 1 is still connected to vertex  $a_{i_{\ell}}$ . Then, on the mth step, an operation on any child of 1 in  $[a_{i_{\ell+2}}, a_{i_{\ell}})$  increases the number of children of vertex 1 in  $[a_{i_{\ell+2}}, a_{i_{\ell}})$ . Now, let N be the number of children of 1 in  $[a_{i_{\ell+2}}, a_{i_{\ell}})$  after  $t_{i_{\ell}} + g_1 - 1$  steps. We thus have

$$N \ge \min\left(\sum_{i=q_1}^{t_{i_\ell}+g_1-1} C_{\ell+2,i}, a_{i_{\ell+2}} - a_{i_\ell}\right).$$

Now, condition on the random variables

- $D_{j,t}$ , for all j,t,
- $D'_{i,t}$ , for  $j \leq \ell$  and  $t \in \mathbb{Z}^+$ ,
- $B_{j,t}$  and  $B'_{i,t}$ , for  $j \geq a_{i_{\ell}}$  and  $t \in \mathbb{Z}^+$ ,
- $C_{i,t}$ , for for  $j \leq \ell$  and  $t \in \mathbb{Z}^+$ .

Let  $E_{6,\gamma_1}$  be the event that  $\sum_{i=g_1}^{t_{i_\ell}+g_1-1} C_{\ell+2,i} > \gamma_1 p t_{i_\ell}$ . By a Chernoff bound, we have that for all  $\gamma_1 \leq 1$ ,

$$\mathbb{P}\left(\sum_{i=g_1}^{t_{i_{\ell}}+g_1-1} C_{\ell+2,i} \le \gamma_1 p t_{i_{\ell}}\right) \le e^{-(1-\gamma_1)^2 p t_{i_{\ell}}/2},$$

$$\mathbb{P}(E_{6,\gamma_1}) \ge 1 - e^{-(1-\gamma_1)^2 p t_{i_{\ell}}/2}.$$

Now, let  $X_0=0$ , and for i>0, let  $X_i$  be the ith smallest index j>0 such that  $D'_{\ell,g_1+t_{i_\ell}+j}=1$ . Let  $Y_i=X_i-X_{i-1}$ . Note that the sequence of  $\{Y_i\}$ 's are distributed as a sequence of mutually independent geometric random variables with parameter p. Thus, conditioning on  $E_{6,1/2}$ , we find that  $N\geq \max(pt_{i_\ell}/2,a_{i_{\ell+2}}-a_{i_\ell})$ . Letting  $R=\max(pt_{i_\ell}/2,a_{i_{\ell+2}}-a_{i_\ell})$ , we find that vertex 1 must receive at least R operations until it is disconnected from all vertices in  $[a_{i_{\ell+2}},a_{i_\ell})$ . Thus

$$t_{i_{\ell+2}} - t_{i_{\ell}} \ge \sum_{i=1}^{R} Y_i.$$

Now, let  $E_{7,\gamma_2}$  be the event that  $\sum_{i=1}^R Y_i \geq \gamma_2 R/p$ . Applying Lemma 2.1 (b), we have that for all  $\gamma_2 \leq 1$ ,

$$\mathbb{P}\bigg(\sum_{i=1}^{R} Y_i < \gamma_2 R/p\bigg) \le e^{-\frac{2Rp(1-\gamma_2)^2}{\gamma_2}},$$

$$\mathbb{P}(E_{7,\gamma_2}) \ge 1 - e^{-\frac{2Rp(1-\gamma_2)^2}{\gamma_2}}.$$

Now, let  $V_{\ell}$  now be the intersection of the events that  $E_{6,1/2}$  and  $E_{7,1/2}$ . Then, we have  $\mathbb{P}(V_{\ell}) \geq (1 - e^{-pt/8})(1 - e^{-\min(pt,n/k^{\ell})/12})$  (where here we use that  $a_{i_{\ell}} - a_{i_{\ell+2}} \geq n/k^{\ell}$ , as previously discussed). Moreover,  $V_{\ell}$  is contained in the event that  $t_{i_{\ell+2}} - t_{i_{\ell}} \geq R/2p = \max(t/4, (a_{i_{\ell+2}} - a_{i_{\ell}})/2p)) \geq \max(t_{i_{\ell}}, n/k^{\ell})/4$ , so we are done.

We now return to the proof of Lemma 5.8. Define  $A_2$  to be the intersection of the events  $V_\ell$  across all even indices  $402 \log \log n \le \ell \le 402 (\log \log n)^2$ . Suppose  $A_1 \cap A_2$  is true. By Lemma 5.9, we know that for all  $\ell \ge 402 \log \log n$ ,  $t_{i_\ell} \ge f(n/2e^{(\log \log n)^2})/1700c_9$ . Thus, by induction, we have that for all even  $402 \log \log n \le \ell \le 402 (\log \log n)^2$ ,

$$\begin{split} t_{i_{\ell}} &\geq \min \left( \frac{n}{4e^{402(\log\log n)^2}}, \frac{f(n/2e^{402(\log\log n)^2})}{1700c_9} \left( \frac{5}{4} \right)^{\frac{\ell - 2\lceil 201\log\log n \rceil}{2}} \right) \\ t_{402(\log\log n)^2} &\geq \min \left( \frac{n}{4e^{402(\log\log n)^2}}, \frac{f(n/2e^{402(\log\log n)^2})}{1700c_9} \left( \frac{5}{4} \right)^{200(\log\log n)^2} \right) \\ &\geq \min \left( \frac{n}{4e^{402(\log\log n)^2}}, \frac{f(n/2e^{402(\log\log n)^2})}{1700c_9} e^{44(\log\log n)^2} \right). \end{split}$$

Adjusting  $c_7$  (and hence  $c_8$ ) appropriately, we find that

$$\min\left(\frac{n}{4e^{402(\log\log n)^2}}, \frac{f(n/2e^{402(\log\log n)^2})}{1700c_9}e^{44(\log\log n)^2}\right) \ge f(n) = \frac{n}{e^{c_7(\log\log n)^3}}.$$

Hence,  $A_1 \cap A_2$  implies that

$$T_n \ge t_{402(\log \log n)^2} + g_1 - 1$$
  
  $\ge f(n) + g_1 - 1.$ 

Now it suffices to compute  $\mathbb{P}(A_1 \cap A_2)$ . Condition on the random variables

- $D_{i,t}$ , for  $j \leq i_{402 \log \log n}$  and  $t \in \mathbb{Z}^+$ , and
- $B_{j,t}$  and  $B'_{j,t}$ , for  $j \ge a_{i_{402 \log \log n}}$  and  $t \in \mathbb{Z}^+$ .

Recall from Lemma 5.9 that these variables determine  $A_1$ . For any even  $\ell \in (402 \log \log n, 402(\log \log n)^2]$ , note that  $V_{2\lceil 201 \log \log n \rceil}, \ldots, V_{\ell}$  all only depend on the random variables

- $D_{i,t}$ , for all j,t,
- $D'_{i,t}$ , for  $j \leq \ell + 2$  and  $t \in \mathbb{Z}^+$ ,
- $B_{j,t}$  and  $B'_{j,t}$ , for  $j \ge a_{i_{\ell+2}}$  and  $t \in \mathbb{Z}^+$ ,
- $C_{i,t}$ , for for  $j \leq \ell + 2$  and  $t \in \mathbb{Z}^+$ .

Hence with an appropriate adjustment of  $c_9$  (to establish the last inequality), we have that

$$\mathbb{P}(V_{\ell+1}: V_{2\lceil 201 \log \log n \rceil}, \dots V_l) \ge (1 - e^{-pt_{402 \log \log n}/8}) (1 - e^{-\min(pt_{402 \log \log n}, n/k^{402 (\log \log n)^2})/12})$$

$$\ge 1 - e^{-p\sqrt{n}}.$$

Thus,

$$\mathbb{P}(A_2|A_1) \ge 1 - 402(\log\log n)^2 e^{-p\sqrt{n}}.$$

By Lemma 5.9 we know  $\mathbb{P}(A_1) \ge 0.9$ . Hence  $\mathbb{P}(A_1 \cap A_2) \ge 0.9 - 402(\log \log n)^2 e^{-p\sqrt{n}}$ , as desired.

# 5.2 Proofs of Theorem 5.7, Theorem 1.6

We now come back to prove Theorems 5.7 and 1.6.

Proof of Theorem 5.7. The statement is trivial for  $n \leq e^{e^{c_8/p^2}}$ . Henceforth assume  $n > e^{e^{c_8/p^2}}$ . Throughout this proof, let  $k = \log n$ . We first claim the following.

Lemma 5.13. We have that

$$\mathbb{P}\bigg(\bigcup_{A \text{ good}} E_A : E_{[1,n],m}\bigg) \ge 1 - \frac{2}{\log\log n}.$$

Proof. Simulate the Ungarian Markov chain using the algorithm from the proof of Lemma 5.8, and then condition on  $E_{[1,n],m}$ . It suffices to show that  $J=\begin{bmatrix} a_1 & a_2 & \cdots & a_l \\ b_1 & b_2 & \cdots & b_l \end{bmatrix}$  is n-good with probability at least  $1-\frac{2}{\log\log n}$ . Now, given any  $i\geq 2$ , we claim that for all reals  $r\in [0,1]$ ,

$$\mathbb{P}\left(\frac{a_{i+1} - 1}{a_i - 2} \ge r : a_i > 2\right) \ge 1 - r.$$

Indeed, if we condition by fixing any value of  $a_i$ , the probability that the ordered set  $\{g_2, \ldots, g_{a_i-1}\}$  is equal to any given  $a_i-2$ -tuple is equal to the probability  $\{g_2, \ldots, g_{a_i-1}\}$  is equal to any permutation of that tuple, and this property is not changed even if we condition on  $E_{[1,n],m}$ . Because  $a_{i+1}$  is defined as the *largest* index such that  $g_{a_{i+1}} = \max(g_2, \ldots, g_{a_i-1})$ , we retrieve the inequality in the case of fixed  $a_i$ ; the inequality above is retrieved from summing over all possible values of  $a_i$ .

Observe that by analogous reasoning we have that

$$\mathbb{P}\left(\frac{a_2 - 1}{n - 1} \ge r\right) \ge 1 - r.$$

Now let the *n*-summary of J be  $J' = \begin{bmatrix} a_1 & a_{i_1} & a_{i_2} & \cdots & a_{i_{l'}} \\ b_1 & b_{i_1} & b_{i_2} & \cdots & b_{i_{l'}} \end{bmatrix}$ . By the claim above we have that  $a_{i_1} \geq n/k$  with probability  $\geq 1 - 1/k$ , and (for a suitably large choice of  $c_8$ ) for each j satisfying that  $a_{i_j} > k^3$ , the probability that  $a_{i_{j+1}}$  exists (i.e. the summary has at least j+2 columns), conditioned on the probability that  $a_{i_j}$  exists, is at least  $1 - \frac{1}{k}$ . Moreover, by definition, for all relevant j we have that  $a_{i_{j+2}} < a_{i_j}/k$ , so thus the maximum index l for which  $a_{i_l} \geq k^3$  satisfies  $l \leq 2\log_k(n) - 7$ . Hence by a union bound we have that

$$\mathbb{P}(J \text{ is good}) \ge 1 - \frac{2\log_k(n) - 6}{k}$$
$$\ge 1 - \frac{2}{\log\log n},$$

as desired.  $\Box$ 

Now, by definition, if for some n-good array  $E_{[1,n],m}$  and  $E_A$  are simultaneously true, then the bottom leftmost entry of A is m. Hence, combining Lemma 5.8 and Lemma 5.13 and applying a union bound gives us that

$$\mathbb{P}(T_n \ge f(n) + m - 1 : E_{[1,n],m}) \ge 0.9 - 402(\log \log n)^2 e^{-p\sqrt{n}} - \frac{2}{\log \log n}$$
  
 
$$\ge \frac{1}{2},$$

as desired.

We now prove Theorem 1.6.

REFERENCES REFERENCES

Proof of Theorem 1.6. Simulate the Ungarian Markov chain on  $\operatorname{Tam}_n$  using the algorithm described in the proof of Lemma 5.8. For any  $i \in [n], M \in \mathbb{Z}^+$ , let  $A_{i,M}$  be the event that  $g_i > M = \max_{j \neq i} (g_j)$ . Suppose  $A_{i,M}$  is true. Then, by Proposition 4.4, we have that the induced subgraph formed by the vertices [i, n] transitions as per the Ungarian Markov chain on  $\operatorname{Tam}_{n-i+1}$ . Hence,

$$\mathcal{E}_p(\operatorname{Tam}_n : A_{i,M}) \ge \mathcal{E}_p(\operatorname{Tam}_{n-i+1} : E_{[1,n-i+1],M})$$

$$\ge \mathbb{E}[T_{n-i+1} : E_{[1,n-i+1],M}]$$

$$\ge \frac{f(n-i+1)}{2},$$

where the colons denote conditioning, and the last inequality is by Theorem 5.7. Now, note that  $\sum_{M=1}^{\infty} \mathbb{P}(A_{i,M})$  is the probability that  $t_i > t_j$  for all  $j \neq i$ , which is precisely  $\frac{\zeta_p(n)}{n}$  (by symmetry). Thus,

$$\mathcal{E}_{p}(\operatorname{Tam}_{n}) \geq \sum_{i=1}^{n} \sum_{M=1}^{\infty} \mathcal{E}_{p}(\operatorname{Tam}_{n} : A_{i,M}) \mathbb{P}(A_{i,M})$$

$$\geq \frac{f(\lceil n/2 \rceil)}{2} \sum_{i=1}^{\lfloor n/2 \rfloor} \sum_{M=1}^{\infty} \mathbb{P}(A_{i,M})$$

$$\geq \frac{(n-1)\zeta_{p}(n)}{4n} f(\lceil n/2 \rceil).$$

This readily implies Theorem 1.6, as desired.

# References

[1] Jinho Baik, Robert Buckingham, and Jeffrey DiFranco. Asymptotics of Tracy-Widom distributions and the total integral of a Painlevé II function. *Comm. Math. Phys.*, 280(2):463–497, 2008.

- [2] F. Thomas Bruss and Colm Art O'Cinneide. On the Maximum and Its Uniqueness for Geometric Random Samples. J. Appl. Probab., 27(3):598–610, 1990.
- [3] Colin Defant and Rupert Li. Ungarian Markov Chains. Electronic Journal of Probability, 28:1–39, 2023.
- [4] Kurt Johansson. Shape Fluctuations and Random Matrices. Comm. Math. Phys., 209(2):437–476,2000.
- [5] Nathan Reading. Cambrian Lattices. Adv. Math., 205(2):313–353, 2006.
- [6] Nathan Reading. Sortable elements and Cambrian lattices. Algebra Universalis, 56(3-4):411-437, 2007.
- [7] Dan Romik. The Surprising Mathematics of Longest Increasing Subsequences, volume 4 of Institute of Mathematical Statistics Textbooks. Cambridge University Press, New York, 2015.
- [8] P. R. Scott. On the Sets of Directions Determined by n Points. Amer. Math. Monthly, 77:502–505, 1970.
- [9] Richard P. Stanley. Enumerative Combinatorics. Vol. I. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Monterey, CA, 1986. With a foreword by Gian-Carlo Rota.
- [10] Dov Tamari. The algebra of bracketings and their enumeration. *Nieuw Arch. Wisk.* (3), 10:131–146, 1962.
- [11] Peter Ungar. 2N Noncollinear points determine at least 2N directions. J. Combin. Theory Ser. A, 33(3):343-347, 1982.