

# HIGHER WEIGHT SPECTRA OF TERNARY CODES ASSOCIATED TO THE QUADRATIC VERONESE 3-FOLD

KRISHNA KAIPA AND PUSPENDU PRADHAN

ABSTRACT. The problem studied in this work is to determine the higher weight spectra of the Projective Reed-Muller codes associated to the Veronese 3-fold  $\mathcal{V}$  in  $PG(9, q)$ , which is the image of the quadratic Veronese embedding of  $PG(3, q)$  in  $PG(9, q)$ . We reduce the problem to the following combinatorial problem in finite geometry: For each subset  $S$  of  $\mathcal{V}$ , determine the dimension of the linear subspace of  $PG(9, q)$  generated by  $S$ . We develop a systematic method to solve the latter problem. We implement the method for  $q = 3$ , and use it to obtain the higher weight spectra of the associated code. The case of a general finite field  $\mathbb{F}_q$  will be treated in a future work.

## 1. INTRODUCTION

Let  $PG(n, q)$  denote the  $n$ -dimensional projective space defined over a finite field  $\mathbb{F}_q$ . Let  $K = (n^2 + 3n + 2)/2$ . The quadratic Veronese  $n$ -fold  $\mathcal{V}_n$  in  $PG(K - 1, q)$  is the image of the quadratic embedding

$$\iota : PG(n, q) \hookrightarrow PG(K - 1, q),$$

defined with respect to coordinates  $(x_0, \dots, x_n)$  on  $PG(n, q)$  and  $(z_0, \dots, z_{K-1})$  on  $PG(K - 1, q)$  by

$$\iota(x_0, \dots, x_n) = (x_0^2, x_0x_1, \dots, x_{n-1}x_n, x_n^2),$$

for some fixed ordering of the  $K$  monomials of degree 2 in the variables  $x_0, \dots, x_n$ . Let  $P_1, \dots, P_N$  where  $N = 1 + q + \dots + q^n$ , be a fixed ordering of the points of  $PG(n, q)$  and let  $\mathbb{G}_n$  denote the matrix of size  $K \times N$  defined by

$$\mathbb{G}_n = [v_1 \ \dots \ v_N],$$

where  $v_j \in \mathbb{F}_q^K$  represents the point  $\iota(P_j)$  of  $PG(K - 1, q)$ . The linear code of length  $N$  and dimension  $K$  generated by the matrix  $\mathbb{G}_n$  is the quadratic Veronese code  $C_n$ , or the second order Projective Reed Muller code of length  $N$  ([5, §5]). A codeword of the code  $C_n$  is a quadratic form  $f$  on  $PG(n, q)$  and the weight of a codeword is the number of points of  $PG(n, q)$  not lying on the corresponding quadric hypersurface defined by the vanishing of  $f$ . An  $r$ -dimensional subcode of  $C_n$  is a linear subspace of  $C_n$  generated by some  $r$  linearly independent quadratic forms  $f_1, \dots, f_r$  on  $PG(n, q)$ . The weight of this subcode is the number of points of  $PG(n, q)$  which do not lie on the common zero set of  $f_1, \dots, f_r$ . Equivalently,  $r$ -dimensional subcodes of  $C_n$  correspond to codimension  $r$  linear subspaces  $\mathcal{L}$  of  $PG(K - 1, q)$ , with the weight of the subcode being the number of points of  $\mathcal{V}_n$  not lying on  $\mathcal{L}$ . The  $r$ -th generalized weight enumerator (in short GWE) of  $C_n$  is the polynomial

$$W^{(r)}(Z) = \sum_{\mathcal{L} \subset PG(K-1, q) \text{ of codim } r} Z^{\#\mathcal{V}_n \setminus \mathcal{L}} = \sum_i A_i^{(r)} Z^i,$$

2020 *Mathematics Subject Classification.* 94B27, 51E20, 05B25.

*Key words and phrases.* Quadratic Veronese varieties, Veronese code, Linear system of quadrics, Generalized weight enumerator polynomial, Extended weight enumerator polynomial.

where  $A_i^{(r)}$  is the number of codimension  $r$  linear subspaces of  $PG(K-1, q)$  which intersect  $\mathcal{V}_n$  in  $N-i$  points.

We also explain the algebraic-geometric significance of the polynomial  $Z^N W^{(r)}(\frac{1}{Z})$ . It gives the answer to the following geometric problem:

**Problem.** *Given natural numbers  $n$  and  $r$ , let  $\mathcal{S}_n^{(r)}$  denote the collection of varieties in  $\mathbb{P}^n$  defined by  $r$  linearly independent quadratic forms  $f_1(X_0, \dots, X_n), \dots, f_r(X_0, \dots, X_n)$  over  $\mathbb{F}_q$ . For  $V \in \mathcal{S}_n^{(r)}$ , let  $\#V(\mathbb{F}_q)$  denote the number of  $\mathbb{F}_q$ -rational points of  $V$ . Determine the generating function*

$$\sum_{V \in \mathcal{S}_n^{(r)}} Z^{\#V(\mathbb{F}_q)} = \sum_{i \geq 0} c_i^{(r)}(n) Z^i,$$

where  $c_i^{(r)}(n)$  is the number of varieties in  $\mathcal{S}_n^{(r)}$  having  $i$  rational points over  $\mathbb{F}_q$ .

The numbers  $c_i^{(r)}(n)$  are just the numbers  $A_{N-i}^{(r)}(C_n)$ .

The first GWE of  $C_n$  is easily calculated (see Theorem 2.1). However, for  $r > 1$ , the  $r$ -th GWEs of  $C_n$  are hard to determine. In the work [2], Johnsen and Verdure determined the  $r$ -GWE's of the code  $C_2$  for all  $r$  and all finite fields  $\mathbb{F}_q$ . In the work [3], they further obtained the  $r$ -th GWEs of the code  $C_3$  for all  $r$  over the binary field  $\mathbb{F}_2$ . In this work, we develop a method to determine all the  $r$ -GWE's of the code  $C_3$  over a general finite field  $\mathbb{F}_q$ . We now explain the gist of our method: Given a linear code  $C$  of dimension  $K$  with a generator matrix  $M$ , let  $B_{j,i}$  denote the number of  $K \times j$  submatrices of  $M$  that have rank  $i$ . As shown by Jurrius-Pellikaan [4], the knowledge of these quantities  $B_{j,i}$  for all  $(j, i)$  suffices to determine the  $r$ -GWE's of the code  $C$  for all  $r$ . For the code  $C_3$ , the quantities  $B_{j,i}$  are the number of configurations of  $j$  points in  $PG(3, q)$  whose images in  $\mathcal{V}_3$  span a  $(i-1)$ -dimensional linear subspace in  $PG(9, q)$ . We treat this problem from a finite geometry perspective, and develop a systematic method to calculate the quantities  $B_{j,i}$  for all  $(j, i)$ , and hence obtain the  $r$ -th GWE of  $C_3$  for all  $r$  and for all fields  $\mathbb{F}_q$ . We execute the method for  $q = 3$  in the present work, and the case of general  $q$  will be treated in a future work.

**Theorem 1.1.** *For  $1 \leq r \leq 10$ , let  $W^r(Z)$  denote the  $r$ -th GWE of the code  $C_3$  over the ternary field  $\mathbb{F}_3$ . We have:*

$$\begin{aligned}
 W^{(1)}(Z) &= 2Z^{18}(195Z^{18} + 4212Z^{12} + 4700Z^9 + 5265Z^6 + 390) \\
 W^{(2)}(Z) &= 13Z^{24}(67797Z^{16} + 243000Z^{15} + 597780Z^{14} + 933120Z^{13} + 1151610Z^{12} + 991440Z^{11} \\
 &\quad + 816480Z^{10} + 428400Z^9 + 182250Z^8 + 116640Z^7 + 43200Z^6 + 14580Z^4 + 760Z^3 + 360). \\
 W^{(3)}(Z) &= 10Z^{26}(408114369Z^{14} + 615347208Z^{13} + 471425994Z^{12} + 230935536Z^{11} + 79511991Z^{10} \\
 &\quad + 20839572Z^9 + 5231304Z^8 + 968760Z^7 + 237978Z^6 + 50544Z^5 + 9360Z^4 + 52Z + 108). \\
 W^{(4)}(Z) &= Z^{27}(304888261329Z^{13} + 152054622720Z^{12} + 37402159860Z^{11} + 5713746480Z^{10} \\
 &\quad + 641358900Z^9 + 69893928Z^8 + 7076160Z^7 + 627120Z^6 + 89505Z^5 + 40). \\
 W^{(5)}(Z) &= 26Z^{33}(49159423665Z^7 + 8099569140Z^6 + 649260360Z^5 + 31541400Z^4 \\
 &\quad + 1366565Z^3 + 73548Z^2 + 2430Z + 120). \\
 W^{(6)}(Z) &= 2Z^{35}(237163797936Z^5 + 12886036860Z^4 + 334019790Z^3 + 4956120Z^2 + 104975Z + 2340). \\
 W^{(7)}(Z) &= 10Z^{36}(1800649161Z^4 + 31769712Z^3 + 252954Z^2 + 936Z + 13). \\
 W^{(8)}(Z) &= 13Z^{38}(5557197Z^2 + 30160Z + 60). \\
 W^{(9)}(Z) &= 4Z^{39}(7371Z + 10). \\
 W^{(10)}(Z) &= Z^{40}.
 \end{aligned}$$

In particular, the generalized Hamming weights of the code  $C_3$  over  $\mathbb{F}_3$  are given by

$$d_1 = 18, d_2 = 24, d_3 = 26, d_4 = 27, d_5 = 33, d_6 = 35, d_7 = 36, d_8 = 38, d_9 = 39, d_{10} = 40.$$

The rest of this paper is organized as follows. In §2, we record the ordinary weight enumerator of the code  $C_n$ . In §3, we explain how the quantities  $B_{j,i}$  determine the  $r$ -GWE's of a code  $C$ , and we develop the geometric method to determine  $B_{j,i}$ 's for the code  $C_3$ . In §4, we determine  $B_{j,1}, B_{j,2}, B_{j,3}, B_{j,4}$  and  $B_{j,5}$  for all  $j$  and for a general finite field  $\mathbb{F}_q$ . In §5, §6, §7 and §8, we determine the quantities  $B_{j,6}, B_{j,7}, B_{j,8}$  and  $B_{j,9}$  respectively for all  $j$  and  $q = 3$ . Finally, in §9 we prove Theorem 1.1.

## 2. ORDINARY WEIGHT ENUMERATOR OF THE CODES $C_n$

We will write down the first generalized weight enumerator  $W_{C_n}^{(1)}(Z)$  of  $C_n$ . The ordinary weight enumerator of  $C_n$  is just  $1 + (q-1)W_{C_n}^{(1)}(Z)$ . We need some notation. Let  $GL_n(q)$  denote the general linear group of invertible  $n \times n$  matrices over  $\mathbb{F}_q$ , The corresponding projective general linear group will be denoted  $PGL_n(q)$ . Let

$$(1) \quad \begin{bmatrix} r \\ j \end{bmatrix}_q = \frac{\prod_{i=0}^{j-1} (q^r - q^i)}{|GL_j(q)|} = \prod_{i=0}^{j-1} \frac{(q^r - 1)(q^{r-1} - 1) \dots (q^{r-j+1} - 1)}{(q^j - 1)(q^{j-1} - 1) \dots (q - 1)},$$

denote the Gaussian binomial coefficient, which is the number of  $j$ -dimensional subspaces in  $\mathbb{F}_q^r$ . Let  $f_0(X, Y)$  denote a fixed quadratic form in the variables  $X, Y$  which is irreducible over  $\mathbb{F}_q$ . We recall from [1, Chapter 1] that the non-degenerate quadratic forms on  $PG(2t, q)$  form a single orbit with representative  $X_0X_1 + X_2X_3 + \dots + X_{2t-2}X_{2t-1} + X_{2t}^2$ . The corresponding quadrics in

$PG(2t, q)$  are known as parabolic quadrics, and have  $(q^{2t} - 1)/(q - 1)$  points. The size of this orbit is  $q^{t^2+t} \prod_{i=1}^t (q^{2i+1} - 1)$ . The non-degenerate quadratic forms on  $PG(2t - 1, q)$  form two orbits represented by  $X_0X_1 + X_2X_3 + \dots + X_{2t-2}X_{2t-1}$  and  $X_0X_1 + X_2X_3 + \dots + X_{2t-4}X_{2t-3} + f(X_{2t-2}, X_{2t-1})$ . These orbits have sizes  $\frac{1}{2}q^{t^2}(q^t + 1) \prod_{i=1}^{t-1} (q^{2i+1} - 1)$  and  $\frac{1}{2}q^{t^2}(q^t - 1) \prod_{i=1}^{t-1} (q^{2i+1} - 1)$  respectively. The corresponding quadrics in  $PG(2t - 1, q)$  are known as hyperbolic and elliptic quadrics respectively, and they have  $(q^{t-1} + 1)(q^t - 1)/(q - 1)$  and  $(q^{t-1} - 1)(q^t + 1)/(q - 1)$  points respectively. Given  $m < n$  and a quadratic form  $f(X_0, \dots, X_m)$  such that the corresponding quadric in  $PG(m, q)$  has  $\nu$  points, then the quadric defined by  $f$  in  $PG(n, q)$  has  $1 + q + \dots + q^{n-m-1} + \nu q^{n-m}$  points. Thus, the weight of  $f$  viewed as a codeword of  $C_n$  is  $q^{n-m}(1 + q + \dots + q^m - \nu)$ . Since there are  $\begin{bmatrix} n+1 \\ m+1 \end{bmatrix}_q$  ways to pick a  $m$ -dimensional linear subspace of  $PG(n, q)$  we conclude that the values that weights of codewords of  $C_n$  can take are as listed below:

(1)  $q^n + q^{n-t}$ , where  $1 \leq t \leq \lfloor (n+1)/2 \rfloor$ . The number of codewords with this weight are

$$A_{q^n+q^{n-t}}^{(1)} = \frac{1}{2} \begin{bmatrix} n+1 \\ 2t \end{bmatrix}_q q^{t^2} (q^t - 1) \prod_{i=1}^{t-1} (q^{2i+1} - 1).$$

(2)  $q^n$ , and the number of codewords with this weight are

$$A_{q^n}^{(1)} = \sum_{t=0}^{\lfloor n/2 \rfloor} \begin{bmatrix} n+1 \\ 2t+1 \end{bmatrix}_q q^{t^2+t} \prod_{i=1}^t (q^{2i+1} - 1).$$

(3)  $q^n - q^{n-t}$ , where  $1 \leq t \leq \lfloor (n+1)/2 \rfloor$ . The number of codewords with this weight are

$$A_{q^n-q^{n-t}}^{(1)} = \frac{1}{2} \begin{bmatrix} n+1 \\ 2t \end{bmatrix}_q q^{t^2} (q^t + 1) \prod_{i=1}^{t-1} (q^{2i+1} - 1).$$

We summarize this:

**Theorem 2.1.** *The ordinary weight enumerator of the code  $C_n$  is given by  $1 + (q - 1)W^{(1)}(C_n)$  where:*

$$(2) \quad W_{C_n}^{(1)}(Z) = A_{q^n}^{(1)} Z^{q^n} + \sum_{t=1}^{\lfloor (n+1)/2 \rfloor} \left( A_{q^n-q^{n-t}}^{(1)} Z^{q^n-q^{n-t}} + A_{q^n+q^{n-t}}^{(1)} Z^{q^n+q^{n-t}} \right),$$

and the quantities  $A_j^{(1)}$  are as above.

In particular, for  $n = 3$  we have 6 orbits of quadratic forms on  $PG(3, q)$ , which we record in Table 1, including their sizes, and weights when regarded as codewords of the code  $C_n$ . Specializing (2) to  $n = 3$ , we get:

$$(3) \quad W_{C_3}^{(1)}(Z) = \frac{1}{2}(q^3 + q^2 + q + 1)(q^3 + q^2 + q) Z^{q^3-q^2} + \frac{1}{2}(q^6 + q^4)(q^3 - 1) Z^{q^3-q} \\ + (q^5 - q^2 + 1)(q^3 + q^2 + q + 1) Z^{q^3} \\ + \frac{1}{2}(q^6 - q^4)(q^3 - 1) Z^{q^3+q} + \frac{1}{2}(q^3 - 1)(q^3 + q) Z^{q^3+q^2}$$

Further specializing this to  $q = 3$ , we get

$$2Z^{18}(195Z^{18} + 4212Z^{12} + 4700Z^9 + 5265Z^6 + 390),$$

which matches with Theorem 1.1.

Orbit	Representative	Size	weight
$\mathcal{O}_1$	$X_0X_1$	$\binom{q^3+q^2+q+1}{2}$	$q^3 - q^2$
$\mathcal{O}_2$	$X_0X_3 - X_1X_2$	$\frac{1}{2}(q^6 + q^4)(q^3 - 1)$	$q^3 - q$
$\mathcal{O}_3$	$X_0^2$	$(1 + q)(1 + q^2)$	$q^3$
$\mathcal{O}_4$	$X_1^2 - X_0X_2$	$(1 + q)(1 + q^2)(q^5 - q^2)$	$q^3$
$\mathcal{O}_5$	$X_2X_3 - f_0(X_0, X_1)$	$\frac{1}{2}(q^6 - q^4)(q^3 - 1)$	$q^3 + q$
$\mathcal{O}_6$	$f_0(X_0, X_1)$	$\frac{1}{2}(q^3 - 1)(q^3 + q)$	$q^3 + q^2$

 TABLE 1. List of  $G$ -orbits of quadratic forms on  $PG(3, q)$ 

### 3. APPROACH TO THE PROBLEM

Let  $C$  be a linear  $[N, K, d]_q$  code over  $\mathbb{F}_q$  of length  $N$ , dimension  $K$  and minimum distance  $d$  with generator matrix  $M_{K \times N}$ . For  $J \subset \{1, \dots, N\}$ , let  $r_J$  be the rank of the submatrix of  $M_{K \times N}$  on the columns indexed by  $J$ . The higher weight spectrum of  $C$  can be obtained just by knowing  $r_J$  for each  $J \subset \{1, \dots, N\}$ . To see this, consider for each  $1 \leq j \leq N$ , the polynomials  $B_j(T) \in \mathbb{Z}[T]$  defined by

$$(4) \quad B_j(T) = \sum_J (T^{K-r_J} - 1),$$

where the sum runs over all subsets  $J \subset \{1, \dots, N\}$  of size  $j$ . We define  $B_0(T) = (T^K - 1)$ . The *extended weight enumerator* polynomial (in short EWE) is the polynomial  $W_C(Z; T) \in \mathbb{Z}[Z, T]$  defined by

$$(5) \quad \begin{aligned} W(Z; T) &= 1 + \sum_{j=0}^N B_j(T)(1 - Z)^j Z^{N-j} \\ &= 1 + (T^K - 1)Z^N + \sum_{j=1}^N B_j(T)(1 - Z)^j Z^{N-j}. \end{aligned}$$

Let  $A_j(T) \in \mathbb{Z}[T]$  be polynomials defined by

$$(6) \quad W(Z; T) = 1 + \sum_{j=1}^N A_j(T)Z^j.$$

The polynomial  $W(Z; q^m)$  is the weight enumerator polynomial of the extended code  $C \otimes \mathbb{F}_{q^m}$  (see [4, Proposition 5.23]). Thus,  $A_j(q^m)$  is the number of weight  $j$  codewords of  $C \otimes \mathbb{F}_{q^m}$ . The  $r$ -th generalized weight enumerator polynomial of  $C$  can be obtained from extended weight enumerator polynomial:

**Theorem 3.1.** [4, Theorem 5.7] *Let  $C$  be a linear code over  $\mathbb{F}_q$  and  $0 \leq r \leq K$ . Then the  $r$ -th generalized weight enumerator polynomial of  $C$  can be obtained from the extended weight enumerator polynomial of  $C$  by:*

$$W^{(r)}(Z) = \frac{1}{|GL_r(q)|} \sum_{j=0}^r \begin{bmatrix} r \\ j \end{bmatrix}_q (-1)^{r-j} q^{\binom{r-j}{2}} W(Z; q^j).$$

From (4), (5) and Theorem 3.1, it follows that determining the quantities  $r_J$  for all  $J \subset \{1, \dots, N\}$  allows us to obtain the  $r$ -th generalized weight enumerator polynomial of  $C$ . It will be convenient to introduce the numbers  $B_{j,i}$  defined to be the number of  $J \subset \{1, \dots, N\}$  such that  $|J| = j$  and  $\text{rank}(G_J) = i$ . In terms of  $B_{j,i}$  we can write

$$B_j(T) = \sum_{i \geq 1} B_{j,i} (T^{K-i} - 1).$$

We note that in the polynomial  $B_j(T)$  the quantities  $B_{j,K}$  are multiplied by  $(T^{K-K} - 1) = 0$ . Therefore, there is no need to determine the quantities  $B_{j,K}$ . In other words, we may re-define

$$(7) \quad B_j(T) = \sum_{i=1}^{K-1} B_{j,i} (T^{K-i} - 1).$$

By [4, Theorem 5.4], we have  $A_0(T) = 1$ , and for  $j > 0$ :

$$(8) \quad A_j(T) = \sum_{i=N-j}^{N-d} (-1)^{N+j+i} \binom{i}{N-j} B_i(T).$$

It will be useful to note that :

$$(9) \quad B_{j,j} = \binom{N}{j} - \sum_{i < j} B_{j,i}, \quad \text{if } j \leq K.$$

Although, in the expression (7) the quantities  $B_{j,K}$  were not required, for later use we note that

$$(10) \quad B_{j,K} = \binom{N}{j} - \sum_{i < K} B_{j,i}, \quad \text{if } j > K.$$

We now return to the code  $C_3$ . Here the Veronese embedding  $\iota : PG(3, q) \rightarrow PG(9, q)$  is given by:

$$(11) \quad \iota(x_0, \dots, x_3) = (x_0^2, x_0x_1, x_1^2, x_0x_2, x_1x_2, x_2^2, x_0x_3, x_1x_3, x_2x_3, x_3^2).$$

The generator matrix  $\mathbb{G}_3$  of  $C_3$  will be denoted as just  $\mathbb{G}$ . We need some more notation and definitions.

- A *configuration*  $S$  is just a set of points of  $PG(3, q)$ .
- The group  $PGL_4(q)$  will be denoted as  $G$ .
- For a homogeneous ideal  $I$  of  $\mathbb{F}_q[X_0, X_1, X_2, X_3]$  generated by quadratic forms  $f_1, \dots, f_r$ , let  $V(I)$  denote the intersection of the quadrics  $f_1 = \dots = f_r = 0$  in  $PG(3, q)$ . For  $S \subset PG(3, q)$ , let  $I_2(S)$  denote the vector space of quadratic forms  $f(X_0, X_1, X_2, X_3)$  vanishing on  $S$ .
- The *rank* of a configuration  $S$  is defined by any one of the following equivalent numbers:
  - (a) one more than the dimension of the linear subspace of  $PG(3, q)$  spanned by  $\iota(S)$ .
  - (b) the rank of the submatrix of  $\mathbb{G}$  on the columns representing the points of  $\iota(S)$ .
  - (c)  $10 - \dim(I_2(S))$

**Definition 3.2.** A configuration  $S \subset PG(3, q)$  of rank  $r$  is a maximal rank  $r$  configuration – in short a  $\mathcal{C}_r$  – if there is no rank  $r$  configuration properly containing  $S$ . Equivalently,  $S$  is a  $\mathcal{C}_r$  if  $V(I_2(S)) = S$  and  $\dim(I_2(S)) = 10 - r$ .

For example, the line  $\mathcal{L}$  given by  $x_2 = x_3 = 0$  has rank 3 because  $\iota(\mathcal{L})$  is the conic  $z_1^2 = z_0z_2$  in the plane  $z_3 = \dots = z_9 = 0$  and the span of this conic is the whole plane, or in other words the vector space  $I_2(\mathcal{L})$  has basis  $X_3^2, X_3X_2, X_3X_1, X_3X_0, X_2^2, X_2X_1, X_2X_0$ . Clearly  $V(I_2(\mathcal{L}))$  is the points of intersection of the hyperplanes  $X_3 = 0$  and  $X_2 = 0$  which is the line  $\mathcal{L}$ . Thus,  $\mathcal{L}$  is a  $\mathcal{C}_3$ . Similarly, the plane  $\mathcal{H}$  given by  $x_3 = 0$ , has rank 6 because  $\iota(\mathcal{H})$  is a Veronese surface in the 5-dimensional linear subspace  $PG(5, q) \subset PG(9, q)$  given by  $z_6 = \dots = z_9 = 0$ , and the linear span of this surface is  $PG(5, q)$ . Here  $I_2(\mathcal{H})$  has basis  $X_3^2, X_3X_2, X_3X_1, X_3X_0$  and clearly  $V(I_2(\mathcal{H}))$  is just the hyperplane  $X_3 = 0$  which is  $\mathcal{H}$ . Thus,  $\mathcal{H}$  is a  $\mathcal{C}_6$ .

We note that there is a monomorphism  $\rho : G \hookrightarrow PGL_{10}(q)$  (see [1, Theorem 4.15]) such that for each  $g \in G$  and  $v \in PG(3, q)$  we have

$$\iota(g \cdot v) = \rho(g)\iota(v),$$

and the image  $\mathcal{G}(\mathcal{V}_3) \subset PGL_{10}(q)$  of this homomorphism is the automorphism group of  $\mathcal{V}_3$  in  $PGL_{10}(q)$ . Therefore, given a line  $\mathcal{L}$  (resp. plane  $\mathcal{H}$ ) of  $PG(3, q)$  we may assume that up to  $G$ -equivalence,  $\mathcal{L}$  is given by  $x_2 = x_3 = 0$  (resp.  $\mathcal{H}$  is given by  $x_3 = 0$ ). Thus, each line of  $PG(3, q)$  is a  $\mathcal{C}_3$  and each plane of  $PG(3, q)$  is a  $\mathcal{C}_6$ .

**Lemma 3.3.** Given a rank  $r$  configuration  $S$ , there is a unique maximal rank  $r$  configuration  $\bar{S} \supset S$ . We have  $\bar{S} = V(I_2(S))$ .

*Proof.* Let  $S' = V(I_2(S))$ . We must show that i)  $S'$  has rank  $r$ , and ii) any rank  $r$  configuration  $T$  containing  $S$ , is contained in  $S'$ .

As for i), it suffices to show  $I_2(S') = I_2(S)$ . Since  $S' \supset S$ , we have  $I_2(S') \subset I_2(S)$ . On the other hand, if  $f \in I_2(S)$  then by definition of  $S'$ , it follows that  $f$  vanishes on  $S'$ , and hence  $f \in I_2(S')$ , i.e.  $I_2(S) \subset I_2(S')$ . Thus,  $I_2(S') = I_2(S)$ .

As for ii), if  $T \supset S$  has rank  $r$ , then the fact that  $\text{rank}(T) = \text{rank}(S) = r$  implies  $I_2(S) = I_2(T)$ , and hence  $T \subset V(I_2(T)) = V(I_2(S)) = S'$ . We define  $\bar{S} = V(I_2(S))$ .  $\square$

The next lemma allows us to build  $\mathcal{C}_r$ 's by adding points to each  $\mathcal{C}_{r-1}$ .

**Lemma 3.4.** If  $S$  is a  $\mathcal{C}_r$ , and  $T \subset S$  then  $\bar{T} \subset S$ . In particular,  $S$  contains a  $\mathcal{C}_{r-1}$  configuration  $T$ .

*Proof.* Since  $T \subset S$ , we get  $I_2(S) \subset I_2(T)$ , and hence  $\bar{T} = V(I_2(T)) \subset V(I_2(S)) = S$ .

We delete points of  $S$  one at a time until the rank drops by one. If  $S'$  is the resulting rank  $(r-1)$  configuration, then  $T = \bar{S}'$  is a  $\mathcal{C}_{r-1}$  configuration contained in  $S$ .  $\square$

In order to determine the rank  $r$  configurations of a given size  $j$ , it suffices to look for size  $j$  and rank  $r$  subsets of each  $\mathcal{C}_r$ . There are two steps in our approach.

Step 1 ) Determine the set  $\mathfrak{C}_r$  of all  $\mathcal{C}_r$ 's. i.e., the collection of all maximal rank  $r$  configurations. We recursively construct all  $\mathcal{C}_r$  configurations by adding points to  $\mathcal{C}_{r-1}$  configurations. We will also obtain a decomposition,

$$\mathfrak{C}_r = \cup_{\alpha \in \{a, b, c, \dots\}} \mathcal{M}_{r\alpha}.$$

Each of the parts  $\mathcal{M}_{r\alpha}$  will be a class of maximal configurations consisting of a union of  $G$ -orbits of  $\mathcal{C}_r$ 's. We will also determine the size of each class  $\mathcal{M}_{r\alpha}$ .

Step 2 ) For each class  $\mathcal{M}_{r\alpha}$  and each integer  $j > r$ , we will determine  $B_{j,r}(\mathcal{M}_{r\alpha})$ , which is the number of rank  $r$  configurations of size  $j$ , contained in a configuration  $T$  representing the class  $\mathcal{M}_{r\alpha}$ .

From Step 1 and Step 2, we calculate  $B_{j,r}$  for  $j > r$  by:

$$(12) \quad B_{j,r} = \sum_{\alpha \in \{a,b,c,\dots\}} |\mathcal{M}_{r\alpha}| \cdot B_{j,r}(\mathcal{M}_{r\alpha}), \quad j > r.$$

The quantity  $B_{r,r}$  is determined by the identity

$$B_{r,r} = \binom{N}{r} - \sum_{i < r} B_{r,i}.$$

We also give a systematic procedure to determine the quantities  $B_{j,r}(\mathcal{M}_{r\alpha})$  recursively with respect to  $r$ :

**Lemma 3.5.** *Let  $T$  be a  $\mathcal{M}_{r\alpha}$  configuration. For each  $i < r$ , let  $\mathfrak{C}_i(T)$  denote the collection of all  $\mathcal{C}_i$  configurations that are contained in  $T$ , and that have at least  $(r + 1)$  points. We can write  $\mathfrak{C}_i(T) = \cup_{\beta \in \{a,b,c,\dots\}} \mathcal{M}_{i\beta}(T)$  where  $\mathcal{M}_{i\beta}(T)$  denotes the class of configurations  $\mathcal{M}_{i\beta}$  that are contained in  $T$  and have at least  $r + 1$  points. We have*

$$(13) \quad B_{j,r}(\mathcal{M}_{r\alpha}) = \binom{|T|}{j} - \sum_{i < r} \sum_{\mathcal{M}_{i\beta}(T) \in \mathfrak{C}_i(T)} |\mathcal{M}_{i\beta}(T)| \cdot B_{j,i}(\mathcal{M}_{i\beta}(T)).$$

*Proof.* In order to obtain  $B_{j,r}(\mathcal{M}_{r\alpha})$ , we must subtract from  $\binom{|T|}{j}$ , the number of  $j$  point configurations  $S \subset T$  having rank at most  $r - 1$ . If  $S \subset T$  is a  $j$  point configuration of rank  $i < r$ , then by Lemma 3.4 and Lemma 3.3,  $\bar{S}$  is the unique  $\mathcal{M}_{i\beta}(T)$  configuration containing  $S$ . Thus, the second term in the right hand side of (13) exactly counts all the  $j$ -point configurations in  $T$  that have rank strictly less than  $r$ . □

In §4, we study configurations of rank at most 5 (for a general finite field  $\mathbb{F}_q$ ), and in §5, §6, §7 and §8, we study configurations of rank 6, 7, 8 and 9 respectively over the field  $\mathbb{F}_3$ . Each of the sections §4 – §8 is divided into two subsections dedicated to Step 1 and Step 2 of the above method.

#### 4. CONFIGURATIONS OF RANK AT MOST 5 FOR A GENERAL FINITE FIELD $\mathbb{F}_q$

##### 4.1. Classification of $\mathcal{C}_r$ , for $r \leq 5$ .

The only configuration of rank 1 is a single point of  $PG(3, q)$ , and the only configuration of rank 2 is a set of two distinct points of  $PG(3, q)$ . We recall that maximal rank  $r$  configurations can be built from maximal rank  $(r - 1)$  configurations by adding points. There are two ways to add a point  $P_3$  to the  $\mathcal{C}_2$  given by  $\{P_0, P_1\}$ : if  $P_2$  lies on the line  $L = \overline{P_0, P_1}$ , then by Lemma 3.3, the unique  $\mathcal{C}_3$  containing this configuration of 3 points is the line  $L$ . All such configurations form one orbit denoted  $\mathcal{M}_{3a} = G \cdot \overline{P_0, P_1}$  of size  $\binom{4}{2}_q = (1 + q^2)(1 + q + q^2)$ .

If  $P_2 \notin L$  then the resulting configuration  $S = \{P_0, P_1, P_2\}$  of 3 points in general position in a plane  $\Pi$  is also a  $\mathcal{C}_3$ : indeed we may assume  $S = \{e_0, e_1, e_2\}$  so that  $I_2(S)$  has basis

$$\{X_0X_1, X_0X_2, X_1X_2, X_0X_3, X_1X_3, X_2X_3, X_3^2\}.$$



It is easily checked that  $V(I_2(S)) = S$ . All such  $\mathcal{C}_3$ 's form a single  $G$ -orbit denoted  $\mathcal{M}_{3b} = G \cdot \{e_0, e_1, e_2\}$ , and of size  $q^3(1+q)^2(1+q^2)(1+q+q^2)/6$ . In summary:

The  $G$ -orbits in  $\mathfrak{C}_r$  for  $r \in \{1, 2, 3\}$  are as follows:

There is one orbit of  $\mathfrak{C}_1$  given by  $\mathcal{M}_{1a}$  of size  $(1+q)(1+q^2)$ . This orbit consists of points of  $PG(3, q)$ .

There is one orbit of  $\mathfrak{C}_2$  given by  $\mathcal{M}_{2a}$  of size  $\binom{(1+q)(1+q^2)}{2}$ . This orbit consists of unordered pairs of points of  $PG(3, q)$ .

There are 2 orbits in  $\mathfrak{C}_3$ :

$\mathcal{M}_{3a}$  of size  $|G(1, 3)|$ . This orbit consists of the lines of  $PG(3, q)$ .

$\mathcal{M}_{3b}$  of size  $q^3(1+q)^2(1+q^2)(1+q+q^2)/6$ . This orbit consists of unordered triples of non-collinear points of  $PG(3, q)$ .

#### 4.1.1. Classification of $\mathcal{C}_4$ 's.

We now turn to determining the set  $\mathfrak{C}_4$ . Let  $S$  be a  $\mathcal{C}_4$  containing  $\mathcal{M}_{3a}$  configuration represented by  $\overline{e_0, e_1}$ . We may assume  $S$  contains  $T = \{e_2\} \cup \overline{e_0, e_1}$ . Here  $I_2(T)$  has basis

$$\{X_0X_2, X_1X_2, X_0X_3, X_1X_3, X_2X_3, X_3^2\},$$

and  $V(I_2(T))$  is clearly equal to  $T$ . Thus,  $T$  is a  $\mathcal{C}_4$ . The set of all such configurations forms a single orbit  $\mathcal{M}_{4a} = G \cdot T$  whose elements consist of a pair  $(L, P)$  where  $L$  is a line not containing the point  $P$ . Since there are  $\binom{4}{2}_q$  ways to pick a line of  $PG(3, q)$  and  $(q^2 + q^3)$  ways to pick a point not on this line, we see that the size of  $\mathcal{M}_{4a}$  is  $\binom{4}{2}_q \cdot (q^2 + q^3)$  which simplifies to  $q^2(1+q)(1+q^2)(1+q+q^2)$ .

Next, we consider  $S$  which is a  $\mathcal{C}_4$  containing a  $\mathcal{M}_{3b}$  configuration represented by  $\{e_0, e_1, e_2\}$ . If  $S$  contains a point  $P$  which lies on one of the 3 lines  $\overline{e_i, e_j}$ , then by maximality,  $S$  contains a  $\mathcal{M}_{3a}$  configuration, which case has already been treated. We now assume  $S$  contains no other point on any line  $\overline{e_i, e_j}$ ,  $0 \leq i < j \leq 2$ . If  $S$  contains another point  $P$  of the plane  $\Pi$  spanned by  $\{e_0, e_1, e_2\}$  then we may assume  $S$  contains  $T = \{e_0, e_1, e_2, e_0 + e_1 + e_2\}$  which has rank 4. Here  $I_2(T)$  has basis

$$\{X_3^2, X_3X_2, X_3X_1, X_3X_0, X_0X_1 - X_0X_2, X_0X_1 - X_1X_2\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_4$ . All such configurations of 4 points in general position in a plane of  $PG(3, q)$  form a single orbit  $\mathcal{M}_{4c} = G \cdot T$ , because given a such a 4-tuple  $(P_1, \dots, P_4)$ , there are  $(q-1)q^3$  elements  $g \in PGL_4(q)$  with the property that  $g \cdot (e_0, e_1, e_2, e_0 + e_1 + e_2) = (P_1, P_2, P_3, P_4)$ . Thus,

$$|\mathcal{M}_{4c}| = \frac{1}{4!} \frac{|PGL_4(q)|}{q^3(q-1)} = q^3(q^4 - 1)(q^3 - 1)(q + 1)/24.$$

On the other hand, if  $S$  contains a point  $P$  not lying on the plane generated by  $\{e_0, e_1, e_2\}$  then we may assume  $S$  contains  $T = \{e_0, e_1, e_2, e_3\}$ . Here  $I_2(T)$  has basis

$$\{X_0X_1, X_0X_2, X_1X_2, X_0X_3, X_1X_3, X_2X_3\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_4$ . Given 4 ordered points  $(P_1, \dots, P_4)$  in general position in  $PG(3, q)$ , there are  $(q-1)^3$  elements  $g \in PGL_4(q)$  with the property that  $G \cdot (e_0, e_1, e_2, e_3) = (P_1, \dots, P_4)$ . Thus, all such configurations form a single orbit which we denote  $\mathcal{M}_{4b} = G \cdot \{e_0, e_1, e_2, e_3\}$  of size

$$|\mathcal{M}_{4b}| = |PGL_4(q)|/(24(q-1)^3) = q^6(1+q)^2(1+q^2)(1+q+q^2)/24.$$

We summarize this:

The  $G$ -orbits of  $\mathfrak{C}_4$  are:

- $\mathcal{M}_{4a}$  of size  $q^2(1+q)(1+q^2)(1+q+q^2)$ . This orbit consists of pairs  $(L, P)$  of a line  $L$  and a point  $P$  of  $PG(3, q)$  such that  $P \notin L$ .
- $\mathcal{M}_{4b}$  of size  $q^6(1+q+q^2+q^3)(1+q+q^2)(q+1)/24$ . This orbit consists of unordered 4-tuples of points in general position in  $PG(3, q)$ .
- $\mathcal{M}_{4c}$  of size  $q^3(q^4-1)(q^3-1)(q+1)/24$ . This orbit consists of unordered 4-tuples of points in general position in some plane of  $PG(3, q)$ .

#### 4.1.2. Classification of $\mathcal{C}_5$ 's.

Next, we determine the set  $\mathfrak{C}_5$ . Let  $S$  be a  $\mathcal{C}_5$  containing a  $\mathcal{M}_{4a}$  configuration represented by  $\{e_2\} \cup \overline{e_0, e_1}$ . If  $S$  contains another point  $P$  of the plane  $\Pi$  spanned by  $\{e_0, e_1, e_2\}$ , then the line  $L = \overline{P, e_2}$  meets the line  $\overline{e_0, e_1}$  in a third point  $Q$ . The configuration  $\{e_2, P, Q\}$  is contained in the  $\mathcal{M}_{3a}$  configuration  $L$ , and hence by Lemma 3.4,  $S$  must contain  $L$ . Thus,  $S$  contains the union of the lines  $L$  and  $\overline{e_0, e_1}$  which we denote  $T$ . We may take  $Q = e_0$ . Here  $I_2(T)$  has basis

$$\{X_1X_2, X_0X_3, X_1X_3, X_2X_3, X_3^2\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_5$ . All such configurations consisting of 2 intersecting lines in  $PG(3, q)$  form a single orbit  $G \cdot T$  which we denote  $\mathcal{M}_{5a}$ . Since  $\mathcal{M}_{5a}$  consists of a pair of lines in a plane, we see that the size of  $\mathcal{M}_{5a}$  is:

$$|\mathcal{M}_{5a}| = (1+q+q^2+q^3) \cdot \binom{1+q+q^2}{2} = q(1+q)^2(1+q^2)(1+q+q^2)/2.$$

If  $S$  contains no other point  $P$  of the plane  $\Pi$  spanned by  $\{e_0, e_1, e_2\}$ , then we may assume  $S$  contains  $T = \{e_2, e_3\} \cup \overline{e_0, e_1}$ . Here  $I_2(T)$  has basis

$$\{X_0X_2, X_0X_3, X_1X_2, X_1X_3, X_3X_2\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_5$ . The set of all configurations consisting of a line  $L$  together with 2 points on a line  $L'$  skew to  $L$  form the class  $\mathcal{M}_{5b}$  of size

$$|\mathcal{M}_{5b}| \cdot q^3/2 = q^5(1+q)(1+q^2)(1+q+q^2)/2.$$

Next, let  $S$  be a  $\mathcal{C}_5$  containing a  $\mathcal{M}_{4c}$  configuration represented by  $S_0 = \{e_0, e_1, e_2, e_0 + e_1 + e_2\}$ . Suppose  $S$  contains another point  $P$  of the plane  $\Pi$  spanned by  $\{e_0, e_1, e_2\}$ . If  $P$  lies on one of the 6 lines between the 4 points of  $S_0$ , then by maximality, the whole line is contained in  $S$ , and hence  $S$  contains a  $\mathcal{M}_{4a}$  configuration, which case has already been treated. If  $P$  does not lie on any of the 6 lines of  $S_0$  (this case is possible only if  $q > 3$ ), then  $S_0 \cup \{P\}$  are 5 points in general position in the plane  $\Pi$  (i.e., no 3 of the points are collinear). For  $q \geq 4$ , given any 5 points in general position of  $PG(2, q)$ , there exists a unique non-degenerate conic passing through these points. Moreover, all the non-degenerate conics of  $PG(2, q)$  are  $PGL_3(q)$  equivalent. Let  $T$  be the conic  $X_1^2 - X_0X_2$  in the plane  $X_3 = 0$ . Here  $I_2(T)$  has basis

$$\{X_3^2, X_3X_2, X_3X_1, X_3X_0, X_1^2 - X_0X_2\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_5$ . Let  $\mathcal{M}_{5e}$  denote this class of  $\mathcal{C}_5$ 's consisting of a non-degenerate conic in a plane of  $PG(3, q)$ . Since there are  $|PGL_3(q)|/|PGL_2(q)| = (q^5 - q^2)$  non-degenerate conics in a plane, and there are  $(1+q)(1+q^2)$  planes in  $PG(3, q)$ , we see that

$$|\mathcal{M}_{5e}| = (1+q)(1+q^2) \cdot |PGL_3(q)|/|PGL_2(q)| = (q^5 - q^2)(1+q^2)(1+q).$$

We now assume  $S$  is a  $\mathcal{C}_5$  containing a  $\mathcal{M}_{4c}$  configuration represented by  $\{e_0, e_1, e_2, e_0 + e_1 + e_2\}$ , and  $S$  contains no other point  $P$  of the plane  $\Pi$  spanned by  $e_0, e_1, e_2$ . We may assume  $S$  contains  $T = \{e_3, e_0, e_1, e_2, e_0 + e_1 + e_2\}$ . Here  $I_2(T)$  has basis

$$\{X_3X_2, X_3X_1, X_3X_0, X_0X_1 - X_0X_2, X_0X_1 - X_1X_2\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_5$ . Let  $\mathcal{M}_{5c}$  denote this class of  $\mathcal{C}_5$ 's consisting of 5 points 4 of which are in general position in a plane (there are  $|\mathcal{M}_{4c}|$  ways to pick these), and the fifth point lies outside this plane (there are  $q^3$  ways to pick this point), Since there are  $|\mathcal{M}_{4c}|$  ways to pick the 4 points, and  $q^3$  ways to pick the fifth point outside the plane, we see that

$$|\mathcal{M}_{5c}| = |\mathcal{M}_{4c}| \cdot q^3 = q^6(q^4 - 1)(q^3 - 1)(q + 1)/24.$$

It remains to consider  $S$  which is a  $\mathcal{C}_5$  containing a  $\mathcal{M}_{4b}$  configuration represented by  $\{e_0, e_1, e_2, e_3\}$ . If  $S$  contains a point  $P$  which lies on one of the 6 lines  $\overline{e_i, e_j}$ , say  $\overline{e_0, e_1}$ , then by maximality,  $S$  contains  $\mathcal{M}_{4a}$  which case has already been treated. If  $P$  does not lie on any of these 6 lines, but it lies on one of the 4 planes formed by  $\{e_0, e_1, e_2, e_3\} \setminus \{e_i\}$ ,  $i = 0, 1, 2, 3$ ,  $S$  will be a  $\mathcal{M}_{5c}$  configuration. Therefore, we assume any  $P \in S \setminus \{e_0, e_1, e_2, e_3\}$  has the property that the points of  $T = \{e_0, e_1, e_2, e_3, P\}$  are in general position. Up to  $G$ -equivalence, we may assume  $P = e_0 + e_1 + e_2 + e_3$ . Here  $I_2(T)$  has basis

$$\{X_0(X_2 - X_1), X_1(X_2 - X_0), X_0(X_3 - X_1), X_1(X_3 - X_0), X_2(X_3 - X_0)\},$$

and  $V(I_2(T)) = T$ . Thus,  $T$  is a  $\mathcal{C}_5$ . Since 5-tuples  $(P_1, \dots, P_5)$  of points in general position in  $PG(3, q)$  correspond bijectively with  $PGL_4(q)$ , we see that the size of the class  $\mathcal{M}_{5d}$  consisting of unordered 5-tuples of points in general position in  $PG(3, q)$  is

$$|\mathcal{M}_{5d}| = |PGL_4(q)|/5! = q^6(q^4 - 1)(q^3 - 1)(q^2 - 1)/120.$$

We summarize this:

The set  $\mathfrak{C}_5$  can be partitioned into the following five classes:

- $\mathcal{M}_{5a}$  of size  $q(1 + q)^2(1 + q^2)(1 + q + q^2)/2$ . This class consists of a pair of intersecting lines of  $PG(3, q)$ .
- $\mathcal{M}_{5b}$  of size  $q^5(1 + q)(1 + q^2)(1 + q + q^2)/2$ . This class consists of a line  $L$  and two points on a line  $L'$  skew to  $L$ .
- $\mathcal{M}_{5c}$  of size  $q^6(q^4 - 1)(q^3 - 1)(q + 1)/24$ . This class consists of four points in general position in a plane, together with a point outside the plane.
- $\mathcal{M}_{5d}$  of size  $q^6(q^4 - 1)(q^3 - 1)(q^2 - 1)/120$ . This class consists of 5 points of  $PG(3, q)$  in general position.
- $\mathcal{M}_{5e}$  of size  $(q^5 - q^2)(1 + q^2)(1 + q)$ . This class consists of a non-degenerate conic in a plane of  $PG(3, q)$ .

#### 4.2. Calculations of $B_{j,i}$ for $j > i$ .

Since a maximal configuration of rank 1 and 2 contains exactly 1 and 2 points of  $PG(3, q)$  respectively. Therefore, in these cases we can see that  $B_{j,i} = 0$  for  $i \in \{1, 2\}$  and  $j > i$ .

##### 4.2.1. $B_{j,3}$ for $j > 3$ .

Let  $T$  be a  $\mathcal{M}_{3a}$  configuration, i.e., a line in  $PG(3, q)$ . Then  $T$  has  $(1 + q)$  points of  $PG(3, q)$ . Since  $j > 3$  and any  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 2$  has at most 2 points. By (13), we have

$$B_{j,3}(\mathcal{M}_{3a}) = \binom{1+q}{j}.$$

A  $\mathcal{M}_{3b}$  configuration has 3 points, so  $B_{j,3}(\mathcal{M}_{3b}) = 0$ . Therefore from (12), we have

$$(14) \quad B_{j,3} = |\mathcal{M}_{3a}| \cdot \binom{1+q}{j} = (1+q^2)(1+q+q^2) \cdot \binom{1+q}{j}.$$

4.2.2.  $B_{j,4}$  for  $j > 4$ .

An  $\mathcal{M}_{4a}$  configuration  $T$  represented by a pair  $(L, P)$  of a line  $L$  and a point  $P$  of  $PG(3, q)$ , has size  $(q+2)$  points. Thus:

$$\mathfrak{C}_1(T) = \mathfrak{C}_2(T) = \emptyset, \quad \mathfrak{C}_3(T) = \mathcal{M}_{3a}(T).$$

Since there is only one line in  $T$ , we have  $|\mathcal{M}_{3a}(T)| = 1$ . By (13), we have,

$$B_{j,4}(\mathcal{M}_{4a}) = \binom{q+2}{j} - \binom{1+q}{j}.$$

Since the  $\mathcal{M}_{4c}$  and  $\mathcal{M}_{4b}$  configurations have exactly 4 points of  $PG(3, q)$  and  $j > 4$ , we see that  $B_{j,4}(\mathcal{M}_{4b}) = B_{j,4}(\mathcal{M}_{4c}) = 0$ . Therefore for  $j > 4$ , by (12), we have

$$(15) \quad B_{j,4} = q^2(1+q)(1+q^2)(1+q+q^2) \cdot \left( \binom{q+2}{j} - \binom{1+q}{j} \right).$$

4.2.3.  $B_{j,5}$  for  $j > 5$ .

A  $\mathcal{M}_{5a}$  configuration  $T$  represented by a pair of intersecting lines of  $PG(3, q)$  contains  $(2q+1)$  points. Thus,

$$\mathfrak{C}_1(T) = \mathfrak{C}_2(T) = \emptyset, \quad \mathfrak{C}_3(T) = \mathcal{M}_{3a}(T), \quad \mathfrak{C}_4(T) = \mathcal{M}_{4a}(T).$$

We also note that  $|\mathcal{M}_{3a}(T)| = 2$  and  $|\mathcal{M}_{4a}(T)| = 2q$ . By (13), we have,

$$B_{j,5}(\mathcal{M}_{5a}) = \binom{2q+1}{j} - 2q \cdot \left( \binom{q+2}{j} - \binom{1+q}{j} \right) - 2 \binom{1+q}{j} = \binom{2q+1}{j} - 2q \cdot \binom{q+2}{j} + 2(q-1) \cdot \binom{1+q}{j}.$$

. A  $\mathcal{M}_{5b}$  configuration  $T$  represented by a line  $L$  and two points  $\{P_1, P_2\}$  on a line  $L'$  skew to  $L$  contains  $(q+3)$  points of  $PG(3, q)$ . Again,

$$\mathfrak{C}_1(T) = \mathfrak{C}_2(T) = \emptyset, \quad \mathfrak{C}_3(T) = \mathcal{M}_{3a}(T), \quad \mathfrak{C}_4(T) = \mathcal{M}_{4a}(T).$$

We also note that  $|\mathcal{M}_{3a}(T)| = 1$  and  $|\mathcal{M}_{4a}(T)| = 2$ . By (13), we have

$$B_{j,5}(\mathcal{M}_{5b}) = \binom{q+3}{j} - 2 \cdot \left( \binom{2+q}{j} - \binom{1+q}{j} \right) - \binom{1+q}{j} = \binom{q+3}{j} - 2 \cdot \binom{2+q}{j} + \binom{1+q}{j}.$$

. A  $\mathcal{M}_{5c}$  configuration  $T$  represented by the  $(q+1)$  points of a non-degenerate conic in a plane of  $PG(3, q)$ . Here

$$\mathfrak{C}_1(T) = \mathfrak{C}_2(T) = \mathfrak{C}_3(T) = \mathfrak{C}_4(T) = \emptyset,$$

because  $T$  does not contain a line which rules out  $\mathcal{M}_{3a}, \mathcal{M}_{4a}$  and configurations in  $\mathfrak{C}_i(T)$  must have at least 6 points which rules out  $\mathcal{M}_{3b}, \mathcal{M}_{4b}, \mathcal{M}_{4c}$ . By (13), we have

$$B_{j,5}(\mathcal{M}_{5c}) = \binom{1+q}{j}.$$

Since each of the  $\mathcal{M}_{5d}$  or  $\mathcal{M}_{5e}$  configurations contains 5 points  $PG(3, 3)$ , we have for  $j > 5$ ,

$$B_{j,5}(\mathcal{M}_{5d}) = B_{j,5}(\mathcal{M}_{5e}) = 0.$$

Therefore for  $j > 5$ , we have

$$(16) \quad \begin{aligned} B_{j,5} = & \frac{1}{2}q(1+q)^2(1+q^2)(1+q+q^2) \cdot \left( \binom{2q+1}{j} - 2q \cdot \binom{q+2}{j} + 2(q-1) \cdot \binom{1+q}{j} \right) \\ & + \frac{1}{2}q^5(1+q)(1+q^2)(1+q+q^2) \cdot \left( \binom{q+3}{j} - 2 \cdot \binom{2+q}{j} + \binom{1+q}{j} \right) \\ & + (q^5 - q^2)(1+q^2)(1+q) \cdot \binom{1+q}{j}. \end{aligned}$$

## 5. CONFIGURATIONS OF RANK 6

For the rest of this paper, we assume  $q = 3$ . The size of the group  $G = PGL_4(q)$  is

$$|G| = 12130560 = 2^8 \cdot 3^6 \cdot 5 \cdot 13.$$

Since an irreducible conic in a projective plane has only 4 points when  $q = 3$ . We see that the class of configuration  $\mathcal{M}_{5e}$  in  $\mathfrak{C}_5$  does not occur when  $q = 3$ . Therefore, the set  $\mathfrak{C}_5$  is partitioned into the following four classes when  $q = 3$ :

$\mathcal{M}_{5a}$  of size  $2^4 \cdot 3 \cdot 5 \cdot 13$ . This class consists of a pair of intersecting lines of  $PG(3, 3)$ .

$\mathcal{M}_{5b}$  of size  $2^2 \cdot 3^5 \cdot 5 \cdot 13$ . This class consists of a line  $L$  and two points on a line  $L'$  skew to  $L$ .

$\mathcal{M}_{5c}$  of size  $2^4 \cdot 3^5 \cdot 5 \cdot 13$ . This class consists of four points in general position in a plane, together with a point outside the plane.

$\mathcal{M}_{5d}$  of size  $2^5 \cdot 3^5 \cdot 13$ . This class consists of 5 points of  $PG(3, 3)$  in general position.

As before, we build the  $\mathcal{C}_6$ 's by adding points to  $\mathcal{C}_5$ 's. We carry this out systematically by running through the classes in  $\mathfrak{C}_5$ .

**Lemma 5.1.** *The set  $\mathfrak{C}_6$  can be partitioned into the following 9 classes:*

$\mathcal{M}_{6a}$  of size  $2^3 \cdot 5$ . It consists of planes of  $PG(3, 3)$ .

$\mathcal{M}_{6b}$  of size  $2^4 \cdot 3^4 \cdot 5 \cdot 13$ . It consists of configurations  $(\{L_1, L_2\}, \{P\})$  where  $\{L_1, L_2\}$  is a pair of intersecting lines, and  $P$  does not lie on the plane generated by the two lines.

$\mathcal{M}_{6c}$  of size  $3^4 \cdot 5 \cdot 13$ . It consists of a pair of skew lines in  $PG(3, 3)$ .

$\mathcal{M}_{6d}$  of size  $2^6 \cdot 3^4 \cdot 5 \cdot 13$ . It consists of configurations  $(L, \{P_1, \dots, P_4\})$  where  $\{P_1, \dots, P_4\}$  are 4 points in general position in a plane  $\Pi$  and the line  $L$  intersects  $\Pi$  in one of these 4 points.

$\mathcal{M}_{6e}$  of size  $2^4 \cdot 3^5 \cdot 5 \cdot 13$ . It consists of configurations  $\{P_1, \dots, P_6\}$  where the 3 lines  $L_i = \overline{P_{2i-1}, P_{2i}}$  are non-coplanar lines meeting at a single point different from  $\{P_1, \dots, P_6\}$ .

$\mathcal{M}_{6f}$  of size  $2^4 \cdot 3^6 \cdot 5 \cdot 13$ , The elements of  $\mathcal{M}_{6f}$  are configurations  $\{P_1, \dots, P_6\}$  where  $P_1, P_2$  are on a line  $L$ , and  $P_3, P_4$  are on a line  $L'$ , and  $P_5, P_6$  are on a line  $L''$  with  $\{L', L''\}$  being a pair of skew lines which meet  $L$  at distinct points  $P, Q \notin \{P_1, P_2\}$ .

*Proof.* Let  $S$  be a  $\mathfrak{C}_6$  configuration.

**Classes  $\mathcal{M}_{6a}, \mathcal{M}_{6b}$ .**

We begin with  $\mathcal{C}_6$  configurations  $S$  containing a  $\mathcal{M}_{5a}$  configuration represented by the pair of lines  $X_1X_2 = 0$  in the plane  $X_3 = 0$ . If  $S$  contains another point  $P$  of the plane  $X_3 = 0$ , then by maximality of  $\mathcal{M}_{5a}$ , the configuration  $\{P\} \cup \mathcal{M}_{5a}$  has rank 6. We have already shown that a plane of  $PG(3, 3)$  is  $\mathcal{C}_6$ , therefore,  $S$  is represented by the plane  $X_3 = 0$ . Let  $\mathcal{M}_{6a}$  denote the class of configurations consisting of planes of  $PG(3, 3)$ . The size of this class is 40.

If  $S$  contains no other point of the plane  $X_3 = 0$ , then we may assume  $S$  contains  $T = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \{e_3\}$ . Here  $I_2(T)$  has basis  $\{X_1X_2, X_0X_3, X_1X_3, X_2X_3\}$  and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_6$ . Let  $\mathcal{M}_{6b}$  denote the class of configurations consisting of a pair of intersecting lines (which

can be chosen in  $|\mathcal{M}_{5a}|$  ways) together with a point not on the plane generated by the pair of lines (which can be chosen in  $3^3$  ways). The size of this class is

$$|\mathcal{M}_{6b}| = |\mathcal{M}_{5a}| \cdot 3^3 = 3^4(1+3)(1+3+3^2)(1+3+3^2+3^3)/2 = 84240.$$

**Classes  $\mathcal{M}_{6c}, \mathcal{M}_{6d}$ .**

Next, we consider  $\mathcal{C}_6$  configurations  $S$  which contain the  $\mathcal{M}_{5b}$  configuration represented by  $S_0 = \overline{e_0, e_1} \cup \{e_2, e_3\}$ . Let  $L = \overline{e_0, e_1}$  and let  $L' = \overline{e_2, e_3}$ . If  $P \in S$  lies on the line  $L' = \overline{e_2, e_3}$ , then by maximality,  $S_0 \cup \{P\}$  has rank 6, and is contained in the configuration  $T = L \cup L' = \overline{e_0, e_1} \cup \overline{e_2, e_3}$ . Here  $I_2(T)$  has basis  $\{X_0X_2, X_0X_3, X_1X_2, X_1X_3\}$  and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_6$ . Let  $\mathcal{M}_{6c}$  denote this class consisting of a pair of skew lines. This class has size

$$|\mathcal{M}_{6c}| = 130 \cdot 3^4/2 = 5265.$$

We now assume  $S \supset S_0$  does not contain any other point of  $L \cup L'$ . Given a point  $P$  of  $S \setminus S_0$ , if one of the lines  $\overline{P, e_2}$  or  $\overline{P, e_3}$  meets the line  $L = \overline{e_0, e_1}$ , then by maximality,  $S$  contains a  $\mathcal{M}_{5a}$  configuration, which has already been treated. Thus, the three points  $P, e_2, e_3$  are non-collinear and  $L$  does not intersect any of the 3 lines generated by  $P, e_2, e_3$ . Up to  $G$ -equivalence, we may assume  $T = \{P\} \cup S_0$  is represented by  $\{e_0, e_1, e_2\} \cup \overline{e_3, e_0 + e_1 + e_2}$ . Here  $I_2(T)$  has basis

$$\{X_3(X_2 - X_0), X_3(X_1 - X_0), X_0(X_1 - X_2), X_1(X_0 - X_2)\},$$

and we verify that  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_6$ . Let  $\mathcal{M}_{6d}$  be the class consisting of 4 points in general position in a plane  $\Pi$  (which can be chosen in  $|\mathcal{M}_{4c}|$  ways), together with a line which intersects  $\Pi$  in one of these 4 points (which can be chosen in  $4 \cdot 3^2$  ways). Thus, the size of this class is

$$|\mathcal{M}_{6d}| = 4 \cdot 3^2 \cdot |\mathcal{M}_{4c}| = 3^5(3^4 - 1)(3^3 - 1)(3 + 1)/6 = 336960.$$

**Classes  $\mathcal{M}_{6e}, \mathcal{M}_{6f}$ .**

Next we consider  $\mathcal{C}_6$  configurations  $S$  which contain the  $\mathcal{M}_{5c}$  configuration  $S_0 = \{e_3, e_0, e_1, e_2, e_0 + e_1 + e_2\}$ . If  $S$  contains a point  $P$  of the plane  $X_3 = 0$ , then by maximality,  $S$  contains a  $\mathcal{M}_{5a}$ , which case has already been treated. So we assume  $S \setminus S_0$  contains no other point of the plane  $X_3 = 0$ . For  $P \in S \setminus S_0$ , let  $T = \{P, e_3, e_0, e_1, e_2, e_0 + e_1 + e_2\}$ .

Let  $Q$  be the intersection point of the line  $\overline{e_3, P}$  with the plane  $X_3 = 0$ . Consider the 6 lines  $L_1, \dots, L_6$  in the plane  $X_3 = 0$  generated by  $\{P_1, \dots, P_4\} = \{e_0, e_1, e_2, e_0 + e_1 + e_2\}$ . Since  $q = 3$ , the union of these 6 lines is the whole plane  $X_3 = 0$ . If a point  $R$  lies on 3 of these lines then  $R \in \{P_1, \dots, P_4\}$ . There are three points  $R_1 = e_0 + e_1, R_2 = e_1 + e_2, R_3 = e_2 + e_0$  which lie on exactly two of these 6 lines. There are three possibilities for the point  $Q$  above: i)  $Q \in \{e_0, e_1, e_2, e_0 + e_1 + e_2\}$ , ii)  $Q \in \{R_1, R_2, R_3\}$ , iii)  $Q$  lies on exactly one of the lines  $L_1, \dots, L_6$ . We consider each of these possibilities one by one.

i) If  $Q \in \{P_1, \dots, P_4\}$ , then by maximality,  $S$  contains a  $\mathcal{M}_{5b}$  configuration, which case has already been treated.

ii) If  $Q \in \{R_1, R_2, R_3\}$  say  $Q = e_0 + e_1$ , then we may assume  $P = \overline{e_0 + e_1 + e_3}$  (by replacing  $e_3$  with  $ce_3$  for a suitable scalar  $c$ ). We note that each of the three lines  $\overline{Q, e_0}, \overline{Q, e_2}, \overline{Q, e_3}$  through  $Q$  contain exactly 2 points of  $T$ , namely  $\{e_0, e_1\}, \{e_2, e_0 + e_1 + e_2\}, \{e_3, e_0 + e_1 + e_3\}$ . Thus,  $T$  is

$G$ -equivalent to the configuration  $\{e_0, e_0 + e_3, e_1, e_1 + e_3, e_2, e_2 + e_3\}$ , with  $e_3$  representing the point of concurrence of the 3 lines  $\overline{e_3, e_i}$  for  $i = 0, 1, 2$ . Here  $I_2(T)$  has basis

$$\{X_0X_1, X_1X_2, X_0X_2, X_3(X_0 + X_1 + X_2 - X_3)\},$$

and clearly  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_6$ . Let  $\mathcal{M}_{6e}$  be this class of configurations consisting of 2 points each on three non-coplanar lines through a point  $P$ . The configuration  $\mathcal{M}_{6e}$  is obtained by i) picking a point  $P$  and three non-coplanar lines through  $P$ , and ii) picking 2 points on each of these 3 lines different from  $P$ . The size of this class is

$$|\mathcal{M}_{6e}| = 40 \cdot \frac{1}{6}(1 + 3 + 3^2)(1 + 3)3^3 \cdot \binom{3}{2}^3 = 252720.$$

For later use, we note that the stabilizer group in  $G$  of the  $\mathcal{M}_{6e}$  configuration  $T$  is group of order 48 which is the wreath product  $Z_2 \wr S_3$  where  $S_3$  permutes the three lines through  $e_3$  (generated by  $\begin{pmatrix} P & 0 \\ 0 & 1 \end{pmatrix}$  where  $P$  is  $3 \times 3$  permutation matrix) and  $Z_2$  interchanges the 2 points on a line, (for example

generated by  $g_0 = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix}$  for the line through  $\overline{e_0, e_3}$ ). In more detail, if  $g_0, g_1, g_2$  are involutions interchanging the points  $\{e_i, e_i + e_3\}$  for  $i = 0, 1, 2$  respectively, then  $g_0, g_1, g_2$  commute with each other and hence generate the group  $Z_2 \times Z_2 \times Z_2$ . The group  $S_3$  above acts on  $Z_2 \times Z_2 \times Z_2$  by permutation action on the three factors. As a corollary, we note that given a point  $Q = ae_0 + be_1 + ce_2 + de_3$  with  $abc \neq 0$ , there exist automorphisms of the  $\mathcal{M}_{6e}$  configuration that allow us to independently change the sign of  $a, b$  and  $c$  (the coordinate  $d$  is not necessarily fixed). Thus, up to  $G$ -equivalence we can assume  $Q = e_0 + e_1 + e_2 + d'e_3$  for some  $d' \in \mathbb{F}_3$ .

iii) If  $Q$  is on exactly one of the lines  $L_1, \dots, L_6$ , say  $\overline{e_0, e_1}$  then  $Q \notin \{R_1, R_2, R_3\}$  and hence  $Q \neq e_0 + e_1$ . Thus,  $P = e_0 + \lambda e_1 + e_3$  for some  $\lambda \in \mathbb{F}_3 \setminus \{0, 1\}$ , i.e.  $\lambda = -1$ . Thus,

$$T = \{e_0, e_1, e_2, e_0 + e_1 + e_2, e_3, e_3 + e_0 - e_1\}.$$

Here  $I_2(T)$  has basis

$$\{X_3X_2, X_3(X_1 + X_0), X_0(X_1 - X_2 + X_3), X_2(X_0 - X_1)\},$$

and it is readily checked that  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_6$ . We will denote this class configuration by  $\mathcal{M}_{6f}$ . We note that 6 points of  $T$  generate 14 planes of which only 2 of them have 4 points: these are the planes  $X_2 = 0$  and  $X_3 = 0$ . Thus, the class  $\mathcal{M}_{6f}$  consists of configurations  $G$ -equivalent to  $T$ . An element of  $\mathcal{M}_{6f}$  can be constructed by i) choosing 2 points  $P_1, P_2$  on the line of intersection of two planes  $\Pi_1, \Pi_2$  ii) 2 points  $P_3, P_4$  on the plane  $\Pi_1$  and iii) 2 points  $P_5, P_6$  with the property that if  $Q_1 = \overline{P_3, P_4} \cap L$  and  $Q_2 = \overline{P_5, P_6} \cap L$  then the points  $P_1, P_2, Q_1, Q_2$  are distinct points of  $L$ . The pair of planes can be chosen in  $\binom{40}{2}$  ways. The points  $P_1, P_2$  can be chosen in  $\binom{3+1}{2}$  ways, and the points  $Q_1, Q_2$  can be chosen in  $(3-1)(3-2)$  ways. The points  $P_3, P_4$  and  $P_5, P_6$  can be chosen in  $(3\binom{3}{2})^2$  ways. Thus, the class  $\mathcal{M}_{6f}$  has size

$$|\mathcal{M}_{6f}| = \binom{40}{2} \cdot \binom{3+1}{2} \cdot (3-1)(3-2) \cdot (3\binom{3}{2})^2 = 758160.$$

For later use, we record the stabilizer group in  $G$  of the  $\mathcal{M}_{6f}$  configuration  $T$ . Consider the following four involutions which preserve  $T$ :

$$g_0 = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}, g_1 = \begin{pmatrix} 1 & -1 & & \\ & -1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}, g_2 = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}, g_3 = \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & -1 \end{pmatrix}.$$

We note that i)  $g_0$  interchanges  $\{P_1, P_2\}$ , ii)  $g_1$  interchanges  $\{P_1, P_2\}, \{P_3, P_4\}$ , iii)  $g_2$  interchanges  $\{P_1, P_2\}, \{P_5, P_6\}$ , and iv)  $g_3$  interchanges  $\{P_3, P_5\}$  and  $\{P_4, P_6\}$ . The matrices  $g_1, g_2$  commute and

generate the group  $Z_2 \times Z_2$ . The matrix  $g_3$  and satisfies  $g_3 g_1 g_3 = g_2$ ,  $g_3 g_2 g_1 = g_2$ . Thus, the group generated by  $g_1, g_2, g_3$  is isomorphic to the dihedral group. The element  $g_0$  commutes with  $g_2, g_3, g_4$ . Thus, the stabilizer group of  $T$  is isomorphic to the direct product  $D_4 \times Z_2$

$$Z_2 \times D_4 : \langle g_0, g_1, g_2, g_3 \mid g_0^2, g_1^2, g_2^2, g_3^2, (g_1 g_2)^2, (g_0 g_1)^2, (g_0 g_2)^2, (g_0 g_3)^2, g_3 g_1 g_3 g_2 \rangle.$$

Finally, we come to the case when  $S$  contains a  $\mathcal{M}_{5d}$  represented by 5 points  $\{P_1, \dots, P_5\}$  in general position in  $PG(3, 3)$ . We may take  $\{P_1, \dots, P_5\} = \{e_0, e_1, e_2, e_3, e_0 + e_1 + e_2 + e_3\}$ . If  $P$  is another point of  $S$ , let  $T = \{P, P_1, \dots, P_5\}$ . Since in  $PG(3, 3)$  there is no collection of 6 points in general position, there are only two possibilities: i) some 3 points of  $T$  are collinear, or ii) no 3 points of  $T$  are collinear but some 4 points of  $T$  are coplanar. In case (i), we may assume  $P$  lies on the line  $\overline{P_4 P_5}$ . In this case by maximality,  $S$  contains a  $\mathcal{M}_{6d}$  configuration, and in case (ii),  $S$  contains a  $\mathcal{M}_{6f}$  configuration. These cases have already been treated above.  $\square$

### 5.1. $B_{j,6}$ for $j > 6$ .

. A  $\mathcal{M}_{6a}$  configuration  $T$  represented by a plane  $\Pi$  contains 13 points of  $PG(3, 3)$ . Since  $j \geq 7$ , we have

$$\mathfrak{C}_1(T) = \dots = \mathfrak{C}_4(T) = \emptyset, \quad \mathfrak{C}_5(T) = \mathcal{M}_{5a}(T).$$

We note that  $|\mathcal{M}_{5a}(T)| = \binom{13}{2} = 78$ . By (13), we have,

$$B_{j,6}(\mathcal{M}_{6a}) = \binom{13}{j} - 78 \binom{7}{j}.$$

. A  $\mathcal{M}_{6b}$  configuration  $T$  of 8 points of  $PG(3, 3)$  represented by a pair of intersecting lines  $L_1, L_2$  and point  $P$  outside the plane  $\langle L_1, L_2 \rangle$ . Here, again:

$$\mathfrak{C}_1(T) = \dots = \mathfrak{C}_4(T) = \emptyset, \quad \mathfrak{C}_5(T) = \mathcal{M}_{5a}(T),$$

and  $|\mathcal{M}_{5a}(T)| = 1$ . By (13), we have,

$$B_{j,6}(\mathcal{M}_{6b}) = \binom{8}{j} - \binom{7}{j}.$$

. A  $\mathcal{M}_{6c}$  configuration  $T$  of 8 points of  $PG(3, 3)$  represented by a pair of skew lines  $L_1, L_2$ . Here  $\mathfrak{C}_i(T) = \emptyset$  for all  $1 \leq i \leq 5$ . By (13), we have,

$$B_{j,6}(\mathcal{M}_{6c}) = \binom{8}{j}.$$

. A  $\mathcal{M}_{6d}$  configuration  $T$  represented by  $(L, \{P_1, \dots, P_4\})$  where  $\{P_1, \dots, P_4\}$  are 4 points in general position in a plane  $\Pi$ , and the line  $L$  intersects  $\Pi$  at  $P_4$ . The configuration  $T$  contains 7 points of  $PG(3, 3)$ . Here again,  $\mathfrak{C}_i(T) = \emptyset$  for all  $1 \leq i \leq 5$ , and hence by (13), we have,

$$B_{j,6}(\mathcal{M}_{6d}) = \binom{7}{j}.$$

We have,  $B_{j,6}(\mathcal{M}_{6e}) = B_{j,6}(\mathcal{M}_{6f}) = 0$  as each of  $\mathcal{M}_{6e}$  or  $\mathcal{M}_{6f}$  configurations contain exactly 6 points of  $PG(3, 3)$ . Therefore, by (12), for  $j > 6$ , we have

$$B_{j,6} = 40 \cdot \left( \binom{13}{j} - 78 \binom{7}{j} \right) + 84240 \cdot \left( \binom{8}{j} - \binom{7}{j} \right) + 5265 \cdot \binom{8}{j} + 336960 \cdot \binom{7}{j},$$

i.e.

$$(17) \quad B_{j,6} = 40 \cdot \binom{13}{j} + 89505 \cdot \binom{8}{j} + 249600 \cdot \binom{7}{j}.$$



6. CONFIGURATIONS OF RANK 7

As before, we build the  $\mathcal{C}_7$ 's by adding points to  $\mathcal{C}_6$ 's. We carry this out systematically by running through the classes in  $\mathfrak{C}_6$ .

**Lemma 6.1.** *The set  $\mathfrak{C}_7$  can be partitioned into the following 7 classes:*

- $\mathcal{M}_{7a}$  of size  $2^3 \cdot 3^3 \cdot 5$ . This class consists of a plane of  $PG(3, 3)$  and a point outside that plane.
- $\mathcal{M}_{7b}$  of size  $2^4 \cdot 3^2 \cdot 5 \cdot 13$ . This class consists of 3 concurrent non coplanar lines of  $PG(3, 3)$ .
- $\mathcal{M}_{7c}$  of size  $2^4 \cdot 3^4 \cdot 5 \cdot 13$ . This class consists of three lines  $L_1, L_2, L_3$  of  $PG(3, 3)$ , where  $L_2$  and  $L_3$  are skew to each other and the line  $L_1$  meets both the lines.
- $\mathcal{M}_{7d}$  of size  $2^5 \cdot 3^5 \cdot 5 \cdot 13$ . This class consists of configurations  $(\{L_1, L_2\}, \{P_1, P_2\})$ , where  $\{L_1, L_2\}$  is a pair of intersecting lines and  $P_1, P_2$  lie outside the plane  $\langle L_1, L_2 \rangle$ . Also, the line  $\overline{P_1, P_2}$  does not meet  $L_1$  and  $L_2$ .
- $\mathcal{M}_{7e}$  of size  $2^5 \cdot 3^6 \cdot 5 \cdot 13$ . This class consists of configurations  $(\{P_1, P_2, P_3, P_4\}, L)$ , where the points  $P_1, P_2, P_3, P_4$  are in general position in  $PG(3, 3)$  and the line  $L$  does not meet any of the 6 lines  $\overline{P_i, P_j}$ , where  $1 \leq i < j \leq 4$ .
- $\mathcal{M}_{7f}$  of size  $2^6 \cdot 3^5 \cdot 5 \cdot 13$ . This class consists of configurations  $\{P_1, \dots, P_6, P_7\}$  where the 3 lines  $L_i = \overline{P_{2i-1}, P_{2i}}$ ,  $i = 1, 2, 3$ , are non-coplanar lines meeting at a single point  $Q$  different from  $\{P_1, \dots, P_6\}$ . Let  $L_4$  be a line through  $Q$  such that no three of the four lines  $\{L_1, L_2, L_3, L_4\}$  are coplanar. Of the three points  $\{Q_1, Q_2, Q_3\}$  on  $L_4$  different from  $Q$ , there is only one point  $Q_1$  which lies on two of the planes spanned by three points of  $\{P_1, \dots, P_6\}$ , whereas the other two points  $\{Q_2, Q_3\}$  lie on three such planes. We take  $P_7 = Q_1$ .
- $\mathcal{M}_{7g}$  of size  $2^2 \cdot 3^5 \cdot 5 \cdot 13$ . This class consists of configurations  $\{P_1, \dots, P_8\}$  where the four lines  $L_i = \overline{P_{2i-1}, P_{2i}}$ ,  $i = 1, 2, 3, 4$ , meet at a single point  $Q$  different from  $\{P_1, \dots, P_8\}$  and no three of these four lines are coplanar. If we arbitrarily pick  $\{P_{2i-1}, P_{2i}\}$  on  $L_i$  for  $i = 1, \dots, 3$ , then the points  $\{P_7, P_8\}$  on  $L_4$  are the pair of points  $\{Q_2, Q_3\}$  in the notation of the description of  $\mathcal{M}_{7f}$  configuration above.

*Proof.* Let  $S$  be a  $\mathcal{C}_7$  configuration.

**Class  $\mathcal{M}_{7a}$ .**

If  $S$  contains a  $\mathcal{M}_{6a}$  configuration represented by the plane  $\Pi$  given by  $X_3 = 0$ , then we may assume  $S$  contains  $T = \{e_3\} \cup \Pi$ . Here  $I_2(T)$  has basis  $\{X_3X_0, X_3X_1, X_3X_2\}$  and  $V(I_2(T))$  clearly equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_7$  which we denote  $\mathcal{M}_{7a}$ . The size of  $\mathcal{M}_{7a}$  is  $3^3 \cdot |\mathcal{M}_{6a}| = 1080$ .

**Classes  $\mathcal{M}_{7b}, \mathcal{M}_{7c}, \mathcal{M}_{7d}$ .**

If  $S$  contains a  $\mathcal{M}_{6b}$  configuration represented by  $S_0 = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \{e_3\}$ , and  $P$  is another point of  $S$ , there are 4 possibilities: i)  $P$  lies on the plane  $\Pi : X_3 = 0$ , ii)  $P$  lies outside  $\Pi$ , and the line  $\overline{P, e_3}$  meets  $\Pi$  at  $e_0$ , iii)  $P$  lies outside  $\Pi$ , and the line  $\overline{P, e_3}$  meets  $\Pi$  at a point  $Q$  of  $S_0$  different from  $e_0$ , iv)  $P$  lies outside  $\Pi$ , and the line  $\overline{P, e_3}$  meets  $\Pi$  at a point  $Q$  not lying on  $S_0$ .

In case i) by maximality,  $S$  contains a  $\mathcal{M}_{6a}$  configuration which case has been treated above.

In case ii) by maximality,  $S$  contains the configuration  $T$  of three concurrent lines  $\overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_0, e_3}$ . Here  $I_2(T)$  has basis  $\{X_3X_2, X_3X_1, X_1X_2\}$  and  $V(I_2(T))$  clearly equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_7$  configuration. We denote the class of configurations  $G$ -equivalent to  $T$  by  $\mathcal{M}_{7b}$ . The size of  $\mathcal{M}_{7b}$  is  $|\mathcal{M}_{6e}| / \binom{3}{2}^3$ , i.e. 9360.

In case iii) by maximality of  $S$ , we may assume  $S$  contains the configuration  $T$  of the three lines  $L_1 \cup L_2 \cup L_3$  where  $L_1 = \overline{e_0, e_1}$ ,  $L_2 = \overline{e_0, e_2}$  and  $L_3 = \overline{e_1, e_3}$ . Here  $I_2(T)$  has basis  $\{X_3X_0, X_3X_2, X_1X_2\}$  and  $V(I_2(T))$  clearly equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_7$ . We denote the class of configurations  $G$ -equivalent to  $T$  by  $\mathcal{M}_{7c}$ . The lines  $L_2$  and  $L_3$  are skew to each other and the line  $L_1$  meets both the lines. We can choose two skew lines  $L_2$  and  $L_3$  in  $PG(3, 3)$  in  $130 \cdot 3^4/2$  ways and pick the line  $L_1$  in  $(3+1)^2$  ways. Thus, the size of the class  $\mathcal{M}_{7c}$  is  $65 \cdot 3^4 \cdot 4^2$ , i.e. 84240.

In case iv) we may assume  $S$  contains the configuration  $T$  consisting of the lines  $L_1 = \overline{e_0, e_1}$  and  $L_2 = \overline{e_0, e_2}$  in the plane  $X_3 = 0$ , and points  $\{e_3, e_3 + e_1 + e_2\}$ . Here  $I_2(T)$  has basis  $\{X_1(X_3 - X_2), X_2(X_3 - X_1), X_3X_0\}$  and  $V(I_2(T))$  clearly equals  $T$ . We denote the class of configurations  $G$ -equivalent to  $T$  by  $\mathcal{M}_{7d}$ . The number of ways to choose  $T$  is the number of ways to pick the configuration  $\mathcal{M}_{6b}$  multiplied by the number of ways to pick the point  $P$  which is  $(3-1)(3^2-3)$  (there are  $(3^2-3)$  ways to pick the point  $Q$  and  $(3-1)$  ways to pick the point  $P$  on the line  $\overline{e_3, Q}$ ). Thus,  $|\mathcal{M}_{7d}| = |\mathcal{M}_{6b}| \cdot \frac{3(3-1)^2}{2}$ , i.e. 505440.

In case  $S$  contains the  $\mathcal{M}_{6c}$  configuration represented  $L_1 = \overline{e_0, e_1}$  and  $L_2 = \overline{e_2, e_3}$ . Given any other point  $P$  of  $S$ , we may assume  $P = e_0 + e_1 + e_2 + e_3$ . In this case the line  $L_3 = \overline{e_0 + e_1, e_2 + e_3}$  also contains  $P$ , and hence by maximality,  $S$  contains  $L_1 \cup L_2 \cup L_3$  which is the configuration  $\mathcal{M}_{7c}$  discussed above.

### Class $\mathcal{M}_{7e}$ .

Next we consider the case when  $S$  contains the  $\mathcal{M}_{6d}$  configuration represented by  $S_0 = \{P_1, \dots, P_4\} \cup L$  where  $\{P_1, \dots, P_4\}$  are in general position in a plane  $\Pi$  and the line  $L$  intersects  $\Pi$  in  $P_4$ . Let  $P$  be another point of  $S$  and let  $T = S_0 \cup \{P\}$ . If  $P \in \Pi$ , then by maximality,  $S$  contains a  $\mathcal{M}_{6b}$  configuration, which cases have been treated above. Similarly, if the plane  $\Pi'$  generated by  $P$  and  $L$  contains a point of  $\{P_1, P_2, P_3\}$ , then by maximality,  $S$  contains a  $\mathcal{M}_{6b}$  configuration, which case has been treated above. So we assume  $\Pi'$  does not contain  $P_1, P_2$  or  $P_3$ . In this case  $\{P_1, P_2, P_3, P\}$  is a  $\mathcal{M}_{4b}$  configuration which we may represent as  $\{e_0, e_1, e_2, e_3\}$ . The line  $L$  does not meet any of the 6 lines  $\overline{e_i, e_j}$ , where  $0 \leq i < j \leq 3$ . We may assume  $L = \overline{e_0 + e_1 + e_2, e_0 + c'e_1 + e_3}$  where  $c' \notin \{0, 1\}$ , i.e.  $c' = -1$ . Here  $I_2(T)$  has basis

$$\{X_3(X_1 + X_0 + X_2), X_0(X_1 - X_2 + X_3), X_1(X_0 - X_2) + X_3(X_2 + X_0)\},$$

and  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a  $\mathcal{C}_7$ . We denote, by  $\mathcal{M}_{7e}$ , the class of configurations which are  $G$ -equivalent to  $T$ . There are  $|\mathcal{M}_{4b}|$  ways to choose the  $\mathcal{M}_{4b}$  configuration represented by  $\{e_0, e_1, e_2, e_3\}$ . Let  $\Pi_i$  be the plane spanned by  $\{e_0, e_1, e_2, e_3\} \setminus \{e_i\}$  and let  $Q_i = \Pi_i \cap L$ ,  $i = 0, 1, 2, 3$ . Writing  $Q_3 = e_0 + ae_1 + be_2$  and  $Q_2 = e_0 + ce_1 + de_3$  the condition for the line  $L = \overline{Q_1, Q_2}$  to not meet any of the 6 lines  $\overline{e_i, e_j}$  is  $abcd \neq 0$  and  $a \neq c$ . Thus, there are  $(3-1)^3(3-2)$  ways to pick  $L$ . Thus,  $|\mathcal{M}_{7e}| = 3^6(3^4-1)(3^3-1)(3^2-1)(3-2)/24$ , i.e. 1516320.

### Classes $\mathcal{M}_{7f}$ and $\mathcal{M}_{7g}$ .

Next, we consider the case when  $S$  contains the  $\mathcal{M}_{6e}$  configuration represented by  $S_0 = \{e_0, e_0 + e_3, e_1, e_1 + e_3, e_2, e_2 + e_3\}$ . Let  $P_{2i+1} = e_i$  and  $P_{2i+2} = e_i + e_3$  for  $i \in \{0, 1, 2\}$ . Let  $P_7$  be another point of  $S$  and let  $T = \{P, P_1, \dots, P_6\}$ . There are 3 possibilities for  $P$ : (i)  $P_7$  lies on any of the lines  $\overline{P_{2i+1}, P_{2i+2}}$ ,  $i \in \{0, 1, 2\}$ , (ii)  $P_7$  is contained in any of the 3 Planes  $\Pi_0 : \langle e_0, e_1, e_3 \rangle$ ,  $\Pi_1 : \langle e_0, e_2, e_3 \rangle$  and  $\Pi_2 : \langle e_1, e_2, e_3 \rangle$ , (iii)  $P_7$  lies in the complement of  $\Pi_0 \cup \Pi_1 \cup \Pi_2$ .

In case (i) and (ii), by maximality,  $S$  contains  $\mathcal{M}_{6b}$ , which cases have already been treated.

In case (iii), we have  $P_7 = ae_0 + be_1 + ce_2 + de_3$  with  $abc \neq 0$ . Here  $I_2(T)$  has basis

$$\{X_1(aX_2 - cX_0), X_0(bX_2 - cX_1), bX_3(X_0 + X_1 + X_2 - X_3) - d(a + b + c - d)X_0X_1\}.$$

A direct calculation shows that

$$\bar{T} = V(I_2(T)) = T \cup \{P'\}, \quad \text{where } P' = ae_0 + be_1 + ce_2 + (a + b + c - d)e_3.$$

Thus,  $T$  is a  $\mathcal{C}_7$  if and only if  $2d = a + b + c$ . Thus, there are 4 choices for  $P_7$ , namely  $\{(1, b, c, 1 + b + c - d) : b, c \in \{\pm 1\}\}$ . We denote the class of configuration  $G$ -equivalent to  $T$  by  $\mathcal{M}_{7f}$ . In this case, we claim that the  $\mathcal{M}_{6e}$  sub-configuration of  $T = \mathcal{M}_{7f}$  is unique, in other words removing a point  $Q$  from  $T$  results in a  $\mathcal{M}_{6e}$  configuration if and only if  $Q = P_7$ : We note that the 6 points  $P_1, \dots, P_6$  represented by  $e_i + \epsilon_i e_3$  where  $i \in \{0, 1, 2\}$  and  $\epsilon_i \in \{0, 1\}$  generate 11 planes of  $PG(3, q)$ . Of these 11 planes there are 3 planes which contain 4 of  $\{P_1, \dots, P_6\}$ , these planes denoted  $\Pi^i$  are generated by  $\{e_3, e_0, e_1, e_2\} \setminus \{e_i\}$  for  $i = 0, 1, 2$ . The remaining 8 planes are denoted  $\Pi_\epsilon$  where  $\epsilon = (\epsilon_0, \epsilon_1, \epsilon_2) \in \{0, 1\}^3$ , and are generated by  $\{e_0 + \epsilon_0 e_3, e_1 + \epsilon_1 e_3, e_2 + \epsilon_2 e_3\}$ . We recall from the discussion about the automorphism group of the  $\mathcal{M}_{6e}$  configuration, that we may assume  $P_7 = e_0 + e_1 + e_2 + de_3$  for some  $d \in \mathbb{F}_3$ . Since  $2d = 1 + 1 + 1 = 0$ , we must have  $P_7 = e_0 + e_1 + e_2$ . Clearly, of the 11 planes above,  $P_7$  is contained only in the planes  $\Pi_\epsilon$  for  $\epsilon = (0, 0, 0)$  and  $\epsilon = (1, 1, 1)$ . Thus,  $P_7$  lies on exactly two planes of the form  $\Pi_\epsilon$  and  $\Pi_\epsilon$ . Therefore, none of the 6 configurations  $\{P_1, \dots, P_7\} \setminus P_i$  for  $i \in \{1, \dots, 6\}$  are  $\mathcal{M}_{6e}$  configurations, Thus, proving that  $\{P_1, \dots, P_6\}$  is the unique  $\mathcal{M}_{6e}$  configuration in  $T$ . Therefore, the number of  $\mathcal{M}_{7f}$  configurations is  $|\mathcal{M}_{6e}| \cdot 4$ , i.e. 1010880. We also note that the automorphism group of  $\mathcal{M}_{7f}$  configuration. We recall that the automorphism group of the  $\mathcal{M}_{6e}$  configuration is  $(Z_2 \times Z_2 \times Z_2) \rtimes S_3$ , where the  $S_3$  factor permutes the lines  $\overline{e_i, e_3}$  for  $i \in \{0, 1, 2\}$  and the factor  $Z_2 \times Z_2 \times Z_2$  has generators  $g_0, g_1, g_2$  where  $g_i$  interchanges the points  $\{e_i, e_i + e_3\}$  for  $i = 0, 1, 2$  respectively. Since  $P_7 = e_0 + e_1 + e_2$ , the automorphism group of the  $\mathcal{M}_{7f}$  configuration is  $Z_2 \times S_3$ , where the  $Z_2$  factor is generated by the involution  $g_0 g_1 g_2 \in Z_2 \times Z_2 \times Z_2$ .

We now consider the case

$$\bar{T} = V(I_2(T)) = T \cup \{P_8\}, \quad \text{where } P_8 = ae_0 + be_1 + ce_2 + (a + b + c - d)e_3.$$

The number of such pairs  $\{P_7, P_8\}$  is 4. We denote the class of configuration  $G$ -equivalent to  $\bar{T}$  by  $\mathcal{M}_{7g}$ . As above, we may assume  $P, P'$  are given by  $e_0 + e_1 + e_2 \pm e_3$ . Thus,  $P_7$  is on the planes  $\Pi_\epsilon$  for  $\epsilon = (1, 0, 0), (0, 1, 0), (0, 0, 1)$  and similarly  $P_8$  is on the planes  $\Pi_\epsilon$  for  $\epsilon = (0, 1, 1), (1, 0, 1), (1, 1, 0)$ . In other words,  $P_7$  is on the planes generated by  $\{P_2, P_3, P_5\}$ ,  $\{P_1, P_4, P_5\}$ , and  $\{P_1, P_3, P_6\}$ , and similarly  $P_8$  is on the planes generated by  $\{P_1, P_4, P_6\}$ ,  $\{P_2, P_3, P_6\}$ , and  $\{P_2, P_4, P_5\}$ . Thus, a list of planes containing 4 points of the configuration  $T'$  is as below. We use the notation  $(ijkl)$  to mean that  $P_i, P_j, P_k, P_l$  are coplanar:

$$(1234), (1256), (3456), (1278), (5678), (3478)(2357), (1457), (1367), (1468), (2368), (2458).$$

Of the  $\binom{8}{2} = 28$  configurations  $T_{ij} = \{P_1, \dots, P_8\} \setminus \{P_i, P_j\}$  obtained from  $T'$  by deleting 2 of the points, we see that there are 16 configurations of type  $\mathcal{M}_{6e}$ :

$$T_{12}, T_{14}, T_{16}, T_{17}, T_{23}, T_{25}, T_{28}, T_{34}, T_{36}, T_{37}, T_{45}, T_{48}, T_{56}, T_{57}, T_{68}, T_{78},$$

and there are 12 configurations of type  $\mathcal{M}_{6f}$ :

$$T_{13}, T_{15}, T_{18}, T_{24}, T_{26}, T_{27}, T_{35}, T_{38}, T_{46}, T_{47}, T_{58}, T_{67}.$$

While there are 4 ways (choices of  $\{P_7, P_8\}$ ) to extend a given  $\mathcal{M}_{6e}$  configuration to a  $\mathcal{M}_{7g}$  configuration, we see that there are 16 choices for the base  $\mathcal{M}_{6e}$  configuration that yield the same  $\mathcal{M}_{7e}$

configuration. Therefore, the number of  $\mathcal{M}_{7g}$  configurations is given by  $\frac{|\mathcal{M}_{6e}| \cdot 4}{16}$ , i.e. 63180.

Next, we consider the case when  $S$  contains the  $\mathcal{M}_{6f}$  configuration represented by

$$S_0 = \{P_1 = e_0, P_2 = e_1, P_3 = e_2, P_4 = e_0 + e_1 + e_2, P_5 = e_3, P_6 = e_3 + e_0 - e_1\}.$$

We use the label  $(ijk)$  for the plane generated by  $P_i, P_j, P_k$ . If  $P_l$  also lies on this plane, the label  $(ijkl)$  will also denote the same plane. The 6 points of  $S_0$  generate 14 planes, of which only the planes (1234) and (1256) contain 4 points of  $S_0$ . These are the only planes containing both  $P_1$  and  $P_2$ . Of the remaining 12 planes, there are 8 of them of the form  $(ijk)$  for  $i \in \{1, 2\}, j \in \{3, 4\}, k \in \{5, 6\}$ , and there are 4 planes (345), (346), (356), (456).

We can write  $PG(3, 3)$  as the union of the 4 planes containing the line  $\overline{P_1, P_2}$ . These are the planes (1234), (1256),  $\Pi_1, \Pi_{-1}$ , where  $\Pi_1$  is the plane generated by  $P_1, P_2, e_2 + e_3$  and  $\Pi_{-1}$  is the plane generated by  $P_1, P_2, e_2 - e_3$ . Let  $P_7$  be another point of  $S$  and let  $T = \{P_7, P_1, \dots, P_6\}$ . There are 3 possibilities for  $P_7$ : (i) some 3 points of  $T$  are collinear, (ii) some 5 points of  $T$  are coplanar, (iii) no 3 points of  $T$  are collinear and no 5 points of  $T$  are coplanar.

In case (i) and (ii), by maximality,  $S$  contains a  $\mathcal{M}_{6b}$  configuration, which cases which have already been treated. In case (iii) we represent  $P_7 = ae_0 + be_1 + ce_2 + de_3$ . The condition that no 5 points of  $T$  are coplanar, implies that  $P_7$  is not on the planes (1234) and (1256), i.e.  $cd \neq 0$ . Therefore, we take  $d = 1$  and  $c \in \{\pm 1\}$ . We recall that the automorphism group of the  $\mathcal{M}_{6f}$  configuration is isomorphic to the group  $Z_2 \times D_4$ . The element  $g_0$  generating the  $Z_2$  factor interchanges the planes  $\Pi_1$  and  $\Pi_{-1}$ , so that we may assume  $c = 1$ . We parametrize the points of the affine plane  $\pi_1 := \Pi_1 \setminus \overline{P_1 P_2}$  as  $xe_0 + ye_1 + (e_2 + e_3)$ .

The 12 planes  $(ijk)$  (not containing both  $P_1, P_2$ ) meet  $\pi_1$  in the twelve lines:

$$\begin{aligned} L_{345} : y = x, \quad L_{346} : y = -x, \quad L_{356} : y = x + 1, \quad L_{456} : y = -x - 1. \\ L_{135} = L_{146} : y = 0, \quad L_{145} : y = 1, \quad L_{136} : y = -1. \\ L_{235} : x = 0, \quad L_{245} = L_{236} : x = 1, \quad L_{246} : x = -1. \end{aligned}$$

The  $D_4$  factor generated by  $g_1, g_2, g_3$  preserves  $\Pi_1$  and the 13 points of  $\Pi_1$  form orbits under this group. One of these orbits consists of the 4 points of  $\overline{P_1, P_2}$ . The remaining 9 points are the points of the affine plane  $\pi_1$ . The four lines  $L_{345}, L_{346}, L_{356}, L_{456}$  in  $\pi_1$  form a pair of parallel lines of slope 1 and another pair of parallel lines of slope  $-1$ , which we refer to as a  $\#$  pattern. These 4 lines account for 8 of the 9 points of  $\pi_1$ . The remaining point is  $(x, y) = (0, 1)$  which forms a single  $D_4$ -orbit. The four points of intersection of the pattern  $\#$  are:  $(x, y) \in \{(0, 0), (1, 1), (1, -1), (-1, 0)\}$  and they form the second orbit. Finally, the 4 points of the  $\#$  pattern which lie on exactly one of the four lines form the third orbit:  $(x, y) \in \{(0, 1), (-1, -1), (0, -1), (-1, 1)\}$ . We summarize the three  $D_4$  orbits on  $\pi_1$ :

- (1) the four points with coordinates  $(x, y) \in \{(0, 0), (1, 1), (1, -1), (-1, 0)\}$  which lie on one of the 4 lines  $\overline{P_i P_j}$  for  $i \in \{3, 4\}, j \in \{5, 6\}$ ,
- (2) the four points with coordinates  $(x, y) \in \{(0, 1), (-1, -1), (0, -1), (-1, 1)\}$ , which lie on the intersection of three planes
  - (a) one of which is in the 4 planes (345), (346), (356), (456)
  - (b) one of which is in the 4 planes (135), (145), (136), (146)
  - (c) one of which is in the 4 planes (235), (245), (236), (246)
- (3) the point with coordinates  $(x, y) = (1, 0)$  which is on the intersection of four planes
  - (a) none of which is in the 4 planes (345), (346), (356), (456)

- (b) two of which are in the 4 planes (135), (145), (136), (146)
- (c) two of which are in the 4 planes (235), (245), (236), (246).

The condition that no 3 points of  $T$  are collinear implies that  $P_7$  lies in  $\pi_1$  and is either in the second or the third orbit above. Thus, we may assume  $P_7$  to be represented by  $(x, y) = (0, 1)$  or  $(x, y) = (1, 0)$ . In the former case,  $P_7$  lies on the planes (356), (145), (235). In this case, the configuration  $T \setminus \{P_6\} = \{P_1, \dots, P_5, P_7\}$  is a  $\mathcal{M}_{6e}$  configuration because it consists of the three non-coplanar lines  $\overline{P_1P_4}, \overline{P_2P_3}, \overline{P_5P_7}$  which are concurrent through the point  $e_1 + e_2$ . Since we have already accounted for the  $\mathcal{C}_7$  configurations arising from  $\mathcal{M}_{6e}$ , we skip this case. Therefore, we are left with  $P_7$  represented by  $e_0 + e_2 + e_3$ .

Here  $I_2(T)$  has basis

$$\{-X_3(X_2 + X_1 + X_0), X_2(X_0 - X_1 - X_3), X_0(X_2 - X_3 - X_1)\}.$$

Next we calculate  $\bar{T} = V(I_2(T))$ . If  $Q = (x, y, z, w) \in \bar{T}$  and  $zw = 0$ , then we just get  $Q \in S_0$ . So we take  $Q = (x, y, z, 1)$  with  $z \in \{\pm 1\}$ . A direct calculation shows that i) if  $z = 1$  then  $Q = P_7$  and ii) if  $z = -1$  then the coordinates  $(x, y, -1, 1)$  of  $Q$  satisfy the conditions  $(x, y) = (0, -1)$ , i.e.,  $Q = P_8 = -e_1 - e_2 + e_3 = g_0P_7$ . The planes containing 4 points of the configuration  $\{P_1, \dots, P_8\}$  are

$$(1234), (1256), (1357), (1467), (2457), (2367), (1458), (1368), (2358), (2468), (3478), (5678).$$

Of the  $\binom{8}{2} = 28$  configurations  $T_{ij} = \{P_1, \dots, P_8\} \setminus \{P_i, P_j\}$  we have 16  $\mathcal{M}_{6e}$  configurations:

$$T_{13}, T_{14}, T_{15}, T_{16}, T_{23}, T_{24}, T_{25}, T_{26}, T_{37}, T_{38}, T_{47}, T_{48}, T_{57}, T_{58}, T_{67}, T_{68},$$

and 12  $\mathcal{M}_{6f}$  configurations:

$$T_{12}, T_{17}, T_{18}, T_{27}, T_{28}, T_{34}, T_{35}, T_{36}, T_{45}, T_{46}, T_{56}, T_{78}.$$

Thus,  $T'$  extends a  $\mathcal{M}_{6e}$  configuration, which has already been treated above (namely  $\mathcal{M}_{7g}$  configuration). □

### 6.1. $B_{j,7}$ for $j > 7$ .

. An  $\mathcal{M}_{7a}$  configuration  $T$  represented by a plane  $\Pi$  and a point  $P$  outside  $\Pi$ , contains 14 points of  $PG(3, 3)$ . Since  $j > 7$ , for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 6$ , the only possible cases are  $S = \mathcal{M}_{6a}, \mathcal{M}_{6b}$ , and  $|\mathcal{M}_{6a}(T)| = 1$ ,  $|\mathcal{M}_{6b}(T)| = \binom{13}{2} = 78$ . By (13), we have,

$$B_{j,7}(\mathcal{M}_{7a}) = \binom{14}{j} - \binom{13}{j} - 78 \binom{8}{j} = \binom{13}{j-1} - 78 \binom{8}{j}.$$

. An  $\mathcal{M}_{7b}$  configuration  $T$  represented by three concurrent lines not contained in a plane contains 10 points of  $PG(3, 3)$ . Since  $j > 7$ , for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 6$ , the only possible case is  $S = \mathcal{M}_{6b}$ , and  $|\mathcal{M}_{6b}(T)| = \binom{3}{2} \cdot 3 = 9$  (as we have  $\binom{3}{2}$  choices for two lines and 3 choices for a point from the remaining line). By (13), we have,

$$B_{j,7}(\mathcal{M}_{7b}) = \binom{10}{j} - 9 \binom{8}{j}.$$

. An  $\mathcal{M}_{7c}$  configuration  $T$  is represented by three lines  $L_1, L_2, L_3$  of  $PG(3, 3)$ , where  $L_2$  and  $L_3$  are skew to each other and the line  $L_1$  meets both the lines. It contains 10 points of  $PG(3, 3)$ . Since  $j > 7$ , the only possible cases for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 6$  are  $S = \mathcal{M}_{6b}, \mathcal{M}_{6c}$ . As there are 2 pairs of intersecting lines ( $\{L_1, L_2\}$  and  $\{L_2, L_3\}$ ) in  $T$ , and a point can be chosen from the remaining line in 3 ways, we have  $|\mathcal{M}_{6b}(T)| = 6$ . It is clear that  $|\mathcal{M}_{6c}(T)| = 1$ , as there is only one configuration of a pair of skew lines and which is  $\{L_1, L_3\}$ . By (13), we have,

$$B_{j,7}(\mathcal{M}_{7c}) = \binom{10}{j} - 6 \cdot \binom{8}{j} - \binom{8}{j} = \binom{10}{j} - 7 \cdot \binom{8}{j}.$$

. A  $\mathcal{M}_{7d}$  configuration  $T$  represented by  $(\{L_1, L_2\}, \{P_1, P_2\})$ , where  $\{L_1, L_2\}$  is a pair of intersecting lines and the line  $\overline{P_1, P_2}$  not contained in the plane does not meet  $L_1$  and  $L_2$ , contains 9 points of  $PG(3, 3)$ . Since  $j > 7$ , the only possible case for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 6$  is  $S = \mathcal{M}_{6b}$ . Clearly,  $|\mathcal{M}_{6b}(T)| = 2$ , as there is one pair of intersecting lines and 2 choices for a point from  $\{P_1, P_2\}$ . By (13), we have,

$$B_{j,7}(\mathcal{M}_{7d}) = \binom{9}{j} - 2 \cdot \binom{8}{j}.$$

. Let  $T$  be a  $\mathcal{M}_{7e}$  configuration represented by  $(\{P_1, P_2, P_3, P_4\}, L)$ , where the points  $P_1, P_2, P_3, P_4$  are in general position in  $PG(3, 3)$  and the line  $L$  does not meet any of the 6 lines  $\overline{P_i, P_j}$ , where  $1 \leq i < j \leq 4$ . The configuration  $T$  has 8 points of  $PG(3, 3)$ . In this case,  $\mathfrak{C}_i(T) = \emptyset$  for  $i \leq 6$ . By (13), we have

$$B_{8,7}(\mathcal{M}_{7e}) = 1.$$

. Let  $T$  be a  $\mathcal{M}_{7g}$  configuration represented by  $\{P_1, \dots, P_6, P_7, P_8\}$  where the 4 lines  $L_i = \overline{P_{2i-1}, P_{2i}}$ ,  $i = 1, 2, 3, 4$ , meet at a single point  $Q$  different from  $\{P_1, \dots, P_6, P_7, P_8\}$  and no three of them are contained in a plane. The configuration  $T$  has 8 points of  $PG(3, 3)$ . Since  $j > 7$ , we have no such  $S$  with  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 6$ . By (13), we have,

$$B_{8,7}(\mathcal{M}_{7g}) = 1.$$

Since a  $\mathcal{M}_{7f}$  configuration contains exactly 7 points  $PG(3, 3)$  and  $j > 7$ , we get  $B_{j,7}(\mathcal{M}_{7f}) = 0$ . Therefore, by (12), for  $j > 7$ , we have

$$\begin{aligned} B_{8,7} &= 1080 \cdot \left( \binom{13}{7} - 78 \binom{8}{8} \right) + 9360 \cdot \left( \binom{10}{8} - 9 \binom{8}{8} \right) \\ &\quad + 84240 \cdot \left( \binom{10}{8} - 7 \cdot \binom{8}{8} \right) + 505440 \cdot \binom{9}{8} - 2 \cdot \binom{8}{8} + 1516320 + 63180, \end{aligned}$$

and, if  $j \geq 9$ ,

$$B_{j,7} = 1080 \cdot \binom{13}{j-1} + 9360 \cdot \binom{10}{j} + 84240 \cdot \binom{10}{j} + 505440 \cdot \binom{9}{j},$$

i.e.

$$(18) \quad B_{8,7} = 9413820, \text{ and } j \geq 9, B_{j,7} = 1080 \cdot \binom{13}{j-1} + 93600 \cdot \binom{10}{j} + 505440 \cdot \binom{9}{j}.$$

## 7. CONFIGURATIONS OF RANK 8

As before, we build the  $\mathcal{C}_8$ 's by adding points to  $\mathcal{C}_7$ 's. We carry this out systematically by running through the classes in  $\mathfrak{C}_7$ . If  $T$  is a  $\mathcal{C}_8$ , then  $I_2(T)$  is a 2 dimensional vectors space, and its projectivization is a pencil of quadratic forms on  $PG(3, 3)$ . If  $S$  denote the set of  $(q+1) = 4$  elements of this pencil, let  $\nu_i(T) = |S \cap \mathcal{O}_i|$ , here  $\mathcal{O}_1, \dots, \mathcal{O}_6$  are the  $G$ -orbits of quadratic forms on  $PG(3, 3)$ , listed in Table 1. We will write down the numbers  $\nu(T) = (\nu_1(T), \dots, \nu_6(T))$ .

**Lemma 7.1.** *The set  $\mathcal{C}_8$  can be partitioned into the following 4 classes:*

$\mathcal{M}_{8a}$  of size  $2^3 \cdot 3^2 \cdot 5 \cdot 13$ . *This class consists of a plane of  $PG(3, 3)$  and a line not contained in that plane.*

$\mathcal{M}_{8b}$  of size  $2^4 \cdot 3^2 \cdot 5 \cdot 13$ . *This class consists of 4 concurrent lines such that no three are contained in a plane.*

$\mathcal{M}_{8c}$  of size  $2^2 \cdot 3^6 \cdot 5 \cdot 13$ . *This class consists of 4 lines  $L_1, L_2, L_3, L_4$  such that  $L_i$  and  $L_j$  intersect if and only if  $i \neq j \pmod{2}$  (and  $L_i$  and  $L_j$  are skew if and only if  $i = j \pmod{2}$ ).*

$\mathcal{M}_{8d}$  of size  $2^6 \cdot 3^4 \cdot 5 \cdot 13$ . *This class consists of  $(\{L_1, L_2\}, \{P_1, P_2, P_3\})$  where  $\{L_1, L_2\}$  is a pair of lines in a plane  $\pi_1$ , and the points  $P_1, P_2, P_3$  are in a different plane  $\pi_2$ . The point  $Q = L_1 \cap L_2$  lies on the line  $L = \pi_1 \cap \pi_2$ , but none of the points  $P_1, P_2, P_3$  lie on  $L$ . The four points  $\{Q, P_1, P_2, P_3\}$  are in general position in  $\pi_2$ .*

$\mathcal{M}_{8e}$  of size  $2^3 \cdot 3^6 \cdot 5 \cdot 13$ . *This class consists of configurations  $\{P_1, \dots, P_6, P_7, P_8\}$  where the 3 lines  $L_i = \overline{P_{2i-1}, P_{2i}}$ ,  $i = 1, 2, 3$ , are non-coplanar lines meeting at a single point  $Q$  different from  $\{P_1, \dots, P_6\}$ . There are four lines  $L$  passing through  $Q$  with the property that no three of the four lines  $\{L_1, L_2, L_3, L\}$  are coplanar. Let  $L_4, L_5$  be two such lines. On each of these lines  $\{L_4, L_5\}$  there is a unique point  $\{Q_1(L_4), Q_1(L_5)\}$  as given in the description of the  $\mathcal{M}_{7f}$  configuration (see Lemma 6.1). We take  $\{P_7, P_8\} = \{Q_1(L_4), Q_1(L_5)\}$ .*

*Proof.* Let  $S$  be a  $\mathcal{C}_8$  configuration.

**Class  $\mathcal{M}_{8a}$ .**

Let  $S$  contain a  $\mathcal{M}_{7a}$  configuration represented by the point set  $T = \{e_3\} \cup \overline{\Pi}$ , where the plane  $\Pi$  given by  $X_3 = 0$ . If  $S$  contains a point  $P \notin T$ , then  $P$  will be on a line  $\overline{e_3, Q}$ , where  $Q$  is a point of  $\Pi$ . Then by maximality,  $\overline{e_3, P}$  is contained in  $S$ . We can assume  $P = e_0 + e_1 + e_3$  and let  $T' = \overline{e_3, e_0 + e_1 + e_3} \cup \overline{\Pi}$ . Here  $I_2(T')$  has basis  $\{X_3(X_0 - X_1), X_3X_2\}$  and  $V(I_2(T'))$  clearly equals  $T'$ . Thus,  $T'$  is a  $\mathcal{C}_8$  which we denote  $\mathcal{M}_{8a}$ . Clearly, all the  $(q + 1) = 4$  elements of the pencil  $\mathbb{P}(I_2(T'))$  are in  $\mathcal{O}_1$  (pairs of planes). Thus,  $\nu(T') = (4, 0, 0, 0, 0, 0)$ . The number of ways to choose the plane  $\Pi$  is 40 and a line not contained in  $\Pi$  is  $13 \cdot 3^2$ . Therefore, the size of  $\mathcal{M}_{8a}$  is  $40 \cdot 13 \cdot 3^2 = 4680$ .

**Class  $\mathcal{M}_{8b}$ .**

Let  $S$  contain a  $\mathcal{M}_{7b}$  configuration represented by  $S_0 = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_0, e_3}$ . Let  $P$  be another point of  $S$ . There are two possibilities for  $P$ : (i)  $P$  lies on one of the planes  $\Pi_1 : X_1 = 0$ ,  $\Pi_2 : X_2 = 0$ ,  $\Pi_3 : X_3 = 0$ , and (ii)  $P$  lies outside  $\Pi_1, \Pi_2$  and  $\Pi_3$ .

In case (i) by maximality,  $S$  contains a  $\mathcal{M}_{7a}$  configuration which case has been treated above.

In case (ii), if  $P = ae_0 + be_1 + ce_2 + de_3$ , then the condition that  $P$  does not lie on the planes  $\Pi_1, \Pi_2$  and  $\Pi_3$  is that  $bcd \neq 0$ . So we can assume  $d = 1$  and  $P = ae_0 + be_1 + ce_2 + e_3$ . Let  $T' = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_0, e_3} \cup \{P\}$ . Here  $I_2(T')$  has basis  $\{X_3(cX_1 - bX_2), X_2(X_1 - bX_3)\}$  and  $V(I_2(T'))$  equals  $T = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_0, e_3} \cup \overline{e_0, P}$ . Thus,  $T$  is a  $\mathcal{C}_8$  which we denote  $\mathcal{M}_{8b}$ . In the pencil  $\mathbb{P}(I_2(T))$ , apart from the mentioned generators, the remaining  $(q - 1) = 2$  elements are in  $\mathcal{O}_4$ . Thus,  $\nu(T) = (2, 0, 0, 2, 0, 0)$ .

The number of ways a  $\mathcal{M}_{7b}$  configuration  $S_0$  can be chosen is  $|\mathcal{M}_{7b}|$ . Since no three lines of a  $\mathcal{M}_{8b}$  configuration are contained in a plane, there are 4 choices for the line  $\overline{e_0, P}$ . The configuration

$T \setminus \overline{e_0, e_i}$ ,  $i = 1, 2, 3$  is  $\mathcal{M}_{7b}$ . Thus, there are 4 different  $\mathcal{M}_{7b}$  configurations contained in a  $\mathcal{M}_{8b}$  configuration. Therefore, the size of  $\mathcal{M}_{8b}$  is  $\frac{|\mathcal{M}_{7b}| \cdot 4}{4} = |\mathcal{M}_{7b}| = 9360$ .

### Class $\mathcal{M}_{8c}$ .

Let  $S$  contain a  $\mathcal{M}_{7c}$  configuration represented by  $S_0 = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_1, e_3}$ . Let  $P$  be another point of  $S$ . There are two possibilities for  $P$ : (i)  $P$  lies on one of the planes  $\Pi_1 : X_3 = 0$ ,  $\Pi_2 : X_2 = 0$ , and (ii)  $P$  lies outside  $\Pi_1, \Pi_2$ . In case (i) by maximality,  $S$  contains a  $\mathcal{M}_{7a}$  configuration which case has been treated above. In case (ii), if  $P = ae_0 + be_1 + ce_2 + de_3$ , then the condition that  $P$  does not lie on the planes  $\Pi_1, \Pi_2$  is that  $cd \neq 0$ . Switching to the basis  $f_0 = e_0, f_1 = e_1, f_2 = ce_2 + ae_0, f_3 = de_3 + be_1$  of  $PG(3, 3)$ , the configuration  $S_0$  is again given by  $S_0 = \overline{f_0, f_1} \cup \overline{f_0, f_2} \cup \overline{f_1, f_3}$ , and  $P = f_2 + f_3$ . Thus, we may take  $T' = S_0 \cup \{P\}$  to be  $T' = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_1, e_3} \cup \{e_2 + e_3\}$ . Here  $I_2(T')$  has basis  $\{X_3X_0, X_2X_1\}$  and  $V(I_2(T'))$  equals  $T = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \overline{e_1, e_3} \cup \overline{e_2, e_3}$ . Thus,  $T$  is a  $\mathcal{C}_8$  which we denote  $\mathcal{M}_{8c}$ . In the pencil  $\mathbb{P}(I_2(T))$ , apart from the mentioned generators, the remaining  $(q-1) = 2$  elements are in  $\mathcal{O}_2$ . Thus,  $\nu(T) = (2, 2, 0, 0, 0, 0)$ . The size of  $\mathcal{M}_{8c}$  is  $3 \cdot |\mathcal{M}_{4b}| = 189540$ , because there are 3 ways to obtain the configuration  $T$  from four points in general position of  $PG(3, 3)$ .

### Class $\mathcal{M}_{8d}$ .

Let  $S$  contain a  $\mathcal{M}_{7d}$  configuration represented by  $S_0 = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \{e_3, e_3 + e_1 + e_2\}$ . If  $P$  is another point of  $S$ , then there are 4 possibilities for  $P$ : (i)  $P$  lies in the plane  $\Pi_0 : \langle e_0, e_1, e_2 \rangle$ , (ii)  $P$  lies in at least one of the planes  $\Pi_1 : \langle e_3, e_0, e_1 \rangle$ ,  $\Pi_2 : \langle e_3, e_0, e_2 \rangle$ ,  $\Pi_3 : \langle e_1 + e_2 + e_3, e_0, e_1 \rangle$ ,  $\Pi_4 : \langle e_1 + e_2 + e_3, e_0, e_2 \rangle$ , (iii)  $P$  lies on the line  $L : \overline{e_3, e_1 + e_2 + e_3}$ , (iv)  $P$  lies in the complement of  $\Pi_0 \cup \Pi_1 \cup \Pi_2 \cup \Pi_3 \cup \Pi_4 \cup L$  in  $PG(3, 3)$ .

In the cases (i) and (iii) by maximality,  $S$  contains a  $\mathcal{M}_{7a}$  configuration, which case has been treated above.

In the case (ii) by maximality,  $S$  contains a  $\mathcal{M}_{7b}$  or  $\mathcal{M}_{7c}$  configuration, which case has been treated above.

In case (iv), let  $P = ae_0 + be_1 + ce_2 + de_3$ . The condition for  $P \notin \Pi_0$  is that  $d \neq 0$ . So we can assume  $d = 1$ . The conditions for  $P \notin \Pi_1, \Pi_2, \Pi_3, \Pi_4$  are  $c \neq 0, 1$  and  $b \neq 0, 1$ . Therefore  $b = c = -1$ . The condition for  $P$  to lie on the line  $L$  is  $a = b - c = 0$ . Thus, we may take  $P$  to be  $ae_0 - e_1 - e_2 + e_3$  with  $a \in \{\pm 1\}$ . Let  $T' = \overline{e_0, e_1} \cup \overline{e_0, e_2} \cup \{e_3, e_3 + e_1 + e_2, P\}$ . Here  $I_2(T')$  has basis

$$\{aX_1(X_3 - X_2) - X_0X_3, X_3X_1 + X_2X_1 + X_3X_2\},$$

and  $V(I_2(T'))$  equals  $T'$ . Thus,  $T'$  is a  $\mathcal{C}_8$  which we denote  $\mathcal{M}_{8d}$ . Of the  $(q+1) = 4$  points of the pencil  $\mathbb{P}(I_2(T'))$ , except for the element  $X_3X_1 + X_2X_1 + X_3X_2$  which is in  $\mathcal{O}_4$ , the remaining  $q = 3$  elements are in  $\mathcal{O}_2$ . Thus,  $\nu(T') = (0, 3, 0, 1, 0, 0)$ .

We will now count the size of  $\mathcal{M}_{8d}$ . This configuration consists of a pair of intersecting lines  $L_1, L_2$  in a plane  $\Pi$ , and three non-collinear points  $P_1, P_2, P_3$  lying outside  $\Pi$ , such that the 3 lines formed by  $P_1, P_2, P_3$  do not meet  $L_1, L_2$ . Since  $q = 3$ , with this condition, the plane  $\langle P_1, P_2, P_3 \rangle$  meets  $\Pi$  in a line  $M$  through  $L_1 \cap L_2$ . A pair of intersecting lines  $L_1, L_2$  in  $PG(3, 3)$  can be chosen in  $\mathcal{M}_{5a}$  ways, the line  $M$  can be chosen in  $\binom{2}{1}$  ways in the plane  $\langle L_1, L_2 \rangle$ , a plane  $\Pi'$  through  $M$  other than  $\Pi$  can be chosen in  $\binom{3}{2}$  ways and on  $\Pi'$  a set of 3 noncollinear points with the property no two are lying on a line through  $L_1 \cap L_2$ , can be chosen in  $(3^3 - 3^2)$  ways. Hence, the size of  $\mathcal{M}_{8d}$  is  $|\mathcal{M}_{5a}| \cdot \binom{2}{1} \cdot \binom{3}{2} \cdot (3^3 - 3^2)$ , i.e. 336960.

Let  $S$  contain a  $\mathcal{M}_{7e}$  configuration represented by  $S_0 = \{e_0, e_1, e_2, e_3\} \cup \overline{e_0 + e_1 + e_2, e_0 - e_1 + e_3}$ . Let us denote  $P_1 = e_0 + e_1 + e_2$  and  $P_2 = e_0 - e_1 + e_3$ . All the points of  $PG(3, q)$  are on the 4



planes  $\Pi_0 : \langle P_1, P_2, e_0 \rangle$ ,  $\Pi_1 : \langle P_1, P_2, e_1 \rangle$ ,  $\Pi_2 : \langle P_1, P_2, e_2 \rangle$ ,  $\Pi_3 : \langle P_1, P_2, e_3 \rangle$ , through  $\overline{P_1, P_2}$ . Let  $P$  be another point of  $S$ . Then  $P$  will lie on one of these 4 planes, say  $\Pi_0$ . In this case, the plane  $\Pi_0$  contains 5 points of  $S$ , and by maximality,  $S$  contains a  $\mathcal{M}_{7d}$  configuration, which case has been treated above.

**Class  $\mathcal{M}_{8e}$ .**

Let  $S$  contain a  $\mathcal{M}_{7f}$  configuration represented by  $S_0 = \{e_0, e_0 + e_3, e_1, e_1 + e_3, e_2, e_2 + e_3, e_0 + e_1 + e_2\}$ . Here,  $I_2(S_0)$  has a basis

$$\{X_1(X_2 - X_0), X_0(X_2 - X_1), X_3(X_0 + X_1 + X_2 - X_3)\}.$$

Let  $P' = ae_0 + be_1 + ce_2 + de_3$  be another point of  $S$ . Then there are three possibilities for  $P'$ : (i)  $P'$  lies on one of the planes  $\Pi_0 : \langle e_0, e_1, e_3 \rangle$ ,  $\Pi_1 : \langle e_0, e_2, e_3 \rangle$  and  $\Pi_2 : \langle e_1, e_2, e_3 \rangle$ ,  $\Pi_3 : \langle e_0, e_1, e_2 \rangle$ ,  $\Pi_4 : \langle e_0 + e_3, e_1 + e_3, e_2 + e_3 \rangle$  containing 4 points of  $S_0$ , (ii)  $P'$  lies on one of the 21 lines containing two points of  $S_0$ , (iii)  $P'$  lies on the line  $\overline{e_3, e_0 + e_1 + e_2}$ , (iv)  $P'$  lies in the complement of these planes and lines in  $PG(3, 3)$ .

In cases (i) and (ii), by maximality,  $S$  contains a  $\mathcal{M}_{6b}$  configuration. From a  $\mathcal{M}_{6b}$  configuration, we have seen earlier that a  $\mathcal{M}_{7b}$ ,  $\mathcal{M}_{7c}$  or  $\mathcal{M}_{7d}$  configuration can be obtained, and these cases have been treated above.

In case (iii), if  $P'$  lies on the line  $\overline{e_3, e_0 + e_1 + e_2}$ , then  $P'$  is either  $e_0 + e_1 + e_2 + e_3$  or  $-e_0 - e_1 - e_2 - e_3$ . Thus, the 8 points of  $S_0 \cup \{P'\}$  lies on 4 concurrent lines  $\overline{e_3e_0}$ ,  $\overline{e_3e_1}$ ,  $\overline{e_3e_2}$ ,  $\overline{e_3e_0 + e_1 + e_2}$ . But these 8 points of  $S_0 \cup \{P'\}$  do not satisfy the condition to be a  $\mathcal{M}_{7g}$  configuration. So, the configuration consisting of points of  $S_0 \cup \{P'\}$  is of rank 8 and contained in a  $\mathcal{M}_{8b}$  configuration represented by the 4 concurrent lines  $\overline{e_3e_0}$ ,  $\overline{e_3e_1}$ ,  $\overline{e_3e_2}$ ,  $\overline{e_3e_0 + e_1 + e_2}$ . By Lemma 3.3,  $S_0 \cup \{P'\}$  is not a maximal configuration and  $S$  is a  $\mathcal{M}_{8b}$  configuration.

Now we will consider case (iv). The number of points contained in the union  $\Pi_0 \cup \Pi_1 \cup \Pi_2 \cup \Pi_3 \cup \Pi_4$  is 35. All the 21 lines formed by the 2 points of  $S_0$  are contained in these 5 planes. Also, there are two points on the line  $\overline{e_3, e_0 + e_1 + e_2}$  which are not in the union of these 5 planes and 21 lines. Thus, there are total 37 points of  $PG(3, 3)$  covered by union of the planes  $\Pi_0, \Pi_1, \Pi_2, \Pi_3, \Pi_4$ , and  $\overline{e_3, e_0 + e_1 + e_2}$ . The remaining 3 points  $\{e_0 - e_1 - e_2 + e_3, -e_0 + e_1 - e_2 + e_3, -e_0 - e_1 + e_2 + e_3\}$  which form a single orbit under the automorphism group  $Z_2 \times S_3$  of the  $\mathcal{M}_{7f}$  configuration. So we may take  $P' = e_0 - e_1 - e_2 + e_3$ . Let  $T = S_0 \cup \{P'\}$ . A basis of the  $I_2(T)$  is given by

$$\{X_1(X_2 - X_0) + X_3(X_0 + X_1 + X_2 - X_3), X_0(X_2 - X_1)\},$$

and it is readily checked that  $V(I_2(T))$  equals  $T$ . Thus,  $T$  is a maximal  $\mathcal{C}_8$ , which we denote as  $\mathcal{M}_{8e}$ . The configuration  $T_i = T \setminus \{P_i\}$  for  $i = 1, \dots, 8$  are  $\mathcal{C}_7$  configurations with 7 points and containing no lines. In the list  $\mathcal{M}_{7a} - \mathcal{M}_{7g}$ , the only configuration with 7 points and containing no lines is  $\mathcal{M}_{7f}$ . Therefore, the size of the class of  $\mathcal{M}_{8e}$  configurations is:

$$|\mathcal{M}_{8e}| = \frac{1}{8}|\mathcal{M}_{7f}| \cdot 3 = |G|/32 = 379080.$$

In the pencil  $\mathbb{P}(I_2(T))$ , there is one element each in  $\mathcal{O}_1$  and  $\mathcal{O}_2$  and the remaining 2 elements are in  $\mathcal{O}_6$ . Thus,  $\nu(T) = (1, 1, 0, 0, 0, 2)$ .

Let  $S$  contain a  $\mathcal{M}_{7g}$  configuration represented by  $S_0 = \{e_0, e_0 + e_3, e_1, e_1 + e_3, e_2, e_2 + e_3, e_0 + e_1 - e_2, e_0 + e_1 - e_2 + e_3\}$ . Let us consider the six planes  $\Pi_0 : \langle e_0, e_1, e_3 \rangle$ ,  $\Pi_1 : \langle e_2, e_0 + e_1 - e_2, e_3 \rangle$ ,  $\Pi_2 : \langle e_0, e_2, e_3 \rangle$ ,  $\Pi_3 : \langle e_0, e_0 + e_3, e_0 + e_1 - e_2 \rangle$ ,  $\Pi_4 : \langle e_1, e_2, e_3 \rangle$ , and  $\Pi_5 : \langle e_1, e_1 + e_3, e_0 + e_1 - e_2 \rangle$ . Each of the plane  $\Pi_i, i \in \{0, \dots, 5\}$ , contain 4 points of  $S_0$ . Now,

$$\begin{aligned} |\cup_{i=0}^5 \Pi_i| &= |\Pi_0| + |\Pi_1 \setminus \Pi_0| + |\Pi_2 \setminus (\Pi_0 \cup \Pi_1)| + |\Pi_3 \setminus (\cup_{i=0}^2 \Pi_i)| + |\Pi_4 \setminus (\cup_{i=0}^3 \Pi_i)| + |\Pi_5 \setminus (\cup_{i=0}^4 \Pi_i)| \\ &= 13 + 9 + 6 + 6 + 3 + 3 = 40. \end{aligned}$$

This shows that the union  $\cup_{i=0}^5 \Pi_i$  contains all the points of  $PG(3, 3)$ . If  $S$  contains another point  $P$ , then  $P$  will be on at least one of the planes  $\Pi_i$  and since  $\Pi_i$  contains 5 points of  $S$ , by maximality, it will contain a  $\mathcal{M}_{5b}$  configuration. From a  $\mathcal{M}_{5b}$  configuration we only get a  $\mathcal{M}_{6b}$  configuration and from a  $\mathcal{M}_{6b}$  configuration we can get a  $\mathcal{M}_{7b}$ , a  $\mathcal{M}_{7c}$  or a  $\mathcal{M}_{7d}$  configuration. So, this case has been treated above.  $\square$

### 7.1. $B_{j,8}$ for $j > 8$ .

. Let  $T$  be a  $\mathcal{M}_{8a}$  configuration of 16 points represented by a plane  $\Pi$  of  $PG(3, 3)$  and a line  $L$  not contained in  $\Pi$ . Let  $L \cap \Pi = \{Q\}$ . Since  $j > 8$ , the possible cases for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 7$  are as follows:

- (i)  $S = \mathcal{M}_{7a}$  consisting of a plane  $\pi$  and a point  $P$  not on  $\pi$ . In this case  $|\mathcal{M}_{7a}(T)| = 3$ , as we must take  $\pi = \Pi$  and there are 3 ways to pick  $P$  from  $L \setminus Q$ .
- (ii)  $S = \mathcal{M}_{7b}$  consisting of 3 non-coplanar lines  $\{\ell_1, \ell_2, \ell_3\}$  through a point  $P$ . In this case,  $|\mathcal{M}_{7b}(T)| = 6$ , as we must take  $P = Q$ ,  $\ell_1 = L$ , and there are  $\binom{4}{2} = 6$  choices for  $\{\ell_2, \ell_3\}$  in  $\Pi$ .
- (iii)  $S = \mathcal{M}_{7c}$  consisting of three lines  $\{\ell_1, \ell_2, \ell_3\}$  where  $\ell_2$  and  $\ell_3$  are skew to each other and the line  $\ell_1$  meets both the lines. We must take  $\ell_3 = L$ ,  $\ell_2$  to be a line in  $\Pi$  not passing through  $Q$ , and  $\ell_1$  a line in  $\Pi$  passing through  $P$ . Thus, there are 1,  $q^2 = 9$  and  $(q + 1) = 4$  choices for  $\ell_1, \ell_2$  and  $\ell_3$  respectively. Therefore,  $|\mathcal{M}_{7c}(T)| = 36$ .
- (iv)  $S = \mathcal{M}_{7d}$  which consists of a pair of lines  $\ell, \ell'$  in a plane  $\pi$ , and a pair of points  $P, P'$  not lying on  $\pi$  such that the line  $\overline{P, P'}$  meets  $\pi$  in a point not lying on  $\ell \cup \ell'$ . In this case  $|\mathcal{M}_{7d}(T)| = 108$ , as there are  $\binom{9}{2} = 36$  choices for the pair  $\ell, \ell'$  of lines not passing through  $Q$  and  $\binom{3}{2} = 3$  choices for the 2 points  $P, P'$  from  $L \setminus Q$ .
- (v)  $S = \mathcal{M}_{6a}$  consisting of a plane  $\pi$ . In this case we must take  $\pi = \Pi$  and hence  $|\mathcal{M}_{6a}(T)| = 1$ .

By (13), we have,

$$B_{j,8}(\mathcal{M}_{8a}) = \binom{16}{j} - 3 \cdot \left( \binom{14}{j} - \binom{13}{j} \right) - 6 \cdot \binom{10}{j} - 36 \cdot \binom{10}{j} - 108 \cdot \binom{9}{j} - \binom{13}{j},$$

i.e.

$$B_{j,8}(\mathcal{M}_{8a}) = \binom{16}{j} - 3 \cdot \binom{13}{j-1} - 42 \cdot \binom{10}{j} - 108 \cdot \binom{9}{j} - \binom{13}{j}.$$

. A  $\mathcal{M}_{8b}$  configuration  $T$  of 13 points represented by 4 concurrent lines  $L_1, \dots, L_4$  through a point  $Q$ , such that no three of these lines are coplanar. Since  $j > 8$ , the possible cases for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 7$  are listed as follows.

- (i)  $S = \mathcal{M}_{7b}$  consisting of three non-planar concurrent lines. In this case  $|\mathcal{M}_{7b}(T)| = 4$ , as there  $\binom{4}{3}$  ways to choose 3 concurrent lines in  $T$ .
- (ii)  $S = \mathcal{M}_{7d}$  which consists of a pair of lines  $L, L'$  in a plane  $\pi$ , and a pair of points  $P, P'$  not lying on  $\pi$  such that the line  $\overline{P, P'}$  meets  $\pi$  in a point not lying on  $L \cup L'$ . In this case  $|\mathcal{M}_{7d}(T)| = \binom{4}{2} \cdot 3 \cdot 3 = 54$ , as there are  $\binom{4}{2}$  ways to choose to pick  $L, L'$  from  $\{L_1, \dots, L_4\}$ , and 9 ways to pick a point each (different from  $Q$ ) from the remaining two lines of  $\{L_1, \dots, L_4\} \setminus \{L, L'\}$ .

By (13), we have,

$$B_{j,8}(\mathcal{M}_{8b}) = \binom{13}{j} - 4 \cdot \binom{10}{j} - 54 \cdot \binom{9}{j}.$$

. Let  $T$  be a  $\mathcal{M}_{8c}$  configuration of 13 points represented by 4 lines  $L_1, L_2, L_3, L_4$  such that  $L_i$  meets  $L_j$  if and only if  $i \not\equiv j \pmod{2}$ . Since  $j > 8$ , the possible cases for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 7$  are as follows.

- (i)  $S = \mathcal{M}_{7c}$  consisting of three lines  $\{\ell_1, \ell_2, \ell_3\}$  where  $\ell_2$  and  $\ell_3$  are skew to each other and the line  $\ell_1$  meets both the lines. In this case  $|\mathcal{M}_{7c}(T)| = 4$ , as we must take  $S$  to be the union of any 3 of the 4 lines  $\{L_1, \dots, L_4\}$ .
- (ii)  $S = \mathcal{M}_{7d}$  which consists of a pair of lines  $L, L'$  in a plane  $\pi$ , and a pair of points  $P, P'$  not lying on  $\pi$  such that the line  $\overline{P, P'}$  meets  $\pi$  in a point not lying on  $L \cup L'$ . In this case, we must take  $L, L'$  to a pair  $L_i, L_j$  with  $i \not\equiv j \pmod{4}$  (of which there are 4 choices). If  $\{L, L'\} = \{L_1, L_2\}$  then there are 2 choices of points on each of the lines  $L_3, L_4$  for the pair  $P, P'$ . Thus,  $|\mathcal{M}_{7d}(T)| = 4 \cdot 2 \cdot 2 = 16$

By (13), we have,

$$B_{j,8}(\mathcal{M}_{8c}) = \binom{12}{j} - 4 \cdot \binom{10}{j} - 16 \cdot \binom{9}{j}.$$

. Let  $T$  be a  $\mathcal{M}_{8d}$  configuration of 10 points, represented by  $(L_1, L_2, \{P_1, P_2, P_3\})$  where  $\{L_1, L_2\}$  is a pair of lines in a plane  $\pi_1$ , and the points  $P_1, P_2, P_3$  are in a different plane  $\pi_2$ . The point  $Q = L_1 \cap L_2$  lies on the line  $L = \pi_1 \cap \pi_2$ , but none of the points  $P_1, P_2, P_3$  lie on  $L$ . The four points  $\{Q, P_1, P_2, P_3\}$  are in general position in  $\pi_2$ . Since  $j > 8$ , the possible case for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 7$  is  $S = \mathcal{M}_{7d}$ . In this case  $|\mathcal{M}_{7d}(T)| = 3$ , as we must take  $S = T \setminus \{P_i\}$  for  $i = 1 \dots 3$ . By (13), we have,

$$B_{j,8}(\mathcal{M}_{8d}) = \binom{10}{j} - 3 \cdot \binom{9}{j}.$$

We have,  $B_{j,8}(\mathcal{M}_{8e}) = 0$ , as an  $\mathcal{M}_{8e}$  configuration contains exactly 8 points  $PG(3, 3)$ . Therefore, by (12), for  $j > 8$ ,

$$B_{j,8} = 4680 \cdot \left( \binom{16}{j} - 3 \cdot \binom{13}{j-1} - 42 \cdot \binom{10}{j} - 108 \cdot \binom{9}{j} - \binom{13}{j} \right) + 9360 \cdot \left( \binom{13}{j} - 4 \cdot \binom{10}{j} - 54 \cdot \binom{9}{j} \right) + 189540 \cdot \left( \binom{12}{j} - 4 \cdot \binom{10}{j} - 16 \cdot \binom{9}{j} \right) + 336960 \cdot \left( \binom{10}{j} - 3 \cdot \binom{9}{j} \right),$$

i.e.,

$$B_{j,8} = 4680 \cdot \binom{16}{j} + 4680 \cdot \binom{13}{j} - 14040 \cdot \binom{13}{j-1} - 655200 \cdot \binom{10}{j} - 5054400 \cdot \binom{9}{j}.$$

## 8. CONFIGURATIONS OF RANK 9

A configuration  $S$  of rank 9 is, by definition, contained in the set of points  $\mathcal{Q}$  of a quadric of  $PG(3, 3)$  defined by some quadratic form  $f(X_0, \dots, X_3)$ . The points of  $\mathcal{Q}$  are a rank 9 configuration if and only if  $I_2(\mathcal{Q})$  is one dimensional. As seen in Table 1, there are six  $G$ -orbits of quadratic forms on  $PG(3, 3)$  represented by

$$\varphi_1 = X_0X_1, \varphi_2 = X_0X_1 - X_2X_3, \varphi_3 = X_1^2 + X_3^2 - X_0X_2, \varphi_4 = X_0X_2 - X_1^2, \varphi_5 = X_0^2, \varphi_6 = X_0^2 + X_1^2.$$

The quadric  $\mathcal{Q}_i = V(\varphi_i)$  is a rank 9 configuration if and only if  $I_2(V(\varphi_i))$  is the one dimensional vector space generated by  $\varphi_i$ . This is not the case for i)  $\varphi_6$ , ii)  $\varphi_5$  and iii)  $\varphi_4$  because: i)  $V(\varphi_6)$  is a line which has rank 3, ii)  $V(\varphi_5)$  is a plane which has rank 6, and iii)  $I_2(V(\varphi_4))$  is 2-dimensional vector space spanned by  $\varphi_4$  and  $X_0(X_2 - X_1)$ .

The quadrics  $\mathcal{Q}_1, \mathcal{Q}_2$  and  $\mathcal{Q}_3$  have 22, 16 and 10 points respectively. The configurations  $\mathcal{M}_{8a}, \mathcal{M}_{8c}$  and  $\mathcal{M}_{8e}$  have 16, 12 and 8 points respectively. We recall from §7, that the pencil of quadrics

defining  $\mathcal{M}_{8a}, \mathcal{M}_{8c}$  and  $\mathcal{M}_{8e}$  configurations contains an element of  $G \cdot \varphi_1$ ,  $G \cdot \varphi_2$  and  $G \cdot \varphi_3$  respectively. Therefore, i)  $\mathcal{Q}_1$  properly contains an  $\mathcal{M}_{8a}$  configuration, ii)  $\mathcal{Q}_2$  properly contains an  $\mathcal{M}_{8c}$  configuration, and iii)  $\mathcal{Q}_3$  properly contains an  $\mathcal{M}_{8e}$  configuration. Since the containment is proper in each case, and the configurations  $\mathcal{M}_{8a}, \mathcal{M}_{8c}, \mathcal{M}_{8e}$  are maximal, it follows that the quadrics  $\mathcal{Q}_1, \mathcal{Q}_2, \mathcal{Q}_3$  are  $\mathcal{C}_9$ 's. We summarize this:

**Lemma 8.1.** *The set  $\mathcal{C}_9$  can be partitioned into the following 3 classes:*

$\mathcal{M}_{9a}$  of size  $2^2 \cdot 3 \cdot 5 \cdot 13$ , consisting of a pair of hyperplanes of  $PG(3, 3)$ .

$\mathcal{M}_{9b}$  of size  $2 \cdot 3^4 \cdot 5 \cdot 13$ , consisting of hyperbolic quadrics of  $PG(3, 3)$ .

$\mathcal{M}_{9c}$  of size  $2^3 \cdot 3^4 \cdot 13$ , consisting of elliptic quadrics of  $PG(3, 3)$ .

### 8.1. $B_{j,9}$ for $j > 9$ .

. Let  $T$  be a  $\mathcal{M}_{9a}$  configuration of 22 points represented by a pair of hyperplanes  $\pi_1, \pi_2$  intersecting in a line  $L$ . Since  $j > 9$ , the possible cases for  $S \in \mathcal{C}_i(T)$ ,  $i \leq 8$  are listed as follows.

- (i)  $S = \mathcal{M}_{8a}$  consisting of a plane  $\pi$  and a line  $\ell$  not contained in  $\pi$ . In this case  $|\mathcal{M}_{8a}(T)| = 2 \cdot 12 = 24$ , as there are 2 choices for  $\pi$ , and 12 choices for  $\ell$ . Therefore,

$$|\mathcal{M}_{8a}(T)| \cdot B_{j,9}(\mathcal{M}_{8a}(T)) = 24 \cdot \left( \binom{16}{j} - 3 \cdot \binom{13}{j-1} - 42 \cdot \binom{10}{j} - \binom{13}{j} \right).$$

- (ii)  $S = \mathcal{M}_{8b}$ . For a  $\mathcal{M}_{8b}(T)$  configuration of 4 concurrent lines  $L_1, \dots, L_4$  through a point  $Q$ , no three of which are coplanar, the point  $Q$  has to be on the line  $L$ , and  $\pi_1, \pi_2$  each contain two of the lines  $\{L_1, \dots, L_4\}$ . The point  $Q$  on  $L$  can be chosen in 4 ways, and the lines  $L_1, \dots, L_4$  through  $Q$  can be chosen in 9 ways. Thus,  $|\mathcal{M}_{8b}(T)| = 4 \cdot 9 = 36$ . Therefore,

$$|\mathcal{M}_{8b}(T)| \cdot B_{j,9}(\mathcal{M}_{8b}(T)) = 36 \cdot \left( \binom{13}{j} - 4 \cdot \binom{10}{j} \right).$$

- (iii)  $S = \mathcal{M}_{8c}$ . The  $\mathcal{M}_{8c}(T)$  configurations can be chosen in the following way: pick a pair of points  $\{P, P'\}$  on the line  $L$ . A pair of lines  $L_1, L_2$  through  $P$ , and another pair of lines  $L'_1, L'_2$  through  $P'$  such that  $L_1, L_2, L'_1, L'_2$  are distinct from  $L$ , and  $L_i, L'_i$  are in the plane  $\pi_i$  for  $i \in \{1, 2\}$ . There are  $\binom{4}{2}$  ways to pick  $\{P, P'\}$  on  $L$ , and there are  $3^4$  ways to choose the four lines  $\{L_1, L'_1, L_2, L'_2\}$ . Thus,  $|\mathcal{M}_{8c}(T)| = 486$ . Therefore,

$$|\mathcal{M}_{8c}(T)| \cdot B_{j,9}(\mathcal{M}_{8c}(T)) = 486 \cdot \left( \binom{12}{j} - 4 \cdot \binom{10}{j} \right).$$

- (iv)  $S = \mathcal{M}_{8d}$  consisting of  $(\{L_1, L_2\}, \{P_1, P_2, P_3\})$  as described in Lemma 7.1. Such an  $S$  can be chosen as follows: Pick a pair of lines  $L_1, L_2$  in one of the planes  $\pi_1, \pi_2$  intersecting at a point  $Q$  of  $L$ . There are  $2 \cdot 4 \cdot 3 = 24$  ways to pick these. In the other plane, there are 3 lines  $L_1, L_2, L_3$  through  $Q$  apart from  $L$ . There are 18 ways to pick  $P_i \in L_i \setminus \{Q\}$  for  $i \in \{1, 2, 3\}$  such that  $P_1, P_2, P_3$  are non-collinear. Thus,  $|\mathcal{M}_{8d}(T)| = 24 \cdot 18 = 432$ . Therefore,

$$|\mathcal{M}_{8d}(T)| \cdot B_{j,9}(\mathcal{M}_{8d}(T)) = 432 \cdot \binom{10}{j}.$$

- (v)  $S = \mathcal{M}_{7a}$  consisting of a plane  $\pi$  and a point  $P$  not on  $\pi$ . There are 2 choices for  $\pi$  namely  $\pi_1, \pi_2$ . Any of the 9 points of the other plane not lying on  $L$  can be chosen as  $P$ . Thus,  $|\mathcal{M}_{7a}(T)| = 2 \cdot 9 = 18$ . Therefore,

$$|\mathcal{M}_{7a}(T)| \cdot B_{j,9}(\mathcal{M}_{7a}(T)) = 18 \cdot \binom{13}{j-1}.$$

- (vi)  $S = \mathcal{M}_{7b}$ . Let us consider a  $\mathcal{M}_{7b}(T)$  configuration represented by 3 concurrent non-coplanar lines  $L_1, L_2, L_3$  through a point  $Q$ . There are two cases depending on whether  $L \in \{L_1, L_2, L_3\}$  or not.

- (a) In the former case, we may assume  $L = L_1$ . The number of way to choose  $Q$  on  $L$  is 4. Then the two other lines  $L_2, L_3$  through  $Q$  lie in different planes, and can be chosen in  $\binom{3}{1}^2 = 9$  ways. This case contributes 36 to  $|\mathcal{M}_{7b}(T)|$ .
- (b) If  $L \notin \{L_1, L_2, L_3\}$ , then two of the lines  $\{L_1, L_2, L_3\}$  lie in the same plane. The number of ways of choosing  $Q$  on  $L$  is 4, the number of ways choosing two lines through  $Q$  in one of the plane  $\pi_1, \pi_2$  is  $2 \cdot \binom{3}{2} = 6$ . The remaining line through  $Q$  in the other plane can be chosen in  $\binom{3}{1}$  ways. Thus, this case contributes  $4 \cdot 2 \cdot 3 \cdot 3 = 72$  to  $|\mathcal{M}_{7b}(T)|$ .

Thus,  $|\mathcal{M}_{7b}(T)| = 36 + 72 = 108$ . Therefore,

$$|\mathcal{M}_{7b}(T)| \cdot B_{j,9}(\mathcal{M}_{7b}(T)) = 108 \cdot \binom{10}{j}.$$

- (vii)  $S = \mathcal{M}_{7c}$  represented by three lines  $L_1, L_2, L_3$ , where  $L_2$  and  $L_3$  are skew to each other and the line  $L_1$  meets both the lines. There are two cases depending on whether  $L \in \{L_1, L_2, L_3\}$  or not.

- (a) If  $L \in \{L_1, L_2, L_3\}$ , say  $L = L_3$ . let  $P_i = L \cap L_i, i = 1, 2$ . The number of way to choose  $P_1$  and  $P_2$  is  $4 \cdot 3$ . Then the two lines  $L_1$  and  $L_2$  can be chosen in  $\binom{3}{1}^2$ . Thus the number of ways of choosing  $L_1, L_2, L_3$  is  $4 \cdot 3 \cdot \binom{3}{1}^2 = 108$ . Thus, this case contributes 108 to  $|\mathcal{M}_{7c}(T)|$ .
- (b) If  $L \notin \{L_1, L_2, L_3\}$ , then exactly one of the two pairs of lines  $\{L_1, L_2\}$  and  $\{L_2, L_3\}$  are coplanar, say  $\{L_1, L_2\}$ . In this case, the pair  $\{L_2, L_3\}$  can be chosen  $4 \cdot 3 \cdot 3$  ways, and  $L_1$  can be chosen in 9 ways. Thus, this case contributes  $2 \cdot 4 \cdot 3 \cdot 3 \cdot 9 = 648$  to  $|\mathcal{M}_{7c}(T)|$ .

Therefore,  $|\mathcal{M}_{7c}(T)| = 108 + 648 = 756$  and

$$|\mathcal{M}_{7c}(T)| \cdot B_{j,9}(\mathcal{M}_{7c}(T)) = 756 \cdot \binom{10}{j}.$$

- (viii)  $S = \mathcal{M}_{6a}$ . In this case  $|\mathcal{M}_{6a}(T)| = 2$ . Therefore,

$$|\mathcal{M}_{6a}(T)| \cdot B_{j,9}(\mathcal{M}_{6a}(T)) = 2 \cdot \binom{13}{j}.$$

By (13), for  $j > 9$ , we have,

$$\begin{aligned} B_{j,9}(\mathcal{M}_{9a}) &= 780 \cdot \left( \binom{22}{j} - 24 \cdot \binom{16}{j} - 3 \cdot \binom{13}{j-1} - 42 \cdot \binom{10}{j} - \binom{13}{j} \right) - 36 \cdot \left( \binom{13}{j} - 4 \cdot \binom{10}{j} \right) \\ &\quad - 486 \cdot \left( \binom{12}{j} - 4 \cdot \binom{10}{j} \right) - 432 \cdot \binom{10}{j} - 18 \cdot \binom{13}{j-1} - 108 \cdot \binom{10}{j} - 756 \cdot \binom{10}{j} - 2 \cdot \binom{13}{j}, \end{aligned}$$

i.e.

$$B_{j,9}(\mathcal{M}_{9a}) = 780 \cdot \left( \binom{22}{j} - 24 \cdot \binom{16}{j} + 1800 \cdot \binom{10}{j} - 486 \cdot \binom{12}{j} + 54 \cdot \binom{13}{j-1} - 62 \cdot \binom{13}{j} \right).$$

. Let  $T$  be a  $\mathcal{M}_{9b}$  configuration represented by a hyperbolic quadric  $\mathcal{H}$ . A hyperbolic quadric in  $PG(3, 3)$  can be written as a disjoint union of 4 skew lines  $H_1, \dots, H_4$  (known as a regulus), and also as a disjoint union of a dual set of 4 skew lines  $V_1, \dots, V_4$  (known as the opposite regulus). The set  $T$  consists of the 16 points  $H_i \cap V_j$  for  $1 \leq i, j \leq 4$ . Since  $j > 9$ , the possible cases for  $S \in \mathfrak{C}_i(T)$ ,  $i \leq 8$  are as follows.

- (i)  $S = \mathcal{M}_{8c}$  consisting of 4 lines  $L_1, \dots, L_4$  with  $L_i, L_j$  being skew if and only if  $i \equiv j \pmod{2}$ . Here,  $|\mathcal{M}_{8c}(T)| = \binom{4}{2}^2 = 36$ , as  $\{L_1, \dots, L_4\}$  consists of 2 lines from  $\{H_1, \dots, H_4\}$  and 2 lines from  $\{V_1, \dots, V_4\}$ . Therefore,

$$|\mathcal{M}_{8c}(T)| \cdot B_{j,9}(\mathcal{M}_{8c}(T)) = 36 \cdot \left( \binom{12}{j} - 4 \cdot \binom{10}{j} \right).$$

- (ii)  $S = \mathcal{M}_{8d}$  consisting of  $(\{L_1, L_2\}, \{P_1, P_2, P_3\})$  as described in Lemma 7.1. We must have  $|\{L_1, L_2\} \cap \{H_1, \dots, H_4\}| = |\{L_1, L_2\} \cap \{V_1, \dots, V_4\}| = 1$ , and hence there are 16 ways to pick  $\{L_1, L_2\}$ . If  $L_1 = H_4$  and  $L_2 = V_4$ , the points  $\{P_1, P_2, P_3\}$  must be  $\{H_1 \cap V_{\sigma(1)}, H_2 \cap V_{\sigma(2)}, H_3 \cap V_{\sigma(3)}\}$  for some permutation  $\sigma$  of  $\{1, 2, 3\}$ . i.e., there are 6 ways to pick  $\{P_1, P_2, P_3\}$ . Thus,  $|\mathcal{M}_{8d}(T)| = 16 \cdot 6 = 96$ . Therefore,

$$|\mathcal{M}_{8d}(T)| \cdot B_{j,9}(\mathcal{M}_{8d}(T)) = 96 \cdot \binom{10}{j}.$$

- (iii)  $S = \mathcal{M}_{7c}$  represented by three lines  $L_1, L_2, L_3$ , where  $L_2$  and  $L_3$  are skew to each other and the line  $L_1$  meets both the lines. We must have  $L_1 \in \{H_1, \dots, H_4\}$  and  $L_2, L_3 \in \{V_1, \dots, V_4\}$  or  $L_1 \in \{V_1, \dots, V_4\}$  and  $L_2, L_3 \in \{H_1, \dots, H_4\}$ . Thus,  $|\mathcal{M}_{7c}(T)| = 2 \cdot \binom{4}{2} \cdot \binom{4}{1} = 48$ . Therefore,

$$|\mathcal{M}_{7c}(T)| \cdot B_{j,9}(\mathcal{M}_{7c}(T)) = 48 \cdot \binom{10}{j}.$$

By (13), for  $j > 9$ , we have,

$$B_{j,9}(\mathcal{M}_{9b}) = \binom{16}{j} - 36 \cdot \left( \binom{12}{j} - 4 \cdot \binom{10}{j} \right) - 96 \cdot \binom{10}{j} - 48 \cdot \binom{10}{j} = \binom{16}{j} - 36 \cdot \binom{12}{j}.$$

. Let  $T$  be a  $\mathcal{M}_{9c}$  configuration represented by an elliptic quadric  $\mathcal{Q}_3 = V(\varphi_3)$  consisting of 10 points. The only size  $j$  subset of  $T$  for  $j > 9$ , is  $T$  itself. Therefore,

$$B_{j,9}(\mathcal{M}_{9c}) = \binom{10}{j}.$$

Finally, for  $j > 9$ , we have by (12):

$$\begin{aligned} B_{j,9} &= 780 \cdot \left( \binom{22}{j} - 24 \cdot \binom{16}{j} + 1800 \cdot \binom{10}{j} - 486 \cdot \binom{12}{j} + 54 \cdot \binom{13}{j-1} - 62 \cdot \binom{13}{j} \right) \\ &\quad + 10530 \cdot \left( \binom{16}{j} - 36 \cdot \binom{12}{j} \right) + 8424 \cdot \binom{10}{j}, \end{aligned}$$

i.e.,

$$(19) \quad B_{j,9} = 780 \cdot \binom{22}{j} - 8190 \cdot \binom{16}{j} - 48360 \cdot \binom{13}{j} - 14025960 \cdot \binom{12}{j} + 1412424 \cdot \binom{10}{j} + 42120 \cdot \binom{13}{j-1}$$

## 9. PROOF OF THEOREM 1.1

We recall from §3, that we have defined  $B_0(T) = (T^K - 1)$ . Using (7),(9),(10),(14)-(19) we calculate the values of  $B_j(T)$ :

$$\begin{aligned}
B_1(T) &= 40(T^9 - 1), \\
B_2(T) &= 780(T^8 - 1), \\
B_3(T) &= 9880(T^7 - 1), \\
B_4(T) &= 130(T^7 + 702T^6 - 703), \\
B_5(T) &= 936(5T^6 + 698T^5 - 703), \\
B_6(T) &= 780(109T^5 + 4812T^4 - 4921), \\
B_7(T) &= 1560(2T^5 + 663T^4 + 11286T^3 - 11951), \\
B_8(T) &= 585(T - 1)(241T^3 + 16333T^2 + 131461T + 131461), \\
B_9(T) &= 520(T - 1)(55T^3 + 5500T^2 + 138016T + 525844), \\
B_{10}(T) &= 104(T - 1)(110T^3 + 8435T^2 + 399125T + 4410326), \\
B_{11}(T) &= 3120(T - 1)(T^3 + 100T^2 + 6211T + 171775), \\
B_{12}(T) &= 260(T - 1)(2T^3 + 326T^2 + 29837T + 1921619), \\
B_{13}(T) &= 40(T - 1)(T^3 + 352T^2 + 61426T + 9659872), \\
B_{14}(T) &= 1080(T - 1)(T^2 + 508T + 230582), \\
B_{15}(T) &= 18720(T - 1)(4T + 7103), \\
B_{16}(T) &= 1170(4T + 49739)(T - 1), \\
B_{17}(T) &= 20540520(T - 1), \\
B_{18}(T) &= 5705700(T - 1), \\
B_{19}(T) &= 1201200(T - 1), \\
B_{20}(T) &= 180180(T - 1), \\
B_{21}(T) &= 17160(T - 1), \\
B_{22}(T) &= 780(T - 1).
\end{aligned}$$

Having calculated the polynomials  $B_j(T)$ , we use them in (8), to determine the polynomials  $A_j(T)$ :

**Theorem 9.1.** *The extended weight enumerator of  $C_3$  is given by*

$$W(X, Y, T) = X^{40} + (T - 1) \sum_{i=1}^{40} a_i(T) X^{40-i} Y^i,$$

where the polynomials  $a_i(T)$  are as follows:

$$\begin{aligned}
a_{18}(T) &= 780, \\
a_{24}(T) &= 1170(4T - 3), \\
a_{26}(T) &= 1080(T - 3)(T - 9), \\
a_{27}(T) &= 40(T^3 - 26T^2 + 442T - 884), \\
a_{28}(T) &= 189540(T - 3), \\
a_{30}(T) &= 936(100T^2 - 600T + 909), \\
a_{31}(T) &= 505440(T - 3)(T - 6), \\
a_{32}(T) &= 5265(T - 3)(17T^2 - 160T + 513), \\
a_{33}(T) &= 3120(T - 3)(T^3 + 84T^2 - 972T + 3000), \\
a_{34}(T) &= 63180(T - 3)(T^3 - 5T^2 - 45T + 249), \\
a_{35}(T) &= 936(T - 3)(5T^4 + 243T^3 - 6408T^2 + 51654T - 142020), \\
a_{36}(T) &= 130(T^6 + 523T^5 - 15635T^4 + 199500T^3 - 1349544T^2 + 4639614T - 6020826), \\
a_{37}(T) &= 9360(T - 3)(T^5 - 30T^4 + 405T^3 - 3132T^2 + 13932T - 27864), \\
a_{38}(T) &= 780(T - 3)(T^6 - 33T^5 + 507T^4 - 4698T^3 + 28242T^2 - 105678T + 189054), \\
a_{39}(T) &= 40(T - 3)(T^7 - 35T^6 + 585T^5 - 6096T^4 + 43320T^3 - 214344T^2 + 695448T - 1128816), \\
a_{40}(T) &= (T - 3)(T^8 - 36T^7 + 633T^6 - 7110T^5 + 56241T^4 - 325134T^3 \\
&\quad + 1371006T^2 - 3936114T + 5856786).
\end{aligned}$$

Having obtained the extended weight enumerator  $W(Z; T)$  of  $C_3$  over  $\mathbb{F}_3$ , we are in a position to compute the quantities  $W(Z; q^j)$  for  $q = 3$ . We calculate the  $r$ -GWE of  $C_3$  listed in Theorem 1.1, by using the quantities  $W(Z; q^j)$  in the result of Theorem 3.1.

#### REFERENCES

1. J. W. P. Hirschfeld and J. A. Thas, *General Galois geometries*, Springer Monographs in Mathematics, Springer, London, 2016. MR 3445888
2. Trygve Johnsen and Hugues Verdure, *Higher weight spectra of Veronese codes*, IEEE Trans. Inform. Theory **66** (2020), no. 6, 3538–3546. MR 4115115
3. ———, *Higher weight spectra of codes from Veronese threefolds*, J. Pure Appl. Algebra **225** (2021), no. 7, Paper No. 106609, 11. MR 4217930
4. Relinde Jurrius and Ruud Pellikaan, *Codes, arrangements and matroids*, Algebraic geometry modeling in information theory, Ser. Coding Theory Cryptol., vol. 8, World Sci. Publ., Hackensack, NJ, 2013, pp. 219–325. MR 3288286
5. Gilles Lachaud, *Projective Reed-Muller codes*, Coding theory and applications (Cachan, 1986), Lecture Notes in Comput. Sci., vol. 311, Springer, Berlin, 1988, pp. 125–129. MR 960714

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, PUNE, MAHARASHTRA, 411008 INDIA.

*Email address:* kaipa@iiserpune.ac.in, puspendupradhan1@gmail.com