

# Extremal oriented graphs avoiding 1-subdivision of an in-star

Zejun Huang\*, Chenxi Yang

School of Mathematical Sciences, Shenzhen University, Shenzhen 518060, China

## Abstract

An oriented graph is a digraph obtained from an undirected graph by choosing an orientation for each edge. Given a positive integer  $n$  and an oriented graph  $F$ , the oriented Turán number  $ex_{ori}(n, F)$  is the maximum number of arcs in an  $F$ -free oriented graph of order  $n$ . In this paper, we investigate the oriented Turán number  $ex_{ori}(n, \overrightarrow{S_{k,1}})$ , where  $\overrightarrow{S_{k,1}}$  is the 1-subdivision of the in-star of order  $k + 1$ . We determine  $ex_{ori}(n, \overrightarrow{S_{k,1}})$  for  $k = 2, 3$  as well as the extremal oriented graphs. For  $k \geq 4$ , we establish a lower bound and an upper bound on  $ex_{ori}(n, \overrightarrow{S_{k,1}})$ .

**Keywords:** in-star; 1-subdivision; oriented graphs; Turán number

**Mathematics Subject Classification:** 05C20; 05C35

## 1 Introduction and main results

All undirected graphs considered in this paper are finite and simple. Let  $G$  be an undirected graph. We denote  $V(G)$  and  $E(G)$  the *vertex set* and the *edge set* of  $G$ , respectively. We call  $|V(G)|$  the *order* and  $e(G) = |E(G)|$  the *size* of  $G$ . Similarly, given a digraph  $D$ , we denote by  $V(D)$ ,  $A(D)$ ,  $a(D)$  the *vertex set*, the *arc set* and the *size* of  $D$ , respectively. We denote by  $K_r$ ,  $\overrightarrow{K_r}$  and  $\overrightarrow{C_r}$  the complete undirected graph, the complete digraph and the directed cycle of order  $r$ , respectively.

Suppose  $D$  is a digraph with  $u, v \in V(D)$ . If  $uv \in A(D)$ , then  $u$  is called an *in-neighbor* of  $v$  and  $v$  is called an *out-neighbor* of  $u$ . The set of in-neighbors (out-neighbors) of  $u$  is called its *in-neighborhood* (*out-neighborhood*), whose cardinality is called the *in-degree* (*out-degree*) of  $u$ . If all vertices of  $D$  have the same in-degree  $k$ , we say  $D$  is an *in-degree  $k$ -regular digraph*. When  $X$  is a vertex set or an arc set of a digraph (or undirected graph)  $D$ , we use  $D[X]$  to denote the subgraph of  $D$  induced by  $X$ .

An oriented graph is a digraph obtained from an undirected graph by choosing an orientation for each edge. The *1-subdivision* of a digraph  $D$  is the digraph obtained from  $D$  by replacing each arc  $uv \in A(D)$  with a directed 2-path  $uvw$  along with a new vertex  $w$ . Denote by  $\overrightarrow{S_k}$  the in-star of order  $k + 1$  and  $\overrightarrow{S_{k,1}}$  the 1-subdivision of  $\overrightarrow{S_k}$ . The *center* of  $\overrightarrow{S_{k,1}}$  is the same as the center of  $\overrightarrow{S_k}$ ; see Figure 1 and Figure 2 for the diagrams of  $\overrightarrow{S_k}$  and  $\overrightarrow{S_{k,1}}$ .

---

\*Corresponding author. Email: mathzejun@gmail.com




Figure 1:  $\overrightarrow{S}_k$  centered at  $v$




Figure 2:  $\overrightarrow{S}_{k,1}$  centered at  $v$

Two graphs  $G_1$  and  $G_2$  are *isomorphic*, written  $G_1 \cong G_2$ , if there is a bijection  $\sigma : V(G_1) \rightarrow V(G_2)$  such that  $uv \in E(G_1)$  if and only if  $\sigma(u)\sigma(v) \in E(G_2)$ . For graphs  $G$  and  $H$ , we say  $G$  contains a *copy* of  $H$  if  $G$  contains a subgraph isomorphic to  $H$ . If  $G$  contains no copy of  $H$ , then  $G$  is said to be  *$H$ -free*. The same conceptions can be defined for digraphs.

Turán problems on undirected graphs consider the maximum size of  $F$ -free graphs of order  $n$  with  $F$  being a single graph or a family of graphs, which have been attracting a lot of studies since Mantel [21] and Turán [25] determined the maximum size of triangle-free and  $K_t$ -free graphs of order  $n$ ; see [2, 23]. The studies of Turán problems on digraphs were initiated by Brown and Harary [6], who determined the maximum sizes of  $\overrightarrow{K}_r$ -free digraphs and  $\overrightarrow{T}_r$ -free digraphs as well as the extremal digraphs, where  $\overrightarrow{T}_r$  is a tournament of order  $r$ . Brown, Erdős and Simonovits [3–5] presented some asymptotic results on extremal digraphs avoiding a family of digraphs. Häggkvist and Thomassen [9], Zhou and Li [28] characterized the extremal  $\overrightarrow{C}_k$ -free digraphs. Bermond, Germa, Heydemann and Sotteau [1] determined the maximum size of digraphs avoiding all directed cycles of length at most  $r$ ; Chen and Chang [7, 8] studied the same problem by posting additional condition on the minimum in-degree and out-degree. The authors of [12–20, 24, 27] studies extremal digraphs avoiding certain distinct walks or paths with the same initial vertex and terminal vertex.

Turán problems on digraphs when the host digraphs are oriented graphs are also interesting. Howalla, Dabboucy and Tout [10, 11] determined the maximum size of oriented digraphs avoiding  $k$  directed paths with the same initial vertex and terminal vertex for  $k = 1, 2, 3$ . Maurer, Rabinovitch and Trotter [22] studied the maximum size of subgraphs of a transitive tournament which contain at most one directed path from  $x$  to  $y$  for any two distinct vertices  $x, y$ .

Given a positive integer  $n$  and a digraph  $F$ , the *oriented Turán number*  $ex_{ori}(n, F)$  is the maximum size of  $F$ -free oriented graphs of order  $n$ . We denoted by  $EX_{ori}(n, F)$  the extremal  $F$ -free oriented graphs of order  $n$  that attain the size  $ex_{ori}(n, F)$ .

We are interested in the following problem.

**Problem 1.** *Let  $n, k$  be positive integers. Determine  $ex_{ori}(n, \overrightarrow{S}_{k,1})$  and  $EX_{ori}(n, \overrightarrow{S}_{k,1})$ .*

In this paper, we solve Problem 1 for the cases  $k = 2$  and  $k = 3$ . For  $k \geq 4$ , we establish bounds on  $ex_{ori}(n, \overrightarrow{S}_{k,1})$ . Our main results states as follows.

**Theorem 1.1.** *Let  $n \geq 16$  be an integer. Then*

$$ex_{ori}(n, \overrightarrow{S}_{2,1}) = \left\lfloor \frac{(n+1)^2}{4} \right\rfloor.$$

*Moreover,  $D \in EX_{ori}(n, \overrightarrow{S}_{2,1})$  if and only if  $V(D)$  has a partition  $V(D) = X \cup Y$  such that*

- (i)  $|X| \in \{[(n-1)/2], \lceil (n-1)/2 \rceil\}$ ;
- (ii)  $xy \in A(D)$  for all  $x \in X, y \in Y$ ;
- (iii)  $D[X]$  is an empty digraph,  $D[Y]$  is an in-degree 1-regular digraph.

**Theorem 1.2.** *Let  $n \geq 40$  be an integer. Then*

$$ex_{ori}(n, \overrightarrow{S_{3,1}}) = \left\lfloor \frac{(n+2)^2}{4} \right\rfloor.$$

Moreover,  $D \in EX_{ori}(n, \overrightarrow{S_{3,1}})$  if and only if  $V(D)$  has a partition  $V(D) = X \cup Y$  such that

- (i)  $|X| \in \{\lfloor n/2 \rfloor - 1, \lceil n/2 \rceil - 1\}$ ;
- (ii)  $xy \in A(D)$  for all  $x \in X, y \in Y$ ;
- (iii)  $D[X]$  is an empty digraph,  $D[Y]$  is an in-degree 2-regular digraph.

**Theorem 1.3.** *Let  $n, k$  be positive integers with  $k \geq 4$  and  $n \geq 3k + 1$ . Then*

$$\left\lfloor \frac{(n+k-1)^2}{4} \right\rfloor \leq ex_{ori}(n, \overrightarrow{S_{k,1}}) \leq \left\lfloor \frac{(n+k-1)^2}{4} \right\rfloor + (k-1)n.$$

We will present some preliminaries in Section 2 and prove the above theorems in Section 3.

## 2 Preliminaries

Firstly we present some definitions and notations. Given a vector  $\mathbf{x} = (x_1, x_2, \dots, x_n)$ , we rearrange its components in decreasing order as  $x_{[1]} \geq x_{[2]} \geq \dots \geq x_{[n]}$ . Given two nonnegative vectors  $\mathbf{x} = (x_1, x_2, \dots, x_s) \in \mathbb{R}^s$ ,  $\mathbf{y} = (y_1, y_2, \dots, y_t) \in \mathbb{R}^t$ , if  $s \geq t$  and  $x_{[i]} \geq y_{[i]}$  for all  $1 \leq i \leq t$ , then we say  $\mathbf{x}$  covers  $\mathbf{y}$ , denoted  $\mathbf{x} \triangleright \mathbf{y}$ . We denote by  $\mathbf{x} \not\triangleright \mathbf{y}$  if  $\mathbf{x}$  does not cover  $\mathbf{y}$ .

The *underlying graph* of a digraph  $D$  is the undirected graph obtained by replacing each arc by an edge with the same ends. A set of vertex disjoint edges (arcs)  $\{u_1v_1, \dots, u_kv_k\}$  in an undirected graph (digraph) is called a *k-matching*. The *matching number* of an undirected graph (digraph) is the maximum cardinality of matchings in it.

Let  $D$  be a digraph with  $E \subseteq A(D)$ , and let  $X, Y$  be disjoint subsets of  $V(D)$  with  $Y = \{y_1, y_2, \dots, y_k\}$ . Suppose  $G$  is the underlying graph of  $D$ . We will need the following notations:

- $N_X^-(u)$ : the in-neighborhood of  $u$  from  $X$ , which is also denoted by  $N^-(u)$  when  $X = V(D)$ ;
- $N_X^+(u)$ : the out-neighborhood of  $u$  from  $X$ , which is also denoted by  $N^+(u)$  when  $X = V(D)$ ;
- $d_X^-(u)$ : the cardinality of  $N_X^-(u)$ , which is also denoted by  $d^-(u)$  when  $X = V(D)$ ;
- $d_X^+(u)$ : the cardinality of  $N_X^+(u)$ , which is also denoted by  $d^+(u)$  when  $X = V(D)$ ;
- $\Delta^-(D)$ : the maximum in-degree of  $D$ , which is also denoted by  $\Delta^-$  when  $D$  is clear from the context;
- $d_X^-(Y)$ :  $(d_X^-(y_1), d_X^-(y_2), \dots, d_X^-(y_k))$ , which is denoted by  $d^-(Y)$  when  $X = V(D)$  and  $\tilde{d}^-(Y)$  when  $X = V(D) - Y$ ;

$[X, Y]$ : the set of arcs from  $X$  to  $Y$ , which also represents the set of edges between  $X$  and  $Y$  in  $G$ ;

$a(X, Y)$ : the cardinality of  $[X, Y]$ , which is abbreviated as  $a(X)$  when  $X = Y$ ;

$\bar{X}$ : the complement of  $X$  in  $V(D)$ , i.e.,  $V(D) - X$ .

$N^-(X)$ : the set of all in-neighbors of the vertices of  $X$  in  $D$ ;

$N_G(X)$ : the set of all neighbors of the vertices of  $X$  in  $G$ ;

For convenience, sometimes we simply write  $v$  for a singleton  $\{v\}$ . If a bipartite graph  $G$  has a vertex partition  $V(G) = X \cup Y$  such that  $G[X]$  and  $G[Y]$  are empty graphs, then we say  $G$  is a  $(X, Y)$ -bigraph, denoted  $G(X, Y)$ .

We will need the following lemmas.

**Lemma 2.1.** [26] *An  $(X, Y)$ -bigraph  $G$  has a matching that saturates  $X$  if and only if  $|N(S)| \geq |S|$  for all  $S \subseteq X$ .*

**Lemma 2.2.** *Let  $D$  be an oriented graph and  $v \in V(D)$  with  $\{u_1, u_2, \dots, u_k\} \subseteq N^-(v)$ ,  $k \geq 2$ . If  $\tilde{d}^-(u_1, u_2, \dots, u_k) \triangleright (k, k-1, \dots, 1)$ , then  $D$  contains an  $\overrightarrow{S_{k,1}}$  centered at  $v$ .*

*Proof.* Denote by  $Y = \{u_1, u_2, \dots, u_k\}$  and  $E = [\bar{Y} \setminus \{v\}, Y]$ . Suppose  $G$  is the underlying graph of  $D[E]$ . Then  $V(G)$  has a bipartition  $V(G) = \tilde{Y} \cup Y'$  with  $\tilde{Y} \subseteq \bar{Y} \setminus \{v\}$  and  $Y' \subseteq Y$  being independent sets in  $G$ . Since  $\tilde{d}^-(u_1, u_2, \dots, u_k) \triangleright (k, k-1, \dots, 1)$ , we have  $Y' = Y$  and  $|N_G(X)| \geq |X|$  for all  $X \subseteq Y$ . Applying Lemma 2.1 on  $G(\tilde{Y}, Y)$ ,  $G(\tilde{Y}, Y)$  has a  $k$ -matching saturated  $Y$ . It follows that  $D$  contains an  $\overrightarrow{S_{k,1}}$  centered at  $v$ .  $\square$

**Lemma 2.3.** *Let  $D$  be an oriented graph and  $v \in V(D)$  with  $N^-(v) = \{u_1, u_2, \dots, u_k\}$ .*

(i) *If  $d^-(u_1, u_2, \dots, u_k) \triangleright (4, 3, 3)$ , then  $D$  contains an  $\overrightarrow{S_{3,1}}$  centered at  $v$ .*

(ii) *If  $d^-(u_1, u_2, \dots, u_k) \triangleright (2, 2)$ , then  $D$  contains an  $\overrightarrow{S_{2,1}}$  centered at  $v$ .*

*Proof.* Without loss of generality, we assume  $u_1 \geq u_2 \geq \dots \geq u_k$ .

(i) Denote by  $S = \{u_1, u_2, u_3\}$ . Suppose  $d^-(u_i) = d_i$  for  $i = 1, 2, 3$ . Since  $D$  is an oriented graph, we have

$$d_S^-(u_i) \leq 2 \quad \text{for } i = 1, 2, 3 \quad (2.1)$$

and

$$d_S^-(u_1) + d_S^-(u_2) + d_S^-(u_3) \leq 3. \quad (2.2)$$

Since  $d^-(u_1, u_2, \dots, u_k) \triangleright (4, 3, 3)$ , we have  $\tilde{d}^-(u_1, u_2, u_3) \triangleright (3, 2, 1)$ . Applying Lemma 2.2 we get the conclusion.

(ii) The proof of (ii) is similar.  $\square$

**Lemma 2.4.** *Let  $D$  be an  $\overrightarrow{S_{3,1}}$ -free oriented graph and  $v \in V(D)$  with  $d^-(v) \geq 3$ . Suppose  $S = \{u_1, u_2, u_3\} \subseteq N^-(v)$ .*

(i) *If  $\tilde{d}^-(S) = (2, 2, 2)$ , then the subgraph of  $D$  induced by  $\{v, u_1, u_2, u_3\} \cup N^-(S)$  contains a spanning subgraph isomorphic to  $H_1$  (Figure 3).*

(ii) If  $d^-(S) = (3, 3, 3)$ , then the subgraph of  $D$  induced by  $\{v, u_1, u_2, u_3\} \cup N^-(S)$  contains a spanning subgraph isomorphic to  $H_2$  (Figure 4).




Figure 3:  $H_1$




Figure 4:  $H_2$

*Proof.* (i) Suppose  $E = [\bar{S}, S]$  and  $G$  is the underlying graph of  $D[E]$ . Since  $D$  is  $\overrightarrow{S_{3,1}}$ -free,  $G$  has no matching saturating  $S$ .

Since  $\tilde{d}^-(u_1, u_2, u_3) = (2, 2, 2)$ , we have

$$|N_G(u_1)| = |N_G(u_2)| = |N_G(u_3)| = 2.$$

Since we have  $|N_G(S')| = 2 \geq |S'|$  for all  $S' \subseteq S$  with  $|S'| \leq 2$ , applying Lemma 2.1, we have  $|N_G(S)| = 2$ , which leads to  $|N^-(S)| = 2$ . Therefore, the subgraph of  $D$  induced by  $\{v, u_1, u_2, u_3\} \cup N^-(S)$  contains a spanning subgraph isomorphic to  $H_1$ .

(ii) Since  $D$  is an oriented graph and  $|S| = 3$ , we have (2.1) and (2.2). On the other hand, since  $D$  is  $\overrightarrow{S_{3,1}}$ -free, applying Lemma 2.2, we have

$$\tilde{d}^-(u_1, u_2, u_3) \not\triangleright (3, 2, 1).$$

Therefore,  $d^-(u_1) = d^-(u_2) = d^-(u_3) = 3$  leads to  $d_{\bar{S}}^-(u_1) = d_{\bar{S}}^-(u_2) = d_{\bar{S}}^-(u_3) = 1$  and  $\tilde{d}^-(S) = (2, 2, 2)$ . By (i), the subgraph  $D_1$  of  $D$  induced by  $\{v, u_1, u_2, u_3\} \cup N^-(S)$  contains a spanning subgraph isomorphic to  $H_1$  and a triangle with vertices  $u_1, u_2, u_3$ . Hence,  $D_1$  contains a spanning subgraph isomorphic to  $H_2$ .  $\square$

**Lemma 2.5.** *Let  $D$  be an  $\overrightarrow{S_{3,1}}$ -free oriented graph and  $v \in V(D)$ . If  $d^-(N^-(v)) \triangleright (3, 3, 3)$ , then*

$$d^-(N^-(v)) = (3, 3, 3, 1, 0, 0, \dots, 0) \quad \text{or} \quad d^-(N^-(v)) = (3, 3, 3, 0, 0, \dots, 0).$$

*Proof.* Suppose  $N^-(v) = \{u_1, \dots, u_k\}$ . Without loss of generality, we may assume  $d^-(u_1) \geq d^-(u_2) \geq \dots \geq d^-(u_k)$ . Applying Lemma 2.3, we have  $d^-(u_1, u_2, u_3) \not\triangleright (4, 3, 3)$ . Since  $d^-(u_i) \geq 3$  for  $i = 1, 2, 3$ , we get

$$d^-(u_1) = d^-(u_2) = d^-(u_3) = 3. \tag{2.3}$$

Applying Lemma 2.4, we have  $|N^-(u_1, u_2, u_3)| = 2$  and  $D[V_1]$  contains a spanning subgraph  $D_1$  isomorphic to  $H_2$ , where  $V_1 = \{v, u_1, u_2, u_3\} \cup N^-(u_1, u_2, u_3)$ . Without loss of generality, we assume  $D_1 = H_3$  with  $N^-(u_1, u_2, u_3) = \{w_1, w_2\}$ ; see Figure 5. Denote by  $V_2 = \{u_1, u_2, u_3, w_1, w_2\}$ .

Firstly we claim

$$d^-(u) = 0 \quad \text{for all} \quad u \in N^-(v) \setminus V_2. \tag{2.4}$$

Otherwise, suppose  $u_4 \in N^-(v) \setminus V_2 \neq \emptyset$  with  $d^-(u_4) \neq 0$  and  $w_3 \in N^-(u_4)$ . Notice that  $w_3 \neq v$ . If  $w_3 \notin \{u_1, u_2, u_3, w_1, w_2\}$ , then  $D$  has a subgraph  $H_4$ ; see Figure 6. If  $w_3 \in \{u_1, u_2, u_3\}$ , without loss of generality, we may assume  $w_3 = u_3$ . Then  $D$  has a subgraph  $H_5$ ; see Figure 7. If  $w_3 \in \{w_1, w_2\}$ , without loss of generality, we may assume  $w_3 = w_2$ . Then  $D$  has a subgraph  $H_6$ ; see Figure 8. Since  $H_i$  contains an  $\overrightarrow{S}_{3,1}$  centered at  $v$  for  $i = 4, 5, 6$ , we get a contradiction.

Combining (2.3) and (2.4), now we only need to prove that at most one vertex in  $N^-(v) \cap \{w_1, w_2\}$  has nonzero in-degree, which equals one.

Suppose  $w_i \in N^-(v)$  with  $i \in \{1, 2\}$ . If  $D$  has a vertex  $w_3 \in N^-(w_i)$  such that  $w_3 \neq w_{3-i}$ , then  $D$  has a subgraph  $H_7$  (Figure 9), which contains an  $\overrightarrow{S}_{3,1}$ , a contradiction. Therefore, we have  $N^-(w_i) = \{w_{3-i}\}$  or  $N^-(w_i) = \emptyset$ . Since  $D$  is an oriented graph,  $D$  contains at most one vertex in  $N^-(v) \cap \{w_1, w_2\}$  with nonzero in-degree, which equals one.  $\square$




Figure 5:  $H_3$




Figure 6:  $H_4$




Figure 7:  $H_5$




Figure 8:  $H_6$




Figure 9:  $H_7$

**Lemma 2.6.** *Let  $n$  and  $k$  be integers such that  $k \geq 2$  and  $n \geq 3k + 1$ . Then*

$$ex_{ori}(n, \overrightarrow{S}_{k,1}) \geq \left\lfloor \frac{(n+k-1)^2}{4} \right\rfloor. \quad (2.5)$$

*Proof.* Let  $D_k$  be an oriented graph with  $n$  vertices and  $V(D_k)$  has a bipartition  $V(D_k) = X \cup Y$  such that

- (i)  $|X| \in \{\lfloor (n-k+1)/2 \rfloor, \lceil (n-k+1)/2 \rceil\}$ ;
- (ii)  $xy \in A(D)$  for all  $x \in X, y \in Y$ ;
- (iii)  $D[X]$  is an empty graph;

(iv)  $D[Y]$  is an in-degree  $(k-1)$ -regular oriented graph.

We remark that the above  $D_k$  always exists. Since  $n \geq 3k+1$ , we have  $|Y| \geq 2k-1$ . Denote by  $Y = \{y_1, y_2, \dots, y_t\}$ . Then  $y_i y_j \in A(D)$  if and only if  $i-j \pmod t \in \{1, 2, \dots, k-1\}$  guarantees an in-degree  $(k-1)$ -regular oriented graph  $D[Y]$ .

Obviously,  $D_k$  is an  $\overrightarrow{S_{k,1}}$ -free oriented graph with  $\lfloor (n+k-1)^2/4 \rfloor$  arcs. Therefore, we have (2.5).  $\square$

### 3 Proofs of the main results

In this section, we present the proofs of our main results.

#### 3.1 Proof of Theorem 1.1

Suppose  $D \in EX_{ori}(n, \overrightarrow{S_{2,1}})$  with  $n \geq 16$ . Partition  $V(D) = X \cup Y$  with

$$X = \{x \in V(D) : d^-(x) \leq 1\}, \quad Y = \{y \in V(D) : d^-(y) \geq 2\}.$$

We have the following claim.

**Claim 3.1.** *For the above  $X$  and  $Y$ , we have*

$$d_Y^-(y) \leq 1 \quad \text{for all } y \in Y, \tag{3.1}$$

and

$$\frac{n}{2} - \sqrt{\frac{n}{2}} < |X| < \frac{n}{2} + \sqrt{\frac{n}{2}}. \tag{3.2}$$

*Proof of Claim 3.1.* By Lemma 2.3, we have (3.1). Since  $D \in EX_{ori}(n, \overrightarrow{S_{2,1}})$ , by Lemma 2.6 we have

$$\begin{aligned} \frac{n^2 + 2n}{4} < \left\lfloor \frac{(n+1)^2}{4} \right\rfloor &\leq a(D) \leq \sum_{v \in X} d_X^-(v) + \sum_{v \in Y} d_Y^-(v) \\ &\leq |X| + (|X|+1)(n-|X|) = -|X|^2 + n|X| + n, \end{aligned}$$

which leads to (3.2).

By Claim 3.1, we have  $\Delta^- \leq |X| + 1$ . Denote by  $s = |X|$ . We distinguish three cases.

*Case 1.*  $\Delta^- \leq s - 1$ . We have

$$\begin{aligned} a(D) &= \sum_{x \in X} d^-(x) + \sum_{y \in Y} d^-(y) \leq s + (s-1)(n-s) \\ &= -s^2 + (n+2)s - n \leq \frac{n^2 + 4}{4} < \left\lfloor \frac{(n+1)^2}{4} \right\rfloor. \end{aligned} \tag{3.3}$$

Case 2.  $\Delta^- = s$ . Then we have

$$a(D) = \sum_{x \in X} d^-(x) + \sum_{y \in Y} d^-(y) \leq s + (n-s)s \leq \left\lfloor \frac{(n+1)^2}{4} \right\rfloor. \quad (3.4)$$

Suppose equality in (3.4) holds. Then we have  $s \in \{\lfloor (n+1)/2 \rfloor, \lceil (n+1)/2 \rceil\}$  and

$$d^-(x) = 1, \quad d^-(y) = s \quad \text{for all } x \in X, y \in Y. \quad (3.5)$$

If  $D[Y]$  is an empty graph. Then

$$N^-(v) = X \quad \text{for all } v \in Y,$$

which implies  $a(X) = s \geq 6$ . Therefore,  $D[X]$  contains a 2-matching, which leads to an  $\overrightarrow{S}_{2,1}$  in  $D$ , a contradiction. If  $D[Y]$  is not an empty graph, choose an arc  $y_1y_2 \in D[Y]$ . Then by (3.5) we have

$$|N_X^-(y_1) \cap N_X^-(y_2)| \geq s - 2 > 3. \quad (3.6)$$

Take a vertex  $x \in N_X^-(y_1) \cap N_X^-(y_2)$  with its unique in-neighbor  $w$ . Then by (3.6) there exists a vertex  $x' \in N_X^-(y_1) \cap N_X^-(y_2) - \{x, w\}$ . It follows that  $D$  has an  $\overrightarrow{S}_{2,1} = \{wxy_2, x'y_1y_2\}$ , a contradiction.

Therefore, we have  $a(D) < \lfloor (n+1)^2/4 \rfloor$  in this case.

Case 3.  $\Delta^- = s + 1$ . Let  $v$  be a vertex with maximum in-degree. Then by (3.1) we have  $N^-(v) = X \cup \{y_1\}$  for some  $y_1 \in Y$ . We distinguish two subcases.

*Subcase 3.1.*  $D[X]$  is not an empty graph. We choose an arbitrary arc  $x_1x_2 \in A(D)$  with  $x_1, x_2 \in X$ . It follows that  $N^-(y_1) \subseteq \{x_1, x_2\}$ ; otherwise a vertex  $w \in N^-(y_1) - \{x_1, x_2\}$  leads to an  $\overrightarrow{S}_{2,1} = \{x_1x_2v, wy_1v\}$ , a contradiction. Since  $d^-(y_1) \geq 2$ , we have  $N^-(y_1) = \{x_1, x_2\}$ , which implies that  $a(X) = 1$ , as  $x_1x_2$  is chosen arbitrarily in  $D[X]$ .

Moreover, we have  $a(y_1, X) = 0$ ; otherwise if  $y_1x_3 \in A(D)$  for some  $x_3 \in X$ , then  $x_3 \notin \{x_1, x_2\}$  and  $D$  contains an  $\overrightarrow{S}_{2,1} = \{x_1x_2v, y_1x_3v\}$ , a contradiction.

Therefore, we have

$$\begin{aligned} a(D) &= a(X) + a(X, Y) + a(Y, X) + a(Y) \\ &= a(X) + [a(X, Y \setminus y_1) + a(Y \setminus y_1, X)] + a(Y) + [a(y_1, X) + a(X, y_1)] \\ &\leq 1 + (n-s-1)s + (n-s) + 2 < \left\lfloor \frac{(n+1)^2}{4} \right\rfloor. \end{aligned}$$

*Subcase 3.2.*  $D[X]$  is an empty graph. We count  $a(D)$  as follows:

$$\begin{aligned} a(D) &= a(X) + a(X, Y) + a(Y, X) + a(Y) \\ &\leq s(n-s) + (n-s) \leq \left\lfloor \frac{(n+1)^2}{4} \right\rfloor. \end{aligned} \quad (3.7)$$



Suppose equality in (3.7) holds. Then  $s \in \{\lfloor (n-1)/2 \rfloor, \lceil (n-1)/2 \rceil\}$ ,

$$d_Y^-(y) = 1 \quad \text{for all } y \in Y \quad (3.8)$$

and

$$|\{xy, yx\} \cap A(D)| = 1 \quad \text{for all } x \in X, y \in Y. \quad (3.9)$$

Next we prove

$$[Y, X] = \emptyset. \quad (3.10)$$

To the contrary, suppose  $[Y, X] \neq \emptyset$ . Denote by  $X_1 = \{v \in X : d^-(x) = 1\}$ . Then we have  $X_1 \neq \emptyset$  and  $N^-(X_1) \subseteq Y$ . Moreover, all vertices in  $X_1$  has the same in-neighborhood, say,  $\{y_1\}$ ; otherwise  $y_1x_1, y_2x_2 \in A(D)$  with  $x_1, x_2 \in X_1$ ,  $y_1, y_2 \in Y$  leads to an  $\overrightarrow{S_{2,1}} = \{y_1x_1v, y_2x_2v\}$ , a contradiction. Therefore, by (3.9), we have

$$X \subset N^-(y) \quad \text{for all } y \in Y \setminus \{y_1\}.$$

On the other hand, by (3.8),  $D[Y]$  has an arc  $y_2y_3$  with  $y_2, y_3 \in Y \setminus \{y_1\}$ . Take two vertices  $x_1 \in X_1$  and  $x_2 \in X \setminus \{x_1\}$ . Then  $D$  contains an  $\overrightarrow{S_{2,2}} = \{y_1x_1y_3, x_2y_2y_3\}$ , a contradiction. Therefore, we have (3.10).

Combining all the above cases, we have  $a(D) \leq \lfloor (n+1)^2/4 \rfloor$ , with equality only if  $V(D)$  has a partition  $V(D) = X \cup Y$  such that the conditions (i), (ii), (iii) in Theorem 1.1 hold.

On the other hand, let  $D$  be an oriented graph whose vertex set  $V(D)$  can be partitioned as  $V(D) = X \cup Y$  such that the conditions (i), (ii), (iii) in Theorem 1.1 hold. Then  $D$  is  $\overrightarrow{S_{2,1}}$ -free with  $a(D) = \lfloor (n+1)^2/4 \rfloor$ .

This completes the proof of Theorem 1.1. □

### 3.2 Proof of Theorem 1.2

Suppose  $D \in EX_{ori}(n, \overrightarrow{S_{3,1}})$  with  $n \geq 40$ . We partition  $V(D) = X \cup Y$  with

$$X = \{u : d^-(u) \leq 2\}, \quad Y = \{u : d^-(u) \geq 3\}.$$

Let  $s = |X|$ . We need the following claim.

**Claim 3.2.** *For the above  $X$  and  $Y$ , we have*

$$d_Y^-(v) \leq 2 \quad \text{for all } v \in V(D) \quad (3.11)$$

and

$$\frac{n}{2} - \sqrt{n} < |X| < \frac{n}{2} + \sqrt{n}. \quad (3.12)$$

*Proof of Claim 3.2.* For a vertex  $w \in V(D)$  with the maximum in-degree, we assert that

$$|\{u \in N^-(w) : d^-(u) \geq 3\}| \leq 2. \quad (3.13)$$

To the contrary, suppose  $|\{u \in N^-(w) : d^-(u) \geq 3\}| \geq 3$ . Applying Lemma 2.5, we have  $d^-(N^-(w)) = (3, 3, 3, 1, 0, 0, \dots, 0)$  or  $(3, 3, 3, 0, 0, \dots, 0)$ . It follows that

$$\begin{aligned} a(D) &\leq \sum_{u \in N^-(w)} d^-(u) + \sum_{u \notin N^-(w)} d^-(u) \leq 10 + \sum_{u \notin N^-(w)} d^-(u) \\ &\leq 10 + \Delta^-(n - \Delta^-) < \left\lfloor \frac{(n+2)^2}{4} \right\rfloor, \end{aligned}$$

which contradicts (2.5).

By (3.13), we have

$$|X| \geq \Delta^- - 2 \quad \text{and} \quad |Y| \leq n - \Delta^-(D) + 2.$$

Suppose  $D$  has a vertex  $v$  such that  $d_Y^-(v) \geq 3$ , i.e.,  $N^-(v)$  contains 3 vertices with in-degree  $\geq 3$ . By (3.13), we have  $d^-(v) \equiv r < \Delta^-$ . Applying Lemma 2.5, we get  $d^-(N^-(v)) = (3, 3, 3, 1, 0, \dots, 0)$  or  $(3, 3, 3, 0, \dots, 0)$ . It follows that

$$|X - N^-(v)| \geq \Delta^- - 2 - (r - 3) = \Delta^- - r + 1 \quad \text{and} \quad |Y - N^-(v)| \leq n - \Delta^- - 1.$$

Therefore, we have

$$\begin{aligned} a(D) &= \sum_{u \in X - N^-(v)} d^-(u) + \sum_{u \in Y - N^-(v)} d^-(u) + \sum_{u \in N^-(v)} d^-(u) \\ &\leq 2(\Delta^- - r + 1) + \Delta^-(n - \Delta^- - 1) + 10 \\ &\leq \frac{n^2 + 2n + 49 - 2r}{4} < \left\lfloor \frac{(n+2)^2}{4} \right\rfloor, \end{aligned}$$

which contradicts (2.5). Hence, we have (3.11).

Since  $D \in EX_{ori}(n, \overrightarrow{S_{3,1}})$ , by Lemma 2.6 we have

$$\begin{aligned} \frac{n^2 + 4n}{4} < \left\lfloor \frac{(n+2)^2}{4} \right\rfloor \leq a(D) &\leq \sum_{v \in X} d_X^-(v) + \sum_{v \in Y} d_Y^-(v) \\ &\leq 2|X| + (|X| + 2)(n - |X|) = -|X|^2 + n|X| + 2n, \end{aligned}$$

which leads to (3.12).

This completes the proof of Claim 3.2.

Suppose  $D[X]$  is empty. Notice that  $a(X, Y) + a(Y, X) \leq s(n - s)$ . We have the following bound on the size of  $D$ :

$$a(D) \leq a(X) + a(X, Y) + a(Y, X) + a(Y) \leq s(n - s) + 2(n - s) \leq \lfloor (n+2)^2/4 \rfloor. \quad (3.14)$$

Suppose equality in (3.14) holds, then  $s \in \{\lfloor n/2 \rfloor - 1, \lceil n/2 \rceil - 1\}$ ,

$$d_Y^-(y) = 2 \quad \text{and} \quad |\{xy, yx\} \cap A(D)| = 1 \quad \text{for all } x \in X, y \in Y. \quad (3.15)$$

Next we prove  $[Y, X] = \emptyset$ .

To the contrary, suppose  $[Y, X] \neq \emptyset$ . We call  $y \in Y$  a *bad vertex* if  $d_X^-(y) = 1, d_X^+(y) = s - 1$ . Then  $Y$  contains at most two bad vertices. In fact, if  $Y$  has three bad vertices  $v_1, v_2, v_3$ , then by the definition of bad vertices we have  $d_X^+(v_i) = s - 1$  for  $i = 1, 2, 3$ , which leads to

$$|N_X^+(v_1) \cap N_X^+(v_2) \cap N_X^+(v_3)| \geq s - 3 \geq 1.$$

It follows that there is vertex  $u \in X$  with  $\{v_1, v_2, v_3\} \subseteq N^-(u)$ , which contradicts the construction of  $X$ .

Suppose  $Y$  has two bad vertices  $v_1, v_2$ . Then  $|N_X^+(v_1) \cap N_X^+(v_2)| \geq s - 2 \geq 1$ . Take  $x' \in N_X^+(v_1) \cap N_X^+(v_2)$  and  $y' \in \{v_1, v_2\}$ . Then  $y'x' \in [Y, X]$ . Now we prove

$$N^+(x') \subseteq N^+(y'). \quad (3.16)$$

Take any  $y_0 \in N^+(x')$ . Since  $D[X]$  is empty, we have  $y_0 \in Y$ . By (3.15) we may denote  $N_Y^-(y_0) = \{y_1, y_2\}$ . Since  $d^-(y_i) \geq 3$ , we have  $N_X^-(y_i) \neq \emptyset$  for  $i = 1, 2$ . We assert that

$$N_X^-(y_i) \setminus \{x'\} \neq \emptyset \quad \text{for } i = 1, 2. \quad (3.17)$$

In fact, if  $N_X^-(y_i) = \{x'\}$  for  $i \in \{1, 2\}$ , then by (3.15) we have  $N_X^+(y_i) = X \setminus \{x'\}$ , which means  $y_i$  is a bad vertex, i.e.,  $y_i \in \{v_1, v_2\}$ . It follows that  $y_i x' \in A(D)$ , which contradicts  $N_X^-(y_i) = \{x'\}$ .

Since  $d_Y^-(y_1) = d_Y^-(y_2) = 2$ ,  $D$  has a matching  $M_1 = \{y_3 y_1, y_4 y_2\}$  with  $y_1, y_2, y_3, y_4 \in Y$ . Take  $x_i \in N_X^-(y_i) \setminus \{x'\}$  for  $i = 1, 2$ , where  $x_1$  and  $x_2$  may be identical. Then  $M_2 = \{x_1 y_1, y_4 y_2\}, M_3 = \{y_3 y_1, x_2 y_2\}$  are matchings in  $D$ .

Since  $D$  is  $\overrightarrow{S_{3,1}}$ -free, for every matching  $M = \{w_1 y_1, w_2 y_2\}$  in  $D$  with  $x' \notin \{w_1, w_2\}$ , we have  $y' \in V(M) = \{w_1, w_2, y_1, y_2\}$ . Therefore,

$$y' \in V(M_1) \cap V(M_2) \cap V(M_3) = \{y_1, y_2\} = N_Y^-(y_0).$$

Hence, we get  $y_0 \in N^+(y')$  and (3.16) holds.

Since  $d^-(x') \leq 2$  and  $d_Y^-(x') + d_Y^+(x') = n - s$ , we have  $d_Y^+(x') \geq n - s - 2$ . Notice that (3.16) implies  $d_Y^+(y') \geq d_Y^+(x')$ . Since  $D$  is an oriented graph and  $d_Y^-(y') + d_Y^+(y') \leq |Y| - 1 = n - s - 1$ , we have  $d_Y^-(y') \leq 1$ , which contradicts  $d_Y^-(y) = 2$ .

If  $Y$  contains one bad vertex  $v$ , take  $y' = v$  and  $x' \in N^+(v)$ ; if  $Y$  contains no bad vertex, choose any  $x' \in X$  and  $y' \in Y$  with  $y'x' \in A(D)$ . Applying the same arguments as above we can deduce a contradiction, noticing that (3.17) is obvious when  $Y$  contains no bad vertex.

Therefore, we have

$$[Y, X] = \emptyset \quad \text{and} \quad [X, Y] = \{uv, u \in X, v \in Y\}.$$

Hence,  $D$  is an oriented graph satisfying the conditions (i), (ii) and (iii) in Theorem 1.2.

On the other hand, any oriented graph  $D$  satisfying the conditions (i), (ii) and (iii) in Theorem 1.2 is  $\overrightarrow{S}_{3,1}$ -free with  $\lfloor (n+2)^2/4 \rfloor$  arcs. Therefore, it suffices to confirm  $a(D) < \lfloor (n+2)^2/4 \rfloor$  when  $D[X]$  is not empty. By Claim 3.2 we have  $\Delta^- \leq s+2$ . We distinguish three cases.

*Case 1.*  $\Delta^- \leq s$ . We count the size of  $D$  as follows:

$$a(D) \leq \sum_{x \in X} d^-(x) + \sum_{y \in Y} d^-(y) \leq 2s + (n-s)s \leq \lfloor (n+2)^2/4 \rfloor. \quad (3.18)$$

Suppose equality in (3.18) holds. Then  $s \in \{\lfloor n/2 \rfloor + 1, \lceil n/2 \rceil + 1\}$ ,

$$d^-(x) = 2 \quad \text{and} \quad d^-(y) = s \quad \text{for all} \quad x \in X, y \in Y. \quad (3.19)$$

We assert that  $D[Y]$  is an empty graph. To the contrary, choose any arc  $y_1 y_0 \in A(D)$  with  $y_0, y_1 \in Y$ . Since  $d^-(y_0) = s$  and  $d_{\overline{Y}}^-(y_0) \leq 2$ , we have

$$d_{\overline{X}}^-(y_0) \geq s-2 \quad \text{and} \quad |X - N^-(y_0)| \leq 2.$$

If  $|X \cap N^-(y_0) - N^+(y_1)| \leq 1$ , then we have

$$|X \cap N^+(y_1)| \geq |X \cap N^-(y_0) \cap N^+(y_1)| \geq |X \cap N^-(y_0)| - 1 = |X| - |X - N^-(y_0)| - 1 \geq s-3.$$

Since  $D$  is an oriented graph and  $d_{\overline{Y}}^-(y_1) \leq 2$ , we have

$$|X \cap N^-(y_1)| \leq |X| - |X \cap N^+(y_1)| \leq 3 \quad (3.20)$$

and

$$d^-(y_1) = d_{\overline{X}}^-(y_1) + d_{\overline{Y}}^-(y_1) = |X \cap N^-(y_1)| + d_{\overline{Y}}^-(y_1) \leq 5 \quad (3.21)$$

which contradicts  $d^-(y_1) = s \geq 14$  by (3.12). Therefore, we have

$$|X \cap N^-(y_0) - N^+(y_1)| \geq 2.$$

Choose  $x_1, x_2 \in X \cap N^-(y_0) - N^+(y_1)$ . Then  $d^-(x_1) = d^-(x_2) = 2$  and we can find a matching  $M = \{w_1 x_1, w_2 x_2\}$  with  $y_1 \notin \{w_1, w_2\}$ . Since  $D$  is  $\overrightarrow{S}_{3,1}$ -free, we have  $N^-(y_1) \subseteq \{w_1, w_2, x_1, x_2\}$  and  $d^-(y_1) \leq 4$ , which contradicts  $d^-(y_1) = s \geq 14$ . Therefore,  $D[Y]$  is an empty graph.

By (3.19), we have

$$N^-(y) = X \quad \text{for all} \quad y \in Y, \quad (3.22)$$

which leads to  $a(Y, X) = 0$ . Hence, given any 2-matching  $M$  in  $D[X]$ , we have

$$N^-(v) \subset V(M) \quad \text{for all } v \in X \setminus V(M), \quad (3.23)$$

since otherwise we can find an  $\overrightarrow{S_{3,1}}$  in  $D$ .

By (3.19) and (3.22),  $D$  contains a matching  $M_0 = \{x_1x_2, x_3x_4\}$  with  $\{x_1, x_2, x_3, x_4\} \subset X$ . Since  $d^-(x_1) = d^-(x_3) = 2$ , there exists a vertex  $x_5 \in X \setminus \{x_1, x_2, x_3, x_4\}$  such that either  $x_5x_1 \in A(D)$  or  $x_5x_3 \in A(D)$ . Without loss of generality, we assume  $x_5x_3 \in A(D)$ .

By (3.19) and (3.23), we have  $|N^-(x_5)| = 2$  and  $N^-(x_5) \subset \{x_1, x_2, x_3, x_4\}$ . We can deduce contradictions for all the six choices of  $N^-(x_5)$ . We take  $N^-(x_5) = \{x_1, x_2\}$  for example, as the argument for other cases are similar. Note that  $D$  has four matchings  $M_1 = \{x_1x_2, x_3x_4\}$ ,  $M_2 = \{x_1x_2, x_5x_3\}$ ,  $M_3 = \{x_1x_5, x_3x_4\}$ ,  $M_4 = \{x_2x_5, x_3x_4\}$ . By (3.23), we have  $N^-(u) \subset V(M_1) \cap V(M_2) \cap V(M_3) \cap V(M_4) = \{x_3\}$  for all  $u \in X \setminus \{x_1, x_2, x_3, x_4, x_5\}$ , which contradicts (3.19) as  $|X| > 5$ .

Therefore, we have  $a(D) < \lfloor (n+2)^2/4 \rfloor$  in this case.

*Case 2.*  $\Delta^- = s + 2$ . Choose a vertex  $y_0 \in Y$  with maximum in-degree. By (3.11) we have  $N^-(y_0) = X \cup \{y_1, y_2\}$  for some  $y_1, y_2 \in Y$ . Recall that  $d^-(y_i) \geq 3$  for  $i = 1, 2$ . We distinguish three subcases.

*Case 2.1.*  $d^-(y_1) \geq 4$  and  $d^-(y_2) \geq 4$ . Since  $D[X]$  is not an empty graph, we choose an arc  $x_1x_2 \in A(D)$  with  $x_1, x_2 \in X$ . Then we have  $\tilde{d}^-(y_1, y_2, x_2) \triangleright (3, 2, 1)$ . Applying Lemma 2.2,  $D$  contains a  $\overrightarrow{S_{3,1}}$  centered at  $y_0$ , a contradiction.

*Case 2.2.* One of  $\{y_1, y_2\}$  has in-degree 3, and the other one has in-degree larger than 3, say,  $d^-(y_1) = 3$  and  $d^-(y_2) \geq 4$ . If there is a vertex  $x \in X$  such that  $N^-(x) - \{y_1, y_2\} \neq \emptyset$  and  $xy_1 \notin A(D)$ , then  $\tilde{d}^-(y_1, y_2, x) \triangleright (3, 2, 1)$ . Applying Lemma 2.2,  $D$  contains an  $\overrightarrow{S_{3,1}}$  centered at  $y_0$ , a contradiction. Therefore, for any  $x \in X$ , we have either

$$N^-(x) \subseteq \{y_1, y_2\} \quad (3.24)$$

or

$$xy_1 \in A(D). \quad (3.25)$$

Since  $d^-(y_1) = 3$ ,  $X$  has at most 3 vertices satisfying (3.25). Hence,  $X$  has at most 3 vertices such that (3.24) does not hold, which leads to  $a(X) \leq 6$ .

Since  $D[X]$  is not empty, there is an arc  $x_1x_2 \in A(D)$  with  $x_1, x_2 \in X$ . Since  $d^-(y_2) \geq 4$ , there is a vertex  $w_1 \in N^-(y_2) - \{x_1, x_2\}$ . It follows that

$$a(y_1, X) + a(X, y_1) \leq 3; \quad (3.26)$$

otherwise, there is a vertex  $x_3 \in N_X^+(y_1) \cup N_X^-(y_1) - \{x_1, x_2, w_1\}$ , which leads to an  $\overrightarrow{S_{3,1}}$  centered at  $y_0$  as the following figures shows, a contradiction.




Figure 10:  $x_3 \in N^-(y_1)$




Figure 11:  $x_3 \in N^+(y_1)$

Therefore, we can count  $a(D)$  as follows:

$$\begin{aligned}
a(D) &= a(X) + a(X, Y) + a(Y, X) + a(Y) \\
&= a(X) + [a(X, Y \setminus y_1) + a(Y \setminus y_1, X)] + [a(X, y_1) + a(y_1, X)] + a(Y) \\
&\leq 6 + (n - s - 1)s + 3 + 2(n - s) \\
&< \left\lfloor \frac{(n + 2)^2}{4} \right\rfloor.
\end{aligned} \tag{3.27}$$

*Case 2.3.*  $d^-(y_1) = d^-(y_2) = 3$ . If there is a vertex  $x \in X$  such that  $x \notin N^-(y_1) \cup N^-(y_2)$  and  $N^-(x) - \{y_1, y_2\} \neq \emptyset$ , then  $\tilde{d}^-(y_1, y_2, x) \triangleright (3, 2, 1)$ . Applying Lemma 2.2,  $D$  contains a  $\overrightarrow{S_{3,1}}$  centered at  $y_0$ , a contradiction. Therefore, for any  $x \in X$ , we have either (3.24) or

$$x \in N^-(y_1) \cup N^-(y_2). \tag{3.28}$$

Since  $|N^-(y_1) \cup N^-(y_2)| \leq 6$ ,  $X$  has at most 6 vertices that do not satisfying (3.24), which leads to  $a(X) \leq 12$ .

Again, since  $D[X]$  is not empty, there is an arc  $x_1x_2 \in a(D)$  with  $x_1, x_2 \in X$ . Notice that  $d^-(y_1) = d^-(y_2) = 3$  and  $D$  is an oriented graph, we have

$$N^-(y_1) - \{x_1, x_2, y_2\} \neq \emptyset \quad \text{or} \quad N^-(y_2) - \{x_1, x_2, y_1\} \neq \emptyset.$$

Without loss of generality, we assume  $N^-(y_2) - \{x_1, x_2, y_1\} \neq \emptyset$  contains a vertex  $w_1$ . Then similarly as in Case 2.2, we have (3.26).

Now using the same calculation as in (3.27) we have  $a(D) < \lfloor (n + 2)^2/4 \rfloor$ .

*Case 3.*  $\Delta^- = s + 1$ . We distinguish two subcases.

*Subcase 3.1.* There exists a vertex  $y_0 \in Y$  with in-degree  $\Delta^-$  such that  $d_X^-(y_0) = s - 1$  and  $d_Y^-(y_0) = 2$ . Let  $N_Y^-(y_0) = \{y_1, y_2\}$  and  $X - N_X^-(y_0) = \{x_0\}$ . Denote by  $X' = N_X^-(y_0)$ . We distinguish three subcases.

*Subcase 3.1.1.*  $d^-(y_1) \geq 4, d^-(y_2) \geq 4$ . If there is a vertex  $x \in X'$  such that  $N^-(x) - \{y_1, y_2\} \neq \emptyset$ , then  $\tilde{d}^-(y_1, y_2, x) \triangleright (3, 2, 1)$ . Applying Lemma 2.2,  $D$  contains an  $\overrightarrow{S_{3,1}}$ , a contradiction. Therefore, we have

$$N^-(x) \subseteq \{y_1, y_2\} \quad \text{for all } x \in X', \tag{3.29}$$

which leads to

$$\sum_{x \in X'} d^-(x) \leq d_{X'}^+(y_1) + d_{X'}^+(y_2).$$

Notices that

$$d_{X'}^+(y_i) + d^-(y_i) \leq d_X^+(y_i) + d_X^-(y_i) + d_Y^-(y_i) \leq s + 2 \quad \text{for } i = 1, 2.$$

We have

$$\begin{aligned} a(D) &= d^-(x_0) + \sum_{x \in X'} d^-(x) + d^-(y_1) + d^-(y_2) + \sum_{y \in Y \setminus \{y_1, y_2\}} d^-(y) \\ &\leq d^-(x_0) + [d_{X'}^+(y_1) + d^-(y_1)] + [d_{X'}^+(y_2) + d^-(y_2)] + \sum_{y \in Y \setminus \{y_1, y_2\}} d^-(y) \\ &\leq 2 + 2(s + 2) + (s + 1)(n - s - 2) \\ &< \left\lfloor \frac{(n + 2)^2}{4} \right\rfloor. \end{aligned} \tag{3.30}$$

*Subcase 3.1.2.* One of  $\{y_1, y_2\}$  has in-degree 3, and the other one has in-degree larger than 3, say,  $d^-(y_1) = 3$  and  $d^-(y_2) \geq 4$ . Using the same arguments as in Case 2.2, for every  $x \in X'$ , we have either (3.24) or (3.25). Since  $d^-(y_1) = 3$ ,  $X'$  has at most 3 vertices such that (3.24) does not hold, which leads to  $\sum_{x \in X'} d_X^-(x) \leq 6$ . It follows that

$$a(X) \leq \sum_{x \in X'} d_X^-(x) + d^-(x_0) \leq 8.$$

If (3.24) holds for all  $x \in X'$ , using the same calculation as in Subcase 3.1.1 we have  $a(D) < \lfloor (n + 2)^2/4 \rfloor$ . If there is a vertex  $x_1 \in X'$  such that  $N^-(x_1) \not\subseteq \{y_1, y_2\}$ , i.e.,  $x_1$  has an in-neighbor  $w_1 \in V(D) \setminus \{y_1, y_2\}$ , then  $d^-(y_2) \geq 4$  implies that  $y_2$  has an in-neighbor  $w_2$  from  $V(D) \setminus \{x_1, y_1, w_1\}$ . It follows that

$$a(X, y_1) + a(y_1, X) \leq 4; \tag{3.31}$$

otherwise there is a vertex  $x_2 \in X' - \{x_1, w_1, w_2\}$  such that either  $x_2 y_1 \in A(D)$  or  $y_1 x_2 \in A(D)$ , which both guarantee an  $\overrightarrow{S_{3,1}}$  in  $D$ , a contradiction. Therefore, similarly as in (3.27), we have

$$\begin{aligned} a(D) &= a(X) + [a(X, Y \setminus y_1) + a(Y \setminus y_1, X)] + [a(X, y_1) + a(y_1, X)] + a(Y) \\ &\leq 8 + s(n - s - 1) + 4 + 2(n - s) \\ &< \left\lfloor \frac{(n + 2)^2}{4} \right\rfloor. \end{aligned}$$

*Subcase 3.1.3.*  $d^-(y_1) = d^-(y_2) = 3$ . Then similarly as in Subcase 2.3, for any  $x \in X'$ , we have either (3.24) or (3.28). If (3.24) holds for all  $x \in X'$ , then we have (3.30) again.

If there is a vertex  $x_1 \in X'$  such that  $N^-(x_1) \not\subseteq \{y_1, y_2\}$ , i.e., there is a vertex  $x_1 \in X'$  which has an in-neighbor  $w_1 \in V(D) \setminus \{y_1, y_2\}$ , then  $d^-(y_1) = d^-(y_2) = 3$  implies that at least one of  $\{y_1, y_2\}$ , say  $y_2$ , has an in-neighbor  $w_2$  from  $V(D) \setminus \{x_1, y_1, y_2, w_1\}$ . Similarly as in Subcase 3.1.2,

we have (3.31). On the other hand, since  $|N^-(y_1) \cup N^-(y_2)| \leq 6$ ,  $X'$  has at most 6 vertices  $x$  such that  $N^-(x) \not\subseteq \{y_1, y_2\}$ , which leads to  $\sum_{x \in X'} d_X^-(x) \leq 12$  and  $a(X) = \sum_{x \in X} d_X^-(x) \leq 14$ . Therefore, similarly as in (3.27), we have

$$\begin{aligned} a(D) &= a(X) + [a(X, Y \setminus y_1) + a(Y \setminus y_1, X)] + [a(X, y_1) + a(y_1, X)] + a(Y) \\ &\leq 14 + s(n - s - 1) + 4 + 2(n - s) \\ &< \left\lfloor \frac{(n + 2)^2}{4} \right\rfloor. \end{aligned}$$

We remark that we need  $n \geq 40$  to guarantee the above inequality.

*Subcase 3.2.*  $d_X^-(y) = s, d_Y^-(y) = 1$  for all  $y \in Y$  with  $d^-(y) = \Delta^-$ . Let  $y_0 \in Y$  be a vertex with maximum in-degree and  $N^-(y_0) = X \cup \{y_1\}$  with  $y_1 \in Y$ . Let  $E = \{ux \in A(D) : x \in X, u \neq y_1\}$ . We distinguish two subcase.

*Subcase 3.2.1.*  $D[E]$  contains a 2-matching  $M_0 = \{w_1x_1, w_2x_2\}$ . Since  $D$  is  $\overrightarrow{S}_{3,1}$ -free and  $N^-(y_0) = X \cup \{y_1\}$ , for every 2-matching  $M = \{u_1v_1, u_2v_2\}$  with  $v_1, v_2 \in X$ , we have

$$N^-(v) \subseteq V(M) \quad \text{for all } v \in X \cup \{y_1\} - V(M). \quad (3.32)$$

We claim that

$$|\{u \in X : d^-(u) = 2\}| \leq 4. \quad (3.33)$$

Otherwise, we have a vertex  $x_3 \in X - \{w_1, w_2, x_1, x_2\}$  with  $d^-(x_3) = 2$ . By (3.32) we have  $N^-(x_3) \subseteq \{w_1, w_2, x_1, x_2\}$ . For each of the six choices of  $N^-(x_3)$ , we can always deduce a contradiction. We take  $N^-(x_3) = \{w_1, x_1\}$  as example, as the arguments for other cases are similar. Note that  $D$  contains three matchings  $M_0 = \{w_1x_1, w_2x_2\}$ ,  $M_1 = \{w_1x_3, w_2x_2\}$ ,  $M_2 = \{x_1x_3, w_2x_2\}$ . By (3.32), we have  $N^-(y_1) \subseteq V(M_0) \cap V(M_1) \cap V(M_2) = \{w_2, x_2\}$ , which contradicts  $y_1 \in Y = \{u \in V(D) : d^-(u) \geq 3\}$ .

Similarly to the argument for (3.26), we have  $a(X, y_1) + a(y_1, X) \leq 4$ , which implies  $d^-(y_1) \leq 6$ . Therefore, by (3.33) we have

$$\begin{aligned} a(D) &= \sum_{u \in X} d^-(u) + \sum_{u \in Y \setminus y_1} d^-(u) + d^-(y_1) \\ &\leq s + 4 + (s + 1)(n - s - 1) + 6 \\ &< \left\lfloor \frac{(n + 2)^2}{4} \right\rfloor. \end{aligned}$$

*Subcase 3.2.2.*  $D[E]$  contains no 2-matching. By the construction of  $E$  we have

$$\sum_{x \in X} d^-(x) = |E| + d_X^+(y_1). \quad (3.34)$$

Since  $d_Y^-(y_1) \leq 2$ , we have

$$d_X^+(y_1) + d^-(y_1) \leq s + 2. \quad (3.35)$$



We assert that

$$|E| + \sum_{u \in Y \setminus y_1} d^-(u) \leq s - 1 + (s + 1)(n - s - 1). \quad (3.36)$$

If  $|E| \leq s - 1$ , the assertion is obvious. Suppose  $|E| \geq s$ . Since  $s \geq 14$  and  $D[E]$  contains no 2-matching, the underlying graph of  $D[E]$  is a star. Moreover, since  $d^-(u) \leq 2$  for all  $u \in X$ ,  $E$  contains at most 2 vertices from  $Y$ , which leads to  $|E| \leq s + 1$ . If  $|E| = s$ , then  $E$  covers at least one vertex  $y' \in Y \setminus y_1$ , which implies  $y'$  is an in-neighbor of some vertex in  $X$ . Therefore, we have  $X \not\subset N^-(y')$  and  $d^-(y') < \Delta^-$ . If  $|E| = s + 1$ , then  $E$  covers two vertices  $v_1, v_2 \in Y \setminus y_1$ , which implies  $X \not\subset N^-(v_i)$  and  $d^-(v_i) < \Delta^-$  for  $i = 1, 2$ . In both cases we have (3.36).

By (3.34),(3.35) and (3.36) we have

$$\begin{aligned} a(D) &= \sum_{u \in X} d^-(u) + \sum_{u \in Y} d^-(u) = |E| + \sum_{u \in Y \setminus y_1} d^-(u) + d_X^+(y_1) + d^-(y_1) \\ &\leq s - 1 + (s + 1)(n - s - 1) + s + 2 < \left\lfloor \frac{(n + 2)^2}{4} \right\rfloor. \end{aligned}$$

This completes the proof of Theorem 1.2.  $\square$

### 3.3 Proof of Theorem 1.3

By Lemma 2.6 we get the lower bound. We prove the upper bound. Suppose  $D$  is an  $\overrightarrow{S_{k,1}}$ -free oriented graph of order at least  $3k + 1$  with  $k \geq 4$ . We assert that

$$|\{u \in N^-(v) : d^-(u) \geq 2k - 1\}| < k \quad \text{for all } v \text{ with } d^-(v) \geq k. \quad (3.37)$$

To the contrary, suppose  $v$  is a vertex with in-degree at least  $k$  and  $S = \{u_1, \dots, u_k\} \subseteq \{u \in N^-(v) : d^-(u) \geq 2k - 1\}$ . Then  $d^-(S) \triangleright (2k - 1, 2k - 1, \dots, 2k - 1)$ . Since  $D$  is an oriented graph, we have

$$d_S^-(u_i) \leq k - 1 \quad \text{for } i = 1, \dots, k,$$

which leads to  $\tilde{d}^-(u_1, u_2, \dots, u_k) \triangleright (k, k - 1, \dots, 1)$ . Applying Lemma 2.2 we get (3.37).

Now we partition  $V(D)$  as  $V(D) = X \cup Y$  with  $X = \{u : d^-(u) \leq 2k - 2\}$ ,  $Y = \{u : d^-(u) \geq 2k - 1\}$ . Let  $s = |X|$ . Then by (3.37) we have

$$d_Y^-(u) \leq k - 1 \quad \text{for all } u \in Y,$$

which leads to

$$d^-(u) \leq s + k - 1 \quad \text{for all } u \in Y.$$

It follows that

$$a(D) = \sum_{x \in X} d^-(x) + \sum_{y \in Y} d^-(y) \leq (2k - 2)s + (s + k - 1)(n - s) \leq \left\lfloor \frac{(n + k - 1)^2}{4} \right\rfloor + (k - 1)n.$$

This completes the proof. □

*Remark.* It would be interesting to determine the precise value of  $ex_{ori}(n, \overrightarrow{S_{k,1}})$  and the extremal digraphs for  $n \geq 4$ . We leave this for future work.

## Acknowledgement

This work was supported by the National Natural Science Foundation of China (No. 12171323) and Guangdong Basic and Applied Basic Research Foundation (No. 2022A1515011995).

## References

- [1] J.C. Bermond, A. Germa, M.C. Heydemann, D. Sotteau, Girth in digraphs, *J. Graph Theory* 4 (1980) 337-341.
- [2] B. Bollobás, Extremal graph theory. *Handbook of combinatorics*, Vol. 1, 2, 1231-1292, Elsevier Sci. B. V., Amsterdam, 1995.
- [3] W.G. Brown, P. Erdős, M. Simonovits, Extremal problems for directed graphs, *J. Combin. Theory Ser. B* 15 (1973) 77-93.
- [4] W.G. Brown, P. Erdős, M. Simonovits, Inverse extremal digraph problems. *Finite and infinite sets*, Vol. I, II (Eger, 1981), 119-156, *Colloq. Math. Soc. János Bolyai*, 37, North-Holland, Amsterdam, 1984.
- [5] W.G. Brown, P. Erdős, M. Simonovits, Algorithmic solution of extremal digraph problems, *Trans. Amer. Math. Soc.* 292 (1985) 421-449.
- [6] W.G. Brown, F. Harary, Extremal digraphs. *Combinatorial theory and its applications*, I-III (Proc. Colloq., Balatonfüred, 1969), pp. 135-198, *Colloq. Math. Soc. János Bolyai*, 4, North-Holland, Amsterdam-London, 1970.
- [7] B. Chen, A. Chang, Turán number of 3-free strong digraphs with out-degree restriction, *Discrete Appl. Math.* 314 (2022) 252-264.
- [8] B. Chen, A. Chang, 3-free strong digraphs with the maximum size, *Graphs Combin.* 37 (2021) 2535-2554.
- [9] R. Häggkvist, C. Thomassen, On pancyclic digraphs, *J. Combin. Theory Ser. B* 20 (1976) 20-40.
- [10] K. Howalla, A.N. Dabboucy, R. Tout, On the maximum number of arcs in some classes of graphs, *Časopis Pěst. Mat.* 107 (1982) 388-392.
- [11] K. Howalla, A.N. Dabboucy, R. Tout, An extremal problem for some classes of oriented graphs, *Časopis Pěst. Mat.* 108 (1983) 53-69.

- [12] Z. Huang, Z. Lyu, 0-1 matrices with zero trace whose squares are 0-1 matrices, *Linear Algebra Appl.* 565 (2019) 156-176.
- [13] Z. Huang, Z. Lyu, Extremal digraphs avoiding an orientation of  $C_4$ , *Discrete Math.* 343 (2020) 111827.
- [14] Z. Huang, Z. Lyu, 0-1 matrices whose  $k$ -th powers have bounded entries, *Linear Multilinear Algebra* 68 (2020) 1972-1982.
- [15] Z. Huang, Z. Lyu, P. Qiao, A Turán problem on digraphs avoiding distinct walks of a given length with the same endpoints, *Discrete Math.* 342 (2019) 1703-1717.
- [16] Z. Huang, X. Zhan, Digraphs that have at most one walk of a given length with the same endpoints, *Discrete Math.* 311 (2011) 70-79.
- [17] Z. Lyu, Digraphs that contain at most  $t$  distinct walks of a given length with the same endpoints, *J. Comb. Optim.* 41 (2021) 762-779.
- [18] Z. Lyu, A note on extremal digraphs containing at most  $t$  walks of length  $k$  with the same endpoints, *Graphs Combin.* 38 (2022) 33.
- [19] Z. Lyu, Extremal digraphs avoiding distinct walks of length 4 with the same endpoints, *Discuss. Math. Graph Theory* 42 (2022) 985-1004.
- [20] Z. Lyu, Extremal digraphs avoiding an orientation of the diamond, *Graphs Combin.* 37 (2021) 1373-1383.
- [21] W. Mantel, Problem 28, *Wiskundige Opgaven* 10 (1907) 60-61.
- [22] S.B. Maurer, I. Rabinovitch, W.T. Trotter, A generalization of Turán's theorem to directed graphs, *Discrete Math.* 32 (1980) 167-189.
- [23] V. Nikiforov, Some new results in extremal graph theory. *Surveys in combinatorics 2011*, 141-181, *London Math. Soc. Lecture Note Ser.*, 392, Cambridge Univ. Press, Cambridge, 2011.
- [24] J. Tuite, G. Erskine, N. Salia, Turán problems for  $k$ -geodetic digraphs, *Graphs Combin.* 39 (2023) 25.
- [25] P. Turán, Eine Extremalaufgabe aus der Graphentheorie, (*Hungarian*) *Mat. Fiz. Lapok* 48 (1941) 436-452.
- [26] D.B. West, *Introduction to graph theory*. Prentice Hall, Inc., Upper Saddle River, NJ, 1996.
- [27] H. Wu, On the 0-1 matrices whose squares are 0-1 matrices, *Linear Algebra Appl.* 432 (2010) 2909-2924.
- [28] W. Zhou, B. Li, The Turán number of directed paths and oriented cycles. *Graph Combin.* 39 (2023) 47.